

Transformations of Minimal Surfaces.

(By LUTHER PFAHLER EISENHART.)

In a Memoir published by CALAPSO in the *Annali* (*), the author has developed at length the theory and transformations of a class of surfaces, discovered by GUICHARD (**), and characterized by the property:

Given a surface N of the class; « there exists a surface N' having the same spherical representation of its lines of curvature as the surface N , and such that if r_1, r_2 are the principal radii of curvature of N , and r'_1, r'_2 the corresponding radii of N' ; one has

$$r_1 r'_2 + r_2 r'_1 = \text{const.}, \quad (\text{I})$$

the constant being different from zero ».

CALAPSO has shown that, if a surface be referred to its lines of curvature, the necessary and sufficient condition that it be a surface N is that either of the following conditions be satisfied:

$$\left(\sqrt{G} \frac{D}{\sqrt{E}} - \sqrt{E} \frac{D''}{\sqrt{G}} \right)^2 = G - E, \quad (\text{II})$$

$$\left(\sqrt{G} \frac{D}{\sqrt{E}} - \sqrt{E} \frac{D''}{\sqrt{G}} \right)^2 = G + E, \quad (\text{III})$$

where the functions E, G, D, D'' are the fundamental quantities of the surface. According as a surface satisfies the first or second of these conditions, it is called by CALAPSO a *surface of GUICHARD of the first kind* or of the *second kind*.

In the present Note we show that one of the surfaces parallel to a min-

(*) *Alcune superficie di GUICHARD e le relative trasformazioni* [Annali, ser. III, vol. XI, p. 201-251, (1905)].

(**) *Sur les Surfaces isothermiques* [Comptes Rendus, vol. 130, p. 159].

imal surface satisfies (II) and that the surface N' associated with it after the manner of the above theorem is a sphere. After deducing these results in § 1, we apply in the subsequent §§ the results of GUICHARD and CALAPSO to this group of surfaces of GUICHARD. We shall refer to these particular surfaces as *surfaces P*.

GUICHARD has shown that from a surface of the first kind one can deduce an infinity of isothermic surfaces referred to their lines of curvature each of which is the locus of a point A situated on an isotropic tangent to the surface N . In § 2 it is found that these transforms of a surface P are minimal surfaces; hence we have a transformation from the minimal surface, S , which is parallel to P , to new minimal surfaces, which for convenience we shall call the surfaces \overline{S} .

In § 3 we apply to the P -surfaces parallel to the surfaces \overline{S} (call them the surfaces \overline{P}) the GUICHARD transformations and find that the determination of all these transformations requires only quadratures. However, these new minimal surfaces are imaginary.

In § 4 we apply to the surfaces \overline{P} the transformation of GUICHARD in which $+i$ has been replaced by $-i$ and vice-versa. Among the infinity of transforms of a surface \overline{P} there is always one real minimal surface other than the original surface S .

In this manner we can obtain from a given minimal surface an infinity of real minimal surfaces, whose determination requires the solution of a pair of RICCATI equations. The relation between the original surface and any one of these transforms is perfectly reciprocal, so that when the transforms S_1 of S have been found, one can get the transforms S'_1 of the surfaces S_1 by quadratures.

CALAPSO has shown (*) that, given any isothermic surface, a surface of GUICHARD can be found by inverting the transformation of GUICHARD. In § 5 we apply this inverted transformation and also its conjugate to the surfaces \overline{S} , and in both cases we find that all the transforms are imaginary.

(*) L. c., pag. 230.

§ 1. SURFACES P .

Consider the minimal surface S with the linear element

$$ds^2 = e^{2\theta} (du^2 + dv^2), \tag{1}$$

and for which the coefficients of the second quadratic form are

$$D = -1, \quad D'' = 1. \tag{2}$$

Now the linear element of the spherical representation is

$$ds'^2 = e^{-2\theta} (du^2 + dv^2), \tag{3}$$

and the GAUSS and CODAZZI equations are satisfied, if θ is a solution of the equation

$$\frac{\partial^2 \theta}{\partial u^2} + \frac{\partial^2 \theta}{\partial v^2} = e^{-2\theta}. \tag{4}$$

If we denote by $X_1, Y_1, Z_1; X_2, Y_2, Z_2; X, Y, Z$ the direction-cosines of the tangents to the curves $v = \text{const.}$, $u = \text{const.}$ and of the normal to the surface respectively, we have (*)

$$\left. \begin{aligned} \frac{\partial x}{\partial u} = e^\theta X_1, \quad \frac{\partial X_1}{\partial u} = -\frac{\partial \theta}{\partial v} X_2 - e^{-\theta} X, \quad \frac{\partial X_2}{\partial u} = \frac{\partial \theta}{\partial v} X_1, \quad \frac{\partial X}{\partial u} = e^{-\theta} X_1, \\ \frac{\partial x}{\partial v} = e^\theta X_2, \quad \frac{\partial X_1}{\partial v} = \frac{\partial \theta}{\partial u} X_2, \quad \frac{\partial X_2}{\partial v} = -\frac{\partial \theta}{\partial u} X_1 + e^{-\theta} X, \quad \frac{\partial X}{\partial v} = -e^{-\theta} X_2, \end{aligned} \right\} \tag{5}$$

and similar equations in the y 's and z 's.

A surface parallel to S at the distance a is given by

$$\xi = x - a X, \quad \eta = y - a Y, \quad \zeta = z - a Z.$$

The fundamental quantities of this surface are found to be

$$\begin{aligned} E &= (e^\theta - a e^{-\theta})^2, & F &= 0, & G &= (e^\theta + a e^{-\theta})^2, \\ D &= -e^{-\theta} (e^\theta - a e^{-\theta}), & D' &= 0, & D'' &= e^{-\theta} (e^\theta + a e^{-\theta}). \end{aligned}$$

(*) BIANCHI, *Lezioni*, vol. II, p. 336.

These expressions satisfy equation (II), if $a = 1$ and only in this case. Hence the surface defined by

$$\xi = x - X, \quad \eta = y - Y, \quad \zeta = z - Z, \quad (6)$$

is a surface of GUICHARD of the first kind; we designate such a surface by P . From above it is seen that the fundamental quantities for P are

$$E = 4 \sinh^2 \theta, \quad G = 4 \cosh^2 \theta, \quad D = -2 e^{-\theta} \sinh \theta, \quad D'' = 2 e^{-\theta} \cosh \theta. \quad (7)$$

CALAPSO has shown (*) that the necessary and sufficient condition that a surface be a surface of GUICHARD is that the linear elements of the surface and its spherical representation be reducible to the respective forms

$$\left. \begin{aligned} d\sigma^2 &= e^{2t} (\sinh^2 \Theta du^2 + \cosh^2 \Theta dv^2) \\ d\sigma'^2 &= (\cosh \Theta + H \sinh \Theta)^2 du^2 + (\sinh \Theta + H \cosh \Theta)^2 dv^2, \end{aligned} \right\} \quad (8)$$

the lines of curvature being parametric, and the functions t , H and Θ being solutions of the system of equations

$$\left. \begin{aligned} \frac{\partial H}{\partial u} &= (H + \coth \Theta) \frac{\partial t}{\partial u}, \quad \frac{\partial H}{\partial v} = (H + \tanh \Theta) \frac{\partial t}{\partial v}, \\ \frac{\partial^2 \Theta}{\partial u^2} + \frac{\partial^2 \Theta}{\partial v^2} + \coth \Theta \frac{\partial^2 t}{\partial u^2} + \tanh \Theta \frac{\partial^2 t}{\partial v^2} - \frac{1}{\sinh^2 \Theta} \frac{\partial \Theta}{\partial u} \frac{\partial t}{\partial u} + \\ &+ \frac{1}{\cosh^2 \Theta} \frac{\partial \Theta}{\partial v} \frac{\partial t}{\partial v} + (\cosh \Theta + H \sinh \Theta) (\sinh \Theta + H \cosh \Theta) = 0. \end{aligned} \right\} \quad (9)$$

In order that (3) may take the form of the second of (8), we must have

$$\cosh \Theta + H \sinh \Theta = e^{-\theta}, \quad \sinh \Theta + H \cosh \Theta = -e^{-\theta},$$

from which it follows that

$$\Theta = \theta, \quad H = -1. \quad (10)$$

Moreover, the first of (8) reduces to (1), if

$$e^t = 2. \quad (11)$$

In consequence of (4) equations (9) are satisfied by the above values (10), (11).

(*) L. c., p. 214.

CALAPSO has shown (*) that the surface N' , which is related to N in the manner stated in the theorem leading to the relation (I), is defined by the forms (8), where now the functions t , H , Θ relating to N' are given by

$$e^t = e^{-t} (1 - H^2), \quad \sinh \Theta = \frac{-1}{1 - H^2} \left[\sinh \Theta (1 + H^2) + 2H \cosh \Theta \right],$$

$$\cosh \Theta = \frac{1}{1 - H^2} \left[\cosh \Theta (1 + H^2) + 2H \sinh \Theta \right].$$

When the surface N is a surface P , the surface N' is the unit sphere upon which the Gaussian representation of P is made.

§ 2. THE TRANSFORMATION OF GUICHARD.

GUICHARD (**) has announced the following theorem:

Given a surface satisfying conditions (8) and (9), a function φ can be determined in such a way that the surface \bar{S} defined by

$$\bar{x} = \xi + e^{\varphi} (X_1 + i X_2), \quad \bar{y} = \eta + e^{\varphi} (Y_1 + i Y_2), \quad \bar{z} = \zeta + e^{\varphi} (Z_1 + i Z_2) \quad (12)$$

is isothermic.

CALAPSO shows (***) that the function φ is determined by the following pair of illimitably integrable equations:

$$\frac{\partial \varphi}{\partial u} + i \frac{\partial \Theta}{\partial v} = -\sinh \Theta \cosh (\varphi - t) -$$

$$- i \tanh \Theta \frac{\partial t}{\partial v} - \frac{1}{2} e^{\varphi-t} (H^2 \sinh \Theta + 2H \cosh \Theta),$$

$$i \frac{\partial \varphi}{\partial v} + \frac{\partial \Theta}{\partial u} = \cosh \Theta \sinh (\varphi - t) -$$

$$- \coth \Theta \frac{\partial t}{\partial u} + \frac{1}{2} e^{\varphi-t} (H^2 \cosh \Theta + 2H \sinh \Theta).$$

(*) L. c., p. 214.

(**) L. c., p. 161.

(***) L. c., p. 224.

When the surface of GUICHARD is a surface P , these reduce to

$$\left. \begin{aligned} \frac{\partial \varphi}{\partial u} + i \frac{\partial \theta}{\partial v} &= \frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \sinh \theta, \\ i \frac{\partial \varphi}{\partial v} + \frac{\partial \theta}{\partial u} &= \frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \cosh \theta. \end{aligned} \right\} \quad (13)$$

In consequence of (5) and (6) we get from (12) by differentiation

$$\left. \begin{aligned} \frac{\partial \bar{x}}{\partial u} &= 2 \sinh \theta X_1 + e^{\varphi} (X_1 + i X_2) \left(\frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \sinh \theta \right) - e^{\varphi-\theta} X, \\ \frac{\partial \bar{x}}{\partial v} &= 2 \cosh \theta X_2 - i e^{\varphi} (X_1 + i X_2) \left(\frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \cosh \theta \right) + i e^{\varphi-\theta} X, \end{aligned} \right\} \quad (14)$$

so that

$$\bar{E} = \Sigma \left(\frac{\partial \bar{x}}{\partial u} \right)^2 = e^{2\varphi}, \quad \bar{F} = \Sigma \frac{\partial \bar{x}}{\partial u} \frac{\partial \bar{x}}{\partial v} = 0, \quad \bar{G} = \Sigma \left(\frac{\partial \bar{x}}{\partial v} \right)^2 = e^{2\varphi}. \quad (15)$$

From (14) we find that the direction-cosines, \bar{X} , \bar{Y} , \bar{Z} , of the normal to the surface \bar{S} are of the form

$$\bar{X} = e^{-\varphi} (X_1 - i X_2) + X. \quad (16)$$

By differentiation we get

$$\frac{\partial \bar{X}}{\partial u} = e^{-2\varphi} \frac{\partial \bar{x}}{\partial u}, \quad \frac{\partial \bar{X}}{\partial v} = -e^{-2\varphi} \frac{\partial \bar{x}}{\partial v}, \quad (17)$$

so that

$$\bar{D} = -\Sigma \frac{\partial \bar{x}}{\partial u} \frac{\partial \bar{X}}{\partial u} = -1, \quad \bar{D}' = 0, \quad \bar{D}'' = 1. \quad (18)$$

Hence \bar{S} is a minimal surface, and the linear element of its spherical representation is

$$d\bar{s}^2 = e^{-2\varphi} (du^2 + dv^2). \quad (19)$$

Denoting by \bar{X}_1 , \bar{Y}_1 , \bar{Z}_1 ; \bar{X}_2 , \bar{Y}_2 , \bar{Z}_2 , the direction-cosines of the tangents to the curves $v = \text{const.}$, $u = \text{const.}$ respectively on \bar{S} , we have

$$\bar{X}_1 = e^{-\varphi} \frac{\partial \bar{x}}{\partial u}, \quad \bar{X}_2 = e^{-\varphi} \frac{\partial \bar{x}}{\partial v},$$

which in consequence of (14) reduce to

$$\left. \begin{aligned} \bar{X}_1 &= \left(\frac{1}{2} e^{\varphi-\theta} + e^{-\varphi} \sinh \theta \right) X_1 + \left(\frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \sinh \theta \right) i X_2 - e^{-\theta} X, \\ i \bar{X}_2 &= \left(\frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \cosh \theta \right) X_1 + \left(\frac{1}{2} e^{\varphi-\theta} + e^{-\varphi} \cosh \theta \right) i X_2 - e^{-\theta} X. \end{aligned} \right\} \quad (20)$$

If e be replaced by ρ , equations (13) take the RICCATI form. Hence there exists a family of imaginary minimal surfaces depending upon a parameter, transforms of P and consequently of S ; and the complete determination of these surfaces requires the integration of a pair of RICCATI equations.

§ 3. GUICHARD TRANSFORMATIONS OF \bar{S} .

Since \bar{S} is a minimal surface, we have a new surface \bar{P} parallel to \bar{S} . From (6), (12) and (16) it follows that this surface is defined by equations of the form

$$\bar{\xi} = \xi + e^{\varphi} (X_1 + i X_2) - e^{-\varphi} (X_1 - i X_2) - X. \quad (21)$$

We apply to \bar{P} a transformation of GUICHARD. From (12) it follows that the transform S_1 is defined by equations of the form

$$x_1 = \bar{\xi} + e^{\varphi_1} (\bar{X}_1 + i \bar{X}_2), \quad (22)$$

where φ_1 is a solution of the equations

$$\frac{\partial \varphi_1}{\partial u} + i \frac{\partial \varphi}{\partial v} = \frac{1}{2} e^{\varphi_1-\varphi} - e^{-\varphi_1} \sinh \varphi, \quad i \frac{\partial \varphi_1}{\partial v} + \frac{\partial \varphi}{\partial u} = \frac{1}{2} e^{\varphi_1-\varphi} - e^{-\varphi_1} \cosh \varphi. \quad (23)$$

In consequence of (20) and (21) equation (22) is reducible to

$$x_1 = \xi + (1 + e^{\varphi_1-\theta}) [e^{\varphi} (X_1 + i X_2) - e^{-\varphi} (X_1 - i X_2)] - (2e^{\varphi_1-\theta} + 1) X. \quad (24)$$

On replacing $\xi + X$ by x in this equation, we get a transformation involving two parameters, which changes the original minimal surface S into another minimal surface S_1 .

If equations (23) be written in the form

$$\begin{aligned}\frac{\partial \varphi}{\partial u} + i \frac{\partial \varphi_1}{\partial v} &= -\frac{1}{2} e^{\varphi - \varphi_1} + \frac{1}{2} e^{-\varphi} \sinh \varphi_1, \\ i \frac{\partial \varphi}{\partial v} + \frac{\partial \varphi_1}{\partial u} &= -\frac{1}{2} e^{\varphi - \varphi_1} + e^{-\varphi} \cosh \varphi_1,\end{aligned}$$

and be compared with (13), it is seen that a solution of them is given by

$$e^{\varphi_1} = -e^{\theta}. \quad (25)$$

Hence, since equations (23) are of the RICCATI type, they are integrated completely by quadratures. Therefore, the complete determination of the doubly-infinite group of surfaces S_1 defined by (24), requires the solution of a pair of RICCATI equations and quadratures.

Evidently the transformation determined by (25) leads to S itself. We inquire whether any of the other surfaces S_1 are real, when S is real.

If we put

$$\varphi = \alpha + i\beta,$$

where α and β are real, equations (13) are replaced by

$$\left. \begin{aligned}\frac{\partial \alpha}{\partial u} &= \left(\frac{1}{2} e^{\alpha - \theta} - e^{-\alpha} \sinh \theta \right) \cos \beta, & \frac{\partial \alpha}{\partial v} &= \left(\frac{1}{2} e^{\alpha - \theta} + e^{-\alpha} \cosh \theta \right) \sin \beta, \\ \frac{\partial \beta}{\partial u} + \frac{\partial \theta}{\partial v} &= \left(\frac{1}{2} e^{\alpha - \theta} + e^{-\alpha} \sinh \theta \right) \sin \beta, \\ \frac{\partial \beta}{\partial v} - \frac{\partial \theta}{\partial u} &= - \left(\frac{1}{2} e^{\alpha - \theta} - e^{-\alpha} \cosh \theta \right) \cos \beta.\end{aligned} \right\} \quad (26)$$

In like manner, if φ_1 is real, equations (23) can be written

$$\begin{aligned}\frac{\partial \alpha}{\partial u} &= \left(\frac{1}{2} e^{\varphi_1 - \alpha} - e^{-\varphi_1} \cosh \alpha \right) \cos \beta, & \frac{\partial \alpha}{\partial v} &= - \left(\frac{1}{2} e^{\varphi_1 - \alpha} + e^{-\varphi_1} \cosh \alpha \right) \sin \beta, \\ \frac{\partial \varphi_1}{\partial u} - \frac{\partial \beta}{\partial v} &= \left(\frac{1}{2} e^{\varphi_1 - \alpha} - e^{-\varphi_1} \sinh \alpha \right) \cos \beta, \\ \frac{\partial \varphi_1}{\partial v} + \frac{\partial \beta}{\partial u} &= - \left(\frac{1}{2} e^{\varphi_1 - \alpha} + e^{-\varphi_1} \sinh \alpha \right) \sin \beta.\end{aligned}$$

In order that the first two equations of these two sets be consistent,

we must have

$$\frac{1}{2} e^{\varphi_1 - \alpha} - e^{-\varphi_1} \cosh \alpha - \frac{1}{2} e^{\alpha - \theta} + e^{-\alpha} \sinh \theta = 0,$$

$$\frac{1}{2} e^{\varphi_1 - \alpha} + e^{-\varphi_1} \cosh \alpha + \frac{1}{2} e^{\alpha - \theta} + e^{-\alpha} \cosh \theta = 0,$$

from which by addition we get (25). Hence of all the surfaces defined by (24) S is the only real one.

§ 4. CONJUGATE GUICHARD TRANSFORMATIONS OF \overline{S} .

We apply now to the surface \overline{P} , defined by (21), the conjugate transformation of GUICHARD, that is, the transformation with $+i$ replaced by $-i$.

Now the equations analogous to (23) are

$$\frac{\partial \theta_1}{\partial u} - i \frac{\partial \varphi}{\partial v} = \frac{1}{2} e^{\theta_1 - \varphi} - e^{-\theta_1} \sinh \varphi, \quad i \frac{\partial \theta_1}{\partial v} - \frac{\partial \varphi}{\partial u} = -\frac{1}{2} e^{\theta_1 - \varphi} + e^{-\theta_1} \cosh \varphi, \quad (27)$$

and the coordinates of the transform S_1 are of the form

$$x_1 = \overline{\xi} + e^{\theta_1} (\overline{X}_1 - i \overline{X}_2),$$

which reduces to

$$x_1 = \xi + e^{\varphi} (X_1 + i X_2) + (e^{\theta_1 + \theta} - 1) e^{-\varphi} (X_1 - i X_2) - X. \quad (28)$$

As in the preceding case, if $\xi + X$ be replaced by x , this gives a transformation from one minimal surface, S , to a double-infinite family of surfaces, S_1 . Later we shall find a particular solution of equations (27); hence the complete determination of all the above transforms of a surface S requires the solution of an equation of RICCATI and quadratures.

We inquire whether any of these transforms are real. On the assumption that θ_1 is real equations (27) can be written

$$\frac{\partial \alpha}{\partial u} = \left(\frac{1}{2} e^{\theta_1 - \alpha} - e^{-\theta_1} \cosh \alpha \right) \cos \beta, \quad \frac{\partial \alpha}{\partial v} = \left(\frac{1}{2} e^{\theta_1 - \alpha} + e^{-\theta_1} \cosh \alpha \right) \sin \beta, \quad (29)$$

$$\left. \begin{aligned} \frac{\partial \theta_1}{\partial u} + \frac{\partial \beta}{\partial v} &= \left(\frac{1}{2} e^{\theta_1 - \alpha} - e^{-\theta_1} \sinh \alpha \right) \cos \beta, \\ \frac{\partial \theta_1}{\partial v} - \frac{\partial \beta}{\partial u} &= \left(\frac{1}{2} e^{\theta_1 - \alpha} + e^{-\theta_1} \sinh \alpha \right) \sin \beta. \end{aligned} \right\} \quad (30)$$

In order that equations (29) be consistent with the first two of (26), we must have

$$\begin{aligned} \frac{1}{2} e^{\theta_1 - \alpha} - e^{-\theta_1} \cosh \alpha - \frac{1}{2} e^{\alpha - \theta} + e^{-\alpha} \sinh \theta &= 0, \\ \frac{1}{2} e^{\theta_1 - \alpha} + e^{-\theta_1} \cosh \alpha - \frac{1}{2} e^{\alpha - \theta} - e^{-\alpha} \cosh \theta &= 0, \end{aligned}$$

which reduce to the single equation

$$e^{\theta_1} = 2 \cosh \alpha \cdot e^{\alpha - \theta}. \quad (31)$$

It is readily found that this value of θ_1 satisfies equations (30).

Substituting in (28) and replacing ξ by $x - X$ we get the equation

$$x_1 = x + 2 e^{\alpha} (\cos \beta X_1 - \sin \beta X_2) - 2 X, \quad (32)$$

and similar expressions for y_1 and z_1 which define a real transform, S_1 , of a given surface, S .

By differentiation we get

$$\left. \begin{aligned} \frac{\partial x_1}{\partial u} &= (e^{2\alpha - \theta} \cos 2\beta - e^{-\theta}) X_1 - e^{2\alpha - \theta} \sin 2\beta X_2 - 2 e^{\alpha - \theta} \cos \beta X, \\ \frac{\partial x_1}{\partial v} &= e^{2\alpha - \theta} \sin 2\beta X_1 + (e^{2\alpha - \theta} \cos 2\beta + e^{-\theta}) X_2 - 2 e^{\alpha - \theta} \sin \beta X. \end{aligned} \right\} \quad (33)$$

From these and similar expressions in y_1 and z_1 we derive the following values for the direction-cosines of the normal to S_1

$$X' = \tanh \alpha X + \frac{1}{\cosh \alpha} (\cos \beta X_1 - \sin \beta X_2). \quad (34)$$

And the direction-cosines of the tangents to the curves $v = \text{const.}$, $u = \text{const.}$ on S_1 are of the respective forms

$$\left. \begin{aligned} X'_1 &= \frac{1}{2 \cosh \alpha} \left[(e^{\alpha} \cos 2\beta - e^{-\alpha}) X_1 - e^{\alpha} \sin 2\beta X_2 - 2 \cos \beta X \right], \\ X'_2 &= \frac{1}{2 \cosh \alpha} \left[e^{\alpha} \sin 2\beta X_1 + (e^{\alpha} \cos 2\beta + e^{-\alpha}) X_2 - 2 \sin \beta X \right]. \end{aligned} \right\} \quad (35)$$

From (32) and (34) we get

$$\Sigma X(x_1 - x) = -2, \quad \Sigma X'(x_1 - x) = 2, \quad (36)$$

and

$$\Sigma (x_1 - x)^2 = 4(e^{2\alpha} + 1).$$

Hence the line joining corresponding points on two surfaces S and S_1 makes equal angles with the normals to the surfaces at these points. As this angle is not a right angle, the transformation is different from the one due to THYBAUT (*).

From (34) it follows that the direction-cosines of the line of intersection of the tangent planes to S and S_1 at corresponding points are

$$\sin \beta X_1 + \cos \beta X_2, \quad \sin \beta Y_1 + \cos \beta Y_2, \quad \sin \beta Z_1 + \cos \beta Z_2.$$

Hence this line is perpendicular to the projection upon the tangent plane to S of the line joining the points of tangency on S and S_1 .

Denote by ξ , η , ζ the coordinates of the point of intersection of these two lines. Since we must have

$$\Sigma X(\xi - x) = 0, \quad \Sigma X'(\xi - x) = 0,$$

if we put

$$\xi = x + \lambda (\cos \beta X_1 - \sin \beta X_2),$$

it is found that

$$\lambda = 2 \cosh \alpha.$$

From the foregoing discussion it follows that each pair of solutions of equations (26) gives a real transform of S . The form of these equations is such that the arbitrary constant entering in the complete solution appears in both α and β . From (32) it is seen that the points, on all the transforms, corresponding to a point on S lie in the plane parallel to the tangent plane to S and at the distance 2 from it.

We inquire whether the normals to the surfaces S and S_1 at corresponding points meet. In order that this may happen two functions λ and μ must exist which are such that

$$x_0 = x + \lambda X = x_1 + \mu X'.$$

and similar equations in y 's and z 's.

(*) BIANCHI, *Lezioni*, vol. II, p. 334.

Annali di Matematica, Serie III, Tomo XIII.

If the above values be substituted, it is found that these equations are satisfied when

$$\lambda = \mu = -2e^\alpha \cosh \alpha.$$

This result suggests that this transformation which we have found coincides with one previously discovered by BIANCHI (*) as an outcome of a theorem of GUICHARD. It is readily found that the two transformations are the same (**). From the investigation of BIANCHI we know that the locus of the point (x_0, y_0, z_0) is a surface applicable to a paraboloid of revolution.

§ 5. INVERSE TRANSFORMATIONS OF GUICHARD.

CALAPSO has shown (***) that the transformation of GUICHARD can be inverted, that is, every isothermic surface can be considered as derived from a GUICHARD surface by means of the construction indicated in the theorem of GUICHARD. We shall apply this inverse transformation to the surfaces \bar{S} .

In the first place we solve equations (16) and (20) for X, X_1, X_2 , getting

$$\left. \begin{aligned} X_1 &= \left(\frac{1}{2} e^{\varphi-\theta} + e^{-\varphi} \sinh \theta \right) \bar{X}_1 - \left(\frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \cosh \theta \right) i \bar{X}_2 + e^{-\varphi} \bar{X}, \\ i X_2 &= - \left(\frac{1}{2} e^{\varphi-\theta} - e^{-\varphi} \sinh \theta \right) \bar{X}_1 + \left(\frac{1}{2} e^{\varphi-\theta} + e^{-\varphi} \cosh \theta \right) i \bar{X}_2 + e^{-\varphi} \bar{X}, \\ X &= -e^{-\theta} (\bar{X}_1 - i \bar{X}_2) + \bar{X}. \end{aligned} \right\} \quad (37)$$

In consequence of these relations equation (12) can be written

$$\bar{x} = \xi + 2 (\sinh \theta \bar{X}_1 + \cosh \theta \cdot i \bar{X}_2 + \bar{X}). \quad (38)$$

If \bar{S} is any minimal surface and the linear element of its representation is written in the form (19), the function φ must satisfy equation (4). But

(*) *Lezioni*, vol. II, p. 333.

(**) The formulae of comparison are

$$\frac{\Phi}{W} = 2e^\alpha \cosh \alpha, \quad \frac{\cos \beta}{\cosh \alpha} = -\frac{2e^{-\theta}}{\Phi} \frac{\partial \Phi}{\partial u}, \quad \frac{\sin \beta}{\cosh \alpha} = 2 \frac{e^{-\theta}}{\Phi} \frac{\partial \Phi}{\partial v}.$$

(***) L. c., p. 230.

this is the condition that equations (13) be consistent in θ . Hence (38) is the transformation from a surface \bar{S} to P as well as equations (12) define the transformation from P to \bar{S} . But since equations (13) are completely integrable in θ for a particular φ , it follows that each of the surfaces defined by

$$\xi_1 = \bar{x} - 2 (\sinh \theta_1 \bar{X}_1 + \cosh \theta_1 i \bar{X}_2 + \bar{X}), \quad (39)$$

where θ_1 is any solution of

$$\frac{\partial \varphi}{\partial u} + i \frac{\partial \theta_1}{\partial v} = \frac{1}{2} e^{\varphi - \theta_1} - e^{-\varphi} \sinh \theta_1, \quad i \frac{\partial \varphi}{\partial v} + \frac{\partial \theta_1}{\partial u} = \frac{1}{2} e^{\varphi - \theta_1} - e^{-\varphi} \cosh \theta_1, \quad (40)$$

is a surface of GUICHARD, P_1 , which is an inverse of the surface \bar{S} , determined by the function φ .

From (37) it is seen that the direction-cosines of the normal to P_1 are of the form

$$X' = -e^{-\theta_1} \bar{X}_1 + e^{-\theta_1} i \bar{X}_2 + \bar{X}. \quad (41)$$

Denoting by S_1 the minimal surface parallel to P_1 , we have

$$x_1 = \xi_1 + X', \quad y_1 = \eta_1 + Y', \quad z_1 = \zeta_1 + Z',$$

which by means of (39), (41), (12) and (20) reduce to

$$x_1 = x + (1 - e^{\theta_1 - \theta}) [e^{\varphi} (X_1 + i X_2) - e^{-\varphi} (X_1 - i X_2) - 2 X], \quad (42)$$

and similar expressions for y_1 and z_1 . Since θ is a solution of (40), the complete determination of the doubly-infinite family of transforms of S defined by (42) requires the solution of one pair of RICCATI equations (13) and quadratures.

If the differential quotients of φ be eliminated from equations (13) and (40), we get

$$\begin{aligned} \frac{\partial \theta_1}{\partial u} - \frac{\partial \theta}{\partial u} &= \left[\frac{1}{2} e^{\alpha} (e^{-\theta_1} - e^{-\theta}) - e^{-\alpha} (\cosh \theta_1 - \cosh \theta) \right] \cos \beta + \\ &\quad + i \left[\frac{1}{2} e^{\alpha} (e^{-\theta_1} - e^{-\theta}) + e^{-\alpha} (\cosh \theta_1 - \cosh \theta) \right] \sin \beta, \\ i \left(\frac{\partial \theta_1}{\partial v} - \frac{\partial \theta}{\partial v} \right) &= \left[\frac{1}{2} e^{\alpha} (e^{-\theta_1} - e^{-\theta}) - e^{-\alpha} (\sinh \theta_1 - \sinh \theta) \right] \sin \beta + \\ &\quad + i \left[\frac{1}{2} e^{\alpha} (e^{-\theta_1} - e^{-\theta}) + e^{-\alpha} (\sinh \theta_1 - \sinh \theta) \right] \sin \beta. \end{aligned}$$

From this it follows that for θ_1 and θ to be real we must have

$$\begin{aligned} \frac{1}{2} e^\alpha (e^{-\theta_1} - e^{-\theta}) + e^{-\alpha} (\cosh \theta_1 - \cosh \theta) &= 0, \\ \frac{1}{2} e^\alpha (e^{-\theta_1} - e^{-\theta}) - e^{-\alpha} (\sinh \theta_1 - \sinh \theta) &= 0, \end{aligned}$$

which reduces to $\theta_1 = \theta$. Hence all the minimal surface transforms (42) of S are imaginary.

We pass finally to the case in which we apply to the surfaces \bar{S} the conjugate inverse transformation. In place of equations (40) we have

$$\frac{\partial \varphi}{\partial u} - i \frac{\partial \theta_1}{\partial v} = \frac{1}{2} e^{\gamma-\theta_1} - e^{-\gamma} \sinh \theta_1, \quad i \frac{\partial \varphi}{\partial v} - \frac{\partial \theta_1}{\partial u} = -\frac{1}{2} e^{\gamma-\theta_1} + e^{-\gamma} \cosh \theta_1,$$

and the equations of transformation are of the form

$$x_1 = x + e^\gamma (X_1 + i X_2) - e^{-\gamma} (e^{\theta_1+\theta} + 1) (X_1 - i X_2) - 2X.$$

Proceeding as in the former case, we find that the condition that θ_1 be real is

$$e^{\theta_1} = -e^{\alpha-\theta} (e^\alpha + e^{-\alpha}),$$

which evidently is impossible.