

done. In the notes furnished by Dr. DaCosta it was stated that a varicose condition of the vessels of the dura was found, and these were ligated. At the same time Dr. Gordon applied a faradic needle to various portions of the cortex. The man made an uninterrupted recovery and the convulsions ceased for about a year. Since then they have returned and are more severe than ever and in addition there is hemiparesis. A second operation was done over the same site and a large sarcomatous growth was found. After the exposure large veins were in evidence which were found to be only part of the tumor. This case is similar to that reported by Dr. Gordon and shows that a varicose condition of the vessels may be only an indication of a graver condition, and it should not be taken for granted that this is present unless a full exposure is made.

It was also, in Dr. Weisenburg's opinion, a mistake to assert that any operation will cause cessation of epileptic convulsions unless the case is observed a number of years, for as in the first instance referred to, the case was reported as apparently cured.

Dr. Gordon said he recalled the case Dr. Weisenburg alluded to distinctly. There was an angiomatous condition of the blood vessels of the brain. The patient presented in addition to the Jacksonian epilepsy other symptoms. He also had vomiting spells and dizziness. The skull was simply opened in the area supposed to have the lesion. The angiomatous condition was found. Dr. Da Costa thought it not wise to examine the brain tissue beneath and simply ligated. Dr. Gordon's patient was perfectly free from any trouble other than the localized convulsions.

A CASE OF CEREBROSPINAL SYPHILIS CAUSING INTERNAL HYDROCEPHALUS AND SYMPTOMS OF CEREBELLAR TUMOR

By S. D. Ingham, M.D.

C. R., white, 42 years; family history negative. Had a chancre at age of 24 years, but gave no history suggesting secondary or tertiary syphilis. Complained for two or three years of pains in arms and legs, unsteadiness in gait, difficulty with micturition, and deafness. His sister stated that he had been mentally dull for five or six years, and had had occasional attacks of headache with nausea and vomiting, vertigo, and sometimes fainting. At the time of his admission to the Philadelphia Hospital his condition was as follows: Muscularly well developed, no motor palsies, no muscular atrophies. Pupils show some inequality and irregularity of outline, reactions present, no choked disc. Tendon reflexes were slightly increased in all the limbs, bilateral ankle clonus, Babinski negative. Slight ataxia in both arms and legs.

The most striking symptom of his condition was the progressive loss of equilibration, at first manifested by inability to walk erect, but leaning backward and quickly coming to a stop. He soon became unable to walk at all, and before his death, about four weeks after admission, could not even sit up in bed, but would invariably fall directly backward, not from weakness, but from loss of equilibrium. The pathological condition found at necropsy was syphilitic cerebrospinal meningitis, especially marked at the base of the brain. The foramen of Magendie was completely occluded by the process, and all of the ventricles of the brain were dilated, especially the fourth, where the pressure had displaced the dentate nuclei and

white matter of the cerebellum. Microscopically, the usual signs of syphilis were found, and, in addition, degeneration of cells in the dentate nuclei, and degeneration and disappearance of many of the Purkinje cells of the cerebellar cortex.

Dr. C. K. Mills stated that Dr. Spiller and he had a case which was reported by them, which presented symptoms very similar to those given by Dr. Ingham. The foramen of Magendie was not closed, as he recalled, but the aqueduct of Sylvius was closed producing symptoms very similar to those in this case.

Dr. T. H. Weisenburg said that aside from the pathological findings this case had been very interesting from the clinical standpoint. The patient had been studied by him a number of months and the most striking feature in the case was the fact that the patient always staggered backward, even when he attempted to sit up. Dr. Weisenburg said that he had never seen in any cerebellar lesion staggering only in the backward direction. This he considered unusual, for while he had seen in other cases backward staggering, it was always in association with a tendency to fall in other directions.

Dr. Spiller stated that he had seen the backward staggering without lateral staggering. In one case in particular, a tumor was found beneath the aqueduct of Sylvius, and the patient had a distinct tendency to stagger backward. When he made an effort to stand up, as in rising from a chair, he would fall backward.

In regard to the case referred to by Dr. Mills the fourth ventricle was not enlarged. The symptoms were caused by pressure on the cerebellum by the dilated lateral ventricles, notwithstanding the tentorium came between.

Dr. Gordon said he had a case where the patient had a tendency to stagger to one side as well as backward. In regard to the cerebellar symptom, called *asynergia*, this is very common. When the patient makes the attempt to move forward, the trunk does not follow the legs and gives the impression of falling backwards. He supposed this had some relation to the remarks made by Dr. Ingham and Dr. Spiller.

A CASE OF SPASTIC PARAPLEGIA DATING FROM CHILDHOOD (LITTLE'S DISEASE?) WITH LITTLE OR NO DEMONSTRABLE LESION OF THE PYRAMIDAL TRACTS

By John H. W. Rhein, M.D.

The patient, a man of 71, presented symptoms of spastic paraplegia dating from early childhood. The arms were not involved. The brain and spinal cord showed nothing of importance, except a fineness of the fibers constituting the crossed pyramidal tracts. There was no evidence of degeneration.

Dr. Rhein asked, how can the spasticity be explained when the pyramidal tracts are not degenerated? The presence of fibers in the pyramidal tract presupposes the presence of intact cells in the cerebrum. The character of the fibers and cells then must have some bearing upon the causation of the spasticity. When fibers of small caliber are called upon to conduct vigorous motor impulses do we not meet with the same condition as when currents of electricity of high potentiality are forced through wires of small caliber, *i. e.*, increased resistance and imperfect