

2. CRANIAL NERVES.

Fuchs, E. UNSTRIPED MUSCLE FIBERS IN INTERNAL OCULAR MUSCLES. [Arch. f. Ophth., 95, 1918, 311.]

The existence of muscular fibers other than those leading to the ciliary muscle has been disputed. The author by means of frontal sections finds them to be present especially in the neighborhood of the papillæ due to attachments of the ocular tendons. They are plain and branched, and a figure of them is given. Whether they help in the maintenance of intra-ocular pressure is discussed. (Phys. Abst.)

Allee, W. C., and Stein, E. R. LIGHT REACTIONS IN MAY-FLY NYMPHS. [Jl. Exp. Zoöl., 26, 1918, No. 3, Phys. Abst.]

The light reaction of the May-fly nymph, *Epeorus*, was reversed by treatment with alcohol, lowered temperature, calcium chloride, and other reagents. Nymphs so reversed had a lower rate of metabolism as measured by resistance to potassium cyanide than normal nymphs. The negative nymph, *Leptophlebia*, was similarly reversed. A negative nymph belonging to the *Heptageniæ* was reversed in its light reactions with accompanying increase or decrease in carbon dioxide production as measured by Tashiro's biometer. The experiments demonstrate that the phototactic reaction is correlated with the metabolic condition, and indicate that changes in metabolism are causal. All nymphs that reversed their light reactions were either stimulated or depressed, but stimulation or depression did not necessarily involve phototactic reversal.

Gordon, Alfred. UNILATERAL PARALYSIS OF THE THIRD NERVE WITH HOMOLATERAL MOTOR AND SENSORY SYMPTOMS. [Phil. Neurological Soc., 1919.]

R. F., girl of 22, was injured by being run over by an automobile. When carried in an unconscious state to a nearby hospital, she was found to have a fracture of the skull. An operation was performed.

When seen by the writer several weeks later she presented besides mental phenomena (confusion, inability of recognizing her own people, extreme restlessness) the following symptoms.

There was a semicircular scar on the right side of the skull extending from the frontal region down back of the ear. Above the temporal region there is a large depression due to removal of a bony portion. The right third nerve was completely involved: superior, inferior and internal recti muscles—all paralyzed; the pupil enlarged and immobile, does not react to light; the upper eyelid in a state of complete ptosis, so that the eye was totally closed. She had difficulty of walking. Ataxia was observed in the right leg; she walked towards the right and her raising and lowering the right leg were very awkward. When in dorsal position the ataxia of the right leg was also marked, as in raising it the

leg would sway from side to side. When asked to touch with the right leg certain portions of the left ataxia was also observed. No paralysis was present. The right knee-jerk was greatly exaggerated. Ankle-clonus was present, but the toe-phenomenon by any of the methods was absent.

The sensations of the right leg were somewhat diminished. The right arm presented also ataxia, dysmetria and intention tremor. Sensations were but slightly diminished, astereognosis was absent.

The patient gradually improved. The mental condition has entirely disappeared. The ocular symptoms have greatly improved with the exception of the pupil which still does not react to light and is dilated.

The slight sensory disturbances of the right arm and leg have totally disappeared. Her gait is still somewhat ataxic in the right leg. The reflexes are the same. The dysmetria, ataxia and intention tremor of the right arm are still present although milder in intensity. Patient states, that frequently objects drop out of her hand and she spills liquids. Besides, her right arm and leg are always animated with jerky movements and sometimes with athetoid movements.

In every other respect the patient is normal. Wassermann test of blood and spinal fluid is negative. There are no changes in the eye-grounds. The mentality is normal.

Comment.—The absence of sensory-motor disturbances on the left side, the presence of cerebellar syndrome elements on the right, viz., ataxia, dysmetria, intention tremor; the absence of paralysis; the involvement of the third nerve,—all these conditions are in favor of a lesion in the right superior cerebellar peduncle below its decussation. There was probably a hemorrhage simultaneously occurring with the fracture of the skull.

The patient has made considerable progress under large doses of iodides with the exception of the pupil which remains dilated and immobile. The cerebellar symptoms have all improved. [Author's Abstract.]

Bourguet, J. GASSERIAN OPERATION FOR TIC DOULOUREUX. [Bull. Acad. de Méd., 81, May 13, 1919.]

This author after having remissions after all forms of alcoholic injections advises gasserectomy through a curved incision in the temporal bone. This exposes the maxillary nerves and leads directly to the ganglion. He detaches the dura and severs the middle meningeal. He then cuts the fibrous connections between the ganglion and the dura. Keratitis develops afterwards, but yields to treatment.

Roumaillac, E. NEURALGIA OF TRIGEMINUS. [Jour. Méd. de Bordeaux, 90, June 25, 1919.]

This patient, a woman 34 years of age, had had very severe trigeminal