

"Under side of anterior wings shining, the costa narrowly, and the outer margin broadly, reddish-yellow, especially at the apex; inner margin rather broadly pale yellow; all the rest uniformly deep black. Posterior wings shining, pale yellow, with the costa and apex broadly irrorate with bright ferruginous scales, and a narrow ill-defined submarginal band; fringes pale yellow.

"Antennæ pale ferruginous, with a slight pubescence. Head and thorax ferruginous. Abdomen cinereous above, darker from numerous black scales below; anal tuft ferruginous. Tarsi annulated.

"Alar expansion, 1.15 inch. Length of body, 0.50 inch.

"*Habitat*.—Nova Scotia. (No. 281, J. M. Jones.) This very pretty little moth may be readily distinguished by its colour and the conspicuous peculiarly-shaped orbicular spot.

"Halifax; rare, taken in July at Ashbourne.—J. M. J."

### LIST OF COLEOPTERA.

TAKEN AT GRIMSBY, ONTARIO, BY J. PETTIT.

(Continued from page 54.)

#### STAPHYLINIDÆ.

FALAGRIA, <i>Mann.</i>	*Rostratus, <i>Lec.</i>	PHILONTHUS, <i>Curtis.</i>
Dissecta, <i>Er.</i>	*Longiceps, <i>Lec.</i>	Cyanipennis, <i>Er.</i>
*Venustula, <i>Er.</i>	QUEDIUS, <i>Steph.</i>	Aeneus, <i>Nord.</i>
LEUCOPARYPHUS, <i>Kraats.</i>	Fulgidus, <i>Er.</i>	Blandus, <i>Er.</i>
*Silphoides, <i>Kr.</i>	*Lævigatus, <i>Er.</i>	Promptus, <i>Er.</i>
COPROPORUS, <i>Kr.</i>	*Capucinus, <i>Er.</i>	Debilis, <i>Er.</i>
Ventriculus, <i>Kr.</i>	Molochinus, <i>Er.</i>	Lomatus, <i>Er.</i>
TACHINUS, <i>Grav.</i>	CREOPHILUS, <i>Steph.</i>	*Fulvipes, <i>Nord.</i>
*Flavipennis, <i>Dej.</i>	Villosus, <i>Kirby.</i>	*Aterrimus, <i>Er.</i>
*Memnonius, <i>Grav.</i>	LEISTOTROPHUS, <i>Perty.</i>	Apicalis, <i>Er.</i>
Fumipennis, <i>Er.</i>	Cingulatus, <i>Kr.</i>	XANTHOLINUS, <i>Serv.</i>
Picipes, <i>Er.</i>	OCYPUS, <i>Kirby.</i>	Fulgidus, <i>Er.</i>
Fimbriatus, <i>Grav.</i>	Ater, <i>Er.</i>	Cephalus, <i>Say.</i>
Limbatus, <i>Mels.</i>	STAPHYLINUS, <i>Linn.</i>	Hamatus, <i>Say.</i>
TACYPORUS, <i>Grav.</i>	Maculosus, <i>Grav.</i>	BAPTOLINUS, <i>Kr.</i>
Jocosus, <i>Say.</i>	*Mysticus, <i>Er.</i>	*Filicornis, <i>Payk.</i>
CONOSOMA, <i>Kr.</i>	Vulpinus, <i>Nord.</i>	LATHROBIUM, <i>Grav.</i>
Crassum, <i>Lec.</i>	Fossator, <i>Grav.</i>	Grande, <i>Lec.</i>
Basale, <i>Lec.</i>	STAPHYLINUS ( <i>contin'd</i> ).	Punctulatum, <i>Lec.</i>
BOLETOBIUS, <i>Leach.</i>	Badipes, <i>Lec.</i>	Angulare, <i>Lec.</i>
*Niger, <i>Er.</i>	Cinnamopterus, <i>Grav.</i>	Concolor, <i>Lec.</i>
*Cincticollis, <i>Er.</i>	Violaceus, <i>Grav.</i>	*Armatus, <i>Say.</i>
Cinctus, <i>Er.</i>	Capitatus, <i>Bland.</i>	*Longiusculus, <i>Grav.</i>

*Collare, <i>Er.</i>	Juno, <i>Fab.</i>	ACIDOTA, <i>Steph.</i>
CRYPTOBIMUM, <i>Mann.</i>	OXYPORUS, <i>Fab.</i>	*Subcarinata, <i>Er.</i>
Bicolor, <i>Er.</i>	Rufipennis, <i>Lec.</i>	*Patruelis, <i>Lec.</i>
Pallipes, <i>Nord.</i>	*Femoralis, <i>Grav.</i>	OLOPHRUM, <i>Er.</i>
*Latebricola, <i>Nord.</i>	Lateralis, <i>Grav.</i>	*Emarginatum, <i>Er.</i>
STILICUS, <i>Latr.</i>	BLEDIUS, <i>Steph.</i>	TRIGONODEMUS, <i>Lec.</i>
*Angularis, <i>Er.</i>	Fumatus, <i>Lec.</i>	Striatus, <i>Lec.</i>
*Dentatus, <i>Er.</i>	PLATYSTETHUS, <i>Mann.</i>	ANTHOBIUM, <i>Steph.</i>
LITHOCHARIS, <i>Er.</i>	Americanus, <i>Er.</i>	Protectum, <i>Lec.</i>
Confluens, <i>Er.</i>	OXYTELUS, <i>Grav.</i>	PROTEINUS, <i>Latr.</i>
SUNIUS, <i>Steph.</i>	Sculptus, <i>Grav.</i>	*Parvulus, <i>Lec.</i>
*Prolixus, <i>Er.</i>	TROGOPHLEUS, <i>Mann.</i>	MEGARTHURUS, <i>Steph.</i>
*Linearis, <i>Er.</i>	*Morio, <i>Er.</i>	*Americanus, <i>Sachse.</i>
SUNIUS ( <i>continued</i> ).	ANTHOPHAGUS, <i>Grav.</i>	Angulicollis, <i>Mäkl.</i>
Longiusculus, <i>Er.</i>	*Cæsus, <i>Er.</i>	OLISTHÆRUS, <i>Er.</i>
PÆDERUS, <i>Grav.</i>	LESTEVA, <i>Latr.</i>	Nitidus, <i>Lec.</i>
Littorarius, <i>Grav.</i>	*Biguttula, <i>Lec.</i>	GLYPTOMA, <i>Er.</i>
STENUS, <i>Latr.</i>		*Costale, <i>Er.</i>
*Colon, <i>Say.</i>		
HISTERIDÆ.		
HOLOLEPTA, <i>Payk.</i>	Coarctatus, <i>Lec.</i>	*Deletus, <i>Lec.</i>
Fossularis, <i>Say.</i>	*Cylindricus, <i>Payk.</i>	TERETRIUS, <i>Er.</i>
HISTER, <i>Linn.</i>	*Gracilis, <i>Lec.</i>	*Americanus, <i>Lec.</i>
Interruptus, <i>Beauv.</i>	PHELISTER, <i>Mars.</i>	PLEGADERUS, <i>Er.</i>
Merdarius, <i>Payk.</i>	*Subrotundus, <i>Mars.</i>	Transversus, <i>Say.</i>
*Harrisii, <i>Kirby.</i>	TRIBALUS, <i>Er.</i>	ACRITUS, <i>Lec.</i>
Fœdatus, <i>Lec.</i>	*Americanus, <i>Lec.</i>	*Exiguus, <i>Lec.</i>
*Cognatus, <i>Lec.</i>	PAROMALUS, <i>Er.</i>	SCAPHIDIIDÆ.
Marginicollis, <i>Lec.</i>	——— ?	SCAPHIDIUM, <i>Oliv.</i>
Depurator, <i>Say.</i>	SAPRINUS, <i>Leach.</i>	4 Pustulatum, <i>Say.</i>
Abbreviatus, <i>Fab.</i>	Distinguendus, <i>Mars.</i>	Piceum, <i>Mels.</i>
*Sedecim striatus, <i>Say.</i>	*Pennsylvanicus, <i>Er.</i>	SCAPHISOMA, <i>Leach.</i>
Americanus, <i>Payk.</i>	*Conformis, <i>Lec.</i>	Convexum, <i>Say.</i>
*Carolinus, <i>Payk.</i>	Assimilis, <i>Er.</i>	TOXIDIUM, <i>Lec.</i>
LeContei, <i>Mars.</i>	*Scupularis, <i>Lec.</i>	*Gammaroides, <i>Lec.</i>
Parallelus, <i>Say.</i>	Fraternus, <i>Lec.</i>	

## MISCELLANEOUS NOTES.

LARVA OF *HELIOTHIS ARMIGERA*.—At a recent meeting of the Entomological Society of London, Eng., "Mr. J. Jenner Weir, exhibited two specimens of *Heliothis armigera*, Linn. (*H. umbrosus*, Grote), bred from larvæ

\*Species marked with an asterisk have not been included in the list of Canadian Coleoptera.