

**ANNALS OF THE UPPER SILESIAN MUSEUM IN BYTOM
ENTOMOLOGY**

Vol. 27 (online 010): 1–13

ISSN 0867-1966, eISSN 2544-039X (online)

Bytom, 24.12.2018

MARCIN WALCZAK^{1,4} , JOLANTA BROŽEK¹ , ŁUKASZ JUNKIERT¹, MAŁGORZATA KALANDYK-KOŁODZIEJCZYK¹ , KRZYSZTOF MUSIK², NATALIA KASZYCA¹, ANNA ŁAZUKA¹, GRZEGORZ GIERLASIŃSKI³ 

***Stictocephala bisonia* KOPP et YONKE, 1977
(Hemiptera: Cicadomorpha, Membracidae) in Poland**

<http://doi.org/10.5281/zenodo.2525549>

¹ University of Silesia in Katowice, Faculty of Biology and Environmental Protection, Department of Zoology,
Bankowa 9, 40-007 Katowice, Poland

² Upper Silesian Nature Heritage Centre, Graniczna 29, 40-543 Katowice, Poland

³ Doliny Miętusiej 27/44, 43-316 Bielsko-Biała, Poland

e-mail: ⁴marcin.walczak@us.edu.pl

Abstract: The aim of this paper is to provide new data on a distribution of alien and invasive species *Stictocephala bisonia* KOPP et YONKE, 1977 in Poland. This species has rapidly spread across Europe and other parts of the world in the last 100 years and has colonized new territories. *S. bisonia* belongs to the group of leafhopper species of Nearctic origin.

Key words: buffalo treehopper, Poland, invasive species, biological invasion, ecology, agriculture, crop losses, economic importance of insects.

INTRODUCTION

Stictocephala bisonia KOPP et YONKE, 1977 is an invasive leafhopper species that is native to the central and eastern part of North America (ARZONE *et al.* 1986, MIFSUD *et al.* 2010). Apart from *Gargara genistae* (FABRICIUS, 1775) and *Centrotus cornutus* (LINNAEUS, 1758) it is one of three species of the family Membracidae in Poland at the moment (ŚWIERCZEWSKI & STROIŃSKI 2011). Before the taxonomic position has changed it was classified as *Ceresa bubalus* (FABRICIUS, 1794) (KOPP & YONKE 1977).

During last 100 years *S. bisonia* has expanded its range in many regions of the world and invaded new territories. Since then, this species has been introduced to the western part of USA, Hawaii, Europe, North Africa, even to the Caucasus region and Central Asia (ŚWIERCZEWSKI & STROIŃSKI 2011). In Europe it was recorded for the first time in Kovin (Serbia) before 1912 (HORVATH 1912) and just after that time in St-Guilhem-le-Désert and Lattes near Montpellier (France) in 1918 (LALLEMAND 1920), resulting in the further colonization of the continent and currently it has been found in Albania, Austria, Belgium, Bulgaria, Czech Republic, Germany, Greece, Italy, Macedonia, Moldavia, Poland, Portugal, Romania, Slovakia, Spain, Switzerland, Ukraine, Turkey and former countries

of Yugoslavia: Bosnia, Herzegovina, Croatia and Slovenia (UVAROV 1939, HOFFRICHTER & TRÖGER 1973, OKALI 1974, DROSOPOULOS 1980, GUNTHART 1980, ARZONE *et al.* 1986, NAST 1987, JANSKÝ *et al.* 1988, D'URSO 1995, LAUTERER 1996, SELJAK 2002, HOLZINGER *et al.* 2003, GJONOV & SHISHINOVA 2014, HOCH 2018). It has been observed, that *S. bisonia* has colonized almost the whole continent excluding the northern areas. First time *S. bisonia* was noticed in Poland in 2007 from Rzeszów and next from Stara Miłosna and Zagorzyce (ŚWIERCZEWSKI & STROIŃSKI 2011). Since that time it has gradually spread all over the country.

BIOLOGY, DISTRIBUTION AND ECONOMIC IMPORTANCE

In the first half of the 20th century *S. bisonia* was treated as widespread, but rather rare species in North America (YOTHERS 1934), although it was recorded in Europe at that time (HORVÁTH 1912). Several features are decisive for the adaptation and speed of spreading of *S. bisonia*. The most important of which are related to the fact this species is polyphagous and eurythermic. KRIŠTÍN *et al.* (1987) observed individuals of this species in Slovakia between August and November, which were alive regardless of the wide range of the ambient temperatures between -1 and +36°C. Most of all specimens were recorded in September. Those insects laid eggs in rather high temperature, about 22°C. *S. bisonia* is able to colonize new areas also due to the condition of being polyphagous species with a wide range of potential food-plants. This species feeds on a several dozen plant species and it has been collected from: *Acer* spp., *Alcea rosea*, *Alnus glutinosa*, *Armeniaca vulgaris*, *Aristolochia* spp., *Artemisia vulgaris*, *Carpinus betulus*, *Castanea sativa*, *Cerasus vulgaris*, *Chrysanthemum indicum*, *Cirsium arvense*, *Clematis vitalba*, *Convolvulus arvensis*, *Cornus sanguinea*, *Crataegus laevigata*, *Erigeron annuus*, *Fraxinus excelsior*, *F. ornus*, *Humulus lupulus*, *Juglans regia*, *Galega officinalis*, *Gossypium* spp., *Ligustrum vulgare*, *Malus pumila*, *Medicago sativa*, *Melilotus alba*, *Morus alba*, *M. nigra*, *M. rubra*, *Onobrychis viciifolia*, *Padus avium*, *Plantago major*, *Persica vulgaris*, *Polygonum aviculare*, *P. persicaria*, *Populus alba*, *P. nigra*, *P. tremula*, *Prunus amygdalus*, *P. domestica*, *P. spinosa*, *Pyrus* spp., *Quercus* spp., *Ribes nigrum*, *R. rubrum*, *Rhamnus* spp., *Robinia pseudacacia*, *Rosa canina*, *Rubus caesius*, *R. idaeus*, *Rumex acetosa*, *R. acetosella*, *R. crispus*, *Salix alba*, *S. caprea*, *S. fragilis*, *S. purpurea*, *Solanum tuberosum*, *Solidago canadensis*, *S. gigantea*, *Tanacetum vulgare*, *Taraxacum officinale*, *Tilia* spp., *Trifolium repens*, *Ulmus minor*, *Urtica dioica*, *Vicia faba*, *V. sativa*, *V. labrusca* and *Vitis vinifera* (YOTHERS 1934, SAAS 1979, KRIŠTÍN *et al.* 1987, JANSKÝ *et al.* 1988, LAUTERER *et al.* 2011). Among these species there are many plants of economic importance. According to YOTHERS (1934), *S. bisonia* was considered as serious pest in USA in the first half of the 20th century. It is generally recognized as perhaps the most injurious of all the Membracidae. This species has caused yield losses in orchards and vineyards in the southern part of Europe over a period of several decades, mainly during more and more frequent gradation phases.

It is an occasional pest of fruit trees, causing injuries to young twigs by cutting the bark and phloem during the oviposition, frequently resulting in their dieback and facilitating a transmission of fungal and bacterial diseases (LAUTERER & ZACHA 1984, LAUTERER 1995, ARZONE *et al.* 1986, SELJAK 2002). Occurrence of *S. bisonia* is manifested by the presence of easily detectable puncture spots and the discoloration of the leaves apexes. These symptoms were observed e.g. in plantations of grape-bearing vines in the area of Tyrol. Damage due to the feeding of adults and larvae has been also reported on grapevine and alfalfa (VIDANO 1963, 1964). The most serious damages to fruit trees and grapevine are caused by this insect pest during its gradation years (SELJAK 2002, LAUTERER *et al.* 2011). These symptoms are

often confused with those caused by Grapevine yellows phytoplasmas (KUNZ *et al.* 2010). It is supposed, that this species does not transmit phytoplasmas astests for ESCA syndrome and grapevine leaf-roll gave negative results (STARÝ *et al.* 2013). However, the monitoring programme of *S. bisonia* is needed because of plant damage and crop losses caused by this species. The attributes like protective coloration and shape, which enable it to blend with its surroundings and afford it protection from observation by its predators, are undoubtedly important in the colonization of new areas by *S. bisonia*.

These insects are considered to be good flyers, so the ability to fly long distances can facilitate invasion of this species. It was observed that specimens of *S. bisonia* covered a distance up to 50 m during 1 hour and about 300 m during twenty four hours (JANSKÝ *et al.* 1988). It spreads most efficiently through watercourses, cultivations of fruit trees, grapevine, lucerne and arteries of traffic (SCHEDL 1991). *S. bisonia* is succesful in spreading to new areas, which is confirmed by its presence in over 20 European countries (MIFSUD *et al.* 2010). This species range includes large part of Europe with almost whole southern, western and central part, an also southern part of Turkey, Ukraine and Caucasus (ARZONE *et al.* 1986, SELJAK 2002, ŚWIERCZEWSKI & STROIŃSKI 2011). Until nineties of 20th century it has been noticed to 50 latitude of north (SCHEDL 1991), the last information from Poland confirm that *S. bisonia* extended its range up to 54 latitude (BRYSZ & SZWEDO 2015). The determination of seasonal fluctuations of these insects abundance is of great importance as this species becomes more and more common. Finding these data are not easy, because the majority of the articles focuses on crop losses caused by *S. bisonia* and information about the single specimens observed are not given. However, there are some articles on localities of this species and data have been published mainly by authors from Austria, Czech Republic, Poland (Fig. 1.), Germany and Bulgaria.




Fig. 1. Seasonal changes in abundance of *Stictocephala bisonia* in central Europe – all observed and caught imago specimens in the territory of Poland (based on new and previously published data: ŚWIERCZEWSKI & STROIŃSKI 2011, KOLAK & TASZAKOWSKI 2013, MUSIK & TASZAKOWSKI 2013, BRYSZ & SZWEDO 2015, DOBOSZ & SZWEDO 2015, TASZAKOWSKI *et al.* 2015, WALCZAK *et al.* 2016, 2018, MUSIK *et al.* 2018), Czech Republic (on the base: LAUTERER *et al.* 2011) and Austria (SCHEDL 1991). Some data contained approximate number of specimens, therefore the graph may contain a small measurement error (no more than 10% for the presented areas).

It was observed, that adult specimens of this species emerged in July and the earliest observations were recorded in Czech Republic on July 12th and in Germany on July 14th, while the latest in Czech Republic on October 10th, in Germany October 27th and in Slovakia on November 11th (KRISTÍN *et al.* 1987, LANDECK 2011, LAUTERER *et al.* 2011). In Poland the adult insects were observed between July 18th an October 26th (see in: New records). To summarize, in the central part of Europe adult specimens of *S. bisonia* are most abundant between second decade of August and first decade of September (Fig. 1.). This conclusion is confirmed by comprehensive information about its occurrence in Poland and Czech Republic and also by observations made in Austria, Bulgaria and Germany, where adults were the most abundant in August and September (SCHEDL 1991, LANDECK 2011, GJONOV & SHISHINIOVA 2014). Information about immature stages of *S. bisonia* is unavailable. According to SCHEDL (1991) first nymphs were found in Austria on May 26th, whereas immatures were collected in Beskid Wschodni (Poland) on May 10th (TASZAKOWSKI *et al.* 2015).

MATERIAL AND METHODS

Specimens of *S. bisonia* were collected between 2011 and 2018. Common collecting methods were used, including a sweep net ($\varnothing = 30$ cm) and visual examination of host plants (WALCZAK *et al.* 2016). Specimens were killed with ethyl acetate or ethanol. In this latter case they were preserved in vials containing solution of ethanol in water. Specimens were identified on the basis of their morphology using the identification keys by BIEDERMANN & NIEDRINGHAUS (2004) and ŚWIERCZEWSKI & STROIŃSKI (2011). The division into zoogeographical region was adopted from „Katalog Fauny Polski” (BURAKOWSKI *et al.* 1973) and the names of regions from GĘBICKI *et al.* (2013).

Specimens are deposited in the Insect Collection of the Department of Zoology (DZUS) (Faculty of Biology and Environmental Protection, University of Silesia, Katowice, Poland). Katarzyna Strzelczyk, who is a student of the Faculty of Biology and Environmental Protection, University of Silesia, was given a permission to legally collect leafhoppers in “Łęczok” Nature Reserve by the Regional Director for Environmental Protection in Katowice (permission number RDOŚ PN 6205.30.2015 MM).

DISTRIBUTION IN POLAND

Published records.

S. bisonia were recorded from 19 localities (in 8 zoogeographical regions) in Poland to date (Fig. 2.):

Baltic Coast: Gdańsk-Oliwa [CF43] (BRYSZ & SZWEDO 2015); **Wielkopolsko-Kujawskia Lowland:** “Łęgi koło Słubic” Nature Reserve [VU60, VT79] (BRYSZ & SZWEDO 2015); **Mazovian Lowland:** Stara Miłosna [EC18] (ŚWIERCZEWSKI & STROIŃSKI 2011); **Upper Silesia :** Wodzisław Śląski [CA14] (KOLAK & TASZAKOWSKI 2013), Rudziniec [CA18] (DOBOSZ & SZWEDO 2015), Piekary Śląskie [CA58], Bukowno [CA86] (MUSIK *et al.* 2018); **Krakowsko-Wieluńska Upland:** Częstochowa-Dźbów [CB62] and Częstochowa-Lisiniec [CB63] (WALCZAK *et al.* 2016); **Malopolska Upland:** Łódź: Widzew, Młynek and Botanic Garden [CC93] (BRYSZ & SZWEDO 2015); **Sandomierska Lowland:** Rzeszów [EA63], Zagorzyce [EA44] (ŚWIERCZEWSKI & STROIŃSKI 2011); **Eastern Beskydy Mountains:** Libusza [EA10], Dobrynia [EV39], Lisów [EA21], Ożenna [EV37], Zdynia [EV28] (MUSIK & TASZAKOWSKI 2013, TASZAKOWSKI *et al.* 2015), Bartne [EV29] (WALCZAK *et al.* 2018).




Fig. 2. Distribution of *Stictocephala bisonia* in Poland: ● – published records from Poland, ● – new localities in Poland.

New records.

This paper presents information about 68 new localities of *S. bisonia* situated in 11 zoogeographical regions in Poland (Fig. 2). *S. bisonia* is a species new to 5 regions: Pomeranian Lake District, Lower Silesia, Eastern Sudetes Mountains, Western Beskyd Mountains and Pieniny Mountains. Data were collected by many researchers, so they may be presented in different ways.

Pomeranian Lake District (reported here as new to this region):

Kaszubski PK, Kamienica Królewska [54°23'32"N, 17°53'19"E; XA83], 24.07.2018, 1 ex., observ. R. Kaźmierczak.

Mazovian Lowland:

Warszawa, Zakole Wawerskie [52°13'47"N, 21°08'08"E; EC08], 01.08.2018, 1 ex., observ. D. Kowalczyk.

Lower Silesia (reported here as new to this region):

Brzeg [50°51'35"N, 17°28'02"E; XS73], 07.08.2014, 1 ex., observ. J. Regner;

Chróścice, gm. Dobrzenie Wielki [50°46'43"N, 17°48'44"E; XS92], 02.09.2018, 1 ex., observ. M. Lisowska;

Gogolin [50°29'31"N, 18°01'16"E; BB89], 06.09.2018, 2 exx., observ. M. Pastrykiewicz;

Gogolin [50°30'17"N, 18°01'35"E; BB89], 10.09.2018, 3 exx., closed quarry near garbage dump, observ. M. Pastrykiewicz;

Górażdże, the Limestone Quarry Górażdże [50°31'58"N, 18°01'15"E; BB90], a xerothermic grasslands, on *Prunus domestica*: 26.07.2018, 1♀, leg. M. Walczak;

Górażdże, the Limestone Quarry Górażdże [50°31'54"N, 18°01'24"E; BB90], a coast of the water reservoir, on *Alnus* sp.: 26.07.2018, 3♂♂, leg. M. Walczak;

Górażdże, the Limestone Quarry Górażdże [50°32'10"N, 18°01'01"E; BB90], a reed bed (*Phragmites*), on *Salix* sp.: 26.07.2018, 2♂♂, 1♀, leg. M. Walczak, 4♀♀, leg. A. Łazuka;

Górażdże, the Limestone Quarry Górażdże [50°31'49"N, 18°01'30"E; BB90], a mixed forest, on shrubs: 26.07.2018, 1♀, leg. A. Łazuka;

Górażdże, the Limestone Quarry Górażdże [50°32'08"N, 18°01'15"E; BB90], a ruderal community with domination *Calamagrostis epigejos*, on young seedlings *Populus nigra*: 26.07.2018, 3♀♀, leg. A. Łazuka.

Upper Silesia:

Będzin, Łagisza [50°20'58"N, 19°08'14"E; CA67], 06.09.2017, 2 ex., observ. S. Skrzypiec;

Chorzów, Wiejska Street, "Żabie Doły" [50°19'12.8"N, 18°57'33.5"E; CA57], a spoil heap: 18.07.2014, 7♂♂, 3♀♀, 04.08.2014, 3♂♂, 4♀♀, 18.08.2014, 2♂♂, 6♀♀, 04.09.2014, 4♂♂, 1♀, leg. M. Gibas;

Chorzów, Wiejska Street, "Żabie Doły" [50°19'13.5"N, 18°57'38.7"E; CA57], an initial forest community dominated by birches that overgrows an edge of a spoil heap: 18.07.2014, 2♂♂, 1♀, 04.08.2014, 6♂♂, 2♀♀, 18.08.2014, 4♂♂, 2♀♀, 04.09.2014, 4♂♂, leg. M. Gibas;

Dąbrowa Górnica, Antoniów, close to Spacerowa Street [50°22'19"N, 19°12'47"E; CA78], a thicket: 12.08.2016, 1♂, leg. A. Potocki;

Gliwice, Łabędy, Fabryczna Street, [50°20'59"N, 18°37'13"E; CA38], garden, on *Cydonia* sp.: 14.09.2016, 2♂♂, 1♀, leg. Ł. Junkiert;

Imielin, Golcówka [50°08'53"N, 19°12'55"E; CA75], a xerothermic grasslands: 13.08.2017, 1♀, leg. N. Kaszyca;

Katowice, Dąbrówka Mała, Szwedzka Street, near Ikea [50°16'11"N, 19°03'29"E; CA67], a ruderal plant community: 17.08.2013, 1♀, leg. L. Kruszelnicki;

Kotlarnia [50°16'13"N, 18°22'26"E; CA07], 12.08.2018, at least 5 exx., Coal Mine, observ. M. Pastrykiewicz;

Miejsce, near Spytkowice [50°01'05"N, 19°30'33"E; CA94], 20.08.2016, 1 ex., observ. G. Kolago;

Mysłowice, a vicinity of Piastów Śląskich Street [50°10'32"N, 19°05'43"E; CA66], a mining waste of "Mysłowice-Wesoła" Coal Mine: 30.08.2014, 1♂, leg. M. Pniok;

Piekary Śląskie, Lotników Street [50°21'59"N, 18°58'14"E; CA58], a post-industrial wasteland covered with a grass community dominated by *Deschampsia caespitosa*: 27.09.2011, 1♀, leg. M. Kalandyk-Kołodziejczyk;

Pilchowice, Stanicka Street [50°12'37"N, 18°32'42"E; CA26], post-agricultural wasteland: 24.09.2012, 1♀, 28.08.2013, 1♂ (Fig. 3.), 14.09.2013, 1♀, leg. Ł. Junkiert;

Puszczew [50°48'09"N, 18°51'21"E; CB43], garden, 14.08.2018, 2♂♂, 1♀, leg. M. Walczak;

Racibórz, Markowice, "Łeżczok" Nature Reserve [50°07'43"N, 18°16'14"E; CA05], a thicket of *Salix* sp.: 17.07.2016 1♂, 22.07.2016 1♂, 14.09.2016 1♀, leg. K. Strzelczyk;

Rogoźnik [50°24'02"N, 18°57'36,6"E; CA58], a thicket at the lakeshore: 18.08.2012, 1♀, leg. K. Musik;

Ruda Kozielska, [50°12'N, 18°24'E; CA16], on *Salix* sp.: 30.08.2013, 1♂, leg. Ł. Junkiert;

Babice near Nędza, Rudzka Street, [50°08'47"N, 18°17'52"E; CA05], wayside shrubs: 17.08.2011, 1♀, leg. Ł. Junkiert;

Siemianowice, Przełajka, Rzeczna Street, [50°20'47"N, 19°03'31"E; CA67], a meadow located near Brynica River: 24.09.2016, 1♂, 6♀♀, leg. J. Brożek;



Fig. 3. Habitus of *Stictocephala bisonia*, dorsal view – male from Pilchowice (collected on 28 August 2013), leg. and det. L. Junkiert.

Sosnowiec, Zagórze, a vicinity of Kielecka Street [50°17'21"N, 19°10'46"E; CA77], urban greenery: 13.08.2016, 1♂, leg. M. Walczak;

Świętochłowice, "Hugona Hill", a vicinity of Śląska Street [50°16'38.8"N, 18°54'48.0"E; CA57], a forest community overgrowing a coal mine heap: 21.07.2014, 2♂♂, 1♀, 1 nymph, 06.08.2014, 2♂♂, 21.08.2014, 3♀♀, leg. M. Gibas;

Trzebinia [50°09'30"N, 19°28'10"E; CA95], 02.10.2012, 1 ex., leg. R. Celadyn, det. & coll. A. Itczak.

Krakowsko-Wieluńska Upland:

Czajowice, quarry [50°11'22"N, 19°48'23"E; DA16], 10.09.2016, 1 ex., observ. G. Kolago;

Kłokoczyn [49°59'42"N, 19°38'14"E; DA03], 17.08.2014, 1 ex., observ. G. Kolago;

Kraków-Czyżyny [50°03'50"N, 20°00'51"E; DA24], wasteland, the remains after garden plots, 05.09.2014, at least 10 exx., observ. G. Kolago;

Kraków-Dębniki, ul. Gronostajowa [50°01'38"N, 19°53'55"E; DA24], *Solidago* sp. and *Urtica dioica* shrubs near Gronostajowa Street, 50°01'42", E19°53'52", 15.09.2015, 7♂♂, 8♀♀, leg. M. Kobiałka & M. Walczak, 04.09.2017, 6♂♂, 6♀♀, leg. A. Michalik, T. Szklarzewicz & M. Walczak;

Kraków-Dębniki, osiedle Bodzów, quarry [50°02'10"N, 19°52'26"E; DA14], 27.08.2015, 1 ex., observ. G. Kolago;

Kraków, Krzemionki [50°02'22"N, 19°56'49"E; DA24], 10.09.2016, 1 ex., observ. G. Kolago;
Kraków, Łąki Nowohuckie [50°04'02"N, 20°02'07"E; DA34], 16.08.2016, 1 ex., observ. G. Kolago;
Kraków, Park Lotników Polskich [50°04'13"N, 19°59'36"E; DA24], 31.08.2016, 1 ex., observ. G. Kolago;
Kraków, Dąbski Pond [50°03'54"N, 19°59'14"E; DA24], 24.07.2017, 1 ex., observ. G. Kolago;
Kraków, Płaszowski Pond [50°02'30"N, 19°58'02"E; DA24], 20.08.2015, 1 ex., observ. G. Kolago;
Przylasek Rusiecki [50°03'33"N, 20°09'30"E; DA34], 26.10.2013, 1 ex., observ. G. Kolago;
“Bonarka” Nature Reserve, vicinity [50°01'47"N, 19°57'31"E; DA24], 04.09.2016, 1 ex., observ. G. Kolago.

Malopolska Upland:

Kalina-Lisiniec, Natura 2000, buffer zone of the floristic Nature Reserve “Kalina-Lisiniec” [50°21'42"N, 20°09'22"E; DA37], 02.08.2018, 1 ex., observ. G. Kolago;
Nowy Koreczyn [50°17'46"N, 20°48'19"E; DA87], a meadow near Wisła River, 07.08.2015, 1 ex., observ. G. Kolago;
Pińczów, a vicinity of Wierciszów [50°30'56"N, 20°33'07"E; DA69], a xerothermic grasslands: 22.07.2017, 3♂♂, leg. Ł. Depa & N. Kaszyca;
Piotrowice [50°11'34"N, 20°39'34"E; DA76], Wisła riverside, 09.08.2014, 1 ex., observ. G. Kolago;
Polana Polichno [50°28'46"N, 20°30'05"E; DA69], 12.08.2018, 1 ex., observ. R. Kaźmierzak;
“Góry Pieprzowe” Nature Reserve [50°41'05"N, 21°47'08"E; EB51], 07.09.2014, a few exx., 30.08.2015, 1 ex., observ. G. Kolago;
“Góry Wschodnie” Nature Reserve [50°22'18"N, 20°44'02"E; DA88], 02.09.2018, 1 ex., observ. G. Kolago
Wola Chorberska [50°23'52"N, 20°31'15"E; DA68], a xerothermic grasslands: 22.07.2017, 1♀, leg. Ł. Depa & N. Kaszyca.

Sandomierska Lowland:

Morsko (Koszycki Obszar Chronionego Krajobrazu) [50°08'34"N, 20°32'59"E; DA65], 26.07.2017, 1 ex., 24.08.2017, 1 ex., observ. G. Kolago.

Eastern Sudetes Mountains (reported here as new to this region):

Opawskie Mountains, Wierzbiec [50°19'19"N, 17°27'25"E; XR77], 30.08.2017, 2♂♂, 1♀, leg. K. Błaszczyk, D. Świerczewski & M. Walczak.

Western Beskydy Mountains (reported here as new to this region):

Bielsko-Biała, Dolina Gościnna [49°47'10"N, 19°01'04"E; CA51], 10.09.2013, 1 ex., leg. & det. G. Gierlasiński;
Bielsko-Biała, Kozia Góra [49°46'04"N, 19°02'21"E; CA51], 29.08.2015, 1 ex., leg. & det. G. Gierlasiński;
Bielsko-Biała, Trzy Lipki [49°50'05"N, 19°01'02"E; CA52], 12.09.2015, 1 ex., leg. & det. G. Gierlasiński;
Buczkowice [49°43'40"N, 19°04'09"E; CA61], 30.07.2014, 1 ex., observ. G. Gierlasiński;
Bystra [49°45'09"N, 19°01'15"E; CA51], 29.09.2017, 1 ex., leg. & det. G. Dubiel;

Cieszyn, ecological use “Łaki na Kopcech” [49°46'09"N, 18°37'00"E; CA21], 18.08.2018, 1 ex., leg. & det. G. Gierlasiński;

Leszna Góra [49°41'32"N, 18°43'49"E; CA30], 22.08.2018, 1 ex., leg. & det. G. Gierlasiński;

Meszna [49°44'46"N, 19°03'47"E; CA61], 17.08.2017, 1 ex., observ. G. Gierlasiński;
Milówka [49°33'33"N, 19°05'12"E; CA69], 09.09.2017, 1 ex., leg. & det. G. Gierlasiński.

Eastern Beskidy Mountains:

Męcina Wielka, close to Gorlice [49°37'20"N, 21°15'53"E; EV19], a mesotrophic grassland communities: 07.08.2018, 2♂♂, leg. N. Kaszycia;

Nowe Sady [49°38'27"N, 22°44'37"E; FA20], 19.08.2018, 1 ex., the shrubs near Wiar Riverside, leg. T. Rutkowski;

Nowosiółki Dydyńskie [49°38'10"N, 22°43'49"E; FV29], the shrubs near Wiar Riverside, 1 ex., 18.08.2018, leg. T. Rutkowski;

Pacław [49°37'31"N, 22°42'24"E; FV29], 1 ex., 22.08.2018, leg. T. Rutkowski.

Pieniny Mountains (reported here as new to this region):

Maniowy [49°27'32"N, 20°15'54"E; DV47], 11.08.2018, 1 ex., leg. & det. G. Gierlasiński.

Moreover another specimens were observed in Łódź in localities previously noticed:
Łódź (CC93), 11.08.2016, 1 ex., 02.09.2017, 1 ex., 18.08.2018, 1 ex., Łódź (DC03),
13.08.2017, 1 ex., observ. R. Kaźmierzak.

DISCUSSION

Biological invasions, which are considered as a global phenomenon that largely decreases global biodiversity, have become a serious ecological and economic problem in the recent years. They can be treated as „side effects” of dynamic development of our civilization.

Our fear of invasive species is caused by their adverse effects on the environment and negative impact on yields. These species can change habitats so the ecosystems they inhabited are supposed to loss their dynamic equilibrium. Food source availability and the lack of predators feeding on them in the new location may cause invasive species population growth. Invasive species outcompete native species for resources such as nutrients, light and water and can crowd out native animals or plants (PERRINGS 2001, BROWN & SAX 2005, MATTSON *et al.* 2007, MIFSUD *et al.* 2010 WILSON *et al.* 2016), moreover they may contribute to many native species extinction (ROTHSCHILD 1907, WILMSHURST *et al.* 2008). Many species of Cicadomorpha and Fulgoromorpha are considered as serious agricultural pests such as *Nilaparvata lugens* and *Sogatella furcifera*, which cause damages to rice crops in Asia and they are among the most important insect pests in the world (GHOSAL *et al.* 2018, RATHEE & DALAL 2018). In Europe crop losses, damages to garden plants and at nurseries caused by alien species of Cicadomorpha and Fulgoromorpha are more and more noticeable. *S. bisonia* is not the only species that has negative impact on cultivated plants. There are other species, attacking plants, with non-european origin: *Erythroneura vulnerata*, *Scaphoideus titanus* and *Metcalfa pruinosa* (ARZONE *et al.* 1986, SELJAK 2002, PAPURA *et al.* 2012, GOGAN *et al.* 2013, GJONOV & SHISHINIOVA 2014, PREDA & SKOLKA 2011). The Auchenorrhyncha species that are non-native to Europe were recently summarized by MIFSUD *et al.* (2010) and, according to this review, a total of 12 species introduced from North America and East Asia have been established in Europe. At least 6 leafhopper species that have invaded Europe and have been recorded recently should be included in this list: *Ricania japonica*, *Erasmoneura vulnerata*, *Osbornellus auronitens*, *Penestragania apicalis*, *Penthimiola bella* and *Sophonia*.

orientalis (GJONOV 2011, SELJAK 2011, NICKEL *et al.* 2013, ZINA *et al.* 2013, TRIVELLONE *et al.* 2017, WILSON *et al.* 2011). Many of them have colonized Central Europe, while 5 of them have been encountered in Poland so far: *Graphocephala fennahi*, *Eupteryx decemnotata*, *Japananus hyalinus*, *Orientus ishidae* (SOIKA & ŁABANOWSKI 2000, MUSIK 2011, WALCZAK *et al.* 2012, LUBIARZ & MUSIK 2015, KLEJDYSZ *et al.* 2017) and mentioned here *S. bisonia*. Our results presented here extend the knowledge of the distribution of *S. bisonia* in Poland and constitute a basis for the further research in Poland and other countries. The research program for this species should include: monitoring of the range expansion, measurement of population size per unit area, account for crop yield losses and gathering information on host plants – especially cultivated plants.

ACKNOWLEDGEMENTS

First of all the authors would like to thank students of Faculty of Biology and Environmental Protection (University of Silesia): Mateusz Gibas, Mariola Pniok, Katarzyna Strzelczyk, Artur Potocki and also Dr. Łukasz Depa – for received specimens of *S. bisonia* gathered especially from the area of Upper Silesia. We are thankful for their information about the locations of the individuals and cooperation with authors. We would like to express our very great appreciation to the students of the Faculty of Biology and Environmental Protection, University of Silesia for their written permissions that have enabled us to publish information about the occurrence of *S. bisonia* included in their bachelor and master's degree thesis. Our special thanks are extended to the staff of Department of Developmental Biology nad Morphology of Invertebrates, Jagiellonian University: Professor Teresa Szklarzewicz, Dr. Anna Michalik and Dr. Michał Kobiałka for their assistance in collecting specimens of *S. bisonia*. Thanks are also due to Dr. Dariusz Świerczewski and MSc. Jakub Błaszczyk, University of Jan Dlugosz in Częstochowa. We would like to thank also other people, mainly users of polish entomological internet forum (entomo.pl): Marek Adamski, Rafał Celadyn, Grzegorz Dubiel, Aneta Itczak, Rafał Kaźmierczak, Grzegorz Kolago, Dariusz Kowalczyk, Lech Kruszelnicki, Monika Lisowska, Monika Pastrykiewicz and Sebastian Skrzypiec for supplying the specimens of *S. bisonia* together with necessary information. Without their help, it would not be possible to collect such an exhaustive data from almost all of Poland.

NOTICES/COMMENTS

The presented work initially concerned only the occurrence of *S. bisonia* in Upper Silesia. The process of its creation was, however, long, and during the preparation in several works was quoted in a less numerous composition and under a different title [WALCZAK M., BROZEK J., JUNKIERT Ł., KALANDYK-KOŁODZIEJCZYK M., MUSIK K. in prep. *Stictocephala bisonia* KOPP et YONKE, 1977 (Hemiptera: Cicadomorpha, Membracidae) in Upper Silesia]. For this reason, the earlier citation should be referred to the article presented here.

REFERENCES

- ARZONE A., VIDANO C., ALMA A. 1986. Auchenorrhyncha introduced into Europe from the Nearctic region: taxonomic and phytopathological problems, In: WILSON M.R. & NAULT L.R. (Eds.), Proceedings of 2nd International Workshop on Leafhoppers and Planthoppers of Economic Importance, Provo, Utah, USA, 28th July-1st August 1986, CIE, London: 3–17.
- BIEDERMANN R., NIEDRINGHAUS R. 2004. Die Zikaden Deutschlands. Bestimmungstafeln für alle Arten. Wissenschaftlich Akademischer Buchvertrieb-Fründ, Scheeßel: 1–409.
- BROWN J.H., SAX D. F. 2005. Short Note. Biological invasions and scientific objectivity: Reply to CASSEY *et al.* (2005). *Australian Ecology* 30: 481–483.

- BRYSZ A., SZWEDO J. 2015. Nowe stanowiska kolcoroga bizoniaka *Stictocephala bisonia* KOPP et YONKE, 1977 (Hemiptera: Cicadomorpha: Membracidae) w Polsce. *Acta entomologica silesiana* 23: 229–230.
- BURAKOWSKI B., MROCKOWSKI M., STEFAŃSKA J. 1973. Chrząszcze Coleoptera. Biegaczowate – Carabidae, cz. 1. *Katalog fauny Polski* 23(2): 1–232.
- DOBOSZ R., SZWEDO J. 2015. Drugie stanowisko *Stictocephala bisonia* KOPP et YONKE (Hemiptera: Cicadomorpha: Membracidae) na Górnym Śląsku. *Acta entomologica silesiana* 2: 225.
- DROSOPOLOUS S. 1980. Hemipterological studies in Greece Part II. Homoptera – Auchenorrhyncha, a catalogue of the reported species. *Biologia Gallo-Hellenica* 9(1): 187–194.
- D’URSO V. 1995. Homoptera Auchenorrhyncha, In: MINELLI A., RUFFO S., LA POSTA S. (eds.). Checklist delle specie della fauna italiana. *Calderini, Bologna* 41: 1–35.
- GĘBICKI C., ŚWIERNIECKI D., SZWEDO J. 2013. Planthoppers and leafhoppers of Poland (Hemiptera: Fulgoromorpha et Cicadomorpha). Systematics. Check-list. Bionomy. *Annals of the Upper Silesian Museum in Bytom, Entomology* 21–22: 5–259.
- GHOSAL A., DOLAI A.K., CHATTERJEE M.L. 2018. Evaluation of a new ready mixed insecticide (Buprofezin 15% +Acephate 35% WP) against BPH (*Nilaparvata lugens* (STAL)) and WBPH (*Sogatella furcifera* (HORVATH)) of rice in West Bengal. *Journal of Crop and Weed* 14(1): 205–211.
- GJONOV I. 2011. *Ricania japonica* MELICHAR, 1898 – a representative of family Ricanidae (Homoptera, Fulgoromorpha), new to the fauna of Bulgaria. *ZooNotes* 23: 1–3.
- GJONOV I., SHISHINOVA M. 2014. Alien Auchenorrhyncha (Insecta, Hemiptera: Fulgoromorpha and Cicadomorpha) to Bulgaria. *Bulgarian Journal of Agricultural Science* 20(Suppl. 1): 151–156.
- GOGAN A., GROZEA I., VIRTEIU A.M., BAGHINA N.G. 2013. Flatid plant hopper (*Metcalfa pruinosa* SAY) a new dangerous insect pest of ornamental and agricultural plants in Romania. *Journal of Food, Agriculture & Environment* 11(3–4): 2065–2070.
- GUNTHART H. 1980. Neuer Fundort und neuer Name für die altbekannte Buffelzikade “*Ceresa bubalus*” (Hom. Auch. Membracidae). *Mitteilungen der entomologischen Gesellschaft Basel* (N.F.) 30: 105–109.
- HOCH H. 2018. Fauna Europaea: Hemiptera: Cicadomorpha: *Stictocephala bisonia*. <http://www.faunaeur.org>. (Accessed on 22.09.2018).
- HOFFRICHTER O., TRÖGER E.J. 1973. *Ceresa bubalus* F. (Homoptera: Membracidae) – Beginn der Einwanderung in Deutschland. *Mitteilungen des Badischen Landesvereins für Naturkunde und Naturschutz* N.F. 11: 33–43.
- HOLZINGER W., KAMMERLANDER I., NICKEL H. 2003. The Auchenorrhyncha of Central Europe – Die Zikaden Mitteleuropas Fulgoromorpha, Cicadomorpha excl. Cicadellidae in Brill Publishers, Leiden, The Netherlands: 1–673.
- HORVÁTH G. 1912. Az amerikai bivalykabócza Magyarországon. *Rovartani lapok* 19: 145–147.
- JANSKÝ V., KRIŠTÍN A., OKÁLÍ I. 1988. Der gegenwärtige Stand der Verbreitung und neue Erkenntnisse über die Bionomie der Art *Stictocephala bisonia* (Homoptera, Membracidae) in der Slowakei. *Biologia* (Bratislava) 43: 527–533.
- KLEJDYSZ T., ZWOLIŃSKA A., WALCZAK M., KOBIALKA M. 2017. First record of a potential pest *Orientus ishidae* (MATSUMURA, 1902) (Hemiptera: Cicadellidae) in Poland. *Journal of Plant Protection Research* 57(2): 107–112.
- KOLAK G., TASZAKOWSKI A. 2013. Nowe stanowisko występowania *Stictocephala bisonia* KOPP et YONKE, 1977 (Hemiptera: Cicadomorpha: Membracidae) w Polsce. *Acta entomologica silesiana* 21: 77.
- KOPP D.D., YONKE T.R. 1977. Taxonomic status of the buffalo treehopper and the name *Ceresa bubalus*. *Annals of the entomological Society of America* 70(6): 901–905.
- KRIŠTÍN A., JANSKY V., OKÁLÍ I. 1987. Is *Stictocephala bisonia* (Membracidae) an invasion species? 6th Auchen. Meeting, Turin, 7–11 Sept. 1987: 417–424.
- KUNZ G., ROSCHAIT C., SCHWEIKOFLER W. 2010. Biodiversity of planthoppers (Auchenorrhyncha) in vineyards infected by the Bois noir phytoplasma. *Gredleriana* 10: 89–108.
- LALLEMAND V. 1920. Un Mambracidae (Hem.) nouveau pour la faune française. *Bulletin de la Société entomologique de France* 3: 56.
- LANDECK I. 2011. Funde der Büffelzikade (*Stictocephala bisonia* KOPP & YONKE, 1977) im südlichen Land Brandenburg (Auchenorrhyncha, Membracidae). Märkischen Entomologischen Nachrichten 13(2): 221–226.
- LAUTERER P. 1995: Ostnöhřbetka *Stictocephala bisonia* nový škůdce na Moravě. *Rostlinolékař* 6(1): 17.
- LAUTERER P. 1996. Faunistic novelties of leafhoppers (Homoptera, Auchenorrhyncha) from the territory of the Czech Republic, with notes on the fauna of Slovakia. *Acta Musei Moraviae. Scientiae naturales* 80: 235–242.
- LAUTERER P., ZACHA V. 1984. Ostnöhřbetka *Stictocephala bisonia* jako škůdce ovočných stromů na jižním Slovensku. *Sborník UV TIZ – Ochrana rostlin* 20[57](1): 39–45.
- LAUTERER P., MALENOVSKÝ I., ŠPRYŇAR P. 2011. Invasion of the treehopper *Stictocephala bisonia* (Hemiptera: Cicadomorpha: Membracidae) in the Czech Republic: current distribution and first records from Bohemia. *Klapalekiana* 47: 67–73.
- LUBIARZ M., MUSIK M. 2015. First record in Poland of the Ligurian leafhopper, *Eupteryx decemnotata* REY, 1891 (Cicadomorpha, Cicadellidae) – an import ant pest of herbs. *Journal of Plant Protecton Research* 55(3): 324–326.

- MATTSON W., VANHANEN H., VETELI T., SIVONEN S., NIEMELA P. Few immigrant phytophagous insects on woody plants in Europe: legacy of the European crucible? *Biological Invasions* 9: 957–974.
- MIFSUD D., COCQUEMPT C., MÜHLETHALER R., WILSON M., STREITO J.C. 2010. Other Hemiptera Sternorrhyncha (Aleyrodidae, Phylloxeroidea and Psylloidea) and Hemiptera Auchenorrhyncha. Chapter 9.4, In: ROQUES A. et al. (Eds.). Alien terrestrial arthropods of Europe. *BioRisk* 4(1): 511–552.
- MUSIK K. 2011. New record of an alien species *Graphocephala fennahi* YOUNG, 1977 (Hemiptera: Cicadomorpha) in Poland. *Acta entomologica silesiana* 19: 41–44.
- MUSIK K., TASZAKOWSKI A. 2013. New data on some rare planthoppers and leafhoppers in Poland (Hemiptera: Auchenorrhyncha). *Acta Musei Moraviae, Scientiae biologicae* 98(2): 265–271.
- MUSIK K., WALCZAK M., KALANDYK-KOŁODZIEJCZYK M., WOJCIECHOWSKI W. 2018. Planthopper and leafhopper communities (Hemiptera: Fulgoromorpha et Cicadomorpha) of selected plant associations of Garb Tarnogórski. *Monographs of the Upper Silesian Museum* 7: 1–244.
- NAST J. 1987. The Auchenorrhyncha (Homoptera) of Europe. *Annales Zoologici* 40: 535–661.
- NICKEL H., CALLOT H., KNOP E., KUNZ G. 2013. *Penestragania apicalis* (OSBORN & BALL, 1898), another invasive Nearctic leafhopper found in Europe (Hemiptera: Cicadellidae, Iassinae). *Cicadina* 13:51–15.
- OKALI I. 1974. *Stictocephala bubalus* (FABRICIUS, 1794) (Homoptera, Membracidae) – eine neue Art für die Fauna der Tschechoslowakei. *Acta Rerum naturalium Musei nationalis slovaca* 20: 257–258.
- PAPURA D., BURBAN C., VAN HELDEN M., GIRESSE X., NUSILLARD B., GUILLEMAUD T., KERDELHUE C. 2012. Microsatellite and Mitochondrial Data Provide Evidence for a Single Major Introduction for the Nearctic Leafhopper *Scaphoideus titanus* in Europe. *PlosOne* 7(5): 1–12.
- PERRINGS C. 2001. The economics of biological invasions. *Land Use and Water Resources Research* 1(3): 1–9.
- PREDA C., SKOLKA M. 2011. Range Expansion of *Metcalfa pruinosa* (Homoptera: Fulgoroidea) in Southeastern Europe. *Ecologia Balkanica* 3(1): 79–87.
- RATHEE M., DALAL P. 2018. Emerging Insect Pests In Indian Agriculture. *Indian Journal of Entomology* 80(2): 267–281.
- ROTHSCHILD W. 1907. Extinct birds. An attempt to unite in one volume a short account of those Birds which have become extinct in historical times – that is, within the last six or seven hundred years. To which are added a few which still exist, but are on the verge of extinction. Hutchinson & Co., London: 1–386.
- SAAS B. 1979. Wirtspflanzen und Schadttigkeit der amerikanischen Büffelzikade (*Ceresa bubalus* F.) in Ungarn und Möglichkeiten der Bekämpfung. *Növényvédelem* 15: 67–73.
- SCHEDL W. 1991. Invasion der Amerikanischen Büffelzikade (*Stictocephala bisonia* KOPP und YONKE, 1977) nach Österreich (Homoptera, Auchenorrhyncha, Membracidae). *Anzeiger für Schädlingskunde, Pflanzenschutz, Umweltschut* 64: 9–13.
- SELJAK G. 2002. Non-European Auchenorrhyncha (Hemiptera) and their geographical distribution in Slovenia. *Acta Entomologica Slovenica* 10(1): 97–101.
- SELJAK, G., 2011. First record of the nearctic leafhopper *Erasmoneura vulnerata* (FITCH, 1851) [Hemiptera, Cicadomorpha: Cicadellidae] in Slovenia. *Acta Entomologica Slovenica* 19(1): 37–42.
- SOIKA G., ŁABANOWSKI G. 2000. Pests of ericaceous plants in the botanical gardens in Poland and proposals for their control, In: WIECH K., ZEMANEK B. (Eds.), Protection of plant collections against pests and diseases. Vol. I, Oficyna Wydawnicza Text, Kraków: 71–76.
- STARÝ M., VÁLOVÁ P., ŠAFÁŘOVÁ D., LAUTERER P., ACKERMANN P., NAVRÁTIL M. 2013. Survey and molecular detection of Bois noir in vineyards of the Czech republic – Short communication. *Horticultural Science* 40(2): 83–87.
- ŚWIERCZEWSKI D., STROIŃSKI A. 2011. The first record of the Nearctic treehopper *Stictocephala bisonia* in Poland (Hemiptera: Cicadomorpha: Membracidae) with some comments on this potential pest. *Polish Journal of Entomology* 80(1): 13–22.
- TASZAKOWSKI A., WALCZAK M., MORAWSKI M., BARAN B. 2015. Piewiki (Hemiptera: Fulgoromorpha et Cicadomorpha) Beskidu Wschodniego. *Acta entomologica silesiana* 23: 83–96.
- TRIVELLONE V., MITROVIĆ M., CHRISTOPHER H. DIETRICH C.H., TOŠEVSKI I. 2017. *Osbornellus auronitens* (Hemiptera: Cicadellidae: Deltoccephalinae), an introduced species new for the Palaearctic region. *The Canadian Entomologist* 149(5): 551–559.
- UVAROV B.P. 1939. An American Membracid in Jugoslavia (Hemiptera). *Proceedings of the Royal Entomological Society of London* 14: 48.
- VIDANO C. 1963: Deviazione trofica ampelofila della *Ceresa bubalus* FABRICIUS e rispondenza reattiva del vegetale. *Atti della Accademia delle Scienze di Torino* 98: 193–212.
- VIDANO C. 1964: Reperti inediti biologici e fitopatologici delle *Ceresa bubalus* FABRICIUS quale nuovo fitomizo della vite. *Rivista di Vinocultura e di Enologia di Conegliano* 17: 457–482.
- WALCZAK M., GĘBICKI C., WOJCIECHOWSKI W., ŚWIERCZEWSKI D. 2016. The fauna of planthoppers and leafhoppers (Hemiptera: Fulgoromorpha et Cicadomorpha) in the city of Częstochowa (southern Poland). The Monograph. *Annals of the Upper Silesian Museum, Entomology* 24: 1–190.

- WALCZAK M., MUSIK K., MOKRZYCKA A. 2012. *Japananus hyalinus* (OSBORN, 1900) – a new leafhopper for Polish fauna (Hemiptera: Cicadomorpha). *Wiadomości Entomologiczne* 31(4): 242–250.
- WALCZAK M., KASZYCA N., TASZAKOWSKI A. 2018. Materiały do znajomości piewików (Hemiptera: Fulgoromorpha, Cicadomorpha) Beskidu Wschodniego. *Acta entomologica silesiana* 26(034): 1–9 [online].
- WILMSHURST J.M., ANDERSON A.J., HIGHAM T.F., WORTHY T.H. 2008. Dating the late prehistoric dispersal of Polynesians to New Zealand using the commensal Pacific rat. *Proceedings of the National Academy of Sciences of the United States of America* 22(105): 7676–7680.
- WILSON J.R.U., GARCÍA-DÍAZ P., CASSEY P., RICHARDSON D.M., PYŠEK P.M., BLACKBURN T.M. 2016. Biological invasions and natural colonisations are different – the need for invasion science. *NeoBiota* 31: 87–98.
- WILSON M., BENSUSAN K., PEREZ C., TORRES J.L. 2011. First records of the exotic leafhopper *Sophonia orientalis* (MATSUMURA, 1912) (Hemiptera: Auchenorrhyncha: Cicadellidae) for the Iberian Peninsula and mainland Europe. *Boletín de la Sociedad Entomológica Aragonesa* 48: 435–436.
- YOTHERS M.A. 1934. Biology and control of Tree Hoppers injurious to fruit trees in the Pacific Northwest .United States Department Of Agriculture Washington.D.C. *Technical Bulletin* 40: 1–46.
- ZINA V., DA SILVA E.B., QUARTAU J.A., FRANCO J.C. 2013. First report of the citrus leafhopper *Penthimiola bella* (STÅL) (Hemiptera, Cicadellidae) in Europe. *Phytoparasitica* 41: 521–527.

Accepted: 13 December 2018; published: 24 December 2018

Licensed under a Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>