



## INDO AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH



### PREPARATION AND IN VITRO EVALUATION OF SOLID DISPERSION OF FINASTERIDE

**Jamal Ali Ashoor, Qasim Allawi Bader, Mohammed Fanokh Abo Khthar, Hasanain Shakir Mahmood**

*College of Pharmacy, Karbala University, Iraq.*

#### ARTICLE INFO

##### Article history

Received 19/04/2017

Available online  
10/05/2017

##### Keywords

FNS,  
HPMC,  
PEG 6000,  
SD.

#### ABSTRACT

Solubility is an important physiochemical element that affect the absorption and effectiveness of desired drug. Failure in formulation development of the drug has been considered as the consequences of its poor aqueous solubility. The reduced solubility of and low dissolution rate in aqueous gastrointestinal tract most often lead to inadequate bioavailability of the drug. In the present study the solubility and dissolution rate of finasteride which show poor solubility has been developed by solid dispersion(SD) technique with different polymers such as polyethylene glycol 6000 (PEG6000) and Hydroxypropyl Methyl Cellulose (HPMC). SD of finasteride (FNS) was prepared by using 60,120 and 200 mg of the polymer for 200 mg of the pure FNS in each formula using solvent evaporation method. Formula containing 200mg of HPMC showed best release rather than other formulas.

#### Corresponding author

**Jamal Ali Ashoor**

College of Pharmacy,  
Karbala University, Iraq.  
qasim\_bdr@yahoo.com

Please cite this article in press as **Jamal Ali Ashoor et al.** Preparation and In Vitro Evaluation of Solid Dispersion of Finasteride. *Indo American Journal of Pharmaceutical Research*.2017;7(04).

Copy right © 2017 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

The term "Solubility" can be defined as the ability of the solid, liquid or gaseous material (solute) to be dissolved in the solvent to make a homogeneous solution [1]. The solubility scope varies from the very soluble to practically insoluble. The term "insoluble" is used when the solute is poorly or very poorly soluble in solvent system [2]. Table 1 shows the main solubility description terms used by USP and BP. Biopharmaceutical Classification System (BCS) accounts three major factors that play significant role in controlling rate and extent of the drug absorption from the oral solid dosage forms which are the dissolution, the solubility and the intestinal permeability [3].


In view of pharmaceutical technology; particle size, molecular size, pressure, temperature, polarity and polymorph characteristics are the main factors that affect solubility [4].

Solid dispersion (SD) technique is one of the most important methods that was used to enhance water solubility of poorly soluble solid drugs in gastrointestinal fluids [5]. The term "solid dispersion" denotes the distribution of single or extra solid active substances (hydrophobic) in a solid matrix (Hydrophilic) both of them can be either in crystalline or amorphous form. Methanol, water, ethanol, chloroform and acetic acid are commonly used as solvents in solid dispersion [6].

Finasteride is synthesized 4-azasteroid compound. It is a particular inhibitor of steroid (type II 5 $\alpha$ - reductase) which is an endogenous enzyme that converts testosterone into 5 $\alpha$  dihydrotestosterone (DHT). For this reason such medication can be used for treatment of prostate gland enlargement [7]. Finasteride is white to off-white, crystalline solid. Melts at about 250°C. Soluble in organic solvents especially chloroform and alcohol; very slightly soluble in aqueous media; its empirical formula is C<sub>23</sub>H<sub>36</sub>N<sub>2</sub>O<sub>2</sub> with M.wt. 372.55 g/mol [8], water solubility about 0.05 mg/ml and 75 mg/ml in ethanol and soluble in methanol and chloroform. In accordance to (BCS); FNS is a class 2 drug (having low solubility and high permeability). It is a weakly acidic drug and has a pK<sub>a</sub> of 15.91. [9]. The main purpose of the research is to develop the aqueous solubility of finasteride by preparing it as SD with different ratios of polyethylene glycol 6000 (PEG6000) and Hydroxypropyl Methyl Cellulose (HPMC).

**Table 1. Solubility descriptive terms.**

Description	Parts of solvent required for one part of solute
Very soluble	Less than 1 portion
Freely soluble	1-10 portions
Soluble	10-30 portions
Sparingly soluble	30-100 portions
Slightly soluble	100-1000 portions
Very slightly soluble	1000-10000 portions
Practically insoluble	More than 10000



**Figure 1. Finasteride chemical structure.**

## MATERIALS

All the materials used in this research were from laboratory grade. FNS was purchased from a local agent of Heter Drug, India. HPMC and PEG6000 also purchased from a local agent for Himedia, India.

## METHODS

### Determination of FNS melting temperature:

Melting point of FNS was measured according to U.S.P by capillary tube procedure [10] by recording the temperature that liquefies the powder as the melting point.

### Determination of FNS $\lambda_{max}$ :

Finasteride solution was prepared by dissolving 10 mg FNS in 2 ml methanol and completed with chloroform up to 100 ml which was scanned by UV visible spectrophotometer (SPUV-26, Germany) from 200 to 400 nm and the maximum absorption of the drug was determined as its  $\lambda_{max}$ .

### Calibration curve of FNS

Was constructed by preparing a serial of dilution of finasteride with different concentration (15, 25, 50 µg/ml) from stock solution containing 10mg/100ml finasteride. The absorbance was then measured at the  $\lambda_{\max}$  of the drug. The measured absorbances were plotted against the respective concentrations.

### Solubility study:

Determination of the solubility is a vital factor for poorly water soluble drugs formulations. The saturated solutions of FNS were prepared by adding excess of the drug in distilled water and shaken by the shaker water bath for 24 h at 30°C, filtered by Whatman filter paper, diluted and analyzed by UV spectrophotometer at its  $\lambda_{\max}$ . This study was done in triplicate [11]

### Preparation of FNS solid dispersion

Formulas by solvent evaporation method[12]. The drug and diverse polymers in dissimilar ratios were dissolved in a measured volumes of water and methanol, poured into petri dish and placed in oven at 50°C overnight to complete evaporation of the solvent. The compositions of formulations are shown in table 2.

### Evaluation of finasteride SD

**Determination of Production Yield of SD** by computing the original weight of the solid components with the ultimate weight of the gained SD as seen in the equation below:

$$\text{SD yield} = \frac{\text{Practical weight of SD}}{\text{Theoretical weight (polymer + drug)}} \times 100$$

### Entrapment efficiency of finasteride SD:

Finasteride content in the SD was determined spectrophotometrically. A saturated mixtures were prepared by addition of excess FNS formula (finasteride+polymer) in distilled water shaking by the shaker water bath for 24h at 30°C. These solutions were clarified through filter paper then the filtrate was dried in oven at temperature 50°C overnight then added chloroform and spectrophotometric absorbance was measured at  $\lambda_{\max}$  of finasteride. The drug content was calculated from the calibration curve and expressed as percent entrapment efficiency as explained in the equation below.

$$\text{Entrapment efficiency} = \frac{\text{Actual weight of finasteride in SD}}{\text{Theoretical weight of finasteride}} \times 100$$

**Table 2. Composition of fenasteride SD formulations.**

Formula code	FNS	PEG6000	HPMC
1	200	60	
2	200	120	
3	200	200	
4	200		60
5	200		120
6	200		200

## RESULTS AND DISCUSSIONS

### Determination of FNS melting temperature

The recorded melting point of FNS was found to be 248°C. This outcome was as the same as references value [13], at the same time this result reflects the purity of the powder used in this study.

Determination  $\lambda_{\max}$  by scanning the solution which contain 10µg/ml of FNS in chloroform by (UV spectrophotometer) at 200-400 nm to produce the spectrum that shown in figure 2.




Figure 2. UV peak spectrum of finasteride.

Calibration curve of FNS in chloroform is shown in figure 3. A trend line was attained by conspiracy of the absorbance versus concentrations which specifies that the curve follows Beer's rule within the series of concentrations used.




Figure 3. Calibration curve of finasteride.

### Solubility study

Saturation Solubility is the extreme amount of a solute that dissolved in a specified solvent; such solution were prepared by adding extra quantities of FNS in distilled water and chloroform and shaken by shaker water bath for 10 min at 30 °C. Filtration of solution was performed to exclude any unsolvable particles that might disturb the results. The results of saturation solubility study are shown in table 3.

Table 3. Solubility data of pure FNS and finasteride SD.

Formula code	Solubility (µg/ml)
Pure drug	11.7
1	27.425
2	43.05
3	48.48
4	30.675
5	41.86
6	55.36

### Entrapment Efficiency

The Entrapment efficiency is an important measure for nano-particulate system. They give an impression about the % drug that successfully entrapped/absorbed in to nanoparticles and idea about production and scale-up capabilities and the encapsulation power of that particular technique. The solid dispersion method has good or excellent entrapment efficiency. The entrapment efficiency was measured to determine exactly how much FNS was entrapped in SD formula which is necessary to calculate the equivalent weight of FNS SD to that of marked product. The result is shown in table 4.

Table4. Percentage yield and entrapment efficiency of FNS SD formula.

Formula	Yield %	Entrapment efficiency%
1	42.3	13.3
2	58	21
3	60	23.65
4	48	15.33
5	61.25	20.93
6	76.25	27.68

## CONCLUSION


Preparation of FNS solid dispersion has been successfully achieved by using HPMC for enhancement the solubility of drug. In vitro solubility studies reveal that there is marked increased in the dissolution profile of F6 when compared to pure FNS.

## REFERENCES

1. L. Lachman, H. Lieberman, and J. L. Kanig, The Theory And Practise of Industrial Pharmacy, Lea & Febiger, 3rd edition, 1986.
2. M. Clugston and R. Fleming, Advanced Chemistry, Oxford Publishing, Oxford, UK, 1<sup>st</sup> edition, 2000.
3. Mehta M ,Biopharmaceutics Classification System (BCS): Development, Implementation, and Growth. Wiley 2016.
4. Sajid A., Chaudhary V.Solubility enhancement methods with importance of Hydrotrophy. Journal of drug discovery and therapeutics,2013; 2(6):96- 101
5. Arunachalam A., Karthikeyan M., Kishore K., Pottabathula H.P., Sethuraman S., Hutoshkumar S.A. Solid Dispersions. Current Pharma Research, 2010.
6. Patel P.B., Prajapati S.T., Patel N.R., Patel N.N., Patel C.N. Solid dispersion: a method of improving bioavailability and taste masking. Inventi Rapid: Pharm Tech, 2012 (4):1-7
7. Bull, H. G., Garcia-Calvo, M., Andersson, S.,: Journal of American chemistry society . 118:2359–2365,1996.
8. Azeem A, Khan ZI, Aqil M, Ahmad FJ, Khar RK, Talegaonkar S (May 2009). "Microemulsions as a surrogate carrier for dermal drug delivery". Drug Development and Industrial Pharmacy. 35 (5): 525–47.
9. Almeida, H. M.; Cabral Marques, H. Physicochemical characterization of finasteride: PEG 6000 and finasteride:Kollidon K25 solid dispersions, and finasteride:β-cyclodextrin inclusion complexes. J. Incl. Phenom. Macro. 2011, 70 (3–4), 397–406.
10. Neelam Indora, Dinesh Kaushik. Design, development and evaluation of ethosomal gel of fluconazole for topical fungal infection, International Journal of Engineering Science Invention Research & Development; Vol. I Issue VIII February 2015.
11. Narasimha Swamy Lakka, Nishant Goswami. Solubility and dissolution profile studies of gliclazide in pharmaceutical formulations by RP-HPLC, Int.Res.Jour. of Pharmacy 2012, 3 (6).
12. Rasenack N, Hartenhauer H, Müller B. Microcrystals for dissolution rate enhancement of poorly water-soluble drugs. Int J Pharm. 2003;254:137-45.
13. PROSCAR<sup>®</sup> Monograph by Merck&CO.,INC.Introduced for FDA Aproval . Issued October 2004 Printed in USA, p13.



54878478451170440



Submit your next manuscript to **IAJPR** and take advantage of:

- Convenient online manuscript submission
- Access Online first
- Double blind peer review policy
- International recognition
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in **ScopeMed** and other full-text repositories
- Redistributing your research freely

Submit your manuscript at: [editorinchief@iajpr.com](mailto:editorinchief@iajpr.com)

