

TITLE: Invasive Siberian chipmunks *Eutamias sibiricus* in Italy: a socio-ecological analysis indicates that they could, and should, be removed

AUTHORS: Jacopo Cerri<sup>1\*</sup>, Emiliano Mori<sup>2</sup>, Rudy Zozzoli<sup>3</sup>, Andrea Gigliotti<sup>4</sup>, Antony Chirco<sup>5</sup>, Sandro Bertolino<sup>6</sup>

#### AUTHORS' INSTITUTIONAL AFFILIATIONS:

1. Istituto Superiore per la Protezione e la Ricerca Ambientale, Piazzale dei Marmi 12, 57123, Livorno, Italy. Email: [jacopo.cerri@isprambiente.it](mailto:jacopo.cerri@isprambiente.it), mobile phone: (+39)3395692346. ORCID ID: 0000-0001-5030-0376. Contribution: major contribution in the conception and design of the study, altogether with the analysis and interpretation of data and with manuscript drafting;
2. Dipartimento di Scienze della Vita, Università degli Studi di Siena. Via P.A. Mattioli 4, 53100, Siena, Italy. Email: [moriemiliano@tiscali.it](mailto:moriemiliano@tiscali.it). Italy. ORCID ID: 0000-0001-8108-7950. Contribution: conception and design of the study, data collection and interpretation of the data. Important contribution in manuscript editing;
3. Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale, Università degli Studi di Parma, Parco Area delle Scienze, 11a, 43124, Parma, Italy. Email: [zozzoli.rudy@gmail.com](mailto:zozzoli.rudy@gmail.com). Contribution: conception and design of the study, data acquisition on the field, intellectual role in manuscript revision;
4. Dipartimento di Scienze delle Produzioni Animali e dell'Ambiente, Università degli Studi di Firenze, Piazzale delle Cascine, 18, 50144 Firenze, Italy. Email: [andrea.gigliotti1988@libero.it](mailto:andrea.gigliotti1988@libero.it). Contribution: data acquisition on the field, data interpretation, intellectual role in manuscript revision;

5. Dipartimento di Biologia, Università degli Studi di Firenze, Via Madonna del Piano, 6, 50019 Sesto Fiorentino FI. Email: [antony.chirco@stud.unifi.it](mailto:antony.chirco@stud.unifi.it). Contribution: data acquisition on the field, data interpretation, intellectual role in manuscript revision;
6. Department of Life Sciences and Systems Biology, University of Turin, Via Accademia Albertina 13, 10123 Torino, Italy. Email: [sandro.bertolino@unito.it](mailto:sandro.bertolino@unito.it). Telephone: (+39)0116704513. ORCID ID: 0000-0002-1063-8281. Contribution: conception and design of the study, analysis and interpretation of data and manuscript drafting;

CORRESPONDING AUTHOR: Jacopo Cerri, Istituto Superiore per la Protezione e la Ricerca Ambientale, Piazzale dei Marmi 12, 57123, Livorno, Italy. Email: [jacopo.cerri@isprambiente.it](mailto:jacopo.cerri@isprambiente.it), mobile phone: (+39)3395692346. ORCID ID: 0000-0001-5030-0376.

## ABSTRACT

**BACKGROUND:** Eradication of invasive alien species is a form of pest control linked to biodiversity conservation that usually involves killing animals. Squirrels are prominent among invasive alien species in Italy with four species introduced. Three of them are included within the list of alien species of European concern and their eradication and control is recommended. However, their local control is not an easy task, being highly appreciated by the general public. We propose a socio-ecological approach to evaluate the feasibility of eradicating Siberian chipmunks (*Eutamias sibiricus*) populations. We performed a structured questionnaire to assess the social perception of invasive Siberian chipmunks in urban parks where they occur and to identify groups of visitors who might oppose eradication. We also carried out geographic profiling to test for the spatial expansion of chipmunk populations.

**RESULTS:** Overall, park visitors regarded chipmunks positively and appreciated to see them, but human-chipmunk interactions were still rare. We did not identify any group of visitors with a strong attachment to chipmunks, who might oppose future control programs. Geographic profiling showed that chipmunks in Valeggio sul Mincio are starting to expand outside of their introduction site.

**CONCLUSIONS:** Data from questionnaires show that chipmunks eradication, coupled with adequate communication initiatives, might be feasible. Moreover, geographic profiling indicates that time for a rapid removal is running out. Socio-ecological approaches, combining the analysis of structured questionnaires administered to stakeholders and statistical modeling of pest observations, could be a valuable tool to decide the feasibility and the urgency of invasive pest control.

**Keywords:** Alien squirrels, eradication, geographic profiling, pest control, socio-ecological model

## 1 INTRODUCTION

The Siberian chipmunk *Eutamias sibiricus* is a widespread species in Russia and the Far East, which has become invasive in some European countries since the 1960s after its widespread trade as a pet species.<sup>1</sup> Chipmunks established viable populations in Italy<sup>1</sup>, chipmunks that escaped from captivity established four viable populations in Northern Italy, and at two urban parks in Rome. Siberian chipmunks are not a mainstream invader, as the Eastern grey squirrel *Sciurus carolinensis*;<sup>2</sup> however, they are considered an invasive alien species of European concern being listed within the EU Regulation 1143/2014. Particularly, chipmunks can act as a vector of tick-borne diseases and zoonoses.<sup>3,4</sup> The European regulation requires member states to eradicate listed species from their territories when it is still possible and this appears the situation in Italy where the species is still localized with small populations.<sup>5,6</sup>

Management interventions aimed at containing or removing invasive alien mammals are more feasible when two ideal conditions occur. First, they are more cost-effective and face a higher success rate whenever the target species are still in the early stages of their invasion.<sup>7-9</sup> Second, eradication initiatives tend to be more feasible when target species have minor interactions with society.<sup>10</sup> Attempts to remove iconic alien mammals could result in strong opposition from some stakeholders.<sup>11,12</sup> For instance, a trial eradication of the gray squirrel from Italy attempted in the 1990s was stopped by animal rights groups who brought the case in front of the court.<sup>13</sup>

Preliminary social impact assessments can tell managers whether control interventions will be opposed by relevant stakeholders and how they can be designed accordingly, to be successful.<sup>14,15</sup> Socio-ecological assessments go one step further, by combining information from relevant stakeholders, obtained through qualitative or quantitative methods from the social sciences, with information about the ecology, distribution and population dynamics of target species<sup>16</sup>. Ecological information can also be spatially explicit, as most ecological processes incorporate a geographical dimension<sup>17</sup>.

Eradicating chipmunks from Italy is required under the European and national legal framework, but, to the best of our knowledge, no study was conducted to verify if the preconditions for a successful eradication occur. Social prerequisites are particular sensitive considering the strong opposition faced by managers aiming to control or eradicate the gray squirrel in this country, even in recent years.<sup>13, 18, 19</sup>

Our research aims to fill this gap, by conduction a socio-ecological analysis combining spatial data of the species altogether with information from a structured surveys administered to a sample of visitors. Our analysis aims to test whether chipmunks became an iconic species and to inspect patterns in their geographical spread over time, to identify at which stage their invasion might be.

## 2 MATERIALS AND METHODS

In this research we focused on the three urban parks where chipmunks established viable populations in Italy. The first one is Sigurtà Garden Park, in Valeggio Sul Mincio, where Korean chipmunks were released in 1978, establishing the largest Italian populations of chipmunks in Italy.<sup>1, 20-22</sup> The latter two areas where two urban parks in Rome, Villa Ada, where chipmunks were introduced at multiple times since the early 1908s and Villa Doria-Pamphili, where chipmunks were observed for the first time in 2018.<sup>6</sup>

We surveyed a sample of visitors, administering a structured questionnaire measuring some their interactions with chipmunks and some psychological drivers of human-chipmunk interactions: attitudes, emotional dispositions, core affect, existence beliefs, social norms and behavioral intention about the presence of chipmunks. Attitudes were measured by means of a Likert scale and they were conceptualized as divided in some beliefs, characterized each one by its strength and the evaluation of its outcome.<sup>23</sup> The attitudinal scale was built up by considering all the potential impacts of a species of ground squirrels living in a park, after a pilot study (S1). Emotions were measured as emotional disposition (joy, fear, surprise, disgust, interest) and core affect, or the extent respondents would have felt positive or negative at the idea of encountering a chipmunk.<sup>24-27</sup> We

measured existence beliefs by asking respondents to rate the importance of chipmunks in the park, both for future generations and per se.<sup>28</sup> We measured social norms about the appropriateness of chipmunk presence in the park, by using three items measuring moral beliefs, empirical and normative expectations, and the willingness to enforce them by reporting the presence of chipmunks to local authorities.<sup>29</sup> Visitors were also asked whether they had ever heard of chipmunks living in the park and if they had ever seen, fed or touched them. A complete list of the various questions adopted in the questionnaire, altogether with their summary is available in Table 1 and a complete copy of the questionnaire at [https://docs.google.com/forms/d/e/1FAIpQLSelkWacunZsTWDB3Qxy3QC YvfaFKg-hhd9LFmhcLcEZAbyfuA/viewform?usp=sf\\_link](https://docs.google.com/forms/d/e/1FAIpQLSelkWacunZsTWDB3Qxy3QC YvfaFKg-hhd9LFmhcLcEZAbyfuA/viewform?usp=sf_link). The questionnaires were administered in Rome in both parks, but the collected data were grouped together as the two parks share the same pool of visitors. The questionnaire was implemented on GoogleForms. Most respondents (93.95%) were recruited on the field and they completed the questionnaire on a tablet. An online version was also administered on some Facebook groups on these urban parks. Questionnaires were confidential and they took approximately 15 minutes to be filled.

We assessed the reliability of our attitudinal scale through McDonald's Omega,<sup>30</sup> and we tested for construct validity through Confirmatory Factor Analysis (CFA), with a Maximum Likelihood estimator with robust standard errors and a Satorra-Bentler scaled test statistic. Both indicator and latent variables were standardized and all the factor loadings were estimated. We adopted two correlated latent variables reflecting the strength of each beliefs and the evaluation of its outcome and we also specified some residual correlations between each couple of items describing a specific impact. We selected the best subset of items and the best latent variable structure by comparing models through likelihood-ratio testing and some fitness indexes. Attitudes were aggregated into a final score by summing the product of each couple of items.<sup>23</sup>

We segmented respondents on the basis of their attitudes, emotional dispositions, core affect and their moral, empirical and normative expectations about the presence of chipmunks in the park.

Segmentation aimed to identify clusters of respondents who strongly supported chipmunk presence and could oppose their eradication. We tested for the presence of clusters in the data through the Hopkins index and we compared k-means, hierarchical and k-medoid cluster analysis<sup>31</sup> to assess which one clustered observations the best.

We carried out Generalized Linear Modeling with a Gamma distribution of the error and an identity link to highlight differences in the two areas, in terms of attitudes scores, core affect, existence beliefs and moral, empirical and normative expectations about the presence of chipmunks in the park. In each model we included a dummy variable assessing whether respondents encountered chipmunks at least once in their lifetime, and an interaction term, to account for the effect of past behavior and belief saliency.

Finally, we tested for chipmunk expansion outside of their introduction sites by means of geographic profiling (GP). GP is common in criminology, where the spatial locations of crimes are used to calculate the probability of occurrence of the offender's residence for each point over a certain geographical area. GP outperforms classical measures of spatial tendency, and many ecologists found it good for tracing back the origin of individuals that could move across space.<sup>32-35</sup> We adopted a Bayesian GP algorithm,<sup>36</sup> requiring only the specification of a distribution parameter, indicating a plausible maximum extent to which individuals could move. Based on available evidence indicating that chipmunks usually disperse within a few hundred meters from their birthplace,<sup>37-38</sup> we opted for a dispersal parameter of 1 km. We used available observations collected in Villa Ada, from 2011 to 2014 (n=26), and in Valeggio sul Mincio, from 1997 to 2018 (n=87), as the input for the GP algorithm. We did not use observations from Villa Doria-Pamphili as the park is embedded in an urbanized matrix, which prevents chipmunks from dispersing around. It is important to note that we were not interested to identify where chipmunks were released, but to reconstruct a probabilistic profile for the origin of the observations: the inspection of its shape told us whether observed chipmunks came from disjoint hotspots, as expected for an expanding invasive alien species, or from a single one, like in the case of a species which is not expanding. Statistical

analyses were carried out with the statistical software R (R Core Team 2018) and a detailed information about statistical analysis, altogether with a reproducible software code is available in the Supplementary Information (S2).

### 3 RESULTS AND DISCUSSION

Respondents had generally positive emotions towards chipmunks. Moreover, they generally agreed with the idea that the presence of chipmunks in the park was important for future generations and that it was important to have chipmunks living in the park even if one does not see them. Finally, most respondents deemed right and common for chipmunks to live in urban parks (Table 1). However, most respondents were not aware of the presence of chipmunks in the park where they were interviewed and about half of them had never observed these animals before. The proportion of respondents who had fed (14.6%) or touched (6.3%) chipmunks were even lower. Moreover, 14.11% of respondents reported to have observed chipmunks, despite they were not aware of their presence (Figure 1).

CFA and McDonald's Omega did not support an overall attitudinal construct, but they identified two separate groups of beliefs. The first one included items about the impact of chipmunks over the quality of recreation at the park: increasing the aesthetic appeal of the park, attracting new visitors and making visitors more prone to visit the park again. The second group included the potential impacts of chipmunks over human health: rummaging garbage from bins, transmitting disease to humans and to visitors' dogs (Table 2; Table 3).

Hierarchical cluster analysis with Euclidean distance and a complete link indicated the presence of a small segment of respondents, characterized by negative attitudes about chipmunk impacts over human health, fear and disgust towards chipmunks (Figure 2). These could be people who are scared or disgusted by rodents and concerned about their impact over hygiene, two aspects that are often related.<sup>39-41</sup>



The two sites differed only in respondents' score about the positive impact of chipmunks over the recreational experience, with Rome having slightly higher scores. Visitors in Rome also agreed slightly more with the idea that most people deemed appropriate for chipmunks to live in urban parks in Italy (Figure 3; Table 4).

Overall, these findings indicate that visitors regard chipmunks positively. Apart from a small cluster of respondents, most of them have positive emotions towards chipmunks, deem appropriate the fact that they live in an urban park and value their presence as something having an intrinsic value. However, this positive perception probably stems from a more general, favorable, disposition towards the presence of wildlife at urban areas. Respondents do not have a coherent system of attitudes about the presence of chipmunks, probably because their real interactions were limited: attitudes are shaped and reinforced by our everyday experience with a certain issue, that make it salient for ourselves.<sup>23,42</sup> On the other hand, visitors had stable beliefs about those impacts of chipmunks that could affect their recreational experience at the park, as well as fears about those impacts that could undermine hygiene. These two sub-dimensions probably indicate that visitors' beliefs are embedded in broader belief networks encompassing different, and more salient, topics.<sup>43</sup> For example, our respondents could have stable belief networks diseases, and they could have tied to them some of their beliefs about chipmunks. Framing experiments, where participants are primed to think about some precise topics and where the effect of this priming over beliefs is measured<sup>44</sup> might be a valuable tool to better investigate how human-wildlife interactions are embedded into broader nomological networks, and influenced by beliefs about relevant social issues. Framing experiments could also be used to test for attitude certainty and strength.<sup>45,46</sup>

The idea that respondents' attitudes were not grounded into experience is reinforced by the limited interactions between visitors and chipmunks: approximately, only half visitors observed chipmunks in the park, 15% of them fed chipmunks and only about 6% of them reported to have touched a chipmunk. Moreover, some visitors who observed chipmunks were not aware of their stable presence in the park: our questionnaire was arguably the first time they were introduced to this

aspect. These superficial interactions are also reflected by the low differences between the two areas. Respondents in Rome and in Valeggio sul Mincio had similar scores for almost all the psychological antecedents of their interactions with chipmunks. They showed minor differences only in their beliefs about the impact of chipmunks over the recreational experience, and in their normative beliefs about the presence of chipmunks in an urban green area. Moreover, hierarchical cluster analysis did not divide respondents into meaningful segments, on the basis of their attitudes, emotions, existence values and social norms towards chipmunks and it did not identify any group of strong supporters of chipmunks.

Taken together, these three points are important for the future management of chipmunks in the study area. Attitudes are an antecedent of human behavior and often a good barometer to forecast an eventual opposition to the management of invasive native<sup>23</sup> and introduced wildlife.<sup>47</sup> As visitors do not have stable attitudes and no segments of highly motivated visitors exist, it is reasonable to say that an eradication campaign would not face any strong opposition from local visitors. Chipmunks at the two sites do not seem to be an iconic species yet like the gray squirrel in many urban parks in the UK.<sup>48</sup> Their interactions with visitors, especially those creating emotional bindings, like feeding, are still limited. However, considering that respondents regard chipmunks favorably and that they value their presence as a legacy for the younger generations, we believe that eradication initiatives should be coupled with an adequate communication strategy, to avoid polarization and the 'backfire effect'. Considering that respondents from the two sites did not show any particular difference, communication actions might be similar for Rome and Valeggio sul Mincio.

Geographic profiling confirmed that invasive chipmunks disperse less than 500 m from the place where they are born. However, while chipmunk observations in Rome come from a single source, observations in Valeggio sul Mincio are likely to have involved individuals coming from two distinct spatial cores (Figure 4). One of these two cores was found to be outside of the boundaries of the urban park where the species was introduced: although slowly, chipmunks are expanding outside Sigurtà Garden Park, their introduction site in Valeggio sul Mincio. As chipmunks increase

their density we expect them to continue their expansion outside of the park. This perspective is not encouraging, because Valeggio sul Mincio is surrounded by a countryside environment and cultivations, that might promote chipmunks dispersal at the landscape scale<sup>20</sup> and maybe even their role as a pest species damaging crops, in the near future. Considered that chipmunks are still distributed over a relatively small area, but there are large areas in Italy suitable for the species<sup>49,50</sup>, we recommend their quick removal, as it will be easy and cost-effective.

#### 4 CONCLUSION

In this research, we showed how spatially-explicit data about a biological invasion and survey data about its social perception can inform decision makers about the feasibility, and the urgency, of management actions. Geographic profiling can be used not only to identify introduction, or to locate dens of invasive pests but also to signal the emergence of source-sink systems. These systems indicate the end of an early invasion stage and the spatial expansion of the invasive alien species, often due to their numerical increase, which can make eradication or control harder and expensive. Moreover, structured surveys could inform conservationists about the interactions between stakeholders and biological invaders, altogether with their social perception.

Our findings indicate that visitors still have limited interactions with invasive chipmunks, at the urban parks in Italy where they have been introduced. They do not have stable attitudes, and there seems not to be any group of visitors who regard chipmunks as an iconic species. Perspectives about chipmunks are positive, but probably weak. At the same time, chipmunks are expanding outside of their introduction site in Northern Italy. We deem that initiatives aimed at removing chipmunks are still feasible, if properly planned, and urgent. Postponing any management intervention could complicate eradications, both for the spatial spread of the species in one of the two areas and for the risk of change in visitors' attitude due to a greater confidence with chipmunks that have become more abundant.

## ACKNOWLEDGMENTS

The “U.O. Manutenzione e Valorizzazione del Verde Urbano” office of the Municipality of Roma provided us with permits to set the survey within urban parks in Rome. Authors would like to thank Davide Sogliani, who kindly helped in the fieldwork.

## 5 REFERENCES

1. Mori E , Zozzoli R, Menchetti M, Global distribution and status of introduced Siberian chipmunks *Eutamias sibiricus*. *Mammal Rev* **48**:139-152 (2018).
2. Bertolino S, Introduction of the American grey squirrel (*Sciurus carolinensis*) in Europe: a case study in biological invasion. *Curr Sci India* **95**:903-906 (2008).
3. Pisanu B, Marsot M, Marmet J, Chapuis JL, Reale D, Vourc'h G, Introduced Siberian chipmunks are more heavily infested by ixodid ticks than are native bank voles in a suburban forest in France. *Int J Parasitol* **40**:1277-1283 (2010).
4. Marsot M, Chapuis JL, Gasqui P, Dozières A, Massèglia S, Pisanu B, Ferquel E, Vourc'h G, Introduced Siberian chipmunks (*Tamias sibiricus barberi*) contribute more to Lyme borreliosis risk than native reservoir rodents. *PLoS One* **8**: e55377 (2013).
5. Benassi G, Bertolino S, Distribution and activity of the introduced Siberian chipmunk (Laxmann 1769) in an urban park in Rome, Italy. *Mammalia* **75**:87–90 (2011).
6. Zozzoli R, Menchetti M, Mori E, Spatial behaviour of an overlooked alien squirrel: the case of Siberian chipmunks *Eutamias sibiricus*. *Behav Process* **153**:107-111 (2018).
7. Pluess T, Cannon R, Jarošík V, Pergl J, Pyšek P, Bacher S, When are eradication campaigns successful? A test of common assumptions. *Biol Invasions* **14**:1365-1378 (2012).
8. Bomford M, O'Brien P, Eradication or control for vertebrate pests? *Wildlife Soc B* **23**:249-255 (1995).

9. Robertson PA, Adriaens T, Lambin X, Mill A, Roy S, Shuttleworth CM, Sutton-Croft M, The large-scale removal of mammalian invasive alien species in Northern Europe. *Pest Manag Sci* **73**:273-279 (2016).
10. McNeely JA, The great reshuffling: human dimensions of invasive alien species The great reshuffling: human dimensions of invasive alien species. Workshop on the Human Dimensions of Invasive Alien Species, Cape Town, ZA, 15-17 September 2000: 213-237 (2001).
11. Blackburn TM, Pettorelli N, Katzner T, Gompper ME, Mock K, Garner TWJ, Altwegg R, Redpath S, Gordon IJ , Dying for conservation: eradicating invasive alien species in the face of opposition. *Anim Conserv* **13**:227-228 (2010).
12. Shine R, Doody JS, Invasive species control: understanding conflicts between researchers and the general community. *Front Ecol Environ* **9**: 400-406 (2011).
13. Bertolino S, Genovesi P, Spread and attempted eradication of the grey squirrel (*Sciurus carolinensis*) in Italy, and consequences for the red squirrel (*Sciurus vulgaris*) in Eurasia. *Biol Conserv* **109**: 351-358 (2003).
14. Crowley SL, Hinchliffe S, McDonald RA, Conflict in invasive species management. *Front Ecol Environ* **15**: 133-141 (2017).
15. Crowley SL, Hinchliffe S, McDonald RA, Invasive species management will benefit from social impact assessment. *J Appl Ecol* **54**: 351-357 (2017).
16. Gálvez N, Guillera-Arroita G, St. John FA, Schüttler E, Macdonald DW, Davies ZG, A spatially integrated framework for assessing socioecological drivers of carnivore decline. *J Appl Ecol* **55**:1393-1405 (2018).
17. Struebig MJ, Linkie M, Deere NJ, Martyr DJ, Millyanawati B, Faulkner SC, Le Comber SC, Mangunjaya FM, Leader-Williams N, McKay JE, St. John FA, Addressing human-tiger conflict using socio-ecological information on tolerance and risk. *Nat Commun* **9**:3455 (2018).

18. Genovesi P, Bertolino S, Human dimension aspects in invasive alien species issues: the case of the failure of the grey squirrel eradication project in Italy. In: McNeely JA (Eds) *The Great Reshuffling: human dimensions of Invasive Alien Species*. IUCN, Gland, Switzerland and Cambridge, UK, Vi + 242 pp: 113-119 (2001).
19. La Morgia V, Paoloni D, Genovesi P, Eradicating the grey squirrel *Sciurus carolinensis* from urban areas: an innovative decision-making approach based on lessons learnt in Italy. *Pest Manag Sci* **73**:354-363 (2017).
20. Mori E, Zozzoli R, Mazza G, Coming in like a wrecking-ball: are native Eurasian red squirrels displacing invasive Siberian chipmunks? A study from an urban park. *Urban Ecosyst* **21**:975–981 (2018).
21. Mori E, Milanesi P, Menchetti M, Zozzoli R, Monaco A, Capizzi D, Nerva L, Genetics reveals that free-ranging chipmunks introduced to Italy have multiple origins. *Hystrix* DOI: 10.4404/hystrix-00116-2018 (2018).
22. Mori E, Menchetti M, Zozzoli R, Milanesi P, The importance of taxonomy in species distribution models at a global scale: the case of an overlooked alien squirrel facing taxonomic revision. *J Zool* **307**:43-52 (2019).
23. Manfredo MJ, *Who cares about wildlife?* Springer, New York, NY, USA: 1-2 (2008).
24. Jacobs MH, Human emotions toward wildlife. *Hum Dimens Wildl* **17**:1-3 (2012).
25. Jacobs MH, Vaske JJ, Roemer JM, Toward a mental systems approach to human relationships with wildlife: the role of emotional dispositions. *Hum Dimens Wildl* **17**:4-15 (2012).
26. Jacobs MH, Vaske JJ, Dubois S, Fehres P, More than fear: role of emotions in acceptability of lethal control of wolves. *Eur J Wildlife Res* **60**:589-598 (2014).
27. Larson LR, Cooper CB, Hauber ME, Emotions as drivers of wildlife stewardship behavior: Examining citizen science nest monitors' responses to invasive house sparrows. *Hum Dimens Wildl* **21**:18-33 (2016).

28. Frank B, Glikman JA, Sutherland M, Bath AJ, Predictors of extreme negative feelings toward coyote in Newfoundland. *Hum Dimens Wildl* **21**:297-310 (2016).
29. Bicchieri C, Norms in the wild: How to diagnose, measure, and change social norms. Oxford University Press, Oxford, UK (2016).
30. Dunn TJ, Baguley T, Brunsten V, From alpha to omega: A practical solution to the pervasive problem of internal consistency estimation. *Brit J Psychol* **105**:399-412 (2014).
31. Kassambara A, Practical guide to cluster analysis in R. CreateSpace: North Charleston, South Carolina, USA (2017).
32. Raine NE, Rossmo DK, Le Comber SC, Geographic profiling applied to testing models of bumble-bee foraging. *J R Soc Interface* **6**:307-319 (2008).
33. Martin RA, Rossmo DK, Hammerschlag N, Hunting patterns and geographic profiling of white shark predation. *Journal of Zoology (London)* **279**: 111-118 (2009).
34. Faulkner SC, Stevenson MD, Verity R, Mustari AH, Semple S, Tosh DG, Le Comber SC, Using geographic profiling to locate elusive nocturnal animals: a case study with spectral tarsiers. *J Zool* **295**: 261-268 (2015).
35. Faulkner SC, Verity R, Roberts D, Roy SS, Robertson PA, Stevenson MD, Le Comber SC, Using geographic profiling to compare the value of sightings vs trap data in a biological invasion. *Divers Distrib* **23**:104-112 (2017).
36. Verity R, Stevenson MD, Rossmo DK, Nichols RA, Le Comber SC, Spatial targeting of infectious disease control: identifying multiple, unknown sources. *Methods Ecol Evol* **5**:647-655 (2014).
37. Marmet J, Pisanu B, Chapuis JL, Home range, range overlap, and site fidelity of introduced Siberian chipmunks in a suburban French forest. *Eur J Wildlife Res* **55**:497-504 (2009).
38. Marmet J, Pisanu B, Chapuis JL, Natal dispersal of introduced Siberian chipmunks, *Tamias sibiricus*, in a suburban forest. *J Ethol* **29**:23-29 (2012).

39. Davey GC, Macdonald BA, Brierley L, The effect of disgust on anxiety ratings to fear-relevant, disgust-relevant and fear-irrelevant stimuli. *J Anxiety Disord* **22**:1347-1354 (2008).
40. Curtis V, Aunger R, Rabie T, Evidence that disgust evolved to protect from risk of disease. *P R Soc B-Biol Sci* **271**:S131-S133 (2004).
41. Prokop P, Usak M, Fančovičová J, Risk of parasite transmission influences perceived vulnerability to disease and perceived danger of disease-relevant animals. *Behav Process* **85**:52-57 (2010).
42. Heberlein TA, *Navigating Environmental Attitudes*. Oxford University Press, Oxford, UK (2012).
43. Nilsson NJ, *Understanding beliefs*. MIT (Massachusetts Institute of Technology) Press, Cambridge, Massachusetts, USA (2014).
44. Chong D, Druckman JN, Framing theory. *Annu Rev Polit Sci* **10**:103-126 (2007).
45. Howe LC, Krosnick JA, Attitude strength. *Annu Rev Psychol* **68**:327-351 (2017).
46. Rucker DD, Petty RE, Briñol P, What's in a frame anyway? A meta-cognitive analysis of the impact of one versus two sided message framing on attitude certainty. *J Consum Psychol* **18**:137-149 (2008).
47. Sharp RL, Larson LR, Green GT, Factors influencing public preferences for invasive alien species management. *Biol Conserv* **144**:2097-2104 (2011).
48. Dunn M, Marzano M, Forster J, Gill RM, Public attitudes towards “pest” management: Perceptions on squirrel management strategies in the UK. *Biol Conserv* **222**:52-63 (2018).
49. Di Febbraro M, Martinoli A, Russo D, Preatoni D, Bertolino S, Modelling the effects of climate change on the risk of invasion by alien squirrels. *Hystrix* **27**:22–29 (2016).
50. Di Febbraro M, Menchetti M, Russo D, Ancillotto L, Aloise G, Roscioni F, Preatoni DG, Loy A, Martinoli A, Bertolino S, Mori E, Integrating climate and land–use change scenarios in modelling the future spread of invasive squirrels in Italy. *Divers Distrib*, DOI: 10.1111/ddi.12890 (2019).



## Figure legends

Figure 1. Sankey plot about visitors-chipmunks interactions.

Figure 2. Dendrogram of the hierarchical cluster analysis.

Figure 3. Marginal effects of the GLMs: average scores of the two areas.

Figure 4. Location of the three study sites (left), heatmap with the posterior probabilities of the origins of observed tamias (center) and posterior probability of the dispersal parameter (right).

## TABLES

Table 1. Questionnaire structure, item summary and completion rate.

Construct	Format	Question	Mean ± SD
Belief strength	Bipolar scale, from “Strongly disagree” (-3) to “Strongly agree” (+3)	<i>Nowadays the presence of chipmunks in an urban park would...</i>	4.55 ± 1.57
		... improve its overall environmental quality.	
		... make it more aesthetically appealing.	5.14 ± 1.50
		... lead people who already visited the park to return there more often.	5.35 ± 1.34
		... attract new visitors.	5.43 ± 1.31
		... make visiting the park a more satisfying experience.	5.69 ± 1.39
		... cause disease transmission to visitors.	2.80 ± 1.64
		... cause disease transmission to the dogs which are walked in the park.	2.95 ± 1.48
		... cause damaging to the vegetation of the park.	2.96 ± 1.57
		... cause nest depredation on those birds that breed there.	3.38 ± 1.68
... affect garbage disposal, as chipmunks could rummage it around while searching for food.	3.12 ± 1.58		
Evaluation of each belief	Unipolar scale from with 7 degrees, from “Bad” to “Good”	<i>“How would you evaluate the following consequences of chipmunks presence in the park?”</i>	5.73 ± 1.86
		The park improves its environmental quality.	
		The park becomes more aesthetically appealing.	6.02 ± 1.40
		People who already visited the park, now return there	5.96 ± 1.48

		more often.	
		New people come to see the park.	5.93 ± 1.30
		Visiting the park becomes a more satisfying experience.	5.76 ± 1.41
		Chipmunks transmit diseases to visitors.	1.79 ± 1.22
		Chipmunks transmit diseases to the dogs which are walked in the park.	1.91 ± 1.18
		The vegetation of the park is damaged by chipmunks.	2.29 ± 1.43
		Chipmunks prey on the nests of the birds that breed in the park.	2.33 ± 1.34
		Chipmunks rummage the garbage around garbage bins, while searching for food.	2.40 ± 1.19
Core affect	Bipolar scale, from “Extremely negative” (-3) to “Extremely positive ” (+3)	<p><i>“Use the following scale to describe how you would feel about living each one of these experience”.</i></p> <p>Spotting one, or more, chipmunks, while strolling in the park.</p>	4.03 ± 0.96
Emotional disposition (Happiness)	Bipolar scale, from “Extremely negative” (-3) to “Extremely positive ” (+3)	<p><i>“Could you please tell us to what extend you agree or disagree with the following statements?”</i></p> <p>I would feel happy, if I spotted some chipmunks when I go to the park. (Happiness)</p>	5.51 ± 1.48
		I would feel scared, if I spotted some chipmunks when I go to the park. (Fear)	1.45 ± 1.11
		I would be surprised, if I spotted some chipmunks when I go to the park. (Surprise)	4.24 ± 1.93
		I would feel disgusted, if I spotted some chipmunks when I go to the park. (Disgust)	1.54 ± 1.18
		I would feel interested, if I spotted some chipmunks	5.48 ± 1.54

		when I go to the park. (Interest)	
Existence beliefs	Bipolar scale, from “Extremely negative” (-3) to “Extremely positive ” (+3)	<p><i>“To what extend do you agree with the following statements?”</i></p> <p>It is important to maintain the presence of chipmunks in the park, for future generations.</p>	5.25 ± 1.83
		<p>Whether or not I would get to see chipmunks in the park, it is important to me that they live there.</p>	5.12 ± 1.74
Moral beliefs	Bipolar scale, from “Extremely negative” (-3) to “Extremely positive ” (+3)	<p><i>“To what extend do you agree with the following statements?”</i></p> <p>Chipmunks should not be allowed to live in an urban green space, like a park.</p>	5.28 ± 1.60
Empirical expectations	Bipolar scale, from “Extremely negative” (-3) to “Extremely positive ” (+3)	<p><i>“To what extend do you agree with the following statements?”</i></p> <p>I expect that most chipmunks living in Italy inhabit urban parks; (empirical expectations).</p>	4.94 ± 1.50
Normative expectations	Bipolar scale, from “Extremely negative” (-3) to “Extremely positive ” (+3)	<p><i>“To what extend do you agree with the following statements?”</i></p> <p>I expect most people to believe that chipmunks should be allowed to lived in an urban green space, like the park; (normative expectations).</p>	5.13 ± 1.37
Perceived willingness to report chipmunks presence to local authorities (other people)	Dichotomous (Yes/No)	I expect most people to report chipmunks to local authorities, if they found they were living in an urban park.	Yes = 4.03%, No = 95.67%
Personal comittment to report chipmunks presence to local	Dichotomous (Yes/No)	I would report chipmunks to local authorities, if they found they were living in an urban park.	Yes = 9.27%, No = 90.73%

authorities			
Awareness about the presence of chipmunks	Dichotomous (Yes/No)	Prior to this survey, have you ever heard that some chipmunks inhabited the park?	Yes = 41.12%, No = 58.87%
Experience with chipmunks	Ordered scale (Multiple times per week; Multiple times per month; Once; Never)	“ <i>Could you please tell us how often did you have the following experiences at the park?</i> ”	Never = 50.40% Once = 27.82% Multiple times per month = 16.12% Multiple times per week = 5.64%
		Spotting one, or more, chipmunks at the park.	
		Feeding the chipmunks at the park.	Never = 85.48% Once = 10.09% Multiple times per month = 2.82% Multiple times per week = 1.61%
		Touching, or caressing, a chipmunks at the park.	Never = 94.76% Once = 4.03% Multiple times per month = 0.40% Multiple times per week = 0.80%
Park usage	Dichotomous (Yes/No)	“ <i>Could you please tell us which of the following activities you usually practice at the park?</i> ”	Yes = 0.28, No = 0.72
		Sport (e.g jogging, cycling, football, volley).	
		Birdwatching.	Yes = 6.85, No = 93.15%
		Strolling.	Yes = 79.43%, No = 20.57%
		Picnicking.	Yes = 22.98%, No = 77.02%
		Walking the dog.	Yes = 24.19%, No = 75.81%
		Painting / photography.	Yes = 25.80%, No = 74.19%
		Playing a musical instrument.	Yes = 4.43%, No = 93.57%
		Feeding animals.	Yes = 6.04%, No = 93.96%
Working or volunteering.	Yes = 2.82%, No = 97.18%		
Age	Ordered scale (15-20 years, 21-30 years, 31-40 years, 41-50 years, 51-65 years, more than 65 years )	Could you please tell us in which year were you born?	15-20 years = 10.5%, 21-30 years = 29.44 % 31-40 years = 22.98 %

			41-50 years = 19.75 % 51-65 years = 10.88 % more than 65 years = 6.5%
Sex	Dichotomous (Male/Female)	<i>We know that sex and gender are not the same concept. Could please just tell us your sex?</i>	Male = 50.81% Female = 49.29%
Education	Ordered scale	<i>Which is your level of education?</i>	Elementary school = 1.61% Secondary school = 8.47 % High school = 41.94 % University degree = 47.98%

Table 2. Confirmatory Factor Analysis, model comparison.

Beliefs about chipmunks' impact over the recreational quality of the park.			
	Full model	Model without the items about the "Environmental quality"	Model without the items about the "Environmental quality" and the "Visitors' satisfaction"
SRMR	0.04213852	0.03863779	0.03520523
RMR	0.04196861	0.03848199	0.03506327
TLI	0.9477260	0.9502502	0.9739350
CFI	0.9663123	0.9733483	0.9913117
IFI	0.9666189	0.9735970	0.9914187
ECVI	0.5632614	0.3781471	0.1772727
BIC	5471.809	4381.852	3500.436
AIC	5380.460	4308.070	3444.221
BIC2	5389.389	4315.282	3449.716
GFI	0.9341394	0.9495378	0.9838887
AGFI	0.8750919	0.8788907	0.9323327
PGFI	0.4925462	0.3956407	0.2342592
MFI	0.8884080	0.9285285	0.9860585
Scaled chi-squared difference test	df = 5, AIC = 3444.2, BIC = 3500.4, chisq = 11.964	df = 15, AIC = 4308.1, BIC = 4381.9, chisq = 51.781, chisq.diff = 28.519, df.diff = 10, p-value = 0.00149	df = 29, AIC = 5380.5, BIC = 5471.8, chisq = 87.689, chisq.diff = 24.250, df.diff = 14, p-value = 0.04274
Cronbach's alpha	0.93	0.92	0.88
McDonald's Omega	Omega Hierarchical = 0.74, Omega H asymptotic = 0.78, Omega Total = 0.95	Omega Hierarchical = 0.76, Omega H asymptotic = 0.81, Omega Total = 0.95	Omega Hierarchical = 0.68, Omega H asymptotic = 0.73, Omega Total = 0.93
Beliefs about chipmunks' impact over the hygiene of the park.			
	Full model	Model without the items about the "Vegetation damaging"	Model without the items about the "Nest depredation"
SRMR	0.06205949	0.05537003	0.03495205
RMR	0.06180925	0.05514676	0.03481111
TLI	0.8835495	0.9502502	0.9742254
CFI	0.9249541	0.9733483	0.9914085
IFI	0.9265352	0.9735970	0.9916257

ECVI	0.5515900	0.3781471	0.1625515
BIC	6467.589	4381.852	3912.691
AIC	6376.239	4308.070	3856.477
BIC2	6385.168	4315.282	3861.971
GFI	0.9386140	0.9495378	0.9888453
AGFI	0.9386140	0.9495378	0.9888453
PGFI	0.4949056	0.3956407	0.2354394
MFI	0.8936076	0.9285285	0.9933433
Scaled chi-squared difference test	df = 5, AIC = 3856.5, BIC = 3912.7, chisq = 8.3128	df = 15, AIC = 5146.0, BIC = 5219.8, chisq = 32.8162 , chisq.diff = 16.867, df.diff = 10 , p-value = 0.077360	df = 29, AIC = 6376.2, BIC = 6467.6, chisq = 84.7943, chisq.diff = 37.521, df.diff = 14, p-value = 0.000615
Cronbach's alpha	0.79	0.74	0.74
McDonald's Omega	Omega Hierarchical = 0.4, Omega H asymptotic = 0.47, Omega Total = 0.85	Omega Hierarchical = 0.42, Omega H asymptotic = 0.51, Omega Total = 0.82	Omega Hierarchical = 0.43, Omega H asymptotic = 0.52, Omega Total = 0.83



Table 3. Confirmatory Factor Analysis: factor loadings, covariances, variances and communalities.


Beliefs about chipmunks' impact over the recreational quality of the park.					
Factor loadings					
		Estimate	S.E	z-value	p-value
Strength of the belief	Aesthetic quality	0.697	0.076	9.188	0.000
	Increased visits in the near future	0.809	0.076	10.589	0.000
	Increase in the number of visitors	0.799	0.081	9.850	0.000
Evaluation of the outcomes	Aesthetic quality	0.874	0.084	10.409	0.000
	Increased visits in the near future	0.840	0.076	10.999	0.000
	Increase in the number of visitors	0.840	0.093	9.036	0.000
Covariances between the strength of each belief and the evaluation of its outcome					
	Latent constructs (Strength x Evaluation)	0.684	0.070	9.807	0.000
	Aesthetic quality	0.017	0.041	0.408	0.683
	Increased visits in the near future	0.093	0.041	2.300	0.021
	Increase in the number of visitors	0.036	0.040	0.893	0.372
Variances					
Strength of the belief	Aesthetic quality	0.510	0.086	5.951	0.000
	Increased visits in the near future	0.340	0.073	4.625	0.000
	Increase in the number of visitors	0.357	0.068	5.250	0.000
Evaluation of the	Aesthetic	0.227	0.074	3.058	0.002

outcomes	quality				
	Increased visits in the near future	0.298	0.059	5.037	0.000
	Increase in the number of visitors	0.296	0.067	4.410	0.000
Communalities					
Strength of the belief: aesthetic quality = 0.4879791	Strength of the belief: increased visits in the near future = 0.6580856	Strength of the belief: increase in the number of visitors = 0.6409658	Evaluation of the outcomes: aesthetic quality = 0.7709439	Evaluation of the outcomes: increased visits in the near future = 0.7031584	Evaluation of the outcomes: increase in the number of visitors = 0.7043100
Beliefs about chipmunks' impact over the hygiene of the park.					
Factor loadings					
		Estimate	S.E	z-value	p-value
Strength of the belief	Disease transmission to visitors	0.878	0.067	13.016	0.000
	Disease transmission to visitors' dogs	0.771	0.062	12.407	0.000
	Garbage rummaging	0.540	0.070	7.672	0.000
Evaluation of the outcomes	Disease transmission to visitors	0.678	0.089	7.664	0.000
	Disease transmission to visitors' dogs	0.865	0.096	9.025	0.000
	Garbage rummaging	0.438	0.096	4.543	0.000
Covariances between the strength of each belief and the evaluation of its outcome					
	Latent constructs (Strength x Evaluation)	0.335	0.084	3.992	0.000
	Disease	0.039	0.042	0.929	0.353


	transmission to visitors				
	Disease transmission to visitors' dogs	-0.025	0.042	-0.596	0.551
	Garbage rummaging	0.057	0.067	0.849	0.396
Variances					
Strength of the belief	Disease transmission to visitors	0.228	0.089	2.553	0.011
	Disease transmission to visitors' dogs	0.404	0.076	5.308	0.000
	Garbage rummaging	0.709	0.077	9.175	0.000
Evaluation of the outcomes	Disease transmission to visitors	0.532	0.153	3.479	0.001
	Disease transmission to visitors' dogs	0.243	0.130	1.870	0.061
	Garbage rummaging	0.802	0.109	7.385	0.000
Communalities					
Strength of the belief: disease transmission to visitors = 0.7721118	Strength of the belief: disease transmission to visitors' dogs = 0.5953521	Strength of the belief: garbage rummaging = 0.2915509	Evaluation of the outcomes: disease transmission to visitors = 0.4637312	Evaluation of the outcomes: disease transmission to visitors' dogs = 0.7546028	Evaluation of the outcomes: garbage rummaging = 0.1927527

Table 4. Output of the Generalized Linear Model

Response variable: positive impacts (aesthetic quality, number of visitors, future visits)				
Predictor	Estimate	S.E.	t-value	p-value
Intercept	110.444	4.887	22.598	< 2e-16 ***
Location	-16.130	6.109	-2.640	0.00882 **
Previous experience with chipmunks	-14.880	6.975	-2.133	0.03389 *
Location * Previous experience	10.072	8.488	1.187	0.23656
Null deviance = 119.71		Residual deviance = 114.77		AIC = 1887
Response variable: normative expectations about chipmunks presence.				
Predictor	Estimate	S.E.	t-value	p-value
Intercept	5.57407	0.19862	28.063	<2e-16 ***
Location	-0.64550	0.25149	-2.567	0.0109 *
Previous experience with chipmunks	0.04131	0.30804	0.134	0.8934
Location * Previous experience	-0.16988	0.37050	-0.459	0.6470
Null deviance = 24.126		Residual deviance = 22.937		AIC = 914.39




## Cluster dendrogram with AU/BP values (%)



Distance: euclidean  
Cluster method: complete

Positive impacts (n. visitors,  
more visits, aesthetics)



Normative expectations

