





Playing with white dwarfs in MESA: Three test cases

Ken Shen (UC Berkeley)


TAs: Jared Brooks & Evan Bauer



Download the materials!

- <http://mesastar.org/teaching-materials/2015-mesa-summer-school/>
- Lecture PDF, minilab #1, minilab #2, maxilab, and maxilab solutions

White dwarfs: Endpoints of stellar evolution for most stars



What are they? Stars held up by electron degeneracy

- Electrons are fermions, can't pack them into the same states: $P_{\text{degeneracy}} \propto n_e^{5/3}$
- Hydrostatic equilibrium: $\frac{dP}{dr} = -\rho \frac{Gm(r)}{r^2}$
- Order of magnitude scaling: $\frac{P}{R} \propto \rho \frac{M}{R^2}$
- Substitute degenerate equation of state: $\rho^{2/3} \propto \frac{M}{R}$
- White dwarf mass-radius relation: $R \propto M^{-1/3}$

Minilab #1: Test the WD mass-radius relation

- In this lab, we'll specify a mass and composition, turn off nuclear burning, and let the resulting star cool down until it becomes a C/O white dwarf. Then, we'll see how our **MESA** models do vs. the simplistic mass-radius relation.
- For various reasons, we can't immediately change the composition of the starting star to C/O. The model star has to be fairly degenerate before we can do that.
 - So, we have to first let it cool down until it reaches some degeneracy criterion and save the model. When we restart it, we can change the composition.


Minilab #1: Test the WD mass-radius relation

- The first step is to read through `inlist_makepurewd`. In its current state, it will start with a solar composition main sequence model of a mass you specify between 0.5 and 1.1 Msol, and cool with nuclear burning turned off until the center reaches $\eta = 5$, where η is the normalized electron chemical potential. This is accomplished by the `eta_center_limit` flag.
- So, choose a random mass, compile, and let it run! When the run ends, it will save a degenerate solar composition star as `intermediate.mod`.

Minilab #1: Test the WD mass-radius relation


- Now that we have a solar composition degenerate model, we can change its composition by turning on the `relax_initial_to_xaccrete` flag. This relaxes to the composition given in the `controls` part of the `inlist`.
- We no longer want to stop at an η limit, so turn off that flag. But we do want to stop when the central temperature reaches 2×10^7 K, so turn on that flag.
- Once the relevant flags are turned on and off, including the `load_saved_model` flag, restart with `./rn`. (The composition flag doesn't work on a `./re` from a photostep, which is why we have to `./rn` from the intermediate saved model.)
- Once the WD cools down and the run terminates, report your mass in M_{sol} units and ending radius in R_{sol} units on the Google spreadsheet (available at http://cococubed.asu.edu/mesa_summer_school_2015/agenda.html), and we'll generate a real-time white dwarf mass-radius relation. (You can just read the radius off of the terminal or `pgstar` output, remembering that the reported quantity is $\log_{10}(R/R_{\text{sol}})$, not R/R_{sol} .)

Pure composition white dwarf mass-radius relation with MESA




Creating white dwarfs in close binary systems

- Stars expand as they evolve onto the giant branch




Creating white dwarfs in close binary systems

- Expansion causes giant to overflow Roche lobe (equipotential surface through L_1), leading to mass transfer



Creating white dwarfs in close binary systems

- Mass transfer from deep convective envelope tends to run away faster than companion can accrete → “common envelope” phase
- Orbital binding energy used to eject this envelope → orbital separation shrinks
- Not well understood, but required to explain close WD systems (cataclysmic variables, double white dwarf binaries, etc.)



(Yungelson '05)

Minilab #2: Create a white dwarf by mimicking a common envelope phase

- For this lab, we'll make a more realistic C/O white dwarf by mimicking a star in a binary system that overflows its Roche lobe when it's on the asymptotic giant branch and loses its hydrogen-rich envelope in a common envelope.
- We'll do this by starting with a main sequence model, rapidly removing mass once the star reaches a certain carbon core mass until the radius shrinks, and then letting the resulting hot core cool off.
- The first step is to read through `inlist_makewdviaRLO`. We'll assign each table an initial main sequence mass and associated carbon core mass limit. Within each table, you'll perform a small convergence study by changing `varcontrol_target` and `mesh_delta_coeff` by factors of 2 or so. Those with faster computers should make these values smaller and vice versa.
- Remember to compile at the beginning!
- When the star's carbon-rich core runs into the `c_core_mass_limit` limit, the run ends, and **MESA** will save the file as **AGB.mod**.


Minilab #2: Create a white dwarf by mimicking a common envelope phase

- Now we want to rapidly remove mass. First, we have to turn off the `c_core_mass_limit` flag and turn on the Roche lobe overflow flags. These remove mass with the given parameters until the photosphere shrinks within `photosphere_r_lower_limit`.
- Then, restart from `AGB.mod` and save the model at the end of this phase as a different model, `donewithRLO.mod`. To do so, turn on and off the appropriate flags in `star_job`, and `./rn`.


Minilab #2: Create a white dwarf by mimicking a common envelope phase

- Now we have a hot exposed AGB core that is starting to cool. Turn off the Roche lobe overflow flags, and turn on the last stopping condition so that the run will stop when the central temperature reaches 2×10^7 K.
- Restart from `donewithRLO.mod`, and wait for the run to complete!
- You'll notice that the resulting C/O WD is not 50/50 C/O like the simple WDs we made in the first minilab. The helium shell is not able to burn completely, so there is a remnant layer of helium on top. Find the `total_mass_he4` at the end of the run in `LOGS/history.data`. Report your total ^4He mass and the final total WD mass on the Google spreadsheet.
- (The terminal outputs masses for `He_core` and `C_core`, which denote the outer mass extent of the He- and C-rich cores, respectively. Due to the shape of the helium abundance profile, the difference between the two is not the same as `total_mass_he4`. If you're waiting for others to finish, find out what **MESA** uses to determine the location of the He- and C-rich cores.)
- (Another note: when you're doing production research runs, turning flags on and off like this is probably not the best thing to do. Instead, you can have different project inlists and reference them from your main inlist.)


Remnant surface helium layer on C/O WDs



Double white dwarf merger evolution



- Double white dwarf binaries emit gravitational waves, spiral closer





(Image credit: Strohmayer & Berry)


(Yungelson '05)

Double white dwarf merger evolution


- Inverse mass-radius relation means less massive WD fills its Roche lobe first
- Mass transfer between WDs usually dynamically unstable





Double white dwarf merger evolution




(Dan+ '11)



Double white dwarf merger evolution



Double white dwarf merger evolution



Maxilab: Constructing and evolving a double white dwarf merger remnant

- The goal of this lab is to construct the remnant of a double white dwarf merger after it has dissipated its rotational kinetic energy and become quasi-spherical. This process converts the disrupted less massive WD into a hot extended envelope on top of the (more or less) undisturbed more massive WD.
- We'll begin by modeling a 0.2 + 0.3 Msol double helium WD merger, so we'll need to construct a 0.5 Msol pure helium object, with a 0.3 Msol degenerate core and a 0.2 Msol puffy envelope that does not exert much pressure on the core.
 - We'll relax the composition using flags in `star_job` and `controls`, similar to what we did in the first minilab. Some of the flags in `controls` are missing, so you will have to fill them in.
 - Entropy relaxation is currently not a standard option in MESA, so we'll have to write our own entropy relaxation routine in `run_star_extras.f`.

Maxilab: Constructing and evolving a double white dwarf merger remnant

- Open `run_star_extras.f` and read through the new energy routine.
- Outside of $M_{\text{transition}} = 0.3 M_{\text{sol}}$, the target entropy is set to something high so that the outer $0.2 M_{\text{sol}}$ gets expanded out.
- Inside of $M_{\text{transition}}$, we want the inner material to approximate an isothermal degenerate WD core. But we have to do this relaxation in two steps. This is because of the negative specific heat of ideal gas stars. When you add heat to a star like the Sun, it expands, and the temperature drops! So cooling the inner core actually makes the temperature increase at first.
 - Instead, we first cool the inner material to a low entropy until it becomes degenerate.
 - When it becomes degenerate “enough”, the specific heat becomes what you’re used to in everyday life, and cooling makes the temperature drop. We parametrize how degenerate is “enough” with $\eta_{\text{transition}}$, where η measures the ratio of the electron chemical potential to the thermal energy.
 - Once the inner material becomes degenerate, we can set a target temperature instead of entropy. This command is left blank for you to fill in.
 - (The `exp_cr` is the exponential function from `crlib` so that the answer is bit-for-bit.)

Maxilab: Constructing and evolving a double white dwarf merger remnant

- The relaxation equation I'm using is not fundamental, but it has the spirit of an entropy equation

$$\epsilon_{\text{heat}} = T \frac{ds}{dt} \sim T \left(\frac{s_{\text{target}} - s}{t_{\text{relax}}} \right)$$

- And something similar for the temperature part. Not physically important, but it helps to use something with the right units to get a roughly correct order of magnitude for easier convergence.

Maxilab: Constructing and evolving a double white dwarf merger remnant

- Once everything is in place, including filling in the missing inlist commands, `./mk` and `./rn`.
- If everything goes smoothly, save the model when it is relaxed and restart from it with the appropriate flags turned on and off. But how do you know when it is relaxed?
 - One thing to check is that the extra heat being added becomes very small.
 - So, make a new profile plot window with `inlist_pgstar` showing the extra heat being added vs. mass.
 - After the model reaches a relaxed state, you should see the extra heat being ~ 0 , except for very near the boundary between the hot envelope and cold core. That's because the relaxation routine is counteracting thermal diffusion across this boundary, which is trying to smooth the sharp transition.
- Once you've reached this state, the model should be relaxed. Now you can set your inlist to save at a recent model number and `./re` from a photostep just before this model so that **MESA** saves the relaxed model.
- Finally, you can turn on and off the relevant flags to evolve your relaxed model without extra heating and with burning and neutrino-cooling back on, and `./rn` the relaxed model.

Maxilab: Constructing and evolving a double white dwarf merger remnant

- Your model should redistribute entropy and start to contract as it cools. The temperature and density of the material at the base of the hot envelope rise until burning is ignited.
- Let's get a better idea of the progress of the burning as it flashes inwards.
 - Specifically, let's track the mass location where the maximum value of burning is taking place and add it to the `pgstar` history window.
 - First, find out if something like that exists in the default `history_columns.list` (search for “nuc” or “eps” using your favorite search method).
 - As it turns out, the variable we want is commented out by default, so we need to work with a non-default copy. So, copy `history_columns.list` over to your work directory (**MESA** looks for this file in your work directory first before using the version in the `defaults` directory), and uncomment the variable back in.
 - Unfortunately, doing so changes the `history.data` file in the `LOGS` directory, so we have to get rid of it and start with a fresh `history.data`. So stop the run and delete the old `LOGS/history.data`. To see the history of the new variable, add it to `inlist_pgstar` as the new `History_Panels1_other_yaxis_name(3)`, and `./re` from a recent photostep.

Maxilab: Constructing and evolving a double white dwarf merger remnant

- How many flashes does it take before the burning wave reaches the center? How long does it take?
- After the burning wave reaches the center, what happens? What does it look like (how bright, how hot), and how long does it live like this?
- Eventually this phase ends, and it cools down as a C/O white dwarf. Let's compare the resulting WD to the WDs from our minilab.
 - Put in a condition to stop the WD when the central temperature reaches 2×10^7 K. You can't do this too early because the temperature is below 2×10^7 K during the burning phase.
 - What is the total ^4He mass at the end of the day? How does it compare to the results from the minilab? Report your values on the Google spreadsheet.

Maxilab: Constructing and evolving a double white dwarf merger remnant

- Now that you've expertly modeled the evolution of a 0.2 + 0.3 Msol double helium WD merger, let's try another set of parameters. For simplicity, let's stick with double helium WDs, so both initial components should be between 0.15 and 0.5 Msol or so, giving total masses between 0.3 and 1.0 Msol.
 - Warning: the higher the mass, the longer it will take to run! If you decide to go above a total mass of 0.7 Msol, use coarser resolution controls.
- Different masses will entail changing the various parameters in the extra energy routine, specifically $M_{\text{transition}}$, $\eta_{\text{transition}}$, `highentropy`, and `lowentropy`. Getting the last 3 correct for your particular choice of masses will probably require some trial and error. Since it's annoying to change them in `run_star_extras.f` and recompile for each trial, let's find a way to change them at run-time, without the need to recompile.
 - This is done by using `x_ctrl(:)`, which you can change in your `inlist` at run-time.
 - You will need to decide which `x_ctrl` matches which variable (e.g., `x_ctrl(1)` might be your $M_{\text{transition}}$), and use them in `run_star_extras.f` by accessing `s%x_ctrl(1)`.
 - Remember to keep careful track of which `x_ctrl` changes which variable!
 - You may find it useful to look at a profile plot of η to pick a good $\eta_{\text{transition}}$. So add this profile to your profile plot of `extra_heat` to help you pick a winner.

Maxilab: Constructing and evolving a double white dwarf merger remnant

- If you've made a couple helium WD merger remnants and there's still time, you can try and make a double C/O WD merger remnant in a similar manner.
 - Unlike with pure helium compositions, relaxing to a pure C/O composition is non-trivial.
 - You'll have to do something similar to what we did in the first minilab. Namely, let the material cool down until it degenerates, and only then relax the composition and entropy to what you desire.
 - Also, because of the slow speed of the inwardly moving burning wave, stick with low total masses $< 1.2 M_{\text{sol}}$ or so (e.g. $0.5 + 0.6 M_{\text{sol}}$).
- As it turns out, the C/O WD merger remnant evolution is very similar to the lab yesterday. Alternatively, you can try to model the remnant of a helium + C/O WD merger (i.e., a puffy helium layer on a cold degenerate C/O core).
 - Relaxing a non-pure composition is a little different. You will need the `relax_initial_composition` flag and its associated flags in `star_job`, and you'll need to give it a file with a compositional profile. This is more complicated, but the comments near the relevant flags in `star_job.defaults` will help explain things.
- How do these remnants evolve differently from double helium WD merger remnants? What different phases do they go through? Are they longer or shorter?