



INDO AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH



ISOLATION AND CHARACTERIZATION OF LIPOLYTIC BACTERIA FROM OIL CONTAMINATED SOIL FROM PETROL BUNK AT SOUTHEAST BANGALORE.

Ajay Kumar Sahu^{1*}, Rahul Nemani^{1*}, Prangya Paramita Acharya¹, Dr. Rupali Sinha¹, Subhranil Sengupta¹, Dr. Kirankumar B¹

¹Dept. of Biotechnology, Sambalpur University, Odisha, INDIA.

²Dept. of Biotechnology, Bangalore University, Bangalore, INDIA.

ARTICLE INFO

Article history

Received 20/03/2019

Available online

02/04/2019

Keywords

Isolate Lipolytic Bacteria,
Oil Contaminated Soil,
Physiological Parameters,
Gravimetric Analysis.

ABSTRACT

The isolation of oil contaminated sites and gravimetric analysis of degradation in which, two bacterial formed maximum clearing zones on minerals salt medium. An increase in cell number indicating that the bacterial isolation was responsible for the oil degradation. The collected oil contaminated sites at kormongala, Indra nagar, MTTC culture, micrococcus spp., Bacillus spp., pseudomonas spp., which are able to utilize the oil in soil as carbon sources, were added to oil contaminated soil sample, the growth profiles were determined by monitoring the optical density, dry weight and Ph of the culture utilizing lubricating oil as sole sources of carbon, Bushnell Haas media supplemented with petrol, kerosene and diesel as sole carbon sources was used for isolation of bacteria capable of degrading these petroleum fractionates. From three soil sample and two water sample, a total of nine bacterial strains were isolated capable of degrading petrol, kerosene and diesel with varying tolerance capacities, the isolates were identify by using standard biochemical test and morphological studies and it was determined that these strains belong to six bacterial genera .the present study suggest that the isolated bacterial species could be employed for bioremediation in environment polluted with petroleum and its products, indigenously from the soil and water contaminated with crude oil in the vicinity of oil drilling well were found to be most efficient crude oil utilize as turbidity observed by spectrophotometrically. In the various study of lipolytic bacteria concluded that the taken of oil contaminated soil from from petroleum bunk and to identify their biochemical characterization by using various sources. How it's helpful for characterize by using of lipolytic bacteria.

Corresponding author

Rahul Nemani

Department of Microbiology,

Bangalore University

Kalyannagar, Banswadi, Bangalore, India.

sahu88506@gmail.com

rahulnemani@gmail.com

Please cite this article in press as *Ajay Kumar Sahu et al. Isolation and Characterization of Lipolytic Bacteria from oil Contaminated Soil from Petrol Bunk at Southeast Bangalore.. Indo American Journal of Pharmaceutical Research.2019:9(03).*

Copy right © 2019 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

In the past years a large number of ecosystem have been changed by the growing influence of human activity, many people have become aware of the need to protect ecosystem as well as to evaluate the damage caused by contamination. Most of the petroleum goes in the ecosystem via leak of coastal oil refiners this fact increased the interest of scientists to investigate the oil distribution and its fate in the environment especially the marine environment. Past analysis of reported oil spills indicated that most of the oil comes from tankers, barges and other vessels as well from land pipeline spills.

Shipping accidents have a serious impacts on the surrounding environment, the consequence include serious, widespread and long term damage to marine ecosystem, terrestrial life, human health and natural resources, it is very important to characterize oil spills as already known sources, this can help environment to predict the behavior of oil and estimate the long term impact on the environment, it is necessary to select an appropriate cleanup methods.

Conventional remediation methods include physical removal of contaminated material; these methods also use chemicals, especially shoreline cleaner which are often organic solvents with or without surfactants.

There is an increased interest in promoting environmental methods in the process of cleaning oil polluted sites, these methods are less expensive and do not introduces additional chemical to environment, compared to physiochemical methods, bioremediation offers a very feasible alternative for an oil spill response , this technique is considered an effectives technology for treatment of oil polluted one reason is that the majority of the molecules in the crude oil and refined produced are biodegradable .

Microbial degradation of petroleum hydrocarbon is one of the major practices in natural decontamination process. Presence of petroleum hydrocarbon has been reported to influence the biodiversity, distribution and pollution of microorganism in an environment, prolonged exposure and high oil concentration may cause the development of liver or kidney disease, possible damage to the bone marrow and an increased risk of cancer.

The indigenous bacteria which degrade hydrocarbon are isolated from oil contaminated petrol bunk sites and screened for their hydrocarbon degradation efficiency, they were further characterized by morphological, cultural and biochemical techniques. Microorganism have enzyme systems enables them to degrade and utilize diesel oil as a sources of carbon energy. The main aim to isolate the lipolytic bacteria from various sites of Bangalore and to identify the biochemical characteristic by using of some biochemical test, so that to know the impact of lipolytic Bacteria from oil contaminated soil of petroleum bunk.

MATERIALS AND METHODS

Isolation of lipolytic bacteria:-

Soil sample

Soil was collected from different areas HRBR LAYOUT, KALYANNAGAR, HEBBAL AND KAMMANHALI.

Soil sample from each sites of petrol pump was taken and transferred into sterile polythene bags to minimize moisture losses during transportation and samples was subjected to bacteriological analysis.

Isolation of Bacteria

- Soil sample was serial diluted up to 10^5 dilution factors.
- A dilution factor 105 was spread plated on BHM media agar.
- Incubated for 24 hour at 37^0 c.

Identification of lipolytic bacteria

Gram staining and motility test was done and observed under microscope.

Biochemical characterization of isolates

Indole test

Peptone broth was prepared and sterilized at 121^0 c for 15 min and inoculated with test organism, incubated the medium at 37^0 c for 24 hours. Added 1 ml of kovac reagent to tubes including control

Methyl Red test

Prepared MR-VP broth in two flasks, inoculate the broth with the test organism and incubated for 24 hours at 37^0 c, after 24 hours of incubation transferred 5 ml of broth into two test tubes. To the each broth culture added 5 drops MR indicator the tubes and shake them.

Voges proskauer test

Prepared MR-VP broth in two flask, inoculates the broth with the test organism and incubated for 24 hours, prepared barritt reagent of A and B, After 24 hours all agar slant were examined for the presence of growth and coloration of the medium.

Citrate utilization test

Prepared citrate agar slant and inoculated each of the test organism into appropriately labeled tubes by means of a loop, the slant was left uninoculated that serve as control, incubated for 24 hours at 37^0 c.

Catalase test

Transferred small quantity of culture from the plates on glass slide, added 1 drop of 3 % H₂O₂ observe bubbles.

Oxidase test

Taken oxidase disc in clean microscopic slide, pasted the culture on the oxidase disc and observed for color changes.

Nitrate test

Prepared nitrate broth and inoculated each of the test organisms into its appropriately labeled tubes means of a loop. The last slant was left uninoculated that serve as control, incubated all culture for 24 hours at 37^oc , after 24 hours add one dropper full of sulfanilic acid and one dropper full of α naphthylamine to each broth.

Starch test

Prepared starch agar and inoculated each of the test organism into its appropriately labeled tubes by means of a loop, the last plates was left un inoculated that serve as control, incubated all culture for 24 hours at 37^o c. after 24 hours all agar slants were examined for the presence of growth and zone formation on the medium.

Gelatin test

Prepared gelatin slant and inoculated each of the test organism into its appropriately labeled tubes by means of a loop. The slant was left un inoculated that serve as control, incubated all culture at the bacterium optimal growth temperature for up to 1 week and checked every day for gelatin liquefaction. Gelatin normally liquefies at 28^oc and above, so to confirm that liquefaction was due to gelatinase activity.

Physiological parameters**Effect of Ph**

Prepared nutrient broth in 5 flasks, which was sterilized and 1 ml of engine oil was added to all the flask except the control flask which was set blank, remaining flask was adjusted to ph 3, 5, 7, 9 and labeled respectively inoculated the test organism and incubate at 37^oc for 72 hours and OD 600 nm was taken at every 24 hours intervals.

Effect of temperature

Prepared nutrient broth in 5 flask, which was sterilized and 1 ml of engine oil was added to the entire flask except the control flask which was set blank, remaining flask was kept at temperature RT 30, water bath 60, incubator 37^oc and fridge 4^oc. Flasks were inoculated with test organism and incubate at 37^oc for 72 hours OD 600 nm was taken at every 24 hour intervals.

To examine the degradation of oil by gravimetric analysis**Oil Degradation**

For examining the degradation of oil, BHM medium supplemented with 5g/l of engine oil was used. About 50 ml medium was dispensed in 250 ml conical flask, the media was inoculated with 0.1 ml of crude oil degrading bacteria and incubated at 28^oc for 7 days on a rotary shakes at 175 rpm.

Extraction of crude oil

For estimation of oil degradation rates by gravimetric analysis 5 ml of n- hexamine was added to above flask, the contents were transferred to a separating funnels and extracted. Extraction was carried out twice to ensure complete recovery of oil, the extract was treated with 0.4 g of anhydrous sodium sulphate to remove the moisture and decanted into a breaker leaving behind sodium sulphate to remove the moisture and decanted into a breaker leaving behind sodium sulphate .the amount of residual oil was measured after extraction of oil from the medium and evaporating it to dryness in warm bath 40^oc, the volume of extracted oil was deducted from the previously weighed breaker.

RESULTS AND DISCUSSION**Table 1 biochemical test.**

| Test | Gramstainin g | Motilit y | Indol e | Methy l red | Voges Proaskaue r | Citrat e | Catalas e | Gelati n | Nitrat e | Oxidas e | Starc h | Tripl e sugar test | Ureas e |
|-------------------|---------------|-----------|---------|-------------|-------------------|----------|-----------|----------|----------|----------|---------|--------------------|---------|
| Pseudomona s spp. | + | -- | -- | -- | -- | + | + | + | + | + | -- | K/N C | + |
| Bacillus spp. | + | -- | -- | -- | + | + | + | + | + | + | + | K/K | -- |

Table5 effect of Ph.

| Organism | PH | DAY1 | DAY2 | DAY3 |
|------------------|----|------|------|------|
| Pseudomonas spp. | 03 | 0.01 | 0.02 | 0.02 |
| | 05 | 0.02 | 0.03 | 0.05 |
| | 07 | 0.39 | 0.42 | 0.50 |
| | 09 | 0.35 | 0.40 | 0.43 |
| Bacillus spp. | 03 | 0.01 | 0.02 | 0.03 |
| | 05 | 0.02 | 0.04 | 0.07 |
| | 07 | 0.14 | 0.20 | 0.50 |
| | 09 | 0.02 | 0.04 | 0.05 |

Table 6 Effect of temperature.

| Organism | Temperature °C | Day1 | Day2 | Day3 |
|------------------|----------------|------|------|------|
| PSEUDOMONAS SPP. | 04 | 0.02 | 0.02 | 0.02 |
| | 30 | 0.26 | 0.32 | 0.48 |
| | 37 | 0.36 | 0.45 | 0.55 |
| | 60 | 0.01 | 0.01 | 0.01 |
| | 04 | 0.02 | 0.02 | 0.02 |
| BACILLUS SPP. | 30 | 0.04 | 0.08 | 0.11 |
| | 37 | 0.15 | 0.25 | 0.36 |
| | 60 | 0.01 | 0.01 | 0.01 |
| | 60 | 0.01 | 0.01 | 0.01 |

Table 7: Oil recovered.

| Isolates | Weight of beaker with oil gm. | Weight of beaker gm. | Recovered oil gm. |
|------------------|-------------------------------|----------------------|-------------------|
| Pseudomonas spp. | 47.55 | 47.25 | 0.30 |
| Bacillus spp. | 48.55 | 48.10 | 0.45 |

Table 8: protein estimation.

| Sample | Concentration µg/ml | O.D (600nm) |
|------------------|---------------------|-------------|
| Pseudomonas spp. | 660 | 0.95 |
| Bacillus spp. | 620 | 0.89 |




Figure 1 :Effect of pH.




Figure 2: Effect of Temperature.




Figure 3: BIOCHEMICAL TESTS.




Figure 4: BIOCHEMICAL TESTS.

CONCLUSION

The ability of the bacterium to withstand the effects of reduce water activity is great ecological importance. Production of valuable metabolites by microorganism has always been one of the main areas of intense scientific research over the years. The present study was undertaken to isolate lipolytic bacteria from contaminated soil sample and subsequent optimization of culture parameters, which include temperature, ph, incubation period, agitation periods for isolation and characterization of lipolytic that effect of carbon source, nitrogen source metals ions were further assessed to determine in maximum lipolytic bacteria from the contaminated soil which was taken from different areas of Bangalore. Selection bacterial culture was achieved by culture in the different media which was taken. The main purpose of isolate lipolytic bacteria from oil contaminated soil which indicate more ph and temperature resistant that help to grown the such bacteria using of such nutrient choosing such compound which helps to identify the particular bacteria by using of biochemical characterization.

ACKNOWLEDGEMENT

First and foremost I would like to offer my prayer to almighty for showering his blessing upon me, I express my deep sense gratitude and indebtedness to respected guide Dr. Kirankumar who introduce the area of research lipolytic bacteria from oil contaminated soil from petroleum bunk in .

Author's contribution

Ajay kumar sahu and Rahul Nemani prepared and conceived designed the experiment and draft the manuscript under the guidance of Dr. Kirankumar , Prangya P. Acharya and Dr. Rupali Sinha prepared all media, reagent for bacterial growth and biochemical test. Subhanil Sengupta participated the work and design some idea and coordination.

Conflict of interest

nil

REFERENCE

1. Ahmed. M, wasim s, hayat m, imran k (2015) identification and characterization of intrinsic petrophilic bacteria from oil contaminated soil and water 338-346 retrieved on February 18, 2015.
2. Bhasheer s.k , umavathi .s, thangavel m. and thangam y. 2014 ' diversity of diesel degrading bacteria from a hydrocarbon contaminated soil ijemas 3(11) 363-398 on April 15, 2015
3. Chithra s and shenpagam H N (2014) biodegradation of crude oil by gravimetric analysis. International journal of advance technology in engineering and science 2(11), 372-376, January 16, 2015.
4. Das and chandaran p 2012 'Microbial degradation of petroleum hydrocarbon contaminant, biotechnology research international, on may 10 2015.
5. Eman A,H Mohamed , naeima M H yousef and azza G farag 2012 ' isolation and molecular identification of polyaromatic hydrocarbon utilizing bacteria from crude petroleum oil sample , 6(49) pp 7479-7484.
6. Ghaima K K, Mohamed A.I and Mohamed M M . effect of some factors on lipase production by bacillus spp isolated from diesel fuel polluted soil,
7. Ghazali F.M rahman , sallah A.B and basri M 2004 .biodegradation of hydrocarbon in soil by microbial consortium .international journal biodegradation 54(1). 61-67 on March 19, 2019.

8. Hamzah a, Rabu a, azmy and yusoff N A. 2010. Isolation and characterization of bacteria degrading sumandak and south angsi oil. 39(21), 161-168. On March 16, 2015.
9. Khan J,A , Rizvi S,H.A and hasan .S 2011, isolation and characterization of microorganism from oil contaminated sites adv appl sci res, 2(3), 455-460.
10. Latha , R and kalaivani R. (2012) bacterial degradation of crude oil by gravimetric analysis, advance applied science research on December 15, 2014.
11. Makut M.D and ishaya p. (2010) ‘ bacterial associated with soils contaminated with used petroleum in keffi town , Nigeria African journal of microbiology research (16), 1698-1702.
12. Pathak hardik , kamna bhatanagar and jaroli p (2013) serratia the 4t engine oil degrader , open access scientific reports 1:117 retrieved on March 19, 2015.
13. Sadhana B. and sarayan J.S (2015) isolation and characterization of soil bacteria from the petrol pump station of Madurai town, 2(32), 60-71.
14. Nikhil T, Deepa v, rohan G and satish B (2013) , isolation and characterization , identification of diesel engine oil bacteria from garage soil and comparison of their bioremediation potential, international journal of environmental science 2(23) 48-52 .
15. Paramaniak D and rajalaxmi G, 2013 ‘biodegradation of petroleum hydrocarbon pollutant in soil using microbial consortium. International journal of plant, animal and environmental science 3(3), 173-178.
16. Vinothi c. sudhakar s, and ravikumar r, 2015 ‘biodegradation of petroleum and crude oil by pseudomonas putida and bacillus cereus. International current microbiology application science 4(1): 318-329.



54878478451190309



Submit your next manuscript to **IAJPR** and take advantage of:

Convenient online manuscript submission

Access Online first

Double blind peer review policy

International recognition

No space constraints or color figure charges

Immediate publication on acceptance

Inclusion in **Scopus** and other full-text repositories

Redistributing your research freely

Submit your manuscript at: editorinchief@iajpr.com

