

Thiamine Biosynthesis


 Thiamine Biosynthesis in *Escherichia coli*: Identification of the Intermediate and By-Product Derived from Tyrosine**

Marco Kriek, Filipa Martins, Martin R. Challand, Anna Croft, and Peter L. Roach*

Thiamine is an essential cofactor in many metabolic pathways and is required by all living organisms. The biosynthesis of thiamine phosphate is convergent, linking two independently biosynthesized heterocyclic precursors (Scheme 1, **6** and **7**) in

(Scheme 1). This hydrolytically unstable intermediate is incorporated into the thiazole **6** in a multistep reaction requiring the thiazole synthase (ThiG),^[7] the sulfur donor ThiFS thiocarboxylate,^[8,9] and 1-deoxyxylulose 5-phosphate (Dxp, **5**). Dehydroglycine **2** has been proposed as the first common intermediate for bacterial thiazole biosynthesis,^[10] but microorganisms growing under anaerobic conditions probably cannot oxidize glycine. Instead anaerobes, such as *Escherichia coli*, utilize an alternative pathway forming dehydroglycine from tyrosine **1** in a ThiH-dependent reaction.^[11,12] This biosynthetic step therefore requires the cleavage of the C_α-C_β bond and release of the aromatic side chain. ThiH shows sequence similarity to the “radical S-adenosylmethionine (AdoMet)” family of proteins,^[13] including conserved ligands to an essential [4Fe-4S] cluster^[14] and has been shown to form a complex with ThiG.^[15] Studies on the reconstitution of the thiazole-forming reaction^[10,16] with purified *E. coli* proteins have shown that ThiGH requires AdoMet

ANAEROBES:



Scheme 1. The biosynthesis of thiamine phosphate **8**.

and a reductant for activity, but (like other proteins in the “radical Adomet” family^[17,18]) it is not catalytically active in vitro. In this paper, we elucidate the reaction products formed in situ hydrolysis of dehydroglycine. Finally, these experiments are shown to be relevant to in vivo metabolism by the detection of *p*-cresol in *E. coli* that are expressing ThiH.

the final step.^[1-4] The biosynthesis of the thiazole moiety of thiamine has been well characterized in the aerobic *Bacillus subtilis*,^[4,5] which obtains the C2-N3 fragment of the thiazole by oxidation of glycine **9** to dehydroglycine **2** using ThiO^[6]

To follow the fate of tyrosine in the reaction, ¹⁴C-labeled tyrosine was incubated with the heterodimeric ThiGH complex, AdoMet and the natural electron-donor system consisting of flavodoxin (FldA), flavodoxin reductase (Fpr), and NADPH. After 2 h, the reaction was analyzed by thin-layer chromatography (TLC) and monitored by autoradiography (Figure 1). By using two TLC systems, two radiolabeled products with very different polarities were observed. The highly polar component co-eluted with glyoxylate (**4** in Scheme 1 and Figure 1A) and comparison of the less polar product with several aromatic standards showed that it co-eluted with *p*-cresol (**3** in Scheme 1 and Figure 1B). This experiment also demonstrated the absolute requirement for AdoMet as the negative control showed no activity.

[*] Dr. M. Kriek, F. Martins, M. R. Challand, Dr. P. L. Roach
 School of Chemistry
 University of Southampton, Highfield
 Southampton SO17 1BJ (UK)
 Fax: (+44) 2380-596-805
 E-mail: plr2@soton.ac.uk
 Homepage: <http://www.soton.ac.uk/~roachweb/home/index.htm>
 Dr. A. Croft
 School of Chemistry
 University of Wales, Bangor
 Gwynedd LL57 2UW (UK)

[**] This work was supported by grants from the Biotechnology and Biological Sciences Research Council (M.K. and M.R.C.), the Engineering and Physical Sciences Research Council (F.T.M.), and the Royal Society (University Research Fellowship to P.L.R.). We thank J. Street, N. Wells, G. J. Langley, and J. Herniman for help with spectroscopy and spectrometry.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.





Figure 1. Autoradiograms of a ThiH assay. Two TLC systems (plates A and B) were used to resolve the products. The complete assay (lane 1 on each plate) contained L-[U-¹⁴C]tyrosine, reconstituted ThiGH, AdoMet, and a reductant. The negative control (lane 2) lacks AdoMet. Spots corresponding to *p*-cresol, tyrosine, and glyoxylate standards are indicated by C, Tyr, and G, respectively.

The formation of these two products was confirmed by ¹³C NMR spectra of reactions containing [U-¹³C]tyrosine. After precipitation of the protein components, the spectrum of the aqueous supernatant showed signals typical of glyoxylate (Figure 2A). The spectrum of an organic extract prepared from this supernatant confirmed the formation of *p*-cresol (Figure 2B). This sample was also analyzed by GCMS, which showed the retention time (13.38 min) and expected mass of *p*-cresol derived from [¹³C]tyrosine. Taken together, the results demonstrate unequivocally that *p*-cresol and glyoxylate were produced from tyrosine during *in vitro* reactions.

The role of ThiH in the reaction can be explained by the mechanism proposed in Scheme 2. The reductive cleavage of AdoMet yields the highly reactive 5'-deoxyadenosyl radical, which can abstract the phenolic hydrogen atom from tyrosine. The resultant radical **10** may react by at least two pathways.



Scheme 2. Proposed mechanism for ThiH. The 5'-deoxyadenosyl radical (DOA[•]) is derived from the reductive cleavage of AdoMet.




Figure 2. ¹³C NMR spectra of ThiGH reactions. The reaction mixtures contained reconstituted ThiGH, L-[U-¹³C]tyrosine, AdoMet, and a reductant, and were incubated at 37°C for 2 h. Panel A shows the ¹³C NMR spectrum of the supernatant obtained after the proteins were removed by acid precipitation and centrifugation. Glyoxylate gives signals at 173 and 86 ppm under these conditions, observed as doublets in here because of the ¹³C labeling. Panel B shows the ¹³C NMR spectrum of an organic extract prepared from the supernatant and is consistent with ¹³C-labeled *p*-cresol [¹³C NMR (100 MHz; CDCl₃) δ = 153.2 (t, *J* = 66 Hz, COH), 130.1 (m, 2 CH, CCH₃), 115.0 (m, 2 CH), 20.4 ppm (m, CH₃)].

Homolytic cleavage of the C_α–C_β bond yields the quinone methide **11** and the radical **12**. The alternative heterolytic pathway leads directly to **2** and the stabilized radical anion **13a** ↔ **13b**. To form *p*-cresol and dehydroglycine by either of these pathways requires further electron-transfer steps, which may be facilitated by the proximity of the intermediates to the [4Fe-4S] cluster in the active site of ThiH. DFT and *ab initio* calculations suggest that the homolytic mechanism is ther-

modynamically more favorable, with increased polarity improving the relative stability of the heterolytic pathway, but not enough for this to be identified as the preferred pathway. Judicious placement of specific residues may therefore result in enough stabilization to tip this balance, and further experimental data will be required to differentiate between these pathways. The subsequent fate of the dehydroglycine **2** depends on the reaction conditions: if the components required for thiazole formation are present (including ThiG, the sulfur donor ThiFS thiocarboxylate, and Dxp), then it will be incorporated into the thiazole phosphate **6**.^[10] If any of these components are absent, the dehydroglycine is likely released from the ThiGH complex into the aqueous medium, where it is rapidly hydrolyzed to glyoxylate and ammonia.

ThiH is not unique in mediating this type of C–C bond cleavage of a phenolic compound. For example, *Clostridium difficile* produces *p*-cresol by the decarboxylation of *p*-hydroxyphenylacetate using a glycol radical enzyme *p*-hydroxyphenylacetate decarboxylase (HPAD).^[19–21] It seems probable that ThiH and HPAD use different carbon-centered radicals to initiate mechanistically related cleavage reactions.

To investigate the kinetics and stoichiometry of product formation, the *p*-cresol and glyoxylate formed in a ThiH reaction were quantified using HPLC over a two-hour period (Figure 3). The reaction produced *p*-cresol and glyoxylate in an approximate 1:1 ratio over the time course and approached completion after 1 h (Table 1). In a subsequent experiment,




Figure 3. Time course of product formation in a ThiH assay. The graph shows the increase in the concentration of *p*-cresol (○, dashed line) and glyoxylate (●, solid line) with time. The assays, which contained reconstituted ThiGH complex, tyrosine, AdoMet, and a reductant, were incubated for the indicated time at 37 °C.

Table 1: Kinetic parameters for product formation. Data from Figure 3 were fitted to first-order processes to determine the calculated final concentration of products and the apparent first-order rate constant. Results are shown with standard errors.

Product	Rate constant [10^{-2} min^{-1}]	Final conc. [μM]
<i>p</i> -cresol	4.6 ± 0.7	176.2 ± 7.8
glyoxylate	6.3 ± 1.1	171.6 ± 7.9

the amount of 5'-deoxyadenosine formed after 2 h was measured as 1.3 mol equivalents relative to *p*-cresol, suggesting some uncoupled turnover of AdoMet had occurred.

To demonstrate the formation of *p*-cresol in vivo, we used a mutant *E. coli* K strain that lacks a genomic copy of *thiH*.^[22] Two derivative strains of this mutant were prepared, one transformed with a ThiH expression plasmid^[15] and the other with a control plasmid containing no thiamine biosynthetic genes. These strains were cultured in defined medium for 24 h, and an organic extract was prepared from the cleared medium. Comparison of the extracts by GCMS analysis (Figure 4) revealed a new peak at 13.38 min, corresponding to *p*-cresol, in the sample derived from the ThiH-expressing cells. In contrast *p*-cresol was not detectable in the negative control. Using cultures of an *E. coli* B strain and [3,3-²H₂]tyrosine, White^[23] reported the isolation of 4-hydroxybenzyl alcohol as a by-product from thiamine biosynthesis, a result that seemingly contradicts our observations. However,




Figure 4. Evidence for the ThiH-dependent formation of *p*-cresol in vivo. A) Section of the GCMS chromatogram of an extract of medium from *E. coli* cultures that express ThiH (solid line, displaced upwards for clarity) and the negative control sample (no ThiH, dashed line). The *p*-cresol peak at 13.38 min is marked. B) The mass spectrum corresponding to the new peak at 13.38 min showing the expected M^+ signal for *p*-cresol (m/z 108.1).

E. coli B strains (but not K strains) express an oxidase that can accept *p*-cresol as a substrate.^[24] It is therefore probable that tyrosine cleavage in *E. coli* produces *p*-cresol, but those strains with a suitable oxidase convert *p*-cresol to *p*-hydroxybenzyl alcohol.

In conclusion, we have characterized the products of tyrosine cleavage in the presence of the “radical Adomet” protein ThiH, AdoMet, and a reductant. The side chain of tyrosine is converted to *p*-cresol during this reaction. The remaining fragment of tyrosine likely forms the intermediate dehydroglycine, which can be combined with Dxp and a sulfur donor during thiazole biosynthesis in vivo. During in vitro experiments that lacked these other thiazole precursors, glyoxylate was observed to accumulate from the hydrolysis of the dehydroglycine.

Received: June 12, 2007

Published online: October 30, 2007

Keywords: biosynthesis · enzyme mechanisms · radicals · thiamine

-
- [1] I. D. Spenser, R. L. White, *Angew. Chem.* **1997**, *109*, 1096; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1032.
- [2] T. P. Begley, D. M. Downs, S. E. Ealick, F. W. McLafferty, A. Van Loon, S. Taylor, N. Campobasso, H. J. Chiu, C. Kinsland, J. J. Reddick, J. Xi, *Arch. Microbiol.* **1999**, *171*, 293.
- [3] B. G. Lawhorn, R. A. Mehl, T. P. Begley, *Org. Biomol. Chem.* **2004**, *2*, 2538.
- [4] J. H. Park, P. C. Dorrestein, H. Zhai, C. Kinsland, F. W. McLafferty, T. P. Begley, *Biochemistry* **2003**, *42*, 12430.
- [5] E. C. Settembre, P. C. Dorrestein, H. Zhai, A. Chatterjee, F. W. McLafferty, T. P. Begley, S. E. Ealick, *Biochemistry* **2004**, *43*, 11647.
- [6] E. C. Settembre, P. C. Dorrestein, J. H. Park, A. M. Augustine, T. P. Begley, S. E. Ealick, *Biochemistry* **2003**, *42*, 2971.
- [7] A. Chatterjee, X. Han, F. W. McLafferty, T. P. Begley, *Angew. Chem.* **2006**, *118*, 3587–3590; *Angew. Chem. Int. Ed.* **2006**, *45*, 3507.
- [8] C. Lehmann, T. P. Begley, S. E. Ealick, *Biochemistry* **2006**, *45*, 11.
- [9] P. C. Dorrestein, H. Zhai, F. W. McLafferty, T. P. Begley, *Chem. Biol.* **2004**, *11*, 1373.
- [10] M. Kriek, F. Martins, R. Leonardi, S. A. Fairhurst, D. J. Lowe, P. L. Roach, *J. Biol. Chem.* **2007**, *282*, 17413.
- [11] R. H. White, F. B. Rudolph, *Biochim. Biophys. Acta Gen. Subj.* **1978**, *542*, 340.
- [12] B. Estramareix, M. Therisod, *Biochim. Biophys. Acta Gen. Subj.* **1972**, *273*, 275.
- [13] H. J. Sofia, G. Chen, B. G. Hetzler, J. F. Reyes-Spindola, N. E. Miller, *Nucleic Acids Res.* **2001**, *29*, 1097.
- [14] N. C. Martinez-Gomez, M. Robers, D. M. Downs, *J. Biol. Chem.* **2004**, *279*, 40505.
- [15] R. Leonardi, S. A. Fairhurst, M. Kriek, D. J. Lowe, P. L. Roach, *FEBS Lett.* **2003**, *539*, 95.
- [16] R. Leonardi, P. L. Roach, *J. Biol. Chem.* **2004**, *279*, 17054.
- [17] N. B. Ugulava, C. J. Sacanell, J. T. Jarrett, *Biochemistry* **2001**, *40*, 8352.
- [18] R. M. Cicchillo, D. F. Iwig, A. D. Jones, N. M. Nesbitt, C. Baleanu-Gogonea, M. G. Souder, L. Tu, S. J. Booker, *Biochemistry* **2004**, *43*, 6378.
- [19] L. Yu, M. Blaser, P. I. Andrei, A. J. Pierik, T. Selmer, *Biochemistry* **2006**, *45*, 9584.
- [20] P. I. Andrei, A. J. Pierik, S. Zauner, L. C. Andrei-Selmer, T. Selmer, *Eur. J. Biochem.* **2004**, *271*, 2225.
- [21] T. Selmer, P. I. Andrei, *Eur. J. Biochem.* **2001**, *268*, 1363.
- [22] T. Baba, T. Ara, M. Hasegawa, Y. Takai, Y. Okumura, M. Baba, K. A. Datsenko, M. Tomita, B. L. Wanner, H. Mori, *Mol. Syst. Biol.* **2006**, DOI: 10.1038/msb4100050.
- [23] R. H. White, *Biochim. Biophys. Acta Gen. Subj.* **1979**, *583*, 55.
- [24] M. A. Prieto, A. Perezaranda, J. L. Garcia, *J. Bacteriol.* **1993**, *175*, 2162.