



INDO AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH



TO ASSESS THE DRUG ADHERENCE, FACTOR AFFECTING ADHERENCE AND MANAGEMENT OF ACUTE ASTHMA CHILDREN'S IN TAMILNADU

Arulprakasam K C^{1*}, Senthilkumar N²

¹Research Scholar, PRIST University, Tanjavuor, Tamilandu.

²JKKMMRFS Annai JKK Sampoorani Ammal College of Pharmacy, Komarapalayam, Namakkal DT, Tamilnadu.

ARTICLE INFO

Article history

Received 05/05/2019

Available online

31/05/2019

Keywords

Asthma,
Adherence,
Health Education,
Non-Adherence,
Self-Action Plan.

ABSTRACT

The prevalence and associated burden of disease due to asthma, it is mandatory to obtain an optimal control of the disease and to improve outcomes for these patients. But it has been observed that there is very poor adherence to the therapy which leads to the suboptimal control of the disease. To study the adherence for therapy in asthma patients and to assess the impact of health education and self action plan in improving the adherence to the therapy. A prospective study was done in a total of 986 asthma patients over a period of 3 years. Once included in the study, the patients were followed up for a total of 12 weeks for calculation of non-adherence to the therapy. In non-adherent patients, we employed various health education strategies to improve the adherence in these cases. A total of 986 patients of asthma who were started on therapy over duration of 6 months were included in the study. At the end of 12 weeks, it was observed that, only 108 patients (10.95%) had regular adherence and 878 patients (89.05%) were non adherence to the therapy as prescribed for asthma. Factors that were associated with poor adherence were: Lower educational level status, poor socioeconomic status, cumbersome regimens, dislike of medication, Fears about side effects, beliefs, changing in regimen, and patient's ill attitudes toward health. After employing the various strategies for improving the adherence in these patients, the adherence increased in patients (61.32%) among the earlier defaulted patients, while the remaining 188 patients (38.68%) were found to be non-adherence even after various educational techniques. Nonadherence in asthma management is a fact of life and no single adherence improving strategy probably will be as effective as a good physician and patient relationship. Optimal self-management allowing for optimization of asthma control by adjustment of medications may be conducted by either self-adjustment with the aid of a written action plan or by regular medical review. Individualized written action plans based on peak expiratory flow are equivalent to action plans based on symptoms.

Corresponding author

Arulprakasam K C

Associate professor

JKKMMRFS Annai JKK Sampoorani Ammal

College of Pharmacy,

Komarapalayam, Namakkal

Tamilnadu, India 638183.

9842778531

kcarul2000@gmail.com

Please cite this article in press as **Arulprakasam K C et al. Maser the Drug Adherence, Factor Affecting Adherence and Management of Acute Asthma Children's In Tamilnadu. Indo American Journal of Pharmaceutical Research.2019;9(05).**

Copy right © 2019 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Asthma is a major public health problem affecting a large number of individuals of all ages. Globally, 100-150 million people suffer from asthma. India has 20-28 million asthmatics and the prevalence amongst children (5-11 years) is 10-15%. Being a chronic medical condition, management of asthma requires continuous medical care. Modern management of asthma requires prolonged medications. A key issue in proper management of asthma is adherence to treatment. Poor adherence to prescribed therapy increases morbidity and mortality and it is increasingly being documented that long-term adherence or adherence to prescribed therapy is difficult to attain [1]. Studies have reported that 50% of patients with a chronic disease do not use their medication at all or do not use it as prescribed [2]. Important reason for poor adherence is that patients with a chronic disease do not have a satisfactory understanding of their condition. The economic burden of asthma to the society is well documented in industrialized countries [4] [3], and is a great burden to the health services. Poor asthma control is responsible for a large proportion of the total cost of the disease, for the patient as well as to the society, and thus responsible for the both direct and indirect cost of therapy. The present study was undertaken to study the factors that influence patient's adherence with prescribed medications, and to assess the impact of health education and self-action plan in improving the adherence in asthma patients.

METHODS

The present study was a prospective study done at a tertiary care hospital over a period of 2 years from June 2015 to September 2018. Children below 15 years of age with the diagnosis of asthma as per Global Initiative for Asthma (GINA) guidelines (GINA 2012) were included in the study. Patients with acute severe asthma, chronic obstructive pulmonary disease (COPD), and cardiac asthma were excluded. All patients were interviewed using a standard interview schedule and requested to maintain a diary regarding the therapy. Apart from a detailed history, physical examination, pulmonary function test, and peak expiratory flow rate were measured during the first visit. All the patients were treated according to GINA guidelines under the supervision of physicians. Regarding the use of medications, the choice was left to the treating physician. All patients were followed-up every 15 days for 12 weeks. At the end of 12 weeks, adherence to treatment was calculated after studying the patient diary noting and frequency of the hospital visits.

Second step in this study was to impart health education in the non-adherent patients. We tried to improve the adherence by imparting the patient education program with the help of health educator. Various strategies which were employed for the patient education included verbal praise, interactive communication skills, tailoring the medications to the patient's routine, conducting asthma awareness camps for the defaulted patients, distribution of literature regarding asthma, and its consequences in the local languages and answering to the family's worries regarding asthma. At this stage, all the non-adherent patients were given detail health education regarding asthma, its allergy status, chronicity of the disease, and duration of treatment, nature of quick relief and long term medications, and preventive aspects of the disease with the help of health educator.

All the patients were asked to keep a diary of any symptoms occurring during this period. Also an action plan was provided by the treating physician for all the non adherence cases, explaining what to do in case of increase in the symptoms, or during any exacerbations, when to take oral steroids, and when to call the doctor. This will help the patient to be more interactive and be more communicable with the doctor. In the second step, all the patients were followed-up for another 12 weeks. Final assessment was done at the end of this period only. The study was approved by the Institutional Ethical Board. The economic status was classified as per modified Prasad classification [5].

STATISTICAL ANALYSIS

Data are expressed as the mean (standard deviation (SD)). Comparison of parameters between two groups was done by Student's *t*-test. Comparisons among three groups were done by one-way analysis of variance (ANOVA) with Bonferroni's multiple comparison tests. Differences in frequency between adherence and nonadherence patients were assessed by the Chi-square test. A *P* value of less than 0.05 was considered significant.

RESULTS

A total of 986 patients were studied during the period. The majority of the patients (42.18%) were in the age group of 5-10 years. The male: female ratio was 3:2. Majority of 361 (36.61%) had secondary education and 212 (21.50%) were illiterate. The majority of the patients belonged to middle socioeconomic status (table 1). Among male patients; regular adherence was observed in 69 (63.88) patients.

Table 1: Baseline characteristics of the patients with adherence.

Characteristic	Adherence n (%)	Non-adherence n (%)	P value
In the total 986respondent	108(10.95)	878(89.05)	0.001
Male	69(63.88)	518(58.99)	0.001
Female	39(36.12)	360(41.00)	0.001
Age			
>5	67(28.69)	223(71.61)	0.001
5-10	48(13.67)	303(86.33)	0.001
10-15	47(11.78)	352(88.22)	0.001
Education of parents			
Illiterate	23(10.85)	189(89.15)	0.001
Primary	25(10.46)	210(89.36)	0.001
Secondary	94(26.04)	267(73.96)	0.001
Degree	72(40.45)	106(59.55)	0.001
Economic status of guardians			
Lower	38(10.92)	310(89.08)	0.001
Middle	44(10.89)	360(89.11)	0.001
Upper	56(23.94)	208(76.06)	0.001

A higher number of male patients missed more doses 518(58.99) as compared to female patients 360(41.00). There was significant correlation between the educational status and the adherence to the therapy. Socioeconomic status of the patient was a significant risk factor associated with the nonadherence to the therapy.

Table 2. Causes for Non-adherence for asthma therapy.

Factors	No(%)
Complexity of medication	304(30.63)
Frequent changes in regimen	627(63.59)
Treatment requiring certain techniques	453(45.78)
Unpleasant side effects	453(45.78)
Duration of therapy	693(70.28)
Lack of immediate benefit of therapy	654(66.33)
Medications with social stigma	461(46.75)
Medication cost	724(73.42)
Lack of family or social support	351(35.60)
beliefs and behavior	670(67.95)
Feeling of well being	548(55.58)

Factors were the key reasons for non-adherent to the therapy (table 2). The major factors associated with poor adherence were: cost of the medication 724(73.42), beliefs 670(67.95), duration of therapy 693(70.28), lack of immediate benefit of therapy 654(66.33). Various strategies were employed with the help of a health educator after 12 weeks of therapy to improve the patient's adherence to the therapy, and these patients were followed-up for a further duration of 12 weeks to evaluate the response to the intervention. We have tried to educate these asthma patients in different ways, so that it will help in improving the adherence to the therapy.

Parents were asked what they do when the child has asthma signs or symptoms or actually has an asthma attack (table 3). Eighty-two percent of the respondents reported using an asthma medicine, and 64.2% went to a clinic or emergency room. The child was told to lie down or rest by 48.0% of parents.

In addition, 15.4% of the parents gave fluids by mouth and 13.4% reported calling a physician as a part of their emergency plan of action. Other responses, such as calling a hospital, performing breathing or relaxing exercises, attempting postural drainage, and using home remedies, were reported,10% of the time. Seventy-two percent of the respondents used a medication or took the child to a physician as the first or second action in their response to an acute asthma event.

Table 3: Parental Report of Management of Acute Asthma.

What do you do when the child is having asthma signs or symptoms or is having an asthma attack? (n=986)	n(%)
Give asthma medicine	805(81.7)
Go to clinic	633(64.2)
Have child lie down or rest	473(48.0)
Give fluids by mouth	151(15.4)
Call physician or Call hospital	224(22.7)
Use home remedies, herbs, or teas	118(12.0)
Ask family or friends for help or advice	118(12.0)
Use medicines or call physician as first or second action	723(73.4)

DISCUSSION

Asthma, a chronic lung disease that affects people of all ages, races, and ethnic groups, is a growing concern throughout the world. There is a need for educating the patient about asthma disease and medications used like DPI/MDI to be taken on regular basis as prescribed. In the study[6] conducted in Trinidad regarding the understanding and use of inhaler medication by asthmatics, it was observed that educating patients with a focus on children and the elderly, inhaler techniques, and reinforcing understanding of asthma medications could improve asthma management to a great extent. Nonadherence to treatment programs is common in patients with asthma. Nonadherence is more common than usually suspected and rates vary from 20% to 80%[7]. True rates of nonadherence are hard to come by because patients do not accurately report and physicians often do not inquire critically. The rule of thumb in chronic nonlethal disease (i.e., asthma, hypertension, etc.) is that one-third of patients are compliant, one-third are somewhat compliant, and one third are non adherence. This is important because compliant patients are significantly less likely to experience exacerbations than less compliant patients[8].

The present study was conducted to know adherence with therapy in asthma patients and reasons for nonadherence. An effort was also made to improve the patient adherence via the patient education program. In the present study a total of 878 patients (89.05%) were observed to be non adherence to the asthma therapy. Gibson *et al.*,¹¹ conducted a study to study the adherence with asthma medications in preschool children. In preschool children, the parents supervise and are responsible for drug administration. In this study it was observed that parental supervision would result in good adherence. It was concluded that adherence with prophylactic therapy is poor in preschool children with asthma whose medication is administered under parental supervision. Nonadherence depends on many factors and they are difficult to sort out Beliefs, perceptions, and experience constitute some of the variables associated with compliant medication taking behavior. It had been suggested that race, crime, age, and other environmental factors are associated with adherence and nonadherence; but these are speculative[9]. Arul studied various factors affecting the adherence in asthma patients and have identified five important factors regarding self-reported adherence with prescribed medications in patients with asthma: Age, gender, length of time with airway problems, whether the staff listen and take into account the patient's views concerning his/her asthma, and whether the patient has received information and education concerning asthma[9].

There were 88 patients with higher education (graduation) and all these patients had regular adherence with the therapy. Patient's having secondary education had a default rate of 62.28%, patients having primary education had a high default rate of 76.14%, while illiterate patients had a higher default rate 80%. Education status was thus a significant factor for the nonadherence to the therapy for asthma medications. Valid educational program for asthmatics can improve the knowledge of the disease and to understand how they look after themselves by careful evaluation of their own symptoms and respiratory function. Patients attending two lessons with helpful training tools can increase significantly asthma knowledge, treatment adherence, and patient self-management. In the present study economic status was significant and there was moderate correlation to adherence and low socioeconomic status. It was observed that there were higher default rates among lower socioeconomic class patients[9].

CONCLUSION

The percentage of regular adherence on therapy in asthma is 8.65%, and nonadherence is 89.05% which is significantly high. Regular adherence is an important aspect in the management and control of asthma, so patients should be advised to take regular and long term therapy for reducing the acute attacks of asthma and maintaining the disease state. Patients who have faith in the physician and the prescribed method of treatment are more likely to adhere to the treatment than patients who have a negative attitude toward treatment. The same is true of the parents of children with asthma. People with asthma should be offered education and written asthma action plans that focus on their individual needs this is a reinforcement of earlier advice.

REFERENCES

1. Chochrane GM. Adherence and outcomes in patients with asthma. *Drugs*, 1996; 52:S12,9.
2. Antonello N. The problem of adherence in the management of asthma: An educational ambulatory course called "The School of Asthma". *Tanaffos*, 2009; 8:14-18.

3. Barnes PJ, Jonsson B, Klim JB. The costs of asthma. *Eur Respir J* 1996; 9:636-642.
4. 2011 (Revised 2011). Available at: <http://www.ginasthma.com> [Accessed 20 Dec 2012].
5. Prasad BG. Changes proposed in the social classification of Indian families. *J Indian Med Assoc*, 1970; 55:198-199.
6. Pinto Pereira LM, Clement Y, Da Silva CK, McIntosh D, Simeon DT. Understanding and use of inhaler medication by asthmatics in specialty care in Trinidad: A study following development of Caribbean guidelines for asthma management and prevention. *Chest* 2002; 121:1833-840.
7. Rand CS, Wise RA. Measuring adherence to asthma medication regimens. *Am J Respir Crit Care Med.*, 1994; 149:S69-S76.
8. Stern L, Berman J, Lwry W, Katz L, Wang L, Rosenblatt L, *et al.* Medication adherence and disease exacerbation in patients with asthma: A retrospective study of managed care data. *Ann Allergy Asthma Immunol*, 2006; 97:402-8.
9. Arulprakasam KC and Senthilkumar N Self-reported Adherence, Management Behavior, and Barriers to Care After Hospital Visit by Children With Asthma, *Int.J. Pharm. and H. Care Res.*, 2017; 5(3):92-102.



54878478451190501



Submit your next manuscript to **IAJPR** and take advantage of:

Convenient online manuscript submission

Access Online first

Double blind peer review policy

International recognition

No space constraints or color figure charges

Immediate publication on acceptance

Inclusion in **ScopeMed** and other full-text repositories

Redistributing your research freely

Submit your manuscript at: editorinchief@iajpr.com

