



CODEN [USA]: IAJPB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

Available online at: <http://www.iajps.com>

Research Article

ASSESSMENT OF DIETARY HABITS OF EXPECTING WOMEN WITH ANEMIA

¹Waleed Javaid ²Asad Sarwar ³Hamza Tariq

¹Medical Officer Rural health centre 153 RB Chak Jhumra Faisalabad

Email ID: wjmalik7@gmail.com

²Medical officer DHQ hospital Faisalabad Email ID: docasadsarwar92@gmail.com

³Medical officer Allied hospital Faisalabad Email ID: hamza-tariq@live.com

Article Received: April 2019

Accepted: May 2019

Published: June 2019

Abstract:

Background: Anemia is a condition categorized by low hemoglobin level. Worldwide communities bear heavy burden especially of pregnant particularly in multigravida females. Among many other factors, dietary intake habits of expecting women contribute mainly to the anemia in such patients. So, a study to probe association of dietary intake and habits with anemia will be helpful for community.

Objective: To access the dietary habits of a multigravida pregnant female with anemia in patients of DHQ hospital, Faisalabad

Materials and Methodology: This cross-sectional study was conducted at District Head Quarter Hospital, Faisalabad from July 2017 to August 2017 in a period of six weeks. The research was carried out at hospitalized patients in Gynecology Unit 2. 120 patients were included after non-probability consecutive sampling and a questionnaire was filled after an interview by the primary investigator.

Results: A total of 120 female were interviewed using a predefined questionnaire during this study. Among them, 63% of women were anaemic and 37% of women were non-anaemic. Among anaemic women 13% were in their first trimester, 23% are in their second trimester and 27% are in their third trimester. About 92% of patients were taking normal 3 meals in a day while the rest of 23% patients had poor dietary habits with less than 3 meals in a day. 76% of women take chicken once a week. Out of 120 women, only 20% were consuming red meat twice a week while majority. 58% of women were taking citrus fruits daily as a source of vitamin C. Green vegetables were consumed by 102 women out of 120. 86% of women were taking beans and nuts as a part of their daily dietary intake.

Conclusion: This study clearly shows that dietary intake and habits of pregnant female directly contributes towards development of anemia.

Key Words: anaemia, multigravida, habits, diet, pregnancy,

Corresponding author:**Waleed Javaid,**

Medical Officer Rural health centre 153 RB Chak Jhumra Faisalabad

Email ID: wjmalik7@gmail.com.

QR code



Please cite this article in press Waleed Javaid et al., *Assessment Of Dietary Habits Of Expecting Women With Anemia.*, Indo Am. J. P. Sci, 2019; 06[06].

INTRODUCTION:

Anaemia is a condition that develops when blood lacks enough healthy red blood cells or haemoglobin. Haemoglobin is an important part of red blood cells and binds oxygen. If the body has very few or abnormal RBCs or the haemoglobin is abnormal or low, the cells will not get enough oxygen [1]. Symptoms of anaemia, like fatigue, occur because organs are not getting what they need to function properly. Anemia is defined by the World Health Organization as haemoglobin level < 11.0 g/dL.[1] Anaemia is the most common blood condition in the united states. It affects about 3.5 million of its population. Women, young children, and people with chronic diseases are at increased risk of anaemia. Women in the childbearing years are particularly prone to develop iron-deficiency anaemia because, during pregnancy, the body produces an increased amount of blood to support the development of the baby. If the mother is not getting enough iron or certain other nutrients due to a certain reason, the body might not be able to produce the required amount of red blood cells to make this additional blood. Older people also may have a greater risk of developing anaemia because of diet deficiencies and other medical conditions.[2]

There are many types of anaemia. All are different in their causes and treatments. Iron-deficiency anaemia, the most common type, is easily treatable with diet changes and iron supplements. Some grades of anemia like the mild anaemia (10.0-10.9 g/dl) that develops during pregnancy are considered normal. However, some anaemias may present lifelong health problems [3]. The overall prevalence of anaemia among the pregnant women was found to be 87.21%.[4] Factors such as religion, level of education of women and of their husbands and socioeconomic status were found to be significantly associated with the prevalence of anaemia in pregnancy ($P < 0.5$ only 0.7% were severely anaemic (haemoglobin < 7.0 g/dL). Non-anaemic women were significantly taller, weighed more, and had a higher body mass index [4].

The survey data showed that 84.9% of pregnant women ($n = 6,923$) were anemic (hemoglobin < 110 g/L); 13.1% had severe anemia (hemoglobin < 70 g/L), and 60.1% had moderate anemia (hemoglobin ≥ 70 to 100 g/L).[2] Among adolescent girls ($n = 4,337$) from 16 districts, the overall prevalence of anemia (defined as hemoglobin < 120 g/L) was 90.1%, with 7.1% having severe anemia (hemoglobin < 70 g/L) [2]. Unfortunately, very little efforts have been done in past in our country to find out relationship between development of anemia and dietary habits in pregnant females. This study aims to

find out the nutritional deficits that are the culprits behind anaemia in pregnancy. So that we can take prompt action to eliminate these causative factors and hence save the child and mother from long-term cardiovascular, muscular and other defects.

OBJECTIVE: To determine the dietary habits of multigravida pregnant females having anemia presenting in Gynecology department, DHQ hospital, University of Faisalabad

OPERATIONAL DEFINATION: None

ANEMIA: Any patient having haemoglobin concentration of less than 110 g/L (less than 11 g/dl) in venous blood was labeled as having anemia.

MATERIALS AND METHODOLOGY:

The cross-sectional research was done from July 2017 to August 2017 in a period of six weeks at Gynecology Ward No 2 of DHQ, Hospital, University of Faisalabad. Inclusion Criteria: 120 pregnant women of either trimester of age 20 to 30 having anemia as per operational definition were included in the study through non-probability sampling. Anemia in pregnancy was defined as a haemoglobin concentration of less than 110 g/L (less than 11 g/dl) in venous blood. Other variables like nutritional status, various dietary habits as mentioned in the predefined questionnaire and socio-economic status were also noted. Exclusion criteria: Patients with diagnosed hemolytic anemia like thalassemia, those with twin pregnancy or having chronic renal failure were excluded from the study. Data collected through questionnaire with interviews had close-ended questions. Study was performed after taking informed consents individually from patients before filling the questionnaire.

Data was entered and later analyzed using SPSS 20.0 version. Frequency and percentages were calculated for qualitative variables and mean with standard deviation was calculated for quantitative variables. Effect modifiers like age, number of pregnancies and socioeconomic status, were stratified and post-stratification chi-square was applied. Percentages were calculated and data were presented in the form of tables.

RESULTS:

A total of 120 female were interviewed over the questionnaire during this study. Pie charts were then drawn according to the results. The total sample size was 120 among them 76 (63%) of women were anaemic and 44 (37%) of women were non-anaemic.

Among anaemic women 15 (13%) were in their first trimester, 28 (23%) were in their second trimester and 33% were in their third trimester.

13% of women were taking caffeine daily. Only 42% of the women were taking 2 eggs as a part of their daily routine. Only 35% of women were taking iron supplements to meet their increased demands of iron.

Figure: no 1 is showing the number and percentage of patients having various dietary habits under our study related to anemia.

When women were asked about pica, 13% have craving for clay, 7% said they consume chalk, 10% said they consume ice, 33% were habitual of eating other items while 30% said that they don't crave for anything like this during pregnancy. 44% of women had a habit of eating raw uncooked rice. The inter-pregnancy difference was found to be >2yr in 37% of people while it was <2yr in 63%.

Table – I: is showing various dietary habits under study among the pregnant anemic patients.

None of them was having a family history of blood disorder like thalassemia or sickle cell anemia. Only 18% had a history of menorrhagia. 23% of women were having twin pregnancies. 42% of women were taking calcium supplements to fulfill the increased need during pregnancy.

Majority of women i.e 77% of women said that they take chicken once a week. Out of 120 women, only 6% were consuming red meat twice a week while majority i.e 94% were not taking it. 20% of women were taking citrus fruits daily as a source of vitamin C. Green vegetables were consumed by 70 women out of 120. 85% of women were taking beans and nuts as a part of their daily dietary intake. 72% of the patients were taking soft drinks daily. Out of 120,




Figure: no 1. Showing the number and percentage of patients having various dietary habits under our study related to anemia

Table – I: Various dietary habits of pregnant anemic patients

Dietary Routine	Status in pregnant females with anemia n =76	
	n	%
Chicken Consumption in a Week	56	73
Red Meat intake Twice a Week	2	2.6
Daily Intake of Citrus Fruit	8	10.5
Green Vegetable Twice a Week	42	55
Beans and Nuts Intake	60	78
Daily Soft Drinks Intake	54	71
Sea Food Consumption	32	42
Daily Caffeine Consumption	16	21
Two Eggs a Day	24	31
Iron Supplements Intake	27	35

DISCUSSION:

Our research based on the effect of dietary habits on the prevalence of anaemia. Anaemia is one of the major health issues that affect the pregnant women and which ultimately has some effect on fetus like IUGR, stillbirth as well as in pregnant women like fatigue and shortness of breath. Anaemia can also increase the risk of Caesarian -section and preterm delivery. There are multiple causes of anaemia like iron deficiency, folate deficiency, malaria, hookworm infestation. Among these causes, nutritional iron deficiency anaemia (IDA) is the commonest (90%) cause of anaemia in pregnancy [9]. Therefore, anaemia in a normal pregnant woman in this environment is usually attributed to iron deficiency, and successful treatment is often achieved with iron and folic acid without further investigations. [10]. Hb level is indicative of anaemia. The normal level of Hb is (12 to 16 g/dl). During pregnancy, this level falls (11.5 to 15 g/dl). This is because the blood volume increases by 50% in the course of pregnancy for providing essential nutrients for the developing baby [11]. The level of haemoglobin falls to 10.5 g/dl is quite normal in pregnancy. When it goes below 10g/dl, then it can cause health issues that affect the mother and child and the normal progression of the pregnancy. On the basis of Hb level anaemia can be mild, moderate and severe.

Based on WHO study 42% of pregnant women have anaemia. Dietary habits have a great effect on the prevalence of anaemia.[1] Our study found that 63% of multigravida are anaemic.

In the present study, the prevalence of anaemia was higher among pregnant women having a meal frequency of fewer than 3 times a day as compared to pregnant women who had a meal frequency of more than 2 times a day. This might be due to the reason

that pregnancy is a critical period with increased energy and nutrient demand for the mother which should be fulfilled with increased meal frequency per day. This result is consistent with other studies conducted in North Western Zone of Tigray, Northern Ethiopia.[3,5] In this study, the consumption of fruit daily was associated with a decreased risk of anaemia. This is because fruits are a rich source of non-heme iron. Fruits are also a rich source of vitamin C which enhances the absorption of iron in the body. And this result is consistent with the study conducted in Pakistan.

In our study women taking red meat twice a week have a high concentration of hemoglobin. Meat is a rich source of iron. This result is the same as a study conducted in Pakistan and Ethiopia. In our study, the prevalence of anaemia was higher among pregnant women who did not take iron and folate supplement during pregnancy as compared to those pregnant women who took their iron and folic acid supplementation. This result is the same as a study conducted in Ethiopia [5] and West Bengal. In our research, women having short interpregnancy difference are anaemic as compared to other women. This is consistent as a study conducted in Pakistan and Ethiopia.

In this study prevalence of anemia was higher in pregnant women in their second (23% out of 63%) and third (27% out of 63%) trimesters This is because during pregnancy the need for calorie and nutrients are increased to support increased maternal metabolism, blood volume and nutrients to fetus and this demand increases during the second and third trimester. In the first trimester, there is a marked decrease in the absorption of iron probably because of lower iron requirements and menstruation stops, saving a median of 0.56 mg Fe/day (160

mg/pregnancy) [13]. However, in the second-trimester iron absorption from a diet of very high iron bioavailability increases by 1.9mg/day and in the last trimester it increases by up to 5.0 mg/day [14]. In this study soft drinks, caffeine, pica has a negative impact on the Hb level and increases the risk of anaemia. In this study green vegetables, beans and nuts and seafood all have a positive impact on the Hb level. The rate of low birth weight babies was high in mothers who were anaemic in their third trimester. Preterm deliveries occurred highly in mothers who were anaemic in their second and third trimesters [15]. Therefore, the efforts should be made to improve the quality of education related to pregnancy and socioeconomic status of females.

CONCLUSION:

This study clearly shows that dietary intake and habits of pregnant female directly contributes towards development of anemia. Patients with anemia also have significant habit of pica. Females with more caffeine intake, short inter-pregnancy period and no iron supplement intake are more prone to have anemia in subsequent pregnancies.

LIMITATION OF THE STUDY:

This study was done on a small number of females because of non-availability of resources and manpower and various effect modifiers were ignored initially. We need to extend this study on a bigger population involving various other hospitals as well.

CONFLICT OF INTEREST:

Authors declare no conflict of interest

REFERENCES:

1. CC Dim, Onah HE. The prevalence of anemia among pregnant women *MedGenMed* 2007;9(3): 11
2. StartReading | Being The Parent <https://www.beingtheparent.com/start-reading/> not found.
3. Kefiyalew F, Zemene E, Asres Y, Gedefaw L. Anemia among pregnant women in Southeast Ethiopia: prevalence, severity and associated risk factors *BMC Res Notes* 2014;7:771.
4. Women attending antenatal clinics in an area with intense and highly seasonal malaria transmission in northern Ghana *tropical medicine and international health* 2009; 14 (6): 688–95.
5. Lebso M, Anato A, Loha E. Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: a community based cross-sectional study. *PLoS ONE* 12(12):e0188783.
6. McMahon LP. Iron deficiency in pregnancy. *Obstet Med* 2010;3(1):17-24
7. Maternal anaemia in various trimesters and its effect on newborn weight
8. Kumar KJ, Asha N, Murthy DS, Sujatha M, Manjunath V. Maternal anemia in various trimesters and its effect on newborn weight and maturity: an observational study *Int J Prev Med.* 2013;4(2):193-9
9. Bondevik GT, Ulstein M, Lie RT, Rana G, Kvåle G. The prevalence of anemia in pregnant Nepali women--a study in Kathmandu *Acta Obstet Gynecol Scand* 2000;79(5):341-9
10. Imdad A, Bhutta ZA Routine iron/folate supplementation during pregnancy *Paediatr Perinat Epidemiol* 2012;1:168-77
11. Pafitis *BMC Public Health* (not found)
12. Folic Acid in Obstetric Practice: A Review: Obstetrical & Gynecological https://journals.lww.com/.../Folic_Acid_in_Obstetric_Practice__A_Review.22.aspx by VS Talaulikar - 2011 - Cited by 37 (not found)
13. Lebso M, Anato A, Loha E. Prevalence of anaemia and associated factors among pregnant women *PLoS ONE* 2017;12(12): e0188783.
14. Toteja GS, Singh P, Dhillon BS, Saxena BN, Ahmed FU, Singh RP, et al. Prevalence of anemia among pregnant women and adolescent girls in 16 districts of India *Food Nutr Bull* 2006;27(4):311-5.
15. Ansari BN, Badruddin SH, Karmaliani R, Harris H, Jehan I, Pasha O, et al. Anaemia prevalence and risk factors in pregnant women in an urban area of Pakistan *Food Nutr Bull* 2008;29(2):132-9.
16. Sharma JB , Soni D, Malhotra M. Effect of dietary habits on the prevalence of anaemia in pregnant women *journal of obstetrics and gynaecol. Research* 2003;29(2):73-8.
17. Anjum A, Manzoor M, Manzoor N, Shakir HA. Prevalence of anaemia during pregnancy in district Faisalabad, Pakistan *Punjab Univ. J. Zool* 2015;30(1):015-20.