

# *Justicia thailandica*, a new species of Acanthaceae from Thailand

Yi Tong<sup>1</sup>, Yunfei Deng<sup>2,3</sup>

**1** School of Chinese Materia Medica, Guangzhou University of Chinese Medicine, Guangzhou 510006, Guangdong, China **2** Key Lab of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, Guangdong, China **3** Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences, Yezin, Nay Pyi Taw 05282, Myanmar

Corresponding author: Yunfei Deng ([yfdeng@scbg.ac.cn](mailto:yfdeng@scbg.ac.cn))

---

Academic editor: Alan Paton | Received 8 February 2019 | Accepted 3 May 2019 | Published 6 June 2019

**Citation:** Tong Y, Deng Y (2019) *Justicia thailandica*, a new species of Acanthaceae from Thailand. *PhytoKeys* 124: 11–22. <https://doi.org/10.3897/phytokeys.124.33745>

---

## Abstract

A new species of *Justicia* (Acanthaceae), *J. thailandica*, is described and illustrated from Thailand. The new species belongs to *Justicia* sect. *Harnieria* and is similar to *J. quadrifaria* and *J. championii*, but differs on account of the obviously densely white indumentum in the inflorescence bracts and calyx, ovate leaf blades with margin usually entire, spatulate inflorescence bracts and length ratio of calyx to mature capsule. It is assessed to be “Near threatened” (NE) according to IUCN Red List Category and Criteria. Pollen and seed morphology characters are also reported. Species of *Justicia* sect. *Harnieria* in Thailand are discussed and a key to the three recognized species is presented.

## Keywords

*Harnieria*, *Calophanoides*, new taxa, taxonomy

## Introduction

*Justicia* L. is the largest genus in the family Acanthaceae and consists of about 600 species distributed in tropical and temperate (to a lesser extent) regions of the world (Graham 1988; Hu et al. 2011; Mabberley 2017). It is characterized by the tubular

and bilabiate corolla with stylar furrow (rugula) in the upper lip, two stamens usually with the lower anther-theca spurred at base, “Knötchenpollen” pollen grains, and 4-(rarely 2-)seeded stalked capsules (Lindau 1894; Graham 1988; Hu et al. 2011). The recent molecular evidence (Deng et al. 2016; Kiel et al. 2017) indicated that the genus *Justicia* in the broad sense is polyphyletic and might be further separated into several independent genera. At the moment, we follow the treatment of Graham (1988) who divided the genus into sixteen sections.

Sect. *Harnieria* (Solms-Laubach) Benth. is characterized by the abbreviated axillary spikes, leaf-like inflorescence bracts, fusiform capsules and tuberculate seeds (Tong et al. 2016). It comprises approximately 76 species distributed in the tropical and subtropical regions of Africa and Asia with two species extending to Australia (Barker 1986; Hedrén 1989; Tong et al. 2016).

In the course of revising *Justicia* sect. *Harnieria* from Asia, some specimens collected from Thailand and identified in herbaria as *J. quadrifaria* (Nees) T. Anderson or *J. championii* T. Anderson appear to represent an undescribed species, which differs from the latter two species by the characters of indumentum, leaves, petiole, calyx, inflorescence bracts and length ratio of calyx to mature capsule.

## Materials and methods

The morphological comparison with related species in *Justicia* sect. *Harnieria* was based on studies of herbarium specimens and information gathered from literature. Pollen grains and seeds were taken from dried specimens (Beusekom *et al.* 3759, MO2366671) and mounted on aluminium stubs coated with gold in a sputter coater after being cleaned in water using ultrasound, and then examined using scanning electron microscopy (SEM; JSM-6360LV). The polar (P) axis and equatorial (E) diameter were measured by imaging analyzer (Smile View 2.1; JEOL Tokyo, Japan). Pollen terminology follows Erdtman (1969) and Punt *et al.* (2007). Seed terminology follows Hedrén (1989) and Rueangsawang *et al.* (2012).

## Taxonomic description


*Justicia thailandica* Y.F.Deng & Y.Tong, *sp. nov.*

urn:lsid:ipni.org:names:77197854-1

Figures 1, 2

**Type.** THAILAND. Kanchanaburi Province, Kanchanaburi District, Huay Bankau, 14°55'00"N, 98°45'00"E, mixed deciduous forest on limestone, 900 m alt., 13 Nov 1971, C. F. van Beusekom, C. Phengkklai, R. Geesink & B. Wongwan 3759 (holotype: MO2366671!; isotypes: BKF!, C!, K!, L!, P!).

**Diagnosis.** The new species is similar to *Justicia quadrifaria* (Nees) T. Anderson, but differs on account of the whole plant being white villous (not pubescent), leaf



**Figure 1.** *Justicia thailandica* **A** habit **B** magnifying the portion of leaf blades showing the tomentum **C** adaxial surface of inflorescence bract **D** abaxial surface of inflorescence bract **E** bract **F** bracteoles **G** calyx **H** corolla **I** dorsal view of the anther **J** frontal view of the anthers **K** pistil with nectary disc **L**, **M** opened capsule **N** capsule with calyx **O** seed. (Drawn by Cui Dinghan from the holotype van Beusekom et al. 3759).



**Figure 2.** *Justicia thailandica* **A** adaxial surface of inflorescence bract **B** abaxial surface of inflorescence bract **C** bract **D** bracteoles **E** stigma **F** pistil with nectary disc **G** calyx **H** dorsal view of anther **I** frontal view of anther **J** opened capsule **K** outside of capsule **L** inside of capsule showing retinacula **M** seed.

blade ovate (not oval, oblong to rarely ovate) with margin usually entire (not slightly undulate), apex shortly caudate or acute (not acuminate), base cuneate (not decurrent), petiole 8–12 mm (not 5–7 mm) long, calyx densely white villous (not pubescent), inflorescence bracts spatulate and villous (not ovate to obovate and sparsely pubescent) and capsule longer than the calyx (not shorter than the calyx). It is also similar to *J. championii*, but differs by the whole plant being white villous (not pubescent), and inflorescence bracts spatulate and villous (not obovate-spatulate with apex emarginate and sparsely pubescent).

Perennial herbs, 20–35 cm tall. Stems cylindrical or sometimes quadrangular, base decumbent and usually rooting at nodes then erect, densely white villous. Leaves opposite; petiole 0.8–1.2 cm long, villous; blades ovate, oval to sometimes lanceolate, 3.5–7.5 × 1.5–4 cm, papery, apex shortly caudate or acute, margin usually entire or slightly undulate, base cuneate to shortly attenuate, both surfaces densely white villous, mid-vein and secondary veins prominent on both surfaces, secondary veins (5

or) 6 on each side of mid-vein, covered with grayish white strip-like cystoliths. Spikes axillary, ca. 1 cm, usually several flowers in a cluster. Inflorescence bracts leaf-like, usually spatulate, rarely obovate, 7–8.5 × 4–5 mm, with a petiole 3–5 mm long, villous, pinnately veined with white strip-like cystoliths, apex round or sometimes obtuse, base decurrent onto petiole. Bracts and bracteoles linear, 1–1.2 mm long, white villous relatively sparsely below the middle. Calyx 7–9 mm, 5-lobed almost to base; lobes linear-lanceolate with conspicuous membranous margins, white villous, especially above the middle, apex acuminate. Corolla ca. 9 mm long, outside white villous, tube and upper lip white, lower lip white with purple spots; tube slightly longer than limb; upper lip triangular, minutely 2-lobed; lower lip 3-lobed, lobes imbricate, suborbicular. Stamens 2, attached to corolla tube, exerted; filaments ca. 2.2 cm long, basally villous; anther bithecos, thecae superposed, upper one smaller and muticous, lower one larger and with a white spur at base, anther connective pubescent. Ovary glabrous, green, fusiform with nectary flower disc bowl-shaped at base; style ca. 5 mm long, sparsely villous at base, stigma slightly 2-lobed. Capsule fusiform, 6–7 mm, glabrous, sometimes pilose at the tip. Seeds 4, compressed, ca. 1.1 × 1 mm, somewhat heart-shaped or obovate, brown or yellowish-brown, testa tuberculate.

**Etymology.** The specific epithet “*thailandica*” is derived from Thailand, where the new species is found.

**Phenology.** The new species was recorded in flowering from August to November and fruiting from October to December.

**Distribution and habitat.** *Justicia thailandica* is only known from Thailand (Fig. 4). It grows in the thickets on the limestone hills or evergreen forest at elevations of 400–900 m.

**Conservation status.** Currently, *Justicia thailandica* is only known from seven locations of Thailand with eight collections and the estimated extent of occurrence more than 20000 km<sup>2</sup>. We speculate that it may be widespread in Thailand, but is likely to qualify for a threatened category in the near future due to its vulnerable karst habitat and human activities leading to a decline in such habitats. It is therefore assessed as “Near threatened” (NE) according to the IUCN Red List categories and criteria (IUCN 2017).


**Additional specimens examined. THAILAND. Chiang Mai:** Chiang Dao District, Doi Chiang Dao, SE foothills near Ban Yang Pong Luang, 575 m alt., 30 Sep 1989, J. F. Maxwell 89-1169 (A, CAS, L). **Kanchanaburi:** Thong Pha Phum District, Krieng Kwia, 420 m alt., 27 Nov 1982, H. Koyama, H. Terao & T. Wongprasert 30402 (BKF, C, K). **Khon Kaen:** Phu Khieo Game Reserve, ca. 80 km east of Phetchabun, 16°50'00"N, 101°58'00"E, 850 m alt., 8 Nov 1984, G. Murata, C. Phengklai, S. Mitsuta, T. Yahara, H. Nagumasu & N. Nantasan T-41809 (A, BKF, TI). **Loei:** Nam Nao National Park, 101°23'00"–28°00"N, 16°48'00"–49°00"E, near check point of road to National Park, 280–350 m alt., 28 Oct 1984, Gen Murata, C. Phengklai, S. Mitsuta, T. Yaahara, G. Nagamasu & N. Nantasan T-51534 (TI); Pha Som Dej-Phataalern, Phu Luang NP, 1000 m alt., 14 Oct 2000, M. Norsaengsri 1075 (QBG). **Nakhon Ratchasima:** Pak Thong Chai District, Salika Forest, 40 km SE from Pak Thong Chai, 14°40'00"N, 102°2'00"E, 400 m alt., 25 Oct 1971, C. F. van Beusekom, Chan Wid

& R. Geesink 3362 (BKF, C, K, L, MO, P). **Phetchabun:** Nam Nao District, Nam Nao National Park, 25 Dec 1982, H. Koyama, H. Terao & T. Wongprasert T-31662 (BKF); Nam Nao, 24 Aug 2006, P. Chantaranothai et al. s.n. (BKF, QBG); Loam Gow District, Nahaw Now, 900 m alt., 17 Nov 1973, J. F. Maxwell 73-614 (AAU).

### Pollen and seed morphology

Pollen grains of *Justicia thailandica* are 2-colporate, bilaterally symmetrical, elliptic in both polar and equatorial view, polar axis [P]= $33.0 \pm 1.55 \mu\text{m}$ , equatorial diameter [E]= $21.6 \pm 0.85 \mu\text{m}$ , P/E=1.52±0.06, aperture area with 2 rows of 6–7 unequal-sized insulae, ornamentation of the insulae microreticulate with few scattered granules between muri (Fig. 3: A–B).

Seeds of *Justicia thailandica* are compressed, somewhat heart-shaped or obovate, brown or yellowish-brown, 1.1–1.2 × 1–1.1 mm, testa densely tuberculate with conspicuous rounded or oblong tubercles, ornamentation of tubercles irregular polygonal (Fig. 3: C–D).



**Figure 3.** Pollen and seed morphology of *Justicia thailandica* under SEM **A** pollen grain in equatorial view **B** exine ornamentation of pollen grain **C** seed **D** seed testa.

## Discussion


Characters of pollen grains and seed testa have proved valuable in sectional delimitation in the genus *Justicia* (Graham 1988). The pollen grains of *Justicia* sect. *Harnieria* are 2-colporate, reticulate exine ornamentation and traversed by 2 rows of unequal-sized insulae (Graham 1988; Hedrén 1989; Hu et al. 2005; Rueangsawang et al. 2013; Tong et al. 2016). Seed testa of *Justicia* sect. *Harnieria* is characterized by rugulose-tuberculate with the apices of the projections pointed (Graham 1988; Hedrén 1989; Rueangsawang et al. 2012; Tong et al. 2016), i.e. “Rugulose-tuberculate” type of Graham (1988). Our observation of the pollen and seed morphology (Fig. 3: A–D) is consistent with that of *Justicia* sect. *Harnieria* (Graham 1988; Hu et al. 2005; Hedrén 1989; Rueangsawang et al. 2012, 2013; Tong et al. 2016) and due to the characters of the abbreviated axillary spikes, leaf-like inflorescence bracts, fusiform capsules, we place the new species into that section.

Only three species of *Justicia* sect. *Harnieria* have been reported from Thailand. Hosseus (1908) reported *J. quadrifaria* from Thailand based on specimen “Hosseus et al. 228”. Subsequently, Imlay (1938) added *J. quadrifaria* var. *salicifolia* (T. Anderson) Imlay based on collections “Marcan 1528 and Lakshnakara 772”. Rueangsawang (2012) and Rueangsawang et al. (2012) recorded three species based on several specimens quoted in text, viz. *J. quadrifaria*, *J. neesiana* (Nees) T. Anderson and *J. championii*.

*Justicia championii* was first recorded in Thailand by Rueangsawang (2012) based on specimens (e.g. Wongprasert et al. 30402, Maxwell 73–614). However, *J. championii*, is currently known only from China and N Vietnam according to our worldwide specimen examination and is very similar to *J. quadrifaria*, but differs by the leaf shape and may be merged with the latter. After detailed comparison between the type specimen of *J. championii* and some Thai specimens (e.g. van Beusekom et al. 3759, Koyama et al. 30402, van Beusekom et al. 3362) identified in herbaria as *J. championii* or *J. quadrifaria*, we found they are obviously different and represent the new species described above. The clearest difference between *J. thailandica* and *J. championii* is that the calyx is densely white villous in *J. thailandica* (not pubescent) and inflorescence bracts are spatulate and villous in *J. thailandica* (not obovate with apex emarginate and sparsely pubescent). The leaves also tend to be different with the leaf blade ovate and white villous in *J. thailandica* (not lanceolate, oblong to oval and pubescent) with margin usually entire (not slightly undulate), apex shortly caudate or acute (not obtuse), base cuneate (not decurrent), (Fig. 5; Tab. 1). However, these leaf differences represent trends in a spectrum of variation rather than clear discontinuities.

Morphologically, *Justicia thailandica* is also similar to *J. quadrifaria*, but it can be easily distinguished from *J. quadrifaria* by the leaf blade being ovate and white villous (not oval, oblong to rarely ovate and pubescent), margin entire (not slightly undulate), petiole 8–12 mm (not 5–7 mm) long, apex shortly caudate or acute (not acuminate), base cuneate (not decurrent onto the petiole), calyx densely white villous (not pubescent), inflorescence bract spatulate and villous (not ovate to obovate and sparsely pubescent) and mature capsule longer than the calyx (not shorter than the calyx).

*Justicia quadrifaria* was recognized to be widely distributed in the tropical region of S to SE Asia from India, Indochina to Malay Archipelago and South China (Ridley



**Figure 4.** Distribution map of *Justicia thailandica* (black circle).

(1923; Hu et al. 2011). However, it is restricted to NE India and Bangladesh based on our worldwide specimen examination. *J. zollingeriana* (Nees) C. B. Clarke was reduced to *J. quadrifaria* or *Calophnodes quadrifaria* by some authors (Clarke 1907; Ridley 1923, Hu et al. 2011). However, it can be easily distinguished from the latter by the plant being glabrous (not pubescent in *J. quadrifaria*), calyx 4–5 mm (not 7–7.3 mm), capsule obviously more than 1.5 times longer than the calyx (not shorter than or subequal to the calyx) and leaves in each pair obviously unequal in size (not equal or subequal). *J. zollingeriana* is distributed in SE Asia from Thai Peninsular to Malay Archipelago. In Thailand, the species was first recorded as a synonym of *J. quadrifaria* by Hosseus (1908) based on specimen “Hosseus et al. 228”, and then some specimen (e.g. Kerr 7375, Garrett 316) of the species were reported under *J. quadrifaria* by Imlay (1938), while some specimen (Marcan 1528 and Lakshnakara 772) were under *J. quadrifaria* var. *salicifolia* by Imlay (1938).

*Justicia neesiana* recorded by Rueangsawang (2012) and Rueangsawang et al. (2012) is very similar to *J. multinodis* in the lanceolate leaves, however, it differs from the latter





**Figure 5.** Comparison between *Justicia thailandica*, *J. quadrifaria*, *J. championii* and *J. salicifolia* **A** holotype of *Justicia thailandica* (Beusekom et al. 3759, MO2366671) **B** isotype of *J. quadrifaria* (Wallich 2479a, GZU000251567) **C** holotype of *J. championii* (Champion 210, K000884038) **D** lectotype of *Justicia barapaniensis* P. Soumya & Sunojk. (a new name (Soumya 2017) for *J. salicifolia* T. Anderson. Hooker & Thomson s.n., K000884122.

by the plant being pubescent (not nearly glabrous in later), leaf 39–47 × 5–7.5 mm (not 22–37 × 2.2–4 mm), base decurrent onto the petiole (not cuneate), lateral leaf vein 5 with veinlet not reticulate (not usually 7 with veinlet obviously reticulate), petiole 5–6 mm (not nearly sessile), axillary spikes usually with 3–5 flowers (not 2–3 flowers) and inflorescence bracts subrotund to oval, persistent (not lanceolate, caducous).

In the course of revising Thailand species of sect. *Harnieria*, we have confirmed there are three species in Thailand, viz. *J. zollingeriana* (Nees) C. B. Clarke, *J. multinodis* R. Benoist and a new species, *J. thailandica*, described here.

A comparison of characters between *Justicia thailandica*, *J. quadrifaria*, *J. championii* and *J. zollingeriana* is provided in Table 1. An identification key to Thai species in sect. *Harnieria* is provided below.

**Table 1.** Comparison of *Justicia thailandica*, *J. quadrifaria*, *J. championii* and *J. zollingeriana*.

	<i>Justicia thailandica</i>	<i>J. quadrifaria</i>	<i>J. championii</i>	<i>J. zollingeriana</i>
<b>Opposite leaf</b>	equal or subequal	equal or subequal	equal or subequal	obviously unequal
<b>Leaf size</b>	3.5-7.5 × 1.5-4 cm	1-2 × 5.5-6.5 cm	1-7(-10.5) × 0.5-2(-3.5) cm	5-10 × 2-3.5 cm
<b>Leaf shape</b>	ovate, oval to sometimes lanceolate with margin entire, slightly undulate shortly caudate or acute	oval, oblong to rarely ovate with margin slightly undulate	lanceolate, oblong to oval and pubescent with margin slightly undulate	oblong to lanceolate with margin slightly undulate
<b>Leaf apex</b>	shortly caudate or acute	acuminate	obtuse	acuminate
<b>Leaf base</b>	cuneate or shortly attenuate	decurrent onto the petiole	decurrent onto the petiole	cuneate
<b>Petiole length</b>	8-12 mm	5-7 mm	5-15 mm	7-15 mm
<b>Lateral leaf vein</b>	(5)6	7(8)	6(7)	5(6)
<b>Indumentum</b>	densely white villous	pubescent	densely pubescent	glabrous
<b>Inflorescence bracts</b>	spathulate and densely villous	ovate to obovate and sparsely pubescent	obovate-spatulate with apex emarginate and sparsely pubescent	spathulate to round and glabrous
<b>Calyx</b>	7-9 mm, densely white long villous	7-7.3 mm, pubescent	7-9.5 mm, pubescent	4-5 mm, glabrous
<b>Calyx-capsule length ratio</b>	0.7-0.82	1.07-1.32	0.89-1.06	0.59-0.67
<b>Flowering</b>	Aug-Nov.	Unknown	Aug-Oct.	Jul-Sep.
<b>Fruiting</b>	Oct-Dec.	Unknown	Aug-Oct.	Jul-Sep.
<b>Distribution</b>	Thailand	India and Bangladesh	South China and north Vietnam	From Thai Peninsular to Malay Archipelago

**Identification key to Thai species in *Justicia* sect. *Harnieria***

- 1 Mature capsule more than 1.5 times longer than the calyx in length, leaves in each pair obviously unequal in size.....***Justicia zollingeriana***
- Mature capsule short than or sub-equal to the calyx, leaves in each pair sub-equal in size ..... **2**
- 2 Inflorescence bract and calyx lobes obviously with dense long villous hairs; leaf large, ovate, 3.5–7.5 × 1.5–4 cm.....***J. thailandica***
- Inflorescence bract and calyx lobes nearly glabrous or with sparsely pubescent hairs; leaf small, narrowly lanceolate, 3–7 × 0.6–0.9 cm.....***J. multinodis***

## Acknowledgements

The authors are very grateful to the curators of the following herbaria for their kind help in research facilities: A, BKF, C, CAS, K, L, MO, P, QBG and TI. Thanks are also given to Mr. Cui Dinghan (IBSC) for preparing the line drawing. This work was supported by the Natural Science Foundation of China (Grant Nos. 31700166, 31470302, 31670191) and Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences (Grant No. Y4ZK111B01).

## References

- Barker RM (1986) A taxonomic revision of Australian Acanthaceae. *Journal of the Adelaide Botanic Gardens* 9: 231–282. <https://www.jstor.org/stable/23873979>
- Clarke CB (1907) Materials for a flora of the Malayan peninsula. *Journal Asiatic Society Bengal* Pt. 2, Vol. LXXIV: 681.
- Deng YF, Gao CM, Xia NH, Peng H (2016) Acanthaceae, a new Chinese endemic genus segregated from *Justicia* (Acanthaceae). *Plant Diversity* 38(6): 312–321. <https://doi.org/10.1016/j.pld.2016.11.010> [PubMed]
- Erdtman G (1969) Handbook of palynology: an introduction to the study of pollen grains and spores. Munksgaard, 486 pp. <https://doi.org/10.1002/fedr.19710810815>
- Graham VAW (1988) Delimitation and infra-generic classification of *Justicia* (Acanthaceae). *Kew Bulletin* 43(4): 551–624. <https://doi.org/10.2307/4129957>
- Hedrén M (1989) *Justicia* sect. *Harnieria* (Acanthaceae) in tropical Africa. *Acta Universitatis Upsaliensis Symbolae Botanicae Upsalienses* 29(1): 1–141.
- Hosseus CC (1908) Die aus Siam bekannten Acanthaceen. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 41: 62–73. <https://www.biodiversitylibrary.org/item/703#page/73/mode/1up>
- Hu CC, Tsui HP, Xi YZ, Zhang YL (2005) Pollen morphology of one genus in Lepidogathideae, two in Andrographideae and eight in Justicieae (Acanthaceae) from China. *Acta Phytotaxonomica Sinica* 43(2): 151–162. <http://www.jse.ac.cn/EN/Y2005/V43/I2/151>
- Hu JQ, Deng YF, Daniel TF (2011) *Justicia*. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* Vol. 19. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 449–461. <http://flora.huh.harvard.edu/china/PDF/PDF19/Justicia.pdf>
- Imlay JB (1938) The taxonomy of the Siamese Acanthaceae. PhD Thesis, University of Aberdeen, Aberdeen.
- IUCN (2017) Guidelines for Using the IUCN Red List Categories and Criteria.version 13. Prepared by the Standards and Petitions Subcommittee of the IUCN Species Survival Commission. <http://cmsdocs.s3.amazonaws.com/RedListGuidelines.pdf> [Accessed 8 February 2019]
- Kiel CA, Daniel TF, Darbyshire I, McDade LA (2017) Unravelling relationships in the morphologically diverse and taxonomically challenging “justicioid” lineage (Acanthaceae: Justicieae). *Taxon* 66(3): 645–674. <https://doi.org/10.12705/663.8>

- Lindau G (1894) Beiträge zur systematik der Acanthaceen. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 18: 36–64. <https://biodiversitylibrary.org/page/203022>
- Mabberley DJ (2017) *Mabberley's Plant-book: a portable dictionary of plants, their classification and uses*. 4<sup>th</sup> edition. Cambridge University Press, Cambridge, pp 1102. <https://doi.org/10.1017/9781316335581>
- Punt W, Hoen PP, Blackmore S, Nilsson S, Thomas AL (2007) Glossary of pollen and spore terminology. *Review of Palaeobotany and Palynology* 143(1–2): 1–81. <https://doi.org/10.1016/j.revpalbo.2006.06.008>
- Ridley HN (1923) *The flora of Malay Peninsula Vol. II*. L. Reeve & Co., Ltd., London, 684 pp.
- Rueangsawang K (2012) Systematics of the genus *Justicia* L. (Acanthaceae) in Thailand. PhD Thesis, Khon Kaen University, 1–155.
- Rueangsawang K, Chantaranothai P, Simpson DA (2012) Contributions to the seed morphology and taxonomy of *Justicia* (Acanthaceae) from Thailand. *Journal of Systematics and Evolution* 50(2): 153–162. <https://doi.org/10.1111/j.1759-6831.2012.00178.x>
- Rueangsawang K, Chantaranothai P, Simpson DA (2013) Pollen morphology of *Justicia* L. (Acanthaceae) from Thailand and its taxonomic value. *Grana* 52(4): 275–288. <https://doi.org/10.1080/00173134.2013.819526>
- Soumya P, Sunojkumar P (2017) *Justicia barapaniensis*, a new name for *Justicia salicifolia* T. Anderson (Acanthaceae) *Phytotaxa* 332(1): 098–100. <https://doi.org/10.11646/phytotaxa.332.1.12>
- Tong Y, Huang YS, Deng YF (2016) *Justicia weihongjinii*, a new species of Acanthaceae from Guangxi, China. *Phytotaxa* 246(2): 127–136. <https://doi.org/10.11646/phytotaxa.246.2.4>