



CODEN [USA]: IAJ PBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**Available online at: <http://www.iajps.com>

Research Article

**ASSESSMENT OF THE OCCURRENCE OF CLINICAL
CHARACTERISTICS OF PATIENTS WITH DIABETIC
KETOACIDOSIS (DKA): A RETROSPECTIVE STUDY**Dr Mahwish Rasheed¹, Dr Sadia Arshad¹, Dr Shazia Hassan²¹ Punjab Medical College, Faisalabad, ² University Medical and Dental College, Faisalabad.

Article Received: April 2019

Accepted: May 2019

Published: June 2019

Abstract:

Objectives: The aim of this study was to assess the occurrence of clinical characteristics of patients with severe Diabetic Ketoacidosis (DKA) who were admitted to the intensive care unit (ICU) and investigating the relationship between para-clinical (glucose, anion gap, and serum bicarbonate) and clinical parameters in patients with severe DKA.

Method: A retrospective chart review of all adult patients with DKA who were admitted to the ICU at Allied hospital, Lahore between January 2018 and December 2018. The data collected include the demographic data, clinical presentation, precipitating factors, duration of hospital stay and mortality rate. The data were analyzed using STAT software.

Results: A total of 60 patients were included. Of these, 50 were men (83.3%). The median age was 23 years (ranging 18-29 years). Newly diagnosed diabetics accounted for 15 (25%) of the cases; the remainder was previously known patients of type1 diabetes on treatment. The main precipitating factors of DKA were insulin treatment cessation (87.5%) and infection/sepsis (39.6%). Serum blood glucose, serum bicarbonate level, and the calculated anion gap did not significantly correlate with clinical parameters of severe DKA.

Conclusion: Most patients with severe DKA who were admitted to the ICU of our institution presented with gastrointestinal symptoms. Non-compliance to insulin therapy was the main precipitating factor of DKA.

Keywords: Intensive Care Unit, Type-1 diabetes, Diabetic ketoacidosis.

Corresponding author:**Dr. Mahwish Rasheed,**

Punjab Medical College, Faisalabad.

QR code



Please cite this article in press Mahwish Rasheed et al., Assessment of the Occurrence of Clinical Characteristics of Patients with Diabetic Ketoacidosis (Dka): A Retrospective Study., Indo Am. J. P. Sci, 2019; 06(06).

INTRODUCTION:

Diabetic ketoacidosis (DKA) is the most common acute hyperglycemic complication of diabetes. According to a recent report,^[1] DKA affects approximately 8 per 1000 diabetics annually. It is associated with significant morbidity and mortality,^[2] with a worldwide mortality rate of 2-10%.^[3] Diabetic ketoacidosis is defined by the triad: hyperglycemia, acidosis and ketonuria. It is the most severe acute metabolic complication of diabetes mellitus.^[4] Current diagnostic criteria and classification of DKA are based on measurements of plasma glucose, arterial pH, serum bicarbonate, anion gap, and mental status alterations.^[5] Most cases of DKA are triggered by cessation of insulin and by infection, especially in developing countries.^[6] Mortality rate is lower in patients that received appropriate treatment: administration of insulin, correction of hypovolemia and electrolytes, and management of precipitating factors.^[7]

The aims of this study were to describe the clinical characteristics of patients with severe DKA who were admitted to the Intensive Care Unit at Allied hospital, Faisalabad. And to investigate the relationship between para-clinical (glucose, anion gap, and serum bicarbonate) and clinical parameters in patients with severe DKA.

METHODS:

This was a retrospective chart review Allied Hospital, Faisalabad between January 1, 2018 and December 31, 2018. The inclusion criteria were critically ill patient, above 13 years of age, with a

confirmed diagnosis of DKA. Diagnosis of DKA was based on latest criteria namely blood glucose concentration > 11 mmol/L, blood pH <7.3, and serum bicarbonate level < 15 mmol/L.^[4] The decision to admit patients with DKA to the ICU was made by the emergency department physician or internist in charge. Patients with a hyperosmolar hyperglycemic state were excluded from the study. The study was approved by the university's ethics committee.

Data collection: The information extracted included from patients' records demographic data, clinical presentation such as fever, abdominal pain, vomiting, dyspnea, altered sensorium and hypotension; as well as precipitating factors, such as cessation of insulin and sepsis, Sepsis being defined according to the Society of Critical Care Medicine / European Society of Intensive Care Medicine / American College of Chest Physicians consensus conference definitions. The length of hospital stay and mortality were also documented mechanical ventilation was accounted for.

Identification of laboratory and clinical parameters to determine the severity of diabetic ketoacidosis: We considered plasma glucose, serum bicarbonate levels and anion gap as indicators of the severity of DKA. The anion gap was calculated using the following formula: $[(Na^+) - (Cl^- + HCO_3^-)]$ (mEq/L). We also determined the severity of DKA based on the patient's level of consciousness (drowsy, stupor or coma) and the need for mechanical ventilation.

Table I: Clinical features of patients admitted to the intensive care unit with diabetic ketoacidosis.

Clinical Characteristics	Frequency (Percent)
Fever	19 (31.6)
Abdominal pain	50 (83.3)
Vomiting	55 (91.7)
Dyspnea	17 (28.3)
Low blood pressure	8 (13.3)
Altered sensorium	22 (36.6)
Cessation of insulin treatment	52 (86.6)
Infection	23 (38.3)
Need for mechanical ventilation	15 (25.0)

Abbreviation: CI, confidence interval.

Statistical analysis: Proportions for dichotomous variables and median (with interquartile range) for continuous variables were calculated to describe patients' characteristics. Associations

between different indicators of DKA (explanatory variables) and outcome variables, alterations in sensorium and need for mechanical ventilation were assessed using regression analysis. Multiple

logistic regressions were performed to control for potential confounding variables (age, sex, and presence of infection). To test for statistical significance, the 95% confidence interval was estimated.

All analyses were performed using STAT (StataCorp, College Station, TX, USA) software, version 12. To test for statistical significance, the 95% confidence interval was estimated. Results are expressed as frequency (percent) and mean \pm standard deviation.

RESULTS:

Patient characteristics: Sixty patients were included in the study; aged 18-29 years (median age is 23 years). Of these, 50 (83.3%) were men (Table-I). Newly diagnosed diabetics accounted

for 15 (25%) of the cases; the remainders were known cases of type 1 diabetes on insulin treatment. Three patients were on insulin pump therapy. The mean glycated haemoglobin of the sample was 10% (range, 8.6– 12.4%).

Clinical presentations: The most common presentation was gastrointestinal symptoms, including vomiting (n=55; 91.7%) and / or abdominal pain (n=50; 83.3%). Other reported symptoms include dyspnea (n=17; 28.3%), fever (n=19; 31.6%) and hypotension (n=8; 13.3%). Altered sensorium (altered consciousness, disorientation, or stupor / coma) was reported in 22 patients (36.6%). The average length of stay in the ICU was 2.5 days (range, 2-3 days). No mortalities occurred in this sample.

Table II: Association between indicators of diabetic ketoacidosis and alterations in sensorium among the sample.

Parameters	Crude OR (95% CI)
Blood Glucose Level	1.00 (0.94 – 1.06)
Serum Bicarbonate Level	1.02 (0.94 – 1.1)
Calculated Anion Gap	0.98 (0.93 – 1.03)

Adjusted for age, sex, and presence of infection. Abbreviations: CI, confidence interval; OR, odds ratio.

Table-III: Association between indicators of diabetic ketoacidosis and the need for mechanical ventilation among the sample.

Parameters	Crude OR (95% CI)	Adjusted* OR (95% CI)
Blood Glucose	1.01 (0.94 – 1.08)	1.01 (0.95 – 1.08)
Level Serum	1.00 (0.91 – 1.10)	0.99 (0.90 – 1.10)
Bicarbonate Level Calculated Anion Gap	1.02 (0.96 – 1.08)	1.05 (0.98 – 1.12)

Adjusted for age, sex, and presence of infection.

Abbreviations: CI, confidence interval; OR, odds ratio. Insulin cessation (n=52; 86.6%). Infection/sepsis were the second most common precipitating factor (n=23; 38.3%).

Correlation between laboratory parameters and clinical predictors of diabetic keto-

acidosis: Serum blood glucose, serum bicarbonate level, and calculated anion gap were independently correlated with the clinical predictors of severe DKA (Tables II and III).

DISCUSSION:

This retrospective study has described the clinical characteristics of patients with severe DKA who were admitted to the ICU of Allied Hospital, Faisalabad. On admission, over four-fifths of our patients presented with gastrointestinal symptoms (vomiting and / or abdominal pain episodes), while one-third presented with altered sensorium (altered consciousness, disorientation, or stupor / coma). These are similar to the reports of other authors that gastrointestinal symptoms and altered sensorium were the commonest presentations of DKA.^[8] However, the rate of gastrointestinal symptoms reported in our study is higher than that reported by these authors. We believe this disparity may be related to differences in sample characteristics. Fifteen patients (25%) of our sample were newly diagnosed type 1 diabetics with DKA as their first presentation, while only 18.2% of the patients in a study by Gavrielatos et al.^[8] were newly diagnosed type 1 diabetics with DKA as a first presentation.

The most commonly used diagnostic criteria for DKA are plasma glucose >250 mg/dL (>13.9 mmol/L), arterial pH <7.3, presence of ketonemia or ketonuria.^[9,10]

Alterations in sensorium and the need for mechanical ventilation are indicators of severe DKA, which has been reported in a retrospective matched cohort study.^[6] Altered.

Mental status in our study was considered to be a predictor of DKA severity. Although the mechanisms underlying the cause of altered sensorium in DKA are unclear, it may be due to different causes, including compromised cerebral blood flow, reduced cerebral glucose utilization, hyperosmolality, high blood glucose concentrations, acidosis, or a direct effect of ketone bodies among many other proposed theories and factors. The need for mechanical ventilation is another predictor of severe DKA in our study. This could be related to sepsis as a precipitating factor. The diagnostic criteria for sepsis were in accordance to the 2003 International Sepsis Definitions Conference,^[11] including altered mental status as a marker of global hypoperfusion. Tissue hypoperfusion is an important factor in the development of multi-organ failure, which is a measure of severity and the need for mechanical ventilation.^[12,13]

In previous studies, infection was the most commonly reported precipitating factor in DKA. Other reported precipitating factors include discontinuation of or inadequate insulin therapy,

cerebrovascular events, cardiovascular events and drugs.^[9] Furthermore, new-onset type 1 diabetes or insulin cessation in established type 1 diabetes typically results in the development of DKA.^[10] In our study, insulin cessation, either deliberately or inadvertently, was the main precipitating factor of DKA. While patients may be non-compliant to treatment for several reasons-financial constraints, non-availability of insulin, or use of alternative treatments-we believe that DKA can be prevented by educating diabetic patients and their family on the importance of optimal insulin therapy, dietary modification, providing psychosocial support, ensuring and monitoring glycemic control. Allied hospital offers free medical care to expatriates and persons with low economic status; this is why many patients who were admitted to the ICU for DKA were unable to afford treatment. Outcomes for our patients were favorable with no

reported deaths in the study group. DKA has a mortality rate of 1-5%.^[10,14] demonstrated in previous studies.

Mortality in DKA is rarely due to the metabolic complications of acidosis, but is mainly related to the underlying precipitating illness.^[9] Our study has a number of limitations that warrant consideration. First, it has all the limitations inherent in retrospective studies second, the relatively small sample size limits the study's statistical power. Finally, data available on the long-term outcomes of our patients after discharge from the ICU were not available.

CONCLUSIONS:

Overall, most patients with DKA who were admitted to the ICU of our institution presented with gastrointestinal symptoms Non-compliance to insulin therapy was the main precipitating factor in these patients, suggesting an important gap in health services delivery. We believe that non-compliant patients might benefit from patient centered strategies that are focused on simplifying access to medical care and education. Consequently, less privileged patients might benefit from a good healthcare system if a multidisciplinary program is created to alleviate the financial and technical burden of treating DKA. This program should also aim to increase compliance to insulin therapy by providing less privileged patients who lack medical insurance or financial stability with the necessary medication, follow up care and education.

REFERENCES:

1. Stipancic G, Sepec MP, Sabolic LL, Radica A, Skrabic V, Severinski S, et al. Clinical characteristics at presentation of type 1 diabetes mellitus in children younger than 15 years in Croatia J *Pediatr Endocrinol Metab*. 2011; 24(9-10): 665-670.
2. Wallace TM, Matthews DR. Recent advances in the monitoring and management of diabetic ketoacidosis. *QJM*. 2004; 97: 773–780 *Pak J Med Sci.*, 2015; 31(6). www.pjms.com.pk.
3. Yehia BR. Diagnosis and management of diabetic ketoacidosis in adults. *Hospital Physician*, 2008; 35: 21-26.
4. Wolfsdorf J, Craig M, Daneman D, Dunger D, Edge JA, Lee WR. ISPAD Clinical Practice Consensus Guidelines 2009 Compendium: diabetic ketoacidosis in children and adolescents with diabetes. *Pediatr Diabetes*, 2009; 10: 118-133.
5. de Vries L, Oren L, Lazar L, Lebenthal Y, Shalitin S, Phillip M. Factors associated with diabetic ketoacidosis at onset of Type 1 diabetes in children and adolescents. *Diabet Med.*, 2013; 30(11): 1360-1366.
6. Azevedo LC, Choi H, Simmonds K, Davidow J, Bagshaw SM. Incidence and long-term outcomes of critically ill adult patients with moderate-to-severe diabetic ketoacidosis. *J Crit Care*, 2014; 29(6): 971-977. doi: 10.1016/j.jcrc.2014.07.034.
7. Hara JS, Rahbar AJ, Jeffres MN, Izuora K E. Impact of a hyperglycemic crises protocol. *EndocrPract*, 2013; 19(6): 953-962. doi: 10.4158/EP13077.OR Gavrielatos G, Ioannidis I, Lionakis N, Avramidis D, Komitopoulos N, Varsamis E. Clinical And Laboratory Characteristics Of Diabetic Ketoacidosis In Adult Diabetic Patients. *Internet J Endocrinol*, 2006; 3(2).
8. Kitabchi AE, Umpierrez GE, Murphy MB, Kreisberg RA. Hyperglycemic crises in adult patients with diabetes: a consensus statement from the American Diabetes Association. *Diabetes Care.*, 2006; 29(12): 2739-2748.
9. Kitabchi AE, Umpierrez GE, Miles JM, Fisher JN. Hyperglycemic crises in adult patients with diabetes. *Diabetes Care.*, 2009; 32(7): 1335-1343. doi: 10.2337/dc09-9032.
10. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D. 2001 SCCM/ESICM/ACCP/ATS /SIS International Sepsis Definitions Conference. *Intensive Care Med.*, 2003; 29: 530-538.
11. Ezeani IU, Eregie A, Ogedengbe OS. Treatment outcome and prognostic indices in patients with hyperglycemic emergencies. *Diabetes Metab SyndrObes.*, 2013; 6: 303-307. doi: 10.2147/DMSO.S44477.
12. Glaser N, Bundros A, Anderson S, Tancredi D, Lo W, Orgain M, et al. Brain cell swelling during hypocapnia increases with hyperglycemia or ketosis. *Pediatr Diabetes*, 2014; 15(7): 484-493.
13. Wang J, Williams DE, Narayan KM, Geiss LS Declining death rates from hyperglycemic crisis among adults with diabetes, U.S., 1985–2002. *Diabetes Care*, 2006; 29(9): 2018–2022.