

A new species and new records of the genus *Alexeter* Förster (Hymenoptera, Ichneumonidae, Ctenopelmatinae) from Beijing with a key to Chinese species

Shu-Ping Sun¹, Tao Wang², Mao-Ling Sheng¹, Shi-Xiang Zong³

1 General Station of Forest and Grassland Pest Management (GSFGPM), National Forestry and Grassland Administration, 58 Huanghe North Street, Shenyang 110034, China **2** Mentougou Forestry Station, Beijing 102300, China **3** College of Forestry, Beijing Forestry University, Beijing 100083, China

Corresponding author: Mao-Ling Sheng (shengmaoling@163.com)

Academic editor: J. Fernandez-Triana | Received 30 March 2019 | Accepted 13 May 2019 | Published 1 July 2019

<http://zoobank.org/7AF35F44-0701-4CD4-904C-091928F3F5A5>

Citation: Sun S-P, Wang T, Sheng M-L, Zong S-X (2019) A new species and new records of the genus *Alexeter* Förster (Hymenoptera, Ichneumonidae, Ctenopelmatinae) from Beijing with a key to Chinese species. ZooKeys 858: 77–89. <https://doi.org/10.3897/zookeys.858.35012>

Abstract

A new species, *Alexeter beijingensis* Sheng, **sp. nov.**, and two new records for China, *A. angularis* (Uchida, 1952) and *A. shakojiensis* Uchida, 1930, collected in Mentougou, Beijing, belonging to the tribe Mesoleiini of the subfamily Ctenopelmatinae (Hymenoptera, Ichneumonidae), are reported. A key to the six species of *Alexeter* known from China is given.

Keywords

Mesoleiini, host, taxonomy

Introduction

Alexeter Förster, 1869, belonging to the tribe Mesoleiini of the subfamily Ctenopelmatinae (Hymenoptera: Ichneumonidae), comprises 32 species (Yu et al. 2016), of which eleven are from the Eastern Palaearctic Region (seven of them also occur in

the Western Palaearctic region) (Šedivý 1971, Uchida 1930, 1952, Yu et al. 2016), 18 from Western Palaearctic (Aubert 1998, Hinz 1996, Heinrich 1949, 1953, Meyer 1936), three from the Neotropical (Gauld et al. 1997), and seven from the Nearctic Region (Yu et al. 2016).

To date, three species are known from China (Chao 1976, Luo et al. 2019, Roman 1936). *Alexeter clavator* (Müller, 1776) was mentioned by Chao (1976) from Gansu, NW China, *A. multicolor* (Gravenhorst, 1829) from Jiangxi and Henan, S China, was reported by Luo et al. (2019), *A. segmentarius* (Fabricius, 1787) was reported from Gansu by Roman (1936) and Chao (1976).

The known hosts of *Alexeter* are sawflies, belonging to Diprionidae and Tenthredinidae (Aubert 1998, 2000, Constantineanu and Istrate 1973, Gauld et al. 1997, Hinz 1961, 1996, Yu et al. 2016).

In this paper we deal with all *Alexeter* species from China, including the description of a new species and a key to all known species from the country.

Materials and methods

Specimens were collected by interception traps (IT) (Li et al. 2012) in the forest of Mentougou, Beijing, P.R. China. The forest of Mentougou is composed of mixed deciduous angiosperms and evergreen conifers (Zong et al. 2013). Images were taken using a Leica M205A stereomicroscope with LAS Montage MultiFocus. Morphological terminology is mostly based on Gauld (1991).

The specimens of *A. clavator* (Müller, 1776), *A. coxalis* (Brischke, 1871), *A. multicolor* (Gravenhorst, 1829), *A. nebulator* (Thunberg, 1822), *A. niger* (Gravenhorst, 1829), *A. rapinator* (Gravenhorst, 1829), *A. segmentarius* (Fabricius, 1787) provided by Dr. Gavin Broad (The Department of Life Sciences, the Natural History Museum, London, UK) (**NHMUK**), were examined. The photos of the types, described by Uchida and deposited in Hokkaido University Museum, Hokkaido University, Japan, taken by Dr. Kyohei Watanabe (Kanagawa Prefectural Museum of Natural History, Odawara, Japan) (**KPMNH**), were examined and compared to the new species by the corresponding author.

Type specimens are deposited in the Insect Museum, General Station of Forest and Grassland Pest Management (**GSFGPM**), National Forestry and Grassland Administration, People's Republic of China.

Alexeter Förster, 1869

Type species. *Mesoleptus ruficornis* Gravenhorst, 1829 (= *segmentarius* Fabricius, 1787).

Diagnosis. (Förster 1869, Gauld et al. 1997, Townes 1970). Clypeus separated from face, apical margin blunt, weakly concave or centrally truncate. Occipital carina

dorsally complete, reaching to hypostomal carina distinctly above base of mandible. Epomia absent. Notaulus usually long and sharp. Mesopleuron with very fine to large punctures. Median longitudinal carina of propodeum usually distinct and complete, pleural carina complete. Fore wing vein 1cu-a opposite or distal of 1-M. Areolet present, or 3rs-m absent (Gauld et al. 1997). Hind wing vein 1-cu longer than cu-a. First tergite slender, median dorsal carina absent, or vestigial or indistinct. Glymma present. Second tergite with fine, weak, or indistinct punctures. Ovipositor short, with a distinct dorsal subapical notch.

Key to species of *Alexeter* known from China

- 1 Body brown to reddish brown. Postocellar line 1.3 times as long as ocular-ocellar line. Malar space approximately 0.2 times as long as basal width of mandible. Median longitudinal carina of propodeum distinct and complete. Third tergite approximately 1.3 times as long as maximum width. ***A. clavator* (Müller)**
- Body, at least mesosoma, black. Other characters variable. **2**
- 2 Median longitudinal carinae of propodeum complete, area superomedia distinctly constricted (Fig. 7) or expanded medially. Metasomal tergites black, or almost entirely black. **3**
- Median longitudinal carinae of propodeum almost parallel (Fig. 14), or divergent posteriorly, or incomplete (Fig. 13). Basal or median tergites, or all tergites brown to red-brown. **5**
- 3 Median longitudinal carinae of propodeum distinctly expanded medially (Fig. 12). Tegula, scutellum and postscutellum pale yellow. ***A. multicolor* (Gravenhorst)**
- Median longitudinal carinae of propodeum distinctly constricted medially. Tegula, scutellum and postscutellum black. **4**
- 4 Fore wing vein 2m-cu connecting to areolet basad of its posterior angle. Ovipositor sheath narrowed backwardly. Middle tarsus and hind leg entirely black. ***A. angularis* (Uchida)**
- Fore wing vein 2m-cu connecting to 4-M slightly distal of posterior angle of areolet. Ovipositor sheath parallel-sided. Subbasal portion of hind tibia widely white. Middle tarsus with at least median portion white or yellowish brown. Hind tarsomeres 2–4 and basal half of 5 white. ***A. beijingensis* sp. nov.**
- 5 Median longitudinal carinae of propodeum almost parallel (Fig. 14). Third tergite distinctly longer than its apical width. Hind femur and second and subsequent tergites reddish brown. Ovipositor sheath whitish yellow. ***A. shakojiensis* Uchida**
- Median longitudinal carinae of propodeum divergent backwardly or incomplete (Fig. 13). Third tergite shorter than its apical width. Hind femur, fifth and subsequent tergites black. Ovipositor sheath black. ***A. segmentarius* (Fabricius)**

***Alexeter beijingensis* Sheng, sp. nov.**

<http://zoobank.org/A36502D6-F805-4493-AFA6-4EAC2B14A464>

Figs 1–11

Etymology. The name of the new species is derived from the type locality.

Material examined. Holotype female, Mentougou, Beijing, 20 August 2004, leg. Tao Wang and Shi-Xiang Zong (GSFGPM).

Diagnosis. Apical portion of clypeus shiny, apical margin weakly and evenly concave. Outer profiles of middle and hind tibiae with distinct spines. Propodeum (Fig. 7) shagreened, area between median longitudinal carinae shiny, almost smooth; posterolateral portion with long and dense grey setae. Head, mesosoma and metasoma almost entirely black. Subbasal portions of all tibiae and median bands of middle and hind tarsi white.

Description. Female. Body length approximately 11.5 mm. Fore wing length 10.0 mm.

Head. Inner margins of eyes weakly indented opposite antennal sockets. Face (Fig. 2) 1.3 times as wide as long, uppermedian portion slightly convex; shagreened, with dense fine indistinct punctures; upper margin with a small median smooth tubercle. Clypeus (Fig. 2) transversely convex medially; basal portion shagreened, with indistinct short transverse wrinkle; apical portion smooth, shiny; apical margin weakly and evenly concave. Mandible with dense dark grey setae, lower tooth slightly longer than upper tooth. Malar space 0.5 times as long as basal mandibular width. Gena almost evenly convergent backward, in dorsal view approximately 0.8 times as long as width of eye. Vertex (Fig. 3) and frons with sculpture as that of face. Postocellar line 0.5 times as long as ocular-ocellar line. Antenna with 50 flagellomeres; ratios of lengths from first to fifth flagellomeres: 4.8:2.2:2.0:1.9:1.7; ultimate flagellomere twice as long as penultimate flagellomere. Occipital carina complete.

Mesosoma. Pronotum shagreened, upper portion with dense fine transverse wrinkles; hind margin of lateral concavity with short transverse oblique wrinkles. Epomia indistinct. Mesoscutum (Fig. 4) finely shagreened, with relatively sparse shallow punctures, distance between punctures mostly subequal to one diameter of puncture. Notaulus weak, present on front half of mesoscutum. Scutoscutellar groove with dense longitudinal wrinkles. Scutellum and postscutellum finely shagreened. Mesopleuron (Fig. 5) with sculpture almost as that of mesoscutum, with indistinct fine punctures. Speculum very small, upper portion shagreened. Mesopleural fovea vestigial. Upper end of epicnemial carina almost reaching anterior margin of mesopleuron, at level of upper 0.6 of pronotum. Metapleuron almost flat, with sculpture as that of mesopleuron; posterior margin with short transverse wrinkles. Submetapleural carina distinct, complete. Wings (Fig. 6) slightly infusate. Fore wing with vein 1cu-a distal to 1-M by 0.3 times length of 1cu-a. Areolet triangular, with long petiole, 0.4 times length of its height. 2m-cu slightly reclivous, connecting to posterior angle of areolet. Hind wing vein 1-cu 1.5 times as long as cu-a. Outer profiles of middle and hind tibiae with relative dense spines. Ratio of length of hind tarsomeres from first to fifth is



Figure 1. *Alexeter beijingensis* sp. nov. Holotype. Female. Habitus (without wings), lateral view.



Figure 2. *Alexeter beijingensis* sp. nov. Holotype. Female. Head, anterior view.



Figure 3. *Alexeter beijingensis* sp. nov. Holotype. Female. Head, dorsal view.



Figure 4. *Alexeter beijingensis* sp. nov. Holotype. Female. Mesoscutum and scutellum, dorsal view.




Figure 5. *Alexeter beijingensis* sp. nov. Holotype. Female. Mesosoma, lateral view.

4.0:2.0:1.5:0.8:1.0. Tarsal claws simple, hind claw strongly thick and curved (Fig. 8). Propodeum (Fig. 7) with distinct posterior transverse and strong complete median longitudinal carinae, latter strongly constricted medially. Area between median longitudinal carinae shiny, with indistinct, irregular transverse oblique fine striae. Area petiolaris




Figure 6. *Alexeter beijingensis* sp. nov. Holotype. Female **6** wings **6a** areolet.



Figure 7. *Alexeter beijingensis* sp. nov. Holotype. Female. Propodeum, dorsal view.


with irregular longitudinal wrinkles. Remainder with sculpture as that of mesopleuron. Posterolateral portion with long dense grey hairs. Propodeal spiracle circular.

Metasoma (Figs 9–11). Tergites shagreened. First tergite (Fig. 9) 2.4 times as long as apical width; median dorsal carina absent; dorsolateral carina indistinct, almost absent; spiracle distinctly convex, located slightly before mid of the tergite. Second tergite (Fig. 10) 1.1 times as long as apical width. Lateral margins of tergites 3 and 5, in dorsal view, almost parallel. Third tergite 1.1 times as long as apical width. Fourth tergite 0.8 times as long as apical width. Ovipositor sheath 0.5 times apical depth of metasoma. Ovipositor (Fig. 11) tapered from base to apex, with a large, deep, almost quadrangular notch.

Coloration (Fig. 1). Black, except for the following. Apical half of clypeus, anterior side and apical portion of fore femur red brown. Dorsoposterior portion of pronotum, fore tibia, basal halves of middle and hind tibiae except bases, mid tarsomeres 3 and 4, and hind tarsomeres 2–5 white. Apicomedian portion of scutellum yellowish white. Pterostigma and wing veins brownish black.



Figure 8. *Alexeter beijingensis* sp. nov. Holotype. Female. Apex of hind tarsus with claw, lateral view.



Figures 9. *Alexeter beijingensis* sp. nov. Holotype. Female. First tergite, dorsal view.



Figures 10. *Alexeter beijingensis* sp. nov. Holotype. Female. Tergites 2-8, dorsal view.



Figures 11. *Alexeter beijingensis* sp. nov. Holotype. Female. Apex of metasoma with ovipositor, lateral view.

Comparative diagnosis. The new species is similar to *A. niger* (Gravenhorst, 1829) in coloration: mesosoma and metasoma black; basal and apical portions of hind tibiae black, median portions white or yellowish white, but can be distinguished from the latter by the following combinations of characters: median longitudinal carina of propodeum complete, strong (absent or indistinct in *A. niger*); fore wing vein 2m-cu connecting to posterior angle of areolet (basad in *A. niger*); antenna, face and tegula black (antenna yellow-brown, face and tegula yellow in *A. niger*).

***Alexeter angularis* (Uchida, 1952)**

Material examined. CHINA: 1 female, Mentougou, Beijing, 29 September 2009, leg. Tao Wang.

Distribution. China, Japan. New record for China.

***Alexeter clavator* (Müller, 1776)**

Material examined. CHINA: 105 females, 121 males, Mt. Liupanshan, Ningxia Hui Autonomous Region, 7 July to 19 September 2005, IT. 1 female, Qinling, Shaanxi province, 5 July 2017, leg. Tao Li.

Distribution. China: Gansu, Ningxia, Shaanxi; Finland; Germany; Netherlands; Sweden; Switzerland.

***Alexeter multicolor* (Gravenhorst, 1829)**

Fig. 12

Material examined. CHINA: 1 female, 2 males, Tianzhu, Guizhou province, April 1996, leg. Yi-Han Li. 1 female, Jiulianshan, Jiangxi province, 5 August 2012, IT. 2 males, Wugongshan, 580 m, Jiangxi province, 16 May 2016, leg. Yu Yao. 1 male, Baotianman National Natural Reserve, 1300–1500 m, Henan province, 12 July 1998, leg. Mao-Ling Sheng.

Distribution. China: Guizhou, Henan, Jiangxi; Europe (Yu et al. 2016).

***Alexeter segmentarius* (Fabricius, 1787)**

Fig. 13

Material examined. CHINA: 1 male, Zhenfengshan, Helong, Jilin province, 2 August 1982, leg. Bing-Zong Ren.

Distribution. China: Gansu, Jilin; Mongolia, Russia, Europe (Yu et al. 2016)

***Alexeter shakojiensis* Uchida, 1930**

Fig. 14

Material examined. CHINA: 6 females, Mentougou, Beijing, 29 August to 22 September 2008, leg. Tao Wang. 1 female, 1 male, Mentougou, Beijing, 4 August to 8 September 2009, leg. Tao Wang. 1 male, Yanqing, Beijing, 12 June 2012, leg. Shi-Xiang Zong.

Distribution. China, Korea. New record for China.



Figure 12. *Alexeter multicolor* (Gravenhorst, 1829). Propodeum, dorsal view.




Figure 13. *Alexeter segmentarius* (Fabricius, 1787). Propodeum, dorsal view.




Figure 14. *Alexeter shakojiensis* Uchida 1930. Propodeum, dorsal view.

Acknowledgements

The authors are deeply grateful to Drs Jose Fernandez-Triana (Canadian National Collection of Insects Ottawa, Canada), Jacek Hilszczanski (Department of Forest Protection, Forest Research Institute, Raszyn, Poland), Andrei E. Humala (Forest Research Institute, Karelian Centre of Russian Academy of Science, Petrozavodsk, Russia), Kyo-

hei Watanabe (KPMNH) and one anonymous referee for reviewing this manuscript. The authors are also indebted to Dr. Gavin Broad (NHM) for his help while the first and corresponding authors were working in the NHM, and also thank Dr. Kyohei Watanabe for taking photographs of several types of *Alexeter* deposited in the Hokkaido University, Japan. This research was supported by Beijing's Science and Technology Planning Project (Z171100001417005) and the National Natural Science Foundation of China (NSFC, No. 31110103062, No. 31010103057).

References

- Aubert JF (1998) Huitième supplément aux ichneumonidés, Scolobatinae, principalement du Musée de Saint-Petersbourg (1). Bulletin de la Société Entomologique de Mulhouse 1998: 17–25.
- Aubert JF (2000) The West Palaearctic ichneumonids and their hosts. 3. Scolobatinae (= Ctenopelmatinae) and supplements to preceding volumes. Litterae Zoologicae 5: 1–310
- Chao HF (1976) An outline of the classification of the Ichneumon-flies of China (Hymenoptera: Ichneumonidae). Scientific Publisher, Beijing, 413 pp.
- Constantineanu MI, Istrate GI (1973) Contributi la cunoasterea Ichneumonidelor parazite i daunatorii molidului (*Picea excelsa* Link) din judetul Suceava. Studii si Comunicari Stiintele Naturii. Muzeul Judetean Suceava 3: 289–307.
- Gauld ID (1991) The Ichneumonidae of Costa Rica, 1. Introduction, keys to subfamilies, and keys to the species of the lower Pimpliform subfamilies Rhyssinae, Poemeniinae, Acaenitinae and Cylloceriinae. Memoirs of the American Entomological Institute 47: 1–589.
- Gauld ID, Wahl D, Bradshaw K, Hanson P, Ward S (1997) The Ichneumonidae of Costa Rica, 2. Introduction and keys to species of the smaller subfamilies, Anomaloninae, Ctenopelmatinae, Diplazontinae, Lycorininae, Phrudinae, Tryphoninae (excluding Netelia) and Xoridinae, with appendices on the Rhyssinae. Memoirs of the American Entomological Institute 57: 1–485.
- Heinrich GH (1949) Ichneumoniden des Berchtesgadener Gebietes. (Hym.). Mitteilungen Münchener Entomologischen Gesellschaft 35–39: 1–101.
- Heinrich GH (1953) Ichneumoniden der Steiermark (Hym.). Bonner Zoologische Beiträge 4: 147–185.
- Hinz R (1961) Über Blattwespenparasiten (Hym. und Dipt.). Mitteilungen der Schweizerischen Entomologischen Gesellschaft 34: 1–29.
- Hinz R (1996) Zur Systematik einiger Ctenopelmatinae (Hymenoptera, Ichneumonidae). Nachrichtenblatt der Bayerischen Entomologen 45(3/4): 75–78.
- Li T, Sheng M-L, Sun S-P, Chen G-F, Guo Z-H (2012) Effect of the trap color on the capture of ichneumonids wasps (Hymenoptera). Revista Colombiana de Entomología 38(2):338–342.
- Luo X-M, Sheng M-F, Ding D-S (2019) A new Chinese record of *Alexeter* Förster (Hymenoptera, Ichneumonidae). South China Forestry Science 47(1): 42–44.
- Meyer NF (1936) Parasitic Hymenoptera of the family Ichneumonidae of the USSR and adjacent countries. Part V. Opredeliteli po faune SSSR, Academy of Sciences of USSR, Leningrad, 340 pp. [In Russian]

- Roman A (1936) Swedish-chinesische wissenschaftliche Expedition nach den Nordwestlichen Provinzen Chinas. 58. Hymenoptera. II. Ichneumoniden. *Arkiv för Zoologi* 27A(40): 1–30.
- Šedivý J (1971) Ergebnisse der mongolisch-tschechoslowakischen entomologisch-botanischen Expeditionen in der Mongolei: 24. Hymenoptera, Ichneumonidae. *Acta Faunistica Entomologica Musei Nationalis Pragae* 14: 73–91.
- Townes HK (1970) The genera of Ichneumonidae, Part 3. *Memoirs of the American Entomological Institute* 13[1969]: 1–307.
- Uchida T (1930) Vierter Beitrag zur Ichneumoniden-Fauna Japans. *Journal of the Faculty of Agriculture, Hokkaido University* 25: 243–298.
- Uchida T (1952) Einige neue oder wenig bekannte Ichneumonidenarten aus Japan. *Insecta Matsumurana* 18: 18–24.
- Yu DS, van Achterberg C, Horstmann K (2016) Taxapad 2016, Ichneumonoidea 2015. Database on flash-drive. Nepean, Ontario. <http://www.taxapad.com>
- Zong S-X, Sun S-P, Sheng M-L (2013) A new species of *Amphibulus* Kriechbaumer (Hymenoptera, Ichneumonidae, Cryptinae) from Beijing with a key to species known from the Oriental and Eastern Palaearctic Regions. *ZooKeys* 312: 89–95. <https://doi.org/10.3897/zookeys.312.5294>