


# Accepted Manuscript

Food cravings in everyday life: An EMA study on snack-related thoughts, cravings, and consumption

Anna Richard, Adrian Meule, Julia Reichenberger, Jens Blechert



PII: S0195-6663(16)30602-X

DOI: [10.1016/j.appet.2017.02.037](https://doi.org/10.1016/j.appet.2017.02.037)

Reference: APPET 3355

To appear in: *Appetite*

Received Date: 19 October 2016

Revised Date: 21 January 2017

Accepted Date: 25 February 2017

Please cite this article as: Richard A., Meule A., Reichenberger J. & Blechert J., Food cravings in everyday life: An EMA study on snack-related thoughts, cravings, and consumption, *Appetite* (2017), doi: 10.1016/j.appet.2017.02.037.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Food cravings in everyday life: An EMA study on snack-related thoughts, cravings, and  
consumption

Anna Richard<sup>1,2</sup>, Adrian Meule<sup>1,2</sup>, Julia Reichenberger<sup>1,2</sup>, & Jens Blechert<sup>1,2</sup>

<sup>1</sup>Department of Psychology, University of Salzburg, Salzburg, Austria

<sup>2</sup>Centre for Cognitive Neuroscience, University of Salzburg, Salzburg, Austria

Correspondence:

Anna Richard, MSc

University of Salzburg

Department of Psychology

Centre for Cognitive Neuroscience

Hellbrunner Straße 34

5020 Salzburg, Austria

Phone: +43 662 8044 5159

Email: [anna.richard@sbg.ac.at](mailto:anna.richard@sbg.ac.at)

**Abstract**

Food craving refers to an intense desire to consume a specific food and is regularly experienced by the majority of individuals. Yet, there are interindividual differences in the frequency and intensity of food craving experiences, which is often referred to as trait food craving. The characteristics and consequences of trait and state food craving have mainly been investigated in questionnaire-based and laboratory studies, which may not reflect individuals' behavior in daily life. In the present study, sixty-one participants completed the Food Cravings Questionnaire-Trait-reduced (FCQ-T-r) as measure of trait food craving, followed by seven days of Ecological Momentary Assessment (EMA), during which they reported snack-related thoughts, craving intensity, and snack consumption at five times per day. Results showed that 86 percent of reported snacks were high-caloric, with chocolate-containing foods being the most often reported snacks. Individuals with high FCQ-T-r scores (high trait food cravers, HCs) thought more often about high-calorie snacks compared to low-calorie snacks whereas no differences were found in individuals with low FCQ-T-r scores (low trait food cravers, LCs). Further, the relationship between craving intensity and snack-related thoughts was stronger in HCs than in LCs. Higher craving intensity was associated with more consumption of snacks and again this relationship was stronger in HCs than in LCs. Finally, more snack-related thoughts were related to more frequent consumption of snacks, independent of trait food craving. Thus, HCs are more prone to think about high-calorie snacks in their daily lives and to consume more snack foods when they experience intense cravings, which might be indicative of a heightened responding towards high-calorie foods. Thus, trait-level differences as well as snack-related thoughts should be targeted in interventions on healthy eating choices and dieting.

**Keywords**

Food craving; Snack foods; Chocolate; Ecological Momentary Assessment; Food Cravings Questionnaire; Ecological validity

**Highlights**

- Food craving and snacking were examined via Ecological Momentary Assessment.
- Chocolate-containing foods were the most frequently desired foods in daily life.
- Thoughts about snacks and state cravings were associated with consumption.
- Trait cravers thought more often about high-calorie snacks than low-calorie snacks.
- Trait cravers consumed more snacks when they experienced intense state cravings.

1

## Introduction

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

In today's obesogenic environment, sugary and fat-rich snack foods are often consumed at quantities that go beyond homeostatic needs (Cleobury & Tapper, 2014; McKiernan, Houchins, & Mattes, 2008), pointing to the relevance of non-homeostatic determinants of food intake (Lowe & Butryn, 2007). One of these determinants is the experience of food craving, which refers to an intense desire to consume a specific food (White, Whisenhunt, Williamson, Greenway, & Netemeyer, 2002) and which can occur in the absence of hunger (Pelchat & Schaefer, 2000). Food cravings are prevalent in societies characterized by abundant food environments (Pelchat, 1997; Weingarten & Elston, 1991) and craved foods are usually high in sugar and fat, with chocolate being the most frequently craved food in Western societies (Rozin, Levine, & Stoess, 1991). Food cravings are quite specific in that they can be satisfied only by the craved—or very similar—food (Bruinsma & Taren, 1999). As food cravings have been associated with past dieting failures (Meule, Westenhöfer, & Kübler, 2011) and prospectively predict increased food intake and weight gain (Boswell & Kober, 2016; Meule, Richard, & Platte, 2017), knowledge about the characteristics of food cravings is relevant for health behaviors in various fields.

17

18

19

20

21

22

23

24

25

Food cravings are multidimensional experiences with cognitive, motivational, and behavioral facets (Cepeda-Benito, Gleaves, Williams, & Erath, 2000). According to the elaborated intrusion theory of desire, craving is a result of a cognitive elaboration of intrusive thoughts about a desired object (Kavanagh, Andrade, & May, 2005; May, Andrade, Panabokke, & Kavanagh, 2004), highlighting the role of food-related thoughts as a prerequisite for the emergence of food cravings. Thus, food cravings are not necessarily triggered by the presence of food stimuli, but can also occur spontaneously through mental imagery of the craved foods (Hallam, Boswell, DeVito, & Kober, 2016). These thoughts can also trigger consumption of the craved food, particularly when cravings are intense

26 (Appelhans, French, Pagoto, & Sherwood, 2016; Hofmann & Van Dillen, 2012; Papies,  
27 Stroebe, & Aarts, 2007), but this link is probably less tight than the one between thoughts and  
28 cravings. There are also marked and stable interindividual differences in the frequency and  
29 intensity of food craving experiences, suggesting that some individuals think more frequently,  
30 crave more intensely and—as a consequence—likely consume more snack foods than others.  
31 Recent theorizing refers to such differences as tonic or *trait food craving* (Boswell & Kober,  
32 2016; Hallam et al., 2016). Importantly, whereas state food cravings are exclusively  
33 experienced as transient states in a particular moment, trait food craving refers to the  
34 experience of food craving in general.

35         Laboratory and questionnaire-based studies revealed that individuals with elevated  
36 levels of trait food craving (i.e., high trait food cravers) seem to have a preference for high-  
37 calorie foods and are more susceptible to experience food cravings spontaneously or when  
38 confronted with external food cues. For instance, relative to low trait food cravers, high trait  
39 food cravers displayed an implicit approach tendency towards high-calorie foods  
40 (Brockmeyer, Hahn, Reetz, Schmidt, & Friederich, 2015) and showed more reward-related  
41 brain activity during food picture viewing (Ulrich, Steigleder, & Grön, 2016) and chocolate-  
42 related thought suppression (Miedl, Blechert, Meule, Richard, & Wilhelm, submitted). High  
43 trait food cravers also reported higher craving intensity when they were asked to imagine their  
44 favorite food (Tiggemann & Kemps, 2005) or were exposed to pictures of palatable foods  
45 (Meule, Hermann, & Kübler, 2014; Meule, Skirde, Freund, Vögele, & Kübler, 2012).  
46 Similarly, high trait chocolate cravers displayed more positive implicit attitudes towards  
47 chocolate (Richard, Meule, Friese, & Blechert, in revision) and had difficulties disengaging  
48 their attention from chocolate cues (Kemps & Tiggemann, 2009).

49         Although experiencing a food craving does not always result in subsequent food  
50 intake, previous studies reported positive associations between state (Meule & Hormes, 2015)

51 and trait food craving (e.g., Martin, O'Neil, Tollefson, Greenway, & White, 2008) with  
52 consumption of the given food in the laboratory. However, various situational and individual  
53 factors can affect whether craved foods are actually consumed (Hill, 2007). For example, the  
54 role of trait food craving on the relationship between state food craving and subsequent  
55 consumption has received little attention. Thus, further research is needed on how closely  
56 state cravings, thoughts about foods, and food consumption are interrelated as a function of  
57 trait food craving in everyday life.

58 Previous studies on the assessment of food cravings in everyday life, however, used  
59 paper-and-pencil methods such as the craving record sheet (Hill, Weaver, & Blundell, 1991).  
60 Here, participants entered craving episodes by hand each time they had felt an urge to  
61 consume a specific type of food regardless of consumption. It was found that momentary  
62 cravings co-occurred with thoughts about craved foods or the presence of these foods and  
63 these cravings were often followed by consumption (Hill & Heaton-Brown, 1994). In a more  
64 recent study, both trait-level and state-level cravings were associated with consumption  
65 (Forman, Hoffman, Juarascio, Butryn, & Herbert, 2013), demonstrating the behavioral  
66 consequences of food cravings as well as the presence of interindividual differences.  
67 Similarly, total energy intake was higher in female trait food cravers than in non-cravers in a  
68 study using a three-day food record (Lafay et al., 2001), which mainly resulted from  
69 consumption of between-meal snack foods.

70 Although these studies provided useful information about food cravings in naturalistic  
71 settings, findings are limited by shortcomings of paper-and-pencil measurements, such as low  
72 compliance rates and under-reporting of craving episodes (Berkman, Giuliani, & Pruitt,  
73 2014), decreases in compliance across the study period (Massey & Hill, 2012), or backfilling  
74 of past events (Stone, Shiffman, Schwartz, Broderick, & Hufford, 2003). Smartphone-based  
75 Ecological Momentary Assessment (EMA) in daily life may solve most of these limitations

76 by including electronic prompting and, thus, appear suitable to address transient phenomena  
77 (i.e., state food cravings). In addition, hierarchical linear modeling was used in the current  
78 study to acknowledge both between-person (here: trait food craving) and within-person  
79 information (here: associations between craving intensity, snack-related thoughts, and  
80 consumption). Specifically, participants completed the Food Cravings Questionnaire-Trait-  
81 reduced (FCQ-T-r), followed by seven days of EMA by use of signal-contingent sampling. At  
82 five times per day, they indicated the amount of thoughts about snacks, craving intensity, and  
83 snack consumption via their smartphone devices.

84         The current study had three aims for characterizing food cravings in everyday life. A  
85 first aim was to examine the type of snack foods that participants craved most frequently.  
86 Based on previous questionnaire-based studies (Rozin et al., 1991; Weingarten & Elston,  
87 1991), it was expected that the majority of craved snacks would be high-caloric and that the  
88 most frequently craved food category would be chocolate. A second aim was to expand the  
89 conceptual understanding of food cravings outside the laboratory. Specifically, the  
90 interrelations between thoughts about snacks, craving intensity, and snack consumption were  
91 examined (arrows A, B, and C in Figure 1A). Given that a cognitive elaboration of food-  
92 related thoughts is essential for the emergence of food craving (Kavanagh et al., 2005; May et  
93 al., 2004), it was hypothesized that episodes with more snack-related thoughts would be  
94 characterized by higher craving intensity. As food cravings usually involve high-calorie  
95 foods, it was expected that the relationship between thoughts about snacks and craving  
96 intensity would be particularly pronounced when energy-dense snack foods were thought  
97 about. As thoughts about food and more intense food cravings are associated with higher  
98 intake of the craved food (Forman et al., 2013; Hill & Heaton-Brown, 1994; Meule &  
99 Hormes, 2015), it was further expected that higher craving intensity (arrow B in Figure 1A)  
100 and more thoughts about snacks (arrow C in Figure 1A) would result in higher snack  
101 consumption. A third aim was to examine associations between trait food craving and type of

102 craved snacks, thoughts about snacks, craving intensity, and consumption of snacks (solid  
103 gray arrows in Figure 1A), which would be indicative of the ecological validity of the concept  
104 of trait food craving. Based on previous laboratory and questionnaire-based studies  
105 (Brockmeyer et al., 2015; Martin et al., 2008; Meule, Hermann, et al., 2014; Richard et al., in  
106 revision), it was expected that individuals with high scores on the FCQ-T-r (i.e., high trait  
107 food cravers) would report more frequent thoughts about high-calorie snacks, crave more  
108 intensely, and also consume more snack foods than low trait food cravers. Finally, it was  
109 explored whether trait food craving moderated associations between thoughts about snacks,  
110 craving intensity, and consumption of snacks (dashed gray arrows in Figure 1A). Specifically,  
111 it was assumed that the relationships between craving intensity, thoughts about snacks, and  
112 snack consumption may be more pronounced in high trait food cravers than in low trait food  
113 cravers.

114 **Methods**115 **Participants**

116 Sixty-six university students were recruited through flyers shared on social media  
117 platforms, bulletin boards, and students' mailing lists. The following inclusion criteria were  
118 defined: a) female, b) aged between 18 and 30 years, c) no self-reported lifetime mental  
119 disorders. Five participants were excluded due to either technical failures with the smartphone  
120 device ( $n = 1$ ), response rates  $< 50\%$  ( $n = 2$ ), or not following the study protocol correctly ( $n =$   
121  $2$ ). The final sample consisted of 61 participants (mean age = 21.6 years,  $SD = 2.47$ ; mean  
122 body mass index = 21.2 kg/m<sup>2</sup>,  $SD = 2.50$ )<sup>1</sup>. All participants signed informed consent before  
123 commencing the study. Ethical approval for the study was granted by the ethics committee of  
124 the University of Salzburg.

125 **Questionnaires**

126 **Food Cravings Questionnaire-Trait-reduced.** The FCQ-T-r (Meule, Hermann, et  
127 al., 2014) was used to measure the frequency of food craving experiences in general. It  
128 consists of 15 items (e.g., "I find myself preoccupied with food.", "If I eat what I am craving,  
129 I often lose control and eat too much."), and responses are scored on a 6-point scale (from 1  
130 [*never/not applicable*] to 6 [*always*]). Higher scores indicate more frequent food cravings.  
131 The FCQ-T-r demonstrated high retest-reliability over six months (Meule, Beck Teran, et al.,

---

<sup>1</sup> Including age as a covariate revealed that it was negatively associated with snack-related thoughts ( $\beta_{10} = -0.03$ ,  $p = .029$ ) and snack consumption ( $\beta_{10} = -0.07$ ,  $p = .043$ ). However, age neither interacted with the predictor variables (i.e., craving intensity, calorie type, snack-related thoughts, or trait food craving; all  $ps \geq .499$ ) nor did it change the pattern and significance of the results. Thus, results are presented without age as covariate.

132 2014), supporting the stability of trait-level food craving experiences. Internal consistency  
133 was  $\alpha = .907$  in the current study.

134 **Craving for high-calorie foods.** Participants were given a list of 10 high-calorie  
135 snacks (chocolate, ice cream, pizza, noodles, pastries, cookies, sweets, chips, French fries,  
136 cake; Meule, Vögele, & Kübler, 2012) to examine frequency of craving experiences for  
137 specific foods. They were asked to indicate how often they generally experience an intense  
138 desire to consume each of the listed foods on a 6-point scale (1 [*never/not applicable*] to 6  
139 [*always*]).

#### 140 **EMA Measures**

141 **Snack-related thoughts, craving intensity, and consumption.** At each of the five  
142 daily prompts, participants indicated how often they had thought about a snack since the last  
143 prompt (i.e., 3 hours) by numerically entering the *number of thoughts* about snack foods. If  
144 participants reported at least one (or more) snack-related thought, they were asked to name the  
145 snack they had thought about in a text box. Snacks were defined as foods that were not  
146 consumed as part of a principal meal (i.e., breakfast, lunch, or dinner). We explicitly did not  
147 constrain snack foods to specific categories, allowing the participants to freely report every  
148 snack they had thought about. Snack-related thoughts should be entered regardless of an  
149 actual consumption. Participants next rated their desire to consume the snacks (i.e., *craving*  
150 *intensity*) they had thought about on a continuous slider (from 0 [*not at all intense*] to 100  
151 [*very intense*]). Further, they specified the number of snack thoughts that had internally and  
152 spontaneously been evoked (i.e., in the absence of external triggers such as the sight or smell  
153 of foods). Lastly, participants reported the *number of consumed snacks* during the respective  
154 period since the last prompt without specifying the specific type they had consumed.

155 **Principal meals and hunger.** Principal meals were assessed so that participants  
156 would be able to differentiate thoughts about snacks (i.e., amount, type, and intensity) from

157 other eating episodes (Massey & Hill, 2012). Participants reported how often they had thought  
158 about a principal meal since the last prompt by numerically entering the number of thoughts  
159 about principal meals. Afterwards, they were asked whether and, if so, what kind of principal  
160 meal (breakfast, lunch, or dinner) they had eaten. Further, participants rated their hunger since  
161 the last prompt on a continuous slider (from 0 [*not at all hungry*] to 100 [*very hungry*]).

## 162 **Procedure**

163 Participants completed a set of questionnaires online, including demographic questions  
164 and questions on trait food craving (i.e., FCQ-T-r and craving for high-calorie foods). Next,  
165 they were trained on the EMA protocol, the usage of the smartphone application, and some  
166 specific concepts (e.g., distinction between principal meals and snacks as well as definition of  
167 food craving as an intense desire to consume a specific type of food) in a face-to-face session  
168 or telephone call. Participants completed seven days of EMA, preceded by a training day  
169 (data discarded). The signal-contingent protocol implemented five daily signal times (set at 10  
170 a.m., 1 p.m., 4 p.m., 7 p.m., and 10 p.m.). Delayed responses to signals triggered reminders  
171 every 10 minutes until 1 hour had elapsed. At each prompt, participants answered questions  
172 regarding their current hunger, thoughts about principal meals and snack foods, craving  
173 intensity, and snack consumption. At the end of the study, participants completed questions  
174 about reactivity and were reimbursed with course credits or €15,-.

175 **Data Analyses**

176 Reported snacks were categorized as high- or low-caloric by examining energy density  
177 (kilocalories per 100 grams).<sup>2</sup> If participants specified more than one snack they had thought  
178 about per prompt, calorie content was averaged over all snacks reported per signal.

179 To analyze the nested, longitudinal structure of the data and to test the conceptual  
180 model (Figure 1A), hierarchical linear models were applied using the software HLM7  
181 (Raudenbush, Byrk, & Congdon, 2011). When participants reported no thoughts about snacks  
182 (indicated by 0), the respective prompts were disregarded because there were no  
183 corresponding data for craving intensity and consumption. To test whether the amount of  
184 thoughts about snacks changed across the study period (i.e., reactivity to the EMA measures),  
185 the effect of measurement point (i.e., days across the study period; 0 = day 1 to 6 = day 7) on  
186 thoughts about snacks was tested at Level 1.

187 To address the first aim (i.e., types of craved snacks foods) and second aim (i.e.,  
188 interrelations between thoughts about snacks, craving intensity, and consumption of snacks)  
189 the following set of analyses was run: snack-related thoughts, craving intensity, and  
190 consumption were modeled within individuals at Level 1 (arrows A, B, and C in Figure 1A).  
191 Specifically, we modeled craving intensity as predictor of thoughts about snacks at Level 1  
192 (Table 3). In a next step, type of snacks (0 = low-calorie, 1 = high-calorie) was modeled as a

---

<sup>2</sup> There were two exceptions to this rule: crispbread (334 kcal/100g) and rusk (365 kcal/100g), which have a high energy density per 100g, were classified as low-calorie snacks as they have a small recommended portion size (e.g., 25g). Low-calorie snacks ranged from tomato (17 kcal/100g) to raw ham (145 kcal/100g). High-calorie snacks ranged from jam sandwich (150 kcal/100g) to peanut butter (626 kcal/100g). Low-calorie snacks ( $M = 59.6$  kcal/100g,  $SD = 25.2$ ) and high-calorie snacks ( $M = 406$  kcal/100g,  $SD = 114$ ) differed in kcal/100g,  $t_{(764)} = 30.1, p < .001$ .

193 further Level 1 predictor (Table 4). Last, we modeled thoughts about snacks, craving  
194 intensity, and their  $z$ -standardized interaction as predictors for consumption of snacks at Level  
195 1 (Table 5). As participants were not asked to specify the snacks they had consumed, type of  
196 snack was not considered as a predictor in the last analysis.

197 To address the third aim (i.e., ecological validity of trait food craving), FCQ-T-r  
198 scores were added to previous analyses at Level 2 to examine main effects (solid gray arrows  
199 in Figure 1A) and moderating effects (dashed gray arrows in Figure 1A) on the interrelations  
200 of craving intensity, snack-related thoughts, and consumption (Tables 3, 4, and 5).  
201 Subsequently, to control for general feelings of hunger, reported hunger level was entered as  
202 additional Level 1 predictor for thoughts about snacks and consumption, respectively. The  
203 five signals per day (Level 1) were nested within participants at Level 2. Slopes and intercepts  
204 were allowed to vary randomly across participants. The  $z$ -standardized predictors as well as  
205 the variables type of snacks and measurement point were entered uncentered into the models.  
206 All other Level 1 predictors were person-mean centered and Level 2 predictors were grand-  
207 mean centered. More detailed information on model equations and their interpretation can be  
208 found in the appendix.

209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232

## Results

### Compliance and Reactivity

Participants responded to 1,870 signals of all possible EMA prompts ( $N = 2,135$ ), reflecting a compliance rate of 87.6% ( $SD = 8.87\%$ , range: 60.0–100%). Among these signals, participants indicated at least one snack-related thought for 768 (41.1%) prompts and consumption of at least one snack for 558 (29.8%) prompts. As participants were allowed to enter more than one snack they had thought about, a total of 1,057 distinguishable snack foods were counted (Table 2). No effect of measurement point was found on the amount of snack-related thoughts ( $\beta_{10} = -0.05$ ,  $p = .155$ ), indicating that individuals did not alter their behavior in response to the EMA measures over time. Similarly, when asked about reactivity at the end of the study, participants reported that the prompts did not change their thoughts about snack foods ( $M = 4.30$ ,  $SD = 2.51$  on a scale ranging from 1 [*not at all*] to 11 [*very much*]).

Descriptive statistics of the included variables are presented in Table 1.

### Aim 1: Types of Craved Snack Foods

Our first aim was to examine the characteristics of momentary cravings (i.e., frequency and specificity) in the individuals' daily routines. On average, individuals thought about 5.75 snacks ( $SD = 5.08$ ) and consumed 2.67 snacks ( $SD = 2.27$ ) per day. In total, 107 (14.0%) prompts contained low-calorie snacks and 659 (86.0%) prompts contained high-calorie snacks, that is, participants generally reported more high-calorie snacks than low-calorie snacks (Figure 2A). Chocolate-containing foods were most frequently reported in both self-report prior to the study and during EMA, accounting for 26.3% of all reported snacks (Table 2).

233 **Aim 2: Interrelations Between Thoughts About Snacks, Craving Intensity, and**  
234 **Consumption of Snacks**

235 Our second aim was to investigate the interrelations of craving intensity, snack-related  
236 thoughts, and consumption within participants. As expected, higher craving intensity co-  
237 occurred with frequent thoughts about snacks (Table 3; arrow A in Figure 1B). When adding  
238 type of snacks to this model, the relation between craving intensity and thoughts about snacks  
239 remained significant and type of snacks additionally predicted thoughts about snacks. As low-  
240 calorie snacks were coded with 0 and high-calorie snacks with 1, the positive coefficient  
241 indicates that individuals generally thought more often about high-calorie snacks than low-  
242 calorie snacks in daily life (Table 4). When modeled separately, both thoughts about snacks  
243 ( $\beta_{10} = 0.22, p < .001$ ) and craving intensity ( $\beta_{20} = 0.02, p < .001$ ) were positively associated  
244 with consumption. Furthermore, thoughts about snacks, craving intensity, and their interaction  
245 (trend level;  $p = .051$ ) predicted consumption of snacks (Table 5; arrows B and C in Figure  
246 1B), that is, when both thoughts about snacks and craving intensity were high, it resulted in  
247 higher consumption of snacks (Figure 2B).

248 **Aim 3: Ecological Validity of Trait Food Craving**

249 Our third aim was to examine main effects and moderating effects of trait food craving  
250 on craving intensity, snack-related thoughts, and consumption. Higher FCQ-T-r scores were  
251 associated with more thoughts about snacks and moderated the effect of craving intensity on  
252 thoughts about snacks (Table 3; gray arrow in Figure 1B). Furthermore, trait food craving  
253 moderated the effect of craving intensity on thoughts about snacks (Table 3; Figure 2C) such  
254 that the association between craving intensity and thoughts about snacks was stronger in  
255 individuals with high FCQ-T-r scores than in those with low scores (dashed arrow on arrow A  
256 in Figure 1B). Trait food craving also moderated the association between type of snacks and  
257 thoughts about snacks (Table 4) such that individuals with high FCQ-T-r scores reported more

258 thoughts about high-calorie snacks than low-calorie snacks, whereas no differences were  
259 found in individuals with low FCQ-T-r scores (Figure 2D). Importantly, when craving  
260 intensity was modeled as an outcome, there was no relation between trait food craving and  
261 craving intensity ( $\beta_{01} = -0.05, p = .746$ ), that is, individuals with high FCQ-T-r scores did not  
262 experience more intense cravings for snack foods in general. Finally, trait food craving  
263 moderated the association between craving intensity and consumption of snacks (Table 5;  
264 Figure 2E) such that the association was more pronounced in individuals with high FCQ-T-r  
265 scores than in those with low scores (dashed arrow on arrow B in Figure 1B). Trait food  
266 craving was neither related to consumption of snacks in general nor did it moderate the  
267 relation between thoughts about snacks and consumption of snacks. There was also no three-  
268 way interaction between trait food craving, thoughts about snacks, and craving intensity  
269 (Table 5).

#### 270 **Controlling for Hunger as a Level 1 Predictor**

271 When modeled separately, hunger was positively related to thoughts about snacks ( $\beta_{10}$   
272 = 0.01,  $p = .030$ ). However, this relation was not significant ( $\beta_{10} = 0.004, p = .141$ ) when  
273 craving intensity and type of snacks were simultaneous predictors for thoughts about snacks  
274 (all  $ps \leq .003$ ). Hunger was unrelated to consumption of snacks when modeled separately  
275 ( $\beta_{10} = 0.0002, p = .873$ ), but there was a negative association between hunger and consumption  
276 ( $\beta_{10} = -0.003, p = .019$ ) when thoughts about snacks and craving intensity were simultaneous  
277 predictors for consumption (all  $ps \leq .001$ ). Importantly, trait food craving was not related to  
278 hunger in general ( $\beta_{01} = -0.01, p = .962$ ) and there were no cross-level interactions of trait  
279 food craving and hunger (all  $ps \geq .293$ ). Effects remained significant for the other cross-level  
280 interactions (i.e., trait food craving with type of snack and craving intensity; all  $ps < .037$ ).

281 **Discussion**

282           The present study aimed at characterizing food craving experiences by means of seven  
283 days of EMA with five daily signals. In addition to characterizing these experiences in terms  
284 of content and frequency, their cognitive and behavioral correlates were of interest as was the  
285 potentiating role of trait food craving.

286           Our first aim of characterizing frequency and specificity of food craving experiences  
287 investigated whether previous findings derived from laboratory or questionnaire-based studies  
288 actually reflect individuals' behavior in everyday life. Such previous studies reported that  
289 chocolate next to other high-calorie snack foods are most frequently desired in Western  
290 societies (e.g., Nicholls & Hulbert-Williams, 2013; Rozin et al., 1991; Weingarten & Elston,  
291 1990). Indeed, our data confirmed this finding: 86% of all reported snack foods were high in  
292 sugar and/or fat and chocolate-containing foods were the most frequently reported high-  
293 calorie snack (26% of all prompts), which was consistently reported across measures (i.e.,  
294 self-report prior to the study and EMA).

295           In line with the hypotheses under the second aim, snack-related thoughts, state  
296 cravings, and consumption were all significantly interrelated and co-occurred in the same time  
297 interval within individuals. However, this does not imply that all thoughts or cravings  
298 necessarily lead to consumption. While the elaborated intrusion theory of desire proposes that  
299 thoughts about tempting foods are essential for the emergence of cravings (Kavanagh et al.,  
300 2005; May et al., 2004) and, thus, making causality between these thoughts and cravings  
301 likely, several processes may moderate whether snack-related thoughts and/or cravings result  
302 into snack consumption. For example, in a "hot" motivational state (e.g., during experiences  
303 of hunger or exposure to attractive foods), food cravings may influence consumption more  
304 directly than in a "cold" motivational state (Appelhans et al., 2016; Hofmann & Van Dillen,  
305 2012), where self-control likely prevents consumption.

306 Our third aim pertained to the question whether high trait food cravers are more  
307 susceptible to snack-related thoughts, cravings, or consumption. Unexpectedly, there were no  
308 direct effects of trait food craving on momentary craving intensity or snack consumption. This  
309 contrasts with laboratory studies in which high trait food cravers reported higher craving  
310 intensity in response to food cues than low trait food cravers (e.g., Hallam et al., 2016; Meule,  
311 Skirde, et al., 2012) and further points to context effects (laboratory vs. daily life). Thus,  
312 while food picture viewing in the laboratory triggers state cravings for food quite consistently  
313 in trait food cravers, such correspondence of state and trait might be contingent on food cue  
314 exposure (e.g., sight or smell). In the natural environment, by contrast, exposure to food cues  
315 might vary considerably across situations and daytimes (Boswell & Kober, 2016; Cepeda-  
316 Benito, Fernandez, & Moreno, 2003), showing that state and trait may be separable to some  
317 extent. Also, it has been reported that, although trait food craving and momentary craving  
318 intensity regularly are related to each other, this relationship is rather small (Cepeda-Benito et  
319 al., 2000; Meule, Hermann, et al., 2014; Meule, Lutz, Vögele, & Kübler, 2012). In sum, the  
320 findings indicate that high trait food cravers do not have a chronic and persistently elevated  
321 level of state craving but rather show an increase in state cravings in certain situations.

322 In contrast, trait food craving was associated with more frequent thoughts about snack  
323 foods. This corresponds well with the respective items of the FCQ-T-r (e.g., “I find myself  
324 preoccupied with food”; Cepeda-Benito et al., 2000; Meule, Hermann, et al., 2014) and  
325 provides ecological validity for the instrument. Furthermore, regarding type of snack food,  
326 high trait food cravers thought more frequently about high-calorie snacks than low-calorie  
327 snacks, whereas no such differences were found in low trait food cravers. Thus, both  
328 frequency and specificity of food craving experiences can be predicted by trait food craving in  
329 the natural environment, revealing a generally greater elaboration of high-calorie foods in trait  
330 food cravers. Again, there is some correspondence with laboratory work: when state craving  
331 was triggered, individuals with high trait food craving showed heightened reactivity on an

332 approach avoidance task (Brockmeyer et al., 2015), implicit measures (Richard et al., in  
333 revision), and greater reward-related brain activation (Miedl et al., submitted; Ulrich et al.,  
334 2016).

335         Although there were no direct effects of trait food craving on craving intensity or  
336 snack consumption, trait food craving moderated the relationships between thoughts about  
337 snacks and craving intensity as well as between craving intensity and snack consumption.  
338 That is, high trait food cravers thought more often about snack foods and consumed more  
339 snack foods, particularly when they experienced intense cravings for these foods. Thus, while  
340 individuals with low trait food craving may also experience intense state cravings for food, it  
341 appears that their consequences differ between high and low trait food cravers. Consequently,  
342 intense food cravings in high trait food cravers might represent “hot” motivational states (cf.,  
343 Appelhans et al., 2016) that go along with both cognitive elaboration and consumption, while  
344 being less consequential in low trait food cravers.

345         Hunger was one of the control variables in the present study that deserves discussion.  
346 Hunger was indeed related to more snack-related thoughts, suggesting that food cravings and  
347 hunger often co-occur (Gilhooly et al., 2007). However, hunger was negatively related to  
348 snack consumption, which is in line with previous research. For instance, snack foods are  
349 often consumed as between-meal groceries in the absence of hunger (Cleobury & Tapper,  
350 2014), and state chocolate craving was a better predictor of chocolate consumption than  
351 hunger (Meule & Hormes, 2015). Importantly, hunger did neither confound the relationships  
352 between snack-related thoughts, cravings, and consumption nor was there a relationship with  
353 trait food craving. Thus, the present investigation may represent a demonstration of the partial  
354 independence of craving and hunger experiences (Blechert, Naumann, Schmitz, Herbert, &  
355 Tuschen-Caffier, 2014; Pelchat & Schaefer, 2000; Richard et al., in revision).

356 **Limitations and Future Directions**

357 The present findings are based on a sample of young, female students, which limits the  
358 generalizability to men and individuals with higher age, higher BMI, or lower education. As  
359 food cravings are thought to be more common in females (Hormes, Orloff, & Timko, 2014)  
360 and young adults (Pelchat, 1997), future studies should replicate findings in samples with  
361 broader range in variables such as age, BMI, or education. Previous research also suggests  
362 that individuals can successfully withhold consumption in the presence of food cravings (Hill,  
363 2007). Hence, future studies should aim at investigating a more comprehensive set of  
364 environmental (e.g., availability of food) and individual characteristics to gain insight into  
365 when food craving experiences result in snack consumption and when this is not the case.  
366 Also, the current investigation does not inform about all types of snacking behavior (which  
367 also includes mindless snacking; Wansink & Sobal, 2007), but only about snack consumption  
368 that can be consciously recalled. However, retrospective recall may be biased even for a short  
369 period of three hours (as used here). A combination of event-contingent sampling (i.e.,  
370 immediate report of each snack) and signal-contingent sampling might at least help in  
371 estimating the size of this bias (through comparison of concurrent and retrospective reports),  
372 while a complete account of snacking prevalence likely requires more continuous  
373 measurements (e.g., chewing- or swallowing-detection; Papapanagiotou et al., 2016).

#### 374 **Conclusions**

375 To conclude, snack-related thoughts, cravings and consumption are highly interrelated  
376 in daily life. This is particularly true for trait food cravers who think more often about high-  
377 calorie snack foods and consume more snack foods in response to intense cravings. To date,  
378 interindividual differences in food craving experiences have received relatively little attention  
379 in interventions that target regulation of thoughts about tempting foods (e.g., May, Kavanagh,  
380 & Andrade, 2015). The present findings give evidence for ecological validity as well as  
381 practical relevance of the concept of trait food craving outside the laboratory.

**382 Acknowledgements**

383 This work was supported by the European Research Council (ERC) under the European  
384 Union's Horizon 2020 research and innovation program (ERC-StG-2014 639445 NewEat)  
385 and by the Austrian Science Fund (FWF): [I 02130-B27]. We thank Julia Maria Pickl, Ann-  
386 Kathrin Arend, and Eleonora Fadel for their assistance in collecting data for this study.

## References

- Appelhans, B. M., French, S. A., Pagoto, S. L., & Sherwood, N. E. (2016). Managing temptation in obesity treatment: A neurobehavioral model of intervention strategies. *Appetite, 96*, 268-279.
- Berkman, E. T., Giuliani, N. R., & Pruitt, A. K. (2014). Comparison of text messaging and paper-and-pencil for ecological momentary assessment of food craving and intake. *Appetite, 81*, 131-137.
- Blechert, J., Naumann, E., Schmitz, J., Herbert, B. M., & Tuschen-Caffier, B. (2014). Startling sweet temptations: hedonic chocolate deprivation modulates experience, eating behavior, and eyeblink startle. *PLoS ONE, 9*, e85679.
- Boswell, R. G., & Kober, H. (2016). Food cue reactivity and craving predict eating and weight gain: a meta-analytic review. *Obesity Reviews, 17*, 159-177.
- Brockmeyer, T., Hahn, C., Reetz, C., Schmidt, U., & Friederich, H.-C. (2015). Approach Bias Modification in Food Craving—A Proof-of-Concept Study. *European Eating Disorders Review, 23*, 352-360.
- Bruinsma, K., & Taren, D. L. (1999). Chocolate: Food or drug? *Journal of the American Dietetic Association, 99*, 1249-1256.
- Cepeda-Benito, A., Fernandez, M. C., & Moreno, S. (2003). Relationship of gender and eating disorder symptoms to reported cravings for food: construct validation of state and trait craving questionnaires in Spanish. *Appetite, 40*, 47-54.
- Cepeda-Benito, A., Gleaves, D. H., Williams, T. L., & Erath, S. A. (2000). The development and validation of the state and trait Food-Cravings Questionnaires. *Behavior Therapy, 31*, 151-173.
- Cleobury, L., & Tapper, K. (2014). Reasons for eating 'unhealthy' snacks in overweight and obese males and females. *Journal of Human Nutrition and Dietetics, 27*, 333-341.

- Forman, E. M., Hoffman, K. L., Juarascio, A. S., Butryn, M. L., & Herbert, J. D. (2013). Comparison of acceptance-based and standard cognitive-based coping strategies for craving sweets in overweight and obese women. *Eating Behaviors, 14*, 64-68.
- Gilhooly, C. H., Das, S. K., Golden, J. K., McCrory, M. A., Dallal, G. E., Saltzman, E., . . . Roberts, S. B. (2007). Food cravings and energy regulation: the characteristics of craved foods and their relationship with eating behaviors and weight change during 6 months of dietary energy restriction. *International Journal of Obesity, 31*, 1849-1858.
- Hallam, J., Boswell, R. G., DeVito, E. E., & Kober, H. (2016). Gender-related Differences in Food Craving and Obesity. *Yale Journal of Biology and Medicine, 89*, 161-173.
- Hill, A. J. (2007). The psychology of food craving. *Proceedings of the Nutrition Society, 66*, 277-285.
- Hill, A. J., & Heaton-Brown, L. (1994). The experience of food craving - A prospective investigation in healthy women. *Journal of Psychosomatic Research, 38*, 801-814.
- Hill, A. J., Weaver, C. F. L., & Blundell, J. E. (1991). Food craving, dietary restraint and mood. *Appetite, 17*, 187-197.
- Hofmann, W., & Van Dillen, L. (2012). Desire: The new hot spot in self-control research. *Current Directions in Psychological Science, 21*, 317-322.
- Hormes, J. M., Orloff, N. C., & Timko, C. A. (2014). Chocolate craving and disordered eating. Beyond the gender divide? *Appetite, 83*, 185-193.
- Kavanagh, D. J., Andrade, J., & May, J. (2005). Imaginary relish and exquisite torture: the elaborated intrusion theory of desire. *Psychological Review, 112*, 446-467.
- Kemps, E., & Tiggemann, M. (2009). Attentional bias for craving-related (chocolate) food cues. *Experimental and Clinical Psychopharmacology, 17*, 425-433.
- Lafay, L., Thomas, F., Mennen, L., Charles, M. A., Eschwege, E., Borys, J. M., & Basdevant, A. (2001). Gender differences in the relation between food cravings and mood in an

- adult community: Results from the Fleurbaix Laventie Ville Santé study. *International Journal of Eating Disorders*, 29, 195-204.
- Lowe, M. R., & Butryn, M. L. (2007). Hedonic hunger: a new dimension of appetite? *Physiology & Behavior*, 91, 432-439.
- Martin, C. K., O'Neil, P. M., Tollefson, G., Greenway, F. L., & White, M. A. (2008). The association between food cravings and consumption of specific foods in a laboratory taste test. *Appetite*, 51, 324-326.
- Massey, A., & Hill, A. J. (2012). Dieting and food craving. A descriptive, quasi-prospective study. *Appetite*, 58, 781-785.
- May, J., Andrade, J., Panabokke, N., & Kavanagh, D. (2004). Images of desire: cognitive models of craving. *Memory*, 12, 447-461.
- May, J., Kavanagh, D. J., & Andrade, J. (2015). The Elaborated Intrusion Theory of desire: A 10-year retrospective and implications for addiction treatments. *Addictive Behaviors*, 44, 29-34.
- McKiernan, F., Houchins, J. A., & Mattes, R. D. (2008). Relationships between human thirst, hunger, drinking, and feeding. *Physiology & Behavior*, 94, 700-708.
- Meule, A., Beck Teran, C., Berker, J., Gründel, T., Mayerhofer, M., & Platte, P. (2014). On the differentiation between trait and state food craving: Half-year retest-reliability of the *Food Cravings Questionnaire-Trait-reduced* (FCQ-T-r) and the *Food Cravings Questionnaire-State* (FCQ-S). *Journal of Eating Disorders*, 2, 1-3.
- Meule, A., Hermann, T., & Kübler, A. (2014). A short version of the *Food Cravings Questionnaire - Trait*: The FCQ-T-reduced. *Frontiers in Psychology*, 5, 1-10.
- Meule, A., & Hormes, J. M. (2015). Chocolate versions of the *Food Cravings Questionnaires*. Associations with chocolate exposure-induced salivary flow and ad libitum chocolate consumption. *Appetite*, 91, 256-265.

- Meule, A., Lutz, A., Vögele, C., & Kübler, A. (2012). Food cravings discriminate differentially between successful and unsuccessful dieters and non-dieters. Validation of the Food Cravings Questionnaires in German. *Appetite*, *58*, 88-97.
- Meule, A., Richard, A., & Platte, P. (2017). Food cravings prospectively predict decreases in perceived self-regulatory success in dieting. *Eating Behaviors*, *24*, 34-38.
- Meule, A., Skirde, A. K., Freund, R., Vögele, C., & Kübler, A. (2012). High-calorie food-cues impair working memory performance in high and low food cravers. *Appetite*, *59*, 264-269.
- Meule, A., Vögele, C., & Kübler, A. (2012). [German translation and validation of the Yale Food Addiction Scale]. *Diagnostica*, *58*, 115-126.
- Meule, A., Westenhöfer, J., & Kübler, A. (2011). Food cravings mediate the relationship between rigid, but not flexible control of eating behavior and dieting success. *Appetite*, *57*, 582-584.
- Miedl, S., Blechert, J., Meule, A., Richard, A., & Wilhelm, F. H. (submitted). Suppressing images of desire: Neural correlates of chocolate-related thoughts in high and low trait chocolate cravers.
- Nicholls, W., & Hulbert-Williams, L. (2013). British English translation of the Food Craving Inventory (FCI-UK). *Appetite*, *67*, 37-43.
- Papapanagiotou, V., Diou, C., Zhou, L., van den Boer, J., Mars, M., & Delopoulos, A. (2016). A novel chewing detection system based on PPG, audio and accelerometry. *IEEE J Biomed Health Inform.*
- Papies, E. K., Stroebe, W., & Aarts, H. (2007). Pleasure in the mind: Restrained eating and spontaneous hedonic thoughts about food. *Journal of Experimental Social Psychology*, *43*, 810-817.
- Pelchat, M. L. (1997). Food cravings in young and elderly adults. *Appetite*, *28*, 103-113.

- Pelchat, M. L., & Schaefer, S. (2000). Dietary monotony and food cravings in young and elderly adults. *Physiology & Behavior*, *68*, 353-359.
- Raudenbush, S. W., Byrk, A. S., & Congdon, R. (2011). HLM7 for Windows [Computer software]. Skokie, IL: Scientific Software International, Inc.
- Richard, A., Meule, A., Friese, M., & Blechert, J. (in revision). Effects of chocolate deprivation on implicit and explicit evaluation of chocolate in high and low trait chocolate cravers.
- Rozin, P., Levine, E., & Stoess, C. (1991). Chocolate craving and liking. *Appetite*, *17*, 199-212.
- Stone, A. A., Shiffman, S., Schwartz, J. E., Broderick, J. E., & Hufford, M. R. (2003). Patient non-compliance with paper diaries. *British Medical Journal*, *324*, 1193-1194.
- Tiggemann, M., & Kemps, E. (2005). The phenomenology of food cravings: The role of mental imagery. *Appetite*, *45*, 305-313.
- Ulrich, M., Steigleder, L., & Grön, G. (2016). Neural signature of the Food Craving Questionnaire (FCQ)-Trait. *Appetite*, *107*, 303-310.
- Wansink, B., & Sobal, J. (2007). Mindless Eating. *Environment and Behavior*, *39*(1), 106-123.
- Weingarten, H. P., & Elston, D. (1990). The phenomenology of food cravings. *Appetite*, *15*, 231-246.
- Weingarten, H. P., & Elston, D. (1991). Food cravings in a college population. *Appetite*, *17*, 167-175.
- White, M. A., Whisenhunt, B. L., Williamson, D. A., Greenway, F. L., & Netemeyer, R. G. (2002). Development and validation of the Food-Craving Inventory. *Obesity Research*, *10*, 107-114.

Table 1

*Descriptive Statistics of Level 1 and Level 2 Variables With Means, Standard Deviations, and Ranges.*

<b>Variable</b>	<b><i>M</i></b>	<b><i>SD</i></b>	<b>Minimum</b>	<b>Maximum</b>
<b>Level 1 (occasions)</b>				
Hunger	39.3	22.3	0.00	100
Thoughts about snacks	2.11	1.59	1.00	15.0
Craving intensity	46.6	21.5	0.00	100
Consumption of snacks	1.02	0.97	0.00	9.00
<b>Level 2 (participants)</b>				
Food Cravings Questionnaire-Trait-reduced	39.5	11.4	22.0	67.0

*Notes.* Descriptive statistics of thoughts about snacks including signals for which participants specified not having thought about snacks were  $M = 0.87$ ,  $SD = 1.46$ , range: 0.00–15.0.

Table 2

*Frequency of Food Cravings for Specific Foods Reported Prior to the Study (Means and Standard Deviations) and Snacks Reported During Ecological Momentary Assessment (Absolute Count of Snacks and Percentages).*

Categories	<i>M</i> ( <i>SD</i> )	Absolute count of snacks	% of total
Chocolate-containing foods	3.49 (1.25)	277	26.2
Sweets	3.26 (1.18)	72	6.81
Pizza	2.82 (1.06)	8	0.76
Pasta	2.70 (1.10)	7	0.66
Cookies	2.61 (0.97)	103	9.74
Pastries	2.52 (0.96)	117	11.1
Others	2.52 (1.50)	141	13.3
Cake	2.49 (0.98)	52	4.92
Chips	2.46 (1.03)	59	5.58
Ice cream	2.30 (0.96)	31	2.92
French fries	2.16 (0.93)	2	0.19
Nuts	–	24	2.26
Beverages	–	25	2.36
Fruits and vegetables	–	139	13.2

*Notes.* Absolute number of reported snacks was  $N = 1057$ . Regarding the amount of snacks per category (in %), the number of specific snacks that participants reported having thought about was divided by the number of all mentioned snacks (differing number of thoughts about snacks per signal not considered here). Frequency of food cravings for specific foods was examined on a scale from 1 (*never/not applicable*) to 6 (*always*). Nuts, beverages, fruits, and vegetables were not assessed prior to the study, but were reported during EMA. “Others” comprises foods that were not classifiable (e.g., cereals, soup, sushi).

Table 3

*Coefficients ( $\beta$ ) with Robust Standard Errors (SE) and p-Values of the Mixed Model With Craving Intensity as a Predictor at Level 1 and Trait Food Craving as a Predictor at Level 2 on Thoughts about Snacks.*

<b>Model</b>	<b>Coefficient <math>\beta</math> (SE)</b>	<b><i>p</i></b>
<b>Thoughts about Snacks with Level 1 predictors <sup>a</sup></b>		
Intercept ( $\beta_{00}$ )	2.01 (0.12)	<b>&lt;.001</b>
Craving intensity ( $\beta_{10}$ )	0.02 (0.004)	<b>&lt;.001</b>
<b>Thoughts about Snacks with Level 1 and Level 2 predictors <sup>b</sup></b>		
<b>Level 1</b>		
Intercept ( $\beta_{00}$ )	2.01 (0.11)	<b>&lt;.001</b>
Craving intensity ( $\beta_{10}$ )	0.02 (0.004)	<b>&lt;.001</b>
<b>Level 2</b>		
Trait food craving ( $\beta_{01}$ )	0.03 (0.01)	<b>.002</b>
<b>Cross-level interactions</b>		
Trait food craving $\times$ craving intensity ( $\beta_{11}$ )	0.001 (0.0004)	<b>.019</b>

*Notes.* *P*-values < .050 are printed in boldface. More details on interpretation of the model equations can be found in the appendix.

<sup>a</sup> Level 1 model equation: thoughts about snacks<sub>*ij*</sub> =  $\pi_{0j}$  +  $\pi_{1j}$ (craving intensity) +  $e_{ij}$ ; Level 2 model equations:  $\pi_{0j}$  =  $\beta_{00}$  +  $r_{0j}$ ;  $\pi_{1j}$  =  $\beta_{10}$  +  $r_{1j}$ .

<sup>b</sup> Level 1 model equation: thoughts about snacks<sub>*ij*</sub> =  $\pi_{0j}$  +  $\pi_{1j}$ (craving intensity) +  $e_{ij}$ ; Level 2 model equations:  $\pi_{0j}$  =  $\beta_{00}$  +  $\beta_{01}$ (trait food craving) +  $r_{0j}$ ;  $\pi_{1j}$  =  $\beta_{00}$  +  $\beta_{11}$ (trait food craving) +  $r_{1j}$ .

Table 4

*Coefficients ( $\beta$ ) with Robust Standard Errors (SE) and p-Values of the Mixed Model With Craving Intensity and Type of Snacks as Predictors at Level 1 and Trait Food Craving as a Predictor at Level 2 on Thoughts about Snacks.*

<b>Model</b>	<b>Coefficient <math>\beta</math> (SE)</b>	<b><i>p</i></b>
<b>Thoughts about Snacks with Level 1 predictors <sup>a</sup></b>		
Intercept ( $\beta_{00}$ )	1.72 (0.13)	<b>&lt;.001</b>
Craving intensity ( $\beta_{10}$ )	0.02 (0.004)	<b>&lt;.001</b>
Type of snacks ( $\beta_{20}$ )	0.34 (0.11)	<b>.004</b>
<b>Thoughts about Snacks with Level 1 and Level 2 predictors <sup>b</sup></b>		
<b>Level 1</b>		
Intercept ( $\beta_{00}$ )	1.72 (0.13)	<b>&lt;.001</b>
Craving intensity ( $\beta_{10}$ )	0.02 (0.003)	<b>&lt;.001</b>
Type of snacks ( $\beta_{20}$ )	0.33 (0.11)	<b>.004</b>
<b>Level 2</b>		
Trait food craving ( $\beta_{01}$ )	0.01 (0.01)	.180
<b>Cross-level interactions</b>		
Trait food craving $\times$ craving intensity ( $\beta_{11}$ )	0.001 (0.0003)	<b>.028</b>
Trait food craving $\times$ type of snacks ( $\beta_{21}$ )	0.02 (0.01)	<b>.035</b>

*Notes.* *P*-values < .050 are printed in boldface.

<sup>a</sup> Level 1 model equation: thoughts about snacks<sub>ij</sub> =  $\pi_{0j}$  +  $\pi_{1j}$ (craving intensity) +  $\pi_{2j}$ (type of snacks) +  $e_{ij}$ ; Level 2 model equations:  $\pi_{0j}$  =  $\beta_{00}$  +  $r_{0j}$ ;  $\pi_{1j}$  =  $\beta_{10}$  +  $r_{1j}$ ;  $\pi_{2j}$  =  $\beta_{20}$  +  $r_{2j}$ .

<sup>b</sup> Level 1 model equation: thoughts about snacks<sub>ij</sub> =  $\pi_{0j}$  +  $\pi_{1j}$ (craving intensity) +  $\pi_{2j}$ (type of snacks) +  $e_{ij}$ ; Level 2 model equations:  $\pi_{0j}$  =  $\beta_{00}$  +  $\beta_{01}$ (trait food craving) +  $r_{0j}$ ;  $\pi_{1j}$  =  $\beta_{10}$  +  $\beta_{11}$ (trait food craving) +  $r_{1j}$ ;  $\pi_{2j}$  =  $\beta_{20}$  +  $\beta_{21}$ (trait food craving) +  $r_{2j}$ .

Table 5

*Coefficients ( $\beta$ ) with Robust Standard Errors (SE) and p-Values of the Mixed Model With Thoughts About Snacks and Craving Intensity as Predictors at Level 1 and Trait Food Craving as a Predictor at Level 2 on Consumption of Snacks.*

<b>Model</b>	<b>Coefficient <math>\beta</math> (SE)</b>	<b><i>p</i></b>
<b>Consumption of snacks with Level 1 predictors <sup>a</sup></b>		
Thoughts about snacks ( $\beta_{10}$ )	0.19 (0.06)	<b>&lt;.001</b>
Craving intensity ( $\beta_{20}$ )	0.20 (0.03)	<b>&lt;.001</b>
Thoughts about snacks $\times$ craving intensity ( $\beta_{30}$ )	0.07 (0.04)	.051
<b>Consumption of snacks with Level 1 and Level 2 predictors <sup>b</sup></b>		
<b>Level 1</b>		
Thoughts about snacks ( $\beta_{10}$ )	0.19 (0.05)	<b>&lt;.001</b>
Craving intensity ( $\beta_{20}$ )	0.19 (0.04)	<b>&lt;.001</b>
Thoughts about snacks $\times$ craving intensity ( $\beta_{30}$ )	0.07 (0.04)	.076
<b>Level 2</b>		
Trait food craving ( $\beta_{01}$ )	0.003 (0.01)	.548
<b>Cross-level interactions</b>		
Trait food craving $\times$ thoughts about snacks ( $\beta_{11}$ )	0.003 (0.004)	.520
Trait food craving $\times$ craving intensity ( $\beta_{21}$ )	0.01 (0.003)	<b>.040</b>
Trait food craving $\times$ thoughts about snacks $\times$ craving intensity ( $\beta_{31}$ )	0.001 (0.003)	.567

*Notes.* *P*-values < .050 are printed in boldface. Predictors at Level 1 were *z*-standardized within each participant before calculating the product term. Intercept of both model equations <sup>a,b</sup>:  $\beta_{00} = 0.97(0.05)$ ,  $p < .001$ .

<sup>a</sup> Level 1 equation: consumption of snacks<sub>*ij*</sub> =  $\pi_{0j}$  +  $\pi_{1j}$ (thoughts about snacks) +  $\pi_{2j}$ (craving intensity) +  $\pi_{3j}$ (thoughts about snacks  $\times$  craving intensity) +  $e_{ij}$ ; Level 2 equations:  $\pi_{0j} = \beta_{00} + r_{0j}$ ;  $\pi_{1j} = \beta_{10} + r_{1j}$ ;  $\pi_{2j} = \beta_{20} + r_{2j}$ ;  $\pi_{3j} = \beta_{30} + e_{3j}$ .

<sup>b</sup> Level 1 equation: consumption of snacks $_{ij} = \pi_{0j} + \pi_{1j}(\text{thoughts about snacks}) + \pi_{2j}(\text{craving intensity}) + \pi_{3j}(\text{thoughts about snacks} \times \text{craving intensity}) + e_{ij}$ ; Level 2 equations:  $\pi_{0j} = \beta_{00} + \beta_{01}(\text{trait food craving}) + r_{0j}$ ;  $\pi_{1j} = \beta_{10} + \beta_{11}(\text{trait food craving}) + r_{1j}$ ;  $\pi_{2j} = \beta_{20} + \beta_{21}(\text{trait food craving}) + r_{2j}$ ;  $\pi_{3j} = \beta_{30} + \beta_{31}(\text{trait food craving}) + r_{3j}$ .

ACCEPTED MANUSCRIPT

**Figure captions**

*Figure 1.* (A) Conceptual model of the interrelations between craving intensity, thoughts about snacks, and consumption as a function of trait food craving. White boxes and black arrows illustrate effects of within-person predictors. The gray box and gray arrows illustrate effects of the between-person predictor (i.e., scores on the Food Cravings Questionnaire-Trait-reduced; FCQ-T-r). Arrows pointing on boxes represent main effects of predictors (e.g., more intense cravings and/or more frequent snack-related thoughts might be related to greater consumption of snacks; arrows B and C). Arrows pointing on arrows represent moderating effects of one predictor on the relation between two others (e.g., the relation between craving intensity and thoughts about snacks might be more pronounced in individuals with high FCQ-T-r scores). The double-headed arrow indicates that variables might be mutually interchangeable (arrow A). Dashed arrows indicate possible cross-level interactions of between- and within-person predictors. Type of snacks refers to the categorization of snacks into high- and low-caloric. (B) Empirical model of the interrelations between craving intensity, thoughts about snacks, and consumption as a function of trait food craving. As opposed to the conceptual model, only significant main and interactive effects are displayed. Craving intensity and type of snack predicted thoughts about snacks (arrow A). Craving intensity and thoughts about snacks predicted consumption of snacks (arrows B and C). High trait food cravers thought more frequently about high-calorie snack foods than low-calorie snack foods (gray arrow). Further, high trait food cravers showed stronger relationships between snack-related thoughts and craving intensity (dashed arrow on arrow A) and between craving intensity and snack consumption (dashed arrow on arrow B).

Figure 2. (A) Simple slopes probing the interaction between type of snacks and trait food craving when predicting the number of snacks reported. Individuals with low FCQ-T-r scores,  $\beta = 7.45$ ,  $SE = 0.94$ ,  $t_{(59)} = 7.89$ ,  $p < .001$ , and individuals with high FCQ-T-r scores,  $\beta = 10.8$ ,  $SE = 0.98$ ,  $t_{(59)} = 11.0$ ,  $p < .001$ , reported more high-calorie snacks than low-calorie snacks, with this effect being slightly more pronounced in individuals with high FCQ-T-r scores (marginally significant cross-level interaction:  $\beta_{11} = 1.67$ ,  $p = .055$ ). (B) Simple slopes probing the interaction between craving intensity and thoughts about snacks when predicting consumption of snacks per signal. Higher craving intensity was particularly associated with more consumption of snacks when participants reported a high number of thoughts about snacks,  $\beta = 0.27$ ,  $SE = 0.05$ ,  $t_{(59)} = 4.85$ ,  $p < .001$ , whereas this relationship was attenuated when participants reported a small number of thoughts about snacks,  $\beta = 0.12$ ,  $SE = 0.05$ ,  $t_{(59)} = 2.46$ ,  $p = .017$ . (C) Simple slopes probing the interaction between craving intensity and trait food craving when predicting thoughts about snacks per signal. Higher craving intensity was associated with more thoughts about snacks in individuals with high FCQ-T-r scores,  $\beta = 0.63$ ,  $SE = 0.10$ ,  $t_{(59)} = 6.52$ ,  $p < .001$ , but not in individuals with low FCQ-T-r scores,  $\beta = 0.19$ ,  $SE = 0.10$ ,  $t_{(59)} = 1.89$ ,  $p = .063$ . (D) Simple slopes probing the interaction between type of snacks and trait food craving when predicting thoughts about snacks per signal. Individuals with high FCQ-T-r scores reported significantly more thoughts about high-calorie snacks than low-calorie snacks,  $\beta = 0.87$ ,  $SE = 0.22$ ,  $t_{(59)} = 4.03$ ,  $p < .001$ , whereas no difference was found in individuals with low FCQ-T-r scores,  $\beta = 0.03$ ,  $SE = 0.22$ ,  $t_{(59)} = 0.14$ ,  $p = .893$ . (E) Simple slopes probing the interaction between craving intensity and trait food craving when predicting consumption of snacks per signal. Higher craving intensity was associated with more consumption of snacks in individuals with high FCQ-T-r scores,  $\beta = 0.26$ ,  $SE = 0.05$ ,  $t_{(59)} = 5.18$ ,  $p < .001$ , whereas this relationship was attenuated in individuals with low FCQ-T-r scores,  $\beta = 0.12$ ,  $SE = 0.05$ ,  $t_{(59)} = 2.27$ ,  $p = .027$ . \*\*\*  $p < .001$ , \*  $p < .050$ .

## Appendix

Model equation for the prediction of thoughts about snacks by craving intensity (modeled at Level 1) and by trait food craving (modeled at Level 2).

### Level 1 (occasions):


$$\text{Thoughts about snacks}_{ij} = \pi_{0j} + \pi_{1j}(\text{craving intensity}_{ij}) + e_{ij}$$

### Level 2 (participants):

$$\pi_{0j} = \beta_{00} + \beta_{01}(\text{trait food craving}) + r_{0j}$$

$$\pi_{1j} = \beta_{10} + \beta_{11}(\text{trait food craving}) + r_{1j}$$

The outcome (participant's  $j$  level of thoughts about snacks) was displayed as a function of an intercept ( $\pi_{0j}$ ) and a slope ( $\pi_{1j}$ ) at Level 1. This shows the effect of craving intensity <sub>$ij$</sub>  (participant's  $j$  level of craving intensity). The intercept  $\pi_{0j}$  represents participant's  $j$  level of thoughts about snacks at an average level of craving intensity. At Level 2, the intercepts ( $\beta_{00}$ ) and ( $\beta_{10}$ ) show the mean level of thoughts about snacks and the mean effect of craving intensity, when trait food craving is on an average level due to grand-mean centering of trait food craving. The regression weights (Level 2 slopes) represent associations between the Level 2 predictor trait food craving and thoughts about snacks ( $\beta_{10}$ ) and craving intensity ( $\beta_{11}$ ), respectively.

**A****B**