

Holothuria (Mertensiothuria) viridiaurantia sp. nov. (Holothuriida, Holothuriidae), a new sea cucumber from the Eastern Pacific Ocean revealed by morphology and DNA barcoding

Giomar Helena Borrero-Pérez¹, María Juliana Vanegas-González¹

¹ Instituto de Investigaciones Marinas y Costeras – INVEMAR, Museo de Historia Natural Marina de Colombia (MHNMC) - Makuriwa, Calle 25 No. 2–55, Playa Salguero, Rodadero, Santa Marta, Colombia

Corresponding author: *Giomar H. Borrero-Pérez* (giomar.borrero@invemar.org.co; giomarborrero@gmail.com)

Academic editor: *A. Martynov* | Received 9 May 2019 | Accepted 16 October 2019 | Published 2 December 2019

<http://zoobank.org/51DD2CEA-0FC9-45D1-861B-15CE8F817061>

Citation: Borrero-Pérez GH, Vanegas-González MJ (2019) *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. (Holothuriida, Holothuriidae), a new sea cucumber from the Eastern Pacific Ocean revealed by morphology and DNA barcoding. ZooKeys 893: 1–19. <https://doi.org/10.3897/zookeys.893.36013>

Abstract

Holothuria (Mertensiothuria) viridiaurantia sp. nov. is described based on specimens from rocky reefs of northern Chocó in the Colombian Pacific Ocean; however, it also occurs along the Eastern Pacific Ocean from Mexico and Panama. Although specimens from Mexico and Panama were previously identified as *Holothuria (Mertensiothuria) hilla* Lesson, 1830 the new species is easily distinguished morphologically and via mtDNA. In terms of morphology, the species can be identified by its olive-green background and white-orange papillae and tentacles, larger tentacles with deep indentations and also by larger buttons on the dorsal and ventral body wall, papillae and tube feet; large, thick and rough tentacle rods, and the absence of ossicles in the longitudinal muscles. The new species is included in the subgenus *Mertensiothuria* considering molecular evidence.

Keywords

16S, COI, Chocó, Colombia, Echinodermata, Cabo Marzo, Gulf of Cupica, Gulf of Tribugá, Holothuroidea, mtDNA, morros, riscales, rocky reefs

Introduction

The family Holothuriidae Ludwig, 1894 currently includes 211 valid species, with the genus *Holothuria* Linnaeus, 1867 being the most diverse, containing 165 formally described species (WoRMS 2019a). Sixteen new *Holothuria* species have been described from different localities around the world since 2000; two of them from the Central and Tropical Eastern Pacific (Laguarda-Figueras and Solís-Marín 2009; Honey-Escandón et al. 2011). The diversity of *Holothuria* will likely continue to grow considering some “cryptic” species currently recognised based on molecular evidence (COI mtDNA) and morphological characteristics, such as colouration, as reported for the *Holothuria* (*Thymiosycia*) *impatiens* complex (Michonneau 2015). In addition, the exploration of poorly known regions could generate new information on the diversity of holothurians and other marine organisms. In particular, an area that warrants further exploration is the north of the Colombian Pacific (Chocó), part of the Tumbes-Magdalena-Chocó biogeographical hotspot that is considered a mega-diverse area (Cortés 1997).

Holothuria (*Mertensiothuria*) Deichmann, 1958, one of the 18 *Holothuria* subgenera, was reviewed by Samyn and Massin (2003), who emended its diagnosis by incorporating ossicles from the longitudinal muscles. According to Samyn and Massin (2003) there were six species in the subgenus; four of them were previously recognised: *H. (M.) albofusca* Cherbonnier, 1988, *H. (M.) fusciorubra* Théel, 1886, *H. (M.) leucospilota* (Brandt, 1835), and *H. (M.) papillifera* Heding in Mortensen, 1938; and two were transferred from the subgenus *Thymiosycia* into *Mertensiothuria*: *H. hilla* Lesson, 1830 and *H. aphanes* Lampert, 1885. Samyn and Massin (2003) also removed four species from the subgenus either because of the absence of ossicles in the longitudinal muscles (*Holothuria arenacava* Samyn et al., 2001 and *Holothuria platei* Ludwig, 1898) or because this characteristic was unknown (*Holothuria artensis* Cherbonnier & Féral, 1984 and *Holothuria exilis* Koehler & Vaney, 1908). According to WoRMS (2019b), *Mertensiothuria* currently includes the same six species accepted by Samyn and Massin (2003), although *H. (M.) papillifera* is considered species inquirenda. In addition, the subgenus includes three more species: *H. (M.) isuga* Mitsukuri, 1912, *H. (M.) arenacava* and *H. (M.) artensis*; the last two species excluded in the revision by Samyn and Massin (2003) have been transferred by WoRMS but with no reference.

Among the species in this subgenus *H. (M.) hilla* Lesson, 1830 is the most widespread species, reported from the Red Sea to Madagascar and across the Indian Ocean and the Pacific Ocean to the Central and Tropical Eastern Pacific (Purcell et al. 2012). It is a common species in the Central and Tropical Eastern Pacific occurring in its common colour morph, comprised of a yellow background and white papillae (Samyn and Massin 2003; Purcell et al. 2012). However, specimens with different colours, such as an olive-green background and white-orange papillae, have been reported by several authors (Solís-Marín et al. 2009, Lam. 30A; Sotelo-Casas et al. 2015: fig. 2E; Molina et al. 2015: fig. 3C). Specimens with yellow and green colour patterns were collected in the Colombian Pacific Ocean in 2016, allowing comparison of the morphology and mitochondrial DNA. The purpose of this paper is to describe a new species of *Holothuria* from the Eastern Pacific and to indicate how it differs from *Holothuria* (*Mertensiothuria*) *hilla*.

Materials and methods

The specimens reviewed were collected as part of the project “Riscales”, developed by the Instituto de Investigaciones Marinas y Costeras – INVEMAR (www.invemar.org.co), seeking to characterise the biodiversity of the rocky reefs (called locally “riscales” and “morros”) located in northern Chocó in the Colombian Pacific Ocean. These ecosystems are important for regional fisheries and conservation (Díaz-Fahrenberger et al. 2016). Specimens were collected by hand using SCUBA diving at three rocky reefs between 10 and 15 m depth, during two sampling events in April and October 2016 (Fig. 1A, B). The specimens were placed in plastic bags with seawater, relaxed using magnesium chloride, fixed and preserved in 96% ethanol. They are deposited at the Museo de Historia Natural Marina de Colombia (MHNMC) – Makuriwa of INVEMAR (INV EQU).

External and internal morphology were reviewed to record standard data for sea cucumbers. Tissue from papillae, dorsal body wall, tube feet, ventral body wall, tentacles, and internal organs (longitudinal muscles, respiratory trees, tentacle ampullae, cloaca, and intestine) was removed and dissolved in fresh household bleach. Ossicles were observed and photographed using light microscopy and, at least, ten ossicles of each type were measured using the software ImageJ (Schneider et al. 2012). Type of ossicles, shape, and size were compared with those described by Lesson (1830) and Samyn and Massin (2003).

Ethanol-fixed tissues of the sea cucumbers collected during the project were processed to obtain sequences of the mitochondrial cytochrome oxidase I (COI) and 16S (large subunit) genes; in this paper, only the data of the specimens of interest are shown. Genomic DNA was extracted using the QIAGEN extraction kit (DNeasy Blood & Tissue Kit) and COI and 16S were amplified using the primers COIceF (ACTGCCACGCCCTAGTAATGATATTTTTTATGGTNATGCC) and COIceR (TCGTGTGTCTACGTCCATTCTACTGTRAACATRTG) (Hoareau and Boissin 2010) and 16SA (CGCCTGTTTATCAAAAACAT) and 16SB (CTCCGGTTT-GAACTCAGATCA) (Palumbi 1996). PCRs were carried out following the conditions described by Hoareau and Boissin (2010). PCR products were purified and sequenced using the BigDye 3.1 (Applied Biosystems) technology. The obtained nucleotide sequences were edited using Mega 7. We analysed a fragment of 443 bp of 16S genes (including gaps) and 439 bp of COI. Sequences of COI were translated into amino acids to ensure their integrity and accuracy. The sequences obtained in the present study were submitted to GenBank (Table 1). Available sequences of *H. (M.) hilla*, *H. (M.) leucospilota*, *H. (T.) arenicola*, and *H. (T.) impatiens* from GenBank were included in the analysis (Table 1). 16S sequences were aligned using the L-INS-i method implemented in MAFFT 6 (Katoh et al. 2002) and COI with Clustal W (Thompson et al. 1994). Distances using Kimura 2 parameters correction were calculated and neighbour-joining trees were generated using Mega 7 (Kumar et al. 2016). Within-group genetic differences were analysed on the species level also based on Kimura 2-parameter distances. The best substitution model was searched using the Akaike information criterion implemented in jModelTest (Posada 2008). Phylogenetic relationships were inferred using Bayesian Inference (BI) and Maximum Likelihood (ML).

BI was performed with MrBayes v. 3.2.6 (Ronquist and Huelsenbeck 2003) using unlinked GTR+G evolutionary model for each gen; the data set was run twice, using four Markov chains for ten million generations for each analysis to estimate posterior probabilities. ML analysis was performed in Mega 7; support was assessed in this case by 1000 bootstrap pseudoreplicates.

Results

Order Holothuriida Miller, Kerr, Paulay, Reich, Wilson, Carvajal & Rouse, 2017

Family Holothuriidae Burmeister, 1837

Genus *Holothuria* Linnaeus, 1767

Subgenus *Mertensiothuria* Deichmann, 1958

***Holothuria* (*Mertensiothuria*) *viridiaurantia* sp. nov.**

<http://zoobank.org/3CF88C56-4A82-4758-B74A-395DF68C7F68>

Figures 1–7; Tables 1–3

Holothuria (*Mertensiothuria*) *hilla* Solís-Marín et al. 2009: 110–111, fig. 30A–G; Santos-Beltrán and Salazar-Silva 2011: fig. 2A; Honey-Escandón et al. 2012; Sotelo-Casas et al. 2015: 3–4, figs 2E, 3(4–8), 4(4–6); Molina et al. 2015: fig. 3C.

Material examined. Holotype: INV EQU4309, one specimen, total length 70 mm; collected in La Viuda rocky reef, Gulf of Cupica, northern Chocó, Colombia (6°37.9812'N, 77°29.985'W), by G. Borrero, 24 Oct 2016; at 15 m depth under rocks and attached; GenBank nucleotide sequences COI MK477997 and 16S MK477991 (Fig. 2A–D).

Paratype: INV EQU4234, one specimen, total length 35 mm; collected in El Faro rocky reef, Cabo Marzo, northern Chocó, Colombia (6°49.4802'N, 77°41.3976'W), by M.J. Vanegas, 24 April 2016; at 13 m depth under rocks and attached; GenBank nucleotide sequences 16S MK477992 (Fig. 2E); **Paratype:** INV EQU4312, one specimen, total length 25 mm; collected in Morromico rocky reef, Gulf of Tribugá, northern Chocó, Colombia (5°52.3194'N, 77°18.6426'W), by G. Borrero, 20 Oct 2016; at 10 m depth under rocks; GenBank nucleotide sequences COI MK477998 (Fig. 2F).

Comparative material *Holothuria* (*Mertensiothuria*) *hilla*: INV EQU4245, one specimen, total length 75 mm; collected in Piedra de Rodrigo rocky reef, Cabo Marzo, northern Chocó, Colombia (6°47.0346'N, 77°41.6148'W), by M.J. Vanegas, 25 April 2016; at 19 m depth under rocks; INV EQU4310, four specimens, total length 70–100 mm; by G. Borrero, 26 Oct 2016; same locality, depth, and habitat as previous; GenBank nucleotide sequences 16S MK477993 (Fig. 2G, H); INV EQU4311, one specimen, total length 65 mm; by G. Borrero, 26 Oct 2016; same locality, depth, and habitat as previous; GenBank nucleotide sequences COI MK477996 and 16S MK477994 (Fig. 2I, J).

Table 1. Specimens of *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. from Colombia and GenBank sequences analysed for the partial cytochrome oxidase subunit 1 (COI) and/or 16S genes. Sequence-Voucher Location column includes the origin of the sequences according to the GenBank references where they have been generated; sequences from this study include the catalogue number at the MHNMC-INVEMAR (INV EQU). An asterisk (*) in the species column indicates changes in the GenBank ID, where the sequences were previously identified as *H. (Mertensiothuria) hilla*.

Species	GenBank Accession number		Sequence-Voucher, Location	Reference
	COI	16S		
<i>H. (M.) viridiaurantia</i> sp. nov.	MK477997	MK477991	Colombia (Pacific) Holotype (INV EQU4309)	This study
	MK477998	–	Colombia (Pacific) Paratype (INV EQU4312)	This study
	–	MK477992	Colombia (Pacific) Paratype (INV EQU4234)	This study
<i>H. (M.) viridiaurantia</i> sp. nov.*	JN207616	JN207515	Mexico	Honey-Escandón et al. 2012
	KP780302	–	India	Deepa and Bijukumar, unpub.
<i>H. (M.) hilla</i>	MK477996	MK477994	Colombia (Pacific) (INV EQU4311)	This study
	–	MK477993	Colombia (Pacific) (INV EQU4310)	This study
	KX874337	KX856783	Mariana Islands, Guam	Miller et al. 2017
	–	EU822442	–	Uthicke and Byrne, unpub.
	–	FJ223856	Malaysia	Kamarul et al. 2006
	–	FJ223864	Malaysia	Kamarul et al. 2006
<i>H. (M.) leucospilota</i>	JN207617	JN207541	Marshall Islands (Majuro)	Honey-Escandón et al. 2012
	KC405566	KY986424	Pangkor Island, Malaysia	Kamarudin and Rehan 2015
	KY986417	KY986423	Pangkor Island, Malaysia	Kamarudin and Rehan 2015
	KC405565	KY986422	Pangkor Island, Malaysia	Kamarudin and Rehan 2015
<i>H. (T.) impatiens</i>	MK477999	MK477995	Colombia (Pacific) (INV EQU4236)	This study
	JN207632	JN207526	Mexico (Pacific)	Honey-Escandón et al. 2012
<i>H. (T.) aff. impatiens</i>	–	FJ223857	Malaysia	Kamarudin et al. 2010
<i>H. (T.) arenicola</i>	JN207608	JN207556	Florida (USA)	Honey-Escandón et al. 2012
<i>Isostichopus fuscus</i>	MK477908	MK477869	Panama (Pacific) IfTa210	This study

Diagnosis. Olive-green background with white-orange dorsal papillae, tube feet and tentacles; buttons >75 µm in length; large tentacles with deep indentations; tentacle rods thick, rough and with some perforations; longitudinal muscles without ossicles.

Description. External appearance: medium-sized species, holotype preserved specimen 70 mm long and 21 mm wide; body loaf like (length < 4× diameter) length/width ratio 2.3. Body shape of living ex situ specimen cylindrical in cross-section (Fig. 2A), tapering posteriorly and widening anteriorly, ending in a large crown of tentacles. Body wall soft and thin (2–3 mm thick). Anus terminal surrounded by small papillae. Mouth directed ventrally in live and preserved specimens, encircled by large papillae (Fig. 2A, D). Large peltate tentacles 20; ca. 5–6 mm total length, and 4–5 mm width shield; with




Figure 1. Maps showing the geographic distribution of *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. **A, B** Detailed distribution in Chocó, Colombia; the triangle indicates the only locality where *H. (Mertensiothuria) hilla* specimens were collected **C** Wider distribution of *H. (M.) viridiaurantia* sp. nov. showing other localities from Panama and Mexico where the species have been identified through photographs, and the locality in India from where the GenBank sequence KP780302 originated. Colombian localities are represented by exact coordinates, and all other localities were derived from specific localities mentioned in the literature.

deep indentions 2–3 mm. Few large, long and slender conical papillae scattered on the dorsal surface, although a vague arrangement into four rows is observed, two of them are lateral, where they are a little larger; smaller papillae scattered among the largest. Ventral tube feet cylindrical, large and thick, densely distributed throughout the ventral surface.

Colour. Background of living specimens olive-green; base of the papillae is a light or whitish green that changes to orange from the middle to the ends, however, the tips of the papillae are whitish. Ventral surface similar to dorsum, with orange tube feet and white suckers; tentacles orange, same colour as papillae and tube feet (Fig. 2A). Dark brownish green in preserved specimens with papillae, tube feet, and tentacles a dark yellow (Fig. 2D–F).




Figure 2. Type specimens of *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. (**A–F**) and comparative material of *Holothuria (Mertensiothuria) hilla* (**G–J**). **A** Dorsal and ventral view of the alive holotype of *H. (M.) viridiaurantia* sp. nov. from Gulf of Cupica, Northern Chocó, Colombia (INV EQU4309, L = 70 mm) **B** calcareous ring and **C** stone canal and madreporite of the Holotype **D** detail of the preserved holotype tentacles **E** paratype from Cabo Marzo; Northern Chocó, Colombia (INV EQU4234, L = 35 mm) **F** smallest paratype from Gulf of Tribugá, Northern Chocó, Colombia (INV EQU4312, L = 25 mm) **G, H** alive specimen of *H. (M.) hilla* from Cabo Marzo, Northern Chocó, Colombia, and detail of tentacles in the preserved specimen (INV EQU4310, L = 100 mm) **I, J** preserved specimen of *H. (M.) hilla*, same locality as **G, H** and detail of tentacles (INV EQU4311, L = 65 mm). Scale bars: 2 mm (**B, C**); 1 cm (**E, F, I**).

Internal anatomy. Square radial plates in the calcareous ring, 3 mm wide and 3 mm high, with three anterior rounded processes, and posterior margin with shallow rounded indentation; interradial plates slender, 1.5 mm high and 2.5 mm wide, pointed anterior margin and rounded posterior margin (Fig. 2B). One free stone canal, 4 mm long, and a helicoidally madreporite, 4 mm long (Fig. 2C). Tubular tentacle ampullae, 3–4 mm long and striped coloured. Tube-like polian vesicle, 17 mm long. Longitudinal muscles pair flat, thinner in the middle of each pair, irregularly wide, 3–4 mm wide each band, or 2–2 mm wide, attached, with narrow free edges. Gonads absent. Cuvierian organ present. Right respiratory tree extending to anterior end; left respiratory tree attached to the intestine until the middle of the body.

Ossicles: Dorsal and ventral body wall include similar tables and buttons, with dorsal tables taller and dorsal buttons larger than ventral (Table 2, Fig. 3A). Tables disc circular to quadrangular in outline; rim of the disc smooth; with four large central perforations and 7–12 smaller peripheral holes, arranged in one ring; spire with four pillars, single crossbeam, spiny crown with a small central hole (Fig. 3A). Dorsal tables 60–81 μm across disc (\bar{x} = 68 μm) and 43–54 μm height (\bar{x} = 49 μm); ventral tables 57–71 μm across disc (\bar{x} = 63 μm) and 40–49 μm height (\bar{x} = 44 μm). Buttons with smooth rim but irregular contour, usually with three pairs of holes, sometimes four pairs or three-four unpaired holes (Fig. 3A). Dorsal buttons 79–115 μm long (\bar{x} = 101 μm); ventral buttons 82–108 long (\bar{x} = 94 μm). Dorsal papillae with tables, buttons, button-like plates, rods and at the very tip one small plate and small rods (Table 2, Fig. 4A). Tables and buttons similar in shape and size to the ones in the body wall, although table's spires are thicker and buttons are larger, up to 130 μm . Rods 167–203 μm long (\bar{x} = 187 μm) with distal or median perforations; small plates at the top of the papillae 99–134 μm (\bar{x} = 187 μm) and small rods 37–58 μm (\bar{x} = 48 μm). Ventral tube feet or pedicels with tables, buttons, plates, and end plates (Table 2, Figs 4A, 5A). Tables and buttons similar in shape and size to the ones in ventral body wall, although buttons are larger, up to 140 μm . Plates 105–133 μm long (\bar{x} = 117 μm) and 72–129 μm wide (\bar{x} = 116 μm); end plates 578–581 μm wide. Tentacles with large and small rods (Table 2; Fig. 5A); large rods are thick plate-like and very rough, usually with perforations at the extremities or along its length, 113–261 μm long and 33–150 μm width. Small rods thin and few spinose (58–107 μm). Longitudinal muscles without ossicles (Fig. 5A), as well as in the other internal organs, such as respiratory trees, tentacle ampullae, cloaca, and intestine.

Paratypes: Juveniles, 35 and 25 mm long, 12 and 8 mm wide respectively (Fig. 2E, F). External morphology different to the holotype, which is much larger at 70 mm long. Small dorsal papillae in the four main rows, as described for the holotype; and three rows of tube feet, two lateral and one in the middle of the ventral side which includes two irregular lines of pedicels (Fig. 2F). Dorsal and ventral body wall buttons are smaller in the juvenile, although there is not a considerable difference in size; however, in shape they are more rounded at the extremities and frequently present more than three pairs of holes (Table 2, Fig. 6A, B). Tables showed more changes during growth in comparison with buttons: the tables spire are taller and narrower, pointed-like without cross beam clearly noted, with few spines around the top; and the tables disc diameter is larger, with peripheral holes less in number and larger in size in the juvenile (Table 2,

Table 2. Comparison of ossicle size in *Holothuria* (*Mertensiothuria*) *viridiaurantia* sp. nov. holotype, juvenile paratype, and *H. (M.) hilla* of similar body size.

Characteristics	<i>H. (M.) viridiaurantia</i> sp. nov. Paratype INV EQU4312		<i>H. (M.) viridiaurantia</i> sp. nov. Holotype INV EQU4309		<i>H. (M.) hilla</i> INV EQU4311	
	L = 25 mm		L = 70 mm		L = 65 mm	
	Range (µm)	Average	Range (µm)	Average	Range (µm)	Average
Dorsal body wall						
Buttons length	88–119	99	79–115	101	47–70	61
Tables disc diameter	62–84	73	60–81	68	51–75	59
Tables height	56–68	62	43–54	49	35–45	41
Width spires	14–15	15	23–31	29	21–26	23
Ventral body wall						
Buttons length	78–104	90	82–108	94	50–73	61
Tables disc diameter	54–80	63	57–71	63	47–61	53
Tables height	45–56	51	40–49	44	34–44	39
Width spires	9–17	14	22–29	25	16–20	18
Dorsal papillae						
Buttons, Buttons-like plates length	80–111	99	106–130	117	64–91	79
Tables disc diameter	51–95	74	60–82	71	48–66	58
Tables height	58–71	64	42–62	52	43–47	45
Width spires	19–28	25	30–42	36	20–25	22
Rods	97–168	143	167–203	187	102–152	129
Plates	–	–	99–134	118	104–107	106
Small Rods	–	–	37–58	48	43–67	57
Ventral tube feet						
Buttons, Buttons-like plates length	93–116	105	91–140	110	52–84	70
Tables disc diameter	54–75	63	53–73	61	41–57	51
Tables height	46–54	50	41–51	45	34–43	38
Width spires	15–23	18	23–31	28	17–24	21
Plates length	66–122	96	105–133	117	76–127	96
Plates width	46–83	65	72–129	116	68–99	84
End Plates	391–401	396	578–581	580	570–584	577
Tentacles						
Large Rods Length	132–259	197	185–261	221	166–267	227
Large Rods Width	21–39	29	33–150	75	10–27	18
Small Rods Length	37–91	56	58–107	74	37–65	47
Longitudinal muscle						
C-O shape ossicles length	NA	NA	NA	NA	13–33	24

Fig. 6A, B). Dorsal papillae and tube feet present similar pattern of change during growth in buttons and tables when comparing the juvenile with the holotype; however, tables in dorsal papillae and tube feet in the juveniles are less pointed-like and one cross beam is clearly noted in most of the tables in comparison with those from the dorsal and ventral body wall (Fig. 6C, D). In addition, rods in dorsal papillae are smaller in size; it was not possible to observe the small plates and rods at the very top of the papillae. Supporting plates and end plates in the tube feet are also smaller in the juvenile (Table 2, Fig. 6D, E). Tentacle rods are not well developed in the paratype, being almost similar in length but less thick than those of the holotype, however, they are thicker than those in the *H. (M.) hilla* individual of 65 mm in length (Table 2, Figs 5A, B; 6F).




Figure 3. Ossicle comparison between *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. and *Holothuria (Mertensiothuria) hilla*. **A** Holotype of *H. (M.) viridiaurantia* sp. nov. (INV EQU4309, L = 70 mm) **B** *H. (M.) hilla* (INV EQU4311, L = 65 mm); showing ossicle set from dorsal body wall (tables, buttons) and ventral body wall (tables, buttons); grey squares indicated in the images are presented enlarged below each image. Scale bar: 100 μ m.




Figure 4. Ossicles comparison between *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. and *Holothuria (Mertensiothuria) hilla*. **A** Holotype of *H. (M.) viridiaurantia* sp. nov. (INV EQU4309, L = 70 mm) **B** *H. (M.) hilla* (INV EQU4311, L = 65 mm); showing ossicle set from dorsal papillae (tables, buttons, rods), dorsal papillae tip (showing the plate and small rods at the tip) and tube feet (tables, buttons and supporting plates); grey squares indicated in some images are presented enlarged below each image. Scale bar: 100 μ m.




Figure 5. Ossicles comparison between *Holothuria* (*Mertensiothuria*) *viridiaurantia* sp. nov. and *Holothuria* (*Mertensiothuria*) *hilla*. **A** Holotype of *H. (M.) viridiaurantia* sp. nov. (INV EQU4309, L = 70 mm) **B** *H. (M.) hilla* (INV EQU4311, L = 65 mm); showing end plates from tube feet and ossicle set from tentacles (large and small rods) and longitudinal muscles (C's and O's ossicles). Scale bars: 100 μ m (except **A** upper).

Etymology. From the Latin *viridis* (green) and *aurantius* (orange-coloured), referring to the living colour with olive-green background and orange-white papillae, tube feet, and tentacles (feminine).

Distribution. *Holothuria* (*Mertensiothuria*) *viridiaurantia* sp. nov. is known and confirmed along the Eastern Pacific from Mexico (as *Holothuria* (*Mertensiothuria*) *hilla*, Solís-Marín et al. 2009; Santos-Beltrán and Salazar-Silva 2011; Honey-Escandón et al. 2012; Sotelo-Casas et al. 2015), Panama (as *H. (M.) hilla*, Molina et al. 2015) and Colombia (present study) (Fig. 1). However, a GenBank sequence of one specimen from Kerala coast, India (Accession number KP780302.1) suggests that the new species could have a wider geographical distribution across the Indian Ocean and the Pacific Ocean to the Central and Tropical Eastern Pacific, like *H. (M.) hilla* (Fig. 1C).




Figure 6. Ossicles of the juvenile paratype of *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. (INV EQU4312, L = 25 mm). **A** Dorsal body wall (tables, buttons) **B** ventral body wall (tables, buttons) **C** dorsal papillae (tables, buttons, rods) **D** tube feet (tables, buttons, supporting plates) **E** tube feet (end plate) **F** tentacles (large and thick rods, small rods). Scale bars: 100 μ m.

However, it was not possible to review the specimen belonging to the sequence, so colouration and morphological characteristics described in the present paper should be reviewed and confirmed. Notably, images of green-coloured *H. (M.) hilla* from the Philippines are presented by Dolorosa et al. (2017; Fig. 2J).

Habitat. *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. is associated with rocky bottoms from the intertidal to 15 m depth (Molina et al. 2015; present study). Specimens collected in Colombia were found attached under medium rocks, differing

from *H. (M.) hilla* which were found under rocks but on a sandy substrate instead of a rocky substrate. *Holothuria (M.) viridiaurantia* sp. nov. was collected in three different rocky reefs in the northern Chocó, whereas *H. (M.) hilla* was found only in one, during both collection trips in 2016. Several specimens of *H. (M.) hilla* were found regenerating new anterior ends, this was not observed in specimens of the new species.

Conservation status. As the specimens of *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. were previously assigned to *H. (M.) hilla*, the conservation status of this species must be considered. Currently *H. (M.) hilla* is included in the IUCN Red List in the category of Least Concern, and in addition, it is classified as a low-value species (about USD3 kg⁻¹ dried in the Philippines) among commercially important sea cucumbers of the world (Purcell et al. 2012). There is no fishery reports of *H. (M.) hilla* in the Eastern Pacific Ocean, however, *H. (M.) hilla* is fished commercially in the Philippines, Indonesia and Madagascar, that could include the new species considering the potential wider distribution of *H. (M.) viridiaurantia* sp. nov.

Remarks. The new species was previously assigned to *Holothuria (Mertensiothuria) hilla* (Solís-Marín et al. 2009; Santos-Beltrán and Salazar-Silva 2011; Honey-Escandón et al. 2012; Molina et al. 2015; Sotelo-Casas et al. 2015), however there is no mention of the distinct and striking colouration of the specimens reported in those papers in comparison with *H. (M.) hilla*. Perhaps the identification of this species was based on the similar external appearance (shape of the body and papillae) and apparent similar ossicles at first sight; without regard to the colouration, which has been traditionally considered to be intra-specific variability in echinoderms. However, recent research demonstrates that it can be a diagnostic characteristic, for example in the species complex *H. (T.) impatiens* (Michonneau 2015); this subject requires careful and exhaustive study, especially the purpose of colouration in sea cucumbers (Clark 1922; Michonneau 2015). In this study, a detailed revision of specimens from the new species and *H. (M.) hilla*, showed not only the colouration as a diagnostic feature, but also the size and shape of the tentacles, which are larger and with deeper indentations in the new species (Fig. 2). In reference to the ossicles, although similar in shape at first sight, a detailed revision showed several diagnostic characteristics: 1) differences in the size of the complete ossicle sets from the dorsal and ventral body wall, dorsal papillae and tube feet; specifically, the tables are taller and thicker with wider discs and the buttons are larger in the new species, in both juvenile and large specimens (Table 2; Figs 3–6); size of the buttons is the most diagnostic trait for the species; 2) the size and shape of the tentacle rods, being wide (plate-like), thick and very rough, and with some perforations in the new species compared to slender rods in *H. (M.) hilla* (Table 2; Figs 5A, B, 6F); 3) longitudinal muscle ossicles are absent in the new species, contrary to *H. (M.) hilla* (Table 2; Fig. 5A, B). In general, the morphological structures of the new species are thicker and stronger than those of *H. (M.) hilla*, which is a more delicate species. Among the morphological characteristics of the new species, the absence of ossicles in the longitudinal muscles, larger size of the perforated plates of the tube feet, and size and shape of the tentacle ossicles, match those considered by Samyn and Massin (2003) for excluding *Holothuria arenacava* and *Holothuria platei* from *Mertensiothuria*. However, the decision for including the new species in this subgenus was made based on the mtDNA evidence.




Figure 7. Bayesian inference tree of mitochondrial genes of the specimens analysed for the present study. **A** COI **B** 16S. The numbers on the nodes indicate Neighbour Joining (bootstrap %) / Maximum Likelihood (bootstrap %) / Bayesian posterior probability. Hyphen (-) indicates nodes not supported in some trees. Sequences from Colombia obtained in this study include the catalogue number at the MHNMC - INVEMAR (INV EQU); GenBank ascension number is included for the other sequences (see Table 1).

Molecular characteristics. We obtained COI and 16S sequence data from three specimens of *Holothuria (Mertensiothuria) viridiaurantia* sp. nov. and two of *Holothuria (Mertensiothuria) hilla* from the rocky reef in northern Chocó, Colombia. Specimens of *H. (M.) viridiaurantia* sp. nov. from Colombia (type specimens) were recovered in a well-supported clade, separated from *H. (M.) hilla* for both, COI and 16S genes (Fig. 7). Two sequences, derived from one specimen from Mexico (GenBank Accession No. JN207616–COI and JN207515–16S) and one from India (KP780302–COI), were recovered in the same clade as type specimens from Colombia. However, different tree topologies for COI and 16S sequence data were recovered. For COI *H. (M.) viridiaurantia* sp. nov. appears sister to *H. (M.) hilla*, with *H. (M.) leucospilota* positioned sister to them (Fig. 7A). However, for 16S *H. (M.) hilla* and *H. (M.) leucospilota* appeared as sister species with *H. (M.) viridiaurantia* sp. nov. as sister clade (Fig. 7B). Species from *Thymiosycia* subgenus appear separated from *Mertensiothuria* subgenus for both genes and all tree reconstruction methods (Fig. 7A, B). Evidence for species status

Table 3. Kimura 2 parameter distances (%) within specimens of *Holothuria* (*Mertensiothuria*) *viridiaurantia* sp. nov. and between the *Holothuria* species included in the analysis. COI distances are below diagonal and 16S distances above. The numbers in bold lettering along the diagonal represent average within species distances for COI and 16S (COI / 16S).

	Species	1	2	3	4	5	6
1	<i>H. (M.) viridiaurantia</i> sp.nov.	1.3 / 0.5	12.5	11.8	15.7	13.5	32.6
2	<i>H. (M.) hilla</i>	16.7	13.8 / 7.9	13.9	19.4	16.3	33.9
3	<i>H. (M.) leucospilota</i>	15.6	17.7	1.9 / 1.6	19.4	17.2	34.8
4	<i>H. (T.) impatiens</i>	21.3	21.5	21.8	0.9 / 8.8	15.5	33.2
5	<i>H. (T.) arenicola</i>	18.0	19.9	17.6	19.4	nc / nc	34.3
6	<i>Isostichopus fuscus</i>	24.4	22.5	24.0	24.5	20.7	nc / nc

of *H. (M.) viridiaurantia* sp. nov. comes from the COI and 16S genetic distances. Inter-specific distances between the two previously recognised *Mertensiothuria* species included in the analysis is 17.7% for COI and 13.9% for 16S; and distances between the new species and them are 16.7 and 15.6% for COI and 12.5 and 11.8% for 16S; inter-specific distances among species of *Mertensiothuria* and *Thymiosycia* showed larger values (Table 3). In addition, intra-specific distances for *H. (M.) viridiaurantia* sp. nov. were 1.31% for COI and 0.5% for 16S, the lowest values in all the species analysed. Intra-specific distances for *H. (M.) hilla* (13.8% for COI and 7.9% for 16S) could be showing a species complex, similar to what was described by Michonneau (2015) for *H. (T.) impatiens*, which is also recovered here with 8.8% for 16S, including one specimen identified as *H. (T.)* aff. *impatiens* (Table 3). Lower intraspecific distance for COI (0.9%) for *H. (T.) impatiens* is explained because the sequence for COI was not available for this specimen. There is, therefore, strong molecular evidence that *H. (M.) viridiaurantia* sp. nov. is an undescribed species different from *H. (M.) hilla*, a finding also supported by the morphological characteristics described previously.

Acknowledgements

We thank researchers who participated in the project “Caracterización biótica de los arrecifes rocosos (Riscales y Morros) en el Chocó Norte, Pacífico Colombiano” for helping during collection of the specimens; Angelica Fernández and Sandy Caldera (MHNMC, INVEMAR) for their collaboration in the molecular lab work; Janneth Beltrán (LABSIS, INVEMAR) for the elaboration of the maps; Crystal Kelehear-Graham for her diligent proofreading of this paper. External reviewers for their suggestions to improve the manuscript. This work was supported by INVEMAR through the projects “Caracterización Biótica de los Arrecifes Rocosos (Riscales y Morros) en el Chocó Norte, Pacífico Colombiano 2016–2018” and “Obtención del código de barras del ADN (Barcoding) para los grupos de fauna y flora marinos– 2016, 2017”; both under the agreement with the Ministry of Environment and Sustainable Development of Colombia, Code: BPIN: 2014011000405 y 2017011000113. INVE-MAR contribution No. CTRB – 1235.

References

- Brandt JF (1835) Echinodermata ordo Holothurina. In: Mertensio H (Ed.) Prodrromus Descriptionis Animalium Ab In Orbis Terrarum Circumnavigatione Observatorum (Fasc. I). Petropoli, 42–62.
- Burmeister H (1837) Handbuch der Naturgeschichte. [Part 2] xii. Enslin, Berlin, 369–858.
- Cherbonnier G, Féral JP (1984) Les holothuries de Nouvelle-Calédonie, Deuxième contribution. Première partie: Synallactidae et Holothuriidae. Bulletin Muséum national d'Histoire naturelle, Paris 4, série 6A, 3: 659–700.
- Cherbonnier G (1988) Echinodermes: Holothurides. Faune de Madagascar. Publié sous les auspices du Gouvernement de la République Malgache, 70, Editions de l'ORSTOM, Paris, 292 pp.
- Clark HL (1922) The holothurians of the genus *Stichopus*. Bulletin of the Museum of Comparative Zoology at Harvard College 65(3): 39–73.
- Cortés J (1997) Biology and geology of Eastern Pacific coral reefs. Coral Reefs 16: 39–46. <https://doi.org/10.1007/s003380050240>
- Deepa RP, Bijukumar A (Unpublished) Barcoding of sea cucumbers along Kerala coast, India. GenBank Reference.
- Deichmann E (1958) The Holothuroidea collected by the VELERO III and IV during the years 1932 to 1954. Part II. Apsidochirota. Allan Hancock Pacific Expeditions 11(2): 253–349.
- Díaz-Fahrenberger A, Neira A, Díaz JM (2016) Manglares y riscales: el sustento ecológico de la pesca. In: Díaz JM, Guillot L, Velandia MC (Eds) La Pesca Artesanal en la Costa Norte del Pacífico Colombiano: Un Horizonteam Bivalente. Fundación MarViva, Bogotá, 25–33.
- Dolorosa RG, Salazar CB, Delfin MYV, Paduga1 JR, Balisco RAT (2017) Sea cucumber fisheries in Rasa Island Wildlife Sanctuary, Narra, Palawan, Philippines. SPC Beche-de-mer Information Bulletin 37: 9–20.
- Heding SG, Mortensen T (1938) Contributions to the study of the development and larval forms of Echinoderms. IV. Kongelige Danske Videnskabernes Selskabs Skrifter (Naturvidenskabelig og matematisk afdeling) (9)7(3): 1–59. [30 figs, 12 pls.]
- Hoareau TB, Boissin E (2010) Design of phylum-specific hybrid primers for DNA barcoding: addressing the need for efficient COI amplification in the Echinodermata. Molecular Ecology Resources 10(6): 960–967. <https://doi.org/10.1111/j.1755-0998.2010.02848.x>
- Honey-Escandón M, Solís-Marín FA, Laguarda-Figueras A (2011) *Holothuria (Selenkothuria) carere*, a new species of sea cucumber (Echinodermata: Holothuroidea) from the Mexican Pacific. Zootaxa 2922: 27–33. <https://doi.org/10.5281/zenodo.202420>
- Honey-Escandón M, Laguarda-Figueras A, Solís-Marín FA (2012) Molecular phylogeny of the subgenus *Holothuria (Selenkothuria)* Deichmann, 1958 (Holothuroidea: Aspidochirotida). Zoological Journal of the Linnean Society 165, 109–120. <https://doi.org/10.1111/j.1096-3642.2011.00794.x>
- Kamarudin KM, Rehan MM (2015) Morphological and Molecular Identification of *Holothuria (Merthensiothuria) leucospilota* and *Stichopus horrens* from Pangkor Island, Malaysia. Tropical life science research 26(1): 87–9.

- Kamarudin KR, Hashim R, Usup G (2010) Phylogeny of Sea Cucumber (Echinodermata: Holothuroidea) as Inferred from 16s Mitochondrial rRNA Gene Sequences. *Sains Malaysiana* 39(2): 209–218.
- Kamarul RK, Gires U, Ridzwan H (2006) Paraphyly of the Genus *Holothuria* (Aspidochirotrida: Holothuriidae) as Inferred from 16S Mitochondrial rRNA Gene Sequences. Proceedings 8th National Symposium on Biology: Indigenous Biological Research for National Development. University Kebangsaan Malaysia (UKM), Malaysian Biotechnology Information Centre (MABIC) and Ministry of Natural Resources and Environment, Putrajaya, BB05, 1–6.
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform (describes the FFT-NS-1, FFT-NS-2 and FFT-NS-i strategies). *Nucleic Acids Research* 30: 3059–3066. <https://doi.org/10.1093/nar/gkf436>
- Koehler R, Vaney C (1908) *Holothuries* recueillis par l'Investigateur dans l'Océan Indien. II. Les *Holothuries* Littorales. Trustees Indian Museum, Calcutta, 54 pp. [3 pls]
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution* 33(7): 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Laguarda-Figueras A, Solís-Marín FA (2009) *Holothuria (Cystipus) casoa* a new species of sea cucumber (Echinodermata: Holothuroidea) from the central-eastern Pacific Ocean. *Scientia Marina* 73(3): 573–578. <https://doi.org/10.3989/scimar.2009.73nn3573>
- Lampert K (1885) Die Seewalzen, eine Systematische Monographie mit Bestimmung und Verbreitungs Tabellen. In: Semper C (Ed.) *Reisen Im Archipel Der Philippinen*. Zweiter teil. (Wissenschaftliche Resultate, Wiesbaden 4(3): 1–311. [1 pl.]
- Lesson RP (1830) *Centurie Zoologique On Choix D'animaux Rares, Nouveaux Ou Inparfaitement Connus*. Levrault, Paris, 244 pp. [80 pls.]
- Linnaeus C (1767) *Systema Naturae – XII edn*, Leipzig.
- Ludwig H (1898) Die *Holothurien* der Sammlung Plate Fauna Chilensis I, *Zoologische Jahrbücher, Supplementband 4*: 431–454. [1 pl.]
- Michonneau F (2015) Cryptic and not-so-cryptic species in the complex *Holothuria (Thymiosycia) impatiens* (Forsskal, 1775) (Echinodermata: Holothuroidea). bioRxiv (The preprint server for biology). <https://doi.org/10.1101/014225>
- Miller AK, Kerr AM, Paulay G, Reich M, Wilson NG, Carvajal JI, Rouse GW (2017) Molecular phylogeny of extant Holothuroidea (Echinodermata). *Molecular Phylogenetics and Evolution* 111: 110–131. <https://doi.org/10.1016/j.ympev.2017.02.014>
- Mitsukuri K (1912) *Studies on Actinopodous Holothurioidea*. Journal College Science, Tokyo Imperial University 29(2): 284 pp. [8 pls.] <https://doi.org/10.5962/bhl.title.37880>
- Molina L, Roca K, Pino Y, Borrero-Pérez GH, Vergara-Chen C (2015) Algunas Especies de Pepinos de Mar (Echinodermata: Holothuroidea) de Zonas Rocosas Intermareales del Pacífico de Panamá. 68th Annual Gulf and Caribbean Fisheries Institute Meeting, 179 pp.
- Palumbi SR (1996) Nucleic acids II: The polymerase chain reaction. In: Hillis DM, Moritz CM, Mable BK (Eds) *Molecular systematics*. Sinauer, Sunderland, 205–247.

- Posada D (2008) jModelTest: Phylogenetic Model Averaging. *Molecular Biology and Evolution* 25: 1253–1256. <https://doi.org/10.1093/molbev/msn083>
- Purcell SW, Samyn Y, Conand C (2012) Commercially important sea cucumbers of the World. FAO Species Catalogue for Fishery Purposes. No. 6. Food and Agriculture Organization of the United Nations, Rome, 223 pp.
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Samyn Y, Massin C (2003) The holothurian subgenus *Mertensiothuria* (Aspidochirotida: Holothuriidae) revisited. *Journal of Natural History* 37(20): 2487–2519. <https://doi.org/10.1080/00222930210144334>
- Samyn Y, Massin C, Muthiga NA (2001) A new species of *Holothuria* (Aspidochirotida, Holothuriidae) from Kenya. Koninklijk Museum Midden-Afrika Tervuren Belgie, *Annalen Zoologische Wetenschappen* 285: 101–110.
- Santos-Beltrán C, Salazar-Silva P (2011) Holoturoideos (Echinodermata: Holothuroidea) de playas rocosas, zona norte de Bahía de Banderas, Nayarit, México. *Ciencia y Mar* 15(45): 3–11.
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nature methods* 9(7): 671–675. <https://doi.org/10.1038/nmeth.2089>
- Solís-Marín FA, Arriaga-Ochoa JA, Laguarda-Figueras A, Frontana-Uribe SC, Duran-Gonzalez A (2009) Holoturoideos (Echinodermata: Holothuroidea) del Golfo de California. Jimenez Editores e Impresores, Distrito Federal, Mexico, 177 pp.
- Sotelo-Casas RC, Cúpul-Magaña AL, Solís-Marín FA, Rodríguez-Troncoso AP (2015) New records of sea cucumbers (Echinodermata: Holothuroidea) at Islas Marietas, Central Mexican Pacific. *Marine Biodiversity Records* 8: 1–8. <https://doi.org/10.1017/S1755267215000810>
- Théel H (1886) Report on the Holothurioidea dredged by H.M.S. ‘Challenger’ during the years 1873–76. *Challenger Reports Zoology* 39: 1–290.
- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673–4680. <https://doi.org/10.1093/nar/22.22.4673>
- Uthicke S, Byrne M (Unpublished) Genetic Barcoding of commercial holothurians. GenBank Reference.
- WoRMS (2019a) Holothuriidae Burmeister, 1837. <http://marinespecies.org/aphia.php?p=taxdetails&cid=731943> [on 2019-01-07]
- WoRMS (2019b) *Holothuria* (*Mertensiothuria*) Deichmann, 1958. <http://www.marinespecies.org/aphia.php?p=taxdetails&cid=241397> [on 2019-01-07]