

RESEARCH ARTICLE

ASSESS THE KNOWLEDGE AND PERCEPTION ON HUMAN BREAST MILK BANK AMONG POSTNATAL MOTHERS.

Sharumathy V¹, Dr.Vetriselvi.P² and Dr.Adhisivam.B³.

.....

- 1. B.SC[N],College of nursing,JIPMER,Puducherry.
- 2. Assistant professor, College of Nursing, JIPMER, Puducherry.
- 3. Additional Professor&HOD, Department of neonatalogy, JIPMER, Puducherry.

Manuscript Info

Manuscript History Received: 16 September 2019 Final Accepted: 18 October 2019 Published: November 2019

Key words:-

Human breastmilk bank, Knowledge, Perception, Postnatal Mothers.

Abstract

..... Background: Human breast milk is considered as the best source of all nutrients, beyond that it the only living food source for infants especially during the first six months of their life. WHO recommends the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding till 6 months of life for all in fan the exclusive breast feeding tin fan the exclusivtsandthensupplemented breastfeedingisrecommended untilat least two yearsofageanditisrecognized asgloballyhighlymore Nutritious. According to the joint statement given by the World Health Organization and United Nations InternationalChildren'sEmergency Fund(UNICEF)in1980. Thebestandessentialfoodforany baby whoseown mother'smilkisnotavailable

is the breast milk from an other donating healthymother. Hence this study focuses on identifying and assessing the knowledge and perception among postnatal mothers about Human Breast Milk Bank establishment and its donation. Itaims on describing

behaviors, beliefs, and feelings behind Human Milk Bank and its donation by identifying donor's individual, social and environmental features that could affect the donation of human milk. The findings will help to develop heal the ducation tools to promote human milk donation by identifying the factors that hinders the human milk donation and rectifying it accordingly.

Objectives:

- 1. Toassessknowledgeand perception on Human MilkBankamongpostnatal mothers.
- 2. Toidentifytheassociationofknowledgeand perception withselected demographic variables.

Methods: The design of the study is cross sectional. Theparticipantswho met the inclusion criteriawereselected byconvenientsamplingtechnique. The total participants included in this study were 100. the data was collected by using structured questionnaire. Statistical methods like frequency and percentage, chisquare test have been used and All statistical analysis has been carried out at 5% level of significant and P<0.05 has considered as significant. **Results:**The study concludedthatonly 19% of the postnatalmothers hadadequate knowledgeandinregardto $perception 70\% of postnat almothers had favorable\ perception.\ There was no significant association$

ofknowledgeandperception with any of thesocio- demographic variables.

Conclusion:The above results stress the importance of organizing health education topostnatalmothers.

Copy Right, IJAR, 2019,. All rights reserved.

Introduction:-

Human breast milk is considered as the best source of all nutrients, beyond that it is the only living food source for infants especially during the first six months of their life.⁽¹⁾

WHOrecommends the exclusive breastfeeding till 6 months of life for all infants and then supplemented breastfeeding is recommended until at least two years of a gean ditisre cognized as globally highly more Nutritious.⁽²⁾

It is vital because of its bio-active substances like free amino acids, nucleotides, cells, growth factors, enzymes, hormones, pro-biotic bacteria, and prebiotic oligosaccharides that are essential for development of immature immune system and gut micro biome of both full-term and high risk newborns like low birth weight and preterm. It also decreases the incidences of necrotizing enterocolitis ,late onset sepsis, retinopathy, food intolerance or malabsorption, immunodeficiency and improvise the neuro-cognitive development. Meanwhile lower the risk of childhood diabetes, obesity compared to those infants fed with formula milk and animal's milk.^(1,3,4,10,12)

Exclusivebreastfeedingisveryessential not only for infants but also for mothers as it prevents the risk of breast cancer and enhances the bond between mothers and babies.⁽¹¹⁾

According to the joint statement given by the World Health Organization and United Nations InternationalChildren'sEmergency Fund(UNICEF)in1980. Thebestandessentialfoodforany baby whoseown mother'smilkisnotavailable isthebreastmilkfromanotherdonatinghealthymother⁽⁵⁾.

BreastMilkDonation is recommended and suggested for hospitalized (sick) or preterm babies in which their mothers don't have sufficient milk production (or) under went any breast surgery in past (or) under any medications like chemotherapy (or) any other condition where mother can't be able to give her breast milk. The Human milk bank receives human milk from donor after screening, then collects, process, pasteurize, store and then distributed to needy.^(6,7,9,12)

But unfortunately, most of the developing countries including our nation which has highest burden of global neonatal mortality and morbidity rate lag in the implementation of donor breastmilk as best alternative choice for vulnerable infants. Breast milk banks aren't a new phenomenon in India, but they are scarce. Eventhough Asia's first milk bank was set up in Mumbai (Lokmanya Tilak Hospital) in 1989. Currently there are only 60 human milk bank. Whereas Brazil, another developing country, is an international leader with 217 banks and 113 collection centers which has helped to reduce brazil's infant mortality rate by 73%.^(1,3,8,9)

Hence this study focuseson identifying andassessing theknowledgeand perceptionamong postnatalmothers aboutHumanBreastMilk Bank establishmentand itsdonation.Itaimsondescribing behaviors.beliefs.andfeelingsbehindHumanMilk Bank anditsdonation bv identifying donor's individual, social and environmental features thatcould affect the donation of human milk. The findings willhelp todevelophealtheducation tools promotehumanmilkdonation by identifying the factors that hinders the human milk donation and rectifying it accordingly.

Objectives:-

- 1. Toassessknowledgeand perception on Human MilkBankamongpostnatal mothers.
- 2. Toidentifytheassociationofknowledgeand perception withselected demographic variables.

Methodology:-

Study design: Cross sectional study

Inclusion criteria: Postnatalmothers inWomen and Children HospitalJIPMER

Exclusion criteria:

Motherswho aresickduringthedatacollection procedure

Sampling population:

Postnatal mothers who are admitted in Women and ChildrenHospitalJIPMER .

Sample size: 100

Sampling technique: Convenience sampling technique.

Tool description:

structured questionnairewasused tocollectdatafrom postnatal mothers.

TheQuestionnairehad 3sections.Theyare:

Section – A: Socio-Demographic Data Section-B: Knowledge on human breast milk bank. Section-C: Perception on human breast milk bank.

Study procedure:

Thestudy wasconducted after approval from IEC (Human studies), JIPMER. The participants who met the inclusion criteria were selected by convenients ampling technique. The total participants included in this study were 100.

Aftergettinginformedwrittenconsentfrom the postnatal mothers, a structured questionnaire was administered to them to assess their knowledge and perception about human breast milk bank. In which it had 16 knowledge questions [multiple choicequestions] and 13 perception questions and instructed themto markthecorrectanswer.

Inknowledgequestionnairethecorrectanswerwasscoredas1andthewronganswer as0.Inperceptionquestionnaire,theresponseoptionsforeachquestionwasstrongly agree,agree,not sure,disagreeand stronglydisagree.

Parameters studied:

Independent variable:

- 1. Age
- 2. Education
- 3. Religion
- 4. Type of family
- 5. Monthly income of family
- 6. Domicile

Outcome variables:

Level of Knowledge and perception on human milk bank.

Confounding and interacting variables: NIL

Statistical test used for data analysis:

The distributions of socio-demographic characteristics were expressed as frequency and percentage. The knowledgelevelwascategorized into inadequate, moderate and adequate based on the scores and it was expressed in

frequency and perception level was categorized into unfavorable, average and favorable based on the scores and it was expressed in frequency and percentage. The association of level of knowledge & perception categories with demographic variables was carried out by using chi-square test. All statistical analysis has been carried out at 5% level of significant and P<0.05 has considered as significant.

Results:-

Table1:-Distribution of socio-demographic**N=100**

Variable	Category	Frequency(N)	Percentage(%)		
Agein years	20-30	90	90%		
	31-40	10	10%		
Domicile	Rural	73	73%		
	Semi-urban	0	0%		
	Urban	27	27%		
Education	Illiterate	23	23%		
	$6^{\text{th}} - 10^{\text{th}}$	20	20%		
	11^{th} - 12^{th}	25	25%		
	Graduate	32	32%		
Type of family	Nuclear	36	36%		
	Joint	64	64%		
	Extended	0	0%		
Monthly income	≤2000	32	32%		
of family	2001-5000	19	19%		
	5001-10,000	21	21%		
	>10,000	28	28%		

Table 2:-Knowledgelevel of postnatal mothers:N=100

Groups	Knowledge score	Frequency(NO)	Percentage(%)
Postnatal	Inadequate (0-50%)	29	29%
mothers	Moderate (51-75%)	52	52%
	Adequate (>75%)	19	19%

Table 3:-Perception levelof postnatal mothers:N=100

Groups	Perceptionscore	Frequency(NO)	Percentage(%)
Postnatal	Unfavorable (0-50%)	1	1%
mothers	Average (51-75%)	29	29%
	Favorable (75-100%)	70	70%

Table 4:-Association between knowledge and perception of postnatal mothers with socio demographic variables: N=100

S	Variables	Ν	Knowledgescore			P Perceptionscore			Р	
I.			inadequ	moder	adequ	valu	unfavora	avera	Favora	val
Ν			ate	ate	ate	e	ble	ge	ble	ue
0										
1	Agein years									
	20-30	9	26	45	19	0.24	1	27	62	0.7
	31-40	0	3	7	0	6	0	2	8	48
		1								
		0								
2	Domicile									
	Rural	7	23	36	14		1	19	53	0.4
	Semi -urban	3	0	0	0	0.06	0	0	0	83
	Urban	0	6	16	5	17	0	10	17	
		2								

		7								
3	Education Illiterate	2	9	9	5		1	7	15	
	6 th -10 th	3	9 6	9 11	3	0.72	0	3	13	0.41
		2	8	13	4	0	0 0	9	16	8
	11^{th} - 12^{th}	0	6	19	7	Ŭ	0	10	22	0
	Graduate	2								
		5								
		3								
		2								
4	Typeoffamily									
	Nucl	3	9	20	7		1	6	29	0.06
	ear	6	20	32	12	0.79	0	23	41	0
	Joint	6	0	0	0	7	0	0	0	
	Exte	4								
	nded	0								
5	Monthlyinco	3								
	me	2								
	offamily	1	12	15	5 5		0	10	22	0.57
	≤2000	9	5	9	5		0	4	15	0
	2001-5000	2	6	10	5	0.69	1	7	13	
	5000-10,000	1	6	18	4	8	0	8	20	
	>10,000	2								
		8								

Discussion:-

The first objective of the study was to assess the knowledge and perception on Human Milk Bank among postnatal mothers.

Amongpostnatalmothersonly19% hadadequateknowledge,52% hadmoderateknowledgeand29% ofthemhadinadequateknowledgeandInregardtoperception70% of the postnatalmothershadfavorableperception,29% of them had average perception and 1% hadun favorable perception.perception,

Theabovefindingsweresupported by thefollowingstudies:

GoodfellowHE, Reimers P,Ballard K,Coutsoudis A(2016)⁽¹³⁾conducted astudy insouthAfrica to determine the mother sperception of breastfeeding and community-based human milkbanks. Through conveniences ampling atotal of 40 mothers were interviewed. The results showed that only 58% of women were willing to do nate the breast milk, only 30% were willing to feed their baby with do nated only 57.5% considered the idea of setting up community based human milkbank.

L,seidlE(2009)⁽¹⁴⁾ Alencar conductedanexploratory, descriptive and cross-sectional study among 36 womenintheagegroupof 14to33yearswho were breastmilk donorsinBrazil. Theobjectivewas to describe the characteristics of donation behavior and to identify the reasons, beliefs and feelings relativetobreastmilk donationpractice. Throughconvenientsampling thesampleswereselectedanddatawere collectedthroughinterviewmethod. The results showed that 91.7% of the mothers thought that breast milkdonationexperience ispositive. The most frequently reported reasons for donating breast milkwere altruismand excessmilkproduction.

Iloh K, Osuorah C, Ndu I, et.all. (2018)⁽¹⁾ conducted a study in south-east Nigeria to determine the perception and acceptability of donor breast milk among mothers. It is a cross- sectional multi-center study in which the data was collected using pretested questionnaire among 1235 selected mothers by using purposive and convenient sampling

method. Results showed that 39% (480/1225) of them heard about it, whereas only 10% had adequate knowledge about the concept and policy of donor milk and 60% of them were willing to donate and use the donor breast milk.

Thesecondobjective of the study was to identify the association of knowledge and perception with selected demographic variables:

The current study findings revealed that there was no significant association of knowledge and perception with any of these lected demographic variables among postnatal mothers.

Theabovefindingsweresupported by the following studies:

Ghuge S, Aghamkar J, Salvi R (2018)⁽⁷⁾ conducted a study in Maharastra to determine the knowledge and attitude towards human milk bank among postnatal mothers. The research design was one group pretest posttest quasi experimental design. Descriptive research approach has been carried out. By using non-probability convenience sampling technique a total of 60 postnatal mothers were included in this study and the data has been collected through structured knowledge and attitude questionnaires. The results concluded that there was no significant association between the baseline variables with knowledge, attitude scores and the demographic variables.

Summary:

Thestudyconcludedconcludedtatonly19%ofmothershadmothershaddequateknowledgeknowledgede

Conclusion:-

The above results stress the importance of organizing health education topostnatalmothers,broadcasting abouthumanmilk bank inantenatalOPD, postnatalward&NICUwillenhancetheirknowledgeandincreasethenumberofdonors.Handouts stressingtheimportanceofhumanmilkdonation should beprovided to allpostnatalmother

References:-

- 1. AlencarLC, SeidlEM. Breastmilkdonation: women's donorexperience. Rev. SaudePublica. 2009;43:70-77.
- 2. Arnold LDW.Humanmilkbanking in North America. JHumLact.1997;13:159-62.
- 3. Barin J, Lotscher M. Contextualizing Human Milk Banking and Milk Sharing Practices and Perception in Switzerland. THE MILK GAP report. 2018 (August)
- 4. DennisCL. Breastfeeding Initiation and Duration. J Obstet Gynecol Neonatal Nurs. 2001;31:12-32.
- 5. Donor breast milk banks: the operation of donor milk bank services. NICE clinical guideline 93. 2018 (July).
- 6. Ghuge S, Aghamkar J, Salvi R. A study to assess the knowledge and attitude regarding donating milk to the human milk bank among postnatal mothers of selected hospital. International Journal of Applied Research. 2018; 4(9): 94-98.
- 7. GoodfellowHE,ReimersP,BallardK,CoutsoudisA.Perceptionsofcommunity-basedhumanmilkbanks beforeand aftertraining ina resource-limited SouthAfrican setting. S Afr J ChildHealth.2016;10(1):83-86.
- 8. Iloh K, Osuorah C, Ndu I, Asinobi I, Anyim I,et al. Perception of donor breast milk and determinants of its acceptability among mothers in a developing community: a cross-sectional multi-center study in south-east Nigeria. International Breastfeeing Journal . 2018; 13:47 : 1-2.
- 9. Kim JH, Unger S. Human Milk Banking. Canadian Pediatric society Nutrition and Gastroenterology Committee.Paediatr ChildHealth .2010;15:595-98.
- 10. Mackenzie C, Hons B, Javanparast S, Newman L, Hons B. Mother's knowledge of and attitudes toward human milk banking in south Australia: A qualitative study. Journal of Human Lactation. 2013; 29(2):222-29.
- 11. McGuireW,AnthonyMY.Donorhumanmilkversusinfantformulaforpreventingnecrotizingenter colitis inpreterminfants:systematicreview. ArchDisChild Fetal NeonatalEd.2003;88:11-14.
- 12. WilliamsAF, FisherC, GreasleyV, TraylerH, WoofrideMW. Humanmilkbanking. JTropPediatr. 1985; 31:185-90.
- 13. Wong'sTextbookofEssential pediatricNursing .7thed.Philadelphia:Elsevier publications;2005.
- 14. YauSY,LeungJCY.PerceptionsofbreastfeedingMothersonBreastMilkDonationandEstablishmentofHuman BreastMilkBankinHong Kong. IntJ NursPract .2015; 2:72-80.