



Systematics, Morphology and Biogeography

Phylogeny of the Augochlora clade with the description of four new species (Hymenoptera, Apoidea)

Rodrigo Barbosa Gonçalves

Universidade Federal do Paraná, Departamento de Zoologia, Curitiba, PR, Brazil



ARTICLE INFO

Article history:

Received 19 September 2018

Accepted 20 December 2018

Available online 3 January 2019

Associate Editor: Rodrigo Feitosa

Keywords:

Augochlorini

Bees

Neotropical

Systematics

Taxonomy

ABSTRACT

The *Augochlora* clade includes four genera: *Augochlora* Smith, *Augochlorella* Sandhouse, *Ceratalictus* Moure, and *Pereirapis* Moure. This is one of the richest and most widespread groups of Augochlorini bees. There are about 150 species, which occur from Argentina to Canada. The species of *Augochlora* clade are considered solitary to facultatively social, except *Ceratalictus* for which nothing is known. Wood nesting behavior arose once in the clade, in *Augochlora* sensu strictu. The objective of this study is to describe four new species and to present a revised phylogenetic analysis of the *Augochlora* clade for the placement of these species. The morphological matrix comprised 77 characters and 42 terminals, and resulted in two most parsimonious trees. The monophyly of the *Augochlora* clade is corroborated. *Ceratalictus* and *Pereirapis* are considered as sister groups and *Ceratalictus inflexus* sp. nov. came as sister to other species of *Ceratalictus*. *Augochlora* and *Augochlorella* are monophyletic and sister groups. Both extant subgenera of *Augochlora* were corroborated as monophyletic. *Augochlorella comis* is considered as sister group to the rest of *Augochlorella* species. All *Augochlorella* new species described belong to the *Augochlorella ephyra* group. *Augochlorella kelliiae* sp. nov. is phylogenetically related to *Augochlorella una*. *Augochlorella procliva* sp. nov. and *Augochlorella mavricera* sp. nov. constitute a clade with *Augochlorella acarinata*. Including the new species, *Augochlorella* has 19 species and *Ceratalictus* 11 species. A revised key for species of *Augochlorella* and *Ceratalictus* is also presented in the Supplementary Information.

© 2019 Sociedade Brasileira de Entomologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Augochlorini is a monophyletic bee tribe and the main lineages divergence dating around 55–20 Ma (Hedtke et al., 2013; Gonçalves, 2016). The tribe comprises 663 described species primarily distributed in the Neotropical region (Moure, 2007). Species of Augochlorini have a remarkable range of behaviors and habits, including solitary, semisocial, primitively eusocial, crepuscular, nesting in the soil and wood, and cleptoparasitism (see Michener, 2007 for a review). These diverse and ecologically interesting habits make the augochlorine bees a good group for the evolution studies of such behaviors. In a recent molecular and morphological phylogeny, Gonçalves (2016) recognized seven suprageneric clades for the tribe and the *Augochlora* clade was recovered sister to the *Megalopta* group (Gonçalves, 2016; Meira and Gonçalves, 2018).

The *Augochlora* clade includes *Augochlora* Smith, *Augochlorella* Sandhouse, *Ceratalictus* Moure, and *Pereirapis* Moure (Eickwort, 1969). The taxonomic treatment of these genera varies among authors, Michener (2007) considered *Ceratalictus* and *Pereirapis*

as subgenera of *Augochlorella*, but others treated them as separate genera (Eickwort, 1969; Danforth and Eickwort, 1997; Engel, 2000; Coelho, 2004; Moure, 2007; Coelho and Gonçalves, 2010; Eickwort and Eickwort, 1973). The late classification is supported by multiples phylogenies (Danforth and Eickwort, 1997; Engel, 2000; Coelho, 2004; Gonçalves, 2015, 2016) in which the genus *Augochlorella* does not form a monophyletic group with *Ceratalictus* and *Pereirapis*. Michener's classification creates a paraphyletic group in relation to *Augochlora*.

Augochlora is the most speciose genus of the group, with 115 species, ranging from Argentina and Chile to Canada (Moure, 2007). *Augochlora* has three subgenera: *Augochlora* s.s. (87 species), *Electraugochlora* Engel (one fossil species), and *Oxystoglossella* Eickwort (27 species). No revisionary or phylogenetic studies have been conducted for the genus. *Pereirapis* is distributed from southern Brazil to Mexico, and it has eight species according to Moure (2007), although Engel (2000) considered the genus to be monotypic with all species synonym with *P. semiaurata* (Spinola, 1853). Coelho (2004) revised *Augochlorella*, recognizing 16 species distributed from Argentina to Canada. The *Augochlorella* phylogeny supports the placement of *Augochlorella* as a sister group to the rest of the *Augochlora* clade (Coelho, 2004). *Ceratalictus* taxonomy

E-mail: goncalvesrb@gmail.com

<https://doi.org/10.1016/j.rbe.2018.12.003>

0085-5626/© 2019 Sociedade Brasileira de Entomologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

was revised by Coelho and Gonçalves (2010) who recognized 10 species. Gonçalves (2015) provided a phylogeny of *Ceratalictus* species and also supported the placement of *Augochlorella* as sister to the remaining genera. Recently, however, Gonçalves (2016) recovered the following different topology using morphology and DNA ((*Augochlora*, *Augochlorella*), (*Ceratalictus*, *Pereirapis*)).

The nesting biology and behavior for some species of all genera are known, except *Ceratalictus*. Studied species of *Augochlora* (*Oxytostiglossella*), *Augochlorella* and *Pereirapis* are soil nesters and can be facultatively eusocial (Eickwort, 1969; Eickwort and Eickwort, 1972; Danforth and Eickwort, 1997 see Michener, 2007 for a review). *Augochlora* s.s. species nest in rotten-wood and are also solitary or facultatively eusocial (Stockhamer, 1966; Dalmazzo and Roig-Alsina, 2012, 2015; Wcislo et al., 2003) and are not exclusively solitary as previously postulated (Stockhamer, 1966; Danforth and Eickwort, 1997). Interestingly, intraspecific variation is also known for some species of the *Augochlora* clade, such as *Augochlora iphigenia* Holmberg, 1899, *Augochlorella comis* (Vachal, 1911) and *Augochlorella aurata* (Smith, 1853) (see Danforth and Eickwort, 1997 for a review).

Since the taxonomic revisions of Coelho (2004) and Coelho and Gonçalves (2010) new species accumulate in *Augochlorella* and *Ceratalictus*. A new *Ceratalictus* was already known by the author but with uncertainty on generic placement, mainly due a peculiar genital capsule. The main goal of this study is to present a revised phylogenetic analysis for the *Augochlora* clade for the placement of four new species. Other objectives are to compile and revise the morphological characters of previous studies into a more complete dataset, and to describe one new *Ceratalictus* and three new *Augochlorella* species. A revised key for species of *Augochlorella* and *Ceratalictus* is also presented in the Supplementary Information.

Material and methods

Phylogenetic analysis

As ingroup I selected seven species of *Augochlora* representing the extant subgenera, one species of *Pereirapis*, and all known species of *Augochlorella* and *Ceratalictus*. The new species of *Augochlorella* and *Ceratalictus* were also included as terminals. The included *Augochlora* species are distributed from Argentina, Brazil and Paraguay. There are several genus level names available for *Augochlora* but they are not tested here. All species were directly examined except some of *Augochlorella aurata* group. For those species, the character coding was the same present by Coelho (2004) with few modifications based on the original drawings and descriptions. As outgroup I selected four species of Augochlorini including two species of the *Megalopta* clade, the sister group of *Augochlora* clade (Gonçalves, 2016; Meira and Gonçalves, 2018). *Pseudaugochlora graminea* (Fabricius, 1804) was used to root the cladogram based on the topologies of Gonçalves (2016) and Meira and Gonçalves (2018). For the character statements, I firstly revised the original datasets of Coelho (2004), and Gonçalves (2015, 2016). The later incorporated previous characters from Eickwort (1969) and Engel (2000). The character coding from the original studies were completely rescored based on direct observation and new interpretations. The second step was the inclusion of new characters statements based on external morphology and male genitalia. The writing of all characters statements followed the recommendations of Sereno (2007). Inapplicable character states for contingent characters are indicated by “-”, missing character states by “?”. The final matrix comprised 77 characters and 42 terminals, both matrix and character list are

provided in the Supplementary Information. Parsimony analyses were carried out in Tree Analysis using New Technology Search (TNT, Goloboff et al., 2008), using equal weights, and the following settings (ratchet weighting probability 5%, 200 iterations), tree-drifting (50 cycles), and tree-fusing (five rounds, minimum length set to be hit 1000 times). All characters were treated as non-additive. Bootstrap (1000 replications) and Bremer support were also calculated with TNT. Trees were viewed and edited with WinClada ver. 1.00.08 (©KC Nixon) and GIMP 2.8.16 (©The GIMP Team).

Taxonomy

The descriptions were organized following the body organization, tagmata, segments, and sclerites, with special attention given to the following: structure (general organization), sculpturing (impressions and elevations on sclerite surface), color and pubescence. Structure terminology follows Eickwort (1969) and Michener (2007) except the basal area of propodeum referred here as the metapostnotum. I use the abbreviations T1, T2, etc., to denote the metasomal terga; S1, S2, etc., to denote metasomal sterna; and F1, F2, etc., to denote the flagellomeres. For examined material the quotation marks surround exact transcriptions of individual labels, while the backslash (\) indicates different lines on the same label. The material cited here is deposited in the following institutions: BMNH – British Museum of Natural History, London, UK; DZUP – Coleção Entomológica Pe. Jesus Santiago Moure, Curitiba, Brazil; MZSP – Museu de Zoologia da Universidade de São Paulo, São Paulo, Brazil; SEMC – Snow Entomological Museum Collection, Lawrence, USA; UFMG – Universidade Federal de Minas Gerais, Belo Horizonte, Brazil; UFV – Universidade Federal de Viçosa, Viçosa, Brazil; USNM – Smithsonian National Museum of Natural History, Washington, USA.

I measured the following: eye notch width (as a measure of head width); head length (distance from base of clypeus to vertex apex); glossa length; clypeus maximum width and length; supraclypeal length (distance from clypeus base to lower tangent of alveoli); distance between upper tangent alveolus and tangent of lower median ocellus; intertegular distance; first tergum maximum width; second tergum maximum width; total length (approximate by measuring head and mesosoma plus metasoma). All measurements are in millimeters. Relative size comparisons used the mid ocellus diameter (=OD) or eye maximum width in lateral view (=EW). The punctures interspaces (i) are described in relation to puncture diameter (pd). Sculpture terminology follows Harris (1979) including conflected for crowded punctures. For some sculpturing I use two terms to describe mixed or transitional patterns, as areolate–rugose. For punctuation, sparsely punctate is used for punctures separated by more than one puncture diameter, coarsely punctate for punctures comparatively coarser than ordinary covering punctures. For microsculpture, visible only in magnifications above 50×, I use the prefix micro-, the term polished is applied for the absence of microsculpture.

Most specimen examination was made with Olympus SZ51 and SZ61 using white led ring illumination. Measurements were made with a Leica Stemi DV4 and an eyepiece micrometer. Photographs were taken with a Nikon D700 and 107 mm Sigma Macro lens using Helicon Remote for controlling image capture. Illumination followed the system of Kawada and Buffington (2016). Image stacking was made with Helicon Focus (version 6.0.18) – render method based on Method C (Pyramid). Final image adjustments (unsharp mask filtering, level control, background color setting) were with GIMP 2.8.16 (©The GIMP Team).




Fig. 1. Phylogeny of *Augochlora* clade (L: 170, Ci: 53, Ri: 84, fast optimization). Black and white circles indicate unique and homoplastic changes, respectively. Red numbers indicate character states not optimized in unambiguous optimization. Numbers above branches: Bootstrap and Bremer support values.

Results

Phylogenetic analysis

The phylogenetic analysis resulted in two most parsimonious trees with 158 steps, the consensus tree is presented in Fig. 1. The *Augochlora* clade is monophyletic and supported by five synapomorphies (Fig. 1). *Ceratalictus* and *Pereirapis* are sister groups with four synapomorphies in fast optimization (one in unambiguous optimization). *Ceratalictus* is supported by seven synapomorphies (six in unambiguous optimization) and *Ceratalictus inflexus* sp. nov. is sister group to the remaining *Ceratalictus* species. *Augochlora* and *Augochlorella* are sister groups sharing four synapomorphies, from these only one supports the group unambiguously. *Augochlora* is monophyletic and the clade shows high support values compared to other genera, its extant subgenera are also corroborated as monophyletic. *Augochlorella comis* is considered as sister group to the remaining *Augochlorella* species. The three new species of *Augochlorella* belong to the "ephyra group" from Coelho (2004). *Augochlorella kelliae* sp. nov. is related to *Augochlorella una*. *Augochlorella procliva* sp. nov. and *Augochlorella mavricera* sp. nov.

constitute a clade with *Augochlorella acarinata*. Based on these results, the generic classification from Moure (2007) was corroborated and the new species are described under *Augochlorella* and *Ceratalictus* at the generic level.

Taxonomy

Augochlorella kelliae sp. nov. (Figs. 2 and 3)

Diagnosis. Females of this species are similar to *Augochlorella una* and can be separated from the remaining *Augochlorella* by the olive green head and mesosoma (not olive in other species), elongate head (wider than longer in other species), and microreticulate posterior and lateral surfaces of the propodeum (not microreticulate in other species) (Fig. 2). From *A. una*, the new species can be separated by the metapostnotum with strong posterior marginal angles, being almost trapezoidal (Fig. 2C), preoccipital ridge carinate, mesoscutum punctures sparser on the median surface (interspaces equal to one puncture diameter on *A. una*, and about two puncture diameter on *A. kelliae* sp. nov.) and the scutellar punctures are not strong and contiguous. The male can be separated from other species by




Fig. 2. Holotype female of *Augochlorella kelliæ* sp. nov. (A) Habitus, (B) frontal view of head, (C) dorsal view of mesosoma. Scale bar: 0.5 mm.




Fig. 3. Paratype male of *Augochlorella kelliæ* sp. nov. (A) Dorsal view; (B) frontal view of head; (C) ventral view of genitalia. Scale bar: A and B at 0.5 mm, C at 0.25 mm.

the combination of clypeus with a transverse yellow band, hind leg with short setae, preoccipital ridge carinate, and metapostnotal carinae almost reaching the posterior margin (Fig. 3).

Description. Holotype female. Measurements (mm). Eye notch width: 1.3; head length: 2.0; glossa length: 1.2; clypeus width: 0.7; clypeus length: 0.5; supraclypeal length: 0.4; clypeus ocellar distance: 0.8; intertegular distance: 1.4; T1 width: 1.8; T2 width: 2.0; body length: 7.5. Head. Labrum basal elevation orbicular, black. Mandible black with subapical amber spot; subapical tooth not strongly produced; long setae (2–3 OD) on outer and ventral margins. Epistomal angle orthogonal. Clypeus sparsely punctate, microreticulate on basal portion; erect setae with 1–1.5 OD; apical surface darkened, most surface olive green. Supraclypeal area sparsely punctulate, microreticulate near clypeus; fine erect setae; olive green. Paraocular area punctate, strongly microreticulate; olive green, some weak copper reflections near eye notch; branched setae with 1–1.5 OD. Antennae: scape dark brown with very weak metallic highlights; flagellomeres amber ventrally,

especially on F9–10; F1–2 and base of F3 setose, with inconspicuous sensilla, remaining flagellomeres with conspicuous sensilla. Frons not depressed; confluent punctate, strongly microreticulate; weak blue iridescence; short and fine setae. Preoccipital ridge carina not reaching the gena. Vertex coarsely punctate becoming rugose near ridge; long plumose (short branches) setae. Gena puncticulate, microlinéate on postgena; green with very weak copper reflections; white plumose setae (1.5–2 OD). Hypostomal ridge carinate. Mesosoma. Pronotum dorsal ridge weakly carinate; lateral angle not projected, rounded; lateral ridge rounded; dorsal surface lineate and puncticulate; green; with few tomentum on dorsal surface. Propleuron brown with green highlights. Prosternum green. Fore leg: coxa dark green, trochanter and femur brown, tibia light brown on outer surface, base of tibia and pretarsus yellow; long white setae on coxa to femur, yellow erect setae on tibia and tarsus. Mesoscutum anterior border weakly acuminate; median sulcus well impressed; parapsidal lines weakly impressed; punctate laterally, more sparse on median surface, microlineations more conspicuous on anterior surface; olive green; short (=OD) and sparse white setae. Scutellum median sulcus very weakly impressed; puncticulate, some scattered punctures, polished; bright olive green; longer setae (=2.5–3 OD) on posterior margin. Mesepisternum rugulose on preepisternum, irregularly punctate on hypoepimeral area, sulci weakly foveate, densely punctate near scrobe, becoming more evenly punctate ventrally; olive green, weak copper reflections; covered with plumose setae (=1.5 OD). Mid leg brown; yellow setae on femur to tarsus. Tegula concentric lineate; dark amber, weak green reflections anteriorly. Fore wing: marginal cell apex rounded; membrane hyaline; black microtrichia. Metanotum median sulcus very weakly impressed; puncticulate to rugulose; green with bluish iridescence; long plumose setae (2–3 OD). Metepisternum lineate above superior metapleural pit, rugulose–punctate below; green with copper and bluish reflections. Hind leg: basitibial plate apex rounded; tibial spur serrate; brown; long plumose setae on ventral coxa to femur; ventral tibial setae short and branched. Hind wing hyaline. Metapostnotum weakly inclined; weakly depressed; apex rounded; less twice as long as metanotum; rugulose on most surface, except a stripe on posterior margin, microreticulate; olive green. Propodeum posterior surface puncticulate, few scattered punctures, microreticulate; olive green; short plumose and some erect (1–2 OD) setae; propodeum lateral surface puncticulate, few scattered punctures, microreticulate; olive green; with thicker and longer (3–4 OD) setae. Metasoma. T1 anterior and dorsal surfaces not angled between; sparsely puncticulate, microlinéate; olive green on anterior and lateral surfaces, dark brown on median surface. T2 puncticulate; olive green on anterior and lateral surfaces, dark brown on median surface; short. T3–T4 puncticulate; olive green with some dark brown on median surface, brown on marginal area; short. T5 punctate; dark brown; covered with black long erect setae. Pygidial plate short, truncate. S1 without tubercle; lineate; amber with green highlights on median surface; very long setae (=3 OD) on median surface. S2–S4 lineate; brown to light brown; subapical setae very long (about 5 OD) and with recurved apices. S5 lineate; brown; long setae (about 4 OD), plumose. S6 lineate; brown; yellowish setae on apical margin.

Paratype male. Measurements (mm). Eye notch width: 1.2; head length: 1.8; clypeus width: 0.6; clypeus length: 0.5; supraclypeal length: 0.3; clypeus ocellar distance: 0.7; intertegular distance: 1.3; T1 width: 1.2; T2 width: 1.3; body length: 9. Head. Labrum and mandible yellow. Clypeus, epistomal angle straight; sparsely punctate, weakly microreticulate; white setae on middle, longer setae on apex; yellow stripe on apical margin, green. Supraclypeal area sparsely punctate, weakly microreticulate; green. Paraocular area sparsely punctate, weakly microreticulate; green, coppery along eye margins; long (>2 OD) branched setae, yellow. Antennae F2 0.7x F1 length, F3 = F1, remaining flagellomeres relatively

longer; scape and pedicel light brown, flagellomeres yellowed ventrally; F11 not flattened; without sensilla placodea patches. Frons not depressed above antennal sockets; coarsely punctate; bright green, few coppery reflections near paraocular; few erect setae. Pre-occipital ridge carinate, some portions lamellate. Vertex punctate becoming rugose near ridge; very long plumose (short branches) setae. Gena puncticulate becoming lineate on postgena; green, some coppery reflections; white plumose setae (1.5–2 OD). Hypostomal ridge carinate. *Mesosoma*. Pronotum dorsal ridge carinate; lateral angle almost orthogonal; lateral ridge rounded; dorsal surface rugose; bright olive green, weakly metallic laterally; with some plumose setae on dorsal surface. Fore leg: coxa to femur bright metallic green, tibia and tarsus light brown; short yellow erect setae. Mesoscutum anterior border slightly projected over pronotum, weakly lobed; median sulcus well impressed; parapsidal lines well impressed; punctate, sparsely on anterior lobe; bright olive green; posterior margin weakly covered with plumose setae. Scutellum median sulcus weakly impressed; punctate, polished; bright olive green; anterior surface covered with plumose setae. Mesepisternum punctate, polished; bright olive green; short plumose setae. Mid leg: coxa brown with metallic reflections, remaining leg light brown; short yellow erect setae. Tegula concentric lineate; brown, green reflections anteriorly. Forewing weakly infuscate on apex. Metanotum median sulcus weakly impressed; puncticulate, polished; bright olive green; anterior surface densely covered with erect setae. Metepisternum lineate above superior metapleural pit, rugulose-punctate below; bright olive green. Hind leg: coxa brown with metallic reflections, remaining leg light brown; short yellow erect setae, shorter than diameter of segments. Metapostnotum weakly inclined and depressed; twice as long as metanotum; apex rounded; carinate, posteriorly only weakly microreticulate; green. Propodeum: posterior surface posterior pit shallow, punctate, microreticulate, olive green, with fine setae; lateral surface punctate, olive green, with thicker setae. *Metasoma*. T1 very sparsely puncticulate; light brown with metallic reflections, yellow marginal areas. T2–5 very sparse puncticulate; light brown with metallic reflections, yellow marginal areas; vestiture white, with weak row of long white setae on basis. T6 green; long setae. T7 black; short black setae. S1 margin straight; sparsely punctate, microlineate; light brown; with very short setae. S2–S5 straight; brown; covered with short white setae on median surface, posterior surface with long setae (>4 OD). S4 margin concave; with median, triangular patch of setae; lateral setae longer. S6 weakly concave; brown; black setae on apical margin. S7 posterior margin sinuose. S8 posterior margin not produced; 0.3 times S8 width. *Genitalia* (Fig. 3C). Gonapophysis broad, longer than gonostylus and without dorsal crest; with large ventral prong and lateral teeth. Gonobase short (0.6 times gonocoxite length); dorsal lobes weakly produced; ventral bridge narrow. Gonostylus basal process reduced; parapenial lobe reduced; ventral process with outer and inner lobes, inner lobe divided and asymmetric, setose; dorsal lobe transversal. Volsella longer than wider; oblique, corners acuminate; without hook.

Comments. Coelho (2004) listed a *A. una* paratype female as “1?m#, Minas Gerais, Serra da Moeda, Zona Metalúrgica, 15.x.1997 (E. A. B. Almeida) (BHMH)”, the author listed all specimens of *A. una* with male symbols but the species is only known from females. As the question mark could represent a dubious identification I speculate that this paratype can be a specimen of *A. kelliiae* sp. nov. In this case *A. una* is known only from Central Brazil and *A. kelliiae* sp. nov. is known only for Minas Gerais.

Type material. Holotype female (DZUP): “Brasil, Minas Gerais, São Tomé das Letras, 21°43'S, 44°58'W, 12.I.2010, K. Ramos and V. Kanamura”. Paratypes: three females (DZUP) with same data of holotype; two females (DZUP) “Brasil, Minas Gerais, Serra do Cipó, X.2018, Letícia Graf leg.”; one male (UFMG) “Abelhas de Canga”.



Fig. 4. Holotype female of *Augochlorella mavricera* sp. nov. (A) Habitus; (B) frontal view of head; (C) dorsal view of mesosoma. Scale bar: 0.5 mm.




Fig. 5. Paratype male of *Augochlorella mavricera* sp. nov. (A) Dorsal view; (B) frontal view of head; (C) ventral view of genitalia. Scale bar: A and B at 0.5 mm, C at 0.25 mm.

Parque Estadual da Serra do Rola-Moça 25745–75506” “Nova Lima MG, BRASIL 14/03/2013 J.E. Santos Jr”.

Etymology. In honor of Dr. Kelli S. Ramos, friend, melittologist, and collector of the type.

Augochlorella mavricera sp. nov.
(Figs. 4–6C,D)

Diagnosis. The female of this species is very similar to *A. acarinata*. The body of *A. mavricera* sp. nov. is mostly dark green with some violet reflections on clypeus and supraclypeal area and black dorsal surfaces of terga (Fig. 4) while *A. acarinata* is bright green with some coppery reflections. Also, *A. mavricera* sp. nov. punctation is sparser than on *A. acarinata* (i>2pd on lower paraocular and i>pd on mesoscutum). The male can be easily separated of *A. acarinata* by the dark body, including legs (Figs. 5 and 6D), compared with yellowish color of *A. acarinata* (Fig. 6A,B). The S4 is slightly bilobed (Fig. 6C) in the new species, while *A. acarinata* males have S4 strongly bilobed (Fig. 6A).

Description. Holotype female. Measurements (mm). Eye notch width: 1.3; head length: 1.6; glossa length: 0.6; clypeus width: 0.7; clypeus length: 0.4; supraclypeal length: 0.3; clypeus ocellar distance: 0.9; intertegular distance: 1.1; T1 width: 1.7; T2 width: 1.8; body length: 7.0. *Head*. Labrum basal elevation orbicular, black. Mandible black with subapical amber spot; subapical tooth




Fig. 6. Ventral view of male metasoma and hind leg. (A) Metasoma of *A. acarinata*; (B) hind leg of *A. acarinata*; (C) metasoma of *A. mavricera* sp. nov.; (D) hind leg of *A. mavricera* sp. nov. S4 apical margin indicated in red. Scale bar: 0.5 mm.

not strongly produced; long setae (2–3 OD) on outer and ventral margins. Epistomal angle orthogonal. Clypeus sparsely punctate, weakly microreticulate on basal portion; erect setae with 1–1.5 OD; apical surface darkened, subapical area violet, remaining surface bright green. Supraclypeal area sparsely punctulate, polished; fine erect setae; green with violet reflections. Paraocular area punctate, strongly microreticulate; olive green, some weak violet reflections near eye notch; branched setae with 1–1.5 OD. Antennae; scape dark brown with very weak metallic highlights, flagellomeres weakly amber ventrally; F1–3 setose, with inconspicuous sensilla, remaining flagellomeres with conspicuous sensilla. Frons not depressed; conflected punctate, strongly microreticulate; olive green; short and fine setae. Preoccipital ridge not carinate. Vertex coarsely punctate becoming rugose near ridge; long plumose (short branches) setae. Gena puncticulate, microliniate on postgena; dark green to black; white plumose setae (1.5–2 OD). Hypostomal ridge carinate. *Mesosoma*. Pronotum dorsal ridge weakly carinate; lateral angle not projected, rounded; lateral ridge rounded; dorsal surface lineate and puncticulate; dark green; with few tomentum on dorsal surface. Propleuron brown with green highlights. Prosternum green. Fore leg: coxa dark green, trochanter and femur brown, tibia brown on outer surface, base of tibia and pretarsus light brown; long yellowish setae on coxa to femur, yellow erect setae on tibia and tarsus. Mesoscutum anterior border weakly acuminate; median sulcus well impressed; parapsidal lines weakly impressed; punctate laterally, more sparse on median surface (i=pd), microlineations more conspicuous on anterior surface; green with violet

reflections; short (=OD) and sparse white setae. Scutellum median sulcus very weakly impressed; punctate to conflected punctate, microreticulate; bright olive green; longer setae (=2.5–3 OD) on posterior margin. Mesepisternum conflected punctate, microreticulate; olive green, weak violet reflections; covered with plumose setae (=1.5 OD). Mid leg brown; yellow setae on femur to tarsus. Tegula concentric lineate; dark amber, weak green reflections anteriorly. Fore wing: marginal cell apex rounded; membrane hyaline; black microtrichia. Metanotum median sulcus very weakly impressed; rugulose, microreticulate; green with bluish iridescence; long plumose setae (2–3 OD). Metepisternum lineate above superior metapleural pit, conflected punctate below; green with violet reflections. Hind leg: basitibial plate apex rounded; tibial spur serrate; brown; long plumose setae on ventral coxa to femur, ventral tibial setae not too long and with short branches. Hind wing hyaline. Metapostnotum weakly inclined; weakly depressed; apex rounded to trapezoidal; less twice as long as metanotum; rugulose on median surface, carinate laterally, microreticulate; dark olive green with some violet reflections; Propodeum posterior surface puncticulate, few scattered punctures, microreticulate to granular; dark olive green; short plumose and some erect (1–2 OD) setae. Propodeum lateral surface puncticulate, microreticulate; olive green; with thicker and longer (3–4 OD) setae. *Metasoma*. T1 anterior and dorsal surfaces not angled between; sparsely puncticulate, microliniate; olive green reflections on lateral surfaces, black on median surface, few copper reflections elsewhere; short. T2 puncticulate; olive green reflections on lateral surfaces, black on median surface, few copper reflections elsewhere; short. T3–T4 puncticulate; olive green reflections on lateral and marginal surfaces, black on median surface, few copper reflections elsewhere; short. T5 punctate; dark brown; covered with black long erect setae. Pygidial plate short, truncate. S1 without tubercle; lineate; amber with green highlights on median surface; very long setae (=3 OD) on median surface. S2–4 lineate; brown to light brown; subapical setae very long (about 5 OD) and with recurved apices. S5 lineate; brown; long setae (about 4 OD), plumose. S6 lineate; brown; yellowish setae on apical margin.

Paratype male. Measurements (mm). Eye notch width: 1.2; head length: 1.6; clypeus width: 0.6; clypeus length: 0.4; supraclypeal length: 0.3; clypeus ocellar distance: 0.9; intertegular distance: 1.2; T1 width: 1.4; T2 width: 1.6; body length: 7.0. **Head.** Labrum and mandible amber; Clypeus: epistomal angle straight; punctate, very weakly microreticulate; white setae on middle, longer setae on apex; yellow marking on latero-apical margin, green. Supraclypeal area sparsely punctate, very weakly microreticulate; green. Paraocular area sparsely punctate, weakly microreticulate; olive green; long (>2 OD) branched setae, yellow. Antennae F2 0.8x F1 length, F3 = F1, remaining flagellomeres relatively longer; scape and pedicel light brown, flagellomeres mostly yellowed; F11 flattened dorso-ventrally; without sensilla placodea patches. Frons not depressed above antennal sockets; coarsely punctate; olive green; few erect setae. Preoccipital ridge not carinate. Vertex punctate becoming rugose near ridge; very long plumose (short branches) setae. Gena puncticulate, almost polished on postgena; olive green, some coppery reflections on postgena; white plumose setae (1.5–2 OD). Hypostomal area, ridge carinate. *Mesosoma*. Pronotum dorsal ridge carinate; lateral angle almost orthogonal; lateral ridge rounded; dorsal surface rugose; bright olive green, weakly metallic laterally; with some plumose setae on dorsal surface. Fore leg: coxa to femur with some metallic reflections, tibia and tarsus dark brown; short yellow erect setae. Mesoscutum anterior border slightly projected over pronotum, weakly lobed; median sulcus well impressed; parapsidal lines well impressed; punctate, sparsely on anterior lobe; bright olive green; posterior margin weakly covered with plumose setae. Scutellum median sulcus weakly impressed; punctate, polished; bright olive green;

anterior surface covered with plumose setae. Mesepisternum punctate, polished; bright olive green; short plumose setae. Mid leg: coxa brown with metallic reflections, remaining leg light brown; short yellow erect setae. Tegula concentric lineate; brown, green reflections anteriorly. Forewing weakly infuscate on apex. Metanotum median sulcus weakly impressed; puncticulate, polished; bright olive green; anterior surface densely covered with erect setae. Metepisternum lineate above superior metapleural pit, punctate below; bright olive green; Hind leg: coxa brown with metallic reflections, remaining leg light brown; apex of ventral surface of femur with apical very long thick setae (7 OD), apex of ventral surface of tibia with long setae (4 OD), ventral surface of basitarsus with tuft of erect setae (3 OD) on a convexity. Metapostnotum weakly inclined; weakly depressed; twice as long as metanotum; apex rounded; rugose, some carina laterally; green. Propodeum posterior surface posterior pit shallow, punctate, microreticulate, olive green, with fine setae; lateral surface punctate, olive green, with thicker setae. *Metasoma*. T1 very sparsely puncticulate; brown with lateral metallic reflections; short. T2 very sparse puncticulate; light brown with metallic reflections, yellow marginal areas; vestiture white, with weak row of long white setae on basis. T3–T5 as T2, some bluish iridescence. T6 green; long setae. T7 black; short black setae. S1 margin straight; sparsely punctate, microliniate; light brown; with very short setae. S2–S3 and S5 straight; brown; covered with few short white setae on median surface, posterior surface with some long setae (>4 OD). S4 weakly bilobed (Fig. 6C); with apical row of thick setae; lateral setae very short. S6 weakly concave; brown; black setae on apical margin. S7 posterior margin sinuose. S8 posterior margin not produced: 0.3 times S8 width. *Genitalia* (Fig. 5C). Gonapophysis broad; longer than gonostylus; without dorsal crest; with large ventral prong; with lateral teeth. Gonobase short (0.6 gonocoxite length); dorsal lobes weakly produced; ventral bridge narrow. Gonostylus basal process reduced; parapenial lobe reduced; ventral process with outer and inner lobes, inner lobe divided and asymmetric, setose; dorsal lobe transversal. Volsella longer than wider; oblique, corners acuminate; without hook.

Type material. Holotype female (DZUP): “Parque Saint-Hilaire\ Brazil: RS, Viamão, 02/03/2017\ Coletado em \ Euphorbia selloi (11–14 h)\ Saibreira.Fogo Recente”. Paratype: one male (DZUP) “Parque Saint-Hilaire\ Brazil: RS, Viamão, 21/12/2015\ Coletado em\ Richardia grandiflora (08–11 h)\ NOVA_Fogo Intermediário”.

Etymology. From Greek terms that mean black, *mavros*, and horn, *ceros*, in reference to the black F11, feminine.

Augochlorella procliva sp. nov.

(Fig. 7)

Diagnosis. Females of *A. procliva* sp. nov. can be separated from the similar *A. ephyra* and *A. iopoecila* by the combination of the weakly carinate preoccipital ridge, green head and mesosoma (Fig. 7), rugulose and declivitous metapostnotum, and the granular posterior and lateral surfaces of propodeum. The carinae of metapostnotum reach the edge posteriorly differently from *A. urania*, other similar species which presents carinate preoccipital ridge. Males are not known.

Description. Holotype female. Measurements (mm). Eye notch width: 1.4; head length: 1.9; glossa length: 0.8; clypeus width: 0.7; clypeus length: 0.7; supraclypeal length: 0.3; clypeus ocellar distance: 0.8; intertegular distance: 1.4; T1 width: 2.0; T2 width: 2.2; body length: 7.6. *Head*. Labrum basal elevation weakly orbicular, amber. Mandible black with subapical and apical amber spot; subapical tooth not strongly produced; long setae (2–3 OD) on outer and ventral margins. Epistomal angle orthogonal. Clypeus sparsely punctate, polished; erect setae with 1–1.5 OD; apical surface darkened, most surface bright green. Supraclypeal area sparsely punctulate, polished; fine erect setae; bright green. Paraocular area




Fig. 7. Holotype female of *Augochlorella procliva* sp. nov. (A) Habitus; (B) frontal view of head; (C) dorsal view of mesosoma. Scale bar: 0.5 mm.

punctate, sometimes conflcted, polished; green; branched setae with 1–1.5 OD. Antenna: scape dark brown with weak metallic highlights, flagellomeres amber ventrally, especially on F9–10; F1–3 setose, with inconspicuous sensilla, remaining flagellomeres with conspicuous sensilla. Frons not depressed; conflcted punctate; bright green. short and fine setae. Preoccipital ridge weakly carinate, carina not reaching the gena. Vertex coarsely punctate becoming rugose near ridge; long plumose (short branches) setae. Gena puncticulate, microliniate on postgena; green with very weak bluish reflections; white plumose setae (1.5–2 OD). Hypostomal ridge carinate. *Mesosoma*. Pronotum dorsal ridge weakly carinate; lateral angle not projected, rounded; lateral ridge rounded; dorsal surface lineate and puncticulate; green; with few tomentum on dorsal surface. Propleura brown with green highlights. Prosternum green. Fore leg coxa dark green, trochanter and femur brown, tibia light brown on outer surface, base of tibia and pretarsus yellow; long white setae on coxa to femur, yellow erect setae on tibia and tarsus. Mesoscutum anterior border weakly acuminate; median sulcus well impressed; parapsidal lines weakly impressed; punctate laterally, more sparse on median surface, microliniate; green, few copper reflections; short (=OD) and sparse white setae. Scutellum median sulcus very weakly impressed; punctate, becoming rugulose posteriorly; bright olive green; longer setae (=2.5–3 OD) on posterior margin. Mesepisternum rugulose, sulci weakly foveate; olive green, weak copper reflections; covered with plumose setae (=1.5 OD). Mid leg brown, tarsus light brown; yellow setae on femur to tarsus. Tegula concentric lineate; dark amber, weak green reflections anteriorly. Fore wing, marginal cell apex rounded; membrane hyaline; black microtrichia. Metanotum median sulcus very weakly impressed; puncticulate to rugulose; green with bluish iridescence; long plumose setae (2–3 OD). Metepisternum lineate above superior metapleural pit, rugulose–punctate below; green with copper and bluish reflections. Hind leg, basitibial plate apex rounded; tibial spur serrate; brown; long plumose setae on ventral coxa to femur, ventral tibial setae not too long and with short branches. Hind wing hyaline. Metapostnotum inclined; weakly depressed; apex rounded to trapezoidal; less twice as long as metanotum; entirely rugulose, some lateral carina extending to propodeum; green. Propodeum




Fig. 8. Holotype female of *Ceratalictus inflexus* sp. nov. (A) Habitus; (B) frontal view of head; (C) dorsal view of mesosoma. Scale bar: 0.5 mm.

posterior surface punctate rugulose, microreticulate; green; short plumose and some erect (1–2 OD) setae. Propodeum lateral surface punctate rugulose, microreticulate; green; with thicker and longer (3–4 OD) setae. Metasoma. T1 anterior and dorsal surfaces not angled between; sparsely punciculate, microliniate; copper green on anterior and lateral surfaces, dark brown on median surface; short. T2 punciculate; copper green on anterior and lateral surfaces, dark brown on median surface; short. T3–T4 punciculate; copper green with some dark brown on median surface, brown on marginal area; short. T5 punciculate; dark brown; covered with amber long erect setae. Pygidial plate short, truncate. S1 without tubercle; lineate; amber with green highlights on median surface; very long setae (=3 OD) on median surface. S2–S4 lineate; brown to light brown; subapical setae very long (about 5 OD) and with recurved apices. S5 lineate; brown; long setae (about 4 OD), plumose. S6 lineate; brown; yellowish setae on apical margin.

Type material. Holotype female (DZUP): “MANAUS – AM\ Brasil I 61\ C. ELIAS leg.”. Paratype: one female (DZUP), with same data as holotype.

Etymology. From Latin term *proclivus*, meaning inclined.

Ceratalictus inflexus sp. nov.

(Figs. 8 and 9)

Diagnosis. *Ceratalictus inflexus* sp. nov. is distinguished from other *Ceratalictus* by the combination of lateral angle of pronotum obtuse; surface of mesoscutal disk weakly microreticulate; carinae of meta-postnotum not reaching edge posteriorly; metapostnotum similar with metanotum in length and shape, and by the metasoma with blue reflections (Figs. 8 and 9). The female has the basal elevation occupying all the longitudinal portion of labrum, the male has the antennae with sensilla placodea patches.

Description. Holotype female. Measurements (mm). Eye notch width: 1.5; head length: 1.7; glossa length: 0.4; clypeus width: 0.8; clypeus length: 0.4; supraclypeal length: 0.3; clypeus ocellar distance: 1.1; intertegular distance: 1.5; T1 width: 2.0; T2 width: 2.2; body length: 8.0. Head. Labrum basal elevation orbicular, black. Mandible black with subapical amber spot; subapical




Fig. 9. Paratype male of *Ceratalictus inflexus* sp. nov. (A) Dorsal view; (B) frontal view of head; (C) ventral view of genitalia. Scale bar: A and B at 0.5 mm, C at 0.25 mm.

tooth not strongly produced; long setae (2–3 OD) on outer and ventral margins. Epistomal angle obtuse. Clypeus coarsely punctate, microreticulate; erect setae with 1–1.5 OD; apical surface darkened, blue green. Supraclypeal area sparsely punctate, microreticulate; fine erect setae; blue green. Paraocular area coarsely punctate becoming conflected, strongly microreticulate; blue green; branched setae with 1–1.5 OD. Antennae, scape dark brown with very weak metallic highlights, flagellomeres amber ventrally, especially on F9–10; F1–2 and base of F3 setose, with inconspicuous sensilla, remaining flagellomeres with conspicuous sensilla. Frons not depressed; conflected punctate, strongly microreticulate; dark blue green; short and fine setae. Preoccipital ridge carinate, carina not reaching the gena. Vertex coarsely punctate becoming rugose near ridge; long plumose (short branches) setae. Gena punciculate, microliniate on postgena; blue green, dark green on postgena; white plumose setae (1.5–2 OD). Hypostomal ridge carinate. Mesosoma. Pronotum dorsal ridge weakly carinate; lateral angle not projected, rounded; lateral ridge rounded; dorsal surface lineate and punciculate; blue green; with few tomentum on dorsal surface. Propleura brown with green highlights. Prosternum green. Fore leg coxa dark green, remaining leg dark brown; long white setae on coxa to femur, yellow erect setae on tibia and tarsus. Mesoscutum anterior border weakly acuminate; median sulcus well impressed; parapsidal lines weakly impressed; punctate laterally, more sparse on median surface, punctures with different diameters intermixed, microliniations only on anterior surface and near median sulcus; blue green; short (=OD) and sparse white setae. Scutellum median sulcus very weakly impressed; irregularly punctate, rugulose posteriorly; blue green; longer setae (2–2.5 OD) on posterior margin. Mesepisternum rugulose on preepisternum, sulci foveate, densely to conflected punctate, microreticulate; blue green, blue ventrally; covered with plumose setae (1.5 OD). Mid leg light brown; yellow setae on femur to tarsus. Tegula concentric lineate; dark amber, weak green reflections anteriorly. Fore wing, marginal cell apex rounded; membrane hyaline; black microtrichia. Metanotum median sulcus very weakly impressed; punciculate to rugulose; blue green; long plumose setae (2–3 OD). Metepisternum lineate above superior metapleural pit, rugulose–punctate below; green with copper and bluish reflections. Hind leg, basitibial plate apex rounded; tibial spur serrate; brown; long plumose setae on ventral coxa to femur, ventral tibial setae not too long and with short branches. Hind wing hyaline. Metapostnotum weakly inclined; weakly depressed; apex rounded; less twice as long as metanotum; carinate to irregularly carinate, carinae not reaching posterior margin, microreticulate; blue green. Propodeum posterior surface punciculate, few scattered setose punctures, microreticulate; blue green; short plumose

and some erect (1–2 OD) setae. Propodeum lateral surface puncticulate, few scattered punctures, microreticulate; blue green; with thicker and longer (3–4 OD) setae. *Metasoma*. T1 anterior and dorsal surfaces not angled between; sparsely punctate, microlineate; blue green with some amber to brown portions on median surface; short. T2 puncticulate, weakly microreticulate; blue green, black on marginal area; short. T3–T4 puncticulate, weakly microreticulate; blue green, black on marginal area; short. T5 punctate; dark brown; covered with black long erect setae. Pygidial plate short, truncate. S1 without tubercle; lineate; amber with green highlights on median surface; very long branched setae (=3 OD) on median surface. S2–S4 lineate; brown to light brown; subapical branched setae very long (about 4 OD). S5 lineate; brown; long setae (about 4 OD), plumose. S6 lineate; brown; yellowish setae on apical margin. **Paratype male.** Measurements (mm). Eye notch width: 1.3; head length: 1.8; clypeus width: 0.6; clypeus length: 0.5; supraclypeal length: 0.3; clypeus ocellar distance: 1.1; intertegular distance: 1.6; T1 width: 1.9; T2 width: 2.0; body length: 8.0. Head. Labrum and mandible brown. Clypeus, epistomal angle obtuse; punctate, weakly microreticulate; white setae on middle, longer setae on apex; yellow stripe on apical margin, green. Supraclypeal area sparsely punctate, microreticulate; green. Paraocular area sparsely punctate, microreticulate; green; long (>2 OD) branched setae, yellow. Antennae F2=F1, remaining flagellomeres relatively longer; scape and pedicel light brown, flagellomeres yellowed ventrally; F11 not flattened; F3–11 with sensilla placodea patches. Frons not depressed above antennal sockets; coarsely punctate; green with bluish iridescence; few erect setae. Preoccipital ridge carinate, some portions lamellate. Vertex punctate becoming rugose near ridge; very long plumose (short branches) setae. Gena puncticulate weakly lineate on postgena; green, copper reflections on postgena; white plumose setae (1.5–2 OD). Hypostomal area, ridge carinate. *Mesosoma*. Pronotum dorsal ridge carinate; lateral angle almost orthogonal; lateral ridge rounded; dorsal surface rugose; green; with some plumose setae on dorsal surface. Fore leg coxa to femur with green reflections, tibia and tarsus brown; short yellow erect setae. Mesoscutum anterior border slightly projected over pronotum, weakly lobed; median sulcus well impressed; parapsidal lines well impressed; punctate, sparsely median surface, microreticulations only on lateral surface; green; posterior margin weakly covered with plumose setae. Scutellum median sulcus weakly impressed; punctate, rugulose on posterior margin, polished; green, blue iridescence posteriorly, weak copper reflections on middle surface; anterior surface covered with plumose setae. Mesepisternum punctate, densely microreticulate; green with some copper reflections; short plumose setae. Mid leg coxa brown with metallic reflections, remaining leg brown; short yellow erect setae. Tegula concentric lineate; brown, green reflections anteriorly. Forewing weakly infuscate on apex. Metanotum median sulcus weakly impressed; puncticulate, polished; bright blue green; anterior surface densely covered with erect setae. Metepisternum lineate above superior metapleural pit, rugulose–punctate below; blue green. Hind leg coxa with metallic reflections, remaining leg brown; short yellow erect setae, shorter than diameter of sclerites. Metapostnotum weakly inclined; weakly depressed; 1.5× as long as metanotum; apex rounded; carinate to irregularly carinate, carinae not reaching posterior margin, microreticulate; blue green. Propodeum posterior surface posterior pit shallow, densely punctate, microreticulate, blue green, with fine setae; lateral surface punctate, blue green, with thicker setae. *Metasoma*. T1 anterior surface not punctate, dorsal surface punctate, polished; blue green, dark marginal area short. T2–5 punctate, microreticulate; dark blue green, marginal area black; vestiture white, with weak row of long white setae on basis. T6 green; long setae. T7 black; short black setae. S1 margin straight; sparsely punctate, microlineate; light brown; with very short setae.

S2–3 straight; brown; covered with short white setae on median surface, posterior surface with long setae (>4 OD). S4 apical margin straight; without median patch of setae; lateral setae short. S5 as S2. S6 weakly concave; brown; black setae on apical margin. S7 with median process. S8 apical margin bilobed. **Genitalia.** Gonapophysis broad; longer than gonostylus; without dorsal crest; with large ventral prong; with lateral teeth. Gonobase short (0.6 times gonocoxite length); dorsal lobes weakly produced; ventral bridge narrow. Gonostylus basal process absent; parapenial lobe present, short; ventral process with outer and inner lobes, inner lobe transversal setose on apex; dorsal lobe transversal. Volsella longer than wider; oblique, corners mostly rounded; without hook.

Variation. Most examined material is blue green but some specimens from Paraná and Espírito Santo are green. The blue reflections vary among specimens, including some violet reflections. The mesoscutum punctuation is sparser in some specimens.

Type material. Holotype female (DZUP), “PEVV\ 0439” “Brasil, Paraná, Parque\ Estadual de Vila Velha,\ 25°14S 49°59'W,\ 12.XII.2003, G.A.R. Melo &\ R. B.Gonçalves”. Paratypes (DZUP), one female, “PEVV\ 1056” “Brasil, Paraná, Parque\ Estadual de Vila Velha,\ 25° 14S 49°59'W,\ 7.iii.2003, G. Melo &\ R. Gonçalves”, one female, “PEVV\ 0072” “Brasil, Paraná, Parque\ Estadual de Vila Velha,\ 25° 14S 49°59'W,\ 05.x.2003, G. Melo &\ R. Gonçalves & A. Aguiar”, two females “Brasil, Paraná, Parque\ Estadual de Vila Velha,\ 25°14S 49°59'W,\ 16.iii.2002, G. Melo & R. B.\ Gonçalves”, one male “PEVV\ 1161” “Brasil, Paraná, Parque\ Estadual de Vila Velha,\ 25°14S 49°59'W,\ 7.iii.2003, G. Melo &\ R. Gonçalves”.

Additional examined material. BRAZIL, Espírito Santo: one female (DZUP), “SANTA TERESA – ES\ BRASIL 22/X/966\ C. and T. Elias leg.”. Minas Gerais: four females (UFV), “Local MEUFV. MG\ Data 19/02/85\ Col. L. Campos”; one female (UFV) with same data except “18/12/85\ A. A. Soares”; one female (UFV), “MEUFV MG\ 11/10/1987\ A. A. Soares”. Paraná: one female (DZUP), “ANTONINA PR\ Brasil 20-I-1966\ Marinoni [and] Azevedo”. Rio de Janeiro: one female (USNM), “RdJaneiro\ Brazil\ X 1938”“YelFevServ\ MESBrazil\ RCShannon”. Santa Catarina: one female (SEMC), “BRAZIL\ Santa Catarina\ Nova Teutonia\ 15 Sept. 1955\ (F. Plaumann)”; two females (SEMC), “Nova Teutonia,\ Santa Catarina,\ Brazil IX-1951\ L. E. Plaumann”. São Paulo: two females (MZSP), “Brasil, São Paulo, Iporanga, 10.ix.2000, B. H. Dietz col.”, one female (DZUP), “Brasil, São Paulo\ Cajuru, Faz. Rio\ Grande, 21°12S\ 47°09'W 02.xii.1999,\ Gabriel A. R. Melo”.

Etymology. From the Latin term *inflexus* means bend, curve, in reference to the form of the inner lobe of ventral process of gonostylus.

Discussion

This study corroborates the monophyly of the *Augochlora* clade recovered by previous studies (Eickwort, 1969; Danforth and Eickwort, 1997; Engel, 2000; Coelho, 2004; Gonçalves, 2015, 2016). The topology of Gonçalves (2016) is corroborated in contrast to previous hypothesis (Eickwort, 1969; Danforth and Eickwort, 1997; Engel, 2000; Coelho, 2004; Gonçalves, 2015) (Fig. 1). In spite of the peculiar genital capsule, *Ceratalictus inflexus* sp. nov. is sister lineage to the rest of the species of the genus. *Ceratalictus* is supported by seven synapomorphies. The phylogenetic relationships among the remaining *Ceratalictus* species were the same as found by Gonçalves (2015). *Ceratalictus* distribution shows biogeographical duplications in southeastern South America (Gonçalves, 2015) and the basal position of the new species reinforces the hypothesis of the origin of the genus in this region.

The topology found for *Augochlorella* is different from that of Coelho (2004) due the placement of *A. comis*. This species is considered here sister to the rest of the genus, in contrast to the previous study where it was considered as related to the “aurata group”. The

male F2 shorter than F1 and the concave S4 are synapomorphies for *Augochlorella* excluding *A. comis*. The convex face and the almost obtuse epistomal angle of *A. comis* are also plesiomorphic features from other *Augochlorella* species and support this new phylogenetic hypothesis. Coelho (2004) postulate a vicariance between *A. ephyra* and *A. aurata* (including *A. comis*) groups. In the present study, the hypothesis point to a different interpretation with the origin of *Augochlorella* in Central America and only later occupying North and South America. More effort is necessary to understand the biogeographical evolution of *Augochlorella*. This study described four new species of the *Augochlora* clade and provided a revised phylogeny for the placement of these species. This was the first attempt to include all species of *Augochlorella* and *Ceratalictus* in a phylogenetic analysis. There is still a lot of systematic work to be done with the clade: *Pereirapis* has never been studied in detail; *Augochlora* taxonomy and systematics is still very poorly investigated due to large number of undescribed species; and the relationships of *Augochlorella* species remains with unresolved nodes.

Funding

This study was partially supported by Brazilian *Conselho Nacional de Desenvolvimento Científico e Tecnológico*, number 443069/2014-0.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

Thanks to Carlos Roberto Brandão and Gabriel Melo for assistance during the execution of this study. I am grateful to five anonymous reviewers for suggestions on the manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.rbe.2018.12.003](https://doi.org/10.1016/j.rbe.2018.12.003).

References

- Coelho, B.W.T., Gonçalves, R.B., 2010. *A taxonomic revision of the augochlorine bee genus Ceratalictus Moure (Hymenoptera, Apoidea)*. Zootaxa 2675, 1–25.
- Dalmazzo, M., Roig-Alsina, A., 2012. Nest structures and notes on social behavior of *Augochlora amphitrite* (Schrottky) (Hymenoptera: Halictidae). J. Hym. Res. 2, 17–29.
- Dalmazzo, M., Roig-Alsina, A., 2015. Social biology of *Augochlora* (*Augochlora*) *phoenoe* (Hymenoptera: Halictidae) reared in laboratory nests. Ins. Soc. 62, 315–323.
- Danforth, B.N., Eickwort, G.C., 1997. The evolution of social behavior in the Augochlorini sweat bees (Hymenoptera: Halictidae) based on a phylogenetic analysis of the genera. In: Choe C, Crespi BJ, organizers. *The Evolution of Social Behavior in Insects and Arachnids*. Cambridge, Cambridge University Press. pp. 270–292.
- Eickwort, G.C., 1969. *A comparative morphological study and generic revision of the augochlorine bees (Hymenoptera: Halictidae)*. Un. Kansas Sci. Bull. 158, 325–524.
- Eickwort, G.C., Eickwort, K.R., 1972. Aspects of the biology of Costa Rican halictine bees. IV. *Augochlora* (*Oxystoglossella*) (Hymenoptera: Halictidae). J. Kansas Entomol. Soc. 45, 18–45.
- Eickwort, G.C., Eickwort, K.R., 1973. Notes on the nests of three wood-dwelling species of *Augochlora* from Costa Rica (Hymenoptera: Halictidae). J. Kansas Entomol. Soc. 46, 17–22.
- Engel, M.S., 2000. Classification of the bee tribe Augochlorini (Hymenoptera: Halictidae). Bull. Am. Mus. Nat. Hist. 250, 1–90.
- Goloboff, P.A., Farris, J.S., Nixon, K.C., 2008. TNT, a free program for phylogenetic analysis. Cladistics 24, 774–786.
- Gonçalves, R.B., 2015. Phylogeny and biogeography of *Ceratalictus* Moure (Apoidea, Augochlorini). In: Aguiar, A.J.C., Gonçalves, R.B., Ramos, K.S., organizers. *Ensaios sobre as abelhas da região neotropical: homenagem aos 80 anos de Danuncia Urban*. Curitiba: Editora UFPR; 221–233.
- Gonçalves, R.B., 2016. A molecular and morphological phylogeny of the extant Augochlorini (Hymenoptera, Apoidea) with comments on implications for biogeography. Syst. Entomol. 41, 430–440.
- Harris, R., 1979. A glossary of surface sculpturing. Occ. Pap. Entomol. 28, 1–31.
- Hedtke, S.M., Patiny, S., Danforth, B.N., 2013. The bee tree of life: a supermatrix approach to apoid phylogeny and biogeography. BMC Evol. Biol. 13, 138.
- Kawada, R., Buffington, M.L., 2016. A scalable and modular dome illumination system for scientific microphotography on a budget. PLoS One 11, e0153426.
- Meira, O.M., Gonçalves, R.B., 2018. The relevance of the mesosomal internal structures to the phylogeny of Augochlorini bees (Hymenoptera: Halictinae). Zool. Scripta (Online first). <http://dx.doi.org/10.1111/zsc.12270>.
- Michener, C.D., 2007. *The Bees of the World*, 2nd ed. Johns Hopkins University Press, Baltimore.
- Moure, J.S., 2007. Augochlorini. In: Moure JS, Urban D, Melo GAR, organizers. *Catalogue of bees (Hymenoptera, Apoidea) in the neotropical region*. Sociedade Brasileira de Entomologia, Curitiba, pp. 677–691.
- Sereno, P., 2007. Logical basis for morphological characters in phylogenetics. Cladistics 23, 565–587.
- Stockhammer, K.A., 1966. Nesting habits and life cycle of a sweat bee, *Augochlora pura* (Hymenoptera: Halictidae). J. Kansas Entomol. Soc. 39, 157–192.
- Wcislo, W.T., Gonzalez, V.H., Engel, M.S., 2003. Nesting and social behavior of a wood-dwelling Neotropical bee, *Augochlora isthmii* (Schwarz), and notes on a new species, *A. alexanderi* Engel (Hymenoptera: Halictidae). J. Kansas Entomol. Soc. 76, 588–602.