

Cloning and preliminary verification of telomere-associated sequences in upland cotton

Yuling Liu¹, Zhen Liu¹, Yangyang Wei¹, Yanjun Wang¹, Jiaran Shuang¹, Renhai Peng¹

¹ Anyang Institute of Technology, Anyang, Henan, 455000, China

Corresponding author: Renhai Peng (aydxprh@163.com)

Academic editor: V. Shneyer | Received 14 December 2019 | Accepted 4 March 2020 | Published 6 April 2020

<http://zoobank.org/632FF29E-E61D-4AB1-89D3-172B79BB09EE>

Citation: Liu Y, Liu Z, Wei Y, Wang Y, Shuang J, Peng R (2020) Cloning and preliminary verification of telomere-associated sequences in upland cotton. *Comparative Cytogenetics* 14(2): 183–195. <https://doi.org/10.3897/CompCytogen.v14i2.49391>

Abstract

Telomeres are structures enriched in repetitive sequences at the end of chromosomes. In this study, using the telomere primer AA(CCCTAAA)₃CCC for the single primer PCR, two DNA sequences were obtained from *Gossypium hirsutum* (Linnaeus, 1753) accession (acc.) TM-1. Sequence analysis showed that the two obtained sequences were all rich in A/T base, which was consistent with the characteristic of the telomere-associated sequence (TAS). They were designated as GhTAS1 and GhTAS2 respectively. GhTAS1 is 489 bp long, with 57.6% of A/T, and GhTAS2 is 539 bp long, with 63.9% of A/T. Fluorescence *in situ* hybridization results showed that both of the cloned TASs were located at the ends of the partial chromosomes of *G. hirsutum*, with the strong signals, which further confirmed that GhTAS1 and GhTAS2 were telomere-associated sequences including highly tandemly repetitive sequences. Results of blast against the assembled genome of *G. hirsutum* showed that GhTAS sequences may be missed on some assembled chromosomes. The results provide important evidence for the evaluation of the integrity of assembled chromosome end sequences, and will also contribute to the further perfection of the draft genomes of cotton.

Keywords

G. hirsutum, telomere-associated sequence, cloning, FISH

Introduction

Telomeres are DNA-protein complexes at the ends of chromosomes (Blackburn 1991). Telomere structures are highly conserved, and vary surprisingly little between organisms (Richards and Ausubel 1988, Ganai et al. 1991, Fajkus et al. 2005, Watson and

Riha 2010). In humans, telomere repeated sequences are composed of conserved a minisatellite sequence unit 5'-TTAGGG-3' (Moyzis et al. 1988), whereas in *Tetrahymena* (Furgason, 1940) each chromosome end has a conserved 5'-TTGGGG-3' telomere repeat unit (Blackburn and Gall 1978). The first plant telomere DNA sequence, 5'-TTTAGGG-3' tandem repeat, was isolated from *Arabidopsis thaliana* (Linnaeus, 1753) (Richards and Ausubel 1988). Subsequent studies have demonstrated that the *Arabidopsis*-type telomeres presented in most plants (Fajkus et al. 2005, Ling et al. 2012, Schrupfová et al. 2019). At the same time, other studies have shown that some plants lacked typical telomere tandem repeat 5'-TTTAGGG-3', which sheds more light on telomere function and how telomeres responded to genetic change (Adams et al. 2001, Sýkorová et al. 2003a, Peška et al. 2015).

Telomere tandem repeats located at the end of chromosomes represent only a part of the end of chromosomes. Telomere-associated sequences (TASs) located directly proximal to telomere tandem repeats (Li et al. 2009) play an important role in telomere maintenance and chromosome stability through epigenetic modification or recombination (Cross et al. 1990, Zhong et al. 1998, Sýkorová et al. 2003b, Tran et al. 2015). In addition, TAS is also a good marker at the end of the genetic linkage map. Three TASs cloned from rice showed high polymorphism at the ends of chromosome arms of different rice varieties based on the results of genetic mapping (Ashikawa et al. 1994). Despite functional importance, the nucleotide sequences in the subtelomere region have not been fully resolved in many sequenced genomes (Lese et al. 1999, Mefford and Trask 2002, Mizuno et al. 2006). So, more work is needed to reveal the structure and function of the subtelomeres.

At present, there is relatively little research on cotton telomere. Combining FISH using the *Arabidopsis*-type telomere sequence amplified from *Arabidopsis* genomic DNA and BAL-31 digestion, Ling et al. (2012) published the first study on cotton telomeres, which proved the *Arabidopsis*-type telomere sequence existed in the cotton genome. *G. hirsutum* is the most important cultivated cotton species. So far, different versions of the genome sequence have been released (Li et al. 2015, Zhang et al. 2015, Wang et al. 2019, Hu et al. 2019), however, high content of repetitive sequences affects the quality of genome assembly (Sýkorová et al. 2013, Liu et al. 2016). TAS occupies a large proportion in subtelomere tandem repeats regions. Therefore, in order to improve the quality of genome assembly, nucleotide sequences in the subtelomere region need to be further analyzed.

Material and methods

Plant materials

The plant material was *G. hirsutum* acc. TM-1 (AADD)₁, which was planted in the experimental field of Anyang Institute of Technology, Henan, China. Genomic DNA was isolated from fresh young leaves using the modified CTAB method (Song

et al. 1998). Root tip material used for *G. hirsutum* chromosome preparation were harvested from the about 6-day seedlings planted in an incubator and pretreated by 25 ppm cycloheximide at 20 °C for 80 min, then fixed in methanol-acetic acid (3:1) and stored at 4 °C for 24 h. Squashes of root tips were prepared according to Liu et al. (2017).

Primers

The eight single primers of the plant telomere repeat were selected from NCBI database (<https://www.ncbi.nlm.nih.gov>) according to the previous studies for single primer PCR (Burr et al. 1992, Gong et al. 1998, Weiss-Schneeweiss et al. 2004, Liu et al. 2005). The primers sequence information is shown in Table 1.

Cloning and sequencing of telomere-associated sequences

The selected single primers of the plant telomere repeat sequence (Table 1) were amplified by single primer PCR using the genomic DNA of *G. hirsutum* as template, to find the suitable conditions for obtaining promising products and candidates for subtelomeric regions. The amplification procedure was as 95 °C for 3 min, followed by 35 cycles of 95 °C for 15 s, 55 °C/60 °C for 15 s, 72 °C for 30 s, and a final extension at 72 °C for 5 min. The amplification products were detected by 1% agarose gel electrophoresis, and the appropriate single primer and annealing temperature were selected based on the above result. Then, PCR amplification was performed using the selected single primer in a 50 µl reaction volume containing 25 µl of 2 × Phanta Max Buffer, 1 µl of Phanta Max Super-Fidelity DNA Polymerase (Vazyme), 0.8 µmol/L of the telomeric single primer, and 10 ng of genomic DNA. The objective band from PCR was recovered by gel extraction kit (SanPrep Column DNA Gel Extraction kit, Sangon Biotech) and was cloned into *Trans1-T1* competent cells by the *pEasy-Blunt* Simple Cloning Vector (TransGen Biotech) according to the manufacturer's instructions. The positive clones were selected for sequencing by Shanghai Sangon.

Table 1. Telomere primer sequence information.

Name	Taxonomic name	Reference	Sequence
TR1	<i>Oryza sativa</i> (Linnaeus, 1753)	Gong et al. 1998	(TTTAGGG) ₃
TR2	<i>Zea mays</i> (Linnaeus, 1753)	Burr et al. 1992	(TTTAGGG) ₄
TR3	<i>Othocallis siberica</i> (Linnaeus, 1753)	Weiss-Schneeweiss et al. 2004	(TTTAGGG) ₅
TR4	<i>Ginkgo biloba</i> (Linnaeus, 1771)	Liu et al. 2005	(CCCTAAA) ₃
TR5	<i>Brassica campestris</i> (Linnaeus, 1753)	Kapila et al. 1996	(CCCTAAA) ₃ CCC
TR6	<i>Othocallis siberica</i>	Weiss-Schneeweiss et al. 2004	AA (CCCTAAA) ₃ CCC
TR7	<i>Zea mays</i>	Burr et al. 1992	(CCCTAAA) ₄
TR8	<i>Othocallis siberica</i>	Weiss-Schneeweiss et al. 2004	(CCCTAAA) ₅

Software and websites for sequences analysis

DNAMAN software was used for extraction and alignment of cloned sequences. Repetitive sequence analysis was performed using the online program CENSOR (<https://www.girinst.org/censor/index.php>). BLAST algorithm blastn (<https://www.cottongen.org/blast>) was used to identify GhTAS from *G. hirsutum* genome database (*Gossypium hirsutum* ZJU v2.1, a1) (Hu et al. 2019). All the above analyses were performed according to the default parameters.

FISH validation

The TAS plasmid DNA was extracted using the TIANprep Mini Plasmid Kit according to the instructions. Then, TAS plasmid DNA was labeled with DIG-Nick Translation Mix (Roche). The 45S rDNA probes derived from *Arabidopsis thaliana* (Gan et al. 2013) were labeled with biotin-Nick Translation Mix (Roche) according to the instructions of the manufacturer. Chromosome preparation and FISH were performed according to the previous methods (Liu et al. 2017).

Results

Optimization of the single primer PCR

According to the melting temperature (T_m) value distribution of the eight candidate single primers (55 °C–62 °C), two annealing temperatures were selected, namely 55 °C and 60 °C. The results of PCR amplification showed that an obvious band of roughly 500 bp was amplified using the single primer TR6 under the two annealing temperatures, especially, the band amplified under annealing temperature of 60 °C showed better specificity and higher brightness (Fig. 1B-6). So, the primer TR6 (AA (CCCTAAA)₃CCC) was chosen for the following PCR amplification.

Cloning of TAS

A single band with a size of roughly 500 bp was amplified using the single primer TR6 under the annealing temperature of 60 °C with Phanta Max Super-Fidelity DNA Polymerase (Fig. 2A-2). After transformation, eight positive clones were obtained after a positive test from transformed clones (Fig. 2B). Then, the eight positive clones were sequenced.




Figure 1. Amplification results of candidate single primers. **M** Marker **A** and **B** the annealing temperature is 55 and 60 °C respectively **1–8** primers TR1–TR8.




Figure 2. Results of cloning and positive test. **A** PCR amplification results **M** Marker **1** Common Taq enzyme **2** High-fidelity enzyme. **B** Positive test of bacterial colony PCR **M** Marker **1–12** the candidate clones.

Sequences analysis

Sequence component analysis

Sequence analysis of the eight positive clones revealed that all clones had the same forward and inverted telomere primer sequence at the two ends. Sequence alignment showed that there were two different internal sequences in eight sequences. So, the two different cloned DNA sequences with different size of 488 bp and 538 bp were selected and named as GhTAS1 and GhTAS2 (Fig. 3). Their sequences had been uploaded to GenBank (accession No. MT078976 and MT078977).

>GhTAS1

AACCCTAAACCCTAAACCCTAAACCCCAAGCACCAAGCCCCAAGCCCTAAACCTCTAACCTCAGAACCTAAACCCCAA
 ACCTAATACCTTAAACCCTAAACCCTAAAAATCCAAAACCTAAATATACATGAAATTGGATATTGATAGAAATGAGATATCC
 CCACCTAGGATATCCAAAATCGGGATA TGCCCGAACCGGGATA TCCATGGATTAGGATCTCTCGACCGGGGATAAGAGAG
 GTAGGTTTCCCGGTT CAGAGACATAAATTA TCTGAACCGGTCACGAAACTATCCAACCTGCAGCTAAGATTCGGACCGG
 GACAGGGTATGAATGGGATATCCCTGAATTGCTTATGCCAAGCGGGAATAGGAATCGAAAATCCCTGAATTGTTAGGGT
 TTGGGTTTAGAGTTTAGGTTTAAGGTTT TAGGGTTTAAAGTTTAGGATTATGGTTTAGGGTTTAGGGTTTAGGGTTAG
 GGTTTAGGGT

>GhTAS2

AACCCTAAACCCTAAACCCTAAACCCAGGGATATCCCAATTCACCGTTAACCCATAGTGGTATATCCGGCCTGGATATAT
 CACATTCTAGGAATATCCCAATTCGCCACTTTTTTTCTCTAATTCCTATACCTTCTTTGTCTTAATTTCTCAAAAAAAA
 AAGTAATTTAAAGAAGGTATTAATTACCATAAATGTTACCGTTGCATATGGGATTAGAAAATTCCTCTCTCCATTTCAT
 AGCTTCTCGTTTCTCACCTATTTCATCTACCTCTTAAATTTATTTTCTTTTATTTCCCTTTTATTTCTGTGTGTTTTT
 CATTCTTGAATCTCTGTAACTCTCTTTTGTCTTTTCTTCTTTTCCCTTCTTCTTCTGTCCTTCGTAAGAAACCTAGA
 CACTTCCACGCCATCAACACCTTCGGCGCTTACCTCCCAATTATCGAAATTTGAATCTCAAAATCTATCGCTTCTATCCA
 TACCACCCATTTAGGGTTTAGGGTTTAGGGTTTAGGGTT

Figure 3. Sequences of the two TASs. The grey shadows are reverse complementary sequences of the telomere primer TR6.

The two sequences were rich in A/T bases, that is, 57.6% and 63.9% respectively. Repeat masking analysis indicated that the tandem repeats content were 31.35% in GhTAS1 and 42.38% in GhTAS2, which mainly consisted of satellite DNA and transposable elements. The above results are consistent with the typical characteristics of telomere-associated sequences (Li et al. 2009).

Homology analysis of GhTASs

Sequence alignment results of DNAMAN shown that GhTAS1 and GhTAS2 had low homology, with the sequence similarity of 38.90%, which may be due to their different chromosomal sources.

After comprehensive comparison of the obtained TASs of *G. hirsutum* and the TASs of *Arabidopsis thaliana*, *Glycine max* (Linnaeus, 1753), *Oryza sativa* (Linnaeus, 1753), *Zea mays* (Linnaeus, 1753), *Larix gmelinii* (Ruprecht, 1920) listed on NCBI, it was found that their similarity was low, ranging from 25% to 50% (Table 2). All these indicated that the cloned telomere-associated sequences had obvious species specificity.

BLAST of GhTAS1 and GhTAS2 against *G. hirsutum* genome

GhTAS1 and GhTAS2 were found using blastn with the latest *G. hirsutum* genome sequence (*Gossypium hirsutum* ZJU v2.1, a1) in Cottongen (<https://www.cottongen.org/>).

Table 2. Similarity of telomere-associated sequences between *G. hirsutum* and other plants.

Species	NCBI accession No.	TASs of <i>G. hirsutum</i>	
		GhTAS1	GhTAS2
<i>Arabidopsis thaliana</i>	AC074298.1	39.60%	36.71%
	AM177016.1	14.08%	12.94%
	AM177019.1	13.52%	13.93%
	AM177060.1	10.88%	10.15%
<i>Glycine max</i>	AF041999.1	20.24%	16.79%
<i>Oryza sativa</i>	U12056.1	28.71%	25.27%
<i>Zea mays</i>	S46927.1	48.70%	41.93%
<i>Larix gmelinii</i>	EF474441.1	31.40%	30.57%

Table 3. Partial blast results of GhTAS1 and GhTAS2 in the *G. hirsutum* genome.

Sequence name	Genomic location	Query matches	Hit matches	Identity (%)
GhTAS1	D06	1–488	65407147–65406660	98.98%
	D03	184–281	26586–26683	78.57%
	A01	171–237	118151185–118151119	82.09%
	D02	138–219	69751633–69751551	79.52%
	Scaffold515-obj	184–281	9914–9817	75.51%
GhTAS2	D01	184–281	64676574–64676477	75.51%
	A06	14–537	126445179–126444656	99.62%
	D11	14–537	71336660–71336138	98.47%
	A13	14–535	47688–48202	94.08%
	A02	25–512	40084–39589	88.15%
	D02	25–512	69751559–69752073	86.68%
	A12	25–512	30186–29672	86.15%
	Scaffold546-obj	46–455	8556–8146	89.07%
	Scaffold515-obj	25–271	31264–31514	89.33%
	A09	25–315	83200103–83200398	86.96%
	A11	25–271	121355653–121355904	88.54%
	scaffold407_obj_A03	59–271	36503–36719	92.24%
	A07	25–271	96580716–96580969	88.24%
	D10	278–455	66830830–66831007	93.26
	A10	25–271	115081227–115081476	87.75%
	A05	285–455	39831434–39831267	93.60%
	A01	278–455	118169784–118169962	91.06%
D08	278–512	69075939–69076196	84.11%	
D03	278–455	23313–23139	91.01%	
D09	278–442	51987281–51987445	91.52%	

Results showed that GhTAS1 was mapped onto five chromosomes and one scaffold of *G. hirsutum*, and GhTAS2 was mapped onto all 26 chromosomes and 14 scaffolds of *G. hirsutum* with different E-value. The partial blast results with lower E-value were listed in Table 3. GhTAS1 was localized at one end of the chromosome D06, with a higher similarity of 98.98%, and was localized at the single end of chromosomes D03, A01, D02 and D01, as well as Scaffold515, with lower similarity (Fig. 4A). GhTAS2 showed




Figure 4. Localization patterns of GhTAS1 and GhTAS2 on *G. hirsutum* partial chromosomes. **A** GhTAS1 **B** GhTAS2.




Figure 5. FISH on *G. hirsutum* chromosome with 45S rDNA (green) and GhTAS1 or GhTAS2 (red) probes, Scale bars: 5 μ m. The white arrows showed co-location **A** GhTAS1 **B** GhTAS2.

higher chromosomes coverage than GhTAS1. Among the all blast results, GhTAS2 was localized at both ends of chromosomes D11, A13, A02 and D02 and at the single end of chromosomes A06, A12 and two scaffolds with higher similarity (Fig. 4B). At the same time, unlike GhTAS1, the GhTAS2 sequence is also mapped to other chromosomal regions in addition to the ends of chromosomes (Fig. 4B1–6).

Chromosome localization of GhTAS1 and GhTAS2 based on FISH

To examine the chromosome physical location of GhTAS1 and GhTAS2, we carried out FISH on *G. hirsutum* metaphase chromosomes using a digoxin-labeled GhTAS probe and a biontin-labeled 45S rDNA probe. The results showed that GhTAS1 had signals at the end of nearly half of the chromosomes of *G. hirsutum*, and most of them were distributed at the single end. The signal intensity on different chromosomes was also different (Fig. 5A-2, A-4). GhTAS2 has signals on both ends of most chromosomes of *G. hirsutum* (Fig. 5B-2, B-4). Three pairs of 45S rDNA signals were detected on the chromosomes of *G. hirsutum* (Fig. 5a-3 and 5B-3 arrows). Two pairs of GhTAS1 signals were collinear with 45S rDNA (Fig. 5A-2 arrows). Three pairs of GhTAS2 signals were collinear with 45S rDNA (Fig. 5B-2 arrows). In addition, the chromosomes carrying GhTAS2 FISH signals were much more than those with GhTAS1 FISH signals (Fig. 5A-2, B-2), which was similar to the blast results (Fig. 4, Table 3).

Discussion

In this study, the telomere primer AA(CCCTAAA)₃CCC was used as a single primer to obtain the TAS sequences of *G. hirsutum* by single primer PCR. The homology of the two TASs is relatively low and with the similarity of 38.90%. Chromosome FISH localization of the two sequences also showed obvious differences in chromosome distribution and signal strength (Fig. 5A, B), which may be due to the differences of chromosome specificity and sequence copy number of the two TASs. In the early study of *Chironomus palidivittatus* (Edwards, 1929) TAS, it was found that there were considerable differences in TAS between species, within species, and even in telomere of the same species (Cohn and Edstrom 1992). Gong et al. cloned six TASs in rice and found high polymorphism of these sequences through RFLP analysis (Gong et al. 1998). From then on, this phenomenon has been found in related studies of other species (Li et al. 2009). Therefore, TASs show great specificity, unlike the more conservative telomere repeated sequences (TR).

Since telomere and adjacent subtelomere regions could not be covered by PAC and BAC clones, sequencing efforts were unable to reveal the structure of these regions. In addition, the discovery of interstitial telomeric sequences (ITSs) makes telomeric minisatellites have double-faced character, which causes more problems in producing genomic assemblies (Richards et al. 1991, Sýkorová et al. 2003). Therefore, nucleotide sequences in the subtelomere regions have not been fully resolved in many genomes that have been sequenced (Mefford and Trask 2002, Mizuno et al. 2006), which greatly affects the quality of genome assembly. FISH localization can reflect the true position of DNA fragments in chromosomes (Jiang and Gill 2006). FISH combined with genomic BLAST can intuitively judge the genomic assembly quality of DNA sequences. Chromosomal locations of 45S rDNA in *G. hirsutum* had been revealed using double-probe FISH, that is, chromosomes A09, D07 and D09 (Gan et al. 2013). In this study, according to the genome BLAST and chromosome FISH localization results of GhTAS and 45S rDNA, it was found that TASs at the end of some chromosomes were not assembled in the genome sequence map. Obviously, results of blastn showed that GhTAS1 was only mapped onto chromosomes D06, D03, A01, D02 and D01 (Table 3, Fig. 4A), but FISH showed more chromosomes carried GhTAS1 signals, including two of the three chromosomes with 45S rDNA A09, D07 or D09, which had not appeared on the blastn results. That is, GhTAS1 sequences may be missed on these assembled chromosomes. The results provide important evidence for the evaluation of the integrity of assembled chromosome end sequences.

Conclusions

We cloned two telomere-associated sequences from *G. hirsutum* acc. TM-1 using the single-primer PCR, and made analysis about the sequence characteristics of two TASs. The two TASs sequences were enriched in A/T, and were flanked by the forward and in-

verted primer sequences at each end respectively. By comparative analysis based on the results of blastn and FISH localization of the two TASs, we found that TASs at the end of some chromosomes were not assembled in the genome sequence map. Our study not only contributes to the analysis of telomere structure of cotton, but also provides intuitive evidence for the evaluation of the integrity of the assembled *G. hirsutum* genome.

Acknowledgements

The research was sponsored by National Key R&D Program of China (No. 2018YFD0100300), Innovation Scientists and Technicians Troop Construction Projects of Henan Province (20IRTSTHN021), Science and Technology Development Program of Anyang City (2018-66-133), Science and Technology Development Project of Henan Province (182102410041).

References

- Adams SP, Hartman TPV, Lim KY, Chase MW, Bennett MD, Leitch IJ, Leitch AR (2001) Loss and recovery of *Arabidopsis*-type telomere repeat sequences 5'-(TTTAGGG)_n-3' in the evolution of a major radiation of flowering plants. *Proceedings of the Royal Society B: Biological Sciences* 268(1476): 1541–1546. <https://doi.org/10.1098/rspb.2001.1726>
- Ashikawa I, Kurata N, Nagamura Y, Minobe Y (1994) Cloning and mapping of telomere-associated sequences from rice. *DNA Research* 1: 67–76. <https://doi.org/10.1093/dnares/1.2.67>
- Blackburn EH (1991) Structure and function of telomeres. *Nature* 350: 569–573. <https://doi.org/10.1038/350569a0>
- Blackburn EH, Gall JG (1978) A tandemly repeated sequence at the termini of the extrachromosomal ribosomal RNA genes in *Tetrahymena*. *Journal of Molecular Biology* 120(1): 33–53. [https://doi.org/10.1016/0022-2836\(78\)90294-2](https://doi.org/10.1016/0022-2836(78)90294-2)
- Burr B, Burr FA, Matz EC, Romeroseverson J (1992) Pinning down loose ends: mapping telomeres and factors affecting their length. *Plant Cell* 4: 953–960. <https://doi.org/10.1105/tpc.4.8.953>
- Cohn M, Edstrom JE (1992) Chromosome ends in *Chironomus pallidivittatus* contain different subfamilies of telomere-associated sequences. *Chromosoma* 101(10): 634–640. <https://doi.org/10.1007/BF00360541>
- Cross S, Lindsey J, Fantes J, McKay S, McGill N, Cooke H (1990) The structure of a subterminal repeated sequence present on many human chromosomes. *Nucleic Acids Research* 18(22): 6649–6657. <https://doi.org/10.1093/nar/18.22.6649>
- Fajkus J, Sýkorová E, Leitch AR (2005) Telomeres in evolution and evolution of telomeres. *Chromosome Research* 13: 469–479. <https://doi.org/10.1007/s10577-005-0997-2>
- Gan YM, Liu F, Chen D, Wu Q, Qin Q, Wang CY, Li SH, Zhang XD, Wang YH, Wang KB (2013) Chromosomal locations of 5S and 45S rDNA in *Gossypium* genus and its phylogenetic implications revealed by FISH. *PLoS ONE* 8(6): e68207. <https://doi.org/10.1371/journal.pone.0068207>

- Ganal MW, Lapitan NL, Tanksley SD (1991) Macrostructure of the tomato telomeres. *Plant Cell* 3: 87–94. <https://doi.org/10.1105/tpc.3.1.87>
- Gong XQ, Gong JM, Liu F, Chen SY (1998) Screening and localization of BAC clones containing telomere-associated sequences in rice. *Science in China (Series C)* 28: 437–443. [In Chinese]
- Hu Y, Chen J, Fang L, Zhang Z, Ma W, Niu Y et al. (2019) *Gossypium barbadense* and *Gossypium hirsutum* genomes provide insights into the origin and evolution of allotetraploid cotton. *Nature Genetics* 51(4): 739–748. <https://doi.org/10.1038/s41588-019-0371-5>
- Jiang JM, Gill BS (2006) Current status and the future of fluorescence in situ hybridization (FISH) in plant genome research. *Genome* 49(9): 1057–1068. <https://doi.org/10.1139/g06-076>
- Kapila R, Das S, Lakshmikumaran M, Srivastava PS (1996) A novel species-specific tandem repeat DNA family from *Sinapis arvensis*: detection of telomere-like sequences. *Genome* 39(4): 758–766. <https://doi.org/10.1139/g96-095>
- Lese CM, Fantes JA, Riethman HC, Ledbetter DH (1999) Characterization of physical gap sizes at human telomeres. *Genome Research* 9: 888–894. <https://doi.org/10.1101/gr.9.9.888>
- Li F, Fan G, Lu C, Xiao G, Zou C, Kohel RJ et al. (2015) Genome sequence of cultivated Upland cotton (*Gossypium hirsutum* TM-1) provides insights into genome evolution. *Nature Biotechnology* 33: 524–530. <https://doi.org/10.1038/nbt.3208>
- Li J, Yang F, Zhu J, He SB, Li LJ (2009) Characterization of a tandemly repeated subtelomeric sequence with inverted telomere repeats in maize. *Genome* 52(3): 286–293. <https://doi.org/10.1139/G09-005>
- Ling J, Cheng H, Liu F, Song GL, Wang CY, Li SH, Zhang XD, Wang YH, Wang KB (2012) The cloning and fluorescence in situ hybridization analysis of cotton telomere Sequence. *Journal of Integrative Agriculture* 11(9): 1417–1423. [https://doi.org/10.1016/S2095-3119\(12\)60141-5](https://doi.org/10.1016/S2095-3119(12)60141-5)
- Liu D, Lu H, Ji FT, Li FL, Guo HH (2005) Cloning and Analysis of telomere-associated Sequences of *Ginkgo biloba* L. *Forestry Studies in China* 7(1): 7–10. <https://doi.org/10.1007/s11632-005-0049-1>
- Liu YL, Liu Z, Peng RH, Wang YH, Zhou ZL, Cai XY, Wang XX, Zhang ZM, Wang KB, Liu F (2017) Cytogenetic maps of homoeologous chromosomes Ah01 and Dh01 and their integration with the genome assembly in *Gossypium hirsutum*. *Comparative Cytogenetics* 11(2): 405–420. <https://doi.org/10.3897/CompCytogen.v11i2.12824>
- Liu YL, Peng RH, Liu F, Wang XX, Cui XL, Zhou ZL, Wang CY, Cai XY, Wang YH, Lin ZX, Wang KB (2016) A *Gossypium* BAC clone contains key repeat components distinguishing sub-genome of allotetraploidy cottons. *Molecular Cytogenetics* 9(1): 27. <https://doi.org/10.1186/s13039-016-0235-y>
- Mefford HC, Trask BJ (2002) The complex structure and dynamic evolution of human subtelomeres. *Nature Reviews Genetics* 3(2): 91–102. <https://doi.org/10.1038/nrg727>
- Mizuno H, Wu J, Kanamori H, Fujisawa M, Namiki N, Saji S, Katagiri S, Katayose Y, Sasaki T, Matsumoto T (2006) Sequencing and characterization of telomere and subtelomere regions on rice chromosomes 1S, 2S, 2L, 6L, 7S, 7L and 8S. *The Plant Journal* 46(2): 206–217. <https://doi.org/10.1111/j.1365-313X.2006.02684.x>
- Moyzis RK, Buckingham JM, Cram LS, Dani M, Deaven LL, Jones MD, Meyne J, Ratliff RL, Wu JR (1988) A highly conserved repetitive DNA sequence, (TTAGGG)_n, present at the telomeres of human chromosomes. *Proceedings of National Academy of Sciences, United States* 85(18): 6622–6626. <https://doi.org/10.1073/pnas.85.18.6622>

- Peška V, Fajkus P, Fojtová M, Dvořáčková M, Hapala J, Dvořáček V, Polanská P, Leitch AR, Sýkorová E, Fajkus J (2015) Characterisation of an unusual telomere motif (TTTTT-TAGGG)_n in the plant *Cestrum elegans* (*Solanaceae*), a species with a large genome. *The Plant Journal* 82(4): 644–654. <https://doi.org/10.1111/tpj.12839>
- Richards EJ, Ausubel FM (1988) Isolation of a higher eukaryotic telomere from *Arabidopsis thaliana*. *Cell* 53: 127–136. [https://doi.org/10.1016/0092-8674\(88\)90494-1](https://doi.org/10.1016/0092-8674(88)90494-1)
- Richards EJ, Goodman HM, Ausubel FM (1991) The centromere region of *Arabidopsis thaliana* chromosome 1 contains telomere-similar sequences. *Nucleic Acids Research* 19: 3351–3357. <https://doi.org/10.1093/nar/19.12.3351>
- Schrumpfová P, Fojtová M, Fajkus J (2019) Telomeres in plants and pumans: not so different, not so similar. *Cells* 8(1): 58. <https://doi.org/10.3390/cells8010058>
- Song GL, Cui RX, Wang KB, Guo LP, Li SH, Wang CY, Zhang XD (1998) A rapid improved CTAB method for extraction of cotton genomic DNA. *Cotton Science* 10(5): 273–275. [In Chinese]
- Sýkorová E, Cartagena J, Horakova M, Fukui K, Fajkus J (2003a) Characterization of telomere-subtelomere junctions in *Silene latifolia*. *Molecular Genetics Genomics* 269(1): 13–20. <https://doi.org/10.1007/s00438-003-0811-9>
- Sýkorová E, Fojtová M, Peška V (2013) A PCR-based approach for evaluation of telomere associated sequences and interstitial telomeric sequences. *Analytical Biochemistry* 439(1): 8–10. <https://doi.org/10.1016/j.ab.2013.03.034>
- Sýkorová E, Lim KY, Chase MW, Knapp S, Leitch IJ, Leitch AR, Fajkus J (2003b) The absence of *Arabidopsis*-type telomeres in *Cestrum* and closely related genera *Vestia* and *Sessea* (*Solanaceae*): first evidence from eudicots. *The Plant Journal* 34(3): 283–291. <https://doi.org/10.1046/j.1365-313X.2003.01731.x>
- Tran TD, Cao HX, Jovtchev G, Neumann P, Novák P, Fojtová M, Vu GTH, Macas J, Fajkus J, Schubert I, Fuchs J (2015) Centromere and telomere sequence alterations reflect the rapid genome evolution within the carnivorous plant genus *Genlisea*. *The Plant Journal* 84(46): 1087–1099. <https://doi.org/10.1111/tpj.13058>
- Wang MJ, Tu LL, Yuan DJ, Zhu D, Shen C, Li JY et al. (2019) Reference genome sequences of two cultivated allotetraploid cottons, *Gossypium hirsutum* and *Gossypium barbadense*. *Nature Genetics* 51(2): 224–229. <https://doi.org/10.1038/s41588-018-0282-x>
- Watson JM, Riha K (2010) Comparative biology of telomeres: where plants stand. *FEBS Letters* 584(17): 3752–3759. <https://doi.org/10.1016/j.febslet.2010.06.017>
- Weiss-Schneeweiss H, Riha K, Jang CG, Puizina J, Scherthan H, Schweizer D (2004) Chromosome termini of the monocot plant *Othocallis siberica* are maintained by telomerase, which specifically synthesizes vertebrate-type telomere sequences. *The Plant Journal* 37: 484–493. <https://doi.org/10.1046/j.1365-313X.2003.01974.x>
- Zhang TZ, Hu Y, Jiang WK, Fang L, Guan XY, Chen JD et al. (2015) Sequencing of allotetraploid cotton (*Gossypium hirsutum* L. acc. TM-1) provides a resource for fiber improvement. *Nature Biotechnology* 33(5): 531–537. <https://doi.org/10.1038/nbt.3207>
- Zhong XB, Franz PF, Wennekes-van Eden J, Ramanna MS, van Kammen A, Zabel P, de Jong JH (1998) FISH studies reveal the molecular and chromosomal organization of individual telomere domains in tomato. *The Plant Journal* 13(4): 507–517. <https://doi.org/10.1046/j.1365-313X.1998.00055.x>