

Mitochondrial discordance in closely related *Theridion* spiders (Araneae, Theridiidae), with description of a new species of the *T. melanurum* group

Marc Domènech¹, Luís C. Crespo¹, Alba Enguïdanos¹, Miquel A. Arnedo¹

¹ Department of Evolutionary Biology, Ecology and Environmental Sciences & Biodiversity Research Institute (IRBio), Universitat de Barcelona, Av. Diagonal 643, E-08028, Barcelona, Spain

<http://zoobank.org/401A609B-8442-4252-BA90-698ADB96FE1A>

Corresponding author: Marc Domènech (marcdomenech@ub.edu)

Academic editor: Danilo Harms ♦ Received 8 January 2020 ♦ Accepted 14 March 2020 ♦ Published 4 May 2020

Abstract

The incorporation of molecular data into current taxonomic practise has unravelled instances of incongruence among different data sets. Here we report a case of mitochondrial discordance in cobweb spiders of the genus *Theridion* Walckenaer, 1805 from the Iberian Peninsula. Morphological examination of samples from a country-wide bioinventory initiative revealed the existence of a putative new species and two nominal species belonging to the *Theridion melanurum* species group. The morphological delineation was supported by the molecular analysis of a nuclear marker but was at odds with the groups circumscribed by a mitochondrial marker. The causes of this discordance remained uncertain, once sample and sequencing errors and the existence of pseudogenes were discarded. The full sorting observed in the alleles of the more slowly evolving nuclear marker ruled out incomplete lineage sorting, while the geographic patterns recovered were difficult to reconcile with ongoing hybridization. We propose that the apparent incongruence observed is most likely the result of old introgression events in a group with high dispersal abilities. We further speculate that endosymbiont-driven cytoplasmic incompatibility could be involved in the fixation of mitochondrial haplotypes across species barriers. Additionally, we describe the new species *T. promiscuum* sp. nov., based on the presence of diagnostic morphological traits, backed up by the nuclear data delimitation. Our study contributes yet another example of the perils of relying on single methods or data sources to summarise the variation generated by multiple processes acting through thousands of years of evolution and supports the key role of biological inventories in improving our knowledge of invertebrate biodiversity.

Key Words

COI, hybridization, Iberian Peninsula, incomplete lineage sorting, introgression, ITS2, Wolbachia

Introduction

The incorporation of DNA sequence information in species delimitation and description has become a gold standard in current taxonomic practise. Under an integrative taxonomy framework (Dayrat 2005; Will et al. 2005; Padial et al. 2010) the use of DNA sequence data allows for a finer species delimitation, reveals population structure, simplifies matching life stages or remnants, and facilitates identification for taxa otherwise difficult to diagnose using only morphological charac-

ters (e.g. cryptic species) (Pires and Marinoni 2010). The comparative analyses of DNA sequences both from different markers and with phenotypic data have unravelled instances of incongruence between data sets, which may point to relevant biological processes in the origin and maintenance of biological diversity (e.g. Sota and Vogler 2001). At the same time, the reported incongruence may challenge some automated approximations to species delimitation and identification, especially when based on single markers (i.e. DNA barcoding) (Will et al. 2005).

Hybridization and incomplete lineage sorting are among the main processes accounting for the apparent incongruence between different sources of taxonomic evidence, namely mitochondrial DNA, nuclear DNA, or the phenotype. Hybridization is defined as the interbreeding of individuals from different species. When this hybridization process involves repeated backcrossing of the hybrids with the parent species, we talk about introgression. The notion that species boundaries are not impermeable but can be porous to introgression is becoming more and more accepted as new evidence appears (Harrison and Larson 2014). The most iconic case might be the recent evidence provided by ancient DNA of Neanderthal introgression with modern humans (Green et al. 2010). Multiple instances of introgression have also been reported within arthropods. Spiders have not been spared from undergoing hybridization and introgression processes. To cite some examples, several studies have revealed hybrid zones among common funnel weaver spiders of the *Eratigena atrica* (C.L. Koch, 1843) group (Agelenidae) in Britain and Continental Europe (Croucher et al. 2007; Oxford and Bolzern 2018), and in deeply divergent lineages of the ground-dweller *Harpactocrates* Simon, 1914 spiders in the Pyrenees (Bidegaray-Batista et al. 2016). In New Zealand, Vink et al. (2008) identified a case of introgression from the Australian black widow *Latrodectus hasseltii* Thorell, 1870 (Theridiidae) into the endemic *L. katipo* Powell, 1871, while Lattimore et al. (2011) found evidence of unidirectional introgression within the fishing spider genus *Dolomedes* Latreille, 1804 (Pisauridae). More recently, Leduc-Robert and Maddison (2018) have revealed that hybridization may have been widespread across phylogenetically distant species of *Habronattus* F.O. Pickard-Cambridge, 1901 jumping spiders (Salticidae).

Introgression between species can be detected by comparing information from mitochondrial (mtDNA) and nuclear DNA (nucDNA). As mtDNA is maternally inherited (Giles et al. 1980), when two individuals of different species hybridize, the offspring's mtDNA will be 100% identical to that of the maternal species, but the nucDNA will be 50% of each parental species. In successive generations, the nucDNA will homogenize through recombination within the population, while the introgressed mtDNA will remain traceable in the population in following generations, unless lost by genetic drift. The hybridization event will be detectable because some individuals will possess the nucDNA and morphology from the paternal population but the mtDNA from the maternal one.

Another source of incongruence among multiple lines of taxonomic evidence in closely related species is incomplete lineage sorting (ILS). ILS is the process by which, as a result of the segregation of an ancestral polymorphism, i.e. the existence of two or more homologous alleles predating the speciation event, the evolutionary relationships between individuals given by the sequences of a certain gene do not match the species phylogeny (Rogers and Gibbs 2014). The occurrence of ILS is especially pervasive in either recently diverged species or

in species with large ancestral effective population sizes (Maddison 1997). Under an ILS scenario, when comparing mtDNA and nucDNA from various species, we would expect mitochondrial gene trees to reflect better the species tree than nuclear gene trees, because, other factors being equal, mtDNA has a smaller effective population size in diploid organisms and is known to have a faster mutation rate than nucDNA (Ballard and Whitlock 2004).

The family Theridiidae is one of the richest and most ecologically diverse spider families, containing 2,516 species grouped in 124 genera (World Spider Catalog 2020). The family is reputed for including, among others, species of medical importance, such as the widow spiders (*Latrodectus* Walckenaer, 1805), or some of the few examples of sociability within spiders (e.g. *Anelosimus* Simon, 1891). Theridiids usually construct tangle webs with gumfoots, i.e. sticky droplets on silk threads radiating from mesh retreats or web hubs attached to the substrate, mainly aimed at capturing pedestrian prey (Blackledge et al. 2011). They are occasional to frequent ballooning, and exhibit a great diversity in morphology, ecology, and behaviour, ranging from solitary web-less hunters, such as the species of the genus *Euryopis* Menge, 1868 (Levi 1954) to elaborate tangled, three-dimensional space webs or even to large colonies of thousands of individuals, like the social spider *Anelosimus eximius* (Keyserling, 1884).

The genus *Theridion* Walckenaer, 1805 contains 584 described species, more than one-fifth of all the theridiids found worldwide (World Spider Catalog 2020). This remarkable species diversity is most likely a taxonomic artifact because the genus has been traditionally used as a dumping ground for theridiids with no trace of colulus that do not fit into better diagnosed genera (Forster et al. 1990). Because of its large size, several smaller species groups have been proposed to further divide *Theridion* (e.g. Levi 1957), some of which have been subsequently erected to genus status. One of these species' complexes, the *Theridion melanurum* species-group (Wunderlich 2011), contains 10 very similar species, often difficult to tell apart. The group contains, along *Theridion melanurum* Hahn, 1831, the species *T. asopi* Vanuytven, 2014, *T. bernardi* Lecigne, 2017, *T. betteni* Wiehle, 1960, *T. bosniense* Wunderlich, 2011, *T. cairoense* Wunderlich, 2011, *T. harmsi* Wunderlich, 2011, *T. malagaense* Wunderlich, 2011, *T. musivivum* Schmidt, 1956, *T. mystaceum* L. Koch, 1870, and *T. semitinctum* Simon 1914.

In the general framework of a biological inventory of the spider communities of white oak woodlands of the Spanish National Park Network (Crespo et al. 2018), we had the opportunity to examine a large sample of *Theridion* specimens and obtained DNA barcoding data for many of these individuals. Comparison between identifications based on morphology and those based on DNA barcoding revealed several instances of incongruence that were subsequently examined and corroborated using the nuclear intron Internal Transcribed Spacer 2 (ITS-2). Here we present the results of the morphological and molecular

analyses, describe a new species based on the combination of the available evidence, and further discuss the implications of these results for our understanding of the diversification of species-rich spider lineages.

Material and methods

Specimens were collected using semi-quantitative methods as part of a larger project that aimed to understand the diversity patterns of Iberian spider communities (Crespo et al. 2018). The sampling design included 16×1 ha plots distributed in white oak forests across six national parks of the Iberian Peninsula. Sampling was conducted using the COBRA protocol (Cardoso 2009) in May and June of 2013 and 2014. The sampling protocol combined timed direct capture, beating, and sweeping with 48 pitfall traps per plot, which were active for two weeks. The distribution of the sampling sites is shown in Figure 1. Detailed collection and genetic information of all specimens used in this work can be found in Suppl. material 1.

Specimens were sorted and identified under a ZEISS Stemi 2000 stereomicroscope. We took photographs using a Leica DFC 450 camera attached to a Leica MZ 16A stereomicroscope, with the software Leica Application Suite v. 4.4. Both the male palp and the female epigyne were excised with the help of entomological needles to facilitate observation under the scope. The muscle tissue of the epigyne was further removed with the needles and digested using immersion in potassium hydroxide (KOH) at a 30% concentration. For SEM examination, palps were excised and cleaned ultrasonically for 1 min and then transferred to 100% ethanol overnight. Palps were submitted to critical point drying, glued to flat-headed rivets and gold sputter coated. Imaging was conducted with the help of a Quanta 200 environmental SEM. We used the Araneae: Spiders of Europe online identification tool to identify most of the species found in our samples (Nentwig et al. 2019). Type specimens were deposited at Museu de Ciències Naturals de Barcelona, Spain (CMCNB) and additional voucher specimens at the Centre de Recursos de Biodiversitat Animal, University of Barcelona (CRBA). Description of the new species followed the format of Vanuytven (2014). All measurements are in millimetres.

For molecular analyses, we aimed at including both sexes and all the sites where each species was collected. We extracted DNA from two to four legs from each specimen using REDEExtract-N-Amp™ Tissue PCR Kit Protocol from Sigma-Aldrich, following the manufacturer's protocol, performed in 96-well plates. We amplified fragments of the animal DNA barcode gene cytochrome c oxidase subunit I (COI) and the nuclear Internal Transcribed Spacer 2 (ITS2). Primers used for amplification are shown in Table 1. We performed polymerase chain reactions in 96-well plates using 8 μ L REDEExtract-N-Amp™ PCR ReadyMix from Sigma-Aldrich, primers forward and reverse, 4 μ L of diluted DNA and ultrapure, distilled water up to a total reaction volume of 20 μ L. PCR condi-




Figure 1. Distribution of the sampled localities in the Iberian Peninsula: **PA** – Aigüestortes i Estany de Sant Maurici; **PC** – Cabañeros; **PM** – Monfragüe; **PO** – Ordesa y Monte Perdido; **PP** – Picos de Europa; **PS** – Sierra Nevada.

tions were as follows: initial denaturing step at 95 °C for 5 min, 35 amplification cycles (94 °C for 30 s, 45 °C for 35 s, 72 °C for 45 s) and a final step at 72 °C for 5 min. PCR products were cycle-sequenced in both directions with the sample amplification primers at MacroGen Inc. (Seoul, South Korea).

We edited and manipulated all sequences using Geneious v. 10.2 (Kearse et al. 2012). For the COI sequences alignment was trivial since there was no evidence of insertions or deletions. Conversely, fragments of the ITS2 sequences showed between and within individual length polymorphisms. Single individual length polymorphisms were manually phased by editing original chromatograms using a sliding window approach. Phased ITS2 sequences were aligned using the automatic algorithm implemented in the online version v. 7.017 of MAFFT (Katoh and Standley 2013) using the G-INS-i strategy. Gaps in the alignment were treated as separate absence/presence characters in subsequent analyses using FastGap v. 1.2 (Borchsenius 2009).

We inferred a Maximum Likelihood tree for the COI with the program RAxML (Stamatakis 2014), using 20 runs of random addition of taxa, using only unique sequences and assessing support by means of MRE bootstrapping. We used the best codon partition and corresponding model retrieved by PartitionFinder v. 2.1 (Lanfear et al. 2012) using the AICc criterion, which was the general time-reversible nucleotide substitution model with Gamma-distributed among-site rate variation and invariant sites (GTR+G+I), and with a separate partition for each codon position. Additionally, we constructed haplotype and allele networks for the COI and ITS2, respectively, using the TCS algorithm (Clement et al. 2002) as implemented in the program PopArt v. 1.7 (Leigh and Bryant 2015). We calculated intraspecific and interspecific p-distances for the COI and the ITS2 for all species using MEGA v. 7.0.26 (Kumar et al. 2016).

We conducted additional species delineation using the barcode index number (BIN) method (Ratnasingham and Hebert 2013) as implemented in the Barcode of Life Data System (BOLD, <http://www.boldsystems.org/>) (Ratnasingham and Hebert 2007).

Table 1. List of primers used in this study.

Target	Primer name	Direction	Sequence	Reference
COI	LCOI1490	Forward	GGTCAACAAATCATAAAGATATTGG	Folmer et al. 1994
COI	HCOI2198	Forward	TAAACTTCAGGGTGACCAAAAAATCA	Folmer et al. 1994
COI	Nancy	Reverse	CCCGGTAAAATTTAAATATAAACTTC	Simon et al. 1994
ITS2	ITS-5.8S	Forward	GGGACGATGAAGAACGGAGC	White et al. 1990
ITS2	ITS-28S	Reverse	TCCTCCGCTTATTGATATGC	White et al. 1990

Results

Morphological study

Among the 8521 adult spiders collected in the inventoring samples, we identified 404 specimens of the genus *Theridion*, corresponding to four nominal species, namely *T. mystaceum* (65 males, 120 females), *T. harmsi* (42 males, 138 females), *T. varians* Hahn, 1833, (14 males, 10 females), and *T. pinastris* L. Koch, 1872 (2 males). In addition, we found two morphotypes that did not fit into any of the nominal species descriptions, tentatively referred to as *T. sp. 6* (seven males, five females) and *T. sp. 15* (one male). The former was found to be a new species to science, hereby described.

Molecular analyses

We obtained the COI sequences of 117 specimens (31 *T. mystaceum*, 59 *T. harmsi*, 18 *T. varians*, two *T. pinastris*, six *T. sp. 6*, and one *T. sp. 15*) and ITS2 sequences of 73 specimens (30 *T. mystaceum*, 30 *T. harmsi*, five *T. varians*, one *T. pinastris*, six *T. sp. 6*, and one *T. sp. 15*). In addition, we downloaded seven COI sequences belonging to the *T. melanurum* group available at NCBI, namely five *T. mystaceum* (KX537283, KY268733, KY269206, KY269434, and KY270001), one *T. melanurum* (EF449609), and one *T. betteni* (KX039404), as well as one ITS2 sequence belonging to *T. varians* (KR526552) (Arnedo et al. 2007; Gregorič et al. 2015; Astrin et al. 2016; Coddington et al. 2016).

The ML tree obtained for the COI (Fig. 2) supported the close relationships between the haplotypes of the *T. melanurum* group but failed to recover monophyly of the different species. Of the four main lineages recovered, albeit some of them with low support, one included all the haplotypes of *T. harmsi*, along with single haplotypes of *T. mystaceum* and *T. sp. 6*, the second included sequences of *T. mystaceum*, *T. sp. 6*, and *T. betteni*, the third two divergent haplotypes of *T. mystaceum* and the fourth haplotypes of *T. mystaceum*, *T. sp. 6*, and *T. melanurum*. The average uncorrected genetic distances within the *T. melanurum* group was 2.8% (SE 0.004). Each of the remaining *Theridion* species sampled were recovered as monophyletic and clearly divergent from the haplotypes of the *T. melanurum* group (9% or above average uncorrected

genetic distance), including the single haplotype of the morphotype *T. sp. 15*.

The COI sequences were split into nine different BINs, one unique and the remaining including additional specimens available in BOLD. The species delineation of the *T. melanurum* complex yielded either BINs with mixed species (Fig. 2) or BINs with single specimens, in all cases identified as *T. mystaceum*. Conversely, in the remaining *Theridion* species, the BINs mirrored the morphological identifications.

The COI haplotype network (Fig. 3) was largely congruent with the ML tree but provided a more detailed view on the relationships and geographic distribution of the *T. melanurum* group haplotypes. Neither a taxonomic nor a geographic structure was readily apparent in the *T. melanurum* group network. All *T. harmsi* haplotypes were highly similar, separated by two or less mutations, and the most common haplotype was widespread all over the sampled parks. The closest haplotype to this group was found in individuals belonging to *T. mystaceum* and *T. sp. 6* from different parks. The bulk of *T. mystaceum* haplotypes were separated by five or less mutations, and also included haplotypes found in *T. sp. 6*, closely related to a haplotype of *T. mystaceum* from the same park. One of the haplotypes of this cluster was shared between two parks but also found in Germany and in a population of *T. betteni* from Britain. A second cluster of *T. mystaceum* haplotypes, separated by six or less mutations, included a haplotype shared with *T. sp. 6* from a different park, and a very closely related haplotype found in *T. melanurum* from Switzerland.

In contrast to the mitochondrial network, the nuclear ITS2 network perfectly matched the morphology-based species delimitations (Fig. 4). The alleles sampled from each species were exclusive and each other's closest relatives. The highest allele diversity was found in *T. harmsi*, whose most common alleles were found across most of the parks. The remaining species yielded two or fewer alleles.

The ITS2 maximum uncorrected intraspecific divergences ranged between 0 and 4.5% (Table 2). Minimum interspecific divergences within the species complex ranged from 2.6% (between *T. mystaceum* and *T. sp. 6*) to 4.3% (between *T. harmsi* and *T. sp. 6*), as seen in Table 3. These values fall within or above the ranges observed within and among closely related spider species across several families (see Agnarsson 2010).




Figure 4. ITS2 allele network. Labels next to the circles indicate the park or country where the specimens were collected: **A** – Aigüestortes i Estany de Sant Maurici; **C** – Cabañeros; **M** – Monfragüe; **O** – Ordesa y Monte Perdido; **P** – Picos de Europa; **S** – Sierra Nevada; **SI** – Slovenia.

Table 2. ITS2 maximum intraspecific sequence divergences in previously published studies. Species in bold correspond to species within the *T. melanurum* group.

Species	Maximum intraspecific sequence divergence	Reference
<i>Latrodectus katipo</i>	0.002	Vink et al. 2008
<i>Latrodectus hasselti</i>	0	Vink et al. 2008
<i>Latrodectus hasselti</i>	0.0027	Zhang et al. 2004
<i>Latrodectus mactans</i>	0.014	Zhang et al. 2004
<i>Anelosimus eximius</i>	0.007	Agnarsson 2010
<i>Anelosimus domingo</i>	0	Agnarsson 2010
<i>Anelosimus tosum</i>	0.008	Agnarsson 2010
<i>Anelosimus studiosus</i>	0.01	Agnarsson 2010
<i>Anelosimus guacamayos</i>	0.002	Agnarsson 2010
<i>Anelosimus octavius</i>	0.007	Agnarsson 2010
<i>Anelosimus baeza</i>	0.02	Agnarsson 2010
<i>Theridion harmsi</i>	0.045	This study
<i>Theridion mystaceum</i>	0	This study
<i>Theridion promiscuum</i> sp. nov.	0.002	This study
<i>Theridion varians</i>	0.004	This study
<i>T. melanurum</i> group	0.154	This study

Table 3. ITS2 minimum interspecific sequence divergences. Values from previously published studies are comparisons between sister species within the corresponding genus. Species in bold correspond to species within the *T. melanurum* group.

Family	Species	Minimum interspecific sequence divergence	Reference
Theridiidae	<i>Anelosimus</i> sp.	0.006	Agnarsson 2010
Theridiidae	<i>Latrodectus</i> sp.	0	Vink et al. 2008
Theridiidae	<i>Latrodectus</i> sp.	0	Zhang et al. 2004
Linyphiidae	<i>Orsonwelles</i> sp.	0.007	Hormiga et al. 2003
Araneidae	<i>Poltys</i> sp.	0.007	Smith 2006
Salticidae	<i>Havaika</i> sp.	0.02	Arnedo and Gillespie 2006
Lycosidae	<i>Pardosa</i> sp.	0.025	Chang et al. 2007
Theridiidae	<i>T. harmsi</i> – <i>T. mystaceum</i>	0.033	This study
Theridiidae	<i>T. harmsi</i> – <i>T. promiscuum</i> sp. nov.	0.043	This study
Theridiidae	<i>T. mystaceum</i> – <i>T. promiscuum</i> sp. nov.	0.026	This study

The combination of diagnostic morphological features both in males and females, similar to those found among other species within the *T. melanaurum* group, and the

additional support of the nuclear data delimitation led us to propose that *T. sp. 6* morphotype actually corresponds to a new species.

Taxonomy

Theridion promiscuum Domènech & Crespo, sp. nov.

<http://zoobank.org/794F29CB-8D96-4135-88EC-E23176914149>

Figs 5, 8

Type material. Holotype ♂: Spain, Parc Nacional d'Aigüestortes i Estany de Sant Maurici, 42.5491, 0.8714, 1739 m. Hand collecting, June 30, 2013 (Marcos Roca-Cusachs leg.). Deposited at CMCNB under the code MZB 2017-3715.

Paratypes: 6♂ and 5♀ from Spain, Parc Nacional d'Aigüestortes i Estany de Sant Maurici, 0.8725, 42.5496, 1760 m. June 29, 2013. Four males and three females deposited at CMCNB under the codes MZB 2019-1687, MZB 2017-3710, MZB 2017-3713, MZB 2017-4570, MZB 2019-1688, MZB 2017-3712, and MZB 2017-3714. Two males and two females deposited at CRBA under the codes CRBA-89454, CRBA-89455, CRBA-89456, and CRBA-89457.

Etymology. The specific name *promiscuum* is derived from the Latin word for “intermingling”, referring to the morphological similarity with other species of the *T. melanurum* group. It also makes reference to the possible introgression between this species and other closely related species here reported.

Diagnosis. Males of *T. promiscuum* sp. nov. can be distinguished from other species in *T. melanurum* group by a bent and twisted embolus, forming a marked angle (Fig. 5e); embolus shorter in *T. promiscuum* sp. nov. than in sympatric *T. mystaceum* (Fig. 6e) and *T. harmsi* (Fig. 7e); tip of the embolus pointing distally in *T. promiscuum* sp. nov. (Fig. 5c) and pointing retrolaterally in remaining species (Figs 6c, 7c); base, prolateral part of the embolus more sclerotized in *T. promiscuum* sp. nov. (Fig. 5e) than in remaining species of the *T. melanurum* group (Figs 6e, 7e); presence of small barbs in embolus base distal border is unique to *T. promiscuum* sp. nov. (only visible in SEM; Fig. 8a, b). Females of *T. promiscuum* sp. nov. can be distinguished from *T. mystaceum* and *T. harmsi* by the lower position of copulatory duct openings in atrium in *T. promiscuum* sp. nov. (Fig. 5f), which are central in *T. harmsi* (Fig. 7f) and close to or under lateral walls of atrium in *T. mystaceum* (Fig. 6f); copulatory ducts shorter, fewer coils in *T. promiscuum* sp. nov. (Fig. 5g) than in *T. mystaceum* (Fig. 6g) and *T. harmsi* (Fig. 7g).

Description. Holotype male: Total length 1.81; abdomen 1.04; cephalothorax length 0.84, width 0.78. Leg length (total, coxa, trochanter, femur, patella, tibia, metatarsus, tarsus): I: 4.46, 0.22, 0.16, 1.18, 0.42, 0.97, 1.02, 0.5; II: 3.38, 0.2, 0.11, 0.91, 0.37, 0.63, 0.75, 0.41; III: 2.49, 0.16, 0.08, 0.69, 0.3, 0.4, 0.53, 0.34; IV: 3.32, 0.21, 0.1, 0.98, 0.38, 0.56, 0.7, 0.39. Leg formula: 1>2>4>3. Eye diameter: AME 0.07; ALE 0.06; PME 0.06; PLE 0.07. Distance from AME to clypeus 0.18. Length of chelicera 0.26, fang 0.14. Dark circles around the eyes, with eye

region darker than the rest of carapace. Carapace yellowish brown, with blackish triangular patch not reaching the eyes (see paratype MZB 2017-3713; Fig. 5b). Margin of the carapace also blackish. Sternum yellowish, paler in the middle and darker in the margins, with thin black border. Legs pale with darker annulations. Dorsal abdomen with light central dented band running longitudinally and bordered with black (see paratype MZB 2017-3713; Fig. 5b). Thinner reddish band running through the middle of the light central band. Rest of the abdomen brown with darker spots. Ventral side with white spots. Spinnerets bordered by thick black line. Colulus absent. Palpal tibia pale. Embolus base wide, well sclerotized basally and prolaterally. Embolus wide, angularly bent, basally well sclerotized (see paratype MZB 2017-3713; Fig. 5e). Presence of small barbs in the distal border of the embolus base, visible in SEM (see paratype MZB 2017-3713; Fig. 8a) but very hardly visible in stereomicroscope given the membranous and translucent nature of these barbs.

Paratype female (MZB 2019-1688): Total length 2.18; abdomen 1.33; cephalothorax length 0.93, width 0.84. Leg lengths (total, coxa, trochanter, femur, patella, tibia, metatarsus, tarsus): I: 4.74, 0.24, 0.14, 1.28, 0.45, 1.01, 1.13, 0.5; II: 3.42, 0.24, 0.1, 0.94, 0.37, 0.61, 0.76, 0.41; III: 2.65, 0.22, 0.08, 0.74, 0.3, 0.39, 0.57, 0.36; IV: 3.7, 0.29, 0.1, 1.08, 0.38, 0.63, 0.82, 0.41. Leg formula: 1>2>4>3. Eye diameters: AME 0.08; ALE 0.07; PME 0.06; PLE 0.07. Distance from AME to clypeus 0.17. Length of chelicera 0.34, fang 0.13. Spine-like bristles in leg I: 2 in patella, 2 in tibia. Relative positions in tibia I: 0.25, 0.68. Thickness of tibia I: 0.1. Serrated bristles on tarsus of fourth leg. Cephalothorax yellowish brown with a black central area reaching the eye region, and broad, dark edge (Fig. 5a). Sternum yellow, paler in the middle and darker in the margins. Abdomen more spherical than in male. Spinnerets bordered by thick black line. Colulus absent. Epigynal cavity wider than long. Openings of the copulatory ducts close to the posterior margin of the epigynal cavity (Fig. 5f). Copulatory ducts relatively short and with few coils (Fig. 5g).

Intraspecific variation. Spine-like bristles in tibia of leg I of paratypes: two in patella, two in tibia. Relative positions of first spine in tibia: 0.25–0.32 (average 0.275). Relative positions of second spine in tibia: 0.7–0.74 (average 0.717). Thickness of tibia I: 0.11–0.13 (average 0.123).

Genetic information. We obtained a 568 bp sequence of the COI mitochondrial gene for the holotype (stored in Genbank under the code **MT215600**) and five paratypes (MZB 2017-3710, MZB 2017-3712, MZB 2017-3713, MZB 2017-3714, and MZB 2017-4570 with GenBank codes **MT215603**, **MT215604**, **MT215602**, **MT215606** and **MT215601**, respectively). We also obtained sequences of the ITS2 of up to 469 bp, including up to 26 bp of the 28S gene and up to 69 bp of the 5.8S gene, for the holotype (GenBank code **MT117179**) and the five paratypes mentioned before (codes **MT117182**, **MT117181**, **MT117180**, **MT117183** and **MT117184**).




Figure 5. *Theridion promiscuum* sp. nov. a) Female paratype MZB 2019-1688, dorsal habitus. b) Male paratype MZB 2017-3713, dorsal habitus. c) Left male palp, ventral. d) Left male palp, retrolateral e) Male embolus f) Female epigynum, ventral g) Female vulva, dorsal. Scale bars: 1 mm (a, b); 0.1 mm (c, d, f, g); 0.05 mm (e).

Distribution and natural history. The holotype and paratypes were collected in an oak forest of *Quercus pubescens* Willd. in Aigüestortes i Estany de Sant Maurici National Park, located in the southern slopes of the Catalan Pyrenees. The specimens were captured either by beating, sweeping or direct sampling, but not by pitfall traps, which suggests that this species is found in the vegetation at a certain height above the ground. Only the three specimens captured by beating method were captured during the day, whereas the sweeping and direct sampling ones were captured at night, which indicates that this species is mostly active at night.

Other taxa analysed. The specimen tentatively referred to as *T. sp. 15* is morphologically similar to the species *T. cinereum* Thorell, 1875, *T. petraeum* L. Koch, 1872, *T. furfuraceum* Simon, 1914, *T. pyrenaicum* Denis, 1944, and *T. wiehlei* Schenkel, 1938. However, the male palp of the single specimen available showed slight differences in the shape of the median apophysis, slightly tilted, and the more pronounced basal curvature of the shorter embolus, which refrained us from assigning it to any of the former species. The DNA barcodes, on the other hand, identified the specimen unambiguously as *T. cinereum*.




Figure 6. *Theridion mystaceum*. a) Female, dorsal habitus b) Male, dorsal habitus. c) Left male palp, ventral. d) Left male palp, retro-lateral e) Male embolus f) Female epigynum, ventral g) Female vulva, dorsal. Scale bars: 1 mm (a, b); 0.1 mm (c, d, f, g); 0.05 mm (e).

Discussion

The genus *Theridion* is already one of the largest genera within spiders, yet new species continue to be described yearly, with no evidence of reaching a plateau. Surprisingly, even the well-known European fauna is still contributing new *Theridion* species. The newest, *T. bernardi*, also belonging to the *T. melanurum* group, was described from Portugal in 2017 (Lecigne 2017). From this perspective, the discovery of yet another new species of *Theridion* from the *T. melanurum* group does not greatly improve our knowledge of this diverse yet poorly defined genus. However, the identification of a clear incongruence between some of the sources providing evidence for

the evolutionary independence of this new species presents some challenges and opens interesting evolutionary questions. This study contributes yet another example of the perils of relying on single methods or data sources to summarise the variation generated by multiple processes acting through thousands of years of evolution. Operational approaches such as DNA barcoding may greatly help to accelerate biological inventories and provide useful information for refining species delimitations and speed up identification. However, the multiple factors involved in the generation and evolution of species – e.g. speciation mechanisms, species ontogeny, gene flow – require the integration of multiple sources of evidence to provide a sound hypothesis of species.




Figure 7. *Theridion harmsi*. a) Female, dorsal habitus. b) Male, dorsal habitus. c) Left male palp, ventral. d) Left male palp, retrolateral e) Male embolus f) Female epigynum, ventral g) Female vulva, dorsal. Scale bars: 1 mm (a, b); 0.1 mm (c, d, f, g); 0.05 mm (e).




Figure 8. Scanning electron microscope image of left male palp. Top: ventral view. Bottom: prolatral view. a, b) *Theridion promiscuum* sp. nov. c, d) *Theridion mystaceum*. e, f) *Theridion harmsi*. Scale bars: 0.1 mm.

Our data clearly shows that while the phenotype and the nuclear data nicely delimit species boundaries in the studied species, the mitochondrial data suggest mixture across members of the *T. melanurum* group. The most obvious explanation for the apparent incongruence would be the human error in the manipulation of specimens or in the lab procedures. We discarded this source of error by re-extracting and resequencing specimens involved at least twice, independently. On the other hand, it is well known that mitochondrial pseudogenes, either as duplications within the mitochondrial genome or as copies inserted into the nuclear genome (e.g. NUMTs) may compromise the recovery of species boundaries using DNA barcoding approaches (Song et al. 2008) and, therefore, be in conflict with other sources of evidence. We examined all mitochondrial sequences generated in our study in search for some evidence of the presence of pseudogenes or NUMTs in our data set, conducting a careful examination of sequence characteristics, including indels, in-

frame stop codons and nucleotide composition. We found neither changes in reading frames nor in the base composition—all sequences showed the characteristic AT bias observed in many arthropod mitogenomes and all passed the composition chi-squared test.

Among the biological processes that may account for inconsistencies among different molecular markers, the incomplete sorting of ancestral polymorphism seems unlikely in our case. At least in spiders, the higher mutation rates and smaller population sizes of mitochondrial markers should ensure sorting in those markers previous to the nuclear ones, an opposite pattern of the one here reported. Although slower evolutionary rates in mitochondrial DNA have been found in some tetrapods (Nabholz et al. 2009) and cnidarians (Shearer et al. 2002), they have never been reported in spiders. Hybridization is the main biological culprit for mitochondrial discordance. However, the individuals sharing the same mitochondrial haplotype in our study were located hundreds of kilometres apart,

which casts some doubts on ongoing gene flow as a main explanation for the incongruence pattern recovered. It could be argued, however, that, like many other spiders, *Theridion* is able to drift long distances by ballooning, i.e. silk mediated airborne dispersal, which partially explains the large distribution ranges of many species. However, the extremely high dispersal rates required to explain the reported patterns of mixture in mtDNA should most likely also affect the nuclear and phenotypic data, which does not seem to be the case. Alternatively, the present day mixing of haplotypes within the species of the *T. melanurum* complex investigated could have been the result of older, hybridization events, possibly at a younger stage of the species ontogeny. The high dispersal rates of the species would have subsequently blurred the geographical signal of the hybridization processes.

Although at the moment we lack any solid evidence, there are also some chances that the mitochondrial incongruence observed could have been maintained by the involvement of endosymbiont driven cytoplasmic incompatibility. It is well known that endosymbiotic bacteria may affect the patterns of mitochondrial variation in invertebrates, which may compromise the inferences made on host evolution from these patterns (Hurst and Jiggins 2005). Although less well characterised than in insects, spiders are also known to host endosymbiotic bacteria (Goodacre et al. 2006; Zhang et al. 2018). The infection frequency may be as high as seventy percent of the individuals and specimens may be infected by multiple bacteria species and strain types (White et al. 2019). Studies on the diversity and structuring of *Wolbachia* endosymbionts in the funnel web spider genus *Agelenopsis* Giebel, 1869 suggest that *Wolbachia* strains induce cytoplasmic incompatibility, which may explain fixation of mtDNA haplotypes across wide geographic distances in these spiders (Baldo et al. 2008). Similarly, endosymbiont driven fixation of mtDNA haplotypes has been invoked to explain the mito-nuclear discordance recently reported in wolf spider species (Ivanov et al. 2018). We propose that cytoplasmic incompatibility driven by endosymbionts could have fixed mitochondrial haplotype mixtures across species in the *T. melanurum* complex originated from ancestral gene flow or ILS. Genotyping the endosymbiont strains hosted by those individuals from different species bearing similar haplotypes will have to be conducted to test our hypothesis.

The genus *Theridion* is a classic example of catch-all genus, a poorly defined group to include species with no trace of colulus that do not match other more precisely diagnosed genera. The redefinition of the genus would require an exhaustive systematic revision of a thorough sample of species, which is beyond the scope of the present paper. However, our results do support the existence of complexes of species within the genus. Specifically, molecular data (Fig. 2) confirms the delimitation of the *T. melanurum* species group, given the short branch lengths within it, in comparison to the branches that separate the remaining species. The *Theridion* species that were included in the tree and that were previously thought to be

long to this species complex (*T. mystaceum*, *T. harmsi*, *T. melanurum*, and *T. betteni*) are confirmed to be part of it. Additionally, the new species *T. promiscuum* sp. nov. is supported as a new member of the *T. melanurum* group. Species complexes, i.e. closely related species with limited morphological variation, are ideally suited to unravel the processes underlying speciation and the factors driving phenotypic change. Assuming that similarity is the result of recent divergence, there are better chances that the patterns of variation exhibited by species in those complexes are directly linked to the causal agents of the diversification of the group, and that they have not been blurred by subsequent processes. Future studies may take advantage of the *melanurum* and other species complexes within the species rich genus *Theridion* to gain better insights on the origins and maintenance of spider diversity in an age of major biodiversity loss.

Conclusions

We investigated here one of the few cases reported of mitochondrial discordance within spiders. A countrywide bioinventorying project revealed the existence of a new, morphologically diagnosable spider species within the *T. melanurum* group. Subsequent mtDNA barcode screening of specimens, however, identified instances of haplotype mixing across closely related species. Molecular information from a nuclear marker, on the other hand, supported the morphological delimitations, including the new specific status. The lack of geographic structure in the shared haplotypes and the lack of sorting in the fastest evolving gene suggests that mechanisms other than ongoing gene flow and deep coalescence are responsible for the observed patterns. We propose that cytoplasmic incompatibility mediated by endosymbionts may have been instrumental in generating mito-nuclear discordance, probably originated from old introgression events. Finally, this study highlights the important role that bioinventories play in improving our knowledge of biodiversity, especially in a time when fieldwork studies that gather new data are becoming less popular than those using pre-existing data, like modelling or meta-analysis studies (Ríos-Saldaña et al. 2018).

Acknowledgements

Collections were conducted under the corresponding permits kindly provided by the following individuals and institutions: Miguel M. de la Hoz (Picos de Europa), Elena Villagrasa (Ordesa), Maria Merced Aniz Montes (Aigüestortes), Angel Rodriguez Martin (Monfragüe), Angel Gómez Manzaneque (Cabañeros), and Blanca Ramos Losada (Sierra Nevada). We are grateful to all the people that contributed in the samplings as well as to all the park rangers that supported us in the field. We also thank Jagoba Malumbres-Olarte for generating the map in Figure 1. Lau-

ra Baldo kindly contributed insights into the interpretation of the results. We thank Cor Vik and Nadine Dupérré for their comments. This work was funded by the Organismo Autónomo de Parques Nacionales (OAPN #485/2012). M.D. thanks the University of Barcelona for the APIF PhD fellowship supporting this research. Additional support was provided by 2017SGR73 from the Catalan Government.

References

- Agnarsson I (2010) The utility of ITS2 in spider phylogenetics: notes on prior work and an example from *Anelosimus*. *Journal of Arachnology* 38: 377–382. <https://doi.org/10.1636/B10-01.1>
- Arnedo MA, Agnarsson I, Gillespie RG (2007) Molecular insights into the phylogenetic structure of the spider genus *Theridion* (Araneae, Theridiidae) and the origin of the Hawaiian *Theridion*-like fauna. *Zoologica Scripta* 36: 337–352. <https://doi.org/10.1111/j.1463-6409.2007.00280.x>
- Arnedo MA, Gillespie RG (2006) Species diversification patterns in the Polynesian jumping spider genus *Havaika* Prószyński, 2001 (Araneae, Salticidae). *Molecular Phylogenetics and Evolution* 41: 472–495. <https://doi.org/10.1016/j.ympev.2006.05.012>
- Astrin JJ, Höfer H, Spelda J, Holstein J, Bayer S, Hendrich L, Huber BA, Kielhorn K, Krammer H, Lemke M, Monje JC, Morini J, Rulik B, Petersen M, Janssen H, Muster C (2016) Towards a DNA barcode Reference Database for Spiders and Harvestmen of Germany. *PLoS ONE* 11: e016264. <https://doi.org/10.1371/journal.pone.0162624>
- Baldo L, Ayoub NA, Hayashi CY, Russell JA, Stahlhut JK, Werren JH (2008) Insight into the routes of *Wolbachia* invasion: High levels of horizontal transfer in the spider genus *Agelenopsis* revealed by *Wolbachia* strain and mitochondrial DNA diversity. *Molecular Ecology* 17: 557–569. <https://doi.org/10.1111/j.1365-294X.2007.03608.x>
- Ballard W, Whitlock M (2004) The incomplete natural history of mitochondria. *Molecular Ecology* 13: 729–744. <https://doi.org/10.1046/j.1365-294X.2003.02063.x>
- Bidegaray-Batista L, Sánchez-Gracia A, Santulli G, Maiorano L, Guisan A, Vogler AP, Arnedo MA (2016) Imprints of multiple glacial refugia in the Pyrenees revealed by phylogeography and palaeodistribution modelling of an endemic spider. *Molecular Ecology* 25: 2046–2064. <https://doi.org/10.1111/mec.13585>
- Blackledge TA, Kuntner M, Agnarsson I (2011) The form and function of spider orb webs: evolution from silk to ecosystems. *Advances in Insect Physiology* 41: 175–262. <https://doi.org/10.1016/B978-0-12-415919-8.00004-5>
- Borchsenius F (2009) FastGap 1.2. Software. http://www.aubot.dk/FastGap_home.htm
- Cardoso P (2009) Standardization and optimization of arthropod inventories—the case of Iberian spiders. *Biodiversity and Conservation* 18: 3949–3962. <https://doi.org/10.1007/s10531-009-9690-7>
- Chang J, Song D, Zhou K (2007) Incongruous nuclear and mitochondrial phylogeographic patterns in two sympatric lineages of the wolf spider *Pardosa astrigera* (Araneae: Lycosidae) from China. *Molecular Phylogenetics and Evolution* 42: 104–121. <https://doi.org/10.1016/j.ympev.2006.06.013>
- Clement M, Snell Q, Walker P, Posada D, Crandall K (2003) TCS: estimating gene genealogies. <https://doi.org/10.1109/IP-DPS.2002.1016585>
- Coddington JA, Agnarsson I, Cheng R-C, Čandek K, Driskell A, Frick H, Gregorič M, Kostanjšek R, Kropf C, Kweskin M, Lokovšek T, Papan M, Videgar N, Kuntner M (2016) DNA barcode data accurately assign higher spider taxa. *PeerJ* 4: e2201. <https://doi.org/10.7717/peerj.2201>
- Crespo LC, Domènech M, Enguídanos A, Malumbres-Olarte J, Moya-Laraño J, Frías-López C, Macías-Hernández N, Mas E De, Mazzuca P, Mora E, Opatova V, Planas E, Ribera C, Ruiz D, Sousa P, Tonzo V, Arnedo MA (2018) A DNA barcode-assisted annotated checklist of the spider (Arachnida, Araneae) communities associated to white oak woodlands in Spanish National Parks. *Biodiversity Data Journal* 6: e29443. <https://doi.org/10.3897/BDJ.6.e29443>
- Croucher PJP, Jones RM, Searle JB, Oxford GS (2007) Contrasting patterns of hybridization in large house spiders (*Teegenaria atrica* group, Agelenidae). *Evolution* 61: 1622–1640. <https://doi.org/10.1111/j.1558-5646.2007.00146.x>
- Dayrat B (2005) Towards integrative taxonomy. *Biological Journal of the Linnean Society* 85: 407–415. <https://doi.org/10.1111/j.1095-8312.2005.00503.x>
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299. <https://doi.org/10.1071/ZO9660275>
- Forster RR, Platnick NI, Coddington JA (1990) A proposal and review of the spider family Synotaxidae (Araneae, Araneoidea), with notes on theridiid interrelationships. *Bulletin of the American Museum of Natural History* 193: 1–116.
- Giles RE, Blanc H, Cann HM, Wallace DC (1980) Maternal inheritance of human mitochondrial DNA. *Proceedings of the National Academy of Sciences of the United States of America* 77: 6715–6719. <https://doi.org/10.1073/pnas.77.11.6715>
- Goodacre SL, Martin OY, Thomas CFG, Hewitt GM (2006) *Wolbachia* and other endosymbiont infections in spiders. *Molecular Ecology* 15: 517–527. <https://doi.org/10.1111/j.1365-294X.2005.02802.x>
- Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, Fritz MHY, Hansen NF, Durand EY, Malaspina AS, Jensen JD, Marques-Bonet T, Alkan C, Prüfer K, Meyer M, Burbano HA, Good JM, Schultz R, Aximu-Petri A, Butthof A, Höber B, Höffner B, Siegemund M, Weihmann A, Nusbaum C, Lander ES, Russ C, Novod N, Affourtit J, Egholm M, Verna C, Rudan P, Brajkovic D, Kucan Ž, Gušić I, Doronichev VB, Golovanova L V., Lalueza-Fox C, De La Rasilla M, Fortea J, Rosas A, Schmitz RW, Johnson PLF, Eichler EE, Falush D, Birney E, Mullikin JC, Slatkin M, Nielsen R, Kelso J, Lachmann M, Reich D, Pääbo S (2010) A draft sequence of the neandertal genome. *Science* 328: 710–722. <https://doi.org/10.1126/science.1188021>
- Gregorič M, Agnarsson I, Blackledge TA, Kuntner M (2015) Phylogenetic position and composition of Zyiellinae and *Caerostris*, with new insight into orb-web evolution and gigantism. *Zoological Journal of the Linnean Society* 175: 225–243. <https://doi.org/10.1111/zoj.12281>
- Harrison RG, Larson EL (2014) Hybridization, introgression, and the nature of species boundaries. *Journal of Heredity* 105: 795–809. <https://doi.org/10.1093/jhered/esu033>
- Hormiga G, Arnedo MA, Gillespie RG (2003) Speciation on a conveyor belt: sequential colonization of the Hawaiian Islands by *Orsonwelles* spiders (Araneae, Linyphiidae). *Systematic Biology* 52: 70–88. <https://doi.org/10.1080/10635150390132786>

- Hurst GDD, Jiggins FM (2005) Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. *Proceedings of the Royal Society B: Biological Sciences* 272: 1525–1534. <https://doi.org/10.1098/rspb.2005.3056>
- Ivanov V, Lee KM, Mutanen M (2018) Mitonuclear discordance in wolf spiders: genomic evidence for species integrity and introgression. *Molecular Ecology* 27: 1681–1695. <https://doi.org/10.1111/mec.14564>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30: 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28: 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33: 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Lanfear R, Calcott B, Ho SYW, Guindon S (2012) PartitionFinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29: 1695–1701. <https://doi.org/10.1093/molbev/mss020>
- Lattimore VL, Vink CJ, Paterson AM, Cruickshank RH (2011) Unidirectional introgression within the genus *Dolomedes* (Araneae: Pisauridae) in southern New Zealand. *Invertebrate Systematics* 25: 70–79. <https://doi.org/10.1071/IS11001>
- Lecigne S (2017) Contribution a la connaissance de l'araneofaune (Araneae) de l'Algarve (Portugal). Description de *Theridion bernardi* n. sp. (Araneae: Theridiidae) et Redécouverte d'*Ozyptila perplexa* Simon, 1875 (Araneae: Thomisidae). *Revista Ibérica de Aracnología* 31: 25–48.
- Leduc-Robert G, Maddison WP (2018) Phylogeny with introgression in *Habronattus* jumping spiders (Araneae: Salticidae). *BMC Evolutionary Biology* 18: 24. <https://doi.org/10.1186/s12862-018-1137-x>
- Leigh JW, Bryant D (2015) POPART: full-feature software for haplotype network construction. *Methods in Ecology and Evolution* 6: 1110–1116. <https://doi.org/10.1111/2041-210X.12410>
- Levi HW (1954) Spiders of the genus *Euryopsis* from North and Central America (Araneae, Theridiidae). *American Museum Novitates* 166: 1–48.
- Levi HW (1957) The spider genera *Enoplognatha*, *Theridion*, and *Paidiscura* in America North of Mexico (Araneae, Theridiidae). *Bulletin of the American Museum of Natural History* 112: 1–124.
- Maddison WP (1997) Gene trees in species trees. *Systematic Biology* 46: 523–536. <https://doi.org/10.1093/sysbio/46.3.523>
- Nabholz B, Glémin S, Galtier N (2009) The erratic mitochondrial clock: variations of mutation rate, not population size, affect mtDNA diversity across birds and mammals. *BMC Evolutionary Biology* 13: 54. <https://doi.org/10.1186/1471-2148-9-54>
- Oxford GS, Bolzern A (2018) Molecules v. Morphology – is *Eratigena atrica* (Araneae: Agelenidae) One Species or Three? *Arachnology* 17: 337–357. <https://doi.org/10.13156/ arac.2017.17.7.337>
- Padial JM, Miralles A, De la Riva I, Vences M (2010) The integrative future of taxonomy. *Frontiers in Zoology* 7: 16. <https://doi.org/10.1186/1742-9994-7-16>
- Pires AC, Marinoni L (2010) DNA barcoding and traditional taxonomy unified through integrative Taxonomy: a view that challenges the debate questioning both methodologies. *Biota Neotropica* 10: 339–346. <https://doi.org/10.1590/S1676-06032010000200035>
- Ratnasingham S, Hebert P (2007) BOLD: the Barcode of Life Data System (<http://www.boldsystems.org>). *Molecular Ecology Notes* 7: 355–364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Ratnasingham S, Hebert P (2013) A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. *PLoS ONE* 8: e66213. <https://doi.org/10.1371/journal.pone.0066213>
- Ríos-Saldaña CA, Delibes-Mateos M, Ferreira CC (2018) Are field-work studies being relegated to second place in conservation science? *Global Ecology and Biogeography* 14: e00389. <https://doi.org/10.1016/j.gecco.2018.e00389>
- Rogers J, Gibbs RA (2014) Comparative primate genomics: emerging patterns of genome content and dynamics. *Nature Reviews Genetics* 15: 347–359. <https://doi.org/10.1038/nrg3707>
- Shearer TL, Oppen MJH Van, Romano SL, Worheire G (2002) Slow mitochondria DNA sequence evolution in the Anthozoa. *Molecular Ecology* 11: 2475–2487. <https://doi.org/10.1046/j.1365-294X.2002.01652.x>
- Simon C, Frati F, Beckenbach A, Crespi B, Liu H, Flook P (1994) Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America* 87: 651–701. <https://doi.org/10.1093/aesa/87.6.651>
- Smith HM (2006) A revision of the genus *Polys* in Australasia (Araneae: Araneidae). *Records of the Australian Museum* 58: 43–96. <https://doi.org/10.3853/j.0067-1975.58.2006.1465>
- Song H, Buhay JE, Whiting MF, Crandall KA (2008) Many species in one: DNA barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are coamplified. *Proceedings of the National Academy of Sciences of the United States of America* 105: 13486–13491. <https://doi.org/10.1073/pnas.0803076105>
- Sota T, Vogler AP (2001) Incongruence of mitochondrial and nuclear gene trees in the carabid beetles *Ohomopterus*. *Systematic Biology* 50: 39–59. <https://doi.org/10.1093/sysbio/50.1.39>
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Vanuytven H (2014) *Theridion asopi* n. sp., a new member of the *Theridion melanurum* group (Araneae: Theridiidae) in Europe. *Arachnology* 16: 127–134. <https://doi.org/10.13156/ arac.2014.16.4.127>
- Vink CJ, Sirvid PJ, Malumbres-Olarte J, Griffiths JW, Paquin P, Paterson AM (2008) Species status and conservation issues of New Zealand's endemic *Latrodectus* spider species (Araneae: Theridiidae). *Invertebrate Systematics* 22: 589–604. <https://doi.org/10.1071/IS08027>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications* 55: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- White JA, Styer A, Rosenwald LC, Curry MM, Welch KD, Athey KJ, Chapman EG (2019) Endosymbiotic bacteria are prevalent and

- diverse in agricultural spiders. *Microbial Ecology* 79: 472–481. <https://doi.org/10.1007/s00248-019-01411-w>
- Will KW, Mishler BD, Wheeler QD (2005) The perils of DNA barcoding and the need for integrative taxonomy. *Systematic Biology* 54: 844–851. <https://doi.org/10.1080/10635150500354878>
- World Spider Catalog World Spider Catalog (2020) version 21.0. Natural History Museum Bern. <http://wsc.nmbe.ch> [Accessed on: 2020-2-27]
- Wunderlich J (2011) Extant and fossil spiders (Araneae). *Beiträge zur Araneologie* 6: 1–640.
- Zhang D, Cook WB, Horner NV (2004) ITS2 rDNA variation of the two black widow species, *Latrodectus mactans* and *Latrodectus hesperus* (Araneae, Theridiidae). *Journal of Arachnology* 32: 349–352. <https://doi.org/10.1636/H02-51>
- Zhang L, Yun Y, Hu G, Peng Y (2018) Insights into the bacterial symbiont diversity in spiders. *Ecology and Evolution* 8: 4899–4906. <https://doi.org/10.1002/ece3.4051>

Supplementary material 1

Additional information on the *Theridion* specimens

Authors: Marc Domènech, Luís C. Crespo, Alba Enguídanos, Miquel A. Arnedo

Data type: Specimen information

Explanation note: Spreadsheet containing the collection site, collection code, and COI and 28S GenBank accession numbers of all the *Theridion* specimens used in this study.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zse.96.49946.suppl1>