

Original Research Article

Physiological role of arginine on wheat (*Triticum aestivum* L.) cultivars grown under water drought condition

¹Ahmed M.A., ²Magda A. F. Shalaby and *¹Reda, E.Essa

Abstract

¹Field Crops Research Depart.,
National Research Centre, Dokki,
Giza, Egypt

²Botany Depart., National Research
Centre, Dokki, Giza, Egypt

*Corresponding Author's E-mail:
mashrera@yahoo.com

Two field experiments were conducted at Agricultural Experimental Station Research Centre of National Research Centre, Nubaria, El-Bahira Governorate, Egypt during two successive seasons 2017/2018 and 2018/2019. Each experiment was arranged in split-split plot design with four replicates. The present study investigated the effect of arginine (Ar) as amino acid priming agent, applied as foliar application spray, on growth, yield and its components, some chemical constituents of yielded grains of two wheat (*Triticum aestivum* L.) cv., Gemiza-9 and Misr-2 under drought. Moreover, the study aims to determine suitable concentrations of (Ar) as exogenous treatment to enhance wheat tolerance to drought. Foliar application spray with three levels of arginine (0, 2 and 4 mM/L) were applied twice after 50 and 60 days from sowing to increase the tolerance drought of wheat cultivars which irrigated with four treatments from water stress (Normal irrigation as a control, Once irrigation at tillering, germination and ripening stages) on two wheat cultivars (Gemiza-9 and Misr-2). The obtained results clearly indicated the role of Ar increasing the tolerance of wheat plants to drought. The highest rates of arginine led to significant increase in growth, yield and its components as well as chemical constituents of the wheat grains in both growing seasons. The magnitude of increments was pronounced in response to wheat cultivar Misr-2 and 4 mM/L of arginine as a foliar spray which led to positive changes in all studied parameters under normal irrigation. All treatments were effective in alleviating the harmful effect of drought.

Keywords: Arginine, Cultivars, Drought, Foliar application, Wheat

INTRODUCTION

World agriculture is facing a lot of challenges like producing 70% more food for an additional population of 2.3 billion by 2050, while at the same time fighting with poverty and hunger, consuming scarce natural resources more efficiently and adapting to climate change (FAO, 2009). The importance of wheat for human main food is the well-known fact, overall the world as well as Egypt, extensive. Effects on wheat are continuously paid for

increasing its productivity by means of vertical and/or horizontal planting. In Egypt wheat crop covers about 3.10 million feddan distributed mainly in the old land and partially in the new land (Egypt stat. Agric. Rep. 2011). In the light of present national water policy using wheat cultivars that produce high yielding under suitable water regime. Egyptian wheat plants are sometimes exposed to drought stress at different period of growth. A possible

Table 1. Physical and chemical analyses of the experimental site

Depth (cm)	Chemical				Mechanical			Texture
	O.M (%)	P ^H (1:2.5)	E.C. (0.5 m ⁻¹)	CaCO ₃ (%)	Course Sand	Fine Sand	Silt & Clay	
0-20	0.65	8.70	0.35	7.01	47.46	49.79	2.49	Sandy
20-40	0.40	8.80	0.33	2.37	56.72	39.57	3.74	
40-60	0.25	9.30	0.45	4.71	36.76	59.39	3.85	
Depth (cm)	SP (%)	FC (%)	WP (%)	AW (%)	Hydraulic conductivity (cm/hr)			
0-20	21.00	10.10	4.68	5.41	22.50			
20-40	19.00	13.50	5.61	7.89	19.00			
40-60	22.00	14.49	4.63	7.89	21.00			

Notes: the data of two growing seasons were combined

approach to minimize drought stress that induces crop losses is the foliar application with arginine on wheat plant (Abdul Qados, 2010 and Ahmed *et al.*, 2015).

In recent years, growth regulating substances played an important role in controlling seed germination, vegetative growth, flowering and yield of several crop plants (Abd El-Monem, 2007). Arginine is one of the essential amino acids, considered the main precursor of polyamines which produced by decarboxylation of arginine via arginine decarboxylase to form putrescine (Bouchereau *et al.*, 1999).

Polyamines and their precursor arginine have been implicated as vital modulators in a variety of growth, physiological and developmental processes in higher plants (Galston and Kaur Sahney, 1990). Polyamines are involved in the control of cell cycle, cell division, morphogenesis in phytochrome and plant hormone mediated process and the control of plant senescence, as well as in plant response to various stress factors (El-Shintinawy and Hassanein, 2001).

The application of arginine significantly promoted the growth and increased the fresh and dry weights, certain endogenous plant growth regulators, chlorophylls a and b and carotenoids in wheat (Abd El-Monem, 2007 and El-Bassiouny *et al.*, 2008). Moreover Hassanein *et al.*, (2008) showed the positive role of arginine in alleviating the inhibition occurs as the result of exposing plants to stress.

Therefore, the objective of the physiological role of foliar treatment with arginine on some wheat cultivars (*Triticum aestivum* L.) under water stress condition in the newly cultivated in sandy soil and to investigate the role of arginine in alleviating the harmful effects of drought.

MATERIALS AND METHODS

Two field experiments were carried out at Agricultural Experimental Station Research Centre of National Research Centre, Nubaria, El-Bahira Governorate, Egypt, during two successive seasons; 2017/2018 and 2018/2019. Each experiment was arranged in split-split

plot design with four replicates. Each experiment was arranged in split-split plot design with four replicates. Cultivars of wheat (*Triticum aestivum* L) cv., Gemiza-9 and Misr-2, were used which obtained from the Ministry of Egyptian Agriculture.

Foliar application spray with three levels of arginine (0, 2 and 4 mM/L) were applied twice after 50 and 60 days from sowing and irrigated with four treatments from water stress (Normal irrigation as a control which irrigated with one-week intervals up to maturity stage (140 days after sowing), Once irrigation at tillering at 49 days after sowing, germination at 98 days after sowing and ripening stage at 105 days after sowing.).

Cultivars were allocated to main plots, while the water treatments ones were allocated to sub-plot and arginine application treatments were distributed in sub-sub plots. The chemical used in the present work was arginine (one of the essential amino acids). The structure formula of this compound is: $\text{NH}_2 - \text{COOH} - \text{CH}(\text{CH}_2) \text{NH}_3 - \text{C} - \text{NH} - \text{NH}_2$. This experiment was carried out to investigate the physiological role of foliar application by arginine on alleviating the harmful effect of drought on wheat cultivars. Table (1), presented the physical and chemical analysis of the soil that determined according to Richards (1954).

The area of experimental unit consisted of 10 rows each of 2.5 m in length at 20 cm apart. Grains were sown in November at the last week for both seasons at grain rates of 40 kg/fad. The recommended cultural practices of growing wheat plant were followed by harvest date as recommended by Wheat Research Department, Agricultural Research Center.

Samples of guarded plants were often taken randomly from each plot for the four replications to measure growth parameters after 110 and 120 days of sowing, where, plant height, number and dried weight of each tiller blades, spikes/plant, flag leaf blade area cm², and blade area cm²/plant were determined according to Bremner and Taha (1966). LAI was estimated according to the method described by Watson (1952).

At harvest date, guarded plants were taken randomly from the middle rows of each plot to determine the

number of spikes per plant, dry weight of spike (g/plant), grain index (weight of 1000 grain/g), straw, biological biomass (above ground) and yields per plant.

In addition, a migration coefficient (i.e. dry weight of spikes/plant/above ground biomass per plant) was estimated according to Abdel-Gawad *et al.*, (1987). Moreover, grains (ton/fed); straw (ton/fed) and above ground biomass in ton/fed were collected from the whole area of each experimental unit and then converted to yield per feddan.

Chemical analysis

Samples were chosen for some chemicals composition, photosynthetic pigments contents in wheat blades (mg/g dry wt.) were extracted by aqueous solution of 85% acetone and calculated using Van Wettstein formula (Van Wettstein, 1957) after 110 and 120 days of sowing. The dried grains were finally grounded and kept for carbohydrates and protein determinations. The total carbohydrates percentages were determined using a phenol sulfuric acid method as described by (Duois *et al.*, 1956) whereas, N % in dry grains was determined according to the method of Pregl (1945), also, crude protein percentages in dry grains were calculated by multiplying N content by 5.75. Phosphorus was extracted and using spectrophotometer according to Jackson (1973), whereas, potassium was assayed using flame photometer according to Allen *et al.*, (1984).

Statistical analysis

Data were statistically analyzed by using analysis of variances according to Snedecor and Cochran (1990) and means were grouped by LSD test at the 5% probability level. Furthermore, combined analysis was made for the data of the two growing seasons as the results were followed a similar trend.

RESULTS

Effect of cultivars

Results in Table (2), showed that significant differences were found between the two wheat cultivars Gemiza-9 and Misr-2 in all growth parameters after 110- and 120-days of two growing seasons. Gemiza-9 cultivar significantly surpassed Misr-2 cultivar in plant height, number of blades/plant and tillers dry weight/plant after 110 and 120 days of sowing, whereas, Misr-2 cultivar gave the highest significant values at number of tillers/plant, number of spikes/plant, blades dry weight/plant, spikes dry weight/plant, flag leaf blade area, blades area/plant and LAI after 110 and 120 days

compared with Gemiza-9 cultivar.

It is worthy, plant height, number of tillers/plant, the number of blades/plant, dry weight of tillers/plant, dry weight of blades/plant, blades area/plant and LAI were decreased with advancing plant age to 110 and 120 days however, number of spikes/plant, dry weight of spikes/plant and the flag leaf blade area seemed to be increased with advancing plant age to 110 and 120 days after sowing.

Regarding, chemical concentration of wheat plant Gemiza-9 and Misr-2 cultivars, table (3) indicated that the cultivars significantly differed in photosynthetic pigment content per wheat blade (i.e. Chl a, Chl b, Chl (a+b) and carotenoids) as well as, total carbohydrates % and crude protein % per grains were significant. Also, photosynthetic pigment content per green wheat blades was decreased after 110 days from sowing. Also, from table (3) Misr-2 in photosynthetic pigments as compared to other wheat cultivars after 110 and 120 days of sowing and total carbohydrates % per grains at harvest date. On the contrary, Gemiza-9 cultivar produced the greatest mean values from crude protein % per grains at harvest date compared with Misr-2 cultivar.

Table (4) showed clearly that Gemiza-9 and Misr-2 cultivars were significantly differed in yield and its components (i.e. Number of spikes/plant, spikes dry weight/plant, mean of spike length, grain index, grains, straw and above ground biomass yields per plant and/or per fed., migration coefficient, crop index and harvest index. Data in the same table clearly indicated that Misr-2 was significantly out weight of Gemiza-9 cultivar in all previous yield and its components except straw yield per plant and/or per fed where Misr-2 cultivar characterized by significant decrease in straw yield compared with Gemiza-9 cultivar.

It is worthy to mention that the differences between wheat cultivars might be attributed to the differences between wheat genotype for mineral elements concentration and to the differences in photosynthetic partitioning and migration photosynthetic among wheat plant organs. Generally, the significant cultivar differences between wheat cultivars in this study are in full agreement with previous results obtained by Hassanain *et al.*, (2014) and El-Metwally *et al.*, (2015).

Effect of drought

Results in Table (2) showed that skipping irrigation at certain developmental stages of growth for wheat plant caused significant decrement in growth parameters, i.e. plant height, number and dried weight of tillers, blades and spikes/plant, flag leaf blade area, blade area/plant and LAI at 110 and 120 days after sowing date (Table 2), photosynthetic pigment content (i.e Chl a, Chl b, Chl (a+b) and carotenoids) per green blades at 110 and 120 days after sowing. In addition, total carbohydrates % and

Table 2. Effect of cultivars, water stress and arginine concentrations on growth parameters of wheat plant (Average of 2017/2018 and 2018/2109 seasons).

Cultivars	Water stress treats.	Ar Con cen (mM /l)	Plant height (cm)	110 days after sowing								LAI
				No./plant			Dry weight (g/plant)			Flag leaf blade area	Blades area (cm ²)/ plant	
				Tiller r	Blade s	Spike s	Tiller + shoot s	Blad es	Spike s			
Gemiza-9			156.29	6.03	40.86	5.17	8.23	5.64	8.09	35.58	958.73	6.39
Misr-2			151.09	6.80	40.58	5.77	7.82	5.83	8.56	40.68	1005.91	6.71
LSD 0.05			1.53	0.29	0.14	0.30	Ns	0.15	0.18	1.27	11.64	0.20
LSD 0.05	Control		160.69	6.94	44.33	6.29	8.62	6.02	8.91	43.58	1026.30	6.85
	Tillering		141.35	5.93	36.52	4.78	7.50	5.42	7.50	31.88	932.37	6.22
	Germinating		149.92	6.14	40.14	5.21	7.82	5.67	8.22	36.70	968.31	6.47
	Ripening		157.58	6.63	43.11	5.70	8.11	5.83	8.68	40.74	1000.30	6.67
			2.88	0.15	1.51	0.23	0.27	0.13	0.60	0.63	1.46	0.06
LSD 0.05		0.00	147.61	6.05	38.17	5.08	7.41	5.50	7.81	34.32	940.50	6.27
		2.00	154.95	6.47	40.60	5.59	8.04	5.70	8.25	38.00	983.73	6.58
		4.00	158.55	6.70	43.39	5.81	8.64	6.00	8.93	42.10	1022.80	6.82
			1.44	0.10	2.01	0.15	0.35	0.16	0.22	2.24	27.46	0.08
120 days after sowing												
Gemiza-9			148.20	5.77	38.70	5.68	7.98	5.08	8.63	38.92	902.30	6.02
Misr-2			141.89	6.36	39.05	6.24	7.38	5.28	9.37	44.32	995.07	6.63
LSD 0.05			3.12	0.25	ns	ns	0.22	0.11	0.36	2.03	17.52	0.41
LSD 0.05	Control		151.71	6.66	42.40	6.55	8.43	5.58	9.78	46.05	1010.05	6.73
	Tillering		137.00	5.29	34.95	5.03	6.67	4.88	8.44	37.13	893.76	5.96
	Germinating		145.51	6.07	38.05	6.00	7.51	5.12	8.81	39.88	927.38	6.18
	Ripening		148.51	6.40	40.10	6.27	7.93	5.25	9.17	43.46	963.52	6.42
			1.06	0.08	0.01	0.13	0.08	0.04	0.31	0.82	9.97	0.08
LSD 0.05		0.00	140.29	5.70	36.33	5.61	7.11	4.91	8.61	38.20	910.73	6.07
		2.00	142.36	6.00	39.00	5.98	7.67	5.18	8.98	41.67	946.70	6.31
		4.00	149.60	6.48	41.28	6.31	8.27	5.45	9.43	45.00	988.14	6.59
			2.01	0.16	1.64	0.20	0.27	0.11	0.12	1.56	30.61	0.14

Table 3. Effect of cultivars, water stress and arginine concentrations on chemical constituents of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivar s	Water stress treats.	Ar Concen . (mM/l)	Photosynthetic pigments content per blades (mg/g) dry weight								Total Carb. (%)	Crude protein(%)
			110 days after sowing				120 days after sowing					
			Chl a	Chl b	Chl (a+b)	Car.	Chl a	Chl b	Chl (a+b)	Car.		
Gemiza-9			2.78	1.10	3.88	1.88	2.63	0.96	3.59	1.75	84.79	11.95
Misr-2			2.89	1.22	4.11	1.95	2.72	1.04	3.76	1.84	85.54	11.75
LSD 0.05			0.07	0.03	0.09	0.05	0.02	0.04	0.08	0.06	0.11	0.10
	Control		3.06	1.35	4.41	2.17	2.89	1.16	4.05	2.01	86.19	11.67
	Tillering		2.61	0.98	3.59	1.70	2.49	0.82	3.31	1.64	83.69	12.31
	Germin ating		2.77	1.12	3.89	1.82	2.61	0.95	3.56	1.72	85.13	11.73
	Ripenin g		2.91	1.19	4.10	1.96	2.73	1.07	3.80	1.83	85.66	11.70
LSD 0.05			0.11	0.08	0.14	0.09	0.12	0.06	0.17	0.13	0.03	0.21
		0.00	2.70	1.08	3.78	1.83	2.55	0.91	3.46	1.68	84.79	11.48
		2.00	2.82	1.14	3.96	1.93	2.68	0.99	3.67	1.83	85.16	11.80
		4.00	2.95	1.28	4.25	1.98	2.80	1.10	3.90	1.90	85.44	11.89
LSD 0.05			0.08	0.07	0.19	0.04	0.10	0.06	0.18	0.04	0.25	0.08

Table 4. Effect of cultivars, water stress and arginine concentrations on yield and its components of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultiva rs	Water stress treats.	Ar Conc en. (mM/ l)	No of spik es/ pla nt	Spi kes dry wei ght (g/pl ant)	Mai n spik es len gth (cm)	Grai n ind ex (g)	Grai n yel d g/pl ant)	Str aw yie ld g/pl ant)	Abo ve grou nd biom ass (g/pl ant)	Grai n yel d (ton /fed)	Stra w yel d (ton /fed)	Abo ve grou nd bio mas s (ton/ fed)	Migr atio n coef ficie nt	Harv est inde x	Cro p inde x
Gemiz a-9			5.79	19.9	15.6	32.9	14.4	21.	35.8	2.83	3.34	6.17	0.56	0.85	0.46
Misr-2			6.44	21.7	17.1	35.1	15.8	20.	36.3	3.01	3.21	6.22	0.60	0.94	0.48
LSD _{0.05}			0.61	0.16	0.13	0.35	0.30	0.8	0.27	0.12	0.07	0.03	0.02	0.05	0.01
	Control		6.68	24.5	24.5	37.5	18.4	24.	42.4	3.20	3.49	6.69	0.58	0.92	0.48
	Tillering		5.13	17.4	13.6	31.3	12.1	17.	29.5	2.55	3.15	5.70	0.59	0.81	0.45
	Germinating		6.12	19.6	15.9	32.6	13.9	20.	34.2	2.90	3.20	6.10	0.57	0.91	0.48
	Ripening		6.37	21.7	17.3	34.5	15.8	22.	38.0	3.04	3.26	6.30	0.57	0.93	0.48
LSD _{0.05}			0.13	0.18	0.49	0.85	0.55	0.5	0.41	0.03	0.15	0.07	0.01	0.01	ns
		0.00	5.71	18.2	14.2	31.9	13.3	19.	32.5	2.81	3.35	6.16	0.56	0.84	0.46
		2.00	6.08	20.0	16.3	33.8	14.8	20.	34.8	2.96	3.39	6.25	0.58	0.90	0.47
		4.00	6.43	24.3	18.6	36.2	17.1	22.	40.0	3.11	3.44	6.55	0.61	0.90	0.47
LSD _{0.05}			0.31	1.82	1.25	0.26	0.22	0.1	0.86	0.09	0.08	0.03	0.02	0.01	0.01

crude protein % per dried grains at harvest date (Table 3), as well as, yield and its components, i.e. number of spikes/plant, dry weight of spike per plant, mean of spike length, grain index, grain, straw and above ground biomass, yields per plant and/or per feddan, migration coefficient, harvest index and crop index (Table 4).

Data collected indicated that plant height, number and the dry weight of tillers + sheaths and blades, blades area/plant and LAI were decreased with advancing plant age after 110 and 120 days of sowing (Table 2), as well as, photosynthetic pigment content (Table 3). On the contrary, the number and dried weight of spikes/plant, flag leaf blade area was increased with advancing plant age after 110- and 120-days. Noticeably, the damaging effect of the harmful effect of drought stress on growth character (Table 2), chemical constituents of wheat plant (Table 3) with yield and its components (Table 4), may be explained on the basis of turgidity loss with effect of cell expansion and ultimate cell size, the loss of turgidity is probably the most sensitive processes to water stress, thus caused decrement in growth rate, stem elongation and enlargement which has been carefully discussed by Kramer and Boyer (1995). As a result, yield and its components were decreased consequently. Our results in Tables (2, 3 and 4) clearly indicated that wheat plants were more sensitive to drought stress at tillering stage, followed by germinating stage, whereas, skipping one irrigation at ripening stage showed the lowest damaging effect on each of growth characters (Tables 2), chemical constituents (Tables 3) and, yield and its components (Tables 4). On the other hand, water stress for wheat plant even water afterward did not recover to their normal behavior to compensate for the adverse effect caused by the exposure to drought conditions (Magda *et al.*, 2014).

Generally, irrigation is recommended due to the highest effect on tiller survival and this included that development and physiological processes in determining final grain yield and the water stress should be avoided at this growth stage. Therefore, the damage in growth characters (Tables 2), chemical constituents (Tables 3), and yield with its components (Tables 4) attributed to water stress at tillering stage was more pronounced, because every plant was subjected to soil moisture at tillering stage and their effect might be attributed to the lack of absorbed water, inadequate uptake of chemical elements, inhibition of meristematic activity and/or reduction in photosynthetic capacity under such unfavorable conditions (Magda, *et al.*, 2018).

Moreover, assimilates translocation to the new developing tillers and to the spike primordial were recorded which were not enough to mention or develop these organs from plants (Magda, *et al.*, 2018). Finally, our results are in good harmony with those obtained by Abdel Salam *et al.*, (2016).

Effect of foliar spray by arginine.

Data illustrated in Table (2) indicated that the growth attributes of wheat plants to the concentration of arginine were significant, where; plant height, number and dried weight/plant of each one of tillers + shoots, blades and spikes/plant, flag leaf blade area and blade area/plant and LAI were significantly affected by foliar application with arginine. In addition, foliar spray with 2.0 mM/l caused a significant increment in all growth parameters after 110 and 120 days of sowing compared to control treatment (Tap water treatment). On the other hand, increasing concentration of arginine from 2.0 to 4.0 mM/l caused another significant increase in growth parameters after 110 and 120 days compared to control treatment and 2.0 mM/l treatment only, respectively.

Furthermore, a significant response of photosynthetic pigment contents after 110 and 120 days of sowing were found due to the foliar application with arginine. Also, carbohydrates % and crude protein % per dry grains at harvest date, where, the application with 2.0 mM/l arginine significantly enhanced the previous chemical composition of wheat plant compared to control plant after 110 and 120 days of sowing, where, increasing arginine concentration from 2.0 to 4.0 mM/l caused further increases in Chl a, Chl b, Chl (a+b) and carotenoids per wheat blades after 110 and 120 days of sowing plus in carbohydrates and crude protein % per dried grains at harvest date compared to 2.0 mM/l concentration.

On the other hand, each to plant height, number of tillers/plant, number of blades/plant, tillers + shoots, dry weight/plant, dried weight of blade/plant, blade area/plant and LAI, as well as, Chl a, Chl b, Chl (a+b) and carotenoids at seemed to be decreased with advancing plant age (110 and 120) days, whereas, the number and dry weight of spikes/plant, as well as, flag leaf blade area were increased with advancing plant age (110 and 120) days.

The effect of arginine concentration on yield and its components of wheat plant were respected significantly at level (5%) in Table (4). Results showed clearly that foliar spray with 2.0 mM/l arginine significantly enhanced the number of spikes/plant; dry weight of spike/plant, the mean of spike length, grain index, grain, straw and above ground biomass/plant, above ground biomass/feddan, migration coefficient, harvest index and crop index compared with control treatment, except the increase in grain and straw yield/fed were not reaching the significant level at 5%.

Moreover, foliar spraying with 4 mM/l arginine on wheat plant caused significant increment in all the yields and its component except the differences in grains and straw yield/feddan, harvest index and crop index between 2 and 4 mM/l arginine that were not in significant level at 5 %.

Table 5. Effect of interaction between cultivars and water stress on growth characters of wheat plant (during 2017/2018 and 2018/2019 seasons).

Cultivar s	Water stress treats.	110 days after sowing									
		Plant height (cm)	No./plant			Dry weight (g/plant)			Flag leaf blade area	Blades area (cm ²)/ plant	LAI
			Tiller	Blade s	Spikes	Tiller + shoots	Blade s	Spikes			
Gemiza-9	Control	167.77	6.77	44.43	6.00	8.87	5.89	8.75	40.82	1003.2	6.69
	Tillering	143.77	5.49	36.63	4.56	7.59	5.30	7.11	29.19	921.60	6.14
	Germin ating	153.11	5.57	40.58	4.83	8.04	5.58	8.08	34.90	939.35	6.26
	Ripenin g	160.43	6.25	41.77	5.28	8.31	5.77	8.36	38.39	970.73	6.47
Misr-2	Control	153.60	7.11	44.22	6.50	8.37	6.14	9.07	46.53	1049.40	7.00
	Tillering	138.93	6.31	36.40	5.00	7.41	5.54	7.83	34.57	943.14	6.29
	Germin ating	146.73	6.64	39.70	5.58	7.60	5.75	8.35	38.90	1000.26	6.67
	Ripenin g	154.73	7.00	42.00	6.12	7.91	5.88	9.00	43.09	1029.87	6.87
LSD _{0.05}		4.55	0.24	2.38	0.37	0.42	0.20	0.10	0.99	23.00	0.10
120 days after sowing											
Gemiza-9	Control	153.60	6.42	42.40	6.33	8.69	5.42	9.26	42.40	965.49	6.44
	Tillering	140.70	5.07	35.20	4.78	6.91	4.73	8.18	34.65	855.39	5.70
	Germin ating	147.93	5.84	37.70	5.70	7.75	5.02	8.55	38.50	877.23	5.85
	Ripenin g	150.58	6.10	39.60	5.93	8.19	5.15	8.93	40.20	911.04	6.07
Misr-2	Control	149.82	6.90	42.40	6.77	8.16	5.72	10.30	49.70	1054.60	7.04
	Tillering	133.27	5.50	34.70	5.28	6.42	5.02	8.69	39.60	932.13	6.21
	Germin ating	142.56	6.30	38.40	6.30	7.26	5.22	9.06	41.25	977.53	6.52
	Ripenin g	146.43	6.70	40.60	6.60	7.67	5.34	9.41	46.72	1016.00	6.37
LSD _{0.05}		1.68	0.12	0.02	0.21	0.12	0.06	0.49	1.29	15.75	0.13

These results may be attributed to the role of PAs (the main group of arginine products), application in alleviating the adverse effect of drought and salt stress by increasing the endogenous PAs contents in wheat plants and proved that arginine treatments significantly increased grains weight per plant, the weight of 1000-grain, straw and biological yields per plant and harvest index in wheat plants (El-Shintinawy and Hassanein, 2001)

It is notable that our results are confirmed with results obtained by Nasibiet *al.*, (2014); Ahemd *et al.*, (2015); Magda *et al.*, (2018) for the connection the main role of arginine products (putrescine, spermidine and spermine) in the stressed plants for the long run, which was to maintain a cation-anion balance in plant tissue by

stabilizing membrane of high external stress (El-Bassiouny, *et al.*, 2008). Foliar spraying with polyamine (end product of arginine to several plant species have been shown to promoting the cell division, cell differentiation and general growth promotion. The arginine can also help in stabilizing the membrane and cell properties (Velikova, *et al.*, 2000) and protecting the plant against environmental stress (Mo and Pua, 2002).

Furthermore, the positive effect of arginine on yield and its components may be attributed to the positive effect of arginine on growth characters (Table 2) and photosynthetic pigment content per wheat green blades (Table 3). Thus, application of arginine can be used to decrease harmful effect of drought stress and wear. The arginine treatment can help in solving the

Table 6. Effect of interaction between cultivars and water stress on chemical constituents of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivar s	Water stress treats.	Photosynthetic pigments content per blades (mg/g) dry weight								Total Carb. (%)	Crude protei n (%)
		110 days after sowing				120 days after sowing					
		Chl a	Chl b	Chl (a+b)	Car.	Chl A	Chl B	Chl (a+b)	Car.		
Gemiza-9	Control	3.01	1.31	4.32	2.11	2.84	1.14	3.98	1.96	86.02	11.64
	Tillering	2.55	0.88	3.43	1.67	2.42	0.77	3.19	1.58	82.92	12.80
	Germin ating	2.68	1.05	3.73	1.79	2.55	0.90	3.45	1.68	84.79	11.69
	Ripenin g	2.87	1.17	4.04	1.94	2.70	1.02	3.72	1.78	85.44	11.66
	Control	3.11	1.38	4.49	2.22	2.94	1.18	4.12	2.05	86.32	11.69
Misr-2	Tillering	2.66	1.08	3.74	1.73	2.55	0.86	3.41	1.69	84.48	11.82
	Germin ating	2.85	1.19	4.04	1.85	2.66	0.99	3.65	1.76	85.47	11.77
	Ripenin g	2.95	1.21	4.16	1.98	2.75	1.11	3.86	1.87	85.87	11.73
	LSD _{0.05}	0.17	0.13	0.22	0.14	0.19	0.10	0.27	0.20	0.05	ns

Table 7. Effect of interaction between cultivars and water stress on yield and its components of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivars	Water stress treats.	No of spikes/plant	Spikes dry weight (g/plant)	Main spikes length (cm)	Grain index (g)	Grain yield g/plant	Straw yield g/plant	Above ground biomass (g/plant)	Grain yield (ton/fed)	Straw yield (ton/fed)	Above ground biomass (ton/fed)	Migration coefficient	Harvest index	Crop index
Gemiza-9	Control	6.45	23.46	18.05	36.33	17.70	24.92	42.62	3.11	3.56	6.67	0.56	0.87	0.47
	Tillering	4.87	16.67	12.98	30.46	11.41	17.94	29.35	2.45	3.23	5.68	0.57	0.76	0.43
	Germinating	5.81	18.92	14.99	31.12	13.21	20.16	33.37	2.81	3.26	6.07	0.57	0.86	0.46
	Ripening	6.04	20.81	16.71	33.79	15.27	22.73	38.00	2.95	3.30	6.25	0.55	0.89	0.47
Misr-2	Control	6.90	25.67	19.31	38.70	19.17	23.09	42.26	3.28	3.42	6.70	0.61	0.96	0.49
	Tillering	5.38	18.24	14.39	32.31	12.97	16.77	29.74	2.65	3.06	5.71	0.61	0.87	0.46
	Germinating	6.42	20.39	16.85	34.19	14.72	20.48	35.20	2.98	3.14	6.12	0.58	0.95	0.49
	Ripening	6.69	22.69	18.04	35.24	16.38	21.63	38.01	3.12	3.22	6.34	0.60	0.97	0.49
LSD _{0.05}		0.20	0.28	0.77	1.35	0.87	0.87	0.65	0.04	0.14	0.11	0.03	0.03	0.01

problems produced by environmental drought stress. The positive effect of arginine products (putrescine, spermidine and spermine) used in our study is in a good harmony with those obtained by Ahmed *et al.*, (2015); Akladios and Hanafy, (2018).

However, the disappearance of some protein bands under stress may be due to the suppression of the genes responsible for protein synthesis as a result of stress. Then, the developed tissues had lost their ability to

synthesize these proteins under stress conditions (Akladios and Abbas, 2014).

Effect of the interaction between cultivars X skipping irrigation

Table (5), indicated that the interaction between wheat cultivars X skipping irrigation at certain developmental

Table 8. Effect of interaction between cultivars and arginine concentrations on growth characters of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivar s	Ar Concen · (mM/l)	Plant height (cm)	110 days after sowing								LAI
			No./plant			Dry weight (g/plant)			Flag leaf blade area	Blades area (cm ²)/ plant	
			Tiller	Blade s	Spikes	Tiller + shoots	Blade s	Spikes			
Gemiza-9	0.00	149.7 1	5.72	37.94	4.75	7.58	5.39	7.42	32.02	918.90	6.13
	2.00	157.4 0	6.03	40.33	5.25	8.33	5.57	8.11	35.21	957.78	6.39
	4.00	161.7 5	6.33	44.30	5.50	8.77	5.95	8.75	39.52	999.50	6.66
Misr-2	0.00	145.6 1	6.38	38.40	5.40	7.23	5.61	8.20	36.61	962.05	6.41
	2.00	152.3 0	6.90	40.86	5.80	7.74	5.80	8.38	40.76	1009.67	6.73
	4.00	155.3 5	7.06	42.48	6.11	8.50	6.08	9.10	44.66	1046.00	6.97
LSD _{0.05}		2.24	0.16	3.14	0.23	0.55	0.25	0.35	3.50	42.84	0.13
120 days after sowing											
Gemiza-9	0.00	143.0 0	5.40	36.12	5.29	7.30	4.83	8.25	35.56	878.90	5.86
	2.00	148.6 7	5.58	38.64	5.65	7.95	5.08	8.55	38.50	891.70	5.98
	4.00	152.9 3	6.32	41.35	6.11	8.70	5.33	9.10	42.71	931.30	6.21
Misr-2	0.00	137.5 7	6.00	36.54	5.92	6.91	4.98	8.96	40.84	942.55	6.28
	2.00	141.8 4	6.44	39.40	6.30	7.38	5.28	9.40	44.83	996.70	6.64
	4.00	146.2 6	6.64	41.20	6.50	7.84	5.57	9.75	47.28	1045.97	6.97
LSD _{0.05}		3.14	0.25	2.56	0.31	0.42	0.17	0.19	2.44	47.75	0.17

stages on growth character after 110 and 120 days of sowing (Table 5), chemical constituents except crude protein per dried grain (Table 6), and yield and its components (Table 7) was significant, where Misr-2 cultivar surpassed Gemiza-9 in all previous determined characters except plant height after 110 and 120 days of sowing, Gemiza-9 was producing the tallest wheat plant. Furthermore, the collected data indicated that wheat plants were more sensitive to skipping irrigation at tillering stage followed by germinating stage, while the skipping irrigation at ripening stage was at the end of list. These results are in agreement with the findings obtained by Magda *et al.*, (2014).

With respect of the interaction of wheat cultivars X arginine concentration, data observed that the interaction was significant on growth characters (Table 8), on chemical constituents (Table 9) and the yield associated with its components (Table 10). Furthermore, data obtained from the previous table, explained that the most effective treatment was Misr-2 cultivar sprayed with 4 mM/l of arginine. These results are in harmony with those obtained by El-Bassiouny, *et al.*, (2008).

Regarding the interaction between skipping irrigation

at certain developmental stages of growth and arginine concentration treatments on growth characters after 110 and 120 days of sowing (Table 11) chemical constituents of the wheat plant (Table 12) and the yield accompanied with its components except for the crop index and harvest index (Table 13) was significant. Also, the most favorable treatment for all the previous characters was the traditional irrigation (without skipping) under foliar spraying with 4 mM/l arginine followed by normal irrigation under spraying with 2 mM/l arginine. These results are in agreement with those obtained by Ahmed *et al.*, (2015) and Magda *et al.*, (2018).

The three-ways interaction among wheat cultivars X drought stress treatment X foliar spraying with arginine clearly showed that the effect on growth characters (Table 14), chemical constituents of the wheat plant, except crude protein per dried grains (Table 15), as well as, yield and its components, except the crop index and harvest index (Table 16) were significant. Generally, the most effective treatment for increasing wheat plant yield and quality are Misr-2 cultivar under normal irrigation using the foliar spraying with 4 mM/l arginine followed by Misr-2 under normal using the foliar spraying with 2 mM/l

Table 9. Effect of interaction between cultivars and arginine concentrations on chemical constituents of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivars	Arginine Concen. (mM/l)	Photosynthetic pigments content per blades (mg/g) dry weight								Total Carb. (%)	Crude protein (%)
		110 days after sowing				120 days after sowing					
		Chl a	Chl b	Chl (a+b)	Car.	Chl a	Chl B	Chl (a+b)	Car.		
Gemiza-9	0.00	2.63	1.00	3.63	1.78	2.51	0.87	3.38	1.64	84.32	11.45
	2.00	2.77	1.08	3.85	1.89	2.64	0.96	3.60	1.76	84.77	11.80
	4.00	2.92	1.22	4.14	1.97	2.72	1.07	3.79	1.85	85.28	11.83
Misr-2	0.00	2.77	1.16	3.93	1.88	2.59	0.95	3.54	1.71	85.26	11.51
	2.00	2.87	1.19	4.06	1.96	2.72	1.02	3.74	1.89	85.55	11.80
	4.00	3.02	1.34	4.36	2.01	2.88	1.13	4.01	1.94	85.79	11.94
LSD _{0.05}		0.12	0.09	0.29	0.04	0.13	0.08	0.24	0.06	0.36	0.10

Table 10. Effect of interaction between cultivars and arginine concentrations on yield and its components of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivars	Ar Conce n. (mM/l)	No of spik es/ plan t	Spik es dry weig ht (g/pl ant)	Main spik es leng th (cm)	Grai n inde x (g)	Grai n yield g/pl ant)	Stra w yield g/pl ant)	Abo ve grou nd bio mas s (g/pl ant)	Grai n yield (ton/ fed)	Stra w yield (ton/ fed)	Abo ve grou nd bio mas s (ton/ fed)	Migr atio n coef ficie nt	Harv est inde x	Cro p inde x
Gemiza-9	0.00	5.39	17.56	13.22	30.92	12.68	19.39	32.07	2.61	3.19	5.80	0.55	0.82	0.45
	2.00	5.75	19.07	15.93	32.75	14.29	21.32	35.61	2.87	3.34	6.21	0.54	0.86	0.46
	4.00	6.23	23.27	17.91	35.16	16.23	23.61	39.84	3.01	3.50	6.51	0.58	0.86	0.46
Misr-2	0.00	6.03	18.92	15.25	32.97	14.09	18.82	32.91	2.78	3.01	5.79	0.57	0.92	0.48
	2.00	6.41	20.98	16.84	34.99	15.36	20.33	35.69	3.05	3.23	6.28	0.59	0.94	0.49
	4.00	6.63	25.36	19.35	37.38	17.99	22.33	40.32	3.21	3.38	6.59	0.63	0.95	0.49
LSD _{0.05}		0.32	2.84	1.95	0.40	0.35	0.28	1.34	0.46	0.11	0.05	0.04	0.01	0.02

Table 11. Effect of interaction between water stress and arginine concentrations on growth characters of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivars	Ar Conce n. (mM/l)	Plant height (cm)	110 days after sowing								LAI
			No./plant			Dry weight (g/plant)			Flag leaf blade area	Blades area (cm ²) /plant	
			Tiller	Blade s	Spikes	Tiller + shoots	Blade s	Spikes			
Control Treat. (No skipping) stage	0.00	159.5 9	6.58	41.04	5.67	8.02	5.62	8.57	39.05	972.73	6.42
	2.00	167.9 4	6.98	44.38	6.31	8.65	5.88	8.72	44.44	1034.60	6.90
	4.00	170.4 8	7.33	47.66	6.75	9.21	6.54	9.45	47.25	1086.90	7.25
Skipping one irrig. tillering stage	0.00	137.0 0	5.67	33.50	4.46	7.00	5.23	6.45	28.71	909.41	6.06
	2.00	141.6 7	5.94	36.70	4.75	7.48	5.43	7.18	31.37	931.23	6.21
	4.00	145.5 4	6.17	38.83	5.05	8.02	5.60	7.45	35.60	956.26	6.38
Skipping one irrig. Germinati ng stage	0.00	143.6 0	5.93	37.80	4.96	7.11	5.48	7.40	33.22	939.87	6.27
	2.00	151.9 0	6.30	40.30	5.23	7.81	5.63	8.04	35.70	967.14	6.45
	4.00	154.8 7	6.45	42.50	5.44	8.56	5.88	8.84	41.20	1002.40	6.68
Skipping one irrig. Ripening stage	0.00	150.9 0	6.10	39.78	5.27	7.49	5.64	8.20	37.50	949.86	6.33
	2.00	158.4 4	6.75	41.20	5.81	8.23	5.80	8.49	40.43	1001.86	6.68
	4.00	163.4 3	7.02	44.66	6.04	8.64	6.04	9.23	44.33	1049.17	6.99
LSD _{0.05}		2.62	0.19	3.69	0.29	0.66	0.29	0.41	4.11	50.40	0.15
120 days after sowing											
Control Treat. (No skipping) stage	0.00	146.7 5	6.12	39.29	6.12	7.83	5.15	9.39	43.24	949.05	6.33
	2.00	151.9 0	6.56	42.23	6.52	8.59	5.43	9.93	45.76	1003.75	6.69
	4.00	156.3 8	6.85	45.25	7.12	8.95	5.82	10.03	50.23	1050.30	7.00
Skipping one irrig. tillering stage	0.00	129.2 0	4.93	32.82	4.64	6.24	4.66	7.96	34.68	876.37	5.84
	2.00	134.7 0	5.40	35.10	5.04	6.55	4.91	8.47	36.67	888.00	5.92
	4.00	140.6 0	5.67	37.30	5.41	7.21	4.82	8.88	39.38	921.24	6.14
Skipping one irrig. germinati ng stage	0.00	140.8 3	5.50	35.83	5.77	7.05	4.85	8.24	36.80	894.82	5.97
	2.00	146.1 0	5.77	38.18	6.04	7.54	5.12	8.91	39.50	925.02	6.17
	4.00	148.9 0	6.12	40.18	6.21	7.94	5.40	9.27	43.30	962.31	6.42
Skipping one irrig. ripening stage	0.00	144.3 0	5.77	37.70	5.90	7.30	5.02	8.81	39.20	922.56	6.15
	2.00	149.3 0	6.21	40.50	6.31	8.10	5.28	9.20	44.00	970.00	6.47
	4.00	152.5 3	6.58	42.40	6.64	8.42	5.56	9.50	47.10	998.01	6.65
LSD _{0.05}		3.69	0.29	3.00	0.37	0.55	0.20	0.23	2.87	51.62	0.21

Table 12. Effect of interaction between cultivars and water stress on growth and arginine concentrations on chemical constituents of wheat plant (Average of 2017/2018 and 2018/2109 seasons).

Water stress treats.	Ar Conce n (mM/l)	Photosynthetic pigments content per blades (mg/g) dry weight								Total Carb. (%)	Crude protein (%)
		110 days after sowing				120 days after sowing					
		Chl a	Chl b	Chl (a+b)	Car.	Chl A	Chl b	Chl (a+b)	Car.		
Control	0.00	2.90	1.22	4.12	2.10	2.78	1.09	3.87	1.91	82.63	11.90
Treat.	2.00	3.04	1.34	4.38	2.16	2.86	1.17	4.03	2.00	82.87	12.21
(No skipping)	4.00	3.25	1.47		2.23	3.04	1.23		2.09	83.17	12.27
stage				4.72				4.27			
Skipping	0.00	2.49	0.89	3.38	1.61	2.35	0.75	3.10	1.45	80.17	12.04
one irrig.	2.00	2.59	0.96	3.55	1.72	2.48	0.80	3.28	1.71	80.62	12.31
tillering	4.00	2.74	1.09		1.78	2.62	0.92		1.76	80.72	12.41
stage				3.83				3.54			
Skipping	0.00	2.66	1.02	3.68	1.76	2.51	0.87	3.38	1.58	81.39	11.94
one irrig.	2.00	2.77	1.08	3.85	1.83	2.62	0.95	3.57	1.79	81.87	12.29
Germinati	4.00	2.85	1.25		1.87	2.69	1.04		1.80	82.41	12.37
ng stage				4.1				3.73			
Skipping	0.00	2.77	1.09	3.86	1.85	2.57	0.93	3.50	1.75	82.04	11.87
one irrig.	2.00	2.87	1.16	4.03	1.99	2.75	1.08	3.83	1.83	81.86	12.25
ripening	4.00	3.05	1.32		2.07	2.86	1.21		1.91	82.81	12.32
stage				4.37				4.07			
LSD 0.05		0.12	0.11	0.31	0.06	0.16	0.09	0.28	0.07	0.40	0.12

Table 13. Effect of interaction between water stress on growth characters and arginine concentrations on yield and its components of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Water stress treats.	Ar Conc en(m M/l)	No of spike s/ plant	Spike s dry weig ht (g/pla nt)	Main spike s lengt h (cm)	Grai n inde x (g)	Grai n yield g/pla nt)	Stra w yield g/pl ant)	Above ground biomas s (g/plant)	Grai n yield (ton/ fed)	Stra w yield (ton/ fed)	Above ground biomas s (ton/fed)	Migr atio n coef ficie nt	Har ves t ind ex	Cr o p in de x
Control	0.00	6.24	22.47	16.45	35.4	16.6	21.6	38.34	2.98	3.35	6.33	0.59	0.8	0.
Treat. (No skipping) stage	2.00	6.68	23.79	19.14	37.5	18.1	23.2	41.54	3.23	3.51	6.74	0.57	0.9	0.
	4.00	6.97	27.46	20.46	39.5	20.5	26.7	47.20	3.38	3.64	7.02	0.58	0.9	0.
Skipping one irrig.	0.00	5.03	14.93	11.68	29.3	10.5	15.6	26.20	2.43	2.98	5.41	0.55	0.8	0.
tillering stage	2.00	5.50	16.78	13.31	31.0	11.7	17.6	29.45	2.57	3.18	5.75	0.57	0.8	0.
	4.00	5.78	21.21	16.06	33.7	14.2	18.9	33.22	2.66	3.27	5.93	0.64	0.8	0.
Skipping one irrig.	0.00	5.60	16.83	13.76	30.3	12.2	18.6	30.82	2.61	3.07	5.68	0.54	0.8	0.
germinatin g stage	2.00	5.87	18.85	15.51	31.9	13.8	20.4	34.26	2.96	3.19	6.15	0.55	0.9	0.
	4.00	6.24	23.29	18.51	34.9	15.8	21.8	37.67	3.12	3.36	6.48	0.62	0.9	0.
Skipping one irrig.	0.00	5.87	19.25	15.08	32.3	14.1	20.3	34.49	2.76	3.02	5.78	0.56	0.9	0.
ripening stage	2.00	6.33	20.77	15.78	34.4	15.5	21.5	37.09	3.06	3.26	6.32	0.56	0.9	0.
	4.00	6.70	25.31	19.47	36.7	17.8	24.6	42.44	3.26	3.50	6.76	0.60	0.9	0.
LSD 0.05		0.37	3.34	2.29	0.29	0.40	1.58	0.54	0.32	0.16	0.29	0.01	ns	ns

Table 14. Effect of three way interaction cultivars X water stress X arginine concentrations on growth characters of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivars	Water stress treats	Ar Conce n (mM/l)	Plant heig ht (cm)	110 days after sowing								LAI
				No./plant			Dry weight (g/plant)			Flag leaf blade area	Blade s area (cm ²)/ plant	
				Tiller	Blade s	Spike s	Tiller + shoot s	Blade s	Spike s			
Gemiza-9	Control Treat. (No skipping) stage	0.00	161.25	6.32	40.50	5.40	8.27	5.47	8.40	36.51	932.52	6.21
		2.00	159.00	6.66	44.50	6.12	8.94	5.65	8.55	40.70	1005.48	6.71
		4.00	173.10	7.29	48.33	6.50	9.40	6.55	9.32	44.66	1070.47	7.14
	Skipping one irrig. tillering stage	0.00	138.25	5.30	34.22	4.30	6.81	5.12	5.82	25.96	901.37	6.00
		2.00	144.10	5.40	36.90	4.50	7.73	5.29	6.81	28.96	926.64	6.18
		4.00	148.90	5.67	38.70	4.90	7.91	5.49	6.96	32.67	936.36	6.24
	Skipping one irrig. Germinati ng stage	0.00	146.50	5.60	37.80	4.50	7.24	5.34	6.89	32.34	914.76	6.16
		2.00	154.60	5.80	40.00	4.90	8.18	5.57	7.99	33.22	934.20	6.23
		4.00	158.20	5.90	44.00	5.17	8.68	5.82	8.55	39.16	949.64	6.46
	Skipping one irrig. Ripening stage	0.00	152.90	5.90	39.25	4.90	7.67	5.61	7.73	35.67	926.91	6.18
		2.00	161.80	6.50	40.00	5.42	8.47	5.77	8.14	37.95	963.68	6.43
		4.00	166.60	6.80	46.12	5.60	8.81	5.91	9.20	41.58	1020.70	6.81
Misr-2	Control Treat. (No skipping) stage	0.00	157.90	6.84	41.60	5.95	7.76	5.78	8.75	41.58	992.95	6.62
		2.00	166.10	7.29	44.30	6.50	8.34	6.11	8.88	48.18	1062.72	7.08
		4.00	167.80	7.40	46.80	7.00	9.01	6.53	9.58	49.83	1103.33	7.35
	Skipping one irrig. tillering stage	0.00	135.80	5.90	33.80	4.60	6.88	5.35	7.07	31.46	917.46	6.11
		2.00	139.20	6.50	36.50	5.04	7.22	5.58	7.55	33.77	935.82	6.24
		4.00	141.80	6.67	38.90	5.25	8.12	5.70	7.93	38.50	976.17	6.51
	Skipping one irrig. Germinati ng stage	0.00	139.90	6.25	37.80	5.40	6.96	5.61	7.90	34.10	964.98	6.43
		2.00	148.80	6.80	40.20	5.60	7.44	5.70	8.08	38.17	1000.08	6.66
		4.00	151.50	7.00	41.00	5.75	8.41	5.94	9.12	43.23	1035.72	6.90
	Skipping one irrig. ripening stage	0.00	148.90	6.50	40.30	5.67	7.31	5.66	8.65	39.27	972.81	6.49
		2.00	155.10	7.00	42.50	6.20	7.97	5.81	8.84	42.90	1040.04	6.93
		4.00	160.20	7.20	44.30	6.50	8.46	6.17	9.26	47.08	1076.76	7.18
LSD 0.05			2.97	0.22	4.18	0.31	0.74	0.32	0.47	4.66	57.02	0.17

Table 14. Continue

120 days after sowing												
Gemiza-9	Control	0.00	148.4	5.80	39.70	5.80	8.10	5.11	8.84	39.1	926.32	6.18
	Treat.		0							8		
	(No	2.00	154.0	6.30	42.30	6.20	8.79	5.37	9.41	41.8	949.75	6.33
	skipping)		0							0		
	stage	4.00	158.5	6.70	45.20	7.00	9.18	5.78	9.54	46.2	1020.6	6.93
			0							0	0	
	Skipping	0.00	132.6	4.70	32.80	4.40	6.33	4.55	7.60	31.9	840.24	5.61
	one irrig.		0							0		
	tillering	2.00	141.7	5.00	35.00	4.70	6.78	4.75	8.40	34.6	861.95	5.75
	stage		0							7		
		4.00	147.7	5.40	37.80	5.25	7.60	4.88	8.55	37.1	872.64	5.82
			0							8		
	Skipping	0.00	144.5	5.40	34.70	5.40	7.21	4.76	7.91	34.3	859.14	5.72
	one irrig.		0							2		
	Germinati	2.00	147.9	5.60	37.80	5.60	7.82	5.01	8.68	37.6	868.86	5.79
	ng stage		0							2		
	4.00	151.5	5.80	40.60	5.90	8.24	5.27	9.04	43.4	903.69	5.64	
		0							5			
Skipping	0.00	146.6	5.60	37.50	5.60	7.53	4.88	8.62	36.8	890.11	5.93	
one irrig.		0							5			
ripening	2.00	151.1	5.90	39.50	5.90	8.40	5.20	8.91	39.6	906.12	6.04	
stage		0							0			
	4.00	154.0	6.50	41.90	6.25	8.65	5.38	9.25	44.0	936.90	6.24	
		0							0			
Misr-2	Control	0.00	145.0	6.50	38.90	6.50	7.54	5.19	9.94	45.1	972.00	6.48
	Treat.		0							0		
	(No	2.00	150.0	6.80	42.10	6.80	8.20	5.50	10.43	49.7	1057.7	6.84
	skipping)		0							2	5	
	stage	4.00	151.0	7.00	45.40	7.20	8.71	5.87	10.52	54.2	1080.0	7.20
			0							6	0	
	Skipping	0.00	126.0	5.20	32.50	4.90	6.15	4.75	8.31	37.4	912.49	6.08
	one irrig.		0							0		
	tillering	2.00	128.0	5.80	35.10	5.40	6.31	5.06	8.55	39.8	914.04	6.09
	stage		0							2		
		4.00	135.0	5.90	36.70	5.60	6.81	5.70	9.22	41.5	969.84	6.47
			0							8		
	Skipping	0.00	137.2	5.58	36.90	6.17	6.89	4.92	8.55	39.2	930.50	6.20
	one irrig.		0							7		
	germinati	2.00	144.4	5.90	38.60	6.30	7.26	5.23	9.13	41.3	981.18	6.54
	ng stage		0							6		
	4.00	146.0	6.50	40.00	6.50	7.64	5.51	9.50	43.1	1020.9	6.80	
		0							2	2		
Skipping	0.00	142.0	5.90	37.90	6.20	7.06	5.04	9.03	41.5	955.00	6.37	
one irrig.		0							8			
ripening	2.00	147.0	6.50	41.10	6.67	7.76	5.37	9.48	48.4	1033.8	6.89	
stage		0							0	8		
	4.00	151.0	6.70	43.00	7.00	8.18	5.74	9.74	50.1	1059.1	7.06	
		0							6	0		
LSD 0.05			4.18	0.33	3.40	0.41	0.55	0.23	0.26	3.25	63.55	0.24

Table 15. Effect of three way interaction cultivars X water stress X arginine concentrations on chemical constituents of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivar s	Water stress treats.	Ar Conc en (mM/l)	Photosynthetic pigments content per blades (mg/g) dry weight								Total Carb. (%)	Crud e protei n (%)
			110 days after sowing				120 days after sowing					
			Chl a	Chl b	Chl (a+b)	Car.	Chl A	Chl b	Chl (a+b)	Car.		
Gemiza- 9	Control	0.00	2.85	1.19	4.04	2.04	2.76	1.07	3.83	1.89	85.70	11.43
	Treat.	2.00	2.99	1.29	4.28	2.08	2.82	1.13	3.95	1.93	86.03	11.73
	(No	4.00	3.20	1.44		2.21	2.91	1.23	4.14	2.05	86.32	11.76
	skipping) stage				4.64							
	Skipping	0.00	2.44	0.81	3.25	1.56	2.29	0.71	3.00	1.42	82.33	11.56
	one irrig.	2.00	2.53	0.88	3.41	1.69	2.45	0.77	3.22	1.63	83.13	11.83
	tillering	4.00	2.68	0.96		1.77	2.53	0.86	3.39	1.69	83.21	11.87
	stage				3.64							
	Skipping	0.00	2.57	0.95	3.52	1.72	2.47	0.81	3.28	1.56	84.16	11.45
	one irrig.	2.00	2.71	1.00	3.71	1.77	2.56	0.90	3.46	1.74	84.71	11.81
	germinat	4.00	2.75	1.18		1.85	2.61	0.99	3.60	1.75	85.48	11.82
	ing stage				3.93							
Skipping	0.00	2.69	1.08	3.77	1.80	2.54	0.89	3.43	1.68	85.05	11.34	
one irrig.	2.00	2.85	1.12	3.97	1.97	2.73	1.02	3.75	1.76	85.18	11.78	
ripening	4.00	3.08	1.30		2.02	2.81	1.17	3.98	1.89	86.08	11.86	
stage				4.38								
Misr-2	Control	0.00	2.94	1.25	4.19	2.17	2.79	1.11	3.90	1.94	86.09	11.45
	Treat.	2.00	3.08	1.38	4.46	2.23	2.89	1.19	4.08	2.07	86.27	11.76
	(No	4.00	3.31	1.50		2.26	3.16	1.22	4.38	2.13	86.59	11.86
	skipping) stage				4.81							
	Skipping	0.00	2.54	0.99	3.53	1.64	2.42	0.78	3.20	1.49	84.36	11.59
	one irrig.	2.00	2.65	1.02	3.67	1.74	2.51	0.82	3.33	1.77	84.48	11.84
	tillering	4.00	2.80	1.21		1.79	2.72	0.97	3.69	1.82	84.61	12.01
	stage				4.01							
	Skipping	0.00	2.76	1.09	3.85	1.80	2.56	0.94	3.50	1.61	85.05	11.51
	one irrig.	2.00	2.84	1.14	3.98	1.85	2.70	0.98	3.68	1.84	85.51	11.82
	germinat	4.00	2.93	1.32		1.88	2.75	1.07	3.82	1.85	85.85	11.97
	ing stage				4.25							
Skipping	0.00	2.85	1.10	3.95	1.89	2.59	0.95	3.54	1.80	85.35	11.47	
one irrig.	2.00	2.89	1.18	4.07	2.01	2.76	1.12	3.88	1.88	85.96	11.79	
ripening	4.00	3.01	1.34		2.10	2.89	1.25	4.14	1.94	86.10	11.92	
stage				4.35								
LSD 0.05			0.16	0.12	0.36	0.06	0.17	0.10	0.31	0.08	0.47	ns

Table 16. Effect of three way interaction cultivars X water stress X arginine concentrations on yield and its components of wheat plant (Average of 2017/2018 and 2018/2019 seasons).

Cultivars	Water stress treatments.	Ar Con. (mM /L)	No of spikes/plant	Spikes dry weight (g/plant)	Main spikes length (cm)	Grain index (g)	Grain yield g/plant)	Straw yield g/plant)	Above ground biomass (g/plant)	Grain yield (ton /fed)	Straw yield (ton /fed)	Above ground biomass (ton /fed)	Migration coefficient	Harvest index	Crop index
Gemiza-9	Control Treat. (No skipping) stage	0.00	5.86	21.57	15.54	34.24	16.21	22.37	38.58	2.89	3.47	6.36	0.56	0.83	0.45
		2.00	6.33	22.51	18.59	36.22	17.75	24.90	42.65	3.14	3.55	6.69	0.53	0.88	0.47
		4.00	7.15	26.20	20.02	38.52	19.14	27.50	46.64	3.29	3.68	6.97	0.56	0.89	0.47
	Skipping one irrig. tillering stage	0.00	4.51	13.79	11.00	28.78	9.51	16.12	25.62	2.34	3.07	5.41	0.54	0.76	0.43
		2.00	4.76	16.14	13.09	29.96	10.86	18.38	29.24	2.46	3.28	5.72	0.55	0.75	0.43
		4.00	5.34	20.02	14.85	32.64	13.88	19.36	33.24	2.56	3.33	5.86	0.60	0.77	0.44
	Skipping one irrig. germinating stage	0.00	5.50	16.31	12.65	29.15	11.53	18.25	29.78	2.55	3.14	5.69	0.55	0.81	0.45
		2.00	5.86	17.90	14.74	30.03	13.32	20.18	33.50	2.89	3.21	6.10	0.53	0.90	0.47
		4.00	6.05	22.55	17.60	33.17	14.78	22.05	36.83	2.99	3.43	6.42	0.61	0.87	0.47
	Skipping one irrig. ripening stage	0.00	5.69	18.43	13.71	31.51	13.47	20.84	34.31	2.67	3.08	6.75	0.54	0.87	0.46
		2.00	6.05	19.76	17.27	33.81	15.23	21.81	37.04	2.97	3.28	6.25	0.53	0.91	0.48
		4.00	6.38	24.24	19.14	36.06	17.10	25.54	42.64	3.20	3.54	6.64	0.57	0.90	0.47
Misr-2	Control Treat. (No skipping) stage	0.00	6.60	23.35	17.34	36.65	17.09	21.02	38.11	3.07	3.22	6.29	0.61	0.95	0.49
		2.00	6.96	25.07	19.69	38.95	18.57	22.37	40.94	3.32	3.46	6.78	0.61	0.96	0.49
		4.00	7.34	28.70	20.90	40.50	21.85	25.89	47.74	3.46	3.60	7.06	0.60	0.96	0.49
	Skipping one irrig. tillering stage	0.00	4.95	14.97	12.35	29.96	11.55	15.23	26.78	2.52	2.89	5.41	0.56	0.87	0.47
		2.00	5.50	17.40	13.53	32.10	12.65	17.01	29.66	2.68	3.07	5.75	0.59	0.87	0.47
		4.00	5.69	22.34	17.27	34.83	14.70	18.07	32.77	2.75	3.21	5.96	0.68	0.86	0.46
	Skipping one irrig. germinating stage	0.00	6.24	17.37	14.86	32.10	12.95	19.10	32.05	2.67	2.98	5.65	0.54	0.90	0.47
		2.00	6.44	19.80	16.28	33.81	14.33	20.69	35.02	3.02	3.15	6.17	0.56	0.96	0.59
		4.00	6.60	24.01	19.42	36.65	16.85	21.63	38.51	3.24	3.28	6.52	0.62	0.99	0.50
	Skipping one irrig. ripening stage	0.00	6.33	20.08	16.45	33.17	14.75	19.91	34.66	2.85	2.97	5.82	0.58	0.96	0.49
		2.00	6.79	21.63	17.88	35.10	15.89	21.25	37.14	3.14	3.24	6.38	0.58	0.97	0.49
		4.00	7.15	26.38	19.80	37.45	18.50	23.73	42.23	3.36	3.45	6.81	0.63	0.97	0.49
LSD 0.05		0.42	3.78	2.59	0.52	0.45	1.12	0.61	0.37	0.08	0.31	0.05	ns	ns	0.42

arginine. It is evidenced to state that these results are in harmony with those obtained by Ahmed, *et al.*, (2015). Also, it can be concluded that the damaging of water stress to wheat plants can be alleviated by using foliar application with 4 mM/l arginine.

DISCUSSION

Based on the obtained results, it can be that the differences between the two wheat cultivars (Gemiza-9 and Misr-2) might be due to the differences between wheat genotype their mineral elements concentration and differences in photosynthetic separations and photo-synthetic movement among wheat plant organ.

The effect of arginine concentration on yield and its components of wheat plant was respected and reached the significance level at 5%. Data showed clearly that foliar application with 2.0 mM/l arginine was significantly the number of spikes/plant; dried weight of spike per plant, the mean of spike length, grain index, grain, straw and above ground biomass/plant, above ground biomass/fed., moving coefficient, harvest index and crop index compared to the control treatment, except the increase in grain and straw yield/feddan were not achieving the significant level at 5%.

Generally, irrigation is recommended due to the highest effect on the tiller survival and this implies that development and physiological processes are critical in determining final grain yield and the water stress should be avoided at this growth stage.

Moreover, foliar spraying with 4 mM/l arginine on wheat plant caused significant increment in all yield and its component except the differences in grain and straw yield/fed., harvest index and crop index between 2 and 4 mM/l arginine that were not reaching the significant level at 5 %.

CONCLUSION

Drought reduced all yield parameters so, the application of foliar application with arginine (Ar) improved productivity of wheat plants and its components and increasing some chemicals composition of wheat grains under drought stress conditions. The Arginine foliar spray with 4.00 mM is the most effective treatment to enhance yield of wheat cv., Misr-2 plants and improve the grain quality under common irrigation compared with the other treatments. Results also indicated that treatment of wheat (*Triticum aestivum* L.) cv., Misr-2 with foliar plant with arginine as an amino acid was the most suitable significant material used to alleviate the harmful effect under drought conditions.

List of abbreviations

O.M	Organic matter
pH	Acidity
CaCO₃	Calcium Carbonates
EC	Electric Conductivity
WP	Wilting point
AW	Available water
SP	Soil penetration
FC	Field capacity
No	Number
LAI	Leaf area index
Concen.	Concentration
Chl a	Chlorophyll a
Chl b	Chlorophyll b
Car.	Carotenoids
Carb.	Carbohydrates
ns	Non-Significant
Cm	Centimeters
mM/L	Milli Mol/liter
G	Gram
Ar	Arginine
LSD	Least significant degree

Authors' contribution

Mohamed Ahmed	Writing the manuscript
	Reviewing the manuscript
	References
Magda Shalaby	Carrying out statistical analysis
	References
	Reviewing the manuscript
Reda Essa	Carrying out the experiments
	Samples analysis
	Carrying out statistical analysis
	Writing the manuscript

Conflict of Interest

The authors declared that present study was performed in absence of any conflict of interest.

ACKNOWLEDGEMENTS

The author(s) are thankful to the Agricultural Experimental Station of NRC, El-Behaira Governorate, Egypt for providing all the necessary facility to complete the paper work.

Authors contributions

All authors significantly contributed in all parts and aspects of the paper

REFERENCES

- Abd El-Monem AA (2007). Polyamines as modulators of wheat growth, metabolism and reproductive development under high temperature stress. Ph.D. Thesis, Ain Shamas Univ., Cairo, Egypt. 155 p.
- Abdel-Gawad A, El-Shouny KA, Saleh SA, Ahmed MA (1987). Partition and migration of dry matter in newly cultivated wheat varieties. Egypt J. Agron., 12(1-2):1-16.
- Abdel-Salam MS, El-Metwally IM, Abdel Lateef EM, Ahmed MA (2016). Effect of weed control and praline treatment on wheat productivity and weed nutrient removal under water stress conditions. Inter. J. of Chem. Tech. Res., 9(7):18-31.
- Abdul-Qados, A.M.S. (2010). Effect of arginine on growth, nutrient composition, yield and nutritional value of mungbean plants grown under salinity stress. Natu. & Sci., 8(7):30-42.
- Ahmed MA, Magda AF, Shalaby and El-komy M.B.A. (2015). Alleviation of water stress effects on corn by polyamine compounds under newly cultivated sandy soil conditions. Int. J. Chemtech Res., 8(12):497-508.
- Akladios SA, Abbas SM (2014). Inter simple sequence repeat (ISSR) markers and some physiological attributes of barley (*Hordeum vulgare* L.) genotypes to drought and potassium nutrition. J. Animal and Plant Sci., 24(2):620-633.
- Akladios SA, Hanafy RS (2018). Alleviation oxidative effect salt stress in white lupine (*lupinus termis* L.) plants by foliar treatment with arginine. J. Animal and Plant Sci., 23(1):165-176.
- Allen M, Lunine JI, Yung YL (1984). Correction to "The vertical distribution of ozone in the mesosphere and lower thermosphere". J. of Geophysical Res., 89.
- Bouchereau A, Aziz A, Larher F, Murting-Tanguy J (1999). Polyamines and development challenges recent development. Plant Sci., 140:103-125.
- Bremner PM, Taha MA (1966). Studies in potato agronomy: 1- the effect of variety, seed size and spacing on growth, development and yield. J. Agric. Sci., 66:241-252.
- Dubois M, Gilles KA, Hamiton JK, Rebers PA, Smith F (1956). Colorimetric method for determination of sugars and related substances. Anal. Chem., 28: 350-356.
- El-Bassiouny HM, Mostafa HA, El-Khawass SA, Hassanein RA, Khalil SI, Abd El-Monem AA (2008). Physiological responses of wheat plant to foliar Treatments with Arginine or Putrescine. Austr. J. of Basic and Appl. Sci., 2(4):1390-1403.
- EL-Metwally IM, Abdel-Raouf RE, Ahmed MA, Mounzer O, Al-Arcon JJ, Abdel-Hamed MT (2015). Response of wheat (*Triticum aestivum* L.) crop and broad-leaved weeds to different water requirements and weed management in sandy soil. Agric. (Poľnohospodárstvo), 61(1):22-32.
- El-Shintinawy F, Hassanein RA (2001). The role of polyamine precursors, arginine, ornithine or methionine in ameliorating the inhibitory effect of NaCl on wheat plant. Egypt J. Biotechnol., 9:328-340.
- FAO (2009). High level expert forum-how to feed the world in 2050. Economic and Social Development Department, Food and Agricultural Organization of the United Nations, Rome.
- Galston AW, Kaur-Sahney (1990). Polyamines in plant physiology. Plant Physiol. 94: 406.
- Hassanain MS, Ahmed MA, Magda AF, Shalaby and Amal, G. Ahmed (2014). Partition and migration of dry matter in newly cultivated wheat cultivars in Egypt. Middle East J. of Agric. Res., 3(2):353-362.
- Hassanein RA, Khalil SI, El-Bassiouny HMS, Mostafa HAM, El-Khawass SA, Abd El-Monem AA (2008). Protective role of exogenous arginine or putrescine treatments on heat shocked wheat plant. 1st International Conference on Biological and Environmental Sciences, Hurghada, Egypt, March 13-16.
- Jackson ML (1973). Soil Chemical Analysis. Prentice- Hall, Inc. Englewood Cliffs. N.J.
- Kramer PJ, Boyer JS (1995). Water Relations of Plants and Soils. Academic Press, San Diego.
- Magda AF, Shalaby Ahmed MA, Ebtihal M. Abd Elhamid (2014). Effect of the Plant Growth Promoter Brassinolide on Barely Cultivars (*Hordeum vulgare* L.) Grown Under Sandy Soil Conditions. Middle East J. of Agric. Res., 3(2):282-287.
- Magda AF, Shalaby and Salem A.K.M. (2014). Growth and productivity barley cv. (Giza-129) cultivar as affected by putrescine and spermidine under stress conditions. Life Sci, J., 11(95):666-671.
- Magda AF, Shalaby, Ahmed MA, Khater MA (2018). Physiological response of some barley cultivars to foliar treatment with arginine under water stress conditions. Middle East J. of Agric. Res., 7(3):1102-1123.
- Mo H, Pua C (2002). Up-regulation of arginine decarboxylase gene expression and accumulation of polyamines in mustard (*Brassica juncea*) in response to stress. Physiol. Planta., 114(3):439-449.
- Nasibi F, Manouchehri Kalantariand Kh, Barand A (2104). Effect of seed pre-treatment with L-arginine on improvement of seedling growth and alleviation of oxidative damage in canola plants subjected to salt stress. Iranian J. of Plant Physiology 5(1):1217-1224.
- Pregl F (1945). Quantitative Organic Microanalysis. 4th ed. J.A. Churchill Ltd. London, p.126-129.
- Richards LA (1954). Diagnosis and Improvement Saline and Alkaline Soils. U.S. Dep. Agr. Handbook 60.
- Snedecor GW, Cochran WG (1990). Statistical Methods. The Iowa State Univ. 7th Ed. pp. 507.
- Van Wettstein D (1957). Chlorophyll lethal faktoren under submikro skopoch formwechsel der plastiden. Exp. cell. Res., (12): 427-433.
- Velikova V, Yordanov I, Edre va A (2000). Oxidative stress and some antioxidant systems in acid rain-treated bean plants protective role of exogenous polyamines. Plant Sci., (115):59-66.
- Watson DJ (1952). The physiological basis of variation in yield. Advances Agron., (4):101-145.