

Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:



<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr



**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2020 (Volume 60): 450 €
<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2018): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme
(Labex Agro: ANR-10-LABX-0001-01)



Acarologia is under free license and distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Phytoseiidae of Vietnam (Acari: Mesostigmata) with description of a new species

Serge Kreiter^a, Marie-Charlotte Bopp^a, Martial Douin^a, Duc Tung Nguyen^b, Kris Wyckhuys^{c,d,e}

^a Montpellier SupAgro, UMR CBGP INRA/ IRD/ CIRAD/ SupAgro, 755 Avenue du Campus Agropolis (Baillarguet), CS 30016, 34988 Montferrier-sur-Lez cedex, France.

^b Vietnam National University of Agriculture, Trau Quy, Gia Lam, Hanoi, Vietnam.

^c Chrysalis Consulting, Hanoi, Vietnam.

^d Institute of Plant Protection, China Academy of Agricultural Sciences, Beijing, China.

^e School of Biological Sciences, University of Queensland, Brisbane, Australia.

Original research

ABSTRACT

Vietnam is a large country of Southeast Asia and a globally recognized hotspot of biodiversity. Until recently, only limited surveys had been carried out on the resident Phytoseiidae fauna and presence of 12 species had been documented, with 11 belonging to the sub-family Amblyseiinae and 1 to the sub-family Typhlodrominae. Here, we present results from 2017 field surveys and add a total of 8 new country records, one of which a newly-described species. At least 5 species (*i.e.*, *Neoseiulus longispinosus*, *Paraphytoseius cracentis*, *P. orientalis*, *Amblyseius cinctus* and *A. herbiculus*) are well-known biological control agents (BCA), while two species (*Euseius ovaloides* and *Gynaeseius liturivorus*) carries ample potential as BCA agents but necessitates further study. In addition to the intrinsic value of phytoseiid mite biodiversity in tropical environments, demonstration of the natural occurrence of efficient BCAs in Southeast Asian countries such as Vietnam is of great agricultural and commercial interest.

Keywords survey; collection; taxonomy; systematics; Vietnam

Zoobank <http://zoobank.org/9062B6C5-A8EF-4DB4-B5DA-30ACCA8B1D4A>

Introduction

Received 21 November 2019
Accepted 11 February 2020
Published 28 February 2020

Corresponding author
Serge Kreiter:
serge.kreiter@supagro.fr

Academic editor
Faraji, Farid

DOI
[10.24349/acarologia/20204362](https://doi.org/10.24349/acarologia/20204362)
ISSN 0044-586X (print)
ISSN 2107-7207 (electronic)

© Copyright
Kreiter S. et al.

Distributed under
Creative Commons CC-BY 4.0

The Phytoseiidae family is widespread all over the world and consists of about 2,500 valid species dispatched in three sub-families and 94 genera (Demite *et al.* 2014, 2020). Several species in the family are important natural enemies of phytophagous mites, insect eggs and small (or immature) insects in natural habitats, arable field crops and protected crops globally (McMurtry and Croft 1997; McMurtry *et al.* 2013). Despite the extensive faunistic surveys carried out for more than 70 years worldwide, the fauna of certain countries and ecosystems remains little explored (Tixier and Kreiter, 2009; Kreiter *et al.* 2020a). Consequently, it is important to conduct surveys in these poorly-investigated areas and gain more information on resident biodiversity, especially in hotspots of biodiversity. Furthermore, in-country surveys of Phytoseiidae can signal local presence of known biological control agents (BCA) and introduce new potential BCAs, especially in the context of international and country regulations including the Nagoya Protocol on Access and Benefit Sharing (Mason *et al.* 2018).

Most areas of Southeast Asia constitute globally-important hotspots of biodiversity (Myers, 1988). These geographical entities are of immediate importance to biodiversity conservation, regularly hold high endemism and have been subject to considerable loss of natural habitats

in recent years (Myers *et al.* 2000). The characterization of the phytoseiid mite diversity in these areas is thus directly contributing to this general topic of conservation. Though Vietnam is a large area in Southeast Asia, scant information exists on the resident phytoseiid mite fauna restricted to only five species listed in the world Phytoseiidae database (Demite *et al.* 2020) [*Amblyseius matinikus* Schicha & Corpuz-Raros, *Graminaseius polisensis* (Schicha & Corpuz-Raros), *Neoseiulus longispinosus* (Evans), *Paraphytoseius orientalis* (Narayanan, Kaur & Ghai), and *Scapulaseius asiaticus* (Evans)], based on only one paper (Nguyen *et al.* 2016) concerning a survey in one city of Vietnam, *i.e.*, Ho Chi Minh City.

Actually, 12 species are known from Vietnam, the previous five and 7 additional species: *Amblyseius tamatavensis* Blommers, *Proprioseiulus dahonagnas* (Schicha & Corpuz-Raros) and an undetermined *Typhlodromus* sp., *Paraphytoseius cracentis* (Corpuz & Rimando), *Amblyseius largoensis* (Muma), *Neoseiulus californicus* (McGregor) and *Proprioseiopsis lenis* (Corpuz & Rimando) (Nguyen & De Clercq, 2018; Nguyen & Dao, 2019; Nguyen *et al.* 2019a; Nguyen *et al.* 2019b).

This paper constitutes the sixth contribution to Vietnam's Phytoseiidae Fauna and reports results of an additional field surveys realised in 2017.

Material and Methods

The survey took place in Yen Bai province, a mountain province located at the northwest part of northern-central Vietnam.

Plant-inhabiting mites were collected in several cultivated plots planted with *Manihot esculenta* and on surrounding wild plants in two villages of Vietnam (Vinh Kien village: 21.74°N, 105.08°E, Yen Binh district, Yen Bai province and Mau Dong village: 21.91°N, 104.64°E, Van Yen district, Yen Bai province).

Mites were directly collected on leaves with a fine brush.

Collected mite specimens were then transferred with a fine brush into small plastic vials containing 70° ethanol. Next, mites were mounted on slides using Hoyer's medium and all identified using a phase contrast and Differential Interference contrast microscope (DMLB, Leica Microsystèmes SAS, Nanterre, France). Characters of specimens were measured using a graduate eyepiece (Leica, see above). Chant and McMurtry's (1994, 2007) concepts of the taxonomy of the family Phytoseiidae and the world catalogue database of Demite *et al.* (2014, 2020) were used to identify mite species. In this paper, the setal nomenclature system adopted was that of Lindquist & Evans (1965) and Lindquist (1994), as adapted by Rowell *et al.* (1978) for the dorsum and by Chant & Yoshida-Shaul (1991) for the venter. The idiosomal setal pattern follows Chant & Yoshida-Shaul (1992). The notation for solenostomes and poroids is based on Athias-Henriot (1975). Numbers of teeth on the fixed and movable cheliceral digits do not include the respective apical tooth. Setae not referred to in the Results section should be considered as absent. All measurements are given in micrometers (μm) in the text and figures and presented as the mean in bold followed by the range in parenthesis. All collected individuals were measured if no precision is given. New measurements added with existing literature records are marked in bold in the tables.

Specimens are deposited in the mite collections of Montpellier SupAgro conserved in UMR CBGP INRA/IRD/CIRAD/SupAgro. All specimens were identified, except for few single males which were not taken into account. The following abbreviations are used in this paper for morphological characters: dsl = dorsal shield length just under *j1* to just below *J5*; dsw = dorsal shield width at the level of *s4*; Per. ext.: peritreme extension; gd = solenostome; Z4 ser., Z5 ser. = Z4, Z5 serrated (if Z4 and Z5 without ser. = not serrated); Metastern. sh. = presence or absence of metasternal shield; lisl = Largest inguinal sigilla (= "metapodal plate") length; lisw = Largest inguinal sigilla (= "metapodal plate") width; sisl = smallest inguinal sigilla (= "metapodal plate") length; gensl = genital shield length; gensw post. cor. = genital shield width posteriorly; vsl = ventrianal shield length; vsw *ZV2* & vsw anus = ventrianal shield width at

ZV2 level and at paranal setae level; Dist. solen. vas = distance between solenostomes on the ventrianal shield; scl = total spermatheca length (calyx + neck or cervix + atrium); scw = calyx width; FD = fixed digit; Fdl = fixed digit length; MD = movable digit; Mdl = movable digit length; No teeth Fd = number of teeth on the fixed digit; No teeth Md = number of teeth on the movable digit; Shaft = length of the shaft of spermatodactyl; BCA = Biological control agents; FCI = French Caribbean Islands; VCW: various countries in the world; im. = immature. The following abbreviations are used in this paper for institution: CBGP = Centre de Biologie pour la Gestion des Populations; CIRAD = Centre International de Recherche Agronomique pour le Développement; INRA = Institut National de la Recherche Agronomique; IRD = Institut de Recherche pour le Développement; MSA = Montpellier SupAgro, France; UMR = Unité Mixte de Recherche.

Results and Discussion

A total of 14 species were found: 11 belonging to Amblyseiinae and to genera *Amblyseius*, *Euseius*, *Graminaseius*, *Gynaeseius*, *Neoseiulus*, *Paraphytoseius* and *Scapulaseius*; Three species are belonging to Phytoseiinae and to genus *Phytoseius*. No species of the sub-family Typhlodrominae has been found.

Thirteen species had been previously described among which 8 are new records for Vietnam, while 1 species is new to science. All results concerning the 14 species are given hereunder.

Subfamily Amblyseiinae Muma

Amblyseiinae Muma, 1961: 273.

Genus *Neoseiulus* Hughes

Neoseiulus Hughes, 1948: 141.

Neoseiulus longispinosus (Evans)

Typhlodromus longispinosus Evans 1952: 413; Evans 1953: 465; Womersley 1954: 177; Ehara 1958: 55.

Typhlodromus (Amblyseius) longispinosus, Chant 1959: 74.

Amblyseius longispinosus, Corpuz & Rimando 1966: 129; Schicha 1975: 103.

Neoseiulus longispinosus, Moraes *et al.* 1986: 85; 2000: 245; Chant & McMurtry 2003a: 37; Moraes *et al.* 2004a: 129; Chant & McMurtry 2007: 29.

Specimens examined: Muoi in P4 plot (aaSL 66 m, lat. 21°54'41" N, long. 104°38'7" E), 1 ♀ on *Styrax tonkinensis* (Styracaceae), 12/V/2017; in P8 plot (aaSL 66 m, lat. 21°54'46" N, long. 104°38'9" E), 1 im. on *Bambusodae* sp. (Poaceae), 17/V/2017; in P1 plot (aaSL 66 m, lat. 21°54' N, long. 104°38' E), 1 ♀ an in plot SP5 plot (aaSL 66 m, lat. 21°54' N, long. 104°38' E), 3 ♀♀, on *Manihot esculenta* (Euphorbiaceae), 17/V/2017; in P11 plot (aaSL 66 m, lat. 21°54'30" N, long. 104°38'14" E), 1 im. on *Mallotus canii* (Euphorbiaceae), and 1 ♀ on *Passiflora foetida* (Passifloraceae), 31/V/2017.

Remarks: this species is distributed in many countries of the world, mainly in tropical areas (Moraes *et al.* 2000; Mailloux *et al.* 2010; Kreiter *et al.* 2013, 2018 a, c; Demite *et al.* 2020). It was rarely encountered before in surveys made in Guadeloupe, Martinique and La Réunion but was recently more captured when screening low vegetation constituting companion plants in citrus orchards (Mailloux *et al.* 2010; Kreiter *et al.* 2013, 2018c; Le Bellec *et al.*, unpub. data). This species appears to be more common on grasses of the lower vegetation with populations of tetranychid mites. *Neoseiulus longispinosus*, a type II phytoseiid predatory mite like *N. californicus* (McMurtry *et al.* 2013), is increasingly being considered for the control of *Eutetranychus*, *Oligonychus*, and *Tetranychus* spider mites in Asia (Nusartlert *et*

Table 1 Character measurements of adult females of *Neoseiulus longispinosus* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam	La Réunion	Comoros	Holotype Indonesia	Sri Lanka	Taiwan	Thailand
	6, this study	18	1	1	3	?	3
Dsl	321 (305 – 350)	320 (278 – 348)	380	332	321 (313 – 338)	-	330 (315 – 340)
Dsw at s4 level	170 (130 – 213)	183 (150 – 205)	192	173	187 (175 – 208)	-	186 (178 – 195)
j1	19 (15 – 23)	18 (15 – 20)	20	14	18 (17 – 19)	17	19 (18 – 21)
j3	57 (50 – 65)	62 (53 – 73)	69	51	62 (61 – 64)	58	61 (56 – 68)
j4	57 (53 – 60)	55 (48 – 60)	70	49	58 (56 – 60)	59	58 (54 – 62)
j5	65 (63 – 70)	66 (58 – 73)	78	59	70 (69 – 71)	-	69 (65 – 72)
j6	69 (65 – 73)	73 (65 – 83)	78	64	70 (68 – 72)	-	71 (68 – 73)
J2	76 (73 – 78)	74 (68 – 83)	88	66	77 (75 – 79)	-	74 (68 – 81)
J5	8	8 (6 – 10)	10	10	8	-	9 (7 – 12)
r3	61 (55 – 65)	59 (48 – 78)	75	54	55 (55 – 56)	-	61 (50 – 67)
R1	56 (55 – 58)	61 (54 – 80)	70	58	60 (59 – 62)	-	63 (55 – 70)
s4	81 (73 – 83)	82 (70 – 88)	-	75	82 (80 – 83)	72	81 (76 – 85)
S2	73 (63 – 80)	75 (68 – 85)	88	67	73 (70 – 79)	70	73 (68 – 80)
S4	61 (55 – 68)	56 (43 – 68)	63	49	59 (57 – 62)	58	61 (56 – 68)
S5	18 (15 – 20)	17 (15 – 18)	18	15	21 (19 – 23)	17	20 (18 – 25)
z2	64 (60 – 73)	67 (56 – 75)	75	58	69 (68 – 70)	62	66 (62 – 70)
z4	69 (63 – 75)	73 (65 – 80)	78	58	73 (73 – 75)	65	71 (68 – 75)
z5	31 (28 – 33)	27 (23 – 33)	38	-	32 (32 – 38)	-	31 (25 – 36)
Z1	74 (68 – 78)	75 (65 – 85)	83	67	77 (76 – 78)	-	75 (71 – 78)
Z4	70 (60 – 73)	70 (63 – 75)	78	68	72 (71 – 73)	65	71 (65 – 76)
Z5	76 (63 – 80)	81 (73 – 88)	85	72	80 (80 – 81)	70	78 (72 – 81)
st1-st1	48 (45 – 48)	47 (43 – 50)	48	-	-	-	-
st2-st2	57 (55 – 60)	58 (55 – 60)	55	-	53 (50 – 55)	-	59 (57 – 60)
st3-st3	70 (63 – 73)	71 (68 – 78)	60	77	-	-	-
st1-st3	54 (53 – 55)	57 (55 – 60)	73	62	55 (53 – 56)	-	63 (57 – 85)
st4-st4	69 (60 – 75)	71 (63 – 83)	85	-	-	-	-
Gensl	116 (103 – 125)	-	-	-	-	-	-
st5-st5	56 (50 – 60)	54 (50 – 58)	58	-	53 (51 – 54)	-	61 (55 – 64)
Gensw post.corn.	75 (70 – 80)	-	-	-	-	-	-
Lisl	29 (25 – 33)	25 (20 – 30)	23	-	-	-	-
Lisw	2	3 (3 – 6)	3	-	-	-	-
Sisl	14 (13 – 18)	12 (10 – 18)	18	-	-	-	-
Vsl	110 (95 – 125)	114 (93 – 125)	125	97	106 (103 – 111)	-	123 (120 – 125)
Vsw ZV2	91 (75 – 100)	86 (80 – 95)	90	87	91 (89 – 93)	-	97 (95 – 100)
Vsw anus	77 (75 – 78)	69 (67 – 73)	78	-	75 (73 – 77)	-	-
JV5	59 (50 – 68)	62 (55 – 70)	73	-	-	-	-
StIV	74 (70 – 75)	79 (75 – 83)	75	80 – 87	68 (68 – 70)	74	74 (72 – 77)
Scl	22 (18 – 23)	24 (13 – 35)	25	30	21 (20 – 21)	17	19 (17 – 22)
Scw	5 (5 – 6)	4 (3 – 10)	5	4	-	-	-
Fdl	23 (20 – 25)	24 (19 – 28)	23	-	22 (21 – 22)	-	23 (22 – 25)
No teeth Fd	-	6	-	-	5	-	-
Mdl	25	24 (23 – 30)	23	-	25 (23 – 25)	-	26 (25 – 27)
No teeth Md	-	2	-	-	2	-	-

Sources of measurements – La Réunion: Kreiter *et al.* 2020 accepted; Grande Comore Island of Comoros Archipelago: Kreiter *et al.* (2018b) (a mistake remains in the paper, 1 single female measured instead of 8 as indicated); Holotype Indonesia: Schicha (1975); Sri Lanka: Moraes *et al.* (2004b); Taiwan: Tseng (1983); Thailand: Oliveira *et al.* (2012); - : not provided.

al. 2011). Its behaviour, feeding, development, prey consumption, cannibalism and intra-guild predation have been extensively studied by several authors (see for example Luong *et al.* 2017), with the ultimate aim to enhance its use for biological control. *Neoseiulus longispinosus* is a well-known BCA sold in several countries and commercially-available for management of spider mites. Under controlled laboratory conditions, *N. longispinosus* exhibits potential as a BCA against the citrus red spider mite *P. citri* (Huyen *et al.*, 2017).

This species was already known from Vietnam and reported as a local BCA in earlier survey work (Nguyen *et al.* 2016) though no measurements were given. Measurements of female specimens collected during this study are given in table 1 and compared with measurements obtained from specimens collected in other countries of Asia and Islands of the Indian Ocean. Measurements match with most of the previous measurements, especially with those obtained on specimens from La Réunion Island. The setal lengths of the present females are longer than those obtained from the holotype, which are in the lower part of the range of data obtained for each seta (probably measurements were obtained from a very small size female).

Genus *Paraphytoseius* Swirski & Shechter

Paraphytoseius Swirski & Shechter 1961: 113; Moraes *et al.* 1986: 104.

Chant & McMurtry 2003b: 216; Moraes *et al.* 2004a: 160; Chant & McMurtry 2007: 49.

Amblyseius (*Paraphytoseius*), Ueckermann & Loots 1987: 221.

Amblyseius (*Ptenoseius*), Pritchard & Baker 1962: 295.

Proprioseius (*Paraphytoseius*), Karg 1983: 302.

Ptenoseius, Schuster & Pritchard 1963: 198.

Paraphytoseius cracentis (Corpuz & Rimando)

Ptenoseius cracentis Corpuz & Rimando 1966: 115.

Amblyseius (*Paraphytoseius*) *multidentatus*, misidentification (S5 illustrated in figures): Ehara & Bhandhu Falck 1977: 79 as mentioned in Ehara *et al.* 2000: 114.

Paraphytoseius cracentis, Moraes *et al.* 1986: 104; Chant & McMurtry 2003b: 220; Moraes *et al.* 2004a: 160; Chant & McMurtry 2007: 53.

Specimens examined: Muoi, in P3 plot (aasl 66 m, lat. 21°54'40" N, long. 104°38'13" E), 1 ♀ + 1 ♂ on *Chromolaena odorata* (Asteraceae) and 1 ♀ on *Rubus alceifolius* (Rosaceae), 11/V/2017; in P5 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'22" E), 3 ♀♀ + 1 ♂ + 1 im. on *C. odorata* (Asteraceae), 1 ♀ + 1 ♂ on *Ficus hispida* (Moraceae) and 1 ♀ + 1 ♂ on *Artemisa vulgaris* (Asteraceae), 13/V/2017; in P6 plot (aasl 66 m, lat. 21°54'55" N, long. 104°38'22" E), 1 ♀ on *Ficus hispida* (Moraceae) and 1 ♀ on *Curculigo orchoides* (Hypoxidaceae), 15/V/2017; in P7 plot (aasl 66 m, lat. 21°54'51" N, long. 104°38'22" E), 1 ♀ on *F. hispida* (Moraceae) and 2 ♀ + 2 ♂ on *Urena lobata* (Malvaceae), 16/V/2017; in P8 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'9" E), 2 ♀♀ on *C. odorata* (Asteraceae) and 1 ♀ on *R. alceifolius* (Rosaceae), 16/V/2017; in plot PM1 plot (aasl 66 m, lat. 21°54' N, long. 104°38' E), 2 ♀♀ on an unknown host plant, 17/V/2017; in P9 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'18" E), 1 ♀ on *U. lobata* (Malvaceae) and 6 ♀♀ + 1 ♂ on *C. odorata* (Asteraceae), 18/V/2017; in P10 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'12" E), 2 ♀♀ + 1 ♂ on *C. odorata* (Asteraceae), 18/V/2017; in P11 plot (aasl 66 m, lat. 21°54'30" N, long. 104°38'14" E), 4 ♀♀ + 1 ♂ on *F. hispida* (Moraceae), 1 ♂ + 1 im. on *Bidens pilosa* (Asteraceae), 4 ♀♀ + 1 ♂ on *C. odorata* (Asteraceae), 1 ♂ on *R. alceifolius* (Rosaceae) and 2 ♀♀ + 1 ♂ on an unknown host plant, 31/V/2017; in P15 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 2 ♀♀ on *F. hispida* (Moraceae), 6 ♀♀ + 2 ♂♂ on *U. lobata* (Malvaceae) and 1 ♀ on *Ch. odorata* (Asteraceae), 03/VI/2019; in P4 plot (aasl 66 m, lat. 21°54'41" N, long. 104°38'7" E), 1 ♀ on *Xanthium strumarium* (Asteraceae), 18/VII/2017; in P5 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'22" E), 1 ♀ on *U. lobata* (Malvaceae), 19/VII/2017; in P6 plot (aasl 66 m, lat. 21°54'55" N, long. 104°38'22" E), 3 ♀♀ on *X. strumarium* (Asteraceae), 20/VII/2017; in P16

plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 1 ♂ on *Ch. odorata* (Asteraceae) and 1 ♀ on *U. lobata* (Malvaceae), 22/VII/2017; in P10 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'12" E), 1 ♀ + 1 ♂ on *Tephrosia candida* (Fabaceae) and 1 ♀ on *Morus alba* (Moraceae), 25/VII/2017; in P14 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 1 ♀ on *X. strumarium* (Asteraceae), 27/VII/2017; in P12 plot (aasl 66 m, lat. 21°54'51" N, long. 104°38'45" E), 2 ♀♀ + 1 ♂ on *U. lobata* (Malvaceae), 29/VII/2017.

Remarks: the predatory mite *P. cracentis* is a natural enemy of thrips found on vegetables such as green bean, cucumber, green squash, eggplant, and pepper in the Red River Delta in Vietnam. *Paraphytoseius cracentis* is feeding on *Frankliniella occidentalis* Pergande (Thysanoptera: Thripidae) and *Carpoglyphus lactis* (L.) (Acari: Acaridae) (Nguyen and De Clercq 2018, paper in which this species was mentioned for the first time for Vietnam. The intrinsic rate of natural increase of *P. cracentis* fed on *C. lactis* (0.215 female/female/day) was however significantly higher than that of mites fed on thrips (0.189). Nguyen and De Clercq (2018) conclude that *P. cracentis* can sustain and increase its population when fed on

Table 2 Character measurements of adult females and males of *Paraphytoseius cracentis* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam (12)	Japan	♀ Papua New Guinea (3 for Dsl & Dsw, 7-8 for other characters)	Philippines	Taiwan	Thailand	Vietnam (12)	Philippines	Taiwan	Thailand 1	Thailand 2
	12, this study		2	2	?	10	12, this study	1	?	5	3
Dsl	293 (273 – 303)	281 (3)	268	266 – 281	264 (251 – 279)	292 (265 – 310)	238 (223 – 268)	220	216 (214 – 219)	320	243 (235 – 255)
Dsw at s4 level	145 (130 – 166)	155 (3)	146	100 – 103	95 (92 – 96)	158 (150 – 168)	116 (103 – 125)	95	86 (85 – 87)	160	143 (140 – 147)
j1	34 (30 – 35)	35 (35 – 37)	33	32 – 38	33 (32 – 34)	33 (33 – 39)	25 (23 – 28)	24	25 (24 – 26)	25	26 (24 – 27)
j3	87 (83 – 93)	87 (85 – 89)	90	84 – 91	92 (87 – 99)	92 (82 – 98)	61 (58 – 68)	68	66 (63 – 70)	59	59 (58 – 60)
j4	5 (5 – 6)	4 (4 – 5)	4	4 – 5	Minute	5 (4 – 6)	4 (3 – 5)	2 – 4	–	4	5
j5	5	4	5	4	Minute	5 (4 – 6)	5 (4 – 5)	2 – 4	–	5	5
j6	6 (5 – 7)	6 (6 – 7)	6	6	minute	7 (5 – 9)	7 (5 – 8)	2 – 4	–	6	6 (6 – 7)
J5	5	4 (4 – 5)	5	2	minute	5 (4 – 6)	4 (3 – 5)	2 – 4	–	4	3
r3	45 (43 – 48)	46 (45 – 47)	37	40 – 42	47 (42 – 53)	48 (44 – 50)	28 (25 – 30)	24	31 (28 – 33)	27	30 (27 – 33)
R1	30 (28 – 33)	34 (33 – 35)	23	26 – 33	30 (27 – 36)	32 (28 – 38)	16 (13 – 20)	12	16 (15 – 17)	14	15 (15 – 16)
s4	120 (113 – 128)	123 (121 – 124)	117	115 – 127	125 (120 – 132)	128 (119 – 134)	84 (78 – 88)	88	89 (80 – 97)	81	88 (85 – 89)
S5	14 (11 – 18)	13 (13 – 14)	14	8 – 13	minute	14 (11 – 17)	8 (6 – 12)	8	–	6	7 (6 – 7)
z2	10 (9 – 11)	10 (9 – 10)	10	8 – 12	minute	10 (8 – 13)	10 (8 – 11)	8	–	9	10 (10 – 11)
z4	9 (8 – 10)	9	10	8 – 12	minute	9 (8 – 12)	9 (8 – 10)	8	–	8	10 (9 – 11)
z5	5	5	4	4	minute	5 (4 – 5)	3 (3 – 4)	2 – 4	–	5	5
Z1	8 (6 – 8)	7	9	7 – 8	minute	7 (6 – 9)	7 (5 – 8)	8	–	6	6 (6 – 7)
Z4	75 (70 – 80)	77 (75 – 78)	71	74 – 82	77 (70 – 83)	80 (75 – 85)	48 (43 – 53)	54	49 (48 – 49)	47	52 (50 – 53)
Z5	100 (90 – 110)	106 (104 – 107)	95	91 – 121	110 (99 – 119)	102 (92 – 105)	54 (50 – 58)	65	55 (48 – 61)	52	59 (55 – 62)
st1-st1	69 (63 – 75)	–	–	–	–	–	56 (53 – 60)	–	–	–	–
st2-st2	68 (65 – 70)	–	–	84 – 85	57 (64 – 69)	73 (71 – 76)	60 (55 – 63)	–	–	–	–
st3-st3	79 (75 – 80)	–	–	–	–	–	68 (63 – 70)	–	69 (68 – 70)	–	–
st1-st3 ♀ / st1-st5 ♂	67 (63 – 70)	–	–	70 – 74	78 (74 – 84)	70 (65 – 75)	115 (110 – 133)	–	115 (113 – 117)	–	–
st4-st4	86 (78 – 90)	–	–	–	–	–	59 (55 – 63)	–	–	–	–
gensl	112 (100 – 123)	–	–	–	–	–			Not applicable		
st5-st5	83 (80 – 85)	–	–	84	86 (78 – 90)	89 (84 – 92)	50 (45 – 63)	–	–	–	–
gensw post. corn.	91 (85 – 95)	–	–	–	–	–			Not applicable		
Lisl	37 (33 – 43)	–	–	24	31 (24 – 36)	–			Not applicable		
Lisw	1	–	–	–	–	–					
Vsl	92 (75 – 103)	–	–	94 – 96	97 (95 – 105)	112 (104 – 120)	91 (83 – 100)	86	95 (89 – 101)	–	102 (101 – 102)
Vsw ZV2	62 (55 – 70)	–	–	58	58 (51 – 64)	66 (53 – 74)	111 (100 – 128)	129	110 (109 – 112)	–	124 (120 – 126)
Vsw anus	51 (48 – 55)	–	–	–	–	–	64 (60 – 70)	–	–	–	–
JV5	74 (60 – 85)	69 (67 – 70)	–	68 – 77	75 (71 – 84)	–	20 (15 – 25)	–	20 (18 – 20)	21	–
SgeII	13 (11 – 15)	–	–	10 – 13	–	14 (12 – 16)	13 (10 – 15)	–	–	–	12
Erected setae femur IV	12 (10 – 13)	16 (15 – 16)	–	–	–	–	8 (7 – 9)	–	–	–	–
SgeIV	30 (28 – 33)	31 (30 – 32)	22	24 – 28	–	31 (27 – 33)	21 (18 – 23)	20	–	21	21 (20 – 22)
StIV	38 (35 – 40)	41 (40 – 42)	34	38 – 39	–	40 (38 – 42)	29 (25 – 30)	30	–	29	30
StIV	46 (43 – 50)	49	38	40 – 41	–	47 (43 – 52)	36 (33 – 38)	36	–	35	37 (35 – 40)
SttIV	45 (43 – 49)	44 (42 – 45)	30	43 – 46	–	47 (45 – 58)	36 (33 – 38)	37	–	36	38 (37 – 40)
ScI	3 (2 – 4)	–	–	?	–	8 (7 – 10)			Not applicable		
Scw	10 (9 – 10)	–	–	–	–	–	21 (20 – 23)	–	–	–	19 (18 – 20)
Fdl	31 (30 – 33)	–	–	29	–	28 (26 – 30)	9 (9 – 10)	12	8	–	–
No teeth Fd	11 (11 – 12)	–	–	12	11	–	22 (20 – 25)	–	–	–	23 (21 – 24)
Mdl	33 (33 – 35)	–	–	33	–	34 (32 – 37)	1	3	1	–	–
No teeth Md	3 (2 – 3)	–	–	3	3	–	15 (13 – 20)	12	–	–	16 (15 – 17)
Shaft			Not applicable				5 (4 – 6)	6	–	–	–
Toe											

Sources of measurements – For ♀♀: Japan: Ehara et al. (2000); Papua New Guinea: McMurry & Moraes (1985); Philippines: Schicha & Corpuz-Raros (1985); Taiwan: Ho & Lo (1989); Thailand: Oliveira et al. (2012). For ♂♂: Philippines: Schicha & Corpuz-Raros (1985); Taiwan: Ho & Lo (1989); Thailand 1: (misidentified as *Paraphytoseius multidentatus* s. setae S5 overlooked as mentioned by Ehara et al. 2000) Ehara & Bhandhufalck (1977); Thailand 2: Oliveira et al. (2012). - : not provided.

F. occidentalis, and that *C. lactis* can be used for mass-rearing of this predatory mite in the perspective to use it in biological control.

This species was already known from Vietnam and reported locally as a BCA from earlier survey work (Nguyen and De Clercq 2018) though no measurements were provided. Measurements are given here in table 2 for females and males and compared with those obtained from specimens collected in other countries of Asia.

All measurements collected during this study match well with most of the previous measurements, especially with measurements of specimens from Thailand.

***Paraphytoseius horrier* (Pritchard & Baker)**

Amblyseius (Ptenoseius) horrier Pritchard & Baker, 1962: 295.

Amblyseius horrier, Meyer & Rodrigues 1966: 30.

Amblyseius (Paraphytoseius) horrier, van der Merwe 1968: 169.

Proprioseius (Paraphytoseius) horrier, Karg 1983: 302.

Paraphytoseius horrier, Moraes *et al.* 1986: 105; Beard 2001: 84; Chant & McMurtry 2003b: 37; Moraes *et al.* 2004a: 152; Chant & McMurtry 2007: 53.

Specimens examined: Muoi, in P11 plot (aasl 66 m, lat. 21°54'30" N, long. 104°38'14" E), 1 ♂ on *Rubus alceifolius* (Rosaceae), 31/V/2017; in P15 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 1 ♀ on *Chromolaena odorata* (Asteraceae), 2/VI/2017 and 1 ♀ on *R. alceifolius* (Rosaceae), 3/VI/2017; Ma, in P1 plot (aasl 66 m, lat. 21°45'53" N, long. 105°1'28" E), 1 ♂ on *Xanthium stumarium* (Asteraceae) and 1 ♀ on *Blumea eberhardtii* (Asteraceae), 14/VII/2017.

Remarks: This species is widely distributed in sub-Saharan Africa and Madagascar. The biology of *P. horrier* remains totally unknown.

This is the first record of this species from Vietnam and from Asia. Measurements are given in the table 3 for females and males and compared with those obtained from specimens collected in other countries of Africa and Indian Ocean. Measurements obtained during this study agree well with most of the previous measurements obtained, especially with measurements of specimens from La Réunion Island.

***Paraphytoseius orientalis* (Narayanan, Kaur & Ghai)**

Typhlodromus (Amblyseius) orientalis Narayanan, Kaur & Ghai 1960: 394.

Paraphytoseius orientalis, Moraes *et al.* 1986: 105; Chant & McMurtry 2003b: 220; Moraes *et al.* 2004a: 162, Chant & McMurtry 2007: 53.

Amblyseius ipomeai El-Banhawy 1984: 126 (synonymy according to Chant & McMurtry 2003b: 216).

Paraphytoseius narayanani Ehara 1967: 67 (synonymy according to Ehara & Ghai, *in* Ehara 1967: 77).

Paraphytoseius multidentatus Swirski & Shechter 1961: 114 (synonymy according to Matthysse & Denmark 1981).

Specimens examined: Muoi, in P4 plot (aasl 66 m, lat. 21°54'41" N, long. 104°38'7" E), 1 ♀ on *Styrax tonkinensis* (Styracaceae), 12/V/2017; in P6 plot (aasl 66 m, lat. 21°54'55" N, long. 104°38'22" E), 1 ♀ on *Curculigo orchoides* (Hypoxidaceae), 15/V/2017; in P4 plot (aasl 66 m, lat. 21°54'41" N, long. 104°38'7" E), 1 ♀ on *Xanthium stumarium* (Asteraceae), 18/VII/2017; in P6 plot (aasl 66 m, lat. 21°54'55" N, long. 104°38'22" E), 1 ♀ on *X. stumarium* (Asteraceae), 20/VII/2017.

Remarks: This species is widely distributed in tropical and subtropical areas in South America, Africa and Asia. It and other members of the genus belong to the large polyphagous generalist group, type III (McMurtry & Croft 1997; McMurtry *et al.* 2013). Navasero and Navasero (2016) have studied the life history of *P. orientalis* on the broad mite, *Polyphago-tarsonemus latus* (Banks), as prey. The authors reported high predation rates on *P. latus* eggs, suggesting its potential use against this pest.

This species has been collected before in Vietnam by Nguyen *et al.* (2016). Exact indications of locations were provided in this paper but without providing measurements of collected specimens. Measurements from this study are provided in the table 4 and compared with those obtained from specimens collected in other countries of Asia and Indian Ocean. Measurements obtained with this study agree well with most obtained from female specimens

Table 3 Character measurements of adult females and males of *Paraphytoseius horrifer* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	♀						♂		
	Vietnam 2, this study	Africa 4	Holotype Zaire 1	La Réunion 12	Senegal 2	South Africa 2	Vietnam 2, this study	La Réunion 1	South Africa 2
Dsl	300 – 308	300 (294 – 310)	298	295 (265 – 330)	300 – 304	318 – 339	225 – 253	250	260
Dsw at s4 level	170 – 175	158 (149 – 166)	166	153 (128 – 175)	163 – 170	165 – 176	113	110	140
j1	35	38 (37 – 40)	38	33 (25 – 40)	35 – 38	39 – 42	23 – 25	15	
j3	90	85 (80 – 88)	84	86 (78 – 100)	83 – 85	-	58 – 60	58	59
j4	5	3	3	5 (3 – 5)	3 – 5	-	3	4	-
j5	5	3	3	5 (3 – 5)	3 – 5	5	5	4	-
j6	5	7 (6 – 8)	6	8 (5 – 10)	5 – 6	9	8	7	-
J5	5	4 (3 – 5)	3	5 (3 – 8)	3 – 5	5	3	3	-
r3	49 – 50	43 (10 – 45)	45	48 (45 – 58)	43 – 45	52	25	30	33
R1	33 – 35	31 (27 – 41)	28	38 (30 – 45)	30 – 32	36	14	18	22
s4	125	131 (122 – 139)	138	126 (110 – 148)	118 – 125	136 – 141	83	80	103
z2	10 – 13	11 (10 – 12)	11	12 (9 – 13)	8 – 10	14	10	10	-
z4	9 – 10	8 (6 – 11)	6	11 (9 – 13)	8 – 10	9	10	10	-
z5	5	5 (4 – 5)	5	5 (3 – 8)	3 – 5	5	3	5	-
Z1	6 – 8	7 (6 – 8)	5	7 (5 – 8)	7 – 9	5	5	7	-
Z4	75	80 (77 – 84)	84	80 (70 – 88)	73 – 75	85 – 89	45 – 48	45	59
Z5	108	114 (112 – 118)	116	109 (93 – 133)	95 – 100	122 – 127	50	50	75
st1-st1	63 – 70	-	-	68 (55 – 73)	-	-	58	53	-
st2-st2	68 – 70	73 (72 – 74)	69	71 (65 – 80)	68 – 69	78	58	55	-
st3-st3	80	-	-	79 (73 – 85)	-	-	68	63	-
st1-st3 ♀ / st1-st5 ♂	70	66 (64 – 69)	63	68 (58 – 75)	68 – 70	92	113 – 133	98	-
st4-st4	85 – 93	-	-	88 (75 – 98)	-	-	58	55	-
gensl	125	-	-	-	-	-	Not applicable		
st5-st5	-	83 (82 – 86)	84	86 (75 – 95)	85 – 88	87	48	48	-
gensw	-	-	-	-	-	-	Not applicable		
Lisl	38	-	-	30 (25 – 38)	-	-	Not applicable		
Lisw	2	-	-	3	-	-	Not applicable		
Sisl	-	-	-	11 (10 – 13)	-	-	Not applicable		
Vsl	-	103 (96 – 110)	115	99 (53 – 118)	113 – 115	117	95 – 100	-	122
Vsw ZV2	-	68 (62 – 74)	75	59 (53 – 75)	68 – 70	70	113 – 118	105	120
Vsw anus	-	62 (59 – 67)	63	61 (55 – 68)	65	-	65 – 70	60	-
JV5	73 – 75	-	-	77 (70 – 85)	-	-	15 – 18	50	28
SgeII	13 – 15	12 (11 – 13)	-	15 (13 – 15)	13 – 15	-	-	13	14
SgeIV	25 – 30	25 (24 – 27)	24	27 (23 – 45)	28 – 30	28	20	18	23
StiIV	37 – 40	37 (35 – 40)	34	36 (30 – 43)	32 – 38	47	25 – 28	23	34
StIV	43	40 (38 – 43)	36	43 (38 – 47)	43 – 45	47	33	30	37
SttIV	45	41 (39 – 42)	-	45 (43 – 48)	33 – 35	-	35	30	-
ScI	2	4 (3 – 5)	4	3 (3 – 5)	-	25	Not applicable		
Scw	8	12 (9 – 13)	-	10 (9 – 11)	-	10	Not applicable		
Fdl	28 – 30	29	-	34 (23 – 50)	-	-	20 – 23	20	22
No teeth Fd	11	10 – 11	-	11	-	-	8	5 ?	8
Mdl	33	34 (33 – 34)	-	32 (25 – 38)	-	-	22 – 23	20	22
No teeth Md	3	2 – 3	-	2	-	-	1	1	1
Shaft	Not applicable						16 – 20	15	18
Toe	Not applicable						5	-	-

Sources of measurements – For ♀♀: Africa (Benin 1♀, Kenya 1♀, Uganda 2♀): Moraes *et al.* (2007); Holotype Zaire : Moraes *et al.* 1989b; La Réunion: Kreiter *et al.* 2020; Senegal: Kade *et al.* (2011); South Africa: van der Merwe (1968). For ♂♂: La Réunion: Kreiter *et al.* 2020; South Africa: van der Merwe (1968); - : not provided.

Table 4 Character measurements of adult females of *Paraphytoseius orientalis* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam 4, this study	Japan 8 for Dsl & Dsw, 10 for other characters	Hong-Kong 7	La Réunion 5	Madagascar ?	Mauritius 1	Paratype Hong-Kong, 1
Dsl	288 (280 – 295)	278 (275 – 281) (8)	259 – 305	266 (260 – 283)	280 – 295	290	306
Dsw at s4 level	128 (123 – 140)	153 (151 – 155) (8)	-	150 (145 – 150)	155 – 160	-	165
j1	36 (33 – 37)	36	36 (30 – 38)	31 (28 – 35)	30 – 36	38	36
j3	78 (75 – 87)	84 (82 – 85)	79 (76 – 81)	83 (81 – 88)	79 – 89	83	81
j4	4 (3 – 5)	3	4 (3 – 5)	5 (4 – 5)	< 10	5	4
j5	5 (3 – 5)	3	4 (3 – 5)	5	< 10	-	4
j6	7 (5 – 8)	6	4 (3 – 5)	8 (5 – 8)	< 10	-	6
J5	4 (3 – 5)	3 (3 – 4)	4 (3 – 5)	5 (4 – 5)	< 10	-	5
r3	45 (43 – 50)	48 (47 – 49)	43 (41 – 46)	38 (35 – 40)	36 – 48	55	45
R1	26 (25 – 28)	26 (25 – 27)	26 (20 – 33)	34 (33 – 35)	23 – 35	33	25
s4	121 (118 – 125)	121 (118 – 123)	117 (107 – 124)	120 (113 – 123)	118 – 130	125	117
z2	10 (9 – 10)	8 (7 – 8)	8 (5 – 11)	11 (10 – 15)	< 10	8	9
z4	9 (7 – 10)	9 (9 – 10)	10 (8 – 13)	10 (5 – 10)	< 10	10	11
z5	5 (3 – 5)	5	4 (3 – 5)	6 (6 – 8)	< 10	-	3
Z1	8 (7 – 8)	7	8 (8 – 10)	9 (6 – 9)	< 10	8	8
Z4	72 (65 – 78)	74 (73 – 75)	74 (69 – 81)	80 (70 – 80)	68 – 80	70	71
Z5	94 (88 – 100)	100 (99 – 101)	82 (76 – 91)	105 (99 – 107)	96 – 116	93	94
st1-st1	65 (60 – 68)	-	-	70	-	-	-
st2-st2	66 (65 – 68)	-	-	68 (65 – 68)	-	-	66
st3-st3	74 (73 – 75)	-	-	75 (75 – 78)	-	-	-
st1-st3	62 (60 – 63)	-	-	65 (64 – 65)	-	65	66
st4-st4	85 (78 – 93)	-	-	85 (85 – 93)	-	-	-
Gensl	107 (105 – 108)	-	-	-	-	-	-
st5-st5	84 (80 – 93)	-	-	85	-	-	79
Gensw	90 (88 – 93)	-	-	-	-	-	-
Lisl	30	-	-	18	-	-	-
Lisw	3	-	-	2	-	-	-
Vsl	96 (88 – 100)	-	93 (84 – 102)	80	102 – 108	113	97
Vsw ZV2	64 (63 – 65)	-	53	65	59 – 63	-	52
Vsw anus	54 (53 – 55)	-	53 – 55	63	-	-	55
JV5	71 (63 – 78)	71 (70 – 72)	69 (64 – 76)	66 (65 – 75)	63 – 82	-	-
SgeI	9 (8 – 10)	9 (8 – 9)	-	-	-	8	6
SgeII	13 (10 – 15)	14 (14 – 15)	-	13	Present	13	13
SgeIV	27 (25 – 29)	28 (27 – 28)	28 (25 – 33)	23 (21 – 28)	22 – 27	30	25
StiIV	34 (31 – 37)	34 (34 – 35)	35 (33 – 38)	38 (30 – 38)	28 – 38	38	35
StIV	46 (45 – 48)	49 (48 – 49)	44 (41 – 46)	41 (38 – 42)	38 – 45	45	43
SttIV	38 (35 – 40)	35 (38 – 39)	33 (30 – 36)	33 (33 – 45)	34 – 42	48	36
Scl	3	-	-	3 (3 – 5)	20	4	-
Scw	10	-	-	10 (8 – 11)	7	6	-
Fdl	25 (25 – 28)	-	25	27 (27 – 30)	-	25	-
No teeth Fd	8	-	7	7 – 8	8	7	-
Mdl	29 (28 – 30)	-	29 (28 – 30)	35 (31 – 35)	-	28	-
No teeth Md	3	-	2	2	3	2	-

Sources of measurements – Japan: (identified as *Paraphytoseius multidentatus*, synonymized by Mathysse & Denmark 1981); Ehara *et al.* (2000); Hong Kong: Swirski & Shechter (1961); La Réunion: Kreiter *et al.* 2020; Madagascar (identified as *Paraphytoseius multidentatus*, synonymized by Mathysse & Denmark 1981); Blommers (1976); Mauritius: Kreiter *et al.* (2018a); Paratype Hong-Kong: Moraes *et al.* (2007); - : not provided.

collected in other countries, especially with measurements of specimens from Hong Kong and La Réunion Island.

Genus *Scapulaseius* Karg & Oomen-Kalsbeek

Scapulaseius Karg & Oomen-Kalsbeek 1987: 132.

Amblyseius (Scapulaseius) Karg & Oomen-Kalsbeek 1987: 132.

newsami group of *Typhlodromus (Amblyseius)*, Chant 1959: 95.

markwelli species group of *Amblyseius*, Schicha 1987: 2.

japonicus species group of *Amblyseius*, Schicha 1987: 26.

oguroi species group of *Amblyseius*, Wu & Ou 1999: 103.

Scapulaseius, Chant & McMurtry 2005b: 331; 2007: 65.

***Scapulaseius asiaticus* (Evans)**

Typhlodromus asiaticus Evans 1953: 461.

Typhlodromus (Amblyseius) asiaticus, Chant 1959: 80.

Amblyseius (Typhlodromopsis) asiaticus, Muma 1961: 289.

Amblyseius (Amblyseius) asiaticus, Ehara 1966: 20; Ehara & Bhandhufalck, 1977: 58.

Amblyseius asiaticus, Carmona 1968: 280; Gupta, 1975: 32.

Amblyseius (Neoseiulus) asiaticus, Ehara 2002: 127.

Typhlodromips asiaticus, Moraes et al. 2004a: 207.

Scapulaseius asiaticus, Chant & McMurtry 2005b: 335; 2007: 67.

Scapulaseius linearis Corpuz-Raros & Rimando 1966: 125 (synonymy according to Schicha & Corpuz-Raros 1992: 60).

Scapulaseius siaki Ehara & Lee 1971: 64 (synonymy according to Ehara & Bhandhufalck 1977: 58).

Specimens examined: Ma, in P1 plot (aasl 63 m, lat. 21°45'53" N, long. 105°1'28" E), 1 ♀ on *Chromolaena odorata* (Asteraceae), 9/V/2017; Muoi, in P7 plot (aasl 66 m, lat. 21°54'51" N, long. 104°38'22" E), 1 ♀ on *Crassocephalum crepidioides* (Asteraceae), 16/V/2017; in P9 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'18" E), 1 ♀ on an unknown plant support, 18/V/2017; in P11 plot (aasl 66 m, lat. 21°54'30" N, long. 104°38'14" E), 1 im. on *C. odorata* (Asteraceae), 31/V/2017; in P15 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 1 ♀ on *C. odorata* (Asteraceae), 31/V/2017; in P8 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'9" E), 2 ♀♀ + 1 im. on *Xanthium strumarium* (Asteraceae), 2/VI/2017; in P13 plot (aasl 66 m, lat. 21°54'50" N, long. 104°38'46" E), 1 ♀ on *X. strumarium* (Asteraceae), 29/VII/2017.

Remarks: Species of this genus are supposed to be of type III (McMurtry and Croft 1997; McMurtry et al. 2013), i.e. a polyphagous generalist predator. However, the biology of *S. asiaticus* remains totally unknown.

This species was already known from Vietnam from results of a previous survey (Nguyen et al. 2016). Exact indications of locations were provided in this paper without providing any morphological data on the collected specimens. Measurements of collected specimens during this study are provided in table 5 and compared with those obtained from specimens collected in other countries of Asia and Indian Ocean. Measurements obtained match with most of previous measurements, especially with measurements of specimens from Thailand.

***Scapulaseius okinawanus* (Ehara)**

Amblyseius (Amblyseius) okinawanus Ehara 1967: 72; Ehara & Bhandhulfack 1977: 58.

Amblyseius okinawanus, Lo 1970: 56.

Neoseiulus okinawanus, Moraes et al. 1986: 91.

Amblyseius (Neoseiulus) okinawanus, Ehara & Amano 1998: 37; Ehara & Amano 2002: 321.

Amblyseius okinawanus, Jung et al. 2003: 189.

Typhlodromips okinawanus, Moraes et al. 1986: 137; 2004a: 220.

Scapulaseius okinawanus, Chant & McMurtry 2005b: 335; 2007: 67.

Table 5 Character measurements of adult females of *Scapulaseius asiaticus* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam 7, this study	Indonesia 3	Mauritius 9	Sri Lanka 1	Thailand 1 ?	Thailand 2 8
Dsl	306 (293 – 313)	300 – 325	311 (304 – 325)	287	330	301 (283 – 318)
Dsw at s4 level	179 (170 – 188)	180 – 195	191 (182 – 199)	182	230	200 (181 – 222)
j1	21 (18 – 23)	16 – 18	22 (20 – 24)	19	20	21 (19 – 23)
j3	14 (13 – 15)	16 – 18	17 (15 – 20)	13	18	15 (10 – 16)
j4	8	8 – 9	5	8	7	8 (7 – 10)
j5	8 (6 – 8)	8 – 9	5	8	7	8 (6 – 10)
j6	9 (8 – 10)	8 – 9	8 (7 – 9)	9	8	10 (9 – 12)
J2	9 (8 – 10)	8 – 9	9 (7 – 9)	9	9	11 (10 – 12)
J5	8 (8 – 9)	8 – 9	8 (7 – 8)	6	6	8 (7 – 8)
r3	13 (13 – 15)	16 – 18	15 (13 – 16)	12	14	13 (11 – 15)
R1	10 (9 – 13)	16 – 18	18 (16 – 19)	14	16	16 (14 – 17)
s4	18 (16 – 20)	16 – 18	23 (22 – 25)	17	21	21 (18 – 23)
S2	15 (13 – 18)	16 – 18	18 (15 – 20)	13	15	16 (13 – 18)
S4	12 (10 – 13)	16 – 18	15 (13 – 17)	12	13	14 (11 – 15)
S5	13 (10 – 15)	16 – 18	14 (13 – 15)	11	13	14 (12 – 19)
z2	12 (10 – 15)	-	14 (13 – 14)	11	13	14 (13 – 15)
z4	13 (10 – 15)	-	11 (10 – 12)	11	11	11 (7 – 13)
z5	8 (7 – 10)	-	7 (6 – 9)	7	9	9 (8 – 11)
Z1	10 (7 – 13)	-	9 (8 – 9)	-	11	11 (10 – 12)
Z4	50 (47 – 55)	44	47 (44 – 52)	45	51	52 (49 – 58)
Z5	77 (75 – 80)	73	73 (65 – 77)	68	73	74 (70 – 83)
st1-st1	50 (48 – 50)	-	-	-	-	-
st2-st2	58 (55 – 60)	-	57 (55 – 60)	52	-	59 (57 – 61)
st3-st3	61 (55 – 65)	-	-	-	-	-
st1-st3	53 (50 – 55)	-	53 (52 – 54)	48	-	54 (50 – 58)
st4-st4	65 (56 – 83)	-	-	-	-	-
Gensl	101 (90 – 113)	-	-	-	-	-
st5-st5	59 (55 – 63)	-	61 (55 – 65)	54	-	61 (58 – 64)
Gensw post. corn.	69 (65 – 80)		71 (60 – 74)	-	-	-
Lisl	16 (13 – 18)	-	-	-	-	-
Lisw	3 (2 – 5)	-	-	-	-	-
Sisl	10	-	-	-	-	-
Vsl	102 (95 – 110)	105	99 (95 – 101)	97	-	103 (97 – 110)
Vsw ZV2	82 (78 – 90)	116	81 (77 – 84)	79	-	84 (80 – 87)
Vsw anus	64 (50 – 73)	-	67 (61 – 74)	65	-	-
JV5	23 (20 – 25)	-	26 (22 – 28)	-	19	-
SgeI	16 (15 – 20)	-	18 (16 – 19)	-	-	17 (15 – 20)
SgeII	12 (10 – 15)	-	12 (10 – 13)	-	-	11 (8 – 13)
SgeIII	17 (15 – 19)	-	15 (13 – 17)	14	-	17 (15 – 18)
StiIII	15 (13 – 18)	-	-	15	-	-
SgeIV	28 (25 – 30)	-	26 (24 – 29)	22	27	27 (24 – 30)
StiIV	22 (20 – 24)	-	20 (18 – 23)	18	22	22 (18 – 26)
StIV	50 (48 – 50)	-	49 (44 – 53)	45	50	49 (44 – 52)
Scl	30 (28 – 35)	-	33 (29 – 34)	27	-	35 (29 – 40)
Scw	5 (4 – 6)	-	-	-	-	-
Fdl, No teeth	25 (23 – 28), 9	-	26 (24 – 27), 9	-	-	22 (21 – 23)
Mdl, No teeth	25 (23 – 28), 3	18	27 (25 – 29), 2-3	-	-	25 (23 – 25)

Sources of measurements – Mauritius: Ferragut & Baumann (2019); Sri Lanka: Moraes *et al.* (2004b); Thailand 1: Ehara & Bhandhufalck (1977); Thailand 2: Oliveira *et al.* (2012); - : not provided.

Specimens examined: Muoi, in PM1 plot (aasl 66 m, lat. 21°54' N, long. 104°38' E), 2 ♂♂ on an unknown host plant, 17/V/2017; in P9 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'18" E), 2 ♀♀ on *Crassocephalum crepidioides* (Asteraceae), 17/V/2017; in P15 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 1 ♀ on *Rubus alceifolius* (Rosaceae), 17/V/2017; in P11 plot (aasl 66 m, lat. 21°54'30" N, long. 104°38'14" E), 3 ♀♀ on *Xanthium strumarium* (Asteraceae), 26/VII/2017.

Remarks: The biology of *S. okinawanus* remains totally unknown.

This is the first mention of this species in Vietnam.

Measurements of specimens collected during this study are provided in table 6 for females and males and compared with measurements obtained from specimens collected in other countries of Asia. Measurements obtained agree well with most of those previously obtained, except setae *j6*, *J2*, and *z5*, and macrosetae *SgeI* and *SgeII* in female specimens and shorter setae of the *j-J*, *S* series and *Z1* of male specimens.

Genus *Amblyseius* Berlese

Amblyseius Berlese, 1914: 143; Garman, 1948: 16; Muma, 1955: 263; Chant, 1957: 528; Kennet, 1958: 474; Muma, 1961: 287; Gonzalez & Schuster, 1962: 8; Pritchard & Baker, 1962: 235; van der Merwe & Ryke, 1963: 89; Corpuz & Rimando, 1966: 116; van der Merwe, 1968: 109; Zack, 1969: 71; Muma and Denmark, 1970: 62; Chant & Hansell, 1971: 703; Denmark & Muma, 1972: 19; Tseng, 1976: 104; Chaudhri *et al.*, 1979: 68; Karg, 1982: 193; Schicha, 1987: 19; Schicha & Corpuz-Raros, 1992: 12; Denmark & Muma, 1989: 4; Chant & McMurtry, 2004a: 188; 2007: 73.

Amblyseius (*Amblyseius*), Karg, 1983: 313.

Amblyseius (*Amblyseialus*), Karg, 1983: 313.

Amblyseius (*Amblyseius*) section *Amblyseius*, Wainstein, 1962: 15.

Amblyseius (*Amblyseius*) section *Italoseius* Wainstein, 1962: 15.

Amblyseius cinctus Corpuz & Rimando

Amblyseius cinctus Corpuz & Rimando 1966: 119.

Amblyseius (*Amblyseius*) *cinctus*, Ehara & Bhandhfalck 1977: 70.

Amblyseius (*Multiseius*) *cinctus*, Denmark & Muma 1989: 103.

Amblyseius cinctus, Moraes *et al.* 1986: 10; 2004a: 20; Chant & McMurtry 2004a: 203; 2007: 78.

Specimens examined: Muoi in P1 plot (aasl 66 m, lat. 21°54' N, long. 104°38' E), 10 ♀ + 1 im. on *Manihot esculenta* (Euphorbiaceae), 17/V/2017; in P5 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'22" E), 1 ♀ on *Urena lobata* (Malvaceae), 19/VII/2017.

Remarks: This species was described from specimens collected in the Philippines on *Panicum* sp. and *Streblus asper*. Life history of *A. cinctus* has been evaluated on *Polyphago-tarsonemus latus* (Banks), an important pest of several cultivated (crop) plants worldwide (Vichitbandha and Chandrapatya 2009) and this species can be considered as a potential BCA of *P. latus*.

This is the first mention of this species in Vietnam.

Measurements of specimens collected during this study are provided in table 7 and compared with measurements obtained from specimens collected in other countries of South-East Asia. Measurements obtained during this study agree well with those obtained from female specimens collected in other countries.

Amblyseius herbicolus (Chant)

Typhlodromus (*Amblyseius*) *herbicolus* Chant 1959: 84.

Amblyseius (*Amblyseius*) *herbicolus*, Muma 1961: 287.

Typhlodromus *herbicolus*, Hirschmann 1962: 23.

Table 6 Character measurements of adult females and males of *Scapulaseius okinawanus* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	♀				♂		
	Vietnam 6, this study	Japan 9	Korea 7	New Hebrides 1?	Vietnam 2, this study	Japan 3	Korea 4
Dsl	324 (310 – 328)	340	306	314	243 – 250	250	-
Dsw at s4 level	201 (180 – 210)	190 – 250	205	173	125 – 128	180	-
j1	20 (18 – 22)	21	20	-	18	18	18
j3	15 (13 – 15)	16	16	-	13 – 14	19	19
j4	9 (8 – 10)	10	9	-	6 – 8	10	9
j5	8 (5 – 10)	10	9	-	6 – 8	9	9
j6	8 (8 – 10)	12	12	-	8	11	10
J2	10 (9 – 10)	15	15	-	8 – 10	12	11
J5	9 (8 – 9)	8	9	-	7 – 8	8	7
r3	13 (10 – 15)	14	13	-	12 – 14	12	10
R1	10 (8 – 13)	11	11	-	11 – 12	11	10
s4	20 (19 – 21)	22	20	-	18 – 19	19	18
S2	19 (18 – 20)	17	18	-	13 – 16	17	18
S4	16 (15 – 18)	16	17	-	13	17	16
S5	16 (14 – 18)	15	18	-	11 – 12	15	17
z2	14 (13 – 15)	15	14	-	13	14	14
z4	15 (13 – 16)	16	15	-	15 – 18	15	15
z5	9 (8 – 10)	11	10	-	6 – 8	10	9
Z1	12 (10 – 13)	14	14	-	5 – 7	12	12
Z4	35 (33 – 37)	35	33	-	30 – 34	30	27
Z5	82 (80 – 88)	78	80	-	53 – 58	57	56
st1-st1	49 (48 – 51)	-	-	-	38 – 43	-	-
st2-st2	58 (55 – 63)	-	-	65	48 – 51	-	-
st3-st3	64 (63 – 68)	-	-	-	47 – 48	-	-
st1-st3 ♀ / st1-st5 ♂	55 (53 – 58)	-	-	55	93 – 95	-	-
st4-st4	70 (60 – 83)	-	-	-	35 – 36	-	-
Gensl	104 (103 – 105)	-	-		Not applicable		
st5-st5	62 (55 – 68)	-	-	-	33 – 34	-	-
Gensw post. corn.	72 (68 – 75)	-	-				
Lisl	16 (15 – 18)	-	-	18			
Lisw	5 (3 – 7)	-	-	-	Not applicable		
Sisl	10 (8 – 13)	-	-	10			
Vsl	104 (98 – 108)	-	-	115	97 – 105	-	-
Vsw ZV2	88 (83 – 90)	-	-	92	100 – 115	-	-
Vsw anus	79 (78 – 80)	-	-	-	58 – 68	-	-
Dist. solen. vas	24 (23 – 25)	-	-		18	20	22
JV5	29 (25 – 30)	28	28	-	18	-	-
SgeI	10 (9 – 13), knob.	-	-	19 – 20 knob.	10	-	-
SgeII	9 (8 – 9), knob.	-	-	15 – 16 knob.	13	-	-
SgeIII	19 (18 – 20), knob.	-	-	21 – 23 knob.	10	-	-
StiIII	17 (13 – 19)	-	-	18	23	20	21
SgeIV	27 (25 – 30), knob.	29	25	31 knob.	15	20	22
StiIV	24 (20 – 25)	24	25	25	43	42	48
StIV	53 (53 – 54), knob.	54	56	55 knob.			
Scl	11 (10 – 13)	-	-	8	Not applicable		
Scw	7 (5 – 8)	-	-	-			
Fdl, No teeth	25 (23 – 25)	-	-	-	23	-	-
No teeth Fd	7	-	-	-	7	-	-
Mdl, No teeth	27 (25 – 28)	-	-	-	23	-	-
No teeth Md	3	-	-	-	1	-	-
Shaft	Not applicable				13 – 15	-	-
Toe					5	-	-

Sources of measurements – For ♀♀: Japan: Ehara (1967); Korea: Ryu & Lee (1992); New Hebrides: Schicha (1982). For ♂♂: Japan Ehara (1967); Korea: Ryu & Lee (1992). - : not provided.

Amblyseius herbicolus, Moraes *et al.* 1986: 14; 1989a: 79; Chant & McMurtry 2004: 208; Moraes *et al.* 2004a: 27; Chant & McMurtry 2007: 78.

Amblyseius deleoni Muma & Denmark 1970: 68 (synonymy according to Daneshvar & Denmark 1982; Denmark & Muma 1989).

Amblyseius giganticus Gupta 1981: 33 (synonymy according to Gupta 1986).

Amblyseius impactus Chaudhri 1968: 553 (synonymy according to Daneshvar & Denmark 1982).

Amblyseius (Amblyseiulus) thermophilus Karg 1991: 12 (synonymy according to El-Banhawy & Knapp 2011).

Typhlodromus (Amblyseius) amitae Bhattacharyya 1968: 677 (synonymy according to Denmark & Muma 1989).

Specimens examined: Muoi in P8 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'9" E), 1 ♀ on *Paspalum* sp. (Poaceae), 17/V/2017.

Remarks: *Amblyseius herbicolus* is widespread in all tropical and subtropical regions of the world. It is the second most abundant phytoseiid mite on *Coffea arabica* in Brazil, and is deemed to be an efficient predator of *Brevipalpus phoenicis* (Geijskes), vector of the coffee ring spot virus (Reis *et al.* 2007). *Amblyseius herbicolus* is also found in association with the broad mite, *Polyphagotarsonemus latus* Banks in crops such as chili pepper (*Capsicum annuum*) in Brazil and has good potential for controlling the latter. Rodriguez-Cruz *et al.* (2013) have demonstrated that *A. herbicolus* was able to develop on broad mites, castor bean pollen (*Ricinus communis*) and sunn hemp pollen (*Crotalaria juncea*). Provision of pollen can enhance populations of this predator and increase biological control (Duarte *et al.* 2015),

Table 7 Character measurements of adult females of *Amblyseius cinctus* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam 11, this study	Philippines 2	Thailand 1 ?	Thailand 2 4	Characters	Vietnam 11, this study	Philippines 2	Thailand 1 ?	Thailand 2 4
Dsl	337 (308 – 363)	314	350	326 (312 – 341)	Gensl	114 (105 – 125)	-	-	
Dsw at s4 level	196 (188 – 207)	175	240	223 (218 – 227)	st5-st5	66 (62 – 69)	74	-	68 (66 – 70)
j1	25 (23 – 30)	24 – 26	25	25 (24 – 25)	Gensw post. corn.	77 (70 – 83)	-	-	-
j3	44 (40 – 48)	42 – 44	44	45 (44 – 46)	Lisl	16 (13 – 18)	16	-	-
j4	5 (4 – 6)	5	5	5 (5 – 6)	Lisw	5 (3 – 6)	-	-	-
j5	5 (4 – 5)	5	4	4	Sisl	9 (7 – 10)	8	-	-
j6	5 (5 – 7)	5	5	6 (5 – 7)	Vsl	109 (100 – 115)	82	-	114 (110 – 118)
J2	6 (5 – 7)	5	5	6 (6 – 7)	Vsw ZV2	81 (75 – 88)	-	-	80 (78 – 84)
J5	5 (5 – 7)	8	6	6	Vsw anus	73 (63 – 78)	-	-	-
r3	13 (10 – 15)	10	13	13 (13 – 14)	Dist. Solen. pva	21 (20 – 23)	-	-	
R1	8 (6 – 8)	9	7	7 (7 – 8)	JV5	59 (55 – 65)	52 – 70	57	-
s4	73 (63 – 82)	68 – 73	69	70 (68 – 77)	Sgel	32 (28 – 35)	33	-	34 (32 – 35)
S2	7 (5 – 8)	6	7	8 (7 – 8)	SgelII	30 (25 – 33)	32	-	33 (32 – 34)
S4	5 (5 – 6)	6	7	7 (6 – 7)	SgelIII	42 (38 – 48)	45	-	44 (42 – 46)
S5	5 (5 – 6)	9	6	6 (5 – 7)	StilII	30 (28 – 33)	34	-	31 (28 – 32)
z2	10 (8 – 13)	7	9	10	SgelIV	85 (80 – 91)	84 – 90	79	84 (80 – 86)
z4	8 (6 – 9)	8	7	7	StilIV	49 (45 – 55)	36 – 53	47	49 (46 – 54)
z5	5 (4 – 5)	4	4	4	StilV	62 (53 – 65)	56 – 60	60	66 (64 – 67)
Z1	6 (5 – 8)	6	6	7 (7 – 8)	Scl	11 (10 – 15)	12 – 16	-	11 (10 – 11)
Z4	88 (83 – 95)	72 – 90	86	95 (88 – 106)	Scw	3 (3 – 4)	3	-	-
Z5	178 (166 – 188)	200	169	172 (168 – 182)	Fdl, No teeth	29 (25 – 33), 9 – 10	27, 9 – 10	-	-
st1-st1	59 (55 – 64)	-	-	-	Mdl, No teeth	30 (28 – 33), 3 – 4	31, 3 – 4	-	-
st2-st2	66 (63 – 68)	60	-	66 (65 – 67)					
st3-st3	74 (70 – 77)	79	-	-					
st1-st3	55 (53 – 58)	-	-	56 (55 – 58)					
st4-st4	76 (68 – 105)	-	-	-					

Sources of measurements – Philippines: Corpuz & Rimando (1966) and Schicha & Corpuz-Raros (1992); Thailand 1: Ehara & Bhandhfalck (1977); Thailand 2: Oliveira *et al.* (2012); - : not provided.

Table 8 Character measurements of adult females of *Amblyseius herbicolus* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam 1, this study	Africa 8	Grande Comore 2	Holotype Portugal, 1	Kenya 46	La Réunion 15	Senegal 2	Thailand 1	Turkey 3
Dsl	388	352 (325 – 368)	385 – 428	369	335	359 (343 – 390)	360 – 365	360	313 – 352
Dsw at s4 level	182	256 (240 – 274)	263 – 275	236	190	240 (225 – 265)	250 – 268	194	196 – 221
j1	38	37 (34 – 40)	40 – 43	38	32	35 (33 – 38)	35 – 38	37	33 – 38
j3	43	49 (38 – 58)	53 – 55	42	37	40 (38 – 45)	38 – 45	38	32 – 39
j4	8	6 (5 – 8)	6	9	4 – 6	7 (5 – 8)	6 – 8	6	8 – 9
j5	5	4 (3 – 5)	5	7	4 – 6	5 (5 – 5)	4	5	7 – 8
j6	8	7 (5 – 8)	5 – 6	11	4 – 6	7 (5 – 8)	8	8	6 – 7
J2	10	10 (8 – 11)	8 – 10	12	4 – 6	8 (8 – 10)	8 – 9	9	8 – 9
J5	10	8 (6 – 10)	10	9	4 – 6	8 (8 – 10)	8 – 10	7	9 – 10
r3	13	14 (11 – 16)	8 – 15	15	10	12 (10 – 13)	10 – 11	13	10 – 16
R1	10	9 (8 – 10)	10 – 13	8	10	10 (10 – 13)	9 – 10	9	8 – 9
s4	105	113 (98 – 130)	120 – 125	100	92	98 (95 – 108)	123 – 135	95	86 – 96
S2	8	12 (8 – 14)	13	11	10	12 (10 – 13)	13 – 15	10	10 – 12
S4	8	11 (8 – 13)	13	13	10	11 (8 – 13)	8 – 10	11	9 – 11
S5	8	9 (8 – 10)	13	11	10	9 (8 – 10)	8 – 10	9	9 – 10
z2	10	11 (8 – 16)	6 – 8	13	6	13 (8 – 18)	8 – 9	13	9 – 12
z4	10	8 (8 – 10)	8	9	6	10 (8 – 13)	10 – 11	10	8 – 12
z5	8	6 (5 – 6)	5	6	6	5 (5 – 8)	7 – 9	7	6 – 7
Z1	8	10 (8 – 13)	8	9	10	12 (10 – 13)	8 – 10	13	9 – 12
Z4	105	126 (101 – 152)	133 – 135	110	90	99 (93 – 108)	163 – 172	94	91 – 99
Z5	288	281 (251 – 306)	288 – 300	236	232	255 (248 – 273)	310 – 345	270	220 – 251
st1-st1	65	-	68	-	-	66 (63 – 68)	-	-	-
st2-st2	70	71 (66 – 75)	73 – 75	-	69	73 (65 – 78)	75 – 78	73	70 – 73
st3-st3	75	-	63 – 65	-	-	77 (73 – 83)	-	-	-
st1-st3	65	65 (58 – 70)	73 – 75	-	62	69 (68 – 73)	63 – 65	67	64 – 69
st4-st4	75	-	78 – 83	-	-	76 (73 – 80)	-	-	-
Gensl									
st5-st5	63	71 (67 – 75)	70 – 78	-	65	65 (63 – 70)	65 – 70	68	58 – 64
Gensw post. C.									
Lisl	20	-	23	-	-	22 (18 – 25)	-	-	-
Lisw	5	-	5	-	-	5 (5 – 5)	-	-	-
Sisl	18	-	13	-	-	14 (10 – 18)	-	-	-
Vsl	110	116 (112 – 118)	120 – 135	-	108	111 (100 – 123)	113 – 115	117	102 – 117
Vsw ZV2	50	57 (53 – 59)	63	-	48	48 (43 – 58)	65 – 69	56	44 – 48
Vsw anus	70	71 (66 – 77)	80	-	69	69 (63 – 78)	75 – 78	70	61 – 69
JV5	55	-	80 – 85	-	52	61 (53 – 78)	-	-	51 – 60
SgeI	55	42 (35 – 48)	45 – 48	-	-	44 (40 – 48)	43 – 48	50	40 – 48
SgeII	40	38 (35 – 42)	38	-	52	37 (33 – 40)	38 – 40	39	34 – 39
SgeIII	53	52 (45 – 59)	53 – 58	-	41	44 (35 – 48)	48 – 52	48	41 – 46
StIII	43	41 (34 – 48)	43	-	35	41 (38 – 48)	43 – 45	40	33 – 40
SgeIV	128	124 (96 – 158)	135	112	110	118 (110 – 123)	160 – 162	110	98 – 128
StIV	Broken	90 (67 – 109)	100	82	76	88 (83 – 93)	102 – 115	85	75 – 89
StIV	70	76 (66 – 86)	80	76	65	72 (65 – 78)	76 – 78	72	63 – 70
Scl	25	-	38 – 40	18	28	31 (28 – 33)	33 – 35	25	24 – 32
Scw	5	-	2 – 3	-	-	5 (5 – 8)	-	-	-
Fdl, No teeth	28, 10+1	31 (30 – 31), 10+1	33, 11 – 12	11 – 12	8	36 (30 – 40), 13	-	34	29 – 36, 12
Mdl, No teeth	33, 3 – 4	35 (34 – 35), 3	30	? , 4	? , 3	33 (28 – 35), 4	-	35	31 – 33, 4

Sources of measurements – Africa (Benin 1♀, Burundi 1♀, Democratic Republic of Congo 1♀, Ghana 1♀, Kenya 3♀, Rwanda 1♀): Zannou *et al.* (2007); Grande Comore Island of the Comoros Archipelago: Kreiter *et al.* (2018b); Portugal (holotype): Denmark & Muma (1989); Kenya: El-Banhawy & Knapp (2011); La Réunion: Kreiter *et al.* 2020; Senegal: Kade *et al.* (2011); Thailand: Oliveira *et al.* (2012); Turkey: Akyazi *et al.* (2016); - : not provided.

especially with cattail pollen (*Typha latifolia*), chilli pepper pollen and bee-collected pollen. *Amblyseius herbicolus* was collected recently in Comoros archipelago (Kreiter *et al.* 2018b). This is the first record of that species in Vietnam.

Measurements of the single specimen collected during this study are provided in table 8 and compared with measurements obtained from specimens collected in other countries of Africa, Asia and Indian Ocean. These measurements match with those obtained previously. The dorsal shield width is less important in the single Vietnam specimen than all order widths of the table, probably because the single female was a small one, all other dimensions being quite similar.

Genus *Graminaseius*

Graminaseius Chant & McMurtry 2004a: 215; 2007: 83.

Graminaseius polisensis (Schicha & Corpus-Raros)

Amblyseius polisensis Schicha & Corpuz-Raros 1992: 39; Moraes *et al.* 2004a: 47.
Graminaseius polisensis, Chant & McMurtry 2004a: 219; 2007: 85.

Specimens examined: Muoi in P6 plot (aasl 66 m, lat. 21°54'55" N, long. 104°38'22" E), 1 ♀ + 1 ♂ on *Xanthium strumarium* (Asteraceae), 20/VII/2017.

Remarks: This species was described from several specimens collected on various plants in Philippines.

The biology of *G. polisensis* remains totally unknown and even the life type of this species and of species of *Graminaseius* are unknown.

Table 9 Character measurements of one adult female and one adult male of *Graminaseius polisensis* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	♀		♂		Characters	♀		♂	
	Vietnam 1, this study	Holotype Philippines, 1	Vietnam 1, this study	Paratype Philippines, 1		Vietnam 1, this study	Holotype Philippines, 1	Vietnam 1, this study	Paratype Philippines, 1
Dsl	325	380	250	299	<i>stl-st3</i> ♀ / <i>st1-st5</i> ♂	58	68	70	68
Dsw at <i>s4</i> level	205	216	166	179	<i>st4-st4</i>	68	-	48	-
<i>j1</i>	15	19	15	16	<i>st5-st5</i>	63	93	38	36
<i>j3</i>	Broken	32	23	25	Lisl	20	29	Not applicable	
<i>j4</i>	10	14	10	12	Lisw	3	-		
<i>j5</i>	10	14	10	12	Sisl	10	14	126	
<i>j6</i>	Broken	18	10	12	Vsl	110	-		
<i>J2</i>	Broken	18	10	15	Vsw ZV2	100	-	105	155
<i>J5</i>	Broken	10	8	8	Vsw anus	83	-	150	155
<i>r3</i>	15	16	Broken	-	<i>JV5</i>	38	38	68	-
<i>R1</i>	14	14	10	-	<i>SgeIV</i>	43	43	30	31
<i>s4</i>	Broken	51	Broken	37	<i>StIV</i>	53	48	43	37
<i>S2</i>	Broken	10	18	19	Scl	27	27	Not applicable	
<i>S4</i>	Broken	10	13	13	Sew	5	4		
<i>S5</i>	Broken	10	13	13	Fdl	33	38	23	-
<i>z2</i>	15	19	13	16	No teeth Fd	-	6	-	-
<i>z4</i>	13	19	13	16	Mdl	33	38	25	-
<i>z5</i>	8	11	Broken	12	No teeth Md	-	1	-	-
<i>Z1</i>	15	19	10	12	Shaft	Not applicable		20	19
<i>Z4</i>	Broken	62	Broken	50	Foot			38	-
<i>Z5</i>	58	57	35	39					
<i>st1-st1</i>	48	-	43	-					
<i>st2-st2</i>	63	82	53	82					
<i>st3-st3</i>	70	-	55	-					

Sources of measurements – For ♀♀: Philippines: Schicha & Corpuz-Raros (1992), probably measurements of the holotype female. For ♂♂: Philippines: Schicha & Corpuz-Raros (1992), probably measurements of the paratype male; - : not provided.

This species was already known from Vietnam from a previous survey (Nguyen *et al.* 2016). Details of locations were provided in this paper though no measurements were provided on the collected specimens. Measurements of specimens collected in this study are provided in table 9 for the single female and the single male and compared with measurements obtained from specimens collected in the Philippines, *i.e.*, country of the original description. Measurements obtained for Vietnamese specimens (both damaged) collected during this study agree well with most obtained from female and male type material from Philippines, most of the setae being shorter in Vietnamese specimens.

Genus *Gynaeseius* Wainstein

Amblyseius (Kampimodromus) section *Gynaeseius* Wainstein 1962: 14.

Indoseiulus Ghai & Menon 1969: 347.

Amblyseius (Indoseiulus) Ehara 1982: 42.

Indoseiulus, Denmark & Kolodochka 1993: 249; Moraes *et al.* 1986: 59; 2004a: 89.

Macmurtryseius Kolodochka & Denmark 1995: 20.

irregularis group Chant 1959: 70.

Gynaeseius, Chant & McMurtry 2006: 22; 2007: 107.

Gynaeseius liturivorus (Ehara)

Amblyseius (Indoseiulus) liturivorus Ehara 1982: 43; McMurtry & Moraes 1984: 29; Ehara 1985: 120.

Amblyseius (Amblyseius) liturivorus, Tseng 1983: 54.

Indoseiulus liturivorus, Moraes *et al.* 1986: 60; Denmark and Kolodochka 1993: 253; Ehara *et al.* 1994: 139; Ehara & Amano 1998: 48; Moraes *et al.*: 2004a: 89.

Amblyseius armellae Schicha and Gutierrez 1985: 175 (synonymy according to Denmark & Kolodochka 1993).

Re-description of the adult female of *Gynaeseius liturivorus* (Ehara)

n = 19 (Figs. 1 a – b, 2 a – c)

Dorsum — (Fig. 1a) Dorsal shield 361 (333 – 375) long and 223 (188 – 250) wide at level of *s*4, smooth, with seven solenostomes (*gd*1, *gd*2, *gd*4, *gd*5, *gd*6, *gd*8, *gd*9), 21 pairs of poroids, 16 pairs of dorsal setae and two pairs of sub-lateral setae off the dorsal shield: *j*1 18 (15 – 20), *j*3 19 (18 – 20), *j*4 14 (12 – 15), *j*5 13 (11 – 15), *j*6 15 (13 – 18), *J*2 18 (15 – 20), *J*5 8 (7 – 8), *z*2 16 (15 – 18), *z*4 18 (15 – 20), *z*5 13 (10 – 15), *Z*1 19 (18 – 20), *Z*4 18 (15 – 20), *Z*5 28 (25 – 30), *s*4 20 (18 – 23), *S*2 19 (15 – 23), *S*5 19 (18 – 20), *r*3 16 (15 – 18), *R*1 19 (17 – 20). All setae minute and smooth.

Peritreme — (Fig. 1a) Extending to a level between *j*1 to *j*3, closer to *j*3.

Venter — (Fig. 1b) All shields smooth. Sternal shield with three pairs of setae and two pairs of lyrifissures; one pair of sternal setae on metasternal shields with a pair of pores; posterior margin with a projection very difficult to see. Distances between *st*1–*st*1 64 (60 – 68), *st*2–*st*2 70 (63 – 78), *st*3–*st*3 81 (75 – 88), *st*1–*st*3 74 (68 – 78), *st*4–*st*4 93 (79 – 106). Genital shield length 130 (115 – 140), width at the level of *st*5 84 (78 – 90), width at the level of the posterior corners 95 (83 – 100). Two pairs of inguinal sigilla of “metapodal shields” 26 (23 – 28) long and 4 (3 – 5) wide for the primary and 14 (13 – 15) long for the secondary shield. Ventrianal shield with three pairs of preanal setae (*JV*1, *JV*2, and *ZV*2), small oblong pre-anal pores. Membrane surrounding ventrianal shield with four pairs of setae (*ZV*1, *ZV*3, *JV*4 and *JV*5), and six pairs of round to oblong poroids; ventrianal shield 109 (91 – 135) long, 66 (63 – 70) wide at level of *ZV*2, and 54 (50 – 60) wide at level of anus. One pair of pore distant to 24 (20 – 33). *JV*5 23 (20 – 25) long, smooth.

Chelicera — (Fig. 2a) Chelicerae are visible by the side with digits open. Consequently, they are drawn. Fixed digit 30 (28 – 33) long with 11 teeth; and movable digit 34 (33 – 36) long with 3 small teeth.




Figure 1 Female of *Gynaesius liturivorus* (Ehara). a – Dorsal shield and peritreme; b – Ventral shields.




Figure 2 Female of *Gynaeseius liturivorus* (Ehara). a – Insemination apparatus; b – Chelicera; c – Genu; tibia and basitarsus of the leg IV.

Spermatheca — (Fig. 2b) Spermatheca with a long, slender tubular calyx 38 (33 – 47) long and 4 (2 – 7) wide, an atrium slightly swollen bifurcate. Ductus minor and ductus major visible in some specimens.

Legs — (Fig. 2c) Macrosetae on all legs, all pointed except on legs IV with macrosetae slightly bulbous; one on genua of legs I, II and III, and three on genu, tibia and basitarsus of leg IV: *SgeI* 35 (30 – 38), *SgeII* 15 (13 – 20), *SgeIII* 29 (25 – 30), *StiIII* 30 (28 – 35), *SgeIV* 42 (37 – 47), *StiIV* 54 (50 – 55), *StiV* 48 (43 – 53). Genu II and III with 7 setae each, chaetotactic formula of genu II: 2-2/0, 2/0-1; genu III: 1-2/1, 2/0-1. A pair of erected setae on femora IV: 30 (28 – 32).

First description of the deutonymphs of *Gynaeseius liturivorus* (Ehara)

n = 4 (Figs 3 a – b)

Dorsum — (Fig. 3a) Dorsal shield 301 (250 – 338) long and 169 (155 – 188) wide at level of *s4* with no solenostomes and no poroids visible. The dorsal shield bears 14 pairs of dorsal setae and 2 pairs of sub-lateral setae: *j1* 28 (25 – 33), *j3* 31 (28 – 35), *j4* 4 (3 – 6), *j5* 4 (3 – 6), *j6* 6 (5 – 8), *J2* 9 (8 – 10), *J5* 5 (4 – 5), *z2* 5 (4 – 7), *z4* 6 (5 – 8), *z5* 4 (3 – 6), *Z4* 12 (10 – 13), *Z5* 31 (28 – 35), *s4* 17 (15 – 18), *S2* 9 (7 – 10), *r3* 8 (6 – 8), *R1* 6 (5 – 8). All setae smooth.

Peritreme — (Fig. 3a) Extending to a level between *z4* and *z2*, closer to *z2*.

Venter — All ventral shields very smooth not reticulated with margins not visible and so no drawings possible. Distances between *st1* – *st1* 44 (40 – 47), *st2* – *st2* 64 (63 – 65), *st3* – *st3* 59 (55 – 63), *st1* – *st3* 80, *st4* – *st4* 53 (50 – 55), *st5* – *st5* 49 (47 – 50). Ventrianal shield with three pairs of pre-anal setae, *JV1*, *JV2*, and *ZV2*, and a pair of small oblong pores 21 (20 – 23) apart. Membrane surrounding ventrianal shield with one pair of setae *JV5* smooth; ventrianal shield 94 (90 – 100) long, 54 (53 – 55) wide at anterior corners and 52 (50 – 53) wide at level of *anus*. *JV5* 15 long, smooth.




Figure 3 Deutonymph of *Gynaeseius liturivorus* (Ehara). a – Dorsal shield and peritreme; b – Chelicera.

Chelicera — (Fig. 3b) Fixed digit 23 (22 – 25) long, with 9 strong teeth and movable digit 25 (23 – 28) long, with 2 small teeth.

Legs — Legs and macrosetae like in females but with different dimensions: *SgeI* 33 (28 – 35), *SgeII* 30 (28 – 35), *SgeIII* 33 (30 – 35), *StI* 34 (30 – 35), *SgeIV* 49 (48 – 50), *StIV* 51 (45 – 55), and *StIV* 54 (53 – 55). Chaetotactic formula of genu II and III similar to that of females.

Specimens examined — Muoi in all experimental plots (aasl 66 m, lat. 21°54' N, long. 104°38' E), 19 ♀♀ + 4 deutonymphs on *Manihot esculenta* (Euphorbiaceae), 20/VII/2017.

Voucher specimens — 19 ♀♀ + 4 deutonymphs in 2 different slides deposited in CBGP, in Montpellier SupAgro Acarology collection, France.

Remarks — This species was found on *Manihot esculenta* in Vietnam. It is a type-III predator and is considered of interest for thrips control on grape infested with *Scirtothrips dorsalis* Hood (Shibao *et al.* 2004) and on various plants with *Frankliniella occidentalis* (Pergande) and *Thrips palmi* Karny (Mochizuki 2009). *Gynaeseius liturivorus* fed on the larvae of these thrips species. Females consumed an average of 7.4 - 19.4 first-stadium larvae of these

Table 10 Character measurements of adult females of *Gynaeseius liturivorus* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam 19, this study	Indonesia 10	Japan 17	Papua-New Guinea ?	Philippines 2	Taiwan 3	Thailand 2
Dsl	361 (333 – 375)	347 (341 – 353)	342	353	313	323 (317 – 326)	368 (362 – 375)
Dsw at s4 level	223 (188 – 250)	235 (231 – 239)	236	194	187	219 (211 – 226)	239 (237 – 242)
j1	18 (15 – 20)	25 (25 – 26)	15 (15 – 16)	27	14 – 20	17	16 (16 – 17)
j3	19 (18 – 20)	20 (19 – 21)	19 (18 – 20)	24	14 – 20	19	18 (17 – 19)
j4	14 (12 – 15)	17 (16 – 17)	13 (13 – 14)	17 – 19	14 – 20	12	17 (16 – 18)
j5	13 (11 – 15)	18 (18 – 19)	14 (13 – 15)	17 – 19	14 – 20	11	15 (15 – 16)
j6	15 (13 – 18)	17 (16 – 18)	15 (14 – 15)	17 – 19	14 – 20	12	13
J2	18 (15 – 20)	19 (18 – 19)	16 (16 – 17)	17 – 19	14 – 20	14	15
J5	8 (7 – 8)	9	7 (6 – 7)	8	14 – 20	12	9 (8 – 10)
r3	16 (15 – 18)	19 (18 – 20)	15 (15 – 16)	21	14 – 20	-	12 (12 – 13)
R1	19 (17 – 20)	20 (19 – 20)	18 (16 – 18)	-	14 – 20	-	17 (17 – 18)
s4	20 (18 – 23)	22 (21 – 22)	20 (20 – 21)	23	14 – 20	17	16
S2	19 (15 – 23)	23 (22 – 23)	21 (21 – 22)	23	14 – 20	16	20 (20 – 21)
S5	19 (18 – 20)	23 (22 – 23)	18 (18 – 19)	24	14 – 20	17	19 (18 – 20)
z2	16 (15 – 18)	19 (19 – 20)	16 (15 – 16)	20	14 – 20	12	16 (16 – 17)
z4	18 (15 – 20)	20 (19 – 20)	16 (16 – 17)	19	14 – 20	16	18 (17 – 19)
z5	13 (10 – 15)	18 (17 – 18)	15 (14 – 15)	17	14 – 20	11	14 (13 – 16)
Z1	19 (18 – 20)	22 (22 – 23)	19 (18 – 20)	25	14 – 20	16	20
Z4	18 (15 – 20)	20 (19 – 20)	17	23	14 – 20	11	19 (18 – 20)
Z5	28 (25 – 30)	30 (29 – 31)	23 (23 – 24)	31	27	24	30
st1-st1	64 (60 – 68)	-	-	-	-	-	-
st2-st2	70 (63 – 78)	-	-	-	-	-	69 (69 – 70)
st3-st3	81 (75 – 88)	-	-	81	-	-	-
st1-st3	74 (68 – 78)	-	-	72	-	-	69 (68 – 70)
st4-st4	93 (79 – 106)	-	-	-	-	-	-
Gensl	130 (115 – 140)	-	-	-	-	-	-
st5-st5	84 (78 – 90)	-	-	-	-	-	84 (81 – 92)
Gensw post.cor.	95 (83 – 100)	-	-	97	-	77	-
Lisl	26 (23 – 28)	-	-	-	-	-	-
Lisw	4 (3 – 5)	-	-	-	-	-	-
Sisl	14 (13 – 15)	-	-	-	-	-	-
Vsl	109 (91 – 135)	-	-	-	-	91	105
Vsw ZV2	66 (63 – 70)	-	-	-	-	65	70
Vsw anus	54 (50 – 60)	-	-	-	-	-	65
Dist. solen. vas	24 (20 – 33)	-	-	-	-	-	-
JV5	23 (20 – 25)	-	24 (23 – 24)	-	-	-	-
SgeI	35 (30 – 38)	-	-	-	-	-	35
SgeII	15 (13 – 20)	-	-	-	-	-	14
SgeIII	29 (25 – 30)	-	-	-	24	-	30
StiIII	30 (28 – 35)	-	-	-	27	-	35
SgeIV	42 (37 – 47)	-	34 (33 – 34)	37	34	34	37
StiIV	54 (50 – 55)	-	46 (45 – 46)	48	41	43	50 (49 – 52)
StIV	48 (43 – 53)	-	45 (45 – 46)	42	37	43	48 (47 – 50)
Scl	38 (33 – 47)	-	-	28	-	-	25
Scw	4 (2 – 7)	-	-	-	-	-	-
Fdl	30 (28 – 33)	-	-	-	-	-	-
No teeth Fd	11	-	-	12 – 13	12	11	-
Mdl	34 (33 – 36)	-	-	-	-	-	-
No teeth Md	3	3	3	1 – 2	3	2	-

Sources of measurements – Indonesia: Ehara (2005) (but identified under the name *Indoseiulus armellae*, synonymized by Denmark & Kolodochka, 1993); Japan: Ehara (1982); Papua New Guinea: Schicha & Gutierrez (1985) (but identified under the name *Indoseiulus armellae*, synonymized by Denmark & Kolodochka, 1993); Philippines: Corpuz-Raros (2005); Taiwan: Tseng (1983); Thailand: Moraes *et al.* (2004b); - : not provided.

thrips within the first 24 hrs and laid an average number of 1.6 - 6.8 eggs within 24-48 hrs. These data suggest that *G. liturivorus* would be a promising BCA against thrips (Mochizuki 2009).

All measurements obtained on female specimens collected during this study (Table 10) agree well with measurements obtained from female specimens collected in other countries, especially with type specimens from Japan and Thailand. Specimens of Taiwan have shorter setae while specimens of Indonesia and Papua New Guinea have longer setae.

It is interesting to notice that in deutonymph specimens some setae are longer than those of female specimens, especially *j1*, *j3*, *Z5*, *SgeII*, *SgeIV*, *StIV*, shorter peritremes and chelicerae and a lower number of teeth than female specimens on both digits of chelicerae.

Genus *Euseius* Wainstein

Amblyseius (*Amblyseius*) section *Euseius*, Wainstein, 1962: 15; *Euseius* De Leon, 1967: 86.

Euseius ovaloides (Blommers)

Amblyseius (*Amblyseius*) *ovaloides* Blommers 1974: 147.

Euseius ovaloides Moraes et al. 1986: 51; 2004b: 78; Chant & McMurtry 2005a: 215; 2007: 121.

Specimens examined: Muoi in P6 plot (aasl 66 m, lat. 21°54'55" N, long. 104°38'22" E), 1 ♀ + 1 ♂ (in too bad state for doing measurements) on *Vernicia montana* (Euphorbiaceae), 20/VII/2017.

Table 11 Character measurements of adult females of *Euseius ovaloides* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam (1) 1, this study	Guadeloupe 4	La Réunion 2	Madagascar 1?	Characters	Vietnam (1) 1, this study	Guadeloupe 4	La Réunion 2	Madagascar 1?
Dsl	343	351 (341 – 360)	324 – 340	335	Gensl	123			
Dsw at s4 level	235	243 (236 – 253)	227 – 258	200	<i>st5-st5</i>	70	79 (76 – 82)	63 – 69	-
<i>j1</i>	33	32 (31 – 33)	28 – 34	27 – 30	Gensw post.cor.	105			
<i>j3</i>	10	9 (8 – 9)	9	8	Lisl	23	-	-	-
<i>j4</i>	5	6	6	5	Lisw	5	-	-	-
<i>j5</i>	5	6 (5 – 6)	6	5	Sisl	13	-	-	-
<i>j6</i>	7	7 (6 – 7)	-	6	Vsl	100	98 (93 – 105)	98 – 104	100
<i>J2</i>	7	8 (8 – 9)	6 – 9	7	Vsw ZV2	40	44 (42 – 47)	44 – 50	-
<i>J5</i>	5	6	6	5	Vsw anus	63	74 (71 – 78)	71 – 74	72
<i>r3</i>	10	10 (9 – 11)	9	8	<i>JV5</i>	25	-	-	22
<i>R1</i>	8	6 (6 – 7)	8 – 9	8	<i>SgeI</i>	19	14 (12 – 16)	-	-
<i>s4</i>	12	11 (11 – 12)	13	9 – 12	<i>SgeII</i>	15	15 (12 – 17)	-	-
<i>S2</i>	10	7 (6 – 8)	8 – 9	8	<i>SgeIII</i>	25	26 (25 – 27)	22 – 28	25
<i>S4</i>	11	8 (7 – 8)	8 – 9	8	<i>StI</i>	23	16 (14 – 19)	19 – 22	20
<i>S5</i>	10	6 (6 – 7)	6 – 8	7	<i>SgeIV</i>	35	40 (37 – 42)	38 – 39	36 – 40
<i>z2</i>	10	8 (7 – 9)	8 – 9	7	<i>StIV</i>	33	36 (34 – 38)	35	29 – 34
<i>z4</i>	8	8 (7 – 9)	8 – 9	8	<i>StIV</i>	55	62 (59 – 66)	47 – 60	52 – 57
<i>z5</i>	5	6	6	7	ScI	14	14 (14 – 15)	9 – 13	11
<i>Z1</i>	8	7 (6 – 8)	8 – 9	8	Scw	2	-	-	1
<i>Z4</i>	9	7 (6 – 8)	6 – 9	8	Fdl	23	29 (27 – 31)	25	28
<i>Z5</i>	48	53 (52 – 54)	50 – 54	44 – 48	No teeth Fd	-	-	2	1
<i>st1-st1</i>	58	-	-	-	Mdl	23	30 (27 – 32)	25	26
<i>st2-st2</i>	53	68 (66 – 68)	61 – 63	-	No teeth Md	-	-	1	1
<i>st3-st3</i>	75	-	-	-					
<i>st1-st3</i>	75	58 (56 – 61)	50	-					
<i>st4-st4</i>	95	-	-	-					

Sources of measurements – Guadeloupe: Moraes et al. (2000); La Réunion: Moraes et al. (2012); Madagascar: Blommers (1976); - : not provided.

Remarks: All measurements obtained on the single female specimen collected during this study agree well with measurements obtained from female specimens collected in other countries.

Euseius ovaloides was described by Blommers (1974) from specimens collected on *Citrus hystrix* and *Persea americana* in Madagascar. Like all *Euseius* species, this species belongs to the type IV (polliniphagous generalist predators) of McMurtry and Croft (1997) and McMurtry *et al.* (2013). The species has been occasionally recorded from Madagascar (Blommers, 1974), Papua-New Guinea (Schicha and Gutierrez 1985), Seychelles (Schicha, 1987), Reunion Island, (Quilici *et al.*, 1997, 2000), Guadeloupe, Martinique and Marie-Galante (Moraes *et al.*, 2000; Kreiter *et al.*, 2006) on various plants, though its biology remains unknown. It is suspected to be a poor predator of tetranychid mites (Gutierrez and Etienne, 1986) but can be considered as a potentially good predator of thrips and of whiteflies. This is one of the more common species on La Réunion Island.

This is the first mention of that species in Vietnam.

Measurements of the single female specimen collected during this study are provided in table 11 and compared with measurements obtained from specimens collected in other countries of Indian Ocean and the Caribbean.

Subfamily Phytoseiinae Berlese

Phytoseiini Berlese 1913: 3; Phytoseiinae, Vitzthum 1941: 768.

Genus *Phytoseius* Ribaga

Phytoseius Ribaga, 1904: 177.

Phytoseius coheni Swirski & Shechter

Phytoseius (Dubininellus) macropilis coheni Swirski & Shechter 1961: 104.

Phytoseius (Phytoseius) macropilis coheni, Ehara 1966: 26.

Phytoseius (Dubininellus) coheni, Swirski & Golan 1967: 226; Wu 1997: 153.

Phytoseius (Phytoseius) coheni, Moraes *et al.* 1986: 219.

Phytoseius coheni, Moraes *et al.* 2004a: 235; Chant & McMurtry 2007: 129.

Phytoseius hawaiiensis Prasad 1968: 1460 (synonymy according to Denmark & Evans 2011: 301).

Phytoseius huangi Ehara 1970: 62 (synonymy according to Ehara 2002: 40).

Phytoseius jianfengensis Chen, Chu & Zhou 1980: 15 (synonymy according to Wu 1997: 153).

Specimens examined: Muoi in PM1 plot (asl 66 m, lat. 21°54' N, long. 104°38' E), 1 ♀ on an unknown plant support, 17/V/2017; in P7 plot (asl 66 m, lat. 21°54'51" N, long. 104°38'22" E), 1 ♀ on an unknown plant support, 21/VII/2017; in P16 plot (asl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 1 ♀ on *Tephrosia candida* (Fabaceae), 22/VII/2017.

Remarks: This species was described from Hong-Kong by Swirski and Shechter (1961) collected on a wide range of plants and very common on citrus.

Although species of the genus *Phytoseius* are considered to belong to the type III (polyphagous generalist predators) of McMurtry and Croft (1997) and McMurtry *et al.* (2013), its specific biology is totally unknown.

This is the first mention of that species in Vietnam.

Measurements of specimens collected during this study are provided in table 12 and compared with measurements obtained from specimens collected in other countries of Asia and Indian Ocean. Measurements obtained during this study agree well with most obtained from female specimens collected in other countries.

Table 12 Character measurements of adult females of *Phytoseius coheni* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	Vietnam 3, this study	Type material Hong-Kong 1	Mauritius 2	Philippines ?	Taiwan ?	Thailand ?
Dsl	293 (275 – 310)	269 – 292	293 (289 – 297)	279	295 – 314	280
Dsw at s4 level	157 (138 – 175)	142	157 (153 – 160)	160	180 – 249	170
j1	26 (25 – 28)	29	26 (24 – 28)	26 – 31	28 – 29	26
j3	20 (18 – 20)	18 – 23	20 (19 – 21)	20 – 23	23 – 25	22
j4	9 (8 – 10)	8	10 (9 – 10)	7 – 12	7 – 8	9
j5	8	8	9	7 – 13	6 – 8	8
j6	10	8	9	8 – 13	8 – 9	9
J5	9 (8 – 10)	13	8 (7 – 8)	8 – 13	8	6
r3	37 (33 – 40)	32	38 (36 – 39)	34 – 39	30 – 41	36
s4	82 (75 – 93)	66 – 81	74 (72 – 76)	78 – 86	90 – 95	80
s6	65 (63 – 70)	58 – 74	60 (57 – 63)	65 – 73	82 – 90	67
z2	12 (10 – 13)	13	12	13 – 14	14 – 15	13
z3	26 (25 – 28)	27	27 (24 – 29)	26 – 28	29 – 30	28
z4	12 (11 – 13)	13	13	12 – 14	15	13
z5	10	10	10 (9 – 10)	10 – 13	8	10
Z4	54 (50 – 58)	51 – 64	55 (50 – 59)	54 – 59	56 – 65	53
Z5	63 (55 – 70)	58 – 71	63 (59 – 66)	70 – 73	70 – 78	61
st1-st1	52 (50 – 55)	-	-	-	-	-
st2-st2	59 (55 – 63)	-	60 (59 – 60)	-	-	-
st3-st3	73 (65 – 80)	-	-	-	-	-
st1-st3	54 (53 – 55)	-	59 (58 – 59)	-	-	-
st4-st4	80 (63 – 98)	-	-	-	-	-
gensl	98 (95 – 100)	-	97	-	-	-
st5-st5	62 (60 – 63)	-	72 (69 – 74)	-	-	-
gensw post. cor.	73 (70 – 75)	-	-	-	-	-
Lisl	27 (25 – 28)	-	28 (27 – 29)	-	16	-
Lisw	2	-	-	-	3	-
Vsl	94 (80 – 103)	86 – 99	98 (95 – 101)	99	98	-
Vsw ZV2	44 (38 – 50)	41 – 48	45 (41 – 49)	44	85	-
Vsw anus	49 (43 – 53)	48 – 53	53 (52 – 53)	51	-	-
JV5	44 (43 – 45)	38 – 46	41 (39 – 43)	47 – 52	53	44
Scl	11 (10 – 13)	-	15 (14 – 16)	-	6	-
Scw	5	-	-	-	-	-
SgeIV	24 (23 – 25)	18 – 23	20 (19 – 20)	20 – 24	15 – 16	21
StiIV	44 (40 – 50)	38 – 51	42 (40 – 43)	42 – 44	45 – 48	41
StIV	24 (23 – 25)	20 – 25	25 (23 – 26)	23 – 27	23 – 40	23
SttIV	20	18 – 25	21 (20 – 21)	20	30	21
Fdl	24 (23 – 25)	23 – 25	-	-	-	-
No teeth Fd	3	3	-	-	-	-
Mdl	24 (23 – 25)	23 – 25	-	-	-	-
No teeth Md	1	1	-	-	-	-

Sources of measurements – Type material Hong-Kong: Swirski & Shechter (1961), Denmark (1966); Mauritius: Ferragut & Baumann (2019); Philippines: Swirski & Golan (1967) and (identified as *Phytoseius hawaiensis*) Corpuz-Raros & Garcia (1994); Taiwan: Lo (1970) & Tseng (1976); Thailand: (identified as *Phytoseius hawaiensis*) Ehara & Bhandhfalck (1977); - : not provided.

***Phytoseius hongkongensis* Swirski & Shechter**

Phytoseius (Phytoseius) hongkongensis Swirski & Shechter 1961: 99; Amitai & Swirski 1966: 22.

Phytoseius (Pennaseius) hongkongensis Ehara 1966: 25; Ehara 1972: 169; Moraes *et al.* 1986: 211.

Phytoseius hongkongensis Moraes *et al.* 2004a: 240; Chant & McMurtry 2007: 129.

Specimens examined: Muoi in P16 plot (aasl 66 m, lat. 21°54'46" N, long. 104°38'18" E), 2 ♀♀ on *Chromolaena odorata* (Asteraceae), 18/V/2017; in P11 plot (aasl 66 m, lat. 21°54'30" N, long. 104°38'14" E), 1 ♀ on *Mallotus canii* (Euphorbiaceae), 31/V/2017; in P14 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'45" E), 2 ♂♂ on *C. odorata* (Asteraceae) and 1 ♂ in an unknown host plant, 2/VI/2017; in P15 plot (aasl 66 m, lat. 21°54'56" N, long. 104°38'46" E), 1 ♀ on *C. odorata* (Asteraceae), 2/VI/2017 and 1 ♀ + 1 ♂ on *Rubus alceifolius* (Rosaceae), 3/VI/2017.

Remarks: This species was described from specimens collected on *Heterosmilax gaudichaudiana* and *Urena lobata* in Victoria Mount Forest, Hong Kong Island, Hong Kong.

Although species of the genus *Phytoseius* are considered to belong to the type III (polyphagous generalist predators) of McMurtry and Croft (1997) and McMurtry *et al.* (2013), its specific biology is totally unknown.

This is the first mention of that species in Vietnam.

Measurements of specimens collected during this study are provided in table 13 for females and males and compared with measurements obtained from specimens collected in other countries of Asia and Africa.

Measurements obtained agree well with most of those obtained previously, with slightly shorter dimensions, especially in male specimens.

***Phytoseius tixierae* Kreiter n. sp.**

Zoobank: [7B2DE1ED-FE36-488E-87B2-969ADBB09507](#)

Figs. 4 a – d, 5 a – b, 6 a – c

Diagnose — *Phytoseius tixierae* Kreiter n. sp. lacks setae *J2* and *R1* and consequently belongs to the *horridus* species group. This species is unique by the following characters: *z2* and *Z4* smooth, *s6* only slightly serrated, setae *j4*, *j5*, *j6*, *J5* also smooth, setae *s6*, *Z4* and *Z5* subequal, *s6* being the longest seta, four macrosetae on leg IV spatulate, *SgeIV* being the shortest and *StlIV* being the longest, with *StlIV* and *SttIV* being intermediate and subequal, setae *j1* and *j3* almost equal, setae *JV5* very short, < 20, peritreme reaching the level of setae *j1*.

Description of the adult female of *Phytoseius tixierae* Kreiter n. sp.

n = 4 (Figs. 4 a – b, Figs. 5 a – b)

(in bold, average measurements)

Dorsum — (Fig. 4a) Dorsal shield rugose **271** (250 – 280) long and **130** (125 – 140) wide, smooth, with no visible solenostome, 8 pairs of poorly visible poroids, 14 pairs of dorsal setae and one pair of sub-lateral setae on the dorsal shield: *j1* **23** (21 – 23), *j3* **18** (15 – 23), *j4* **10** (10 – 11), *j5* **10** (10 – 11), *j6* **11** (10 – 13), *J5* **9** (8 – 9), *z2* **12** (10 – 13), *z3* **20** (18 – 25), *z4* **19** (18 – 23), *z5* **12** (9 – 13), *Z4* **57** (48 – 63), *Z5* **53** (48 – 55), *s4* **44** (43 – 45), *s6* **59** (55 – 63), *r3* **30** (28 – 30). Setae *z2*, *j4*, *j5*, *j6*, *z5* and *Z4* smooth, setae *s6* very slightly serrated at the distal third of setae, setae *j1*, *j3*, *z3*, *z4*, *r3*, *s4* and *Z5* serrated.

Peritreme — (Fig. 4a) Extending to the level of setae *j1*.

Venter — (Fig. 4b) All shields smooth.

Sternal shield with three pairs of setae and two pairs of lyrifissures; one pair of sternal setae on metasternal shields with a pair of pores; posterior margin straight. Distances between *st1-st1* **46** (43 – 50), *st2-st2* **53** (50 – 55), *st3-st3* **59** (43 – 68), *st1-st3* **58** (55 – 60), *st4-st4* **62** (50 – 75).

Genital shield length **101** (98 – 105), *st5-St5* **56** (50 – 60), width at the level of the posterior corners of the genital shield **80** (78 – 82).

Table 13 Character measurements of adult females and males of *Phytoseius hongkongensis* collected in this study and those reported in previous studies (localities followed by the number of specimens measured).

Characters	♀					♂	
	Vietnam 5, this study	Africa 4	Type material Hong-Kong, 2	Taiwan ?	Thailand 5	Vietnam 4, this study)	Hong-Kong 1
	Dsl	285 (260 – 300)	274 (262 – 285)	264 – 277	304	274 (264 – 294)	217 (210 – 228)
Dsw at s4 level	125 (115 – 130)	138 (133 – 145)	134 – 140	196	145 (135 – 150)	115 (110 – 123)	
j1	25 (23 – 25)	24 (22 – 26)	25 – 28	30	28 (25 – 30)	19 (19 – 20)	20
j3	65 (63 – 68)	67 (59 – 71)	64 – 69	70	72 (70 – 75)	45 (45 – 46)	53
j4	5	5 (5 – 6)	3 – 8	9	5 (4 – 5)	4	4 – 8
j5	5	5 (5 – 6)	3 – 8	9	4 (4 – 5)	4	4 – 8
j6	5	6 (5 – 7)	3 – 8	9	6 (5 – 6)	4	4 – 8
J2	9 (8 – 10)	9 (8 – 10)	9 – 12	9	9 (7 – 10)	5	-
J5	6 (5 – 8)	7 (6 – 8)	6 – 9	-	6 (6 – 7)	5	4 – 8
r3	35 (30 – 40)	42 (40 – 43)	36 – 44	45	45 (43 – 47)	30 (29 – 30)	36
R1	13 (10 – 15)	16 (14 – 17)	13 – 18	9	16 (15 – 17)	9 (8 – 10)	12
s4	95 (90 – 98)	96 (90 – 105)	91 – 102	95	103 (102 – 106)	65 (63 – 68)	76
s6	75 (75 – 78)	81 (77 – 86)	75 – 77	91	84 (80 – 90)	45 (43 – 45)	51
z2	13 (13 – 15)	12 (10 – 13)	10 – 15	13	17 (16 – 18)	13 (11 – 14)	10
z3	44 (43 – 45)	45 (45 – 46)	38 – 40	58	44 (39 – 47)	32 (30 – 33)	36
z4	10 (8 – 10)	8	10 – 13	18	10 (8 – 12)	10 (9 – 10)	9
z5	5	4 (3 – 5)	3 – 8	-	4	3	4 – 8
Z4	72 (68 – 75)	74 (67 – 82)	76 – 90	75	79 (75 – 82)	39 (37 – 40)	53
Z5	69 (67 – 70)	74 (67 – 85)	55 – 75	83	80 (75 – 84)	32 (30 – 33)	36
st1-st1	61 (58 – 63)	-	-	-	-	50 (48 – 53)	-
st2-st2	68 (65 – 70)	66	-	-	69 (66 – 70)	55 (55 – 56)	-
st3-st3	73 (70 – 75)	-	-	-	-	63 (63 – 64)	-
st1-st3 ♀ / st1-st5 ♂	59 (58 – 60)	60 (58 – 63)	-	-	61 (60 – 62)	103 (100 – 108)	-
st4-st4	80 (73 – 83)	-	-	-	-	54 (53 – 57)	-
Gensl	113 (110 – 115)					Not applicable	
st5-st5	61 (58 – 63)	60 (58 – 62)	-	-	64 (63 – 66)	41 (38 – 43)	-
Gensw post. corn.	64 (60 – 68)						
Lisl	19 (18 – 20)	-	-	50	-	Not applicable	
Lisw	2	-	-	2	-		
Vsl	92 (83 – 100)	89 (85 – 93)	-	93	97 (95 – 100)	82 (78 – 90)	-
Vsw ZV2	50 (49 – 50)	53 (50 – 60)	-	53	55 (52 – 58)	128 (125 – 130)	-
Vsw anus	46 (43 – 48)	44 (43 – 45)	-	-	46 (44 – 49)	52 (50 – 55)	-
JV5	48 (45 – 50)	-	46 – 61	63	-	19 (15 – 23)	25
Scl	15 (13 – 18)	12 (10 – 18)	-	10	10 (9 – 12)	18 (18 – 19)	23
Scw	14 (10 – 18)	-	-	-	-	19 (18 – 20)	23
SgeIV	26 (25 – 28)	25 (24 – 26)	26 – 33	30	29 (28 – 31)	23	29
StiIV	30	31 (29 – 32)	29 – 33	30	32 (31 – 32)	25	-
StIV	25 (23 – 25)	24 (22 – 24)	21 – 25	30	26 (24 – 28)		
SttIV	29 (28 – 33)	24 (22 – 25)	26	30	27 (25 – 28)	Not applicable	
Fdl	27 (25 – 30)	25	-	-	26 (25 – 27)	17 (15 – 18)	-
No teeth Fdl	2	2	-	-	-	2	-
Mdl	28 (25 – 30)	26	-	-	29 (29 – 30)	19 (18 – 20)	-
No teeth Mdl	1	1	-	-	-	1	-
Shaft			Not applicable			14 (13 – 15)	-
Toe						6 (4 – 8)	-

Sources of measurements – For ♀♀: Africa (Benin: 1♀, Kenya: 2♀, Malawi: 1♀): Ueckermann *et al.* (2007); Hong-Kong: Swirski & Shechter (1961), Denmark (1966) & Ehara & Lee (1971); Taiwan: Tseng (1976). For ♂♂: Hong-Kong: Swirski & Shechter (1961). - : not provided.




Figure 4 Female of *Phytoseius tixierae* Kreiter n. sp.. a – Dorsal shield and peritreme; b – Ventral shields.

One pair of narrow metapodal shields **24** (20 – 28) long and **2** wide.

Ventrianal shield with three pairs of preanal setae (*JV1*, *JV2* and *ZV2*) and a pair of pores. Membrane surrounding ventrianal shield with two pairs of setae (*ZV3* and *JV5*), and no visible poroids in our specimens; ventrianal shield **92** (85 – 98) long, **51** (50 – 53) wide at level of anterior corners (*ZV2*), and **44** (43 – 45) wide at level of *anus*.

JV5 short, **19** (15 – 20) long, and serrated.

Chelicera — Chelicerae not clearly visible and therefore not drawn. Fixed digit **17** (14 – 20) long with apparently 2 teeth; and movable digit **18** (15 – 20) long with apparently 1 tooth.

Spermatheca — (Fig. 5a) Spermatheca with calyx pocular (Denmark and Evans 2011), 10 long and 5 wide, and a small atrium. Ductus minor not visible but large ductus major well visible in all specimens.

Legs — (Fig. 5b) Four macrosetae on leg IV, all spatulate: *SgeIV* 10, *StIIV* **39** (38 –




Figure 5 Female of *Phytoseius tixierae* Kreiter n. sp.. a – Insemination apparatus; b – Genu, tibia and basitarsus of the leg IV.

40), *StIV* 21 (20 – 23), *SttIV* 19 (18 – 20). Genu II with 7 setae and Genu III with 6 setae, chaetotactic formula of genu II: 2-2/0, 2/0-1; genu III: 1-2/0, 2/0-1.

Description of the adult male of *Phytoseius tixierae* Kreiter n. sp.

n = 1 (Figs 6 a – c)

Dorsum — (Fig. 6a) Dorsal shield rugose 200 long and 113 wide, with no visible solenostome. The dorsal shield bears 14 pairs of dorsal setae and 1 pair of sub-lateral setae situated on the dorsal shield: *j1* 20, *j3* 16, *j4* 10, *j5* 8, *j6* 9, *J5* 6, *z2* 10, *z3* 18, *z4* 18, *z5* 10, *Z4* 30, *Z5* 28, *s4* 35, *s6* 35, *r3* 25. Setae *z2*, *j4*, *j5*, *j6*, *z5* and *Z4* smooth, setae *s6* very slightly serrated, setae *j1*, *j3*, *z3*, *z4*, *r3*, *s4* and *Z5* serrated, like in the female.

Peritreme — (Fig. 6a) Extending to the level of setae *j1*.

Venter — (Fig. 6b) Sternal shield smooth. Distances between *st1* – *st1* 43, *st2* – *st2* 48, *st3* – *st3* 50, *st1* – *st5* 90, *st4* – *st4* 43, *st5* – *st5* 35. Ventrianal shield with three pairs of pre-anal setae, *JV1*, *JV2* and *ZV2*. Pre-anal pores absent. Membrane surrounding ventrianal shield with one pair of setae *JV5*; ventrianal shield 75 long, 90 wide at anterior corners and 50 wide at level of paranal setae. Short setae *JV5* serrated, 10 long.

Chelicera — Fixed digit 15 long, with 2 teeth visible and movable digit 15 long with 1 tooth visible. Spermatodactyl circumflex accent shaped, shaft (Fig. 6c) 11, toe 9.

Legs — Legs IV with four spatulate macrosetae like in the female: *SgeIV* 5, *SttIV* 10, *StIV* 15, *SttIV* 15. Chaetotactic formula of genu II and III similar to that of females.

Specimens examined — Muoi in P4 experimental plot (aasl 66 m, lat. 21°54'41" N, long. 104°38'7" E), 1 ♀ + 1 ♂, 12/05/2017 and in P12-P13 plots (aasl 66 m, lat. 21°54'50" N, long. 104°38'45" E), 3 ♀♀ on *Bambusodae* sp. (Poaceae).

Type material — The holotype female, 3 paratype females, and 1 paratype male on 3 slides deposited in CBGP, in Montpellier SupAgro Acarology collection, France.

Etymology — The name “*tixierae*” refers to our colleague Professor-Dr Marie-Stéphane Tixier (Montpellier SupAgro) to whom this species is dedicated for her huge contribution to




Figure 6 Male of *Phytoseius tixierae* Kreiter n. sp.. a – Dorsal shield and peritreme; b – Ventral shields; c – Spermatodactyl.

the knowledge of mites and especially to the taxonomy of Phytoseiidae and her great assistance with this paper.

Remarks — Measurements of specimens collected during this study are provided in table 14 for females and for the single male and compared with measurements obtained from specimens of the closest described species.

Females of *Phytoseius tixierae* Kreiter n. sp. (Table 14) resembles that of *P. huaxiensis* Xin, Liang and Ke but differs in having *s6* only slightly serrated, shorter *s4*, *s6*, *Z4*, *Z5* and *JV5*. It also resembles *P. indicus* Bhattacharya but differs in having *s6* only slightly serrated, shorter

s4, s6, Z4, JV5, Z4 not serrated but *z4* serrated and longer. It is also related to *P. kazusanus* Ehara but differs in having *s6* only slightly serrated, *z4* serrated, longer *Z5, JV5* and *StIV* and *Z4* not serrated. It is also very close to *P. longchuanensis* Wu but differs from this species by having longer setae except *s4* and *Z4*, *s6* only slightly serrated, *z4* serrated, longer *JV5* and 4 macrosetae on the leg IV instead of zero. It is also very similar to *P. rimandoi* Corpuz but differs from this species in having shorter *s6, Z4* smooth. Finally, the closest species is *P. nudus* Wu and Li but the new species differs from this species in having *z4* strongly serrated (only slightly serrated in the former), and the dorsal shield rugose (smooth in the former).

Males of *Phytoseius tixierae* Kreiter n. sp. (Table 14) also resembles to the males of *P. huaxiensis* and *P. kazusanus* that are the only species for which males have been described. But males of the new species differ from males of the former species in having *s6* only slightly

Table 14 Character measurements of adult females and one adult male of *Phytoseius tixierae* Kreiter n. sp. collected in this study and close species reported in previous studies (localities followed by the number of specimens measured).

Characters	♀							♂		
	<i>Phytoseius tixierae</i> Kreiter n. sp.	<i>Phytoseius</i> <i>huaxiensis</i>	<i>Phytoseius</i> <i>indicu</i> s	<i>Phytoseius</i> <i>kazusanus</i>	<i>Phytoseius</i> <i>longchuanensis</i>	<i>Phytoseius</i> <i>nudus</i>	<i>Phytoseius</i> <i>rimandoi</i>	<i>Phytoseius</i> Kreiter n. sp.	<i>Phytoseius</i> <i>huaxiensis</i>	<i>Phytoseius</i> <i>kazusanus</i>
	4	Xin, Liang & Ke	1?	8	2	?	1?	1	2	?
Dsl	271 (250 – 280)	-	260 – 299	284	268	285	332	200	222	228
Dsw at <i>s4</i> level	130 (125 – 140)	-	136 – 145	153	196	145	186	113	118	144
Ds	rugose	rugose	rugose	rugose	rugose	smooth	rugose	smooth	?	?
<i>j1</i>	23 (21 – 23)	28	20 – 27	25 (25 – 26)	17	23	26	20	21	19
<i>j3</i>	18 (15 – 23)	20	22 – 27	19 (18 – 19)	17	18	16	16	21	17
<i>j4</i>	10 (10 – 11)	10	8 – 9	9	7	13	8 to 12	10	9	8
<i>j5</i>	10 (10 – 11)	8	6 – 12	8 (8 – 9)	7	13	8 to 12	8	7	8
<i>j6</i>	11 (10 – 13)	11	9 – 12	10	7	13	8 to 12	9	10	8
<i>J5</i>	9 (8 – 9)	9	8 – 9	9 (9 – 10)	7	9	8 to 12	6	7	7
<i>r3</i>	30 (28 – 30)	36	30 – 36	31 (31 – 32)	24	33	36	25	28	26
<i>s4</i>	44 (43 – 45)	57	60 – 72	48 (47 – 49)	55	43	48	35	43	37
<i>s6</i>	59 (55 – 63) sl. ser.	70 ser.	60 – 72 ser.	59 (58 – 61) ser.	58 ser.	61 ser.	58 ser.	35	48	36
<i>z2</i>	12 (10 – 13) not ser.	13 not ser.	10 – 12 not ser.	14 (13 – 14) not ser.	14 not ser.	13 not ser.	8 to 12 not ser.	10	11	13
<i>z3</i>	20 (18 – 25)	35	26 – 29	29 (28 – 29)	34	20	24	18	31	23
<i>z4</i>	19 (18 – 23) ser.	17 ser.	10 not ser.	16 (16 – 17) ser.	10 not ser.	20 serrul.	22 ser.	18	19	14
<i>z5</i>	12 (9 – 13)	11	8 – 9	11 (10 – 11)	7	13	8 to 12	10	10	9
<i>Z4</i>	57 (48 – 63) not ser.	72 not ser.	48 – 52 ser.	63 (62 – 64) ser.	69 not ser.	53 not ser.	56 ser.	30	43	37
<i>Z5</i>	53 (48 – 55)	69	52 – 65	62 (61 – 62)	38 – 41	56	52	28	37	35
<i>st1-st1</i>	46 (43 – 50)	-	-	-	-	-	-	43	-	-
<i>st2-st2</i>	53 (50 – 55)	-	-	-	-	-	-	48	-	-
<i>st3-st3</i>	59 (43 – 68)	-	63	-	-	-	-	50	-	-
<i>st1-st3 ♀ / st1-st5 ♂</i>	58 (55 – 60)	-	63	-	-	-	-	90	-	-
<i>st4-st4</i>	62 (50 – 75)	-	-	-	-	-	-	43	-	-
Gensl	101 (98 – 105)	-	-	-	-	-	-	Not applicable		
<i>st5-st5</i>	56 (50 – 60)	-	-	-	-	-	-	35	-	-
Gensw post. corn.	80 (78 – 82)	-	-	-	-	-	-	Not applicable		
Lisl	24 (20 – 28)	-	-	-	-	-	-	Not applicable		
Lisw	2	-	-	-	-	-	-	-		
Vsl	92 (85 – 98)	-	90	-	-	-	-	75	-	-
Vsw ZV2	51 (50 – 53)	-	45	-	-	-	-	90	-	-
Vsw anus	44 (43 – 45)	-	-	-	-	-	-	50	-	-
<i>JV5</i>	19 (15 – 20) ser.	44	36 ser.	41 (40 – 41)	10	20	-	10	19	16
<i>SgeIV</i>	10 spat.	-	12 – 20 spat.	10 (9 – 10) spat.	-	14 spat.	6 spat.	5	-	8
<i>StIIV</i>	39 (38 – 40) spat.	-	32 – 40 spat.	31 (31 – 32) spat.	No macrosetae	43 spat.	42 spat.	7	-	11
<i>StIV</i>	21 (20 – 23) spat.	-	18 – 20 spat.	26 (26 – 27) spat.		21 spat.	24 spat.	13	-	19
<i>StIV</i>	19 (18 – 20) spat.	-	22 spat.	21 (20 – 21) spat.		21 spat.	24 spat.	15	-	16
Scl	10	-	-	-	-	-	-	Not applicable		
Scw	5	-	-	-	-	-	-	-		
Fdl	17 (14 – 20)	-	-	-	-	-	-	15	-	-
No teeth Fd	2	-	-	-	-	-	-	3	2	-
Mdl	18 (15 – 20)	-	-	-	-	-	-	15	-	-
No teeth Md	1	-	-	-	-	-	-	1	1	-
Shaft	Not applicable							11	-	-
Toe	Not applicable							9	-	-

Sources of measurements – For ♀♀: *Phytoseius huaxiensis*: Xin et al. (1982); *P. indicus* (= *P. neglectus* Gupta): Bhattacharya (1968) and Gupta (1969); *P. kazusanus*: Ehara in Ehara et al. (1994); *P. longchuanensis*: Wu (1997); *P. nudus*: Wu & Li (1984); *P. rimandoi*: Corpuz (1966). For ♂♂: *P. huaxiensis*: Xin et al. (1982); *P. kazusanus*: Ehara in Ehara et al. (1994). - : not provided.

serrated, and shorter *s4*, *s6*, *z3*, *Z4*, *Z5* and *JV5*. Males of the new species differ from males of *P. kazusanus* in having *s6* only slightly serrated, shorter *Z4*, *Z5* and *JV5*, *z4* serrated and longer and *Z4* not serrated.

Conclusion

The fauna of Phytoseiidae of Vietnam is now composed of 20 species, 12 already known from previous studies and 8 added in this paper. As this number of species is still far below the anticipated real number of phytoseiid species present in this tropical biodiversity hotspot, follow-up survey work is urgently needed.

Acknowledgements

We are very grateful to Nguyễn Tuân Cường for identification of plant species on which phytoseiid mites were collected. The International Center for Tropical Agriculture (CIAT, CGIAR) graciously provided logistical support for the in-country visit of M.-C. Bopp and facilitated the necessary permits for field visits. Financial support for the field collections in Vietnam was provided through the CGIAR research program on Roots, Tubers and Bananas - CRP-RTB and the CGIAR research program on Climate Change, Agriculture and Food Security - *i.e.*, CRP-CCAFS. We are also very grateful to Professor-Dr Marie-Stéphane Tixier (Montpellier SupAgro) for very valuable comments of an earlier version of this manuscript.

References

- Akyazi R., Ueckermann E.A., Soysal M. 2016. The new distribution of *Amblyseius herbicolus* in Turkey (Parasitiformes, Phytoseiidae) with a key of *Amblyseius* species found in Turkey. *Acarologia*, 56(2): 237-244. doi:[10.1051/acarologia/20162241](https://doi.org/10.1051/acarologia/20162241)
- Amitai S., Swirski E. 1966. Illustrations of spermathecae in several previously described phytoseiid mites (Acarina) from Hong Kong and Israel. *The Israel J. Agric. Res.*, 16: 19-24.
- Athias-Henriot C. 1975. Nouvelles notes sur les Amblyseiini. II. Le relevé organotaxique de La face dorsale adulte (Gamasides protoadéniques, Phytoseiidae). *Acarologia*, 17(1): 20-29.
- Beard J.J. 2001. A review of Australian *Neoseiulus* Hughes and *Typhlodromips* De Leon (Acari: Phytoseiidae: Amblyseiinae). *Invert. Tax.*, 15: 73-158. doi:[10.1071/IT99017](https://doi.org/10.1071/IT99017)
- Berlese A. 1913. Systema Acarorum genera in familiis suis disposita. *Acaroteca Italica*, 1-2: 3-19.
- Berlese A. 1914. Acari nuovi. *Manipulus IX*. *Redia*, 10: 113-150.
- Bhattacharyya S.K. 1968. Two new phytoseiid mites from eastern India (Acarina: Phytoseiidae). *J. Bombay Nat. Hist. Soc.*, 65(3): 677-680.
- Blommers L. 1974. Species of the genus *Amblyseius* Berlese, 1914, from Tamatave, east Madagascar (Acarina: Phytoseiidae). *Bull. Zool. Mus. Univ. Amsterdam*, 3: 143-155.
- Blommers L. 1976. Some Phytoseiidae (Acarina: Mesostigmata) from Madagascar, with descriptions of eight new species and notes on their biology. *Bijdragen tot Dierkunde*, 46(1): 80-106. doi:[10.1163/26660644-04601005](https://doi.org/10.1163/26660644-04601005)
- Carmona M.M. 1968. Contribuicao para o estudo de alguns acaros fitofagos e depredadores, de Angola. *Agron. Lusit.*, 29: 267-288 + 12 plates.
- Chant D.A. 1957. Note on the status of some genera in the family Phytoseiidae (Acarina). *Can. Entomol.*, 89(11), 528-532. doi:[10.4039/Ent89528-11](https://doi.org/10.4039/Ent89528-11)
- Chant D.A. 1959. Phytoseiid mites (Acarina: Phytoseiidae). Part I. Bionomics of seven species in southeastern England. Part II. A taxonomic review of the family Phytoseiidae, with descriptions of thirty-eight new species. *Can. Entomol.*, 61(12): 1-166.
- Chant D.A., Hansell R.I.C. 1971. The genus *Amblyseius* (Acarina: Phytoseiidae) in Canada and Alaska. *Can. J. Zool.*, 49(5), 703-758. doi:[10.1139/z71-110](https://doi.org/10.1139/z71-110)
- Chant D.A., McMurtry J.A. 1994. A review of the subfamilies Phytoseiinae and Typhlodrominae (Acari: Phytoseiidae). *Intern. J. Acarol.*, 20(4): 223-310. doi:[10.1080/01647959408684022](https://doi.org/10.1080/01647959408684022)
- Chant D.A., McMurtry J.A. 2003a. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part I. Neoseiulini new tribe. *Intern. J. Acarol.*, 29(1): 3-46. doi:[10.1080/01647950308684319](https://doi.org/10.1080/01647950308684319)
- Chant D.A., McMurtry J.A. 2003b. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part II. The tribe Kampimodromini Kolodochka. *Intern. J. Acarol.*, 29(3): 179-224. doi:[10.1080/01647950308684331](https://doi.org/10.1080/01647950308684331)
- Chant D.A., McMurtry J.A. 2004. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part III. The tribe Amblyseiini Wainstein, subtribe Amblyseiina n. subtribe. *Intern. J. Acarol.*, 30(3): 171-228. doi:[10.1080/01647950408684388](https://doi.org/10.1080/01647950408684388)

- Chant D.A., McMurtry J.A. 2005a. A review of the subfamily Amblyseiina Muma (Acari: Phytoseiidae): Part V. Tribe Amblyseiini, subtribe Proprioseiopsina Chant & McMurtry. Intern. J. Acarol., 31(1): 3-22. doi:[10.1080/01647950508684412](https://doi.org/10.1080/01647950508684412)
- Chant D.A., McMurtry J.A. 2005b. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae) Part VI. The tribe Euseiini n. tribe, subtribes Typhlodromalina n. subtribe, Euseiina n. subtribe, and Ricoseiina n. subtribe. Intern. J. Acarol., 31(3): 187-224. doi:[10.1080/01647950508684424](https://doi.org/10.1080/01647950508684424)
- Chant D.A., McMurtry J.A. 2006. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part VIII. The tribes Macroseiini Chant, Denmark and Baker, Phytoseiulini n. tribe, Afroseiulini n. tribe and Indoseiulini Ehara and Amano. Intern. J. Acarol., 32(1), 13-25. doi:[10.1080/01647950608684439](https://doi.org/10.1080/01647950608684439)
- Chant D.A., McMurtry J.A. 2007. Illustrated keys and diagnoses for the genera and subgenera of the Phytoseiidae of the world (Acari: Mesostigmata). Indira Publishing House, West Bloomfield, 219pp.
- Chant D.A., Yoshida-Shaul E. 1991. Adult ventral setal patterns in the family Phytoseiidae (Acari: Gamasina). Intern. J. Acarol., 17(3): 187-199. doi:[10.1080/01647959108683906](https://doi.org/10.1080/01647959108683906)
- Chant D.A., Yoshida-Shaul E. (1992) Adult idiosomal setal patterns in the family Phytoseiidae (Acari: Gamasina). Intern. J. Acarol., 18(3): 177-193. doi:[10.1080/01647959208683949](https://doi.org/10.1080/01647959208683949)
- Chaudhri W.M. 1968. Six new species of mites of the genus *Amblyseius* (Phytoseiidae) from Pakistan. Acarologia, 10: 550-562.
- Chaudhri W.M., Akbar S., Rasool A. 1979. Studies on the predatory leaf inhabiting mites of Pakistan. University of Agriculture, Faisalabad, Pakistan, 243 pp.
- Chen S.-W., Chu C.-M., Zhou F.-W. 1980. On the phytoseiid mites of Guangdong (Acarina: Phytoseiidae). Journal of Jiangxi University, 4(1), 15-20 [in Chinese with English abstract]
- Corpuz L.A. 1966. Seven new species of mites of the genera *Typhlodromus* and *Phytoseius* (Phytoseiidae: Acarina). Philipp. Agricul., 50, 729-738.
- Corpuz L.A., Rimando L. 1966. Some Philippine Amblyseiinae (Phytoseiidae: Acarina). Philippine Agric., 50: 114-136.
- Corpuz-Raros L.A. 2005. Some new species records, discovery of males in two species and first report of *Wolbachia* infection in predatory mites (Phytoseiidae, Acari) from the Philippines. Philipp. Agric. Sc., 88(4), 431-439.
- Corpuz-Raros L.A., Garcia R.C. 1994. Two additional records and a key to the species of *Phytoseius* (Phytoseiidae, Acari) from the Philippines. Philippine Agric., 77(4): 473-479.
- Daneshvar H., Denmark H.A. 1982. Phytoseiids of Iran (Acarina: Phytoseiidae). Intern. J. Acarol., 8(1): 3-14. doi:[10.1080/01647958208683272](https://doi.org/10.1080/01647958208683272)
- De Leon D. 1967. Some mites of the Caribbean Area. Part I. Acarina on plants in Trinidad, West Indies. Allen Press Inc., Lawrence, Kansas, USA: 1-66.
- Demite P.R., McMurtry J.A., Moraes G.J. de. 2014. Phytoseiidae Database: a website for taxonomic and distributional information on phytoseiid mites (Acari). Zootaxa, 3795 (5): 571-577. doi: [10.11646/zootaxa.3795.5.6](https://doi.org/10.11646/zootaxa.3795.5.6)
- Demite P.R., Moraes G.J. de, McMurtry J.A., Denmark H.A., Castilho R.C. 2020. Phytoseiidae Database. Available from: www.lea.esalq.usp.br/phytoseiidae (last access 10/01/2020)
- Denmark H.A. 1966. Revision of the genus *Phytoseius* Ribaga, 1904 (Acarina: Phytoseiidae). Fla Dep. Agric. Bull., 6: 1-105.
- Denmark H.A., Evans G.A. 2011. Phytoseiidae of North America and Hawaii (Acari: Mesostigmata). Indira Publishing House, West Bloomfield, USA, 451 pp.
- Denmark H.A., Kolodochka L.A. 1993. Revision of the genus *Indoseiulus* Ehara (Acari: Phytoseiidae). Intern. J. Acarol., 19(3): 249-257. doi:[10.1080/01647959308683551](https://doi.org/10.1080/01647959308683551)
- Denmark H.A., Muma M.H. 1972. Some Phytoseiidae of Colombia (Acarina: Phytoseiidae). Fla Entomol., 55(1), 19-29. doi:[10.2307/3493637](https://doi.org/10.2307/3493637)
- Denmark H.A., Muma M.H. 1989. A revision of the genus *Amblyseius* Berlese, 1914 (Acari: Phytoseiidae). Occasional Papers of the Florida State Collection of Arthropods, USA, 4, 149 pp.
- Duarte M.V.A., Venzon M., Bittencourt M.C.de S., Rodriguez-Cruz F.A., Pallini A., Janssen A. 2015. Alternative food promotes broad mite control on chilli pepper plants. BioControl, 60: 817-825. doi:[10.1007/s10526-015-9688-x](https://doi.org/10.1007/s10526-015-9688-x)
- Ehara S. 1958. Three predatory mites of the genus *Typhlodromus* from Japan (Phytoseiidae). Annot. Zool. Japon., 31: 53-57.
- Ehara S. 1966. A tentative catalogue of predatory mites of Phytoseiidae known from Asia, with descriptions of five new species from Japan. Mushi, 39: 9-30.
- Ehara S. 1967. Phytoseiid mites from Okinawa Island (Acarina: Mesostigmata). Mushi, 40(6): 67-82.
- Ehara S. 1970. Phytoseiid mites from Taiwan (Acarina: Mesostigmata). Mushi, 43(6), 55-63.
- Ehara S. 1972. Some phytoseiid mites from Japan, with descriptions of thirteen new species (Acarina: Mesostigmata). Mushi, 46(12): 137-173.
- Ehara S. 1982. Two new species of phytoseiid mites from Japan (Acarina: Phytoseiidae). Appl. Entomol. Zool., 17: 40-45. doi:[10.1303/aez.17.40](https://doi.org/10.1303/aez.17.40)
- Ehara S. 1985. Five species of phytoseiid mites from Japan with descriptions of two new species (Acarina: Phytoseiidae). Zool. Sci., 2(1): 115-121.
- Ehara S. 2002. Some phytoseiid mites (Arachnida: Acari: Phytoseiidae) from west Malaysia. Species Div., 7: 29-46. doi:[10.12782/specdiv.7.29](https://doi.org/10.12782/specdiv.7.29)
- Ehara S. 2005. A collection of phytoseiid mites (Acarai: Phytoseiidae) from Java with description of a new species. Acta Arachnol., 54(1), 31-39. doi:[10.2476/asjaa.54.31](https://doi.org/10.2476/asjaa.54.31)
- Ehara S., Amano H. 1998. A revision of the mite family Phytoseiidae in Japan (Acari: Gamasina), with remarks on its biology. Species Div., 3(1): 25-73. doi:[10.12782/specdiv.3.25](https://doi.org/10.12782/specdiv.3.25)
- Ehara S., Amano H. 2002. Some Japanese phytoseiid mites (Acarai: Phytoseiidae) mostly from Ishigaki and Taketomi Islands. Entomol. Sci., 5(3): 321-329.

- Ehara S., Bhandhu Falck A. 1977. Phytoseiid mites of Thailand (Acarina: Mesostigmata). J. Fac. Educ., Tottori Univ., Natur. Sci., 27(2): 43-82.
- Ehara S., Gotoh T., Amano, H. 2000. Two Japanese species of the genus *Paraphytoseius* Swirski and Shechter (Acaria: Phytoseiidae). J. Acarol. Soc. Japan, 9(2): 113-118. doi:[10.2300/acari.9.113](https://doi.org/10.2300/acari.9.113)
- Ehara S., Lee L.H.Y. 1971. Mites associated with plants in Hong Kong. J. Fac. Educ., Tottori Univ., Natur. Sci., 22(2): 61-78.
- Ehara S., Okada Y., Kato H. 1994. Contribution to the knowledge of the mite family Phytoseiidae in Japan (Acaria: Gamasina). J. Fac. Educ., Tottori Univ., Natur. Sci., 42(2): 119-160.
- El-Banhawy E.M. 1984. Description of some phytoseiid mites from Brazil (Acarina: Phytoseiidae). Acarologia, 25(2): 125-144.
- El-Banhawy E.M., Knapp M. 2011. Mites of the family Phytoseiidae Berlese from Kenya (Acaria: Mesostigmata). Zootaxa, 2945: 1-176. doi:[10.11646/zootaxa.2945.1.1](https://doi.org/10.11646/zootaxa.2945.1.1)
- Evans G.O. 1952. A new typhlodromid mite predaceous on *Tetranychus bimaculatus* Harvey in Indonesia. Ann. Mag. Natur. Hist., 5: 413-416. doi:[10.1080/00222935208654311](https://doi.org/10.1080/00222935208654311)
- Evans G.O. 1953. On some mites of the genus *Typhlodromus* Scheuten, 1857, from S. E. Asia. Ann. Mag. Natur. Hist., 6, 449-467. doi:[10.1080/00222935308654444](https://doi.org/10.1080/00222935308654444)
- Ferragut F., Baumann J. 2019. New phytoseiid mites (Mesostigmata: Phytoseiidae) of Mauritius, with the description of two new species. Syst. Appl. Acarol., 24(5): 825-856. doi:[10.11158/saa.24.5.8](https://doi.org/10.11158/saa.24.5.8)
- Garman P. 1948. Mite species from apple trees in Connecticut. Connect. Agric. Exp. St., Bull., 520, 1-27.
- Ghai S., Menon M.G.R. 1969. Taxonomic studies on Indian mites of the family Phytoseiidae (Acarina). II. Two new genera and species of Phytoseiidae. Orient. Ins., 3, 347-352. doi:[10.1080/00305316.1969.10433928](https://doi.org/10.1080/00305316.1969.10433928)
- Gonzalez R.H., Schuster R.O. 1962. Especies de la familia Phytoseiidae en Chile I. (Acarina: Mesostigmata). Bol. Tecn. Est. Exp. Agron., 16, 1-35.
- Gupta S.K. 1969. Three new species of the genus *Phytoseius* (Acarina: Phytoseiidae) from India. Isr. J. Agric. Res., 19: 115-120.
- Gupta S.K. 1975. Mites of the genus *Amblyseius* (Acarina: Phytoseiidae) from India with descriptions of eight new species. Intern. J. Acarol., 1(2): 26-45. doi:[10.1080/01647957508683746](https://doi.org/10.1080/01647957508683746)
- Gupta S.K. 1981. On a collection of Phytoseiidae (Acaria: Mesostigmata) from Himachal Pradesh (India), with descriptions of two new species. Ind. J. Acarol., 5, 32-36.
- Gupta S.K. 1986. *Fauna of India (Acaria: Mesostigmata) Family Phytoseiidae*. Zoological Survey of India, Calcutta, India, 350 pp.
- Gutierrez J., Etienne J. 1986. Les Tetranychidae de l'île de la Réunion et quelques-uns de leurs prédateurs. Agron. Trop., 41(1): 84-91.
- Hirschmann W. 1962. Gangsystematik der Parasitiformes. *Acarologie Schriftenreihe fur Vergleichende Milbenkunde, Hirschmann-Verlag*, Furth/Bay, 5(5-6), 80 pp.+ 32 plates.
- Ho C.C., Lo K.C. 1989. Contribution to the knowledge of the genus *Paraphytoseius* Swirski and Shechter (Acarina: Phytoseiidae) in Taiwan. J. Agric. Res. China, 38(1): 88-99.
- Hughes A.M. 1948. *The mites associated with stored food products*. Ministry of Agriculture and Fisheries, H. M. Stationary Office, London, 168 pp.
- Huyen L.T., Tung N.D., Lan D.H., Chi C.V., De Clercq P., Dinh N.V. 2017. Life table parameters and development of *Neoseiulus longispinosus* (Acaria: Phytoseiidae) reared on citrus red mite, *Panonychus citri* (Acaria: Tetranychidae) at different temperatures. Syst. Appl. Acarol., 22(9): 1316-1326. doi:[10.11158/saa.22.9.3](https://doi.org/10.11158/saa.22.9.3)
- Jung C., Kim, S., Lee S.W., Lee J.-H. 2003. Phytoseiid mites (Acaria: Phytoseiidae) from Korean apple orchards and their ecological notes. Korean J. Appl. Entomol., 42(3): 185-195.
- Kade N., Gueye-Ndiaye A., Duverney C., Moraes G.J. de. 2011. Phytoseiid mites (Acaria: Phytoseiidae) from Senegal. Acarologia, 51(1): 133-138. doi:[10.1051/acarologia/20112001](https://doi.org/10.1051/acarologia/20112001)
- Karg W. 1982. Diagnostic and systematics of predatory mites of the family Phytoseiidae Berlese in orchards. Zool. Jahr. Syst., 109, 188-210.
- Karg W. 1983. Systematische untersuchung der Gattungen und Untergattungen der Raubmilbenfamilie Phytoseiidae Berlese, 1916, mit der beschreibung von 8 neuen Arten. Mitteil. Zool. Mus. Berlin, 59(2): 293-328. doi:[10.1002/mmzn.4830590203](https://doi.org/10.1002/mmzn.4830590203)
- Karg W. 1991. Die Raubmilbenarten der Phytoseiidae Berlese (Acarina) Mitteleuropas sowie angrenzender Gebiete. Zool. Jahr. Syst., 118(1): 1-64.
- Karg W., Oomen-Kalsbeek F. 1987. Neue Raubmilbenarten der Gattung *Amblyseius* Berlese (Acarina, Parasitiformes, Phytoseiidae) - Antagonisten der unechten Spinnmilbe *Brevipalpus phoenicis* Geijskes. Zool. Jahr. Syst., 114(1): 131-140.
- Kennett C.E. 1958. Some predacious mites of the subfamilies Phytoseiinae and Aceosejinae (Acarina: Phytoseiidae, Aceosejidae) from central California with descriptions of new species. Ann. Entomol. Soc. Amer., 51, 471-479. doi:[10.1093/aesa/51.5.471](https://doi.org/10.1093/aesa/51.5.471)
- Kolodochka L.A. 1998. Two new tribes and the main results of a revision of Palearctic phytoseiid mites (Parasitiformes, Phytoseiidae) with the family system concept. Vest. Zool., 32(1-2): 51-63 [in Russian].
- Kolodochka L.A., Denmark H.A. 1995. A new genus of phytoseiid mites (Acaria, Phytoseiidae). Zh. Ukrainskogo Entomol. Tovar., 1(3-4), 19-26 [in Russian].
- Kreiter S., Amiri K., Douin M., Bohinc T., Trdan S., Tixier M.-S. 2020a. Phytoseiid mites of Slovenia (Acaria: Mesostigmata): new records and first description of the male of *Amblyseius microorientalis*. Acarologia 60(2): 203-242. doi:[10.24349/acarologia/20204364](https://doi.org/10.24349/acarologia/20204364)
- Kreiter S., Fontaine O., Payet R.-M. 2018a. New records of Phytoseiidae (Acaria: Mesostigmata) from Mauritius. Acarologia; 58(4): 773-785. doi:[10.24349/acarologia/20184273](https://doi.org/10.24349/acarologia/20184273)

- Kreiter S., Mailloux J., Tixier M.-S., Le Bellec F., Douin M., Guichou S., Etienne J. 2013. New phytoseiid mites of the French West Indies, with description of a new species, and new records (Acar: Mesostigmata). *Acarologia*, 53(3): 285-303. doi:[10.1051/acarologia/20132095](https://doi.org/10.1051/acarologia/20132095)
- Kreiter S., Payet R.-M., Douin M., Fontaine O., Fillâtre J., Le Bellec F. 2020b. Phytoseiidae of La Réunion Island (Acar: Mesostigmata): three new species and two males described, new synonymies, and new records. *Acarologia*, 60(1): 111-195. doi:[10.24349/acarologia/20204361](https://doi.org/10.24349/acarologia/20204361)
- Kreiter S., Payet R.-M., Fillâtre J., Abdou Azali H. 2018b. First records of Phytoseiidae from one island of the Comoros Archipelago. *Acarologia*, 58(3): 529-545. doi:[10.24349/acarologia/20184256](https://doi.org/10.24349/acarologia/20184256)
- Kreiter S., Tixier M.-S., Etienne J. 2006. New records of phytoseiid mites (Acar: Mesostigmata) from the French Antilles, with description of *Neoseiulus cecileae* sp. nov. *Zootaxa*, 1294: 1-27. doi:[10.11646/zootaxa.1294.1.1](https://doi.org/10.11646/zootaxa.1294.1.1)
- Kreiter S., Zriki Z., Ryckewaert P., Pancarte C., Douin M., Tixier M.-S. 2018c. New phytoseiid mites of Martinique, with redescription of four species and new records. *Acarologia*, 58(2): 366-407. doi:[10.24349/acarologia/20184248](https://doi.org/10.24349/acarologia/20184248)
- Lindquist E.E. 1994. Some observations on the chaetotaxy of the caudal body region of gamasine mites (Acar: Mesostigmata), with a modified notation for some ventrolateral body setae. *Acarologia*, 35: 323-326.
- Lindquist E.E., Evans G.W. 1965. Taxonomic concepts in the Ascidae, with a modified setal nomenclature for the idiosoma of the Gamasina Acarina: Mesostigmata. *Mem. Entomol. Soc. Canada*; 47: 1-64. doi:[10.4039/entm9747fv](https://doi.org/10.4039/entm9747fv)
- Lo P.K.C. 1970. Phytoseiid mites from Taiwan (I) (Acar: Mesostigmata). *Bull. Sun Yatzen Cult. Found.*, 5: 47-62.
- Luong T.H., Nguyen D.T., Dang H.L., Cao V.C., De Clercq P., Nguyen V.D. 2017. Life table parameters and development of *Neoseiulus longispinosus* (Acar: Phytoseiidae) reared on citrus red mite, *Panonychus citri* (Acar: Tetranychidae) at different temperatures. *Syst. Appl. Acarol.*, 22(9): 1316-1326. doi:[10.11158/saa.22.9.3](https://doi.org/10.11158/saa.22.9.3)
- Mailloux J., Le Bellec F., Kreiter S., Tixier M.-S., Dubois P. 2010. Influence of ground cover management on diversity and density of phytoseiid mites (Acar: Phytoseiidae) in Guadeloupean citrus orchards. *Exp. Appl. Acarol.*, 52: 275-290. doi:[10.1007/s10493-010-9367-7](https://doi.org/10.1007/s10493-010-9367-7)
- Mason P.G., Cock M.J.W., Barratt B.I.P., Klapwijk J.N., Van Lenteren J.C., Brodeur J., Hoelmer K.A., Heimpel, G.E. 2018. Best practices for the use and exchange of invertebrate biological control genetic resources relevant for food and agriculture. *BioControl*, 63(1): 149-154. doi:[10.1007/s10526-017-9810-3](https://doi.org/10.1007/s10526-017-9810-3)
- Matthysse J.G., Denmark H.A. 1981. Some phytoseiids of Nigeria (Acarina: Mesostigmata). *Fla Entomol.*, 64: 340-357. doi:[10.2307/3494585](https://doi.org/10.2307/3494585)
- Meyer M.K.P., Rodrigues M.C. 1966. Acari associated with cotton in southern Africa (with reference to other plants). *Garcia de Orta, Rev. Junta Invest. Ultramar*, 13(2): 195-226.
- McMurtry J.A., Croft B.A. 1997. Life-styles of phytoseiid mites and their roles in biological control. *Ann. Rev. Entomol.*, 42: 291-321. doi:[10.1146/annurev.ento.42.1.291](https://doi.org/10.1146/annurev.ento.42.1.291)
- McMurtry J.A., Moraes G.J. de. 1984. Some phytoseiid mites from the South Pacific, with descriptions of new species and a definition of the *Amblyseius largoensis* species group. *Intern. J. Acarol.*, 10: 27-37. doi:[10.1080/01647958408683347](https://doi.org/10.1080/01647958408683347)
- McMurtry J.A., Moraes G.J. de. 1985. Some phytoseiid mites (Acar: Phytoseiidae) of Papua New Guinea, with descriptions of six new species. *Intern. J. Acarol.*, 11: 75-88. doi:[10.1080/01647958508683399](https://doi.org/10.1080/01647958508683399)
- McMurtry J.A., Moraes G.J. de., Sourassou N.F. 2013. Revision of the life styles of phytoseiid mites (Acar: Phytoseiidae) and implications for biological control strategies. *Syst. Appl. Acarol.*, 18: 297-320. doi:[10.11158/saa.18.4.1](https://doi.org/10.11158/saa.18.4.1)
- Mochizuki M. 2009. Development, Reproduction and Prey Consumption of the Thrips Predator *Gynaeseius liturivorus* (Ehara) (Acar: Phytoseiidae). *J. Acarol. Soc. Japan*, 18(2):73-84. doi:[10.2300/acari.18.73](https://doi.org/10.2300/acari.18.73)
- Moraes G.J. de, Castro T.M.G. de, Kreiter S., Quilici S., Gondim Jr. M.G.C., Sá L.A. 2012. Search for natural enemies of *Raoiella indica* Hirst in Réunion Island (Indian Ocean). *Acarologia*, 52(2), 129-134. doi:[10.1051/acarologia/20122043](https://doi.org/10.1051/acarologia/20122043)
- Moraes G.J. de, Kreiter S., Lofego A.C. 2000. Plant mites (Acar: Phytoseiidae) of the French Antilles. 3. Phytoseiidae (Gamasida). *Acarologia*, 40(3): 237-264.
- Moraes G.J. de, Lopes P.C., Fernando C.P. 2004b. Phytoseiid mite (Acar: Phytoseiidae) of coconut growing areas in Sri Lanka, with descriptions of three new species. *J. Acarol. Soc. Japan*, 13(2): 141-160. doi:[10.2300/acari.13.141](https://doi.org/10.2300/acari.13.141)
- Moraes G.J. de, McMurtry J.A., Denmark H.A. 1986. *A catalog of the mite family Phytoseiidae. References to taxonomy, synonymy, distribution and habitat*. EMBRAPA - DDT, Brasilia, Brazil, 353 pp.
- Moraes G.J. de, McMurtry J.A., Denmark H.A., Campos C.B. 2004a. A revised catalog of the mite family Phytoseiidae. *Zootaxa*, 434: 1-494. doi:[10.11646/zootaxa.434.1.1](https://doi.org/10.11646/zootaxa.434.1.1)
- Moraes G.J. de, McMurtry J.A., van den Berg H., Yaninek J.S. 1989a. Phytoseiid mites (Acar: Phytoseiidae) of Kenya, with descriptions of five new species and complementary descriptions of eight species. *Intern. J. Acarol.*, 15(2): 79-93. doi:[10.1080/01647958908683829](https://doi.org/10.1080/01647958908683829)
- Moraes G.J. de, McMurtry J.A., Yaninek, J.S. 1989b. Some phytoseiid mites (Acar: Phytoseiidae) from tropical Africa with description of a new species. *Intern. J. Acarol.*, 15(2): 95-102. doi:[10.1080/01647958908683830](https://doi.org/10.1080/01647958908683830)
- Moraes G.J. de, Zannou I.D., Ueckermann E.A., Oliveira A.R., Yaninek J.S., Hanna R. 2007b. Phytoseiid mites of the tribes Afroseiulini, Kampimodromini and Phytoseiulini, and complementary notes on mites of the tribes Euseiini and Neoseiulini (Acar: Phytoseiidae) from sub-Saharan Africa. *Zootaxa*, 1628: 1-22. doi:[10.11646/zootaxa.1628.1.1](https://doi.org/10.11646/zootaxa.1628.1.1)

- Muma M.H. 1955. Phytoseiidae (Acarina) associated with citrus in Florida. Ann. Entomol. Soc. Amer., 48, 262-272. doi:[10.1093/ae/48.4.262](https://doi.org/10.1093/ae/48.4.262)
- Muma M.H. 1961. Subfamilies, genera, and species of Phytoseiidae (Acarina: Mesostigmata). Fla St. Mus. Bull., 5(7): 267-302
- Muma M.H., Denmark H.A. 1970. Phytoseiidae of Florida. Arthropods of Florida and neighbouring land areas, 6. *Florida Department of Agriculture and Consumer Services, Division of Plant Industry*, Gainesville, USA, 150 pp.
- Myers N. 1988. Threatened biotas: hostspots in tropical forests. *Environmentalist*, 8: 187-208. doi:[10.1007/BF02240252](https://doi.org/10.1007/BF02240252)
- Myers N., Mittermeier R.A., Mittermeier C.G., Da Fonseca G.A., Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature*, 403, 853-858. doi:[10.1038/35002501](https://doi.org/10.1038/35002501)
- Narayanan E.S., Kaur R.B.N., Ghai S. 1960. Importance of some taxonomic characters in the family Phytoseiidae., with new records and descriptions of species. Proceed. Nat. Inst. Sc. India, 26B: 384-394.
- Navasero M.M., Navasero M.V. 2016. Biology of *Paraphytoseius orientalis* (Narayanan *et al.*) reared on the broad mite, *Polyphagotarsonemus latus* (Banks) (Acari: Phytoseiidae, Tarsonemidae) in the Philippines. *Philip. Entomol.*, 30 (1): 21-28.
- Nguyen D.T., Dao T.L. 2019. Effect of temperatures and diets on biological characteristics of predatory mite *Amblyseius largoensis* (Acari: Phytoseiidae). *Sci. Technol. J. Agric. Rural Dev.*, 11: 66-72.
- Nguyen D.T., De Clercq P. 2018. Some morphological and biological characters of predatory mite *Paraphytoseius cracentis* (Acari: Phytoseiidae) reared on two different preys. *Vietnam J. Agric. Sci.*, 16 (2): 95-104 (in Vietnamese).
- Nguyen D.T., Than A.T., Jonckeere W., Nguyen V.H., Van Leeuwen T., De Clercq P. 2019a. Life tables and feeding habits of *Proprioseiopsis lenis* (Acari: Phytoseiidae) and implications for its biological control potential in Southeast Asia. *Syst. Appl. Acarol.*, 24 (5): 857-865. doi:[10.11158/saa.24.5.9](https://doi.org/10.11158/saa.24.5.9)
- Nguyen T.P.T., Tran N.V., Tran T.T.A. 2016. Species occurrence of phytophagous and predatory mites (Acari: Tetranychidae, Phytoseiidae) on fruit vegetables in Ho Chi Minh City, Vietnam. *J. Acarol. Soc. Japan*, 25(S1): 133-136. doi:[10.2300/acari.25.Suppl_133](https://doi.org/10.2300/acari.25.Suppl_133)
- Nguyen V.H., Jonckheere W., Nguyen D.T., Moraes G.J. de, Van Leeuwen T., De Clercq P. 2019b. Phytoseiid mites prey effectively on thrips eggs: evidence from predation trials and molecular analyses. *Biological Control*, 137: 1-7. doi:[10.1016/j.bioc.2019.104012](https://doi.org/10.1016/j.bioc.2019.104012)
- Nusartlert N., Vichitbandha P., Baker G., Chandrapatya A. 2011. Pesticide-induced mortality and prey dependent life history of the predatory mite *Neoseiulus longispinosus* (Acari: Phytoseiidae). *Trends in Acarology*. Netherlands, Springer: 495-498. doi:[10.1007/978-90-481-9837-5_83](https://doi.org/10.1007/978-90-481-9837-5_83)
- Oliveira D.C., Charanasri V., Kongchuensin M., Konvipasruang P., Chandrapatya A., Moraes G.J. de. 2012. Phytoseiidae of Thailand (Acari: Mesostigmata), with a key for their identification. *Zootaxa*, 3453: 1-24. doi:[10.11646/zootaxa.3453.1.1](https://doi.org/10.11646/zootaxa.3453.1.1)
- Prasad, V. (1968b) Some phytoseiid mites from Hawaii. *Annals of the Entomological Society of America*, 61, 1459-1462
- Pritchard A.E., Baker E.W. (1962) Mites of the family Phytoseiidae from Central Africa, with remarks on the genera of the world. *Hilgardia*; 33(7): 205-309. doi:[10.3733/hilg.v33n07p205](https://doi.org/10.3733/hilg.v33n07p205)
- Quilici S., Kreiter S., Ueckermann E. A., Vincenot D. 1997. Predatory mites (Acari) from various crops on Réunion Island. *Intern. J. Acarol.*, 23(4): 283-291. doi:[10.1080/01647959708683578](https://doi.org/10.1080/01647959708683578)
- Quilici S., Ueckermann E. A., Kreiter S., Vayssières J.-F. 2000. Phytoseiidae (Acari) of La Réunion Island. *Acarologia*, 41(1-2): 97-108.
- Reis P.R., Teodoro A.V., Pedro Neto M., Da Silva E.A. 2007. Life history of *Amblyseius herbicolus* (Chant) (Acari: Phytoseiidae) on coffee plants. *Neotrop. Entomol.*, 36(2): 282-287. doi:[10.1590/S1519-566X2007000200016](https://doi.org/10.1590/S1519-566X2007000200016)
- Ribaga C. (1904) Gamasidi planticoli. *Riv. Patol. Veg.*, 10: 175-178.
- Rodriguez-Cruz F.A., Venzon M., Pinto C.M.F. 2013. Performance of *Amblyseius herbicolus* on broad mites and on castor bean and sunnhemp pollen. *Exp. Appl. Acarol.*, 60: 497-507. doi:[10.1007/s10493-013-9665-y](https://doi.org/10.1007/s10493-013-9665-y)
- Rowell H.J., Chant D.A., Hansell R.I.C. 1978. The determination of setal homologies and setal patterns on the dorsal shield in the family Phytoseiidae (Acarina: Mesostigmata). *Can. Entomol.*, 110: 859-876. doi:[10.4039/Ent110859-8](https://doi.org/10.4039/Ent110859-8)
- Ryu M.O., Lee W.K. 1992. Ten newly recorded phytoseiid mites (Acarina: Phytoseiidae) from Korea. *Korean J. Entomol.*, 22(1): 23-42. doi:[10.2476/asjaa.40.23](https://doi.org/10.2476/asjaa.40.23)
- Schicha E. 1975. A new predacious species of *Amblyseius* Berlese from strawberry in Australia, and *A. longispinosus* (Evans) redescribed (Acari: Phytoseiidae). *J. Austral. Entomol. Soc.*, 14: 101-106. doi:[10.1111/j.1440-6055.1975.tb02010.x](https://doi.org/10.1111/j.1440-6055.1975.tb02010.x)
- Schicha E. 1982. A new species of *Amblyseius* Berlese from the New Hebrides with notes on closely related species from Australia, Japan and China (Acari: Phytoseiidae). *Austral. Entomol. Mag.*, 9(2-3): 33-38.
- Schicha E. 1987. *Phytoseiidae of Australia and neighbouring areas*. Indira Publishing House, West Bloomfield, Michigan, USA, 187 pp.
- Schicha E., Corpuz-Raros L.A. 1985. Contribution to the knowledge of the genus *Paraphytoseius* Swirski and Shechter (Acarina: Phytoseiidae). *Intern. J. Acarol.*, 11(2): 67-73. doi:[10.1080/01647958508683398](https://doi.org/10.1080/01647958508683398)
- Schicha E., Corpuz-Raros L.A. 1992. *Phytoseiidae of the Philippines*. Indira Publishing House, West Bloomfield, Michigan, USA, 190 pp.
- Schicha E., Gutierrez J. 1985. Phytoseiidae of Papua New Guinea, with three new species, and new records of Tetranychidae (Acari). *Intern. J. Acarol.*, 11(3): 173-181. doi:[10.1080/01647958508683412](https://doi.org/10.1080/01647958508683412)
- Schuster R.O., Pritchard A.E. 1963. Phytoseiid mites of California. *Hilgardia*, 34: 191-285. doi:[10.3733/hilg.v34n07p191](https://doi.org/10.3733/hilg.v34n07p191)

- Shibao M., Ehara S., Hosomii A., Tanaka H. 2004. Seasonal fluctuation in population density of phytoseiid mites and the yellow tea thrips, *Scirtothrips dorsalis* Hood (Thysanoptera: Thripidae) on grape and predation of the thrips by *Euseius sojaensis* (Ehara) (Acarina: Phytoseiidae). *Appl. Entomol. Zool.*, 39, 727-730. doi:[10.1303/aez.2004.727](https://doi.org/10.1303/aez.2004.727)
- Swirski E., Golan Y. 1967. On some phytoseiid mites (Acarina) from Luzon Island (Phillipines). *Israel J. Agric. Res.*, 17: 225-227.
- Swirski E., Shechter R. 1961. Some phytoseiid mites (Acarina: Phytoseiidae) of Hong-Kong, with a description of a new genus and seven new species. *Israel J. Agric. Res.*, 11: 97-117.
- Tixier M.-S., Kreiter S. 2009. Arthropods in biodiversity hotspots: the case of the Phytoseiidae (Acarina: Mesostigmata). *Biodivers. Conserv.*, 18:507-527. doi:[10.1007/s10531-008-9517-y](https://doi.org/10.1007/s10531-008-9517-y)
- Tseng Y.H. 1976. Systematics of the mite family Phytoseiidae from Taiwan, with a revised key to genera of the world (II). *J. Agric. Assoc. China, N. Ser.*, 94: 85-128.
- Tseng Y.H. 1983. Further study on phytoseiid mites from Taiwan (Acarina: Mesostigmata). *Chin. J. Entomol.*, 3: 33-74.
- Ueckermann E.A., Loots G.C. 1987. Two species of *Amblyseius (Paraphytoseius)* Swirski & Shechter from Africa (Acarina: Phytoseiidae). *Acarologia*, 28(3): 221-226.
- Ueckermann E.A., Zannou I.D., Moraes G.J. de, Oliveira A.R. de, Hanna R., Yaninek J.S. 2007. Phytoseiidae mites of the subfamily Phytoseiinae (Acarina: Phytoseiidae) from sub-Saharan Africa. *Zootaxa*, 1658: 1-20. doi:[10.11646/zootaxa.1658.1.1](https://doi.org/10.11646/zootaxa.1658.1.1)
- van der Merwe G.G. 1968. A taxonomic study of the family Phytoseiidae (Acarina) in South Africa with contributions to the biology of two species. *Entomol. Mem., South Africa Dep. Agric. Techn. Serv.*, 18: 1-198.
- van der Merwe G.G., Ryke P.A.J. 1963. Key characteristics of the predacious family Phytoseiidae with special reference to the genus *Amblyseius* Berlese (Acarina). *J. Entomol. Soc. South Africa*, 26, 82-93.
- Vichitbandha P., Chandrapatya A. 2009. Life history of *Amblyseius cinctus* Corpuz and Rimando (Acarina: Phytoseiidae) on broad mites larvae, *Polyphagotarsonemus latus* (Banks) (Acarina: Tarsonemidae) and its predation response on broad mites at different predators: prey ratios in laboratory and greenhouse conditions. *J. Intern. Soc. Southeast Asian Agric. Sci.*
- Vitzthum H. von. 1941. Acarina. In: Bronns, H.G. (Ed.), *Klassen und Ordnungen des Tierreichs 5*, Akademischer Verlag, Leipzig, Germany: 764-767.
- Wainstein B.A. 1962. Révision du genre *Typhlodromus* Scheuten, 1857 et systématique de la famille des Phytoseiidae (Berlese 1916) (Acarina: Parasitiformes). *Acarologia*, 4: 5-30.
- Womersley H. 1954. Species of the subfamily Phytoseiinae (Acarina: Laelaptidae) from Australia. *Austral. J. Zool.*, 2: 169-191. doi:[10.1071/Z09540169](https://doi.org/10.1071/Z09540169)
- Wu W.N. 1997. A review of taxonomic studies of the genus *Phytoseius* (Acarina: Phytoseiidae) from China. *Syst. Appl. Acarol.*, 2: 149-160. doi:[10.11158/saa.2.1.21](https://doi.org/10.11158/saa.2.1.21)
- Wu W.N., Li Z.Q. 1984. Three new species of the genus *Phytoseius* from south China (Acarina: Phytoseiidae). *Acta Entomol. Sinica*, 27(4): 457-461 [in Chinese].
- Wu W.N., Ou J.F. 1999. A new species group of the *Amblyseius*, with descriptions of two new species (Acarina: Phytoseiidae) from China. *Syst. Appl. Acarol.*, 4: 103-110. doi:[10.11158/saa.4.1.15](https://doi.org/10.11158/saa.4.1.15)
- Xin J.L., Liang L.R., Ke L.S. 1982. Two new species of *Phytoseius* Ribaga from Yunnan and Guizhou (Acarina: Phytoseiidae). *Zool. Res., Suppl.*, 3, 57-60 [in Chinese].
- Zack R.E. 1969. Seven new species and records of phytoseiid mites from Missouri (Acarina: Phytoseiidae). *J. Kansas Entomol. Soc.*, 42(1), 68-80.
- Zannou I.D., Moraes G.J. de, Ueckermann E.A., Oliveira A.R., Yaninek J.S., Hanna R. 2007. Phytoseiid mites of the subtribe Amblyseiina (Acarina: Phytoseiidae: Amblyseiini) from sub-Saharan Africa. *Zootaxa*, 1550: 1-47. doi:[10.11646/zootaxa.1550.1.1](https://doi.org/10.11646/zootaxa.1550.1.1)