



CODEN (USA): IAJ PBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

Available online at: <http://www.iajps.com>

Research Article

FREQUENCY, CLINICAL PRESENTATION AND MANAGEMENT OF DIABETIC FOOT: (THE MULTI-INSTITUTIONAL STUDY)

Dr. Madiha Shah¹, Dr. Aqeel Ahmed Channa², Dr. Nisar Ahmed Shah^{1*},
Dr. Hamid Nawaz Ali Memon³, Dr Imran Karim¹ and Dr. Zulfiqar Ali Qutrio Baloch⁴

¹Department of Medicine, Liaquat University Hospital Hyderabad / Jamshoro.

²Government Lyari General Hospital Karachi.

³Zulekha Hospital, Dubai United Arab Emirates.

⁴Brandon Regional Hospital, Brandon, Florida.

Received: 27 February 2016

Accepted: 11 March 2017

Published: 14 March 2017

Abstract:

OBJECTIVE: To determine the frequency, clinical presentation and management of diabetic foot at tertiary care hospitals

PATIENTS AND METHODS: This multi-institutional one year study descriptive case series was conducted on diabetic population had foot complications at Liaquat University Hospital & Govt Lyari general hospital Karachi. The inclusion criteria of the study were all the patients, ≥ 12 years of age, either gender with diabetes mellitus (type 1 or type 2) presented with foot ulcers, infection and gangrene of foot. The conservative management includes strict glycemic control, meticulous dressing and if needed approach for major surgical interventions. The data was recorded on pre-designed proforma while analyzed in SPSS 16. The frequency, percentages and mean \pm SD calculated for study variables.

RESULTS: During one year study period total seventy two patients of diabetic foot were enrolled and evaluated. The mean age \pm SD for whole population was 52.85 ± 8.93 while the mean \pm SD for hemoglobin A1C, fasting and random blood sugar was 9.52 ± 2.84 , 169 ± 7.74 and 276.92 ± 8.96 respectively. Thirty subjects had a history of trauma before the onset of foot lesion and majority (75%) were belonged to rural populated areas and had history of diabetes for more than five years (77.7%). The common pathogen identified was staphylococcus aureus 38 (52.8%) and common procedures were slough excision and regular dressing 26 (36.1%) and split skin graft 14 (19.4%).

CONCLUSION: Commonest presenting lesion was ulcer, cellulitis and Gangrene. Conservative treatment consists of control of diabetes along with antibiotics while the surgical interventions are other mode of treatment.

Keywords: Diabetes mellitus, Foot ulcer and Diabetic foot

Corresponding Author:

Dr. Nisar Ahmed Shah*,

M.D, Dip:Card,

Department of Medicine,

Liaquat University Hospital Hyderabad.

Email: zulfiqar229@hotmail.com

QR code



Please cite this article in press as Nisar Ahmed Shah et al, *Frequency, Clinical Presentation and Management of Diabetic Foot*, Indo Am. J. P. Sci, 2017; 4(03).

INTRODUCTION:

The diabetic foot ulcer is a worldwide problem and major complications of diabetes mellitus [1]. Majority of the victims required long term management including amputations [2,3]. Diabetes associated foot disorders remains most frequent cause of diabetes specific hospitalizations [4]. Each year 5% of diabetic patients acquire foot ulcer while around 15% of diabetics develop some foot issues during their illness [5]. Former studies had shown that Type 1 diabetic foot complications are also common causes for amputations [6,7]. In the Jain AKC, et al [6] all major amputations performed had history of type 1 diabetes and foot complications with prevalence ranges from 8.89% to 13.39% [6,7]. The etio-pathogenesis of diabetic foot lesion is variable; diabetic vasculopathy, neuropathies, uncontrolled diabetes and infections are few of them [8-11]. The reasons for involvement of foot in diabetic population because of the expose part of body for injury and infections usually neglected by patient while neuropathy and vasculopathy site is also the foot [12,13]. The complications are more likely among poor population due to illiteracy, poverty and negligence [14]. The reason for conducting the present study and evaluating the diabetes in detail because it affects younger age to elderly subjects leads to significant burden in their life style; it is commonest systemic diseases in the society so the complication should be known at community level and also to prevent the complications, the longstanding uncontrolled diabetes leads to vascular and neurological changes which play key role in foot ulcerations, infectious, deformities, systemic complications and amputations. Thus the early and

appropriate effective measures can save the diabetic individuals to acquire various complications.

PATIENTS AND METHODS:

This one year descriptive case series was conducted on diabetic population had foot complications at Liaquat University Hospital Hyd / Jamshoro & Govt: Lyari general hospital Karachi. The inclusion criteria of the study were all the patients, ≥ 12 years of age, either gender had diabetes mellitus (type 1 or type 2) for ≥ 3 years presented with foot ulcers, infection and gangrene of foot while the exclusion criteria were patients with foot infections without diabetes, subjects with vasculitis, immune-compromised and non cooperative individuals. The informed consent and detail history was taken, specific clinical examination was performed, and routine and relevant investigations were advised. The conservative management includes strict glycemic control, meticulous dressing (care of foot) and if needed approached for surgical interventions. The data was recorded on pre-designed proforma while analyzed in SPSS 16. The frequency, percentages and mean \pm SD calculated for study variables.

RESULTS:

Total seventy two patients of diabetic foot were enrolled and evaluated during study period. The mean age \pm SD for whole population was 52.85 ± 8.93 while the mean \pm SD for hemoglobin A1C, fasting and random blood sugar was 9.52 ± 2.84 , 169 ± 7.74 and 276.92 ± 8.96 respectively. Thirty subjects had a history of trauma before the onset of foot lesion and majority (75%) were belonged to rural populated areas. The results of the study are shown in Table 1-5.

Table 1: The Age and Gender Distribution

	AGE (yrs)	GENDER		Total
		Male	Female	
	12-19	2	1	3
		3.9%	4.8%	4.2%
	20-29	4	2	6
		7.8%	9.5%	8.3%
	30-39	12	0	12
		23.5%	.0%	16.7%
	40-49	8	7	15
		15.7%	33.3%	20.8%
	50-59	13	9	22
		25.5%	42.9%	30.6%
	60+	12	2	14
		23.5%	9.5%	19.4%
	Total	51	21	72
		100.0%	100.0%	100.0%

Table 2: The Gender and Clinical Presentation

		GENDER		Total
PRESENTATION		Male	Female	
	Ulcer	24	10	34
		47.1%	47.6%	47.2%
	Gangrene	7	6	13
		13.7%	28.6%	18.1%
	Abscess	8	2	10
		15.7%	9.5%	13.9%
	Cellulitis	12	3	15
		23.5%	14.3%	20.8%
Total		51	21	72
		100.0%	100.0%	100.0%

Table 3: The Gender and Duration of Diabetes Mellitus

		DURATION (yrs)		Total
GENDER		≤5 years	> 5	
	Male	12	39	51
		75.0%	69.6%	70.8%
	Female	4	17	21
		25.0%	30.4%	29.2%
Total		16	56	72
		100.0%	100.0%	100.0%

Table 4: The Gender In Relation To Culture and Sensitivity

		GENDER		Total
CULTURE & SENSITIVITY		Male	Female	
	Staphylococcus aureus	23	15	38
		45.1%	71.4%	52.8%
	Pseudomonas aeruginosa	9	1	10
		17.6%	4.8%	13.9%
	Klebsiella	13	2	15
		25.5%	9.5%	20.8%
	Coliform	4	2	6
		7.8%	9.5%	8.3%
	Proteus	2	1	3
		3.9%	4.8%	4.2%
Total		51	21	72
		100.0%	100.0%	100.0%

Table 5: The Gender and Surgical Interventions

	TREATMENT	Gender		Total
		Male	Female	
	Slough excision and regular dressing	16	10	26
		31.4%	47.6%	36.1%
	Split skin graft	11	3	14
		21.6%	14.3%	19.4%
	Fasciotomy	10	1	11
		19.6%	4.8%	15.3%
	Disarticulation	6	2	8
		11.8%	9.5%	11.1%
	Transmetatarsal amputation	4	0	4
		7.8%	.0%	5.6%
	Below knee amputation	2	2	4
		3.9%	9.5%	5.6%
	Above knee amputation	2	3	5
		3.9%	14.3%	6.9%
	Total	51	21	72
		100.0%	100.0%	100.0%

DISCUSSION:

The present study had 72 diabetic individuals (both type 1 and 2) and when compared to Jain AKC [15] and Shahbazian H, et al [16] no age group difference was reported. In current study 51 (70.8%) were males and 21 (29.1%) were female cases. The incidence is more in males might be because of house holders of the family and outdoor working routine that creates risk for trauma and sequelae. The observation is consistent with the study by Dinh T, et al [17]. The thirty individuals in this series had a history of trauma before the onset of foot lesion, the findings was also detected by study published in 2004 [18]. The commonest pathophysiological alterations responsible for diabetic foot are neuropathy, ischaemia and infection. Majority of individuals (59 patients) had > 5 years duration of diabetes. Few cases were diagnosed as diabetics following admission and investigations. The neuropathy changes were observed in 25 subjects while the Ischemic complication was detected in 12 individuals whereas the infective complication of foot observed in all patients. The observation was also identified formerly by Lipsky BA, et al [19]. The commonest organism cultured was staphylococcus aureus 38 (58.2%) cases, followed by klebsiella 15(20.8%) and were sensitive to antibiotics as gentamycin, ampicillin, cloxacillin, amikacin, ciproflaxacin and cephalosporin, etc. The role of antibiotics in diabetic foot infections were also discussed by Edmonds M, et al and Abbas M, et al [20,21]. In current study, 26 (36.1%) cases were treated by slough excision, 14(19.4%) with skin graft, 8(11.1%) by

disarticulation of single or multiple toes, fasciotomy 11 (15.4%), below knee amputation was done in 4 (5.6%) and above knee amputation were done in 5(6.9%) subjects. The surgical intervention s were also reported by Setacci C, et al and Armstrong DG, et al respectively [22,23]. Proper control of diabetes and foot care is very important in diabetic foot management. Fasting and random blood sugar estimations should be under control with annual ophthalmologist and nephrologist visit [24]. Infection should be treated with broad spectrum antibiotics and pentoxifylline was should also considered in ischemic lesions [25,26]. In present series the amputation rate is lower compared to former literature [27,28]. Might be due to, better glycaemic control, patient education regarding diet and foot care, antibiotics usage, extensive debridement along with regular dressing. In current series after amputation, wound had good healing process and the subjects were referred to rehabilitation centers for prosthesis.

CONCLUSION:

This study comprised of 72 subjects with diabetic foot patients with emphasis on surgical management with predominance of male population. Commonest presenting lesion was ulcer, cellulitis and Gangrene. Conservative treatment consisting of control of diabetes along with antibiotics, wound debridement slough excision followed by dressing resulting in healing process while split skin grafts, transmetatarsal amputation, disarticulation,

amputation below and above knee were the other modes of treatment.

REFERENCES:

1. Adeghate E, Schattner P, Dunn E. An update on the etiology and epidemiology of diabetes mellitus. *Ann N Y Acad Sci.* 2006 Nov;1084:1-29.
2. Gadsby R. Epidemiology of diabetes. *Adv Drug Deliv Rev.* 2002 Nov 5;54(9):1165-72.
3. Tan MH, MacLean DR. Epidemiology of diabetes mellitus in Canada. *Clin Invest Med.* 1995 Aug;18(4):240-6.
4. Riaz M, Miyan Z, Zaidi SI, Alvi FD, Fawwad A, Ahmadani MY, et al. Characteristics and Outcomes of Subjects With Diabetic Foot Ulceration. *Diabetes Care.* 2012 Sep; 35(9): e63.
5. Shah SF, Hameed S, Khawaja Z, Abdullah T, Waqar SH, Zahid MA. Evaluation and management of diabetic foot: A multicentre study conducted at Rawalpindi, Islamabad. *Ann Pak inst Med Sci.* 2011;7(4):233-237.
6. Jain AKC. A new classification of diabetic foot complications: a simple and effective teaching tool. *J Diab Foot Comp.* 2012;4(1):1-5.
7. Jain AKC, Viswanath S. Distribution and analysis of diabetic foot. *OA Case Reports.* 2013;2(21):117.
8. Jeffcoate WJ, Harding KG. Diabetic foot ulcers. *Lancet.* 2003 May 3;361(9368):1545-51.
9. Ricco JB, Thanh Phong L, Schneider F, Illuminati G, Belmonte R, Valagier A, et al. The diabetic foot: a review. *J Cardiovasc Surg (Torino).* 2013 Dec;54(6):755-62.
10. Edmonds M. Diabetic foot ulcers: practical treatment recommendations. *Drugs.* 2006;66(7):913-29.
11. Vuorisalo S, Venermo M, Lepäntalo M. Treatment of diabetic foot ulcers. *J Cardiovasc Surg (Torino).* 2009 Jun;50(3):275-91.
12. Hobizal KB, Wukich DK. Diabetic foot infections: current concept review. *Diabet Foot Ankle.* 2012; 3: 10.
13. Bowering CK. Diabetic foot ulcers. Pathophysiology, assessment, and therapy. *Can Fam Physician.* 2001 May; 47: 1007–1016.
14. Janisse D, Janisse E. Pedorthic management of the diabetic foot. *Prosthet Orthot Int.* 2015 Feb;39(1):40-7
15. Jain AKC. Type 1 diabetic foot complications. *The Journal of Diabetic Foot Complications.* 2016;8(1):17-22
16. Shahbazian H, Yazdanpanah L, Latifi SM. Risk assessment of patients with diabetes for foot ulcers according to risk classification consensus of International Working Group on Diabetic Foot (IWGDF). *Pak J Med Sci.* 2013 May-Jun; 29(3): 730–734.
17. Dinh T, Veves A. The influence of gender as a risk factor in diabetic foot ulceration. *Wounds.* 2008 May;20(5):127-31.
18. Boddenberg U. Healing time of foot and ankle fractures in patients with diabetes mellitus: literature review and report on own cases. *Zentralbl Chir.* 2004 Dec;129(6):453-9.
19. Lipsky BA, Berendt AR, Deery HG, Embil JM, Joseph WS, Karchmer AW, et al. Diagnosis and treatment of diabetic foot infections. *Plast Reconstr Surg.* 2006 Jun;117(7 Suppl):212S-238S.
20. Edmonds M, Foster A. The use of antibiotics in the diabetic foot. *Am J Surg.* 2004 May;187(5A):25S-28S.
21. Abbas M, Uçkay I, Lipsky BA. In diabetic foot infections antibiotics are to treat infection, not to heal wounds. *Expert Opin Pharmacother.* 2015 Apr;16(6):821-32
22. Setacci C, Sirignano S, Mazzitelli G, Setacci F, Messina G, Galzerano G. Diabetic Foot: Surgical Approach in Emergency. *Int J Vasc Med.* 2013; 2013: 296169.
23. Armstrong DG, Lipsky BA. Diabetic foot infections: stepwise medical and surgical management. *Int Wound J.* 2004 Jun;1(2):123-32.
24. Torreguitart MV. Diabetic foot care. importance of education. *Rev Enferm.* 2011 May;34(5):25-30.
25. Rewale V, Prabhakar KR, Chitale AM. Pentoxifylline: a new armamentarium in diabetic foot ulcers. *J Clin Diagn Res.* 2014 Jan;8(1):84-6.
26. Adler PF. Assessing the effects of pentoxifylline (Trental) on diabetic neurotrophic foot ulcers. *J Foot Surg.* 1991 May-Jun;30(3):300-3.
27. Weledji EP, Fokam P. Treatment of the diabetic foot - to amputate or not?. *BMC Surg.* 2014; 14: 83.
28. Wong KL, Nather A, Liang S, Chang Z, Wong TT, Lim CT. Clinical outcomes of below knee amputations in diabetic foot patients. *Ann Acad Med Singapore.* 2013 Aug;42(8):388-94.