



REVIEW ARTICLE

## Dorcadionini of Turkey (Coleoptera: Cerambycidae)

Hüseyin Özdkmen

Science Faculty, Department of Biology, Gazi University, Ankara, Turkey

### ABSTRACT

The Dorcadionini fauna of Turkey is reviewed, and a complete list with type information and provincial distributions is presented. Two hundred and seventy-eight species-group taxa are catalogued in total. This represents approximately 35% of the known Palaearctic Dorcadionini fauna. The Turkish Dorcadionini fauna is characterised by 227 endemic species-group taxa. So its endemism ratio is very high (approximately 82%). Distributions of the taxa in Turkey are given based on available literature records. Taxa in the list are arranged systematically under tribe, genus, subgenus (when recognised), species and subspecies (when recognised). The species group taxa whose occurrences in Turkey need conformation are not included in the list but are given separately. Additionally, two taxa are removed from the list: *Dorcadion (Cribridorcadion) investitum* Breuning in Braun, 1978a as nomen nudum and *Dorcadion (Cribridorcadion) rosinae* K. Daniel in Bodemeyer, 1900 as nomen nudum; and two taxa are given as status nov. and comb. nov.: *Dorcadion (Cribridorcadion) atritarse* Pic, 1931a and *Dorcadion (Cribridorcadion) equestris bernhauerorum* Peks, 2010. Moreover, a checklist of Dorcadionini of Turkey is presented in an appendix.

### ARTICLE HISTORY

Received 18 September 2015  
Accepted 20 May 2016

### KEYWORDS

Coleoptera; Cerambycidae;  
Dorcadionini; list; Turkey

## Introduction

Turkey is adjacent to large bodies of water to the south, west and north; it has continental properties. Turkey is the centre of origin of many taxa, and its exceptionally diverse topography has provided refugia in which many species have survived in spite of harsh geological and climatic changes. The great biological importance of Turkey is evident from the remarkable variety of arthropods in Turkey. Nevertheless the fauna of Cerambycidae of Turkey have not been thoroughly studied and documented (Özdkmen and Koçak 2015).

Also, Turkey is one of the most diverse areas of the west Palaearctic region with its unique geographic position as a land bridge between Asia, Europe and North Africa (also linking to the Ethiopian Region via the Arabian Peninsula) and accumulating natural complexes typical of several climatic zones. Three of four hotspots of Europe and Central Asia recognised by Conservation International are partly included within the boundary of Turkey: Caucasus, Irano-Anatolian and Mediterranean Basin hotspots.

Hence, Turkey contains a rich flora and fauna with a considerable number of endemic species (Myers et al. 2000; Çiplak 2003, 2004; Konstantinov et al. 2009; Ekiz et al. 2013; Conservation International c. 2015).

The aim of this work is to determine the current status of the tribe Dorcadionini in Turkey. This work on Dorcadionini of Turkey is meant to realise a study on approximately one-third of the Turkish Cerambycidae fauna. By the same token, the Turkish Dorcadionini represent more than one-third of the Dorcadionini in the whole world.

Hence, Dorcadionini of Turkey is a very significant group. This importance originates from its high endemism ratio (approximately 82%), and also the large number of species-group taxa (278). However, the information on all Turkish Dorcadionini is not sufficient.

The first large work on Dorcadionini species was published by Thomson (1867). Another large work on Dorcadionini species was carried out by Ganglbauer (1884). He evaluated a total of 152 species (30 species for Turkey) of the genus *Dorcadion* and 13 species of the genus *Neodorcadion* (four species for Turkey). Then, the number of Turkish *Dorcadion* species exceeded 50 with species described by various authors. In 1962, the largest work on Dorcadionini was accomplished by Breuning. He mentioned 353 *Dorcadion* species (92 species for Turkey) and nine *Neodorcadion* species (five species for Turkey) in his study. After Breuning (1962a), many new *Dorcadion* species had been described by various authors from Turkey until 1990. In 1990 and 1991, a study on Turkish *Dorcadion* fauna was carried out by Önalp. She stated a total of 52 species for Turkey. However, her work was far from determining the real status of Turkish Dorcadionini since it lacked a great deal of information for Turkey. After Önalp's work, many new *Dorcadion* species were also described by various authors from Turkey. Recently, Löbl and Smetana (2010) recorded a total of 155 species for Dorcadionini of Turkey while Özdkmen (2010) reported a total of 192 species for Turkey.

According to Breuning (1962a), the tribe Dorcadionini included only the genera *Dorcadion* Dalman, 1817, *Neodorcadion* Ganglbauer, 1884 and *Eodorcadion* Breuning, 1947. *Iberodorcadion* described by Breuning (1943) as a subgenus of the genus *Dorcadion*, was ranked by Vives (1976) as a separate genus. Also, Danilevsky et al. (2005) accepted it as a genus on the basis of endophallic structures. So, recently, Löbl and Smetana (2010) and Danilevsky (2015a) listed it as a separate genus. *Politodorcadion* was described by Danilevsky (1996) as a subgenus of *Dorcadion*. Recently, Löbl and Smetana (2010) and Danilevsky (2015a) listed it as a separate genus. The genus *Trichodorcadion* Breuning, 1942 from India and Nepal was transferred in Dorcadionini by Danilevsky and Kasatkin (2007) on the base of endophallic structures. However, Löbl and Smetana (2010) and Danilevsky (2015a) gave *Trichodorcadion* in the tribe Morimopsini. As a result of this, Dorcadionini included five genera as *Dorcadion* Dalman, 1817, *Neodorcadion* Ganglbauer, 1884, *Iberodorcadion* Breuning, 1943, *Eodorcadion* Breuning, 1947 and *Politodorcadion* Danilevsky (1996). Recently, Özdkmen and Kaya (2015g) proposed new arrangements for some Dorcadionini. After this work, the tribe Dorcadionini includes six genera and 17 subgenera (including nominate subgenera) worldwide now.

The tribe Dorcadionini has a Palaearctic (North Africa and Western Europe to China) chorotype. *Iberodorcadion* Breuning, 1943 has a European (chiefly Iberian) chorotype (Spain to Poland). *Eodorcadion* Breuning, 1947 has an East Palaearctic chorotype (Russia, Siberia, Mongolia and China). *Politodorcadion* Danilevsky (1996) has an Asian chorotype

(Kazakhstan, West Siberia and China). These genera are not represented in Turkey. After Özdkmen and Kaya (2015g), the tribe Dorcadionini of Turkey is represented by the genera *Dorcadion* Dalman, 1817, *Megalodorcadi* Pesarini and Sabbadini, 1998 and *Neodorcadion* Ganglbauer, 1884.

## Material and methods

This list is the consequence of careful reviews of the available literature on Turkish Dorcadionini. The genera, subgenera, species and subspecies are listed alphabetically. Information in the list is as follows: the name of the species or subspecies and author(s), type information, synonyms, provincial distributions in Turkey, the literature(s) in which the species or subspecies is cited (in parentheses), and distribution in the world. The type information for each species or subspecies is based on Tavakilian (2015). The provincial distributions of taxa in Turkey are coded as in Table 1. A map showing the administrative provinces of Turkey is provided (Figure 1). It was not possible to give the exact provinces of some species or subspecies, so they are recorded as Asian Turkey (TR-A) or European Turkey (TR-E). For global distributional data, Löbl and Smetana (2010) and Danilevsky (2015a) are chiefly used in the text. Distributional abbreviations for the work are available in Löbl and Smetana (2010). The endemic species are marked with an asterisk. A checklist of Dorcadionini of Turkey is given in Appendix.

**Table 1.** Abbreviations of localities (administrative divisions) of Turkey used in the text and in Figure 1.

Adana (AD)	Erzincan (ER)	Muğla (MG)
Adiyaman (ADY)	Erzurum (EZ)	Muş (MU)
Afyon (AF)	Eskişehir (ES)	Nevşehir (NE)
Agri (AG)	Gaziantep (GA)	Niğde (NI)
Aksaray (AK)	Giresun (GI)	Ordu (OR)
Amasya (AM)	Gümüşhane (GU)	Osmaniye (OS)
Ankara (AN)	Hakkâri (HA)	Rize (RI)
Antalya (ANT)	Hatay (HT)	Sakarya (SA)
Ardahan (AR)	Iğdır (IG)	Samsun (SM)
Artvin (ART)	Isparta (IP)	Siirt (SI)
Aydin (AY)	İçel (IC)	Sinop (SN)
Balıkesir (BL)	İstanbul (IS)	Sivas (SV)
Bartin (BR)	İzmir (IZ)	Şanlıurfa (SU)
Batman (BA)	Kahramanmaraş (KA)	Şırnak (SK)
Bayburt (BY)	Karabük (KR)	Tekirdağ (TE)
Bilecik (BL)	Karaman (KM)	Tokat (TO)
Bingöl (BN)	Kars (KAR)	Trabzon (TB)
Bitlis (BT)	Kastamonu (KS)	Tunceli (TU)
Bolu (BO)	Kayseri (KY)	Uşak (US)
Burdur (BU)	Kırıkkale (KI)	Van (VA)
Bursa (BS)	Kırklareli (KK)	Yalova (YA)
Çanakkale (CA)	Kırşehir (KIR)	Yozgat (YO)
Çankiri (CN)	Kılıç (KL)	Zonguldak (ZO)
Çorum (CO)	Malatya (MA)	European
Denizli (DE)	Manisa (MN)	Turkey = Thracia (TR-E)
Diyarbakır (DL)	Mardin (MR)	Asian
Düzce (DU)	Kocaeli (KO)	Turkey = Anatolia (TR-A)
Edirne (ED)	Konya (KN)	
Elaaziğ (EL)	Kütahya (KU)	Turkey (TR)



**Figure 1.** Map showing the administrative provinces of Turkey.

## Taxonomy

All species-group taxa of the Dorcadionini of Turkey are presented as follows:

## **Tribe DORCADIONINI** Swainson and Shuckard, 1840

## **Genus *DORCADION* Dalman, 1817: 397**

[type species *Cerambyx glicyrrhizae* Pallas, 1773]

**Subgenus *D.* (*CARINATODORCADION*) Breuning, 1943: 524**

[type species *Cerambyx carinatus* Pallas, 1771]

**D. aethiops** Scopoli, 1763: 53 (Cerambyx)

**D. aethiops aethiops** Scopoli, 1763: 53 (Cerambyx)

Type information: Holotype, ex collection G.A. Scopoli [type locality 'Carniola: Circa Labacum (Ljubljana)' (Slovenia)]

*morio* Fabricius, 1787: 141 (*Lamia*) [Austria]

*scopolii* Gmelin, 1790: 1835 (*Cerambyx*) [Austria]

*bergmanni* Tippmann, 1958: 151 (*D. aethiops* ssp.) [Austria]

Provincial distribution in Turkey: TR-A: ES; TR-E: ED ([Önalp 1990](#); [Pesarini and Sabbadini 2007](#); [Özdikmen 2010](#); [Danilevsky 2015a](#))

Range: E (Europe): AL AU BU CR CZ GR HU MC MD RO SK SL TR UK YU A (Asia): TR

Other known subspecies: *D. aethiops asenovi* Danilevsky, 2014: 253 E: BU; *D. aethiops majoripenne* Pic, 1926: 13 E: GR; *D. aethiops propinquum* Breuning, 1962a: 517 E: GR; *D. aethiops strumense* Danilevsky, 2014: 252 E: BU

**D. fulvum** Scopoli, 1763: 53 (Cerambyx)

**D. fulvum erythropterum** Fischer von Waldheim, 1823a: table L-7 (*D. erythropteron*)

Type information: Holotype ♂, ex collection Gotthelf Fischer von Waldheim, Zoological Museum, University of Moscow [type locality 'Rossia meridionalis' (Russia)]

*canaliculatum* Fischer von Waldheim, 1823b: 240 [S Russia]

*frontale* Mulsant and Rey, 1863: 154 [Turkey: Istanbul]

Provincial distribution in Turkey: TR-E: IS (Mulsant and Rey 1863; Özdīkmen 2010; Danilevsky 2015a)

Range: E: AL BH BU CR MC MD RO PL TR UK YU

Other known subspecies: *D. fulvum fulvum* Scopoli, 1763: 53 (*Cerambyx*) E: AU CZ HU SK SL UK

***D. hybridum*** Ganglbauer, 1884: 441

***D. hybridum hybridum*** Ganglbauer, 1884: 441

Type information: Holotype ♂, Magyar Természettudományi Múzeum, Budapest [type locality 'Rumelia' (Greece)]

Provincial distribution in Turkey: TR-E (Breuning 1962a; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdi̇kmen 2010; Danilevsky 2015a)

Range: E: GR TR

Other known subspecies: *D. hybridum hedvigae* Jureček, 1933: 127 E: GR

\****D. ingeae*** Peks, 1993: 3

Type information: Holotype ♂, collection Heinz Peks, Schwanfeld [type locality 'Keşan' (Turkey: Edirne)]

Provincial distribution in Turkey: TR-E: ED (Peks 1993; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: E: TR

**Subgenus *D. (Cribridorcadion)*** Pic, 1901: 12

[type species *Dorcadiion mniszechi* Kraatz, 1873]

*Pedestredorcadion* Breuning, 1943: 526 [type species *Lamia pedestris* Poda von Neuhaus, 1761]

*Dzhungarodorcadion* Danilevsky, 1993: 47 [type species *Dorcadiion jacobsoni* Jakovlev, 1899]

*Autodorcadion* Plavilstshikov, 1958: 45 [type species *Cerambyx arenarius* Scopoli, 1763]

*Bergerianum* Pesarini and Sabbadini, 2004: 150 [type species *Dorcadiion chrysochroum* Breuning, 1943]

\****D. abstersum*** Holzschuh, 1982: 75

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Sat Mt.' (Turkey: Hakkari)]

Provincial distribution in Turkey: TR-A: HA (Holzschuh 1982; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\****D. accola*** Heyden, 1894: 87

Type information: Holotype, ex collection L. von Heyden, Natur-Museum und Forschungs-Institut Senckenberg, Frankfurt am Main [type locality 'Mardin' (Turkey)] *glabrolineatum* Pic, 1927: 6 [?Macedonia]

Provincial distribution in Turkey: TR-A: MR SU (Heyden 1894; Breuning 1962a; Braun 1978a; Önalp 1991; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\****D. afflictum*** Pesarini and Sabbadini, 1998: 54

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Çorum' (Turkey)]

Provincial distribution in Turkey: TR-A: CO (Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. akpinarensis* Bernhauer and Peks, 2010: 196**

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Anamur: Akpınar' (Turkey: İçel)]

Provincial distribution in Turkey: TR-A: IC (Bernhauer and Peks 2010; Özdkmen 2012; Danilevsky 2015a)

Range: A: TR

**\**D. albolineatum* Küster, 1847: 86**

Type information: Holotype, ex collection Jacob Sturm, Zoologische Staatssammlung des Bayerischen Staates, München [type locality 'İstanbul' (Turkey)]

*byzantinum* J. Thomson, 1867: 69 [Turkey]

*abbreviatolineatum* Breuning, 1946: 132 (*D. albolineatum* m.) [Turkey]

Provincial distribution in Turkey: TR-A: BI DI IS KO (Küster 1847; Kraatz 1873; Breuning 1962a; Tuatay et al. 1972; Özdkmen et al. 2005; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. albonotatum* Pic, 1895a: 39**

Type information: Holotype, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Bitlis' (Turkey)]

Provincial distribution in Turkey: TR-A: BT (Pic 1895a; Plavilstshikov 1958; Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. altinyaylaense* Bernhauer and Peks, 2014a: 106**

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'W Altınyayla' (Turkey: Sivas)]

Provincial distribution in Turkey: TR-A: SV (Bernhauer and Peks 2014a; Danilevsky 2015a)

Range: A: TR

**\**D. amanense* Breuning, 1943: 94**

Type information: Holotype ♀, ex collection Josef Breit > G. Frey, Naturhistorisches Museum Basel [type locality 'Amanos Mts.' (Turkey: Hatay)]

Provincial distribution in Turkey: TR-A: HT (Breuning 1943, 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. anatolicum* Pic, 1900a: 12**

Type information: Syntypes ♂ and ♀, ex collection H. Tournier > M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Konya' (Turkey)]

*ininterruptum* Pic, 1900a: 13 [HN] (*D. anatolicum* var.) [Turkey: Konya]

*mediointerruptum* Pic, 1901: 73 (*D. semilineatum* var.) [Turkey: Konya]

*brevipenne* Pic, 1902b: 10 [Turkey: Konya]

*chehirense* Pic, 1902b: 10 (*D. brevipenne* var.) [Turkey: Konya]  
*subuniforme* Pic, 1902c: 27 [Turkey]  
*humeraliinterruptum* Breuning, 1946: 108 (*D. anatolicum* m.) [Turkey: Konya]  
*postapertum* Breuning, 1946: 108 (*D. anatolicum* m.) [Turkey: Konya]  
*pseudomerkli* Breuning, 1946: 108 (*D. anatolicum* m.) [Turkey: Ankara]  
*flavosignatum* Breuning, 1946: 108 (*D. anatolicum* m.) [Turkey: Isparta]  
 Provincial distribution in Turkey: TR-A: AD AN ANT CA IP KA KN (Pic 1900a; Breuning 1946, 1962a; Demelt 1963; Braun 1978a; Adlbauer 1988; Önalp 1991; Özdi̇kmen and Hasbenli 2004a; Özdi̇kmen and Okutaner 2006; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Turgut and Özdi̇kmen 2010; Okutaner et al. 2011; Küçükayakçı et al. 2013; Danilevsky 2015a)

Range: A: TR

\****D. andirinense*** Bernhauer and Peks, 2011a: 218

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Andırın (Altınyayla)' (Turkey: Kahramanmaraş)]

*amaliae* Pesarini and Sabbadini, 2013: 55 [Turkey: Kahramanmaraş: Andırın: Altınyayla]  
 Provincial distribution in Turkey: TR-A: KA (Bernhauer and Peks 2011a, 2013b; Özdi̇kmen 2015; Danilevsky 2015a)

Range: A: TR

\****D. apicerufum*** Breuning, 1943: 91 [RN]

Type information: Holotype, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris as *D. sanguinolentum* [type locality 'Armenia']

*sanguinolentum* J. Thomson, 1865: 548 [HN] [Armenia]  
*thomsonianum* Plavilstshikov, 1948: 126 [RN] [Armenia]  
*lineiferum* Breuning, 1946: 125 (*D. apicerufum* m.) [Armenia: Novo Bajaset]  
 Provincial distribution in Turkey: TR-A: EZ KAR TB (Plavilstshikov 1958; Breuning and Villiers 1967; Löbl and Smetana 2010; Özdi̇kmen 2010; Danilevsky 2015a)  
 Range: A: AB AR GG ?IN TR

\****D. aracense*** Bernhauer and Peks, 2011b: 204

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Araç' (Turkey: Kastamonu)]

Provincial distribution in Turkey: TR-A: KS (Bernhauer and Peks 2011b, 2013b; Danilevsky 2015a)

Range: A: TR

\****D. arcivagum*** J. Thomson, 1867: 121

Type information: Holotype, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Gülek' (Turkey: İçel)]

*cilicum* Pic, 1904: 4 (*D. arcivagum* var.) [Turkey: Cilicia]  
 Provincial distribution in Turkey: TR-A: AD IC (Thomson 1867; Ganglbauer 1884; Pic 1904; Breuning 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)  
 Range: A: TR

**\**D. ardahense*** Breuning, 1975: 10

Type information: Holotype ♀, Muséum National d'Histoire Naturelle, Paris) [type locality 'Ardahan' (Turkey)]

Provincial distribution in Turkey: TR-A: AR (Breuning 1975; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. atritarse*** Pic, 1931a: 10 (*D. pedestre* var.) status nov., comb. nov.

Type information: Syntypes ♂ and ♀, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Greece']

Provincial distribution in Turkey: TR-A: CA; TR-E: ED (Adlbauer 1988; Althoff and Danilevsky 1997; Özdkmen and Demir 2006)

Range: E: GR TR A: TR

**\**D. avgadiense*** Bernhauer and Peks, 2011a: 215

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Erdemli: Avgadi' (Turkey: İçel)]

Provincial distribution in Turkey: TR-A: IC (Bernhauer and Peks 2011a; Danilevsky 2015a)

Range: A: TR

***D. axillare*** Küster, 1847a: 88***D. axillare axillare*** Küster, 1847: 88

Type information: Holotype, ex collection Heinrich Carl Küster > Johann Menzel, Naturhistorisches Museum, Nürnberg [type locality 'Balkans' (Bulgaria)]

*kraatzi* Tournier, 1872: 281 (*D. pedestre* var.) [Turkey]

*varnanum* Pic, 1926: 13 (*D. litigiosum* var.) [Bulgaria: Varna]

Provincial distribution in Turkey: TR-E (Tournier 1872)

Range: E: BU TR

Other known subspecies: *D. axillare moldavicum* Dascalu and Fusu, 2012: 39 E: RO

**\**D. bangi*** Heyden, 1894: 89**\**D. bangi bangi*** Heyden, 1894: 89

Type information: Syntypes 4, ex collection L. von Heyden, Natur-Museum und Forschungs-Institut Senckenberg, Frankfurt am Main [type locality 'Kastamonu' (Turkey)]

Provincial distribution in Turkey: TR-A: AN BO KR KS (Heyden 1894; Breuning 1946, 1962a; Demelt 1967; Sama 1982; Önalp 1991; Adlbauer 1992; Özdkmen et al. 2005; Özdkmen 2007, 2010, 2012; Löbl and Smetana 2010; Danilevsky 2015a)

Range: A: TR

**\**D. bangi heinzorum*** Braun, 1975: 17 (*D. heinzorum*)

Type information: Holotype ♂, collection Walter Braun, Tübingen [type locality 'İskilip' (Turkey: Çorum)]

Provincial distribution in Turkey: TR-A: CO KS (Braun 1975, 1978a, 1979; Adlbauer 1992; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. bangi roridum* Pesarini and Sabbadini, 1998: 55**

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Gerede' (Turkey: Bolu)]

Provincial distribution in Turkey: TR-A: BO (Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdi̇kmen 2010, 2011, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. banjkovskiyi* Plavilstshikov, 1958: 147**

Type information: Syntypes 4♂♂ and 3♀♀, ex collection N. N. Plavilstshikov, Zoological Museum, University of Moscow [type locality 'Erzurum' (Turkey)]

Provincial distribution in Turkey: TR-A: EZ (Plavilstshikov 1958; Breuning 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. basale* Kraatz, 1873: 86**

Type information: Holotype ♂, ex collection Jacob Sturm, Zoologische Staatsammlung des Bayerischen Staates, München [type locality 'Armenien', but undoubtedly Turkey: ? Kars]

*kagyzmanicum* Suvorov, 1915: 120 [Turkey: Kars: Kağızman]

Provincial distribution in Turkey: TR-A: KAR (Kraatz 1873; Suvorov 1915; Plavilstshikov 1958; Breuning 1962a; Löbl and Smetana 2010; Özdi̇kmen 2012; Danilevsky 2015a)

Range: A: TR

**\**D. beloni* Pic, 1891: lxxvii**

Type information: Holotype ♂, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Siberia?' but undoubtedly mislabeled, probably (Turkey: Hatay)]

Provincial distribution in Turkey: TR-A: HT (Pic 1891; Löbl and Smetana 2010; Özdi̇kmen 2012; Danilevsky 2015a)

Range: A: TR

**\**D. bisignatum* Jakovlev, 1899: 66**

Type information: Lectotype ♂, ex collection P. Semenov, Collection of Zoological Institute of the Russian Academy of Sciences, Saint-Petersbourg [type locality 'Artvin' (Turkey)]

Provincial distribution in Turkey: TR-A: ART (Jakovlev 1899; Plavilstshikov 1958; Breuning 1962a; Breuning and Villiers 1967; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Bernhauer and Peks 2013b; Danilevsky 2015a)

Range: A: TR

***D. bistriatum* Pic, 1898: 55**

Type information: Holotype, ex collection Théry > M. Pic, Muséum National d'Histoire Naturelle, Paris) [type locality 'Caucasus']

Provincial distribution in Turkey: TR-A: IG (Breuning 1962a; Özdi̇kmen 2010)

Range: A: AB AR TR

**\**D. bithyniense*** Chevrolat, 1856: 88

Type information: Holotype ♂, ex collection C. Bowring-Chevrolat, The Natural History Museum, London) [type locality 'Bursa' (Turkey)]

*serotinum* J. Thomson, 1865: 549 [Turkey: İzmir]

*dorsoplagiatum* Breuning, 1946: 120 (*D. bithyniense* m.) [Turkey: Bursa]

Provincial distribution in Turkey: TR-A: BL BS IS IZ KN KS SA (Chevrolat 1856; Thomson 1865; Ganglbauer 1884; Breuning 1946; 1962a; Demelt and Alkan 1962; Demelt 1963; Breuning and Villiers 1967; Adlbauer 1988; Önalp 1991; Özdkmen 2010; 2012; Löbl and Smetana 2010; Danilevsky 2015a)

Range: A: TR

***D. blanchardi*** Mulsant and Rey, 1863: 147

Type information: Syntypes, Muséum National d'Histoire Naturelle, Paris [type locality 'Iran']

Provincial distribution in Turkey: TR-A: KA MA (Heyden 1888; Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010; Bernhauer and Peks 2011a; Danilevsky 2015a).

Range: A: IN TR

**\**D. blandulum*** Holzschuh, 1977: 131

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Kuruca pass' (Turkey: Bingöl)]

Provincial distribution in Turkey: TR-A: BN (Holzschuh 1977; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2012, 2015a)

Range: A: TR

**\**D. bodemeyeri*** K. Daniel, 1900: 140

Type information: Holotype ♂, ex collection K. Daniel, Deutsches Entomologisches Institut, Eberswalde [type locality 'Asia Minor (Lycaonia)' (Turkey: ?Konya)]

*rubricolle* Breuning, 1946: 125 (*D. bodemeyeri* m.) [Turkey: Eskişehir]

Provincial distribution in Turkey: TR-A: AF AM AN BO ES IZ KN (K. Bodemeyer 1900; Daniel 1900; Breuning 1946, 1962a; Demelt 1963; Braun 1978a; Sama 1982; Önalp 1990; Löbl and Smetana 2010; Özdkmen 2010, 2011, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. boluense*** Breuning, 1962b: 38

**\**D. boluense boluense*** Breuning, 1962b: 38

Type information: Holotype ♀, collection H. Muche [type locality 'Bolu Mts.' (Turkey: Bolu)]

*nigrobasipenne* Breuning, 1962b: 38 (*D. boluense* m.) [Turkey: Bolu]

Provincial distribution in Turkey: TR-A: BO (Breuning 1962b; Braun 1978a; Adlbauer 1992; Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2011, 2012; Danilevsky 2015a)

Range: A: TR



**\**D. boluense corallinum*** Pesarini and Sabbadini, 1998: 53

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Kızılıcahamam: Çamlıdere' (Turkey: Ankara)]

Provincial distribution in Turkey: TR-A: AN (Braun 1978a; Sama 1982; Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. boluense imitator*** Pesarini and Sabbadini, 1998: 53

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Çubuk: Karagöl' (Turkey: Ankara)]

Provincial distribution in Turkey: TR-A: AN CN (Braun 1978a; Pesarini and Sabbadini 1998; Özdkmen 2006, 2010, 2012; Özdkmen and Demir 2006; Özdkmen et al. 2009; Löbl and Smetana 2010; Danilevsky 2015a)

Range: A: TR

**\**D. boszdaghense*** Fairmaire, 1866: 275

Type information: Syntypes, ex collection L. Fairmaire, Muséum National d'Histoire Naturelle, Paris [type locality 'Ovacık' (Turkey: İzmir)]

*grammophilum* J. Thomson, 1867: 70 ['Armenia', undoubtedly mislabeled]

*rufobasale* Breuning, 1946: 107 (*D. boszdaghense* m.) [Turkey: İzmir: Bozdağ]

Provincial distribution in Turkey: TR-A: IZ (Fairmaire 1866; Thomson 1867; Ganglbauer 1884; Breuning 1946; 1962a; Özdkmen 2010, 2012; Löbl and Smetana 2010; Danilevsky 2015a)

Range: A: TR

**\**D. bouilloni*** Breuning and Ruspoli, 1975: 116

Type information: Holotype ♀, ex collection M. Ruspoli, Museo Civico di Storia Naturale 'Giacomo Doria', Genova [type locality 'Mesudiye' (Turkey: Ordu)]

Provincial distribution in Turkey: TR-A: OR SM SV (Breuning and Ruspoli 1975; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. brauni*** Breuning, 1979: 92

Type information: Holotype ♂, collection W. Heinz, Schwanfeld [type locality 'Tavas Valley' (Turkey: Denizli)]

Provincial distribution in Turkey: TR-A: DE (Breuning 1979; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a)

Range: A: TR

**\**D. bremeri*** Breuning, 1981: 181

Type information: Holotype ♂, collection H. J. Bremer, Naturkundemuseum Erfurt [type locality 'Akyarma' (Turkey: Afyonkarahisar)]

Provincial distribution in Turkey: TR-A: AF (Breuning 1981; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. breunigi*** Heyrovský, 1943: 78

Type information: Syntypes ♂♂ and ♀♀, ex collection Leo Heyrovský, Musée National de Prague [type locality 'Thrazién: Umgebung von Xanthi, Ferre und Badoma' (Greece)]  
 Provincial distribution in Turkey: TR-E (Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdi̇kmen 2010; Danilevsky 2015a)

Range: E: GR TR

**\**D. bulgharmaadense*** Breuning, 1946: 116

Type information: Holotype ♀, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Bulghar-Maaden [Bolkar Mts.-Maden vill. (Çamardı)]' (Turkey: Niğde)]

Provincial distribution in Turkey: TR-A: NI (Breuning 1946, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. cachinno*** J. Thomson, 1867: 120

Type information: Syntypes, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Caramania' (Turkey: Karaman)]

*immersum* Tournier, 1872: 281 [Turkey]

*fuscoflavum* Breuning, 1946: 110 (*D. cachinno* m.) [Turkey]

Provincial distribution in Turkey: TR-A: KM US (Thomson 1867; Breuning 1946, 1962a; Demelt 1963; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. caglayanceritense*** Bernhauer and Peks, 2013a: 314

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Çağlayancerit' (Turkey: Kahramanmaraş)]

Provincial distribution in Turkey: TR-A: KA (Bernhauer and Peks 2013a; Danilevsky 2015a)

Range: A: TR

**\**D. carinipenne*** Pic, 1900a: 13

Type information: Holotype, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Akbez' (Turkey: Hatay)]

*quinquevittatum* Breuning, 1946: 127 (*D. carinipenne* m.) [Turkey: Gaziantep]

Provincial distribution in Turkey: TR-A: GA HT (Pic 1900a; Breuning 1946, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. carolisturanii*** Breuning and Ruspoli, 1971: 128

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Kümbet' (Turkey: Giresun)]

*baiburtense* Breuning, 1975: 10 (*D. carolisturanii* m.) [Turkey: Bayburt]

Provincial distribution in Turkey: TR-A: BY GI OR (Breuning and Ruspoli 1971; Breuning 1975; Braun 1978a; Sama 1982; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR



**D. catenatum** Waltl, 1838: 469

**D. catenatum catenatum** Waltl, 1838: 469

Type information: Holotype ♀, ex collection Imre Frivaldszky, Magyar Természettudományi Múzeum, Budapest [type locality 'Turkei' (Turkey)]

*divisum* Germar, 1839: 15 [Turkey]

*mancum* Gistel, 1848: 431 [?Syria]

*subobliteratum* T. Pic, 1899: 351 (*D. divisum* var.) [Turkey]

*uninterruptum* T. Pic, 1899: 351 (*D. divisum* var.) [Turkey: İzmir]

*dorsale* Pic, 1907: 179 (*D. divisum* var.) [Turkey: İzmir]

*sparsedivisum* Pic, 1911: 185 (*D. divisum* var.) [Greece: Lesbos (Mytilene) Island]

*smyrnense* Pic, 1917a: 10 (*D. divisum* var.) [Turkey: İzmir]

*thebesianum* Pic, 1942d: 1 (*D. divisum* var.) [Greece: Thebes]

*bonyi* Pic, 1942a: 1 (*D. divisum* var.) [?Syria]

*smyrnanum* Breuning, 1946: 106 [RN] [Turkey: İzmir]

*postmediomaculatum* Breuning, 1946: 106 (*D. divisum* m.) [Turkey: İzmir]

*chioticum* Breuning, 1946: 106 (*D. divisum* m.) [Greece: Chios Island]

*tenebripes* Breuning, 1946: 106 (*D. divisum* m.) [Greece: Samos Island]

*parteinterruptum* Breuning, 1962a: 384 (*D. divisum* m.) [Turkey: Manisa: Demirci]

*subatrum* Breuning, 1962a: 385 (*D. divisum* m.) [Turkey: Balıkesir]

*peratrum* Breuning, 1962a: 385 (*D. divisum* m.) [Greece: Lesbos Island: Ayassos]

*submytilinense* Breuning, 1962a: 386 (*D. divisum* m.) [Greece: Lesbos Island]

*dorsaloides* Breuning, 1966a: 20 (*D. divisum* m.) [Turkey: İzmir: Bozdağ]

Provincial distribution in Turkey: TR-A: BL BS BU ES IP IZ KN KU MN (Küster 1846a; Kraatz 1873; Ganglbauer 1884; Pic 1899, 1917a; Breuning 1946; 1962a, 1966a; Demelt 1963; Sama 1982; Önalp 1991; Özdi̇kmen 2010; 2012; Löbl and Smetana 2010; Şenyüz and Özdi̇kmen 2013; Danilevsky 2015a)

Range: E: GR A: TR

**\*D. catenatum dissimile** Ganglbauer, 1884: 458 (*D. divisum* var.)

Type information: Syntypes, ex collection L. Ganglbauer, Naturhistorisches Museum Wien [type locality 'European Turkey']

Provincial distribution in Turkey: TR-E: CA ED (Ganglbauer 1884; Pesarini and Sabbadini 2009; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Sama et al. 2010; Danilevsky 2015a)

Range: E: TR

**\*D. catenatum intercicum** Kraatz, 1873: 66 (*D. latevittatum* var.)

Type information: Holotype, ex Musaeo Mniszech > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Kleinasién' (Turkey: Diyarbakır)]

Provincial distribution in Turkey: TR-A: AD ADY IC MR NI OS SV (Kraatz 1873; Bodemeyer 1900; Braun 1978a; Önalp 1990; Özdi̇kmen 2006, 2010, 2012; Löbl and Smetana 2010; Danilevsky 2015a)

Range: A: TR

**\*D. catenatum loratum** J. Thomson, 1867: 123 (*D. loratum*)

Type information: Holotype ♂, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Smyrne' (Turkey: İzmir)]

Provincial distribution in Turkey: TR-A: CA IZ (Thomson 1867; Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. catenatum mytilenense*** Kraatz, 1873: 66 (*D. latevittatum* var.)

Type information: Holotype, ex Musaeo Mniszech > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Kleinasiens' (Turkey)]

*latevittatum* Kraatz, 1873: 66 [Turkey]

Provincial distribution in Turkey: TR-A: IZ (Kraatz 1873; Löbl and Smetana 2010; Özdkmen 2010)

Range: E: GR A: TR

**\**D. catenatum oedemischense*** Heyrovský, 1932: 104 (*D. divisum* var.)

Type information: Syntypes 5, Zoologische Staatssammlung des Bayerischen Staates, München [type locality 'Ödemiş' (Turkey: İzmir)]

Provincial distribution in Turkey: TR-A: IZ (Heyrovský 1932; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. catenatum subdivisum*** Breuning, 1955: 263 (*D. divisum* ssp.)

Type information: Holotype ♂, Magyar Természettudományi Múzeum, Budapest) [type locality 'Ankara' (Turkey)]

*submediomaculatum* Breuning, 1955: 263 (*D. divisum subdivisum* m.) [Turkey: Ankara]

Provincial distribution in Turkey: TR-A: AN (Breuning 1955; Fuchs and Breuning 1971; Özdkmen 2006, 2010, 2012; Löbl and Smetana 2010; Danilevsky 2015a)

Range: A: TR

Other known subspecies: *D. catenatum bleusei* Pic, 1899: 300 E: GR

**\**D. cinctellum*** Fairmaire, 1866: 272

Type information: Syntypes ♂♂, ex collection L. Fairmaire, Muséum National d'Histoire Naturelle, Paris [type locality 'Kisilgye-Aole (Ödemiş: Kızılca köyü)' (Turkey: İzmir)]

*aureofasciatum* Breuning, 1947: 168 ['Kisilgye-Aole', Turkey: İzmir: Ödemiş: Kızılca]

*quadrifuscoplagiatum* Breuning, 1947: 168 (*D. cinctellum* m.) [Turkey: İzmir: Ödemiş]

*reductum* Breuning, 1947: 168 [Turkey: İzmir: Ödemiş]

Provincial distribution in Turkey: TR-A: HT IZ KN (Fairmaire 1866; Ganglbauer 1884; Pic 1917a; Breuning 1947, 1962a; Adlbauer 1992; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. cingulatum*** Ganglbauer, 1884: 484

Type information: Syntypes 2♀♀, ex collection L. Ganglbauer, Naturhistorisches Museum Wien [type locality 'Persien' (Iran)]

*invicinum* Pic, 1902a: 93 [Turkey: Diyarbakır]

*persianum* Breuning, 1943: 94 [Iran]

Provincial distribution in Turkey: TR-A: DI (Pic 1902a; Breuning 1962a; Villiers 1967; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a)



Range: A: IN TR

**\**D. coiffaiti*** Breuning, 1962a: 392

Type information: Holotype ♂, ex collection Coiffait, Muséum National d'Histoire Naturelle, Paris [type locality 'Manyas' (Turkey: Balıkesir)]

Provincial distribution in Turkey: TR-A: BL (Breuning 1962a; Pesarini and Sabbadini 2009; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. condensatum*** Küster, 1852: 92

Type information: Syntypes ♂♂ and ♀♀, ex collection Kiesenwetter, Zoologische Staatssammlung des Bayerischen Staates, München [type locality 'Turkei' (Turkey)]

*punctatissimum* Ganglbauer, 1884: 94 (*D. condensatum* var.) [Turkey]

*adampolum* Pic, 1917b: 6 (*D. condensatum* var.) [Turkey: İstanbul: Polonezköy (Adampol)]

*inlineatum* Pic, 1917b: 6 (*D. condensatum* var.) [Turkey]

Provincial distribution in Turkey: TR-A: IS KO TR-E: IS (Ganglbauer 1884; Bodemeyer 1906; Pic 1917b; Demelt and Alkan 1962; Breuning 1962a; Demelt 1963; Braun 1978a; Önalp 1990; Adlbauer 1992; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: E: BU TR A: TR

**\**D. confluens*** Fairmaire, 1866: 274

Type information: Syntypes ♂♂ and ♀♀, ex collection L. Fairmaire, Muséum National d'Histoire Naturelle, Paris [type locality 'Bozdağ' (Turkey: İzmir)]

*subobscuripes* Pic, 1914: 9 [Turkey: İzmir: Bozdağ]

Provincial distribution in Turkey: TR-A: IZ (Fairmaire 1866; Pic 1914; Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. conspersum*** Pesarini and Sabbadini, 2013: 56

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Bayramgazi' (Turkey: Afyonkarahisar)]

Provincial distribution in Turkey: TR-A: AF (Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

***D. cretosum*** 1864: 479

Type information: Syntypes ♂♂ and ♀♀, Naturhistorisches Museum Wien [type locality 'Achalzik' (Georgia)]

*niveisparsum* J. Thomson, 1865: 548 [Caucasus]

*sobrinum* Pic, 1898: 55 (*D. niveisparsum* var.) [Caucasus]

*postobliteratum* Pic, 1942c: 1 (*D. niveisparsum* var.) [Caucasus]

Provincial distribution in Turkey: TR-A: EZ MN (Plavilstshikov 1958; Sama 1982; Önalp 1990; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a)

Range: A: GG TR

**\**D. crux* Billberg, 1817: 175 (*Lamia*)**

Type information: Holotype, ex Musaeo Schönherr, Naturhistoriska Riksmuseet, Stockholm [type locality unknown but very likely İzmir (Turkey)]

- smyrnensis* Linnaeus, 1757: 411 (*Cerambyx*) [Turkey: İzmir]
- graecum* Waltl, 1838: 470 [Greece]
- cruciator* Gistel, 1848: 131 [?Syria]
- bifidum* Chevrolat, 1882: 60 [Turkey: İzmir]
- lunulatum* Pic, 1900a: 12 (*D. graecum* var.) [Turkey: İzmir]
- integrofasciatum* K. Daniel, 1900: 140 (*D. crux* var.) [Turkey: İzmir: Bozdağ]
- niveopictum* K. Daniel, 1900: 140 (*D. crux* var.) [Turkey: İzmir: Bozdağ]
- biinterruptum* Pic, 1901: 19 (*D. crux* var.) [Turkey]
- grandevittatum* Pic, 1904: 7 [Turkey: İmbey]
- subfurcatum* Pic, 1914: 9 (*D. cinctellum* var.) [Turkey]
- brevedisjunctum* Pic, 1929: 9 (*D. crux* var.) [Turkey]
- lateargenteum* Pic, 1929: 9 (*D. crux* var.) [Turkey]
- quadriinterruptum* Breuning, 1946: 101 (*D. smyrnense* m.) [Turkey: İzmir]
- subniveopictum* Breuning, 1946: 101 (*D. smyrnense* m.) [Turkey: Eskişehir]
- holodisjunctum* Breuning, 1946: 101 (*D. smyrnense* m.) [Turkey: Eskişehir]
- rufulum* Breuning, 1946: 106 (*D. smyrnense* m.) [Turkey: İzmir]
- dorsoabbreviatum* Breuning, 1946: 106 (*D. smyrnense* m.) [Turkey: Konya: Akşehir]
- semiinterruptum* Breuning, 1946: 106 (*D. smyrnense* m.) [Turkey: Eskişehir]
- lunulatooides* Breuning, 1948: 58 (*D. smyrnense* m.) [Turkey]
- bodemeyerianum* Breuning, 1962a: 279 (*D. smyrnense* m.) [Turkey: İzmir: Bozdağ]

Provincial distribution in Turkey: TR-A: BI BL BS ES IS IZ KN KU US (Ganglbauer 1884; Bodemeyer 1900, 1906; Pic 1900a; Breuning 1946, 1962a; Demelt 1963; Tuatay et al. 1972; Braun 1978a; Adlbauer 1988; Önalp 1991; Özdi̇kmen et al. 2005; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. culminicola* J. Thomson, 1867: 122**

Type information: Holotype ♂, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Caramania' (Turkey: Karaman)]

Provincial distribution in Turkey: TR-A: KM (Thomson 1867; Ganglbauer 1884; Breuning 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. delagrangei* Pic, 1894: 110**

Type information: Syntypes ♂ and ♀, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Akbez' (Turkey: Hatay)]

- mardinense* Pic, 1900b: 6 (*D. accola* var.) [Turkey: Mardin]
- fuscopubescens* Pic, 1900a: 12 (*D. delagrangei* var.) [Turkey: Hatay: Akbez]

Provincial distribution in Turkey: TR-A: GA HT KL MR OS (Pic 1894, 1897, 1900a, 1900b; Breuning 1962a; Braun 1978a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Bernhauer and Peks 2011a; Danilevsky 2015a)

Range: A: TR



**\**D. deyrollei*** Ganglbauer, 1884: 482

**\**D. deyrollei deyrollei*** Ganglbauer, 1884: 482

Type information: Syntypes ♂♂, ex collection L. Ganglbauer, Naturhistorisches Museum Wien [type locality 'Bitlis' (Turkey)]

*subalbipenne* Breuning, 1946: 116 (*D. deyrollei* m.) [Turkey: Van]

Provincial distribution in Turkey: TR-A: BT VA (Ganglbauer 1884; Breuning 1946, 1962a; Plavilstshikov 1958; Fuchs and Breuning 1971; Holzschuh 1980; Önalp 1990; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. deyrollei opacum*** Pesarini and Sabbadini, 2013: 54

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Ahır Mt.' (Turkey: Kahramanmaraş)]

*melas* Breuning, 1946: 116 (*D. deyrollei* m.) [Turkey: Kahramanmaraş]

Provincial distribution in Turkey: TR-A: GA KA (Breuning 1946, 1962a; Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

***D. dimidiatum*** Motschulsky, 1838: 186

***D. dimidiatum dimidiatum*** Motschulsky, 1838: 186

Type information: Syntypes ♂♂ and ♀♀, ex collection V. I. Motschulsky, Zoological Museum, University of Moscow [type locality 'Armenia']

*erivanicum* Pic, 1901: 91 [Armenia: Erevan]

*reductemaculatum* Breuning, 1946: 123 (*D. dimidiatum* m.) [Transcaucasia]

*flavescens* Breuning, 1946: 124 (*D. dimidiatum* m.) [Armenia: Erivan]

*nigrovittatum* Breuning, 1946: 124 (*D. dimidiatum* m.) [Armenia: Erivan]

*nigroreductum* Breuning, 1946: 124 (*D. dimidiatum* m.) [Armenia: Erivan]

*unimaculatum* Breuning, 1946: 124 (*D. dimidiatum* m.) [Armenia: Erivan]

*degaullei* Pic, 1948: 13 (*D. dimidiatum* var.)

*transitorium* Pic, 1948: 13

Provincial distribution in Turkey: TR-A: AG ER EZ MA (Thomson 1867; Plavilstshikov 1958; Villiers 1967; Gfeller 1972; Özbek 1978; Braun 1978a; Önalp 1991; Özdi̇kmen and Hasbenli 2004b; Löbl and Smetana 2010; Özdi̇kmen 2010; Danilevsky 2015a)

Range: A: AB AR GG IN TR

**\**D. dimidiatum kelkiticum*** Özdi̇kmen and Hasbenli, 2004b: 23

Type information: Holotype ♂, Zoological Museum of Gazi University, Ankara [type locality 'Kelkit' (Turkey: Gümüşhane)]

Provincial distribution in Turkey: TR-A: GU (Özdikmen and Hasbenli 2004b; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. dimidiatum korgei*** Breuning, 1966a: 21

Type information: Holotype ♀, Institut Royal des Sciences Naturelles de Belgique [type locality 'near Pülümür' (Turkey: Tunceli)]

Provincial distribution in Turkey: TR-A: TU (Breuning 1966a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. dobrovjanskii* Suvorov, 1915: 116**

Type information: Holotype ♂, ex collection G. Suvorov, Zoological Museum, University of Moscow [type locality 'Ervansk. g.' (should be 'NE Turkey')]

Provincial distribution in Turkey: TR-A: North-Eastern TR (Suvorov 1915; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. drusoides* Breuning, 1962a: 375**

Type information: Holotype ♂, ex collection Lepesme, Musée d'Histoire Naturelle de Lyon [type locality 'Amanos Mts.' (Turkey: Hatay)]

Provincial distribution in Turkey: TR-A: HT (Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. eflaniense* Bernhauer and Peks, 2011b: 205**

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Eflani: Kırıklär' (Turkey: Karabük)]

Provincial distribution in Turkey: TR-A: KR KS (Bernhauer and Peks 2011b; Danilevsky 2015a)

Range: A: TR

**\**D. elazigi* Fuchs and Breuning, 1971: 439**

Type information: Holotype ♂, Anatolianexpedition 1966, Naturhistorisches Museum Wien [type locality 'Buğlan pass' (Turkey: Muş)]

*holzschuhi* Breuning, 1974a: 148 [Turkey: Muş: Buğlan pass]

Provincial distribution in Turkey: TR-A: BN EL MU (Fuchs and Breuning 1971; Breuning 1974a; Holzschuh 1980; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. enricisturanii* Breuning and Ruspoli, 1971: 127**

**\**D. enricisturanii densepunctatum* Braun, 1978a: 107**

Type information: Holotype ♂, collection Walter Braun, Tübingen [type locality 'Vezirköprü' (Turkey: Samsun)]

Provincial distribution in Turkey: TR-A: SM (Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. enricisturanii enricisturanii* Breuning and Ruspoli, 1971: 127**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'SE Tosya' (Turkey: Kastamonu)]

Provincial distribution in Turkey: TR-A: CO KS (Breuning and Ruspoli 1971; Braun 1975, 1978a; Adlbauer 1992; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR



**D. equestris** Laxmann, 1770: 596 (*Cerambyx equestris*)

\***D. equestris bernhauerorum** Peks, 2010: 213 status nov., comb. nov.

Type information: Holotype ♂, collection Heinz Peks, Schwalbach [type locality 'İznik: Haciosman village' (Turkey: Bursa)]

Provincial distribution in Turkey: TR-A: BS (Peks 2010; Özdi̇kmen 2012; Danilevsky 2015a)

Range: A: TR

**D. equestris bisuturale** Jureček, 1933: 128 (*D. equestris* var.)

Type information: Syntypes 7 ♂♂ and ♀♀, ex collection Štepán Jureček, Musée National de Prague [type locality 'Griechenland, Bos-Dagh-Gebirge bei Drama' (Greece)]

Provincial distribution in Turkey: TR-E (Breuning 1962a; Althoff and Danilevsky 1997; Danilevsky 2015a)

Range: E: BU GR TR

\***D. equestris festivum** Pesarini and Sabbadini, 2013: 58

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Sündiken Mts.: Karatepe' (Turkey: Eskişehir)]

Provincial distribution in Turkey: TR-A: ES (Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

\***D. equestris nogelli** Fairmaire, 1866: 270 (*D. nogelli*)

Type information: Syntypes ♂♂ and ♀♀, ex collection L. Fairmaire, Muséum National d'Histoire Naturelle, Paris [type locality 'Bozdağ et Ovacık' (Turkey: İzmir)]

*nogelli* Thomson, 1867: 58 [?Armenia]

*exclamationis* Thomson, 1867: 58 (*D. nogelli* var.) [?Armenia]

Provincial distribution in Turkey: TR-A: IZ EZ (Fairmaire 1866; Ganglbauer 1884; Breuning 1962a; Önalp 1991; Pesarini and Sabbadini 1998, 2013, Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Peks 2010; Danilevsky 2015a)

Range: A: TR

Other known subspecies: *D. equestris equestris* Laxmann, 1770: 596 (*Cerambyx equestris*) E: CT ST UK A: GG; *D. equestris reclinatum* Kraatz, 1892: 173 E: AL GR MC YU; *D. equestris transsilvanicum* Ganglbauer, 1884: 462 E: MD RO

\***D. erdemi** Özdi̇kmen, Kaya and Al-Hamadani, 2014: 179

Type information: Holotype ♂, collection H. Özdi̇kmen, Zoological Museum of Gazi University, Ankara [type locality 'Sungurlu road' (Turkey: Çorum)]

Provincial distribution in Turkey: TR-A: CO (Özdikmen et al. 2014; Danilevsky 2015a)

Range: A: TR

\***D. eskipazarensis** Bernhauer and Peks, 2011b: 203

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Eskipazar' (Turkey: Karabük)]

Provincial distribution in Turkey: TR-A: KR (Bernhauer and Peks 2011b; Danilevsky 2015a)

Range: A: TR

***D. ferruginipes* Ménétriés, 1836: 151**

Type information: Holotype, Collection of Zoological Institute of the Russian Academy of Sciences, Saint-Petersbourg [type locality 'İstanbul' (Turkey)]

*thracicum* Küster, 1846a: 90 [Turkey: İstanbul]

*sublineatum* Pic, 1909b: 99 (*D. ferruginipes* var.) [Turkey]

*subfuscopubens* Pic, 1911: 7 (*D. ferruginipes* var.) [Turkey: İstanbul: Alem Mt.]

Provincial distribution in Turkey: TR-A: IS KO SA; TR-E: IS (Ménétriés 1836; Küster 1846a; Ganglbauer 1884; Breuning 1962a; Braun 1978a; Önalp 1991; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdiikmen 2010, 2012; Danilevsky 2015a)

Range: E: GR TR A: TR

**\**D. formosum* Kraatz, 1871: 411**

**\**D. formosum formosum* Kraatz, 1871: 411**

Type information: Syntypes, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Caucasus' but undoubtedly mislabeled, should be Turkey]

*inincisum* Breuning, 1946: 124 (*D. formosum* m.) ['Ain-Tab', Turkey: Gaziantep]

Provincial distribution in Turkey: TR-A: GA (Kraatz 1871; Breuning 1946, 1962a; Löbl and Smetana 2010; Özdiikmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. formosum ponticum* Breuning, 1964a: 1**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Eğribel pass' (Turkey: Giresun)]

Provincial distribution in Turkey: TR-A: GI TO (Breuning 1962a, 1964a; Braun 1978a; Löbl and Smetana 2010; Özdiikmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. gallipolitanum* J. Thomson, 1867: 59**

***D. gallipolitanum gallipolitanum* J. Thomson, 1867: 59**

Type information: Holotype ♂, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Gallipoli' (Turkey: Çanakkale)]

*suturaalba* Desbrochers des Loges, 1870: 125 [Turkey: Çanakkale]

*gandolphi* Tournier, 1872: 288 [Turkey: Çanakkale]

*potamense* Jurecek, 1931: 124 (*D. gallipolitanum* ab.) [Turkey: Sinop: Dedeağaç env.]

*biapicevittatum* Breuning, 1946: 110 (*D. gallipolitanum* m.) [Turkey: İstanbul]

*latesuturale* Breuning, 1946: 110 (*D. gallipolitanum* m.) [Greece: Thrace]

*subregulare* Breuning, 1948: 59 (*D. gallipolitanum* m.) [Turkey: Çanakkale (European Turkey part)]

*stramentosopubescens* Breuning, 1948: 59 (*D. gallipolitanum* m.) ['Turkey: Port Baclar', Turkey: ?İstanbul: ?Bağcılar]

*plurivittatum* Breuning, 1948: 59 (*D. gallipolitanum* m.) ['Turkey: Port Baclar', Turkey: ?İstanbul: ?Bağcılar]

Provincial distribution in Turkey: TR-A: CA IS SN; TR-E: CA ED IS (Thomson 1867; Ganglbauer 1884; Breuning 1946, 1948, 1962a; Braun 1978a; Adlbauer 1988; Önalp 1990; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdiikmen 2010, 2012; Danilevsky 2015a)

Range: E: GR TR A: TR

Other known subspecies: *D. gallipolitanum fumidum* Pesarini and Sabbadini, 2010: 186 E: GR; *D. gallipolitanum samothrakeanum* Breuning, 1962a: 326 E: GR

**\**D. gebzeense*** Breuning, 1974a: 148

Type information: Holotype ♂, Collection Carolus Holzschuh, Villach [type locality 'Gebze' (Turkey: Kocaeli)]

*tosyense* Breuning in Sama, 1982: 223 [Turkey: Kastamonu: Tosya]

Provincial distribution in Turkey: TR-A: KO KS (Breuning 1974a; Braun 1975, 1978a; Sama 1982; Özdi̇kmen 2010, 2012; Özdi̇kmen and Kaya 2015e)

Range: A: TR

**\**D. gencense*** Bernhauer and Peks, 2014a: 110

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach) [type locality 'SW Genç' (Turkey: Bingöl)]

Provincial distribution in Turkey: TR-A: BN (Bernhauer and Peks 2014a; Danilevsky 2015a)

Range: A: TR

**\**D. glabrimolle*** Breuning, 1943: 92

Type information: Holotype ♀, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Anatolia' (Turkey)]

Provincial distribution in Turkey: TR-A (Breuning 1943, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. goelbasiense*** Bernhauer and Peks, 2013a: 310

**\**D. goelbasiense goelbasiense*** Bernhauer and Peks, 2013a: 310

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Gölbaşı' (Turkey: Adiyaman)]

Provincial distribution in Turkey: TR-A: ADY (Bernhauer and Peks 2013a; Danilevsky 2015a)

Range: A: TR

**\**D. goelbasiense karadutense*** Bernhauer and Peks, 2013a: 311

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Kahta: Karadut' (Turkey: Adiyaman)]

Provincial distribution in Turkey: TR-A: ADY (Bernhauer and Peks 2013a; Danilevsky 2015a)

Range: A: TR

**\**D. goelbasiense kathaense*** Bernhauer and Peks, 2013a: 312

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Kahta' (Turkey: Adiyaman)]

Provincial distribution in Turkey: TR-A: ADY (Bernhauer and Peks 2013a; Danilevsky 2015a)

Range: A: TR

**\**D. goelbasiense oeksuezdagense*** Bernhauer and Peks, 2013a: 311

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Öksüz Mt.' (Turkey: Kahramanmaraş)]

Provincial distribution in Turkey: TR-A: KA (Bernhauer and Peks 2013a; Danilevsky 2015a)

Range: A: TR

**\**D. guzeldereense*** Bernhauer and Peks, 2012: 213

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Başkale: Güzeldere pass' (Turkey: Van)]

*guezeldereense* Bernhauer and Peks, 2013b: 336 [unjustified emendation]

Provincial distribution in Turkey: TR-A: VA (Bernhauer and Peks 2012, 2013b; Danilevsky 2015a)

Range: A: TR

**\**D. hacgedigiense*** Bernhauer and Peks, 2011a: 216

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Haçgediği' (Turkey: İçel)]

Provincial distribution in Turkey: TR-A: IC (Bernhauer and Peks 2011a, 2013b; Danilevsky 2015a)

Range: A: TR

***D. haemorrhoidale*** Hampe, 1852: 313

Type information: Syntypes, ex collection Moritz Wagner, Museum für Naturkunde der Humboldt-Universität zu Berlin [type locality 'Armenia']

*sublineiferum* Breuning, 1946: 125 (*D. haemorrhoidale* m.) [Armenia: Erivan]

*vittidorsum* Plavilstshikov, 1958: 113 (*D. haemorrhoidale* m.) [?Armenia]

Provincial distribution in Turkey: TR-A: AG AN EZ (Plavilstshikov 1958; Gfeller 1972; Braun 1978a; Önalp 1990; Löbl and Smetana 2010; Özdiğmen 2010; Danilevsky 2015a)

Range: A: AR IN TR

***D. halepense*** Kraatz, 1873: 72 (*D. saulcyi* var.)

***D. halepense halepense*** Kraatz, 1873: 72 (*D. saulcyi* var.)

Type information: Syntypes, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Aleppo' (Syria)]

*internesignatum* Pic, 1914: 79 (*D. aleppense* var.) [Syria: Aleppo]

*latealbum* Pic, 1926: 13 (*D. aleppense* var.) [Syria: Aleppo]

*latebivittatum* Pic, 1931a: 1 (*D. aleppense* var.) [?]

*internenotatum* Pic, 1931c: 9 (*D. aleppense* var.) [Syria: Aleppo]

*brunneosericans* Breuning, 1946: 101 (*D. halepense* m.) [Syria: Aleppo]

*akbesianum* Breuning, 1946: 101 (*D. halepense* m.) [Turkey: Hatay: Akbez]

*pseudosimile* Breuning, 1946: 102 (*D. halepense* m.) [Turkey: Hatay: Akbez]

*pseudomardinense* Breuning, 1946: 102 (*D. halepense* m.) [Turkey: Hatay: Akbez]

*dorsoinvittatum* Breuning, 1948: 58 (*D. halepense* m.) [Turkey: Hatay: Akbez]

Provincial distribution in Turkey: TR-A: HT (Breuning 1946, 1948; Önalp 1991; Löbl and Smetana 2010; Özdiğmen 2010; Danilevsky 2015a)

Range: A: SY TR



**\**D. halepense sehitkamilense*** Özdkmen, Mercan and Cihan, 2012a: 604

Type information: Holotype ♂, collection H. Özdkmen, Zoological Museum of Gazi University, Ankara [type locality 'Şehitkamil' (Turkey: Gaziantep)]

Provincial distribution in Turkey: TR-A: GA KL (Breuning 1962a; Braun 1978a; Özdkmen 2010, 2012; Özdkmen et al. 2012a; Danilevsky 2015a)

Range: A: TR

**\**D. hampii*** Mulsant and Rey, 1863: 157

**\**D. hampii aureovittatum*** Kraatz, 1873: 81 (*D. aureovittatum*)

Type information: Syntypes ♂♂, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Kleinasiens' (Turkey)]

Provincial distribution in Turkey: TR-A: SV (Kraatz 1873; Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. hampii hampii*** Mulsant and Rey, 1863: 157

Type information: Holotype, ex collection Petri Pellet, Muséum d'Histoire Naturelle de Perpignan [type locality 'La Perse' undoubtedly mislabeled, should be Turkey]

*claresuturale* Breuning, 1946: 109 (*D. hampei* m.) [Turkey: Amasya]

*discofasciatum* Breuning, 1946: 110 (*D. hampei* m.) [Turkey: Amasya]

Provincial distribution in Turkey: TR-A: AM EZ KAR TO (Breuning 1946, 1962a; Plavilstshikov 1958; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a)

Range: A: TR

**\**D. hanneloreae*** Bernhauer and Peks, 2014a: 107

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'S Bulanık' (Turkey: Muş)]

Provincial distribution in Turkey: TR-A: MU (Bernhauer and Peks 2014a; Danilevsky 2015a)

Range: A: TR

**\**D. heinzi*** Breuning, 1964a: 1 (*D. cinerarium* ssp.)

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Eğribel pass' (Turkey: Giresun)]

Provincial distribution in Turkey: TR-A: GI (Breuning 1964a; Braun 1978a, 1979; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. hellmanni*** Ganglbauer, 1884: 486

Type information: Syntypes ♂♂ and ♀♀, ex collection L. Ganglbauer, Naturhistorisches Museum Wien [type locality 'Persien' (Iran)]

*ingens* Breuning, 1946: 100 (*D. hellmanni* m.) [Turkey: Van]

*postjunctum* Breuning, 1946: 100 (*D. hellmanni* m.) [Turkey: Van]

Provincial distribution in Turkey: TR-A: HA MR MU TU VA (Breuning 1946; Plavilstshikov 1958; Fuchs and Breuning 1971; Holzschuh 1980; Löbl and Smetana 2010; Özdkmen 2010; Sama et al. 2012; Danilevsky 2015a)

Range: A: IN IQ TR

**\**D. holtzi*** Pic, 1905a: 115 (*D. culminicola* ssp.)

Type information: Holotype, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Taurus' (Turkey: ?İçel)]

Provincial distribution in Turkey: TR-A: IC (Pic 1905a; Bodemeyer 1906; Breuning 1962a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. iconiense*** K. Daniel, 1900: 140

Type information: Syntypes ♂♂, ex collection K. Daniel, Deutsches Entomologisches Institut, Eberswalde [type locality 'Asia Minor' (Turkey)]

*semisetosum* Jakovlev, 1901: 85 [Turkey: Konya]

*fulvovestitum* Pic, 1903a: 5 (*D. iconiense* var.) [Turkey]

*albicolle* Breuning, 1943: 89 [Turkey: Konya]

*posticeapertum* Breuning, 1946: 109 (*D. iconiense* m.) [Turkey: Konya]

Provincial distribution in Turkey: TR-A: AK CO KIR KN KY YO (Daniel 1900; Jakovlev 1901; Pic 1903a; Breuning 1943, 1946, 1962b; Braun 1978a, 1979; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Lazarev 2011; Danilevsky 2015a; Özdīkmen and Kaya 2015a)

Range: A: TR

**\**D. infernale*** Mulsant and Rey, 1863: 158

**\**D. infernale asperatum*** Breuning, 1947: 169

Type information: Holotype ♀, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Diyarbakır' (Turkey)]

Provincial distribution in Turkey: TR-A: DI (Breuning 1947, 1962a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. infernale edremitense*** Breuning, 1966a: 20

Type information: Holotype ♂, Institut Royal des Sciences Naturelles de Belgique [type locality 'Edremit' (Turkey: Balıkesir)]

Provincial distribution in Turkey: TR-A: BL (Breuning 1966a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. infernale infernale*** Mulsant and Rey, 1863: 158

Type information: Holotype, ex collection Petri Pellet, Muséum d'Histoire Naturelle de Perpignan [type locality 'La Perse' undoubtedly mislabeled, should be Turkey]

*rugosum* J. Thomson, 1867: 46 [?Russia: Siberia]

*revestitum* K. Daniel, 1900: 140 (*D. infernale* var.) [Turkey]

*immutatum* Pic, 1903b: 171 (*D. infernale* var.) [Turkey: İzmir: Bozdağ]

*costidorsum* Heyrovský, 1932: 103 (*D. infernale* ab.) [Turkey: Eskişehir]

*subrevestitum* Breuning, 1946: 115 (*D. infernale* m.) [Turkey: Bilecik]

*murinum* Breuning, 1946: 115 (*D. infernale* m.) [Turkey: İzmir: Bozdağ]

*irrevestitum* Breuning, 1946: 115 (*D. infernale* m.) [Turkey: Bilecik]

Provincial distribution in Turkey: TR-A: AM AN ANT BI BU CO ES IC IZ KN KU NI SV US; TR-E: KK (Ganglbauer 1884; Bodemeyer 1900; 1906; Daniel 1900; Pic 1903b; Daniel and Daniel 1903; Heyrovský 1932; Breuning 1946; 1962a; Demelt 1963; Braun 1978a; Önalp

1990; Adlbauer 1992; Özdi̇kmen and Hasbenli 2004a; Özdi̇kmen 2006; 2010, 2012; Özdi̇kmen et al. 2009; Löbl and Smetana 2010; Sama et al. 2012; Şenyüz and Özdi̇kmen 2013; Danilevsky 2015a).

Range: E: TR A: TR

**\**D. infernale miminfernale*** Breuning, 1970: 98 (*D. miminfernale*)

Type information: Holotype ♂, Muséum National d'Histoire Naturelle, Paris [type locality 'Binboğa Mt.: Göksun' (Turkey: Kahramanmaraş)]

Provincial distribution in Turkey: TR-A: KA KY (Breuning 1970; Braun 1978a; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. inikliense*** Bernhauer and Peks, 2010b: 208

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'İznik' (Turkey: Bursa)]

Provincial distribution in Turkey: TR-A: BS (Bernhauer and Peks 2010; Özdi̇kmen 2012; Danilevsky 2015a)

Range: A: TR

**\**D. inopinatum*** Pesarini and Sabbadini, 2013: 51

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Bozdağ' (Turkey: İzmir)]

Provincial distribution in Turkey: TR-A: IZ (Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

**\**D. inspersum*** Holzschuh, 1982: 76

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Kuzgunkırın pass' (Turkey: Van)]

Provincial distribution in Turkey: TR-A: VA (Holzschuh 1982; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. intrusum*** Pesarini and Sabbadini, 2009: 26

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'SW Gelibolu' (Turkey: Edirne)]

Provincial distribution in Turkey: TR-E: ED (Pesarini and Sabbadini 2009; Özdi̇kmen 2012; Danilevsky 2015a)

Range: E: TR

**\**D. ispartense*** Breuning, 1960: 1

Type information: Holotype ♀, ex collection H. Coiffait, Muséum National d'Histoire Naturelle, Paris [type locality 'Bozanönü' (Turkey: Isparta)]

Provincial distribution in Turkey: TR-A: IP (Breuning 1960, 1962a; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. ispirense*** Bernhauer and Peks, 2013b: 328

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'İspir' (Turkey: Erzurum)]

Provincial distribution in Turkey: TR-A: EZ (Bernhauer and Peks 2013b; Danilevsky 2015a)

Range: A: TR

**\**D. ivani*** Pesarini and Sabbadini, 2011: 47

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Kardeşler' (Turkey: Muş)] A: TR

Provincial distribution in Turkey: TR-A: BN MU (Pesarini and Sabbadini 2011; Özdīkmen 2012; Rapuzzi and Sama 2012; Danilevsky 2015a)

Range: A: TR

**\**D. jacovleviellum*** Plavilstshikov, 1951: 120

Type information: Lectotype ♂, ex collection N. N. Plavilstshikov, Zoological Museum, University of Moscow [type locality 'Erzurum env.' (Turkey)]

Provincial distribution in Turkey: TR-A: EZ KAR (Plavilstshikov 1951; Breuning 1962a; Braun 1978a; Danilevsky 2009, 2015a; Löbl and Smetana 2010; Özdīkmen 2010, 2012)

Range: A: TR

**\**D. janatai*** Kadlec, 2006: 11

Type information: Holotype ♂, collection S. Kadlec, Litvinov [type locality 'Karaoğlan Mt.' (Turkey: Tunceli)]

Provincial distribution in Turkey: TR-A: TU (Kadlec 2006; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. javeti*** Kraatz, 1873: 91

Type information: Syntypes ♂♂ and ♀♀, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Syria: Akbez', should be 'Akbez' (Turkey: Hatay)]

*destinoi* Fairmaire, 1884: 173 [Turkey: Hatay: Akbez]

*univestitum* Pic, 1903a: 5 (*D. destinoi* var.) [Turkey: Hatay: Akbez]

*tabense* Pic, 1931a: 1 (*D. aleppense* var.) [Turkey: Gaziantep ('Ain-Tab'), not Syria]

Provincial distribution in Turkey: TR-A: AD GA HT IC OS (Fairmaire 1884; Ganglbauer 1884; Pic 1903a, 1931a; Breuning 1946, 1962a; Fuchs and Breuning 1971; Önalp 1991; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a; Özdīkmen and Kaya 2015b)

Range: A: SY TR

***D. johannisfranci*** Pesarini and Sabbadini, 2007: 40

Type information: Holotype ♂, Museo Civico di Storia Naturale, Milano [type locality 'Tracia: Monastraki' (Greece)]

Provincial distribution in Turkey: TR-E: ED (Pesarini and Sabbadini 2007; Löbl and Smetana 2010; Özdīkmen 2010; Danilevsky 2015a)

Range: E: GR TR



**\**D. karacaorenense*** Bernhauer and Peks, 2012: 211

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'İmrani: Karacaören' (Turkey: Sivas)]

*karacaorenense* Bernhauer and Peks, 2013b: 336 [unjustified emendation]

Provincial distribution in Turkey: TR-A: ER SV (Bernhauer and Peks 2012, 2013b; Danilevsky 2015a)

Range: A: TR

**\**D. karamanense*** Özdi̇kmen and Koçak, 2013: 672

Type information: Holotype ♂, collection Özgür Koçak, Karaman [type locality 'Kılbasan-Karadağ: Halisgömü plateau' (Turkey: Karaman)]

Provincial distribution in Turkey: TR-A: KM (Özdi̇kmen and Koçak 2013; Danilevsky 2015a)

Range: A: TR

**\**D. karsense*** Suvorov, 1915: 118

Type information: Holotype ♂, ex collection G. Suvorov, Zoological Museum, University of Moscow [type locality 'Kağızman' (Turkey: Kars)]

Provincial distribution in Turkey: TR-A: AR KAR (Suvorov 1915; Plavilstshikov 1958; Breuning 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. kartaldagense*** Bernhauer and Peks, 2013a: 314

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Kartaldağ' (Turkey: Çanakkale)]

Provincial distribution in Turkey: TR-A: CA (Bernhauer and Peks 2013a; Danilevsky 2015a)

Range: A: TR

***D. kasikoporanum*** Pic, 1902b: 10

Type information: Holotype ♂, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Kazkoparan' (Turkey: İğdır)]

Provincial distribution in Turkey: TR-A: AR IG (Pic 1902b; Plavilstshikov 1958; Breuning 1962a; Lazarev 2009; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: AR TR

**\**D. kazanciense*** Bernhauer and Peks, 2010: 197

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'N Anamur: N Akpinar, Kazancı' (Turkey: İçel)]

Provincial distribution in Turkey: TR-A: IC (Bernhauer and Peks 2010; Özdi̇kmen 2012; Danilevsky 2015a)

Range: A: TR

**\**D. kindermannii*** Waltl, 1838: 470

Type information: Holotype ♂, ex collection Imre Frivaldszky, Magyar Természettudományi Múzeum, Budapest [type locality 'Turkei' (Turkey)]

*griseum* Waltl, 1838: 470 [Turkey]

*fuscoquadriplagiatum* Breuning, 1946: 110 (*D. kindermannii* m.) [Turkey: İzmir]

*fuscoreductum* Breuning, 1946: 110 (*D. kindermannii* m.) [Turkey: Çanakkale]

Provincial distribution in Turkey: TR-A: AN CA IZ US (Küster 1846a; Ganglbauer 1884; Breuning 1946, 1962a; Demelt 1963; Önalp 1991; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. kizildagense*** Bernhauer and Peks, 2012: 212

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'İmrani̇: Kızıldağ' (Turkey: Sivas)]

Provincial distribution in Turkey: TR-A: ER SV (Bernhauer and Peks 2012, 2013b; Danilevsky 2015a)

Range: A: TR

**\**D. konradi*** Bernhauer and Peks, 2012: 210

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Kuruca pass' (Turkey: Bingöl)]

Provincial distribution in Turkey: TR-A: BN (Bernhauer and Peks 2012, 2013b; Danilevsky 2015a)

Range: A: TR

**\**D. kraetschmeri*** Bernhauer, 1988: 98

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Akseki' (Turkey: Antalya)]

Provincial distribution in Turkey: TR-A: ANT (Bernhauer 1988; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. kurdistanum*** Breuning, 1944: 12

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Diyarbakır' (Turkey)]

*rufulipes* Breuning, 1963: 36 (*D. kurdistanum* m.) [Turkey: Batman: Sason Mts.]

Provincial distribution in Turkey: TR-A: ADY BA BN DI (Breuning 1944, 1962a, 1963; Fuchs and Breuning 1971; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Sama et al. 2012; Danilevsky 2015a)

Range: A: TR

**\**D. kurucanum*** Holzschuh, 2007: 249

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Kuruca pass' (Turkey: Elazığ)]

Provincial distribution in Turkey: TR-A: EL (Holzschuh 2007; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. ladikanum*** Braun, 1976a: 173 [RN]

Type information: Holotype ♂, Muséum National d'Histoire Naturelle, Paris [type locality 'Ladik' (Turkey: Samsun)]

*ladikense* Breuning, 1975: 10 [HN] [Turkey: Samsun: Ladik]

*paraladikense* Breuning, 1977: 60 [RN] [Turkey: Samsun: Ladik]

Provincial distribution in Turkey: TR-A: SM (Breuning 1975, 1977; Braun 1976a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Bernhauer and Peks 2013b; Danilevsky 2015a)

Range: A: TR



**D. laeve** Faldermann, 1837: 278

\***D. laeve micula** Plavilstshikov, 1937: 26 (*D. micula*)

Type information: Lectotype, ex collection Dobrovl. > N. N. Plavilstshikov, Zoological Museum, University of Moscow [type locality 'Oltu' (Turkey: Erzurum)]

Provincial distribution in Turkey: TR-A: EZ (Plavilstshikov 1937; Breuning 1962a; Danilevsky and Murzin 2009; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

Other known subspecies: *D. laeve laeve* Faldermann, 1837: 278 A: IN; *D. laeve hyrcanum* Jakovlev, 1899b: 64 A: IN; *D. laeve vladimiri* Danilevsky and Murzin, 2009: 13 A: AR

\***D. lameeri** Théry, 1896: 109

Type information: Syntypes 2♂♂ and 2♀♀, ex collection A. Théry > M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Kastamonu' (Turkey)]

Provincial distribution in Turkey: TR-A: KS (Théry 1896; Breuning 1946, 1962a; Demelt 1967; Önalp 1991; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\***D. ledouxi** Breuning, 1974b: 65

Type information: Holotype ♀, Muséum National d'Histoire Naturelle, Paris [type locality 'İllica' (Turkey: Erzurum)]

Provincial distribution in Turkey: TR-A: EZ (Breuning 1974b; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\***D. linderi** Tournier, 1872: 285

Type information: Holotype ♀, ex collection H. Tournier > M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Spain' but undoubtedly mislabeled, should be Turkey]

*kollari* Kraatz, 1873: 87 [Turkey: Amasya]

*henrici* Pic, 1905b: 156 [Turkey: Tokat]

*anticepunctatum* Breuning, 1946: 109 (*D. kollari* m.) [Turkey: Tokat]

*quadripunctatum* Breuning, 1946: 109 (*D. kollari* m.) [Turkey: Tokat]

*unipunctum* Breuning, 1946: 109 (*D. kollari* m.) [Turkey: Tokat]

*sublinderi* Breuning, 1966a: 20 (*D. kollari* m.) [Turkey: Samsun]

*dorsoimmaculipenne* Breuning, 1966b: 147 (*D. kollari* m.) [Turkey: Çorum: Boğazkale: Yazılıkaya]

Provincial distribution in Turkey: TR-A: AM CO SM TO (Kraatz 1873; Ganglbauer 1884; Pic 1905b; Breuning 1946, 1962a, 1966a, 1966b; Perissinotto and Luchini 1966; Sama 1982; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**D. lineatocolle** Kraatz, 1873: 57 (*D. femoratum* var.)

Type information: Syntypes ♂♂ and ♀♀, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Salonica' (Greece)]

*atrofemoratum* Pic, 1917b: 7 (*D. lineatocolle* var.) [Macedonia]

*sinuaticolle* Pic, 1927: 6 [Greece: Salonica]

*riveti* Pic, 1927: 162 (*D. lineatocolle* var.) [Macedonia]  
 Provincial distribution in Turkey: TR-E (Önalp 1991; Özdiikmen 2010)  
 Range: E: AL BU GR MC TR YU

\**D. lodosi* Sabbadini and Pesarini, 1992: 29  
*\*D. lodosi abanozense* Bernhauer and Peks, 2010: 199

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Anamur: Abanoz' (Turkey: İçel)]  
 Provincial distribution in Turkey: TR-A: IC (Bernhauer and Peks 2010; Özdiikmen 2012; Danilevsky 2015a)  
 Range: A: TR

\**D. lodosi lodosi* Sabbadini and Pesarini, 1992: 29

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Bucaklışa' (Turkey: Karaman)]  
 Provincial distribution in Turkey: TR-A: KM (Sabbadini and Pesarini 1992; Bernhauer and Peks 2010; Löbl and Smetana 2010; Özdiikmen 2010, 2012; Danilevsky 2015a)  
 Range: A: TR

\**D. lohsei* Braun, 1976b: 254

Type information: Holotype ♂, collection Walter Braun, Tübingen [type locality 'Karaman' (Turkey)]  
 Provincial distribution in Turkey: TR-A: IC KN KM (Braun 1976b, 1978a; Sama 1982; Adlbauer 1992; Löbl and Smetana 2010; Özdiikmen 2010, 2012; Turgut and Özdiikmen 2010; Pesarini and Sabbadini 2013; Danilevsky 2015a)  
 Range: A: TR

\**D. longulum* Breuning, 1943: 89

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Bozdağ' (Turkey: İzmir)]  
 Provincial distribution in Turkey: TR-A: IZ (Breuning 1943, 1962a; Löbl and Smetana 2010; Özdiikmen 2010, 2012; Danilevsky 2015a)  
 Range: A: TR

*D. lugubre* Kraatz, 1873: 41  
*D. lugubre lugubre* Kraatz, 1873: 41

Type information: Lectotype ♂, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Salonica' (Greece)]  
*salonicum* Pic, 1916: 22 [Greece: Salonica]  
*discosulcatum* Pic, 1926: 1 (*D. salonicum* var.) [Macedonia]  
*sparsepilosum* Pic, 1926: 13 (*D. lugubre* var.) [Greece: Salonica]  
*pseudolugubre* Breuning, 1943: 96 [Macedonia: 'Karatschköi' = Dioni]  
 Provincial distribution in Turkey: TR-A: AM SV; TR-E (Önalp 1990; Özdiikmen and Hasbenli 2004a; Özdiikmen 2010)  
 Range: E: ?BU GR MC TR A: TR



Other known subspecies: *D. lugubre meteorum* Breuning, 1969: 42 E: GR; *D. lugubre minkovae* Heyrovský, 1962: 225 E: BU; *D. lugubre parinfernale* Breuning, 1975: 10 E: GR; *D. lugubre pelionense* Breit, 1923: 145 E: GR; *D. lugubre thessalicum* Pic, 1916: 22 E: GR

**\**D. maceki*** Holzschuh, 1995: 41

Type information: Holotype ♂, collection J. Macek [type locality 'Abant' (Turkey: Bolu)]  
 Provincial distribution in Turkey: TR-A: BO (Breuning 1943, 1962a; Holzschuh 1995; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. margheritae*** Breuning, 1964b: 32

Type information: Holotype ♂, collection Dr. Franco Tassi, Rome [type locality 'Epire (Ioánnina): Pinde Mts.' (Greece)]

*katarense* Breuning, 1969: 42 (*D. margheritae* m.) [Greece: Katara]

Provincial distribution in Turkey: TR-A: CA; TR-E (Adlbauer 1988; Althoff and Danilevsky 1997; Özdi̇kmen 2010; Danilevsky 2015a)

Range: E: GR TR A: TR

**\**D. martini*** Bernhauer, 1988: 100

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Nallıhan' (Turkey: Ankara)]

Provincial distribution in Turkey: TR-A: AN (Bernhauer 1988; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. menradi*** Holzschuh, 1989: 172

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Afşin' (Turkey: Kahramanmaraş)]

Provincial distribution in Turkey: TR-A: KA MA (Holzschuh 1989; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

**\**D. merkli*** Ganglbauer, 1884: 506

Type information: Syntypes ♂ and ♀, ex collection L. Ganglbauer, Naturhistorisches Museum Wien [type locality 'nordwestlichen Kleinasiens: Akdağ (?Simav)' (Turkey: ?Kütahya)]

*humerointerruptum* Breuning, 1946: 109 (*D. merkli* m.) [Turkey: ?Kütahya: ?Simav: Akdağ]

Provincial distribution in Turkey: TR-A: KU (Ganglbauer 1884; Breuning 1946, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. mesopotamicum*** Breuning, 1944: 12

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Mesopotamia: Ras Al-Ain' (Iraq)]

*submarinense* Breuning, 1946: 109 (*D. mesopotamicum* m.) [Iraq: Mesopotamia: Ras Al-Ain]

Provincial distribution in Turkey: TR-A: MR SU (Breuning 1962a; Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a)

Range: A: IQ TR

\**D. micans* J. Thomson, 1867: 61

\**D. micans micans* J. Thomson, 1867: 61

Type information: Holotype ♂, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Armenien' undoubtedly mislabeled, should be Turkey]

*macropus* Kraatz, 1873: 99 [Turkey: Amasya]

*sericatum* Kraatz, 1873: 98 [?Caucasus, Turkey: Ankara, Bursa]

*obscurans* Pic, 1892a: 91 (*D. macropus* var.) [Turkey: Amasya]

*amasinum* Pic, 1898: 58 [Turkey: Amasya]

*atipes* Reitter, 1900: 88 (*D. sericatum* var.) [Turkey: Amasya]

*tokatense* Pic, 1901: 12 (*D. impressicolle* var.) [Turkey: Tokat]

*corallicorne* Pic, 1904: 4 (*D. sericatum* var.) [Turkey: Ankara]

*subreductum* Pic, 1942a: 2 (*D. sericatum* var.) [Turkey]

*subobesum* Pic, 1942a: 2 [Turkey: Amasya]

Provincial distribution in Turkey: TR-A: AM AN BS CA CO IZ TO YO (Kraatz 1873; Pic 1892a, 1898, 1901, 1904, 1942a; Reitter 1900; Breuning 1962a; Demelt 1963; Perissinotto and Luchini 1966; Breuning and Villiers 1967; Fuchs and Breuning 1971; Gül-Zümreoglu 1975; Holzschuh 1980; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a; Özdkmen and Kaya 2015d)

Range: A: TR

\**D. micans susheriense* Breuning, 1970: 97 (*D. sinopense* ssp.)

Type information: Holotype ♂, Muséum National d'Histoire Naturelle, Paris [type locality 'Suşehri' (Turkey: Sivas)]

Provincial distribution in Turkey: TR-A: SV (Breuning 1970; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a; Özdkmen and Kaya 2015d)

Range: A: TR

*D. mniszechi* Kraatz, 1873: 39

*D. mniszechi mniszechi* Kraatz, 1873: 39

Type information: Syntypes ♂ and ♀, ex Musaeo Mniszech > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Caucasus']

*rugulosum* Breuning, 1946: 96 (*D. mniszechi* m.) [Turkey: İğdir: Kazkoparan]

Provincial distribution in Turkey: TR-A: AG EZ IG KAR SV (Breuning 1946; Plavilstshikov 1958; Özdkmen and Hasbenli 2004a; Löbl and Smetana 2010; Özdkmen 2010; Lazarev 2014; Danilevsky 2015a)

Range: A: AR TR

*D. mniszechi cavernosum* Lazarev, 2014: 701

Type information: Holotype ♂, ex collection M. L. Danilevsky, Collection of Zoological Institute of the Russian Academy of Sciences, Saint-Petersbourg [type locality Arteni (Armenia)]

Provincial distribution in Turkey: TR-A: ART IG KAR (Breuning 1946; Lazarev 2014; Danilevsky 2015a)



Range: A: AR TR

Other known subspecies: *D. mniszechi georgianum* Lazarev, 2014: 701 A: GG

**\**D. muchei* Breuning, 1962b: 38**

Type information: Holotype ♂, collection H. Muche [type locality 'Çankırı' (Turkey)]

*plurivittipenne* Breuning, 1962b: 39 (*D. muchei* m.) [Turkey: Çankırı]

*parescherichi* Breuning, 1966b: 146 [Turkey: Çorum: Boğazkale]

Provincial distribution in Turkey: TR-A: CN CO KS KY YO (Breuning 1962a, 1962b, 1966b;

Perissinotto and Luchini 1966; Demelt 1967; Braun 1978a, 1979; Holzschuh 1980;

Özdikmen 2010, 2012; Özdikmen and Kaya 2015a)

Range: A: TR

**\**D. multimaculatum* Pic, 1932: 22**

Type information: Holotype, ex collection Leo Heyrovský, Musée National de Prague [type locality 'Ödemiş' (Turkey: İzmir)]

*postbiplagiatum* Breuning, 1946: 107 (*D. multimaculatum* m.) [Turkey: İzmir: Ödemiş]

Provincial distribution in Turkey: TR-A: IZ (Pic 1932; Breuning 1946, 1962a; Löbl and Smetana 2010; Özdikmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. murzini* Danilevsky, 2015a: 221**

Type information: Holotype ♂, ex collection M. L. Danilevsky, Collection of Zoological Museum, University of Moscow [type locality 'İspir: Yukarıözbağ' (Turkey: Erzurum)]

Provincial distribution in Turkey: TR-A: EZ (Danilevsky, 2015b)

Range: A: TR

**\**D. musense* Bernhauer and Peks, 2014a: 109**

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'S Bulanık' (Turkey: Muş)]

Provincial distribution in Turkey: TR-A: MU (Bernhauer and Peks 2014a; Danilevsky 2015a)

Range: A: TR

**\**D. narlianum* Özdikmen, Mercan and Cihan, 2012b: 550**

Type information: Holotype ♂, collection H. Özdikmen, Zoological Museum of Gazi University, Ankara [type locality 'Narlı' (Turkey: Kahramanmaraş)]

Provincial distribution in Turkey: TR-A: KA (Özdikmen 2012; Özdikmen et al. 2012b, 2012c; Danilevsky 2015a)

Range: A: TR

**\**D. nigrostriatum* Adlbauer, 1982: 104**

Type information: Holotype ♂, collection Karl Adlbauer, Graz [type locality 'N Tunceli' (Turkey)]

Provincial distribution in Turkey: TR-A: TU (Adlbauer 1982, 1988; Löbl and Smetana 2010; Özdikmen 2010, 2012; Sama et al. 2012; Danilevsky 2015a)

Range: A: TR

**\**D. nihalae* Rapuzzi and Sama, 2012: 671**

Type information: Holotype ♂, collection Gianfranco Sama, Cesena [type locality 'Zara' (Turkey: Sivas)]

Provincial distribution in Turkey: TR-A: SV (Özdikmen 2012; Rapuzzi and Sama 2012; Danilevsky 2015a)

Range: A: TR

**\**D. niksarensis* Bernhauer and Peks, 2013b: 327**

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Niksar' (Turkey: Tokat)]

Provincial distribution in Turkey: TR-A: TO (Bernhauer and Peks 2013b; Danilevsky 2015a)

Range: A: TR

***D. nitidum* Motschulsky, 1838: 185**

Type information: Holotype, ex collection V. I. Motschulsky, Zoological Museum, University of Moscow [type locality 'Armenia']

*suturatum* Ferrari, 1864: 481 [Georgia: Achalzik]

Provincial distribution in Turkey: TR-A: ART EZ KAR YO (Plavilstshikov 1958; Breuning 1962a; Özbek 1978; Adlbauer 1988; Önalp 1991; Özdikmen and Hasbenli 2004a; Löbl and Smetana 2010; Özdikmen 2010; Danilevsky 2015a)

Range: A: AB AR GG TR

**\**D. nobile* Hampe, 1852: 313**

Type information: Syntypes, ex collection Moritz Wagner, Museum für Naturkunde der Humboldt-Universität zu Berlin [type locality 'W Persien', undoubtedly mislabeled, should be 'Erzurum env.' (Turkey)]

*maculatum* Breuning, 1946: 119 (*D. nobile* m.) [Turkey: Kurdistan]

Provincial distribution in Turkey: TR-A: BT EZ VA (Breuning 1946, 1962a; Plavilstshikov 1958; Villiers 1967; Braun 1978a; Sama 1982; Önalp 1991; Özdikmen and Hasbenli 2004a; Löbl and Smetana 2010; Özdikmen 2010, 2012; Pesarini and Sabbadini 2011; Danilevsky 2015a)

Range: A: TR

***D. nodicorne* Tournier, 1872: 287**

Type information: Syntypes 2♂♂ and ♀, ex collection H. Tournier > M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Taurus' (Turkey)]

*maculereductum* Breuning, 1962a: 532 (*D. quadrimaculatum* m.) [Greece: Lesbos Is.]

*subnodicorne* Breuning, 1962a: 532 (*D. quadrimaculatum* m.) [Greece: Lesbos Is.]

Provincial distribution in Turkey: TR-A: CA IZ Taurus Mts.; TR-E (Breuning 1962a; Krätschmer 1985; Önalp 1990; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdikmen 2010, 2012; Danilevsky 2015a)

Range: E: GR TR A: TR

***D. obsoletum* Kraatz, 1873: 78 (*D. graecum* var.)**

Type information: Holotype ♂, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Bursa' (Turkey), but not Bursa, according to Kraatz with İstanbul (Turkey) and Greece]

*obsoletoides* Breuning, 1946: 111 (*D. obsoletum* m.) [Greece: Thrace: Ferre]

Provincial distribution in Turkey: TR-A: ?BS IS; TR-E: ED IS KK TE (Kraatz 1873; Bodemeyer 1906; Fuchs and Breuning 1971; Braun 1978a; Althoff and Danilevsky 1997; Özdikmen 2010, 2012; Danilevsky 2015a; Özdikmen and Kaya 2015c)

Range: E: BU GR TR A: TR



**\**D. obtusum*** Breuning, 1944: 13

**\**D. obtusum marashense*** Breuning, 1948: 59

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Kahramanmaraş' (Turkey)]

*dorsointerruptum* Breuning, 1948: 60 (*D. obtusum* m.) [Turkey: Kahramanmaraş]

Provincial distribution in Turkey: TR-A: KA (Breuning 1948, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. obtusum obtusum*** Breuning, 1944: 13

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Anatolia' (Turkey)]

Provincial distribution in Turkey: TR-A: ADY DI HA VA (Breuning 1944, 1962a; Fuchs and Breuning 1971; Braun 1978a; Rezek and Hoskovec 1999; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. oezdurali*** Önalp, 1988: 361

Type information: Holotype ♂, collection Birsen Önalp [type locality 'Ahır Mt.' (Turkey: Kahramanmaraş)]

Provincial distribution in Turkey: TR-A: ADY KA (Önalp 1988, 1991; Rezek and Hoskovec 1999; Pesarini and Sabbadini 1998, 2013; Özdi̇kmen and Okutaner 2006; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Sama et al. 2012; Özdi̇kmen and Koçak 2013; Danilevsky 2015a)

Range: A: TR

**\**D. olympicum*** Ganglbauer, 1882: 228

**\**D. olympicum convexum*** Breuning, 1943: 90

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Achu Dagh (Simav: Akdağ)' (Turkey: Kütahya)]

Provincial distribution in Turkey: TR-A: KU (Breuning 1943, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a; Özdi̇kmen and Kaya 2015c)

Range: A: TR

**\**D. olympicum flavosuturale*** Krätschmer, 1987: 340

Type information: Holotype ♂, collection Otto Erich Krätschmer, München [type locality 'Mustafakemalpaşa: Gölyazı: Uluabat Lake' (Turkey: Bursa)]

Provincial distribution in Turkey: TR-A: BS (Krätschmer 1987; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a; Özdi̇kmen and Kaya 2015c)

Range: A: TR

**\**D. olympicum olympicum*** Ganglbauer, 1882: 228

Type information: Lectotype ♂, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Bursa' (Turkey)]

*subalpinum* Kraatz, 1873: 78 [Turkey: Bursa: Osmangazi: Uludağ]

*oreophilum* Ganglbauer, 1884: 500 (*D. olympicum* var.) [Turkey: Bursa: Osmangazi: Uludağ]

*brussense* Breuning, 1946: 112 (*D. olympicum* m.) [Turkey: Bursa]  
*quinquefasciatum* Breuning, 1946: 112 (*D. olympicum* m.) [Turkey: Bilecik]  
 Provincial distribution in Turkey: TR-A: AN BI BS (Kraatz 1873; Ganglbauer 1882, 1884;  
 Breuning 1946, 1962a; Demelt 1963; Braun 1978a; Adlbauer 1988; Önalp 1991; Sama  
 1999; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a; Özdi̇kmen and  
 Kaya 2015c)  
 Range: A: TR

**\**D. ortrudae* Braun, 1978b: 185**

Type information: Holotype ♂, collection Walter Braun, Tübingen [type locality 'Göksun'  
 (Turkey: Kahramanmaraş)]

Provincial distribution in Turkey: TR-A: KA (Braun 1978b; Adlbauer 1988; Löbl and  
 Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. paracinerarium* Breuning, 1974a: 149**

Type information: Holotype ♂, Collection Carolus Holzschuh, Villach [type locality 'Mut:  
 Sertavul pass' (Turkey: İçel)]

Provincial distribution in Turkey: TR-A: IC KM SU (Breuning 1962a, 1974a; Gül-Zümreoglu  
 1972; Braun 1978a, 1979; Özdi̇kmen 2012; Özdi̇kmen and Kaya 2015d)

Range: A: TR

**\**D. pararufipenne* Braun, 1976b: 257**

**\**D. pararufipenne pararufipenne* Braun, 1976b: 257**

Type information: Holotype ♂, collection Walter Braun, Tübingen [type locality 'Abant'  
 (Turkey: Bolu)]

Provincial distribution in Turkey: TR-A: BO (Braun 1976b, 1978a; Özdi̇kmen 2006, 2010,  
 2012; Özdi̇kmen and Demir 2006; Löbl and Smetana 2010; Bernhauer and Peks 2011b;  
 Danilevsky 2015a)

Range: A: TR

**\**D. pararufipenne rassei* Braun, 1976b: 261**

Type information: Holotype ♂, collection Walter Braun, Tübingen [type locality 'Gerede'  
 (Turkey: Bolu)]

Provincial distribution in Turkey: TR-A: AN BO (Braun 1976b, 1978a; Adlbauer 1992; Löbl  
 and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. parilis* Pesarini and Sabbadini, 2013: 52**

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano  
 [type locality 'Oluközü' (Turkey: Yozgat)]

Provincial distribution in Turkey: TR-A: YO (Pesarini and Sabbadini 2013; Danilevsky  
 2015a)

Range: A: TR

**\**D. pavesii* Pesarini and Sabbadini, 1998: 57**

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano  
 [type locality 'Yüksekova' (Turkey: Hakkari)]



Provincial distribution in Turkey: TR-A: HA (Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. pedestre*** Poda von Neuhaus, 1761: 34 (*Cerambyx*)

***D. pedestre pedestre*** Poda von Neuhaus, 1761: 34 (*Cerambyx*)

Type information: Syntypes, ex collection N. Poda [type locality 'ad Graecium' (Greece)]

Provincial distribution in Turkey: TR-E: IS (Küster 1846a; Önalp 1990; Özdkmen 2010)

Range: E: AL AU BH BU CR CZ HU MC MD PL RO SK SL TR UK YU

Other known subspecies: *D. pedestre kaszabi* Breuning, 1956: 404 E: HU

**\**D. peksi*** Bernhauer, 2010: 190

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Anamur' (Turkey: İçel)]

Provincial distribution in Turkey: TR-A: IC (Bernhauer 2010; Özdkmen 2012; Danilevsky 2015a)

Range: A: TR

**\**D. pergamenum*** Pesarini and Sabbadini, 2009: 22

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Pergama (Bergama)' (Turkey: İzmir)]

Provincial distribution in Turkey: TR-A: IZ (Pesarini and Sabbadini 2009; Özdkmen 2012; Danilevsky 2015a)

Range: A: TR

**\**D. petrovitzi*** Heyrovský, 1964: 97

Type information: Holotype ♂, ex collection Carl von Demelt, Staatliches Museum für Naturkunde, Stuttgart [type locality 'Çamlıayla (Namrun)' (Turkey: İçel)]

Provincial distribution in Turkey: TR-A: IC (Heyrovský 1964; Demelt 1967; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. pilosipenne*** Breuning, 1943: 90

**\**D. pilosipenne kazdagense*** Özdkmen in Küçükaykı et al., 2013: 56

Type information: Holotype ♂, Department of Biology at Eskişehir Osmangazi University, Eskişehir [type locality 'Edremit, Yayla' (Turkey: Balıkesir)]

Provincial distribution in Turkey: TR-A: BL (Özdikmen in Küçükaykı et al. 2013; Danilevsky 2015a)

Range: A: TR

Other known subspecies: *D. pilosipenne pilosipenne* Breuning, 1943: 90 E: GR

**\**D. piocardi*** Kraatz, 1873: 85

Type information: Syntypes ♂ and ♀, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Samsun-Amasya: Akdağ' (Turkey)]

*quadripustulatum* Ganglbauer, 1884: 457 [Turkey: Çanakkale]

*elongatum* Pic, 1898: 58 (*D. piocardi* var.) [Turkey: Taurus Mt.]

*pelops* Jakovlev, 1901: 83 [Turkey: Amasya]

*theophilei* Pic, 1901: 32 [HN] (*D. piochardi* var.) [Turkey]  
*roberti* Pic, 1905c: 11 (*D. piochardi* var.) ['Syria', should be Turkey]  
*mauricei* Plavilstshikov, 1927: 51 [RN]  
*varievittatum* Pic, 1942a: 1 [S Russia, undoubtedly mislabeled]  
*apicedisparatum* Breuning, 1946: 109 (*D. piochardi* m.) [Turkey: Amasya]  
*ladikense* Breuning and Villiers, 1967: 60 (*Pedestredorcadion piochardi* var.) [Turkey: Samsun: Ladik]

Provincial distribution in Turkey: TR-A: AM CA CO SM SN YO (Kraatz 1873; Ganglbauer 1884; Jakovlev 1901; Breuning 1962a; Breuning and Villiers 1967; Braun 1978a, 1979; Önalp 1991; Adlbauer 1992; Löbl and Smetana 2010; Özdkmen 2010, 2012; Lazarev 2011; Danilevsky 2015a)

Range: A: TR

\****D. pittinorum*** Pesarini and Sabbadini, 1998: 48

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Çorum' (Turkey)]

Provincial distribution in Turkey: TR-A: CO (Pesarini and Sabbadini 1998, 2013; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\****D. pluto*** J. Thomson, 1867: 47

Type information: Holotype ♂, ex collection J. Thomson > R. Oberthür, 1952, Muséum National d'Histoire Naturelle, Paris [type locality 'Siberia' but undoubtedly mislabeled, should be Turkey]

Provincial distribution in Turkey: TR-A: AM ES EZ SA SM (Ganglbauer 1884; Breuning and Villiers 1967; Braun 1978a; Önalp 1990; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\****D. poleti*** Breuning, 1948: 59

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Caria: Babadağ' (Turkey: Denizli)]

Provincial distribution in Turkey: TR-A: DE (Breuning 1948, 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\****D. postalbosuturale*** Özdkmen and Koçak, 2015: 439

Type information: Holotype ♂, collection Ö. Koçak, Karaman [type locality 'Elmaşehir district' (Turkey: Karaman)]

Provincial distribution in Turkey: TR-A: KM (Özdkmen and Koçak 2015)

Range: A: TR

\****D. praetermissum*** Pesarini and Sabbadini, 1998: 47

\****D. praetermissum mikhaili*** Özdkmen, 2010: 457

Type information: Holotype ♂, Entomology Department of Eastern Mediterranean Forestry Research Institute, İçel [type locality 'Abant' (Turkey: Bolu)]

Provincial distribution in Turkey: TR-A: BO (Özdkmen 2010, 2011, 2012; Danilevsky 2015a)

Range: A: TR



**\**D. praetermissum praetermissum*** Pesarini and Sabbadini, 1998: 47

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Uludag' (Turkey: Bursa)]

Provincial distribution in Turkey: TR-A: BS (Pesarini and Sabbadini 1998, 2013; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. preissi*** Heyden, 1894: 86

Type information: Syntypes 2 ♂ and ♀, ex collection L. von Heyden, Natur-Museum und Forschungs-Institut Senckenberg, Frankfurt am Main [type locality 'Kastamonu' (Turkey)]

Provincial distribution in Turkey: TR-A: KS (Heyden 1894; Breuning 1962a; Braun 1978a; Önalp 1991; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Bernhauer and Peks 2011b; Danilevsky 2015a)

Range: A: TR

**\**D. pseudarcivagum*** Breuning, 1943: 91

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Amasya' (Turkey)]

*brevidorsale* Breuning, 1946: 109 (*D. pseudarcivagum* m.) [Turkey: Samsun]

Provincial distribution in Turkey: TR-A: AM SM (Breuning 1943, 1946, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. pseudoinfernale*** Breuning, 1943: 94

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Amanos Mts.' (Turkey: Hatay)]

*pseudimmutatum* Breuning, 1948: 61 (*D. pseudoinfernale* m.) [Turkey: Hatay: Amanos Mts.]

Provincial distribution in Turkey: TR-A: HT (Breuning 1943, 1948, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. pseudobithyniense*** Breuning, 1962a: 465

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'W Balıkesir' (Turkey)]

Provincial distribution in Turkey: TR-A: BL CA (Breuning 1962a; Braun 1978a; Adlbauer 1988; Pesarini and Sabbadini 2009; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. pseudocinctellum*** Breuning, 1943: 90

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Akşehir' (Turkey: Konya)]

*fuscoplagiatum* Breuning, 1946: 101 (*D. pseudocinctellum* m.) [Turkey: Konya: Akşehir]

*fuscoconjunctum* Breuning, 1946: 101 (*D. pseudocinctellum* m.) [Turkey: Konya: Akşehir]

Provincial distribution in Turkey: TR-A: KN US (Breuning 1943, 1946, 1962a; Demelt 1963; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Lingafelter et al. 2014; Danilevsky 2015a)

Range: A: TR

**\**D. pseudoholosericeum* Breuning, 1962a: 300**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Sinop' (Turkey)]

Provincial distribution in Turkey: TR-A: SN (Breuning 1962a; Braun 1978a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. pseudopreissi* Breuning, 1962a: 411**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'S Sinop' (Turkey)]

Provincial distribution in Turkey: TR-A: AM SN (Breuning 1962a; Adlbauer 1992; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. punctipenne* Küster, 1852: 94**

Type information: Holotype, ex collection Heinrich Carl Küster > Johann Menzel, Naturhistorisches Museum, Nürnberg [type locality 'Kleinasien' (Turkey)]

*distinguendum* Pic, 1931c: 10 (*D. olympicum* var.) [Greece]

Provincial distribution in Turkey: TR-A: IS; TR-E: IS (Küster 1852; Breuning 1962a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a; Özdīkmen and Kaya 2015e)

Range: E: GR TR A: TR

**\**D. punctulatum* Pesarini and Sabbadini, 2013: 57**

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Tepeköy' (Turkey: Manisa)]

Provincial distribution in Turkey: TR-A: MN (Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

**\**D. punctulicolle* Breuning, 1944: 12**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Diyarbakır' (Turkey)]

Provincial distribution in Turkey: TR-A: DI (Breuning 1944, 1962a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. pygmaeum* Breuning, 1947: 168**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Anatolia' (Turkey)]

*presuturesignatum* Breuning, 1947: 169 (*D. pygmaeum* m.) [Turkey]

Provincial distribution in Turkey: TR-A: Kurdistan (?DI) (Breuning 1947, 1962a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. quadripustulatum* Kraatz, 1873: 88**

Type information: Holotype ♀, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Dardanellen' (Turkey: Çanakkale)]

*tenuefasciatum* Breuning, 1948: 60 (*D. quadripustulatum* m.) [Turkey: Çanakkale]



Provincial distribution in Turkey: TR-A: CA (Kraatz 1873; Breuning 1948, 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**D. regulare** Pic, 1931b: 6 (*D. gallipolitanum* var.)

**D. regulare regulare** Pic, 1931b: 6 (*D. gallipolitanum* var.)

Type information: Holotype, ex collection Dr. Buresch > M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Keşan' (Turkey: Edirne)]

*angustesuturale* Breuning, 1946: 111 (*D. regulare* m.) [Greece: Thrace: Mandra]

Provincial distribution in Turkey: TR-E: ED KK (Pic 1931b; Breuning 1962a; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: E: GR TR

Other known subspecies: *D. regulare dramaticum* Pesarini and Sabbadini, 2010: 190 E: GR

**\*D. resadiyeense** Bernhauer and Peks, 2011a: 219

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Reşadiye' (Turkey: Malatya)]

Provincial distribution in Turkey: TR-A: MA (Bernhauer and Peks 2011a, 2013b; Danilevsky 2015a)

Range: A: TR

**\*D. rigattii** Breuning, 1966b: 145

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Zara' (Turkey: Sivas)]

*atracampliatum* Breuning, 1966b: 145 (*D. rigattii* m.) [Turkey: Sivas: Zara]

Provincial distribution in Turkey: TR-A: AM CO SV (Perissinotto and Luchini 1966; Breuning 1966b; Braun 1978a; Adlbauer 1992; Pesarini and Sabbadini 1994, 2013; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\*D. rizeanum** Breuning and Villiers, 1967: 60 (*Pedestredorcadion*)

Type information: Holotype ♂, The Natural History Museum, London [type locality 'Sivrikaya' (Turkey: Rize)]

*ikizderense* Breuning, 1973: 54 [Turkey: Rize: İkizdere]

Provincial distribution in Turkey: TR-A: RI (Breuning and Villiers 1967; Breuning 1973; Holzschuh 1977; Braun 1978a; Özdkmen 2007, 2010, 2012; Löbl and Smetana 2010; Danilevsky 2015a)

Range: A: TR

**\*D. robustum** Ganglbauer, 1884: 500

Type information: Holotype ♂, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Amasya' (Turkey)]

Provincial distribution in Turkey: TR-A: AM (Ganglbauer 1884; Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. rolandmenradi* Peks, 1992:** 197

Type information: Holotype ♂, collection Heinz Peks, Schwanfeld [type locality 'Ballısaray' (Turkey: Bursa)]

Provincial distribution in Turkey: TR-A: BS KU (Peks 1992; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. rosti* Pic, 1900c:** 82

Type information: Holotype, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Caucasus']

*kalinowskyi* König, 1901: 10 [Georgia: Sekarpasse]

Provincial distribution in Turkey: TR-A: DI (Plavilstshikov 1958; Özdīkmen and Hasbenli 2004a; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Danilevsky 2015a)

Range: A: GG TR

**\**D. rufoapicipenne* Breuning, 1946:** 125

Type information: Holotype [type locality 'Bayburt' (Turkey)]

*rufoapicale* Breuning, 1943: 91 [HN] [Turkey: Bayburt]

*rufobrunneotomentosum* Breuning, 1946: 125 (*D. rufoapicenne* m.) [Turkey: Bayburt]

Provincial distribution in Turkey: TR-A: AN BY DU GI RI SV (Breuning 1943, 1946, 1962a; Braun 1978a, 1979; Sama 1982; Özdīkmen and Demir 2006; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Özdīkmen et al. 2012d; Danilevsky 2015a)

Range: A: TR

**\**D. sabanoezueense* Bernhauer and Peks, 2013b:** 326

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Şabanözü' (Turkey: Çankırı)]

Provincial distribution in Turkey: TR-A: CN (Bernhauer and Peks 2013b; Danilevsky 2015a)  
Range: A: TR

**\**D. sarkislaense* Bernhauer and Peks, 2014a:** 105

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'S Şarkışla' (Turkey: Sivas)]

Provincial distribution in Turkey: TR-A: SV (Bernhauer and Peks 2014a; Danilevsky 2015a)  
Range: A: TR

***D. saulcyi* J. Thomson, 1865:** 549

Type information: Holotype ♂, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Syria']

*simile* Kraatz, 1884: 234 [Turkey: Malatya]

*fenestratum* Pic, 1895b: cclxxxiv [Turkey: Gaziantep ('Ain-Tab'), not Syria]

*adanense* Pic, 1904: 4 (*D. destinoi* var.) [Turkey: Adana]

*bilatevittatum* Pic, 1927: 167 (*D. saulcyi* var.) [Syria: Aleppo]

*subelongatum* Pic, 1931a: 1 (*D. destinoi* var.) [Turkey: Hatay: Akbez]

Provincial distribution in Turkey: TR-A: AD GA HT IC MA OS (Kraatz 1884; Ganglbauer 1886; Pic 1895b, 1904, 1931a; Breuning 1962a; Demelt 1963; Fuchs and Breuning 1971; Braun 1978a; Adlbauer 1992; Özdīkmen et al. 2005, 2010; Löbl and Smetana 2010; Özdīkmen 2010, 2012; Cihan et al. 2013; Danilevsky 2015a; Özdīkmen and Kaya 2015b)

Range: A: SY TR



***D. scabricolle* Dalman, 1817: 174 (*Lamia*)**

\****D. scabricolle balikesirensis* Breuning, 1962a: 460**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Balıkesir' (Turkey)]

Provincial distribution in Turkey: TR-A: BL (Breuning 1962a; Braun 1978a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\****D. scabricolle caramanicum* K. Daniel and J. Daniel, 1903: 332**

Type information: Syntypes ♂ and ♀, ex collection K. Daniel, Deutsches Entomologisches Institut, Eberswalde [type locality 'Bulghar-Maaden und Bulghar-Magara (Bolkar Mts.-Çamardı)' (Turkey: Niğde)]

*bulghardaghense* Breuning, 1946: 121 (*D. scabricolle* m.) ['Bulghar-Maaden' (Turkey: Bolkar Mts.: Niğde: Çamardı)]

Provincial distribution in Turkey: TR-A: NI (Bodemeyer 1900; Daniel and Daniel 1903; Breuning 1946, 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\****D. scabricolle crassofasciatum* Özdi̇kmen in Küçükaykı et al., 2013: 56**

Type information: Holotype ♂, Department of Biology at Eskişehir Osmangazi University, Eskişehir [type locality 'Edremit: Sarıkız hill' (Turkey: Balıkesir)]

Provincial distribution in Turkey: TR-A: BL (Özdi̇kmen in Küçükaykı et al. 2013; Danilevsky 2015a)

Range: A: TR

\****D. scabricolle paphlagonicum* Breuning, 1962a: 459**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Kastamonu' (Turkey)]

Provincial distribution in Turkey: TR-A: CN KS (Breuning 1962a; Braun 1978a; Önalp 1991; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Al-Hamadani and Özdi̇kmen 2014; Danilevsky 2015a)

Range: A: TR

***D. scabricolle scabricolle* Dalman, 1817: 174 (*Lamia*)**

Type information: Holotype, ex Musaeo Schönherr, Naturhistoriska Riksmuseet, Stockholm [type locality 'Georgia']

*modestum* Tournier, 1872: 338 [Azerbaijan: Annenfeld]

*lutescens* Kraatz, 1873: 50 [Caucasus]

*aeruginosum* Breuning, 1946: 121 (*D. scabricolle* m.) [Turkey: Konya: Akşehir]

Provincial distribution in Turkey: TR-A: AD AF AG AN ANT AR BI CO ER EZ GU IC IP KA KAR KN KU KY MA SV US VA YO (Heyden 1888; Bodemeyer 1900; Breuning 1946, 1962a; Demelt 1963; Perissinotto and Luchini 1966; Breuning and Villiers 1967; Fuchs and Breuning 1971; Gfeller 1972; Tuatay et al. 1972; Özbek 1978; Braun 1978a; Sama 1982; Adlbauer 1988; Önalp 1991; Özdi̇kmen and Hasbenli 2004a; Özdi̇kmen et al. 2005, 2009, 2010; Özdi̇kmen and Demir 2006; Özdi̇kmen and Okutaner 2006; Löbl and Smetana 2010; Özdi̇kmen 2010; Turgut and Özdi̇kmen 2010; Okutaner et al. 2011; Sama et al. 2012; Cihan et al. 2013; Şenyüz and Özdi̇kmen 2013; Danilevsky 2015a)

Range: A: AB AR GG TR

**\**D. scabricolle uludaghicum*** Breuning, 1970: 98

Type information: Holotype ♂, Muséum National d'Histoire Naturelle, Paris [type locality 'Uludağ' (Turkey: Bursa)]

Provincial distribution in Turkey: TR-A: BS (Breuning and Villiers 1967; Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

Other known subspecies: *D. scabricolle corpulentum* Ménétriés, 1832: 226 A: AB IN; *D. scabricolle gilanense* Lazarev, 2013: 227 A: IN; *D. scabricolle elisabethpolicum* Suvorov, 1915: 119 A: AB; *D. scabricolle nakhiczevanum* Danilevsky, 1999: 28 A: AB; *D. scabricolle paiz* Danilevsky, 1999: 28 A: AB; *D. scabricolle sagezense* Lazarev, 2013: 227 A: IN; *D. scabricolle skoupyi* Lazarev, 2013: 226 A: IN; *D. scabricolle sevangense* Reitter, 1889: 41 A: AR; *D. scabricolle subcorpulentum* Breuning, 1946: 121 A: IN

**\**D. schultzei*** Heyden, 1894: 86

Type information: Syntypes 2 ♂♂ and ♀, ex collection L. von Heyden, Natur-Museum und Forschungs-Institut Senckenberg, Frankfurt am Main [type locality 'Mardin' (Turkey)]

Provincial distribution in Turkey: TR-A: MR (Heyden 1894; Breuning 1962a; Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. scrobicolle*** Kraatz, 1873: 97

**\**D. scrobicolle morulum*** Holzschuh, 1995: 42

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Gökdere' (Turkey: Tokat)]

*fuscosubrevestitum* Breuning, 1962a: 372 (*D. scrobicolle* m.) [Turkey: Tokat: Niksar]

Provincial distribution in Turkey: TR-A: TO (Breuning 1962a; Fuchs and Breuning 1971; Adlbauer 1992; Holzschuh 1995; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. scrobicolle scrobicolle*** Kraatz, 1873: 97

Type information: Syntypes ♂ and ♀, ex Musaeo Mniszech > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Kleinasiens' (Turkey)]

Provincial distribution in Turkey: TR-A: AM YO (Kraatz 1873; Ganglbauer 1884; Breuning 1962a; Fuchs and Breuning 1971; Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. sebenense*** Bernhauer and Peks, 2011b: 202

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Nallıhan' (Turkey: Ankara)]

Provincial distribution in Turkey: TR-A: AN BO (Bernhauer and Peks 2011b; Danilevsky 2015a)

Range: A: TR



\**D. semibrunneum* Pic, 1903b: 170 (*D. mniszechi* var.)

\**D. semibrunneum mediocreimpressum* Pic, 1909a: 123 (*D. mniszechi* var.)

Type information: Holotype ♀ [type locality 'Anamas Gbg. (Dedegöl Mt.)' (Turkey: Isparta)]

*medioimpressum* Pic, 1910: 5 (*D. semibrunneum* var.) [wrong posterior spelling – unavailable]

*anamasum* Pic, 1934: 33 (*D. mniszechi* ssp.) [Turkey: Isparta: Dedegöl Mt.]

Provincial distribution in Turkey: TR-A: AF IP (Pic 1909a, 1934; Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Pesarini and Sabbadini 2013; Lazarev 2014; Danilevsky 2015a)

Range: A: TR

\**D. semibrunneum notatum* Pesarini and Sabbadini, 2013: 62

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Bala' (Turkey: Ankara)]

Provincial distribution in Turkey: TR-A: AN (Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

\**D. semibrunneum semibrunneum* Pic, 1903b: 170 (*D. mniszechi* var.)

Type information: Syntype, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Bozdağ' (Turkey: İzmir)]

*boszdaghana* Breuning, 1948: 61 (*D. mniszechi* ssp. *semibrunneum* m.) [Turkey: İzmir: Bozdağ]

Provincial distribution in Turkey: TR-A: ES IZ (Pic 1903b; Bodemeyer 1906; Breuning 1948, 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

\**D. semilineatum* Fairmaire, 1866: 273

Type information: Syntypes ♂♂ and ♀♀, ex collection L. Fairmaire, Muséum National d'Histoire Naturelle, Paris [type locality 'Bozdağ' (Turkey: İzmir)]

*medioincisum* Breuning, 1946: 108 (*D. semilineatum* m.) [Turkey: İzmir: Bozdağ]

Provincial distribution in Turkey: TR-A: IZ KN (Fairmaire 1866; Ganglbauer 1884; Breuning 1946, 1962a; Tuatay et al. 1972; Özdkmen et al. 2005; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

\**D. semivelutinum* Kraatz, 1873: 82

Type information: Holotype ♀, ex Musaeo Mniszech > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Kleinasién' (Turkey)]

Provincial distribution in Turkey: TR-A: SV (Kraatz 1873; Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Rapuzzi and Sama 2012; Danilevsky 2015a)

Range: A: TR

*D. septemlineatum* Waltl, 1838: 469

\**D. septemlineatum abanti* Braun, 1976c: 54

Type information: Holotype ♂, collection Walter Braun, Tübingen [type locality 'Abant' (Turkey: Bolu)]

Provincial distribution in Turkey: TR-A: AN BO CN DU KR (Braun 1976c, 1978a; Sama 1982; Önalp 1991; Adlbauer 1992; Özdkmen and Hasbenli 2004a; Özdkmen et al. 2005, 2012d; Özdkmen 2007, 2010, 2012; Löbl and Smetana 2010; Al-Hamadani and Özdkmen 2014; Danilevsky 2015a)

Range: A: TR

**\**D. septemlineatum demirciense* Breuning, 1966a: 19**

Type information: Holotype ♂, Institut Royal des Sciences Naturelles de Belgique [type locality 'Demirci' (Turkey: ?Manisa)]

*subapicale* Breuning, 1966a: 20 (*D. septemlineatum* ssp. *demirciense* m.) [Turkey: ? Manisa: Demirci]

Provincial distribution in Turkey: TR-A: MN (Breuning 1966a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. septemlineatum novemlineatum* Kraatz, 1873: 61**

Type information: [type locality 'Bilecik, Eskişehir' (Turkey)]

*forcipiferum* Kraatz, 1873: 60 ['Palestina', undoubtedly mislabeled]

*dorsobasireductum* Breuning, 1946: 106 (*D. septemlineatum* ssp. *novemlineatum* m.) [Turkey: Eskişehir]

Provincial distribution in Turkey: TR-A: AF BI BU ES GA IP KN KU (Bodemeyer 1900; Breuning 1946, 1962a; Braun 1978a; Adlbauer 1988; Önalp 1991; Özdkmen et al. 2005; Löbl and Smetana 2010; Özdkmen 2010, 2012; Şenyüz and Özdkmen 2013; Danilevsky 2015a)

Range: A: TR

**\**D. septemlineatum octolineatum* Kraatz, 1873: 61**

Type information: [type locality 'Bursa' (Turkey)]

*pseudapiceconjunctum* Breuning, 1946: 105 (*D. septemlineatum* ssp. *octolineatum* m.) [Turkey: Bursa]

Provincial distribution in Turkey: TR-A: BL BS CA IS KO SA (Heyden 1884; Breuning 1946, 1962a; Fuchs and Breuning 1971; Braun 1978a; Adlbauer 1988; Önalp 1991; Löbl and Smetana 2010; Özdkmen 2010, 2012; Tekin and Özdkmen 2015; Danilevsky 2015a)

Range: A: TR

***D. septemlineatum septemlineatum* Waltl, 1838: 469**

Type information: Syntypes, ex collection Ménétriés > Joseph Waltl, Naturhistorisches Museum Wien [type locality 'Turkei' (Turkey)]

*apicale* J. Thomson, 1865: 549 [Turkey]

*rufoapicale* Pic, 1942a: 2 (*D. septemlineatum* var.) [Greece: Salonica]

*basilineatum* Breuning, 1946: 105 (*D. septemlineatum septemlineatum* m.) [Turkey]

Provincial distribution in Turkey: TR-E: IS (Küster 1846b; Kraatz 1873; Breuning 1946, 1962a; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: E: BU GR TR

**\**D. sinopense* Breuning, 1962a: 299**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Bergen' (Turkey: S Sinop)]



*reductevittipenne* Breuning, 1962a: 300 (*D. sinopense* m.) [Turkey: S Sinop: Bergen]  
*drannadaghense* Breuning, 1970: 98 (*D. sinopense* m.) [Turkey: Sinop: Dranaz Mt.]  
*nigrovittipenne* Breuning, 1974b: 65 (*D. sinopense* m.) [Turkey: Sinop: Dranaz Mt.]  
 Provincial distribution in Turkey: TR-A: SN (Breuning 1962a, 1970, 1974b; Braun 1978a;  
 Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)  
 Range: A: TR

**\**D. sinuatevittatum* Pic, 1937: 6**

Type information: Holotype ♂, ex collection Heyrovský > M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Taurus: Salbakos Mt. (Babadağ)' (Turkey: Denizli)]  
*brevemediolineatum* Pic, 1942b: 6 [Turkey: Taurus Mt.]

Provincial distribution in Turkey: TR-A: BU DE KA (Pic 1937, 1942b; Breuning 1962a;  
 Özdkmen et al. 2005; Özdkmen and Okutaner 2006; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. skoupyorum* Bernhauer and Peks, 2014b: 191 [RN]**

Type information: Holotype ♂, collection Heinz Peks, Schwanfeld as *D. skoupyi* [type locality 'Şavşat and Çam pass' (Turkey: Artvin)]  
*skoupyi* Bernhauer and Peks, 2013b: 326 [HN]

Provincial distribution in Turkey: TR-A: ART (Bernhauer and Peks 2013b, 2014b;  
 Danilevsky 2015a)

Range: A: TR

***D. sodale* Hampe, 1852: 313**

**\**D. sodale blumenthali* Breuning, 1966a: 20**

Type information: Holotype ♂, Institut Royal des Sciences Naturelles de Belgique [type locality 'Giresun' (Turkey)]

Provincial distribution in Turkey: TR-A: GI (Breuning 1966a; Löbl and Smetana 2010;  
 Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. sodale sodale* Hampe, 1852: 313**

Type information: Syntypes, ex collection Moritz Wagner, Museum für Naturkunde der Humboldt-Universität zu Berlin) [type locality 'Kurdistan' (Turkey)]

Provincial distribution in Turkey: TR-A: EZ (Plavilstshikov 1958; Gfeller 1972; Braun 1978a;  
 Önalp 1990; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: ?GG TR

**\**D. sodale trapesanticum* Breuning, 1946: 124**

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Trabzon' (Turkey)]

Provincial distribution in Turkey: TR-A: BY TB (Breuning 1946, 1962a; Villiers 1959;  
 Breuning and Villiers 1967; Demelt 1967; Adlbauer 1988; Löbl and Smetana 2010;  
 Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. sonjae* Peks, 1993:** 5

Type information: Holotype ♂, collection Heinz Peks, Schwanfeld [type locality 'Elmapınarı' (Turkey: Elazığ)]

Provincial distribution in Turkey: TR-A: EL (Peks 1993; Pesarini and Sabbadini 2004; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. steineri* Holzschuh, 1977:** 129

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Çamlıhemşin: Hacı Vanak plateau' (Turkey: Rize)]

Provincial distribution in Turkey: TR-A: RI (Holzschuh 1977; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. striolatum* Kraatz, 1873:** 93

Type information: Syntypes ♂♂ and ♀♀, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Caucasus']

*obesum* Tournier, 1872: 340 [HN] [Georgia: Surami]

*masculinum* Plavilstshikov, 1927: 51 [RN] (*D. striolatum* ab.) [Georgia: Surami]

*suramense* Breuning, 1946: 98 [RN] (*D. striolatum* ssp.) [Georgia: Surami]

Provincial distribution in Turkey: TR-A: AG (Kraatz 1873; Plavilstshikov 1958; Villiers 1967; Önalp 1991; Löbl and Smetana 2010; Özdi̇kmen 2010; Danilevsky 2015a)

Range: A: ?AB AR GG ?IN TR

***D. sturmii* Frivaldszky von Frivald, 1837:** 179

Type information: Syntypes, ex collection Imre Frivaldszky, Magyar Természettudományi Múzeum, Budapest [type locality 'Türkei' (Turkey)]

Provincial distribution in Turkey: TR-E: ED (Frivaldszky von Frivald 1837; Waltl 1838; Thomson 1867; Önalp 1991; Özdi̇kmen 2010; Danilevsky 2015a)

Range: E: BU MC TR

**\**D. subatritarse* Breuning, 1966b:** 146

Type information: Holotype ♂, collection Antonello Perissinotto [type locality 'Boğazkale' (Turkey: Çorum)]

Provincial distribution in Turkey: TR-A: CO KIR (Perissinotto and Luchini 1966; Breuning 1966b; Braun 1978a; Özdi̇kmen 2010, 2012; Özdi̇kmen and Kaya 2015a)

Range: A: TR

**\**D. subcinctellum* Breuning, 1962a:** 275

Type information: Holotype ♂, ex collection Pierre Lepesme, Musée d'Histoire Naturelle de Lyon [type locality 'Yazihan: Hamidiye' (Turkey: Malatya)]

Provincial distribution in Turkey: TR-A: MA (Breuning 1962a; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. subinterruptum* Pic, 1900a:** 12 (*D. divisum* var.)

Type information: Holotype, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Konya' (Turkey)]



*indivisum* Pic, 1900a: 12 (*D. divisum* var.) [Turkey: Konya]

*interruptevittatum* Breuning, 1946: 106 (*D. divisum* m.) [Turkey: Afyonkarahisar]

*exiguum* Breuning, 1946: 108 (*D. divisum* m.) [Turkey: Konya]

*posticeconunctum* Breuning, 1946: 108 (*D. divisum* m.) [Turkey: Konya]

Provincial distribution in Turkey: TR-A: AF ES IZ ?IC KN; TR-E: ED (Pic 1900a; Breuning 1946, 1962a; Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Sama et al. 2010; Danilevsky 2015a)

Range: E: TR A: TR

**\**D. subsericatum* Pic, 1901: 12**

**\**D. subsericatum majus* Breuning, 1962a: 371** (*D. rufipenne* ssp. *major*)

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Sinop' (Turkey)]

Provincial distribution in Turkey: TR-A: SN (Breuning 1962a; Braun 1978a; Sama 1982; Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. subsericatum rufipenne* Breuning, 1946: 118** (*D. rufipenne*)

Type information: Holotype ♂, ex collection Vogt [type locality 'Ilgaz Mt.' (Turkey: Kastamonu-Çankırı)]

*cankiriense* Breuning, 1962b: 39 (*D. rufipenne* m.) [Turkey: Çankırı]

*antenigrescens* Breuning, 1962b: 39 (*D. rufipenne* ssp. *cankiriense* m.) [Turkey: Çankırı]

*univittatum* Breuning, 1962a: 371 (*D. rufipenne* m.) [Turkey: Kastamonu: Ilgaz Mt.]

*multivittipenne* Breuning, 1962a: 371 (*D. rufipenne* m.) [Turkey: Kastamonu: Ilgaz Mt.]

Provincial distribution in Turkey: TR-A: AN CN KS (Breuning 1946, 1962a; Demelt 1967; Braun 1978a; Sama 1982; Adlbauer 1988, 1992; Pesarini and Sabbadini 1998; Özdkmen 2006, 2010, 2012; Löbl and Smetana 2010; Al-Hamadani and Özdkmen 2014; Danilevsky 2015a)

Range: A: TR

**\**D. subsericatum subsericatum* Pic, 1901: 12**

Type information: Holotype, ex collection Théry > M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Syria' but undoubtedly mislabeled, should be Turkey]

*paphlagoniense* Breuning, 1944: 12 [Turkey: Paphlagonia: Kastamonu]

Provincial distribution in Turkey: TR-A: KS (Pic 1901; Breuning 1944, 1962a; Braun 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. subsericatum vulneratum* Pesarini and Sabbadini, 1998: 51**

Type information: Holotype ♂, collection Carlo Pesarini and Andrea Sabbadini, Milano [type locality 'Masruf pass' (Turkey: Kastamonu)]

Provincial distribution in Turkey: TR-A: KS (Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. subvestitum* K. Daniel, 1900: 140**

Type information: Syntypes ♂♂ and ♀♀, ex collection K. Daniel, Deutsches Entomologisches Institut, Eberswalde [type locality 'Bozdağ' (Turkey: İzmir)]

Provincial distribution in Turkey: TR-A: AM AN ES IZ KN MA NI (Daniel 1900; Breuning 1962a; Gfeller 1972; Önalp 1990; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. sulcipenne*** Küster, 1847: 87

***D. sulcipenne argonauta*** Suvorov, 1913: 74 (*D. argonauta*)

Type information: Syntypes ♂♂ and ♀♀, ex collection G. L. Suvorov, Zoological Museum, University of Moscow [type locality 'Erivan' (Armenia)]

*kurda* Suvorov, 1913: 76 [Turkey: Kars: Sarıkamış]

Provincial distribution in Turkey: TR-A: EZ KAR (Suvorov 1913; Plavilstshikov 1958; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a)

Range: A: AR GG TR

Other known subspecies: *D. sulcipenne plyushchi* Danilevsky, 2014: 673 A: GG; *D. sulcipenne gubini* Danilevsky, 2014: 674 A: GG; *D. sulcipenne borzhomiense* Danilevsky, 2014: 677 A: GG; *D. sulcipenne zekariense* Danilevsky, 2014: 678 A: GG; *D. sulcipenne paki* Danilevsky, 2014: 674 A: GG; *D. sulcipenne demokidovi* Suvorov, 1915: 115 A: GG; *D. sulcipenne caucasicum* Küster, 1847: 98 A: AR GG; *D. sulcipenne goktshanum* Suvorov, 1915: 117 A: AR; *D. sulcipenne sulcipenne* Küster, 1847: 87 A: GG

\****D. syriense*** Breuning, 1943: 94

Type information: Holotype ♀, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Syria' but undoubtedly mislabeled, should be Amanos Mts. (Turkey)]

Provincial distribution in Turkey: TR-A: ?HT (Breuning 1943, 1962a; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

***D. tauricum*** Waltl, 1838: 468

***D. tauricum pittinoi*** Pesarini and Sabbadini, 2010: 185

Type information: Holotype ♂, Museo Civico di Storia Naturale, Milano [type locality 'Doxa' (Greece)]

Provincial distribution in Turkey: TR-E: ED (Pesarini and Sabbadini 2010; Danilevsky 2015a)

Range: E: GR TR

***D. tauricum tauricum*** Waltl, 1838: 468

Type information: Syntypes, ex collection Imre Frivaldszky, Magyar Természettudományi Múzeum, Budapest [type locality 'Türkei' (Turkey)]

*nigritarse* Kraatz, 1873: 77 [?]

Provincial distribution in Turkey: TR-A: IS SA; TR-E: ED TE (Waltl 1838; Thomson 1867; Breuning 1962a; Önalp 1991; Pesarini and Sabbadini 2010; Özdkmen 2010; Löbl and Smetana 2010; Danilevsky 2015a)

Range: E: BU GR MD RO TR UK A: TR

\****D. theophilei*** Pic, 1898: 57

Type information: Holotype ♂, ex collection M. Pic, Muséum National d'Histoire Naturelle, Paris [type locality 'Trabzon' (Turkey)]

*costiferum* Pic, 1898: 57 (*D. theophilei* var.) [Turkey: Trabzon]

Provincial distribution in Turkey: TR-A: ART GU TB (Pic 1898; Plavilstshikov 1958; Villiers 1959; Breuning 1962a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)  
Range: A: TR

**\**D. turicum*** Breuning, 1963: 35

Type information: Holotype ♂, ex collection Hauser, Naturhistorisches Museum Wien [type locality 'Paşalimanı' (Turkey: Balikesir)]

Provincial distribution in Turkey: TR-A: BL (Breuning 1963; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. ullrichi*** Bernhauer, 1988: 102

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Malatya' (Turkey)]

Provincial distribution in Turkey: TR-A: MA (Bernhauer 1988; Löbl and Smetana 2010; Özdkmen 2010, 2012; Pesarini and Sabbadini 2013; Danilevsky 2015a)

Range: A: TR

***D. urmianum*** Plavilstshikov, 1937: 27

Type information: Holotype ♂, ex collection N. N. Plavilstshikov, Zoological Museum, University of Moscow [type locality 'Urmia' (Iran)]

Provincial distribution in Turkey: TR-A: HA (Braun 1975, 1978a; Löbl and Smetana 2010; Özdkmen 2010, 2012; Sama et al. 2012; Danilevsky 2015a)

Range: A: IN TR

***D. variegatum*** Ganglbauer, 1884: 502

Type information: Syntypes ♂ and ♀, ex collection G. Kraatz, Deutsches Entomologisches Institut, Eberswalde [type locality 'Syria']

*apicedisjunctum* Breuning, 1946: 126 (*D. variegatum* m.) ['Ain-Tab', Turkey: Gaziantep]

*nubilatum* Breuning, 1946: 126 (*D. variegatum* m.) ['Ain-Tab', Turkey: Gaziantep]

Provincial distribution in Turkey: TR-A: GA KA (Breuning 1946, 1962a; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a)

Range: A: SY TR

***D. wagneri*** Küster, 1846a: 87

Type information: Holotype, ex collection Heinrich Carl Küster > Johann Menzel, Naturhistorisches Museum, Nürnberg [type locality 'Ararat Mt.' (Armenia)]

*solyzinosum* Pic, 1942b: 2 [Solyzino]

Provincial distribution in Turkey: TR-A: AG EZ KAR (Kraatz 1873; Ganglbauer 1884; Plavilstshikov 1958; Önalp 1991; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: AR GG TR

**\**D. weyersii*** Fairmaire, 1866: 271

Type information: Syntypes ♂♂ and ♀♀, ex collection L. Fairmaire, Muséum National d'Histoire Naturelle, Paris [type locality 'Kisilgye-Aole (?Ödemiş: Kızılca köyü)' (Turkey: ?İzmir)]

*aolense* Pic, 1942c: 1 (*D. weyersi* var.) [Turkey]

*brunneicolor* Breuning, 1946: 121 (*D. weyersi* m.) [Turkey: İzmir: Ödemiş]

*subbrunneicolor* Breuning, 1946: 121 (*D. weyersi* m.) [Turkey: İzmir: Ödemiş]

Provincial distribution in Turkey: TR-A: BU IZ MN (Fairmaire 1866; Ganglbauer 1884; Pic 1942c; Breuning 1946, 1962a; Güll-Zümreoglu 1972; Adlbauer 1988; Önalp 1991; Löbl and Smetana 2010; Özdimek 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. yahyaliense*** Bernhauer and Peks, 2011a: 217

Type information: Holotype ♂, collection Dieter Bernhauer, Bad Schwalbach [type locality 'Yahyalı' (Turkey: Kayseri)]

Provincial distribution in Turkey: TR-A: KY (Bernhauer and Peks 2011a; Danilevsky 2015a)

Range: A: TR

**Subgenus *D. (Maculatodorcadion)*** Breuning, 1943: 525

[type species *Dorcadion quadrimaculatum* Küster, 1848]

**\**D. janssensi*** Breuning, 1966a: 21

Type information: Holotype ♂, Institut Royal des Sciences Naturelles de Belgique [type locality 'Tatos Mts.' (Turkey: Rize-Erzurum)]

Provincial distribution in Turkey: TR-A: EZ RI (Breuning 1966a; Löbl and Smetana 2010; Özdimek 2010, 2012; Danilevsky 2015a)

Range: A: TR

**\**D. phrygicum*** Peks, 1993: 8

Type information: Holotype ♂, collection Heinz Peks, Schwanfeld [type locality 'Kaş: Ova' (Turkey: Antalya)]

Provincial distribution in Turkey: TR-A: ANT BS CA DU (Breuning 1962a; Peks 1993; Löbl and Smetana 2010; Özdimek 2010, 2012; Özdimek et al. 2012d; Danilevsky 2015a; Özdimek and Kaya 2015f)

Range: A: TR

***D. quadrimaculatum*** Küster, 1848: 79

Type information: Syntypes ♂♂ and ♀♀, ex collection Heinrich Carl Küster > Johann Menzel, Naturhistorisches Museum, Nürnberg [type locality 'Türkei' (Turkey)]

*bimaculatum* Pic, 1898: 56 (*D. quadrimaculatum* var.) [Greece: Attica]

*innotatum* Pic, 1908: 2 (*D. quadrimaculatum* var.) [Greece]

*clementinum* Pic, 1918: 11 (*D. quadrimaculatum* var.) [Greece: Clementi]

Provincial distribution in Turkey: TR-E (Küster 1848; Ganglbauer 1884; Breuning 1962a; Krätschmer 1985; Althoff and Danilevsky 1997; Löbl and Smetana 2010; Özdimek 2010, 2012; Danilevsky 2015a)

Range: E: GR TR

**\**D. triste*** Frivaldszky von Frivald, 1845: 184

Type information: Syntypes ♂♂ and ♀♀, ex collection Imre Frivaldszky, Magyar Természettudományi Múzeum, Budapest [type locality 'Smyrna' (Turkey: Izmir)]

*vittipenne* Breuning, 1946: 97 (*D. triste* m.) [Turkey: Izmir]

*postmedioreductum* Breuning, 1946: 97 (*D. triste* m.) [Turkey: Izmir]

Provincial distribution in Turkey: TR-A: BL BS IZ (Frivaldszky von Frivald 1845; Ganglbauer 1884; Breuning 1946, 1962a; Demelt 1963; Güll-Zümreoglu 1972; Braun 1978a; Krätschmer 1985; Önalp 1990; Peks 1993; Löbl and Smetana 2010; Özdimek 2010, 2012; Danilevsky 2015a; Özdimek and Kaya 2015f)



Range: A: TR

\****D. wolfi*** Krätschmer, 1985: 24

Type information: Holotype ♂, collection O.E. Krätschmer [type locality 'Burhaniye: Ören' (Turkey: Balıkesir)]

Provincial distribution in Turkey: TR-A: BL (Krätschmer 1985; Adlbauer 1992; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a)

Range: A: TR

**Genus *Megalodorcadiion*** Pesarini and Sabbadini, 1998: 58

[type species *Dorcadion ledereri* J. Thomson, 1865]

**Subgenus *M. (Anatolodorcadiion)*** Özdkmen and Kaya, 2015g: 3

[type species *Dorcadion dombilicoides* Özdkmen and Kaya, 2013]

\****M. dombilicoides*** Özdkmen and Kaya, 2013: 494 (*Dorcadion*)

Type information: Holotype ♂, collection H. Özdkmen, Zoology Museum of Gazi University, Ankara [type locality 'Gölünyazı' (Turkey: Çorum)]

Provincial distribution in Turkey: TR-A: CO (Özdkmen and Kaya 2013, 2015g; Danilevsky 2015a)

Range: A: TR

\****M. glabrofasciatum*** K. Daniel, 1900: 140 (*Dorcadion*)

Type information: Holotype ♂, ex collection K. Daniel, Deutsches Entomologisches Institut, Eberswalde [type locality 'Bilecik' (Turkey)]

*imparivittatum* K. Daniel, 1900: 140 (*D. glabrofasciatum* var.) [Turkey]

*parivittatum* Breuning, 1946: 98 (*D. glabrofasciatum* m.) [Turkey: İzmir: Bozdağ]

*glabroseparatum* Breuning, 1962a: 310 (*D. glabrofasciatum* m.) [Turkey: Eskişehir]

Provincial distribution in Turkey: TR-A: AF BI ES IZ US (K. Bodemeyer 1900, 1906; Daniel 1900; Breuning 1946, 1962a; Demelt 1963; Braun 1978a; Önalp 1991; Pesarini and Sabbadini 2004; Löbl and Smetana 2010; Özdkmen 2010, 2012; Özdkmen and Kaya 2013, 2015g; Danilevsky 2015a)

Range: A: TR

\****M. nudefasciatum*** Özdkmen and Şenyüz, 2015: 647

Type information: Holotype ♂, collection Y. Şenyüz, Dumlupınar University, Kütahya [type locality 'Dumlupınar University' (Turkey: Kütahya)]

Provincial distribution in Turkey: TR-A: KU (Özdkmen and Şenyüz 2015)

Range: A: TR

**Subgenus *M. (Fusodorcadiion)*** Özdkmen and Kaya, 2015g: 3

[type species *Dorcadion parallelum* Küster, 1847]

\****M. parallelum*** Küster, 1847: 79 (*Dorcadion*)

Type information: Holotype ♂, ex collection Jacob Sturm, Zoologische Staatssammlung des Bayerischen Staates, München [type locality 'Türkei' (Turkey)]

*rufinimembre* Breuning, 1946: 132 (*D. parallelum* m.) [?Syria, should be Turkey]

Provincial distribution in Turkey: TR-A: AM AN CO TO YO (Küster 1847; Ganglbauer 1884; Braun 1978a; Önalp 1991; Adlbauer 1992; Özdkmen and Hasbenli 2004a; Löbl and

Smetana 2010; Özdkmen 2010, 2012; Özdkmen and Kaya 2013, 2015g; Danilevsky 2015a)

Range: A: TR

**Subgenus *M.* (*Megalodorcadion*)** Pesarini and Sabbadini, 1998: 58

[type species *Dorcadion ledereri* J. Thomson, 1865]

\****M. escherichi*** Ganglbauer, 1897: 54 (*Dorcadion*)

Type information: Syntypes ♂♂ and ♀, ex collection K. L. Escherich, Deutsches Entomologisches Institut, Eberswalde [type locality 'Ankara' (Turkey)]

*egregium* Ganglbauer, 1897: 56 [Turkey: Ankara]

*angorense* Ganglbauer, 1897: 57 [Turkey: Ankara]

*oblique-signatum* Pic, 1900a: 12 (*D. escherichi* var.) [Turkey: Konya]

*posticedisjunctum* Pic, 1909b: 99 (*D. escherichi* var.) [Turkey: Ankara]

Provincial distribution in Turkey: TR-A: AM AN BI KN TO (Ganglbauer 1897; Bodemeyer 1900; Pic 1900a, 1909b; Breuning 1962a; Braun 1978a; Önalp 1991; Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Özdkmen and Kaya 2013, 2015g; Danilevsky 2015a)

Range: A: TR

\****M. ledereri*** J. Thomson, 1865: 548 (*Dorcadion*)

Type information: Syntypes, ex collection J. Thomson > R. Oberthür, Muséum National d'Histoire Naturelle, Paris [type locality 'Turkey']

*cappadocicum* Breuning, 1946: 132 (*D. ledereri* m.) [Turkey: Cappadocia]

*discoseparatum* Breuning, 1946: 132 (*D. ledereri* m.) [Turkey: Amasya]

*preconjugatum* Breuning, 1946: 132 (*D. ledereri* m.) [Turkey: Amasya]

*presuturenigrum* Breuning, 1970: 98 (*D. ledereri* m.) [Turkey: Tokat]

Provincial distribution in Turkey: TR-A: AM CO SM TO (Thomson 1865, 1867; Ganglbauer 1884; Breuning 1946, 1962a, 1970; Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Özdkmen and Kaya 2013, 2015g; Danilevsky 2015a)

Range: A: TR

\****M. walteri*** Holzschuh, 1991: 55 (*Dorcadion*)

Type information: Holotype ♂, collection Carolus Holzschuh, Villach [type locality 'Alput Bey village' (Turkey: Bolu)]

Provincial distribution in Turkey: TR-A: BO (Holzschuh 1991; Pesarini and Sabbadini 1998; Löbl and Smetana 2010; Özdkmen 2010, 2012; Anichtchenko and Shavrin 2013; Özdkmen and Kaya 2013, 2015g; Danilevsky 2015a)

Range: A: TR

**Genus *Neodorcadion*** Ganglbauer, 1884: 437

[type species *Lamia bilineata* Germar, 1824]

**Subgenus *N.* (*Neodorcadion*)** Ganglbauer, 1884: 437

[type species *Lamia bilineata* Germar, 1824]

***N. bilineatum*** Germar, 1824: 485 (*Lamia*)

Type information: Syntypes, ex collection Germar, Museum für Naturkunde der Humboldt-Universität zu Berlin [type locality 'Austria']

*macedonicum* Pic, 1917b: 5 (*N. bilineatum* var.) [Macedonia]



Provincial distribution in Turkey: TR-E: ED IS TE (Heyden 1884; Ganglbauer 1884; Plavilstshikov 1958; Breuning and Villiers 1967; Braun 1978a; Althoff and Danilevsky 1997; Özdkmen 2010; Danilevsky 2015a; Özdkmen and Kaya 2015g)

Range: E: AL BH BU CR GR HU MC MD RO SK TR UK YU

**\**N. exornatoides*** Breuning, 1962a: 68

Type information: Holotype ♂, ex collection S. Breuning, Muséum d'Histoire Naturelle de Genève [type locality 'Balıkesir' (Turkey)]

Provincial distribution in Turkey: TR-A: BL (Breuning 1962a; Pesarini and Sabbadini 2008; Löbl and Smetana 2010; Özdkmen 2010, 2012; Danilevsky 2015a; Özdkmen and Kaya 2015g)

Range: A: TR

***N. exornatum*** Frivaldszky von Frivald, 1835: 268 (*Dorcadion*)

Type information: Holotype, ex collection Imre Frivaldszky, Magyar Természettudományi Múzeum, Budapest [type locality 'Balkans']

*labyrinthicum* J. Thomson, 1865: 549 (*Dorcadion*) [Ukraine: Crimea]

*balcanicum* Tournier, 1872: 284 (*Dorcadion*) [?Balkans 'Orient']

*anticejunctum* Pic, 1917b: 6 (*N. balcanicum* var.) [Greece: Salonica]

*subdiversum* Pic, 1917b: 6 (*N. balcanicum* var.) [Balkans]

*basisseparatum* Breuning, 1947: 170 (*N. exornatum* m.) [Turkey]

*latefasciatum* Breuning, 1948: 58 (*N. exornatum* m.) [Turkey: İstanbul: Alem Mt.]

Provincial distribution in Turkey: TR-A: IS; TR-E (Ganglbauer 1884; Breuning 1947, 1948, 1962a; Althoff and Danilevsky 1997; Pesarini and Sabbadini 2008; Löbl and Smetana 2010; Özdkmen 2010; Danilevsky 2015a; Özdkmen and Kaya 2015g)

Range: E: BU GR RO TR A: TR

***N. laqueatum*** Waltl, 1838: 469 (*Dorcadion*)

Type information: Holotype, ex collection Imre Frivaldszky > Joseph Waltl, Naturhistorisches Museum Wien [type locality 'İstanbul' (Turkey)]

*interruptum* Mulsant and Rey, 1863: 150 (*Dorcadion*) [Turkey: İstanbul]

*sparsum* Mulsant and Rey, 1863: 152 (*Dorcadion*) [Turkey: İstanbul]

*abeillei* Tournier, 1872: 208 (*Dorcadion*) [Turkey]

*anceyi* Pic, 1913: 186 (*N. laqueatum* var.) [Turkey: İstanbul]

*bulgaricum* Pic, 1913: 186 (*N. laqueatum* var.) [Bulgaria]

*insparsum* Pic, 1914: 8 (*N. laqueatum* var.) [Turkey: İstanbul]

*lemonulti* Pic, 1914: 8 (*N. laqueatum* var.) [Turkey: İstanbul]

*diversesparsum* Pic, 1914: 8 (*N. laqueatum* var.) [Turkey: İstanbul]

*sparsenotatum* Pic, 1914: 8 (*N. laqueatum* var.) [Turkey: İstanbul]

*subinterruptum* Pic, 1917b: 6 (*N. laqueatum* var.) [Turkey]

*servilorum* Pic, 1917b: 6 (*N. laqueatum* var.) [Turkey: İstanbul]

*kadiense* Pic, 1917a: 10 (*N. balcanicum* var.) ['Kadi Kei', Turkey: İstanbul: Kadıköy]

*rufobrunneum* Breuning, 1947: 170 (*N. fallax* m.) [Turkey: İstanbul: Alem Mt.]

*nigrimembre* Breuning, 1947: 170 (*N. laqueatum* m.) [Turkey: İstanbul: Alem Mt.]

Provincial distribution in Turkey: TR-A: AM BS IS KO; TR-E: IS (Waltl 1838; Küster 1848; Mulsant and Rey 1863; Ganglbauer 1884; Bodemeyer 1906; Pic 1913, 1914, 1917a, 1917b;

Breuning 1947, 1962a; Demelt and Alkan 1962; Demelt 1963; Braun 1978a; Althoff and Danilevsky 1997; Pesarini and Sabbadini 2008; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a; Özdi̇kmen and Kaya 2015g)

Range: E: BU TR A: TR

\**N. orientale* Ganglbauer, 1884: 510 (*N. balcanicum* var.)

Type information: Syntypes ♀♀, ex collection L. Ganglbauer, Naturhistorisches Museum Wien [type locality 'Kleinasiens' (Turkey)]

*flaschneri* Pic, 1890: clxxv [Turkey: İstanbul: Beykoz]

*dispar* Pic, 1892b: xxvii (*N. flaschneri* var.) [Turkey]

*merkli* Pic, 1893: cxcvi (*N. flaschneri* var.) [Turkey]

*eduardi* Pic, 1900c: 82 (*N. flaschneri* var.) [Turkey]

Provincial distribution in Turkey: TR-A: IS IZ KO (Ganglbauer 1884; Pic 1890, 1892b, 1893, 1900c; Bodemeyer 1906; Demelt and Alkan 1962; Breuning 1962a; Demelt 1963; Braun 1978a; Pesarini and Sabbadini 2008; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a; Özdi̇kmen and Kaya 2015g)

Range: A: TR

***N. pelleti* Mulsant and Rey, 1863: 149 (*Dorcadion*)**

Type information: Holotype, ex collection Petri Pellet, Muséum d'Histoire Naturelle de Perpignan [type locality 'Smyrne' (Turkey: İzmir)]

*segne* Mulsant and Rey, 1863: 155 (*Dorcadion*) [Turkey: İzmir]

*disjunctum* Pic, 1917b: 6 (*N. segne* var.) ['?Serbia and Montenegro: Beograd' probably mislabeled]

*credulum* Pic, 1917a: 10 (*N. segne* var.) [Turkey]

Provincial distribution in Turkey: TR-A: IS IZ; TR-E: IS (Mulsant and Rey 1863; Ganglbauer 1884; Bodemeyer 1906; Pic 1917a; Villiers 1959; Breuning 1962a; Pesarini and Sabbadini 2008; Löbl and Smetana 2010; Özdi̇kmen 2010, 2012; Danilevsky 2015a; Özdi̇kmen and Kaya 2015g)

Range: E: GR TR A: TR

#### ***Dorcadionini species with uncertain status in Turkish fauna***

Some Dorcadionini species that were recorded from Turkey erroneously and whose occurrences in Turkey need confirmation are excluded from the list and included here.

#### ***Dorcadion (Carinatodorcadion)***

***Dorcadion carinatum* (Pallas, 1771)** – This species has been recorded by a few authors from NE Anatolia. It has five subspecies from European Russia, Kazakhstan, Ukraine and Azerbaijan. The subspecies structure of this species was revised by Danilevsky (1998). According to Danilevsky (1998) distribution patterns of *D. carinatum* never reach Turkey in the South. Therefore, the presence of this species in Turkey is uncertain, in my opinion.

#### ***Dorcadion (Cribridorcadion)***

***Dorcadion arenarium* (Scopoli, 1763)** – This species has been recorded by a few authors from NC and NW Anatolia. It has 11 subspecies from France, Italy, Croatia,



Bosnia and Herzegovina, Slovenia and Albania. Therefore, the presence of this species is impossible for Turkey.

***Dorcadion atticum Kraatz, 1873*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Greece now. It is endemic to Greece. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion auratum Tournier, 1872*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Georgia now. It is endemic to Georgia. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion beckeri Kraatz, 1873*** – This species has been recorded by a few authors from N and E Anatolia. It has two subspecies from South European Russia and Azerbaijan. Therefore, the presence of this species in Turkey is impossible.

***Dorcadion cinerarium (Fabricius, 1787)*** – This species has been recorded by various authors from different parts of Anatolia. It has 19 subspecies from European Russia, Ukraine, Moldavia and Azerbaijan. According to Lazarev (2009), *Dorcadion* populations from Turkey were wrongly regarded as different forms of *D. cinerarium* (*D. micans* Thomson, 1867, *D. sericatum* Kraatz, 1873 and *D. macropus* Kraatz, 1873). He also stated that *Dorcadion* populations that were attributed before to *D. cinerarium* from Turkey (as subspecies, variations, morphs, forms or aberrations) belong to another partly undescribed species. Therefore, the presence of this species in Turkey is doubtful.

***Dorcadion decipiens (Germar, 1824)*** – This species recorded by Önalp (1991) and Lodos (1998) from Turkey without any exact locality. It is known from Hungary, Moldavia, Romania, Ukraine, Yugoslavia and Bulgaria. Önalp (1991) mentioned a male specimen in the Senckenberg Museum and one female specimen in her collection from Turkey. Therefore, the presence of this species in Turkey is uncertain.

***Dorcadion elegans Kraatz, 1873*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from South European Russia, Ukraine and Kazakhstan now. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion etruscum (Rossi, 1790)*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Italy now. It is diffuse in Italy, Greece and Albania. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion glaucum Faldermann, 1837*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Iran now. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion holosericeum Krynicki, 1832*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. It has two subspecies from European Russia, Kazakhstan, Belarus, Moldavia, Poland, Romania, Ukraine and Georgia. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion impressicolle Kraatz, 1873*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Syria now. Therefore its presence in Turkey is uncertain.

***Dorcadion indutum Faldermann, 1837*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Armenia and Azerbaijan now. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion investitum Breuning in Braun, 1978a: 111*** – This species, as attributed to Breuning, has been recorded only by Braun (1978a) from Erzurum province in NE Anatolia. It is a nomen nudum. Therefore, it is removed from the list.

***Dorcadion litigiosum Ganglbauer, 1884*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. It has two subspecies from Romania, ?Moldavia and Ukraine now. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion mediterraneum Breuning, 1942*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Italy now. It is endemic to Italy. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion minutum Kraatz, 1873*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. It has two subspecies from Greece. It is endemic to Greece. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion murrayi Küster, 1847*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Romania and Yugoslavia now. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion pusillum Küster, 1847*** – This species recorded only by Lodos (1998) from Turkey in his unrealistic list without any exact locality. It has three subspecies from Moldavia, Romania, Ukraine and South European Russia now. Therefore, the presence of this species in Turkey is impossible.

***Dorcadion reitteri Ganglbauer, 1884*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded only from Georgia now. It is endemic to Georgia. Therefore, it is impossible for Turkey.



***Dorcadion rosinae K. Daniel in Bodemeyer, 1900: 127*** – This species, as attributed to K. Daniel, has been recorded only by Bodemeyer (1900) from Konya province (Akşehir, Sultan Mt.) in C Anatolia. It is a nomen nudum. Therefore, it is removed from the list.

***Dorcadion sareptanum Kraatz, 1873*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. It has four subspecies from South European Russia, Kazakhstan and Ukraine now. Therefore, it is impossible for it to occur in Turkey.

***Dorcadion scopolii (Herbst, 1784)*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species has been recorded from Bulgaria, Croatia, Czech Republic, Hungary, Moldavia, Poland, Romania, Slovakia, Ukraine and Yugoslavia now. Therefore, the presence of this species in Turkey is uncertain.

***Dorcadion semilucens Kraatz, 1873*** – This species has been recorded by a few authors from Anatolia. However, the species is known only from Armenia and Azerbaijan now. Therefore, the presence of this species in Turkey is uncertain.

***Dorcadion seminudum Kraatz, 1873*** – This species has been recorded only by Plavilstshikov (1958) and Lodos (1998) from NE Anatolia. However, the species is known only from Armenia and Azerbaijan now. Therefore, the presence of this species in Turkey is uncertain.

***Dorcadion sericatum Sahlberg, 1823*** – This species has been recorded by various authors from different parts of Anatolia. It is endemic to Crimea now. Therefore, the presence of this species in Turkey is impossible.

***Dorcadion tuerki Ganglbauer, 1884*** – This species has been recorded only by Demelt (1963) from Amanos Mountains in South Anatolia. It is endemic to Iran now. Therefore, the presence of this species in Turkey is impossible.

#### ***Neodorcadion (s. str.)***

***Neodorcadion fallax (Kraatz, 1873)*** – This species has been reported only by Lodos (1998) for Turkey in his unrealistic list without any exact locality. However, the species is recorded only from Bulgaria, Greece and Macedonia now. Therefore, the presence of this species in Turkey is uncertain.

## **Results and discussion**

This work provides essential information to understand and conserve the existing biodiversity of Turkey (a particular region). Although the Dorcadionini of Turkey is not well studied, there is always a need to present updated information because of changes in the taxonomic nomenclature and many surveys resulting in new descriptions, records and localities. This is a crucial attempt to present a synthesised and updated list of Dorcadionini of Turkey. In all, 278 species-group taxa [Dorcadion: 265 species-group taxa; Megalodorcadion: seven species;

and *Neodorcadion*: six species) are listed. *Eodorcadion*, *Iberodorcadion* and *Politodorcadion* have no species in Turkey. The Dorcadionini fauna of Turkey presents a remarkable endemism with 227 species group taxa, which constitute about 82% of the total number. This is surely a result of hotspots included in the area. We suggest future investigations in localities that are not well collected. Forthcoming surveys will most likely contribute to the Dorcadionini fauna of Turkey with new records, and new species and subspecies.

## Disclosure statement

No potential conflict of interest was reported by the author.

## References

- Adlbauer K. 1982. Eine neue *Dorcadion*-Art aus Kleinasien (Coleoptera, Cerambycidae). Nachrichtenblatt der Bayerischen Entomologen. 31:104–107.
- Adlbauer K. 1988. Neues zur Taxonomie und Faunistik der Bockkäferfauna der Türkei (Coleoptera, Cerambycidae). Entomofauna. 9:257–297.
- Adlbauer K. 1992. Zur Faunistik und Taxonomie der Bockkäferfauna der Türkei II (Coleoptera, Cerambycidae). Entomofauna. 13:485–509.
- Al-Hamadani NDS, Özdi̇kmen H. 2014. Longicorn beetles of Çankırı province in Turkey (Coleoptera: Cerambycidae). Munis Entomol Zool. 9:931–941.
- Althoff J, Danilevsky ML. 1997. A check-list of longicorn beetles (Coleoptera, Cerambycoidea) of Europe. Ljubljana: Slovensko Entomološko Društvo Štefana Michelija; p. 64.
- Anichtchenko AV, Shavrin AV. 2013. The type specimens of Coleoptera (Insecta) deposited in the Beetle collection (DUBC) of Daugavpils University, Latvia. Acta Biologica Universitatis Daugavpiliensis. 13:1–10.
- Bernhauer D. 1988. Türkische Dorcadien (1. Beitrag) (Coleoptera: Cerambycidae: Lamiinae). Entomologische Zeitschrift (Essen). 98:97–112.
- Bernhauer D. 2010. Eine neue Art der Gattung *Dorcadion* Dalm. aus der Türkei, nördlich von Anamur (Coleoptera, Cerambycidae, Lamiinae). 2. Beitrag zur Kenntnis der türkischen *Dorcadion*. Coleoptera, Schwanfelder Coleopterologische Mitteilungen. 14:189–194.
- Bernhauer D, Peks H. 2010. Zwei neue Arten der Gattung *Dorcadion* Dalm. aus der Türkei, nördlich von Anamur (Coleoptera, Cerambycidae, Lamiinae), sowie eine neue Unterart zu *Dorcadion lodosi* Sabbadini und Pesarini. 3. Beitrag zur Kenntnis der türkischen *Dorcadion*. Coleoptera, Schwanfelder Coleopterologische Mitteilungen. 14:195–206.
- Bernhauer D, Peks H. 2011a. Fünf neue Arten der Gattung *Dorcadion* Dalm. Aus dem mittleren und östlichen Taurus. (Coleoptera, Cerambycidae, Lamiinae). 6. Beitrag zur Kenntnis der türkischen *Dorcadion*. Coleoptera, Schwanfelder Coleopterologische Mitteilungen. 15:215–227.
- Bernhauer D, Peks H. 2011b. Vier neue Arten der Gattung *Dorcadion* Dalm. aus der Nord-West-Türkei. (Coleoptera, Cerambycidae, Lamiinae). 5. Beitrag zur Kenntnis der türkischen *Dorcadion*. Coleoptera, Schwanfelder Coleopterologische Mitteilungen. 15:201–214.
- Bernhauer D, Peks H. 2012. Vier neue Arten der Gattung *Dorcadion* Dalm. aus der Osttürkei. (Coleoptera, Cerambycidae, Lamiinae). 7. Beitrag zur Kenntnis der türkischen *Dorcadion*. Entomologische Blätter und Coleoptera. 108:209–221.
- Bernhauer D, Peks H. 2013a. Drei neue Arten der Gattung *Dorcadion* Dalman, 1817 aus der Südost-Türkei, sowie drei neue Unterarten von *D. (Cribridorcadion) goelbasiense* sp.n. (Coleoptera, Cerambycidae, Lamiinae). 8. Beitrag zur Kenntnis der türkischen *Dorcadion*. Entomologische Blätter und Coleoptera. 109:309–323.
- Bernhauer D, Peks H. 2013b. Vier neue Arten der Gattung *Dorcadion* Dalman, 1817 aus der Nordtürkei, (Coleoptera, Cerambycidae, Lamiinae). 9. Beitrag zur Kenntnis der türkischen *Dorcadion*. Entomologische Blätter und Coleoptera. 109:325–336.



- Bernhauer D, Peks H. 2014a. Fünf neue Arten der Gattung *Dorcadion* Dalman, 1817 aus der Zentral- und Osttürkei (Coleoptera, Cerambycidae, Lamiinae). 10. Beitrag zur Kenntnis der türkischen *Dorcadion*. Entomologische Blätter und Coleoptera. 110:101–118.
- Bernhauer D, Peks H. 2014b. *Dorcadion skoupyorum* nom. nov. for *Dorcadion skoupyi* Bernhauer & Peks, 2014 (Coleoptera, Cerambycidae). Human Space Int Almanac. 3:191–192.
- Bilberg GJ. 1817. [New taxa]. In: Schönherr CJ, editor. Appendix ad C. J. Schönherr Synonymiam Insectorum. Descriptiones Novarum Specierum Insectorum. Scaris, Lewerentziana. Stockholm. 1:1–266.
- Bodemeyer HEV. 1900. Quer durch Klein Asien, in den Bulghar Dagh; Eine Naturwissenschaftliche studien-Reise. Emmendingen: Coleopterologisches; p. 196.
- Bodemeyer HEV. 1906. Beiträge zur Käferfauna von Klein Asien. Deutsche Entomologische Zeitschrift. 2:417–437.
- Braun W. 1975. Beitrag zur Kenntnis der Gattung *Dorcadion* (Col., Cerambycidae, Lamiinae). Entomologische Zeitschrift. 85:17–21.
- Braun W. 1976a. 3. Beitrag zur Kenntnis der Dorcadien Anatoliens. *Dorcadion ladikanum* n. nom. Und Bemerkungen zur Kenntnis des *Dorcadion ardahense* Breuning (Col., Cerambycidae). Entomologische Zeitschrift (Stuttgart). 86:173–175.
- Braun W. 1976b. 4. Beitrag zur Kenntnis der Dorcadien Anatoliens. Neue Dorcadien-Arten und Bemerkungen zur Variationsbreite des *D. mesopotamicum* Breun. (Col., Cerambycidae). Entomologische Zeitschrift (Stuttgart). 86:254–264.
- Braun W. 1976c. 2. Beitrag zur Kenntnis der Dorcadien Anatoliens (Col., Cerambycidae). Entomologische Zeitschrift (Stuttgart). 86:54–60.
- Braun W. 1978a. Die Dorcadienausbeute der Forschungsreisen von W. Heinz 1963–1977, Faunistische Aufstellung, Beschreibung einer neuen Unterart und Bemerkungen zur Systematik wenig bekannter Arten (Coleoptera, Cerambycidae). Nachrichtenblatt der Bayerischen Entomologen. 27:101–116.
- Braun W. 1978b. Ein neue Art der Gattung *Dorcadion* aus Anatolien (Col.: Cerambycidae). Entomologische Zeitschrift (Stuttgart). 88:185–187.
- Braun W. 1979. Beitrag zur Kenntnis der Gattung *Dorcadion* Systematisch neu bewertete *Dorcadion*-Formen (Col., Cerambycidae). Nachrichtenblatt der Bayerischen Entomologen. 28:81–86.
- Breit J. 1923. Beitrag zur Kenntnis der *Dorcadion*-Arten des Balkans. Wiener Entomologische Zeitung. 40:145–149.
- Breuning S. 1942. Novae species Cerambycidarum. XI. Folia Zoologica et Hydrobiologica. 11:113–175.
- Breuning S. 1943. Nouveaux Cérambycides paléarctiques (1re note). Miscellanea Entomologica. 40:89–104.
- Breuning S. 1944. Nouveaux Cerambycides paléarctiques (2e note). Miscellanea Entomologica. 41:11–16.
- Breuning S. 1946. Nouvelles formes de *Dorcadion* (Col. Cerambycidæ). Miscellanea Entomologica. 43:91–132.
- Breuning S. 1947. Nouvelles formes de *Dorcadion* (Col., Cerambycidæ) 2e note. Miscellanea Entomologica. 43:149–172.
- Breuning S. 1948. Nouvelles formes de *Dorcadion* (Col., Cerambycidae). (3e note). Miscellanea Entomologica. 45:57–64.
- Breuning S. 1955. Beschreibung einer neuen Rasse des *Dorcadion divisum* Germ. (Ceramb., Coleoptera). Annales Historico-Naturales Musei Nationalis Hungarici, Budapest (Ser. Nov.). 6:263.
- Breuning S. 1956. Deux nouveaux Lamiaires de Sumatra et deux nouvelles formes de *Dorcadion* du Musée de Budapest (Coleoptera, Cerambycidae). Annales Historico-Naturales Musei Nationalis Hungarici (series nova). 7:403–404.
- Breuning S. 1960. Nouvelles formes de Lamiaires. (douzième partie). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique (Bruxelles). 36:1–30.

- Breuning S. 1962a. Revision der Dorcadionini. Entomologische abhandlungen und berichte aus dem staatliche museum für tierkunde in Dresden. 27:665.
- Breuning S. 1962b. Neue und interessante Formen der Gattungen *Carabus* L. und *Dorcadion* Muls. aus Anatolien. Reichenbachia (Dresden). 1:37–40.
- Breuning S. 1963. Weiterer Beitrag zur Kenntnis der Gattung *Dorcadion* Dalm. (Col. Ceramb.). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen. 15:35–36.
- Breuning S. 1964a. Nouvelles formes de Lamiaires. (Seizième partie). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique (Bruxelles). 40:1–8.
- Breuning S. 1964b. Quatre nouvelles espèces du genre *Dorcadion* Dalm. (Coleoptera Cerambycidae). Bollettino dell'Associazione Romana di Entomologia. 19:31–32.
- Breuning S. 1966a. Nouvelles formes de Lamiaires (dix-septième partie). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique (Bruxelles). 42:1–22.
- Breuning S. 1966b. Deux nouvelles espèces du genre *Dorcadion* Dalm. d'Anatolie (Coleoptera, Cerambycidae). Bollettino della Società Entomologica Italiana (Genova). 96:145–147.
- Breuning S. 1969. Quelques formes nouvelles du genre *Dorcadion* Dalm. de Grèce (Coleoptera Cerambycidae). Bollettino dell'Associazione Romana di Entomologia. 24:42.
- Breuning S. 1970. Nouveaux *Dorcadion* des collections du Muséum de Paris. L'Entomologiste (Paris). 26:97–101.
- Breuning S. 1973. Beschreibung zweier neuer Arten der Gattung *Dorcadion* (Col., Cerambycidae, Lamiinae). Entomologische Zeitschrift (Frankfurt an Main). 83:54–55.
- Breuning S. 1974a. Beschreibung neuer Arten der Gattung *Dorcadion* Dalm. (Coleoptera: Cerambycidae). Mitteilungen der Entomologischen Gesellschaft (Basel, NF). 24:148–154.
- Breuning S. 1974b. Descriptions de Dorcadions nouveaux d'Anatolie (Coleoptera, Lamiinae). Bulletin de la Société Entomologique de Mulhouse [octobre-novembre-décembre]. 1974:65.
- Breuning S. 1975. Descriptions de Dorcadions nouveaux d'Anatolie (Coleoptera Lamiinae). Bulletin de la Société Entomologique de Mulhouse. 1975:10.
- Breuning S. 1977. Quelques rectifications systématiques sur les Lamiaires (Col. Cérambycidae). Bulletin de la Société Entomologique de Mulhouse [Octobre-Décembre]. 1977:59–60.
- Breuning S. 1979. Beschreibung einer neuen species der Gattung *Dorcadion* (Col.: Cerambycidae). Entomologische Zeitschrift (Stuttgart). 89:92–93.
- Breuning S. 1981. Descriptions de quelques nouvelles espèces de Lamiinae (Col., Cerambycidae) de la coll. Bremer. Entomologische Blätter für Biologie und Systematik der Käfer. 77:181–182.
- Breuning S, Ruspoli M. 1971. Description de deux espèces nouvelles de Cerambycidae (Col.) d'Anatolie. Entomops (Nice). 20:127–128.
- Breuning S, Ruspoli M. 1975. Description d'une espèce nouvelle de Cerambycidae (Col.) d'Anatolie. Entomops (Nice). 36:116.
- Breuning S, Villiers A. 1967. Cérambycides de Turquie (2. note). L'Entomologiste. 23:59–63.
- Chevrolat LAA. 1882. Espèces nouvelles de Longicornes européens et circa-méditerranéens et Remarques diverses. Annales de la Société Entomologique de France, Paris. 2:57–64.
- Chevrolat LAA. 1856. Coléoptères nouveaux. Revue et Magasin de Zoologie (Paris). 8:84–89.
- Cihan N, Özdiikmen H, Aytar F. 2013. Contributions of the longhorned beetles knowledge of Turkey by the subfamilies Prioninae, Lepturinae, Dorcadioninae and Lamiinae (Coleoptera: Cerambycidae). Munis Entomol Zool. 8:883–894.
- Çiplak B. 2003. Distribution of Tettigoniinae (Orthoptera, Tettigoniidae) bush-crickets in Turkey: the importance of the Anatolian Taurus Mountains in biodiversity and implications for conservation. Biodiv Conserv. 12:47–64.
- Çiplak B. 2004. Biogeography of Anatolia: the marker group Orthoptera. Mem Soc Entomol Ital. 82:357–372.
- Conservation International: Biodiversity Hotspots [Internet].c. 2015. Arlington, Virginia: Center for Applied Biodiversity Science at Conservation International; [cited 2015 Aug 15]. Available from: <http://www.biodiversityhotspots.org/>
- Dalman JW. 1817. Descriptiones novarum specierum insectorum. In: Schönherr CJ, editor. Synonymia insectorum, oder: Versuch einer Synonymie aller bisher bekannten Insecten; nach



- Fabricii Systema Eleutheratorum & c. geordnet. Eleutherata oder Käfer. Lewerentzischen Buch., Skara. 1: 506 pp.
- Daniel K. 1900. Vorläufige Diagnosen. Societas Entomologica. 15:139–140.
- Daniel K, Daniel J. 1903. Nova, von Hauptmann v. Bodemeyer in Kleinasien gesammelt. 2. Serie. Münchener Koleopterologische Zeitschrift. 1:319–333.
- Danilevsky ML. 1993. New species of Cerambycidae (Coleoptera) from East Asia with some new records. Annales Historica-Naturales Musei Nationalis Hungarici. 84:111–116.
- Danilevsky ML. 1996. New taxa of the genus *Dorcadion* Dalman from Asia (Coleoptera, Cerambycidae). Lambillionea. 96:407–420.
- Danilevsky ML. 1998. A revue of subspecific configuration of *Dorcadion* (*Carinatodorcadion*) *carinatum* (Pallas, 1771) with a description of new subspecies (Coleoptera: Cerambycidae). Acta Entomologica Slovenica. 6:135–142.
- Danilevsky ML. 1999. Two new subspecies of *Dorcadion scabricolle* Dalman, 1817 close to *D. scabricolle* *sevangense* Reitter, 1889 from Transcaucasia (Coleoptera, Cerambycidae). Lambillionea. 99:25–29.
- Danilevsky ML. 2009. Species group taxa of longhorned beetles (Coleoptera, Cerambycidae) described by N. N. Plavilstshikov and their types preserved in the zoological museum of the Moscow State University and in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg. Entomol Rev (Washington). 89:689–720.
- Danilevsky ML. 2012. Additions and corrections to the new catalogue of palaearctic Cerambycidae (Coleoptera) edited by I. Löbl and A. Smetana, 2010. Part. III. Munis Entomol Zool. 7:109–173.
- Danilevsky ML. 2014. Two new subspecies of *Dorcadion aethiops* (Scopoli, 1763) from Bulgaria (Coleoptera, Cerambycidae). Human Space Int Almanac. 3:251–254.
- Danilevsky ML. 2015a. Catalogue of Palaearctic Cerambycoidea [cited 2015 Mar 8]. Available from: <http://www.cerambycidae.net/catalog.pdf>
- Danilevsky ML. 2015b. A new species of genus *Dorcadion* Dalman, 1817 (Coleoptera, Cerambycidae) from Turkey (Ispır, Erzurum). Human Space Int Almanac. 4:220–223.
- Danilevsky ML, Kasatkina DG. 2007. Further investigation of Dorcadionini (Coleoptera: Cerambycidae) endophallus with a revision of taxonomical position of the genus *Trichodorcadion* Breuning, 1942. Russ Entomol J. 15:401–407.
- Danilevsky ML, Kasatkina DG, Rubenyan AA. 2005. Revision of the taxonomic structure of the tribe Dorcadionini (Coleoptera, Cerambycidae) on the base of endophallic morphology. Russ Entomol J. 13:127–149.
- Danilevsky ML, Murzin SV. 2009. A revue of *Dorcadion* Dalman, 1817 species of “laeve-group” Part I (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes. 92:1–20.
- Dascalu M-M, Fusu L. 2012. *Dorcadion axillare* Küster, 1847 (Coleoptera, Cerambycidae): distribution, morphometrics, karyotype and description of a new subspecies from Romania. Zootaxa. 3322:35–48.
- Demelt CV. 1963. Beitrag zur Kenntnis der Cerambycidenfauna Kleinasiens und 13. Beitrag zur Biologie palaearkt. Cerambyciden, sowie Beschreibung einer neuen *Oberea*-Art. Entomologische Blätter. 59:132–151.
- Demelt CV. 1967. Nachtrag zur Kenntnis der Cerambyciden-Fauna Kleinasiens. Entomologische Blätter. 63:106–109.
- Demelt CV, Alkan B. 1962. Short information of Cerambycidae Fauna of Turkey. Bitki Koruma Bülteni. 2:49–56.
- Desbrochers des Loges J. 1870. Descriptions de Coléoptères nouveaux d'Europe et confins. L'Abeille. 7:97–135.
- Ekiz AN, Şen İ, Aslan EG, Gök A. 2013. Checklist of leaf beetles (Coleoptera: Chrysomelidae) of Turkey, excluding Bruchinae. J Nat Hist. 47:2213–2287.
- Fairmaire LMH. 1866. Notice sur les Coléoptères récoltés par M. J. Léderer sur le Bosz-Dagh (Asie Mineure). Annales de la Société Entomologique de France (Paris). 6:249–280.
- Faldermann F. 1837. Fauna entomologica Transcaucasica. Coleoptera 2. Longicornes vel Capricornes. Nouveaux Mémoires de la Société Impériale des Naturalistes de Moscou. 5:1–433.

- Ferrari GA. 1864. Zwei neue *Dorcadion*-Arten vom Caucasus beschrieben vom süddeutschen Entomologen. Wiener entomologische Monatschrift. 8:479–483.
- Fischer von Waldheim G. 1823a. Entomographie de la Russie, et genres des Insectes. Entomographia Imperii Rossici, sue Caesareae Majestati Alexandro I dicata. 2: plates 18–50.
- Fischer von Waldheim G. 1823b. Entomographie de la Russie, et genres des Insectes. Entomographia Imperii Rossici, sue Caesareae Majestati Alexandro I dicata. 2: 264 pp.
- Fairmaire LMH. 1884. Liste des Coléoptères recueillis par M. l'abbé David à Akbès (Asie-Mineure) et Descriptions des Espèces Nouvelles. Annales de la Société Entomologique de France. 4:165–180.
- Frivaldszky von Frivald I. 1835. Á Magyar Tudós Társaság' Évkönyvei. Második Kötet. Á Magyar Tudós Társaság' Évkönyvei. 2:251–276.
- Frivaldszky von Frivald I. 1837. Ujabb Közlései az általa eszközölt, balkányi természettudományi utazásról. Á Magyar Tudós Társaság' Évkönyvei. 3:156–184.
- Frivaldszky von Frivald I. 1845. A Királyi Magyar természettudományi társulat évkönyvei I. Annalen der Ungarischen Naturforschenden Gesellschaft, Pesten. 1:161–187.
- Fuchs E, Breuning S. 1971. Die Cerambycidenausbeute der Anatolianexpedition 1966–67 des Naturhistorischen Museums, Wien. Annalen Naturhistorischen Museum Wien. 75:435–439.
- Ganglbauer L. 1882. Über *Dorcadion graecum* Waltl. Wiener Entomologische Zeitung. 1:228.
- Ganglbauer L. 1884. Bestimmungs-Tabellen der europaeischen Coleopteren. VIII. Cerambycidae. Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien. 33:437–586.
- Ganglbauer L. 1886. Neue und weniger bekannte Longicornier des paläarktischen Faunengebietes. Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien. 35:515–524.
- Ganglbauer L. 1897. Cerambycidae. Bestimmt und beschrieben von L. Ganglbauer. In K. Escherich: zoologische Ergebnisse einer von Dr. K. Escherich und Dr. L. Kathariner nach Central-Kleinasien unternommenen Reise. IV. Theil. Coleopteren. Stettiner Entomologische Zeitung. 58:1–69.
- Germar EF. 1824. Insectorum species novae aut minus cognitae, descriptionibus illustratae. Coleopterorum Species. Hendelii et filii, Halae Volumen primum. Coleoptera: xxiv + 1–624.
- Germar EF. 1839. Fauna Insectorum Europae. Fasc. 21. Halae: C. A. Kümmelii; p. 1–25.
- Gfeller W. 1972. Cerambycidae (Coleoptera) der Türkei-Persienexpedition 1970 der Herren Dr. H. c. W. Wittmer und U. v. Botmer. Mitteilungen der Entomologischen Gesellschaft Basel. 22:1–8.
- Gistel JNFX. 1848. Naturgeschichte des Thierreichs für höhere Schulen. Hoffmann, Stuttgart. 216 pp.
- Gül-Zümreoğlu S. 1972. Catalogue of Insect and common pests (1928–1969). Bornova: T. C. Publications of Agriculture Ministry; p. 119.
- Gül-Zümreoğlu S 1975. Investigations on taxonomy, host plants and distribution of the long-horned beetles (Cerambycidae-Coleoptera) in Aegean Region. T. C. Ministry of Food, Agriculture and Stockbreeding, No: 28. Izmir: İstiklal Press; 208 pp.
- Hampe C. 1852. Verzeichniß der von mir im Kaukasus, in Trans-Kaukasien, Armenien, Kurdistan und West-Persien gesammelten und von Dr. Hampe bestimmten Koleopteren. In Moritz Wagner, editor. Reise nach Persien und dem Lande der Kurden. 2:302–315.
- Herbst JFW. 1784. Kritisches Verzeichniß meiner Insektsammlung. Archiv der Insectengeschichte. 5:1–151.
- Heyden L. 1884. Beitrag zur Coleopterenfauna der Insel Askold und anderer Theile des Amurgebietes. Deutsche entomologische Zeitschrift. 28:273–300.
- Heyden L. 1888. Neue und interessante Coleopteren aus Malatia in Mesopotamien. Deutsche Entomologische Zeitschrift. 32:72–78.
- Heyden L. 1894. Ueber Meloe Olivieri Chevrolat und sechs neue Longicornen aus Klein-Asien. Deutsche Entomologische Zeitschrift. 1894:85–90.
- Heyrovský L. 1932. Weitere neue palaearktische Cerambyciden-formen. Entomologisches Nachrichtenblatt (Troppau). 6:103–104.
- Heyrovský L. 1943. Príspěvek k faune tribu Dorcadiini Thrácie (Col., Ceramb.). Casopis České Společnosti Entomologické. 40:77–80.
- Heyrovský L. 1962. Zwei neue *Dorcadion*-Arten aus der Balkanhalbinsel. Bulgarische Akademia der Wissenschaften. Mitteilungen des zoologischen Institutes und Museums. 12:225–226.



- Heyrovský L. 1964. Eine neue *Dorcadion*-Art aus Kleinasien (Coleoptera, Cerambycidae). Reichenbachia (Dresden). 3:97–98.
- Holzschuh C. 1977. Neue Bockkäfer aus Anatolien und Iran (Col., Cerambycidae). Koleopterologische Rundschau (Wien). 53:127–136.
- Holzschuh C. 1980. Revision einer Cerambycidenausbeute des Naturhistorischen Museums Wien (Coleoptera). Annalen des Naturhistorischen Museums in Wien. 83:573–574.
- Holzschuh C. 1982. Elf neue Bockkäfer aus Europa und Asien (Col.: Cerambycidae). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen. 33:65–76.
- Holzschuh C. 1989. Beschreibung neuer Bockkäfer aus Europa und Asien (Cerambycidae, Col.). Koleopterologische Rundschau (Wien). 59:153–183.
- Holzschuh C. 1991. 63 neue Bockkäfer aus Asien, vorwiegend aus China und Thailand (Coleoptera: Disteniidae und Cerambycidae). Schriftenreihe der Forstlichen Bundesversuchsanstalt (Fbva-Berichte) (Wien). 60:5–71.
- Holzschuh C. 1995. Beschreibung von 65 neuen Bockkäfern aus Europa und Asien, vorwiegend aus Thailand und China (Coleoptera: Disteniidae und Cerambycidae). Schriftenreihe der Forstlichen Bundesversuchsanstalt (Fbva-Berichte) (Wien). 84:1–63.
- Holzschuh C. 2007. Beschreibung von 80 neuen Bockkäfern aus der orientalischen und palaearktischen Region, vorwiegend aus China, Laos und Borneo (Coleoptera, Cerambycidae). Entomologica Basiliensa et Collectionis Frey. 29:177–286.
- Jakovlev AI. 1901. Deux nouvelles espèces de *Dorcadion* Dalm. De l'Asie mineure (Coleoptera, Cerambycidae). Revue Russe d'Entomologie (Saint-Petersbourg). 1:83–87.
- Jakovlev BE. 1899. Nouvelles espèces du genre *Dorcadion* Dalm. Horae Soc Entomol Ross (St. Petersbourg). 34:59–70.
- Jurecek Š. 1931. Dve nové formy tesariku z Recka a Malé Azie. Casopis České Společnosti Entomologické. 28:124.
- Jureček Š. 1933. Noví tesaříci palearktické fauny. Casopis České Společnosti Entomologické. 30:127–130.
- Kadlec S. 2006. Two new species of the genus *Dorcadion* Dalman, 1817 from Iran and Turkey (Col.: Cerambycidae: Lamiinae). Animma.x. 12:8–15.
- König EG. 1901. Erster Beitrag zur Coleopteren-Fauna des Kaukasus. Wiener Entomologische Zeitung. 20:9–10.
- Konstantinov AS, Korotyaev BA, Volkovitsh MG. 2009. Insect biodiversity in the Palearctic region. Chapter 7. In: Foottit RG, Adler PH, editors. Insect biodiversity: science and society. Oxford: Wiley-Blackwell; p. 107–162.
- Kraatz G. 1871. *Dorcadion formosum* n. sp. Berliner Entomologische Zeitschrift. 14:411–412.
- Kraatz G. 1873. Die Käfer Europas. Nach der Natur beschrieben von Dr. G. Kraatz im Anschluss an die Käfer Europa's von Dr. H. C. Küster. In Küster, 1873. Die Käfer Europas nach der Natur beschrieben. 29:1–101.
- Kraatz G. 1884. Neu Käfer-Arten aus Malatia im südlichen Kleinasien. Deutsche Entomologische Zeitschrift (Berlin). 28:231–234.
- Kraatz G. 1892. Die Varietäten des *Dorcadion equestre* Laxman. Deutsche entomologische Zeitschrift. 36:173–174.
- Krätschmer OE. 1985. Eine neuer *Maculatodorcadion*. Beitrag zur Kenntnis der *Dorcadion*-Fauna Anatoliens (Coleoptera: Cerambycidae: Lamiinae). Entomologische Zeitschrift. 95:24–27.
- Krätschmer OE. 1987. Beitrag zur Kenntnis der *Dorcadion*-Fauna Griechenlands und der Türkei (Coleoptera: Cerambycidae: Lamiinae). Entomologische Zeitschrift (Essen). 97:337–340.
- Krynicki J. 1832. Enumeratio Coleopterorum Rossiae meridionalis et praecipue in Universitatibus Caesareae Charkoviensis circulo obvenientium, quae annorum 1827–1831 spatio observavit. Bulletin de la Société Impériale des Naturalistes de Moscou. 5:65–179.
- Küçükaykı EC, Şirin Ü, Çalışkan H, Şenyüz Y. 2013. Preliminary work on longhorned beetles fauna of Kaz Dağları (Ida Mountain) and near with two new subspecies (Coleoptera: Cerambycidae). Munis Entomol Zool. 8:50–62.
- Küster HC. 1846a. Die Käfer Europa's. Nach der Natur beschrieben. Mit Beiträgen mehrerer Entomologen. Nürnberg, Bauer & Raspe. 5:1–100.

- Küster HC. 1846b. Die Käfer Europa's. Nach der Natur beschrieben. Mit Beiträgen mehrerer Entomologen. Nürnberg, Bauer & Raspe. 6:1–100.
- Küster HC. 1847. Die Käfer Europa's. Nach der Natur beschrieben. Mit Beiträgen mehrerer Entomologen. Nürnberg, Bauer & Raspe. 8:1–100.
- Küster HC. 1848. Die Käfer Europa's. Nach der Natur beschrieben. Mit Beiträgen mehrerer Entomologen. Nürnberg, Bauer & Raspe (Julius Merz). 15:1–100.
- Küster HC. 1852. Die Käfer Europa's. Nach der Natur beschrieben. Mit Beiträgen mehrerer Entomologen. Nürnberg, Bauer & Raspe. 25:1–100.
- Laxmann E. 1770. Novaes Insectorum species. Novi commentarii Academiae Scientiarum Imperialis Petropolitanae. 14:593–604.
- Lazarev MA. 2009. Armenian *Dorcadiion* (Coleoptera: Cerambycidae) of “cinerarium-group”. Stud Rep District Mus Prague-East Taxon Ser. 5:197–219.
- Lazarev MA. 2011. Species group taxa of longhorned beetles (Coleoptera, Cerambycidae) described by B. E. Jakovlev and their types preserved in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg, and in the Siberian Zoological Museum, Novosibirsk. Entomol Rev (Washington). 91:634–644.
- Lazarev MA. 2013. New subspecies of *Dorcadiion scabricolle* (Dalman, 1817) of Iran and Azerbaijan (Coleoptera, Cerambycidae). Human Space Int Almanac. 2:222–234.
- Lazarev MA. 2014. New subspecies of *Dorcadiion (Cribridorcadion) mniszechi* Kraatz, 1873 from Georgia, Armenia and Turkey (Coleoptera, Cerambycidae). Human Space Int Almanac. 3:698–709.
- Lingafelter SW, Nearns EH, Tavakilian GL, Monné MÁ, Biondi M. 2014. Longhorned woodboring beetles (Coleoptera: Cerambycidae and Disteniidae) primary types of the Smithsonian Institution. Washington (DC): Smithsonian Institution Scholarly Press; p. v–xviii + 1–390.
- Löbl I, Smetana A. 2010. Catalogue of palaearctic coleoptera. Volume 6. Chrysomeloidea. Löbl I, Smetana A, editors. Stenstrup: Apollo Books; p. 1–924.
- Lodos N. 1998. Entomology of Turkey VI (general, aplied and faunistic). Ege Ü. Ziraat Fak. Yayınları No: 529. İzmir: E. Ü. Faculty of Agriculture Press; p. 300.
- Ménétriés E. 1832. Catalogue raisonné des objets de Zoologie recueillis dans un voyage au Caucase et jusqu'aux frontières actuelles de la Perse entrepris par ordre de S. M. L' Empereur. Saint-Pétersbourg. 4:1–271.
- Ménétriés E. 1836. Insectes nouveaux de la Turquie. Bulletin Scientifique de l'Académie Impériale des Sciences de Saint-Petersbourg. 1:149–151.
- Motschulsky VI. 1838. Coléoptères du Caucase et des Provinces Transcaucasiennes, recueillis et décrits par T. Victor. Bulletin de la Société Impériale des Naturalistes de Moscou. 11:175–188.
- Mulsant É, Rey C. 1863. Longicornes nouveaux ou peu connus. Annales de la Société Linnéenne de Lyon (Nouvelle Série). 10:144–184.
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. Nature. 403:853–858.
- Okutaner AY, Özdiğmen H, Yüksel E, Koçak Y. 2011. Some cytogenetic observations of two *Dorcadiion* Dalman, 1817 species (Coleoptera: Cerambycidae: Lamiinae: Dorcadiini). Munis Entomol Zool. 6:866–876.
- Önalp B. 1988. Eine neue *Dorcadiion*-Art aus der Türkei (Coleoptera: Cerambycidae: Lamiinae). Entomologische Zeitschrift (Stuttgart). 98:361–363.
- Önalp B. 1990. Systematic researches on *Dorcadiion* Dalman, 1817 species in Turkey (Coleoptera, Cerambycidae: Lamiinae) I. HÜ Eğitim Fakültesi Dergisi. 5:57–102.
- Önalp B. 1991. Systematic researches on *Dorcadiion* Dalman, 1817 species in Turkey (Coleoptera, Cerambycidae: Lamiinae) II. HÜ Eğitim Fakültesi Dergisi. 6:191–227.
- Özbek H. 1978. *Hylotrupes bajulus* (L.) Serville in Erzurum and the near, and some others longhorn beetles. Ataturk Üniversitesi Ziraat Fakultesi Dergisi. 9:31–44.
- Özdikmen H. 2006. Contribution to the knowledge of Turkish longicorn beetles fauna (Coleoptera: Cerambycidae). Munis Entomol Zool. 1:71–90.
- Özdikmen H. 2007. The longicorn beetles of Turkey (Coleoptera: Cerambycidae) part I – black sea region. Munis Entomol Zool. 2:179–422.



- Özdikmen H. 2010. The Turkish Dorcadiini with zoogeographical remarks (Coleoptera: Cerambycidae: Lamiinae). *Munis Entomol Zool.* 5:380–498.
- Özdikmen H. 2011. Longhorned beetles of Bolu province in Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 6:210–240.
- Özdikmen H. 2012. The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) part III – Cerambycidae (Dorcadioninae). *Munis Entomol Zool.* 7:759–779.
- Özdikmen H. 2015. A new synonym of *Dorcadion (Cribridorcadion) andirinense* Bernhauer & Peks, 2011 (Cerambycidae). *Munis Entomol Zool.* 10:289–290.
- Özdikmen H, Demir H. 2006. Notes on longicorn beetles fauna of Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 1:157–166.
- Özdikmen H, Güven M, Gören C. 2010. Longhorned beetles fauna of Amanos Mountains, Southern Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 5:1141–1167.
- Özdikmen H, Hasbenli A. 2004a. Contribution to the knowledge of longhorned bettles (Coleoptera, Cerambycidae) from Turkey, Subfamily Lamiinae. *J Ent Res Soc.* 6:25–49.
- Özdikmen H, Hasbenli A. 2004b. A New Subspecies of *Dorcadion dimidiatum* Motschulsky, 1838 (Coleoptera, Cerambycidae, Lamiinae) from Turkey. *J Ent Res Soc.* 6:23–27.
- Özdikmen H, Kaya G. 2013. *Dorcadion (Megalodorcadion)* Pesarini & Sabbadini, 1998 with a new species from Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 8:493–501.
- Özdikmen H, Kaya G. 2015a. *Dorcadion muchei* Breuning, 1962 rest. nov. and *Dorcadion subatritarse* Breuning, 1962 rest. nov. (Cerambycidae). *Munis Entomol Zool.* 10:272–276.
- Özdikmen H, Kaya G. 2015b. *Dorcadion (Cribridorcadion) javeti* Kraatz, 1873 rest. nov. (Cerambycidae). *Munis Entomol Zool.* 10:166–168.
- Özdikmen H, Kaya G. 2015c. *Dorcadion (Cribridorcadion) obsoletum* Kraatz, 1873 rest. nov. (Cerambycidae). *Munis Entomol Zool.* 10:222–224.
- Özdikmen H, Kaya G. 2015d. *Dorcadion (Cribridorcadion) paracinerarium* Breuning, 1974 rest. nov. (Cerambycidae). *Munis Entomol Zool.* 10:252–254.
- Özdikmen H, Kaya G. 2015e. *Dorcadion (Cribridorcadion) gebzeense* Breuning, 1974 rest. nov. (Cerambycidae). *Munis Entomol Zool.* 10:186–187.
- Özdikmen H, Kaya G. 2015f. *Dorcadion (Maculatodorcadion) phrygicum* Peks, 1993 stat. nov. (Cerambycidae). *Munis Entomol Zool.* 10:199–200.
- Özdikmen H, Kaya G. 2015g. A propose to new arrangement on some Dorcadionini (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 10:1–10.
- Özdikmen H, Kaya G, Al-Hamadani N. 2014. A new species of *Dorcadion* Dalman, 1817 (Coleoptera: Cerambycidae) from Turkey. *Acta Zoologica Bulgarica.* 66:179–180.
- Özdikmen H, Koçak Ö. 2013. A new species of *Dorcadion* Dalman, 1817 from Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 8:672–676.
- Özdikmen H, Koçak Ö. 2015. A new species of *Dorcadion (Cribridorcadion)* (Coleoptera: Cerambycidae) in Turkey. *Florida Entomol.* 98:439–441.
- Özdikmen H, Mercan N, Cihan N. 2012a. A new subspecies of *Dorcadion halepense* (Kraatz, 1873) from South Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 7:604–606.
- Özdikmen H, Mercan N, Cihan N. 2012b. A new species of *Dorcadion* Dalman, 1817 from Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 7:550–554.
- Özdikmen H, Mercan N, Cihan N. 2012c. Male genitaliae of *Dorcadion narlianum* Özdikmen et al., 2012 and *Dorcadion halepense sehitkamilese* Özdikmen et al., 2012 (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 7:1276–1277.
- Özdikmen H, Mercan N, Tunç H. 2012d. Longhorned beetles of Düzce province in Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 7:714–731.
- Özdikmen H, Okutaner AY. 2006. The longhorned beetles fauna (Coleoptera, Cerambycidae) of Kahramanmaraş province. *G U J Sci.* 19:77–89.
- Özdikmen H, Özdemir Y, Turgut S. 2005. Longhorned beetles collection of the Nazife Tuatay Plant Protection Museum, Ankara, Turkey (Coleoptera, Cerambycidae). *J Ent Res Soc.* 7:1–33.
- Özdikmen H, Şenyüz Y. 2015. A new species of *Megalodorcadion (Anatolodorcadion)* (Coleoptera: Cerambycidae) in Turkey. *Florida Entomol.* 98:647–650.

- Özdikmen H, Turgut S, Güzel S. 2009. Longhorned beetles of Ankara region in Turkey (Coleoptera: Cerambycidae). *Munis Entomol Zool.* 4:59–102.
- Pallas PS. 1771. Reise durch verschiedene Provinzen des Rußischen Reichs. Erster Theil. St. Petersburg: Kayserliche Academie der Wißenschaften; p. 3–504.
- Pallas PS. 1773. Reise durch verschiedene Provinzen des Rußischen Reichs. Zweyter Theil, zweytes Buch vom Jahr 1771. St. Petersburg: Kayserliche Academie der Wißenschaften; p. 369–744.
- Peks H. 1992. Eine neue *Dorcadion*-Art aus Anatolien (Coleoptera: Cerambycidae). *Entomologische Zeitschrift* (Stuttgart). 102:197–200.
- Peks H. 1993. Drei neue Taxa der Gattung *Dorcadion* aus Anatolien (Coleoptera, Cerambycidae, Lamiinae). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen.* 1:3–11.
- Peks H. 2010. Eine neue Art der Gattung *Dorcadion* Dalm. aus der Türkei, Umgebung Iznik (Coleoptera, Cerambycidae, Lamiinae). *Coleoptera.* 14:213–220.
- Perissinotto A, Luchini SR. 1966. Coleotteri Raccolti Nel Vicino e Medio Oriente Nota I. *Dorcadion* Dalm. (Coleoptera, Cerambycidae). *Bollettino della Società Entomologica Italiana.* 96:147–149.
- Pesarini C, Sabbadini A. 1994. Catalogo dei tipi del Museo Civico di Storia Naturale di Milano X. I tipi dei Coleotteri Cerambicidi. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano.* 133:217–244.
- Pesarini C, Sabbadini A. 1998. Osservazioni sistematiche su alcuni *Dorcadion* della fauna anatolica, con descrizione di 9 nuovi taxa (Coleoptera, Cerambycidae). *Annali del Museo civico di Storia naturale di Ferrara.* 1:45–61.
- Pesarini C, Sabbadini A. 2004. Ricerche sui Dorcadiini di Grecia. I. Le specie del Peloponneso (Coleoptera Cerambycidae). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano.* 145:133–153.
- Pesarini C, Sabbadini A. 2007. Ricerche sui Dorcadiini di Grecia. II. Le specie della Grecia centro-meridionale e quelle del gruppo di *Dorcadion kozanii* (Coleoptera Cerambycidae). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano. 2007 Gennaio.* 148:35–85.
- Pesarini C, Sabbadini A. 2008. Ricerche sui Dorcadiini di Grecia. III. Le specie di *Neodorcadion Ganglbauer, 1884*, quelle del gruppo di *Dorcadion ljubetense* e descrizione della nuova specie *Dorcadion ariannaee* (Coleoptera Cerambycidae). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano.* 149:109–124.
- Pesarini C, Sabbadini A. 2009. Sei nuovi taxa di Cerambycidae della fauna turca e greca (Coleoptera). *Annali del Museo Civico di Storia Naturale di Ferrara.* 12:15–32.
- Pesarini C, Sabbadini A. 2010. Ricerche sui Dorcadionini di Grecia. IV. Le specie della Macedonia e della Tracia (Coleoptera Cerambycidae). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano.* 151:179–216.
- Pesarini C, Sabbadini A. 2011. Note su Cerambycidae di Grecia e Turchia, con descrizione di tre nuove specie e una nuova sottospecie (Coleoptera). *Annali del Museo Civico di Storia Naturale di Ferrara.* 13:41–59.
- Pesarini C, Sabbadini A. 2013. Note su *Dorcadion turchi*, con descrizione di cinque nuove specie e tre nuove sottospecie (Coleoptera Cerambycidae). *Annali del Museo Civico di Storia Naturale di Ferrara.* 14:51–64.
- Pic M. 1890. Séance du 11 septembre 1889. [Description d'une espèce nouvelle de Longicorne du genre *Neodorcadion* Gangl. et communication sur *Vadonia bitliensis* Chevr.]. *Annales de la Société Entomologique de France (Paris).* 9:clxxv–clxxvi.
- Pic M. 1891. Séance du 13 mai 1891. [description d'un Longicorne nouveau]. *Bulletin de la Société Entomologique de France (Paris).* 60:lxxvii–lxxviii.
- Pic M. 1892a. Descriptions. L'Échange, Revue Linnéenne. 8:91.
- Pic M. 1892b. Séance du 27 janvier 1892. [Observations complémentaires sur mon *Neodorcadion Flachneri*]. *Bulletin de la Société Entomologique de France (Paris).* 61:xxvii.
- Pic M. 1893. Séance du 27 juillet 1892. [*Neodorcadion Flachneri*, v. *Merkli*, var. n.]. *Bulletin de la Société Entomologique de France (Paris).* 61:cxcvi.



- Pic M. 1894. Descriptions de deux Coléoptères de la Turquie d'Asie. L'Échange, Revue Linnéenne. 10:110–111.
- Pic M. 1895a. Descriptions de Longicornes d'Arménie et régions voisines. L'Échange, Revue Linnéenne. 11:38–40.
- Pic M. 1895b. [Diagnoses de plusieurs Coléoptères rapportés de Syrie par M. C. Delagrange]. Bulletin de la Société Entomologique de France (Paris). 63:cclxxxiv–cclxxxv.
- Pic M. 1897. Voyage de M. Ch. Delagrange dans la Haute Syrie. Liste des Anthicides et supplément aux Longicornes. Annales de la Société entomologique de France (Paris). 66:389–392.
- Pic M. 1898. Supplément. Matériaux pour servir à l'étude des Longicornes. 2:54–58.
- Pic M. 1900a. Descriptions. Matériaux pour servir à l'étude des Longicornes. 3:11–16.
- Pic M. 1900b. Notes diverses. Matériaux pour servir à l'étude des Longicornes. 3:6–7.
- Pic M. 1900c. Contribution à l'étude des Longicornes. L'Échange, Revue Linnéenne. 16:81–83.
- Pic M. 1901. Descriptions. Matériaux pour servir à l'étude des Longicornes. 3:9–14.
- Pic M. 1902a. Description d'un *Dorcadion*, de la Turquie d'Asie (Col.). Bulletin du Muséum National d'Histoire Naturelle de Paris. 8:93–94.
- Pic M. 1902b. Notes diverses et diagnoses. Matériaux pour servir à l'étude des Longicornes. 4:8–11.
- Pic M. 1902c. Coléoptères paléarctiques présumés nouveaux. L'Échange, Revue Linnéenne. 18:25–27.
- Pic M. 1903a. Notes diverses, diagnoses, synonymies. Matériaux pour servir à l'étude des Longicornes. 4:4–9.
- Pic M. 1903b. Espèces et variétés nouvelles de Coléoptères [2me article]. L'Échange, Revue Linnéenne. 19:169–171.
- Pic M. 1904. Notes diverses et diagnoses. Matériaux pour servir à l'étude des Longicornes. 5:3–6.
- Pic M. 1905a. Descriptions abrégées et notes diverses (2e article). L'Échange, Revue Linnéenne. 21:113–115.
- Pic M. 1905b. Nouveaux coléoptères d'Algérie & d'Asie occidentale. L'Échange, Revue Linnéenne. 21:153–156.
- Pic M. 1905c. Descriptions et Notes Diverses. Matériaux pour servir à l'étude des Longicornes. 5:5–15.
- Pic M. 1907. Deux nouvelles variétés du "*Dorcadion divisum*" Germ. L'Échange, Revue Linnéenne. 23:179.
- Pic M. 1908. Descriptions ou diagnoses et notes diverses. L'Échange, Revue Linnéenne. 24:2.
- Pic M. 1909a. Descriptions ou diagnoses et notes diverses – suite. L'Échange, Revue Linnéenne. 25:121–123.
- Pic M. 1909b. Descriptions ou diagnoses et notes diverses – suite. L'Échange, Revue Linnéenne. 25:97–100.
- Pic M. 1910. Notes diverses et diagnoses. Matériaux pour servir à l'étude des Longicornes. 7:2–6.
- Pic M. 1911. Descriptions ou diagnoses et notes diverses (suite.). L'Échange, Revue Linnéenne. 27:185–186.
- Pic M. 1913. Notes diverses, descriptions et diagnoses (suite.). L'Échange, Revue Linnéenne. 29:185–187.
- Pic M. 1914. Notes diverses et diagnoses. Matériaux pour servir à l'étude des Longicornes. 9:3–11.
- Pic M. 1916. Notes diverses, descriptions et diagnoses (suite.). L'Échange, Revue Linnéenne. 32:21–23.
- Pic M. 1917a. Notes diverses, descriptions et diagnoses (suite.). L'Échange, Revue Linnéenne. 33:9–11.
- Pic M. 1917b. Notes diverses et diagnoses. Matériaux pour servir à l'étude des Longicornes. 10:3–10.
- Pic M. 1918. Notes diverses, descriptions et diagnoses (suite.). L'Échange, Revue Linnéenne. 34:9–11.
- Pic M. 1926. Notes diverses, descriptions et diagnoses (suite.). L'Échange, Revue Linnéenne. 42:13–14.
- Pic M. 1927. Notes diverses, descriptions et diagnoses (suite.). L'Échange, Revue Linnéenne. 43:5–7.
- Pic M. 1929. Notes diverses, nouveautés. L'Échange, Revue Linnéenne. 45:9–10.

- Pic M. 1931a. Notes diverses, nouveautés. L'Échange, Revue Linnéenne. 47:1–2.
- Pic M. 1931b. Notes diverses, nouveautés (suite.). L'Échange, Revue Linnéenne. 47:5–6.
- Pic M. 1931c. Notes diverses, nouveautés (suite.). L'Échange, Revue Linnéenne. 47:9–10.
- Pic M. 1932. Notes diverses, nouveautés (suite.). L'Échange, Revue Linnéenne. 48:21–22.
- Pic M. 1934. Nouveautés asiatiques. Matériaux pour servir à l'étude des Longicornes. 11:33–40.
- Pic M. 1937. Notes diverses, nouveautés (suite.). L'Échange, Revue Linnéenne. 53:5–6.
- Pic M. 1942a. Opuscula martialis VII. L'Echange, Revue Linnéenne Numéro spécial. 7:1–16.
- Pic M. 1942b. Coléoptères du globe (suite.). L'Échange, Revue Linnéenne. 58:5–8.
- Pic M. 1942c. Opuscula martialia VIII. L'Echange, Revue Linnéenne Numéro spécial. 8:1–13.
- Pic M. 1942d. Opuscula martialis VI. L'Echange, Revue Linnéenne. Numéro spécial 6:1–16.
- Pic M. 1948. Coléoptères du globe (suite.). L'Échange, Revue Linnéenne. 64:6–16.
- Pic T. 1899. Über *Dorcadion divisum* Germ. und dessen Varietäten. Entomologisches Nachrichtenblatt (Troppau). 25:349–352.
- Plavilstshikov NN. 1927. Coleoptera II. Addenda et corrigenda concernant le Coleopterorum Catalogus, parties 73 et 74 (Lamiinae) de Chr. Aurivillius. Paul Lechevalier, Paris. Encyclopédie Entomologique (série B). 1:49–68.
- Plavilstshikov NN. 1937. Descriptions des espèces nouvelles de genres *Dorcadion* Dalm. et *Neodorcadion* Ganglb. (Coleoptera, Cerambycidae). Sborník Entomologického Oddelení Národního Musea v Praze. 15:25–34.
- Plavilstshikov NN. 1948. A key for Longicorn beetles of Armenia. Erevan. 232 pp. [in Russian].
- Plavilstshikov NN. 1951. New species of timber-beetles of Palaearctic fauna (Coleoptera, Cerambycidae). Archives of the Zoological Museum of Moscow State University. 7:113–122.
- Plavilstshikov NN. 1958. Faune de l'URSS. Insects Coléptères. V.23 (1). Cerambycidae (P.3). Sousfamille Lamiinae. p.1. Moscou, Leningrad; p. 592.
- Poda von Neuhaus N. 1761. Insecta Musei Graecensis, quae in ordines, genera et species Juxta systema naturae Caroli Linnaei digessit. Widmanstad: Graecii. 12 + 127 + 18 pp.
- Rapuzzi P, Sama G. 2012. New taxa and new records of longhorn-beetles from Eastern Mediterranean Region (Coleoptera: Cerambycidae). Munis Entomol Zool. 7:663–690.
- Reitter E. 1889. Neue Coleopteren aus Europa, den angrenzenden Ländern und Sibirien, mit Bemerkungen über bekannte Arten. Sechster Theil. Deutsche entomologische Zeitschrift. 33:17–44.
- Reitter E. 1900. Beitrag zur Coleopteren-Fauna von Europa und den angrenzenden Ländern. Deutsche entomologische Zeitschrift (Berlin). 1900:81–88.
- Rejzek M, Hoskovec M. 1999. Cerambycidae of Nemrut Dağı National Park (Anatolia, South-East Turkey). Biocosome Mésogéen (Nice). 15:257–272.
- Sabbadini A, Pesarini C. 1992. Due nuove specie dei generi *Parmena* Latr. e *Dorcadion* Dalm. appartenenti alla fauna Turca (Coleoptera Cerambycidae). Bollettino Della Società Entomologica Italiana (Genova). 124:27–32.
- Sahlberg CR. 1823. Periculi entomographici, species insectorum nondum descriptas propositur. Aboae. 4:46–64.
- Sama G. 1982. Contributo allo studio dei coleotteri Cerambycidae di Grecia e Asia Minore. Fragmenta Entomologica (Roma). 16:205–227.
- Sama G. 1999. Notes on some Cerambycidae names published by Kraatz and Chevrolat (Coleoptera, Cerambycidae). Entomologische Zeitschrift (Stuttgart). 109:290–295.
- Sama G, Dascalu M-M, Pesarini C. 2010. Description of *Dorcadion gashtarovi* n. sp. (Coleoptera, Cerambycidae) from Romania and Bulgaria with review of the closely related species. North-Western J Zool. 6:286–293.
- Sama G, Rapuzzi P, Özdkmen H. 2012. Preliminary report of the entomological surveys (2010, 2011) of G. Sama and P. Rapuzzi to Turkey (Coleoptera: Cerambycidae). Munis Entomol Zool. 7:22–45.
- Scopoli JA. 1763. Entomologia Carniolica exhibens insecta Carnioliae indigena et distributa in ordines, genera, species, varietates. Methodo Linnaeana, Vindobonae. 1–421.
- Şenyüz Y, Özdkmen H. 2013. A contribution to the knowledge of Turkish longicorn beetles fauna (Coleoptera: Cerambycidae). Munis Entomol Zool. 8:571–577.

- Suvorov GL. 1913. Beschreibung neuer Cerambyciden-Arten (Coleoptera). Revue Russe d'Entomologie (Saint-Petersbourg). 13:66–81.
- Suvorov GL. 1915. Espèces nouvelles des genres *Dorcadion* et *Compsodorcadion* (Coleoptera, Cerambycidae). Revue Russe d'Entomologie (Saint-Petersbourg). 15:115–121.
- Swainson W, Shuckard WE. 1840. On the history and natural arrangement of insects. London: Longman, Brown, Green & Longman's; 406 pp.
- Tavakilian G. 2015. Base de données Titan sur les Cerambycidés ou Longicornes [cited 2015 Jan 16]. Available from: <http://lis-02.snv.jussieu.fr/titan/>
- Tekin K, Özdkmen H. 2015. A contribution of Turkish longhorned beetles fauna from Bursa (Coleoptera: Cerambycidae). Munis Entomol Zool. 10:122–130.
- Théry A. 1896. Séance du 26 février 1896. Description de quelques Cérambycides paléarctiques [Col.]. Bulletin de la Société Entomologique de France (Paris). 1896:108–110.
- Thomson J. 1865. Diagnoses d'espèces nouvelles qui seront décrites dans l'appendix du systema cerambycidarum. Mémoires de la Société Royale des Sciences de Liège. 19:541–578.
- Thomson J. 1867. VI. Supplément à la Révision de la sous-tribu des Dorcadionites (P. 10). Physis Recueil d'Histoire Naturelle (Paris). 1:118–125.
- Tippmann FF. 1958. *Dorcadion*-Belustigungen am Leitha-Gebirge (Austria inferior). Mitteilungen der Münchener entomologischen Gesellschaft. 48:130–165.
- Tournier H. 1872. Catalogue des Longicornes récoltés par M. Théophile Deyrolle, en Imerétie, Mingrélie et Géorgie, et description des espèces nouvelles. Revue et Magasin de Zoologie pure et appliquée (Paris). 23:276–292.
- Tuatay N, Kalkandelen A, Aysev N. 1972. Bitki Koruma Müzesi Böcek Kataloğu (1961–1971). Ankara: T.C. Tarım Bakanlığı; p. 53–55.
- Turgut S, Özdkmen H. 2010. New data for Turkish longhorned beetles fauna from Southern Turkey (Coleoptera: Cerambycidae). Munis Entomol Zool. 5:859–889.
- Villiers A. 1959. Cérambycides de Turquie. L'Entomologiste. 15:7–11.
- Villiers A. 1967. Contribution à la faune de l'Iran. I. – Coléoptères Cerambycidae. Annales de la Société Entomologique de France (NS). 3:327–379.
- Vives E. 1976. Contribución al conocimiento de los *Iberodorcadion* Breuning (Col. Cerambycidae). Miscellanea Zoologica. 111:163–168.
- Waltl J. 1838. Beyträge zur Kenntniss der Coleopteren der Türken. Isis Von Oken (Leipzig). 31:449–472.

## **Appendix 1.** A checklist of Dorcadionini of Turkey. The endemic species are marked with an asterisk

- Tribe DORCADIONINI Swainson and Shuckard, 1840
- Genus *DORCADION* Dalman, 1817: 397
- Subgenus *CARINATODORCADION* Breuning, 1943: 524
- D. aethiops* Scopoli, 1763: 53 (Cerambyx)  
*D. aethiops aethiops* Scopoli, 1763: 53 (Cerambyx)
- D. fulvum* Scopoli, 1763: 53 (Cerambyx)  
*D. fulvum erythropterum* Fischer von Waldheim, 1823a: table L–7 (*D. erythropteron*)
- D. hybridum* Ganglbauer, 1884: 441  
*D. hybridum hybridum* Ganglbauer, 1884: 441
- \**D. ingeae* Peks, 1993: 3
- Subgenus *Cribridorcadion* Pic, 1901: 12
- \**D. abstersum* Holzschuh, 1982: 75
- \**D. accola* Heyden, 1894: 87
- \**D. afflictum* Pesarini and Sabbadini, 1998: 54
- \**D. akpinarensense* Bernhauer and Peks, 2010: 196
- \**D. albolineatum* Küster, 1847: 86
- \**D. albonotatum* Pic, 1895a: 39

- \**D. altinyaylaense* Bernhauer and Peks, 2014a: 106
- \**D. amanense* Breuning, 1943: 94
- \**D. anatolicum* Pic, 1900a: 12
- \**D. andirinense* Bernhauer and Peks, 2011a: 218
- \**D. apicerufum* Breuning, 1943: 91 [RN]
- \**D. aracense* Bernhauer and Peks, 2011b: 204
- \**D. arcivagum* J. Thomson, 1867: 121
- \**D. ardahense* Breuning, 1975: 10
- D. atritarse* Pic, 1931a: 10 (*D. pedestre* var.)
- \**D. avgadiense* Bernhauer and Peks, 2011a: 215
- D. axillare* Küster, 1847a: 88
  - D. axillare axillare* Küster, 1847: 88
- \**D. bangi* Heyden, 1894: 89
  - \**D. bangi bangi* Heyden, 1894: 89
  - \**D. bangi heinzorum* Braun, 1975: 17 (*D. heinzorum*)
  - \**D. bangi roridum* Pesarini and Sabbadini, 1998: 55
- \**D. banjkovskyi* Plavilstshikov, 1958: 147
- \**D. basale* Kraatz, 1873: 86
- \**D. beloni* Pic, 1891: lxxvii
- \**D. bisignatum* Jakovlev, 1899: 66
- D. bistriatum* Pic, 1898: 55
- \**D. bithyniense* Chevrolat, 1856: 88
- D. blanchardi* Mulsant and Rey, 1863: 147
- \**D. blandulum* Holzschuh, 1977: 131
- \**D. bodemeyeri* K. Daniel, 1900: 140
- \**D. boluense* Breuning, 1962b: 38
  - \**D. boluense boluense* Breuning, 1962b: 38
  - \**D. boluense corallinum* Pesarini and Sabbadini, 1998: 53
  - \**D. boluense imitator* Pesarini and Sabbadini, 1998: 53
- \**D. boszdaghense* Fairmaire, 1866: 275
- \**D. bouilloni* Breuning and Ruspoli, 1975: 116
- \**D. brauni* Breuning, 1979: 92
- \**D. bremeri* Breuning, 1981: 181
- D. breuningeri* Heyrovský, 1943: 78
- \**D. bulgharmaadense* Breuning, 1946: 116
- \**D. cachinno* J. Thomson, 1867: 120
- \**D. caglayanceritense* Bernhauer and Peks, 2013a: 314
- \**D. carinipenne* Pic, 1900a: 13
- \**D. carolistanii* Breuning and Ruspoli, 1971: 128
- D. catenatum* Waltl, 1838: 469
- D. catenatum catenatum* Waltl, 1838: 469
  - \**D. catenatum dissimile* Ganglbauer, 1884: 458 (*D. divisum* var.)
  - \**D. catenatum intercicum* Kraatz, 1873: 66 (*D. latevittatum* var.)
  - \**D. catenatum loratum* J. Thomson, 1867: 123 (*D. loratum*)
  - D. catenatum mytilinense* Kraatz, 1873: 66 (*D. latevittatum* var.)
  - \**D. catenatum oedemischense* Heyrovský, 1932: 104 (*D. divisum* var.)
  - \**D. catenatum subdivisum* Breuning, 1955: 263 (*D. divisum* ssp.)
- \**D. cinctellum* Fairmaire, 1866: 272
- D. cingulatum* Ganglbauer, 1884: 484
- \**D. coiffaiti* Breuning, 1962a: 392
- D. condensatum* Küster, 1852: 92
- \**D. confluens* Fairmaire, 1866: 274
- \**D. conspersum* Pesarini and Sabbadini, 2013: 56
- D. cretosum* Ferrari, 1864: 479



- \**D. crux* Billberg, 1817: 175 (*Lamia*)
- \**D. culminicola* J. Thomson, 1867: 122
- \**D. delagrangei* Pic, 1894: 110
- \**D. deyrollei* Ganglbauer, 1884: 482
  - \**D. deyrollei deyrollei* Ganglbauer, 1884: 482
  - \**D. deyrollei opacum* Pesarini and Sabbadini, 2013: 54
- D. dimidiatum* Motschulsky, 1838: 186
  - D. dimidiatum dimidiatum* Motschulsky, 1838: 186
  - \**D. dimidiatum kelkiticum* Özdkmen and Hasbenli, 2004b: 23
  - \**D. dimidiatum korgei* Breuning, 1966a: 21
- \**D. dobrovljanskii* Suvorov, 1915: 116
- \**D. drusoides* Breuning, 1962a: 375
- \**D. eflaniense* Bernhauer and Peks, 2011b: 205
- \**D. elazigi* Fuchs and Breuning, 1971: 439
- \**D. enricisturanii* Breuning and Ruspoli, 1971: 127
  - \**D. enricisturanii densepunctatum* Braun, 1978a: 107
  - \**D. enricisturanii enricisturanii* Breuning and Ruspoli, 1971: 127
- \**D. erdemii* Özdkmen, Kaya and Al-Hamadani, 2014: 179
- D. equestre* Laxmann, 1770: 596 (*Cerambyx*)
  - \**D. equestre bernhauerorum* Peks, 2010: 213
  - D. equestre bisuturale* Jureček, 1933: 128 (*D. equestre* var.)
  - \**D. equestre festivum* Pesarini and Sabbadini, 2013: 58
  - \**D. equestre nogelli* Fairmaire, 1866: 270 (*D. nogelli*)
- \**D. eskipazarense* Bernhauer and Peks, 2011b: 203
- D. ferruginipes* Ménétriés, 1836: 151
- \**D. formosum* Kraatz, 1871: 411
  - \**D. formosum formosum* Kraatz, 1871: 411
  - \**D. formosum ponticum* Breuning, 1964a: 1
- D. gallipolitanum* J. Thomson, 1867: 59
  - D. gallipolitanum gallipolitanum* J. Thomson, 1867: 59
- \**D. gebzeense* Breuning, 1974a: 148
- \**D. gencense* Bernhauer and Peks, 2014a: 110
- \**D. glabricalle* Breuning, 1943: 92
- \**D. goelbasiense* Bernhauer and Peks, 2013a: 310
  - \**D. goelbasiense goelbasiense* Bernhauer and Peks, 2013a: 310
  - \**D. goelbasiense karadutense* Bernhauer and Peks, 2013a: 311
  - \**D. goelbasiense kathaense* Bernhauer and Peks, 2013a: 312
  - \**D. goelbasiense oeksuezdagense* Bernhauer and Peks, 2013a: 311
- \**D. guzeldereense* Bernhauer and Peks, 2012: 213
- \**D. hacgedigiense* Bernhauer and Peks, 2011a: 216
- D. haemorrhoidale* Hampe, 1852: 313
- D. halepense* Kraatz, 1873: 72 (*D. saulcyi* var.)
  - D. halepense halepense* Kraatz, 1873: 72 (*D. saulcyi* var.)
  - \**D. halepense sehitkamilense* Özdkmen, Mercan and Cihan, 2012a: 604
- \**D. hampii* Mulsant and Rey, 1863: 157
  - \**D. hampii aureovittatum* Kraatz, 1873: 81 (*D. aureovittatum*)
  - \**D. hampii hampii* Mulsant and Rey, 1863: 157
- \**D. hanneloreae* Bernhauer and Peks, 2014a: 107
- \**D. heinzi* Breuning, 1964a: 1 (*D. cinerarium* ssp.)
- D. hellmanni* Ganglbauer, 1884: 486
- \**D. holtzi* Pic, 1905a: 115 (*D. culminicola* ssp.)
- \**D. iconiense* K. Daniel, 1900: 140
- \**D. infernale* Mulsant and Rey, 1863: 158
  - \**D. infernale asperatum* Breuning, 1947: 169

- \**D. infernale edremitense* Breuning, 1966a: 20
- \**D. infernale infernale* Mulsant and Rey, 1863: 158
- \**D. infernale miminfernale* Breuning, 1970: 98 (*D. miminfernale*)
- \**D. inikliense* Bernhauer and Peks, 2010b: 208
- \**D. inopinatum* Pesarini and Sabbadini, 2013: 51
- \**D. inspersum* Holzschuh, 1982: 76
- \**D. intrusum* Pesarini and Sabbadini, 2009: 26
- \**D. ispartense* Breuning, 1960: 1
- \**D. ispirene* Bernhauer and Peks, 2013b: 328
- \**D. ivani* Pesarini and Sabbadini, 2011: 47
- \**D. jacobleviellum* Plavilstshikov, 1951: 120
- \**D. janatai* Kadlec, 2006: 11
- D. javeti* Kraatz, 1873: 91
- D. johannisfranci* Pesarini and Sabbadini, 2007: 40
- \**D. karacaorenense* Bernhauer and Peks, 2012: 211
- \**D. karamanense* Özdīkmen and Koçak, 2013: 672
- \**D. karsense* Suvorov, 1915: 118
- \**D. kartaldagense* Bernhauer and Peks, 2013a: 314
- D. kasikporanum* Pic, 1902b: 10
- \**D. kazanciense* Bernhauer and Peks, 2010: 197
- \**D. kindermannii* Waltl, 1838: 470
- \**D. kizildagense* Bernhauer and Peks, 2012: 212
- \**D. konradi* Bernhauer and Peks, 2012: 210
- \**D. kraetschmeri* Bernhauer, 1988: 98
- \**D. kurdistanum* Breuning, 1944: 12
- \**D. kurucanum* Holzschuh, 2007: 249
- \**D. ladikanum* Braun, 1976a: 173 [RN]
- D. laeve* Faldermann, 1837: 278
  - \**D. laeve micula* Plavilstshikov, 1937: 26 (*D. micula*)
- \**D. lameerei* Théry, 1896: 109
- \**D. ledouxi* Breuning, 1974b: 65
- \**D. linderi* Tournier, 1872: 285
- D. lineatocolle* Kraatz, 1873: 57 (*D. femoratum* var.)
- \**D. lodosi* Sabbadini and Pesarini, 1992: 29
  - \**D. lodosi abanozense* Bernhauer and Peks, 2010: 199
  - \**D. lodosi lodosi* Sabbadini and Pesarini, 1992: 29
- \**D. lohsei* Braun, 1976b: 254
- \**D. longulum* Breuning, 1943: 89
- D. lugubre* Kraatz, 1873: 41
  - D. lugubre lugubre* Kraatz, 1873: 41
- \**D. maceki* Holzschuh, 1995: 41
- D. margheritae* Breuning, 1964b: 32
- \**D. martini* Bernhauer, 1988: 100
- \**D. menradi* Holzschuh, 1989: 172
- \**D. merkli* Ganglbauer, 1884: 506
- D. mesopotamicum* Breuning, 1944: 12
- \**D. micans* J. Thomson, 1867: 61
  - \**D. micans micans* J. Thomson, 1867: 61
  - \**D. micans susheriense* Breuning, 1970: 97 (*D. sinopense* ssp.)
- D. mniszechi* Kraatz, 1873: 39
  - D. mniszechi mniszechi* Kraatz, 1873: 39
  - D. mniszechi cavernosum* Lazarev, 2014: 701
- \**D. muchei* Breuning, 1962b: 38
- \**D. multimaculatum* Pic, 1932: 22



- \**D. murzini* Danilevsky, 2015b: 221
- \**D. musense* Bernhauer and Peks, 2014a: 109
- \**D. narlianum* Özdi̇kmen, Mercan and Cihan, 2012b: 550
- \**D. nigrostriatum* Adlbauer, 1982: 104
- \**D. nihalae* Rapuzzi and Sama, 2012: 671
- \**D. nikssarenske* Bernhauer and Peks, 2013b: 327
- D. nitidum* Motschulsky, 1838: 185
- \**D. nobile* Hampe, 1852: 313
- D. nodicorne* Tournier, 1872: 287
- D. obsoletum* Kraatz, 1873: 78 (*D. graecum* var.)
- \**D. obtusum* Breuning, 1944: 13
  - \**D. obtusum marashense* Breuning, 1948: 59
  - \**D. obtusum obtusum* Breuning, 1944: 13
- \**D. oezdurali* Önalp, 1988: 361
- \**D. olympicum* Ganglbauer, 1882: 228
  - \**D. olympicum convexum* Breuning, 1943: 90
  - \**D. olympicum flavosuturale* Krätschmer, 1987: 340
  - \**D. olympicum olympicum* Ganglbauer, 1882: 228
- \**D. ortrudeae* Braun, 1978b: 185
- \**D. paracinerarium* Breuning, 1974a: 149
- \**D. pararufipenne* Braun, 1976b: 257
  - \**D. pararufipenne pararufipenne* Braun, 1976b: 257
  - \**D. pararufipenne rassei* Braun, 1976b: 261
- \**D. parilis* Pesarini and Sabbadini, 2013: 52
- \**D. pavesii* Pesarini and Sabbadini, 1998: 57
- D. pedestre* Poda von Neuhaus, 1761: 34 (*Cerambyx*)
  - D. pedestre pedestre* Poda von Neuhaus, 1761: 34 (*Cerambyx*)
- \**D. peksi* Bernhauer, 2010: 190
- \**D. pergamenum* Pesarini and Sabbadini, 2009: 22
- \**D. petrovitzi* Heyrovský, 1964: 97
- D. pilosipenne* Breuning, 1943: 90
  - \**D. pilosipenne kazdagense* Özdi̇kmen in Küçükaykı et al., 2013: 56
- \**D. piocardi* Kraatz, 1873: 85
- \**D. pittinorum* Pesarini and Sabbadini, 1998: 48
- \**D. pluto* J. Thomson, 1867: 47
- \**D. poleti* Breuning, 1948: 59
- \**D. postalbosuturale* Özdi̇kmen and Koçak, 2015: 439
- \**D. praetermissum* Pesarini and Sabbadini, 1998: 47
  - \**D. praetermissum mikhaili* Özdi̇kmen, 2010: 457
  - \**D. praetermissum praetermissum* Pesarini and Sabbadini, 1998: 47
- \**D. preissi* Heyden, 1894: 86
- \**D. pseudarcivagum* Breuning, 1943: 91
- \**D. pseudinfernale* Breuning, 1943: 94
- \**D. pseudobithyniense* Breuning, 1962a: 465
- \**D. pseudocinctellum* Breuning, 1943: 90
- \**D. pseudoholosericeum* Breuning, 1962a: 300
- \**D. pseudopreissi* Breuning, 1962a: 411
- D. punctipenne* Küster, 1852: 94
- \**D. punctulatum* Pesarini and Sabbadini, 2013: 57
- \**D. punctulicolle* Breuning, 1944: 12
- \**D. pygmaeum* Breuning, 1947: 168
- \**D. quadripustulatum* Kraatz, 1873: 88
  - D. regulare* Pic, 1931b: 6 (*D. gallipolitanum* var.)
- D. regulare regulare* Pic, 1931b: 6 (*D. gallipolitanum* var.)

- \**D. resadiyeense* Bernhauer and Peks, 2011a: 219
- \**D. rigattii* Breuning, 1966b: 145
- \**D. rizeanum* Breuning and Villiers, 1967: 60 (*Pedestredorcadion*)
- \**D. robustum* Ganglbauer, 1884: 500
- \**D. rolandmenradi* Peks, 1992: 197
- D. rosti* Pic, 1900c: 82
- \**D. rufoapicipenne* Breuning, 1946: 125
- \**D. sabanoezueense* Bernhauer and Peks, 2013b: 326
- \**D. sarkislaense* Bernhauer and Peks, 2014a: 105
- D. saulcyi* J. Thomson, 1865: 549
- D. scabricolle* Dalman, 1817: 174 (*Lamia*)
  - \**D. scabricolle balikesirensis* Breuning, 1962a: 460
  - \**D. scabricolle caramanicum* K. Daniel and J. Daniel, 1903: 332
  - \**D. scabricolle crassofasciatum* Özdi̇kmen in Küçükaykçı et al., 2013: 56
  - \**D. scabricolle paphlagonicum* Breuning, 1962a: 459
  - D. scabricolle scabricolle* Dalman, 1817: 174 (*Lamia*)
  - \**D. scabricolle uludaghicum* Breuning, 1970: 98
- \**D. schultzei* Heyden, 1894: 86
- \**D. scrobicolle* Kraatz, 1873: 97
  - \**D. scrobicolle morulum* Holzschuh, 1995: 42
  - \**D. scrobicolle scrobicolle* Kraatz, 1873: 97
- \**D. sebenense* Bernhauer and Peks, 2011b: 202
- \**D. semibrunneum* Pic, 1903b: 170 (*D. mniszechii* var.)
  - \**D. semibrunneum mediocreimpressum* Pic, 1909a: 123 (*D. mniszechii* var.)
  - \**D. semibrunneum notatum* Pesarini and Sabbadini, 2013: 62
  - \**D. semibrunneum semibrunneum* Pic, 1903b: 170 (*D. mniszechii* var.)
- \**D. semilineatum* Fairmaire, 1866: 273
- \**D. semivelutinum* Kraatz, 1873: 82
- D. septemlineatum* Waltl, 1838: 469
  - \**D. septemlineatum abanti* Braun, 1976c: 54
  - \**D. septemlineatum demirciense* Breuning, 1966a: 19
  - \**D. septemlineatum novemlineatum* Kraatz, 1873: 61
  - \**D. septemlineatum octolineatum* Kraatz, 1873: 61
  - D. septemlineatum septemlineatum* Waltl, 1838: 469
- \**D. sinopense* Breuning, 1962a: 299
- \**D. sinuatevittatum* Pic, 1937: 6
- \**D. skoupyorum* Bernhauer and Peks, 2014b: 191 [RN]
- D. sodale* Hampe, 1852: 313
  - \**D. sodale blumenthali* Breuning, 1966a: 20
  - D. sodale sodale* Hampe, 1852: 313
  - \**D. sodale trapesunticum* Breuning, 1946: 124
- \**D. sonjae* Peks, 1993: 5
- \**D. steineri* Holzschuh, 1977: 129
- D. striolatum* Kraatz, 1873: 93
- D. sturmii* Frivaldszky von Frivald, 1837: 179
- \**D. subatritarse* Breuning, 1966b: 146
- \**D. subcinctellum* Breuning, 1962a: 275
- \**D. subinterruptum* Pic, 1900a: 12 (*D. divisum* var.)
- \**D. subsericatum* Pic, 1901: 12
  - \**D. subsericatum majus* Breuning, 1962a: 371 (*D. rufipenne* ssp. *major*)
  - \**D. subsericatum rufipenne* Breuning, 1946: 118 (*D. rufipenne*)
  - \**D. subsericatum subsericatum* Pic, 1901: 12
  - \**D. subsericatum vulneratum* Pesarini and Sabbadini, 1998: 51
- \**D. subvestitum* K. Daniel, 1900: 140

- D. sulcipenne* Küster, 1847: 87  
*D. sulcipenne argonauta* Suvorov, 1913: 74 (*D. argonauta*)  
*\*D. syriense* Breuning, 1943: 94  
*D. tauricum* Waltl, 1838: 468  
   *D. tauricum pittinoi* Pesarini and Sabbadini, 2010: 185  
   *D. tauricum tauricum* Waltl, 1838: 468  
*\*D. theophilei* Pic, 1898: 57  
*\*D. turicum* Breuning, 1963: 35  
*\*D. ullrichi* Bernhauer, 1988: 102  
*D. urmianum* Plavilstshikov, 1937: 27  
*D. variegatum* Ganglbauer, 1884: 502  
*D. wagneri* Küster, 1846a: 87  
*\*D. weyersii* Fairmaire, 1866: 271  
*\*D. yahyaliense* Bernhauer and Peks, 2011a: 217  
Subgenus *Maculatodorcadion* Breuning, 1943: 525  
*\*D. janssensi* Breuning, 1966a: 21  
*\*D. phrygicum* Peks, 1993: 8  
*D. quadrimaculatum* Küster, 1848: 79  
*\*D. triste* Frivaldszky von Frivald, 1845: 184  
*\*D. wolfi* Krätschmer, 1985: 24  
Genus *Megalodordorcadion* Pesarini and Sabbadini, 1998: 58  
Subgenus *Anatolodordorcadion* Özdi̇kmen and Kaya, 2015g: 3  
*\*M. dombilicoides* Özdi̇kmen and Kaya, 2013: 494 (*Dorcadion*)  
*\*M. glabrofasciatum* K. Daniel, 1900: 140 (*Dorcadion*)  
*\*M. nudefasciatum* Özdi̇kmen and Şenyüz, 2015: 647  
Subgenus *Fusodordorcadion* Özdi̇kmen and Kaya, 2015g: 3  
*\*M. parallelum* Küster, 1847: 79 (*Dorcadion*)  
Subgenus *Megalodordorcadion* Pesarini and Sabbadini, 1998: 58  
*\*M. escherichi* Ganglbauer, 1897: 54 (*Dorcadion*)  
*\*M. ledereri* J. Thomson, 1865: 548 (*Dorcadion*)  
*\*M. walteri* Holzschuh, 1991: 55 (*Dorcadion*)
- Genus *Neodorcadion* Ganglbauer, 1884: 437  
Subgenus *Neodorcadion* Ganglbauer, 1884: 437  
*N. bilineatum* Germar, 1824: 485 (*Lamia*)  
*\*N. exornatoides* Breuning, 1962a: 68  
*N. exornatum* Frivaldszky von Frivald, 1835: 268 (*Dorcadion*)  
*N. laqueatum* Waltl, 1838: 469 (*Dorcadion*)  
*\*N. orientale* Ganglbauer, 1884: 510 (*N. balcanicum* var.)  
*N. pelleti* Mulsant and Rey, 1863: 149 (*Dorcadion*)