

CO₂ storage by cyanobacteria induced biomineralization in presence of basaltic glass

Thomas Ferrini¹, Olivier Grandjouan¹, Olivier Pourret^{1,#}, Raul E. Martinez^{2,*}

¹UniLaSalle, AGHYLE, 19 rue Pierre Waguet, 60026 Beauvais cedex, France

²Institut für Geowissenschaften, Universität Freiburg Alberstraße
23b, D-79104 Freiburg, Germany

*Present address: Max Planck Institute for Biogeochemistry

#Corresponding author: olivier.pourret@unilasalle.fr.

Abstract

Cyanobacteria induced biomineralization of atmospheric CO₂ is a natural process leading to the formation of carbonates by spontaneous precipitation or through the presence of nucleation sites, under supersaturated conditions. . Basaltic glass was chosen in this study to test its ability to release cations needed for carbonate formation in presence of *Synechococcus* sp. cyanobacteria. Active cyanobacteria were expected to generate an alkaline environment through photosynthetic metabolism. This process produces oxygen and hydroxide ions as waste products, raising the pH of the immediate cell surface vicinity and indirectly enhancing the carbonate CO₃²⁻ concentration and providing the a degree of saturation that can lead to the formation of calcite CaCO₃ or magnesite MgCO₃. In presence of active cells, the saturation index (SI) increased from -10.56 to -9.48 for calcite and from -13.6 to -12.5 for magnesite, however they remained negative due to the low Ca²⁺ and Mg²⁺ activities. Dead cells were expected to act as nucleation sites by the stepwise binding of carbonate and Ca²⁺ and Mg²⁺ on their surface. In the presence of inactive cells, SI values were closer to 0 but still negative due to the low pH and cation concentrations.

Keywords: biomineralization, cyanobacteria, basaltic glass, carbonate, saturation index

1. Introduction

The raise of CO₂ concentration in the atmosphere is one of the most important environmental problems on Earth today arising from anthropic activity (Gerlach, 2011). Carbon storage methods, or carbon concentration mechanisms (CCM), need to be developed urgently in order to reduce the CO₂ impact on Earth's terrestrial and aquatic environments. The ability of cyanobacteria to induce the biomineralization of CO₂ is an option which is still in its infancy and could possibly be used in the near-future to decrease atmospheric CO₂ levels. Previous studies have proved that the link between cyanobacteria photosynthesis and the presence of metallic cations could lead to carbonate precipitation in the presence of atmospheric CO₂ and under suitable solution conditions (Jansson and Northen, 2010; Martinez *et al.*, 2016; Obst *et al.*, 2009).

Photosynthesis and CCM are the first steps to biomineralization of CO₂. Cyanobacteria are autophototrophic microorganisms. They are primary producers, which generate their own nutrients by carrying out the oxygenic photosynthesis (Martinez *et al.*, 2016). This metabolism needs carbon a source of inorganic carbon in the form of dissolved CO₂, natural light as source of energy and water to provide electrons needed in the redox transformation of carbon (Schubert *et al.*, 1997). The Calvin-Benson cycle permits the assimilation of the inorganic carbon thanks to the initial CO₂-fixing-enzyme of photosynthesis: RuBisCO (Ribulose Biphosphate Carbolxylase Oxygenase) (Fridlyand and Scheibe, 1999; Spreitzer and Salvucci, 2002). Chemical and photochemical reactions take place in photosystems located on the thylakoid membrane of the cyanobacteria cell. ATP (Adenosine triphosphate) and NADPH (Nicotinamide adenine dinucleotide phosphate) are chemical substances which permit CO₂ assimilation. A constant supply of inorganic carbon (Ci) is required to carry out photosynthesis.

Dissolved CO₂, in the form of bicarbonate HCO₃⁻ is the inorganic form of carbon used by cyanobacteria as a source of C_i. A metabolic system is used to attain inorganic carbon from the aqueous environment. The CCM is an environmental adaptation of cyanobacteria which permits to attract C_i inside the cell in the form of HCO₃⁻ and convert it to CO₂ (Price *et al.*, 2008). Molecules are attracted and stocked inside the cell with the help of C_i transporters located in both plasma and outer membranes. HCO₃⁻ transport is enhanced by the extra-ATP generated during photosynthesis. The inorganic carbon is transported into the micro-compartment carboxysome where the bicarbonate molecules are transformed into CO₂ by the enzyme carbonic anhydrase. The remaining CO₂ is then assimilated by the enzyme RuBisCO for photosynthesis. A proton H⁺ is consumed by each of these reactions, leading to the release of a hydroxide ion (OH⁻) outside the cell as a waste product along with O₂. This allows the generation of an alkaline pH within the micro-environment around the cyanobacteria cell. This alkaline pH permits the deprotonation of the external bicarbonate HCO₃⁻ to carbonate ions CO₃²⁻. The presence of both metal cations (e.g. Ca²⁺/Mg²⁺) and carbonate (CO₃²⁻) permits the nucleation of carbonates in the form of calcite (CaCO₃) and magnesite (MgCO₃) (Martinez *et al.*, 2016; Riding, 2006). However, the amount of metal cations and carbonate has to be high sufficient to reach supersaturation with respect to these carbonate minerals.

In this study, basaltic glass was chosen as a source of metallic cations due to its composition and abundance on the Earth's surface. Its importance in the carbon cycle has been proven on several previous studies (Dessert *et al.*, 2003; Grimm *et al.*, 2019; Schlesinger and Bernhardt, 2013; Schott *et al.*, 2012). A high rate of dissolution of basaltic glass would be required to release significant metallic cation concentrations to achieve supersaturation with respect to carbonate phases. Varying an electrolyte NaNO₃ background concentration has been shown to affect the process of simple silicate mineral dissolution and the rate of release of metallic cations as well as their activity (Martinez *et al.*, 2014; Martinez *et al.*, 2016). An

optimal concentration is needed in order to reach Ca^{2+} and Mg^{2+} activities, which together with carbonate ion activities would lead to precipitation of carbonates. The *Synechococcus* cyanobacteria species chosen for this study is representative of seawater cyanobacteria as along with *Prochlorococcus* sp., it is responsible for approximately 30% of the photosynthesis on Earth. *Synechococcus* sp. is a unicellular coccoid cyanobacteria containing outermost surface layers but no cell covering sheath (Martinez *et al.*, 2010; Martinez *et al.*, 2008)(). The goal of this study is to determine whether basaltic glass is an efficient source of metal cations needed for CO_2 mineralization and then to quantify the rates of carbonate precipitation in presence of this primary silicate and *Synechococcus* cyanobacteria species so that optimal conditions in terms of basaltic glass dissolution and cyanobacteria biomass can be found.

2. Material and methods

A 20 mL aliquot of active *Synechococcus* sp. cyanobacteria stock at the stationary phase was placed in a 5 L Schott® bottle. *Synechococcus* sp. cultures were grown in Cyanobacteria BG-11 Freshwater Medium (Sigma-Aldrich C3061) for three weeks to the stationary growth phase (Martinez *et al.*, 2010). Active cyanobacteria stocks were prepared in order to achieve photosynthesis in presence of basaltic glass, in order to quantify the influence of cyanobacteria on the dissolution of the mineral could be observed and the supersaturation conditions reached. Stock cultures of cyanobacteria were kept at room temperature ($23\pm 1^\circ\text{C}$) under constant cool white fluorescent light illumination (5000 lx) and a constant filtered air injection (3 L/min) in order to provide cyanobacteria with needed inorganic carbon. The air inflow was used to keep the culture in a constant movement needed for optimal growth. For each experiment, 5 L of 1/500 diluted cyanobacteria growth medium inoculated with a 20 mL aliquot of regular 1/50 BG-11 (Table 1) cyanobacteria stock was placed on a rotary shaker at 250 rpm (Martinez *et al.*,

2008). Dead cyanobacteria were prepared by autoclaving an aliquot of active stock at 121°C for 20 min. To measure the exact cyanobacteria mass, a 25 mL sample was collected. Vacuum filtration was used to filter the sample through a 0.2 µm pore size cellulose nitrate filter (Whatman®). The filtered sample was placed in an oven at 60°C for 30 min and then weighed.

2.1. Flow Through open reactors system experimental design

Basaltic glass dissolution experiments were performed at 23±1°C by mixing two different stock solutions in flow through reactors with a peristaltic pump with a fixed inflow rate of 1.4 µL/min. The active cyanobacteria and inorganic stocks solutions used are described in **Tables 1 and 2**.

To quantify the rates of basaltic glass and carbonate precipitation in presence of *Synechococcus* sp. cyanobacteria, duplicate flow through reactors were used, each connected to the inorganic and active cyanobacteria stocks by a peristaltic pump (Active cyanobacteria Flow Through open reactor, AFT; **Table 1**).

Table 1 Composition of flow through reactors initial solution for AFT experiment.

5L of active cyanobacteria stock	
Reagent	Quantity
aliquot BG-11	10 mL
<i>Synechococcus</i> sp. at the stationary phase	20 mL
NaNO ₃	0.04 mol/L
MgSO ₄	0.003 mol/L

"Inorganic" stock

Basaltic glass	20 g
NaNO ₃	0.04 mol/L
MgSO ₄	0.003 mol/L

To assess the role of bio-mineralization by dead cells a calcium chloride stock solution was used instead of basaltic glass as the source of metal cations (e.g. Ca²⁺). For these experiments, a flow through reactor was connected through the peristaltic pump to both the calcium chloride and dead cyanobacteria stocks previously washed by centrifuging at 10,000 rpm for 20 min (Calcium Flow Through reactor, CFT; **Table 2**).

Table 2 Composition of flow through reactors initial solution for CFT experiment.

5 L dead cyanobacteria "autoclaved" stock	
Reagent	Quantity
Active cyanobacteria "autoclaved" (20 min at 121°C)	60 mL
NaNO ₃	0.04 mol/L
MgSO ₄	0.003 mol/L
Calcium chloride stock	
CaCl	0.09 mol/L
NaNO ₃	0.04 mol/L
MgSO ₄	0.003 mol/L




Figure 1 AFT experimental set up




Figure 2 CFT experimental set up.

For all experiments, Tygon R 3603 tubes were used to link both stocks to 200 mL flow through reactors with the help of a 4 channel peristaltic pump with a low flow rate of 1.4 $\mu\text{L}/\text{min}$.

2.2. Sample analysis

In order to monitor the chemical evolution and the cyanobacteria biomass concentration of the solution inside the flow through reactor, the pH, conductivity, alkalinity, optical density, and Ca^{2+} and Mg^{2+} concentrations were measured from samples collected in 50 mL polypropylene Falcon® tubes from the output of the flow through reactors, once daily. Time zero was taken to be the time of this first sampling of the flow through reactor solution (Martinez *et al.*, 2016). Reactors were placed in a water bath at 22°C in order to keep their contents solutions at a constant temperature. A 20 mL sample was taken from each reactor for all experiments. An aliquot of 7 mL of sample was used to measure the optical density of the solution at 750 nm wavelength, to approximate the concentration of cyanobacteria biomass over time. The remaining 13 mL were used to measure pH, total alkalinity and conductivity. The pH and alkalinity were measured with an automatic titration system (Metrohm® 719 Titrino automatic titrator). In this study, dominated by the presence of dissolved CO_2 , the total alkalinity (TA) provides an estimate of the concentrations of bicarbonate and carbonate species in solution. The TA is determined through an acid titration (alkalinity titration) where the final concentration of the acid, at the second end-point of the carbonate system, $\text{pH} = 4.3$, represents the total alkalinity of the solution. Carbonate alkalinity (CA) is determined by driving the alkalinity titration to the first end-point for the carbonate system at $\text{pH} = 8.3$. From the carbonate alkalinity, the CO_3^{2-} concentration in solution, which is part of total alkalinity, can be determined:

$$TA = [HCO_3^-] + 2 [CO_3^{2-}] + [OH^-] - [H^+] \quad (1a)$$

$$CA = 2 [CO_3^{2-}] + [OH^-] \quad (1b)$$

As $[CO_3^{2-}]$, $[H^+]$ and $[OH^-]$ are negligible for neutral pH conditions, total alkalinity can be re-defined to:

$$TA = [HCO_3^-] \quad (2)$$

Ca^{2+} and Mg^{2+} concentrations were measured by flame atomic adsorption spectroscopy (FAAS) with an Analytik Jena Vario 6 flame atomic adsorption spectrometer. The conductivity of sample aliquots was measured using a conductivity meter where conductivity is measured in mS/cm (Harris, 2010). Conductivity measurements for a particular solution allow calculation of the ionic strength (IS). The IS is proportional to the concentration of all dissolved salts in a solution. The ionic strength is required to calculate cation activity (a_c) as shown below:

$$a_c = [C] \gamma_C \quad (3)$$

where $[C]$ is the species concentration and γ_C is the activity coefficient. The \log_{10} of γ_C can be calculated as per the Debye-Hückel equation Eq. 4 as a function of the solution IS determined from conductivity measurements:

$$\log \gamma_C = (-0.51 * z^2 * \sqrt{IS}) / (1 + (\alpha * \sqrt{IS} / 305)) \quad (4)$$

where z represents the charge of the ion (e.g. for Mg^{2+} , $z=2$), α the radius of the ion in picometers (e.g. for Ca^{2+} , $r_{Ca^{2+}} = 600$ pm) (Harris, 2010; Stumm and Morgan, 1996).

To predict the formation of carbonate phases in this study, it is first necessary to quantify the degree of solution supersaturation with respect to Ca or Mg carbonates. This requires knowledge of the saturation index for each of these mineral phases. However, calculation of

This article is a preprint and is not peer-reviewed

saturation state is first necessary. The saturation state (Ω) represents the degree of saturation of a solution with respect to a particular mineral phase (Langdon *et al.*, 2000) and is defined as follows:

$$\Omega = (aCa^{2+} * aCO_3^{2-}) / K_{sp} \quad (5)$$

Where K_{sp} is the solubility product of the mineral phase that can potentially form in solution (e.g. for calcite, $K_{sp}=10^{-8.48}$) and, aCa^{2+} and aCO_3^{2-} are the activities of Ca^{2+} and CO_3^{2-} respectively. The saturation index (SI) is then defined as:

$$SI = \log_{10}(\Omega) \quad (6)$$

A positive value of SI would indicate supersaturation with respect to a mineral phase and suggests that precipitation of the mineral would occur. However this depends further on solution conditions (e.g. mineral nucleation, spontaneous nucleation/precipitation) (Langdon *et al.*, 2000).

In the case of precipitation, the rate (R) of calcite and magnesite formation in the experiments with cyanobacteria can be measured as a function of the saturation state (Pechukas, 1981):

$$R = k_+ (\Omega - 1)^n \quad (7)$$

where k_+ and n represent the kinetic constant and reaction order respectively, and are regression parameters.

3. Results

Temporal evolution of the pH, the total alkalinity and the cations concentrations has been measured. These values were used to calculate saturation index with respect to calcite and magnesite. Related values for AFT and CFT are presented in Figures 3 and 4 and Figures 5 and 6, respectively. For all conditions, experiments were carried out in duplicates.




Figure 3 Evolution of pH and total alkalinity through time experiment (0.04 mol/L NaNO₃; AFT2 experiment).




Figure 4 Evolution of saturation index through time experiment (0.04 mol/L NaNO₃; AFT2 experiment).




Figure 5 Evolution of pH and total alkalinity through time experiment (0.004 mmol/L NaNO₃; CFT experiment).




Figure 6 Evolution of Saturation index through time experiment (0.004 mmol/L NaNO₃; CFT experiment)

3.1. AFT experiment results

In this experiments, the pH remained alkaline, starting at 10.9 ± 1 and declining until a pH of 10.1 at 150 h. Total alkalinity increased from 1 mmol/L to 1.8 mmol/L in 242 h for both reactors. $[\text{Ca}^{2+}]$ and $[\text{Mg}^{2+}]$ remained constant for the duration of the experiments at an average of 3.5 mg/L for $[\text{Ca}^{2+}]$ and 1.4×10^{-4} mg/L for $[\text{Mg}^{2+}]$. A slight decrease in these concentrations was observed for the experiment in the first reactor. SI values for both calcite and magnesite remained negative for the time of the experiment. The $\text{SI}_{(\text{calcite})}$ increased from -10.6 to -9.4 over a period of 145 h whereas the $\text{SI}_{(\text{magnesite})}$ raised from -13.6 to -12.5. Both SI curves reached a constant threshold at 150 h.

3.2. CFT experiment results

The pH and the total alkalinity increased through the 300 h of the experiments. A total alkalinity of 0.2 mmol/L was recorded at the start of experiments to reach a final value of 0.75 mmol/L. The pH was for the most part neutral but raised from 6.3 to a last value of 6.75. The calcium chloride used permitted to obtain a rather constant $[\text{Ca}^{2+}]$ concentration at an average of 280 mg/L. The calculated $\text{SI}_{(\text{calcite})}$ is negative but close to the solution supersaturation with respect to calcite. These values are remained within the range of -4.6 to -4.9 for the duration of the experiment.

4. Discussion

None of both experiments could achieve supersaturation state with respect to calcite or magnesite. The SI remain negative but fluctuate as a function of the pH, the cation concentration and the total alkalinity. This confirms that saturation index depends on metal cations and CO_3^{2-} activities and further suggests the inability of basaltic glass dissolution to provide the necessary amount of cations required for solution supersaturation with respect to carbonates. The AFT experiment shows constant $[\text{Ca}^{2+}]$ and $[\text{Mg}^{2+}]$ concentrations. In this case an increase in the CO_3^{2-} activity would be required to achieve supersaturated conditions with respect to carbonate, this would be reflected in a positive value of the SI for these mineral phases. As the concentration of cations remained constant, a higher concentration of carbonate or a higher degree of bicarbonate deprotonation from a higher pH would be required. This means an enhanced deprotonation of bicarbonate HCO_3^- and the consumption of hydroxide ion (OH^-). This then explains the decreasing pH and the increasing total alkalinity as per eq.1. According to the previously mentioned definition of the carbonate alkalinity a pH higher than 8.3 would be required in order to induce bicarbonate ion deprotonation for production of CO_3^{2-} in significant concentrations. A higher pH could be reached by active cyanobacteria. Indeed, cyanobacteria consume bicarbonate ions by photosynthesis which leads to the higher pH values, the formation of carbonate (CO_3^{2-}) and a raising SI in the presence of sufficient metal cations for supersaturation. In the CFT case, increasing pH rates can be explained by the presence of dead cyanobacteria mass and their interaction of their deprotonated cell surface groups with dissolving CO_2 from air injection (Weber and Martinez, 2017). However, SI remains negative because of the near-neutral pH which prevents calcite precipitation as bio-mineralization needs a pH higher than 8.3 to commence the generation of carbonate ions.

It has been shown that active cyanobacteria have a protective mechanism against carbonate precipitation on their surface, which is missing for dead cyanobacteria implying that this is the consequence of the metabolic activity of these primary producers (Martinez *et al.*, 2010; Martinez *et al.*, 2008). Dead cyanobacteria cells or cellular debris may act as nucleation sites by binding Ca^{2+} and Mg^{2+} cations on their surface. This could explain why the SI results become less negative in the CFT experiment in presence of dead cells. Therefore, the presence of both active and dead cyanobacteria, as in the natural environment, contributes to controlling the mode of carbonate mineral precipitation, through the generation of an alkaline pH or due to the efficient sorption of metal cations leading to the nucleation of mineral phases (Martinez *et al.*, 2010; Martinez *et al.*, 2016; Weber and Martinez, 2017).

Another important factor which is needed for supersaturation is the metal cations concentrations. CFT results proved that high metal cations concentrations from an efficiently dissolving source, such as calcium chloride, lead to SI values which are closer to solution supersaturation with respect to carbonates. AFT [Ca^{2+}] and [Mg^{2+}] concentrations from basaltic glass dissolution were sufficient to achieve carbonate precipitation. This implies that the use of basaltic glass for carbonation and CO_2 mineralization would be inefficient if employed at large scale, requiring the use faster dissolving simple silicate minerals such as olivine, or finding a way to enhance the release of metal cations from simple and more complicated alumino-silicate minerals. The slight decrease in metal cation concentrations in these experiments, occasionally observed, can be explained by the need for more testing and development of the flow through reactor used, where a fast flow rate would cause dilution of metal cations. A slower pump rate would have to be used. Moreover, higher metal cations dissolution rates would have to be reached, however, another silicate mineral would need to be tested which can release metal cations at a faster rate. The background electrolyte concentration of 0.04 mol/L NaNO_3 could be modified in order to find the optimal concentration and optimize the dissolution rate and the

metal cation activities. Smaller basaltic glass particles can be used, this would produce and increase in mineral surface area which would lead to an increase in the basaltic glass dissolution rate.




Figure 7 Schematic interpretation of optimal conditions leading to cyanobacteria induced bio-mineralization in presence of basaltic glass.

Figure 7 presents all the parameters which could contribute to achieving supersaturation state with respect to carbonates and subsequent precipitation in the presence of cyanobacteria. Several points have to be improved in order to achieve carbonate nucleation and precipitation. *Synechococcus* cyanobacteria have to be grown in proper conditions with enough space and time to get to cell population saturation (at least 2 weeks). Active cells lead to an alkaline (high pH) environment by the photosynthesis process, and indirectly to the formation of carbonate CO_3^{2-} . The “flow through reactor” open system permits a constant CO_2 supply and other nutrients required for photosynthesis and cell growth. On another hand, basaltic glass has to be dissolved in optimal conditions so as to reach high aqueous $[\text{Ca}^{2+}]$ and $[\text{Mg}^{2+}]$ concentrations.

Providing all these conditions, calcite or magnesite precipitation could be expected in the flow through reactors. Using the rate model in Eq.11 an approximation of the total carbonate precipitation amount could be calculated for potential implementation of these systems at the larger scale. SEM and EDX analyses could then help to the analysis and identification of carbonates to confirm the presence of calcite or aragonite polymorphs or that of magnesite. By finding optimal conditions for cyanobacteria induced bio-mineralization, this method could be used to stock significant quantities of CO₂ at a large scale.

Acknowledgments

The authors would like to thank the Institut Polytechnique UniLaSalle, for funding part of this study. In addition the authors would like to thank Ms. Sigrid Hirth-Walther of the Institute of Geosciences at the University of Freiburg for F-AAS measurements. Furthermore, the authors would like to extend their gratitude to Dr. Christian Grimm and Dr. Eric H. Oelkers of the GET-CNRS in Toulouse, France, for providing the basaltic glass samples used in this study.

References

- Dessert, C., Dupré, B., Gaillardet, J., François, L. M. and Allègre, C. J. (2003) Basalt weathering laws and the impact of basalt weathering on the global carbon cycle. *Chem. Geol.* **202**, 257-273.
- Fridlyand, L. E. and Scheibe, R. (1999) Regulation of the Calvin cycle for CO₂ fixation as an example for general control mechanisms in metabolic cycles. *Biosystems.* **51**, 79-93.
- Gerlach, T. (2011) Volcanic versus anthropogenic carbon dioxide. *Eos, Transactions American Geophysical Union.* **92**, 201-202.

- Grimm, C., Martinez, R. E., Pokrovsky, O. S., Benning, L. G. and Oelkers, E. H. (2019) Enhancement of cyanobacterial growth by riverine particulate material. *Chem. Geol.* **525**, 143-167.
- Harris, D. C. (2010) *Quantitative Chemical Analysis, Eight Edition*. MacMillan Publishing
- Jansson, C. and Northen, T. (2010) Calcifying cyanobacteria—the potential of biomineralization for carbon capture and storage. *Current Opinion in Biotechnology.* **21**, 365-371.
- Langdon, C., Takahashi, T., Sweeney, C., Chipman, D., Goddard, J., Marubini, F., Aceves, H., Barnett, H. and Atkinson, M. J. (2000) Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef. *Global Biogeochem. Cy.* **14**, 639-654.
- Martinez, R. E., Gardés, E., Pokrovsky, O. S., Schott, J. and Oelkers, E. H. (2010) Do photosynthetic bacteria have a protective mechanism against carbonate precipitation at their surfaces? *Geochim. Cosmochim. Acta.* **74**, 1329-1337.
- Martinez, R. E., Pokrovsky, O. S., Schott, J. and Oelkers, E. H. (2008) Surface charge and zeta-potential of metabolically active and dead cyanobacteria. *J. Colloid Interface Sci.* **323**, 317-325.
- Martinez, R. E., Weber, S. and Bucher, K. (2014) Quantifying the kinetics of olivine dissolution in partially closed and closed batch reactor systems. *Chem. Geol.* **367**, 1-12.
- Martinez, R. E., Weber, S. and Grimm, C. (2016) Effects of freshwater *Synechococcus* sp. cyanobacteria pH buffering on CaCO₃ precipitation: Implications for CO₂ sequestration. *Appl. Geochem.* **75**, 76-89.
- Obst, M., Wehrli, B. and Dittrich, M. (2009) CaCO₃ nucleation by cyanobacteria: laboratory evidence for a passive, surface-induced mechanism. *Geobiology.* **7**, 324-347.

- Pechukas, P. (1981) Transition State Theory. *Annual Review of Physical Chemistry*. **32**, 159-177.
- Price, G. D., Badger, M. R., Woodger, F. J. and Long, B. M. (2008) Advances in understanding the cyanobacterial CO₂-concentrating-mechanism (CCM): functional components, Ci transporters, diversity, genetic regulation and prospects for engineering into plants. *Journal of Experimental Botany*. **59**, 1441-1461.
- Riding, R. (2006) Cyanobacterial calcification, carbon dioxide concentrating mechanisms, and Proterozoic–Cambrian changes in atmospheric composition. *Geobiology*. **4**, 299-316.
- Schlesinger, W. H. and Bernhardt, E. S. (2013) Chapter 5 - The Biosphere: The Carbon Cycle of Terrestrial Ecosystems. *Biogeochemistry (Third Edition)* (Schlesinger, W. H. and Bernhardt, E. S. eds.), 135-172, Academic Press.
- Schott, J., Pokrovsky, O. S., Spalla, O., Devreux, F., Gloter, A. and Mielczarski, J. A. (2012) Formation, growth and transformation of leached layers during silicate minerals dissolution: The example of wollastonite. *Geochim. Cosmochim. Acta*. **98**, 259-281.
- Schubert, W.-D., Klukas, O., Krauß, N., Saenger, W., Fromme, P. and Witt, H. T. (1997) Photosystem I of *Synechococcus elongatus* at 4 Å resolution: comprehensive structure analysis. *Journal of Molecular Biology*. **272**, 741-769.
- Spreitzer, R. J. and Salvucci, M. E. (2002) RUBISCO: Structure, Regulatory Interactions, and Possibilities for a Better Enzyme. *Annual Review of Plant Biology*. **53**, 449-475.
- Stumm, W. and Morgan, J. J. (1996) *Aquatic Chemistry*. Wiley Intersciences
- Weber, S. and Martinez, R. E. (2017) Effects of *Synechococcus* sp. cyanobacteria inert biomass on olivine dissolution: Implications for the application of enhanced weathering methods. *Appl. Geochem.* **84**, 162-172.