


Age	Ichnotaxon	State	Unit	Depositional environment	References
Lopingian	<i>Pachypes dolomiticus</i>	Italy	Arenaria di Val Gardena	marginal marine floodplain	Leonardi et al., 1975; Conti et al., 1977; Valentini et al., 2009; Marchetti et al., 2019a
Lopingian	<i>Pachypes loxodactylus</i>	Scotland	Cornockle	aeolian dune	McKeever and Haubold, 1996; Marchetti et al., 2019b
Lopingian	<i>Pachypes loxodactylus</i>	Scotland	Locharbriggs	aeolian dune	Dudgeon, 1878; McKeever and Haubold, 1996; Marchetti et al., 2019b
Lopingian	<i>Pachypes</i> isp.	Russia	Vyatkan horizon	continental floodplain	Surkov et al., 2007; Valentini et al., 2009
Lopingian	<i>Pachypes</i> 'primus'	Russia	Poldarsa	continental floodplain	Gubin et al., 2003; Valentini et al., 2009
Lopingian	cf. <i>Pachypes</i> isp.	Germany	Cornberg	aeolian dune	Schmidt, 1959; Marchetti et al., 2019b
Guadalupian/Lo pingian	<i>Pachypes</i> isp.	Morocco	T2 of Argana Basin	continental floodplain	Voigt et al., 2010
Guadalupian/Lo pingian	<i>Pachypes</i> isp.	Niger	Moradi	continental floodplain	Smith et al., 2015
Guadalupian/Lo pingian	aff. <i>Pachypes</i> isp.	Germany	Mammendorf horizon	continental floodplain	Buchwitz et al., 2017
Guadalupian/Lo pingian	<i>Pachypes?</i> isp.	Spain	Upper Red Unit	ephemeral lacustrine/playa-lake	Mujal et al., 2017
Guadalupian	<i>Pachypes ollieri</i>	France	Le Muy	continental floodplain	Demathieu et al., 1992; this work
Guadalupian	cf. <i>Pachypes</i> isp.	France	Pradineaux	continental floodplain	Gand et al., 1995; this work
Guadalupian	<i>Pachypes ollieri</i>	France	La Lieude	continental floodplain	Ellenberger, 1983a, 1983b; Gand et al., 2000; this work
Cisuralian	<i>Pachypes ollieri</i>	France	Rabejac	continental floodplain	this work
Cisuralian	<i>Pachypes ollieri</i>	Spain	Peranera (Lower Red Unit)	continental floodplain, meandering fluvial	Mujal et al., 2016; this work
Cisuralian	<i>Pachypes ollieri</i>	Texas, USA	San Angelo	continental floodplain	Lucas and Hunt, 2005; this work
Cisuralian	<i>Pachypes ollieri</i>	Arizona, USA	Hermit	continental floodplain	Marchetti et al., 2020; this work

Suppl. Table 1. *Pachypes* occurrences.



Suppl. Fig. 1. cf. *Pachypes* isp. Guadalupian, Pradineaux Formation, southern France. MASR 13, plaster cast of FS-SR 7. A, trackway section. B, enlargement of A, right pes-manus couple. C, false-colour depth map of B. D, left pes-manus couple. E, false-colour depth map of D. F, right pes-manus couple of the holotype of '*Lunaepes fragilis*', FS-SR 7. Roman numbers indicate the digit imprint number. Values in the black boxes refer to the preservation scale.

References

- BUCHWITZ, M., LUTHARDT, L., MARCHETTI, L., TROSTHEIDE, F., VOIGT, S. and SCHNEIDER, J.W. 2017. A Middle to Late Permian tetrapod tracksite from northern Germany. In: Bordy, E. (Ed.), 2nd Conference of Continental Ichnology, Nuy Valley, Western Cape, South Africa, 1st–4th October 2017. Abstract Book, p. 15.
- CONTI, M.A., LEONARDI, G., MARIOTTI, N. and NICOSIA, U. 1977. Tetrapod footprints of the Val Gardena Sandstone (North Italy). Their paleontological, stratigraphic and paleoenvironmental meaning. *Palaeontographia Italica*, **70**, 1–91.
- DEMATHIEU, G., GAND, G. and TOUTIN-MORIN, N. 1992. La palichnofaune des bassins permiens provençaux. *Geobios*, **25**, 19–54.
- DUDGEON, P. 1878. Note on a new fossil footprint from the Permian sandstone of Dumfriesshire. *Proceedings of the Royal Society of Edinburgh*, **9**, 154–155.
- ELLENBERGER, P. 1983a. Sur l'intérêt paléontologique de la dalle à pistes de la Lieude (commune de Mérifons, Hérault, France). *Société de la Protection de la Nature du Languedoc Roussillon*, **1**, 11–20.
- 1983b. Sur la zonation ichnologique du Permien moyen (Saxonien) du bassin de Lodève (Hérault). *Comptes-rendus des séances de l'Académie des sciences. Série 2, Mécanique-physique, chimie, sciences de l'univers, sciences de la terre*, **297**, 553–558.
- GAND, G., DEMATHIEU, G. and BALLESTRA, F. 1995. La palichnofaune de vertébrés tétrapodes du Permien supérieur de l'Esterel (Provence, France). *Palaeontographica Abteilung A*, **235**, 97–139.
- GARRIC, J., DEMATHIEU, G. and ELLENBERGER, P. 2000. La palichnofaune de vertébrés tétrapodes du Permien supérieur du bassin de Lodève (Languedoc-France). *Palaeovertebra*, **29**, 1–82.
- Gubin et al., 2003
- LEONARDI, P., CONTI, M.A., LEONARDI, G., MARIOTTI, N. and NICOSIA, U. 1975. *Pachypes dolomiticus* n. gen. n.sp.; Pareiasaur footprint from the 'Val Gardena Sandstone' (Middle Permian) in the western Dolomites (N. Italy). *Atti della Accademia Nazionale dei Lincei*, **57**, 221–232.
- LUCAS, S.G. and HUNT, A.P. 2005. Permian tetrapod tracks from Texas. *New Mexico Museum of Natural History and Science Bulletin*, **30**, 202–206.
- MARCHETTI, L., VOIGT, S. and KLEIN, H. 2019a. Revision of Late Permian tetrapod tracks from the Dolomites (Trentino-Alto Adige, Italy). *Historical Biology*, **31**, 748–783.
- VOIGT, S. and LUCAS, S.G. 2019b. An anatomy-based study of the Lopingian eolian tracks of Germany and Scotland reveals the first evidence of the end-Guadalupian mass extinction at low paleolatitudes of Pangea. *Gondwana Research*, **73**, 32–53.
- FRANCISCHINI, H., LUCAS, S.G., VOIGT, S., HUNT, A.P. and SANTUCCI, V.L. 2020. Paleozoic Vertebrate Ichnology of Grand Canyon National Park. In: Santucci, V.L. and Tweet, J.S. (Eds.), *Grand Canyon National Park: Centennial paleontological resource inventory (non-sensitive version)*. Natural Resource Report NPS/GRCA/NRR—2020/2103. National Park Service, Fort Collins, Colorado, p. 333–379.
- MCKEEVER, P.J. and HAUBOLD, H. 1996. Reclassification of vertebrate trackways from the Permian of Scotland and related forms from Arizona and Germany. *Journal of Paleontology*, **70**, 1011–1022.
- MUJAL, E., FORTUNY, J., OMS, O., BOLET, A., GALOBART, À. and ANADÓN, P. 2016. Palaeoenvironmental reconstruction and early Permian ichnoassemblage from the NE Iberian Peninsula (Pyrenean Basin). *Geological Magazine*, **153**, 578–600.

- FORTUNY, J., PÉREZ-CANO, J., DINARÈS-TURELL, J., IBÁÑEZ-INSA, J., OMS, O., VILA, I., BOLET, A. and ANADÓN, P. 2017. Integrated multi-stratigraphic study of the Coll de Terrers late Permian–Early Triassic continental succession from the Catalan Pyrenees (NE Iberian Peninsula): A geologic reference record for equatorial Pangaea. *Global and Planetary Change*, **159**, 46–60.
- SCHMIDT, H., 1959. Die Cornberger Fahrten im Rahmen der Vierfusler. *Entwicklung Abhandlungen des Hessischen Landesamtes für Bodenforschung*, **28**, 1–137.
- SMITH, R.M.H., SIDOR, C.A., TABOR, N.J. and STEYER, J.S. 2015. Sedimentology and vertebrate taphonomy of the Moradi Formation of northern Niger: a Permian wet desert in the tropics of Pangaea. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **440**, 128–141.
- SURKOV, M.V., BENTON, M.J., TWITCHETT, R. J., TVERDOKHLEBOV, V.P. and NEWELL, A.J. 2007. First occurrence of footprints of large therapsids from the Upper Permian of European Russia. *Palaeontology*, **50**, 641–652.
- VOIGT, S., NICOSIA, U. and CONTI, M.A. 2009. A re-evaluation of *Pachypes*, a pareiasaurian track from the Late Permian. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, **251**, 71–94.
- HMINNA, A., SABER, H., SCHNEIDER, J.W. and KLEIN, H. 2010. Tetrapod footprints from the uppermost level of the Permian Ikakern Formation (Argana basin, western High Atlas, Morocco). *Journal of African Earth Sciences*, **57**, 470–478.