



CODEN [USA]: IAJPBB

ISSN : 2349-7750

## INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Research Article

### MANUAL INSTRUMENTATION AND ROTARY TECHNIQUE IN TERM OF INSTRUMENTATION TIME AND QUALITY OF ROOT CANAL OBTURATION IN SINGLE ROOTED TEETH

<sup>1</sup>Dr Muhammad Hadee Aziz, <sup>2</sup>Dr Roheen Fatima

<sup>1,2</sup> Rawal Institute of Health Sciences, Islamabad

**Article Received:** October 2020      **Accepted:** November 2020      **Published:** December 2020

**Abstract:**

**Objective:** Root canal obturation is an essential stage of root canal treatment aimed to seal the root canal in order to prevent future bacterial contamination/ recontamination of the canal space. Aim of this study was to compare the outcomes of manual procedure with rotary technique in term of quality of root canal obturation in patients presented with single rooted teeth.

**Study Design:** This was a randomized controlled trial study.

**Place and Duration of Study:** This study was conducted at the Rawal Institute of Health Sciences, Islamabad, for the duration of six months from January 2020 to July 2020.

**Materials and Methods:** Eighty teeth of male and female patients with ages 20 to 60 years were enrolled and divided equally into two groups. Group I consist of 40 teeth and rotary method was applied. Group II with 40 teeth and manual instrumentation was done. Post obturation radiographs were done to examine the difference in length, density and taper of root canal filling by using T-score. Time taken for instrumentation and canal filling was recorded and compare the findings between both groups. SPSS version 24.0 was used for data analysis.

**Results:** There were 50 (62.5%) females and 30 (37.5%) male patient's teeth with mean age  $32.25 \pm 11.68$  years. We found a significant difference in term of obturation quality between both groups with p-value 0.008. In group I, 15 (37.5%) patients had T-score 2 and 20 (50%) had T-score 3 while in group II, 13 (32.5%) patients had T-score 2 and 8 (20%) patients had T-score 3, a significant difference was observed between both techniques with p-value  $< 0.05$ . Instrumentation time was higher in group II as compared to group I (20.2 min Vs 10.6 min) with p-value  $< 0.05$ .

**Conclusion:** Rotary method in term of quality of root canal obturation was better as compared to manual technique.

**Key Words:** Root canal, Rotary technique, Manual method, Obturation quality

**Corresponding author:**

**Dr Muhammad Hadee Aziz,**

Rawal Institute of Health Sciences, Islamabad

QR code



Please cite this article in press Muhammad Hadee Aziz et al, *Awareness And Knowledge Of Nursing Care In Ent Disorders In Pakistan.*, Indo Am. J. P. Sci, 2020; 07(12).

## INTRODUCTION:

Root canal obturation is an essential stage of root canal treatment aimed to seal the root canal in order to prevent future bacterial contamination/recontamination of the canal space.<sup>1</sup> Many obturation methods have been introduced over the years, each attempting to provide a better seal of the root canal.<sup>2</sup> All have in common the assumption that the root canal is properly cleaned and shaped before the obturation stage. It is assumed by all that if the root canal is not adequately prepared and if tissue remnants and debris are present along the walls, proper sealing may be jeopardized, even with the best root canal filling method.<sup>3,4</sup> When simple, narrow, straight root canals with round cross-sections are considered, most current rotary nickel-titanium file systems will adequately clean and shape the canal with favorable results. The case is different in oval, flat, or curved root canals. In flat root canals, rotary file systems often fail to adequately clean and shape the canal, leaving “fins” that may have not been prepared.<sup>2-4</sup> In such a case, even warm gutta-percha obturation methods will fail to adequately seal the root canal (4). Clinical mesiodistal radiographs will fail to detect such discrepancy. Quality of obturation is one of the characteristic determinants in the prognosis of root canal treatment. One of the ways to judge the quality of endodontic treatment is by periapical radiographic evaluation which is the most common method used for assessment so far. Radiographic quality of the endodontic treatment can be evaluated on the basis of three parameters which include length, homogeneity and taper of the root canal filling visible on radiographs.<sup>5,6</sup>

Although several researches have been conducted among the undergraduates, graduates and postgraduates for the evaluation of the obturation quality using different methods of canal preparation (manual/rotary), but the results are quite variable.<sup>7-10</sup>

## MATERIALS AND METHODS:

This study was conducted at the Rawal Institute of Health Sciences, Islamabad, for the duration of six months from January 2020 to July 2020. A total of 80 patients of both genders with ages 20 to 60 years required root canal treatment for single rooted were included in this study. Patients detailed demographic were recorded after taking informed written consent.

Patients with multi-rooted teeth, patients with apical pathology and those root canals with curvature more than 30 degrees were excluded. All the patients selected from OPD were randomly divided into two groups using computer generated randomization scheme. Group I consist of 40 teeth and rotary method (Universal Protaper Niti files, Dentsply Maillefer) followed by F1, F2, or F3 Gutta Percha (Dentsply Maillefer) was applied. Group II with 40 teeth and manual instrumentation (circumferential filing technique) with K and H files (Mani, Japan) followed by cold lateral condensation technique was done. The sealer was calcium hydroxide based (Apexit-plus, Ivoclar Vivadent AG, Germany) and it was same for both groups. Post procedure intraoral periapical radiograph with paralleling device was done to examine the length, density and taper of root canal filling. T-score scoring system was applied, 0 score for inadequate and 1 for adequate. Patients with all three parameters were adequate marked as score 3, patients with two parameters were adequate marked as score 2, patients with any one parameter were adequate marked as score 1 and those with none of parameter was adequate marked as score 0. Time taken for instrumentation and canal filling was recorded. All procedures were done by same operator to reduce operator related bias. The periapical radiographs were evaluated by two different operators separately who were blinded to the procedure type. The intra observer differences were not significant. Data was analyzed by SPSS 24. Chi square test was done to compare the T-score and instrumentation time between both groups with p-value <0.05 was taken as significant.

## RESULTS:

In Group I, 28 (70%) patients were females and 12 (30%) were males with mean age  $31.95 \pm 10.64$  years and in group II, 22 (55%) patients were females and 18 (45%) patients were males with mean age  $30.38 \pm 9.45$  years. No significant difference was observed between both groups regarding age and gender (Table 1). According to the instrumentation time taken it was higher in group II as compared to group I (20.2 min Vs 10.6 min) with p-value <0.05. Mean filling time was also higher in group II as compared to group I (3.25 min Vs 1.6 min) with p-value <0.05 (Table 2).

**Table No.1: Age and gender wise distribution between both groups**

Variable	Group I	Group II	P-value
Age (years)	31.95±10.64	30.38±9.45	0.07
<b>Gender</b>			
Male	12 (30%)	18 (45%)	N/S
Females	28 (70%)	22 (55%)	N/S

**Table No.2: Comparison of instrumentation time and canal filling time between both groups**

Variable	Group I	Group II	P-value
Instrumentation Time	10.6	20.2	0.001
Filling Time	1.6	3.2	0.01

**Table No.3: Comparison of length, density and taper of root canal filling between both groups**

Variable	Group I	Group II	P-value
<b>Length</b>			
Adequate	31 (77.5%)	30 (75%)	N/S
Inadequate	9 (22.5%)	10 (25%)	
<b>Density</b>			
Adequate	34 (85%)	28 (70%)	N/S
Inadequate	6 (15%)	12 (30%)	
<b>Taper</b>			
Adequate	34 (85%)	11 (27.5%)	0.0001
Inadequate	6 (15%)	29 (72.5%)	

**Table No. 4: Quality of obturation regarding T-score between both groups**

T-score	Group I	Group II	P-value
3	20 (50%)	8 (20%)	0.0001
2	15 (37.5%)	13 (32.5%)	
1	5 (12.5%)	16 (40%)	
0	0	3 (7.5%)	

According to the post obturation quality of root canal we found that 31 (77.5%) in group I and 30 (75%) patients in group II showed adequate length of root canal filling while 9 (22.5%) and 10 (25%) patients had inadequate in group I and II. No significant difference was observed regarding length of root canal filling(RFC) between both groups with p-value 0.2. No significant difference was observed regarding density of RCF between both groups (p-value >0.05), in group I 34 (85%) patients and in group II 28 (70%) patients were adequate while 6 (15%) and 12 (30%) patients showed inadequacy in group I and II. We found a significant difference regarding taper of root canal filling between both groups with p-value 0.0001 (34 (85%) in group I and 11 (27.5%) in group II had adequate findings while 6 (15%) and 29 (72.5%) had inadequacy in group I and II (Table 3). In group I, 15 (37.5%) patients had T-score 2, 20 (50%) had T-score 3, 5 (12.5%) had T-score 1 and none of patient had T-score 0. In group II 13 (32.5%) patients had T-score 2, 8 (20%) patients had T-score 3, 16 (40%) had score 1 and 3 (7.5%) had score 0. A significant difference was observed between both groups regarding T-score with p-value 0.01 (Table 4)

## DISCUSSION:

In present study 80 patients of both genders were enrolled to compare the outcomes of rotary method with manual K and H file instrumentation in term of quality of obturation. There were 50 (62.5%) female and 30 (37.5%) male patient's teeth with mean age  $32.25 \pm 11.68$  years. These results were similar to the study by Jalees et al<sup>11</sup> regarding comparison of rotary procedure versus manual method and they reported female patients were high in numbers 56.67% as compared to males and average age of patients was  $33.3 \pm 7.4$  years in group I and  $37.6 \pm 12.9$  years in group II.

In present study we found that patients treated with rotary method had less instrumentation time as compared to manual k-file technique with p-value <0.001. A study conducted by Babaji et al<sup>16</sup> reported that manual technique taking higher instrumentation time as compared to rotary method. They reported a significant difference between both techniques with p-value <0.05. In our study we found no significant difference was observed regarding length of root canal filling between both groups with p-value 0.2. 31 (77.5%) in group I and 30 (75%) patients in group II showed adequate length of root canal filling while 9 (22.5%) and 10 (25%) patients had inadequate in

group I and II. We found no significant difference was observed regarding density of RCF between both groups ( $p$ -value  $>0.05$ ), in group I, 34 (85%) patients and in group II 28 (70%) patients were adequate. These results were similar to many of previous study in which no significant difference was observed between rotary and manual method regarding length of Root canal filling and density of RCF.<sup>12-14</sup>

We found a significant difference regarding taper of root canal filling between both groups with  $p$ -value 0.0001 (34 (85%) in group I and 11 (27.5%) in group II had adequate findings while 6 (15%) and 29 (72.5%) had inadequacy in group I and II). These results were similar to the study by Jalees et al<sup>11</sup>, in which they reported a significant difference regarding taper of RCF between both methods with  $p$ -value  $<0.05$ . Many of other studies showed significant improvement regarding taper of root canal filling after applying rotary methods and manual technique. These studies were reported that rotary method was much better and effective as compared to manual technique.<sup>15-17</sup>

In present study we used scoring system (T-score) to compare the quality of root canal obturation between both procedures and we found a significant difference between both procedures with  $p$ -value 0.0001. We found that 87.5% patients who received rotary method had T-score 2 and 3 and none of patient had score 0 while in patients whom were received manual technique 13 (32.5%) patients had T-score 2, 8 (20%) patients had T-score 3, 16 (40%) had score 1 and 3 (7.5%) had score 0. These results were similar to several previous study in which rotary method showed better quality of root canal obturation as compared to manual technique.<sup>18-21</sup> A study by Samady et al<sup>22</sup> reported rotary method had better obturation quality as compared to manual K-files technique.

### CONCLUSION:

Rotary method in term of quality of root canal obturation was better as compared to manual technique. We found no significant difference regarding length and density of root canal filling between both procedures however, regarding taper of RCF a significant better result was observed in rotary method as compared to manual technique.

### REFERENCES:

1. Walton RE, Torabinejad M. Principles and Practice of Endodontics. 3<sup>rd</sup> ed. Philadelphia, Pa; London: Saunders Company; 2002.p.222.
2. Azar MR, Safi L, Nikaein A. Comparison of the cleaning capacity of Mtwo and pro taper rotary

- systems and manual instruments in primary teeth. Dent Res J (Isfahan) 2012;9:146-51.
3. Madan N, Rathnam A, Shigli AL, Indushekar KR. K-file vs. ProFiles in cleaning capacity and instrumentation time in primary molar root canals: *Anin vitro* study. J Indian Soc Pedod Prev Dent 2011;29:2-6.
4. Schafer, E, Erler M, Dammaschke T. Comparative study on the shaping ability and cleaning efficiency of rotary Mtwo instruments. Part b. Cleaning effectiveness and shaping ability in severely curved root canals of extracted teeth. Int Endod J 2016; 39(3): 203-12.
5. Azar SN. Cleaning ability of primary root canals by M two and Pro Taper rotary systems and manual instruments in primary teeth. Dent Res J 2012; 9(2): 146-151.
6. Alvesa RAA, Souzaa JB, GonçalvesAlencara AH, Pécoraa JD, Estrela C. Detection of procedural errors with stainless steel and NiTi instruments by undergraduate students using conventional radiograph and cone beam computed tomography. Iran Endod J 2013; 8(4): 161-5.
7. AbuMostafa A, Ahmed I, Alenezy G, Alzoman H. Quality of root canal filling performed by undergraduate students in Saudi Dental College. J Dent Oral Hyg 2015; 7(5): 64-70.
8. Makarem A, Ravandeh N, Ebrahimi M, Radiographic assessment and chair time of rotary instruments in the pulpectomy of primary second molar teeth: a randomized controlled clinical trial. J Dent Res Dent Prospects 2014; 8(2): 84-9.
9. Zinge PR, Patil J. Comparative evaluation of effect of rotary and reciprocating single-file systems on pericervical dentin: A cone-beam computed tomography study. J Conserv Dent 2017;20(6): 424-8.
10. Román-Richon S, Faus-Matoses V, Alegre-Domingo T, Faus- Llácer VJ. Radiographic technical quality of root canal treatment performed ex vivo by dental students at Valencia University Medical and Dental School, Spain. Medicina Oral, Patología Oral Cirugía Bucal 2014;19(1): 93-7.
11. Jalees M, Noor N, Javed K, Qamar R, Abbasi S, Ahmad M. Comparison of quality of root canal obturation in single rooted teeth prepared by manual and rotary method. Pak Oral Dent J 2018; 38(3).
12. Govindaraju L, Jeevanandan G, Subramanian E. Clinical evaluation of quality of obturation and instrumentation time using two modified rotary file systems with manual instrumentation in primary teeth. J Clin Diagn Res 2017; 11(9): 55-8.

13. Zinge PR. Comparative evaluation of effect of rotary and reciprocating single-file systems on pericervical dentin: a cone-beam computed tomography study. *J Conserv Dent* 2017;20(6): 424-8.
14. Arun Kumar S, Lanke RB. An invitro-comparative evaluation of cleaning efficacy of hyflex and hero shaper rotary niti file systems scanning electron microscope study. *Int J Recent Sci Res* 2015;6(9): 6320- 24.
15. Edionwe JI, Shaba OP, Umesi DC. Single visit root canal treatment: A prospective study. *Niger J Clin Pract* 2014;17:276-81
16. Babaji P, Mehta V, Manjooran T. Clinical evaluation of rotary system over manual system in deciduous molars: A clinical trial. *Int J Pedod Rehabil* 2019;4:13-6.
17. Bane K, Faye B, Sarr M, Niang SO, Ndiaye D, Machtou P. Root canal shaping by single-file systems and rotary instruments : a laboratory study. *Iran Endod J* 2015; 10(2): 135-9.
18. Stavileci M, Hoxha V, Görduysus Ö, Tatar I, Laperre K, Hostens J, et al. Evaluation of root canal preparation using rotary system and hand instruments assessed by micro-computed tomography. *Med Sci Monit Basic Res* 2015; 21: 123-30.
19. Schäfer E, Schulz-Bongert U, Tulus G. Comparison of hand stainless steel and nickel titanium rotary instrumentation: a clinical study. *Int Endod J* 2004; 30: 432-5.
20. Azar MR, Mokhtare M. Rotary M two system versus manual K-file instruments: Efficacy in preparing primary and permanent molar root canals. *Ind J Dent Res* 2011;22:363.
21. Guelzow A, Stamm O, Martus P, Kielbassa AM. Comparative study of six rotary nickel-titanium systems and hand instrumentation for root canal preparation. *Int Endod J* 2005;38:743-52.
22. Samadi F, Jaiswal JN, Saha S, Garg N, Chowdhary S, Samadi F, et al. A comparative evaluation of efficacy of different obturation techniques used in root canal treatment of anterior teeth: an in vitro study. *Int J Clin Pediatr Dent* 2014; 7(1): 1-5.