



CODEN [USA]: IAJPBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF  
**PHARMACEUTICAL SCIENCES**

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Research Article

## ASSESSMENT OF INFLUENCE MORBIDITY AND MORALITY RATE OF FOURNIER'S GANGRENE AND ETIOLOGICAL PREDISPOSING ASPECTS

<sup>1</sup>Dr Ali Hussain Bokhari, <sup>2</sup>Dr Muhammad Hashim, <sup>3</sup>Dr Asma Farooq

<sup>1</sup>Akhter Saeed Trust Hospital Lahore, <sup>2</sup>Akhter Saeed Trust Hospital Lahore,

<sup>3</sup>Lahore General Hospital, Lahore.

**Article Received:** October 2020

**Accepted:** November 2020

**Published:** December 2020

**Abstract:**

**Objective:** To confirm the FG (Fournier's Gangrene) over duration of past eight years and to evaluate the prompting aspects. Which can influence mortality rate was the main objective of this research work. **Methodology:** We retrospectively reviewed the anatomical spot of the infective gangrene, prompting and etiological Aspects and outcomes. Total 65 patients who got admission in the Emergency Surgical Department of Lahore General Hospital, Lahore suffering from FG were the participants of this research work. This research work started from December 2010 and lasted up to December 2018.

**Results:** In this research study, there were total eight females and fifty seven males. Mean age of the male patients was  $51.0 \pm 13.90$  with a range from 19 to 75 years and mean age of the females was  $63.0 \pm 10.50$  with a range of age from 52 to 76 years. Mean duration of hospitalization was  $9.20 \pm 6.60$  with a range of age from 5 to 25 days. Most common comorbidity was DM (Diabetes Mellitus) and most common etiology was the perianal abscess. We performed colostomy in eleven patients; two patients underwent orchiectomy and cystostomy in only 2 patients. It is much important to note that all of eight patients who died because of FG were also suffering from DM and they were from low socioeconomic class. There was requirement of more than single debridement in 6 patients who died of FG.

**Conclusions:** There is very high rate of mortality of FG, which is a serious surgical emergency. Low socioeconomic condition, DM and greater than single debridement perform main role in the rate of morbidity as well as mortality.

**Key Words:** Surgery, outcome FG, debridement, prompting, emergency, diabetes mellitus.

**Corresponding author:**

**Dr. Ali Hussain Bokhari,**

Akhter Saeed Trust Hospital Lahore.

QR code



Please cite this article in press Ali Hussain Bokhari et al, Assessment Of Influence Morbidity And Morality Rate Of Fournier's Gangrene And Etiological Predisposing Aspects., Indo Am. J. P. Sci, 2020; 07(12).

**INTRODUCTION:**

In this research work, we assessed the patients who got admission in the Emergency Surgery Department of our institute because of FG in terms of site of infective gangrene, prompting and etiological Aspects and final outcome. FG is a fulminate necrotizing contagion of periurethral and perianal tissues that can disseminate even at the tissues of abdomen or thigh following planes of the penis & dartos fascia of scrotum, colle's fascia and scarpa's fascia [1]. Most important predisposing Aspects are increases age, low social and economic class, abscess, primary genitourinary infection, leakage of urine, DM, local trauma, perineal surgery, alcohol abuse and immune-suppression [2-4]. The patients who are present with adverse condition of general health, are highly prone to this complication. These poor conditions include obesity or malnutrition, diseases of liver and kidney and some other condition which are the main cause of immune-suppression [5,6]. The cultures of wound generally yield variety of organisms, implicating anaerobic and aerobic synergy [7].

**METHODOLOGY:**

Review of these clinical records included age of patient, sex, etiological agents, predisposing Aspects, period from start of symptoms and first debridement, total amount of surgical interventions, clinical outcome, culture findings and total stay in the hospital. With hindsight it is assessed that the hospital files of sixty five patients from December 2010 to December 2018. We calculated the 90-day rate of mortality. FG's diagnosis was depending upon the clinical signs and symptoms as necrosis, swelling, erythema, crepitus and rash in regions of perianal, perineal or genital. There was collection of samples of wound and healthy tissue with swab margin. Once, there was diagnosis of FG by physical checkup, all the patients underwent treatment with parenteral broad spectrum triple antimicrobial agents, utilizing penicillin G, metronidazole, and amino glycoside and obtained the hemodynamic support at the time of requirement. All the patients experienced extensive debridement under SA (Spinal Anesthesia) or GA (General Anesthesia). Debridement of the necrotic areas carried out and the removal of the necrotic tissues carried out in different stages so that bleeding tissues must be apparent.

Cleaning of the wound carried out with betadine, saline and 2.0% oxygenated water and then covering of the wound carried out with the nitrofurazone dressing. There was conduction of this method each other day. Cultures of tissues were gathered at regularly at debridement time for the identification of the microorganisms which are main causative agents. There was accomplishment of cystostomy in the patients suffering from urethral or/and penile pathology. There was accomplishment of colostomy in the patients present with fecal contamination of severe nature. SPSS V.20 was in use for the statistical analysis of the collected information. Various risk aspects that are able to have impact on disease prognosis and demographic traits, were compared according to the state of survival. We used T-test for the comparisons. P value of less than 0.050 was significant.


**RESULTS:**

Statistically, we found a significant difference between sex and age of the patients ( $P < 0.050$ ). There were total sixty five patients who were diagnosed with suffering from FG. There were 12.30% (n: 8) female and 87.70% (n: 57) male patients. Mean age of the male patients was  $51.0 \pm 13.90$  with a range from 19 to 77 years. Mean age of females was  $63.0 \pm 10.50$  years with a range of age from 52 to 76 years. Average duration from the start of symptoms to the time of admission was  $3.740 \pm 2.090$  days with a range from 1 to 10 days. Furthermore, average duration of hospitalization was  $9.20 \pm 6.60$  days with a range from 5 to 25 days. There was development of sepsis in 20.0% (n: 13) patients. Average debridement number was 2.50. There was requirement of greater than single debridement in 18.40% (n: 12) patients and six patients among them died. The average age of the 12.30% (n: 8) patients who met their death was  $57.83 \pm 6.70$  years, whereas average age of the survivors was  $51.67 \pm 14.90$  years ( $P = 0.340$ ). There was not much difference in the mean stay at hospital for dead and survivors ( $7.0 \pm 1.70\%$  vs  $10.90 \pm 4.70$ ,  $P = 0.060$ ). Most frequent factor of etiology was anorectal conditions followed by urogenital abnormalities and trauma. There was no observance of etiologic aspects for complication of FG 20.0% (n: 13) patients and the classification of these patients carried out as idiopathic FG. FG's etiology and predisposing agents are present in Table-1.

**Table-I: Etiology of FG and predisposing aspects in patients.**

Etiological Aspects	DM	MALG	CARD	COPD	UD	Total
Ischiorectal abscess	3.0	1.0	-	1.0	2.0	7.0
Posthemorrhoidectomy	1.0	-	-	-	1.0	2.0
Prostatik biopsy	1.0	-	-	-	1.0	2.0
Urethral stricture	1.0	-	1.0	-	1.0	3.0
Perineal trauma	3.0	2.0	1.0	-	3.0	9.0
Scrotal abscess	4.0	1.0	-	-	-	5.0
Fistula to the rectum	1.0	1.0	-	1.0	-	3.0
Sinuses to the skin	-	-	-	-	1.0	1.0
Perianal abscess	8.0	3.0	1.0	1.0	3.0	16.0
Perineal soft tissue infection	2.0	-	1.0	1.0	-	4.0
Idiopathic	-	1.0	3.0	8.0	1.0	13.0
Total	29.0	8.0	7.0	12.0	13.0	65.0

DM: Diabetes mellitus, MALG: Malignancy, CARD: Cardiac disorders, COPD: Chronic Obstructive pulmonary disease, UD: Undefined



Reconstructive surgical intervention carried out in eleven patients after the treatment of FG. There was no use of vacuum assisted closure in any patient. The prevalence of DM in the patients of this research work was 44.60%. It is important to note that the patients who were died, they were also suffering from DM. We obtained the culture results of thirty six patients. The most common cultured bacterial organisms from the surface of wound were Escherichiacoli(n: 9, 25.0%), anaerobic species of Streptococcus(n:4, 11.10%), Staphylococcus(n:4, 11.10%),

Enterobacterspp. (n:2, 5.60%),Bacteroides(n:2, 5.60%), P. aeruginosa(n:2,5.60%), Proteus(n:2, 5.60%), Clostridia(n:2, 5.60%) and mixed organisms in nine cultures (25.0%).There was application of multiple resection methods in 80.0% (n: 52) patients. Colostomy carried out in eleven patients, diverting to cystostomy in only two patients.

#### DISCUSSION:

In this recent research work, Bacterial over growth can create nitrogen & oxygen that gather in the tissues

resulting crepitation [6]. Most of these patients have more than twenty year of age [7,8 and 9]. Some research work revealed the dominance of male gender in FG as compared to the female gender [2,10]. There was dominance of male gender (87.70%). There are few research works in this field which showed the FG in pediatric patients [11]. There was no pediatric patient in this research work. Urgent surgery, hyperbaric oxygen and antibiotic treatment are the best options for management [12]. Sterile dressing was in use for the open wounds after debridement [13]. In one other research work, there was comparison of the effectiveness of the management of wound with the povidoneiodine dressing and Dakin's solution which has high antimicrobial effectiveness against microorganisms. Various authors discovered that hospitalization period was much lower in the patients who underwent management by Dakin's solution in comparison with the iodine dressing [14].

Honey also has the ability to prevent microbial treatment in accordance with some studies [15]. The rate of mortality of this complication is very high regardless of the advances in the facilities of health care for FG. The range of rate of mortality of this research work is from 4.0% to 80.0% [2,6,8, and 16]. The rate of mortality in this research work is 12.30%. Some authors stated that there was no association of DM with the high rate of morbidity as well as mortality [17]. High duration of hospitalization, high debridement numbers lead to high rate of morbidity as well as mortality [18, 19]. In this research work, anorectal diseases were most important reason (n:33, 50.80%), followed by idiopathic (n: 13, 20.0%) and some others which are in consistent with other research works [20, 21].

*E. coli* has been stated to be the most frequent isolated organism from cultures of wound (43.0% to 80.0%).

There is less isolation of the anaerobes than expectations which can be the result of technical faults [22,23]. There are rare reports about some other isolated microorganisms as *Candida Albicans* & *Lactobacillus Gasseri* [12,24, and 25]. There is requirement of colostomy to reduce the fecal contamination [26,27] but in some situations, there is need of cystostomy [27]. In one research work conducted by Ayan, he reviewed the records of forty one patients retrospectively, he carried out bilateral orchidectomy in 9.70% (n: 4) patients & unilateral orchidectomy in 12.10% (n: 5) patients because of necrosis [28].

## CONCLUSIONS:

The standard treatment is aggressive debridement and use of antibiotics but still the rate of mortality is very high. FG is much severe surgical emergency. DM, Low social as well as economic status and greater than single debridement perform a main role in increasing the rate of morbidity as well as mortality.

## REFERENCES:

1. Eke N. Fournier's gangrene: a review of 1726 cases. *Br J Surg.* 2000;87:718-728. doi: 10.1046/j.1365-2168.2000.01497.x
2. Canbaz H, Caglikulekci M, Altun U, Dirlik M, Turkmenoglu O, Taşdelen B, et al. Fournier's gangrene: analysis of risk Aspects affecting the prognosis and cost of therapy in 18 cases. *Ulus TravmaAcilCerrahiDerg.* 2010;16(1):71-76.
3. Villanueva-Saens E, Martinez Hernandez-Magro P, Valdes Ovalle M, et al. Experience in management of Fournier's gangrene. *Tech Coloproctol.* 2002;6(1):5-10. doi: 10.1007/s101510200001
4. Altunoluk B, Resim S, Efe E, Eren M, Can B, Kankilic N, et al. Fournier's gangrene: conventional dressings versus dressings with Dakin's solution. *ISRN Urol* 2012; 762340. doi: 10.5402/2012/762340
5. Haidari M, Nazer MR, Ahmadinejad M, Almasi V, Khorramabadi MS, Pournia Y. Honey in the treatment of Fournier's gangrene as an adjuvant: a cross sectional study. *J Pak Med Assoc.* 2014;64(5):571-573.
6. Sugihara T, Yasunaga H, Horiguchi H, Fujimura T, Ohe K, Matsuda S, et al. Impact of surgical intervention timing on the case fatality rate for Fournier's gangrene: an analysis of 379 cases. *BJU Int.* 2012;110(11 Pt C):E1096-1100. doi: 10.1111/j.1464-410X.2012.11291.x
7. Chawla SN, Gallop C, Mydlo JH. Fournier's Gangrene: An Analysis of Repeated Surgical Debridement. *Eur Urol.* 2003;43:572-575. doi:10.1016/S0302-2838(03)00102-7
8. Rouzrokh M, Tavassoli A, Mirshemirani A. Fournier's Gangrene in Children: Report on 7 Cases and Review of Literature. *Iran J Pediatr.* 2014;24(5):660-661.
9. Sánchez Mazzaferri F, FadilIturralde JL, Provenzal O, Damiani H, Milman A, Bragagnolo J, et al. Fournier's gangrene. Our experience over 10 years. A review of the literature. *Arch Esp Urol.* 1999;52(7):721-727.
10. Benjellounel B, Souiki T, Yakla N, Ousadden A, Mazaz K, Louchi A, et al. Fournier's gangrene: our experience with 50 patients and analysis of

- Aspects affecting mortality. *World J Emerg Surg.* 2013;8(1):13. doi: 10.1186/1749-7922-8-13
11. Clayton MD, Fowler JE, Sharifi R, Pearl RK. Causes, presentation and survival of fifty-seven patients with necrotizing fasciitis of the male genitalia. *SurgGynecol Obstet.* 1990;170(1):49–55.
  12. Hejase MJ, Simonin JE, Bihrl R, Coogan CL. Genital Fournier's gangrene: Experience with 38 patients. *Urology.* 1996;47(5):734-739. doi:10.1016/S0090-4295(96)80017-3
  13. Baskin LS, Carroll PR, Cattolica EV, McAninch JW. Necrotising soft tissue infections of the perineum and genitalia: Bacteriology, treatment and risk assessment. *Br J Urol.* 1990;65:524-529.
  14. Chennamsetty A, Khourdaji I, Burks F, Killinger KA. Contemporary diagnosis and management of Fournier's gangrene. *TherAdv Urol.* 2015;7(4):203-215. doi: 10.1177/1756287215584740
  15. Koukouras D, Kallidonis P, Panagopoulos C, Al-Aown A, Athanasopoulos A, Rigopoulos C, et al. Fournier's gangrene, a urologic and surgical emergency: presentation of a multi-institutional experience with 45 cases. *Urol Int.* 2011;86(2):167-172. doi: 10.1159/000321691
  16. Tleyjeh I M, Routh J, Qutub MO, Lischer G, Liang KV, Baddour LM. *Lactobacillus gasseri* causing Fournier's gangrene. *Scand J Infect Dis.* 2004;36:501-503.
  17. McCormack M, Valiquette AS, Ismail S. Fournier's gangrene: A retrospective analysis of 26 cases in a Canadian hospital and literature review. *Can UrolAssoc J.* 2015;9(5-6):E407-410. doi: 10.5489/cuaj.2445
  18. Ruiz-Tovar J, Córdoba L, Devesa JM. Prognostic Aspects in Fournier gangrene. *Asian J Surg.* 2012;35(1):37-41. doi: 10.1016/j.asjsur.2012.04.006
  19. Yaghan RJ, Al-Jaberi TM, Bani-Hani I. Fournier's gangrene: changing face of the disease. *Dis Colon Rectum.* 2000;43:1300-1308.
  20. Johnin K, Nakatoh M, Kadowaki T, Kushima M, Koizumi S, Okada Y. Fournier's gangrene caused by *Candida* species as the primary organism. *Urology.* 2000;56(1):153. doi: 10.1016/S0090-4295(00)00527-6
  21. Oguz A, Gumus M, Turkoglu A, Bozdog Z, Ülger BV, Agacayak E, et al. Fournier's Gangrene: A summary of 10 years of clinical experience. *Int Surg.* 2015;100(5):934-941. doi: 10.9738/INTSURG-D-15-00036.1
  22. Ayan F, Sunamak O, Paksoy SM, Polat SS, As A, Sakoglu N, et al. Fournier's gangrene: A retrospective clinical study of forty-one patients. *Anz J Surg.* 2005;75(12):1055-1058. doi: 10.1111/j.1445-2197.2005.03609.x
  23. Kouadio K, Kouame YJ, Bi KL, Turquin HH. Perineal gangrene: Report of 30 cases observed at Abidjan. *Med Trop.* 1998;58(3):245-248.
  24. Kara C, Bozkurt OF, Unsal A. A Fournier's gangrene after transurethral resection of the prostate. *ActaChir Belg.* 2011;111:236-237.
  25. Quatan N, Kirby RS. Improving outcomes in Fournier's gangrene. *BJU Int.* 2004;93:691-692. doi: 10.1111/j.1464-410X.2003.04753.x
  26. Wróblewska M, Kuzaka B, Borkowski T, Kuzaka P, Kawecki D, Radziszewski P. Fournier's Gangrene – Current Concepts. *Pol J Microbiol.* 2014;63(3):267-673.
  27. Korkut M, Icoz G, Dayangac M, Akgun E, Yeniay L, Erdogan O, et al. Outcome analysis in patients with Fournier's gangrene: Report of 45 cases. *Dis Colon Rectum.* 2003;46:649-652.
  28. Kuo CF, Wang WS, Lee CM, Liu CP, Tseng HK. Fournier's gangrene: ten-year experience in a medical center in northern Taiwan. *J. MicrobiolImmunol Infect.* 2007;40(6):500–506.