

The Electromagnetic Drive: cyclotron motion at the geostationary orbit of Jupiter

Alberto Caballero¹

Abstract: In this research note it is proposed how the equilibrium between the Lorentz force and the centrifugal force in a crewed spacecraft acting as a tidally stabilised skyhook, and orbiting the geostationary orbit of Jupiter, could prevent the spacecraft from deorbiting and reaching unbearable accelerations. A pair of superconducting electromagnetics would produce a constantly-increasing Lorentz Force, which would cancel the centrifugal force while producing a constant acceleration of 4 g, the limit that a crew could withstand during long periods of time. The acceleration phase would last 2.9 months, time in which the crew would be hibernating inside Faraday suits electrically charged. The proposed drive forms part of the Solar One project, a program intended to design the first crewed interstellar spacecraft.

Key-words: Lorentz force, centrifugal force, electromagnetic propulsion, skyhook, space tether.

Space tethers are long cables which can be used for propulsion, momentum exchange, stabilization and attitude control, as well as maintaining the relative positions of the components of a large dispersed satellite or spacecraft sensor system.

One type of space tether is the skyhook. Specifically, a non-rotating skyhook is a vertical gravity-gradient stabilized tether.

Here I wish to discuss how a balance between the Lorentz force (which would act as the centripetal force) and the centrifugal force could provide a high yet bearable acceleration to a crewed skyhook-style spacecraft orbiting Jupiter at geostationary altitude.

The escape velocity of Jupiter is 59.5 km/s. Any spacecraft orbiting at this speed would leave the geostationary orbit, and the Lorentz force would gradually weaken to the point of disappearing.

A possible balance between forces could be achieved with the proper configuration of a system formed by a pair of superconducting electromagnets, one on each side of the spacecraft. These elements would provide the Lorentz force to the spacecraft while cancelling the centrifugal force.

¹ * alberto.caballero@uvigo.es




Figure 1: Balance between the Lorentz force (Fl) and the centrifugal force (Ff)

The electromagnetic system would constantly increase the Lorentz force in order to both prevent the spacecraft from deorbiting while it is being accelerated, and from producing an unbearable G-force to the crew. In theory, if the spacecraft is kept under a controlled constant acceleration, relativistic speeds could fall within the realm of plausibility.

Current flux would need to be adjusted once the speed of 42,5 km (first cosmic velocity of Jupiter) is reached, in order to balance the forces and produce a constant acceleration. At this point, if we have a spacecraft with a mass of 1,000 tons at the geostationary orbit of Jupiter, the centrifugal force would be the following:

$$F = m \cdot v^2 / r$$

$$F = 1,000,000 \cdot 42,500^2 / 159,722,000$$

$$F = 11,308,711 \text{ N}$$

Now, to calculate the Lorentz force needed to keep the spacecraft in orbit and produce the accelerate desired, we first need to determine the field strength at the geostationary orbit. Considering that Jupiter's Magnetic Dipole Moment is $1.55e+20 \text{ T}\cdot\text{m}^3$, we obtain the following field strength:

$$B = M / R^3$$

$$B = 1.55e + 20 / 159,722,000^3$$

$$B = 3.8e - 5 \text{ tesla}$$

So, for example, at 0.5c, the centrifugal force would produce an unbearable acceleration of 14.3 million g's. To cancel this, the electromagnetic system would have to produce an opposite Lorentz force equally strong. However, the Lorentz force has to be higher than the centrifugal force in order to

maintain the spacecraft in orbit and produce the constant acceleration desired. Considering that the crew could withstand a maximum of 4 g during a prolonged period of time (Poljak, 2018), the Lorentz force would be the following:

$$F = q \cdot v \cdot B \cdot \sin(\alpha)$$

$$F = 6,865,000 \cdot 150,000,000 \cdot 3.8e-5 \cdot \sin(90)$$

$$F = 1.4e+14 \text{ N}$$

If we used batteries with a power output equivalent to those on board the ISS (around 7.68 Kw/h), a voltage of around 0.001 V could produce a charge of 6.8 million Ah.

Considering an initial speed of 10,000 m/s, the amount of time that the crew would have to spend orbiting Jupiter in order to reach the speed of light is the following:

$$a = \frac{v_1 - v_0}{t}$$

$$t = 300,000,000 - 10,000 / 39.2$$

$$t = 7,652,806 \text{ sec} = 2.9 \text{ months}$$

The crew would have to be hibernating during the 2.9 months of the acceleration of phase. NASA has been working on a cryo-sleep design called TORPOR (Torpor Inducing Transfer Habitat For Human Stasis To Mars) (NASA, 2013), which could be used in this case. The crew would also need to be protected with Faraday suits, which would be electrically charged as the rest of the spacecraft. Faraday suits could be able to withstand an amperage similar to that of lightnings: around 20,000 amperes during 0.2 seconds. Considering a necessary amperage of 6.8 million Ah, which equals to 22,667 amperes every 0.2 seconds, Faraday suits could protect the crew from such continuous amperage.

Finally, instead of using Jupiter to accelerate the spacecraft, it could be possible to use the Sun for the same purpose. The magnetic dipole moment of the Sun is $3.5e+24 \text{ T} \cdot \text{m}^3$. Some NASA scientists have discussed the possibility of sending a crew as close to 6.4 million kilometres from the Sun (Mancini, 2018). The field strength at such distance would be slightly lower than Jupiter: $1.33e-5$ tesla. This means that more power would be required to achieve any acceleration around the Sun in comparison with an orbit around Jupiter. Moreover, the threat from radiation would increase at such small distances from the Sun.

References:

Feynman, Richard Phillips; Leighton, Robert B.; Sands, Matthew L. (2006). The Feynman lectures on physics (3 vol.). Pearson / Addison-Wesley. ISBN 0-8053-9047-2.: volume 2.

Mancini, Mark (2018) 'How Close Can We Get to the Sun?'. HowStuffWorks.

Murad, Paul (1995) 'An electromagnetic rocket hyper-light stellar drive'.

National Aeronautics and Space Administration (2013) 'Torpor inducing transfer habitat for human stasis to Mars'.

Poljak et. al (2018) 'Effects of exoplanetary gravity on human locomotor ability'.

Serway, Raymond A.; Jewett, John W., Jr. (2004). Physics for scientists and engineers, with modern physics. Belmont, [CA.]: Thomson Brooks/Cole. ISBN 0-534-40846-X.

Srednicki, Mark A. (2007). Quantum field theory. Cambridge, [England]; New York [NY.]: Cambridge University Press. ISBN 978-0-521-86449-7.