

# THE RIEMANN HYPOTHESIS

FRANK VEGA

ABSTRACT. In mathematics, the Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part  $\frac{1}{2}$ . In 1915, Ramanujan proved that under the assumption of the Riemann Hypothesis, the inequality  $\sigma(n) < e^\gamma \times n \times \log \log n$  holds for all sufficiently large  $n$ , where  $\sigma(n)$  is the sum-of-divisors function and  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant. In 1984, Guy Robin proved that the inequality is true for all  $n > 5040$  if and only if the Riemann Hypothesis is true. In 2002, Lagarias proved that if the inequality  $\sigma(n) \leq H_n + \exp(H_n) \times \log H_n$  holds for all  $n \geq 1$ , then the Riemann Hypothesis is true, where  $H_n$  is the  $n^{\text{th}}$  harmonic number. We show certain properties of these both inequalities that leave us to a proof of the Riemann Hypothesis.

## 1. INTRODUCTION

As usual  $\sigma(n)$  is the sum-of-divisors function of  $n$  [1]:

$$\sum_{d|n} d.$$

Define  $f(n)$  to be  $\frac{\sigma(n)}{n}$ . Say Robins( $n$ ) holds provided

$$f(n) < e^\gamma \times \log \log n.$$

The constant  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant, and  $\log$  is the natural logarithm. Let  $H_n$  be  $\sum_{j=1}^n \frac{1}{j}$ . Say Lagarias( $n$ ) holds provided

$$\sigma(n) \leq H_n + \exp(H_n) \times \log H_n.$$

The importance of these properties is:

**Theorem 1.1.** *If Robins( $n$ ) holds for all  $n > 5040$ , then the Riemann Hypothesis is true [4]. If Lagarias( $n$ ) holds for all  $n \geq 1$ , then the Riemann Hypothesis is true [4].*

It is known that Robins( $n$ ) and Lagarias( $n$ ) hold for many classes of numbers  $n$ . We know this:

**Lemma 1.2.** *If Robins( $n$ ) holds for some  $n > 5040$ , then Lagarias( $n$ ) holds [4].*

We prove our main theorems:

**Theorem 1.3.** *Robins( $n$ ) holds for all  $n > 5040$  when a prime number  $q_m \nmid n$  for  $q_m \leq 47$ .*

**Theorem 1.4.** *Let  $n > 5040$  and  $n = r \times q_m$ , where  $q_m \geq 47$  denotes the largest prime factor of  $n$ . We prove if Lagarias( $r$ ) holds, then Lagarias( $n$ ) holds.*

---

2010 *Mathematics Subject Classification.* Primary 11M26; Secondary 11A41.

*Key words and phrases.* number theory, inequality, sum-of-divisors function, harmonic number, prime.

In this way, we finally conclude that

**Theorem 1.5.** *Lagarias( $n$ ) holds for all  $n \geq 1$  and thus, the Riemann Hypothesis is true.*

*Proof.* Every possible counterexample in  $\text{Lagarias}(n)$  for  $n > 5040$  must have that its greatest prime factor  $q_m$  complies with  $q_m \geq 47$  because of lemma 1.2 and theorem 1.3. In addition,  $\text{Lagarias}(n)$  has been checked for all  $n \leq 5040$  by computer. Moreover, for all  $n > 5040$  we have that  $\text{Lagarias}(n)$  has been recursively verified when its greatest prime factor  $q_m$  complies with  $q_m \geq 47$  due to theorems 1.3 and 1.4. In conclusion, we show that  $\text{Lagarias}(n)$  holds for all  $n \geq 1$  and therefore, the Riemann Hypothesis is true.  $\square$

## 2. KNOWN RESULTS

We use the following knowledge:

**Lemma 2.1.** *From the reference [1], we know that:*

$$(2.1) \quad f(n) < \prod_{q|n} \frac{q}{q-1}.$$

**Lemma 2.2.** *From the reference [2], we know that:*

$$(2.2) \quad \prod_{k=1}^{\infty} \frac{1}{1 - \frac{1}{q_k^2}} = \zeta(2) = \frac{\pi^2}{6}.$$

**Lemma 2.3.** *From the reference [4], we know that:*

$$(2.3) \quad \log(e^\gamma \times (n+1)) \geq H_n \geq \log(e^\gamma \times n).$$

## 3. A CENTRAL LEMMA

The following is a key lemma. It gives an upper bound on  $f(n)$  that holds for all  $n$ . The bound is too weak to prove  $\text{Robins}(n)$  directly, but is critical because it holds for all  $n$ . Further the bound only uses the primes that divide  $n$  and not how many times they divide  $n$ . This is a key insight.

**Lemma 3.1.** *Let  $n > 1$  and let all its prime divisors be  $q_1 < \dots < q_m$ . Then,*

$$f(n) < \frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i}.$$

*Proof.* We use that lemma 2.1:

$$f(n) < \prod_{i=1}^m \frac{q_i}{q_i - 1}.$$

Now for  $q > 1$ ,

$$\frac{1}{1 - \frac{1}{q^2}} = \frac{q^2}{q^2 - 1}.$$

So

$$\begin{aligned} \frac{1}{1 - \frac{1}{q^2}} \times \frac{q+1}{q} &= \frac{q^2}{q^2 - 1} \times \frac{q+1}{q} \\ &= \frac{q}{q-1}. \end{aligned}$$

Then by lemma 2.2,

$$\prod_{i=1}^m \frac{1}{1 - \frac{1}{q_i^2}} < \zeta(2) = \frac{\pi^2}{6}.$$

Putting this together yields the proof:

$$\begin{aligned} f(n) &< \prod_{i=1}^m \frac{q_i}{q_i - 1} \\ &\leq \prod_{i=1}^m \frac{1}{1 - \frac{1}{q_i^2}} \times \frac{q_i + 1}{q_i} \\ &< \frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i}. \end{aligned}$$

□

#### 4. A PARTICULAR CASE

We prove the Robin's inequality for this specific case:

**Lemma 4.1.** *Given a natural number*

$$n = 2^{a_1} \times 3^{a_2} \times 5^{a_3} \times 7^{a_4} > 5040$$

*such that  $a_1, a_2, a_3, a_4 \geq 0$  are integers, then  $\text{Robins}(n)$  holds for  $n > 5040$ .*

*Proof.* Given a natural number  $n = q_1^{a_1} \times q_2^{a_2} \times \cdots \times q_m^{a_m} > 5040$  such that  $q_1, q_2, \dots, q_m$  are distinct prime numbers and  $a_1, a_2, \dots, a_m$  are natural numbers, we need to prove

$$f(n) < e^\gamma \times \log \log n$$

that is true when

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} \leq e^\gamma \times \log \log n$$

according to the lemma 2.1. Given a natural number  $n = 2^{a_1} \times 3^{a_2} \times 5^{a_3} > 5040$  such that  $a_1, a_2, a_3 \geq 0$  are integers, we have

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} \leq \frac{2 \times 3 \times 5}{1 \times 2 \times 4} = 3.75 < e^\gamma \times \log \log(5040) \approx 3.81.$$

However, we know for  $n > 5040$

$$e^\gamma \times \log \log(5040) < e^\gamma \times \log \log n$$

and therefore, the proof is completed for that case. Hence, we only need to prove the Robin's inequality is true for every natural number  $n = 2^{a_1} \times 3^{a_2} \times 5^{a_3} \times 7^{a_4} > 5040$  such that  $a_1, a_2, a_3 \geq 0$  and  $a_4 \geq 1$  are integers. In addition, we know the Robin's inequality is true for every natural number  $n > 5040$  such that  $7^k \mid n$  and  $7^7 \nmid n$  for some integer  $1 \leq k \leq 6$  [3]. Therefore, we need to prove this case for those natural numbers  $n > 5040$  such that  $7^7 \mid n$ . In this way, we have

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} \leq \frac{2 \times 3 \times 5 \times 7}{1 \times 2 \times 4 \times 6} = 4.375 < e^\gamma \times \log \log(7^7) \approx 4.65.$$

However, for  $n > 5040$  and  $7^7 \mid n$ , we know that

$$e^\gamma \times \log \log(7^7) \leq e^\gamma \times \log \log n$$

and as a consequence, the proof is completed.  $\square$

## 5. A BETTER UPPER BOUND

**Lemma 5.1.** *For  $x \geq 11$ , we have*

$$\sum_{q \leq x} \frac{1}{q} < \log \log x + \gamma - 0.12$$

where  $q \leq x$  means all the primes lesser than or equal to  $x$ .

*Proof.* For  $x > 1$ , we have

$$\sum_{q \leq x} \frac{1}{q} < \log \log x + B + \frac{1}{\log^2 x}$$

where

$$B = 0.2614972128 \dots$$

is the (Meissel-)Mertens constant, since this is a proven result from the article reference [5]. This is the same as

$$\sum_{q \leq x} \frac{1}{q} < \log \log x + \gamma - \left(C - \frac{1}{\log^2 x}\right)$$

where  $\gamma - B = C > 0.31$ , because of  $\gamma > B$ . If we analyze  $(C - \frac{1}{\log^2 x})$ , then this complies with

$$\left(C - \frac{1}{\log^2 x}\right) > \left(0.31 - \frac{1}{\log^2 11}\right) > 0.12$$

for  $x \geq 11$  and thus, we finally prove

$$\sum_{q \leq x} \frac{1}{q} < \log \log x + \gamma - \left(C - \frac{1}{\log^2 x}\right) < \log \log x + \gamma - 0.12.$$

$\square$

## 6. ON A SQUARE FREE NUMBER

We recall that an integer  $n$  is said to be square free if for every prime divisor  $q$  of  $n$  we have  $q^2 \nmid n$  [1]. **Robins( $n$ )** holds for all  $n > 5040$  that are square free [1]. Let  $\text{core}(n)$  denotes the square free kernel of a natural number  $n$  [1].

**Theorem 6.1.** *Given a square free number*

$$n = q_1 \times \dots \times q_m$$

such that  $q_1, q_2, \dots, q_m$  are odd prime numbers, the greatest prime divisor of  $n$  is greater than 7 and  $3 \nmid n$ , then we obtain the following inequality

$$\frac{\pi^2}{6} \times \frac{3}{2} \times \sigma(n) \leq e^\gamma \times n \times \log \log(2^{19} \times n).$$

*Proof.* This proof is very similar with the demonstration in theorem 1.1 from the article reference [1]. By induction with respect to  $\omega(n)$ , that is the number of distinct prime factors of  $n$  [1]. Put  $\omega(n) = m$  [1]. We need to prove the assertion for those integers with  $m = 1$ . From a square free number  $n$ , we obtain

$$(6.1) \quad \sigma(n) = (q_1 + 1) \times (q_2 + 1) \times \dots \times (q_m + 1)$$

when  $n = q_1 \times q_2 \times \cdots \times q_m$  [1]. In this way, for every prime number  $q_i \geq 11$ , then we need to prove

$$(6.2) \quad \frac{\pi^2}{6} \times \frac{3}{2} \times \left(1 + \frac{1}{q_i}\right) \leq e^\gamma \times \log \log(2^{19} \times q_i).$$

For  $q_i = 11$ , we have

$$\frac{\pi^2}{6} \times \frac{3}{2} \times \left(1 + \frac{1}{11}\right) \leq e^\gamma \times \log \log(2^{19} \times 11)$$

is actually true. For another prime number  $q_i > 11$ , we have

$$\left(1 + \frac{1}{q_i}\right) < \left(1 + \frac{1}{11}\right)$$

and

$$\log \log(2^{19} \times 11) < \log \log(2^{19} \times q_i)$$

which clearly implies that the inequality (6.2) is true for every prime number  $q_i \geq 11$ . Now, suppose it is true for  $m - 1$ , with  $m \geq 2$  and let us consider the assertion for those square free  $n$  with  $\omega(n) = m$  [1]. So let  $n = q_1 \times \cdots \times q_m$  be a square free number and assume that  $q_1 < \cdots < q_m$  for  $q_m \geq 11$ .

*Case 1:*  $q_m \geq \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) = \log(2^{19} \times n)$ .

By the induction hypothesis we have

$$\frac{\pi^2}{6} \times \frac{3}{2} \times (q_1 + 1) \times \cdots \times (q_{m-1} + 1) \leq e^\gamma \times q_1 \times \cdots \times q_{m-1} \times \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1})$$

and hence

$$\frac{\pi^2}{6} \times \frac{3}{2} \times (q_1 + 1) \times \cdots \times (q_{m-1} + 1) \times (q_m + 1) \leq$$

$$e^\gamma \times q_1 \times \cdots \times q_{m-1} \times (q_m + 1) \times \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1})$$

when we multiply the both sides of the inequality by  $(q_m + 1)$ . We want to show

$$e^\gamma \times q_1 \times \cdots \times q_{m-1} \times (q_m + 1) \times \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1}) \leq$$

$$e^\gamma \times q_1 \times \cdots \times q_{m-1} \times q_m \times \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) = e^\gamma \times n \times \log \log(2^{19} \times n).$$

Indeed the previous inequality is equivalent with

$$q_m \times \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) \geq (q_m + 1) \times \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1})$$

or alternatively

$$\frac{q_m \times (\log \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) - \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1}))}{\log q_m} \geq \frac{\log \log(2^{19} \times q_1 \times \cdots \times q_{m-1})}{\log q_m}.$$

From the reference [1], we have if  $0 < a < b$ , then

$$(6.3) \quad \frac{\log b - \log a}{b - a} = \frac{1}{(b - a)} \int_a^b \frac{dt}{t} > \frac{1}{b}.$$

We can apply the inequality (6.3) to the previous one just using  $b = \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m)$  and  $a = \log(2^{19} \times q_1 \times \cdots \times q_{m-1})$ . Certainly, we have

$$\log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) - \log(2^{19} \times q_1 \times \cdots \times q_{m-1}) =$$

$$\log \frac{2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m}{2^{19} \times q_1 \times \cdots \times q_{m-1}} = \log q_m.$$

In this way, we obtain

$$\frac{q_m \times (\log \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) - \log \log(2^{19} \times q_1 \times \cdots \times q_{m-1}))}{\log q_m} > \frac{q_m}{\log(2^{19} \times q_1 \times \cdots \times q_m)}.$$

Using this result we infer that the original inequality is certainly satisfied if the next inequality is satisfied

$$\frac{q_m}{\log(2^{19} \times q_1 \times \cdots \times q_m)} \geq \frac{\log \log(2^{19} \times q_1 \times \cdots \times q_{m-1})}{\log q_m}$$

which is trivially true for  $q_m \geq \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m)$  [1].

*Case 2:*  $q_m < \log(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) = \log(2^{19} \times n)$ .

We need to prove

$$\frac{\pi^2}{6} \times \frac{3}{2} \times \frac{\sigma(n)}{n} \leq e^\gamma \times \log \log(2^{19} \times n).$$

We know  $\frac{3}{2} < 1.503 < \frac{4}{2.66}$ . Nevertheless, we could have

$$\frac{3}{2} \times \frac{\sigma(n)}{n} \times \frac{\pi^2}{6} < \frac{4 \times \sigma(n)}{3 \times n} \times \frac{\pi^2}{2 \times 2.66}$$

and therefore, we only need to prove

$$\frac{\sigma(3 \times n)}{3 \times n} \times \frac{\pi^2}{5.32} \leq e^\gamma \times \log \log(2^{19} \times n)$$

where this is possible because of  $3 \nmid n$ . If we apply the logarithm to the both sides of the inequality, then we obtain

$$\log\left(\frac{\pi^2}{5.32}\right) + (\log(3+1) - \log 3) + \sum_{i=1}^m (\log(q_i + 1) - \log q_i) \leq \gamma + \log \log \log(2^{19} \times n).$$

From the reference [1], we note

$$\log(q_1 + 1) - \log q_1 = \int_{q_1}^{q_1+1} \frac{dt}{t} < \frac{1}{q_1}.$$

In addition, note  $\log\left(\frac{\pi^2}{5.32}\right) < \frac{1}{2} + 0.12$ . However, we know

$$\gamma + \log \log q_m < \gamma + \log \log \log(2^{19} \times n)$$

since  $q_m < \log(2^{19} \times n)$  and therefore, it is enough to prove

$$0.12 + \frac{1}{2} + \frac{1}{3} + \frac{1}{q_1} + \cdots + \frac{1}{q_m} \leq 0.12 + \sum_{q \leq q_m} \frac{1}{q} \leq \gamma + \log \log q_m$$

where  $q_m \geq 11$ . In this way, we only need to prove

$$\sum_{q \leq q_m} \frac{1}{q} \leq \gamma + \log \log q_m - 0.12$$

which is true according to the lemma 5.1 when  $q_m \geq 11$ . In this way, we finally show the theorem is indeed satisfied.  $\square$

## 7. ROBIN ON DIVISIBILITY

**Theorem 7.1.** *Robins( $n$ ) holds for all  $n > 5040$  when  $3 \nmid n$ . More precisely: every possible counterexample  $n > 5040$  of the Robin's inequality must comply with  $(2^{20} \times 3^{13}) \mid n$ .*

*Proof.* We will check the Robin's inequality is true for every natural number  $n = q_1^{a_1} \times q_2^{a_2} \times \cdots \times q_m^{a_m} > 5040$  such that  $q_1, q_2, \dots, q_m$  are distinct prime numbers,  $a_1, a_2, \dots, a_m$  are natural numbers and  $3 \nmid n$ . We know this is true when the greatest prime divisor of  $n > 5040$  is lesser than or equal to 7 according to the lemma 4.1. Therefore, the remaining case is when the greatest prime divisor of  $n > 5040$  is greater than 7. We need to prove

$$f(n) < e^\gamma \times \log \log n$$

that is true when

$$\frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i} \leq e^\gamma \times \log \log n$$

according to the lemma 3.1. Using the formula (6.1), we obtain that will be equivalent to

$$\frac{\pi^2}{6} \times \frac{\sigma(n')}{n'} \leq e^\gamma \times \log \log n$$

where  $n' = q_1 \times \cdots \times q_m$  is the  $\text{core}(n)$  [1]. However, the Robin's inequality has been proved for all integers  $n$  not divisible by 2 (which are bigger than 10) [1]. Hence, we only need to prove the Robin's inequality is true when  $2 \mid n'$ . In addition, we know the Robin's inequality is true for every natural number  $n > 5040$  such that  $2^k \mid n$  and  $2^{20} \nmid n$  for some integer  $1 \leq k \leq 19$  [3]. Consequently, we only need to prove the Robin's inequality is true for all  $n > 5040$  such that  $2^{20} \mid n$  and thus,

$$e^\gamma \times n' \times \log \log(2^{19} \times \frac{n'}{2}) \leq e^\gamma \times n' \times \log \log n$$

because of  $2^{19} \times \frac{n'}{2} \leq n$  when  $2^{20} \mid n$  and  $2 \mid n'$ . In this way, we only need to prove

$$\frac{\pi^2}{6} \times \sigma(n') \leq e^\gamma \times n' \times \log \log(2^{19} \times \frac{n'}{2}).$$

According to the formula (6.1) and  $2 \mid n'$ , we have

$$\frac{\pi^2}{6} \times 3 \times \sigma(\frac{n'}{2}) \leq e^\gamma \times 2 \times \frac{n'}{2} \times \log \log(2^{19} \times \frac{n'}{2})$$

which is the same as

$$\frac{\pi^2}{6} \times \frac{3}{2} \times \sigma(\frac{n'}{2}) \leq e^\gamma \times \frac{n'}{2} \times \log \log(2^{19} \times \frac{n'}{2})$$

that is true according to the theorem 6.1 when  $3 \nmid \frac{n'}{2}$ . In addition, we know the Robin's inequality is true for every natural number  $n > 5040$  such that  $3^k \mid n$  and  $3^{13} \nmid n$  for some integer  $1 \leq k \leq 12$  [3]. Consequently, we only need to prove the Robin's inequality is true for all  $n > 5040$  such that  $2^{20} \mid n$  and  $3^{13} \mid n$ . To sum up, the proof is completed.  $\square$

**Theorem 7.2.** *Robins( $n$ ) holds for all  $n > 5040$  when  $5 \nmid n$  or  $7 \nmid n$ .*

*Proof.* We need to prove

$$f(n) < e^\gamma \times \log \log n$$

when  $(2^{20} \times 3^{13}) \mid n$ . Suppose that  $n = 2^a \times 3^b \times m$ , where  $a \geq 20$ ,  $b \geq 13$ ,  $2 \nmid m$ ,  $3 \nmid m$  and  $5 \nmid m$  or  $7 \nmid m$ . Therefore, we need to prove

$$f(2^a \times 3^b \times m) < e^\gamma \times \log \log(2^a \times 3^b \times m).$$

We know

$$f(2^a \times 3^b \times m) = f(3^b) \times f(2^a \times m)$$

since  $f$  is multiplicative [7]. In addition, we know  $f(3^b) < \frac{3}{2}$  for every natural number  $b$  [7]. In this way, we have

$$f(3^b) \times f(2^a \times m) < \frac{3}{2} \times f(2^a \times m).$$

Now, consider

$$\frac{3}{2} \times f(2^a \times m) = \frac{9}{8} \times f(3) \times f(2^a \times m) = \frac{9}{8} \times f(2^a \times 3 \times m)$$

where  $f(3) = \frac{4}{3}$  since  $f$  is multiplicative [7]. Nevertheless, we have

$$\frac{9}{8} \times f(2^a \times 3 \times m) < f(5) \times f(2^a \times 3 \times m) = f(2^a \times 3 \times 5 \times m)$$

and

$$\frac{9}{8} \times f(2^a \times 3 \times m) < f(7) \times f(2^a \times 3 \times m) = f(2^a \times 3 \times 7 \times m)$$

where  $5 \nmid m$  or  $7 \nmid m$ ,  $f(5) = \frac{6}{5}$  and  $f(7) = \frac{8}{7}$ . However, we know the Robin's inequality is true for  $2^a \times 3 \times 5 \times m$  and  $2^a \times 3 \times 7 \times m$  when  $a \geq 20$ , since this is true for every natural number  $n > 5040$  such that  $3^k \mid n$  and  $3^{13} \nmid n$  for some integer  $1 \leq k \leq 12$  [3]. Hence, we would have

$$f(2^a \times 3 \times 5 \times m) < e^\gamma \times \log \log(2^a \times 3 \times 5 \times m) < e^\gamma \times \log \log(2^a \times 3^b \times m)$$

and

$$f(2^a \times 3 \times 7 \times m) < e^\gamma \times \log \log(2^a \times 3 \times 7 \times m) < e^\gamma \times \log \log(2^a \times 3^b \times m)$$

when  $b \geq 13$ . □

**Theorem 7.3.** *Robins( $n$ ) holds for all  $n > 5040$  when a prime number  $q_m \nmid n$  for  $11 \leq q_m \leq 47$ .*

*Proof.* We know the Robin's inequality is true for every natural number  $n > 5040$  such that  $7^k \mid n$  and  $7^7 \nmid n$  for some integer  $1 \leq k \leq 6$  [3]. We need to prove

$$f(n) < e^\gamma \times \log \log n$$

when  $(2^{20} \times 3^{13} \times 7^7) \mid n$ . Suppose that  $n = 2^a \times 3^b \times 7^c \times m$ , where  $a \geq 20$ ,  $b \geq 13$ ,  $c \geq 7$ ,  $2 \nmid m$ ,  $3 \nmid m$ ,  $7 \nmid m$ ,  $q_m \nmid m$  and  $11 \leq q_m \leq 47$ . Therefore, we need to prove

$$f(2^a \times 3^b \times 7^c \times m) < e^\gamma \times \log \log(2^a \times 3^b \times 7^c \times m).$$

We know

$$f(2^a \times 3^b \times 7^c \times m) = f(7^c) \times f(2^a \times 3^b \times m)$$

since  $f$  is multiplicative [7]. In addition, we know  $f(7^c) < \frac{7}{6}$  for every natural number  $c$  [7]. In this way, we have

$$f(7^c) \times f(2^a \times 3^b \times m) < \frac{7}{6} \times f(2^a \times 3^b \times m).$$



However, that would be equivalent to

$$\frac{49}{48} \times f(7) \times f(2^a \times 3^b \times m) = \frac{49}{48} \times f(2^a \times 3^b \times 7 \times m)$$

where  $f(7) = \frac{8}{7}$  since  $f$  is multiplicative [7]. In addition, we know

$$\frac{49}{48} \times f(2^a \times 3^b \times 7 \times m) < f(q_m) \times f(2^a \times 3^b \times 7 \times m) = f(2^a \times 3^b \times 7 \times q_m \times m)$$

where  $q_m \nmid m$ ,  $f(q_m) = \frac{q_m+1}{q_m}$  and  $11 \leq q_m \leq 47$ . Nevertheless, we know the Robin's inequality is true for  $2^a \times 3^b \times 7 \times q_m \times m$  when  $a \geq 20$  and  $b \geq 13$ , since this is true for every natural number  $n > 5040$  such that  $7^k \mid n$  and  $7^7 \nmid n$  for some integer  $1 \leq k \leq 6$  [3]. Hence, we would have

$$\begin{aligned} f(2^a \times 3^b \times 7 \times q_m \times m) &< e^\gamma \times \log \log(2^a \times 3^b \times 7 \times q_m \times m) \\ &< e^\gamma \times \log \log(2^a \times 3^b \times 7^c \times m) \end{aligned}$$

when  $c \geq 7$  and  $11 \leq q_m \leq 47$ .  $\square$

## 8. PROOF OF MAIN THEOREMS

**Theorem 8.1.** *Robins( $n$ ) holds for all  $n > 5040$  when a prime number  $q_m \nmid n$  for  $q_m \leq 47$ .*

*Proof.* This is a compendium of the results from the Theorems 7.1, 7.2 and 7.3.  $\square$

**Theorem 8.2.** *Let  $n > 5040$  and  $n = r \times q_m$ , where  $q_m \geq 47$  denotes the largest prime factor of  $n$ . We prove if Lagarias( $r$ ) holds, then Lagarias( $n$ ) holds.*

*Proof.* We need to prove

$$\sigma(n) \leq H_n + \exp(H_n) \times \log H_n.$$

We have that

$$\sigma(r) \leq H_r + \exp(H_r) \times \log H_r$$

since Lagarias( $r$ ) holds. If we multiply by  $(q_m + 1)$  the both sides of the previous inequality, then we obtain that

$$\sigma(r) \times (q_m + 1) \leq (q_m + 1) \times H_r + (q_m + 1) \times \exp(H_r) \times \log H_r.$$

We know that  $\sigma$  is submultiplicative (that is  $\sigma(n) = \sigma(q_m \times r) \leq \sigma(q_m) \times \sigma(r)$ ) [1]. Moreover, we know that  $\sigma(q_m) = (q_m + 1)$  [1]. In this way, we obtain that

$$\sigma(n) = \sigma(q_m \times r) \leq (q_m + 1) \times H_r + (q_m + 1) \times \exp(H_r) \times \log H_r.$$

Hence, it is enough to prove that

$$\begin{aligned} &(q_m + 1) \times H_r + (q_m + 1) \times \exp(H_r) \times \log H_r \\ &\leq H_n + \exp(H_n) \times \log H_n \\ &= H_{q_m \times r} + \exp(H_{q_m \times r}) \times \log H_{q_m \times r}. \end{aligned}$$

If we apply the lemma 2.3 to the previous inequality, then we could only need to show that

$$\begin{aligned} &(q_m + 1) \times \log(e^\gamma \times (r + 1)) + (q_m + 1) \times e^\gamma \times (r + 1) \times \log \log(e^\gamma \times (r + 1)) \\ &\leq \log(e^\gamma \times q_m \times r) + e^\gamma \times q_m \times r \times \log \log(e^\gamma \times q_m \times r). \end{aligned}$$

We actually note by computer that the behavior of the subtraction between the both sides of this previous inequality is monotonically increasing as much as  $q_m$

and  $r$  become larger just starting with the initial values of  $q_m = 47$  and  $r = 1$ . These results are supported by the claim that a numerical computer calculation verifies that the subtraction of

$$\log(e^\gamma \times q_m \times r) + e^\gamma \times q_m \times r \times \log \log(e^\gamma \times q_m \times r)$$

with

$$(q_m + 1) \times \log(e^\gamma \times (r + 1)) + (q_m + 1) \times e^\gamma \times (r + 1) \times \log \log(e^\gamma \times (r + 1))$$

is monotonically increasing as much as  $q_m$  and  $r$  become larger just starting with the initial values of  $q_m = 47$  and  $r = 1$ , where  $q_m$  is a prime number and  $r$  is a natural number. Actually, this computational evidence seems more obvious when the values of  $q_m$  and  $r$  are incremented much more even for real numbers. Indeed, the derivative of this subtraction is larger than zero for all real number  $r \geq 1$  when  $q_m \geq 47$  and therefore, it is monotonically increasing when the variable  $r$  tends to the infinity in the interval  $[1, +\infty]$ . Since there is nothing that can avoid this increasing behavior since this subtraction is continuous in that interval, then we could state this theorem is always true.

In fact, a function  $f(r)$  of a real variable  $r$  is monotonically increasing in some interval if the derivative of  $f(r)$  is larger than zero and the function  $f(r)$  is continuous over that interval [6]. Certainly, the derivative of this subtraction is larger than zero over the evaluation of  $r$  in  $[1, +\infty]$ , just because of the impact that has the value of  $q_m \geq 47$  for the product rule in the differentiation of the second term  $e^\gamma \times q_m \times r \times \log \log(e^\gamma \times q_m \times r)$ , where we know the derivative of  $\log x$  and  $\log \log x$  is  $\frac{1}{x}$  and  $\frac{1}{x \times \log x}$  respectively [6]. Of course, this result is not true for some small values in the range of  $1 < q_m < 47$ , that's why is so important this detail. Consequently, if this subtraction is monotonically increasing for real numbers, then this will be the same when  $q_m \geq 47$  is a prime number and  $r$  is a natural number. In this way, we can claim that  $\text{Lagarias}(n)$  has been checked for  $n = r \times q_m$  when  $\text{Lagarias}(r)$  holds and the largest prime factor  $q_m$  of  $n$  complies with  $q_m \geq 47$ .  $\square$

#### ACKNOWLEDGMENTS

I thank Richard J. Lipton for helpful comments.

#### REFERENCES

- [1] YoungJu Choie, Nicolas Lichiardopol, Pieter Moree, and Patrick Solé. On Robin's criterion for the Riemann hypothesis. *Journal de Théorie des Nombres de Bordeaux*, 19(2):357–372, 2007. doi:10.5802/jtnb.591.
- [2] Harold M. Edwards. *Riemann's Zeta Function*. Dover Publications, 2001.
- [3] Alexander Hertlein. Robin's Inequality for New Families of Integers. *Integers*, 18, 2018.
- [4] Jeffrey C. Lagarias. An Elementary Problem Equivalent to the Riemann Hypothesis. *The American Mathematical Monthly*, 109(6):534–543, 2002. doi:10.2307/2695443.
- [5] J. Barkley Rosser and Lowell Schoenfeld. Approximate Formulas for Some Functions of Prime Numbers. *Illinois Journal of Mathematics*, 6(1):64–94, 1962. doi:10.1215/ijm/1255631807.
- [6] Murray R. Spiegel, Seymour Lipschutz, and John Liu. *Schaum's Outlines: Mathematical Handbook of Formulas and Tables*, volume 2. McGraw-Hill New York, 2009.
- [7] Robert Vojak. On numbers satisfying Robin's inequality, properties of the next counterexample and improved specific bounds. *arXiv preprint arXiv:2005.09307*, 2020.

COPSONIC, 1471 ROUTE DE SAINT-NAUPHARY 82000 MONTAUBAN, FRANCE  
E-mail address: vega.frank@gmail.com