

1 **Abundance and types of plastic pollution in surface waters in the Eastern Arctic (Inuit**
2 **Nunangat) and the case for reconciliation science**

3
4 Max Liboiron^{1,2}

5 Alex Zahara^{1,2}

6 Kaitlyn Hawkins¹

7 Christina Crespo^{1,3}

8 Bárbara de Moura Neves⁴

9 Vonda Wareham-Hayes⁴

10 Evan Edinger²

11 Charlotte Muise¹

12 Mary Jane Walzak⁵

13 Rebecca Sarazen⁵

14 Jillian Chidley¹

15 Carley Mills¹

16 Lauren Watwood¹

17 Hridisha Arif¹

18 Elise Earles¹

19 Liz Pijogge⁶

20 Jamal Shirley⁷

21 Jesse Jacobs⁸

22 Paul McCarney⁶

23 Louis Charron¹

24

25 ¹ Civic Laboratory for Environmental Action Research (CLEAR), Memorial University, St. John's,
26 Newfoundland and Labrador, Canada

27 ² Department of Geography, Memorial University, St. John's, Newfoundland and Labrador,
28 Canada

29 ³ Department of Anthropology, University of Georgia, Athens, USA

30 ⁴ Department of Fisheries and Oceans, St. John's, Newfoundland and Labrador, Canada

31 ⁵ Surface Science Western, University of Western Ontario, London, Ontario, Canada

32 ⁶ Lands and Natural Resources, Nunatsiavut Government, Nain, Newfoundland and Labrador,
33 Canada

34 ⁷ Nunavut Research Institute, Nunavut Arctic College, Iqaluit, Nunavut, Canada

35 ⁸ Nunavut Arctic College - Memorial University Partnership, Memorial University, St. John's,
36 Newfoundland and Labrador, Canada

37

38 **Abstract:** Plastics are not only an environmental concern but also an issue of justice in the
39 Arctic, particularly in Inuit Nunangat (Inuit homelands), as plastics and other contaminants that
40 originate in the south accumulate in the north and have implications for Inuit sovereignty and
41 wellness. This collaborative study finds an average of 0.018 plastics/m² in surface waters in two
42 sites in the eastern Arctic (Tasiujarjuaq in Nunavut near Iqaluit and southwest Greenland
43 offshore from Qaqortoq and Narsaq). A comparison with other studies shows this abundance of
44 plastics is lesser than abundances reported further north in the Arctic, but greater than adjacent
45 waters further south. However, within and across study areas at similar latitudes, there does not
46 appear to be a significant difference in plastic abundance. Some characteristics of recovered
47 plastics such as morphology and colour support local origins, while others support long range
48 transport. Research moving forward should consider relative scales in spatial trends of plastic
49 abundance. The discussion concludes by reflecting on the methods and findings in terms of
50 their role in Inuit governance and research relationships, including elements of research
51 personnel, permitting, categorization, measurement, and reporting findings. Our goal is to
52 provide insights of where we, as scientists, may choose to intentionally move our scientific work
53 towards reconciliation while we produce knowledge about environmental pollution in Inuit
54 Nunangat and the Arctic broadly.

55

56 **Keywords:** Arctic; plastic; surface water; marine debris; colonialism; Greenland; Nunavut; Inuit
57 Nunangat; reconciliation science

58

59 **Highlights:**

- 60 • Reports plastics in surface water in SW Greenland (0.026/m²) & Tasiujarjuaq, Nunavut
61 (0.014/m²)
- 62 • Recovered plastics show indications of both long-range and local sources
- 63 • Surface water plastic research in Inuit Nunangat and Greenland is led by southerners
64 and non-Inuit
- 65 • “Reconciliation science” requires changes in personnel, methods, and communicating
66 results

67

68

69 **1. Introduction**

70 Plastics are not only an environmental concern, but also an issue of justice in the Arctic,
71 particularly in Inuit Nunangat (Inuit homelands), as plastics and other contaminants that
72 originate in the south accumulate in the north and have implications for Inuit sovereignty and
73 wellness (Bourdages et al. 2020; Sudlovenick 2019; Watt-Cloutier 2015; Furgall et al. 2005; Van
74 Oostam et al. 2005). Inuit Nunangat contains four regions: the Inuvialuit Settlement Region
75 (northern Northwest Territories), Nunavut, Nunavik (northern Quebec), and Nunatsiavut
76 (northern Labrador). It includes 53 communities and encompasses roughly 35 percent of
77 Canada's land mass and 50 percent of its coastline, but no manufacturing infrastructure for the
78 production of plastics (Royal Canadian Geographical Society 2018). Yet studies have shown a
79 clear trend where plastic has been found to accumulate in Arctic waters and ecosystems
80 compared to waters further south with higher populations and plastic manufacturing
81 infrastructure (Obbard 2018; Cózar et al 2017). The hypothesis is that as the global
82 Thermohaline Circulation (THC) actively moves "warm surface water from low to high latitudes
83 across the North Atlantic Ocean to the Arctic, it could collect buoyant plastic from highly
84 populated latitudes, leading to accumulation in the Greenland and Barents seas, where the
85 landmasses, together with the polar ice cap, would constitute a dead end for the surface
86 transport of floating debris" (Cózar et al 2017: 1; see also Lusher et al 2015). Southerly
87 movement of plastics to Arctic locations also takes place through atmospheric transport of
88 microplastics such as microfibers (Evangelidou et al 2020; Bergmann et al 2019), the release of
89 plastics deposited by Pacific Ocean waters through melting sea ice (Peeken et al 2018; Obbard
90 et al 2014), and the biotransport of plastics via seasonally migrating animals, such as seabirds
91 (Bourdages et al. In press; Provencher et al 2010; Mallory et al 2006).

92
93 At the same time, the Arctic is relatively understudied from a scientific perspective (AMAP/EU-
94 PolarNet 2020), meaning more work is needed to describe the patterns in plastic accumulation
95 in the North. Existing scientific studies of plastic pollution in the Arctic are framed in terms of
96 baseline figures (Lusher et al 2015; Mallory 2008) or plastics' environmental effects (Kanhai et
97 al 2018; Provencher et al 2010). Our study continues this scientific trajectory by providing
98 abundance measures (number of plastics/m²) and an analysis of types of plastic pollution in
99 surface waters near the capital city of Iqaluit in Tasiujarjuaq (Frobisher Bay), Nunavut, and
100 offshore from Qaqortoq and Narsaq in southwest Greenland. We compare these findings to
101 other results in the Arctic generally and Inuit Nunangat in particular.

102

103 This study is the result of a collaboration between four different groups: a scientific team based
104 in St. John's, Newfoundland and Labrador, that collected all samples in Nunavut and
105 Greenland; CLEAR, an interdisciplinary plastic pollution laboratory also based in St. John's that
106 studies marine plastics as well as colonialism in science that conducted analysis and writing;
107 scientists at Surface Science Western Lab at the University of Western Ontario that conducted
108 spectrometry work; and a group of Inuit and non-Inuit research professionals who have lived
109 and worked in Inuit Nunangat who provided expert insights, context, and validation for analyzing
110 findings and recommendations. This partnership was not planned in advance but rather came
111 together as research needs arose and has resulted in a unique extension of the scientific study
112 in the discussion section of the paper on reconciliation science. In addition to discussing
113 research findings for trends in plastic pollution, we also provide critical reflection on research
114 methods and findings in terms of Inuit-based research, reconciliation, and governance
115 relationships (Loseto et al 2020; Pedersen et al 2020; Wong et al 2020; ITK 2018; Pfeifer 2018;
116 Moffitt et al 2015; ITK & NRI 2007). Such reflections are crucial for ongoing and future research
117 in Inuit Nunangat to ensure our work as scientists and research partners is part of reconciliation.
118 We are calling this "reconciliation science." Rather than dividing these reflections into a separate
119 "opinion" piece or social science paper, we make the case that existing and ongoing Indigenous
120 research relations should never be divided from scientific study and reporting, and our goal is to
121 provide one model for how reconciliation science might be done.

122

123 **2. Materials and Methods**

124 *2.1 Sample Collection*

125 Samples for this study were collected in 2018 by a team of settler, non-Inuit authors based in St.
126 John's, Newfoundland and Labrador, who were already traveling to Inuit Nunangat and
127 southwestern Greenland for research unrelated to plastic pollution for seafloor mapping and
128 biodiversity (via the Amundsen Science Leg 2c, 2018). Sample collection took place in July and
129 August of 2018 near the capital city of Iqaluit in Tasiujarjuaq (Frobisher Bay), Nunavut, and in
130 the Labrador Sea offshore from Qaqortoq and Narsaq in southwest Greenland aboard the
131 Canadian Coast Guard Vessel (CCGS) *Amundsen* (Figure 1).

132

133 Sample collection sites were designed to answer two questions. (H1) First, we aimed to discern
134 whether plastics that might have originated in the Iqaluit landfill, which burned for 178 days in
135 2014, were moving from land into Tasiujarjuaq, a local hypothesis that resulted in an invitation
136 to do related plastic pollution research in the area (also see Watson 2014, Nunatsiaq News

137 2015, Varga 2015). Sampling locations in Tasiujarjuaq were grouped close to Iqaluit and further
138 out in the bay to test that hypothesis, and plastics were inspected for signs of burning or
139 melting. (H2) Secondly, the site in southwest Greenland was used as a comparison to
140 determine whether the abundance and types of plastics in Tasiujarjuaq were markedly different
141 than those of a location in another current but at a similar latitude. Tasiujarjuaq is macrotidal
142 (11m tide range) with particularly strong tidal currents through the mid-bay islands, and
143 southwest Greenland is dominated by the West Greenland current, bringing surface waters
144 around Greenland from the northeast Atlantic Ocean (see S13). This sample collection design
145 provided a snapshot of plastic profiles in two key locations and cannot be used for wider
146 generalizations of geographic patterns of plastic pollution, given its relatively low sample size
147 and limited duration of sampling.

148
149 Researchers used a Manta surface water trawl with a net mesh size of 335 μ m, and each trawl
150 was conducted for 30 minutes. The mouth of the trawl is 0.53m, determining the maximum size
151 of plastic we would have collected. A flowmeter was attached to the trawl to determine the
152 amount of water sampled. Entire cod ends containing samples were placed in sample bags with
153 20mL of hydrogen peroxide (3%) for storage and frozen at -20 °C for later laboratory analysis.
154 Contamination samples were taken each day from mittens, gloves, scarfs, toques (winter hats),
155 jackets, and other fabrics of all personnel in contact with the trawl to ensure potential sources of
156 contamination during data collection could be identified. The colour of the ship's paint, sampling
157 gear, and ropes were also recorded as potential sources of contamination.

158 159 *2.2 Permits*

160 Permits, licenses, and/or permissions are required before conducting research in most areas,
161 including the Arctic. Following research permitting protocols is particularly important in
162 Indigenous land claim areas and traditional homelands as part of Indigenous sovereignty— the
163 right of Indigenous peoples to govern Indigenous lands. Permits were sought and obtained by
164 Amundsen Science, including: Nunavut Research Institute Scientific Research License #
165 0501318R-M; Department of Fisheries and Oceans License to Fish for Scientific Purposes in
166 the waters of Nunavut # S-18/19-1012-NU; Vessel Clearance to conduct scientific work in
167 Greenland waters; Danish Ministry of Foreign Affairs file # 2018-15931; and Government of
168 Greenland Survey License # G18-028. For an overview of research licensing bodies in Inuit
169 Nunangat, see ITK 2018: 15-16 and Table S1.

170

171 *2.3 Lab Analysis*

172 After collection, samples were transferred to a team at the Civic Laboratory for Environmental
173 Action Research (CLEAR) based in St. John's (Beothuk homelands) for processing and
174 analysis. CLEAR team co-authors are non-Inuit: some are settlers and some are Indigenous
175 from other places. Samples were transferred from the freezer to a bleach solution for 18-20
176 hours to discolour any organics to aid in identifying plastics. Once bleached, samples were
177 sieved to eliminate items smaller than 425µm. This size cut off was chosen to align with existing
178 plastic monitoring occurring in Nunatsiavut, another region of Inuit Nunangat, and because this
179 size allows accurate visual identification (Song et al. 2015), a more accessible method for future
180 comparisons (Government of Canada 2018). We visually examined sieve contents using a
181 stereo microscope (Olympus SZ61, model SZ2-ILST) with a magnification range of 0.5–12x.
182 Suspected plastics were extracted from the sample and placed into folded filter paper to dry for
183 a minimum of 4 days, until a consistent weight was observed.

184
185 After the drying period, suspected plastics were re-examined under the microscope to further
186 confirm plastic identification. Visually confirmed plastics were classified based on morphology,
187 colour, size class, and whether and what types of erosion were present, including burning and
188 melting (Table 1. See Table S2-S6 for detailed results and category definitions). Mass, length,
189 width, and height of plastics were also recorded. Plastics were classified as microplastics
190 (<5mm), mesoplastics (5-20mm) or macroplastics (>20mm) following other studies in the region
191 for comparability (Baak et al. 2020; Avery-Gomm et al. 2018). Plastics were transferred into
192 labeled scintillation jars for storage and then glass slides for fourier-transform infrared (FTIR)
193 polymer analysis.

194
195 *2.4 Contamination Measures and Controls*

196 To mitigate contamination of samples by airborne microfibers, all equipment (petri dishes,
197 sieves, tools, and the microscope) was rinsed or wiped down with paper kimwipes and tap water
198 filtered by a 333µm mesh identical to that in trawl nets. Cotton lab coats were worn, and hair
199 was tied back. Pinches of clothing fibres were part of contamination samples to account for
200 possible contamination from our clothing. Separate blanks were taken for each sampling
201 day. We used blanks to account for plastic contamination that may come from the air, our
202 clothing, and/or lab equipment. Blanks were taken each day.

203

204 To eliminate contamination plastics from our analysis, sample plastics were compared with
205 those in blanks from both the lab and the ship and any plastics in our sample that were identical
206 to those in the control were eliminated. Criteria for elimination included identical: color,
207 thickness, and “kinkiness” (shape characteristics) for microplastics, and color, morphology, and
208 erosion patterns for non-microplastics such as paint chips or ropes. Any plastics confirmed as
209 contamination were recorded and removed from the sample analysis.

210

211 *2.5 Spectroscopy*

212 Samples were sent to the Surface Science Western Lab at the University of Western Ontario on
213 the homelands of the Anishinaabek, Haudenosaunee, Lūnaapéewak and Attawandaron peoples
214 for FTIR spectroscopy to determine the polymer type of plastics and to validate visual
215 identification. This technique involves shining a beam of many frequencies of light at an object,
216 producing a spectral graph of absorption. Outputs from individual plastics were compared to
217 outputs of known polymer types to determine identity.

218

219 *2.6 Statistical analysis*

220 Two statistical analyses were performed to address two questions: (1) to determine whether the
221 total plastic abundance differed between the two research of Tasiujarjuaq and SW Greenland;
222 and (2) to test the effect of local-scale latitude on plastic abundance within each site.

223

224 To determine whether the difference in abundance between the two sites was significant, we
225 performed a t-test (two-sided) with a 95% confidence interval using the function `t.test` (R
226 v.3.6.0). To determine the power of the t-test we performed a power analysis, using the function
227 `pwr.t2n.test` (R v.3.6.0) for unequal sample sizes. We also conducted a power analysis to
228 predict the number of samples that should be taken in future research to detect a difference in
229 abundance. The power analysis was performed using the function `pwr.t.test` (R v.3.6.0) for
230 equal sample sizes, with a power of 0.8 (standard), a significance level of 0.05, and the effect
231 size of this study (Cohen’s $d = 0.7335$; the absolute mean difference divided by the standard
232 deviation). The certainty (probability) of not committing an error (type 1 or 2) in this test is
233 represented by the “power” of the test, which is low in this case: 0.19. This is primarily due to
234 the small sample size. We performed an ANOVA on total plastic abundance as a function of
235 latitude within each site independently of one another.

236

237 *2.7 Literature review*

238 To compare our findings to other plastic research in the region, the CLEAR research group
239 conducted an exhaustive literature review for all English-language, peer reviewed publications
240 on plastic pollution conducted in Inuit Nunangat and Greenland. To do so, we used Web of
241 Science Core Collection and Scopus for the following terms in topic and title searches: Inuit
242 Nunangat plastic, Arctic plastic, Arctic plastic Canada, Arctic plastic pollution, Arctic
243 microplastic. Using Scopus, the following search terms were used in the title, keywords, and
244 abstract category search feature: 'Arctic AND plastic', 'Inuit Nunangat AND plastic', 'Nunavut
245 AND plastic', 'Nunatsiavut AND Plastic', 'Inuvialuit AND plastic', 'Nunavik AND 'plastic',
246 'Greenland AND plastic', 'Northwest Territories AND plastic', 'Yukon AND plastic', and 'Labrador
247 AND plastic'.

248
249 Titles and abstracts were examined to ensure studies were scientific research on plastic
250 pollution. We also contacted the Nunatsiavut Government and the Nunavut Research Institute
251 for unpublished data sets. Duplicates and papers whose study areas did not fit the geographic
252 area of interest or were not primary studies on plastic pollution were eliminated and all abstracts
253 were reviewed to ensure fit.

254
255 We recovered a total of 18 peer reviewed papers and one grey literature report on plastic
256 pollution from across regions of Inuit Nunangat, Greenland, and adjacent waters. These existing
257 studies were organized based on region and location of study, type of plastic pollution study
258 conducted (e.g. ingestion, surface trawl, benthic, ice core), plastic pollution findings (Table S7)
259 and statements regarding research licensing (Table S8). Because a number of different
260 methods were used in each study, we included a measurement unit reported as well as
261 minimum detection limit so comparison between studies account for key methodological
262 similarities and differences.

263 264 *2.8 Analysis of methods for reconciliation science*

265 A unique feature of this study is its framing of scientific methods and findings in terms of Inuit
266 relationships and Northern governance, one aspect of reconciliation. We analyzed the scientific
267 methods and findings according to key concepts and texts, supported by fluency in literature in
268 the social studies of science (STS), decolonizing research and science in particular (e.g. Wilson
269 2008; Tuck 2009; Smith 2012; Kimmerer 2013), and settler colonial studies (e.g. Byrd 2011;
270 Snelgrove et al. 2014). Key concepts include reconciliation, colonialism and research
271 sovereignty (e.g. Liboiron 2021; Loseto et al. 2020; Carroll et al 2019; Pfeifer 2018; Rodriguez-

272 Lonebear 2016; Coulthard 2014; Walter and Anderson 2013; Gaudry and Lorenz 2018). To
273 avoid a pan-Indigenous analysis that homogenizes thousands of different Indigenous cultures,
274 we foreground locally salient Inuit texts such as Inuit Tapiriit Kanatami's publications on *National*
275 *Inuit Strategy on Research* (2018) and *Negotiating Research Relationships with Inuit*
276 *Communities* (2007), among others.

277

278 2.9 Community Peer Review

279 When CLEAR conducts plastic pollution research on Inuit lands and food webs, we engage in
280 community peer review of findings before we disseminate to academic venues (Liboiron et al.
281 2018. Also see Loseto et al. 2020 and Wong et al. 2020). We shared a manuscript draft with key
282 personnel in research institutions in Nunavut (Nunavut Arctic College, Nunavut Research
283 Institute) and Nunatsiavut (Nunatsiavut Research Centre). Many reviewers provided substantial
284 feedback and are co-authors.

285

286 2.10 Author Order

287 The method to determine author order followed Liboiron et al. (2017). The forms of labour in the
288 study were discussed and ranked, and the people who performed those types of labour were
289 listed. Forms of labour that attended to accountability were ranked highest for this report. Where
290 two or more researchers were ranked identically, we considered issues of equity to order them.
291 All authors as well as the full lab were invited to be part of these discussions, which were
292 recorded. This process was revisited during review and subsequent edits as necessary.

293

294 3. Results

295 3.1 Abundance of plastics in surface waters


296 A total of 12 marine surface water trawls were deployed (Figure 1 & Table S2). Plastic pollution
297 was found in all surface trawls. Plastic abundance ranged between 0.005 and 0.063 plastics/m²
298 and averaged 0.018 plastics/m² across all samples (Figure 1). For lay reference, this represents
299 an average of about 27 pieces of plastic in a body of water the size of a National Hockey
300 League (NHL) ice rink. Southwestern Greenland samples averaged the highest plastic
301 abundance at 0.026 (\pm 0.026 SD) plastics/m² compared to Tasiujarjuaq at 0.014 (\pm 0.008 SD)
302 plastics/m². The highest abundance tow was the northernmost trawl (Trawl 9) in the Davis
303 Straight/Labrador Sea by Qaqortoq, Greenland at 0.063 plastics/m² (Figure 1 & Table S2).

304

305 **Table 1: Abundance of plastics in surface water in Tasiujarjuaq, Nunavut and Southwest**
 306 **Greenland**

Site	Sample size	Total Plastic Abundance		
		Median (#/m ²)	Mean (#/m ²)	SD
Tasiujarjuaq, Nunavut	8	0.0126	0.0143	0.0082
Southwest Greenland	4	0.0182	0.0262	0.0260

307



308
309

Figure 1. Abundance of plastics in surface water near Iqaluit and Qaqortoq. The map shows the

310 relative abundance of plastics (particles/m²) for each surface water trawl. For a more detailed
 311 map of trawl sites near Iqaluit, see S10.

312
 313

314 *3.2 Characteristics of plastics*

315 Samples yielded a total of 42 different plastics, 22 of which were from Tasiujarjuaq and 20 of
 316 which came from southwest Greenland. Plastics were arranged by site and categorized by size,
 317 morphology, polymer type, colour, and erosion. The majority of plastics were microplastics
 318 (81.0%, n=34) between 0.45 and 5mm in size (Table 1). This size trend was consistent across
 319 both study regions, including 82% of plastics from Nunavut (n=18) and 80% of plastics from
 320 Greenland (n=16). Mesoplastics between 5-20mm comprised 14.3% of all plastics (n=6), the
 321 majority of which were threads. Only two macroplastic pieces larger than 20mm were collected:
 322 one green polyacrylonitrile thread collected from southern Greenland and one clear
 323 polypropylene film from Tasiujarjuaq, Nunavut.

324

325 **Table 2: Morphology and size of plastics near Iqaluit in Tasiujarjuaq (Frobisher Bay),**
 326 **Nunavut and southwest Greenland near Qaqortoq**

Location	Film	Foam	Fragments	Microbeads	Microfibers	Pellets	Threads	Macro	Meso	Micro
Tasiujarjuaq, Nunavut	2	5	11	1	0	0	3	1	3	18
Southwest Greenland	2	0	10	0	0	0	8	1	3	16
Total	4	5	21	1	0	0	11	2	6	34

327
 328

329 Fragments comprised the majority of all plastic types collected (50%, n= 21), followed by
 330 threads (26%, n=11), foam (12%, n=5), film (10%, n= 4), and a single microbead (2%). No
 331 microfibers were recovered, which is common when using a 0.45mm cut off size. While both
 332 regions had similar percentages of fragments and film, they varied in the abundance of other
 333 morphologies. All foam (n=5) was found in Frobisher Bay, for example, and more threads were
 334 found near Greenland (Table S2).

335

336 The most abundant colour was red (26%, split evenly between the two study sites). White and
 337 black were the second most plentiful at 12% of the total (Table S3). All plastics (100%, n= 42)
 338 had signs of erosion or wear. More than one erosion pattern can occur on a single plastic, and

339 75 erosion notes were gathered on the 42 plastics (Table S4). The most common erosion type
 340 was fraying, which was related to the high number of plastic threads recovered. Among plastics,
 341 19% were discoloured, 19% were pitted, 14% were fragmented, and 9% were stretched. All
 342 other erosion patterns were noted on 5% of plastics or less. Four, all from Tasiujarjuaq, were
 343 melted and/or burned.

344

345 All suspected plastics were submitted for FTIR analysis to determine polymer type (Table 2).
 346 Four threads were cellulosic and likely from non-plastic textile sources (Athey et al. 2020). Two
 347 plastics could not be identified, and 39 plastics were identified as synthetic polymers. The most
 348 common polymer was polyethylene (21%, n=8), followed by epoxy blends (18%, n = 7), and
 349 polystyrene (15%, n = 6), though all polystyrene was recovered from Tasiujarjuaq. Table 2
 350 shows polymer types at each location. We also include common uses of each polymer type in
 351 anticipation of diverse users of this data.

352

353 **Table 3: Polymer types of plastics recovered in surface water in Tasiujarjuaq near Iqaluit**
 354 **and the Labrador Sea near Qaqortoq (SW Greenland)**

Polymer type	Common uses	Tasiujarjuaq (count)	Tasiujarjuaq (%)	SW Greenland (count)	SW Greenland (%)	Total (count)	Total (%)
Acrylonitrile butadiene styrene (ABS)	Automotive components, pipe fittings, consumer goods (like Lego)	0	0%	0	0%	0	0%
Acrylate	Cosmetics, paints, diapers, textiles	0	0%	1	5%	1	3%
Alkyd Resin	Paints and varnishes, furniture and architectural coatings	0	0%	0	0%	0	0%
Acrylonitrile styrene acrylate (ASA)	Automotive components, home appliances	0	0%	1	5%	1	3%
Polyamide/Nylon (PA)	Textiles, carpets, automotive components, fishing gear	1	5%	0	0%	1	3%
Polyacrylonitrile (PAN)	Textiles, cement reinforcement	0	0%	3	16%	3	8%
Polyethylene (PE)	Packaging, trash bags, wire and cable insulation	6	30%	2	11%	8	21%
Polyethylene terephthalate (PET)	Packaging, soda pop bottles, textiles	0	0%	2	11%	2	5%
Polypropylene (PP)	Packaging, textiles, medical applications, fishing gear	2	10%	3	16%	5	13%

Polystyrene (PS)	Foam packaging and coolers, medical applications	6	30%	0	0%	6	15%
Polyurethane (PUR)	Coatings, spray and rigid foams common in construction, transportation, and furniture	0	0%	0	0%	0	0%
Silicone	Sealants and adhesives, medical applications	0	0%	0	0%	0	0%
Epoxy blends	Sealants and adhesives	2	10%	5	26%	7	18%
Other polymer	-	3	15%	2	11%	5	13%
Totals		20	100%	19	100%	39	100%

355
356

357 3.3. Spatial distribution of plastics between and within study sites

358 There was no significant difference in total plastic density between both locations ($p = 0.4335$.
359 $DF = 3.3022$). In short, the difference in averages between the abundance of plastics in
360 Tasiujarjuaq and southeast Greenland cannot be differentiated from chance (H_2 is null).
361 However, this test had a power of 0.19, indicating a high potential for type 2 error. This result is
362 somewhat surprising given that the currents and land use patterns for each site are different. A
363 power analysis shows that a minimum sample size of 30.2 samples (or 30 trawl tows per site)
364 would allow future studies to detect a significant difference from these findings.

365

366 The ANOVA test also showed that abundance within each site was not related to local-scale
367 latitude, which was also somewhat unanticipated, especially given that in Tasiujarjuaq, the
368 location of the City of Iqaluit is located further north in the bay and thus is anticipated to be a
369 source of urban plastic pollution (H_1 is null) (p -value was > 0.05 . It was 0.662 for Tasiujarjuaq
370 and 0.04738 for SW Greenland). The F value was 0.2112 for Tasiujarjuaq and 0.7658 for SW
371 Greenland.

372

373 3.4 Comparison of findings with existing studies in the literature

374 The systemic literature search for all research on plastic pollution in the Arctic resulted in 1,116
375 title results, of which 1,099 were excluded because they were not primary research on plastic
376 pollution (merely mentioning the Arctic and plastics in passing), or were outside of the study
377 area, or were duplicates. One additional grey literature report was found by contacting Inuit
378 governments.

379

380 The final 18 English-language, published studies on plastic pollution in Inuit Nunangat and
381 Greenland (Table S7) were dominated by ingestion studies (66%, $n = 12$), almost exclusively in

382 birds (61%, n=11). Ingestion studies comprised all of the studies in Inuit Nunangat regions of
 383 Nunavut (100%, n= 8), Nunatsiavut (100%, n= 1) and Inuvialuit (100%, n=1), while a greater
 384 variety of plastic pollution research has been conducted outside of Inuit Nunangat in Greenland,
 385 including ingestion (50%, n= 3), subsurface water trawl (33%, n= 2) and ice core studies (17%,
 386 n=1). Studies conducted in polar areas adjacent to Inuit Nunangat (e.g. Labrador Sea, Arctic
 387 Central Basin) likewise included more diverse study types including ingestion (33%, n=2),
 388 surface water (17%, n= 1), subsurface water (17%, n=1), ice core (17%, n=1), and benthic
 389 studies (17%, n=1). Amongst all studies examined, over half of sample locations were in
 390 Nunavut (51.2%, n= 32) and in the Qikiqtani Inuit Region in particular (38%, n=24), which
 391 include prominent migratory bird monitoring sites such as Prince Leopold Island (Provencher et
 392 al. 2009, 2010; Poon et al. 2017).

393
 394 We conducted a comparison of plastic abundance among water studies (i.e. surface water,
 395 subsurface water, ice) in all of Inuit Nunangat and the nearby Arctic and subarctic in relation to
 396 the findings in this study (Table 4, S11). Findings from published studies show a greater
 397 abundance of plastics at higher latitudes in the Arctic, and particularly high concentrations in
 398 sea ice (S11). The average abundance in our study is 0.018 plastics/m², lower than an average
 399 abundance of 0.063 plastics/m² found by Cózar et al. (2017) in the Greenland and Barents Seas
 400 north of our study sites using similar surface water methods. Subsurface water studies in
 401 Ukalqarteq (Amelineau et al. 2016), Northeastern Greenland (Morgana et al. 2018), and the
 402 Arctic Central Basin (Kanhai et al. 2018) also found higher abundances further north of our
 403 study area but used different methods and are not directly comparable (Table 4). Studies find a
 404 lower abundance in plastics south of our sites: published studies that used identical methods to
 405 our own shows Nunatsiavut to the south averaged 0.008 plastics/m² (Liboiron et al. 2020),
 406 followed by the more southerly region of Newfoundland and Labrador outside of the Arctic,
 407 which averaged 0.007 plastics/m² of surface water (Liboiron et al. 2020).

408
 409 **Table 4. Plastic density figures from surface water, subsurface water, and sea ice core**
 410 **studies in waters surrounding Inuit Nunangat and Greenland**

Citation	Year	Environment	Sample depth (m)	Location	n	(#/unit)	Unit	Detection limit (mm)
Cózar et al., 2017	2013	surface water	0.15	"Rest of Arctic Ocean"	42	0.0000	m ²	0.5
Liboiron et al. 2020	2017	surface water	surface	Ramah Bay	1	0.0000	m ³	0.425
Liboiron et al. 2020	2017	surface water	surface	L'Anse aux Meadows	3	0.0049	m ²	0.425
Liboiron et al. 2020	2017	surface water	surface	Indian Island	2	0.0074	m ²	0.425

Liboiron et al. 2020	2017	surface water	surface	Grady Island,	3	0.0084	m ²	0.425
Liboiron et al. 2020	2017	surface water	surface	Nunajnguk (Nain)	4	0.0093	m ²	0.425
Liboiron et al. 2020	2017	surface water	surface	Tongait Kakkasuangita SilakKijapvinga	1	0.0094	m ²	0.425
Liboiron et al. this study	2018	surface water	surface	Frobisher Bay Nunavut	8	0.0144	m ²	0.425
Liboiron et al. this study	2018	surface water	surface	East Greenland	4	0.0263	m ²	0.425
Cózar et al., 2017	2013	surface water	0.15	"Greenland and Barents Seas"	42	0.0630	m ²	0.5
Kanhai et al. 2018	2016	subsurface water	8.5	"Arctic Central Basin"	58	0.9700	m ³	0.25
Amelineau et al., 2016	2005	subsurface water	50	Ukalqearteq	18	0.9900	m ³	0.5
Amelineau et al., 2016	2014	subsurface water	50	Ukalqearteq	20	2.3800	m ³	0.5
Morgana et al. 2018	2018	subsurface water	6	Northeastern Greenland	7	2.4000	m ³	0.08
Peeken et al., 2018	2014	sea ice	-	Fram Strait	1	4.1000	m ²	0.011
Obbard et al. 2014	2014	sea ice	1.07	"Arctic Ocean"	1	50	L	-
Obbard et al. 2014	2014	sea ice	1.35	"Arctic Ocean"	1	238	L	-

411

412

413 4. Discussion

414 4.1 Spatial trends of plastics abundance

415 Arctic and subarctic regions (north of 60° latitude), which include many Inuit communities, are
416 less populated than locations further south and there are no sources of plastic production in the
417 Arctic. Yet, both models (van Sebille et al. 2012) and published studies (Obbard et al. 2014;
418 Lusher et al. 2015; Bergmann et al. 2016; Cozar et al. 2017) find high plastic accumulation in
419 the northern and easternmost areas of the Greenland and Barents seas and the Arctic Polar
420 Circle generally. For this reason, we chose two sites at relatively similar latitudes (between 60-
421 63°) to compare the abundance of plastics in surface water in each, and to locations further
422 north and south (via other published literature).

423

424 We did not find a statistically significant spatial trend of abundance within either site, even
425 though in Tasiujarjuaq one cluster of trawl tows were close to Iqaluit and its landfill, which is
426 located on a cliff overlooking the ocean, and the others ran out into the bay including two tows
427 near the mouth of the bay. We expected a statistically significant increase of abundance close
428 to urban centers (Iqaluit population is 7,740) (City of Iqaluit, 2021) and its landfill as per H1, but
429 this was not the case. While proximity to urban centers and latitude are not synonymous, in the

430 specific case of Tasiujarjuaq they have an equivalency. Based on our statistics, differences in
431 abundance between trawls, as a function of latitude, was likely due to chance.

432

433 There was also no a statistically significant difference in the two abundance measures between
434 Tasiujarjuaq and southwest Greenland (p -value = 0.4335) (H_2). However, the number of
435 samples in this study makes the certainty of these measures relatively low. More samples (30
436 per site, based on power analysis) would allow not only more certainty, but would be able to
437 show a demonstrable change in future studies. It should be highlighted that deploying 30 trawls
438 per site, at 30 minutes each, would represent an important time commitment during any
439 expedition, particularly during inter-disciplinary surveys, and in practice it might be challenging
440 to obtain such elevated number of samples.

441

442 The literature review allowed us to situate our findings within other studies of plastic abundance
443 in water and ice further north and south of our sites. Our study found abundances of
444 microplastics that are similar to those found in Arctic surface waters at comparable latitudes,
445 such as off the coast of Svalbard, Norway (0.028 plastics/ m^2), though this study sampled
446 subsurface water and had a lower cut off size than our study (Lusher et al. 2015). As mentioned
447 above, our results show a lower abundance of plastics than in the suspected plastic
448 accumulation zone further north near the Greenlandic and Barents seas using similar surface
449 water methods and cut off sizes (0.0630 plastics/ m^2) (Cózar et al. 2017; also see van Sebille
450 2012 for a model). Subsurface water studies in Ukalqarteq (Amelineau et al. 2016),
451 Northeastern Greenland (Morgana et al. 2018), and the Arctic Central Basin (Kanhai et al. 2018)
452 also found higher abundances further north of our study area but used different methods and
453 are not directly comparable (Table 4). However this trend in higher abundance at northern
454 latitudes is complicate by the study by Cózar et al. (2017), which included 42 sites across the
455 Arctic Circle, 37% of surface net tows were free of plastic, while none of the tows in this study
456 were plastic-free.

457

458 The average abundance in both our locations was higher than those in adjacent southern Arctic,
459 subarctic and North Atlantic regions including an average of 0.008 plastics/ m^2 in Nunatsiavut
460 and 0.005 plastics/ m^2 near L'Anse aux Meadows just south of Labrador in studies that used
461 identical methods and cut off sizes (Liboiron et al. 2020). Indeed, the highest abundance found
462 in this study (trawls 3 and 6 in Tasiujarjuaq and trawls 9 and 12 in southwest Greenland) were
463 closer to those found in more heavily populated south Pacific waters (0.027 plastics/ m^2) with

464 similar methods (Eriksen et al. 2013). Caution should be exercised when doing these sorts of
465 comparisons, however, as sampling and laboratory methods are not standardized across all
466 studies. For this reason, Table 4 includes the environment sampled, sample depth, number of
467 samples, and the lowest detection size, all of which can influence findings. We have consistently
468 noted where results are more or less comparison due to these different methods in the text
469 above.

470
471 Such a comparison leads us to believe that the effects of latitude on plastic abundance occur
472 over large scales rather than those of single bays or study areas, and our findings do not
473 invalidate research that shows the high Arctic as a sink for long-range transport of plastics that
474 originate further south. Thus, we propose that spatial nuance in studies is important, as studies
475 that consider localities and scales that have meaning to local Arctic residents may well have
476 different spatial patterns than those that consider entire oceans or regions of the Arctic. In the
477 future, a greater number and more even coverage of sampling within a bay such as Tasiujarjuaq
478 is recommended for making meaningful data about local plastic conditions.

479
480 *4.2 Trends of plastics type (morphology, polymer, erosion) and relationships to local or long-*
481 *range sources*

482
483 One of the core questions in a study like this is whether plastics might be local or from long-
484 range sources. This is a difficult question to answer, as it is impossible to definitively determine
485 the source of plastics in nearly all cases. In this study, there are conflicting indications of local
486 and long-range plastics, and it is likely that both sources are represented here.

487
488 The polymer type most commonly found in this study was polyethylene (21%), which likewise
489 comprised the largest percentage of polymers found in studies of surface water in Northwest
490 Greenland (41%) (Morgana et al. 2018) and in Arctic sea ice cores (48%) (Peeken et al. 2018).
491 A similar percentage was found in the Greenland Sea (23%), though in this last case polyester
492 was more abundant (53%) (Amélineau et al. 2016). Polyethylene is commonly found in plastic
493 packaging materials and other 'user' plastics, though this does not indicate that these plastics
494 are from local sources. The prevalence of polyethelene is in line with many other studies and
495 may indicate long-range sources, our percentage (21%) was comparatively low. Uniquely, the
496 polymer types identified in this study were more diverse that other studies in water and ice in the
497 Arctic and we found an unusually high prevalence of epoxy blends (18% of the overall study,

498 and 26% of those in SW Greenland), which are not noted in other Arctic water and ice studies in
499 our literature review. Likewise, polystyrene is rarely or never mentioned in these other studies,
500 but it was our second most abundant category (15%), all of which was found in Tasiujarjuaq.

501

502 We found five burned and/or melted erosion patterns on plastics, all from Tasiujarjuaq. As the
503 dump in Iqaluit has been a source of both controlled and uncontrolled burning of plastics in the
504 past (Watson 2014, Varga 2015, Zahara 2015, Zahara 2018), this may indicate the plastic is
505 local in origin. However, other studies have found burned plastics in regions further south where
506 burning is not a part of official waste management (Saliu et al 2018). Avery-Gomm et al., (2016),
507 for example, found that 37% of plastics ingested by dovekies (*alle alle*) on the island of
508 Newfoundland south of this study were burned or melted. It is likely these are local plastics,
509 though we cannot be sure.

510

511 Finally, most studies of Arctic sea water and ice found that blue plastics are the most prevalent
512 (e.g. Obbard et al. 2014; Morgana et al. 2018, Kanhai et al. 2018), but we found that red was
513 the most prevalent (26% of plastics overall, split evenly between sites), followed by white and
514 black (both 12%). Taken together, these findings on polymer type and erosion patterns may
515 indicate a unique profile of plastics in these regions or at this latitude.

516

517 At the same time, there are indications that some of the plastics are likely from long-range
518 transport. We found the presence of erosion on all plastics (100%, n= 42). Following Cozar et
519 al. (2017: 1), “an abundant presence of aged debris” is a possible indicator that plastics
520 “originated from distant sources.” Of course, aged plastics do not necessarily mean they are
521 from distant sources as they could accumulate in local areas over time, and indeed, the freezing
522 and thawing cycles at high latitudes can contribute to fragmentation and other erosion patterns,
523 as can warm and sunny shorelines (Cooper and Corcoran, 2010).

524

525 Cozar et al. (2017) and others have used the high abundance of plastics in the Arctic in
526 juxtaposition to low local populations to argue that Arctic plastics are likely not local. Our
527 findings on abundance are consistent with this hypothesis. Abundance compared to population,
528 ubiquitous erosion patterns, and the lack of statistical evidence that areas closer to Iqaluit have
529 a higher abundance of plastics appear to support the idea that many Arctic plastics are not
530 originating locally (van Sebille et al. 2012), though very likely some are. More research that

531 considers plastics along entire potential pathways into local regions would be able to shed more
532 light on the issue.

533

534 There are several explanations for long-range transport of plastics from the south to the Arctic.
535 Studies have predicted hotspots of plastic accumulation in Arctic waters caused by converging
536 currents (van Sebille et al. 2012), for example. The West Greenland Current is a northerly
537 flowing current with inputs of southern waters (Yang et al. 2016) and is a potential source of
538 plastics found in our samples collected off the coast of Greenland. The West Greenlandic
539 Current merges with the East Greenlandic Current and may indirectly receive plastics from
540 highly populated regions of northern Europe (Morgana et al. 2018) in addition to those brought
541 north from Atlantic regions (Cózar et al. 2017). In Tasiujarjuaq, southerly flowing water from the
542 Baffin Current may also be a source of plastics, given high abundance of plastics found in
543 nearby sea ice (Obbard et al. 2014; Peeken et 2018), likely deposited from Pacific waters. Yet,
544 other studies hypothesize that plastics in the Greenland Sea originated from the higher Arctic
545 via melting sea ice traveling on the East Greenland current rather than originating from the
546 south directly (Amelineau et al. 2016). See S13 for a map of currents.

547

548 Other potential sources of plastic pollution include atmospheric transport (Bergmann et al. 2019)
549 and biotransport via migrating animals (Provencher et al. 2010; Mallory et al. 2006), which
550 disproportionately deposit plastics in Arctic regions (Bourdages et al. in press; Evangelidou et al.
551 2020). At the same time, there are complex oceanographic factors at play in the circulation of
552 plastics in and to the Arctic. This includes the reduced buoyancy of material in cold water and
553 Enders et al., (2015) have suggested a strong dispersal of small plastics throughout the surface
554 mixed layers. Cozar et al., (2017) have noted that forces impacting the density of water,
555 including melting ice creating a freshwater layer and differences in salinity, may impact transport
556 as well as where in the water column plastics accumulate. Future studies might investigate both
557 surface and subsurface water for plastics and consider a full range of indicators of local and
558 long-range transport.

559

560 *4.3 Reflection on research relationships to Inuit Nunangat in this study*

561 Inuit Tapiriit Kanatami is an Inuit-led non-profit organization that protects and advances the
562 rights and interests of Inuit in Canada, including in research. Their *National Inuit Strategy on*
563 *Research* (2018) makes the legacy and context of research in Inuit Nunangat clear: “The
564 relationship between Inuit and the research community is replete with examples of exploitation

565 and racism. Research has largely functioned as a tool of colonialism, with the earliest scientific
566 forays into Inuit Nunangat serving as precursors for the expansion of Canadian sovereignty and
567 the dehumanization of Inuit. Early approaches to the conduct of research in Inuit Nunangat cast
568 Inuit as either objects of study or bystanders. This legacy has had lasting impact on Inuit and it
569 continues to be reflected in current approaches to research governance, funding, policies, and
570 practices” (2018: 5. Also see Smith 2012). We and many other researchers understand that this
571 legacy is not in the past and signs of it exist in this study. This is not a legacy that individual
572 researchers or research organizations can opt out of, even if we are respectful in other ways
573 such as using local place names or fostering anti-racist cultures in our workplaces. The moment
574 researchers decide to pursue research in Indigenous territories and homelands, we inherit these
575 legacies and we work within this context (O’Brien 1993; McGregor 2004; Tuck 2009; Smith
576 2012; Moffitt et al. 2015). For this reason, the discussion below highlights the ways that normal
577 aspects of research, from permitting to categorization of plastics, are engaged in these
578 relationships, usually in ways that are difficult to see because they are part of scientific norms.
579 We aim to bring some of these contexts and legacies to light so that we, as settler and non-Inuit
580 researchers, can actively make choices to conduct reconciliatory science that align with Inuit
581 Tapiriit Kanatami’s call to change research in Inuit Nunangat.

582

583 4.3.1 *Permissions and permitting*

584 Nearly all areas in Inuit Nunangat have research permitting and permission processes in place
585 (Table S1). Of the 18 peer reviewed articles collected in our systematic literature review, seven
586 (39%) included a statement regarding whether a research permit was obtained (Table S8). Two
587 of these were in the methods section rather than the acknowledgments. Of these, five (28%)
588 mentioned whether a permit from an Inuit Nunangat research centre was obtained. Others
589 included statements about federal permits, boating permits, or statements of community
590 consultations and partnerships. A third (n= 6, 33%) of all papers did not include any type of
591 statement on research permitting or permissions of any kind. This does not mean permits and
592 permissions were not sought and obtained, but it does indicate that the formal protocols of
593 permissions, permits, and consent as highlighted by Inuit Tapiriit Kanatami (2018) and Inuit
594 researchers (Pederson et al. 2020; Bull and Hudson 2019) are not a core part of research
595 discussions in one of the most authoritative places that research is discussed: peer reviewed
596 publications.

597

598 Though all researchers in this study obtained required permits and permissions, CLEAR has
599 been asked by others to process samples collected in Inuit Nunangat without permits or to
600 reanalyze samples when reanalysis was not part of the original permit. We decline to process
601 samples obtained without permit. It is clear to us that permits and permission for research in
602 Indigenous homelands is not to be taken for granted as common knowledge or practice. We
603 encourage researchers to report Indigenous permits in methods sections to formalize their
604 critical nature in doing research, rather than relegating them to a “helping” role as per
605 acknowledgements (Loseto et al. 2020).

606

607 Following research permits and permission processes is a simple step establishing good
608 research relations, but at the same time it must be stressed that in many parts of Inuit
609 Nunangat, permits and permissions are carried out by territorial or other settler state bodies that
610 do not necessarily have Inuit staff, nor do they necessarily consult with Inuit. If they do, there is
611 often no formal requirement to heed Inuit requests in the permit. Indeed, there are cases where
612 research permits were granted in Nunavut despite the protests of Inuit (Qikiqtani Inuit
613 Association. V. Canada 2010; Riddell-Dixon 2011). The role of Inuit in decision-making around
614 research in their homelands is uneven across Inuit Nunangat and is in constant development.

615

616 4.3.2 Personnel

617 Inuit Tapiriit Kanatami argues that “the primary beneficiaries of Inuit Nunangat research
618 continue to be [non-Inuit] researchers themselves, in the form of access to funding, data and
619 information, research outcomes, and career advancement” (ITK 2018: 5). This observation held
620 for this study: we mapped the locations (“research bases”) of first authors in all papers in our
621 comparative analysis of plastics in water in the Eastern Arctic (Figure 3). 100% of leads are
622 based in the south.

623

624

Places involved in marine plastic surface water research in and around Inuit Nunangat & the Arctic



625
 626 Figure 2: Places associated with marine surface water plastic research in the Eastern Arctic,
 627 including research sites, researcher home bases for first authors on published research on
 628 plastics in water in the region, and settlements in homelands. Inuit Nunangat is coloured for
 629 reference with different regions of Inuit Nunangat in different shades (Nunatsiavut in orange,
 630 Nunavut in yellow, and Nunavik in green), and settlements in Inuit Nunangat are coloured
 631 orange. The map shows a clear trend of researchers in the south producing all research in the
 632 north germane to this study.

633
 634

635 This study only brought in Inuit and Northerners at the end, rather than at the beginning, which
 636 is not ideal. As contributors to this study, some consented to be co-authors. Several studies in
 637 our literature review mention working with Inuit hunters and guides, but few are listed as co-
 638 authors or appear to be considered researchers despite these activities generating samples and
 639 knowledge crucial to the success of scientific studies and despite the existence and excellence
 640 of Inuit researchers (Sawatzky et al. 2020). Our author list addresses the way northern and Inuit
 641 intellectual labour is often dismissed and unnamed in dominant scientific culture.

642

643 An investigation into the impacts of research personnel in plastic pollution research in the
644 eastern Arctic yielded an unexpected outcome: while the majority of studies in the circumpolar
645 Arctic are on surface water, in Inuit Nunangat they are on plastic ingestion (of 65 sites, 77%),
646 and mainly in birds (54% of all species and sites)(Table S7). We hypothesize that this is due to
647 the low diversity of published researchers working in this area, resulting in an overall regional
648 skew of knowledge in the direction of particular research team's or leads' interests and skills,
649 even when they work with Inuit partners.

650

651 *4.3.3 Categorization of plastics*

652 Scientific categories can seem benign, exempt from social relationships by simply describing
653 the natural world. However, categories and their standardization dictate what is counted and
654 what is not, what is considered the best unit of measurement, and how different things are
655 grouped together into measurable entities through categorization (Pine & Liboiron 2015; Bowker
656 2000; O'brien 1993; Porter 1996). For example, the categories of "user" and "industrial" plastics
657 are common in the literature (Eriksen et al 2020; Provencher et al. 2010) but when used in Inuit
658 Nunangat, they infer that plastics are locally sourced and users are Inuit despite the vast
659 majority of these likely arriving from hundreds or thousands of kilometers away (Obbard 2018).
660 Moreover, research on waste and disposability argues that the production of "user" plastics are
661 expressions of industry production goals and their circulation are not usually based on user
662 behaviour, but rather on waste infrastructure and investment (MacBride 2011; Lepawsky 2018).
663 All plastics have roots in industrial extraction, design, production, and circulation regardless of
664 consumer choice (Liboiron 2016; Amélineau et al. 2016). This mismatch between the agency of
665 industry versus end consumers as sources of plastic pollution is amplified in the North, as
666 consumer choice, the ability to choose between disposable packaging or other forms of plastic
667 waste or not, and recycling and composting are often unavailable (Keskie et al. 2018; Liboiron
668 2018b ECCC 2017; exp Services 2014; Dawley 2013; Eisted & Christensen 2013; Cantin et al.
669 2012; Arktis Solutions 2011). Categorization should match this reality.

670

671 This may seem like an insignificant detail, but Inuit Tapiriit Kanatami opens their case for a
672 *National Inuit Strategy on Research* with the argument that evidence provided by research is
673 foundational to strong public policies, decision making, and governance (2018: 4).


674 Measurements and their categories are related to governance and therefore to sovereignty.

675 Sovereignty is not only about the right to self-determination and self-governance, but also the

676 ability to access the resources, including data, to govern well (Carroll et al. 2019; Rodriguez-
677 Lonebear 2016). The use of user and industrial categories make implicit arguments about
678 sources and thus intervention. Sociologist Stephen Lukes refers to this latent or potential
679 shaping of governance through the measurements it depends on as a “mobilization of bias,” or
680 the ability to shape agendas before overt political conflict even emerges (in Scott 1998: 58).
681 This view is summed up in the truism, “You can’t manage what you can’t measure” and its
682 inverse, “you can only manage what you do measure.” Indeed, for this reason we have included
683 a large number maps, detailed supplementary material, lay understandings of measures
684 (number of plastics in a hockey rink), checklists, and transparency in data—we anticipate that if
685 Inuit groups wish to use this data for decision-making, it will have to be complete enough for
686 such uses.

687
688 The relationship between scientific categories and evidence-based governance is why “threads”
689 appear as a morphological category for plastics in this study. While plastic pollution research in
690 the 1970s included the category of threads, more recent morphological standardizations do not
691 (e.g. Rochman et al. 2019). For many places this makes sense. Yet, threads (Figure 3) are the
692 second highest category of plastic type in this study, accounting for 26% of plastics overall
693 (Table 1). Threads typically originate from fishing gear (Saturno 2020, Richardson et al., 2019).
694 As such, our lab includes the category in all studies occurring in areas with high fishing activity
695 to guide the governance of likely sources of plastic pollution.

696



Trawl 12, SW Greenland

Trawl 8, Frobisher Bay

697

698 Figure 3: Examples of threads from this study, which likely originate from fishing gear. Threads
 699 are sturdier than microfibers, even though they may be small. They are less kinked and often
 700 occur in woven bundles. Fraying of the ends is a common erosion pattern. Squares in images
 701 are 1mm x 1mm.

702

703


704 *4.4 Research as a source of waste and pollution*

705

706 Research activities, usually conducted with gear and personnel from the south (Figure 2), are a
 707 known source of plastic pollution and waste transport in Inuit Nunangat. In their baseline study
 708 of surface water plastics in the Southern Ocean, Suaria et al. (2020) found that more than half
 709 (58%) of plastics collection via neuston samples were paint fragments from survey ships. They
 710 explain, “Although most presumably came from our survey vessels, paint chips are continuously
 711 generated during ongoing repair, maintenance and cleaning of all ship decks and
 712 superstructures, including research vessels, cruise liners and fishing boats” (p. 6). A study by
 713 Caylarde et al. (2021) found that while paint fragments “were 30 times more abundant than
 714 other plastic particles in the surface waters around the Antarctic peninsula,” they are often not
 715 included in estimates of plastics abundance, making them a potential underestimated and
 716 understudied source of plastics (3). Other studies have found that the paint from research
 717 vessels are a source of pollution stemming from research activity (Lacerda et al. 2019; Song et
 718 al. 2014), and Peeken et al. (2018) found 27.2% of all microplastics in their study of the Arctic
 719 Ocean were acrylic or varnish, likely indicative of paint. In this study we found paint chips in both
 720 study locations; two acrylate fragments found in southwest Greenland did not match any

721 sources of plastics on the research vessel, but four red fragments that FTIR classified as “other
722 commercial polymers” were red on one side, with yellow or rust colours on the other (Figure 4).
723 The *Amundsen’s* hull is red and they may have originated from our research vessel. While none
724 of the chips were identical to the ones on our contamination samples/blanks, the chips we
725 archived account for only a small portion of the ship’s paint. Any materials brought on research
726 expeditions from the south to Inuit Nunangat left in the environment, from trash to sewage, is
727 also part of the transport of waste from the south to the north through research activities. As
728 one co-author based in Iqaluit can attest, researchers routinely leave plastic research
729 disposables such as baggies and vials as well as toxic research materials such as acetone in
730 Nunavut, where there are no safe methods of disposal. Other researchers have noted the
731 extreme carbon footprint of Arctic research based in the south (Brook 2009).

732



733

734 Figure 4: Plastics fragments from surface water trawls likely to be paint chips. The squares in
735 each image are 1mm x 1mm.

736

737 We suggest that researchers include research-based waste, pollution, and contamination as a
738 form of harm within their research applications and designs. We also recommend researchers
739 take samples of paint from their research vessels, not to eliminate paint as a form of
740 contamination in samples but to compare to trawl samples as a potentially identifiable point
741 source of marine plastic pollution.

742

743 **5. Conclusion**

744 This paper has two distinct, yet inextricably interrelated and dependant parts: (1) a traditional
745 scientific study of plastic pollution in surface waters that aligns with other studies, finding that
746 abundance of plastics are greater at higher latitudes in the Arctic; and (2) a reflective analysis of
747 methods and findings with an eye to Inuit relationships and governance in the research area.

748

749 Focusing on these usually unintentional relationships gives us, in the scientific community, the
750 opportunity to recognize that colonial relationships often exist in our work so we can then
751 change relationships towards those characterized by reconciliation and respecting Indigenous
752 sovereignty. While good intentions are a prerequisite to address colonial aspects of research,
753 good intentions alone are not sufficient to challenge embedded colonial research practices; this
754 requires a deliberate and rigorous change in behaviour, practices, and institutional paradigms
755 and policies. Research methodologies are one such route for such change (Wilson 2008;
756 Kovach 2010; Smith 2012; Walter and Andersen 2013; Wong et al. 2020; Liboiron 2021).

757

758 Ensuring we gain permits and permissions to access Indigenous homelands, stepping back to
759 allow and foster Inuit-led researchers and thus to decrease the need for outsider access to Inuit
760 homelands, paying attention to the way categories, metrics, and standards are used in research
761 and their ties to Inuit governance and sovereignty, and framing research findings for use by Inuit
762 are all aspects of doing reconciliatory science (see also S8 for a summary of
763 recommendations). This list should only be understood as a compliment to already existing
764 documents such as the Inuit Tapiriit Kanatami's *National Inuit Strategy on Research* (2018) and
765 *Negotiating Research Relationships with Inuit Communities* (2007) that clearly lay out desired
766 researcher-Inuit relationships, principles, and protocols from an Inuit perspective. Research
767 relations to Indigenous homelands will vary between homelands— what works for Inuit in Inuit
768 Nunangat will almost certainly be different for Kumeyaay in US-Mexico borderlands, for
769 example. Indeed, what works in Nunatsiavut will be different than what is needed in Nunavut,
770 even though both are part of Inuit Nunangat (See Obed 2017; Cunsolo Willox et al. 2012). But
771 we do believe that some of the insights outlined here generalize to other Indigenous homelands
772 in the Arctic and beyond.

773

774 Finally, the framing of action and good relations here falls squarely within definitions of
775 reconciliation, based in “the promise of redemption and resolution” (Stein 2018: 156) in a way
776 that largely leaves existing land relations intact (Daigle 2019). Gaudry and Lorenz (2018)
777 articulate a spectrum of definitions, goals, and actions for indigenization. The first is Indigenous
778 inclusion, into which our recommendation in section 4.3.2. on foregrounding Inuit research
779 personnel falls (also see Anonymous 2019). The second is Indigenous reconciliation, which
780 “locates indigenization on common ground between Indigenous and Canadian ideals, creating a
781 new, broader consensus on debates such as what counts as knowledge, how should

782 Indigenous knowledges and European-derived knowledges be reconciled, and what types of
783 relationships academic institutions should have with Indigenous communities” (219). This is
784 where we see this article falling. Finally, they describe decolonial indigenization, which would
785 reorient “knowledge production based on balancing power relations between Indigenous
786 peoples and Canadians, transforming the academy [and science] into something dynamic and
787 new” (219). Our recommendations fall short of this final goal. That is, our framing of
788 reconciliation science is more a call to reform of existing, dominant system rather than
789 decolonization (Tuck and Yang 2012, Gaudry and Lorenz 2018), #Land Back (Briarpatch 2020;
790 Yellowhead Institute 2019), or even allyship (Indigenous Action 2014). This is a limitation. As
791 such, we understand the reconciliatory moves in scientific research outlined here as the bare
792 minimum of what might count as taking up land acknowledgements and meeting existing
793 requirements of Inuit governance (Wong et al. 2020). Future projects may think about what
794 other relations can take Arctic science further.

795

796 **6. Acknowledgements**

797 We acknowledge the land upon which samples were collected as the homelands of diverse Inuit
798 groups in Inuit Nunangat. The land upon which samples were processed and this article was
799 written is the traditional homelands of the Beothuk, and the larger island of Newfoundland is the
800 homelands of the Mi'kmaq and the Beothuk. Analysis of polymer types conducted by SSW
801 occurred on the traditional homelands of the Anishinaabek, Haudenosaunee, Lūnaapéewak and
802 Attawandaron peoples.

803

804 Thank you to Amundsen Science, chief scientist Dr. Philippe Archambault, and the captain and
805 crew of CCGS *Amundsen* for a successful expedition. Thank you to participants in the equity in
806 author order meeting (Liboiron et al. 2017): Dominique David-Chavez, Jesse Jacobs, Katherine
807 Crocker, Taylor Thompson, Lydia Jennings, Taylor Hess, Noah Hutton, Nadia Duman, Natasha
808 Healey, Molly Rivers, Arif Abu, Doménica Lombeida, John Atkinson, Nicole Power, and Charles
809 Mather. A special thank you to those that reviewed drafts of this article to ensure it was in good
810 relations and standing: Norm Catto and Jackie Price.

811

812 This research was supported by ArcticNet, a Network of Centres of Excellence Canada; an
813 NSERC ship time grant to Edinger et al.; a Social Science and Humanities Research Council
814 (SSHRC) Insight grant to Liboiron; Memorial University's Undergraduate Career Experience

815 Program (MUCEP); and Memorial University's International Student Work Experience Program
816 (ISWEP).

817

818 **References**

819 AMAP/ EU-PolarNet. (2020). *AMAP/ EU-PolarNet International Stakeholder Workshops on*
820 *Research Needs*. Tromsø, Norway: Arctic Monitoring and Assessment Programme.

821 Amélineau, F., Bonnet, D., Heitz, O., Mortreux, V., Harding, A. M. A., Karnovsky, N., Walkusz,
822 W., Fort, J., & Gremillet, D. (2016). Microplastic pollution in the Greenland Sea:

823 Background levels and selective contamination of planktivorous diving

824 seabirds. *Environmental Pollution*, 219, 1131-1139. DOI: 10.1016/j.envpol.2016.09.017

825 Anonymous. (2020). Indigenization is Indigenous. *The Gazette*, February 12.

826 <https://gazette.mun.ca/campus-and-community/indigenization-is-indigenous/>

827 Arktis Solutions. (2011). *Report on Current State of Solid Waste Management and Facilities in*
828 *Nunavut and Cost-Benefit Analysis of Selected Solid Waste Management Approaches*

829 (GN-CGS PROJECT #2010-27). Retrieved from:

830 <https://assembly.nu.ca/library/Gnedocs/2011/000359-e.pdf>.

831 Athey, S. N., Adams, J. K., Erdle, L. M., Jantunen, L. M., Helm, P. A., Finkelstein, S. A., &
832 Diamond, M. L. (2020). The Widespread Environmental Footprint of Indigo Denim

833 Microfibers from Blue Jeans. *Environmental Science & Technology Letters* 7, 840-847.

834 DOI: 10.1021/acs.estlett.0c00498

835 Avery-Gomm, S., Provencher, J. F., Liboiron, M., Poon, F. E., & Smith, P. A. (2018). Plastic

836 pollution in the Labrador Sea: an assessment using the seabird northern fulmar (*Fulmarus*

837 *glacialis*) as a biological monitoring species. *Marine Pollution Bulletin*, 127, 817-822. DOI:

838 10.1016/j.marpolbul.2017.10.001.

839 Baak, J. E., Linnebjerg, J. F., Barry, T., Gavrilov, M. V., Mallory, M. L., Price, C., & Provencher, J.

840 F. (2020). Plastic ingestion by seabirds in the circumpolar Arctic: A review. *Environmental*

841 *Reviews*, 28(4), 506-516. DOI: 10.1139/er-2020-0029

842 Bergmann, M., Sandhop, N., Schewe, I., & D'Hert, D. (2016). Observations of floating

843 anthropogenic litter in the Barents Sea and Fram Strait, Arctic. *Polar Biology*, 39(3), 553-

844 560. DOI: 10.1007/s00300-015-1795-8

845 Bergmann, M., Mützel, S., Primpke, S., Tekman, M. B., Trachsel, J., & Gerdt, G. (2019). White

846 and wonderful? Microplastics prevail in snow from the Alps to the Arctic. *Science*

847 *Advances*, 5(8), eaax1157. DOI: 10.1126/sciadv.aax1157

848 Bourdages, M.P.T., Provencher, J.F., Sudlovenick, E., Ferguson, S.H., Young, B.G., Pelletier,
849 N., Murphy, M.J.J., D'Addario, A., & Vermaire, J.C. (2020). No plastics detected in seal
850 (Phocidae) stomachs harvested in the eastern Canadian Arctic. *Marine Pollution Bulletin*,
851 150, 110772. DOI: 10.1016/j.marpolbul.2019.110772

852 Bourdages, M. P., Provencher, J. F., Baak, J. E., Mallory, M. L., & Vermaire, J. C. (In Press).
853 Breeding seabirds as vectors of microplastics from sea to land: Evidence from colonies in
854 Arctic Canada. *Science of The Total Environment*, 142808. DOI:
855 10.1016/j.scitotenv.2020.142808

856 Bowker, G.C. (2000). Biodiversity Datadiversity. *Social Studies of Science*, 30, 643–683. DOI:
857 10.1177/030631200030005001

858 Briarpatch Magazine. (2020). *The Land Back Issue*. September/October.

859 Brook, R. K. (2009). Ignoring the elephant in the room: the carbon footprint of climate change
860 research. *Arctic*, 62(2), 253-255. DOI: 10.14430/arctic144

861 Bull, J., & Hudson, A. (2019). Research governance in NunatuKavut: engagement, expectations
862 and evolution. *International Journal of Circumpolar Health*, 78(2), 1556558. DOI:
863 10.1080/22423982.2018.1556558

864 Byrd, J. A. (2011). *The Transit of Empire: Indigenous Critiques of Colonialism*. Minneapolis, MN:
865 University of Minnesota Press.

866 Cantin, P., Bury, D., & Vaillancourt, P. (2012). *Inventory and Feasibility Assessment of*
867 *Electronic Waste Recovery in the Northwest Territories: Final Report (068-P-0000378-*
868 *0100-MR-R0100-01)*. Retrieved from: www.dessau.com

869 Carroll, SR, Rodriguez-Lonebear, D and Martinez, A. (2019). Indigenous Data Governance:
870 Strategies from United States Native Nations. *Data Science Journal*, 18(31), 1-15. DOI:
871 10.5334/dsj-2019-031

872 City of Iqaluit. (2021). Demographics. [https://www.iqaluit.ca/visitors/explore-](https://www.iqaluit.ca/visitors/explore-iqaluit/demographics)
873 [iqaluit/demographics](https://www.iqaluit.ca/visitors/explore-iqaluit/demographics)

874 Cooper, D. A., & Corcoran, P. L. (2010). Effects of mechanical and chemical processes on the
875 degradation of plastic beach debris on the island of Kauai, Hawaii. *Marine pollution*
876 *bulletin*, 60(5), 650-654. DOI: [10.1016/j.marpolbul.2009.12.026](https://doi.org/10.1016/j.marpolbul.2009.12.026)

877 Coulthard, G. S. (2014). *Red Skin, White Masks: Rejecting the Colonial Politics of Recognition*.
878 Minneapolis, MN: University of Minnesota Press.

879 Cózar, A., Martí, E., Duarte, C.M., García-de-Lomas, J., Van Sebille, E., Ballatore, T.J., Eguíluz,
880 V.M., González-Gordillo, J.I., Pedrotti, M.L., Echevarría, F. & Troublè, R. (2017). The
881 Arctic Ocean as a dead end for floating plastics in the North Atlantic branch of the

882 Thermohaline Circulation. *Science Advances*, 3(4), e1600582. DOI:
883 10.1126/sciadv.1600582

884 Cunsolo Willox, A., Harper, S. L., Ford, J. D., Landman, K., Houle, K., & Edge, V. L. (2012).
885 "From this place and of this place." Climate change, sense of place, and health in
886 Nunatsiavut, Canada. *Social Science and Medicine*, 75(3), 538–547. DOI:
887 10.1016/j.socscimed.2012.03.043

888 Daigle, M. (2019). The spectacle of reconciliation: On (the) unsettling responsibilities to
889 Indigenous peoples in the academy. *Environment and Planning D: Society and*
890 *Space*, 37(4), 703-721. DOI: 10.1177/0263775818824342

891 Dawley, D. (2013). *Municipal Hazardous Waste Inventory Beaufort Delta/Arctic Pilot Project*
892 (SC437531). Retrieved from:
893 https://www.enr.gov.nt.ca/sites/enr/files/haz_waste_inventory_report.pdf.

894 Eisted, R., & Christensen, T. H. (2011). Waste management in Greenland: Current situation and
895 challenges. *Waste Management and Research*, 29(10), 1064–1070. DOI:
896 10.1177/0734242X10395421

897 Enders, K., Lenz, R., Stedmon, C. A., & Nielsen, T. G. (2015). Abundance, size and polymer
898 composition of marine microplastics $\geq 10 \mu\text{m}$ in the Atlantic Ocean and their modelled
899 vertical distribution. *Marine pollution bulletin*, 100(1), 70-81.
900 doi.org/10.1016/j.marpolbul.2015.09.027

901 Environment and Climate Change Canada (ECCC) (2017). *Solid Waste Management for*
902 *Northern and Remote Communities*. Gatineau, QC: Government of Canada.

903 Eriksen, M., Borgogno, F., Villarrubia-Gómez, P., Anderson, E., Box, C., & Trenholm, N. (2020).
904 Mitigation strategies to reverse the rising trend of plastics in Polar Regions. *Environment*
905 *International*, 139, 105704. DOI: 10.1016/j.envint.2020.105704

906 Eriksen, M., Maximenko, N., Thiel, M., Cummins, A., Lattin, G., Wilson, S., Hafner, J., Zellers,
907 A., & Rifman, S. (2013). Plastic pollution in the South Pacific subtropical gyre. *Marine*
908 *Pollution Bulletin*, 68(1-2), 71-76. DOI: 10.1016/j.marpolbul.2012.12.021

909 Evangeliou, N., Grythe, H., Klimont, Z., Heyes, C., Eckhardt, S., Lopez-Aparicio, S., & Stohl, A.
910 (2020). Atmospheric transport is a major pathway of microplastics to remote regions.
911 *Nature Communications*, 11(1), 3381. DOI: 10.1038/s41467-020-17201-9

912 exp Services Inc. (2014). Nunavut Solid Waste Management Plan. Retrieved from:
913 <http://www.cgs.gov.nu.ca/rfp/Past%20Documents%20Sanikiluaq/Nunavut%20Solid%20Waste%20Management%20Plan.pdf>.

915 Furgal, C.M., Powell, S., & Myers, H. (2005). Digesting the message about contaminants and
916 country food in the Canadian North: a review and recommendations for future research
917 and action. *Arctic* 58(2): 103-114. DOI: 10.14430/arctic404.

918 Gaudry, A., & Lorenz, D. (2018). Indigenization as inclusion, reconciliation, and decolonization:
919 Navigating the different visions for indigenizing the Canadian Academy. *AlterNative: An*
920 *International Journal of Indigenous Peoples*, 14(3), 218-227. DOI:
921 [10.1177/1177180118785382](https://doi.org/10.1177/1177180118785382)

922 Gaylarde, C. C., Neto, J. A. B., & da Fonseca, E. M. (2021). Paint fragments as polluting microplastics: A
923 brief review. *Marine Pollution Bulletin*, 162, 111847.

924 Goldstein, M. C., Titmus, A. J., & Ford, M. (2013). Scales of spatial heterogeneity of plastic
925 marine debris in the northeast Pacific Ocean. *PLoS one*, 8(11), e80020. DOI:
926 10.1371/journal.pone.0080020

927 Government of Canada. (2018). *Community Based Monitoring and Research*. Northern
928 Contaminants Funded Projects 2018-2019. Retrieved from:
929 https://www.ic.gc.ca/eic/site/063.nsf/eng/h_97916.html?Open=1&wbdisable=true#9

930 Indigenous Action. (2014) *Accomplices Not Allies: Abolishing the Ally Industrial Complex*.
931 Indigenous Action Media. Retrieved from: [http://www.indigenousaction.org/accomplices-](http://www.indigenousaction.org/accomplices-not-allies-abolishing-the-ally-industrial-complex/)
932 [not-allies-abolishing-the-ally-industrial-complex/](http://www.indigenousaction.org/accomplices-not-allies-abolishing-the-ally-industrial-complex/)

933 Inuit Tapiriit Kanatami (ITK) (2018). *National Inuit Strategy on Research*. Ottawa, Canada: Inuit
934 Tapiriit Kanatami.

935 Inuit Tapiriit Kanatami and Nunavut Research Institute (ITK& NRI) (2007). *Negotiating Research*
936 *Relationships with Inuit Communities: A Guide for Researchers*. Nickels, S., Shirley, J. &
937 Laidler, G. (eds). Ottawa & Iqaluit: Inuit Tapiriit Kanatami and Nunavut Research Institute.

938 Kanhai, L.K., Gårdfeldt, K., Lyashevskaya, O., Hassellöv, M., Thompson, R.C., O'Connor, I.
939 (2018). Microplastics in sub-surface waters of the Arctic Central Basin. *Marine Pollution*
940 *Bulletin*, 130, 8-18. DOI: 10.1016/j.marpolbul.2018.03.011

941 Keske, C. M., Mills, M., Godfrey, T., Tanguay, L., & Dicker, J. (2018). Waste management in
942 remote rural communities across the Canadian North: Challenges and opportunities.
943 *Detritus*, 2(1), 63-77. DOI: 10.31025/2611-4135/2018.13641

944 Kimmerer, R. W. (2013). *Braiding Sweetgrass: Indigenous Wisdom, Scientific Knowledge and*
945 *the Teachings of Plants*. Minneapolis, MN: Milkweed Editions.

946 Kovach, M. (2010). *Indigenous Methodologies: Characteristics, Conversations, and Contexts*.
947 Toronto, ON: University of Toronto Press.

948 Lacerda, A.L.D.F., Rodrigues, L.D.S., Van Sebille, E., Rodrigues, F.L., Ribeiro, L., Secchi, E.R.,
949 Kessler, F. & Proietti, M.C. (2019). Plastics in sea surface waters around the Antarctic
950 Peninsula. *Scientific Reports*, 9(1), 1-12. DOI: 10.1038/s41598-019-40311-4.

951 Law, K. L., Morét-Ferguson, S., Maximenko, N. A., Proskurowski, G., Peacock, E. E., Hafner, J.,
952 & Reddy, C. M. (2010). Plastic accumulation in the North Atlantic subtropical
953 gyre. *Science*, 329(5996), 1185-1188. DOI: 10.1126/science.1192321

954 Lepawsky, J. (2018). *Reassembling Rubbish: Worlding Electronic Waste*. Cambridge, MA: MIT
955 Press.

956 Liboiron, M. (2016). Municipal versus Industrial Waste: Questioning the 3-97 ratio. *Discard*
957 *Studies*, March 2. Retrieved from: [https://discardstudies.com/2016/03/02/municipal-](https://discardstudies.com/2016/03/02/municipal-versus-industrial-waste-a-3-97-ratio-or-something-else-entirely/)
958 [versus-industrial-waste-a-3-97-ratio-or-something-else-entirely/](https://discardstudies.com/2016/03/02/municipal-versus-industrial-waste-a-3-97-ratio-or-something-else-entirely/) Accessed 10/18/2020.

959 Liboiron, M. (2018a). Waste colonialism. *Discard Studies*, November 11. Retrieved from:
960 <https://discardstudies.com/2018/11/01/waste-colonialism/> Accessed 8/03/2020.

961 Liboiron, M. (2018b). How plastic is a function of colonialism. *Teen Vogue*, December 21.
962 Retrieved from: <https://www.teenvogue.com/story/how-plastic-is-a-function-of-colonialism>
963 Accessed 8/18/2020.

964 Liboiron, M. (2021). *Pollution is Colonialism*. Durham, NC: Duke University Press.

965 Liboiron, M., Ammendolia, J., Winsor, K., Zahara, A., Bradshaw, H., Melvin, J., Mather, C.,
966 Dawe, N., Wells, E., Liboiron, F., Fürst, B., Coyle, C., Saturno, J., Novacefski, M.,
967 Westscott, S., & Liboiron, G. (2017). Equity in author order: a feminist laboratory's
968 approach. *Catalyst: Feminism, Theory, Technoscience*, 3(2), 1-17. DOI:
969 10.28968/cftt.v3i2.28850

970 Liboiron, M., Zahara, A., & Schoot, I. (2018). Community peer review: A method to bring
971 consent and self-determination into the sciences. *Preprints*, 2018060104. DOI:
972 10.20944/preprints201806.0104.v1

973 Liboiron, M., Duman, N., Bond, A., Charron, L., Liboiron, F., Ammendolia, J., Hawkins, K.,
974 Wells, E., Melvin, J., Dawe, N., Novacefski, M. (2020). *Regional Report on plastic pollution*
975 *in Newfoundland and Labrador, 1962-2019*. St. John's, NL: Civic Laboratory for
976 Environmental Action Research (CLEAR), Memorial University.

977 Loseto, L.L., Breton-Honeyman, K., Etiendem, D.N., Johnson, N., Pearce, T.D., Allen, J., Amos,
978 A., Arqviq, J., Baak, J., Bélanger, É. and Bourdages, M. (2020). Indigenous participation in
979 peer review publications and the editorial process: Reflections from a workshop. *Arctic*
980 *Science*, 6(3), 352-360. DOI: 10.1139/AS-2020-0023.

981 Lusher, A. L., Tirelli, V., O'Connor, I., & Officer, R. (2015). Microplastics in Arctic polar waters:
982 the first reported values of particles in surface and sub-surface samples. *Scientific*
983 *Reports*, 5, 14947. DOI: 10.1038/srep14947

984 MacBride, S. (2011). *Recycling Reconsidered: the Present Failure and Future Promise of*
985 *Environmental Action in the United States*. Cambridge, MA: MIT Press.

986 Mallory, M. L. (2008). Marine plastic debris in northern fulmars from the Canadian high Arctic.
987 *Marine Pollution Bulletin*, 56: 1501-1504. DOI: 10.1016/j.marpolbul.2008.04.017

988 Mallory, M. L., Roberston, G. J., & Moenting, A. (2006). Marine plastic debris in northern fulmars
989 from Davis Strait, Nunavut, Canada. *Marine Pollution Bulletin*, 52(7), 813-815. DOI:
990 10.1016/j.marpolbul.2006.04.005

991 McGregor, D. (2004). Coming full circle: Indigenous knowledge, environment, and our future.
992 *The American Indian Quarterly* 28(3), 385-410. DOI: 10.1353/aiq.2004.0101

993 Moffitt, M., Chetwynd, C., & Todd, Z. (2015). Interrupting the Northern research industry: Why
994 Northern research should be in Northern hands. *Northern Public Affairs*, December 3.
995 Retrieved from: [http://www.northernpublicaffairs.ca/index/interrupting-the-northern-](http://www.northernpublicaffairs.ca/index/interrupting-the-northern-research-industry-why-northern-research-should-be-in-northern-hands/)
996 [research-industry-why-northern-research-should-be-in-northern-hands/](http://www.northernpublicaffairs.ca/index/interrupting-the-northern-research-industry-why-northern-research-should-be-in-northern-hands/) Accessed
997 8/18/2020.

998 Morgana, S., Ghigliotti, L., Estévez-Calvar, N., Stifanese, R., Wieckzorek, A., Doyle, T.,
999 Christiansen, J., Faimali, M., & Garaventa, F. (2018). Microplastics in the Arctic: A case
1000 study with sub-surface water and fish samples off Northeast Greenland. *Environmental*
1001 *Pollution*, 242, 1078-1086. DOI: 10.1016/j.envpol.2018.08.001

1002 Nunatsiaq News. (2015). Photo: Garbage collectors: volunteers planning Iqaluit clean up. July
1003 23.
1004 [https://nunatsiaq.com/stories/article/65674photo_garbage_collectors_volunteers_planning](https://nunatsiaq.com/stories/article/65674photo_garbage_collectors_volunteers_planning_iqaluit_clean_up/)
1005 [_iqaluit_clean_up/](https://nunatsiaq.com/stories/article/65674photo_garbage_collectors_volunteers_planning_iqaluit_clean_up/)

1006 Obbard, R. W. (2018). Microplastics in polar regions: the role of long range transport. *Current*
1007 *Opinion in Environmental Science & Health*, 1, 24-29. DOI: 10.1016/j.coesh.2017.10.004

1008 Obbard, R. W., Sadri, S., Wong, Y. Q., Khitun, A. A., Baker, I., & Thompson, R. C. (2014).
1009 Global warming releases microplastic legacy frozen in Arctic Sea ice. *Earth's Future*, 2(6),
1010 315-320. DOI: 10.1002/2014EF000240.

1011 Obed, D. (2017). *Illiniavugut Nunami: Learning from the Land Envisioning an Inuit-centered*
1012 *Educational Future*. Master's thesis, Halifax, NS: Saint Mary's University.

1013 O'Brien, M. H. (1993). Being a scientist means taking sides. *BioScience*, 43(10), 706-708. DOI:
1014 10.2307/1312342

1015 Pedersen, C., Otokiak, M., Koonoo, I., Milton, J., Maktar, E., Anaviapik, A., Milton, M., Porter,
1016 G., Scott, A., Newman, C., & Porter, C. (2020). SciQ: an invitation and recommendations
1017 to combine science and Inuit Qaujimajatuqangit for meaningful engagement of Inuit
1018 communities in research. *Arctic Science*, 6(3), 326-339. DOI: 10.1139/AS-2020-0015

1019 Peeken, I., Primpke, S., Beyer, B., Gütermann, J., Katlein, C., Krumpfen, T., Bergmann, M.,
1020 Hehemann, L. and Gerdts, G (2018). Arctic sea ice is an important temporal sink and
1021 means of transport for microplastic. *Nature communications*, 9(1): 1-12. DOI:
1022 10.1038/s41467-018-03825-5

1023 Pfeifer, P. (2018). From the credibility gap to capacity building: An Inuit critique of Canadian
1024 Arctic research. *Northern Public Affairs*, 6(1), 29-34.

1025 Pine, K. H., & Liboiron, M. (2015). The politics of measurement and action. In *Proceedings of*
1026 *the 33rd Annual ACM Conference on Human Factors in Computing Systems*, 3147-3156.
1027 DOI: 10.1145/2702123.2702298

1028 Poon, F. E., Provencher, J. F., Mallory, M. L., Braune, B. M., & Smith, P. A. (2017). Levels of
1029 ingested debris vary across species in Canadian Arctic seabirds. *Marine Pollution Bulletin*,
1030 116(1-2), 517-520. DOI: 10.1016/j.marpolbul.2016.11.051

1031 Porter, T.M. (1996). *Trust in Numbers: The Pursuit of Objectivity in Science and Public Life*.
1032 Princeton, NJ: Princeton University Press.

1033 Provencher, J. F., Gaston, A. J., & Mallory, M. L. (2009). Evidence for increased ingestion of
1034 plastics by northern fulmars (*Fulmarus glacialis*) in the Canadian Arctic. *Marine Pollution*
1035 *Bulletin*, 58(7), 1092-1095. DOI: 10.1016/j.marpolbul.2009.04.002

1036 Provencher, J. F., Gaston, A. J., Mallory, M. L., O'hara, P. D., & Gilchrist, H. G. (2010). Ingested
1037 plastic in a diving seabird, the thick-billed murre (*Uria lomvia*), in the eastern Canadian
1038 Arctic. *Marine Pollution Bulletin*, 60(9), 1406-1411. DOI: 10.1016/j.marpolbul.2010.05.017

1039 Qikiqtani Inuit Association. v. Canada (Minister of Natural Resources). (2010). NUCJ 12. 08-10-
1040 42-CVC. Iqaluit, Nunavut.

1041 Richardson, K., Hardesty, B. D., & Wilcox, C. (2019). Estimates of fishing gear loss rates at a
1042 global scale: A literature review and meta-analysis. *Fish and Fisheries*, 20(6), 1218–1231.
1043 DOI: 10.1111/faf.12407

1044 Riddell-Dixon, E. (2011). Seismic testing in Lancaster Sound: Lessons Learned. *Policy*
1045 *Opinions*, October 1. Retrieved from: [https://policyoptions.irpp.org/magazines/the-new-](https://policyoptions.irpp.org/magazines/the-new-normal-majority-government/seismic-testing-in-lancaster-sound-lessons-learned/)
1046 [normal-majority-government/seismic-testing-in-lancaster-sound-lessons-learned/](https://policyoptions.irpp.org/magazines/the-new-normal-majority-government/seismic-testing-in-lancaster-sound-lessons-learned/)
1047 Accessed 12/02/2020.

1048 Rochman, C.M., Brookson, C., Bikker, J., Djuric, N., Earn, A., Bucci, K., Athey, S., Huntington,
1049 A., Mcllwraith, H., Munno, K., De Frond, H., Kolomijeca, A., Erdle, L., Grbic, J., Bayoumi,
1050 M., Borelle, S.B., Wu, T., Santoro, S., Webowski, L.M, Zhu, X., Giles, R.K., Hamilton, B.M,
1051 Thaysen, C., Kaura, A., Klasios, N., Ead, L., Kim, J., Sherlock, C., Ho, A., & C. Hung.
1052 (2019). Rethinking microplastics as a diverse contaminant suite. *Environmental*
1053 *Toxicology and Chemistry*, 38(4), 703-711. DOI: 10.1002/etc.4371

1054 Rodriguez-Lonebear, D. (2016). 'Building a data revolution in Indian country'. In Kukuta, T. &
1055 Taylor, J. (eds.) *Indigenous Data Sovereignty: Toward an Agenda*, 253-272. Canberra,
1056 AU: ANU Press.

1057 Royal Canadian Geographical Society. (2018). *Inuit: Indigenous peoples atlas of Canada/ Inuit:*
1058 *Atlas des peuples autochtones du Canada*. Retrieved from:
1059 <https://indigenouspeoplesatlasofcanada.ca/section/inuit/> Accessed 12/02/2020.

1060 Saliu, F., Montano, S., Garavaglia, M. G., Lasagni, M., Seveso, D., & Galli, P. (2018).
1061 Microplastic and charred microplastic in the Faafu Atoll, Maldives. *Marine pollution*
1062 *bulletin*, 136, 464-471. DOI: [10.1016/j.marpolbul.2018.09.023](https://doi.org/10.1016/j.marpolbul.2018.09.023)

1063 Saturno, J. (2020). *Investigating the Role of Fishing Gear on Plastic Pollution: The Occurrence*
1064 *of Fishing Gear-Related Plastic Ingested by Atlantic Cod (Gadus morhua) and the*
1065 *Fragmentation of Polymer Ropes*. Master's thesis, St. John's, NL: Memorial University of
1066 Newfoundland.

1067 Sawatzky, A., Cunsolo, A., Jones-Bitton, A., Gillis, D., Wood, M., Flowers, C., Shiwak, I., The
1068 Rigolet Inuit Community Government, & Harper, S. L. (2020). "The best scientists are the
1069 people that's out there": Inuit-led integrated environment and health monitoring to respond
1070 to climate change in the Circumpolar North. *Climatic Change*, 160, 45-66.

1071 Scott, J. C. (1998). *Seeing like a State: How Certain Schemes to Improve the Human Condition*
1072 *have Failed*. New Haven, CT: Yale University Press.

1073 Smith, L. T. (2012). *Decolonizing Methodologies: Research and Indigenous Peoples*. New York,
1074 NY: Zed Books.

1075 Snelgrove, C., Dhamoon, R. K., & Corntassel, J. (2014). Unsettling settler colonialism: The
1076 discourse and politics of settlers, and solidarity with Indigenous nations. *Decolonization:*
1077 *Indigeneity, Education & Society*, 3(2): 1-32

1078 Song, Y. K., Hong, S. H., Jang, M., Kang, J-H., Kwon, O. Y., Han, G. M., & Shim, W. J. (2014).
1079 Large accumulation of micro-sized synthetic polymer particles in the sea surface
1080 microlayer. *Environmental Science & Technology*, 48(16), 9014-9021. DOI:
1081 10.1021/es501757s.

1082 Song, Y.K., Hong, S.H., Jang, M., Han, G.M., Rani, M., Lee, J., Shim, W.J. (2015). A
1083 comparison of microscopic and spectroscopic identification methods for analysis of
1084 microplastics in environmental samples. *Marine Pollution Bulletin*, 93, 202–209. DOI:
1085 10.1016/j.marpolbul.2015.01.015.

1086 Suaria, G., Perold, V., Lee, J. R., Lebouard, F., Aliani, S., & Ryan, P. G. (2020). Floating macro-
1087 and microplastics around the Southern Ocean: Results from the Antarctic
1088 Circumnavigation Expedition. *Environment International*, 136, 105494. DOI:
1089 10.1016/j.envint.2020.105494

1090 Sudlovenick, E. (2019). *A serological survey and Inuit Qaujimagatuqangit of Ringed seals*
1091 *(Nattiit) in Frobisher Bay, Nunavut*. Master's thesis, Charlottetown, PEI: University of
1092 Prince Edward Island.

1093 Stein, S. (2020). 'Truth before reconciliation': the difficulties of transforming higher education in
1094 settler colonial contexts. *Higher Education Research & Development*, 39(1), 156-170.

1095 Tirelli, V., Suaria, G., & Lusher, A. L. (2020). 'Microplastics in Polar Samples' in Rocha-Santos,
1096 T., Costa, M., Mouneyrac, C. (eds.) *Handbook of Microplastics in the Environment*, 1-42.
1097 Cham, Switzerland: Springer Nature Switzerland AG. DOI: 10.1007/978-3-030-10618-8_4-
1098 1

1099 Tuck, E. (2009). Suspending damage: A letter to communities. *Harvard Educational Review*,
1100 79(3), 409-428. DOI:10.17763/haer.79.3.n0016675661t3n15

1101 Tuck, E., & Yang, K. W. (2012). Decolonization is not a metaphor. *Decolonization: Indigeneity,*
1102 *Education & Society*, 1(1), 1-40.

1103 Van Oostdam, J., Donaldson, S.G., Feeley, M., Arnold, D., Ayotte, P., Bondy, G., Chan, L.,
1104 Dewailly, É., Furgal, C.M., Kuhnlein, H., Loring, E., Muckle, G., Myles, E., Receveur, O.,
1105 Tracy, B., Gill, U., & Kalhok, S. (2005). Human health implications of environmental
1106 contaminants in Arctic Canada: a review. *Science of the Total Environment*, 351, 165-246.
1107 DOI: 10.1016/j.scitotenv.2005.03.034

1108 Van Sebille, E., England, M. H., & Froyland, G. (2012). Origin, dynamics and evolution of ocean
1109 garbage patches from observed surface drifters. *Environmental Research Letters*, 7(4),
1110 044040. DOI: 10.1088/1748-9326/7/4/044040

1111 Varga, P. (2015). Iqaluit tries new waste management practices at overloaded dump. *Nunatsiaq*
1112 *News*, May 4.
1113 https://nunatsiaq.com/stories/article/65674iqaluit_tries_new_waste_management_practices_at_overloaded_dump/
1114

1115 Walter, M., & Andersen, C. (2013). *Indigenous Statistics: A Quantitative Research Methodology*.
1116 Walnut Creek, CA: Left Coast Press.

1117 Watson, A. (2014). Iqaluit's long-smouldering 'dumpcano' garbage fire is finally out. *The Globe*
1118 *and Mail*, September 16. [https://www.theglobeandmail.com/news/national/iqaluits-long-](https://www.theglobeandmail.com/news/national/iqaluits-long-smouldering-dumpcano-garbage-fire-finally-out/article20620273/)
1119 [smouldering-dumpcano-garbage-fire-finally-out/article20620273/](https://www.theglobeandmail.com/news/national/iqaluits-long-smouldering-dumpcano-garbage-fire-finally-out/article20620273/)

1120 Watt-Cloutier, S. (2015). *The Right to be Cold: One Woman's Story of Protecting her Culture,*
1121 *the Arctic and the Whole Planet*. Toronto, ON: Penguin Canada.

1122 Wilson, S. (2008). *Research is Ceremony: Indigenous Research Methods*. Winnipeg, MB:
1123 Fernwood Books.

1124 Wong, C., Ballegooyen, K., Ignace, L., Johnson, M., & Swanson., H. (2020). Towards
1125 reconciliation: 10 Calls to Action to natural scientists working in Canada. *FACETS*, 5, 769-
1126 783. DOI: 10.1139/facets-2020-0005

1127 Yang, Q., Dixon, T. H., Myers, P. G., Bonin, J., Chambers, D., van den Broeke, M. R.,
1128 Ribergaard, M. H., & Mortensen, J. (2016). Recent increases in Arctic freshwater flux
1129 affects Labrador Sea convection and Atlantic overturning circulation. *Nature*
1130 *Communications*, 7(1), 10525. DOI: 10.1038/ncomms10525

1131 Yellowhead Institute (2019). *Land Back: A Yellowhead Institute Red Paper*. Toronto, ON:
1132 Yellowhead Institute.

1133 Zahara, A. (2015). *The Governance of Waste in Iqaluit, Nunavut* (Queen's University, CA:
1134 Master's thesis).

1135 Zahara, A. (2018). On sovereignty, deficits and dump fires: Risk governance in an Arctic
1136 'dumpcano. *Inevitably Toxic: Historical Perspectives on Contamination, Exposure, and*
1137 *Expertise*, 259-283.

1138
1139
1140
1141
1142
1143
1144

Supplementary material

Table S1: Research permitting bodies in Inuit Nunangat

Inuit Nunangat region	Licensing Body	Type of research	Governing Legislation	Mandatory
Inuvialuit Settlement Region	Aurora Research Institute & Inuvialuit Land Administration	All research	Inuvialuit Final Agreement and Government of Northwest Territories <i>Scientist Act</i> (double check)	Yes
Nunavut	Nunavut Research Institute	All research	Nunavut Land Claim Agreement & Nunavut <i>Scientist Act</i>	Yes
Nunavik	Nunavik Research Centre	All research	Consultation with institute expected	No
Nunavik	Nunavik Marine Region Wildlife board	Wildlife research in Nunavik Marine Region	Nunavik Inuit Land Claims Agreement	Yes
Nunatsiavut	Nunatsiavut Research Centre	All research	Labrador Inuit Land Claims Agreement & Nunatsiavut Research procedure	Yes

Table S2. Details of all surface water trawls covered in this study.

Location	Year	latitude	longitude	Sample ID	total plastics	Abundance m ²	film	foam	fragments	microbeads	microfibers	pellets	threads	macro	meso	micro	Acrylonitrile butadiene styrene (ABS)	Acrylate	Alkyd Resin	Acrylonitrile styrene acrylate (ASA)	Polyamide (PA)	Polyacrylonitrile (PAN)	Polyethylene (PE)	Polyethylene terephthalate (PET)	Polypropylene (PP)	Polystyrene (PS)	Polyurethane (PUR)	Silicone	Epoxy blends	Other polymer	
Frobisher Bay, Nunavut	2018	63.64688	-68.5221	Trawl-1	3	0.0183	0	0	2	0	0	0	1	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	2	0	
Frobisher Bay, Nunavut	2018	63.67578	-68.5341	Trawl-2	3	0.01409	0	1	2	0	0	0	0	0	0	3	0	0	0	0	0	0	2	0	0	1	0	0	0	0	
Frobisher Bay, Nunavut	2018	63.67017	-68.5981	Trawl-3	4	0.02066	0	3	1	0	0	0	0	0	1	3	0	0	0	0	0	0	1	0	0	3	0	0	0	0	
Frobisher Bay, Nunavut	2018	63.64192	-68.5866	Trawl-4	2	0.01122	0	1	1	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	
Frobisher Bay, Nunavut	2018	63.13268	-67.4407	Trawl-5	1	0.004973	0	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	
Frobisher Bay, Nunavut	2018	62.68736	-66.7709	Trawl-6	6	0.02928	2	0	2	1	0	0	1	1	1	4	0	0	0	0	0	0	3	0	1	0	0	0	0	0	1
Frobisher Bay, Nunavut	2018	62.43105	-65.895	Trawl-7	2	0.01056	0	0	2	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1
Frobisher Bay, Nunavut	2018	62.43132	-65.8976	Trawl-8	1	0.005308	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	
SW Greenland	2018	60.41441	-48.4548	Trawl-9	12	0.06312	2	0	7	0	0	0	3	0	1	1	0	0	0	1	0	1	0	1	1	0	0	0	5	2	
SW Greenland	2018	60.38912	-48.4532	Trawl-10	1	0.005144	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	
SW Greenland	2018	60.36307	-48.4563	Trawl-11	2	0.01109	0	0	2	0	0	0	0	0	0	2	0	0	0	0	0	0	2	0	0	0	0	0	0	0	
SW Greenland	2018	60.33705	-48.4586	Trawl-12	5	0.02535	0	0	1	0	0	0	4	0	2	3	0	1	0	0	0	1	0	1	2	0	0	0	0	0	

Table S3: Colour of recovered plastics

Colour	Nunavut	Greenland	Total plastics	Percentage Total Plastics
Red	6	5	11	26.2%
Brown	1	0	1	2.4%
White	5	0	5	11.9%
Grey	0	1	1	2.4%
Clear	5	2	7	16.7%
Orange	0	0	0	0.0%
Black	2	3	5	11.9%
Yellow	1	3	4	9.5%
Blue	1	3	4	9.5%
Green	1	3	4	9.5%
Totals	22	20	42	100.0%

Table S4: Erosion patters of recovered plastics

Erosion Type	Nunavut	Greenland	Total	% total
Pitted	8	6	14	18.7%
Stretched	2	5	7	9.3%
Frayed	11	13	24	32.0%
Melted	4	0	4	5.3%
Fragmented	7	4	11	14.7%
Discolored	8	6	14	18.7%
Burnt	1	0	1	1.3%
Cracked	0	0	0	0.0%

Table S5: Definitions of plastic morphologies

Morphological category	Definition
Film	Sheet plastics such as plastic bags. However, not synonymous with plastic bags
Foam	Plastics such as Styrofoam. Bounces back to the touch, has air pockets
Fragment	Hard plastic fragments, though they can be flexible. Paint chips included in this category, which are often very brittle
Industrial Pellets	Industrial pre-production pellets or nurdles
Microbead	Small spheres often from cosmetics, usually brightly colored
Microfibre	Thinner and kinked plastics compared to threads. Usually from synthetic fabrics, often too small for erosion patterns
Thread/Fibre Bundle*	Tightly interwoven threads or fibres that are bundled together and are difficult to separate, often noted as conglomerates in the note section in past studies.
Thread	Thick filaments, such as fishing line, large enough to see erosion patterns

*Term from Rochman et al. 2019.

Table S6: Definitions of erosion patterns

Erosion pattern	Definition
Burnt	Burnt parts charred from flame along with visible melting around the burns
Cracked	Fissures in plastic particles without full separation into pieces
Discoloured	Stained or 'dirty' looking parts of plastic particles. Parts of plastics a different colour i.e. not uniform
Fragmented	Splitting into fragments, newly split pieces will have smooth shiny edges where it fragmented
Frayed	Worn at the edges with pieces fraying off
Melted	Visible parts of the plastic are melted and disformed (shiny and shrunken)
Stretched	Visible parts that have been 'stretched' or are thinner than other parts. Common in threads.
Pitted	Often found in fragments, grooves or pits in the plastic's surface, where chunks have been scrapped off

Table S7: Published plastic pollution research conducted in Inuit Nunangat and Greenland

Region	Year	Study type	Location	Lat.	Long.	Environment	Measurement	Plastic contamination	Detection Limit (mm)	Citation
Nunavut										
	2007	Ingestion	Coats Island	63.63	-82.00	Thick-billed murre	% FO	4% (n=25)	N/A	Provencher et al. 2010
	2008	Ingestion	Akpatok Island	60.97	-68.13	Thick-billed murre	% FO	23% (n= 31)	N/A	Provencher et al. 2010
	2008	Ingestion	Digges Sound	62.55	-77.58	Thick-billed murre	% FO	17% (n=30)	N/A	Provencher et al. 2010
	2007	Ingestion	Minarets	67.00	-62.33	Thick-billed murre	% FO	6% (n=30)	N/A	Provencher et al. 2010
	2008	Ingestion	Minarets	67.00	-62.33	Thick-billed murre	% FO	10% (n=20)	N/A	Provencher et al. 2010
	2008	Ingestion	Prince Leopold Island	74.03	-90.00	Thick-billed murre	% FO	8% (n=32)	N/A	Provencher et al. 2010
	2008	Ingestion	Prince Leopold Island	74.03	-90.00	Thick-billed murre	% FO	13% (n=18)	N/A	Provencher et al. 2010
	2008	Ingestion	Devon Island	76.25	-89.25	Northern fulmar	% FO	31% (n=102)	N/A	Mallory 2008
	2008	Ingestion	Arviat	61.11	-96.04	Ringed seal	% FO	0% (n=32)	0.425	Bourdages et al., 2020
	2008	Ingestion	Arviat	61.11	-96.04	Bearded seal	% FO	0% (n=5)	0.425	Bourdages et al., 2020
	2008	Ingestion	Arviat	61.11	-96.04	Harbour seal	% FO	0% (n=1)	0.425	Bourdages et al., 2020
	2018	Ingestion	Arviat	61.11	-96.04	Ringed seal	% FO	0% (n=1)	0.425	Bourdages et al., 2020
	2008	Ingestion	Naujaat	66.53	-86.24	Ringed seal	% FO	0% (n=38)	0.425	Bourdages et al., 2020
	2007	Ingestion	Sanikiluaq	56.54	-79.22	Ringed seal	% FO	0% (n=33)	0.425	Bourdages et al., 2020
	2007	Ingestion	Sanikiluaq	56.54	-79.22	Bearded seal	% FO	0% (n=1)	0.425	Bourdages et al., 2020

2008	Ingestion	Sanikiluaq	56.54	-79.22	Ringed seal	% FO	0% (n=11)	0.425	Bourdages et al., 2020
2018	Ingestion	Iqaluit	63.75	-68.52	Ringed seal	% FO	0% (n=18)	0.425	Bourdages et al., 2020
2019	Ingestion	Iqaluit	63.75	-68.52	Ringed seal	% FO	0% (n=2)	0.425	Bourdages et al., 2020
2008	Ingestion	Prince Leopold Island	74	-90	Northern fulmar	% FO	80% (n=10)	1	Poon et al., 2017
2013	Ingestion	Prince Leopold Island	74	-90	Northern fulmar	% FO	89% (n= 9)	1	Poon et al., 2017
2008	Ingestion	Prince Leopold Island	74	-90	Thick-billed murre	% FO	0% (n=10)	1	Poon et al., 2017
2013	Ingestion	Prince Leopold Island	74	-90	Thick-billed murre	% FO	0% (n= 10)	1	Poon et al., 2017
2013	Ingestion	Prince Leopold Island	74	-90	Black-legged kittiwake	% FO	9% (n= 11)	1	Poon et al., 2017
2013	Ingestion	Prince Leopold Island	74	-90	Black guillemot	% FO	0% (n= 3)	1	Poon et al., 2017
2008	Ingestion	Nuvuttiq (Cape Searle)	67.25	-62.58	Northern fulmar	% FO	87% (n= 15)	N/A	Provencher et al., 2009
2008	Ingestion	Prince Leopold Island	74	-90	Northern fulmar	% FO	80% (n= 10)	N/A	Provencher et al., 2009
2002	Ingestion	Davis Strait	68.43	-61.81	Northern fulmar	% FO	36% (n=42)	N/A	Mallory et al., 2006
1998	Ingestion	Sanikiluaq (Belcher Island)	N/A	N/A	Common eider	% FO	0% (n=388)	0.5	Provencher et al. 2014
2000	Ingestion	Kinngait (Cape Dorset)	N/A	N/A	Common eider	% FO	0% (n= 108)	0.5	Provencher et al. 2014
2011	Ingestion	Kinngait (Cape Dorset)	N/A	N/A	Common eider	% FO	1% (n=100)	0.5	Provencher et al. 2014

2006	Ingestion	Akpatordjuark (Coats Island)	N/A	N/A	Thick-billed murre	% FO	0% (n=15)	0.5	Provencher et al. 2014
2007	Ingestion	Nasaruvaalik Island	N/A	N/A	Arctic tern	% FO	0% (n= 41)	0.5	Provencher et al. 2014
Nunatsiavut									
2006	Ingestion	Nunainguk (Nain)	N/A	N/A	Surf scoter	% FO	0% (n= 38)	0.5	Provencher et al. 2014
2006	Ingestion	Gannet Islands	N/A	N/A	Thick-billed murre	% FO	0% (n=15)	0.5	Provencher et al. 2014
2006	Ingestion	Gannet Islands	N/A	N/A	Common murre	% FO	0% (n=15)	0.5	Provencher et al. 2014
Inuvialuit									
2017	Ingestion	Hendrickson Island	69	-133	Beluga whale	% FO	100% (n=4)	0.02	Moore et al., 2020
2018	Ingestion	Hendrickson Island	69	-133	Beluga whale	% FO	100% (n=3)	0.02	Moore et al., 2020
Greenland									
1991	Excrement	Nansen Land, North Greenland	83	-43	Arctic wolf feces	% FO	3% (n= 110)	N/A	Marquard-Petersen, 1998
1991	Excrement	Hold with Hope, East Greenland	74	-22	Arctic wolf feces	% FO	0% (n=351)	N/A	Marquard-Petersen, 1998
2005	Ingestion	Ukaleqarteq, East Greenland	74.73	-21.58	Little auk	% FO	100% (n=26)	0.5	Amelineau et al., 2016
2014	Ingestion	Ukaleqarteq, East Greenland	74.73	-21.58	Little auk	% FO	100% (n=18)	0.5	Amelineau et al., 2016
1988	Ingestion	Southwest Greenland	N/A	N/A	Thick-billed murre	% FO	6% (n=202)	0.5	Provencher et al. 2014
1988	Ingestion	Nuuk	N/A	N/A	Dovekie	% FO	0% (n= 19)	0.5	Provencher et al. 2014
1997	Ingestion	Hakluyt Island	N/A	N/A	Thick-billed murre	% FO	0% (n= 40)	0.5	Provencher et al. 2014
1999	Ingestion	Nuuk	N/A	N/A	Common eider	% FO	0% (n= 241)	0.5	Provencher et al. 2014

2006	Ingestion	Nuuk	N/A	N/A	Thick-billed murre	% FO	0% (n=15)	0.5	Provencher et al. 2014
2012	Ingestion	Nuuk	N/A	N/A	Common eider	% FO	0% (n= 135)	0.5	Provencher et al. 2014
2005	Trawl	Ukalqarteq	74.73	-21.58	Subsurface water	Plastics / m ³	0.99	0.5	Amelineau et al., 2016
2014	Trawl	Ukalqarteq	74.73	-21.58	Subsurface water	Plastics/ m ³	2.38	0.5	Amelineau et al., 2016
2014	Ice core	Fram Straight	78.27	-14.71	Sea ice	Plastics / N x m ³	4.1 ± 2.0 × 10 ⁶	0.011	Peeken et al., 2018
2018	Trawl	Northeastern Greenland	75.50	-10.87	Subsurface water	Plastics/ m ³	2.4	0.08	Morgana et al., 2018
2018	Ingestion	Northeastern Greenland	76.14	-9.03	Bigeye sculpin	% FO	34% (n= 71)	0.07	Morgana et al., 2018
2018	Ingestion	Northeastern Greenland	76.14	-9.03	Polar cod	% FO	18% (n= 85)	0.07	Morgana et al., 2018
<hr/>									
Other									
2013	Trawl	Greenland and Barents Seas	60-80	NA	Surface water	Plastics/ m ²	0.063	0.5	Cózar et al., 2017
2013	Trawl	'Rest of Arctic Ocean'	60-80	NA	Surface water	Plastics/ m ²	0	0.5	Cózar et al., 2017
2014	Ice core	Arctic Ocean	78.29	-176.7	Sea ice	Plastics/ m ³	0.050	N/A	Obbard et al. 2014
2014	Ice core	Arctic Ocean	84.31	-149.1	Sea ice	Plastics/ m ³	0.250	N/A	Obbard et al. 2014
2014	Ingestion	Labrador Sea	55	-58	Northern fulmar	% FO	64% (n=39)	1	Avery-Gomm et al., 2018
2015	Ingestion	Labrador Sea	55	-58	Northern fulmar	% FO	97%(n=31)	1	Avery-Gomm et al., 2018
2015	Ingestion	Labrador Sea	55	-58	Northern fulmar	% FO	47% (n=30)	1	Provencher et al. 2018
2016	Sediment	Arctic Central Basin	87.4	-110.5	Benthos	Plastics/ kg	117	0.100	Kanhai et al., 2019
2016	Trawl	Arctic Central Basin	N/A	N/A	Subsurface water	Plastics/ m ³	0.7	0.250	Kanhai et al., 2018

Table S8: Research licensing information provided in plastic pollution research articles from Inuit Nunangat, Greenland and Canadian polar waters

Region	Citation	Study type	Indigenous Licensing stated	Wording of research licensing	Location in paper
Nunavut	Mallory et al., 2006	Ingestion	Unclear	Forty-two northern fulmars were retrieved between 15 August and 10 September 2002, by a fisheries observer stationed aboard a Norwegian longliner fishing for Greenland halibut, <i>Reinhardtius hippoglossoides</i> , operating under licence in Canadian waters (North Atlantic Fisheries Organization Division 0A).	Main text
Nunavut	Mallory 2008	Ingestion	Yes	Collections were in accordance with Canadian Council on Animal Care guidelines, and were conducted under the following permits: research (NUN-SCI-03-02, WL000190, WL000714) animal care (2003PNR017, 2004PNR021, 2005PNR- 021), and land use (59A/7-2-2).	Acknowledgements
Nunavut	Provencher et al. 2009	Ingestion	Yes	Scientific studies and collections were conducted in accordance with guidelines from the Canadian Council on Animal Care, and under appropriate territorial and federal research permits.	Acknowledgements

Nunavut	Provencher et al. 2010	Ingestion	Yes	Scientific studies and collections were conducted in accordance with guidelines from the Canadian Council on Animal Care, and under appropriate territorial and federal research permits.	Acknowledgements
Nunavut	Poon et al., 2017	Ingestion	Unclear	The fieldwork component of this project was supported by Environment and Climate Change Canada, Acadia University (48-0-504807) and the Northern Contaminants Program (58-0-205554) (Indigenous and Northern Affairs Canada).	Acknowledgements
Nunavut	Bourdages et al., 2020	Ingestion	Yes	Tissues from the seals were harvested with permits from the Department of Fisheries and Oceans (DFO) (DFO LFSP S-17/181005-NU), approval from the Nunavut Wildlife Management Board (NWMB), and support from the Arviat, Arviq, Sanikiluaq, and Amaruq Hunter and Trapper Organizations.	Introduction/ Methods

Inuvialuit	Moore et al., 2020	Ingestion	Unclear	<p>For over 30 years, morphometric measurements, observations, and tissues of beluga have been collected throughout the Inuvialuit Settlement Region as part of a community-partnered harvest-based monitoring program... In this study, we worked with the Hendrickson Island Beluga Monitoring Program and the community of Tuktoyaktuk to secure samples from beluga whales with the goal of characterizing presence and degree of potential contamination by microplastics in this cetacean population... The authors thank staff at Fisheries and Oceans Canada (Pacific Region and Central & Arctic Region), the Canada-Inuvialuit Fisheries Joint Management Committee, the Tuktoyaktuk Hunters and Trappers Committee, and the Tuktoyaktuk Community Corporation for their support... We thank the Tuktoyaktuk harvesters for allowing us to sample from their harvests.</p>	Introduction & Acknowledgements
Greenland	Marquard-Petersen, 1998	Excrement analysis	No	No statement given	N/A

Greenland	Amelineau et al., 2016	Ingestion and sub-surface trawl	Yes	All field work in East Greenland was conducted with the permission of the Greenland Home Rule Government, Ministry of Environment and Nature (Danish Polar Center Scientific Expedition Permit 512-240 and 2014-098814), and under permits granted by the Ethics Committee of the French Polar Institute (MP/12/24/05/ 05).	Methods
Greenland	Peeken et al., 2018	Ice core	No	No statement given	N/A
Greenland	Morgana et al., 2018	Ingestion and subsurface trawl	No	No statement given	N/A
Arctic ocean	Obbard et al. 2014	Ice core	No	No statement given	N/A
Various	Provencher et al. 2014	Ingestion	Unclear	[Samples] were collected by hunters as part of larger studies led by Environment Canada, U.S. Fish and Wildlife Service, and/or the Sea Duck Joint Venture in Canada, the Greenland Institute of Natural Resources in Greenland and the Natural Museum in the Faroe Island	Main text
Arctic ocean	Cózar et al. 2017	Surface water	Unclear	We are also grateful to the French Ministry of Foreign Affairs for supporting the expedition and to the	Acknowledgements


				countries that granted sampling permissions.	
Labrador sea	Avery-Gomm et al. 2018	Ingestion	Yes	The collection of fulmars was permitted by the Canadian Wildlife Service, Environment and Climate Change Canada under the Migratory Birds Regulations (Scientific Permit # SC2790)	Acknowledgements
Labrador sea	Provencher et al. 2018	Ingestion	Yes	Collections for this work were done in conjunction with an Environment and Climate Change Canada project funded by the Environmental Studies Research Fund (ESRF). Scientific Permit from Environment and Climate Change Canada: SC2790.	Acknowledgements
Arctic ocean	Kanhai et al. 2018	Subsurface water	No	No statement given	N/A
Arctic ocean	Kanhai et al. 2019	Sediment	No	No statement given	N/A

S9: Summary of interventions into research to foreground good relations with Inuit governance and sovereignty in Inuit Nunangat

1. Researchers should be familiar with research reports, guidelines, priorities and other issues articulated by Inuit Tapiriit Kanatami and other Inuit organizations before research begins in order to align research design and partnerships to calls in those documents.
2. Researchers should be familiar with Indigenous and anticolonial research methodologies, using texts such as Kovach's (2010) *Indigenous Methodologies*, Smith's (2012) *Decolonizing Methodologies*, Walter and Andersen's (2013) *Indigenous Statistics*, Liboiron's (2021) *Pollution is Colonialism*, or Wilson's (2008) *Research is Ceremony*.
3. Researchers should be fluent in the land relations they are engaged with, both in the Arctic in terms of communities, organizations, and permits, but also at home in terms of whose land and relations they are already engaged in.
4. Researchers should obtain permits and permissions from all required bodies, including Inuit governments and communities. Researchers should be aware that obtaining state permits does not guarantee Inuit consultation or consent.
5. Waste that researchers bring into the north, both in terms of solid waste and disposables but also in terms of unintentional waste from gear wear, microfibres releasing, and other forms of microplastic pollution, should be listed as a potential form of harm on permit applications.
6. Permits and permissions should be listed in the methods section.
7. Researchers should endeavour to support Inuit-led research by Inuit researchers rather than consistently launching research in the North from the south.
8. Research teams should include Inuit researchers.
9. Ensure maps and other place descriptors show Inuit and other settlements and land use, rather than depicting the Arctic as an empty space.
10. Inuit researchers, including all personnel who contribute value, samples, or knowledge to a study or study design, should be recognized as co-authors.
11. Categories and measures of plastic should reflect the needs of Inuit users, including local and regional governments. This may include using morphological categories such as "threads" when fishing gear is predominant in a region, translating abundance and other measures into lay terms, conducting brand audits on macroplastics, documenting when plastics are burned or are likely paint chips, and other markers that local experts can use to identify or posit sources.
12. Use Inuktitut place names.
13. Research should take samples of plastics in research gear and on vessels to identify when their work is a source of plastic pollution uncovered in their study, rather than eliminating it as contamination antithetical to findings.
14. Raw data should be accessible to Northern and Inuit governments and use Indigenous data sovereignty and OCAP® principles.
15. Engage in community peer review before submission of papers to academic venues.

S10: Detailed map of plastics in Tasiujarjuaq, Nunavut

Plastics in surface water in Tasiujarjuaq near Iqaluit, Nunavut



S11: Map of plastic abundance from published literature and this study



Plastic abundance in water in the eastern Arctic from published literature, grey literature reports, and this study. Several studies reported a regional average without indicating density or abundance figures for specific trawls or subareas, so several points on the map represent an average density or abundance by general location. Some locations are reported in quotation marks based on location descriptors used in the articles. See Table 3 for more details on each study. Data from surface water measured in plastics/m², including this study, are marked with circles; data from subsurface water measured in plastics/m³ are marked with downward-facing triangles; and sea ice studies measured with plastics/litre, which contain heavily concentrated plastics, are marked with a snowflake. Settlements and cities are indicated with triangles. Overall, these data show that higher abundance of plastics are found at higher latitudes and in sea ice.

1 **S12: References for Supplementary Material**

2
3 Amélineau, F., Bonnet, D., Heitz, O., Mortreux, V., Harding, A.M., Karnovsky, N., Walkusz, W.,
4 Fort, J. and Gremillet, D (2016). Microplastic pollution in the Greenland Sea: Background levels
5 and selective contamination of planktivorous diving seabirds. *Environmental Pollution*, 219:
6 1131-1139.

7
8 Avery-Gomm, S., Provencher, J. F., Liboiron, M., Poon, F. E., & Smith, P. A. (2018). Plastic
9 pollution in the Labrador Sea: an assessment using the seabird northern fulmar *Fulmarus*
10 *glacialis* as a biological monitoring species. *Marine pollution bulletin*, 127, 817-822.

11
12 Bourdages, M.P., Provencher, J.F., Sudlovenick, E., Ferguson, S.H., Young, B.G., Pelletier, N.,
13 Murphy, M.J., D'Addario, A. and Vermaire, J.C. (2020). No plastics detected in seal (*Phocidae*)
14 stomachs harvested in the eastern Canadian Arctic. *Marine Pollution Bulletin*, 150: 110772.

15
16 Cózar, A., Martí, E., Duarte, C.M., García-de-Lomas, J., Van Sebille, E., Ballatore, T.J., Eguíluz,
17 V.M., González-Gordillo, J.I., Pedrotti, M.L., Echevarría, F. and Troublè, R. (2017). The Arctic
18 Ocean as a dead end for floating plastics in the North Atlantic branch of the Thermohaline
19 Circulation. *Science advances*, 3(4): p.e1600582.

20
21 Inuit Tapiriit Kanatami (ITK) (2018). *National Inuit Strategy on Research*. Ottawa, Canada: Inuit
22 Tapiriit Kanatami.

23
24 Inuit Tapiriit Kanatami and Nunavut Research Institute (ITK& NRI) (2007). *Negotiating Research*
25 *Relationships with Inuit Communities: A Guide for Researchers*. Nickels, S., Shirley, J. &
26 Laidler, G. (eds). Ottawa & Iqaluit: Inuit Tapiriit Kanatami and Nunavut Research Institute.

27
28 Kanhai, L.K., Gårdfeldt, K., Lyashevskaya, O., Hassellöv, M., Thompson, R.C., O'Connor, I.
29 (2018). Microplastics in sub-surface waters of the Arctic Central Basin. *Marine Pollution Bulletin*
30 130, 8-18.

31
32 Kanhai, L.K., Johansson, C., Frias, J.P.G.L., Gardfeldt, K., Thompson, R.C., O'Connor, I.
33 (2019). Deep sea sediments of the Arctic Central Basin: A potential sink for microplastics. *Deep*
34 *Sea Research Part 1: Oceanographic Research Papers* 145: 137-142.

35
36 Kovach, M. (2010). *Indigenous Methodologies: Characteristics, Conversations, and Contexts*.
37 Toronto, ON: University of Toronto Press.

38
39 Liboiron, M. (2021). *Pollution is Colonialism*. Durham, NC: Duke University Press.


40
41 Mallory, M. L (2008). Marine plastic debris in northern fulmars from the Canadian high Arctic.
42 *Marine Pollution Bulletin* 56: 1501-1504

43
44 Mallory, M. L., Roberston, G. J., & Moenting, A. (2006). Marine plastic debris in northern fulmars
45 from Davis Strait, Nunavut, Canada. *Marine Pollution Bulletin*, 52(7), 813-815.

46
47 Marquard-Petersen, U. (1998). Food habits of arctic wolves in Greenland. *Journal of*
48 *Mammalogy*, 79(1), 236-244.

49
50 Morgana, Silvia, Laura Ghigliotti, Noelia Estévez-Calvar, Roberto Stifanese, Alina Wieckzorek,
51 Tom Doyle, Jørgen S. Christiansen, Marco Faimali, and Francesca Garaventa. "Microplastics in

52 the Arctic: A case study with sub-surface water and fish samples off Northeast Greenland."
53 Environmental pollution 242 (2018): 1078-1086.
54
55 Moore, R. C., L. Loseto, M. Noel, A. Etemadifar, J. D. Brewster, S. MacPhee, L. Bendell, and P.
56 S. Ross. (2020) Microplastics in beluga whales (*Delphinapterus leucas*) from the Eastern
57 Beaufort Sea. Marine pollution bulletin 150: 110723.
58
59 Obbard, R. W., Sadri, S., Wong, Y. Q., Khitun, A. A., Baker, I., & Thompson, R. C. (2014).
60 Global warming releases microplastic legacy frozen in Arctic Sea ice. *Earth's Future*, 2(6), 315-
61 320.
62
63 Peeken, I., Primpke, S., Beyer, B., Gütermann, J., Katlein, C., Krumpfen, T., Bergmann, M.,
64 Hehemann, L. and Gerdts, G (2018). Arctic sea ice is an important temporal sink and means of
65 transport for microplastic. *Nature communications*, 9(1): 1-12.
66
67 Poon, F. E., Provencher, J. F., Mallory, M. L., Braune, B. M., & Smith, P. A. (2017). Levels of
68 ingested debris vary across species in Canadian Arctic seabirds. *Marine pollution bulletin*,
69 116(1-2), 517-520.
70
71 Provencher, J. F., Gaston, A. J., & Mallory, M. L. (2009). Evidence for increased ingestion of
72 plastics by northern fulmars (*Fulmarus glacialis*) in the Canadian Arctic. *Marine pollution*
73 *bulletin*, 58(7), 1092-1095.
74
75 Provencher, J. F., Gaston, A. J., Mallory, M. L., O'hara, P. D., & Gilchrist, H. G. (2010). Ingested
76 plastic in a diving seabird, the thick-billed murre (*Uria lomvia*), in the eastern Canadian Arctic.
77 *Marine Pollution Bulletin*, 60(9), 1406-1411
78
79 Provencher, J.F., Bond, A.L., Hedd, A., Montevecchi, W.A., Muzaffar, S.B., Courchesne, S.J.,
80 Gilchrist, H.G., Jamieson, S.E., Merkel, F.R., Falk, K. and Durinck, J., (2014). Prevalence of
81 marine debris in marine birds from the North Atlantic. *Marine Pollution Bulletin*, 84(1-2): 411-
82 417.
83
84 Provencher, J. F., Vermaire, J. C., Avery-Gomm, S., Braune, B. M., & Mallory, M. L. (2018).
85 Garbage in guano? Microplastic debris found in faecal precursors of seabirds known to ingest
86 plastics. *Science of the Total Environment*, 644, 1477-1484.
87
88 Wilson, S. (2008). *Research is Ceremony: Indigenous Research Methods*. Winnipeg, MB:
89 Fernwood Books.
90
91 Walter, M., & Andersen, C. (2013). *Indigenous Statistics: A Quantitative Research Methodology*.
92 Walnut Creek, CA: Left Coast Press.
93



S13: Currents near sample sites. In addition to those depicted, the sample site in southwest Greenland is additionally influenced by the three cross sections of current, the Paamiut Section, Cape Desolation Section, and Cape Farewell Section. Currents and sections based on Yang et al. 2015.