

The Riemann Hypothesis

Frank Vega

Abstract. Let's define $\delta(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log x - B)$, where $B \approx 0.2614972128$ is the Meissel-Mertens constant. The Robin theorem states that $\delta(x)$ changes sign infinitely often. Let's also define $S(x) = \theta(x) - x$, where $\theta(x)$ is the Chebyshev function. A theorem due to Erhard Schmidt implies that $S(x)$ changes sign infinitely often. Using the Nicolas theorem, we prove that when the inequalities $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied for $x \geq 127$, then the Riemann Hypothesis should be false. However, the Mertens second theorem states that $\lim_{x \rightarrow \infty} \delta(x) = 0$. In addition, we could modify the well-known result $\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1$ as $\lim_{x \rightarrow \infty} S(x) = 0$. In this way, this work could mean a new step forward in the direction for finally solving the Riemann Hypothesis.

1 Introduction

In mathematics, the Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$ [1]. Let $N_n = 2 \times 3 \times 5 \times 7 \times 11 \times \dots \times p_n$ denotes a primorial number of order n such that p_n is the n^{th} prime number. Say Nicolas(p_n) holds provided

$$\prod_{q|N_n} \frac{q}{q-1} > e^\gamma \times \log \log N_n.$$

The constant $\gamma \approx 0.57721$ is the Euler-Mascheroni constant, \log is the natural logarithm, and $q | N_n$ means the prime q divides to N_n . The importance of this property is:

Theorem 1.1 [6], [7]. *Nicolas(p_n) holds for all prime $p_n > 2$ if and only if the Riemann Hypothesis is true.*

In mathematics, the Chebyshev function $\theta(x)$ is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

where $p \leq x$ means all the prime numbers p that are less than or equal to x . We use the following property of the Chebyshev function:

Theorem 1.2 [9]. *For $x > 1$:*

$$\theta(x) = (1 + \varepsilon(x)) \times x$$

where $\varepsilon(x) < \frac{1}{2 \times \log x}$.

2010 Mathematics Subject Classification: Primary 11M26; Secondary 11A41, 11A25.

Keywords: Riemann Hypothesis, Nicolas theorem, Prime numbers, Chebyshev function.

Besides, in the Grönwall paper appears this:

Theorem 1.3 [3].

$$\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1.$$

Let's define $S(x) = \theta(x) - x$. Nicolas also proves that

Theorem 1.4 [7]. For $x \geq 121$:

$$\log \log \theta(x) \geq \log \log x + \frac{S(x)}{x \times \log x} - \frac{S(x)^2}{x^2 \times \log x}.$$

From the paper of Schmidt, then we can deduce that:

Theorem 1.5 [10]. $S(x)$ changes sign infinitely often.

The famous Mertens paper provides the statement:

Theorem 1.6 [5].

$$\log \left(\prod_{q \leq x} \frac{q}{q-1} \right) = \sum_{q \leq x} \frac{1}{q} + \gamma - B - \frac{1}{2} \times \sum_{q > x} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > x} \frac{1}{q^3} - \dots$$

where $B \approx 0.2614972128$ is the Meissel-Mertens constant.

Let's define:

$$\delta(x) = \left(\sum_{q \leq x} \frac{1}{q} - \log \log x - B \right),$$

Robin theorem states the following result:

Theorem 1.7 [8]. $\delta(x)$ changes sign infinitely often.

In addition, the Mertens second theorem states that:

Theorem 1.8 [5].

$$\lim_{x \rightarrow \infty} \delta(x) = 0.$$

Putting all together yields the proof that when the inequalities $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied for $x \geq 127$, then the Riemann Hypothesis should be false.

2 Central Lemma

Lemma 2.1 For $x \geq 127$:

$$\frac{S(x)}{x} < 1.$$

Proof By the theorem 1.2, for all $x \geq 127$:

$$\begin{aligned}
 \frac{S(x)}{x} &= \frac{\theta(x) - x}{x} \\
 &= \frac{(1 + \varepsilon(x)) \times x - x}{x} \\
 &= \frac{x \times ((1 + \varepsilon(x)) - 1)}{x} \\
 &= (1 + \varepsilon(x) - 1) \\
 &= \varepsilon(x) \\
 &< \frac{1}{2 \times \log x} \\
 &< 1.
 \end{aligned}$$

■

3 Main Theorem

Theorem 3.1 *If the inequalities $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied for $x \geq 127$, then the Riemann Hypothesis should be false.*

Proof For $x \geq 127$, suppose that simultaneously the inequalities Nicolas(p), $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied, where p is the greatest prime such that $p \leq x$. If Nicolas(p) holds, then

$$\prod_{q \leq x} \frac{q}{q-1} > e^\gamma \times \log \theta(x).$$

We apply the logarithm to the both sides of the inequality:

$$\log \left(\prod_{q \leq x} \frac{q}{q-1} \right) > \gamma + \log \log \theta(x).$$

We use that theorem 1.6:

$$\log \left(\prod_{q \leq x} \frac{q}{q-1} \right) = \sum_{q \leq x} \frac{1}{q} + \gamma - B - \frac{1}{2} \times \sum_{q > x} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > x} \frac{1}{q^3} - \dots.$$

Besides, we use that theorem 1.4:

$$\log \log \theta(x) \geq \log \log x + \frac{S(x)}{x \times \log x} - \frac{S(x)^2}{x^2 \times \log x}.$$

Putting all together yields the result:

$$\begin{aligned} & \sum_{q \leq x} \frac{1}{q} + \gamma - B - \frac{1}{2} \times \sum_{q > x} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > x} \frac{1}{q^3} - \dots \\ & > \gamma + \log \log \theta(x) \\ & \geq \gamma + \log \log x + \frac{S(x)}{x \times \log x} - \frac{S(x)^2}{x^2 \times \log x}. \end{aligned}$$

Let distribute it and remove γ from the both sides:

$$\begin{aligned} & \sum_{q \leq x} \frac{1}{q} - \log \log x - B - \frac{1}{2} \times \sum_{q > x} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > x} \frac{1}{q^3} - \dots > \\ & \frac{1}{\log x} \times \left(\frac{S(x)}{x} - \frac{S(x)^2}{x^2} \right). \end{aligned}$$

We know that $\delta(x) = \sum_{q \leq x} \frac{1}{q} - \log \log x - B$. Moreover, we know that

$$\left(\frac{S(x)}{x} - \frac{S(x)^2}{x^2} \right) \geq 0.$$

Certainly, according to the lemma 2.1, we have that $\frac{S(x)}{x} < 1$. Consequently, we obtain that $\frac{S(x)}{x} \geq \frac{S(x)^2}{x^2}$ under the assumption that $S(x) \geq 0$, since for every real number $0 \leq x < 1$, the inequality $x \geq x^2$ is always satisfied. To sum up, we would have that

$$\delta(x) - \frac{1}{2} \times \sum_{q > x} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > x} \frac{1}{q^3} - \dots > 0$$

because of

$$\frac{1}{\log x} \times \left(\frac{S(x)}{x} - \frac{S(x)^2}{x^2} \right) \geq 0.$$

However, the inequality

$$\delta(x) - \frac{1}{2} \times \sum_{q > x} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > x} \frac{1}{q^3} - \dots > 0$$

is never satisfied when $\delta(x) \leq 0$. By contraposition, $\text{Nicolas}(p)$ does not hold when $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied for $x \geq 127$, where p is the greatest prime such that $p \leq x$. In conclusion, if $\text{Nicolas}(p)$ does not hold for a prime $p \geq 127$, then the Riemann Hypothesis should be false due to the theorem 1.1. \blacksquare

4 Discussion

The Riemann Hypothesis has been qualified as the Holy Grail of Mathematics [4]. It is one of the seven Millennium Prize Problems selected by the Clay Mathematics Institute to carry a US 1,000,000 prize for the first correct solution [2]. In the theorem 3.1, we show that if the inequalities $\delta(x) \leq 0$ and

$S(x) \geq 0$ are satisfied for $x \geq 127$, then the Riemann Hypothesis should be false. Nevertheless, the well-known theorem 1.8 states that

$$\lim_{x \rightarrow \infty} \delta(x) = 0.$$

In the same way, we can restate the theorem 1.3 as

$$\lim_{x \rightarrow \infty} S(x) = 0.$$

Indeed, we think this work could help to the scientific community in the global efforts for trying to solve this outstanding and difficult problem.

References

- [1] Peter B. Borwein, Peter Borwein, Stephen Choi, Brendan Rooney, and Andrea Weirathmueller. *The Riemann Hypothesis: A Resource for the Afficionado and Virtuoso Alike*, volume 27. Springer Science & Business Media, 2008.
- [2] Keith J. Devlin. *The Millennium Problems: The Seven Greatest Unsolved Mathematical Puzzles Of Our Time*, volume 100. Basic Books New York, NY (USA), 2002.
- [3] Thomas H. Grönwall. Some asymptotic expressions in the theory of numbers. *Transactions of the American Mathematical Society*, 14(1):113–122, 1913. doi:10.2307/1988773.
- [4] Gilles Lachaud. L'hypothèse de Riemann: le Graal des mathématiciens. *La Recherche Hors-Série*, (20), 2005.
- [5] Franz Mertens. Ein Beitrag zur analytischen Zahlentheorie. *J. reine angew. Math.*, 1874(78):46–62, 1874. URL: <https://doi.org/10.1515/crll.1874.78.46>, doi:10.1515/crll.1874.78.46.
- [6] Jean-Louis Nicolas. Petites valeurs de la fonction d'Euler et hypothèse de Riemann. *Séminaire de Théorie des nombres DPP, Paris*, 82:207–218, 1981.
- [7] Jean-Louis Nicolas. Petites valeurs de la fonction d'Euler. *Journal of number theory*, 17(3):375–388, 1983. doi:10.1016/0022-314X(83)90055-0.
- [8] Guy Robin. Sur l'ordre maximum de la fonction somme des diviseurs. *Séminaire Delange-Pisot-Poitou Paris*, 82:233–242, 1981.
- [9] J. Barkley Rosser and Lowell Schoenfeld. Approximate Formulas for Some Functions of Prime Numbers. *Illinois Journal of Mathematics*, 6(1):64–94, 1962. doi:10.1215/ijm/1255631807.
- [10] Erhard Schmidt. Über die Anzahl der Primzahlen unter gegebener Grenze. *Mathematische Annalen*, 57(2):195–204, 1903.

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France
 Email: vega.frank@gmail.com