




# CLIMATE AND ENVIRONMENTAL CHANGE IN THE MEDITERRANEAN BASIN

Current situation and risks for the future

First Mediterranean Assessment Report

by **MedECC** (Mediterranean Experts on Climate and environmental Change)



# CLIMATE AND ENVIRONMENTAL CHANGE IN THE MEDITERRANEAN BASIN

First Mediterranean Assessment Report (MAR1)

## Edited by

**Wolfgang Cramer**

MedECC Coordinator  
CNRS, France

Mediterranean Institute for terrestrial and  
marine Biodiversity and Ecology (IMBE)

**Joël Guiot**

MedECC Coordinator  
CNRS, France

Centre Européen de Recherche et d'Enseignement  
des Géosciences de l'Environnement (CEREGE)

**Katarzyna Marini**

MedECC Science Officer  
MedECC Secretariat  
Plan Bleu



Mediterranean  
Action Plan  
Barcelona  
Convention



Union for the Mediterranean  
Union pour la Méditerranée  
الإتحاد من أجل المتوسط



Sweden  
Sverige

## MEDECC

The Mediterranean Experts on Climate and environmental Change (MedECC) are an independent network of scientists, founded 2015. MedECC assesses the best available scientific knowledge on climate and environmental change and associated risks in the Mediterranean Basin in order to render it accessible to policymakers, stakeholders and citizens.

To date (September 2020), MedECC counts more than 600 scientific members, all contributing in individual capacity and without financial compensation. MedECC scientists are based in 35 countries, including 19 countries registered as Contracting Parties to the Convention for the Protection of the Marine Environment and the Coastal Region of the Mediterranean (Barcelona Convention) and 23 countries that are members of the Union for the Mediterranean.

The UNEP/MAP – Barcelona Convention Secretariat, through its Plan Bleu Regional Activity Center, and the Secretariat of the Union for the Mediterranean are working in partnership to support MedECC, and to contribute to establishing a sound and transparent scientific assessment process.

MedECC reports are prepared for use of policymakers and a broader audience. They are developed on the basis of scientific criteria only; their validity is therefore the responsibility of MedECC Report Authors alone. The available knowledge concerning the risks studied by MedECC has significant certain gaps, often due to limited monitoring systems or scientific research capacity – these have been indicated as clearly as possible. Despite best efforts, errors and omissions are nevertheless not unlikely.

## THIS REPORT

MedECC has prepared the First Mediterranean Assessment Report (MAR1) on the current state and expected risks of climate and environmental change in the Mediterranean Basin. The report includes a Summary for Policymakers (SPM), which comprises the key messages of the MAR1. A first draft of the MAR1 was prepared in 2019 and underwent expert peer review.

The second draft, revised to take into account review comments, and now supplied with a SPM, has undergone broad consultation with governments, decision-makers and stakeholders in 2020. The particular aim of the consultation was to ascertain that MAR1 findings, as presented in the SPM, are fully comprehensible and unambiguous. While chapter drafts were also supplied with the SPM under review, they served as background information only and were not part of the review.

The MedECC coordinators are very grateful for the expertise, rigor and dedication shown by the volunteer Coordinating Lead Authors and Lead Authors, working across scientific disciplines in each chapter of the report, with essential help from many Contributing Authors.

At the end of the stakeholder consultation, 453 comments for the SPM had been received and were used to revise the SPM wherever this was possible. The stakeholder review was concluded during the online Plenary Session of stakeholders held on September 22, 2020.

MedECC Authors and Coordinators want to thank all reviewers for their time and effort. A record is being kept on the responses to all review comments.

This publication has been made possible through the collaboration of 190 contributors, who are listed in the full report.

**Editors:** Wolfgang Cramer, Joël Guiot, Katarzyna Marini.

**Editorial Committee:** Semia Cherif (Tunisia), Wolfgang Cramer (France), Carlo Giupponi (Italy), Joël Guiot (France), Manfred Lange (Cyprus/Germany), Piero Lionello (Italy), Katarzyna Marini (France), Maria Snoussi (Morocco), Andrea Toreti (Italy), Elena Xoplaki (Greece/Germany).

Reproduction is authorised provided the source is acknowledged. An online version of this work is published at <https://www.medecc.org/first-mediterranean-assessment-report-mar1/> which permits re-use, distribution and reproduction in any medium for non-commercial purposes providing appropriate credit to the original work is given. All versions of this work may contain content reproduced under license from third parties. Permission to reproduce this third-party content must be obtained from these third-parties directly.

**Preferred citation:**

MedECC (2020) Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer, W., Guiot, J., Marini, K. (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, 632pp.  
ISBN: 978-2-9577416-0-1

**Cover design:** Pandaroo (Péronnas)

**Graphs (re)designing and layout:** Zen design studio (Marseille)

**Copy-editing** by Connected Language Services

The UNEP/MAP – Barcelona Convention Secretariat, through its Plan Bleu Regional Activity Center, and the Secretariat of the Union for the Mediterranean are working in partnership to support MedECC. The MedECC Secretariat is supported and funded by UfM, through a grant provided by the Swedish International Development Cooperation Agency (SIDA), hosted by Plan Bleu in Marseille, France.

**Supporting institutions**



Union for the Mediterranean  
Union pour la Méditerranée  
الإتحاد من أجل المتوسط



Mediterranean  
Action Plan  
Barcelona  
Convention



**With financial support from**



Sweden  
Sverige



RÉPUBLIQUE  
FRANÇAISE  
Liberté  
Égalité  
Fraternité



ADEME  
AGENCE DE LA  
TRANSITION  
ÉCOLOGIQUE



Gouvernement Princier  
PRINCIPAUTÉ DE MONACO



Generalitat de Catalunya  
Consell Assessor  
per al Desenvolupament Sostenible



Aix  
Marseille  
Provence



PAYS D'AIX



acterra  
ENVIRONNEMENT  
CLIMATE CHANGE



Institut de Recherche  
pour le Développement  
FRANCE  
French National Research Institute - Sustainable Development



MINISTÈRE  
DE LA TRANSITION  
ÉCOLOGIQUE  
Liberté  
Égalité  
Fraternité

The content and views expressed in this document are purely those of the authors and may not, in any circumstances, be interpreted as stating an official position of the supporting institutions. Neither the supporting institutions nor any person acting on their behalf may be held responsible for the use, which may be made of the information contained therein.

The supporting institutions does not guarantee the accuracy of the information included in this document, nor does it accept any responsibility for any use thereof. Reference herein to any specific products, specifications, processes or services by trade name, trademark, manufacturer or otherwise does not necessarily constitute or imply its endorsement, recommendation or favoring by the supporting institutions.



# Table of contents

<b>Front Matter</b> .....	<b>III</b>
Foreword .....	<b>VIII</b>
Acknowledgments .....	<b>X</b>
<b>Summary for Policymakers</b> .....	<b>11</b>
<b>Chapter 1 - Introduction</b> .....	<b>41</b>
<b>Chapter 2 - Drivers of change</b> .....	<b>59</b>
2.1 Introduction .....	<b>66</b>
2.2 Climate change .....	<b>67</b>
2.3 Pollution .....	<b>94</b>
2.4 Land and sea use change .....	<b>109</b>
2.5 Non-indigenous species .....	<b>116</b>
2.6 Interactions among drivers .....	<b>126</b>
2.7 Mediterranean socioeconomic scenarios .....	<b>134</b>
<b>Chapter 3 - Resources</b> .....	<b>181</b>
3.1 Water .....	<b>184</b>
3.2 Food .....	<b>237</b>
3.3 Energy transition in the Mediterranean .....	<b>265</b>
<b>Chapter 4 - Ecosystems</b> .....	<b>323</b>
4.1 Marine ecosystems .....	<b>329</b>
4.2 Coastal ecosystems .....	<b>352</b>
4.3 Terrestrial and freshwater ecosystems .....	<b>375</b>
<b>Chapter 5 - Society</b> .....	<b>469</b>
5.1 Development .....	<b>472</b>
5.2 Health .....	<b>493</b>
5.3 Human security .....	<b>515</b>
<b>Chapter 6 - Managing future risks and building socio-ecological resilience</b> .....	<b>539</b>
6.1 Introduction .....	<b>543</b>
6.2 Human health impacts .....	<b>544</b>
6.3 Water security .....	<b>547</b>
6.4 Agricultural drought .....	<b>550</b>
6.5 Wildfires .....	<b>553</b>
6.6 Soil erosion, degradation, and desertification .....	<b>555</b>
6.7 Heat waves .....	<b>558</b>
6.8 River and pluvial flooding .....	<b>559</b>
6.9 Sea-level rise: coastal erosion and flooding, saltwater intrusion .....	<b>560</b>
6.10 Seawater temperature anomalies and extremes .....	<b>563</b>
6.11 Ocean acidification .....	<b>564</b>
6.12 Non-indigenous species: marine, freshwater, and terrestrial .....	<b>566</b>
6.13 Interactions of hazards, synergies and trade-offs between adaptation strategies and mitigation .....	<b>568</b>
<b>Appendix A - Appendix to Chapter 1 - Introduction</b> .....	<b>589</b>
<b>Appendix B - Maps of seasonal temperature and precipitation changes for the Mediterranean Basin</b> .....	<b>599</b>
<b>Appendix C - List of acronyms, chemical symbols and scientific units</b> .....	<b>613</b>
<b>Appendix D - ISO2 country codes</b> .....	<b>621</b>
<b>Appendix E - Lists of Figures, Tables and Boxes</b> .....	<b>623</b>



## FOREWORD

Climate change is possibly the most crucial challenge for the future of the Mediterranean: the pace and magnitude of climate change and its impacts in the region can turn it into a serious stability risk if untamed and not integrated organically in national and regional policy.

In accordance with their respective mandates, the Secretariats of the Union for the Mediterranean (UfM) and of the United Nations Environment Programme / Mediterranean Action Plan – Barcelona Convention (UNEP/MAP) have joined efforts to support the work undertaken by the network of Mediterranean Experts on Climate and environmental Change (MedECC) and its Secretariat, which is hosted by Plan Regional Activity Centre (UNEP/MAP) in Marseille, France.

Despite the pressing need to better understand the current state of play and the present and future risks of climate and environmental change in the Mediterranean, there was until recently a dearth of reliable, robust and scientific data combined with a deficit in knowledge sharing with decision-makers and stakeholders.

Thanks to MedECC, the First Mediterranean Assessment Report (MAR1) now bridges the gap.

The methodology used by MedECC is based on key principles of the Intergovernmental Panel on Climate Change (IPCC) and of the Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES). This approach is innovative, but its application was fraught with challenges specific to the Mediterranean region. We commend MedECC for the results achieved.

For the UfM Secretariat, it is clear that MAR1 is a major deliverable of the 1st UfM Ministerial Declaration on Environment and Climate Change; it is also a cornerstone between the first and the second UfM Ministerial Declarations. From the UNEP/MAP perspective, the support to MedECC is a major achievement under the Mediterranean Strategy for Sustainable Development (MSSD), which identifies the establishment of a regional science-policy interface on climate change as a priority (Flagship Initiative). MAR1 will take centre stage in the context of the 22nd Meeting of the Contracting Parties to the Barcelona Convention, which will adopt the UNEP/MAP Medium-Term Strategy (MTS) 2022-2027.


At a time when an unprecedented health crisis having severe impacts on Mediterranean societies and economies, regional cooperation and multilateralism underpinned by robust science have never been more sorely needed. In this vein, MedECC offers a unique synthesis on best available scientific knowledge on climate and environmental changes in the Mediterranean, thus allowing for an evidence-based, basin-wide policy response.

This is also a report of great importance to our partners in the Mediterranean region as MAR1 will inform integrated programming in ways that can improve cooperation to address urgent needs and to inform long-term climate and environmental action. Its findings constitute a foundation for meeting sustainable development needs whilst informing strategic planning and investment at the regional level.

We acknowledge with appreciation MedECC's efforts, which also have an important impact in terms of capacity-building. UfM and UNEP/MAP reiterate their commitment to bolstering MedECC and its linkages with global processes. We will deploy additional support for the best-possible use of MAR1 in the context of the enforcement of the three UN Rio conventions in the region.



**Mr. Nasser KAMEL**  
*Secretary General*  
*Union for the Mediterranean*



**Mr. Gaetano LEONE**  
*Coordinator*  
*UNEP/MAP – Barcelona Convention*



Within the mandate given by the Contracting Parties to the Barcelona Convention to Plan Bleu in its role of Regional Activity Centre of the UNEP/MAP system, and in the framework of an agreement with the Secretariat of the Union for the Mediterranean, Plan Bleu has hosted the MedECC Secretariat and supported directly its activities since its creation.

We would like to commend all the scientists who contributed to this report, MedECC Steering Committee and especially its Coordinators and Scientific Secretary for the delivery of this outstanding report. This work would not have been possible without the financial support of the Swedish International Development Cooperation Agency (SIDA), the Principality of Monaco, the French Agency for Ecological Transition (ADEME), the French Ministry of Ecological Transition and Solidarity, the French National Centre for Scientific Research (CNRS) and the French Research Institute for Development (IRD) via the MISTRALS programme, the Aix-Marseille University via the Laboratory of Excellence (Labex) OT-Med, the Advisory Council for the Sustainable Development of Catalonia of the Government of Catalonia (CADS) and the Aix-Marseille Provence Metropole.

Plan Bleu will maintain its support to the MedECC network to disseminate the key MAR1 findings to all stakeholders, including to policymakers through UfM and UNEP/MAP processes; and to dive in the assessment of thematic issues in its next phase of activities.



**Mr. Thierry LAVOUX**  
*Président, Plan Bleu*

## ACKNOWLEDGEMENTS

The MedECC report is the outcome of the work of many people, supported by many institutions. While we cannot mention all of them here, the editors and authors wish to acknowledge and thank the following individuals and institutions who provided their help at different stages of the process: Plan Bleu director, its current François Guerquin, as well as his predecessors Elen Lemaître-Curri and Anne France-Didier, also the Plan Bleu team, in particular Antoine Lafitte, Lina Tode, Sandra Dulbecco and Anna Goubert; the Union for the Mediterranean, its Climate Adviser Arnault Graves, its Senior Deputy Secretary General Grammenos Mastrojeni, its previous Climate Adviser Nicolas Debaisieux, the team of the Union for the Mediterranean Energy and Climate Action, as well as the UfM Communication and Public Affairs Unit; the UN Environment Programme/Mediterranean Action Plan Coordinating Unit, its Coordinator Gaetano Leone and its Programme Management Officer Julien Le Tellier.

We also wish to thank wholeheartedly the reviewers of various drafts of the report – all their comments have been considered carefully even if it was not always possible to accommodate them. The many participants of the Conference “Climate Our Common Future” in Paris 2015 and numerous following scoping meetings made very important contributions to the report.

We received important support from the hosting institutions of several workshops: Mohammed V University in Rabat (Morocco), University of Palermo (Italy), Venice International University (VIU) (Italy), Scientific Centre of Monaco (CSM), Oceanographic Museum of Monaco, National Research Council CNR Congress Center in Milan (Italy) and Aix-Marseille University (France). The Summary for Policymakers (SPM) was crucially refined during a day-long discussion with focal point representatives of the main stakeholders, chaired by Valentina Mauriello and Grammenos Mastrojeni. The report preparation was also supported by the team of the Labex OT-Med (including administrative staff: Sophie Pékar, Barbara Bourlion), the team of AIR Climat (Aurore Aubail and Antoine Nicault), Acterra (Stéphane Simonet), Service Informatique Pythéas (OSU Institut Pythéas, France), Stéphanie Wicha, Aurore Pfitzmann and Marie-Aimée Gros-Rosanvallon.





# SUMMARY FOR POLICYMAKERS


## Text as approved during Plenary Session of MedECC Stakeholders on September 22, 2020

It will be presented to government representatives for endorsement without further changes, other than editorial and technical corrections.

### Drafting Authors:

Wolfgang Cramer (France), Joël Guiot (France), Katarzyna Marini (France), Brian Azzopardi (Malta), Mario V Balzan (Malta), Semia Cherif (Tunisia), Enrique Doblas-Miranda (Spain), Maria dos Santos (Portugal), Philippe Drobinski (France), Marianela Fader (Germany), Abed El Rahman Hassoun (Lebanon), Carlo Giupponi (Italy), Vassiliki Koubi (Greece/Switzerland), Manfred Lange (Cyprus), Piero Lionello (Italy), Maria Carmen Llasat (Spain), Stefano Moncada (Malta), Rachid Mrabet (Morocco), Shlomit Paz (Israel), Robert Savé (Spain), Maria Snoussi (Morocco), Andrea Toreti (Italy), Athanasios T. Vafeidis (Germany/Greece), Elena Xoplaki (Germany)

*This document should be cited as: MedECC 2020 Summary for Policymakers. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/ MAP, Marseille, France, pp 11-40.*



# Table of contents

Summary for Policymakers.....	11
Executive Summary:	
Climate and environmental change in the Mediterranean Basin .....	14
Background and key findings of the First Mediterranean Assessment Report .....	16
<b>1. Background for the assessment .....</b>	<b>16</b>
<b>2. Drivers of environmental change in the Mediterranean Basin .....</b>	<b>17</b>
2.1 Climate change.....	17
2.2 Pollution .....	20
2.3 Land and sea use change.....	22
2.4 Non-indigenous species.....	22
<b>3. Resources .....</b>	<b>23</b>
3.1 Water.....	23
3.2 Food.....	26
3.3 Energy transition in the Mediterranean.....	27
<b>4. Ecosystems.....</b>	<b>29</b>
4.1 Marine ecosystems .....	29
4.2 Coastal ecosystems.....	31
4.3 Terrestrial ecosystems .....	32
<b>5. Society.....</b>	<b>34</b>
5.1 Development.....	34
5.2 Human health.....	36
5.3 Human security .....	38
<b>6. Managing future risks and building socio-ecological resilience in the Mediterranean.....</b>	<b>39</b>



## Executive Summary: Climate and environmental change in the Mediterranean Basin

**Virtually all sub-regions of the Mediterranean Basin, on land and in the sea, are impacted by recent anthropogenic changes in the environment. The main drivers of change include climate (temperature, precipitation, atmospheric circulation, extreme events, sea-level rise, sea water temperature, salinity and acidification), population increase, pollution, unsustainable land and sea use practices and non-indigenous species. In most areas, both natural ecosystems and human livelihoods are affected. Due to global and regional trends in the drivers, impacts will be exacerbated in the coming decades, especially if global warming exceeds 1.5 to 2°C above the pre-industrial level. Significantly enhanced efforts are needed in order to adapt to inevitable changes, mitigate change drivers and increase resilience.**

Due to anthropogenic emissions of greenhouse gases, climate is changing in the Mediterranean Basin, historically and projected by climate models, faster than global trends. Annual mean temperatures on land and sea across the Mediterranean Basin are 1.5°C higher than during pre-industrial times and they are projected to rise until 2100 by an additional 3.8 to 6.5°C for a high greenhouse gas concentration scenario (RCP8.5) and 0.5 to 2.0°C for a scenario compatible with the long-term goal of the UNFCCC Paris Agreement to keep the global temperature well below +2°C above the pre-industrial level (RCP2.6). On land and in the sea, heat waves will intensify in duration and peak temperatures. Despite strong regional variations, summer rainfall will likely be reduced by 10 to 30% in some regions, increasing existing water shortages, desertification and decreasing agricultural productivity.

It is virtually certain that sea surface warming will continue during the 21st century by 1 to 4°C depending on the scenario (low or high greenhouse gas emissions) and likely that deep waters will warm more in the Mediterranean than in other oceans in the world. Rising carbon dioxide (CO<sub>2</sub>) concentrations lead to seawater acidification, and this trend will continue. The Mediterranean mean sea level has risen by 6 cm over the past 20 years. This trend is likely to accelerate (with regional differences) by the global rate of 43 to 84 cm until 2100, but possibly more than 1 m in the case of further ice-sheet destabilization in Antarctica.

Most impacts of climate change are exacerbated by other environmental challenges such as changing land use, increasing urbanization and tourism, agricultural intensification, overfishing, land degradation, desertification, and pollution (air, land, rivers and ocean). Sulphur dioxide (SO<sub>2</sub>) and nitrogen oxide (NO<sub>x</sub>) have recently increased drastically, mainly because of shipping activity. Tropospheric ozone (O<sub>3</sub>) concentrations increase due to pollution and warming, and high-level episodes will be more frequent in the future. Saharan dust transport is likely to also increase. The Mediterranean Sea is heavily polluted by multiple substances including plastic, emerging contaminants, heavy metals, fecal bacteria and viruses, all with expected increase in the future.

The Mediterranean Sea is invaded by many non-indigenous species, particularly from the Red Sea but also through the Strait of Gibraltar, maritime transport and aquaculture. On land, non-indigenous species are particularly present in regions with high infrastructure and commerce development, including accidentally introduced phytophagous pests which cause damage to crops and forests. These trends are expected to continue in the future.

Agriculture is the largest user of water in the Mediterranean region. Climate change impacts water resources in combination with demographic and socio-economic drivers, reducing runoff and groundwater recharge, water quality, increasing conflicts among users, ecosystem degradation and groundwater salinization in coastal aquifers. Demand for irrigation is expected to increase by 4 to 18% by 2100. Demographic change, including the growth of large urban centers, could enhance this demand by 22 to 74%. There is adaptive potential in the improvement of water use efficiency and reuse. Other important adaptations are changing agriculture practices and promoting the traditional Mediterranean diet, local production and reduction of food waste.

Land and seafood production activities are strongly impacted by climate change, more frequent and intense extreme events, together with higher soil salinization, ocean acidification and land degradation. Crop yield reductions are projected for the next decades in most current areas of production and for most crops. This will potentially be worsened by emerging pests and pathogens. There is large

adaptation potential in changing farming practices and management to agroecological methods, also providing important potential for climate change mitigation by increased carbon storage in soils. Marine food production is threatened by unsustainable fishing practices, non-indigenous species, warming, acidification and water pollution, which together may affect species distribution and trigger local extinction of more than 20% of exploited fish and marine invertebrates by 2050. Adaptation will require more rigorous management of fisheries in the Mediterranean. The sustainability of the Mediterranean food sector (from the land and the ocean) also depends on population growth, regional consumer behavior (diet) and the global food markets (which may be affected by environmental crisis elsewhere).

Marine ecosystems and their biodiversity are also impacted by overfishing, warming, acidification and the spread of non-indigenous species from tropical waters. Expected consequences include increased jellyfish outbreaks, mucilage and algal bloom outbreaks, reduced commercial fish stocks, and general biodiversity loss due to altered physiology and ecology of most marine organisms. There is potential for mitigating these impacts through improved conservation within and beyond marine protected areas, more sustainable fishing practices and by reducing pollution from agriculture, urban areas and industry. In coastal systems, sea level rise will impact most infrastructure, aquifers, coastal crops, world heritage and other protected sites, notably in river deltas and estuaries. Increasing nutrient flows towards the sea increase the number and frequency of plankton blooms and jellyfish outbreaks, with negative impacts on fisheries, aquaculture and human health. The multiple levels of land-sea interactions could benefit from the implementation of new approaches of ecosystem-based Integrated Coastal Zone Management and conservation planning.

Land biodiversity changes in multiple ways. In countries of the northern rim, forest area is increasing at the expense of extensive agriculture and grazing, while ecosystems in southern countries are still at risk of fragmentation or disappearance due to clearing and cultivation, overexploitation of firewood and overgrazing. Over the past 40 years, biodiversity changes and species loss have led to homogenization and a general simplification of biotic interactions. Half of wetland area has been lost or degraded, and this trend is expected to continue. Dryland extension and an increase in areas burnt during more frequent wildfires are expected. Adaptation options for land biodiversity

include preservation of natural flow variability in Mediterranean rivers and the protection of riparian zones, reduction of water abstraction, modified silvicultural practices, and the promotion of climate-wise landscape connectivity.

Human health is already impacted by high temperatures as well as air and water pollution in the Mediterranean Basin. The combined impacts of expected environmental changes (notably air pollution and climate) increase risks to human health from heat waves, food and water shortages, vector-borne, respiratory and cardio-vascular diseases. These health risks particularly impact disadvantaged or vulnerable populations, including the elderly, children, pregnant women and people with low income. Human security faces new risks from extreme events, particularly along coastal areas. Conflicts caused by scarce resources and human migration are likely to increase due to drought and degrading agricultural and fisheries resources, although socio-economic and political factors are likely to still play a major role.

Mediterranean cities are growing due to increasing population and socio-economic change, notably on the coasts of southern countries. Due to increasing heat stress, the planning and management of cities around the Mediterranean will need to focus more on human health and resilience to environmental change. Impacts of climate change on urban areas are expected to be disproportionately high due to a concentration of population and assets – especially in high-risk prone areas – in combination with hazard-amplifying conditions (e.g., increased runoff resulting from soil sealing, or urban heat island effects). Tourism will likely be affected by climate change through reduced thermal comfort, degradation of natural resources, including freshwater availability, and coastal erosion due to sea level rise and urban development. The net economic effect on tourism will depend on the country and the season.

All Mediterranean countries have significant potential to mitigate climate change through an accelerated energy transition. This will involve phasing down fossil fuel and accelerated development of renewable energies. This ambitious energy transition, reaching beyond the plans and targets announced by governments and policymakers in line with contributions made for the UNFCCC Paris Agreement, requires a significant transformation of energy policies and economic models in Mediterranean countries. While northern rim countries advance towards this transition by gradually diversifying their energy mix, improving energy efficiency

cy and increasing the share of renewable energies, despite investments, some eastern and southern rim countries need support, funding, technology transfer and capacity-building in the framework of the UNFCCC Paris Agreement. Around 2040, the share of renewable energies could triple to reach 13 to 27% under current transition scenarios. Enhanced regional energy market integration and cooperation are crucial to unleashing cost-effective climate change mitigation.

More effective policy responses to climate and environmental changes will require both strengthened mitigation of the drivers of environmental change, such as greenhouse gas emissions, as well as enhanced adaptation to impacts. Poverty, inequalities and gender imbalances presently

hamper the achievement of sustainable development and climate resilience in Mediterranean countries. Culture is a key factor to the success of adaptation policies in the highly diverse multicultural setting of the Mediterranean Basin. Aimed at supporting local and vulnerable communities, policies for climate adaptation and environmental resilience need take into account concerns such as justice, equity, poverty alleviation, social inclusion, and redistribution. To support policies for sustainable development with scientific evidence about climate and environmental change, a synthesis of current scientific knowledge, covering most relevant disciplines, sectors and sub-regions is presented by the First Mediterranean Assessment Report (MAR1).

## BACKGROUND AND KEY FINDINGS OF THE FIRST MEDITERRANEAN ASSESSMENT REPORT

### 1 - Background for the assessment

**1.1** Global environmental change exacerbates existing challenges for the population living around the Mediterranean Sea, through climate change, land use changes, increasing urbanization and tourism, agricultural intensification, pollution, declining biodiversity, resource competition, and socio-economic trends. Environmental, socio-economic and cultural conditions are highly heterogeneous across the Mediterranean Region (*Section 1.1.1*), resulting in different manifestations of regional environmental change that require specific adaptation measures as well as enhanced capacity-building. To account for these specificities, a comprehensive risk assessment approach encompassing the entire Mediterranean Basin is needed to provide adequate and timely information as well as data needed for decision makers to design effective mitigation and adaptation strategies. (*Section 1.1.1*).

**1.2** Despite major research efforts across many disciplines and regions, to date, there has been no comprehensive assessment of risks posed by climate and environmental changes in the Mediterranean Basin. Most countries of the Middle East and North Africa (MENA) are likely to face potentially greater risks from climate and environmental changes than other parts of the Mediterranean Basin, but they have limited capacity to monitor important environmental parameters

or carry out adequate risk analyses. Effective mitigation and adaptation require integrative studies that go beyond the current knowledge. The main challenges for the Mediterranean are to fill data and knowledge gaps across countries, and to foster the development of high-level climate services, including early warning systems. More research is needed for short- and medium-term projections, as well as large scale programs at the Mediterranean scale to address pressing challenges. (*Section 1.1.2*).

**1.3** The 1st Mediterranean Assessment Report (MAR1) has been developed and drafted in order to provide science-based guidance to multiple actors involved in coming up with a response to climate and environmental changes and to reduce associated risks to communities and natural ecosystems in the Mediterranean region (*Section 1.3.1.4*). The report was developed by the scientific community, based on publications in scientific journals, for policymakers and other stakeholders through the conclusions in its Summary for Policymakers (SPM), as well as for a broader audience of experts through its detailed technical chapters supporting the SPM. The report is also intended to be communicated more broadly to the public through additional efforts of communication and participatory actions. (*Section 1.3.2*).



**1.4** The report assesses risks for the entire Mediterranean Basin (land and sea), associated with four main drivers of environmental change: climate, pollution, land and sea use and non-indigenous species. Throughout the report,

scientific confidence in its findings is indicated based on the consistency of evidence and the degree of agreement of the scientific community, using the terms “high”, “medium” and “low”. (Section 1.3.3).

## 2 - Drivers of environmental change in the Mediterranean Basin


### 2.1 Climate change

Anthropogenic climate change has been observed for many variables in the Mediterranean Basin during recent decades. For the future, the region is expected to remain among the regions most affected by climate change, particularly when it comes to precipitation and the hydrological cycle.


**2.1.1** There is robust evidence that the Mediterranean region has significantly warmed. Basin-wide, annual mean temperatures are now 1.54°C above the 1860-1890 level for land and sea areas, i.e. 0.4°C more than the global average change (*high confidence*). (Fig. SPM.1) (Section 2.2.4.1; Box 2.1).

**2.1.2** Multi-model sets of climate simulations show that widespread warming will continue in the Mediterranean during the 21st century (*high confidence*). (Section 2.2.4.2, Table 2.1).

**2.1.2.1** Over land, warming will likely be in the range of 0.9 to 1.5°C or 3.7 to 5.6°C during the 21st century, for low (RCP2.6) or high greenhouse gas emissions (RCP8.5), respectively (*high confidence*). Future regional average warming will exceed the global mean value by 20% on an annual basis and 50% in summer (*high confidence*). (Fig. SPM.2) (Section 2.2.4.2).



**Figure SPM.1 | Observed changes in temperature and rainfall.** Recent trends in temperature [A and B, °C decade<sup>-1</sup>] and rainfall [C and D, mm day<sup>-1</sup> decade<sup>-1</sup>] in the Mediterranean Basin over land. Panels A & C average for the period 1950-2018, panels B & D for 1980-2018 (Fig. 2.5 and 2.8).



**Figure SPM.2 | Projected warming in the Mediterranean Basin over land.** Projected changes in annual temperature relative to the recent past reference period (1980-1999), based on the EURO-CORDEX 0.11° ensemble mean, A: simulations for pathways RCP2.6 and RCP8.5, B: warming at the end of the 21st century (2080-2099) for RCP2.6, C: idem for RCP8.5.

**2.1.2.2** In the future, warm temperature extremes will increase and heat waves will intensify in duration and peak temperatures. For 2°C of global warming above the pre-industrial value, maximum daytime temperatures in the Mediterranean will likely increase by 3.3°C. With 4°C global warming, nearly all nights will be tropical (nighttime temperature for at least five days above a location-dependent threshold) and there will be almost no cold days (below a location-dependent threshold) (*high confidence*). (Section 2.2.4.2).

**2.1.3** The sign and magnitude of observed land precipitation trends show pronounced spatial variability, depending on the time period and season considered (*medium confidence*) (Section 2.2.5.1), so that the confidence in the detection of anthropogenic trends in rainfall for the historical past is low.

**2.1.3.1** The most evident observed trend is a decrease in winter precipitation over the central and southern portions of the basin since the second half of the 20th century (*medium confidence*). (Section 2.2.5.1).

**2.1.4** Models project a consistent decrease in precipitation during the 21st century, for the entire Mediterranean Basin during the warm season (April through September, with the highest magnitude in summer) and in winter for most of Mediterranean, except for the northernmost regions (e.g., the Alps),


where wetter conditions are projected (*medium confidence*). (Fig. SPM.3) (Section 2.2.5.2).

**2.1.4.1** The mean rate of land precipitation decrease among models is 4% per each degree of global warming, which would determine a reduction in the range of 4 to 22% depending on scenario at the end of the 21st century (*medium confidence*) (Section 2.2.5.2). The magnitude of this decrease varies across models, rendering sub-regional projections uncertain.

**2.1.4.2** Future climate projections indicate a predominant shift towards a precipitation regime of higher interannual variability, higher intensity and greater extremes (especially in winter, spring and fall, but not in the southern areas, *low confidence*), decreased precipitation frequency and longer dry spells (especially in summer and in the southern countries) (*medium confidence*). (Section 2.2.5.2).

**2.1.5** There are no significant trends in the number of observed cyclones in recent decades (*low/medium confidence*) (Section 2.2.2.3). Most future climate projections indicate a decrease in cyclones, especially in winter (*medium confidence*). (Section 2.2.2.3).

**2.1.5.1** There is insufficient information for assessing past trends of “medicane” (Mediterranean hurricanes), but projections indicate decreasing frequency and increasing intensity (*medium confidence*). (Section 2.2.2.3).



**Figure SPM.3 | Projected rainfall change in the Mediterranean Basin.** Projected changes in annual rainfall relative to the recent past reference period (1980-1999), based on the EURO-CORDEX 0.11° ensemble mean, A: simulations for pathways RCP2.6 and RCP8.5, B: rainfall anomalies at the end of the 21st century (2080-2099) for RCP2.6, C: idem for RCP8.5.

**2.1.5.2** Projections of future wind speeds converge on a limited wind speed reduction over most of the Mediterranean Sea, with the exception of an increase over the Aegean Sea and northeastern land areas (*medium confidence*). (Section 2.2.2.4).

**2.1.5.3** Projections suggest a general decrease in mean significant wave height, as well as in the number and intensity of wave extremes, over a large part of the Mediterranean Sea, especially in winter, and storm surges along the coasts (*medium confidence*), but with no consensus on the most extreme events. (Section 2.2.8.2).

**2.1.6** Surface solar radiation in the Mediterranean Basin decreased from the 1950s to the 1980s (between  $-3.5$  and  $-5.2$   $W\ m^{-2}$  decade $^{-1}$ ) and recovered thereafter (between  $+0.9$  and  $+4.6$   $W\ m^{-2}$  decade $^{-1}$ ), consistent with global trends (*very high confidence*). (Section 2.2.3.1). In future climate projections, anthropogenic aerosol loads over the Mediterranean are expected to continue to decrease (*high confidence*), leading to an increase in surface solar radiation (*medium confidence*). (Section 2.2.3.2).

**2.1.7** Observations and most model projections indicate a trend towards drier conditions over the Mediterranean Basin, especially in the warm season and over the southern areas (*medium/high confidence*). (Section 2.2.5.3).

**2.1.7.1** Across the Mediterranean Sea, net fresh water loss (evaporation minus precipitation and river runoff) has increased since the last decades of the 20th century (*medium confidence*) (Section 2.2.5.3). The main cause is the strong evaporation increase due to local warming (the estimated rate of evaporation change in relation to warming is about  $0.7$  mm day $^{-1}$  °C $^{-1}$  (or 25% °C $^{-1}$ ) over the period of 1958-2006).

**2.1.7.2** Net water loss from the sea is expected to increase in the future due to a decrease in precipitation and river runoff and an increase in evaporation (*high confidence*). (Section 2.2.5.3).

**2.1.8** In the 20th century a significant reduction in the area and volume of glaciers across high mountains of the Mediterranean has occurred. Deglaciation has generally accelerated in recent decades (*high confidence*). (Section 2.2.6.1).

**2.1.8.1** Warming has shifted the occurrence of periglacial processes to higher elevations and degraded permafrost in high mountain environments. Glaciers in the Mediterranean region are projected to continue losing mass in the 21st century until complete disappearance of most mountain glaciers by the end of the century (*very high confidence*). (Section 2.2.6.2).

**2.1.8.2** At lower elevation, the snow water equivalent is projected to decline by 25% (10 to 40%) from 1986-2005 to 2031-2050, regardless

of the scenario. This will continue with a 30% decrease at the end of the 21st century for a low emission scenario to 80% for high emission scenario (*high confidence*). [Section 2.2.6.2].

**2.1.9** Mediterranean Sea surface waters are warming and deep waters are becoming saltier (*high confidence*). [Section 2.2.7.1].

**2.1.9.1** Since the beginning of the 1980s, average Mediterranean Sea surface temperatures have increased throughout the basin, but with large sub-regional differences in the range between +0.29 and +0.44°C per decade, with stronger trends in the eastern basins (Adriatic, Aegean, Levantine and north-east Ionian Sea), marine heat waves have become longer and more intense (*high confidence*). [Section 2.2.7.1].

**2.1.9.2** The water mass temperature and salinity changes of the water outflowing from the Mediterranean Sea through the Strait of Gibraltar are 0.077°C decade<sup>-1</sup> and 0.063 psu (practical salinity unit) decade<sup>-1</sup>, respectively, compared to 2004 (*high confidence*). [Section 2.2.7.1].

**2.1.10** Widespread sea surface temperature increase will continue in the 21st century (*very high confidence*).

**2.1.10.1** During the 21st century, the basin mean sea surface temperature is expected to warm by 2.7 to 3.8°C and 1.1 to 2.1°C under the RCP8.5 and the RCP4.5 scenarios, respectively (*very high confidence*). The sign of future basin average sea surface salinity change remains largely uncertain and its changes will likely be spatially and temporally heterogeneous (*medium confidence*). [Section 2.2.7.2].

**2.1.10.2** Marine heat waves will very likely increase in spatial extent, become longer, more intense and more severe than today (*medium confidence*). Under the high emission scenario, the 2003 marine heat wave may become a regular event for the period 2021-2050 and a weak event at the end of the 21st century (*medium confidence*). [Section 2.2.7.2].

**2.1.11** Mediterranean Sea waters have acidified and will continue to acidify along with the global ocean (*medium confidence*). The Mediterranean Sea is able to absorb relatively more anthropogenic CO<sub>2</sub> per unit area than the global ocean because it is more alkaline and because deep waters are ventilated over shorter timescales (*medium confidence*). [Section 2.2.9].

**2.1.11.1** Sea water surface pH has decreased by -0.08 units since the beginning of the 19th century, similar to the global ocean, with deep waters exhibiting a larger anthropogenic change in pH than typical global ocean deep waters because ventilation times are faster (*medium confidence*). [Section 2.2.9.1].

**2.1.11.2** In 2100, reduction of pH might reach 0.462 and 0.457 units for the western and for the eastern basins, respectively (*low confidence*). [Section 2.2.9.2].

**2.1.12** Mediterranean sea level is rising, similar to global trends, with strong spatial and temporal variation and expected acceleration (*medium confidence*). [Section 2.2.8.1].

**2.1.12.1** Averaged across the Mediterranean Basin, mean sea level has risen by 1.4 mm yr<sup>-1</sup> during the 20th century and has accelerated to 2.8 mm yr<sup>-1</sup> recently (1993–2018) (*high confidence*). [Section 2.2.8.1].

**2.1.12.2** Mostly due to global ocean and ice-sheet dynamics, Mediterranean mean sea level rise is projected to accelerate further throughout the 21st century (*high confidence*). Around 2100, depending on the scenario, the basin mean sea level will likely be 37-90 cm higher than at the end of the 20th century, with a small probability of being over 110 cm (*medium confidence*). [Section 2.2.8.2].

**2.1.12.3** Sea level rise will increase the frequency and intensity of coastal floods and erosion (*high confidence*). [Section 2.2.8.2].

## 2.2 Pollution

**2.2.1** Across the Mediterranean Basin, ocean and inland pollution are transboundary, ubiquitous, diverse and increasing in both quantity and in the

number of pollutants, due to demographic pressure, enhanced industrial and agricultural activities, and climate change (*high confidence*). [Section 2.3.1].

## Fertilizer use and nitrogen release in the Mediterranean region



Figure SPM.4 | Fertilizer use and nitrogen release in the Mediterranean Sea (UNEP/MAP/MED POL, 2013).

### 2.2.2 Pollution of sea water

**2.2.2.1** Mediterranean waters are generally oligotrophic (low nutrient), with decreasing levels from Gibraltar eastwards to the Levantine Sea. Several coastal regions are hotspots of human-induced nutrient inputs (Lagoons of Venice and Bizerte, Gulfs of Lion and Gabès, eastern Adriatic and western Tyrrhenian Sea, North Lake of Tunis, Algerian-Provençal Basin and the Gibraltar Strait) (*high confidence*) (Fig. SPM.4). (Section 2.3.3.1).

**2.2.2.2** Nutrient enrichment causes eutrophication and may provoke harmful and toxic algal blooms, trends which will likely increase. Harmful algal blooms may cause negative impacts on ecosystems (red-tide, mucilage production, anoxia) and may present serious economic threats for fisheries, aquaculture and tourism. They may also harm human health, since 40% of blooming microalgae are able to produce toxins responsible of different human intoxications. Harmful algal blooms can also occur in freshwater environments. (Section 2.3.4).

**2.2.2.3** Emerging contaminants (related to recently discovered chemicals or materials) are prevalent across the Mediterranean Basin, and enhanced by increasing inflow of untreated wastewater. These substances may cause disorders of the nervous, hormonal and reproductive system (*high confidence*). (Section 2.3.3.5).

**2.2.2.4** The increasing frequency of extreme precipitation events in the north of the Mediterranean increases the supply of faecal bacteria and viruses to the coastal zone (*medium confidence*). (Section 2.3.4).

**2.2.2.5** The Mediterranean Sea is one of the most polluted large water bodies globally in terms of plastic and the level of this pollution is expected to increase in the future (*medium confidence*). (Section 2.3.2.3). Even with rigorous reduction of use, plastic debris and their dissolved derivatives will remain a problem since they can take 50 or more years to fully decompose (*medium confidence*) (Section 2.3.2.3).

### 2.2.3 Air pollution

**2.2.3.1** The Mediterranean Basin is among the regions in the world with the highest concentrations of gaseous air pollutants ( $\text{NO}_2$ ,  $\text{SO}_2$  and  $\text{O}_3$ ). Its dry and sunny climate, and specific atmospheric circulation patterns enhance air pollution levels (*high confidence*). (Section 2.3.3.2) Emissions of aerosols and particulate matter (PM) into the atmosphere arise from a variety of anthropogenic activities (transport, industry, biomass burning, etc.), but also from natural sources (volcanic eruptions, sea salt, soil dust suspension, natural forest fires, etc.). (Section 2.3.2.1).

**2.2.3.2** Ships are among the major emitters of  $\text{SO}_2$  and  $\text{NO}_x$ , along with road traffic. Their contribution to transport sector emissions and general air pollution in the Mediterranean Basin is increasing (*medium confidence*). (Section 2.3.3.2).

**2.2.3.3** Tropospheric ozone ( $\text{O}_3$ ) concentrations observed in the summer across this region are among the highest in the northern Hemisphere and still increasing in average and with more frequent high-level episodes. They are influenced by Volatile Organic Compounds (VOCs),  $\text{NO}_x$  emissions and

climate. This trend will likely be enhanced by future warming (*medium confidence*). (Section 2.3.3.2).

**2.2.3.4** Particular meteorological conditions and natural sources, including the proximity of the Sahara Desert, create specific patterns of aerosol

concentrations that may influence particulate matter (PM) concentrations. The occurrence of critically high PM concentrations associated with dust outbreaks is higher in the southern Mediterranean (>30 % of annual days) than in the northern area (<20% of annual days) (*high confidence*). (Section 2.3.2.1).

## 2.3 Land and sea use change

**2.3.1** Landscapes and their use have changed over millennia in the Mediterranean Basin, however the rate of change has increased substantially since the second half of the 20th century (*high confidence*). (Section 2.4.1.1).

**2.3.1.1** Urban and peri-urban areas are growing rapidly all over the Mediterranean, especially along the coasts. Urbanization is a major driving force of biodiversity loss and biological homogenization causing landscape fragmentation, loss of open habitats and of the land use gradient, replacing agricultural systems and natural vegetation (*high confidence*). (Section 2.4.1.2).

**2.3.1.2** Outside urban areas and areas with intensive agriculture, forest and shrub encroachment, as a consequence of abandoned agro-pastoralism, mainly affects marginal lands, arid and mountain regions, primarily in the north (*high confidence*). (Section 2.4.1.1).

**2.3.1.3** In many regions of North Africa and the Middle East (but also on some Mediterranean islands), the dominant land use change process is forest degradation caused by land overexploitation. From the 1980's to the 1990's deforestation has increased by 160% (*high confidence*). (Section 2.4.1.1 and 2.4.1.2).

**2.3.1.4** Future land use trends depend strongly on regional policies for urbanization, ag-

riculture, forestry and nature conservation. Grassland and pastures will likely continue to further decrease in extension due to rural abandonment, often due to insufficient job opportunities and public services in marginal areas (*medium confidence*). (Section 2.4.1.3).

**2.3.2** Marine resource overexploitation and unsustainable fishing practices are the main driver of marine species population decline. (Section 2.4.2).

**2.3.2.1** Fishing efforts have increased over long periods, but particularly so since the 1990's due to new technologies and higher capacity vessels (*high confidence*). (Section 2.4.2.1).

**2.3.2.2** In 2010, the cumulative percentage of collapsed and overexploited stocks exceeded 60% across the Mediterranean Sea (*medium confidence*). The eastern Mediterranean is the most overexploited sub-basin with the highest number of collapsed species (*medium confidence*). (Section 2.4.2.2).

**2.3.2.3** Sustainable management of marine resources requires reduced fishing pressure. The implementation of an ecosystem-based approach may ensure the recovery of both high and low trophic levels and support both ecosystem health and resilience against sea warming (*high confidence*). (Section 2.4.2.3).

## 2.4 Non-indigenous species

**2.4.1** The Mediterranean Sea (and particularly the Levantine Basin) is a hotspot for the establishment of many non-indigenous species (*high confidence*). (Section 2.5.1).

**2.4.1.1** Among known marine non-indigenous species introduced over the last 30 years, invertebrates dominate with >58% (mostly mollusks and

decapods), primary producers follow with approx. 23% and vertebrates with 18% (mostly fish) (*high confidence*). (Section 2.5.1.1).

**2.4.1.2** Most marine non-indigenous species arrive from the Red Sea and Atlantic Ocean, but the highest impact is attributed to those introduced by ships and aquaculture (*high confidence*). (Section 2.5.1.2).

**2.4.1.3** The increase in non-indigenous species can be linked to decrease or collapse in populations of native species, and to other ecological changes to the marine ecosystem (*high confidence*). (Section 2.5.1.2).

**2.4.1.4** The number and spread of non-indigenous species will likely increase further with increasing shipping activity and the impacts of climate on the ocean (*medium evidence*). Forecasting future establishment of non-indigenous species using species distribution models is challenging. (Section 2.5.1.3).

**2.4.2** On land, there is a high number of non-indigenous species in human-modified ecosystems and in regions with high infrastructure development (*high confidence*). (Section 2.5.2.1).

**2.4.2.1** On land, most non-indigenous species in the region are plants (introduced intention-

ally as ornamentals), followed by invertebrates. Phytophagous pests, which cause damages to crops and forests, dominate non-indigenous species all over the Mediterranean Basin, accounting for more than a half of the invertebrate species. The main pathways of introduction for vertebrates are accidental escapes (*medium evidence*). (Section 2.5.2.1).

**2.4.2.2** With warming, current major non-indigenous species are predicted to shift northwards by 37 to 55 km decade<sup>-1</sup>, leaving a window of opportunity for new non-indigenous species adapted to xeric conditions. The trend has recently shifted towards increasing numbers of introduced invertebrates and vertebrates. This pattern will very likely continue in the near future, due to increasing air and maritime cargo, where these taxa can be easily transported as stowaways (*medium confidence*). (Section 2.5.2.3).

## 3 - Resources

### 3.1 Water

**3.1.1** Water resources in the Mediterranean are scarce: resources are limited, unevenly distributed and in some areas not accessible, often mismatching human and environmental needs. (Section 3.1.1).

**3.1.1.1** Renewable water resources are unevenly distributed among Mediterranean regions (72 to 74% are located in the northern Mediterranean) and so is the spatial distribution of water needs, but with opposite trends. As a consequence, 180 million people in the southern and eastern Mediterranean countries suffer from water scarcity (<1,000 m<sup>3</sup> capita<sup>-1</sup> yr<sup>-1</sup>) and 80 million people from extreme water shortage (<500 m<sup>3</sup> capita<sup>-1</sup> yr<sup>-1</sup>) (*high confidence*). (Section 3.1.1.1).

**3.1.1.2** River discharge is characterized by high temporal - seasonal and inter-annual - variability and groundwater is the main source of freshwater for some Mediterranean countries (Libya, Malta, Palestine, Israel) (Section 3.1.1.2). In several cases in southern Mediterranean countries, groundwater resources are drawn from fossil aquifers, i.e. non-renewable resources (*high confidence*). (Section 3.1.1.3).


**3.1.1.3** Sustainable water management is complicated by the transboundary nature of many river basins and aquifers, common in Mediter-

anean countries (18% of total renewable water resources originate outside the territories of the southern Mediterranean, 27% in eastern Mediterranean countries (*high confidence*). (Section 3.1.1.1).

**3.1.2** Due to the general scarcity of water resources, conflicts arise from different sectors of water use (agriculture, tourism, industry, people, also biodiversity conservation) (*medium confidence*). (Section 3.1.2).

**3.1.2.1** The spatial distribution of water use per sector in the Mediterranean area is heterogeneous. In southern and eastern countries, agricultural use reaches 76-79%. In the northern part, the four sectors are much more balanced (18-36%, Fig. SPM.5), with differences between countries. (Section 3.1.2.1).

**3.1.2.2** The percentage of irrigated land of the total cultivated area in the Mediterranean is about 25% (but more than 70% in Egypt, Israel, Lebanon, Greece), with a strong increase (21%) in recent years (Section 3.1.2.2). The trend towards more efficient irrigation systems does not always generate absolute water savings due to the introduction of more water demanding crops (e.g. vegetables) (*medium confidence*). (Section 3.1.2.2).



**Figure SPM.5 | Total water consumption** rates across four main sectors and three sub-regions (data source: AQUASTAT).

**3.1.2.3** Tourism activity is at its highest in summer, coinciding with peak demands by irrigated agriculture, creating tensions for water, and this will likely be exacerbated in the future due to climate change (*medium confidence*). (Section 3.1.2.3).

**3.1.2.4** Municipal water use is already constrained in several Mediterranean countries affected by water scarcity, exacerbated by demographic and migratory phenomena, as well as by the limits and obsolescence of water distribution infrastructure (*medium confidence*). Several northern countries have managed to reduce their municipal withdrawal in absolute values while several southern and eastern countries have the opposite trend (*medium confidence*). (Section 3.1.2.5).

**3.1.2.5** Water-related intersectoral conflicts are likely to be exacerbated in the future because of the interactions between climate change (increasing droughts) and ongoing socio-economic and demographic trends (*medium/high confidence*). (Section 3.1.5.2).

**3.1.3** Disastrous flash floods are frequent in many countries including Italy, France and Spain, affecting mainly the coastal areas, in particular, where population and urban settlements are growing in flood-prone areas. These will likely become more frequent and/or intense due to climate change and surface-sealing (*medium confidence*). (Section 3.1.3.3).

**3.1.4** Climate change, in interaction with other drivers (mainly demographic and socio-economic developments including unsustainable agricultural practices), is likely to impact most of the Mediterranean Basin, through reduced runoff and groundwater recharge, increased water requirements for crops, increased conflicts among users, and increased risk of overexploitation and degradation (*high confidence*). (Section 3.1.4.1).

**3.1.4.1** Impacts of even moderate (1.5 to 2°C) global warming and associated socio-economic pathways are expected to stem from reduced precipitation associated with increased evaporation, leading to a decline in runoff water (Section 3.1.4.1). In many regions, this will likely increase low flow periods in summer and the frequency of no-flow events, and higher drought risks (Section 3.1.4.1). More urban populations are likely to be exposed to severe droughts, and the number of affected people will essentially scale with the temperature increase (*high confidence*). (Section 3.1.4.1).

**3.1.4.2** Aquifer recharge will be strongly impacted by warming and reduced rainfall, particularly in semi-arid areas. At current extraction rates, overexploitation of groundwater is likely to continue having a greater impact on decreasing groundwater levels than climate change (*high confidence*). (Section 3.1.4.1).

**3.1.4.3** Important challenges to groundwater quality in coastal areas are likely to arise from salt-water intrusion driven by enhanced extraction of coastal groundwater aquifers and sea-level rise, as well as from increasing water pollution in the southern and eastern Mediterranean (*medium confidence*). (Section 3.1.4.1).

**3.1.4.4** Impacts of global warming levels higher than 1.5 to 2°C on water resources by the end of the 21st century will be significantly stronger, generating substantially increased risks in the Mediterranean region (Section 3.1.4.2). The probability of more extreme and frequent meteorological, hydrological and agricultural droughts will likely increase substantially, with 5 to 10 times more frequent droughts in many Mediterranean regions (*high confidence*). (Section 3.1.4.2).

**3.1.5** The combined dynamics of climate and socio-economic changes suggest that despite an important potential for adaptation to reduce freshwater resource vulnerability, climate change exposure cannot be fully and uniformly counterbalanced. In many regions, socio-economic developments will have greater impact on water availability com-



pared to climate-induced changes (*low confidence*). (Section 3.1.4.2).

**3.1.5.1** Strategies and policies for water management and climate change adaptation are strongly interconnected with all other sectors (e.g., the water-energy-food nexus). Most adaptation and water management strategies rely on the principles of Integrated Water Resources Management (IWRM), which is based on economic efficiency, equity and environmental sustainability, also considering the nexus with agriculture (food production in particular) and energy for building the resilience needed to adapt to climate change. (Section 3.1.5.1).

**3.1.5.2** Technical solutions are available to improve water availability and the efficient use of water resources. Seawater desalination is increasingly used to reduce (potable) water scarcity in arid and semi-arid Mediterranean countries, despite known drawbacks in terms of environmental impacts on near-coastal marine ecosystems and energy requirements with associated CO<sub>2</sub> emissions. Promising new (solar) technologies are under development, potentially reducing both greenhouse gas emissions and costs (*medium confidence*). (Section 3.1.5.2).

**3.1.5.3** Technology is also expected to contribute significantly to the reduction of wastewater volume, its reclamation and reuse and the reduction of impacts on sea water quality. Agricultural, industrial and watering activities present together approx. 70% water reuse potential. The proposal has been made to recharge aquifers with treated wastewater, but critical issues in terms of water quality remain to be resolved (*medium confidence*). (Section 3.1.5.2).

**3.1.5.4** Inter-basin transfer of water has been implemented in several large-scale schemes, with high social and environmental costs, and risks of conflict (*low confidence*). (Section 3.1.5.2).

**3.1.5.5** Dams for water storage or hydropower exist in most countries, and rivers are diverted for water management in some countries. Large dams often generate social and environmental impacts, such as the destruction of river and wetland ecosystems and the loss of aquatic biodiversity, forced relocation of people and loss of cultural resources. Reductions of these impacts are possible, for example through constructed wetland habitats, and management of fishing and other recreational

opportunities and enhanced coordination among countries sharing the same water resources (*low confidence*) (Section 3.1.5.2). Technological developments also allow for the use of underground- or subsurface dams, to contribute to sustainable management of groundwater. (Section 3.1.5.2).

**3.1.5.6** The strategy of trading commodities (in particular from agriculture) that cannot be produced due to lacking water (virtual water trade) can be considered a form of adaptation. Most Mediterranean countries (e.g., Portugal, Spain, Italy, Greece, Israel, Turkey) have high footprints in terms of national consumption (above 2000 m<sup>3</sup> yr<sup>-1</sup> capita<sup>-1</sup>) (*low confidence*). (Section 3.1.5.1).

**3.1.5.7** Water demand management, i.e. methods used to save (high quality) water, may reduce water consumption or water losses. This includes technical, economic, administrative, financial and/or social measures, with priority for increases in water use efficiency, in particular in the tourism and food sectors and with case-specific solutions integrating traditional knowledge with modern technical achievements (*high confidence*). (Section 3.1.5.1).

**3.1.5.8** The reduction of water losses in all sectors of water use in the Mediterranean is crucial for sustainable management and adaptation strategies. Leakage in urban distribution networks and inefficient irrigation technologies are in urgent need of being addressed (*high confidence*). (Section 3.1.5.1).

**3.1.5.9** Maintaining the traditional Mediterranean diet and shifting back to a locally produced Mediterranean food in conjunction with a reduction of food waste, could generate water savings in comparison to the present increasingly meat-based diet: 753 l for a locally produced diet and 116 l for less waste of water per capita and per day, in addition to benefits for health (obesity, diabetes) (*high confidence*). (Box 3.1.2).

## 3.2 Food

**3.2.1** Warmer and drier climate conditions, with more frequent and intense extreme events, in combination with higher soil salinization, ocean acidification and land degradation, sea level rise and the emergence of new pathogens pose a threat to most elements of the food production system in the Mediterranean Basin (*high confidence*).

**3.2.1.1** Climate extremes pose a threat to the entire agricultural sector. Crop yield reductions are projected for the coming decades in most current areas of production and for most crops if no adaptation takes place. (*Section 3.2.2.1*).

**3.2.1.2** Maize is the crop most affected by climate change, projected to decline in yield by up to 17% in some countries by around 2050 under RCP8.5 scenario and assuming current agricultural practices (*medium confidence*); it could become infeasible in regions with limited access to irrigation water (*medium confidence*) (*Section 3.2.2.1*). Wheat yield losses of 5% to 22% are also projected because of decreased resilience of production and higher inter-annual variability in 2021-2050 under RCP8.5 scenario with no adaptation. Other water demanding crops, e.g., tomatoes, are also at risk. The production of some currently rainfed crops, such as olives, could become infeasible without irrigation (*medium confidence*). (*Section 3.2.2.1*).

**3.2.1.3** Increasing atmospheric CO<sub>2</sub> concentrations may help offset yield losses for some crops, such as wheat and barley, but this effect could impact nutritional quality. Beneficial effects of CO<sub>2</sub> are likely limited by water stress conditions as well as by nutrient availability (*low confidence*). (*Section 3.2.2.1*).

**3.2.1.4** Climate extremes, such as heat stress, droughts, and floods, can cause crop yield losses/failures, crop quality reduction and impacts on livestock (*high confidence*) (*Section 3.2.1.4*). These events can also induce long-term socio-economic and landscape changes (*medium confidence*). (*Section 3.2.1.4*).

**3.2.1.5** Sea level rise will likely impact the agricultural sector by a direct impact on (or loss of) agricultural areas in coastal zones (e.g., in Egypt), along with up to a three-fold increase in the salinity of irrigation water and soil, and retention of sediments that do not reach the coast (*high confidence*). (*Section 3.2.2.1*).

**3.2.1.6** New and/or re-emerging pests and pathogens may contribute to larger than estimated losses in the agricultural sector. Food quality and

security may also be affected by mycotoxigenic fungal pathogens and a higher level of contamination (*medium confidence*). (*Section 3.2.2.1*).

**3.2.1.7** Total landings from Mediterranean fisheries have declined by 28% from 1994 to 2017 (*Section 3.2.1.3, Fig. 3.22*). Climate change is projected to heavily affect marine resources in the coming decades. Warming, acidification and water pollution are likely to reduce marine productivity, affect species distribution and trigger local extinction of more than 20% of exploited fish and marine invertebrates by 2050 (*high confidence*). (*Section 3.2.2.2*).

**3.2.1.8** Perturbations in global markets for agricultural and marine products, potentially caused by environmental change elsewhere, may exacerbate the local impacts of climate change, especially because most Mediterranean countries are net importers of cereal and fodder/feeding products (*high confidence*). (*Section 3.2.1.5*).

**3.2.2** Adaptation to environmental change will be of key importance to limit and partially offset the impacts of climate change in the food sector (*high confidence*).

**3.2.2.1** Projected yield losses in most crops may be reduced by targeted adaptation strategies, such as crop diversification, adapting the crop calendar and use of new varieties adapted to evolving climate conditions. Strategies based on increased irrigation will have limited applicability in the region. Thus, adapted production of crops such as maize will depend on more drought-resistant varieties (*medium confidence*). (*Section 3.2.3.1*).

**3.2.2.2** Successful adaptation strategies are based on combining different approaches, i.e. on farming practices (e.g., varieties, rotational patterns, crop diversity, agroforestry) and management (e.g., diversification of income, modifying irrigation practices). Sectoral co-designed climate services may help reduce risks linked to unfavorable climate conditions and extremes (*medium confidence*). (*Section 3.2.3.1*).

**3.2.3** The food production system on land has the capacity to contribute to greenhouse gas mitigation strategies through nitrogen fertilization optimization, improved water management, better storage of soil organic carbon and carbon sequestration, management of crop residues and agroindustry by-products (*high confidence*). (*Section 3.2.3.2*).

**3.2.3.1** N<sub>2</sub>O emissions in Mediterranean agro-ecosystems can potentially be mitigated by 30 to 50%, through adjusted fertilization (rate and timing). Replacing mineral nitrogen with organic fertilization provides soil and crops not only with nitrogen, phosphorus, potassium and micronutrients, but also enhances organic carbon when using solid fertilizers (i.e., solid manure, compost, etc.), this would be beneficial in many Mediterranean soils with low organic carbon contents (*medium confidence*). (Section 3.2.3.2).

**3.2.3.2** Optimized irrigation techniques may decrease greenhouse gas emissions from Mediterranean regions in perennial crops and intensive vegetable cropping systems on paddy soils (water table management) (*medium confidence*). (Section 3.2.3.2).

**3.2.3.3** Soil organic carbon content in Mediterranean croplands is responsive to management changes such as organic amendments, cover crops and tillage reductions. There is high potential to enhance soil organic carbon storage through land restoration (as proposed by the “4‰ initiative” proposed 2015 by France during the UN-FCCC COP21). Organic fertilizers, tillage reduction and residue retention are effective practices in herbaceous systems. Woody systems, in which the carbon storage potential is higher, can benefit from maintaining a soil cover and use of agro-industry byproducts, such as composted olive mill waste, as a source of organic matter (*medium confidence*). (Section 3.2.3.3).

## 3.3 Energy transition in the Mediterranean

**3.3.1** From 1980 to 2016, primary energy consumption in the Mediterranean Basin steadily increased by approx. 1.7% yr<sup>-1</sup>, mostly due to changing demographic, socio-economic (lifestyle and consumption) and climate conditions (*high confidence*). (Section 3.3.2.1: Fig. 3.25).

**3.3.1.1** The current level of Mediterranean greenhouse gas emissions is approx. 6% of global emissions, close to its proportion of the world population. International climate policy agreements demand an accelerated energy transition in the countries of this region to enable secure, sustainable and inclusive development. (Section 3.3.1).

**3.3.1.2** The contribution of oil to energy production has remained stable between 1995 and 2016, while that of coal has gradually decreased. Primary energy production from natural gas has doubled, while the contribution of nuclear power and renewable energy sources contribution has risen by about 40% (*high confidence*). (Section 3.3.2.1, Fig. 3.28).

**3.3.1.3** While northern rim countries advance towards the transition by gradually diversifying their energy mix, improving energy efficiency and increasing the share of renewable energies, despite recent investments, some eastern and southern rim countries lag behind in these developments (*high confidence*). (Section 3.3.3.2).

**3.3.2** Projected trajectories for energy demand over the next few decades in the Mediterranean Basin differ significantly between the northern and the eastern/southern rim countries (*high confidence*). (Section 3.3.3.2).

**3.3.2.1** Energy demand in the north has decreased by 8% since 2010, due to moderate population growth, increasing efficiency and a stable economy, and is expected to continue to decrease. In 2040, northern Mediterranean energy demand would be 22%, 10% and 23% lower than 2015 levels, for three stylized energy policy scenarios (“transition” - TS, “reference” - RS, and “proactive” - PS), respectively (*medium confidence*). (Section 3.3.3.2).

**3.3.2.2** Southern Mediterranean countries have undergone sustained economic and population growth over recent decades. Energy demand is thus expected to continue increasing and to reach 55% (TS), 118% (RS) and 72% (PS) by 2040 when compared to 2005 (*medium confidence*). (Section 3.3.3.2).

**3.3.3** Climate change in the Mediterranean is expected to impact energy production (due to impacts on infrastructure) and energy use (by decreased heating demand and increased cooling needs). (Section 3.3.2.3).

**3.3.3.1** Losses in power generation are projected due to warming in the region, with only

marginal impact if global warming does not exceed 2°C (losses <5%), but rapid deterioration beyond 2°C (losses >5% reaching 10% at specific locations) *(low confidence)*. (Section 3.3.3.5).

**3.3.3.2** Traditional hydropower and thermo-electric power usable capacity is expected to decline, due to decreased streamflow and increased water temperature, leading to a 2.5 to 7% decrease in hydropower by 2050 and 10 to 15% decrease in thermopower by 2050 (ranges indicate RCP2.6 vs. RCP8.5 estimates vs 1971-2000) *(high confidence)*. (Section 3.3.3.5).

**3.3.3.3** Weather and climate variability, as well as extreme events, cause significant impacts on the availability and magnitude of renewable energy generation. With the increase of the share of renewable energies, the electricity transmission system will be more exposed to weather variations and may be threatened by specific weather conditions that are usually not considered as extremes *(medium confidence)*. (Section 3.3.2.3).

**3.3.3.4** With warming, all Mediterranean countries will experience a net increase in energy demand for cooling. The change in average daily peak electric load from 2006-2012 to 2080-2099 under RCP4.5 climate change scenarios is up to 4-6% (Balkans) and 8-10% under RCP8.5 (Balkans, Spain, Portugal) *(high confidence)*. (Section 3.3.3.6, Fig. 3.50).

**3.3.4** The Mediterranean Basin has significant potential for additional renewable energy production, on land and in the ocean. These include wind, solar, hydro, geothermal and bioenergy as well as energy generation by waves and currents *(high confidence)* (Section 3.3.2.2). There is also potential for high energy efficiency gains *(high confidence)*. (Section 3.3.3.2).

**3.3.4.1** Thermal energy from biomass (mainly wood residues and waste) currently exceeds use of all other renewable energies, mainly for the production of heat or fuel (less for electricity). Overall production of energy from solid biomass is currently 1.56 PW, varying considerably between countries and mainly concentrated on the northern rim. The production of firewood has increased by about 90% in north Africa over the last 60 years and has recently returned to its 1960's level in southern Europe, after a significant reduction from 1973 to 2009 *(medium confidence)*. (Section 3.3.2.2).

**3.3.4.2** Although fossil fuels are expected to remain the dominant component of the energy

mix until 2040, renewable energies will overtake natural gas and coal and become the second most used energy source in the Mediterranean Basin. In 2040, the share of renewable energies would triple to reach 27% in TS, 13% in the RS and 24% in PS (scenarios "transition" - TS, "reference" - RS, and "proactive" - PS) *(high confidence)*. (Section 3.3.3.3).

**3.3.4.3** Among the various renewable energy technologies, solar is expected to grow at the fastest pace in both sub-regions. End usage of solar thermal energy, in particular solar water heaters, has high potential in the south and is efficient with a good return on investment *(medium confidence)*. (Section 3.3.3.3).

**3.3.4.4** The potential for energy efficiency enhancements is substantial in the Mediterranean Basin, particularly in the south *(high confidence)*. Overall, energy intensity is decreasing in the region, largely related to shifts in the buildings, industry and transport sector *(high confidence)*. (Section 3.3.3.2).

**3.3.5** By further improving energy efficiency and deploying renewable energies on a large scale, the entire Mediterranean region can reduce tensions on energy security for importing countries, improve opportunities for exporting ones and reduce energy costs and environmental damage for the whole region. Embarking on an energy transition path will also help improve social welfare in the region and contribute to job creation, among other positive externalities *(medium confidence)*. (Section 3.3.3).

**3.3.5.1** Given socio-economic development and climate change, an important gap between energy supply and demand is expected, particularly in southern and eastern rim countries. This challenge can be met by rapid restructuring of the energy sector, and particularly further accelerated integration of renewable energies *(medium confidence)*. (Section 3.3.4.2).

**3.3.5.2** Advantages/asures of the energy transition include: (i) drastic reduction of per capita greenhouse gas emissions, (ii) return on investment in renewable energies, which may lead to savings of up to 54% in energy costs for a given country, and (iii) establishment of a CO<sub>2</sub> emissions trading market which will provide economic incentives for investments in renewable energies *(medium confidence)*. (Section 3.3.4.2).

**3.3.5.3** Despite electrification rates of almost 100% in southern and eastern rim countries,

the energy dynamics of these countries are largely unsustainable in the long term, as a result of a highly subsidized electricity market (with some exceptions, e.g., Turkey) leading to a systemic misallocation of resources, population growth, increasing urbanization and expected socio-economic changes in the region, and global warming (*high confidence*). (Section 3.3.4.3).

**3.3.5.4** A change in domestic energy policies, including reforming the energy pricing mechanisms, and/or the introduction of tax and regulatory incentives may be needed in some southern and eastern rim countries to reduce the cost disadvantage of renewable energies compared to fossil fuels (*medium confidence*). (Section 3.3.4.2).

**3.3.5.5** Regional energy market integration and cooperation are needed to unleash cost-effective climate change mitigation. (Section 3.3.4.5). Cross-border regulations require the convergence of national regulations to allow interconnections to work effectively. Investment regulation requires the design and development of infrastructure that will be needed for promoting international com-

plementarities and technical standards (*high confidence*). (Section 3.3.4.5).

**3.3.6** Mediterranean islands experience specific threats, challenges and opportunities in the context of global change and energy transition. Geographical and socio-economic singularities of Mediterranean islands put additional pressure on water and energy, leading to resource depletion and environmental degradation, threatening sustainable development, especially during the high touristic season when population doubles for some (*high confidence*). (Box 3.3.2).

**3.3.6.1** On most islands, energy demand is set to increase, due to socio-economic trends including tourism, but also due to expected increase in the use of energy-intensive desalination techniques (*medium confidence*). (Box 3.3.2).

**3.3.6.2** Enhancement of hydropower is limited on most Mediterranean islands, but there is important potential for wind power and hydrogen generation (*medium confidence*). (Box 3.3.2).

## 4 - Ecosystems

### 4.1 Marine ecosystems

**4.1.1** Mediterranean marine ecosystems are unique due to their high number of endemic species, but they are also highly vulnerable to local and global pressures including environmental change. (Section 4.1.1.1).

**4.1.1.1** The Mediterranean Sea represents the highest proportion of threatened marine habitats in Europe (32%, 15 habitats) with 21% being listed as vulnerable and 11% as endangered. This threat includes several valuable and unique habitats (e.g., seagrasses and coralligenous), supporting an extensive repository of biodiversity. Despite covering only 0.82% of the planet's ocean surface, the Mediterranean Sea hosts 18% of all known marine species (*high confidence*). (Section 4.1.1.1).


**4.1.1.2** Over millennial time-scales, productivity in the overall oligotrophic Mediterranean Sea responds rapidly to short and long-term changes in nutrient input, either from rivers, winds or upwelling activity, all of which modify the benthic-pelagic ecosystems by extending into the entire food chain (*high confidence*). (Section 4.1.1.2).

**4.1.1.3** Tropical non-indigenous species are spreading into the Mediterranean through current warming trends, causing "tropicalization" of marine fauna and flora (*medium confidence*). (Section 4.1.1.1).

**4.1.1.4** Acidification in Mediterranean waters will likely impact the marine trophic chain, from its primary producers (i.e., coccolithophores and foraminifera) to corals and coralline red algae (*medium confidence*). (Section 4.1.1.1).

**4.1.1.5** Climate change and direct human activities impact the integrity of marine ecosystems by disturbing plankton ecology, increasing jellyfish outbreaks, reducing fish stocks, and more generally causing changes in physiology, growth, reproduction, recruitment and behavior in marine organisms (*medium confidence*). (Section 4.1.1.1).

**4.1.2** The combination of various ongoing climate drivers of environmental change (e.g., sea warming, ocean acidification, and sea level rise) has numerous detectable effects on marine organisms



**Figure SPM.6 | Climate change drivers potentially affecting marine pelagos and benthos in the Mediterranean Sea.**

acting at individual, population, and ecosystem scales. Expected future impacts include major reorganizations of the biota distribution, species loss, decrease in marine productivity, increase in non-indigenous species, and potential species extinctions (*medium confidence*) (Fig. SPM.6). (Section 4.1.2.1).

**4.1.2.1** Projections for high emission scenarios show that endemic assemblages will be modified by 2041–2060 and among 75 Mediterranean endemic fish species, 31 will likely extend their geographical range, while 44 will likely reduce it (*medium confidence*).

**4.1.2.2** Alterations of natural habitats for commercially valuable species are likely to occur, resulting in many repercussions on marine ecosystem services such as tourism, fisheries, climate regulation, coastal protection, and ultimately on human health (*medium confidence*). (Section 4.1.2.2).

**4.1.2.3** In general, small pelagic species, thermophilic and/or exotic species of smaller size and of low trophic levels, could benefit from envi-

ronmental change. Large-sized species, often with commercial interest may find conditions for survival reduced (*medium confidence*). (Section 4.1.2.1).

**4.1.3** Adaptation strategies to reduce environmental change impacts on marine ecosystems need to occur in conjunction with climate mitigation and pollution reduction policies and actions. (Section 4.1.3.4).

**4.1.3.1** Due to the diversity of marine community responses to climate change and other stressors in different sub-basins, wider monitoring coverage is needed to improve knowledge of the different adaptation processes that characterize and best suit each zone (*high confidence*). (Section 4.1.3.1).

**4.1.3.2** All measures that improve marine ecosystem health, resilience or biodiversity have the potential to delay and reduce the adverse effects of climate drivers. These include more sustainable fishing practices, reducing pollution from agricultural activity, sustainable tourism and more effective waste management (*high confidence*). (Section 4.1.3.4).

**4.1.3.3** Marine protected areas can provide an “insurance” role for biodiversity if they are placed in locations with limited vulnerability to ocean acidification and climate change (*medium confidence*) (Section 4.1.3.4). While marine protected areas cannot halt climate change and its consequences, such as ocean acidification, they are an important tool for enhancing the resilience and adaptive capacity of ecosystems (*high confidence*). (Section 4.1.3.2).

**4.1.3.4** Developing practical management actions that take into consideration the uniqueness of each species and their responses towards different drivers is crucial to increasing their resilience and plasticity in the context of climate change (*high confidence*). (Section 4.1.3.3).

## 4.2 Coastal ecosystems

**4.2.1** The coastal zone, i.e. the area in which the interaction between marine systems and the land dominate ecological and resource systems, is a hotspot of risks, especially in the MENA region (*high confidence*). (Section 4.2.1.1).

**4.2.1.1** Alterations of coastal ecosystem regimes (lagoons, deltas, salt marshes, dune systems, etc.) due to climate change and human activities affect the flow of nutrients to the sea, the magnitude, timing and composition of plankton blooms, significantly increase the number and frequency of jellyfish outbreaks, and could have negative impacts on fisheries (*high confidence*). (Section 4.2.1.1).

**4.2.1.2** In addition to hosting a wide diversity of wild faunal and floral species, coastal ecosystems are also often used as aquaculture platforms (i.e., fish, shellfish cultures, etc.), and the pressures on them may have significant consequences on their usages (*medium confidence*). (Section 4.2.1.1).

**4.2.1.3** Seagrass meadows in the Mediterranean Sea cover 1.35 to 5 million hectares, between 5 and 17% of the worldwide seagrass habitat. The current loss rate of seagrass is approx. 5% in the Mediterranean. Even in the remaining *Posidonia* meadows, almost half of the surveyed sites have suffered net density losses of over 20% in 10 years (*medium confidence*). (Section 4.2.1.1).

**4.2.1.4** The rapid spread of non-indigenous fish species represents a serious problem for trophic networks and fisheries in coastal areas, due to the local extinction of species that are preys of these generalist fish (*high confidence*). (Section 4.2.1.1).

**4.2.2** In the future, environmental change, particularly warming, decreasing nutrient replenishment, and ocean acidification, are expected to cause changes in plankton communities at differ-

ent levels, from phenology and biomass to community structure (*medium confidence*) (Section 4.2.2.1). Negative impacts are also expected to affect fish, corals and seagrass meadows, while non-indigenous species are expected to be favored (*medium confidence*). (Section 4.2.2.1).

**4.2.2.1** Sea level rise impacts coastal wetlands and estuaries, while reduced precipitation and prolonged droughts will reduce the water discharge and sediments flow of Mediterranean rivers and catchments. Mobile coastlines are likely to retreat or disappear because of the effects of erosion due to the accelerated rise in sea level, with the most severe impacts affecting the least mobile species (*medium confidence*). (Section 4.2.1.1 and 4.2.2.2).

**4.2.2.2** Mediterranean coasts are expected to suffer further severe disturbance due to intensive urbanization and other land uses, which could worsen as land availability decreases and population growth continues. In the future, coastal storms and floods, probably more frequent and intense, will have adverse impacts on ecological balances, as well as human health and well-being, particularly in Mediterranean coastal cities (*medium confidence*). (Section 4.2.2.3).

**4.2.3** Developing more integrated approaches would support adaptation policies for the entire Mediterranean, involving ecosystem-based management of coastal areas, identifying synergies and conflicts, as well as integrating local knowledge and institutions. (Section 4.2.3.6).

**4.2.3.1** Suitable adaptation policies include (i) reducing pollution from runoff, both from agriculture, industry and waste management, (ii) defining policies to limit or prevent acidification and (iii) moving aquaculture operations to areas pro-

tected from critical acidification levels (*high confidence*). [Section 4.2.3.1].

**4.2.3.2** Early Detection and Rapid Response has been recognized as a key aspect for non-indigenous species management. Efficient public

awareness campaigns disseminating information to local communities may help to quickly detect unwanted non-indigenous species, together with formalized early warning systems (*medium confidence*). [Section 4.2.3.3].

## 4.3 Terrestrial ecosystems

**4.3.1** Terrestrial biodiversity changes in the Mediterranean Basin over the past 40 years have occurred more quickly and extensively than in most other regions in the world. Urbanization and the loss of grasslands are key factors in ecosystem degradation across the region. Since 1990, agricultural abandonment has led to a general increase in forested area of 0.67% yr<sup>-1</sup> across the basin, with significant variations between northern and southern shores of the Mediterranean. [Section 4.3.1.2].

**4.3.1.1** Since about 1980, biodiversity changes have occurred more quickly and extensively in different Mediterranean species groups and habitats than before. Species loss is marked by a general trend of homogenization (loss of vulnerable and rare species) recorded in several species groups, and also by a general simplification of biotic interactions (loss of specialized relationships) (*high confidence*) [Section 4.3.1.2].

**4.3.1.2** In all Mediterranean mountain regions, subalpine species move to higher altitudes wherever this is possible (*medium confidence*). [Section 4.3.1.2].

**4.3.1.3** Almost all countries in the northern sub-region have undergone increase in forest area due to the decline of extensive agriculture and agro-pastoral systems, with rates around 1% yr<sup>-1</sup> in Italy, France and Spain. In the southernmost areas, semi-natural ecosystems are more at risk of fragmentation or disappearance due to human pressure from clearing and cultivation, overexploitation of firewood and overgrazing (*high confidence*). [Section 4.3.1.2].

**4.3.1.4** Agro-system biodiversity has declined dramatically since the early 1950s due to the intensification of agriculture, leading to an increase of highly modified agroecosystems and simplified agricultural landscapes (*high confidence*). Traditional and extensive agricultural practices, including agro-ecological methods, gener-

ally help maintain high biodiversity levels (*medium confidence*). [Section 4.3.1.2].

**4.3.1.5** Over the last five decades, agricultural production has increasingly been impacted by loss of pollinators, with an increase by a factor of three in the number of crops requiring the intervention of pollinators (*medium confidence*). [Section 4.3.1.2].


**4.3.1.6** Mediterranean drylands have a significant and specific biodiversity value, with most plants and animals highly adapted to water-limited conditions. [Section 4.3.1.2]. European Mediterranean drylands are undergoing an overall increase in the percent of arid area in response to climate change and extensive land abandonment. Almost 15% of the humid Mediterranean domain has been replaced by more arid area since the 60s, while arid area has remained stable (*medium confidence*). [Section 4.3.1.2].

**4.3.1.7** Freshwater ecosystems offer many important ecosystem services (e.g., water supply for drinking, agriculture and industries, water purification, erosion control, recreation, tourism and flood mitigation) [Section 4.3.1.2: freshwater ecosystems]. 48% of Mediterranean wetlands were lost between 1970 and 2013, with 36% of wetland-dependent animals in the Mediterranean threatened by extinction (*high confidence*). [Section 4.3.1.2].

**4.3.2** Drier climate and increased human pressure are expected to cause significant impacts on terrestrial biodiversity, forest productivity, burnt area, freshwater ecosystems and agro-systems during the 21st century (*medium confidence*). [Section 4.3.2].

**4.3.2.1** All factors considered, a general reduction of forest productivity in the medium- and long-term is likely associated with higher mortality and dieback, particularly for species or populations growing in water-limited environments,





**Figure SPM.7 | Distribution of drylands and their subtypes** based on observations for the 1981–2010 period. Areal cover of drylands per subtype is estimated within the boundaries of the Mediterranean SREX region (dashed line). (B, C, D) Distribution of projected dryland transitions for three Global Warming Levels (GWLs: +1.5°C, +2°C and +4°C above preindustrial levels), relative to the baseline period. Grey shaded areas in (B), (C) and (D) are drylands of the baseline period. Chord diagrams denote the areal extent of projected transitions in each dryland subtype for each GWL (proportional to the total extent of land changing to drier types) (see Section 4.3.2.4, Fig. 4.15)

which constitute the majority of Mediterranean forests (*medium confidence*). (Section 4.3.2.1).

**4.3.2.2** An increase in wildfires, and hence burnt area is projected in Mediterranean Europe under most global warming scenarios. Burnt area could increase across the region by up to 40% for 1.5°C warming and up to 100% from current levels for 3°C warming at the end of the 21st century (*high confidence*). (Section 4.3.2.1).

**4.3.2.3** Most Mediterranean drylands will likely become drier and their extent is expected to increase across the region. Global warming projections of 1.5°C, 2°C and 4°C above pre-industrial levels correspond to 12%, 20% and 41% increases in dryland area respectively (*medium confidence*) (Fig. SPM.7). (Section 4.3.2.3).

**4.3.2.4** For freshwater systems, projections suggest decreased hydrological connectivity, increased concentration of pollutants during droughts, changes in biological communities as a result of harsher environmental conditions, and a decrease in biological processes like nutrient uptake, primary production, or decomposition. Increased pressure by users on the shrinking water resources will likely aggravate impacts on river ecosystems (*medium confidence*). (Section 4.3.2.5).

**4.3.3** For most ecosystems, management options exist that can enhance resilience under environmental change. (Section 4.3.3).

**4.3.3.1** Promotion of "climate-wise connectivity" through permeability of landscapes, conservation or creation of dispersal corridors and habitat networks may all facilitate the upward migration of lowland species to mountains in order to adapt to new climate change conditions (*medium confidence*). (Section 4.3.3.2).

**4.3.3.2** Promotion of more adequate forest management taking into account local conditions and future projections can improve the adaptation of Mediterranean forests to warmer climates (e.g., mixed-species forest stands, thinning, management of understory). The management of spatial heterogeneity in landscapes can help reduce fire extent under climate warming (*low confidence*). (Section 4.3.3.1).

**4.3.3.3** Preserving the natural flow variability of Mediterranean rivers and streams and wide riparian zones, along with reductions in water demand may assist adaptation of freshwater ecosystems to future environmental change (*medium confidence*). (Section 4.3.3.5).

## 5 - Society

### 5.1 Development

**5.1.1** For this report, sustainable development seeks to address the needs of current and future generations, utilizing natural resources in ways that preserve and sustain them, and ensure equitable access to them in the present and the future. If losses in well-being are to be avoided for future generations, sustainability strategies will need to improve well-being and environmental sustainability at the same time. (Section 5.1.1.1).

**5.1.2** Due to the growing impact of climate change on population, institutional response is increasingly needed, at a local, national and international level. This means mitigating, adapting and regulating the action of business and other multinational enterprises, and taking into account human rights issues. (Section 5.1.1.2).

**5.1.2.1** Climate-proofing infrastructure across the entire Mediterranean region is necessary to with-

stand present and future climate change impacts in the coming decades. Investments in research and development greatly reduce the costs of adaptation (*high confidence*). (Section 5.1.1.3).

**5.1.2.2** The Mediterranean has a rich history as well as exceptional natural and cultural landscapes, which attracted more than 360 million tourists in 2017. In the past 20 years, the gross domestic product contribution from the tourism sector has steadily increased by 60% in Mediterranean countries. Climate change will likely impact the thermal comfort of tourists during the main season. Sea-level rise will likely affect beaches and cultural heritage sites (*high confidence*) (Section 5.1.1.3).

**5.1.2.3** A significant part of Mediterranean tourism is oriented towards outdoor activities, which if unmitigated, are at risk of further degrad-

ing natural resources, including freshwater availability (*high confidence*). (Section 5.1.1.3).

**5.1.2.4** Mediterranean tourism has a major role for employment throughout the region, and has the potential to become more resilient to climate change than the overall economy. Sustainable tourism can secure significant employment and help offset the negative economic impact of climate change (*medium confidence*). (Section 5.1.1.3).

**5.1.3** Poverty, inequalities and gender imbalances relate both directly and indirectly to the achievement of sustainable development in Mediterranean countries. The presence of these imbalances, both relative and absolute, hampers economic development, de facto blocking parts of society from the benefits of higher standards of living (Section 5.1.1.3).

**5.1.3.1** The loss to human development due to inequality over the past few years (2010 to 2017) is consistently more significant in southern Mediterranean countries than northern Mediterranean countries (*high confidence*). (Section 5.1.1.3; Box 5.1.1).

**5.1.3.2** Gender inequalities are significant in Mediterranean countries, ranked between the 18th position and the 159th (out of 164) in the global ranking of the Gender Development Index (*high confidence*). (Section 5.1.1.3; Box 5.1.2).

**5.1.3.3** Climate change education means active participation of the community, especially children and youth as agents of change and enhanced collaboration between education policymakers and researchers to set the basis of educational policy and actions in scientific knowledge and expertise (*medium confidence*). (Section 5.1.1.4).

**5.1.4** The expected increasingly extreme climate conditions and pollution of the Mediterranean Basin are likely to result in economic vulnerabilities and risks of higher intensity than in other European regions. (Section 5.1.2).

**5.1.4.1** Higher intensity and more recurrent flash-floods with higher mortality in the eastern Mediterranean directly affect agriculture, commerce, tourism and industry (*medium confidence*). (Section 5.1.2).

**5.1.4.2** The effect of sea level rise, together with changes in storm features is likely to seriously affect port operations, slowing down trade operations and productivity levels (*medium confidence*). (Section 5.1.2).

**5.1.4.3** The economic impact on tourism depends on the country and the season. Some adaptation to warming can be achieved by spreading out tourism offers to the spring and autumn. Northern Mediterranean regions could experience climate-induced tourism revenue decreases of up to -0.45% of gross domestic product per year by 2100 (*medium confidence*). (Section 5.1.2).

**5.1.4.4** Economic costs due to droughts (e.g., on food security) may exceed those caused by earthquakes or floods (*low confidence*). (Section 5.1.1.3).

**5.1.5** The success of adaptation strategies will involve consideration of the specific regional climate conditions, in sectoral, political and socio-economic contexts by ensuring dialogue between stakeholders, through cooperative structures, knowledge transfer and monitoring progress to support regular reviews of policy objectives and the inclusion of new scientific information when it becomes available. (Section 5.1.3).

**5.1.5.1** The variants of sustainable urban growth represented by sustainable cities, resilient cities, green cities or low carbon cities bring opportunities to create pathways for transformative and sustainable urban development (*high confidence*). (Section 5.1.3.1).

**5.1.5.2** Stronger pollution and greenhouse gas emissions control instruments can be deployed. Institutional approaches may facilitate internalization of externalities. Command and control instruments may have an action on production inputs, emission outputs, location or production techniques. Economic incentive (market-based) instruments include taxes, liability payments, emission permits, subsidies etc. (Section 5.1.3.2, Table 5.3).

## 5.2 Human health

**5.2.1** Environmental change has already led to a wide range of impacts on human health in Mediterranean countries, and most trends are likely to continue. (Section 5.2.1.1).

**5.2.1.1** Direct impacts are related to exposure to extreme events as heat waves and cold spells, floods and storms. Interaction with environmental systems leads to indirect impacts such as changes in water availability and quality, in food availability and quality, rising air pollution including pollution from forest fires, and changing patterns of vector-, food- and water-borne diseases (high confidence). (Section 5.2.1.1).


**5.2.1.2** Population vulnerability to the impacts of environmental and climate change is strongly influenced by population density, level of economic development, food availability, income level and distribution, local environmental conditions, pre-existing health status, and the quality

and availability of public health care (high confidence). (Section 5.2.2).

**5.2.1.3** Vulnerable Mediterranean populations include the elderly, the poor, and people with pre-existing or chronic medical conditions, displaced people, pregnant women and babies. People who are disadvantaged due to a lack of shelter, clean water, energy or food are more at risk from extreme events (high confidence). (Section 5.2.2).

**5.2.2** Heat waves are responsible for high mortality rates causing tens of thousands of premature deaths, especially in large cities and among the elderly. Heat-related morbidity and mortality has been partially reduced in recent years by more efficient protection of people (high confidence) (Fig. SPM.8). (Section 5.2.3.1).

**5.2.2.1** Most Mediterranean cities are compact and densely populated and have experienced



**Figure SPM.8 | Attributable fraction of heat-related deaths during summer with different climate scenarios** by country in Europe. A) RCP4.5 in 2050; B) RCP8.5 in 2050, C) RCP4.5 in 2085 and D) RCP8.5 in 2085 (Kendrovski et al., 2017).

strong impacts from extremely high temperatures on their population (*medium confidence*). (Section 5.2.3.1).

**5.2.2.2** In recent decades, mortality rates due to heat stress have been reduced through national plans and alert systems that have raised risk awareness and avoidance among the population (*high confidence*). (Section 5.2.3.1).

**5.2.2.3** The European population at risk for heat stress is expected to increase (4% annually) in the coming years and could increase to 20 to 48% by 2050, depending on different combinations of socio-economic scenarios. Vulnerability varies between regions and the Mediterranean region will be among the most affected. Annual mortality attributable to heat in Mediterranean Europe will increase by a factor of 1.8 and 2.6 for moderate (RCP4.5) or high (RCP8.5) global warming levels, respectively, by the middle of the 21st century, while by the end of the century the increase will be by a factor of 3 and 7 respectively (*high confidence*). (Section 5.2.5.2).

**5.2.2.4** The impact of heat on mortality will be more influenced by socio-economic factors due to the impacts on vulnerability than by the exposure to high temperatures (*medium confidence*). (Section 5.2.5.2).

**5.2.3** Despite the rise in mean temperature, cold waves are not likely to disappear (*high confidence*). Moderate cold-related risk will remain a temperature-related risk throughout the 21st century, in combination with risks due to pathogenic agents (*low confidence*). (Section 5.2.5.3 and 5.2.3.4).

**5.2.4** Environmental changes in the Mediterranean Basin will likely exacerbate risks for vector-borne disease outbreaks in the Mediterranean region, since warmer climate and changing rainfall patterns (together with landscape management) may create hospitable environments for mosquitoes, ticks, and other climate-sensitive vectors, particularly for the West Nile Virus, Chikungunya and Leishmaniasis (*medium confidence*). (Section 5.2.3.3).

**5.2.4.1** Projections for 2025 show an elevated risk for vector-borne diseases in the Mediterranean. By 2050, the West Nile Virus high-risk areas are expected to expand further and the transmission seasons will extend significantly (*medium confidence*). (Section 5.2.5.4).

**5.2.4.2** Future changes in the habitability of the Mediterranean Basin for vector-based disease vectors and pathogens vary geographically and

will significantly modify the extent and transmission patterns in the area. A significant reduction of habitat suitability for the tiger mosquito *Aedes albopictus* (vector for chikungunya and dengue) is projected for the middle of the 21st century in southern Europe and the Mediterranean related to significant increase in summer temperatures (*high confidence*). (Section 5.2.5.4).

**5.2.4.3** With rising average temperatures and increasing frequency and length of heat waves, a rising number of cases of food-borne illness must be expected for business-as-usual scenarios, unless education, epidemiological surveillance and enforcement (related to food safety) are intensified (*high confidence*). (Section 5.2.5.4).

**5.2.5** Every year, around one million fatalities are attributed to outdoor and indoor air pollution in the European and eastern Mediterranean regions. (Section 5.2.4.1).

**5.2.5.1** Synergistic impacts are observed between ozone levels, particulate matter concentrations and climate, especially during heat wave days, with high temporal and spatial variability with a 1.66% increase in mortality for each 1°C temperature increase on low ozone level days and an increase of up to 2.1% on days with high ozone levels. Reducing the exposure to particulate matter improves the life expectancy of Europeans by about 8 months (*high confidence*). (Section 5.2.4.1).

**5.2.5.2** Exposure to forest fire smoke and pollutants of natural origin, such as Saharan dust, is related to increased mortality, respiratory and cardiovascular diseases with variable impacts depending on age (*medium confidence*). (Section 5.2.4.2).

**5.2.5.3** Ozone-related morbidity and mortality is expected to increase by 10-14% from 2021 to 2050 in several Mediterranean countries. The combined influence of O<sub>3</sub> and PM<sub>2.5</sub> (particulate matter with a diameter of less than 2.5 µm) will increase European mortality by 8-11% in 2050 and by 15-16% in 2080 compared to the year 2000 (*medium confidence*). (Section 5.2.5.5).

**5.2.6** Climate change and extreme events have a negative impact on mental health for people who experience loss of homes, destruction of settlements and damage to community infrastructure (*medium confidence*) (Section 5.2.4.3). Displacement may lead to adverse health outcomes, especially for vulnerable population groups as well as those who suffer from chronic diseases (*medium confidence*). (Section 5.2.4.4).

**5.2.7** Prevention plans related to human health should be developed further by specifically considering climate change risks. Most mitigation and adaptation measures for climate change offer synergies with other public health issues, notably air pollution. Mediterranean countries need to en-

hance cross-border collaboration, as adaptation to many of the health risks (e.g., vector-borne diseases, droughts, migration) requires collaboration across borders and also across the different parts of the basin (*low confidence*). (Section 5.2.6.2).

## 5.3 Human security

**5.3.1** Human security is a condition that exists when the vital core of human lives is protected, and where people have the freedom and capacity to live with dignity (*medium confidence*). (Section 5.3.1.1).

**5.3.1.1** Environmental and climate change constitutes a threat to the enjoyment of economic, social and cultural rights, acting as a risk multiplier and a key crosscutting issue for multiple aspects of human rights and international justice. (Section 5.3.2.2).

**5.3.1.2** There is a substantial divide between Mediterranean countries when it comes to individual circumstances and the specific impacts of environmental change on security, which depend on climate but also geographical, social, cultural, economic and political conditions. (Section 5.3.1.1).

**5.3.2** Recent human migration (mostly within southern and eastern countries of the Mediterranean Basin but also between the South and the North) can partially be attributed to environmental change, but other drivers such as economic and political factors are usually more important. While slow-onset environmental and climate-related events have significantly affected human well-being in some areas, adaptation is usually possible for reducing the need for human migration. In contrast, fast-onset events with associated environmental degradation (such as storms and floods) have likely led to migration, mostly temporary and over short-distances (*medium confidence*). (Section 5.3.2.3).

**5.3.3** Climate fluctuations have likely played a role in the decline or collapse of ancient civilizations, probably involving situations of increased violent conflicts. For the contemporary period, several studies indicate a link between armed conflict and environmental change, but other scholars disagree (*low confidence*). (Section 5.3.2.4; Box 5.3.1)

**5.3.3.1** Negative weather shocks such as dry spells occurring during the crop growing season by reducing agricultural production and income may increase the continuation and intensity rather than the

outbreak of civil conflicts, especially in regions with agriculturally-dependent and politically excluded groups. Several recent studies identify a link between higher food prices caused by climate change and urban social unrest in Africa. Rising food prices are considered to have played a significant role in the Arab Spring unrest across North Africa and the Middle East in 2011, although such forms of violence are mostly triggered by a complex set of political and economic factors rather than only by higher food prices caused by climatic change (*low confidence*). (Section 5.3.2.4).

**5.3.3.2** For conflict, the impact of expected future environmental change remains rather speculative. However, recent historical experience makes it likely that severe and rapid climate change could further exacerbate political instability in the poorest parts of the Mediterranean Basin (*medium confidence*). (Section 5.3.3.2).

**5.3.3.3** Knowledge is limited regarding how natural disasters interact with and/or are conditioned by socio-economic, political, and demographic contexts to cause conflict. Future research remains necessary. (Section 5.3.5).

**5.3.4** Parts of the rich Mediterranean cultural heritage, notably many UNESCO World Heritage Sites, are directly threatened by sea-level rise or other aspects of environmental change. There is an urgent need for mitigation and adaptation as a large number of world heritage sites are already at risk today. By 2100, flood risk may increase by 50% and erosion risk by 13% across the Mediterranean region (*high confidence*). (Section 5.3.3.1).

**5.3.5** Culture is a key factor to the success of environmental change adaptation policies in the highly diverse multicultural setting of the Mediterranean Basin. Climate adaptation policies have the potential to infringe on human rights in the Mediterranean region if they are disconnected from concerns such as justice, equity, poverty alleviation, social inclusion, and income redistribution (*high confidence*). (Section 5.3.4.1).

## 6 - Managing future risks and building socio-ecological resilience in the Mediterranean

**6.1** Although national governments have an important role to play in reducing the burden of climate change on human health, it is at the local scale that most actions and measures are taken. These measures include (but are not limited to) the improvement of housing and infrastructure, the education and awareness-raising of the most vulnerable communities, the implementation of early warning systems, the strengthening of local emergency and healthcare services, and the general improvement of the adaptive capacity of the community and local institutions (*high confidence*). (Section 6.2.2).

**6.2** Sustainable water security measures require integrated approaches which include water saving technologies, such as new equipment in irrigation agriculture and households, often complemented by improved water efficiency, multi-scale storages, use of unconventional water sources stemming from recharging wastewater or sea water desalinization. Some of these measures may cause environmental impacts due to soil contamination, energy consumption or coastal ecosystem degradation (*high confidence*). (Section 6.3.3).

**6.3** Adaptation of Mediterranean agriculture to water scarcity will benefit from more sustainable approaches. Many studies on no tillage and agroforestry in the Mediterranean show that these practices may have positive effects on the soil by keeping more water, therefore enhancing yields, especially in water-stressed years (Section 6.4.3). These strategies also have benefits for climate mitigation, since conservation agriculture emits less greenhouse gases and enhances soil carbon sequestration and storage (*medium confidence*). (Section 6.4.2).

**6.4** Anticipated changes in fire regimes can have significant impacts on natural and social systems. These impacts can be exacerbated by some of the current fire suppression policies, such as deployment of prescribed fire over large tracts of land (Section 6.5.3). Transformative changes in fire management practices in the Mediterranean countries are necessary for reducing risk and vulnerability and increasing natural and societal resilience, e.g., development of socio-economic sustainable activities that ensure low overall landscape risk (*medium confidence*). (Section 6.5.4).

**6.5** Land Degradation Neutrality is a conceptual framework to halt the loss of land due to unsustainable management and land use changes. Its purpose

is to maintain the land resource base so that it can continue to supply ecosystem services while enhancing the resilience of the communities that depend on the land. This concept, endorsed by the UNCCD Parties and the sustainable development goals (SDG), just starts to be applied, but could beneficially be extended to further Mediterranean areas (*low confidence*). (Section 6.6.4).

**6.6** Interconnections between hazards may result in consecutive and compound events that can lead to non-linear increases in the magnitude of individual events, thus challenging the resilience of populations living in floodplains. Good practices in flood management include development of dedicated early warning systems, construction of check dams, improvement of drainage systems in urbanized areas, emergency management plans in addition to urban planning for resilience and strategic retreat and nature-based solutions, such as reforestation in upstream areas, floodplain restoration and bank erosion protection, and adequate agricultural practices for retaining water (*medium confidence*). (Section 6.8.2).

**6.7** Sea-level rise will lead to increases in coastal-flood and erosion risk along the entire Mediterranean coast. Proactive adaptation to these hazards is essential for maintaining the functions of coastal zones. Coastal adaptation practices can be classified in the following broad categories: Protect, accommodate, advance, and retreat. Nature-based protection solutions, i.e. beach and shore nourishment as well as dune or wetland restoration, is becoming a more common alternative to hard structures. Flood fatalities are reduced as societies are learning to live with flood hazards (*medium confidence*). (Section 6.9.2).

**6.8** Tourism and recreation, red coral extraction, and fisheries (both capture and aquaculture production) are the sectors that are most vulnerable to sea acidification (Section 6.11.1). Recruitment and seed production present possible bottlenecks for shellfish aquaculture in the future since early life stages are vulnerable to acidification and warming (Section 6.11.1). As an example, seagrasses may provide “refugia” from ocean acidification for associated calcifying organisms, as their photosynthetic activity may raise pH above the thresholds for impacts on calcification and/or limit the time spent below some critical pH thresholds (*medium confidence*). (Section 6.11.4).

**6.9** Although the level of non-indigenous species arrivals will likely remain high in northern countries

in the coming decades, their presence will likely increase substantially in southern and eastern countries where biodiversity may be high but capacity to manage non-indigenous species is low. In such places, unmanaged non-indigenous species may threaten human livelihoods (Section 6.12.1). Only few non-native species succeed in establishing in their new locations and gaining importance, but those that do can result in billions of dollars in costs (medium confidence). (Section 6.12.2).

**6.10** Only few Mediterranean cities have local climate plans that consider mitigation and adaptation in a joint manner. There is an urgent need for more integrated local climate plans. Cities, in particular, need to become more resilient to environmental change as impacts will be disproportionately high in these locations due to a concentration of population and assets in combination with hazard-amplifying conditions (e.g., increased runoff through soil sealing, urban heat island effect). This requires knowledge exchange and promotion of ambitious action against climate and environmental change and new approaches to urban development (medium confidence). (Section 6.13).





# 1 INTRODUCTION

**Coordinating Lead Authors:**

Manfred A. Lange (Cyprus), Maria Carmen Llasat (Spain), Maria Snoussi (Morocco)


**Lead Authors:**

Arnault Graves (Spain/France), Julien Le Tellier (Greece/France), Arnau Queralt (Spain), Grazia Maria Vagliasindi (Italy)

**Contributing Authors:**

Elen Lemaitre-Curri (France), Piero Lionello (Italy), Katarzyna Marini (France), Cyril Moulin (France)

*This chapter should be cited as: Lange MA, Llasat MC, Snoussi M, Graves A, Le Tellier J, Queralt A, Vagliasindi GM 2020 Introduction. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 41-58.*



## Table of contents

<b>1.1</b>	<b>The Mediterranean Basin: a region affected by accelerating climate and environmental change</b> .....	<b>44</b>
1.1.1	What do we know about the Mediterranean Basin and what do we need to know?.....	44
1.1.2	An integrated Mediterranean risk assessment for sustainability.....	46
<b>1.2</b>	<b>The Mediterranean Experts on Climate and environmental Change (MedECC)</b> .....	<b>47</b>
1.2.1	Goals, basic structure and first accomplishments of MedECC.....	47
1.2.2	Principles and processes of work in MedECC.....	47
<b>1.3</b>	<b>The First Mediterranean Assessment Report (MAR1): A synthesis of knowledge on risks from climate and environmental change in the Mediterranean Basin</b> .....	<b>48</b>
1.3.1	Report scope and objectives.....	48
	1.3.1.1 <i>Geographic scope</i> .....	48
	1.3.1.2 <i>Scenarios and reference periods</i> .....	50
	1.3.1.3 <i>Adapting to climate and environmental change</i> .....	51
	1.3.1.4 <i>Toward a systemic approach</i> .....	51
	1.3.1.5 <i>MedECC MAR1 as a policy support instrument</i> .....	52
1.3.2	Methodology.....	52
1.3.3	Communicating uncertainties and results.....	53
1.3.4	Report structure.....	54
	<b>References</b> .....	<b>55</b>
	<b>Information about authors</b> .....	<b>58</b>

## 1.1 The Mediterranean Basin: a region affected by accelerating climate and environmental change

### 1.1.1 What do we know about the Mediterranean Basin and what do we need to know?

The Mediterranean Basin is considered one of the cradles of civilization (Abulafia 2011; Sağlamer 2020). Its rich cultural and scientific heritage dates back to the Egyptian and Mesopotamian empires, followed by the remarkable accomplishments of Greek thinkers and philosophers and the no less ingenious scientists, engineers and architects of the Roman Empire. Throughout the history of the last several millennia, people, communities and societies have learned to adapt to and master environmental conditions that have often been adverse, and which frequently caused severe damage to human lives and environmental integrity. But it has not only been their ability to adapt to changes in natural conditions that has protected Mediterranean societies from severe impacts. As time has gone on, there has been a steadily growing body of knowledge and understanding of the underlying causes and the usual course of such changes, based on careful and long-term observations and analysis. It is no surprise then that we now know a lot about the changes and variations in environmental conditions that we observe today. But neither the level of observation nor the level of knowledge is the same, whether for the entire region, or for all environmental issues, or for their integration with the socio-economic system. Added to this lack of homogeneity is the complexity of the region, both in terms of climate and environment, and culture and socio-economics (Zamora Acosta and Maya Álvarez 1998; Woodward 2009; IEMed 2015).

During the second half of the 20th century and up to the present day, numerous scientific projects have studied the changing environment of the Mediterranean Basin and its adjacent regions (for a compilation of such research activities, see *Appendix A.4.1*). The capacity for systematic observation of the environment has increased substantially, followed by the development of conceptual and numerical models of the changing atmosphere and ocean, as well as of other environmental changes and their impacts. Through the work done by large numbers of scientists, we have gained not only a more complete understanding of past changes but have also learned to generate projections of future climate and environmental change with increasing confidence in the reliability

of the results. Projections of such changes into the future are key for the mitigation of risks to human livelihoods (Bolle 2003; Lionello 2012; Navarra and Tubiana 2013; Luterbacher et al. 2016).

As a first assessment, based on various sources of information, Cramer et al. (2018) have shown that the Mediterranean Basin is at risk of suffering from levels and rates of climate and environmental changes now and in the foreseeable future that exceed global mean values. This applies to changes in temperatures, precipitation and the frequency and magnitude of extreme weather events, but it also implies changes in land and sea use, pollution, air quality and other factors. Average annual mean temperatures in the Mediterranean Basin have risen by 1.5°C since pre-industrial times (1861-1890), approx. 0.4°C above the global average (*Chapter 2*), due to a combination of local drivers (e.g., land use changes) and changes on a global scale that affect the Mediterranean Basin through various modes of tele-connectivity (Lionello et al. 2014).

Although droughts have been a common experience for most of the history of the Mediterranean, the recently observed decline of seasonal precipitation for parts of the Basin (*Chapter 2*) presents new and significant challenges for Mediterranean communities. While most climate models show remarkable agreement in the expected decrease in Mediterranean rainfall, there are also some results indicating significant differences in magnitude and sign for these changes. Even if future global warming is limited to 2°C, as prescribed by the UNFCCC Paris Agreement, summer rainfall risks being reduced by 10-30% in some regions. Such a decrease will enhance existing water shortages and increase irrigation demand for agricultural productivity, particularly in countries on the southern rim of the Basin (Vautard et al. 2014). Climate change, population growth, increasing domestic needs and pressure from tourism, new industries and urban sprawl may see irrigation demand rise by 26-92% by the end of the 21st century (Fader et al. 2016). Although frequently addressed by policymakers, irrigation and food security remain a sensitive issue that needs additional research (Grafton et al. 2018; WWAP 2019). Research into the extent to which local landraces can cope with projected climate changes without significant loss of productivity should be a priority (FAO 2015). More aridity

exacerbates ongoing desertification, with strong socio-economic impacts on farmers (UNEP/MAP 2016). The interlinkages between resources have been addressed through the concept of the Water-Energy-Food Nexus (e.g., Hoff 2011; Kennou et al. 2019), but they still present unresolved challenges. Since the impacts of climate change are likely to include the degradation of agricultural water resources and loss of fertile soils, enhanced efforts to adapt agricultural and other land systems to climate change are necessary to ensure food security and rural livelihoods (UNEP/MAP 2016). Despite this reduction, extreme precipitation events are expected to intensify in a large part of the region. Therefore, flood socio-economic and environmental impacts should be added to those produced by water scarcity (Tramblay and Somot 2018).

Primarily caused by global processes, including thermal expansion of sea water and accelerated melting of ice sheets in Greenland and Antarctica, sea level is projected to increase more strongly than previously estimated (DeConto and Pollard 2016; IPCC 2019). This will have repercussions for the Mediterranean Sea, as well. For the recent past, in which sea level has been monitored by satellite altimetry (1993-2018), Mediterranean sea level has increased up to  $2.8 \pm 0.1 \text{ mm yr}^{-1}$ , which is consistent with global sea level trend ( $3.1 \pm 0.4 \text{ mm yr}^{-1}$ ) (Cazenave and WCRP Global Sea Level Budget Group 2018). At a sub-regional level, by the end of the 21st century (2080-2099) the projected rise in the average sea level of the Mediterranean Basin with respect to the present climate (1980-1999), is estimated to be 37 cm, 45 cm, 62 cm and 90 cm under RCP2.6, RCP4.5, RCP8.5 and high-end greenhouse gas emission scenarios, respectively (Somot et al. 2016; Jordà et al. 2020).

Coastal regions around the Mediterranean Basin are densely populated. Due to the near absence of tides in much of the Mediterranean, many cities and coastal infrastructures are built close to current mean sea levels. They are therefore particularly vulnerable to future sea level rise. Paired with an increasing frequency of storm surges, these cities and infrastructures face enhanced risks of flooding. Sea level rise will lead to loss of arable lowlands, notably in intensively used river deltas such as those of the Nile, the Po and elsewhere, with adverse consequences for agricultural activities and food security. A related problem is the intrusion of seawater into coastal aquifers (seawater intrusion), which renders these aquifers unsuitable for human consumption and for most agricultural purposes (Hegazi et al. 2005). Enhanced uptake of atmospheric CO<sub>2</sub> has led to

a significant increase in seawater acidity of the Mediterranean, which is set to continue (Tsimplis et al. 2013; Palmiéri et al. 2015). Acidification has serious consequences for organisms that produce carbonate shells and skeletons and for marine ecosystems throughout the Mediterranean Basin (Gattuso et al. 2015; Palmiéri et al. 2015).

Beyond climate change, the Mediterranean Basin also experiences environmental challenges due to changing land and sea use, agricultural intensification and urban sprawl, increasing pollution and declining biodiversity. Due to drought, land use change, and high temperatures, the area affected by forest fires could increase by approx. 40% up to ~100% relative to recent levels, generally proportional to warming (Turco et al. 2018). Likewise, warming in combination with overfishing risk causing the local extinction of more than 20% of exploited fish and marine invertebrates by 2050 (Jones and Cheung 2015).

These examples show that climate change, in combination with other challenges, will likely not only affect ecosystems on land and in the ocean, but also create risks for the services they provide and therefore ultimately the ecological basis for the well-being of people in the Mediterranean Basin. Combined with current changes in lifestyle, e.g., the switch to a more urbanized life-style and a more processed animal-based diet, in particular southern Mediterranean countries in particular are at risk of increasing their dependence on food imports and trade from elsewhere (CIHEAM 2014; UNEP/MAP 2016). Landraces are likely to be lost as farmers replace them with other landraces, or improved varieties, that are better adapted to the new conditions (FAO 2015). Some scholars argue that this trend could be mitigated by boosting a return to the traditional Mediterranean diet, with significant health benefits for all Mediterranean people including their visitors (Serra-Majem et al. 2011).

Climate and environmental changes and their impacts imply risks for human security in the Mediterranean region (Karmaoui 2016; Rigaud et al. 2018). These changes are added to escalating conflict and insecurity in some African and Eastern countries that are leading thousands of people to flee, taking their chances on unseaworthy boats across the Mediterranean. Public health is already affected by multiple facets of climate and environmental change, including enhanced and more frequent (urban) heat waves, increasing air pollution (higher risk of cardiovascular or respiratory diseases), and increased spread

of disease vectors (West Nile virus, dengue, chikungunya) (Kuglitsch et al. 2010; Negev et al. 2015; Orru et al. 2017). Environmental change is also increasingly recognized as a relevant factor for socio-economic risks (e.g., famines) in situations of instability and conflict. Synergistic effects between societal, economic and environmental factors should also be considered, as well as the relevance of globally connected socio-economic structures (Le Roy Ladurie 2004, 2006). For instance, droughts, floods or other extreme events in agricultural regions elsewhere may lead to market disturbances and may affect prices, trade, production and security in the Mediterranean too.

### 1.1.2 An integrated Mediterranean risk assessment for sustainability

Considering this incomplete compilation of examples for current understanding of global and environmental changes in the Mediterranean Basin, one might get the impression that much is known already. However, impressive as it may seem, most environmental research conducted so far in the region is primarily driven by disciplinary and sectoral investigations. A more comprehensive, systemic and holistic approach to interrelated processes and components would likely make useful contributions to environmental decision-making in the Mediterranean Basin. So far, an adequate and comprehensive assessment of risks posed by climate and environmental changes in the Mediterranean Basin is lacking (Cramer et al. 2018).

The absence of integrated studies comes in addition to the painful lack of monitoring and risk analysis capacity in southern and eastern countries of the Mediterranean Basin. There are few, but strong indications that these countries potentially face larger risks from climate and environmental changes, compared to northern countries, while commanding significantly scarcer financial resources to effectively adapt to their impacts (IPCC 2014).

This report cannot replace a full and integrated research-based risk assessment in the terms outlined above. It rather aims at providing an assessment of current knowledge such as it emerges out of the existing body of research. While the assessment is designed and carried out as a regional study, its results may also provide useful conclusions on a global scale. Global environmental change inevitably has distinct regional manifestations. Similarly, processes and changes on the regional level will have consequences for global processes. The MedECC network sees this re-

gional assessment as a possible “bridge” between the global and the national to local scale, essential for the advancement of mitigation and adaptation strategies. Such a bridge appears to have particular potential in the Mediterranean context, since this region is at a crossroads between Africa, Europe and (Western) Asia. The importance of the regional dimension has been emphasized in the context of the implementation of the 2030 Agenda and the Sustainable Development Goals (SDGs). The 2030 Agenda recognizes regional dimension and regional governance as playing a crucial role in translating sustainable development policies into concrete actions at the national level (UN 2015; UN-SDSN 2018).

Mitigation of the processes underlying climate and environmental change and adaptation to their unavoidable impacts represent a priority for public and private decision makers concerned with the future of communities and environmental integrity in the Mediterranean Basin. Effective mitigation and adaptation require investigations that go beyond current knowledge. Looking at the current state of the Mediterranean, we note the following challenges and needs:

- A substantial gap between knowledge and understanding of climate and environmental change between the northern-rim countries and most of the southern- and eastern-rim countries of the Mediterranean Basin.
- A disparity of observational data and monitoring systems between the North versus the South and the East, which calls for an intensification of observations and the creation of observational networks, notably in the MENA region.
- Research on short- and mid-term weather and climate predictions, including seasonal forecasting to better manage water resources and agriculture, is being carried out in some Mediterranean countries. The results of such studies have the potential to be applied more broadly.
- Despite some initiatives (e.g., the Mediterranean Climate Outlook Forum - MedCOF), the level of climate services offered by the scientific communities in most Mediterranean countries remains insufficient. Such services can be decisive in providing vital information on short-term to intermediate climate trends to planners and decision makers involved in agricultural and water policies.
- The implementation of more advanced early warning systems may enable better preparation for extreme events and other climate-related risks that usually affect the Mediterranean region. Such systems need to be based on

and accompanied by an adequate societal and individual risk awareness (see the recommendations of UNISDR 2015).

- Large-scale programs are needed (or should be strengthened) in eastern and southern Mediterranean countries to address pressing multi-factorial challenges such as land degradation and desertification, ultimately focusing on increased resilience to change as well as the reinforcement of ecological transitions to more sustainable resource use.
- In order to address these issues, a comprehensive scientific synthesis and assessment report is required. This report should include recent trends, likely future developments, and consequences of climate and environmental changes on natural systems, the economy, and human well-being in the Mediterranean Basin.

The overarching goal of this report is to provide such a synthesis, based on existing knowledge in the scientific literature. The work has been carried out by the Mediterranean Experts on Climate and environmental Change (MedECC) aiming for a comprehensive synthesis of current scientific knowledge that covers all relevant disciplines, sectors and sub-regions. The assessment considers three major interconnected domains, namely resources (water, food & energy), ecosystems (marine, coastal and land), and society (development, health and security). Although the target audience of this assessment is decision makers and policymakers, anyone interested in the Mediterranean can benefit from it. After completion of this first assessment, there is a desire to develop a platform for constructive science-society-policy dialogue.

## 1.2 The Mediterranean Experts on Climate and environmental Change (MedECC)

### 1.2.1 Goals, basic structure and first accomplishments of MedECC

MedECC's main, overarching goal is to provide a state-of-the-art risk assessment on climate and environmental changes and their impacts across the Mediterranean, based on existing scientific knowledge, with the following specific objectives:

- To activate and engage the scientific community working on environmental and climate changes in the Mediterranean Basin;
- To provide comprehensive updated and consolidated scientific knowledge on these changes and make it accessible to policymakers, key stakeholders and the general public in a process which facilitates ownership of scientific knowledge;
- To identify possible gaps in the current understanding of environmental and climate changes and their impacts in the Mediterranean;
- To help build capacity of scientists from southern and eastern Mediterranean countries to international levels and standards by encouraging training, research and development efforts in these countries in the context of the Paris Agreement;
- To bridge the gap between research and decision-making, contributing to the improvement of policies at all levels;

- To contribute to future reports of the Intergovernmental Panel on Climate Change (IPCC), the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) or similar assessments in the Mediterranean Basin. MedECC has an important role to play in the work of the IPCC, as it is contributing to the Sixth Assessment Report (AR6) with a cross-chapter paper dedicated for the first time to the Mediterranean.

MedECC's work therefore focuses on two complementary directions:

- Publishing a scientifically robust assessment and synthesis of environmental and climate changes and their impacts in the Mediterranean Basin, based on currently available research;
- Building a science-policy interface on these changes and their impacts in the Mediterranean and thereby providing a scientifically sound basis for decision-making.

### 1.2.2 Principles and processes of work in MedECC

Since the founding of MedECC, two scientists at the French National Centre for Scientific Research (CNRS) coordinate the development and work of the network. Central to the governance of MedECC is its Steering Committee (SC, *Fig. 1.1*) which





Figure 1.1 | Structure and functioning of MedECC.

met for the first time in April 2016 in Barcelona, Spain. The SC is currently composed of 20 members from 11 countries (Appendix A.4.1), including 16 scientists in a personal capacity (representing environmental sciences, political sciences and economics) and 4 representatives of policymaking bodies: the United Nations Environment Programme/Mediterranean Action Plan – Barcelona Convention Secretariat (UNEP/MAP) and its Plan Bleu Regional Activity Centre, the Secretariat of the Union for the Mediterranean (UfM), and the Advisory Council for the Sustainable Development of the Government of Catalonia (CADS). More details on the work and accomplishments of the SC can be found in Appendix A.4.1.2.

MedECC work is carried out by an open and inclusive network of scientific experts to support decision-making through accurate and accessible information on current and future environmental and climate changes and their impacts in the Mediterranean Basin. The network currently comprises more than 600 scientists from 35 countries who contribute in a personal capacity on a voluntary basis. Members of the network do not primarily represent an institution or country.



(country code: Appendix D)

Figure 1.2 | Distribution of MAR1 Coordinating Lead Authors and Lead Authors by country.

The added value of the work of MedECC lies in the large geographical and thematic scope addressed (see below) and the comprehensive involvement of more than 86 Coordinating Lead Authors (CLAs) and Lead Authors (LAs) from 21 countries participating in the drafting of the report (Fig. 1.2). 27 (31%) are based in southern and eastern Mediterranean countries and 31 (36%) are women. In total, 190 authors from 25 countries contributed to the report (CLAs, LAs and Contributing Authors - CAs). Appendix A.4 provides more information about the steps of MAR1 as well as the duties of the different players involved in report preparation. By providing information on the assessment/synthesis process underlying MAR1, we aim to address both institutional stakeholders/organizations as well as individuals interested in the organization of MedECC and its work.

### 1.3 The First Mediterranean Assessment Report (MAR1): A synthesis of knowledge on risks from climate and environmental change in the Mediterranean Basin

#### 1.3.1 Report scope and objectives

##### 1.3.1.1 Geographic scope

Located at the crossroads of three continents, the Mediterranean region is unique in its historical and geographical specificities, as well as its natural and

cultural heritage. While the Mediterranean Sea is a well-defined water body, limited by the Strait of Gibraltar, the Dardanelles Strait and the Suez Canal (Fig. 1.3), various definitions are used for the land boundaries of the Mediterranean region. In political terms, the United Nations Environment Program Mediterranean Action Plan (UNEP/MAP<sup>1</sup>)

<sup>1</sup> <https://www.unep.org/uneppmap/>





**Figure 1.3 | Geography, physiography and landscapes of the Mediterranean Basin**  
(Source: GRID-Arendal)<sup>2</sup>

covers 21 riparian countries, which are Contracting Parties (CPs) to the Barcelona Convention (excluding Jordan and Portugal, which are not riparian countries *sensu stricto*, and Gibraltar/UK and Palestine, which are not currently CPs but have observer status). The population of this so-defined Mediterranean region is about 480 million inhabitants (EEA 2020). From a geopolitical point of view, the Union for the Mediterranean (UfM<sup>3</sup>), created in 2008, embodies a much wider spatial scope of 43 countries (all the countries of the European Union and 15 countries in the southern and eastern Mediterranean).

For much of the physical assessment by MedECC, we adopt a simple regular latitude-longitude box (29°N to 47.5°N and 10°W to 39°E, *Fig. 2.1*), which includes some regions with non-Mediterranean climates, such as the Alps, the Eastern Balkans or part of the Sahara. This definition of the Mediterranean region is similar to the MED zone adopted in IPCC-AR4 (IPCC 2007), and slightly larger than in IPCC-AR5 (IPCC 2013) and the ongoing AR6 assessment.

The Mediterranean Sea is a relatively small, semi-enclosed sea with limited exchange to the global

ocean through the Strait of Gibraltar, located at the western end of the Mediterranean, the linkup between the Mediterranean and the Black Sea through the Bosphorus and Dardanelles and the connection to the Red Sea through the Suez Canal in the South-East (*Fig. 1.3*). The Mediterranean Sea can be divided into two sub-basins: the Western and the Eastern Mediterranean, which are in turn made up of a series of various small basins (Amblàs et al. 2004). The Mediterranean region includes 75 coastal watersheds and 224 coastal administrative regions, with a total of 46,000 km of coastline (UNEP/MAP 2016).

The topographic, geographic and socio-economic structures of Mediterranean landscapes are heterogeneous. Their current shapes are the result of centuries of interactions between natural forcings and diverse human activities, both past and present.

The coastal area comprises a large set of ecosystems that deliver valuable services to people, including lagoons, estuaries, deltas, coastal plains, wetlands, rocky shores and nearshore coastal areas, seagrass meadows, coral communities, frontal systems and upwellings, seamounts, and pelagic

<sup>2</sup> <https://www.grida.no/resources/5931>

<sup>3</sup> <https://ufmsecretariat.org/>

systems. Most of these systems are very sensitive to human and climate forcing. Wetlands represent 1.7-2.4% of the total area of Mediterranean countries (Tour du Valat 2012). The most extensive coastal wetlands are found in estuaries like that of the Po (Italy), Nile (Egypt), Rhône (France) and Ebro (Spain) rivers.

Other elements of Mediterranean heritage are the nearly 15,000 islands and islets dotted throughout the basin, some of which are particularly vulnerable. The largest islands are Sicily, Sardinia, Corsica, Cyprus, and Crete, and the major island groups include the Balearics off the coast of Spain and the Ionian, Cyclades, and Dodecanese islands of Greece. In total, the islands' coastlines comprise around 19,000 km, or more than 41% of the Mediterranean coastline (Emmanouilidou 2015). In terms of land surface, islands represent only 4% of the land area of the whole Mediterranean Sea Basin (Kolodny 1974). The small islands (less than 10 km<sup>2</sup>), are particularly valuable in terms of biodiversity. Since many of them are uninhabited or weakly impacted by human activities, they constitute valuable "life laboratories" for assessing the sole impacts of climate change. Since 2005, the Mediterranean Small Islands Initiative (PIM<sup>4</sup>) has been working to improve knowledge and management of these territories, as well as raising awareness of the importance of local populations for the preservation of these fragile territories.

The Mediterranean is also complex in terms of its socio-political settings. For millennia, the Mediterranean has been a unique geographical space but – except for the Roman Empire – it has been a politically, economically and culturally divided region. While its geographical scope has remained unchanged over the past 5,000 years of human history, the modes of government in countries and regions have been in permanent flux (Brauch 2010).

Approximately one-third of the Mediterranean population is concentrated along its coastal regions. Meanwhile, about 250 million people reside in coastal hydrological basins. In the southern region of the Mediterranean, around 120 million inhabitants are concentrated in coastal hydrological basins, where environmental pressures have increased (EEA 2020). In addition, around 360 million international tourists visited the Mediterranean countries in 2017 (UNWTO 2019). Approximately half of these arrivals - 170 million - are in Mediterranean coastal areas,

exacerbating the human pressures in coastal zones, and generating a 40% increase in marine litter, particularly during the summer season (Galgani et al. 2014).

The variety of cultures, policy and governance approaches and the diversity of social systems have led to very different levels of socio-economic development and ecological footprints of the Mediterranean states between the north and the south (Raleigh et al. 2008; IPCC 2014; Rigaud et al. 2018; GRID 2019). Per capita income levels are three to five times higher in southern European countries (France, Italy and Spain, in particular), compared to countries on the southern and eastern shores of the Mediterranean Sea. While important progress has been made in the South and the East over the last twenty years, instability and significant inequalities persist. This north-south dichotomy is at the heart of the climate change issue, as it exacerbates imbalances by having greater impacts on lesser-developed countries, which have limited capacities to deal with the impacts and possible adaptation measures.

Due to the complexities and heterogeneities of topographic, geographic and socio-economic structures in the Mediterranean Basin, regional, sub-regional, national or local scales are considered where appropriate and where data and information are available. For instance, the spatial separation of drivers has been considered at the level of ecosystems (open ocean, deltas, river basins, wetlands, drylands, etc.). Impacts of environmental change are sometimes quite localized but concern a large number of domains. They are complex to understand and require studies and simulations at reduced spatial scales, associated with high degrees of uncertainty. Best practices with regard to mitigation and adaptation measures are usually reported at the local level, as adaptation measures are generally implemented at a territorial scale where end-users and decision makers are more engaged.

### 1.3.1.2 Scenarios and reference periods

Different periods and time windows have been selected by climatologists and adopted by IPCC to monitor and record changes in climate conditions throughout Earth history along the different reports. In order to quantify human impacts with respect to an "unperturbed" reference state (ideally the climate just before human activities started to

<sup>4</sup> <http://initiative-pim.org/>

demonstrably change the environment at global scale) a practical approach to identifying impacts and their characteristics is needed (*Chapter 2*).

Instrumental observations of temperature are available mainly since the second half of the 18th century but only in some European countries. In the IPCC AR5 report, the 1850-1900 period is considered as the best approximation for an unperturbed state, but, it is not named pre-industrial. It has nonetheless been kept as the reference for the pre-industrial period in the IPCC Special Report on the Impacts of Global Warming at 1.5°C (SR15) (IPCC 2018) (*Section 2.2.1*).

Climate modelling and climate projections adopted the practice of describing future changes with respect to a recent baseline (during which validation of models is supported by a large amount of instrumental observations). Because of the long residence time of carbon dioxide in the atmosphere, the human influences on the current trajectory of a changing climate appear to be irreversible for decades to centuries, even if significant mitigation measures are implemented immediately (Millar et al. 2006). Thus, given the dynamics of the natural climate system and the superimposed changes humans are causing, the 21st century is an important transitional time for undertaking both mitigation and adaptation actions. In order to be able to propose future climate projections considering various possible socio-economic trajectories and climate policy pathways, we follow the IPCC scenario approach. Although results based on multiple IPCC scenarios are reported in MAR1, we mostly focus on two options which encompass the range of IPCC-AR5, CMIP5 and CORDEX simulations: the “business as usual” scenario (RCP8.5, for an explanation of the RCPs see *Box 2.2*) and the optimistic scenario closest to the UNFCCC Paris Agreement target (RCP2.6). These scenarios were also chosen due to model projection availability constraints at the regional scale. Where more recent studies are not available, the assessment also considers studies based on the older IPCC SRES approach (Nakićenović 2000) (*Section 2.2.1*).

### 1.3.1.3 Adapting to climate and environmental change

Climate change adaptation is a necessity, possibly an opportunity, but many definitions exist and the nature and effectiveness of adaptive responses is critically influenced by the framing of adaptation responses (Wise et al. 2014). The IPCC AR5 has framed adaptation as a “process of adjustment

to actual or expected climate and its effects in order to either lessen or avoid harm or exploit beneficial opportunities” and also emphasizes that there is “increasing recognition that an adequate adaptive response will mean acting in the face of continuing uncertainty about the extent of climate change and the nature of its impacts” (IPCC 2014). Other authors emphasize how the increasing climate pressure recalls the need to shift rapidly from incremental to transformative adaptation (Rickards and Howden 2012; Vermeulen et al. 2013). Adaptation can be understood as a set of actions which “adjust to” a new situation (e.g., change date of seeding), which recalls a “technical fix” approach, or as a “structural coupling dynamic process”, where social learning is the main process informing decisions at any critical point of an ongoing adaptive pathway (Collins and Ison 2009). In the latter framing, the assumption is that no single group has clear access to understanding the issues and their resolutions, hence the difficulties in securing the active and broad-based engagement of stakeholders, and the facilitated spaces for “learning to adapt” become crucial. Adaptation can be operationalized as part of pathways of change and response, which implies the reorganization of institutional structures that are likely to lead to more sustainable trajectories (Rickards and Howden 2012).

The design of adaptation pathways can emerge from the integration of capacity-based (i.e., bottom-up) approaches with impact-based (i.e., top-down) approaches (Vermeulen et al. 2013). The same authors show that when the signal-to-noise ratio of the changing climate is low, i.e., the noise associated with natural weather variability is higher than the changing climate signals, the capacity to respond can basically rely on “no-regret” or “win-win” approaches resulting in incremental adaptation practices, such as short-term investments on higher water storage capacity or use of water-saving technologies, which can also respond to weaknesses emerging from the usual climate uncertainty. When climate shifts are stronger, impact-based decisions can more effectively inform systemic or transformative adaptation pathways, even when uncertainty of future predictions is relatively high (Vermeulen et al. 2013).

### 1.3.1.4 Toward a systemic approach

As previously emphasized in this chapter, a large amount of scientific material (data, publications, reports, etc.) is available for various Mediterranean areas (from localities to countries or marine sub-

basins), for different periods of various durations, for different compartments (atmosphere, soil, continental or sea water, etc.) and on different topics (chemical composition, physical characteristics, ecosystems, human activities, etc.). New data from satellite providers are essential for those regions with poor instrumental coverage, but also to provide a complete overview of the Mediterranean Region (e.g., GEO-Cradle, geocradle.eu, the COPERNICUS and Digital Africa systems, or the Integrated Geospatial information Frameworks, IGIF). All of these elements contribute to the overall understanding of the impacts of climate and environmental changes in the Mediterranean. The fact that advanced research is so complex that it requires very focused and specialized studies makes it difficult to achieve the objective of having an overall integrated understanding of the functioning of the Mediterranean environment, of the modifications due to local human activities and of the evolutions related to global changes. A holistic approach is however of major importance in order to better understand the interactions between these multiple elements, and then provide a robust scientific basis to develop and implement sustainable and effective policies.

This systemic approach has been successfully adopted for decades by scientists contributing to the IPCC by developing more and more refined global climate models. Nevertheless, these models are not designed to address all processes that determine the evolution of the Mediterranean environment and socio-ecosystems under both climate change and other environmental processes caused by human activity.

The development of an integrated Mediterranean model with the spatial and temporal resolution suitable for comprehensively resolving the relevant processes has to be a long-term target for the scientific community. Such an approach could provide answers to the concerns of policymakers related to sustainable development strategies in the Mediterranean region. At present, MedECC aims to initiate the process by providing a first synthesis of the state of knowledge in the various scientific fields that need to be considered and brought together in order to move one step forward in this direction.

### **1.3.1.5 MedECC MAR1 as a policy support instrument**

Given the stated goals and objectives of MedECC, MAR1 is intended to support policies in deriving mitigation and adaptation strategies, particularly in the context of Mediterranean cooperation under

policies of the European Union, Arab League, EU-Africa and EU-Asia cooperation, North African Unions, Maghreb and Mashriq. To provide adequate support, MAR1 has been inspired by other science-policy interfaces such as the IPCC and IPBES, aiming to provide an unbiased, scientific view of climate and environmental change, its various, multi-sectoral impacts and the risks they imply for society.

**The MedECC MAR1, by summarizing existing findings and results, aims to highlight their policy relevance without being policy-prescriptive.**

The MedECC MAR1 is designed to address the needs of multiple actors involved in providing a response to climate and environmental changes and risks in the Mediterranean region. The primary target users of the report are governments and policymakers at all levels, the UfM and the UNEP/MAP at the regional level, and more broadly, the comprehensive system of intergovernmental processes pursuing different aims, including the three "Rio Conventions" (UNFCCC, CBD, UNCCD) and the Ramsar Convention (and their financial instruments), as well as IPCC, IPBES and the Commission on Generic Resources for Food and Agriculture (CGRFA). Other important end-users include the scientific community, major economic decision makers and the private sector, the education sector, civil society and non-governmental organizations (CSOs and NGOs). MedECC aims to build close relationships with various media in order to help guide the interpretation of its report and to ensure that the public is provided with objective and unbiased information about MAR1.

MAR1 contributes to meeting the need for an advancement and implementation of regulatory instruments aiming to reduce greenhouse gas emissions and emissions of pollutants to mitigate climate and environmental changes in the Mediterranean Basin. The scoping and drafting of the report involved ample consultation of completed or ongoing assessments of comparable nature like the IPCC and IPBES assessment reports and builds on the existing relations between MedECC and other groups. Findings of MAR1 will be directly employed for the drafting of a Mediterranean Cross-Chapter Paper in the forthcoming IPCC Sixth Assessment Report (*Appendix A.1*).

### **1.3.2 Methodology**

The drafting of MAR1 entailed a collective and iterative review, synthesis, analysis and judgment

of available scientific knowledge. The entire assessment is supported by scientific references; no additional research has been undertaken by MedECC. In some cases, a new analysis of data was conducted using existing models to address specific questions and to identify knowledge gaps to be addressed by other initiatives and research programs.

The report is primarily based on peer-reviewed literature (in English or other languages) but selected non-peer reviewed literature was also considered (such as institutional or government reports, national statistics, etc.) in which case the authors carefully checked the quality of the references included to justify their inclusion. A scientific literature database was maintained by the MedECC Secretariat and was made available to all report authors.

Drafts of the report have been subject to a dual scientific review allowing suggestions and amendments by scientific experts. A first internal review of the First Order Draft (FOD) involving the SC and the authors of the report was carried out, and the Second Order Draft (SOD) was submitted to external scientific reviewers.

The main body of MAR1 is accompanied by a Summary for Policymakers (SPM), which undergoes an approval procedure organized with the UNEP/MAP – Barcelona Convention Secretariat and its Plan Bleu Regional Activity Centre, through their Focal Points and/or the Members of the Mediterranean Commission on Sustainable Development (MCSD), as well as with the UfM Member State representatives within the regional Climate Change Expert Group (CCEG).

### 1.3.3 Communicating uncertainties and results

Communication of the findings of MAR1 aims to also adequately communicate scientific uncertainties and confidence in the material used. For this purpose, three different target groups are differentiated:

- the scientific community; all of the main conclusions will be supported by robust literature and/or evidence following the AR5 IPCC (IPCC 2013) criteria to communicate the uncertainty of findings;
- policymakers and stakeholders; conclusions for them will be summarized in the MAR1 Summary for Policymakers; considering the strong relationships with key institutions such


as UNEP/MAP and UfM, main questions and key messages will be discussed with them prior to final publication, in order to take their views and the advice of the MedECC SC into account;

- the public at large, mainly, but not exclusively, those living in Mediterranean countries; the main challenges and opportunities will be communicated to them; in so doing, we will progress from MAR1's strictly "informative" role towards a more "participatory" and "responsible" one, in order to further understanding and acceptance of measures aimed to cope better with climate and environmental changes in the Mediterranean region.

The approach of the MAR1 report meets the call made by the United Nations program on the Sustainable Development Goals as well as the concessions proposed after the COP 21 under the UNFCCC (Paris Agreement), in particular SDG 17 "Strengthen the means of implementation and revitalize the global partnership for sustainable development" (Partnerships for the Goals).

Following the AR5 IPCC (IPCC 2013), the metrics for communicating the degree of certainty in key findings (notably on climate drivers) will be the following:

- **Confidence:** confidence in the validity of a finding, will be based on the type, amount, quality, and consistency of evidence (e.g., data, mechanistic understanding, theory, models, expert judgment) and the degree of agreement; confidence is expressed qualitatively and its level will be based on the evidence (robust, medium and limited) and the agreement (high, medium and low). A combination of different methods, e.g., observations and modelling, is important



**Figure 1.4 | The basis for the confidence level is given as a combination of evidence (limited, medium, robust) and agreement (Low, medium and high) (Mastrandrea et al. 2011).**

for evaluating the confidence level. There is flexibility in this relationship; for a given evidence and agreement statement, different confidence levels could be assigned, but increasing levels of evidence and degrees of agreement are correlated with increasing confidence. *Fig. 1.4* shows how the combined evidence and agreement results in five levels for the confidence level used in this assessment. Confidence should not be interpreted probabilistically, and is distinct from “statistical confidence”.

Term	Likelihood of the outcome
Virtually certain	>99% probability
Extremely likely	>95% probability
Very likely	>90% probability
Likely	>66% probability
More likely than not	>50% probability
About as likely as not	33-66% probability
Unlikely	<33% probability
Extremely unlikely	<5% probability
Exceptionally unlikely	<1% probability

**Table 1.1 | Likelihood terms associated with outcomes used in MAR1** (from IPCC 2013).

- Uncertainty:** quantified measures of uncertainty in a finding will be expressed probabilistically, i.e., based on a statistical analysis of observations or model results, or on expert judgement. The qualifier “likelihood” provides calibrated language for describing quantified uncertainty. It can be used to express a probabilistic estimate of the occurrence of a single event or of an outcome, for example, a change in a given climate parameter, an observed trend, or a projected change lying in a given range. Statements made using the likelihood scale may be based on statistical or modelling analyses, elicitation of expert views, or other quantitative analyses. Where sufficient information is available, it is preferable to avoid the likelihood qualifier in favor of the full probability distribution or the appropriate probability range. *Table 1.1* shows the list of “likelihood” qualifiers to be used in this report.

### 1.3.4 Report structure

The outline for the MAR1 report was approved by the MedECC Scientific Committee during its meeting on May 24, 2018 in Marseille, France. MAR1 consists of a Summary for Policymakers, six main chapters and several appendices, as follows:

- Summary for Policymakers, including an Executive Summary.
- This chapter, Chapter 1, “**Introduction**”, frames the motivation and main components of the MedECC Assessment.
- Chapter 2, “**Drivers of change**”, focuses on the physical, bio-chemical and human drivers of climate and environmental changes, distinguishing between climate, pollution, land/sea use and management, and invasive species. Based on these drivers, the analyses in Chapters 3-5 all consider past trends and current situation, projections, vulnerabilities and risks, adaptation, knowledge gaps and research needs.
- Chapter 3, “**Resources**”, assesses the state of knowledge for major resource challenges: water, food and energy in three sub-chapters describing each of these resources.
- Chapter 4, “**Ecosystems**”, assesses the state of knowledge for marine, coastal and terrestrial ecosystems.
- Chapter 5, “**Society**”, addresses major issues of development, health and human security under climate and environmental change.
- Chapter 6, “**Managing future risks and building socio-ecological resilience**”, discusses options for more sustainable policies given the risks identified in *Chapters 3-5*. It describes the future risks associated with climate change in Mediterranean countries, and critically reviews a range of examples of adaptation and mitigation, promoting their synergies, as well as cooperation and networking among Mediterranean countries for building resilience.
- Supplementary information is given by the appendices, which include the information on MedECC partners and related research activities, the institutional context of MedECC, the main steps in MAR1 preparation, maps of projected temperature and precipitation changes for the Mediterranean Basin, the lists of acronyms and country codes.

## References

- Abulafia D 2011 *The Great Sea: A Human History of the Mediterranean*, ed. Penguin.
- Amblàs D, Canals M, Lastras G, Berné S, Loubrieu B 2004 Imaging the seascapes of the Mediterranean. *Oceanography* 17, 144–155. doi: [10.5670/oceanog.2004.11](https://doi.org/10.5670/oceanog.2004.11)
- Bolle H-J 2003 *Mediterranean Climate - Variability and Trends*. Berlin Heidelberg: Springer-Verlag. doi: [10.1007/978-3-642-55657-9](https://doi.org/10.1007/978-3-642-55657-9)
- Brauch HG 2010 Climate change and Mediterranean security. International, national, environmental and human security impacts for the Euro-Mediterranean region during the 21<sup>st</sup> century. Proposals and perspectives.
- Cazenave A, WCRP Global Sea Level Budget Group 2018 Global sea-level budget 1993–present. *Earth Syst. Sci. Data* 10, 1551–1590. doi: [10.5194/essd-10-1551-2018](https://doi.org/10.5194/essd-10-1551-2018)
- CIHEAM 2014 *Mediterra 2014: Logistics and Agro-Food Trade: A Challenge for the Mediterranean*. Paris: Les Presses de Sciences Po.
- Collins K, Ison R 2009 Jumping off Arnstein's ladder: social learning as a new policy paradigm for climate change adaptation. *Environ. Policy Gov.* 19, 358–373. doi: [10.1002/eet.523](https://doi.org/10.1002/eet.523)
- Cramer W, Guiot J, Fader M, Garrabou J, Gattuso J-P et al. 2018 Climate change and interconnected risks to sustainable development in the Mediterranean. *Nat. Clim. Chang.* 8, 972–980. doi: [10.1038/s41558-018-0299-2](https://doi.org/10.1038/s41558-018-0299-2)
- DeConto RM, Pollard D 2016 Contribution of Antarctica to past and future sea-level rise. *Nature* 531, 591–597. doi: [10.1038/nature17145](https://doi.org/10.1038/nature17145)
- EEA 2020 Mediterranean Sea region briefing - The European environment - state and outlook 2015.
- Emmanouilidou P 2015 Le statut juridique des îles de la Méditerranée: un droit fragmenté. *HAL* 01385765.
- Fader M, Shi S, Von Bloh W, Bondeau A, Cramer W 2016 Mediterranean irrigation under climate change: More efficient irrigation needed to compensate for increases in irrigation water requirements. *Hydrol. Earth Syst. Sci.* 20, 953–973. doi: [10.5194/hess-20-953-2016](https://doi.org/10.5194/hess-20-953-2016)
- FAO 2015 Coping with climate change – the roles of genetic resources for food and agriculture. Rome.
- Galgani F, Claro F, Depledge M, Fossi C 2014 Monitoring the impact of litter in large vertebrates in the Mediterranean Sea within the European Marine Strategy Framework Directive (MSFD): Constraints, specificities and recommendations. *Mar. Environ. Res.* 100, 3–9. doi: [10.1016/j.marenvres.2014.02.003](https://doi.org/10.1016/j.marenvres.2014.02.003)
- Gattuso J-P, Magnan A, Billé R, Cheung WWL, Howes EL et al. 2015 Contrasting futures for ocean and society from different anthropogenic CO<sub>2</sub> emissions scenarios. *Science* (80-. ). 349. doi: [10.1126/science.aac4722](https://doi.org/10.1126/science.aac4722)
- Grafton RQ, Williams J, Perry CJ, Molle F, Ringler C et al. 2018 The paradox of irrigation efficiency. *Science* (80-. ). 361, 748–750. doi: [10.1126/science.aat9314](https://doi.org/10.1126/science.aat9314)
- GRID 2019 Global report on internal displacement.
- Hegazi AM, Afifi MY, Elwan AA, Shorbagy MAE, El-Deimerdashe S 2005 Egyptian National Action Program to Combat Desertification. <http://extwpr-legs1.fao.org/docs/pdf/egy165179.pdf>
- Hoff H 2011 Understanding the nexus. Background paper for the Bonn2011 Conference: The Water, Energy and Food Security Nexus. Stockholm.
- IEMed 2015 Macroeconomic and monetary policies in the Mediterranean: management in a context of uncertainty.
- IPCC 2007 *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* , eds. Solomon S, Qin D, Manning M, Chen Z, Marquis M et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- IPCC 2013 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* , eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. Cambridge, United Kingdom and New York, NY, USA.
- IPCC 2014 Summary for Policymakers, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 1–32.
- IPCC 2018 *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change.* eds. Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J et al. In press.
- IPCC 2019 *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate.* , eds. Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M et al.
- Jones MC, Cheung WWL 2015 Multi-model ensemble projections of climate change effects on global marine biodiversity. *ICES J. Mar. Sci.* 72, 741–752. doi: [10.1093/icesjms/fsu172](https://doi.org/10.1093/icesjms/fsu172)
- Jordà G, Gomis D, Adloff F 2020 Vulnerability of marginal seas to sea-level rise: The case of the Mediterranean Sea. (*in Rev.*

- Karmaoui A 2016 Environmental Vulnerability to Climate Change in Mediterranean Basin: Socio-Ecological Interactions between North and South, in *Handbook of Research on Climate Change Impact on Health and Environmental Sustainability*, ed. Dinda S, 105–138. doi: [10.4018/978-1-4666-8814-8.ch006](https://doi.org/10.4018/978-1-4666-8814-8.ch006)
- Kennou H, Soer G, Menichetti E, Lakhdari F, Quagliarotti D 2019 The Water-Energy-Food Security Nexus in the Western Mediterranean. Barcelona, Spain.
- Kolodny EY 1974 *La Population des îles de la Grèce. Essai de géographie insulaire en Méditerranée orientale*. Edisud. Aix-en-Provence.
- Kuglitsch FG, Toreti A, Xoplaki E, Della-Marta PM, Zerefos CS et al. 2010 Heat wave changes in the eastern Mediterranean since 1960. *Geophys. Res. Lett.* 37, 1–5. doi: [10.1029/2009GL041841](https://doi.org/10.1029/2009GL041841)
- Le Roy Ladurie E 2004 *Histoire humaine et comparée du climat Tome 2 : Disettes et révolutions 1740-1860*. , ed. Fayard.
- Le Roy Ladurie E 2006 *Histoire humaine et comparée du climat Tome 3: Le réchauffement de 1860 à nos jours*. ed. Fayard.
- Lionello P 2012 *The Climate of the Mediterranean Region: From the Past to the Future*. Elsevier Science. doi: [10.1016/C2011-0-06210-5](https://doi.org/10.1016/C2011-0-06210-5)
- Lionello P, Abrantes FG, Gačić M, Planton S, Trigo RM et al. 2014 The climate of the Mediterranean region: research progress and climate change impacts. *Reg. Environ. Chang.* 14, 1679–1684. doi: [10.1007/s10113-014-0666-0](https://doi.org/10.1007/s10113-014-0666-0)
- Luterbacher J, Werner JP, Smerdon JE 2016 European summer temperatures since Roman times. *Environ. Res. Lett.* 11, 2 12. doi: [10.1088/1748-9326/11/2/024001](https://doi.org/10.1088/1748-9326/11/2/024001)
- Mastrandrea MD, Mach KJ, Plattner G-K, Edenhofer O, Stocker TF et al. 2011 The IPCC AR5 guidance note on consistent treatment of uncertainties: a common approach across the working groups. *Clim. Change* 108, 675–691. doi: [10.1007/s10584-011-0178-6](https://doi.org/10.1007/s10584-011-0178-6)
- Millar C, Neilson R, Bachelet D, Drapek R, Lenihan J 2006 Climate change at multiple scales, in *Forests, Carbon and Climate Change: A Synthesis of Science Findings*, eds. Salwasser H, Cloughesy M (Portland, Oregon, USA: Oregon Forest Resources Institute), 31–60.
- Nakićenović N 2000 *Special report on emissions scenarios: A special report of Working Group III of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- Navarra A, Tubiana L 2013 *Regional Assessment of Climate Change in the Mediterranean*. Berlin, Heidelberg, New York: Springer.
- Negev M, Paz S, Clermont A, Pri-Or NG, Shalom U et al. 2015 Impacts of climate change on vector borne diseases in the Mediterranean Basin - implications for preparedness and adaptation policy. *Int. J. Environ. Res. Public Health* 12, 6745–6770. doi: [10.3390/ijerph120606745](https://doi.org/10.3390/ijerph120606745)
- Orru H, Ebi KL, Forsberg B 2017 The interplay of climate change and air pollution on health. *Curr. Environ. Heal. Reports* 4, 504–513. doi: [10.1007/s40572-017-0168-6](https://doi.org/10.1007/s40572-017-0168-6)
- Palmiéri J, Orr JC, Dutay J-C, Béranger K, Schneider A et al. 2015 Simulated anthropogenic CO<sub>2</sub> storage and acidification of the Mediterranean Sea. *Biogeosciences* 12, 781–802. doi: [10.5194/bg-12-781-2015](https://doi.org/10.5194/bg-12-781-2015)
- Raleigh C, Jordan L, Salehyan I 2008 *Assessing the Impact of Climate Change on Migration and Conflict*. Washington, DC.
- Rickards L, Howden SM 2012 Transformational adaptation: Agriculture and climate change. in *Crop and Pasture Science*. doi: [10.1071/CP11172](https://doi.org/10.1071/CP11172)
- Rigaud KK, de Sherbinin A, Jones B, Bergmann J, Clement V et al. 2018 *Groundswell: Preparing for Internal Climate Migration*. Washington, DC. <https://openknowledge.worldbank.org/handle/10986/29461>
- Sağlam G 2020 The Mediterranean Sea: Cradle of Civilization. <https://www.un.org/en/chronicle/article/mediterranean-sea-cradle-civilization> [Accessed July 19, 2020]
- Serra-Majem L, Bach-Faig A, Miranda G, Clapes-Badrinas C 2011 Foreword: Mediterranean diet and climatic change. *Public Health Nutr.* 14, 2271–2273. doi: [10.1017/s1368980011002503](https://doi.org/10.1017/s1368980011002503)
- Somot S, Jordà G, Harzallah A, Darmaraki S 2016 The Mediterranean Sea in the future climate projections, in *The Mediterranean Region under Climate Change. A Scientific Update*, eds. Thiébaud S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 93–104.
- Tour du Valat 2012 *Mediterranean Wetlands: Outlook. First Mediterranean Wetlands Observatory report - Technical report*.
- Tramblay Y, Somot S 2018 Future evolution of extreme precipitation in the Mediterranean. *Clim. Change* 151, 289–302. doi: [10.1007/s10584-018-2300-5](https://doi.org/10.1007/s10584-018-2300-5)
- Tsimplis MN, Calafat FM, Marcos M, Jordà G, Gomis D et al. 2013 The effect of the NAO on sea level and on mass changes in the Mediterranean Sea. *JGR Ocean.* 118, 944–952. doi: [10.1002/jgrc.20078](https://doi.org/10.1002/jgrc.20078)
- Turco M, Jerez S, Doblas-Reyes FJ, AghaKouchak A, Llasat MC et al. 2018 Skilful forecasting of global fire activity using seasonal climate predictions. *Nat. Commun.* 9, 2718. doi: [10.1038/s41467-018-05250-0](https://doi.org/10.1038/s41467-018-05250-0)
- UN-SDSN 2018 *The World's Knowledge Network for the Sustainable Development Goals*. Paris, France and New York, USA.



- UN 2015 Transforming our world: the 2030 Agenda for Sustainable Development. New York.  
[https://www.un.org/ga/search/view\\_doc.asp?symbol=A/RES/70/1&Lang=E](https://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E)
- UNEP/MAP 2016 Mediterranean Strategy for Sustainable Development 2016-2025. Valbonne.
- UNISDR 2015 Sendai Framework for Disaster Risk Reduction 2015-2030.
- UNWTO 2019 Yearbook of Tourism Statistics, Data 2013-2017. Madrid, Spain  
 doi: [10.18111/9789284420414](https://doi.org/10.18111/9789284420414)
- Vautard R, Gobiet A, Sobolowski SP, Kjellström E, Stegehuis A et al. 2014 The European climate under a 2°C global warming. *Environ. Res. Lett.* 9, 34006. doi: [10.1088/1748-9326/9/3/034006](https://doi.org/10.1088/1748-9326/9/3/034006)
- Vermeulen SJ, Challinor AJ, Thornton PK, Campbell BM, Eriyagama N et al. 2013 Addressing uncertainty in adaptation planning for agriculture. *Proc. Natl. Acad. Sci. U. S. A.* 110, 8357–8362.  
 doi: [10.1073/pnas.1219441110](https://doi.org/10.1073/pnas.1219441110)
- Wise RM, Fazey I, Stafford Smith M, Park SE, Eakin HC et al. 2014 Reconceptualising adaptation to climate change as part of pathways of change and response. *Glob. Environ. Chang.* 28, 325–336.  
 doi: [10.1016/J.GLOENVCHA.2013.12.002](https://doi.org/10.1016/J.GLOENVCHA.2013.12.002)
- Woodward J 2009 (ed.) 2009. *The Physical Geography of the Mediterranean*. Oxford University Press.
- WWAP 2019 The United Nations World Water Development Report 2019: Leaving No One Behind. Paris.
- Zamora Acosta E, Maya Álvarez P 1998 Relaciones interétnicas y multiculturalidad en el Mediterráneo Occidental. in *V Centenario de Melilla, Melilla*.

## Information about authors

### Coordinating Lead Authors

Manfred A. Lange:

*The Cyprus Institute, Nicosia, Cyprus*

Maria Carmen Llasat:

*University of Barcelona, Barcelona, Spain*

Maria Snoussi:

*Mohammed V University, Rabat, Morocco*

### Lead Authors

Arnault Graves:

*Union for the Mediterranean, Barcelona, Spain*

Julien Le Tellier:

*United Nations Environment Programme /  
Mediterranean Action Plan (UNEP/MAP) –  
Barcelona Convention Secretariat, Athens, Greece*

Arnault Queralt Bassa:

*Advisory Council for the Sustainable  
Development of Catalonia of the Government  
of Catalonia (CADS), Barcelona, Spain*

Grazia Maria Vagliasindi:

*University of Catania, Catania, Italy*

### Contributing Authors

Elen Lemaître-Curri:

*Centre International de Hautes Études Agronomiques  
Méditerranéennes (CIHEAM), Mediterranean  
Agronomic Institute of Montpellier, (IAM), Montpellier,  
France*

Piero Lionello:

*University of Salento, Lecce, Italy*

Marini Katarzyna:

*MedECC, Plan Bleu, Marseille, France*

Cyril Moulin:

*National Institute for Earth Sciences and Astronomy  
(CNRS-INSU), Paris, France*



# 2 DRIVERS OF CHANGE

#### Coordinating Lead Authors:

Semia Cherif (Tunisia), Enrique Doblás-Miranda (Spain), Piero Lionello (Italy)

#### Lead Authors:

Carlos Borrego (Portugal), Filippo Giorgi (Italy), Ana Iglesias (Spain), Sihem Jebari (Tunisia), Ezzeddine Mahmoudi (Tunisia), Marco Moriondo (Italy), Olivier Pringault (France), Gil Rilov (Israel), Samuel Somot (France), Athanassios Tsikliras (Greece), Montserrat Vilà (Spain), George Zittis (Cyprus)

#### Contributing Authors:

Giovanni Argenti (Italy), Marie-Anne Auger-Rozenberg (France), Ernesto Azzurro (Italy), Corina Basnou (Spain), Sophie Bastin (France), Mustapha Béjaoui (Tunisia), Lorenzo Brilli (Italy), Martina Carrete (Spain), Emma Cebrian (Spain), Hanene Chaabane (Tunisia), Silvia Coelho (Portugal), Renato Colucci (Italy), Styliani Dafka (Germany), Sofia Darmaraki (Greece/Canada), Camilla Dibari (Italy), Donna Dimarchopoulou (Greece), Jean-Claude Dutay (France), Monia El Bour (Tunisia), Antonietta Elia (Spain), Elena Georgopoulou (Greece), Sylvaine Giakoumi (France), Juan Jesús González Alemán (Spain), Pablo González-Moreno (UK), Madeleine Goux (France), Olivier Grünberger (Tunisia), Ivan Güttler (Croatia), Nathalie Hilmi (Monaco), Gabriel Jordà (Spain), Stelios Katsanevakis (Greece), Mehdi Lahlou (Morocco), Manfred A. Lange (Cyprus), Luisa Leolini (Italy), Myriam Lopes (Portugal), Annarita Mariotti (USA), Ana Isabel Miranda (Portugal), Meryem Mojtafid (France), Alexandra Monteiro (Portugal), Samuel Morin (France), Pierre Nabat (France), Anika Obermann-Hellhund (Germany), Tuğba Öztürk (Turkey), Androniki Pardalou (Greece), Sandra Rafael (Portugal), Francesca Raffaele (Italy), Lena Reimann (Germany), Alain Roques (France), Asma Sakka Hlaili (Tunisia), Alberto Santini (Italy), Giuseppe Scarcella (Italy), Katrin Schroeder (Italy), Isla Simpson (USA), Nicolina Staglianò (Italy), Meryem Tanharte (Morocco), Rob Tanner (France), Rémi Thiéblemont (France), Yves Trambly (France), Marco Turco (Italy), Nassos Vafeidis (Germany), Martin Wild (Switzerland), Elena Xoplaki (Germany), Argyro Zenetos (Greece)

*This chapter should be cited as: Cherif S, Doblás-Miranda E, Lionello P, Borrego C, Giorgi F, Iglesias A, Jebari S, Mahmoudi E, Moriondo M, Pringault O, Rilov G, Somot S, Tsikliras A, Vilà M, Zittis G 2020 Drivers of change. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 59-180.*



# Table of contents

<b>2. Drivers of change</b> .....	<b>64</b>
Executive summary.....	64
<b>2.1 Introduction</b> .....	<b>66</b>
<b>2.2 Climate change</b> .....	<b>67</b>
2.2.1 Framing.....	67
2.2.1.1 <i>Observations and reanalyses</i> .....	68
2.2.1.2 <i>Modelling</i> .....	68
2.2.2 General circulation and dynamics.....	70
2.2.2.1 <i>General circulation</i> .....	70
<i>Observed trends</i> .....	70
<i>Future changes</i> .....	70
2.2.2.2 <i>Teleconnection patterns</i> .....	72
2.2.2.3 <i>Extratropical cyclones and medicanes</i> .....	72
2.2.2.4 <i>Regional winds</i> .....	73
2.2.3 Radiation, clouds and aerosols.....	74
2.2.3.1 <i>Observed change in surface radiation</i> .....	74
2.2.3.2 <i>Projected change in surface radiation</i> .....	75
2.2.4 Temperature and related extremes.....	75
2.2.4.1 <i>Observed temperature changes</i> .....	75
2.2.4.2 <i>Future temperatures</i> .....	76
2.2.5 Precipitation, related extremes and the water cycle.....	79
2.2.5.1 <i>Observed trends in precipitation</i> .....	79
2.2.5.2 <i>Future precipitation</i> .....	81
<i>Mean precipitation changes</i> .....	81
<i>Variability and extremes</i> .....	82
2.2.5.3 <i>Changes in evaporation, net water losses over sea and over land</i> .....	83
2.2.6 The cryosphere.....	84
2.2.6.1 <i>Observed trends in the cryosphere</i> .....	84
2.2.6.2 <i>Future conditions in the cryosphere</i> .....	85
2.2.7 Ocean hydrology.....	86
2.2.7.1 <i>Observed change in marine waters</i> .....	87
2.2.7.2 <i>Future change in marine waters</i> .....	88
<i>Air-sea and land-sea exchanges</i> .....	88
<i>Sea surface temperature</i> .....	89
<i>Sea surface salinity</i> .....	90
<i>Surface circulation and exchanges across straits</i> .....	90
<i>Deep water characteristics</i> .....	90
2.2.8 Sea level, storm surges and wave heights.....	91
2.2.8.1 <i>Observed change in sea-level</i> .....	91
2.2.8.2 <i>Future sea-levels</i> .....	91
2.2.9 Acidification of the Mediterranean Sea.....	92
2.2.9.1 <i>Observed change in acidity</i> .....	94
2.2.9.2 <i>Future change in acidity</i> .....	94
<b>2.3 Pollution</b> .....	<b>94</b>
2.3.1 Introduction.....	94

2.3.2	Physical pollutants.....	94
2.3.2.1	Particulate matter (PM) levels and sources.....	95
2.3.2.2	Particulate matter (PM) chemical profiles.....	96
2.3.2.3	Plastics (macro/micro/nano).....	97
2.3.3	Chemical pollutants.....	98
2.3.3.1	Nutrients.....	98
2.3.3.2	Gaseous pollutants: nitrogen dioxide, sulphur dioxide, ozone.....	100
2.3.3.3	Trace metallic elements.....	101
2.3.3.4	Organic pollutants.....	103
	Polycyclic Aromatic Hydrocarbons (PAHs).....	103
	Pesticides.....	104
2.3.3.5	Emerging contaminants.....	105
2.3.4	Biological pollutants.....	107
<b>2.4</b>	<b>Land and sea use changes.....</b>	<b>109</b>
2.4.1	Land use changes.....	109
2.4.1.1	Past trends and recent dynamics.....	109
	Mediterranean landscapes.....	109
	Recent changes in Mediterranean landscapes.....	110
2.4.1.2	Principal impacts of land use changes.....	110
	Forestry and other natural resources.....	110
	Crops and livestock.....	112
	Different land uses as a single driver of change.....	113
2.4.1.3	Future projections.....	113
2.4.2	Sea use changes.....	114
2.4.2.1	Trends in fisheries exploitation.....	114
2.4.2.2	Current status of marine fisheries resources.....	115
2.4.2.3	The future of marine resources.....	115
<b>2.5</b>	<b>Non-indigenous species.....</b>	<b>116</b>
2.5.1	Non-indigenous species in the Mediterranean Sea.....	116
2.5.1.1	Spatiotemporal trends, sources and vectors of introduction.....	117
2.5.1.2	Non-indigenous species as drivers of biodiversity and ecosystem change.....	117
2.5.1.3	Further introductions, monitoring and managing non-indigenous species.....	119
2.5.2	Terrestrial non-indigenous species and pests.....	120
2.5.2.1	Spatial patterns and temporal trends.....	120
	Degree of introduction across Mediterranean-type ecosystems and geographical areas.....	120
	Temporal trends of non-indigenous species and pests.....	122
	Pathways of introduction (intentional and accidental) of non-indigenous species and pests.....	123
2.5.2.2	Non-indigenous species as drivers of biodiversity and ecosystem change.....	123
2.5.2.3	Further introductions, spread and impacts of non-indigenous species and pests.....	125
	Existing tools for predicting the risk of introduction and research needs.....	125
	Non-indigenous species likely to be introduced into the Mediterranean in the next 20-50 years.....	125
<b>2.6</b>	<b>Interaction among drivers.....</b>	<b>126</b>
2.6.1	Drivers impacting other drivers.....	126
2.6.2	Pairs of interacting drivers.....	127
2.6.2.1	Climate change effects on pollution.....	127
2.6.2.2	Pollution effects on climate change.....	127
2.6.2.3	Impact of climate on land and sea use.....	127
	Effects of climate change on land use.....	127
	Climate change and variability drives dynamics of marine species.....	128
2.6.2.4	Effects of land use on climate change.....	129

2.6.2.5	<i>Links between trends in non-indigenous species and climate change</i> .....	<b>129</b>
	<i>Impact of climate change on marine non-indigenous species</i> .....	<b>129</b>
	<i>Impacts of climate change on terrestrial non-indigenous species</i> .....	<b>131</b>
2.6.2.6	<i>Impacts of pollution on land and sea use</i> .....	<b>131</b>
2.6.2.7	<i>Impacts of land and sea use change on pollution</i> .....	<b>131</b>
2.6.2.8	<i>Pollution effects on non-indigenous species</i> .....	<b>132</b>
2.6.2.9	<i>Effects of land and water use on non-indigenous species</i> .....	<b>132</b>
2.6.2.10	<i>Effects of non-indigenous species on land and sea use</i> .....	<b>132</b>
2.6.3	<b>More complex interactions among drivers</b> .....	<b>132</b>
2.6.3.1	<i>Floods</i> .....	<b>132</b>
2.6.3.2	<i>Desertification</i> .....	<b>132</b>
2.6.3.3	<i>Wildfires</i> .....	<b>132</b>
2.7	<b>Mediterranean socioeconomic scenarios</b> .....	<b>134</b>
Box 2.1	<b>How much has the Mediterranean Basin warmed since the pre-industrial period?</b> .....	<b>137</b>
Box 2.2	<b>Representative Concentration Pathways (RCPs)</b> .....	<b>138</b>
References	.....	<b>139</b>
Information about authors	.....	<b>179</b>

## 2 Drivers of change

### Executive summary

#### *Climate drivers*

During recent decades, observations of several variables provide evidence of the ongoing anthropogenic climate change in the Mediterranean region, particularly increase of mean and extreme temperatures, and dry environmental conditions. Climate projections show that the region will among the most affected regions by climate change, specifically regarding precipitation and the hydrological cycle, but also mean warming and heat extremes (in both the terrestrial and marine environment), sea level rise and sea water acidification.

Basin-wide, annual mean temperatures are now 1.5°C above the preindustrial level. In the last decades dry conditions have become more frequent and a large reduction of glaciers across high mountains of the Mediterranean has occurred at a progressively increasing pace. Mediterranean Sea waters have become warmer and saltier, Mediterranean sea level has risen at a rate (1.4 mm yr<sup>-1</sup>) similar to the global trend at centennial scale.

In the future, the regional average warming will exceed the global mean value by 20% and it might reach 5.6°C at the end of the 21st century in the RCP8.5 high emission scenario. Heat waves and warm temperature extremes will intensify. Total annual precipitation is expected to decrease over most of the region (the average reduction rate is approximately 4% per each degree of global warming). However, magnitude and spatial distribution of changes are uncertain, because of differences among models. Dry conditions will be further enhanced by increasing evapotranspiration over land. At the same time, the inter-annual variability of the hydrological cycle will increase, with longer dry spells especially in the southern areas. Extreme precipitation events will become more intense over large parts of the northern Mediterranean areas.

Mediterranean mean sea level is projected to be at the end of the 21st century in the range from 20 to 110 cm higher than at the end of the 20th century, depending on the level of anthropogenic emissions. Sub-regional and local relative sea level rise will be further modulated by vertical land motions and regional circulation features (with deviations in the

order of 10 cm from the basin average). Therefore, though in the future milder marine storms are expected, coastal hazards, floods and erosion will increase, because of mean sea level rise.

Widespread seawater warming will continue. Annual mean surface temperature will increase 2.7-3.8°C and 1.1-2.1°C in one century under the RCP8.5 and the RCP4.5 scenarios, respectively. Marine heat waves will become longer, more intense than today and their spatial extent will increase. Seawater acidification will continue, with a pH reduction that might larger than 0.4 units at the end of the 21st century.

#### *Pollution*

Across the Mediterranean Basin, ocean and inland pollution are ubiquitous, diverse and increasing in both quantity and in the number of pollutants, due to demographic pressure, enhanced industrial and agricultural activities, and climate change.

Mediterranean seawater is generally oligotrophic (low nutrient), with decreasing levels from Gibraltar eastwards to the Levantine Sea. Several coastal regions are hotspots of human-induced nutrient inputs. This nutrient enrichment causes eutrophication and may provoke harmful and toxic algal blooms, whose frequency will likely increase. Harmful algal blooms may cause negative impacts on ecosystems and may represent serious economic threats for fisheries, aquaculture, tourism and human health.

Emerging contaminants are well present across the Mediterranean Basin, and enhanced by increasing inflow of untreated wastewater. These substances may cause disorders of the nervous, hormonal and reproductive system. And the increasing frequency of extreme precipitation events in the north of the Mediterranean increases the supply of fecal bacteria and viruses to the coastal zone. The Mediterranean Sea is one of the most polluted large water bodies globally in terms of plastic and the level of this pollution is expected to increase in the future.

The Mediterranean Basin is among the regions in the world with the highest concentrations of gaseous air pollutants (NO<sub>2</sub>, SO<sub>2</sub> and O<sub>3</sub>). Its dry and sunny climate, and specific atmospheric circulation patterns enhance air pollution levels. Ships are among the major causes of increasing SO<sub>2</sub> and NO<sub>x</sub> emissions in this region.



Emissions of aerosols and particulate matter (PM) into the atmosphere arise from a variety of anthropogenic activities. Particular meteorological conditions and natural sources, including the proximity of the Sahara Desert, create particular patterns of aerosol concentrations that may influence particulate matter PM concentrations. The occurrence of critically high PM concentrations associated with dust outbreaks is higher in the southern Mediterranean (>30% of the annual days) than in the northern area (<20% of the annual days).

### ***Land and sea use change***

Landscapes and their use have changed over millennia in the Mediterranean Basin, however the rate of change has increased substantially since the second half of the 20th century, with rapid growth of urban and peri-urban areas leading to loss of biodiversity and habitats. Abandonment of agro-pastoralism (which will likely continue in the future) is causing unmanaged shrubs forest development in marginal lands, arid and mountain areas in European countries, while land overexploitation is causing widespread forest degradation in areas of North Africa and the Middle East. Future land use trends depend strongly on regional policies for urbanization, agriculture, forestry and nature conservation.

Marine resource overexploitation and unsustainable fishing practices have increased in time and are the main drivers of the population decline of several species. Presently, more than 60% of marine stocks have collapsed or are overexploited. Sustainable management of marine resources requires the reduction of fishing pressure.

### ***Non-indigenous species***

The Mediterranean Sea (and particularly the Levantine Basin) is a hotspot for the establishment of many non-indigenous species (invertebrates, primary producers, and vertebrates), whose arrival and increase are linked to the decrease or collapse in populations of native species. Most marine non-indigenous species enter the Mediterranean Sea from the Red Sea and Atlantic Ocean, but those introduced by ships and aquaculture produce the largest impact on the environment. The number and spread of non-indigenous species will likely further increase with increasing shipping activity and impacts of climate change on the Mediterranean water masses.

Mediterranean land areas currently host a high number of non-indigenous species (mostly plants and invertebrates) in human-modified ecosystems and in regions with high infrastructure development. Most invertebrate species are phytophagous pests that cause damages to crops and forests. Future warming is predicted to induce a northward shift at a speed of 37-55 km decade<sup>-1</sup> of current major non-indigenous species and determine a window of opportunity for new non-indigenous species adapted to dry environments. The presently increasing trend of the numbers of introduced invertebrates and vertebrates (the latter generally caused by accidental escapes) will very likely continue, as they can be easily transported also as stowaways in air and maritime cargo.

### ***Interaction among drivers***

When ecosystems and societal sectors are threatened by multiple, co-occurring drivers, climate change, pollution, land and sea use change, and non-indigenous species can interact. Interactions cause effects that can be additive/cumulative, synergistic or antagonistic and result in alteration, intensification, and even in generation of new impacts. Examples of new threats are increase of flood events, due to a combination of climatic and land use changes, desertification, which is the result of increasing aridity and exploitation of resources, and wildfires, affected by forest encroachment and heat waves, among many other interactions.

## 2.1 Introduction

This chapter describes characteristics and evolution of human-induced and natural factors that cause changes in the Mediterranean Basin ecosystems and human systems. In order to cover most major risks for people and biodiversity, four broad domains of change drivers are considered: climate change and variability, pollution, land and sea use changes and non-indigenous species. These factors correspond to the concept of “direct drivers”, which was introduced in the Millennium Ecosystem Assessment (MEA 2005; Nelson et al. 2006), that unequivocally influence processes in ecosystems and can be identified and measured to differing degrees of accuracy.

Anthropogenic climate change is already affecting the environment and societies in the Mediterranean region. Warming is unequivocal, and there are emerging signs of changes of the hydrological cycle and other climate variables (*Section 2.2*). Climate models indicate a trend towards a warmer and drier environment, seawater warming, with more intense warm extremes both over land and in the sea, and regional increase of sea level (*Section 2.2*). The Mediterranean region is likely very vulnerable to climate change and many components of its terrestrial and marine environment are already under stress (*Section 2.2*).

Atmospheric and water pollution can be driven by many factors, which affect all the compartments of the environment: water, air and soil/sediments. Pollutants can migrate from one media to another. There is a wide range of pollutants that can be biological (e.g., bacteria or insects), chemical (e.g., pesticides, trace metals) or physical (e.g., particulate matter) (*Section 2.4-6 and Chapters 3 and 4 of this report*).

Changes in land and sea use changes are considered among the major direct drivers of environmental change worldwide, but their characteristics vary, depending on each region, even at very local scale. Mediterranean terrestrial landscapes and ecosystems show different patterns of change on northern and southern shores, due to urbanization, coastal development, evolving agricultural and farming practices, including their abandonment. The overexploitation of the Mediterranean Sea resources poses a particular threat due to its intrinsic geographical limits (*Section 2.4*).

Non-indigenous species are profoundly affecting terrestrial and marine ecosystems in the

Mediterranean and their impact is not only measurable in biodiversity alterations, but also in human health and economic damages (*Section 2.5*).

There is no strict consensus of the grouping of the drivers into the categories that have been adopted in this report. More condensed or more articulated lists can be found in the scientific literature. Our four categories include all physical, chemical and biological factors that directly act on the Mediterranean environment, with a substantial correspondence with those used in the recent IPBES 2018 regional reports (Bustamante et al. 2018; Elbakidze et al. 2018; Nyingi et al. 2018; Wu et al. 2018b).

The level or rate of change of direct drivers can be influenced or altered by indirect drivers (MEA 2005; Nelson et al. 2006). Indirect drivers are grouped in categories such as demographic, economic, sociopolitical, cultural, religious, technological, legislation and financial drivers. In turn, indirect drivers are distinguished in “endogenous” and “exogenous” drivers, whose magnitude can and cannot be influenced/alterd by the decision-makers, respectively. Whether a driver is exogenous or endogenous depends on the organizational level and on the spatial and temporal scale. The concepts of indirect and direct drivers roughly match those of driving forces and pressures in the Drivers–Pressures–State–Impacts–Responses (DPSIR) framework, which was initially developed by the European Environmental Agency (EEA 1999).

Understanding of indirect drivers is essential for the benefit of the environment. It is the action on indirect drivers by policymakers and stakeholders that can effectively manage the risks posed to the environment and human societies by climate change, pollution, land and sea use changes and arrival of non-indigenous species. However, this is a quite different topic with respect to the content of this chapter. Mechanism and tools influencing direct drivers are considered in other parts of this report. The scope of this chapter is the assessment of the state of knowledge of physical, chemical, and biological factors, of their present status, past and future evolution in the Mediterranean Basin.


## 2.2 Climate change

### 2.2.1 Framing

There is no universal definition of the land boundaries of the Mediterranean region (*Chapter 1*). In this section, and also for much of this report, we adopt a simple regular latitude-longitude box (29°N to 47.5°N and 10°W to 39°E, *Fig. 2.1*), which includes some regions with other than Mediterranean climates, such as in the Alps, the Eastern Balkans or part of the Sahara. This definition of Mediterranean region is similar to the MED zone adopted in IPCC-AR4 (IPCC 2007), and slightly larger than in IPCC-AR5 (IPCC 2013a).

delimitations (*Fig. 2.1*), the Western Mediterranean Sea (MEDW) between the Gibraltar Strait and the Sicily Strait, the Adriatic Sea (ADR) north of the Otranto Strait, the Aegean Sea (AEG) north of the Cretan Arc Straits and the Eastern Mediterranean Sea (MEDE) for the remaining.

When assessing future climate change, it is important to specify the reference period to which climate projections are compared, along with future “time slices”. In MAR1 we use 20-year long time periods following standard IPCC practice. This length of time period is sufficient to smooth part



**Figure 2.1 | Mediterranean coastline, topography over land and bathymetry over the sea plus the box definition.** Relief data are derived from the ETOPO1 1 Arc-Minute Global Relief Model<sup>5</sup>. Sub-regions are defined as West Maghreb (WM), East Maghreb (EM), Levant (LE), Anatolia (AN), Central Mediterranean (CM), Iberia (IB), France (FR), Alps (AL) and Balkans (BA). The Mediterranean Sea is divided into 4 sub-basins, the Western Mediterranean Sea (MEDW), the Eastern Mediterranean Sea (MEDE), the Aegean Sea (AEG) and the Adriatic Sea (ADR).

In order to provide a spatially refined assessment, we define sub-regions over land and over sea by using smaller rectangular boxes. Over land, the European sub-regions follow the definition proposed during the PRUDENCE European project (Christensen et al. 2002) (*Fig. 2.1*), and we define new sub-regions (inspired by Nabat et al. 2015b) for the Middle East and Africa. Over the sea, 4 sub-regions are defined using the natural strait

of the high-frequency natural climate variability that can otherwise mask the forced trend, but it is short enough to assume that climate does not change much during the 20 years covered. For the reference period, we choose the latest years of the 20th century (1980-1999). This choice is a compromise related to the observation and model data availability at the Mediterranean scale. IPCC has traditionally chosen the pre-industrial

<sup>5</sup> <http://www.ngdc.noaa.gov/mgg/global/global.html>

period (around 1850 or 1900) as a reference but regional climate models and regional high-quality observations are not available for that period. This choice (2 full decades at the end of the 20th century) also targets to facilitate the repeatability of the MAR1 computations made in future Mediterranean studies or reports.

In addition to the reference period, we define also the “present-climate” period (1995-2014), which defines the current climate conditions. The reference period is fixed in different reports to be able to intercompare results of simulations across different model generations, whereas the present-climate will move from one report to another. For example, the IPCC-AR4 (IPCC 2007) defined 1981-2000 as present climate, whereas the IPCC-AR5 (IPCC 2013a) used 1986-2005. For the assessment of past changes, the longest period available (generally 1950-2019 or 1900-2019, if possible) in the observations is used in this report and units such as °C per decade are used in order to compare past trends computed over different past periods.

For the future, we keep 20-year time slices in order to sample the same level of internal variability as in the reference period. We divide the 21st century in 20-year time slices with a present-climate period (2000-2019), a near-future period (2020-2039), a mid-term period centered in 2050 (2040-2059) and a far-future period close to the end of the 21st century (2080-2099). The mid-21st century period is arguably of particular interest for many stakeholders, especially for mid-term adaptation. The end of the 21st century period is also of interest for stakeholders working on mitigation targets and involved in very long-term planning (e.g., for the design and planning of dams, forests, cities).

For the future climate change assessment, an important part of the uncertainty is related to the future evolution of socio-economic development. To be able to propose future climate projections considering various possible socio-economic trajectories and climate policy pathways, we follow the IPCC scenario approach. Where more recent studies are not available, the assessment also considers studies based on the older IPCC SRES approach (Nakićenović 2000). Although results based on multiple IPCC scenarios are reported in the MAR1, we mostly focus on two options which encompass the range of IPCC-AR5, CMIP5 and CORDEX simulations: the “business as usual” scenario (RCP8.5, for an explanation of the RCPs see *Box 2.2*) and the optimistic scenario closest to the UNFCCC Paris Agreement target (RCP2.6). These scenarios have been chosen also due to

model projection availability constraints at the regional scale.

Detecting trends and attributing regional climate change to human influence is challenging due to natural climate variability and the strong spatio-temporal dependency of different climate variables. For projections, uncertainty estimates are provided where this is possible. For small model ensembles, the total range is also given. For larger ensemble, 90% confidence intervals are used as much as possible or else “whisker plots” describing the various statistics of the distribution (median, 25th and 75th percentile, 90% interval, minimum and maximum values).

### 2.2.1.1 Observations and reanalyses

More and more observation datasets have become available at regional scales, either from satellite or *in-situ* observations, or from reconstructions and reanalyses. This new generation of observation-based products are (1) long and homogeneous enough to allow trend studies (ESA-CCI) (Ribes et al. 2019); (2) of sufficiently high spatial resolution to capture complex topography and land-sea mask (SAFRAN, EURO-4M), thereby allowing regional to local studies; and (3) of sufficiently high temporal resolution (daily or hourly, COMEPHORE) (Fumière et al. 2019), to allow the study of regionally-relevant extreme events. High-resolution gridded products have also become available for southern Mediterranean countries e.g., Cyprus (Camera et al. 2017) or Tunisia (Tramblay et al. 2019).

Finding the best fit-for-purpose observation dataset is becoming a new challenge, given the large number of available products, often characterized by substantial differences. Results of past trend studies and model evaluations are sensitive to the choice of the reference dataset (Flaounas et al. 2012; Prein and Gobiet 2017; Zittis 2018; Fumière et al. 2019; Kotlarski et al. 2019; Peña-Angulo et al. 2020). Long-term, accessible, gridded, well-calibrated and homogeneous in time and space *in-situ* data are nonetheless still lacking, especially for the ocean or the high-frequency variables over land. In addition, regional model-based reanalyses are still rare.

Various observation datasets are used to assess the past evolution of the different components of the Mediterranean climate system. Atmospheric dynamics are mostly assessed against atmospheric reanalyses (ERA-Interim, ERA20C, 20CR) (*Section 2.2.2*). For aerosols, clouds and surface radiation, both satellite products and station data (BRSN,

GEBA) are used to estimate past evolutions (Section 2.2.3). The trend evaluation over land relies on high-resolution observation-based gridded products, i.e., CRU, E-OBS for temperature (Section 2.2.4) and CRU, E-OBS, U. Del, GPCC for precipitation (Section 2.2.5). The other water cycle components are evaluated against reconstructed products blending *in-situ* observations, satellite and models of river networks (Ludwig et al. 2009; Pellet et al. 2019; Wang and Polcher 2019) (Section 2.2.6), while satellite data are used for sea surface temperature (Marullo et al. 2010; Pisano et al. 2016; Pastor et al. 2018) (Section 2.2.7), and *in-situ* data for the deep water characteristics (Houpert et al. 2016; Schroeder et al. 2017; Testor et al. 2018; von Schuckmann et al. 2018).

### 2.2.1.2 Modelling

Complex and realistic global and regional climate models (GCMs and RCMs), based on fundamental physics, chemistry and biology equations are currently the standard tools to simulate the future evolution of the regional climate system. Different types of climate models are available to study the Mediterranean climate (past and future), often organized in large coordinated multi-model initiatives under the World Climate Research Programme (WCRP) umbrella CORDEX (Giorgi et al. 2009) and CMIP5 (Taylor et al. 2012). Combining the various sources of information or extracting the most credible (actionable) information is a new challenge, sometimes called the “distillation” problem (Hewitson et al. 2014; Fernández et al. 2019).

The MAR1 climate assessment concerning future climate evolution is based on four climate model ensembles, chosen for their good representation of the Mediterranean climate and for their good coverage of the various sources of uncertainty in future climate projections:

- CMIP3 and CMIP5 GCM ensembles with resolution ranging approximately from 300 to 100 km: they are the largest GCM multi-model ensembles available so far. They cover at a relatively low resolution all the uncertainty sources and can provide data for all the components of the climate system (atmosphere, land, ocean, marine biogeochemistry and aerosols). Some of the participating models share components and this may result in a redundancy in the ensemble results (Knutti et al. 2017).
- The Euro-CORDEX RCM ensemble is a large high-resolution ensemble at 12 km resolution (Jacob et al. 2014), which has clearly improved the representation of climate variables compared

to coarse resolution GCMs over land, e.g., for extreme precipitation (Fantini et al. 2018), regional winds (Obermann et al. 2018), mountain climate (Torma et al. 2015; Torma 2019), and over the sea, e.g., for regional winds (Herrmann et al. 2011) and extreme winds (“medicanes”) (Gaertner et al. 2018). An example for the strong modification of the future climate change signal by high-resolution RCM compared to GCMs has been found for summer precipitation over the Alps (Giorgi et al. 2016). Although the EURO-CORDEX ensemble is a large high-resolution multi-model dataset, it does not cover the entire uncertainty space of the CMIP5 ensemble.

- The Med-CORDEX RCM ensemble (Ruti et al. 2016) is a relatively small ensemble which does not cover the CMIP5 uncertainty range particularly well, but is the best data source available to study the future evolution of the Mediterranean Sea (Darmaraki et al. 2019b; Soto-Navarro et al. 2020), its ecosystems (Moullec et al. 2019) and atmosphere-ocean interactions. The models of this ensemble have high-resolution in both the atmosphere (resolution range: 25-50 km) and the ocean (resolution range: 6-30 km) component of the regional climate system (Somot et al. 2018).
- The CORDEX FPS-convection CPRCM ensemble (Coppola et al. 2020): this mini-ensemble provides the highest spatial resolution (2-3 km) for the greater Alpine region, reaching convection resolving scales. It yields, in particular, a strong improvement in the representation of extreme precipitation at sub-daily time scales (Kendon et al. 2014; Ban et al. 2015; Fosser et al. 2015; Berthou et al. 2018; Fumière et al. 2019). Convection resolving models are very promising tools to study the future evolution of extreme precipitation associated with thunderstorms, medicanes or mesoscale convective systems in the Mediterranean region (Lenderink et al. 2019) or urban-climate interactions.

A large variety of downscaling methods are available to study the Mediterranean climate (COST VALUE) (Maraun et al. 2019), including on-line tools on web processing servers (Cofiño et al. 2007). Among these methods, correcting climate change simulations using statistical tools (bias correction methods) allows to improve the present-climate statistics of climate simulations, with clear benefits for studying threshold-dependent extremes or for forcing impact models. All statistical methods require long-term observations (stations, gridded-products or satellite data calibrated for climate applications) for the learning phase and their application is therefore limited in regions where observations are lacking.

In the MAR1 report, CMIP ensembles are mostly used for the assessment of atmospheric dynamics, aerosol, cloud and radiation, water cycle, sea hydrology, sea level and acidification, whereas the Euro-CORDEX ensemble is used for the assessment of wind, clouds, temperature, precipitation and the cryosphere. The Med-CORDEX coupled regional models are used for sea hydrology and sea level, and the FPS-convection ensemble for the assessment of extreme precipitation.

Despite the continuous improvement of climate model ensembles by increased resolution and ensemble size, increased complexity and improved model physics, these still suffer from deficiencies and systematic errors. In particular their poor representation of some key regional phenomena may limit confidence for some aspects of the MAR1 assessment. This is especially true for coastal sea level, medicanes, tornadoes, hail phenomena, lightning, city climate, sub-daily precipitation, glaciers, clouds or cloud-aerosol interactions, human influence on land and water use.

## 2.2.2 General circulation and dynamics

### 2.2.2.1 General circulation


The proximity of the Mediterranean to the Atlantic and Indian Oceans and the surrounding massive land areas, with diverse climatic characteristics, places the area at the crossroads of many global climate patterns and processes of tropical and extra-tropical origin (Xoplaki et al. 2003a; Lionello et al. 2006; Lelieveld et al. 2012; Lionello et al. 2012a; Lionello 2012; Ulbrich et al. 2012). Its location on the eastern edge of the Atlantic Ocean means it is particularly affected by variability and change in the North Atlantic jet stream (or Polar Front Jet) in both winter and summer (Düneloh and Jacobbeit 2003; Hurrell et al. 2003; Athanasiadis et al. 2010; Bladé et al. 2012) and by stationary blocking patterns (Tyrlis and Hoskins 2008). The Mediterranean Basin is also influenced by semi-permanent large-scale anticyclones (e.g., the Azores anticyclone in the west during summer and the cold Siberian anticyclone in the northeast during winter), while mobile anticyclones play also important role throughout the whole year (Hatzaki et al. 2014). During the summer, the climate of the Mediterranean is further influenced by circulation patterns set up by the Asian summer monsoon (Rodwell and Hoskins 1996) and local orography (Simpson et al. 2015).

### Observed trends

The wintertime large-scale circulation has exhibited a long-term trend toward increased sea-level pressure and anticyclonic circulation over the Mediterranean (*Fig. 2.2a and b*) (Mariotti and Dell'Aquila 2012). Aside from this long-term trend, the historical record has also exhibited sizable multi-decadal variability. This is illustrated for the sea-level pressure anomalies in *Fig. 2.2e* and has also been discussed widely in the contexts of trends in the North Atlantic Oscillation (NAO) and associated Mediterranean drying that occurred over the latter half of the 20th century (Hurrell 1995) in which internal variability is thought to play an important role (Kelley et al. 2012). CMIP5 models suggest that the externally forced contribution to sea-level pressure trends since 1900 consist of a pattern that resembles that of the observed trends but with a magnitude that is considerably smaller (*Fig. 2.2c*). It is likely that both external forcing and internal variability have contributed to the observed long-term trends (Hoerling et al. 2012). During summer, it is challenging to assess the observed long-term trends, as there is no strong agreement in the pattern of sea-level pressure change (*Fig. 2.2f and g*). There are indications of a summertime decline in sea-level pressure over North Africa and the southern Mediterranean and, indeed, the CMIP5 models suggest that external forcings have contributed to a decline in sea-level pressure in this region over the 20th century (*Fig. 2.2h*).

### Future changes

Under rising greenhouse gas concentrations, climate models project that the Hadley Cell circulation will change, the tropics will expand and the mid-latitude westerlies and associated storm tracks will likely shift poleward (*medium/high confidence*) (Yin 2005; Lu et al. 2007b, 2007a; Chang et al. 2012; Barnes et al. 2013; Shaw et al. 2016; D'Agostino et al. 2017, 2020). This is expected to enhance subsidence and reduce storminess at the latitudes of the Mediterranean region, with a resulting reduction in precipitation (*medium confidence*). While there is considerable inter-model spread in the magnitude of these projected changes and the forced signal can be small compared to internal variability (Woollings and Blackburn 2012; Barnes et al. 2013; Zappa et al. 2015; Quan et al. 2018; Grise et al. 2019), the Mediterranean could be influenced by additional local circulation anomalies, leading to pronounced hydroclimate changes (Seager et al. 2014; D'Agostino and Lionello 2020). Future projections suggest that the wintertime trend toward increased anticyclonic circulation



**Figure 2.2 | Trends in sea level pressure (SLP).** (a)–(e) show the DJF (December–January–February) season. (a) and (b) show 1900–2018 trends in SLP from ERA20C and 20CR reanalyses, respectively. Grey = not significantly different from zero at the 95% level. Significance is calculated by resampling, with replacement, the residuals of the linear trend, adding the resampled residuals to the linear trend and re-calculating the linear trend. This is repeated 1,000 times to obtain the probability (p-value) at each grid point that the trend is significantly different from zero. Spatial autocorrelation is accounted for using the False Discovery Rate method of Wilks (2016) with control value = 0.1. Stippling depicts grid points where the magnitude of the trend is larger than in any of the individual CMIP5 ensemble members. (c) shows the CMIP5 multi-model mean trend for 1900–2018. The ensemble mean for each model is calculated, then the linear trend is obtained before calculating the average trend across all models. Grey depict regions where less than 3/4 of the models agree on the sign of the change. (d) is as (c) but for future trends from 2019–2099. (e) shows time series of 20 year running mean SLP averaged over the Mediterranean (green box in c) for the two re-analyses, all individual ensemble members from all models and the CMIP5 multi-model mean. (f)–(j) are as (a)–(e) but for JJA (June–July–August).

over the Mediterranean will continue at an accelerated pace (*Fig. 2.2d*) (Giorgi and Coppola 2007). This is accompanied by a strengthening of the westerly winds and increased storminess over northern Europe, reduced westerlies over north Africa

and decreasing storminess over the Mediterranean (Woollings and Blackburn 2012; Rojas et al. 2013; Zappa et al. 2013). Climate models predict a summertime poleward shift of the North Atlantic jet (Simpson et al. 2015) and a summertime de-

crease/wintertime increase in sea-level pressure centered over the Mediterranean (Fig. 2.2i) (Giorgi and Coppola 2007; Bladé et al. 2012), with the reduction in Mediterranean sea-level pressure continuing at an accelerated pace over the coming decades (Fig. 2.2i), albeit with a large inter-model spread (Fig. 2.2j). This was argued to be dominated by a heat-low response to drier soils in the Mediterranean region (Haarsma et al. 2009).

### 2.2.2.2 Teleconnection patterns

The influence of teleconnection patterns (linkages between weather changes occurring in widely separated regions of the globe) on Mediterranean climate has been extensively studied (Corte-Real et al. 1995; Hurrell and Van Loon 1997; Wibig 1999; Pozo-Vázquez et al. 2001; Quadrelli et al. 2001; Xoplaki et al. 2003a, 2003b, 2004; Hatzaki et al. 2009; Toreti et al. 2010; Ulbrich et al. 2012; Tyrlis and Lelieveld 2013; Rousi et al. 2015; Sen et al. 2019). Particularly during winter, the region is prone to the impacts of the westerly flow and the teleconnection patterns of North Atlantic Oscillation (NAO), Eastern Atlantic/Western Russia (EA/WR) and Scandinavian (SCAN) (Barnston and Livezey 1987).

The NAO is, for parts of the region, one of the most important modes of internal climate variability. It affects especially the storm-tracks and cyclogenesis over parts of the basin (Trigo et al. 2000, 2004; Reale and Lionello 2013) and mainly precipitation over the western-central Mediterranean in winter (Lamb and Pepler 1987; Rodríguez-Fonseca and de Castro 2002; Xoplaki et al. 2004). NAO is also found to have some influence on winter precipitation in parts of the eastern Mediterranean, however this influence is smaller (Düneloh and Jacobeit 2003; Xoplaki et al. 2004; Feliks et al. 2010; Felis and Rimbu 2010; Nissen et al. 2010). A lesser but distinct influence is detected between NAO and the Mediterranean surface air temperature (Hurrell 1995; Cullen and DeMenocal 2000; Ben-Gai et al. 2001; Pozo-Vázquez et al. 2001; Sáenz et al. 2001; Castro-Díez et al. 2002; Trigo et al. 2002b; Türkeş and Erlat 2003; Xoplaki et al. 2003b; Toreti et al. 2010).

Observed trends of NAO are not monotonic and are difficult to assess since decadal oscillations are too large to reach a conclusion with an acceptable level of confidence. Nevertheless, mostly negative trends prevail since the early 1990s following a positive trend starting in the 1960s (Ulbrich and Christoph 1999; Mariotti et al. 2002b; Türkeş and Erlat 2003; Trigo et al. 2004, 2006; Xoplaki et al.

2004; Pinto and Raible 2012; Saffioti et al. 2016; Iles and Hegerl 2017).

Climate projections mostly suggest a weak positive NAO trend in a warmer future climate (*low/medium confidence*), accompanied by a small northeastward displacement of its centers-of-action by the end of the 21st century (Ulbrich and Christoph 1999; Gillett et al. 2003, 2013; Hu and Wu 2004; Stephenson et al. 2006; Bacer et al. 2016; Deser et al. 2017; Barcikowska et al. 2020). Some studies indicate no significant trends (Fyfe et al. 1999; Dorn et al. 2003; Rauthe et al. 2004; Fischer-Bruns et al. 2009), or even decreasing trends (Osborn et al. 1999).

Studies on the effect of El Niño Southern Oscillation (ENSO) phenomenon on Mediterranean precipitation have shown that links exist, particularly during autumn and spring in the western Mediterranean and during winter in the Eastern Mediterranean. However, results are not conclusive concerning their evolution and robustness. In fact, the ENSO signal is difficult to be isolated, because of the dominating mid-latitude dynamics, the sign of its correlation with total precipitation depends on season and it is not stationary (Rodó et al. 1997; Rodó 2001; Mariotti et al. 2002b, 2005; Knippertz et al. 2003; Hasanean 2004; Alpert et al. 2006; López-Parages and Rodríguez-Fonseca 2012; Kalimeris et al. 2017).

### 2.2.2.3 Extratropical cyclones and medicanes

The Mediterranean is one of the main cyclogenetic areas of the world (Petterssen 1956; Hoskins and Hodges 2002; Wernli and Schwerz 2006), with much of the high-impact weather (e.g., strong winds and heavy precipitation) associated with cyclonic structures. Cyclogenesis areas such as the north-western Mediterranean, North Africa, the north shore of the Levantine Basin, the seasonality (Alpert et al. 1990a, 1990b; Trigo et al. 1999, 2002a; Lionello et al. 2006, 2016; Campins et al. 2011), as well as the occurrence of explosive cyclogenesis (Kouroutzoglou et al. 2011; Reale et al. 2019) are well documented in the literature. Within Mediterranean cyclones, there is a sub-group of hybrid depressions of extratropical cyclogenesis, the so-called 'medicanes' (Mediterranean hurricanes) or tropical-like cyclones (Rasmussen and Zick 1987; Reale and Atlas 2001; Emanuel 2005). These are mesoscale maritime extratropical cyclones that can physically emulate tropical characteristics at a certain point of their life cycle (Emanuel 2005; Miglietta 2019). Such features can include a cloud-



free area at the center (the “eye”), spiral bands with deep convection around it, intense surface winds and a warm-core and symmetric structure (Miglietta et al. 2015). These events can pose serious societal and ecological threat to the affected coastal regions (Nastos et al. 2018).

During the recent past there is an absence of strong trends in cyclone numbers affecting the Mediterranean (Lionello et al. 2016), however when trends are detected these are mostly negative (*low/medium confidence*) (Trigo et al. 2000; Maheras et al. 2001; Flocas et al. 2010; Nissen et al. 2010). Similarly, the number of explosive Mediterranean cyclones has likely decreased, but this reduction is not statistically significant (Kouroutzoglou et al. 2010). The statistical record of medicanes has limited reliability and sample size, given their maritime characteristics, small size and infrequent occurrence. Thus, it has not been possible to derive an objective climatology. Observational studies cannot be used to identify trends because the identification is commonly subjective (Miglietta et al. 2013; Tous and Romero 2013; Nastos et al. 2018). Dynamical downscaling methods have been used to build a medicane climatology, but only negligible trends were obtained (Cavicchia et al. 2014).

For the future, climate models project a reduction in the number of cyclones (*medium/high confidence*) especially in winter (Lionello et al. 2002; Geng and Sugi 2003; Bengtsson et al. 2006; Leckebusch et al. 2006; Lionello and Giorgi 2007; Pinto et al. 2007; Löptien et al. 2008; Ulbrich et al. 2009; Raible et al. 2010; Zappa et al. 2013; Nissen et al. 2014). There is some uncertainty, as the spread in the model responses appears to be quite large (Ulbrich et al. 2008, 2009; Harvey et al. 2012). With respect to cyclone intensity, climate projections are more controversial, as some models suggest a decrease in the frequency of the most intense systems (Pinto et al. 2007; Raible et al. 2010), while other models show more extreme events or increases in the intensity of extreme cyclones (Lionello et al. 2002; Gaertner et al. 2007). For medicanes, climate projections indicate a decreasing response in frequency but increasing intensity (Gaertner et al. 2007; Romero and Emanuel 2013, 2017; Cavicchia et al. 2014; Walsh et al. 2014; Tous et al. 2016; Romera et al. 2017; González-Alemán et al. 2019).


#### 2.2.2.4 Regional winds

Surface wind speed and its changes on different temporal and spatial scales are governed by driving

and drag forces, where all relevant contributions are difficult to estimate and disentangle (Wu et al. 2018a). Surface wind climate studies are less common than air-temperature and precipitation studies for example, and more work is needed to explain historical wind speed evolution and precisely estimate different sources of uncertainty in the future projections. This variable is now becoming more important, since parts of the region, both inland and offshore, have high potential for the production of wind energy (Balog et al. 2016; Onea et al. 2016) (*Chapter 3.3*).

Observation-based studies of winds over the Mediterranean are rare, and depend on the availability of homogenized and long time series. In most regions, wind trends were found non-monotonic over the past decades and, concrete conclusions are difficult to be established (Pirazzoli and Tomasin 2003; Vautard et al. 2010; Azorin-Molina et al. 2014). An additional source of information are reanalysis datasets, but robust trends have been identified over only a few regions in the Mediterranean (Nissen et al. 2010; Donat et al. 2011; Bett et al. 2013). Climate model simulations over historical periods can also be used in assessing and understanding past trends (Knippertz et al. 2000).

Despite the uncertainties in future projections (Shepherd 2014; Belušić Vozila et al. 2019), there is a general agreement for a limited wind speed reduction over most of the Mediterranean, with the exception of the Aegean Sea and north eastern land areas (*Fig. 2.3, Section 2.2.8*) (*medium confidence*) (Somot et al. 2006; McInnes et al. 2011; Dobrynin et al. 2012; Planton et al. 2012; Belušić Vozila et al. 2019). Changes in the local winds (such as Bora, Mistral, Tramontane, Sirocco and Etesians) may have more complex responses involved, depending on the changes in their underlying feedbacks (Grisogono and Belušić 2009; Ulbrich et al. 2012). Regional projections over the Adriatic reveal strong sensitivity in the climate change signal of the local Bora and Sirocco winds (Belušić Vozila et al. 2019). In particular, the frequency of winter Bora events is projected to increase while the frequency of Sirocco events is expected to decrease. Overall, the mean wind speed during Bora and Sirocco events is expected to be reduced, with the exception of Bora in northern Adriatic. RCM projections of Mistral and Tramontane show small changes in the former and significant decrease in the frequencies of the latter (Obermann-Hellhund et al. 2018). Etesian winds over the Aegean Sea is one of the few exceptions since increases in the wind speed are expected for the future (Ezber 2018; Dafka et al. 2019). In general, RCM projections have the



**Figure 2.3 | Projected changes (%) in surface wind speed** based on Med-CORDEX simulations [Ruti et al. 2016] for the end of the 21st century (2071-2100) relative to the period 1961-1990, for pathways RCP4.5 (top panel) and RCP8.5 (bottom panel). Seven RCMs were used for RCP4.5 and eight RCMs were used for RCP8.5. Dotted areas show differences that passed a 95% significance test.

tendency to simulate decrease of the wind energy density over the Mediterranean, with the exception of the Aegean Sea [Hueging et al. 2013; Tobin et al. 2015; Moemken et al. 2018].

Regional climate simulations indicate changes in wind speed over land regions as well (Fig. 2.3). Most pronounced changes, consistent for both RCP4.5 and RCP8.5 pathways, are an increase of wind speeds over the Balkans and a decrease over the Alps.

### 2.2.3 Radiation, clouds and aerosols


The amount of solar radiation reaching the Earth's surface is a key determinant of the spatio-temporal variations of climate on our planet and is the ultimate energy source for all processes relevant for climate and life. The main factors explaining the variability of surface solar radiation over the Mediterranean region for different time scales (daily, seasonal, interannual, past and future trends) are clouds [Pyrina et al. 2015] and aerosols [Nabat et al. 2015a, 2015b]. The daily variability in clouds and aerosols is strongly influ-

enced by weather regimes [Rojas et al. 2013; Nabat et al. 2020] and climate oscillations such as the North Atlantic Oscillation, NAO [Chiacchio and Wild 2010].

Aerosols in the Mediterranean come from various and numerous sources [Lelieveld 2002], both natural, notably dust and sea-salt, and anthropogenic, notably sulfates, nitrates and black carbon [Section 2.3]. Their interactions with radiation and clouds are essential in understanding climate in this region.

#### 2.2.3.1 Observed change in surface radiation

The long-term solar radiation records taken at widespread locations around the globe underwent substantial multidecadal variations, characterized by a reduction of surface solar radiation from the 1950s to the 1980s, known as “global dimming” and a partial recovery, thereafter, referred to as “brightening” [Wild 2009, 2012; Wild et al. 2017]. This dimming/brightening pattern is also observed



**Figure 2.4 | Past and future evolution of Surface Downwelling Shortwave Radiation (RSDS in  $W m^{-2}$ ), Total Cloud cover (CLT in %) and Aerosol Optical Depth (AOD) from 1950 to 2100 averaged over the Mediterranean area in CMIP5 Global Climate Models (GCM) and 12-km Euro-CORDEX Regional Climate Models (RCM) ensembles. Figures based on Bartok et al. (2017), redrawn and extended using published datasets.**

in the Mediterranean area (*very high confidence*), both in all-sky and clear-sky conditions, documented with many ground-based and satellite observations after careful data quality assessment and homogenization (Sánchez-Lorenzo et al. 2007, 2013, 2017; Zerefos et al. 2009; Kambezidis et al. 2016; Manara et al. 2016; Alexandri et al. 2017; Pfeifroth et al. 2018), as well as climate simulations (Folini and Wild 2011; Zubler et al. 2011; Nabat et al. 2014). The surface solar radiation trends averaged over the Mediterranean have been estimated in climate model simulations between  $-3.5$  and  $-5.2 W m^{-2}$  per decade for the dimming period (1953-1968), against for the brightening period between  $+0.9$  and  $+4.6 W m^{-2}$  per decade in 1989-2004 (Folini and Wild 2011) and  $2.3 W m^{-2}$  per decade in 1980-2012 (Nabat et al. 2014). The surface solar radiation anomalies calculated by global climate models for the periods 1950-1969 and 2000-2019 against the reference period of 1980-1999 are positive, showing respectively the dimming and the brightening effects, since the reference period refers to the period where surface solar radiation was the lowest (Fig. 2.4).

In parallel to the brightening period, a decrease in aerosol loads has been observed since 1980 both in ground-based stations (Li et al. 2014) and in satellite data (Floutsi et al. 2016). This decrease, corresponding to a trend in aerosol optical depth of  $-0.03 decade^{-1}$ , is mainly due to reductions in anthropogenic emissions, leading to a decrease in anthropogenic aerosol concentrations such as sulfate (Nabat et al. 2013). These aerosol trends have been shown to be the main explanation of the dimming-brightening phenomenon in the Mediterranean area (*high confidence*) through attribution model studies (Folini and Wild 2011; Zubler et al. 2011; Nabat et al. 2014), and with the direct aerosol effect responsible for about 80% of the simulated brightening. This phenomenon is qualitatively reproduced by most GCMs, but only by a few regional climate models, due to different treatments of aerosols in models (Fig. 2.4). The evolution of natural aerosols is more uncertain over the same period.

Concerning clouds, a decrease in cloud cover of 0.63% per decade since the 1970s has also been detected from different observations datasets over

the Mediterranean area (Sánchez-Lorenzo et al. 2017). This trend mainly concerns low and mid cloud layers (Kambezidis et al. 2016) (*medium confidence*). The spread between models that capture these trends is high, because of their difficulties to capture cloud characteristics (Fig. 2.4). Clouds may also have played a significant role in the past trend of surface solar radiation, at least locally (*low confidence*). Stronger positive trends in surface solar radiation are detected in spring over western Mediterranean Basin, explained by a decrease in cloud optical depth for this season over this basin (Kambezidis et al. 2016), and despite an averaged positive trend of surface solar radiation observed over the eastern basin, more uncertainty exists due to the lack of observations of both clouds and aerosols (Alexandri et al. 2017).

### 2.2.3.2 Projected change in surface radiation

In future climate projections, anthropogenic aerosol loads over the Mediterranean are expected to keep on decreasing (*high confidence*) because of decreases in anthropogenic emissions in Europe (Shindell et al. 2013). The decrease is expected to be more pronounced in the near future with an aerosol optical depth anomaly between -0.12 and -0.03 (5-95% uncertainty range) for the 2040-2059 period with respect to the reference period 1980-1999 (Fig. 2.4), and will slow down in the far future (between -0.18 and -0.04, 2080-2099 vs 1980-1999). The evolution of natural aerosols is more uncertain, due to current unknown future evolution of the desert dust (Section 2.3.2).

Total cloud cover is also expected to decrease during the 21st century over the Mediterranean (*medium confidence*) (Boé and Terray 2014; Enriquez-Alonso et al. 2016; Bartók et al. 2017; Hentgen et al. 2019). This is consistent with the northward expansion of the Hadley cell (Sánchez-Lorenzo et al. 2017; D'Agostino et al. 2020) (Section 2.2.2) and with enhanced lower tropospheric drying (Hentgen et al. 2019). The expected anomaly in cloud cover for the mid-21st century ranges from -4.9 to -0.2% in the RCP8.5 (5-95% uncertainty range, Fig. 2.4), because of the difficulty of models to capture the spatial variability of the cloudiness evolution (Bartók et al. 2017).

As projected by GCMs, surface solar radiation is expected to continue increasing in the 21st century, especially in the near future (*medium confidence*). The anomaly is between 0.6 and 7.7 W m<sup>-2</sup> for the period 2020-2039 in the RCP8.5 (5-95% uncertainty


range), in line with the decrease in anthropogenic aerosols (Boé et al. 2020; Gutiérrez et al. 2020) (Fig. 2.4). However, this evolution is not shared by all RCMs (Bartók et al. 2017; Gutiérrez et al. 2020). Only regional climate models which consider aerosol dynamics simulate the increase in surface solar radiation as the global models (Boé et al. 2020; Gutiérrez et al. 2020). This increase in future surface solar radiation is reinforced by an expected decrease in cloud cover (Enriquez-Alonso et al. 2016), despite a compensational effect of increased absorption in clear sky conditions due to higher water vapor content in the atmosphere (Haywood et al. 2011) (Fig. 2.4).

## 2.2.4 Temperature and related extremes

### 2.2.4.1 Observed temperature changes

Climate reconstructions, ground-based observations, reanalysis and remote-sensing datasets all corroborate the transition to warmer conditions during the 20th century and that warming has accelerated during the last decades (*high confidence*). Basin-wide, annual mean temperatures are now 1.5°C above late 19th century levels (Box 2.1, Fig. 2.33). Particularly after the 1980s, regional warming has accelerated and increases at a higher rate than the global average (Lelieveld et al. 2012; Lionello et al. 2012a; Zittis and Hadjinicolaou 2017; Cramer et al. 2018; Lionello and Scarascia 2018; Zittis et al. 2019). These studies present a strong consensus that the recent observed warming is robust throughout the region analysis, though magnitude and level of significance of the observed temperature trends in the Mediterranean varies depending (a) on the region, country or station under consideration, (b) on the type of data set investigated and (c) on the season and period of analysis.

Solar forcing and large volcanic eruptions are found to have a strong influence on the Mediterranean temperature variability over the last centuries (Trouet 2014). A combination of climate reconstructions, documentary sources and observed data suggests that looking at the long-term timescale (e.g., over the last 500 years), warm periods are not exceptional for the Mediterranean, which is characterized by a sequence of warming-cooling cycles (Luterbacher and Xoplaki 2003; Camuffo et al. 2010; Lelieveld et al. 2012). A study of summer temperature since Roman times shows that although the mean 20th century European (including the northern part of the Mediterranean Basin) was not significantly warmer than some



**Figure 2.5 | Observed temperature trends (Left panels) and time-series of temperature over land for the Mediterranean based on the Climatic Research Unit (CRU) (Harris et al. 2020) gridded observations. Time-series refer to the whole Mediterranean as defined in the left panels and for three Mediterranean sub-regions (Fig. 2.1): North (FR, ALBA), Central (IB, CM, AN) and South Mediterranean (WM, EM, LE).**

earlier centuries, there are no earlier 30-year periods found to be warmer than the most recent 3 decades (Luterbacher et al. 2016).


Recent climatic trends are clear, particularly after the 1980s. Over these last decades, according to different type of observations, significant positive trends of the order of 0.1-0.5°C per decade have been identified (Fig. 2.5) (Nasrallah and Balling 1993; Saaroni et al. 2003; Feidas et al. 2004; Brunetti et al. 2006; El Kenawy et al. 2009; Tanarhte et al. 2012; Lelieveld et al. 2012; Lionello 2012; Donat et al. 2014; Mariotti et al. 2015; Lionello and Scarascia 2018; Bilbao et al. 2019). In addition, for parts of the basin, there is some evidence that the diurnal temperature range has also changed (Price et al. 1999; Bilbao et al. 2019; Sun et al. 2019).

Besides mean values, hot and cold extremes have also become warmer, while in particular there is strong evidence and consensus that heat waves have become more frequent and severe. Various climatic indicators show significant increasing trends of extreme heat events characteristics (e.g., duration, frequency and intensity). The number of warm and tropical nights has also increased over most Mediterranean locations including Iberia,

north Africa, Italy, Malta, Greece, Anatolia and the Levant (Kostopoulou and Jones 2007; Bartolini et al. 2008; Kuglitsch et al. 2010; El Kenawy et al. 2011; Galdies 2012; Donat et al. 2014; Filahi et al. 2015; Lelieveld et al. 2016; Ceccherini et al. 2017; Nashwan et al. 2018; Tolika 2019). Parts of the region were impacted by some of the most severe record-breaking weather events of the last decade, mainly related with summer heat extremes (Coumou and Rahmstorf 2012). Considering only winter, some studies that suggest a different behavior of hot and cold extremes between the eastern and western parts of the Mediterranean, with negative temperature trends in the former and positive trends in the latter (Hertig et al. 2010; Efthymiadis et al. 2011), but these studies do not include the most recent warm decades.

### 2.2.4.2 Future temperatures

According to future projections, the greater Mediterranean Basin is among the most responsive regions to global warming. Previous studies have identified the region as one of the most prominent climate change hot spots (Giorgi 2006; Lionello et al. 2006; Giorgi and Lionello 2008; Diffenbaugh and Giorgi 2012; Lionello 2012; Lionello and Scarascia




**Figure 2.6 | Projected changes in annual temperature between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean of EURO-CORDEX 0.11° simulations for pathways RCP2.6 (top panels) and RCP8.5 (bottom panels).**

2018). Multi-model ensembles of climate simulations indicate that widespread warming will almost certainly occur in the Mediterranean in the 21st century (*high confidence*), though, climate models might overestimate actual values in warm and dry conditions (Boberg and Christensen 2012).

The warming level of warming strongly depends on the reference period definition, the future time horizon and the level of greenhouse gas forcing (Christensen et al. 2007; Giorgi and Lionello 2008; Collins et al. 2013; Dubrovský et al. 2014; Jacob et al. 2014; Mariotti et al. 2015; Ozturk et al. 2015; Lionello and Scarascia 2018; Zittis et al. 2019). A quantitative estimation based on state-of-the-art EURO-CORDEX regional simulations is presented in Table 2.1 and Fig. 2.6-2.7. Over land regions, a robust and significant warming of the range of 0.9-5.6°C (with respect to the reference period 1980-1999) is suggested for the future. The robustness and significance of the climate signal is much higher for air temperature rather than other variables such as precipitation (Knutti and Sedláček 2012; Lelieveld et al. 2016). There are strong indications and a general consensus that regional warming will continue faster than the global average and will exceed the global mean value by 20% on an annual basin and 50% in summer (*high confidence*) (Vautard et al. 2014; Dosio and Fischer 2018; Lionello and Scarascia 2018; Nikulin et al. 2018). Daytime temperatures are expected to increase more than nighttime temperatures, indicating an increase of the amplitude of the diurnal temperature range (Lionello and Scarascia 2018).

Changes in the occurrence of extreme events closely follow changes in inter-annual variability. Therefore, such changes can be also considered as a proxy measure of seasonal extremes (Schär et al. 2004; Giorgi 2006). The intensity of extreme temperature is projected to increase more rapidly than the intensity of more moderate temperatures over the continental interior due such increases in temperature variability (Beniston et al. 2007).

Projected changes in extreme temperature indicators suggest that the frequency and severity of heat waves will increase (*high confidence*) (Diffenbaugh et al. 2007; Goubanova and Li 2007; Giorgi and Lionello 2008; Fischer and Schär 2010; Diffenbaugh and Giorgi 2012; Sillmann et al. 2013; Russo et al. 2014; Jacob et al. 2014; Kostopoulou et al. 2014; Zittis et al. 2016; Lelieveld et al. 2016; Ouzeau et al. 2016; Lionello and Scarascia 2020). According to projections for a business-as-usual scenario, summer daily maximum temperature is expected to increase up to 7°C by the end of the 21st century in comparison with the recent past (Sillmann et al. 2013; Lelieveld et al. 2016). Besides warmer daytime temperature maxima, parts of the Mediterranean will likely face an increase of more than 60% in the number of tropical nights. Increase of warm temperature extremes will be dramatic particularly in summer and with a 4°C global warming almost all nights will be warm and there will be no cold days (Sillmann et al. 2013; Dosio and Fischer 2018; Lionello and Scarascia 2020).



**Figure 2.7 | Time-series of simulated mean annual (top panel), summer (middle panel) and winter (bottom panel) temperature averaged over the Mediterranean based on EURO-CORDEX 0.11° simulations** for historical times (black curve) and future pathways RCP2.6 (blue curve) and RCP8.5 (red curve). Solid lines indicate the ensemble means and shaded areas the spread of the simulations. Box-plots represent the averages over the decade 2091-2100 in terms of model spread.

Warming is projected to be milder in winters and much stronger during summers. This is mainly attributed to land-atmosphere interactions and the transition to drier conditions (Seneviratne et al. 2006; Jaeger and Seneviratne 2011; Quesada et al. 2012; Zittis et al. 2014). Another important feedback, particularly for southern Mediterranean, is the coupling of longwave radiation between the desert soil surface and lower atmosphere which


amplifies warming and intensifies the summertime heat low over the Sahara (Cook and Vizy 2015; Evan et al. 2015). The exceptional summertime warming over parts of the region is also likely associated with a thermal low, which is explained by the widening of the Persian trough that extends from South Asia to the eastern Mediterranean, and is projected to expand westward and combine with the intensifying thermal low over the Sahara (Lelieveld et al. 2016).

## 2.2.5 Precipitation, related extremes and the water cycle


### 2.2.5.1 Observed trends in precipitation

Observed precipitation trends during the full or portions of the 20th century up to present day, covering the full or portions of the Mediterranean Basin, are available in gridded format from various sources, such as CRU, UDEL, E-OBS, EURO4M. Annual, DJF (December-January-February) and JJA (June-July-August) precipitation trends from the CRU dataset for different time periods, 1950-2018 and 1980-2018 are shown in Fig. 2.8. Fig. 2.9 shows the temporal evolution of land precipitation (1901-2018) averaged over the full Mediterranean area and its northern, central and southern portions (defined as the sum of the three northern, central and southern regions of Fig. 2.8, respectively).

The sign of the observed precipitation trend exhibits pronounced spatial variability and depends on the time period and season considered (Fig. 2.8). For example, the period 1950-2018 shows a prevailing decreasing trend over most of the Mediterranean Basin of annual and winter precipitation, which is reversed over large portions of the basin if we only consider the period 1980-2018. This is because of the marked multidecadal variability of precipitation in the Mediterranean, which may actually mask trends induced by greenhouse gas emissions. The prominent role of multidecadal variability is also evident when precipitation is regionally averaged (Fig. 2.9). In this case the most evident trend is a decrease of winter precipitation over the central and southern portions of the basin since the second half of the 20th century. Overall, because of the marked multidecadal variability of precipitation and the small magnitude of trends, the confidence in the detection of trends from greenhouse gas emissions for the historical past is low (Lelieveld et al. 2012; Lionello et al. 2012a; Peña-Angulo et al. 2020; Vicente-Serrano et al. 2020).



**Figure 2.8 | Observed annual, DJF, JJA precipitation trends from the CRU dataset.** Left and right panels consider the 1950-2018 and 1980-2018 periods, respectively.



**Figure 2.9 | Time series of annual, DJF, and JJA precipitation over land from the CRU dataset.** Time-series refer to the whole Mediterranean as defined in the left panels and three Mediterranean sub-regions (Fig. 2.1): North (FR, ALBA), Central (IB, CM, AN) and South Mediterranean (WM, EM, LE).



### 2.2.5.2 Future precipitation

#### Mean precipitation changes


21st century precipitation projections for the Mediterranean region have been produced based on GCM and RCM ensembles of experiments. Analyses of GCM projections over the Mediterranean region have been conducted for CMIP3 (Giorgi and Coppola 2007; Giorgi and Lionello 2008; Mariotti et al. 2008) and CMIP5 (Mariotti et al. 2015; Lionello and Scarascia 2018). Several generations of RCM-based projections for the EURO-Mediterranean region are also available from projects such as PRUDENCE (Déqué 2007), ENSEMBLES (Déqué et al. 2012) and EURO-CORDEX (Jacob et al. 2014). In addition, projections based on coupled regional atmosphere-ocean models have been conducted as part of the CIRCE (Gualdi et al. 2013) and MED-CORDEX (Ruti et al. 2016) projects.

A consistent dominant signal emerges from these projections, consisting of a predominant drying through the entire Mediterranean Basin in the warm seasons (April through September, with largest magnitude in JJA), drying in most central and southern areas along with wetting in the northernmost regions (e.g., the Alps) in the winter season (*medium confidence*). This large-scale pattern of change is illustrated for the Euro-CORDEX dataset, at the annually averaged scale, in Fig. 2.10 for the RCP2.6 and RCP8.5 scenarios

and different future time slices. Table 2.1 provides quantitative values of precipitation change for different scenarios and model ensembles over the Mediterranean region.

In general, the patterns of change intensify in magnitude from the near future to the far future time slices and from the low to high greenhouse gas emission scenarios, i.e., they intensify with the anthropogenic forcing and resulting global warming. As a result, for example, at the Mediterranean scale, the CMIP5 ensemble yields a decrease of annual precipitation over the Mediterranean area of about 4% per degree of global warming (Lionello and Scarascia 2018).

The magnitude and pattern of precipitation decrease vary widely across models. For example, the summer precipitation reduction in the CMIP3 and CMIP5 datasets for the high-end greenhouse gas emission scenarios (roughly equivalent to the RCP8.5) varied from less than 10% to over 40% across models (IPCC 2007; Giorgi and Lionello 2008; Lionello and Scarascia 2018). Although qualitatively consistent, different ensembles show different sensitivities over the Mediterranean. The CMIP5 GCM ensemble produced a less pronounced summer drying than the CMIP3 one, when expressed in terms of change per degree of global warming (IPCC 2013b). RCM-based projections, e.g., as part of Euro-CORDEX yield a lower drying than GCM-based ones, with reduced areas



**Figure 2.10 | Maps showing EURO-CORDEX-based change in annual, winter and summer precipitation change, for the RCP2.6 and RCP8.5 scenarios and the near-future, mid-term and far-future with respect to the reference period.**


of precipitation decrease in the future compared to the GCMs. Considering a range of global warming of 0.9-5.6°C (Section 2.2.4.2) at the end of the 21st century (with respect to the reference period 1980-1999) and a decrease of 4% per degree of global warming, this gives a reduction between 4 and 22% for the Mediterranean annual precipitation (on land).

Fig. 2.11 shows the temporal evolution of Mediterranean scale precipitation (RCP2.6 and RCP8.5) from 1950 to 2100 in the EURO-CORDEX dataset, and presents both mean and inter-model range of data. The large inter-model spread includes some positive values and two scenarios start to separate, at least in an ensemble mean sense, only after the mid of the 21st century. Therefore, while it is possible to assess that precipitation will likely decrease over the Mediterranean Basin, at least under the higher end scenarios, it is difficult to assign robust quantitative values, especially at sub regional scale.

The uncertainty in projections is even larger as we move to the local scale because of the effects of local forcings, e.g., topography and coastlines. For example, focusing on the high elevations of the Alpine chain, Giorgi et al. (2016) found that the RCM projections at 12 km resolution exhibit an increase of summer precipitation in areas where the GCMs project a decrease. This is due to the occurrence of increased convection related to high elevation warming and heating. As another example, increases of cold season precipitation in the upwind side of mountain chains and decreases in the lee side have been found by the high-resolution simulations of Gao et al. (2006) in response to the topographically-forced precipitation shadowing effect. These results were confirmed by the analysis of the EURO-CORDEX ensemble projections (Kotlarski et al. 2019). In other words, high resolution RCM projections suggest that care needs to be taken when extending large scale patterns to the local scale, since local scale changes can be heavily affected by topography and coast lines.

**Variability and extremes**

Several studies have assessed changes in interannual variability of precipitation over the Mediterranean region in GCM-based projections, mostly using as a measure of variability the coefficient of variation (i.e., the interannual standard deviation divided by the mean), which removes the strong dependence of precipitation standard deviation from the mean (Räisänen 2002;



**Figure 2.11 | Time-series of simulated mean annual (top panel), wet (middle panel) and dry (bottom panel) season over the Mediterranean land areas based on EURO-CORDEX 0.11° simulations** for historical times (black curve) and future pathways RCP2.6 (blue curve) and RCP8.5 (red curve). Solid lines indicate the ensemble means and shaded areas the spread of the simulations. Box-plots represent the averages over the decade 2091-2100 in terms of model spread.

Giorgi and Bi 2005; Giorgi and Coppola 2009; Giorgi et al. 2019). They all found a prevailing increase in precipitation variability over the Mediterranean, especially over areas showing strong precipitation decreases, which thus appears to be a robust response in global climate projections. However, this result may not apply if the metrics used to measure variability is the standard deviation.

Giorgi et al. (2011, 2014) carried out an analysis of changes in different hydroclimatic indices from an ensemble of CMIP5 projections and, on an annual basis, consistently found an increase in mean daily precipitation intensity and 95th percentile of daily precipitation, a decrease in precipitation frequency and wet spell length and an increase in the number of dry days and dry spell length (*medium confidence*). The hydroclimatic intensity index introduced by Giorgi et al. (2011), which is essentially the product of precipitation intensity and mean dry spell length, shows a consistent increase throughout the Mediterranean Basin. These conclusions based on GCM projections, were essentially confirmed by an analysis of high-resolution EURO-CORDEX projections by Jacob et al. (2014), except for a slight decrease of 95th percentile daily precipitation over some areas of the Iberian, Italian and Hellenic peninsulas and southern France in summer. A recent study (Lionello and Scarascia 2020), based on CMIP5, shows that global warming will further increase the existing difference in intensity of precipitation and hydrological extremes between North and South Mediterranean areas. Both the daily precipitation intensity the total precipitation during extreme events are already larger in the North than in the South Mediterranean areas, and differences will increase with global warming. The projected increase of dry spell length is larger in the south than in the North Mediterranean (*medium confidence*).

In conclusion, (*high confidence*) both global and regional climate projections indicate a predominant shift towards a precipitation regime of higher interannual variability (when measure by the coefficient of variation), higher intensity of precipitation and greater extremes (especially in winter, spring and fall, but not in the southern areas), decreased precipitation frequency and longer dry spells (especially in summer). This hydroclimatic response to global warming is greater for the RCP8.5 than the RCP2.6 scenario and for the far future vs. the near future time slices (*high confidence*).

### 2.2.5.3 Changes in evaporation, net water losses over sea and over land

Evaporation in the Mediterranean not only provides moisture locally, but also results in a net export of water to neighboring areas, primarily to the South and East (Mariotti et al. 2002a; Nieto et al. 2006). The Mediterranean Sea is the dominant regional evaporation source, and changes in Mediterranean Sea evaporation impact the

sea's water, salt and heat budgets. Large-scale internal climate variability and greenhouse gas forced global change have been primary drivers of Mediterranean evaporation changes during the 20th century and into the 21st century, while local processes have acted to modulate those effects (Mariotti and Dell'Aquila 2012). Overall, the net surface water loss (evaporation minus precipitation over the sea) has increased over most of the Mediterranean surface, mainly due to a decrease of precipitation during the period 1960-1990 and a strong evaporation increase since the mid-seventies due to local warming (Mariotti 2010; Sevault et al. 2014; Mariotti et al. 2015; Skliris et al. 2018). The freshwater discharge due to the river runoff has also decreased (Ludwig et al. 2009). Projected regional warming trends point to continuing increases in Mediterranean Sea evaporation, land drying in southern areas during summer and a net regional water loss.

Observations-based estimates of Mediterranean Sea evaporation from the OAFflux Programme starting in 1958 point to decadal variations with a minimum around 1965-1975, and an overall positive trend of about 10% decade<sup>-1</sup> (0.06 mm day<sup>-1</sup> decade<sup>-1</sup>) (Mariotti 2010). Since the mid-1970s, there is a substantial evaporation increase (0.1-0.2 mm day<sup>-1</sup> decade<sup>-1</sup>) with a tendency toward higher rates of increase during the 1990s. Much of the evaporation increase since the mid-1970s has been in early winter, especially in the Ligurian Sea, Adriatic Sea, and southeastern Mediterranean. The evaporation increase has resulted in a rate of increase in freshwater fluxes during 1979-2006 estimated in the range of 0.1-0.3 mm day<sup>-1</sup> decade<sup>-1</sup>. Increases in sea surface temperatures have primarily driven these evaporation changes via changes in the surface humidity gradient. Based on OAFflux data, the estimated Mediterranean mean rate of evaporation change in relation to the warming is about 0.7 mm day<sup>-1</sup> K<sup>-1</sup> (or 25% K<sup>-1</sup>) over the period of 1958-2006. An increase in net Gibraltar water flux to compensate for the overall increase in fresh water loss has been derived (Fenoglio-Marc et al. 2013).

For the land surrounding the Mediterranean Sea, past evapotranspiration changes are regionally and seasonally dependent and largely follow precipitation trends, since soil moisture availability is a primary limiting factor. Increasing soil-moisture limitations seem to have driven recent global evapotranspiration decline and increased drought tendencies over the Mediterranean region (Sheffield and Wood 2008; Vicente-Serrano et al. 2014; Samaniego et al. 2018). Evapotranspiration

estimates from French National Centre for Meteorological Research (CNRM) (Douville et al. 2013) display a tendency for evapotranspiration to increase during winter since the 1970. For the summer there is a progressive decrease (Mariotti et al. 2015).

The future evolution of the Mediterranean Sea physical characteristics is strongly related to the evolution of the air-sea and land-sea exchanges of water and heat. For the Mediterranean Sea, the net surface water loss by the sea is constituted by the combination of the evaporation over the sea, the precipitation over the sea, the river runoff and the Bosphorus Strait net transport. Increase in the net surface water loss by the sea is expected in the future due to a decrease in precipitation and in river runoff and an increase in evaporation (Mariotti et al. 2008, 2015; Sánchez-Gomez et al. 2009; Elguindi et al. 2011; Dubois et al. 2012; Planton et al. 2012; Adloff et al. 2015). Relative to the 20th century, this increase ranges from +8 to +35% for the mid-21st century (2020-2049) and from +20 to +60% at the end of the 21st century (2070-2099) in the medium-range A1B socio-economic scenarios (Planton et al. 2012).

To a first order, CMIP5 projections are largely similar to those based on CMIP3 (Mariotti et al. 2008) and consistent with those based on regional model downscaling (Sánchez-Gomez et al. 2009; Dell'Aquila et al. 2018). By 2071-2098, the Mediterranean Sea evaporation is projected to increase during all seasons and especially in winter (projected annual-mean increase is  $0.25 \pm 0.08$  mm day<sup>-1</sup>) (Mariotti et al. 2015). Note that future change in the Nile freshwater inflow remains unknown due to the impossibility so far to accurately model the influence of regional water- and land-use anthropogenic activities on its past and future evolution (Somot et al. 2006; Dubois et al. 2012).

Over land, evapotranspiration projections present mixed changes, with a precipitation-driven increase in winter over Northern areas, and a decrease in summer over many land areas, especially over Spain, western Northern Africa and Turkey. Evapotranspiration increase will be also driven by increase of atmospheric evaporative demand (Vicente-Serrano et al. 2015). These evapotranspiration changes have been linked to a projected northward expansion of the Mediterranean land type (Alessandri et al. 2015) and regional surface vegetation changes (Anav and Mariotti 2011).

Changes in precipitation and evaporation over the Mediterranean Basin will lead to changes in

drought occurrence. Drought can be of different types, such as meteorological, hydrological and agricultural drought, which can often be difficult to separate. Here we focus on meteorological drought, essentially measured by indices of monthly, seasonal up to annual precipitation deficits, such as the precipitation index (PI) or the standardized precipitation index (SPI). The Mediterranean Basin, is impacted by frequent drought episodes due to the strong inter-annual variability of rainfall in this region, and a trend towards drier conditions and increased meteorological drought occurrence after the 1970s over the Mediterranean Basin was found based on analyses of observations (Vicente-Serrano et al. 2011; Hoerling et al. 2012; Spinoni et al. 2015; Caloiero et al. 2018). Due to the pronounced interannual and decadal variability of Mediterranean precipitation, the robustness of this result needs to be confirmed, and may differ for different areas of the Mediterranean.

Concerning projections, since most model simulations indicate a trend towards drier conditions over the Mediterranean, especially in the warm season and over the southern areas, it is expected that the frequency and intensity of meteorological drought will increase under warmer climates. This has been confirmed (*high confidence*) by extensive analyses of precipitation projections with both global and regional climate models (Giorgi and Lionello 2008; Mariotti et al. 2008; Dai 2013; Dubrovský et al. 2014; Spinoni et al. 2015, 2018; Stagge et al. 2015; Quintana-Seguí et al. 2016; Naumann et al. 2018; Lionello and Scarascia 2020).

## 2.2.6 The cryosphere

### 2.2.6.1 Observed trends in the cryosphere

After the peak of the "little ice age" (~1,400-1,860 AD, Ivy-Ochs et al. 2009) increasing summer and mean annual air temperature led to a dramatic reduction in the area and volume of glaciers across high mountains of the Mediterranean (Hughes 2018). Short glacier readvances were observed in the 1890s, 1920s, 1970s and 1980s (Zemp et al. 2008). Deglaciation rate generally accelerated in recent decades (Rabatel et al. 2013), although the patterns of glacier retreat were complicated by the sensitivities of glaciers to different climatic regimes (Hughes 2018). As glacier retreats to cirque headwalls, it becomes more dominated by local topo-climatic controls, especially avalanching snow. Nevertheless, a complete loss of glaciers in some low-latitude mountain ranges has already

occurred (Rabatel et al. 2013), accompanied by a shorter duration of seasonal snow cover (Brown and Mote 2009). Several small cirque glaciers existing in the southern Dinaric Alps, Balkan Peninsula, Turkey, Pyrenees, Sierra Nevada and the Apennines disappeared across the 20th century and in the last decades (Hughes 2018). In the Alps, glaciers covered 4,470 km<sup>2</sup> in 1850, 2,909 km<sup>2</sup> in the 1970s and 2,270 km<sup>2</sup> in 2000, meaning a 50% loss from 1850 to 2000 (Zemp et al. 2008). Few very small glaciers still exist in mountains of Montenegro and Albania. Elsewhere, perennial ice and snow patches still survive and attest to how close some Mediterranean mountains are to supporting small glaciers even where the equilibrium line altitude (ELA) is located above the highest peaks (Hughes 2018). ELA raised by about 170 m in the western Alps over the period 1984-2010 (Rabatel et al. 2013) while in the southeastern Alps change in the ELA was in the order of about +250 m between the 1980s and 2010 (Colucci and Žebre 2016).

Temperature increase led to a shift of periglacial processes to higher elevations as well as degradation of mountain permafrost in high mountain environments (Oliva et al. 2018). In the western and central Mediterranean, permanently frozen ground is now rarely found below 2,500 m. Alpine permafrost belt is detected above 2,630 m in northern aspects and 2,800 m in southern ones and in the Pyrenees, above 2,400 m in the Southern Alps, above ~2,350 m on Rila Mountain and ~2,700 m on Mount Olympus and above 2,800-3,400 m in north-eastern Turkey and central Anatolia (Oliva et al. 2018). No permafrost belt is found in the highest mountains in southern Europe (Sierra Nevada) and northern Africa (Atlas) where permanent frozen conditions are only found in the form of isolated patches at the highest elevations at 3,000-3,100 m (Oliva et al. 2016) and 3,800 m (Vieira et al. 2017), respectively. Certain climate conditions (i.e., reduced snow cover) can favour the presence of permafrost patches at relatively low elevations in the Central Apennines or by lithological conditions (i.e., volcanic sediments, karst lithology), as detected in the highest active European volcano (Mt. Etna) at elevations above 2,900 m (Maggi et al. 2018) or in limestone dominated mountains across the Mediterranean (Colucci and Guglielmin 2019).

### 2.2.6.2 Future conditions in the cryosphere

Mountain glaciers in the Mediterranean region are projected to continue losing mass in the 21st century until complete disappearance of most

mountain glaciers by the end of the century (*very high confidence*). A recent multi-model projection exercise (GlacierMIP, Hock et al. 2019) indicates that relative volume losses by 2100 (average of model runs  $\pm 1$  standard deviation) are of the order of  $69 \pm 19\%$  for RCP2.6 and  $93 \pm 10\%$  for RCP8.5. This indicates that, even under scenarios with strong reduction in greenhouse gas emissions, only glaciers at the highest elevation will persist at the end of the 21st century. For mid-century, changes depend far less on the climate scenario, with reductions of the order of  $50 \pm 20\%$  for RCP2.6 and  $60 \pm 20\%$  for RCP8.5.

Projected changes of the mountain snow cover are studied based on climate model experiments, either directly from GCM or RCM output, or following downscaling and the use of snowpack models. Future changes in snow conditions are mostly driven by changes in meteorological drivers. The projections generally do not specifically account for future changes in the deposition rate of light absorbing particles on snow and associated changes in snow albedo. At lower elevation, under the current multi-annual mean rain/snow transition elevation, the water mass of the snow cover is projected to decline by 25% likely range: (10-40%, between the recent past period (1986-2005) and the near future (2031-2050), regardless of the climate scenario). By the end of the 21st century (2081-2100), reductions of up to 80% (likely range 50-90%) are expected under RCP8.5, 50% (likely range from 30-70%) under RCP4.5 and 30% (likely range 10-40%) under RCP2.6 (Beniston et al. 2018; Hanzer et al. 2018; Verfaillie et al. 2018). At higher elevations, projected reductions are smaller (*high confidence*), as temperature increases at higher elevations affect the ablation component of snow mass evolution (in particular, melt and sublimation), rather than the onset and accumulation component. The strong interannual variability of snow conditions is projected to remain a key feature of this cryospheric component throughout the 21st century (*high confidence*).

In the Mediterranean domain, permafrost is only located in the mountains, often patchy and confined to areas of rugged topography, including cliffs. In contrast to glacier and snow cover, climate projections of the ground thermal regime have not been performed in a comprehensive manner, using a cascade of climate models and impact models. Evidence stems from small-scale studies, but all studies points towards increased permafrost thaw in mountain environments, following surface air temperature changes (Marmy et al. 2016; Beniston et al. 2018). Future changes in mountain


permafrost have major implication for natural hazards (slope instabilities).

### 2.2.7 Ocean hydrology

The Mediterranean Sea can be considered as a laboratory of the global ocean (Lacombe 1990; Béthoux et al. 1999) as it shows many key and interesting oceanic physical processes, such the open-sea deep convection occurring in some areas (Gulf of Lion, South Adriatic, and the Cretan Sea) leading to the formation of cold and salty deep-water masses (Tsimplis et al. 2006; Schroeder et al. 2012). The dominance of evaporation in the Mediterranean Sea and intermediate water formation (in the Rhodes Gyre) leads to an anti-estuarine

thermohaline circulation with a surface layer with the Atlantic Water (AW, comparatively fresh and warm) and a layer with the Levantine Intermediate Water (LIW, very salty and comparatively cold) entering and exiting across the Gibraltar Strait, simultaneously.

In addition, the Mediterranean Sea is surrounded by various and complex topography channeling regional winds (Mistral, Tramontane, Bora, Meltem, Sirocco, Etesians) that define local circulations. The presence of complex coastlines, islands, narrow and shallow straits require adapted observation strategies and high-resolution modeling tools. Further, the Mediterranean Sea is also known to impact the Atlantic Ocean through the Mediter-



**Figure 2.12 | A) Local amplification factor of the Mediterranean sea surface warming using 1980-1999 as reference period.** Local sea surface warming values are divided by the basin-averaged warming value. Reddish (resp. blueish) colors mean that the area is warming more (resp. less) than the basin-average. The top panel is the ensemble mean of five Med-CORDEX coupled regional climate system models, further averaging warming rates of four 20-year long time periods (2000-2019, 2020-2039, 2040-2059, 2080-2099) and 2 scenarios (RCP4.5, RCP8.5). The bottom is based on the CMEMS observations [2000-2017]. **B) Fraction of the Mediterranean Sea surface (in %) experiencing a given sea surface temperature change value (in °C),** compared to the reference period [1980-1999] for various periods [2000-2019 in blue, 2020-2039 in orange, 2040-2059 in purple, 2080-2099 in red] and for the scenarios RCP4.5 and RCP8.5 using the envelope of the results of 5 Med-CORDEX coupled regional climate system models. The change in the CMEMS observations is added for the period 2000-2017 (in black). Information is first aggregated at the yearly scale.

anean Outflow Waters that flow into the Atlantic at about 1,000 m depth and are considered as a source of salt and heat for the Atlantic Ocean (Artale et al. 2006).

### 2.2.7.1 Observed change in marine waters

There is increasing evidence that Mediterranean water masses are becoming warmer and deep water masses saltier. This assertion is supported both by direct measurements (Béthoux and Gentili 1999; Rixen et al. 2005; Vargas-Yáñez et al. 2010, 2017) and by numerical simulations (Beuvier et al. 2010; Harzallah et al. 2018; Somot et al. 2018).

Since the 1980s upper layer temperature has increased (Rivetti et al. 2017; Vargas-Yáñez et al. 2017) as well as sea surface temperature (Marullo et al. 2010; Pastor et al. 2018), with acceleration since the 1990s (Macías et al. 2013). Since the beginning of the 1980s, the sea surface warming rate ranges between  $+0.29$  and  $+0.44^{\circ}\text{C decade}^{-1}$  on average over the whole Mediterranean Sea, depending on the studied period and on the reference data sets (Nabat et al. 2014; CEAM 2019; Darmaraki et al. 2019a). In the period 2000-2017 with respect to 1980-1999, all Mediterranean Sea areas show a positive yearly-mean sea surface temperature anomaly of at least  $+0.2^{\circ}\text{C}$ . The sea surface warming has not been uniform, but mostly bimodal (Fig. 2.12b) with stronger trends in the eastern basin (Adriatic, Aegean, Levantine and North-East Ionian Seas, Fig. 2.12a), where some areas warmed by  $+1.2^{\circ}\text{C}$ . Very local places in the Levantine Basin have warmed 50% more rapidly than the Mediterranean Sea average whereas a spot in the Ionian Sea has warmed 50% less than the basin average (Fig. 2.12a). Note that the climate models currently underestimate the observed sea surface warming (blue and black lines in Fig. 2.12b) (Nabat et al. 2014; Sevault et al. 2014; Dell'Aquila et al. 2018).

In the Mediterranean Sea, periods of abnormally warm sea surface, also called "marine heat waves" have become more frequent, more intense, spatially more extended and more severe over the last decades (Oliver et al. 2018; Darmaraki et al. 2019a). To illustrate this trend, the most severe marine heat waves detected since 1982 are 2003, 2012, 2015, and 2017 (Bensoussan et al. 2019). In addition, 14 marine heat waves occurred during the 2008-2017 10-year period whereas only 2 occurred during the 1982-1991 period (Darmaraki et al. 2019a). Contrary to sea surface temperature, a corresponding sea surface salinity evolution has


not been reported (Rixen et al. 2005; Sevault et al. 2014), except for specific locations (Ozer et al. 2017; Vargas-Yáñez et al. 2017).

Long-term trends in the Mediterranean Sea intermediate and deep hydrology have been detected, though they are affected by marked variability on decadal time scales, the Eastern Mediterranean Transient (EMT, Roether et al. 2007), Western Mediterranean Transition (WMT, Schroeder et al. 2016) and BiOS (Gačić et al. 2010) being probably the most known manifestations of this.

Since the mid 1990s the temperature and the salinity of the Levantine Intermediate Water (LIW) have increased by  $0.53^{\circ}\text{C}$  (Fig. 2.13, "Sicily Channel" panel) and  $0.13$  psu, i.e., with rates of  $0.024^{\circ}\text{C yr}^{-1}$  and  $0.006$  psu  $\text{yr}^{-1}$ , respectively (Schroeder et al. 2017). Such trends are at least one order of magnitude greater than those reported for the global ocean intermediate layer (Schroeder et al. 2017). The western basin deep waters are shown to have gradually increased their temperature and salinity since the 1950s (Rohling and Bryden 1992; Béthoux et al. 1998; Rixen et al. 2005; Marty and Chiavérini 2010), with an acceleration after the mid 1980s and an even stronger rate since 2005 due to an abrupt WMT (Marty and Chiavérini 2010; Borghini et al. 2014; Schroeder et al. 2016). Deep-water trends of  $0.04^{\circ}\text{C} \pm 0.001^{\circ}\text{C decade}^{-1}$  and  $0.015 \pm 0.003$  psu  $\text{decade}^{-1}$  (since 1961) have been reported by comparing time series of deep CTD (conductivity-temperature-depth) casts (Borghini et al. 2014) (Fig. 2.13, bottom panel).

Changes in the Mediterranean water mass characteristics have a signature also in the water outflowing from the Mediterranean Sea through the Strait of Gibraltar (Millot et al. 2006; Naranjo et al. 2017). Mooring observations collected since 2004 (Fig. 2.13, "Gibraltar Strait" panel) show a positive trend in temperature and salinity of  $7.7 \times 10^{-3}^{\circ}\text{C yr}^{-1}$  and  $0.63 \times 10^{-3}$  psu  $\text{yr}^{-1}$ , respectively (von Schuckmann et al. 2018). Since 2012 a noticeable increase of these trends is interpreted as the signal of the WMT (Naranjo et al. 2017). In addition, no significant changes in the strait transports (net exchange: Fenoglio-Marc et al. 2013; Boutov et al. 2014; Soto-Navarro et al. 2015) and surface circulation (Pascual et al. 2014) have been detected.

No significant trends in frequency of dense water formation events have been detected (Beuvier et al. 2010; Houpert et al. 2016; Somot et al. 2018; Dunić et al. 2019), although a strong interannual



**Figure 2.13 | Warming stripes in the Intermediate Water (from east to west) and the Deep Water (in the Gulf of Lion).** Each stripe refers to a single year and covered periods differ depending on variable. Values have been computed using yearly potential temperature averages in different locations where long-term mooring data are available at different depths. Data from the eastern Ionian come from the HCMR Pylos deep Observatory (Velaoras et al. 2013), and have been downloaded from CMEMS. Data from the Sicily Channel come from the CNR-ISMAR mooring (Schroeder et al. 2017) and can be downloaded from CMEMS. Data from the Gulf of Lion (500 m and 2,300 m) come from the LION Observatory of the MOOSE Network (Houpert et al. 2016) and can be downloaded from SEANOE (Testor et al. 2019). Data from the Gibraltar Strait come from the IEO (Spanish Oceanographic Institute) mooring (von Schuckmann et al. 2018) and can be downloaded from the Copernicus Marine Environment Monitoring Service (CMEMS). Note that colour scales are different for each panel.

variability is reported for each of the dense water formation sites, the northwestern Mediterranean (Marty and Chiavérini 2010; Houpert et al. 2016; Somot et al. 2018; Waldman et al. 2018), the Adriatic Sea (Dunić et al. 2019), and the Aegean Sea (Roether et al. 2007; Beuvier et al. 2010).

### 2.2.7.2 Future change in marine waters

#### Air-sea and land-sea exchanges

The future evolution of the Mediterranean Sea physical characteristics is strongly related to the evolution of the air-sea and land-sea exchanges of water and heat. For the Mediterranean Sea, the net surface water loss by the sea is constituted by the combination of the evaporation over the sea, the precipitation over the sea, the river runoff and the Bosphorus Strait net transport. The net surface heat loss by the sea consists of the shortwave radiation, the longwave radiation, the latent heat and the sensible heat fluxes (these variables are assessed in more detail in Sections 2.2.3, 2.2.4 and 2.2.5, as well as in Section 3.1.3.2).

In addition to the changes in radiation, the future evolution of the Mediterranean Sea physical characteristics is strongly related to the evolution of the other air-sea heat fluxes. Under present-climate conditions, the net heat loss by the Mediterranean Sea surface (namely the sum of shortwave radiation, longwave radiation, latent heat and sensible heat fluxes) is positive meaning that sea is losing heat by its surface over a long period of time. In the future, the net heat loss by the sea surface is expected to decrease (Somot et al. 2006, 2008; Dubois et al. 2012; Gualdi et al. 2013; Adloff et al. 2015; Soto-Navarro et al. 2020), because the increase in shortwave, net longwave and sensible heat fluxes will dominate the increase in latent heat loss (Dubois et al. 2012). With respect to the end of the 20th century and based on coupled regional climate models, the decrease in the net heat loss could reach between  $-1.8$  and  $-5.5 \text{ W m}^{-2}$  by 2050 following the medium-range A1B scenario (Dubois et al. 2012) and between  $-2.1$  and  $-6.4 \text{ W m}^{-2}$  (resp.  $-1.0$  and  $-3.7 \text{ W m}^{-2}$ ) at the end of the 21st century following high-range RCP8.5 (resp. medium-range RCP4.5) scenarios (Soto-Navarro et al. 2020). This implies that the atmosphere could even start to warm the Mediterranean Sea from the mid-21st century instead of cooling it in the present-day climate according to some models (*medium confidence*). Changes in the Mediterranean Sea surface heat budget depend to a great extent on the socio-econom-



ic scenario chosen: the higher the greenhouse gas emissions, the greater the response of the budget.

To summarize, an increase in the net surface water loss by the sea is expected in the future due to a decrease in precipitation and in river runoff and an increase in evaporation (Sections 2.2.5 and 3.1.1). In addition, a decrease in the net surface heat loss by the sea is expected in the future (Sections 2.2.3 and 2.2.5) because the increase in shortwave, net longwave and sensible heat will dominate the increase in latent heat loss. In particular, this means that, from the middle of the 21st century, some models predict that the atmosphere could, in average, warm the Mediterranean Sea instead of cooling it in the present-day climate.

Future changes in the wind strength over the sea will likely remain low even at the end of the 21st century in pessimistic scenarios (Section 2.2.2).

### Sea surface temperature

In future climate change scenarios based on both GCMs and RCMs including the Mediterranean Sea representation, a significant warming of the Mediterranean Sea surface temperature is projected (*very high confidence*) (Somot et al. 2006; Planton et al. 2012; Shaltout and Omstedt 2014; Adloff et al. 2015; Mariotti et al. 2015; Alexander et al. 2018; Darmaraki et al. 2019b). The warming rate depends at the first order on both the temporal horizon and the greenhouse gas emission scenario (*very high confidence*) (Adloff et al. 2015; Mariotti et al. 2015; Darmaraki et al. 2019b). The sea warming will generally remain below that of the air over surrounding land (*high confidence*) due to ocean thermal inertia, probably leading to an increase in land-sea temperature contrast (Somot et al. 2008). With respect to the end of the 20th century, the annual-mean and basin-mean sea surface temperature is expected to increase by 0.6-1.3°C before the mid-21st century and by 2.7-3.8°C (resp. 1.1-2.1°C) at the end of the 21st century period under the pessimistic RCP8.5 (resp. medium RCP4.5) scenario (Darmaraki et al. 2019b). The upper values of those warming ranges are possibly underestimated as higher warming are obtained in CMIP5 GCMs (Mariotti et al. 2015; Darmaraki et al. 2019b).

Future warming will be roughly homogeneous in space (*medium confidence*) with the Balearic Sea, the North Ionian Sea, the Northeast Levantine Sea and the Adriatic Sea identified as potential hotspots of maximum warming (*low confidence*) (Fig. 2.12a)

(Adloff et al. 2015; Darmaraki et al. 2019b; Soto-Navarro et al. 2020). This hotspot pattern however does not match well with the observed warming pattern (Fig. 2.12a), illustrating that climate change related evolution is likely still hidden by natural variability. Spatially, the future warming of the sea surface is bimodal as it has been in the past (Fig. 2.12b). For the near-future (2020-2039) with respect to the end of the 20th century, local annual-mean sea surface temperature change is everywhere positive and can reach locally +1.6°C at maximum whatever the scenarios whereas at the end of the 21st century, the local annual-mean warming spreads from 2 to 5°C for scenario RCP8.5 (resp. from 0.5-3.0°C for RCP4.5).

Warming is not projected to be constant all year round. Stronger warming is expected in summer and weaker warming in winter (*medium confidence*), resulting in substantial increase in warm extremes and a decrease in cold extremes (Alexander et al. 2018). As an illustration, under RCP8.5, maximum monthly-mean sea surface temperature anomalies could reach +3°C over 2040-2059 and more than +5°C over 2080-2099 (median of the CMIP5 models) averaged over the Mediterranean Sea (Alexander et al. 2018). In addition, from the period 2040-2069, the 30-year mean sea surface temperature will always be warmer than the warmest year during the period 1976-2005 (*medium confidence*). This will already be the case in about 50% of the years for the 2010-2039 period (Alexander et al. 2018).

Marine heat waves will very likely increase in spatial coverage, become longer, more intense and more severe than today (*medium confidence*). The intensity of this evolution strongly depends on the temporal horizon and on the socio-economic scenario (Frölicher et al. 2018; Darmaraki et al. 2019b). By 2021-2050, it is expected that marine heat wave frequency increases by a factor 1.5, duration by 2.4-2.7, mean intensity by 1.5 and severity by 5-7 with values largely independent from the socio-economic scenarios (Darmaraki et al. 2019b). By 2100, models project at least one long-lasting marine heat wave occurring every year under RCP8.5 up to 3 months longer, and about 4 times more intense and 42 times more severe than today's events. Their occurrence is expected between June and October, affecting at peak, the entire Mediterranean Basin (Darmaraki et al. 2019b). Under a RCP8.5 scenario, the 2003 marine heatwaves may become a normal event for the period 2021-2050 and a weak event at the end of the 21st century (*medium confidence*) (Darmaraki et al. 2019b).

The warm extreme sea surface temperature changes at the end of the 21st century is likely due to a combination of three factors: a mean sea surface warming, an amplification of the seasonal cycle and an increase in the interannual and day-to-day variability (Alexander et al. 2018; Darmaraki et al. 2019b).

### Sea surface salinity

The future evolution of sea surface salinity of the Mediterranean Sea remains largely uncertain as its sign of change (Adloff et al. 2015; Soto-Navarro et al. 2020). Any change will likely be spatially and temporally inhomogeneous (*medium confidence*) due to the primary role of the river and near-Atlantic freshwater inputs (Adloff et al. 2015; Soto-Navarro et al. 2020). For the end of the 21st century, basin-scale surface salinity anomalies range from -0.18 to +0.16 psu (resp. -0.25 to 0.25 psu) for the pessimistic RCP8.5 (resp. RCP4.5) scenario (Soto-Navarro et al. 2020). However, a surface salinity increase in the eastern Mediterranean Basin is more likely than not whereas the western basin may see an increase or a decrease in its surface salinity (Adloff et al. 2015; Soto-Navarro et al. 2020).

### Surface circulation and exchanges across straits

Change in sea surface circulation has not been deeply assessed yet in the literature (Adloff et al. 2015; Macías et al. 2018), despite their strong capacity to locally modulate the future sea surface temperature and salinity anomalies. In particular, it is likely that the surface circulation changes affect the local sea surface warming hotspots listed above. Noticeable surface circulation changes have been reported for the end of the 21st century in the Balearic Sea and in the North Ionian Sea independently from the scenario choice (Adloff et al. 2015).

At the strait of Gibraltar, the net heat transport towards the Mediterranean Sea will likely increase due to near-Atlantic warming as well as the net mass transport due to increased sea surface water deficit (Somot et al. 2006; Marcos and Tsimplis 2008; Carillo et al. 2012; Adloff et al. 2015; Soto-Navarro et al. 2020). The future evolution of the net salt transport at the strait is unclear, because it depends on the salinity change in the near-Atlantic Ocean surface layer entering the Mediterranean Sea (Marcos and Tsimplis 2008; Adloff et al. 2015; Soto-Navarro et al. 2020). This means that it is currently unclear if the salt transport from the

Atlantic will increase or decrease in the future leading to large uncertainty for the salinity change in the Mediterranean Sea.

### Deep water characteristics

Due to the contrasting effects of increase in sea surface temperature and salinity, the future evolution of the sea surface density is uncertain. Generally, scenarios with strong greenhouse gas concentration increase project a decrease in surface density associated to an increase in vertical stratification of the water column. Increase in density (thus a decrease in stratification) is still possible in scenarios with low level of warming (Adloff et al. 2015). Due to its active thermohaline circulation, the surface climate change signal may be propagated efficiently towards the deeper layers of the Mediterranean Sea (Somot et al. 2006; Carillo et al. 2012) and lead to larger deep warming rates than in other oceans in the world.

The warming and saltening rates of the deep layers is very uncertain as it depends on various factors such as the surface signal, the intensity of the present and future Mediterranean thermohaline circulation (MTHC). This means in particular that the socioeconomic scenario is not the main source of uncertainty in future changes of the deep layers (Adloff et al. 2015). At the end of the 21st century, water masses deeper than 600 m may warm between +0.03 and +1.38°C, and their salinity may increase or decrease with a large uncertainty range, depending on the model (-0.05; +0.51) psu (Adloff et al. 2015; Soto-Navarro et al. 2020).

All published studies agree on a long-term weakening of the open-sea deep convection, the winter deep water formation and the related branch of the thermohaline circulation in the western Mediterranean Sea in high emission scenarios (Thorpe and Bigg 2000; Somot et al. 2006; Adloff et al. 2015; Soto-Navarro et al. 2020). However, natural variability may lead to increase in deep water formation with respect to today's situation during short periods in the future (Macías et al. 2018). For the end of the 21st century and the A2 scenario, decrease of the maximum mixed layer depth reached in the northwestern Mediterranean Sea reach between -17% and -82% depending on the model choice (Somot et al. 2006; Adloff et al. 2015). The picture in the eastern Mediterranean Sea is more contrasted with weakening in some simulations but enhanced convection and thermohaline circulation in others (Somot et al. 2006; Adloff et al. 2015; Soto-Navarro et al. 2020). Some simulations (but not all) project that EMT-like

situation may become the new normal situation for the eastern basin (Adloff et al. 2015).

### 2.2.8 Sea level, storm surges and wave heights

A particular characteristic of the Mediterranean Sea in terms of sea level variability is that it is a semi-enclosed domain linked to the global ocean through the Strait of Gibraltar. This implies that changes in the nearby Atlantic are quickly transferred into the Mediterranean as a basin-wide barotropic signal. At the same time, basin-wide sea level anomalies caused by local forcing (e.g., thermal expansion, evaporation) tend to be transferred to the global ocean in a way that the Mediterranean is in balance with the nearby Atlantic. As a consequence, the Mediterranean basin-wide variations, especially at low frequencies, closely follow the variations in the nearby Atlantic (Calafat et al. 2012; Adloff et al. 2018). The exception to this is the part of the variability related to changes in the atmospheric mechanical forcing (i.e., wind and atmospheric pressure), which can produce Mediterranean-Atlantic differences of few cm even at multidecadal time scales (Menemenlis et al. 2007; Jordà et al. 2012).

The Mediterranean Sea is a microtidal region, with tidal range mostly below 15 cm and relatively low sea levels with 50-year return values below 60 cm over most of the basin (Marcos et al. 2009). These values are small compared to other European Seas. The two exceptions are the North Adriatic and the Gulf of Gabes, where storm surge levels are estimated to be several times per year above 50 cm, with 5-year return values around 90 cm and 70 cm, respectively (Conte and Lionello 2013).

The wave climate in the region is milder than in the Atlantic with smaller mean wave heights (1-1.5 m) and shorter periods (5-6 s) and presents an important spatial variability due to the complex orography and coastline surrounding the basin (Menéndez et al., 2014). Its variability is connected to NAO and the Indian Monsoon index (Lionello and Sanna 2005) and other northern hemisphere teleconnection indices, particularly the East Atlantic Pattern (Lionello and Galati 2008). Annual maxima along the coastlines are largest (above 5 meters) at the northwestern coast of Africa, but high values well above 4 meters occur in several parts (Lionello et al. 2017).

#### 2.2.8.1 Observed change in sea-level

During the 20th century, coastal tide gauges around the Mediterranean have recorded a rise

in the mean sea level. Once tide gauge data have been corrected for the vertical land motion, the sea level trend is very consistent among sites being  $\sim 1.4 \text{ mm yr}^{-1}$  (Wöppelmann and Marcos 2012). This trend is superimposed on interannual and decadal variability that can temporarily mask the sea level rise. The clearest example is the period 1960-1980 during which Mediterranean sea level showed a decreasing trend because a higher than usual atmospheric pressure (Tsimplis et al. 2005). After that period, the atmospheric pressure returned to the typical values and sea level continued to follow the global evolution. For the more recent period, in which sea level has been monitored by satellite altimetry (1993-2018), Mediterranean sea level trend has increased up to  $2.8 \pm 0.1 \text{ mm yr}^{-1}$ , consistent with global sea level trend ( $3.1 \pm 0.4 \text{ mm yr}^{-1}$ ) (Cazenave and WCRP Global Sea Level Budget Group 2018). The rise at global scale is mainly the result of a combination of water thermal expansion and land-based ice melting. During the 20th century both factors contributed equally, although during the last decades, glacier melt is dominating (Cazenave and WCRP Global Sea Level Budget Group 2018).

Analyses of tide gauge data have revealed an increase in the magnitude and duration of the extreme sea level events in the region during the last decades, caused by the rise in the relative mean sea level (for the northern Adriatic Sea: Lionello et al. 2012b; Marcos et al. 2015). In general, wave observational records are too short for assessing multidecadal trends, with the exception of the Northern Adriatic Sea, where one among the worldwide longest instrumental time series (1979 to present) shows an increase in the number of storms, but a decrease of the extreme wave heights (Pomaro et al. 2017).


#### 2.2.8.2 Future sea-levels

The modeling of Mediterranean mean sea level future variations is not straightforward. With their coarse spatial resolution, present-day global climate models (GCMs) are not able to reproduce the regional processes in the basin, although, they are better suited to represent the connection to the global ocean (Calafat et al. 2012). Conversely, regional climate models (RCMs) can capture part of the regional variability but are usually not designed to reproduce the connection with the global ocean, and thus missing a key part of the variability (Adloff et al. 2018). Therefore, sea level rise projections solely based on RCMs have missed that component and only should be considered for the regional patterns, which can cause local

spatial deviations from the basin average by up to +10 cm (Carillo et al. 2012; Adloff et al. 2015, 2018). In conclusion, accounting for all components shows that the Mediterranean sea level rise will be close to the northeastern Atlantic, where future sea level will be similar (difference lower than 5%) to the global mean sea level because regional differences produced by changes in the circulation and mass redistribution almost compensate each other (Slangen et al. 2017). This leads to estimate that the basin mean sea level will likely be 37-90 cm higher than at the end of the 20th century, with a small probability to be above 110 cm. Main contributions to basin-average sea level changes are coming from terrestrial ice melting and the northeastern Atlantic dynamics (Jordà et al. 2020, Fig. 2.14). A different computation based on the sea level projections in the SROCC (Oppenheimer et al. 2019) and accounting for the uncertainty calculation method of the AR5 (Church et al. 2013), confirms that the likely range of the Mediterranean Sea level will be approximately in the range from

a function of time and of the emission scenario, reaching a value in the range from -5% and -10% at the end of the 21st century in the RCP8.5 scenario (Lionello et al. 2017). In any case future sea level rise will become the dominant factor and it will lead to an increase frequency and intensity level of coastal floods (Lionello et al. 2017; Vousdoukas et al. 2017).

Regarding future changes in waves, they will be determined by changes in the wind field over the Mediterranean Sea (Section 2.2.2). Published studies point towards a generalized reduction of the mean significant wave height field over a large fraction of the Mediterranean Sea, especially in winter (Lionello et al. 2008, 2017; Perez et al. 2015). Similarly, the wave extremes are expected to decrease in number and intensity, although there is no consensus whether very large extreme events, associated with very strong winds, would also decrease (Gaertner et al. 2007; Romera et al. 2017; Romero and Emanuel 2017).



**Figure 2.14 | Projected Mediterranean sea level rise averaged in (2080-2099) with respect to present climate (1980-1999) under scenario RCP8.5.** Results based on CMIP5 and Med-CORDEX outputs for the dynamical components and Slangen et al. (2017) for other components. (a) Sum of all contributors (b) Range of projected values for the different contributors: NE Atlantic dynamics (Atl Dyn), Mediterranean dynamics (Med Dyn), Atmospheric Loading (AL), Glacial Isostatic Adjustment (GIA), Ground Water (GW) and Terrestrial Ice Melting (TIM) and Total.

20 to 110 cm higher (depending on scenario) at the end of the 21st century than at the end of the 20th century (Le Cozannet et al. 2019; Thiéblemont et al. 2019).

RCMs and GCMs do not model extreme sea level events and specific 2D simulations forced by high frequency atmospheric forcing are needed. Published studies point towards a reduction on the average number of positive surges throughout the 21st century (Marcos et al. 2011; Conte and Lionello 2013; Lionello et al. 2017). Overall, the results indicate small progressive reduction in comparison with their present-day magnitude as

## 2.2.9 Acidification of the Mediterranean Sea

Human activities are responsible for an increase in atmospheric CO<sub>2</sub> since the beginning of the industrial era. The input of anthropogenic carbon in the Mediterranean is caused by the flux at the air-sea interface, but also results for the Mediterranean Sea, from exchange with the Atlantic Ocean across the Strait of Gibraltar. Approximately 30% of anthropogenic carbon is absorbed by the oceans (Sabine et al. 2004) and leads to decrease of pH in ocean water masses. The Mediterranean Sea is able to absorb relatively more anthropogenic CO<sub>2</sub> per unit area than the

	RECENT CHANGE			PROJECTED CHANGES RCP2.6									PROJECTED CHANGES RCP8.5								
				20-YEAR PERIODS									20-YEAR PERIODS								
	2000-2019			NEAR-FUTURE (2020-2039)			MID-CENTURY (2040-2059)			END OF THE 21ST CENTURY (2080-2099)			NEAR-FUTURE (2020-2039)			MID-CENTURY (2040-2059)			END OF THE 21ST CENTURY (2080-2099)		
<b>CHANGE IN SURFACE TEMPERATURE (°C, MEAN, VALUE, LAND-ONLY)</b>																					
	BASED ON CRU			BASED ON EURO_CORDEX 12 KM ENSEMBLE																	
	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA
MED	0.8	0.41	1	1.1	1.1	1.2	1.3	1.2	1.5	1.3	1.4	1.3	1.3	1.3	1.5	2.2	2	2.5	4.6	4.2	5.3
WEST MAGHREB	0.5	0.1	0.8	1.3	1.1	1.3	1.4	1.1	1.6	1.3	1.3	1.4	1.4	1.2	1.6	2.4	2.1	2.7	5	4.2	4.6
EAST MAGHREB	0.6	0.3	0.6	1.2	1	1.2	1.4	1.1	1.6	1.4	1.3	1.4	1.4	1.2	1.6	2.3	2	2.6	4.6	4	5.3
LEVANT	0.9	0.7	1	1.2	1	1.2	1.3	1.3	1.4	1.3	1.3	1.3	1.3	1.3	1.4	2.1	1.9	2.4	4.3	3.9	4.8
ANATOLIA	0.9	0.4	1.2	1.2	1.2	1.3	1.4	1.4	1.6	1.5	1.6	1.5	1.4	1/4	1.6	2.3	2.1	2.6	4.6	4.3	5.3
CENTRAL MED	0.8	0.3	1	1.1	1	1.3	1.3	1.2	1.4	1.2	1.4	1.2	1.3	1.3	1.5	2.1	1.9	2.5	4.3	3.9	5.1
IBERIA	0.5	0.2	0.8	1	0.9	1.2	1.1	0.9	1.3	1	1	1.1	1.1	1	1.3	2	1.7	2.4	4.3	3.6	5.3
FRANCE	0.7	0.4	0.7	0.9	0.9	1	1.1	0.9	1.1	0.9	1.1	0.8	1	1	1.1	1.8	1.7	2	3.9	3.6	4.7
ALPS	0.8	0.5	1	1.1	1.1	1.1	1.3	1.2	1.3	1.1	1.4	1	1.2	1.3	1.3	2.1	2.1	2.2	4.5	4.5	5
BALKANS	1.1	0.8	1.4	1.1	1.1	1.1	1.3	1.4	1.3	1.2	1.7	1	1.3	1.4	1.4	2.1	2.3	2.2	4.4	4.6	4.8
<b>CHANGE IN SURFACE TEMPERATURE (°C, MEAN, VALUE, LAND-ONLY)</b>																					
	BASED ON CRU			BASED ON EURO_CORDEX 12 KM ENSEMBLE																	
	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY
MED	1.3	2.5	-0.4	0.8	2.3	-1	1	2.3	-0.6*	3.9	6	1.1*	0	1.5	-2.1*	-1.9	0	-4.5	-6.8	-2	-13.3
WEST MAGHREB	2.7	1.3	6.2	-6.7	-7.8	-5.2	-6.2	-5.7	-6.9	-1.7*	0.8	-4.9	-11.2	-9.4	-13.7	-16.8	-20.3	-11.8	-31.2	-33	-28.7
EAST MAGHREB	-7.6	-11.1	5.9	-10.1	-11	-8.7	-5.4	-3.4*	-8.4	-7.8	-8.2	-7.2	-9.7	-8.6	-11.4*	-11.8	-11.7	-11.9	-23.6	-24.1	-22.7
LEVANT	-1.6	-3.5	12.2	-5.8	-5.1	-8.4	-5.8	-4.7	-10.1	-5.8	-4.3	-12	-4.6	-5.1	-2.4*	-10.7	-10	-13.5	-23.5	-23.6	-23.1
ANATOLIA	5.7	3.7	10	0.5*	3.1*	-4.1	-0.1*	2*	-3.9*	2.6	5.3	-2.4	0.2*	2.2	-4.1	-2.4	-0.2	-6.5	-8.7	-4.4	-16.4
CENTRAL MED	2.8	1.3	6.3	0.1*	1.8	-2.4	0.8*	2.1*	-1.2*	3.8	6.3	-0.8	-0.5*	1.5*	-4*	-3.3	-0.7	-7.6	-9.7	-4.7	-18.1
IBERIA	4.9	10	-4.1	-0.1	2.3*	-3.6	0.3*	2.6	-3.2*	5	10.1	-2.8*	-1.5*	0.5*	-4.9*	-6	-2.2	-12.4	-15.1	-7.5	-27.7
FRANCE	-1.8	0.7	-4.4	2.5	4.3	0.3*	2.4	4.2	0.2*	4.1	5.2	2.8	1.6	3.1	-0.3*	0.1	4.4	-5.2	-2.5*	6.7	-13.9
ALPS	-0.5	2.2	-2.8	3.5	6.3	0.7*	4.6	6	3.2	6.2	7.1	5.3	2.9	5	0.9*	3.1	6.6	-0.4*	1.5	9.2	-6.3*
BALKANS	2.5	6.7	-0.4	5.3	8.8	2.4	4.5	6	3.3	8.3	11.8	5.4	3.9	6.6	1.5	4.7	8.1	1.6*	4.9	14.4	-3.7*
<b>CHANGE IN SURFACE SOLAR RADIATION (W/M², 90% INTERVAL BASED ON CMIP5 SIMULATIONS, LAND+SEA)</b>																					
	ANN			ANN			ANN			ANN			ANN			ANN			ANN		
MED	0.3 ; 5.2			0.8 ; 8.6			0.8 ; 10.7			0.5 ; 12.6			0.6 ; 7.7			1.0 ; 10.5			0.6 ; 12.4		
<b>CHANGE IN CLOUD COVER (% , 90% INTERVAL BASED ON CMIP5 SIMULATIONS, LAND+SEA)</b>																					
MED	-2.1 ; 0.2			-3.6 ; 0.2			-4.4 ; 0.5			-4.3 ; 0.5			-3.3 ; 0.0			-4.9 ; -0.2			-7.4 ; -0.8		
<b>CHANGE IN AOD (- , 90% INTERVAL BASED ON CMIP5 SIMULATIONS, LAND+SEA)</b>																					
MED	-0.06 ; -0.02			-0.12 ; -0.04			-0.15 ; -0.04			-0.18 ; -0.05			-0.10 ; -0.03			-0.12 ; -0.04			-0.14 ; -0.04		
<b>CHANGE IN SURFACE TEMPERATURE (°C)</b>																					
	CMEMS (2000-2017)			TOTAL RANGE BASED ON MED-CORDEX RCSM																	
MEDSEA	0.74 (+0.23 ; +0.87)			X			X			X			0.59 ; 1.18			1.30 ; 2.07			2.86 ; 4.10		
<b>CHANGE IN SEA LEVEL (CM, BASED ON BLENDED MULTIPLE DATABASE, SEE THE TEXT, CM, VERY LIKELY RANGE)</b>																					
MEDSEA										(+42, +82)									(+70, +110)		

**Table 2.1 | Climate change as a function of time period and Representative Concentration Pathway for the land sub-regions in Fig. 2.1 and the whole Mediterranean Sea area.** All changes are with respect to the (1980-1999) reference period. For temperature and precipitation, recent changes are based on the Climate Research Unit CRU-TS, future changes are based on the Euro-Cordex regional model simulations. For surface solar radiation, cloud cover and aerosol optical depth values are based on CMIP5 global simulations. For sea surface temperature, recent changes are based on CMEMS observations (2000-2017), future changes on the Med-Cordex regional simulations. For sea level rise, future changes are based on blended multiple databases (*see text*).

global ocean for two reasons. It is more alkaline, thus giving it greater chemical capacity to take up anthropogenic CO<sub>2</sub>, and deep waters are ventilated on shorter timescales (Schneider et al. 2010), thus allowing rapid penetration of CO<sub>2</sub> in its interior.

### 2.2.9.1 Observed change in acidity

Concerning the past trends of anthropogenic carbon absorption by the Mediterranean Sea, the presence of natural CO<sub>2</sub> prevents to determine it from direct measurements in the water column. Estimations of anthropogenic CO<sub>2</sub> from data-based approaches are limited and with large uncertainties that provides concentrations that disagree by more than a factor of two in the Mediterranean Sea (Schneider et al. 2010; Touratier and Goyet 2011). These large differences further result in even opposing estimates for the net CO<sub>2</sub> transport across the Strait of Gibraltar.

In this context, the modeling approach using high-resolution regional model provided some insights on the information resulting from the data based-estimates and quantification of processes responsible of anthropogenic CO<sub>2</sub> storage and acidification of the Mediterranean Sea (Palmiéri et al. 2015). 25% of total anthropogenic carbon inventory in the Mediterranean Sea is due to net exchange at the Strait of Gibraltar, while the remaining 75% is from the air-sea flux. It confirms that the 10% higher mean total alkalinity of the Mediterranean Sea is responsible for a 10%

increase in anthropogenic carbon inventory. The higher alkalinity acts to neutralize acidification and simulated average surface pH change is thus similar for the Mediterranean Sea and the global ocean (-0.08 units), with deep waters exhibiting a larger anthropogenic change in pH than typical global ocean deep waters because ventilation times are faster (between -0.005 and -0.06 units) (Palmiéri et al. 2015).

### 2.2.9.2 Future change in acidity

The published literature concerning specifically the future acidification of the Mediterranean Sea is quite limited leading to low confidence in the assessment. Based on thermodynamic equations of the CO<sub>2</sub>/carbonate system chemical equilibrium in seawater, Goyet et al. (2016) calculated the variation of pH ( $\Delta$ pH) as a function of theoretical anthropogenic CO<sub>2</sub> concentrations. Under the most optimistic SRES scenario, the results indicate that in 2100, pH may decrease down to 0.245 in the western basin and down to 0.242 in the eastern basin (compared to the pre-industrial pH). Whereas for the most pessimistic SRES results for 2100 project a pH decrease down to 0.462 and 0.457, for the western and for the eastern basins, respectively (Goyet et al. 2016). However, these estimates do not consider that the warming of seawater will decrease exchanges across ocean-atmosphere interface and penetration of anthropogenic Jungcarbon, thus they tend to overestimate future acidification.

## 2.3 Pollution

### 2.3.1 Introduction

The 2030 Agenda for Sustainable Development pledges to “ensure that all human beings can enjoy prosperous and fulfilling lives and that economic, social and technological progress occurs in harmony with nature” (UN 2015). Pollution puts at risk the possibility of achieving these outcomes and hence health and well-being. Pollution touches all parts of the planet. It is affecting our health through the food we eat, the water we drink and the air we breathe. Approximately 19 million premature deaths are estimated to occur annually as a result of the way we use natural resources and impact the environment to support global production and consumption. By definition, “pollutant” shall mean any substance that is introduced into the environment that has undesired

effects, or adversely affects the usefulness of a natural resource (air, soil, water and ecosystems). Pollutants can take many forms: (i) physical, substances that are not necessarily involved in chemical or biological reactions, e.g., dust; (ii) chemical, substances that are involved in chemical reactions, e.g., pesticides; and (iii) biological, e.g., bacteria.

### 2.3.2 Physical pollutants

Particulate matter (PM) levels have been monitored during the past decades, mainly because of their effects on health and climate. Aerosols into the atmosphere arise from a variety of anthropogenic activities (transport, industry, biomass burning, etc.) as well as natural sources (volcanic eruptions, sea salt, soil dust suspension, natural forest fires,

etc.) (Seinfeld and Pandis 2006). Both sources result in direct emission of PM (primary PM) and emission of gaseous aerosol precursors (leading to secondary PM). A number of epidemiological studies have examined the impact of PM on human health, expressed as increased mortality and morbidity varying according to the physical (size, shape, etc.) and chemical (composition) characteristics of PM (Van Dingenen et al. 2004).

The PM impact on climate is primarily a cooling effect due to increased scattering to space as the atmospheric aerosol burden increases. The overall cooling by aerosols might be equivalent to a radiative forcing of up to  $2.5 \text{ W m}^{-2}$ , counterbalancing global warming by greenhouse gases (Gillett et al. 2013; Knutson et al. 2013). More important than this direct effect may be the indirect effect that aerosols have on climate, acting as cloud condensation nuclei (CCN) (Levin et al. 2003; Gerasopoulos et al. 2006). Moreover, the particles have a large effect in reducing visibility as well as play a significant role in the deterioration of monuments and buildings (Gerasopoulos et al. 2006). Several studies conducted over the Mediterranean Basin revealed a distinct spatial inhomogeneity (Gerasopoulos et al. 2006), with PM levels increasing from north to south and west to east of the basin (Querol et al. 2009), and distinct sources. PM analysis through the years allowed the identification and classification of PM episodes as follows: (i) local urban PM pollution events (mostly in the cold season), (ii) regional PM pollution episodes (warm season) and (iii) dust outbreaks (Rodríguez et al. 2003).

### 2.3.2.1 Particulate matter (PM) levels and sources

Several factors favor the occurrence of high PM concentrations in the Mediterranean Basin. First, the abrupt topography, coupled with the characteristic synoptic scale patterns, results in low mean wind speeds that hinder the air mass renovations and favor the accumulation of PM in the surrounds of emission regions – leading to the so-called Atlantic/Northern clean air advections events (Rodríguez et al. 2007). Second, the low precipitation in the Mediterranean Basin favors the long residence time of PM in the atmosphere, leading to higher background PM levels (Rodríguez et al. 2007; Querol et al. 2009). The joint influence of low precipitation rates and traffic-forced resuspension of road (which is strongly enhanced by the dust accumulation in streets and roads), construction and demolition dust promoted the local urban episodes in the Mediterranean Basin

(Rodríguez et al. 2007; Talbi et al. 2018). This factor, combined with the high percentage of water coverage of the area, especially in the East region of the basin, has a great contribution of the sea-salt aerosols to the PM levels and composition (Im 2013).

PM observations from monitoring networks, in the period 2007-2009, were analyzed in order to characterize particulate pollution and its health effects across Mediterranean countries (Karanasiou et al. 2014). It was concluded that the average concentrations for PM across the Mediterranean Basin are within the range of annual means typical of European sites and according to the monitoring site characteristics (traffic and urban background sites) (Querol et al. 2004; Putaud et al. 2010). The regional patterns mentioned in previous studies were highlighted, with higher PM concentrations in Italian and Greek cities, and lower levels in the Western Mediterranean (Barcelona, Marseille, Madrid, Huelva). PM10 levels at the traffic sites showed a quite similar variation. In Turin, as in the other cities of the Po valley (Bologna, Milan, Parma, Modena and Reggio Emilia), the combination of stagnant air conditions with high emissions and high population density is the main cause of very strong pollution episodes (Cyrus et al. 2012). Similarly, the air pollution problems in Athens and Thessaloniki are the result of the high population density and the accumulation of air pollutants over the city, due to topography (basin surrounded by mountains), narrow and deep street canyons and adverse meteorological conditions (Karanasiou et al. 2009; Kassomenos et al. 2011). Thermal inversions, followed by accumulation of air pollutants in the lower layers of the atmosphere are also very common in different locations like Athens (Karanasiou et al. 2014) or Beirut (Saliba et al. 2006), increasing the evening concentrations of ambient PM10.

In most countries of the southern Mediterranean, air pollution is not sufficiently monitored (Naidja et al. 2018). Emission inventories are less precise than that available in the northern Mediterranean since they are generally based on surveys and questionnaires. Because of that, local scientific articles were relatively scarce and hard to find. However, most of the available studies show that PM concentrations in this Mediterranean region are much higher than the limit values given in WHO guidelines (Naidja et al. 2018). Emissions from road traffic, resuspension of road dust, especially on unpaved roads, and natural contributions have been found to be an important source of fine particles and play a key role on the

observed levels and exceedances (Mahmoud et al. 2008; Abderrahim et al. 2016; Naidja et al. 2018). Cairo (Egypt) is an example of a city where road traffic emissions are hugely important in PM10 concentration, and, according to WHO is ranked in the 33rd position on the list of the most polluted cities by PM10 (Mahmoud et al. 2008; Lowenthal et al. 2014; Naidja et al. 2018).

Since it is expected that the majority of the Mediterranean population will continue to live in cities, especially in the eastern and southern part of the basin, with a tendency to growth, higher anthropogenic pressure in a context of climate change will occur (Rafael et al. 2015; Naidja et al. 2018). Most PM exceedances were registered in regional background sites (Escudero et al. 2007), with more than 70% of them being attributed to dust outbreaks (Escudero et al. 2007; Mitsakou et al. 2008). Compared with the central and northern Europe, the occurrence of higher PM concentrations associated with dust outbreaks is higher in the Mediterranean Basin (Rodríguez et al. 2007), and are more frequent and more intense in the central and eastern than those in the western Mediterranean Basin. These episodes have been studied on an 11-yr period (2001–2011) (Pey et al. 2013). Dust outbreaks are very frequent in the southern Mediterranean, where they occur more than 30% of the days, while in northern Mediterranean this value is below 20%. The central Mediterranean appears as a transitional area, with a decreasing south to north gradient of dust outbreaks, with slightly higher frequency of dust episodes in its south, when compared to west and east sides of the basin, for similar latitudinal positions (Pey et al. 2013).

Regarding intensity characteristics and seasonality patterns, significantly high contributions are common in autumn-spring in the eastern Mediterranean,

with occurrence of many severe episodes (daily dust averages over  $100 \mu\text{g m}^{-3}$  in PM10) throughout the year. However, in the western Mediterranean a clear summer prevalence is noticed, with low occurrence of severe episodes; and no seasonal trend is detected in the central region, with moderate-intensity episodes (Pey et al. 2013). The contribution of dust outbreaks to PM concentrations reveals a downward trend in the period between 2006 and 2011, a period in which there was also a decrease of the NAO index for the summer period. Therefore, it can be concluded that a sharp change in the atmospheric circulation have affected the number of dust episodes and, consequently, the annual dust inflows to PM10 (Fig. 2.15) observed in the Mediterranean Basin (Pey et al. 2013).

The low PM2.5/PM10 ratio (approximately 0.25) in the eastern Mediterranean region also indicates that the particle size distribution has a large contribution of coarse particles which are either affected by a background level of naturally occurring dust (dust outbreaks from the Saharan Desert and sea salt particles from the Mediterranean Sea itself) or that the region is characterized by high levels of primary coarse PM emissions (Koçak et al. 2007b, 2007a). Even though the PM2.5/PM10 ratio showed seasonal variations, the values remained lower than 0.5 in most cases (Koçak et al. 2007a; Asaf et al. 2008), a value that is least two times lower than those of the western Mediterranean (Saliba and Massoud 2010).

### 2.3.2.2 Particulate matter (PM) chemical profiles

Regarding the chemical composition of PM, different species can be found such as carbonaceous compounds, inorganic ions and metals (Galindo et al. 2018). Although they are present at extremely low levels, some components such as trace metals are relevant in air quality studies because of



Figure 2.15 | PM10 concentration above the annual limit value of  $40 \mu\text{g}\cdot\text{m}^{-3}$  (based on EU Directive 2008/50/CE).



their toxicity and environmental persistence (Roig et al. 2013). Recent clinical and toxicological studies demonstrate the link between exposure to airborne metal through inhalation and pulmonary and cardiovascular effects, genotoxic and carcinogenic outcomes and increased daily mortality (Gottipolu et al. 2008; Lippmann and Chen 2009; Tchounwou et al. 2012).

Cooling metal concentrations are considered as good tracers of specific pollution sources, both natural and anthropogenic (Arhami et al. 2017; Diapouli et al. 2017). The main natural sources include wind-blown dust and sea-spray (Chen et al. 2008; Engelbrecht and Jayanty 2013) including elements such as calcium, aluminum, iron, potassium, sodium and magnesium. Desert dust contribute to PM composition and have a high influence on climate in the North as well as in the South of the Mediterranean (Kchih et al. 2015; Kaskaoutis et al. 2019). Specific meteorological circulations and natural sources like the Mediterranean Sea and the proximity of Sahara create specific patterns of aerosol concentrations that could influence not only the particulate concentrations through Europe but also the global climate due to the transport of dust from the Sahara (Ganor et al. 2010).

Regarding anthropogenic activities, exhaust and non-exhaust vehicle emissions, coal combustion and a variety of industrial processes, like metal works and smelters, are the major sources of heavy metals such as zinc, copper, nickel or chromium (Thorpe and Harrison 2008; Pant and Harrison 2013). In the last decades, emissions of some heavy metals in Mediterranean Basin have dropped significantly, in particular from industrial facilities due to improvement of abatement techniques (Dayan et al. 2017). In the case of lead, a drastic reduction in ambient concentrations has been observed since the introduction of unleaded gasoline (Cho et al. 2011; Salvador et al. 2012). The influence of traffic and dust outbreak intrusions on PM levels and metal content have been studied (Galindo et al. 2018), showing that the PM coarse fraction was affected more by variations in traffic intensity than the submicron fraction: the highest decreases during the weekends due to the reduction in traffic induced resuspension. That dust outbreaks had a greater impact on the levels of other metals such as titanium and lead, significantly affecting their seasonal variability. High concentrations of vanadium and nickel compared with the values found at larger urban areas were observed. This could be attributed to a significant contribution from soils, dust outbreaks (Galindo et al. 2018) and even ship emissions (Monteiro et al. 2018a; Russo et al. 2018).

Another issue related to PM composition is its radionuclide content. Radionuclides in the atmosphere rapidly attach on submicron-sized aerosols, and their variability in ground-level air is driven by the behavior of aerosols (Povinec et al. 2012; Hirose and Povinec 2015). Atmospheric radionuclides are deposited from the air onto the land and sea surface by wet and dry deposition. In this way, the terrestrial and marine environments are labeled by natural and anthropogenic radionuclides that can be used as tracers of environmental processes (Pham et al. 2017). Radionuclide content can pose a health hazard following an accident involving nuclear material (Baeza et al. 2016). However, the occurrence of anthropogenic radionuclides in aerosols is also due to erosion and resuspension processes, as well as the emission and transport of particulate matter due to biomass burning as consequence of wild fires (Strode et al. 2012; Evangelidou et al. 2014), and dust transport due to dust outbreaks (Hernández et al. 2005). Due to these processes, the anthropogenic radionuclide concentration in near surface atmosphere is variable. Naturally occurring radionuclides are also present in airborne particles as they are also present in soil particles able to be eroded, re-suspended or transported by the processes previously described, and also due to the radon exhalation from soil, which is especially significant to lead-210 and polonium-210 (Baeza et al. 2016).

### 2.3.2.3 Plastics (macro/micro/nano)

We live in the plastic age, since synthetic polymers are present in most aspects of human life both in developing and industrialized countries. The worldwide production for plastics increased annually by 10% since the 1950s, reaching 300 Mt in 2015 (Geyer et al. 2017). As of 2015 approx. 6,300 Mt of plastic waste had been generated, around 9% of which had been recycled, 12% was incinerated, and 79% was accumulated in landfills or the natural environment (Geyer et al. 2017). Synthetic thermoplastics constitute the most abundant and still growing component of anthropogenic debris entering the Earth's oceans (Ivar do Sul and Costa 2014). Up to 80%, or sometimes more, of the waste that accumulates on land, shorelines, the ocean surface or seabed is plastic (Barnes et al. 2009). The smallest form of plastic litter is called micro-plastic (<5 mm) and can represent up to 335,000 items km<sup>-2</sup> or 5 kg km<sup>-2</sup> in marine waters, and up to 25 kg km<sup>-2</sup> in coastal sediments (Koutsodendris et al. 2008; Ryan et al. 2009). Plastic debris, their dissolved derivatives, as well as, the adsorbed organic pollutants (Hirai et al. 2011) pose a direct risk to human and marine ecosystem health (Galloway 2015; Koelmans et al. 2017). As a rule, widely used plastics do not rapidly degrade

naturally when released into the environment, it can take 50 or more years for plastic to fully decompose (Müller et al. 2001).

In the Mediterranean Sea, the average density of plastic (1 item per 4 m<sup>2</sup>), as well as its frequency of occurrence (100% of the sites sampled), are comparable to the accumulation zones described for the five subtropical ocean gyres (Cincinelli et al. 2019), increasing the impact for marine biota with hotspots for the risk of plastic ingestion across multiple taxa especially in the coastal zone (Compa et al. 2019). Plastic debris in the Mediterranean surface waters (Fig. 2.16) was dominated by millimeter-sized fragments, but showed a higher proportion of large plastic objects than that present in oceanic gyres, reflecting the closer connection with pollution sources (Cózar et al. 2015). Multi-annual simulations of advected surface passive debris depict the Tyrrhenian Sea, the northwestern Mediterranean sub-basin and the Gulf of Sirte as possible retention areas (Mansui et al. 2015). No permanent structure able to retain floating items in the long-term were found, as the basin circulation variability brings sufficient anomalies to alter the distribution (Mansui et al. 2015).


Beyond the concern with “traditional” PM effects, an emergent research issue worldwide has been

focused in the occurrence of microplastics in the atmospheric compartment (MP; plastic particles with a longest dimension < 5 mm). MP may undergo photo-oxidative degradation in the environment, along with wind shear and/or abrasion against other ambient particulates, eventually fragmenting into fine particles (Gasperi et al. 2018). The risk of inhaling fibrous MP following widespread contamination within different environmental compartments deserves special attention owing to both the scale of their worldwide production and their potential to fragment into smaller, more bioavailable fibers. Human exposure to MP could also occur through ingestion, for example fibrous MP can settle on the floor; children, owing to crawling and frequent hand-to-mouth contact, ingest daily settled dust (Gasperi et al. 2018). Two studies have demonstrated the presence of MP in the atmosphere (Dris et al. 2016, 2017), thereby suggesting potential human exposure (none of these studies has been conducted in the Mediterranean Basin).

### 2.3.3 Chemical pollutants

#### 2.3.3.1 Nutrients

Nutrients (mainly nitrogen, N, and phosphorous, P) constitute an important factor controlling marine




**Figure 2.16 | Averaged 2013–2017 map of plastic debris concentration (g km<sup>-2</sup>) at the sea surface.** Some geographical names used in the text are given: the (1) Gulf of Arzew, (2) Gulf of Tunis, (3) Gulf of Sidra, (4) Abu Qir Bay, (5) Cilician Sea, (6) Izmir, the (7) Thermaic Gulf, (8) Saronic Gulf, (9) Buna-Bojana, (10) NW Adriatic, (11) Taranto Gulf, (12) Gulf of Naples, (13) Gulf of Genoa, (14) Gulf of Lion, (15) Catalan Sea, and (16) Malaga Bay (Liubartseva et al. 2018).

primary producers, as they control phytoplankton growth, biomass and species composition (Sakka Hlaili et al. 2006). According to the nutrient concentrations, marine waters are characterized as oligotrophic (low nutrient concentrations), mesotrophic (nutrient enriched water), or eutrophic (nutrient rich water). The Mediterranean Sea is characterized by oligotrophic off-shore waters, with decreasing levels of nutrients eastwards from Gibraltar to the Levantine Sea (Ignatiades et al. 2009; Tanhua et al. 2013). The Eastern Mediterranean Sea is the most oligotrophic region, with very low nitrate concentrations ( $< 0.5 \mu\text{M}$ ) and phosphorous ( $< 0.2 \mu\text{M}$ ) (Pujo-Pay et al. 2011). Pronounced phosphorous limitation, with N/P ratio  $> 30$ , is observed for the south of Levantine Sea and Ionian Sea (Kress et al. 2003; Pujo-Pay et al. 2011).

However, Mediterranean coastal areas, which are highly populated, are experiencing increasing N and P loading from anthropogenic activities, such as urban effluents, industrial discharges, agricultural runoffs, aquaculture activities and riverine inputs from a drainage area of  $1.5 \times 10^6 \text{ km}^2$  (UNEP/MAP 2017). The overall inputs of N and P in these areas are about 1.5-4.5 and 0.1-0.4 Mt  $\text{yr}^{-1}$ , respectively. The main sources of N in Mediterranean are urban wastewater treatment (45%) and livestock farming (24%). Fertilizer use can also bring nitrogen and inputs can exceed  $10^6 \text{ kg yr}^{-1}$  (Fig. 2.17). Aquaculture contributed also to the emission on N (10%). For P, the main emitters are manufacture of fertilizers (40%), farming of animals (39%) and urban wastewater treatment (13%) (UNEP/MAP 2012b).

Some coastal regions are known as hotspots of nutrient inputs. In the North of Mediterranean Sea, the Lagoon of Venice and the Gulf of Lion sustained high nitrate levels,  $18 \mu\text{M}$  and  $9 \mu\text{M}$ , respectively (Aciri et al. 2004; Severin et al. 2014). Nitrate rich waters characterize also the Eastern Adriatic Sea ( $4 \mu\text{M}$ ) (Skejic et al. 2017) and the Western Tyrrhenian Sea ( $6.5 \mu\text{M}$ ) (Astraldi et al. 2002). In the southern Mediterranean, The Gulf of Gabès is a main region known for P enrichment ( $1-11.2 \mu\text{M}$ ), since Tunisia is an important producer country of P fertilizers. Nitrate ( $6-6.5 \mu\text{M}$ ) and ammonia ( $\sim 4 \mu\text{M}$ ) showed also pronounced levels in this Gulf (Dira et al. 2016). High nitrate concentrations were often measured in other Tunisian coastal systems, such as the Lagoon of Bizerte ( $\text{NO}_3^-$ :  $1-6.3$ ,  $\text{NH}_4^+$ :  $20-30 \mu\text{M}$ ) (Sahraoui et al. 2012) and the North Lake of Tunis ( $\text{NO}_3^-$ :  $7.5-198 \mu\text{M}$ ) (Armi et al. 2011). In the Algerian-Provençal Basin and the Gibraltar Strait, enrichment of water with nitrate has been reported ( $9.5-10 \mu\text{M}$ ) (Béthoux et al. 1998; Gómez et al. 2000).

Nutrient enrichment of Mediterranean Sea may result in a high increase in phytoplankton growth and biomass, leading to the eutrophication. The impacts of eutrophication include hypoxia or anoxia and may provoke harmful algal blooms, some of them toxic. Harmful algal blooms (HABs) cause human illness and mortality and have socio-economic impacts related to toxicity of harvested fish and shellfish, loss of aesthetic value of coastal ecosystems, and reduced water quality impacting tourism (Section 2.3.4).



**Figure 2.17 | Fertilizer use and nitrogen release in Mediterranean Sea** (UNEP-GRID Arendal 2013).

### 2.3.3.2 Gaseous pollutants: nitrogen dioxide, sulphur dioxide, ozone


The Mediterranean Basin is one of the regions in the world where high concentrations of gaseous air pollutants (nitrogen dioxide – NO<sub>2</sub>, sulphur dioxide – SO<sub>2</sub>, and ozone – O<sub>3</sub>) have been reported frequently (Dayan et al. 2017). The elevated concentrations observed are attributed to the combination of diverse emission sources affecting the Mediterranean Basin including industry, traffic and shipping emissions (Karanasiou et al. 2014). The Mediterranean climate is characterized by arid conditions as well as many hours of sunshine and specific atmospheric recirculation patterns that significantly enhance air pollution levels (Karanasiou et al. 2014; Querol et al. 2018).

Commonly, NO<sub>2</sub> concentrations in North Mediterranean countries are higher than those observed in northern Europe (Cyrus et al. 2012). This fact was attributed to the transport sector, and to the higher conversion of nitric oxide (NO) to NO<sub>2</sub> caused by high temperatures and O<sub>3</sub> concentrations (Schembari et al. 2012; Karanasiou et al. 2014). The spatial pattern of N deposition varies across the Mediterranean Basin (Fig. 2.18). In Iberia, dry deposition is an important component of the total atmospheric N input to natural habitats (García-Gómez et al. 2018; Oliveira et al. 2020).

Ships are among the major emitters of air pollutants such as SO<sub>2</sub> and NO<sub>x</sub>, their contribution to the emissions from the transport sector (Schembari et al. 2012) and to the air pollution in the Mediterranean Basin (Monteiro et al. 2018b; Russo et al. 2018) is growing. Several studies have also shown that NO<sub>2</sub> exceedances (yearly and hourly) in cities of the Mediterranean Basin are caused by road traffic emissions (Borrego et al. 2012; Belhout et al. 2018) (Fig. 2.19).

The large variety of Volatile Organic Compounds (VOCs), NO<sub>x</sub> emissions and the climate conditions of Mediterranean Basin influences O<sub>3</sub> formation and destruction (Sahu and Saxena 2015; Sahu et al. 2016). These factors result in higher O<sub>3</sub> concentrations and frequent tropospheric O<sub>3</sub> episodes recorded across the Mediterranean Basin, with different frequencies in the East and West (Sicard et al. 2013).

The western Mediterranean Basin is characterized by frequent sea breezes, driven by sea-land thermal contrast. These sea-land breezes play an important role for the O<sub>3</sub> concentrations since they transport air masses, including O<sub>3</sub> precursor gases, from urban agglomerations located in coastal areas, towards inland suburban and rural areas (Millán et al. 2000).



**Figure 2.18 | Modelled nitrogen deposition for the Mediterranean region** based on the European Monitoring and Evaluation Programme (EMEP) model at 0.1x0.1° longitude-latitude resolution (EMEP MSC-W chemical transport model version rv4.7)<sup>6</sup>. Modelled N deposition is based on 2013 emissions data. (A) Total N deposition [oxidized+reduced; dry+wet], (B) percentage of dry deposition, (C) percentage of wet deposition, (D) percentage of oxidized deposition and (e) percentage of reduced deposition (Ochoa-Hueso et al. 2017).

High O<sub>3</sub> episodes in this region are linked to the combination of one or several of these mechanisms: (i) local/regional photochemical production and surface transport from coastal to inland regions;

<sup>6</sup> [www.emep.int](http://www.emep.int)



**Figure 2.19 | Nitrogen dioxide (NO<sub>2</sub>) concentrations above the annual limit value of 40 µg·m<sup>-3</sup>** (based on EU Directive 2008/50/CE).

(ii) O<sub>3</sub> transport from higher-altitude atmospheric layers, due to air mass re-circulation in the previous days; and (iii) long-range transport of O<sub>3</sub> and its precursor gases (Querol et al. 2018). In the eastern Mediterranean Basin, the O<sub>3</sub> episodes depends on the relative strength of the high-pressure system covering the eastern Mediterranean and Balkan area: (i) strong pressure gradient with northerly winds, creating good ventilation in the Athens Basin (Kallos et al. 2014); (ii) weak pressure gradient with local/regional O<sub>3</sub> events prevail; and (iii) stratospheric O<sub>3</sub> contributions to increase surface O<sub>3</sub> concentrations during specific meteorological scenarios (Zanis et al. 2014; Kalabokas et al. 2015). Tropospheric O<sub>3</sub> concentrations observed in the summer over this region (Fig. 2.20) are among the highest over the Northern Hemisphere (Dayan et al. 2017).

### 2.3.3.3 Trace metallic elements

Metal trace elements (MTE, or heavy metals in the old designation) whose main ones are cadmium (Cd), chromium (Cr), copper (Cu), lead (Pb), nickel (Ni),

zinc (Zn) and mercury (Hg) are common elements in the earth's crust (Navarro-Pedreño et al. 2008). They are also generated by human activities (Hassanien and Abdel-Latif 2008; Tovar-Sánchez et al. 2016; Merhaby et al. 2018). Urban and industrial wastewaters, atmospheric deposition and run-off from metal contaminated sites constitute the major sources of toxic metals (UNEP/MAP 2012a).

High MTE levels have been found in various soils (vegetated soils, dikes, waste rock and slag) at mining sites in Morocco. These mining activities in addition to agricultural and pastoral practices constitute a way of entry of the MTEs into the food chain and thus increase the risk of contamination of the population. Several plant species are adapted to these high levels of MTE and thus represent an important potential for the development of mining site rehabilitation strategies (Smouni et al. 2010). In agricultural soils of the Argolida Basin (Peloponnese, Greece), the MTE concentrations are high, following a decreasing order: Fe > Mn > Ni > Zn ~ Cr > Cu > Co ~ Pb > As > Cd (Kelepertzis 2014).



**Figure 2.20 | Number of days (more than 25) above ozone (O<sub>3</sub>) limit value of 120 µg·m<sup>-3</sup>** (based on EU Directive 2008/50/CE).

	Al	Cr	Cu	Fe	Hg	Mn	Ni	Pb	Ti	Zn
Min	36,866	75.2	42.4	26,313	0.04	552	47.9	44.7	2,343	86.8
Max	72,020	102.8	52.3	36,098	0.41	2,826	60.7	74.8	3,876	129.0
Mean	58,564	85.9	47.4	31,566	0.12	893	53.3	57.8	3,065	102.3


**Table 2.2 | Metal concentrations ( $\mu\text{g g}^{-1}$  dry weight) in marine sediment of Taranto Gulf (Ionian Sea, southern Italy) (Buccolieri et al. 2006).**

In marine ecosystems, the hotspots of lead, mercury and cadmium were essentially located on the north-central and southeastern shores of the Mediterranean Basin (Fig. 2.21) (UNEP/MAP 2012b). Polluted surface samples on the Barcelona city continental shelf taken in 1987 reached enrichment factors of up to 490 for Hg, about 40 for Pb and Cd, and about 17 for Zn, Cr and Cu. In 2008, the data showed a decline with enrichment factors between 20 and 30 for Hg and Cd and between 5 and 12 for Zn, Cr, Pb and Cu (Palanques et al. 2017). In the Taranto Gulf (Ionian Sea, Southern Italy) (Table 2.2), MTE distribution is principally influenced by industrial and urban wastes. River discharges and prevailing anticlockwise marine currents are further factors influencing metal accumulation in sediments (Buccolieri et al. 2006). In surface sediments in Lebanon (eastern Mediterranean Sea), trace metals (Cd, Pb, Zn, and Cr) contamination at Beirut Port was classified as “the most highly polluted” and as “moderately polluted” at Tripoli Port (Merhaby et al. 2018).

Freshwater ecosystems are also affected by MTE pollution. In the Ichkeul Lake Basin (northeastern Tunisia) MTE showed concentrations in the sediment samples following the order: Fe > Mn > Zn > Pb > Ni > Cr > Cu > Cd (Touaylia et al. 2016). The concentrations of Fe, Cd, Ni and Cr in the bottom sediments of the Lower Litani River Basin (Lebanon) were higher in the dry season (Nehme et al. 2014). In river sediments from a semi-arid Mediterranean Basin (Algeria), MTEs were grouped by their level of contamination: high (Pb, Cd, Zn, Cu) and low (Al, Fe, Cr, Co, Ni). Sources of this contamination were essentially industrial, agricultural and domestic waste, as well as very specific ones (gasoline station) and diffuse pollution from atmospheric deposition (gasoline, ores, aerosols) (Benabdelkader et al. 2018).

MTE are known for their toxicity, persistence, and bioaccumulation in human and animal tissues, and biomagnify (concentrate at successively higher levels in tissues) in food chains (UNEP/MAP 2012b). In the northwestern Mediterranean Sea, the concentrations of 21 trace elements showed great variability in three species; fish

(sea bass, *Dicentrarchus labrax*), mussels (*Mytilus galloprovincialis*) and oysters (*Crassostrea gigas*). The essential elements (Cu, Mn and Zn) were highest in oysters, but Fe, Cr, Ni, Se, Co and Mo levels were highest in mussels. Fish had the lowest concentrations for all trace elements, which were at least one order of magnitude lower than in



**Figure 2.21 | Mean concentrations of principal trace metals in coastal sediments of the Mediterranean Basin (UNEP/MAP 2012b).**

bivalves. The maximum values set by European regulations for Hg, Cd and Pb were never exceeded (Squadrone et al. 2016). The levels of As, Hg and Pb measured in some commercially key species from Sicilian coasts suggest relatively reduced pollution levels for fish resources in this part of the Mediterranean region (Traina et al. 2019).

### 2.3.3.4 Organic pollutants

#### Polycyclic Aromatic Hydrocarbons (PAHs)

The contamination of the marine environment by polycyclic aromatic hydrocarbons (PAHs) has received considerable attention since the early 1980s after occurrence of tragic marine oil spills like the Amoco Cadiz oil spill (March 1979, Brest, France;  $223 \times 10^3$  t crude oil). PAHs consist of two or more fused benzene rings. PAH distribution is controlled by multiple and inter-dependent parameters. Some of these parameters are linked to intrinsic physicochemical properties of these compounds, mainly hydrophobicity which controls their partition between dissolved and particulate phases. Others are related to the hydrological and biogeochemical characteristics of the environment including water agitation and turbidity, sediment granulometry and particulate or dissolved organic matter content. They are issued from unburned petroleum or oil-derived products (petrogenic PAHs) or from the incomplete combustion of fossil fuels and biomass (pyrogenic PAHs). Petrogenic PAHs are characterized by lighter compounds (phenanthrene, naphthalene, fluorene) and their alkylated derivatives (methyl, dimethyl etc.) and have high affinity for the dissolved phase. Pyrogenic PAHs are characterized by unsubstituted higher molecular weight compounds that exhibit more affinity for suspended particles. Distribution of petrogenic-like and pyrogenic-like compounds help identification of PAH sources in the marine environment. Eight PAHs belong to the list of priority substances: Anthracene, Benzo (a) Pyrene, Benzo (b) Fluoranthene, Benzo (ghi) Perylene, Benzo (k) Fluoranthene, Indeno (123cd) Pyrene, Fluoranthene and Naphthalene. The top 5 are listed as priority hazardous substances due to their potential toxic, mutagenic and carcinogenic effects on organisms and human health (Hussein et al. 2016).

The diagnosis of the Mediterranean with regard to these substances is governed by international and national environmental agencies (i.e., the US Environmental Protection Agency (US-EPA), the European Environmental Commission, the United Nations Environment Program (UNEP), The National Observatory of the Environment and Sustainable Development, ONEDD (Algeria), National Environ-

mental Protection Agency, ANPE (Tunisia), Egyptian Environmental Affairs Agency, EEAA (Egypt) that recommend the monitoring of 16-19 PAHs with special emphasis on PAHs micropollutants in marine matrices.

The overall data set suggests that both petrogenic and pyrogenic PAHs contribute to the PAH pool in Mediterranean coastal waters. Pyrogenic inputs increase in urbanized areas due to large atmospheric inputs and deposition in winter (Guigue et al. 2014; Barhoumi et al. 2018). Total PAH inputs from the Rhône river represent  $50 \text{ t yr}^{-1}$  whereas inputs from sewage treatment plant are much lower ( $<1 \text{ t yr}^{-1}$ ) (Witkowski et al. 2017). Due to their hydrophobicity, PAHs are considered to be preferentially associated with particles in coastal marine waters and sediments (Adhikari et al. 2015). Dissolved PAHs concentrations in the water column may be 4-fold higher than in suspended particles (Guigue et al. 2011). Dissolved PAH concentrations are in the range  $0.158\text{--}3.655 \text{ ng L}^{-1}$  ( $\Sigma 18$  PAHs) (Berrojalbiz et al. 2011) in Mediterranean open sea waters whereas higher concentrations characterize coastal urbanized areas, up to  $560 \text{ ng L}^{-1}$  ( $\Sigma 32$  PAHs) in Marseille-Gulf of Fos (France) (Guigue et al. 2011),  $12\text{--}267 \text{ ng L}^{-1}$  ( $\Sigma 17$  PAHs) in Venice lagoon (Italy) (Manodori et al. 2006),  $13\text{--}120 \text{ ng L}^{-1}$  ( $\Sigma 7$  PAHs) in Alexandria coastal waters (Egypt) (El Nemr and Abd-Allah 2003). In coastal sediments, concentrations range from  $10\text{--}200 \text{ ng g}^{-1}$  d.w. (dry weight) (Asia et al. 2009) in Marseille Bay, France (Gogou et al. 2000), Cretan Sea, Greece (Zaghden et al. 2005), Sfax, Tunisia (Cannarsa et al. 2014), Liguria, Italy (Merhaby et al. 2015), Tripoli harbor, Lebanon (Emara et al. 2008) and Eastern harbor, Egypt (Emara et al. 2008).

PAH concentration gradients are pronounced along coast-open sea transects. PAH concentrations rapidly decrease from the vicinity of rivers, estuaries and small effluents towards coastal and offshore waters. The contamination of estuaries of large rivers and that of harbors in the vicinity of big industrial and urban centers indicates a poor quality environment where a potential risk to the local population may occur (Barhoumi et al. 2018). Otherwise in most Mediterranean coastal waters, data reflect contamination levels from slightly polluted to polluted. Industrial areas near the cities of Sfax and Gabès (Tunisia) were reported moderate-to-highly impacted by hydrocarbons compared to other Mediterranean coastal environments (Fourati et al. 2018b, 2018a). They remain globally higher than those reported in the northern Gulf of Mexico and much lower than those recorded in Chinese coastal environments (Zhou and Maskouei 2003; Fourati et al. 2018b).

In some cases, PAH concentrations are influenced by physical circulation processes that can lead to deposits of contaminants an order of magnitude higher offshore than those near the source of pollutants. Episodic processes of pollutants redistribution may also significantly affect the pollution status of marine areas. For example, the physical accumulation of PAH at air/sea interface ( $\times 200-1,000$ ) during microlayer formation in absence of wind (Wurl and Obbard 2004), followed by PAH scattering during microlayer disruption by wind blow recovery, can locally enhance PAHs concentrations and impact the biota. Similarly, PAH sediment remobilization during resuspension events may greatly modify their potential harmful effects on marine biota (Guigue et al. 2017). In Toulon bay (France), the resuspension of highly contaminated surface sediments (concentration of  $\Sigma 34$  PAHs =  $38.2 \times 10^3$  ng g<sup>-1</sup>) led to a 10-fold increase of dissolved  $\Sigma 34$  PAH concentrations in the water above. The remobilization in seawater was higher for 4-6 ring PAHs, especially benzo(g,h,i) perylene, whose concentration exceeded the authorized limit values of the European Water Framework Directive (Guigue et al. 2017). It is important to monitor pollutants not only at active industrial facility sites but also in disused industrial areas close to the sea border where remnant pollution can produce chronic adverse effects on marine biota.

Species feeding on particles and phytoplankton may bioaccumulate and/or bioamplify PAH concentrations in their body tissues. Measuring accumulation of PAHs in mussel bivalves from the *Mytilus edulis* complex has become a European Commission control strategy of marine waters quality in the Mediterranean (Olenycz et al. 2015; Sire and Amouroux 2016). Different metabolites may be measured in fish and shells and considered as markers of exposure to PAH. Research on the relationships with emergent contaminants is at its beginning and cocktail effects have not been much studied yet. Microplastics have a high potential to adsorb these hydrophobic contaminants and to transfer them throughout the food web to the deep ocean for longer sequestration time.

### Pesticides

Pesticide Active Ingredients (PAI) can be considered as a contaminant as well as a pollutant, in the compartments where they are detected. Mainly originated by agricultural activities, water pollution by PAI is a concern for continental water resources (rivers, lakes and aquifers) and coastal and marine environment of the Mediterranean Sea. Studies

carried out in southern Europe showed the high leaching of herbicides in Mediterranean weather conditions (Louchart et al. 2001), allowing the contamination of groundwater resources. In the other side, by runoff process, surface waters would be contaminated by PAI and their metabolites or degradation products. A large number of pesticide active ingredients (PAI) (over 1,300) are presently used or were used until non-approval or non-renewal in Europe (European Food Safety Authority 2011) and even modern screening methods limit the number of PAI analyzed in one sample ( $< 450$ ) (Rousis et al. 2017). In water bodies, the maximum allowable concentrations are  $2 \mu\text{g L}^{-1}$  for each PAI and  $5 \mu\text{g L}^{-1}$  for all quantified PAIs. For drinkable water these limits are  $0.1 \mu\text{g L}^{-1}$  and  $0.5 \mu\text{g L}^{-1}$  respectively.

Concentrations of these molecules in water bodies (surface and underground waters) were qualified and quantified in nearly all the countries around the Mediterranean Sea. But a recent review at world scale (Stehle and Schulz 2015) stated the difference of availability of referenced data sets for insecticide concentrations in water: notably in the North and the North-East Africa poor information was available.

In European countries, particularly in France (Dubois et al. 2010; Lopez et al. 2015), Italy (Onorati et al. 2006; Meffe and de Bustamante 2014), Spain (Balaguer et al. 2018) and Greece (Lekkas et al. 2004; Konstantinou et al. 2006) implementation of the EU Water Framework Directive (WFD) produced large public data sets for pesticide concentrations in surface waters and aquifers by state administrations. According to the statistical office of the European Union, Spain and Italy are the countries with most use of pesticide. As a result, in those countries, pesticides are one of the most frequently detected classes of micro-pollutants in water<sup>7</sup>. It is not possible to give in this report an exhaustive account of all PAI mentioned in the literature as encountered in Mediterranean waters. Thus, we decided to focus on the most frequently detected PAI and give some examples of maximum concentrations measured.

In water bodies most frequently mentioned PAI-insecticides already in use are chlorpyrifos ( $18.8 \mu\text{g L}^{-1}$ ) (Ccanccapa et al. 2015), dimethoate ( $0.640 \mu\text{g L}^{-1}$ ) (Campo et al. 2013), malathion ( $0.048 \mu\text{g L}^{-1}$ ) (Yurtkuran and Saygı 2013), imidacloprid ( $0.350 \pm 0.433 \mu\text{g L}^{-1}$ ) (Herrero-Hernández et al. 2013) and diazinon ( $14.5 \mu\text{g L}^{-1}$ ) (Youssef et al. 2015). Prohibited PAI-insecticides mentioned are DDT (dichlorodiphenyltrichloroethane) and its metabolites ( $0.40$  to  $3.22 \mu\text{g L}^{-1}$ ) (Dahshan et al. 2016), HCB (hexa-

<sup>7</sup> <https://www.eea.europa.eu/airs/2018/environment-and-health/pesticides-sales>



chlorobenzene) ( $1.1 \mu\text{g L}^{-1}$ ) (Youssef et al. 2015) and endosulfan ( $0.247 \mu\text{g L}^{-1}$ ) (El Bakouri et al. 2008).

The most frequently mentioned PAI-herbicides in use are simazin ( $3.18 \mu\text{g L}^{-1}$ ) (Konstantinou et al. 2006), terbuthylazin ( $0.0219 \mu\text{g L}^{-1}$ ) (Ricart et al. 2010), linuron ( $13.13 \mu\text{g L}^{-1}$ ), 2,4-D ( $20 \mu\text{g L}^{-1}$ ) and glyphosate with its metabolite AMPA ( $167 \mu\text{g L}^{-1}$ ) (Meffe and de Bustamante 2014). Most prohibited PAI-herbicides mentioned in studies are atrazine-desethyl ( $0.158 \mu\text{g L}^{-1}$ ) (Campo et al. 2013), metolachlor ( $1.120 \mu\text{g L}^{-1}$ ) (Konstantinou et al. 2006), DEA (diethyl-atrazine) ( $1.98 \mu\text{g L}^{-1}$ ) (Hildebrandt et al. 2008), diuron ( $0.0169 \mu\text{g L}^{-1}$ ) (Robles-Molina et al. 2014), DIA (deisopropyl-atrazine) ( $8 \mu\text{g L}^{-1}$ ) (Shomar et al. 2006), alachlor ( $0.213 \mu\text{g L}^{-1}$ ) (Stamatis et al. 2013), isoproturon ( $7 \mu\text{g L}^{-1}$ ) (Ricart et al. 2010) and molinate ( $0.026 \mu\text{g L}^{-1}$ ) (Gómez-Gutiérrez et al. 2006).

Concentrations of PAI-fungicides are less mentioned than other PAI. Metalaxyl ( $0.49 \mu\text{g L}^{-1}$ ) (Hildebrandt et al. 2008) and carbendazim ( $1.81 \mu\text{g L}^{-1}$ ) (Licciardello et al. 2011) have been reported.

For the future, in most southern Mediterranean countries, prohibitions of particular PAIs for agricultural uses are applied within a few years delay from European Commission decisions. Many countries developed programs to reduce pesticide use that have uncertain effects like in France (Guichard et al. 2017). New proposed PAI are characterized by shorter standard half-lives and lower dose requirements, nevertheless there is no assurance that these "new" PAI once used widely in many different contexts will have virtuous environmental spreading behaviors.

Climate changes may have conversational effects on PAI water contamination. Firstly, some authors predict an increase use of pesticides to compensate increases of abundance and seasonal activity of bioaggressors (Boxall et al. 2009), although temperature enhancement has been declared to decrease PAI efficiency by conditional resistance of bioaggressors towards herbicides, insecticides and fungicides (Matzrafi 2019). Under these conditions, doses applied should be increased to guarantee the same protection with unseen consequences in environment spreading. Daily temperature fluctuation may increase PAI toxicity like for chlorpyrifos (Verheyen and Stoks 2019), but temperature enhancement should boost degradation of most known PAI probably shortening their half-life.

Change in precipitation patterns with increased occurrence of extreme precipitation events should also modify agricultural patterns. For instance,

rained barley Mediterranean production yields would decrease (Verheyen and Stoks 2019) fueling some changes in the crop systems. Implementation of more intensive systems (Malek and Verburg 2018) can imply an increase of treatments. Increasing weight of extreme events would increase erosion (Raclot et al. 2018) and then promote displacement from land to sea for the PAIs easily adsorbed on soil particles like clay and organic matter.

### 2.3.3.5 Emerging contaminants

In the Mediterranean Basin, 63% of coastal settlements with more than 2,000 inhabitants operate a wastewater treatment plant, while 37% do not. Secondary treatment is mostly used (67%) in Mediterranean treatment plants, while 18% of the plants have only primary treatment (Chatha et al. 2017). As a consequence of this technical, social and environmental issue, different types of chemical substances are released into the environment (Gros et al. 2010; Ratola et al. 2012; Moreno-González et al. 2015; Paluselli et al. 2018b, 2018a). Among these substances, emerging contaminants (ECs) are a category that has received special attention over the last 25 years. ECs are defined as "contaminants of emerging concern that are naturally occurring, manufactured or man-made chemicals or materials which have now been discovered or are suspected present in various environmental compartments and whose toxicity or persistence are likely to significantly alter the metabolism of a living being" (Sauvé and Desrosiers 2014).

There are different classifications of these contaminants due to their usage or origin and effects. For example, (1) antibiotics, (2) antimicrobials, (3) detergent metabolites, (4) disinfectants, (5) disinfection byproducts, (6) estrogenic compounds, (7) fire or flame retardants, (8) fragrances, (9) insect repellants, (10) PAHs (polyaromatic hydrocarbons), (11) personal care products, and (12) pesticides or insecticides (13) pharmaceuticals, (14) plasticizers, (15) reproductive hormones, (16) solvents, (17) steroids and (18) surfactants (Singh and Kumar 2017).

ECs that are very soluble in water (tetracycline, sulfamethoxazole, carbamazepine, and erythromycin, etc.) receive more attention than others because of their impact on the environment (Klaper and Welch 2011). Potential ECs sources and pathways of ground and surface water pollution are shown in Fig. 2.22. Typically, the route of these compounds towards a water body begins with the excretion of the metabolites and parent compounds and their disposal to the wastewater treatment plants (Barrios-Estrada et al. 2018).





Figure 2.22 | Potential sources and pathways for grounds and surface water pollution (Barrios-Estrada et al. 2018).

The endocrine disruptors (EDs) are a subdivision of ECs with chemicals that interfere with the body's endocrine system and produce adverse developmental, reproductive, neurological, and immune effects in humans, abnormal growth patterns and neurodevelopmental delays in children (Moreno-González et al. 2015). The majority of the EDs come from products used to fight undesirable wildlife and agricultural threats (Moreno-González et al. 2015), for example, pesticides, fungicides and rodenticides, synthetic products used in plastic industry (bisphenols or phthalates) and a variety of buildings materials, isolation materials (polychlorinated biphenyl and metals). A list of common sources of EDs is shown in Fig. 2.23.

The presence of ECs in Mediterranean Basin is well documented. Phthalic Acid Esters (PAEs), including dimethyl phthalate (DMP), diethyl phthalate (DEP), di-isobutyl phthalate (DiBP), di-n-butyl phthalate (DnBP), benzylbutyl phthalate (BzBP) and diethylhexyl phthalate (DEHP), with total concentrations ranging from 130 to 1,330 ng L<sup>-1</sup> were found in Marseille Bay (northwestern Mediterranean Sea) (Paluselli et al. 2018b). High concentrations of PAEs were also observed in deep waters offshore (310.2 ng L<sup>-1</sup>) as well as in the Rhône River (615.1 ng L<sup>-1</sup>) (Paluselli et al. 2018a).

A total of 20 pharmaceuticals in sea water and 14 in sediments were found at concentrations from

low ng L<sup>-1</sup> up to 168 ng L<sup>-1</sup> (azithromycin) in sea water and from low ng g<sup>-1</sup> up to 50.3 ng g<sup>-1</sup> (xylazine) in sediments of Mar Menor lagoon located in the South East of Spain (Moreno-González et al. 2015). Pharmaceutically active compounds (PhACs) were detected in the Evrotas River (Southern Greece) waters. The diuretics and the analgesics/anti-inflammatory class were the most abundant, followed by antihypertensives, psychiatric drugs, β-blocking agents and antibiotics and the concentration levels ranged from 0.31 ng L<sup>-1</sup> up to 51 ng L<sup>-1</sup> (Mandarić et al. 2019). Antibiotics were detected in more than 90% of the water samples collected from a Mediterranean river (Llobregat, Spain) and the concentration levels ranging from 0.3 ng L<sup>-1</sup> (flumequine) to 907.6 ng L<sup>-1</sup> (sulfamethoxazole) (Proia et al. 2013). Triclosan (an antimicrobial) was reported to be a contaminant of the Llobregat and Ebro rivers (Spain) and the concentrations in some samples were higher than 150 ng L<sup>-1</sup>. These concentrations should be considered significant considering the toxicity of these compounds and their expected ability to be a precursor of other highly toxic compounds such as dioxins (Kantiani et al. 2008). Azithromycin (antibiotic) was measured at 16,633 ng L<sup>-1</sup> in a tributary of El Albujòn (Spain) (Moreno-González et al. 2015) and acetaminophen was detected at 3,000 ng L<sup>-1</sup> off Thessaloniki, Greece (Nödler et al. 2014). Some drug classes, such as analgesics, antibiotics and betablockers, were still quantified at levels between 0.3 (metoprolol) and hundreds of ng L<sup>-1</sup> (azithromycin) in seawater. In



**Figure 2.23 | Common sources of endocrine disruptors in the environment** (Barrios-Estrada et al. 2018).


particular, six substances (e.g., azithromycin, amoxicillin, venlafaxine, salicylic acid, acetaminophen and ibuprofen) were measured at median concentrations higher than 20 ng L<sup>-1</sup>. Pharmaceuticals were also found in coastal and oceanic waters adjacent to the Strait of Gibraltar (Biel-Maeso et al. 2018).

The impacts of exposure to some ECs have caused concern for both people and wildlife. Many of these substances may cause disorders of the nervous, hormonal and reproductive system, thus posing adverse health outcomes (Rezg et al. 2014; Bilal et al. 2018, 2019; Hernandez-Vargas et al. 2018; Ullah et al. 2018; Rasheed et al. 2019). These synthetic chemicals exhibit geno- or cytotoxic activity and can cause obesity, diabetes, cardiovascular and reproductive disorder or even leads to cancer (Tiwari et al. 2012). Etteieb et al. (2016) have shown that the components responsible of cytotoxicity in water samples from Medjerda river in Tunisia were mainly cyclopentasiloxane, decamethyl (D5), cyclohexasiloxane, dodecamethyl (D6), D-limonene, and ergoline-8-methanol, 8,9-didehydro-6-methyl. Some ECs, such as 17- $\alpha$ -estradiol, bisphenol A and phthalates were reported to alter marine community structure (Essid et al. 2013; M'Rabet et al. 2019).

### 2.3.4 Biological pollutants

Numerous viruses or bacteria of human or animal origin can spread in the environment and infect peo-

ple via water, air and food, mostly through ingestion and occasionally through skin contact. These viruses and bacteria are released into the environment by various routes including water run-offs and aerosols. Furthermore, they can infect humans exposed to contaminated surface waters and ground water used for agriculture irrigation with severe consequences for human health (de Giglio et al. 2017). In most semi-arid areas, groundwater and surface waters constitutes an important and strategic resource, particularly as water stress increases and water resources of good quality become scarce (El Ayni et al. 2013). Molecular epidemiology and regular surveillance are necessary to elucidate the public health hazards associated with exposure to environmental viruses and bacteria (Cabral 2010; Rodriguez-Lazaro et al. 2012) especially in the South region where the water contamination combined to water scarcity strongly have large socio-economic impacts (UN-Water 2014). Climate projections for the Mediterranean climate areas estimate general warming and changes in precipitation distribution. Mediterranean coastal rivers are subject to flash floods during extreme events that transport the majority of the annual loads of bacteria and other contaminants (Chu et al. 2011). The frequency of extreme summer precipitation events increased over large regions of the Mediterranean (Giorgi and Lionello 2008) (Sections 2.2.5 and 3.1.3.3), increasing so the supply of fecal bacteria and viruses to the coastal zone. In a global context, wastewater management will be the key to preventing environmental dispersion of human



**Figure 2.24 | Harmful algal blooms** recorded before 1980 (A) and between 1980 and 2015 (B) along the Mediterranean coast (Cecchi et al. 2016).

fecal pathogens in future climate change scenarios (Rusiñol et al. 2015).

Harmful Algal Blooms (HABs) are sporadic phenomena triggered by massive proliferations of phytoplankton species reaching high cell concentrations (Sellner et al. 2003) that are suspected to have greater occurrences due to global warming (Hallegraeff 2010). HABs have environmental impacts (red-tide, mucilage production, anoxia) (Rodger et al. 2011; D'Silva et al. 2012) and represent serious economic threat for fisheries, aquaculture and tourism (Hoagland et al. 2002). They may also harm human health, since 40% of blooming microalgae are able to produce toxins responsible of different human intoxications (Santi Delia et al. 2015). HABs can occur in both freshwater and marine environments. In the Mediterranean Basin, marine ecosystems have received more attention although occurrence of HAB in freshwater lakes are reported (Cook et al. 2004; Romo et al. 2013) and are suspected to increase with climate change in the region (Romo et al. 2013) with consequences for potable water supply (Lévesque et al. 2014). Before 1980, HABs were rarely documented in the Mediterranean Sea. Since then, adverse events and several toxic episodes have been reported in different coastal regions (Fig. 2.24) (Cecchi et al. 2016; Garrido et al. 2016). Nowadays, harmful and toxic phytoplankton species become dominant in many coastal environments of the Mediterranean Sea.

Most toxic events in the Mediterranean Sea are mainly related to the dinoflagellates *Alexandrium* (known to induce Paralytic Shellfish Poisoning - PSP) and *Dinophysis* (producers of toxins causing the Diarrhetic Shellfish Poisoning - DSP) (Vila et al. 2001; Penna et al. 2007). *A. minutum* is the most observed dinoflagellate, with recurrent blooms ( $10^3$ - $10^8$  cells  $l^{-1}$ ) in the Ebro Delta and the Gulf of Gabès, inducing significant fish mortality (Abdennadher et al. 2012; Garcés and Camp 2012). Blooms of *A. catenella* ( $10^4$  cells  $l^{-1}$ ) have been observed both in the northern (Thau Lagoon) and the southern Med-

iterranean Sea (Bizerte Lagoon, Tunisia), causing sometimes shellfish contamination (Laabir et al. 2013). The epi-benthic dinoflagellate *Prorocentrum lima* (producer of DSP toxins) has been detected on most Mediterranean coasts (Garcés and Camp 2012), sometimes with high densities ( $>10^7$  cells  $l^{-1}$ ) (Aissaoui et al. 2014; Moncer et al. 2017). Blooms of *Karenia selliformis* ( $10^3$ - $10^5$  cells  $l^{-1}$ ), associated with intensive fish mortality, were reported for many years in the Gulf of Gabès (Feki et al. 2013). Recently, human health problems are caused by blooms of *Ostreopsis cf. ovata* ( $10^3$ - $10^4$  cells  $l^{-1}$ ) in Italy, Spain, Algeria and France (Ciminiello et al. 2014). New records of *Gambierdiscus* and *Fukuyoa* in the West and East of the Mediterranean Sea increase the risk of Ciguatera intoxication (Laza-Martinez et al. 2016).

Toxic diatoms *Pseudo-nitzschia* (producers of toxins causing the Amnesic Shellfish Poisoning - ASP) showed also blooms ( $10^3$ - $10^7$  cells  $l^{-1}$ ) in Mediterranean coasts of Spain, France, Greece, Italy and Tunisia, and contamination of mussels by ASP were reported (Sakka Hlaili et al. 2016). Mussel contamination by ASP toxin were reported in Spanish and Tunisian waters and have been linked to the blooms of *Pseudo-nitzschia* spp. and *Nitzschia bizertensis*, respectively (Giménez Papiol et al. 2013; Bouchouicha-Smida et al. 2015).

Toxic and harmful algal blooms continue to increase in magnitude, frequency and geographical distribution around the world and over the Mediterranean Sea (Hallegraeff 2010). The chronic eutrophication and climatic change including global warming are reported as significant factors involved in this global increase of HABs and toxic events (Anderson et al. 2012; Sakka Hlaili et al. 2016).

## 2.4 Land and sea use changes

### 2.4.1 Land use changes

#### 2.4.1.1 Past trends and recent dynamics

##### *Mediterranean landscapes*

The climatic conditions of Mediterranean Basin have played a key role for drawing its landscapes, which result from the interaction between human activity, a complex topography, and an extreme varied soil and climate (Pinto-Correia and Vos 2004). Altitudinal gradients and the distance from the sea are the main factors differentiating the landscapes of the Mediterranean Basin (Pinto-Correia and Vos 2004), and the main ecosystems contributing to outline these landscapes include the Mediterranean shores with a high number of habitats such as areas of dunes, rocks and wet areas. Moving away from the coast, Mediterranean maquis (a typical dense and closed shrub vegetation, mainly constituted by sclerophyllous species) and forests evolve, which includes evergreen shrubs or trees fully adapted to xeric conditions. In many cases, the maquis derives from the evergreen Mediterranean forest that develops in less water limiting growing conditions. Mediterranean steppic prairies represent the typical degradation of maquis that advances desertification. Finally, the Mediterranean deciduous forest extends where climates gradually shifts from typical Mediterranean to inland.

The modifications occurred in the Mediterranean landscapes are the result of land use practices that increased in intensity since the Neolithic, when in the Middle east domestication of plant and animals took the place of hunting and gathering (Blondel 2006). Human pressure on the natural ecosystem greatly varied depending on societal evolution developed over the area. In any case, wood exploitation has been key in localized periods of empires expansion. Forest destruction was the first step consequence of human pressure on the natural habitat as the effect of increasing agricultural activity and livestock and wood exploitation in the Mediterranean Sea trade cycle in order to cover the needs of the maritime empire (Barkaoui 2003). There is a long history of urbanization, reflected in countless archeological and historical settlements over the region (Diappi 2015).

As a result, the Mediterranean landscape is a complex mosaic of alternating semi-natural hab-

itats. Grasslands and pastures represent one of the most spread land use in European Mediterranean areas in the plain and low hilly areas (Cosentino et al. 2014) and they can provide forage for grazing animals or hay for conservation. The use of terraces in hilly areas permitted the cultivation on slopes of olive groves, vineyard and sowing crops while at the same their cultivation represented a means to reduce soil erosion, prevent run-off and increase water saving (Blondel 2006). The Dehesa-Montado System (typical of Spain and Portugal) is characterized by low density trees (represented by evergreen Mediterranean oaks) combined with crop production or pastoral activities, mainly represented by animal grazing. This system integrates the three main rural activities (forest/cork product harvesting, livestock husbandry, and agriculture) within a single landscape that consists of grass and trees. This system that combines extensive grazing of natural pastures, cereal cultivation and harvest of wood products, has shown remarkable stability, biodiversity, and sustained productivity over 800 years or longer as the result of the maintenance of botanically rich mosaic-like herbaceous plant layers (Joffre and Rambal 1993). In addition to agricultural production, these complex systems contributed, at the same time, to several ecosystem services such as preservation of the environment and its natural resources securing the sustainability of the system (Blondel 2006; Hao et al. 2017).

Despite small-scale traditional farming systems has been practiced for long time and it is still adopted in many parts of the Mediterranean region, huge changes in agricultural practices have taken place during the last 50-100 years across several Mediterranean areas as driven by increasing profitability of new agricultural systems (Debolini et al. 2018). The greatest changes involved uprooting of ancient and small-scale vineyards, orchards and olive groves, which turned into industrial scale fruit or olive plantations. Similarly, mixed rotational farming systems were simplified and replaced usually by intensive monocultures, requiring high inputs (i.e., high fertilization rate and water requirement) (Debolini et al. 2018). Intensive and large-scale farming systems also required the creation of new infrastructure, such as basins to cope with water shortage, which contributed to change the natural landscape. All these changes caused unsustainable pressure on the surrounding environment, resulting in either loss of wildlife-rich habitats and socio-economic

viability over large parts of the region due to land-abandonment of small-scale farmers.

### **Recent changes in Mediterranean landscapes**

The rate of change of landscapes in the Mediterranean Basin has increased since the second half of the 20th century. Many regions in Mediterranean Europe experienced the abandonment of marginal lands, especially in arid and mountain areas (Lasanta et al. 2017) (Section 3.2.3.1), and the following development by shrubs and tree species. Many studies show the abandonment of typical features like farming terraces, olive orchards, and upland grasslands leading to non-managed reforestation; for example, in 29% of the Iberian Peninsula for the 1989-2004 period (Hill et al. 2008); all-over Italy during 1990-2013, small forest patches cover increased a 27.4% (Sallustio et al. 2018), while 24% of pastures were turned into forests in areas of Tuscany from 1954 to 2005 (Amici et al. 2017); in Italian mountains, 16.3% of agricultural areas disappeared during the 1990's (Conti and Fagarazzi 2015); similarly in pre-alpine France during 1956-1991 (Taillefumier and Piégay 2003); in areas from the Eastern Mediterranean region of France, 14.2% of crops and 78.2% of pastures were converted into forests from 1958 to 2010 (Abadie et al. 2018); a 35.6% of vineyard areas in Serbia was transformed into meadows and pastures during 1985-2013 (Perović et al. 2018); Mediterranean islands as Elba, where up to 52% of agricultural areas were abandoned during the 1954-2000 period (Carta et al. 2018), and Lesvos, showing recent (2001-2011) slight decreases in agricultural land (Van der Sluis et al. 2016).

The remaining agricultural systems have generally become more intensive, with a shift towards livestock production and an increase of industrial inputs (fertilizers and pesticides), as it has been reported for Spain, especially since the 1960's to 2008 (Guzmán et al. 2018), and punctual areas of Greece and Italy, from 2001 to 2011 (Van der Sluis et al. 2016) (Section 3.1.2.1).

Conversely, scarce and mainly pre-21st century studies from North Africa show extreme land degradation due to overexploitation (Le Houérou 1995), principally by grazing pressure but also by forest conversion to agriculture and fuel-wood recollection. For the total North Africa and Middle East regions, the rate of deforestation increased from the 1980's to the 1990's by 160%, the fastest increase worldwide (Hansen and DeFries 2004). For example, in Morocco, it has been registered forest regression and degradation from 1962 to

1992 (Rejdali 2004), while in Rogassa (Algeria) a long-term experiment (1975-1993) demonstrated the main role of overgrazing in such degradation (Slimani and Aidoud 2004). Recent studies based in new observations and remote sensing show similar desertification trends in most parts of the Maghreb (Hirche et al. 2018). Grazing has been also the principal factor of forest degradation in northern Mediterranean islands as Crete during the 1977-1996 period (Hostert et al. 2003). Moreover, increases over 10% in livestock density have been observed between 2001 and 2011 in Portofino (Italy) and Lesvos (Greece) (Van der Sluis et al. 2016), and up to 40.1% in Nisyros (Greece) from 1991 to 2001 (Petanidou et al. 2008), showing that land abandonment does not always result in forest encroachment.

Changes in Mediterranean landscapes have been particularly intense in metropolitan areas and their surroundings. These landscapes are growing very rapidly all-over the Mediterranean (17% between 1990 and 2000) (Underwood et al. 2009) but especially in coastal areas. Urbanization mainly occurred at the cost of arable land (e.g., in Barcelona) (Basnou et al. 2013) and forested areas, as generally reported in the Mediterranean (Gerard et al. 2010). However, in some periurban areas, agricultural land has also increased following the growth of cities, as in Murcia (Spain) between 1995 and 2007 (García-Ayllón 2018). Examples could be found also in eastern Mediterranean, such as in Erdemli (Turkey), where the total length of the roads increased 23.6% between 2004 and 2015 following the growth of periurban agriculture (Alphan 2018).

### **2.4.1.2 Principal impacts of land use changes**

#### **Forestry and other natural resources**

New forests after land abandonment could alter biodiversity patterns in the Mediterranean (Fabbio et al. 2003), as has been demonstrated for the range of certain bird species (Gil-Tena et al. 2010). New forests are established by tree species whose dynamics have been favoured by recent land use changes, even more than climate (Améztegui et al. 2010). Meanwhile, other species as oaks are less benefited by such changes (Acácio et al. 2017), although general patterns show in fact successional dynamics towards *Quercus* dominance at the expense of *Pinaceae* (Alfaro-Reyna et al. 2018). Fire regimes are also altered, as fuel continuity is increased facilitating fire spread, which in turn could result in more landscape

homogeneity [Loepfe et al. 2010]. Also, forest continuity could increase forest insect pest spread [Hódar and Zamora 2004]. The effect of increased forest cover in the diminution of water resources is more conflictive although the pattern is clearer in catchments with records of large and rapid forest expansion [Gallart et al. 2011] (*Section 3.1.1.3*). During the first stages of land abandonment there is also a great risk of land degradation due to soil and nutrient loss [Thornes 2009]. Nevertheless, the contribution of these new forests to carbon storage is certain [Vilà-Cabrera et al. 2018].

Despite the natural forest recovery after land abandonment, forests in the Mediterranean Basin are still interested by many different threatening factors, such as wildfires, overgrazing, incorrect management and extreme climate and meteorological events (drought, windstorms) and this in turn could lead eventually to desertification [Vilà-Cabrera et al. 2018]. Wildfires become more and more frequent due to drier summers coupled with wood expansion due to land abandonment [Pausas and Fernández-Muñoz 2012]. Although evidence indicates that fires are decreasing due to increased efforts in fire suppression all-over Mediterranean Europe in more recent periods [Turco et al. 2016], there is large potential risk of mega-fires in the near future [Loepfe et al. 2011]. Changes of traditional silvicultural schemes able to produce high productive goals to a more extensive management or forest abandonment has led to deep modifications of composition and structure.

Another major factor affecting degradation of Mediterranean forests is grazing by domestic animals that utilize understory especially in period of reduced forage availability in pastures (such as summer or, in certain cases, winter). The traditional grazing in forest formation should not be banned as when the stocking rate is adequate, this result in a proper sustainable forest management that can reduce potential fuel biomass and can preserve this tradition landscape [Kairis et al. 2015] while, on the contrary, overgrazing can produce erosion, reduction in soil cover, losses of nutrients and in this way is it can be considered one of the most important factors of desertification in Mediterranean areas [Papanastasis and Kazaklis 1998]. Droughts, heat waves or windstorms can have a negative effect on many forests in the Mediterranean Basin producing reductions in forest growth and of forest declines [Vayreda et al. 2012] that prelude to land degradation and, in turn, desertification. All these factors produce stresses to forests that can be exacerbated in the next decades by climate change [Valladares et al. 2014].

Land degradation is the principal consequence of plant cover loss (by uncontrolled forestry, overgrazing, fires, etc.), which could lead to desertification in combination with increasing aridity and extreme climatic events [Thornes 2009]. In mountain environments, forest and understory cover loss, principally by overgrazing, have been clearly associated to higher erosion rates [Cheggour et al. 2012], as well as recent increases in forest cover were linked to the opposite [Barreiro-Lostres et al. 2017]. Soil erosion is a serious problem throughout the Tunisian Dorsal (the easternmost part of the Atlas mountain range). It has been estimated that 7% of the area is badly damaged by erosion and 70% of the area is moderately damaged [DG/ACTA 1993]. This degradation is an accumulated effect of agricultural strategies adopted over the Tunisian semi-arid areas during the last three millennia [Jebari 2009]. If we only consider short-term effects, the degradation can be partly attributed to the building of large dams during the 1960s and 1970s. This was done without giving sufficient attention to proper management of upper catchment areas [Jebari et al. 2010]. However, the contribution of the specific bioclimatic conditions of the Mediterranean climate in this degradation should not be underestimated. The soils are better characterized by the degradation of rock material rather than their organic matter content [Cerdan et al. 2004; Cudennec et al. 2007]. Consequently, they are not well developed and often shallow. The human influence is crucial on catchment scale in terms of landscape degradation that affects the hydrological regime [Jebari et al. 2010]. In the last hundred years, continuous changes were undertaken by the introduction of new crop cover (in fact, the most important change leading to better water management in the southern part of the Mediterranean may come from improving water efficiency in agricultural irrigation) [Berndtsson et al. 2016], deforestation, urbanization, river network modification, dam buildings and embankment. Fortunately, better water resource planning, reservoir maintenance, shortage, flood management, and hydro-agricultural infrastructure design are currently promoted [Verkerk et al. 2017] (*Section 4.5.1*).

Urbanization is considered a major driving force of biodiversity loss and biological homogenization [Grimm et al. 2008], causing landscape fragmentation, dramatic loss of open habitats and of the land use gradient, replacing adjacent land uses such as agricultural and more natural vegetation. Urbanization is also one of the main drivers of introduction of non-indigenous species, generating high propagule pressure, and frequent and intense

disturbance with complex consequences for biodiversity (Basnou et al. 2015; Clotet et al. 2016). The strong human pressure has also contributed to increase water shortages, pollution, forest fires, and the abandonment of ancient pastoral regimes. Recent studies also demonstrate the negative consequences of new artificial areas in coastal dunes habitats, both affecting carbon stocks (Carranza et al. 2018), and generating more complex effects when stabilizing the natural changing dynamics of dune systems (Manzano et al. 2019).

### **Crops and livestock**

Food production in the Mediterranean region is changing rapidly, due to multiple local and global social and environmental changes. The increased number of urban and displaced people increase demand for food in urban areas, with limited agricultural production and with great water restrictions (FAO 2017a). Scarce resources, such as fisheries, are being exhausted (see next section). Evidence for the limited capacity to cope with these challenges can be documented in recent history. For example, water reserves were not able to cope with extensive droughts in the last two decades in Spain, Morocco and Tunisia, causing many irrigation dependent agricultural systems to cease production (Faurès et al. 2002; Garrido et al. 2006; FAO 2015).

Livestock production, mainly located in semi-arid and arid lands has shifted from extensive modes to systems heavily dependent on feed grain (32% of total food imports), inducing high poverty rates and rural exodus and rendering production sensitive to climatic shifts elsewhere (Sections 3.1.2.1 and 4.5.1).

Besides soil erosion, the major land degradation processes in the Mediterranean Basin are soil sealing, compaction mainly due to agricultural intensification, salinization, and contamination due to industrial activities. Soil organic carbon stocks tend to decrease when transforming grasslands, forest or other native ecosystems to croplands and to increase when restoring native vegetation on former croplands or by restoring organic soils to their native condition. Permanent and traditional woody cultivation such olive tree and grapevine may compensate this trend due to their positive contribution of their carbon uptake.

Human society will have to rely on an increase of output per unit area in agriculture and forestry. Intensively used agricultural systems are often N-saturated and the augmented use of fertiliz-

er increases the leaching of N into aquifers and aquatic ecosystems and thus carries costs to environmental services such as water quality. In summer irrigated crops conditions are propitious for high N<sub>2</sub>O losses. Emission factors for N<sub>2</sub>O, distinguishing the effects of water management, crop type, and fertilizer management. Mediterranean agricultural soils produce large CH<sub>4</sub> emissions in flooded crops (e.g., rice) through methanogenesis, representing 6% of all CH<sub>4</sub> production from agricultural sources (Section 3.2.3.2).

During the past two decades, rural areas were reshaped by technological improvements in resources exploitation, the accelerating abandonment of traditional rural life and an increase in the mobility of individuals (Pinilla et al. 2008; Domon 2011). This pathway has led in many developed countries to a particular land-cover change pattern that consists in low plains and coastal areas that are being increasingly utilised for human activities due to their higher potential for agricultural productivity, while mountain or marginal areas are being abandoned because no more economically viable for production (Statuto et al. 2016; Nori 2018). Considering the increasing demand of food for an ever-growing population, leading to an increased productivity and intensification efforts in producing areas (Phelps and Kaplan 2017), this pattern has noticeable effects on the patchwork alternating semi-natural habitats that characterized the Mediterranean environment and the relevant natural ecosystem services. Management practices of grassland in hilly Mediterranean areas were progressively reduced because of reduction of animal grazing and abandonment. This produced remarkable effects on floristic simplification, loss of biodiversity, reduction of habitat for wildlife or to the survivor itself of the resource due to shrubs encroaching (Papanastasis 2004; Argenti et al. 2011). Perennial cultivation, such as olive tree grove, originally planted in marginal areas on terraces and representing a quite stable ecosystem managed with few chemical inputs, were replaced by intensive modern plantations managed under an intensive and highly mechanized system. This intensification resulted into a progressive abandonment of marginal areas because of their low economic viability leading to a shift of land use to pasture for sheep and goats, since the land is not suitable for any other kind of cultivation (Loumou and Giourga 2003).

Conversely, the intensification process towards highly mechanized and high-density plantation is boosting farmers' income but it is also causing some



environmental drawbacks. The combined effect of a more intensive management and cultivation extension is rising the issue of contamination by excess in the use of synthetic fertilizers and other agrochemicals to increase land productivity (Beaufoy and Pienkowski 2000; Beaufoy 2001). Intensification results into a degradation of habitats and landscapes and the exploitation of scarce water resources, thus putting the naturally scarce resources of olive growing areas to an edge. Drip irrigation, associated to an improved efficiency of irrigation, often has in fact no effect on efficiency but increases global water consumption. The use of this technique decreases the need for human power, allowing the increase of cultivation area and fostering multiple cropping (Kuper et al. 2017; Molle and Tanouti 2017). Soil erosion is also a growing issue due to the widening of cultivated area, as in tree crops a significant fraction of the soil is vulnerable to the action of rainfall and runoff. The intensification of agricultural land use has therefore raised the question of the long-term sustainability of agroecosystems (Liebig et al. 2004).

### ***Different land uses as a single driver of change***

Demography, technology, socio-economic factors and climate change have gradually transformed Mediterranean landscapes. Land uses are in fact the combination between the use of natural and food resources, constructions, road networks, etc., and land use changes result in the balance of their consequences. Natural habitats as affected by human activities are consequently reduced in size and continuity finally resulting into a loss of connectivity, i.e., the capability of the landscape to help or to prevent movements of organisms across habitat components (Taylor et al. 1993). A reduced or lack of connectivity may have consequences on biodiversity through losses of ecological fluxes between habitat patches and therefore trends in connectivity across different landscapes should be evaluated to consider proper actions to counteract potentially negative impact of human pressure on animal or vegetal biodiversity (Hernández et al. 2015). Over the Mediterranean Basin, two main trends, reforestation following the abandonment of agricultural areas in hilly and mountainous areas and the relevant expansion of agricultural and artificial areas in the coastal areas, plains and valleys, had different impacts on connectivity. In particular, reforestation after agricultural abandonment was correlated to a slight improvement in the connectivity of the European forest in the period 1990-2000 (Saura et al. 2011). At the same time, the spread of monoculture in

the plain resulted in a de-fragmentation process. Both these trends produced a simplification and homogenization of the landscape, in terms of number, dimensions and typology of the patches that shape the agro-ecological territory.

### ***2.4.1.3 Future projections***

Land use change is expected to have different consequences on the productivity of several ecosystems and the carbon balance. The expected warmer and dryer conditions on southern Mediterranean Basin will likely shift crop cultivation to North, where water deficit is projected to be less harsh (Ceglar et al. 2019). Whilst the less adaptable crop systems will likely suffer changed pedological and climatic conditions, the most resilient crops, such as olive tree and grapevine may have the potential to resist to this trend due to their high adaptability to cope with high temperatures and water scarcity. Moriondo et al. (2013) and Hannah et al. (2013) predicted a gradual northward shift of this cultivation in the medium term (Tanasijevic et al. 2014). Increasing temperature joint with the expected reduce rainfall rate may lead detrimental consequences for those cultivations such as corn, rice and spring wheat requiring wetter conditions. Some crops may be replaced with more resilient crops such as barley, sorghum and hay, which, however, may not well fit with the market demand and the production chain of the Mediterranean area. For most of the main Mediterranean vegetable and cereals, a decrease in production is expected in the absence of specific high input agronomic strategies such as fertilization and irrigation (Bregaglio et al. 2017; Ruiz-Ramos et al. 2018; Brilli et al. 2019) whose requirement is expected to increase (Tanasijevic et al. 2014; Saadi et al. 2015).

Climate change impacts are expected to also affect managed forests, leading to shift in typical forest communities to higher altitudes (Gitay et al. 2001). These impacts will likely affect the whole woody spinneret, influencing timber extraction and plantations, and management practices. These latter practices (e.g., fuelwood collection, forest grazing, and road expansion) can degrade the forest ecosystem conditions, particularly when applied over new forest area. Accordingly, depredated soil and forestry systems may indirectly favor the introduction of pests and pathogens, changing fire-fuel loads, changing patterns and frequency of ignition sources, and changing local meteorological conditions (Nepstad et al. 1999). Grassland and pastures will likely experience a further decrease in extension due to a progressive

rural abandonment and emigration to urban areas, often associated with low-income level in mountain areas and lack of job opportunity (Sturaro et al. 2013), and the impacts of climate change. This latter will particularly impact natural pastures, which are acknowledged to be very sensitive and vulnerable to climate conditions.

The predicted climate warming is also expected to lead changes of grasslands structure and composition, increasing the soil-water competition with trees that will be found to place at higher altitude as effect of higher temperatures. In temperate climate, warming may lengthen the forage growing season but decrease forage quality, with important variations due to rainfall changes (Craine et al. 2010; Hatfield et al. 2011; Izaurre et al. 2011), whilst Mediterranean pastures will likely show production decrease due to prolonged drought conditions. Modern agricultural practice can partly overcome expected production due to changed cultivation areas, but plant adaptation would require more time to be able to adapt to the new climate. Also, higher input needed to cope with changed agronomic conditions can lead to wrong perspective of the crop, forest and grassland production trend as well as extensive environmental damage. More specifically, a larger use of fertilizer or other high inputs may result in short-term increases in food production for long-term losses in ecosystem services, such as water quality degradation (Zalidis et al. 2002; Malagó et al. 2019), soil erosion, reduced fertility, or overgrazing (Wood et al. 2000). All these changes are expected to affect the carbon balance.

Land use is an important control of carbon storage; therefore, ecosystem shifts and harsher climatic conditions may lead to different forms of stress (i.e., water and nutrient stress, pedological stress, climatic stress, abiotic, etc.) which, in turn, reduce the potentiality of the different ecosystems in terms of carbon storage (i.e., biomass reduction, less growth and development). All these changes can have negative consequences in the perspective of CO<sub>2</sub> mitigation capacity (Foley et al. 2005), since is expected a decrease of total carbon sequestration capacity from agro and forestry systems, increased carbon fluxes from soils due to quicker decomposition process and lower carbon mineralized in soil.

### 2.4.2 Sea use changes

Fisheries (over) exploitation is the main driver of marine population decline and has led to the bad

state of most highly commercial stocks and the low abundance of top predators. Climate change and variability may be responsible for catch fluctuations of some stocks (especially the small pelagic fishes), for distribution shifts but also for altering catch composition in favour of warm-water species. Recent theory predicts fish size decreases in response to increased sea surface temperature and low oxygen supply. Excessive exploitation will certainly lead to even lower stock biomasses, especially for top predators. Further sea warming will very likely lead to a higher percentage of warm-water species in the catch and smaller fish sizes.

#### 2.4.2.1 Trends in fisheries exploitation

In the Mediterranean Sea, which together with the Black Sea, constitutes FAO Major Fishing Area 37, fishing has been practiced since antiquity. Today, Mediterranean fisheries are diverse among areas and the fishing vessels and techniques vary geographically as a result of different environmental, oceanographic, biological, climatic, cultural and socio-economic conditions prevailing in each area (Papaconstantinou and Farrugio 2000), with a strong contrast between the northern and southern coastlines. The high number of islands, ports and shelters across the Mediterranean and the contrast between north and south renders the enforcement of fisheries regulations and management very difficult. Mediterranean fisheries are highly multispecies in nature targeting over 200 fish and invertebrate species (Dimarchopoulou et al. 2017) and are operated through a large number of small sized and low tonnage fishing vessels with no large industrial fleets (Stergiou et al. 2016). The number of small-scale coastal vessels operating in Mediterranean EU waters is about 86% of the total (around 72600 vessels) with the remaining 9% being trawlers and 5% being purse seiners (Colloca et al. 2017). Although the number of all types of Mediterranean EU fishing vessels declined since 1991, the actual fishing effort has been increasing due to new technologies and higher capacity vessels (Colloca et al. 2017).

Small pelagic fisheries operate all year round but in many Mediterranean Sea subareas they show a strong seasonality that is reflected upon their catches and is derived from fishing regulations and consumer habits. According to the monthly distribution of landings and fishing effort, the main fishing season in most areas is concentrated in spring and summer months (Lloret et al. 2004b).

### 2.4.2.2 Current status of marine fisheries resources

In the Mediterranean Sea, recent publications based on scientific surveys, stock assessments and catch data, generally agree that the majority of Mediterranean fisheries stocks are declining in biomass as a result of their overexploitation (Colloca et al. 2013, 2017; Vasilakopoulos et al. 2014; Tsikliras et al. 2015). Local reports also confirm the bad status of Mediterranean fisheries, e.g., in Greek seas (Tsikliras et al. 2013b) and in the Ligurian Sea (Abella et al. 2010), often attributed to inadequate management practices (Tsikliras 2014; Cardinale et al. 2017). The long-lasting overexploitation of the Mediterranean Sea has been driving the decline in biomass of most commercial fish and invertebrate stocks across the basin and the near depletion of several of them (Froese and Kesner-Reyes 2002; Vasilakopoulos et al. 2014; Osio et al. 2015; Tsikliras et al. 2015; Stergiou et al. 2016; Colloca et al. 2017; Froese et al. 2018). The overall stock status is rather uniform across the Mediterranean ecoregions with low stock biomass being the common characteristic. However, the stock specific biomass levels vary among ecoregions (Froese et al. 2018).

The catch history of Mediterranean Sea stocks unmasked the overexploitation of many stocks since the 1950s, when about 40% of them were declining in biomass (Froese and Kesner-Reyes 2002). Recent literature reveals that fisheries overexploitation occurs across the entire area (Tsikliras et al. 2013a) and locally, e.g., in Greek Seas (Tsikliras et al. 2013b). Several Mediterranean stocks have been reported overfished based on data from landings (Tsikliras et al. 2013b, 2013a), scientific surveys (Stergiou and Tsikliras 2011), or stock assessments (Colloca et al. 2013). Other studies confirm that almost all species targeted by the fishing fleets are being overexploited in the Mediterranean Sea (Cardinale et al. 2017; Fernandes et al. 2017).

Based on the catch-based method, the cumulative percentage of collapsed and overexploited stocks appeared to exceed 60% across the Mediterranean Sea in 2010 (Tsikliras et al. 2013a) with the exploitation pattern differing among the Mediterranean subareas (Tsikliras et al. 2015). The western Mediterranean has been reported to be in a better state with less overexploited and collapsed stocks and more developing ones compared to the central and eastern parts of the sea (Stergiou et al. 2016). Similarly, based on various fisheries indicators, the western and central Mediterranean are in

better condition compared to the eastern part of the sea (Tsikliras et al. 2015). According to the official stocks assessments that were then available, the percentage of overexploited stocks exceeds 90% in most areas (Colloca et al. 2013) and even reaching 95% in some (Osio et al. 2015). The model approach of Osio et al. (2015) estimates that 98% of the unassessed demersal fish species are potentially overexploited in most areas. Cardinale et al. (2017) reported that the stocks of all target species that have been assessed are overexploited with the average ratio of  $F/F_{MSY}$  (actual fishing mortality to the level that would provide maximum sustainable yield) ranging from 1.7 (giant red shrimp *Aristaeomorpha foliacea*) to 8.1 (hake *Merluccius merluccius*). Steadily increasing exploitation rates and deteriorating gear selectivity have been recently reported as two conditions that lead to shrinking fish stocks (Vasilakopoulos et al. 2014). The most recent assessment of 169 Mediterranean stocks showed that 126 of them (75%) were subject to ongoing overfishing (Froese et al. 2018).

### 2.4.2.3 The future of marine resources

The Gill-Oxygen Limitation Theory (GOLT) predicts a reduction in the size of fish due to their inability to compensate, via their gill surface, for the increased metabolic rate that results from higher temperatures. Fish individuals that survive are expected to shrink in size (Cheung et al. 2013a). The Mediterranean Sea is among the semi-enclosed areas where local species extinctions and range shifts were predicted to be most common (Cheung et al. 2009). The GOLT theory may also explain the poleward shift of marine organisms (Cheung et al. 2013b) and their expansion to deeper waters (Perry et al. 2005) both of which occur in the Mediterranean Sea (Tsikliras and Stergiou 2014) and may have an impact on Mediterranean fisheries in terms of catch and revenue (Cheung et al. 2010).

Besides fish distribution shifts and declines in local fish stocks, scientific projections suggest that marine resources and biodiversity will suffer increasing stress if temperatures are not held below 2°C above preindustrial levels (Gattuso et al. 2015). Sea warming and deoxygenation combined with fishing pressure and other stresses could affect growth, and distribution of fish populations, resulting in changes in the potential yield of exploited marine species and economic losses (Sumaila et al. 2011) as fisheries are expected to decline (Cheung et al. 2010). Reaching the goals of the UNFCCC Paris Agreement would benefit ocean life and economies by protecting millions of metric

tons of high valued catch with 75% of maritime countries benefiting from this protection (Sumaila et al. 2019).

Fisheries in the Mediterranean will not be sustainable in the future unless the marine exploited populations are fished less, i.e., if they are allowed to recover and rebuild their biomass (Pauly and Zeller 2016) through the reduction of the

fishing pressure that is applied upon them (Froese et al. 2018). Ecosystem-based approach has also an important role that will ensure that both higher and the lower trophic levels are rebuilding (Pikitch et al. 2004) and fully marine protected areas are a key management tool to accomplish rebuilding of the biomass of marine populations, ensure ecosystem health and resilience against sea warming (Roberts et al. 2017).

## 2.5 Non-indigenous species

The human-aided introduction of non-indigenous species into new biogeographic regions – has been one of the main increasing global drivers of ecological change for over a millennium (Elton 1958). Non-indigenous species homogenize biodiversity across the globe, resulting in shifted and sometimes more simple ecological communities (Mooney and Hobbs 2000; Rilov and Crooks 2009). They can displace native species out of their natural habitats through competition, consumption, or parasitism. This displacement results in affected communities with reduced native species diversity, altered species composition as well as in major changes in ecosystem functioning (Vilà et al. 2011; Cameron et al. 2016). Moreover, these changes alter supporting, provisioning, regulating ecosystem services that people depend upon, impacting human well-being (Katsanevakis et al. 2014b; Vilà and Hulme 2017).

Rates of introduction and impacts have accelerated dramatically in the past few decades due to various human activities and related pathways of non-indigenous species introduction (Carlton 1989; Crooks and Suarez 2006). In the marine environment, the growth of seaborne trade with its huge fleets facilitates the dispersal of organisms attached to the hulls of ships and inside ballast water (Crooks and Suarez 2006; Katsanevakis et al. 2013). Aquaculture, live marine seafood and bait, and aquarium trade have also become important vectors for the introduction of marine non-indigenous species, as well as the artificial connections of water bodies with very different biotas (Rilov and Crooks 2009), and major catastrophes like tsunamis (Carlton et al. 2017). In terrestrial ecosystems, intentional introductions prevail over unintentional. For non-indigenous plants, ornamental and horticultural introductions escaped from cultivation account for the highest number (Lambdon et al. 2008) and are increasing steadily (Van Kleunen et al. 2018). Terrestrial non-

indigenous vertebrates follow similar patterns as plants. However, most terrestrial non-indigenous invertebrates have been introduced accidentally; many are major pests in forestry and agriculture (Roques et al. 2010).

### 2.5.1 Non-indigenous species in the Mediterranean Sea

The first introduction of marine species into the Mediterranean Sea dates back to the late 18th century (Poli 1791). Since then, due to maritime shipping expansion and after three centuries of accumulating human pressures, this basin has become a hotspot of introduction of non-indigenous species (Rilov and Galil 2009; Coll et al. 2010). Non-indigenous species in the Mediterranean Sea mostly arrive from the Indo-Pacific region either directly (by swimming or drifting) or indirectly as foulers or as hitchhikers inside ballast water in a process called Lessepsian migration (Por 1978). But there are many other vectors that deliver non-indigenous species into the Mediterranean and have varying importance depending on the region (Rilov and Galil 2009). Today, the total




Figure 2.25 | Number of new non-indigenous species per 6 years in the Mediterranean since 1988 (Zenetos 2019).

known number of non-indigenous species is still debated, but the last count puts it close to a thousand species [Zenetos et al. 2017]. For some of the species, the impact on local biodiversity, and possibly also ecosystem functions and services, seems obvious, but in most cases, the impact is unknown because of lack of research [Katsanevakis et al. 2014b]. The Mediterranean Sea is warming rapidly [Nykjaer 2009; Sisma-Ventura et al. 2014] (Section 2.2.4). Therefore, it is quite possible that the establishment of thermophilic non-indigenous species is strongly facilitated by climate change [Stachowicz et al. 2002].

### 2.5.1.1 Spatiotemporal trends, sources and vectors of introduction

The most recent assessment of marine non-indigenous species in the Mediterranean (November 2018) counts 957 species, including Foraminifera [Zenetos 2019]. Among the introduced species during the last 30 years (491 taxa), invertebrates dominate with >58% (287 species) represented mostly by molluscs and decapods. Primary producers follow with approximately 114 species among which macroalgae, especially rhodophytes, prevail. Vertebrates (mostly fishes) follow with 90 species.


The trend in introduction of species in the Mediterranean (Fig. 2.25), which culminated in the 2000-2005 period with more than 20 new species per year (122 in total), appears to be overall decreasing after 2005 (Fig. 2.26). As opposed to invertebrates and primary producers, vertebrates continue increasing, with 34 species detected in the 2012-2017 period vs 26 species in the period 2006-2011. The overall decreasing rate in non-indigenous species is evident across the Mediterranean Marine Strategy Framework Directive (MSFD) areas, except for the Central Mediterranean where an increase is observed, attributed to vertebrates (Fig. 2.26). 17 new fish species were detected in the central Mediterranean in the period 2012-2017 vs. 8 fish species in the period 2006-2011. These species are either spreading from the eastern Mediterranean to the central region or are newly introduced species in the area. Vertebrates (fish only) are dominated by Lessepsian immigrants but over the last decade the number of fish species related to aquarium trade which have been intentionally released to the wild (classified as escapees from confinement) is increasing [Zenetos et al. 2016; Marcelli et al. 2017; Deidun 2018].

Regarding the spatial distribution of non-indigenous species, the number of Lessepsian species is very high on the eastern Mediterranean coastline, reaching 129 species per 100 km<sup>2</sup>, and declines toward the north and west [Katsanevakis et al. 2014a]. The distribution of species introduced by shipping is strikingly different, with several hot-spot areas occurring throughout the Mediterranean Basin. Two main hotspots for aquaculture-introduced species have been identified (the Thau and Venice lagoons). Certain taxonomic groups were mostly introduced through specific pathways—fish through sea corridors, macrophytes by aquaculture, and invertebrates through sea corridors and by shipping [Katsanevakis et al. 2014a]. Hence, the local taxonomic identity of the non-indigenous species is greatly dependent on the dominant maritime activities/interventions and the related pathways of introduction. The composition of non-indigenous species assemblages differs among Mediterranean ecoregions; such differences are greater for Lessepsian and aquaculture-introduced species.

### 2.5.1.2 Non-indigenous species as drivers of biodiversity and ecosystem change

The introduction of non-indigenous species in the Mediterranean Sea have caused modifications in biodiversity patterns. One of the best documented and most profound impacts of the introduction of non-indigenous species on native Mediterranean ecosystems is the deforestation of algal forests and the creation of extent barrens (i.e., areas with bare rock and encrusting calcified algae; Fig. 2.27) by the overgrazing activity of two non-indigenous herbivore rabbitfishes: *Siganus luridus* and *S. rivulatus*. These species have become dominant in the ichthyofauna of shallow rocky habitats in the eastern Mediterranean, and have caused ecosystem-wide changes by creating and maintaining areas denuded of canopy algae (an important habitat for many coastal fishes) [Cheminée et al. 2013]. This form of "deforestation" is associated with a dramatic reduction in biodiversity, biomass, and algal growth, and effects that move up the food chain to the local fisheries [Sala et al. 2011; Vergés et al. 2014b].

In the Levantine Sea, the catch of commercial fisheries is now dominated by non-indigenous species [Edelist et al. 2013; Katsanevakis et al. 2018], reflecting the decline of native biota and its replacement by thermophilic non-indigenous species [Arndt et al. 2018]. In Turkish coastal waters in the Levantine Sea, non-indigenous



**Figure 2.26 | Change in the number of new non-indigenous species per 6 years in the Mediterranean MSFD regions** (Adriatic, Western Mediterranean, Central Mediterranean, Eastern Mediterranean Seas) since 1988. Notice that the scale is different for the eastern Mediterranean (Zenetos 2019).

fish species biomass exceeds 50% of the total fish biomass and 90% of the total herbivore fish biomass (Giakoumi et al. 2019). Fishers in the same area (Kaş, Turkey) perceived the introduction of non-indigenous species as the most important reason for the current fish stock depletion.

Yet, it is uncertain whether the decline of native biota in the Levantine Sea is driven mainly by biotic interactions with non-indigenous species or climate change, or both, as the Lessepsian species are thermophilic and their establishment is assisted by climate change (see discussion below). A recent meta-analysis suggests that the decline of native fish can be mostly attributed to ocean warming and not to negative interactions with non-indigenous species (Givan et al. 2017a). Multi-species collapses of native species are also at least partially attributed to climate change (Rilov 2016), with the collapse of the sea urchin *Paracentrotus lividus* experimentally demonstrated to be related to the fast ocean warming (Yeruham et al. 2015).

Analysis focused on shallow reef fish shows that non-indigenous species are very diverse ecologically, and they considerably increase the total com-

munity trait diversity of the Mediterranean (Givan et al. 2017b). Furthermore, trait similarity between non-indigenous and indigenous Mediterranean species was lower than expected, indicating that non-indigenous fish tend to occupy relatively vacant niches within the Mediterranean. Temporally, non-indigenous fish species display increased trait similarity to native Mediterranean species, suggesting that forecasting future establishment may be challenging. Givan et al. (2017b) conclude that the Mediterranean, at least in fish, is transforming into an extension of the Red Sea in terms of trait and species composition. Such biological trait analysis is also required for other taxonomic groups.

There is a serious research gap in assessing and quantifying the impacts of non-indigenous species on marine ecosystems in the Mediterranean Sea. Impact assessment is mostly based on expert judgment or correlational studies, while manipulative or natural experiments are largely lacking for assessing the impacts of most non-indigenous species in the region (Katsanevakis et al. 2014b). Disentangling the role of the introduction of non-indigenous species and climate change or other local or global stressors to derive cause-



**Figure 2.27 | Ecosystem shift from algal forests to barrens due to the overgrazing activity of non-indigenous herbivore rabbitfish.** Algal forests host high fish, invertebrate, and algae biodiversity whereas barrens are associated with low levels of biodiversity across all taxonomic groups. *Rabbit fish photo: Murat Draman.*

effect pathways is inherently difficult and would probably necessitate a combination of experimental and modelling approaches.

Efforts are made to assess non-indigenous species impacts with existing knowledge. Based on a conservative additive model, which downgrades reported impacts of low inferential strength, an index of the Cumulative IMPacts of invasive ALien species (CIMPAL) on marine habitats in the Mediterranean has been developed and estimated (Katsanevakis et al. 2016). The estimation of CIMPAL was based on assessments of impacts for every combination of 60 non-indigenous species and 13 habitats, and their distributions in the Mediterranean (i.e., presence in 10x10 km cells). It showed strong spatial heterogeneity in impacts. Spatial patterns varied depending on the pathway of introduction of the non-indigenous species in the Mediterranean Sea. Species introduced by shipping gave the highest impact scores and impacted a much larger area than those introduced by aquaculture and through sea corridors. Overall, non-indigenous macroalgae had the highest impact among all taxonomic groups, when estimated as the sum of impact scores across the entire Mediterranean Sea, i.e., accounting not only for the severity of the impacts but also their spatial extent. The most impactful non-indigenous species was *Caulerpa cylindracea*, which has become dominant over large areas of shallow waters in the western Mediterranean and compete with native species (Piazzi et al. 2016). Negative impacts of *C. cylindracea* have been documented on al-

gal, sea grass, and sponge communities (Ceccherelli et al. 2002; Piazzzi et al. 2005; Piazzzi and Ceccherelli 2006; Baldaconi and Corriero 2009).

### 2.5.1.3 Further introductions, monitoring and managing non-indigenous species

The introduction of species in the Mediterranean Sea is a continuous process and it is very likely that it will continue for years to come. In most of the cases, these species fail to thrive but, evidently, some become numerically and ecologically dominant in their new environment, generating new and sometimes severe impacts of the introductions. As a consequence, in the past two decades research interests in non-indigenous species have increased, mostly stimulated by evidence on their ecological and socio-economic impacts in the Mediterranean region. This has also raised the urgency of innovative approaches to forecast, track and manage these species (Corrales et al. 2018).

One of the most recent and potentially damaging non-indigenous species for this basin is the common lionfish (*Pterois miles*), which increasingly appears in many parts of the eastern Mediterranean in the last few years (Bariche et al. 2013). Due to its rapid increase in abundance (Kletou et al. 2016) and fast geographical expansion (Azzurro et al. 2017), this harmful species has become emblematic for raising concern on Mediterranean non-indigenous species introductions but it also well illustrates a process of developing monitoring capabilities and

management strategies within the Mediterranean region.

Documenting the spread of this non-indigenous species can greatly benefit by the participation of resource users, a partnership which can support monitoring objectives (Azzurro and Bariche 2017) and be used to reduce, at least locally, the abundance of non-indigenous species (Kleitou et al. 2019). More generally, participatory approaches benefit the scientific consensus, which is a key element for both documenting and responding to these introductions (Scyphers et al. 2015). This is also reflected in some of the guiding principles on the management of non-indigenous species adopted by key regional and international bodies/legislative frameworks, concerning non-indigenous species, such as those provided by the EU (Regulation 1143/2014); UNEP-MAP and by FAO-GFCM, which are converging towards finding common strategies to face Mediterranean species introductions. In this regard, the Integrated Monitoring and Assessment Programme and related Assessment Criteria (IMAP) adopted through Decision IG.22/7 by the 19th Ordinary Meeting (COP 19, Athens, Greece, 9-12 February 2016) of the Contracting Parties to the Barcelona Convention, stress the need of comprehensive monitoring and coordinated transnational actions to face the common issue of Mediterranean non-indigenous species introductions. A recent expert assessment of management options of marine non-indigenous species prioritized 11 management actions for controlling 12 model species according to their dispersion capacity, distribution, and taxonomic identity (Giakoumi et al. 2019). The actions were assessed using five criteria (effectiveness, feasibility, acceptability, impacts on native communities, and cost), combined in an "applicability" metric. Raising public awareness and encouraging the commercial use of non-indigenous species gained the highest priority, and biological control was considered the least applicable (Giakoumi et al. 2019).

To predict future change in the distribution of native species and the spread of non-indigenous species, species distribution models (SDM) are regularly used. In these models, climate matching is calculated between the area of origin (donor) and the area of potential spread (recipient). However, a recent study that matched the native range of Red Sea fish and their new range in the Mediterranean Sea showed poor matching, and thus indicated that SDMs may underestimate the potential spread of non-indigenous species (Parravicini et

al. 2015). The authors call for caution in employing such models for forecasting the introduction of non-indigenous species and their response to environmental change, as uncertainty is large. Better knowledge of the fundamental niche of species (their physiological performance under different environmental conditions) can potentially improve the prediction of spread of non-indigenous species in the Mediterranean. Furthermore, an analysis combining ship movements with port environmental conditions and biogeography can be used to quantify the probability of new primary introductions through ballast water (Seebens et al. 2013).

## 2.5.2 Terrestrial non-indigenous species and pests

### 2.5.2.1 Spatial patterns and temporal trends


#### *Degree of introduction across Mediterranean-type ecosystems and geographical areas*

The information available on terrestrial non-native species in the Mediterranean Basin countries is not comprehensive, and the number of non-indigenous species is underestimated due to the incompleteness of collected data and the monitoring bias towards some taxonomic groups. For example, Abellán et al. (2016) reported more than 370 non-indigenous birds for Spain and Portugal, more than twice the number listed by DAISIE (2009) for the whole Europe.

Most non-indigenous species in the Mediterranean Basin are plants, followed by invertebrates (Fig. 2.28). Natural habitats in the Mediterranean Basin host more than 400 non-indigenous plant species (Arianoutsou et al. 2013). The taxonomic similarity of the non-native flora among Mediterranean Basin countries is very low. For example, less than 30 species are common across 4 Mediterranean European countries (Arianoutsou et al. 2010), and only 10 species are common in 8 major Mediterranean islands (Lloret et al. 2004a). However, non-native plants share similar traits, as shown by the high proportion of perennial herbs with a long flowering period, which are pollinated and dispersed by the wind (Lloret et al. 2004a, 2005).

The major part of non-indigenous invertebrate species are arthropods, especially insects (Roques 2010), with a low representation of nematodes and flatworms (Naves et al. 2016; Justine et al. 2018). Phytophagous pest species are largely dominating among non-indigenous species all





**Figure 2.28 | Number and proportion of terrestrial non-indigenous plant, invertebrate and vertebrate species per country** (EASIN)<sup>a</sup>.

over the Mediterranean Basin, accounting for more than a half of the invertebrate species and about one-third are associated with woody plants (Matošević and Pajač Živković 2013; Roques 2015; Avtzis et al. 2017). Among them, Hemipterans, mostly scales and aphids, constitute the dominant group, accounting for 40-75% of the non-indigenous species in any part of the Mediterranean Basin (Roll et al. 2007; Inghilesi et al. 2013; Matošević and Pajač Živković 2013; Seljak 2013; Avtzis et al. 2017). This over-representation of Hemiptera in non-indigenous species seems related to their small size and easier of transportation with infested imported plants.

A complete plant pathogens database is not available. However, a European database of non-indigenous forest and woody pathogens indicate a list of 123 plant pathogens. In Mediterranean countries, Ascomycota is the most numerous group, while Oomycota and Basidiomycota represent 21% and 9% of the total, respectively (Santini et al. 2013). With regard to vertebrates, most non-indigenous species are birds, followed by mammals, while the number of reptiles and amphibians is low.

<sup>a</sup> <https://easin.jrc.ec.europa.eu>

The impact of non-indigenous species varies largely among countries. Countries with higher Human Developmental Indexes and imports host a large density of non-native plants (Vilà and Pujadas 2001). Between countries and within countries, the density of non-native plants is related to the length of terrestrial transport networks. Areas with extensive road and rail networks, high anthropogenic disturbance, low altitude, short distance to the coastline and dry, hot climate show higher richness of non-indigenous plant species (Gassó et al. 2012). Most affected landscapes are those highly urbanized and with high population densities (Sobrino et al. 2002; González-Moreno et al. 2013). Accordingly, the most affected ecosystems by non-indigenous plants are human-modified such as ruderal, waysides or agricultural fields (Vilà et al. 2007; Hulme et al. 2008; Arianoutsou et al. 2010).

As for other non-indigenous taxa, most pests are introduced in managed habitats, such as agricultural lands and parks and gardens, forests being less affected (Matošević and Pajač Živković 2013). Among the infested plant species, non-native ornamental plants (palms, legume trees),

*Citrus* and *Eucalyptus* are slightly more colonized than native species (Roques 2015). Vertebrates also tend to occupy anthropic habitats and, to a lower extent, woodlands (at least for birds) (Kark et al. 2009). The degree of introduction of non-indigenous invertebrate species also varies among countries, Italy and France showing much more established non-indigenous species than any other European country (Roques 2010). The same trend is reported also for plant pathogens (Santini et al. 2013).

Mediterranean islands and islets host a large number of non-indigenous species, mainly plants (Brundu 2013). For example, an analysis of 37 small Italian islands showed that they are affected by 203 non-native plants, with a remarkable increase of acacias and succulents in the last decades (Celesti-Grappo et al. 2016). The main determinants of non-indigenous plant species richness in small islands are tourist development and the percentage of artificial land-cover. However, at the local scale, para-oceanic island ecosystems such as the Balearic Islands have a relatively lower number of non-native plants than their mainland counterparts (Vilà et al. 2010). Yet, some species can be introduced into more ecosystem types in islands than in the mainland (Gimeno et al. 2006). In general, there are large differences in the taxonomic composition of non-indigenous insect assemblages between islands and continental countries, e.g., France and Corsica, Italy and Sicily (Liebhold et al. 2016), and Greece and Crete (Avtzis et al. 2017).

### **Temporal trends of non-indigenous species and pests**

The rate at which humans have moved species beyond their native ranges has tremendously increased over the last 150–200 years (di Castri 1991; Reichard and Hamilton 1997), and more so in the last decades (Genovesi et al. 2009; Cardador et al. 2019). Although all taxonomic groups have shown a general rise during this period consistent with the exponential increase in trade and travel (Jeanmonod et al. 2011; Seebens et al. 2017; Cardador et al. 2019), little is known about how temporal dynamics of non-indigenous species varies among taxa. Where time series are available, the number of non-indigenous species established in Europe has increased exponentially in terrestrial ecosystems (Jeschke and Strayer 2005; Hulme et al. 2008; Lambdon et al. 2008; Santini et al. 2013). Abellán et al. (2016) analyzed data on bird introductions in Spain and Portugal since 1912 and found that most of them (99.9%) were recorded

from 1955 onwards, with a sharp increase after the 1980s that mirrors the number of non-native birds imported into these countries. Cage birds (mainly Passeriformes and Psittaciformes) constitute the bulk of the species introduced during the last 40 years through escapes of individuals kept in captivity as pets. Although the information is less detailed, and reptiles and amphibians have smaller numbers of recorded non-indigenous species than birds, both groups have also increased their numbers during the 20th century in parallel with the rise in human immigration into Europe (Jeschke and Strayer 2006) and the international trade (Jenkins 1999).

The rate of establishment of non-indigenous insect species has also increased during the last decades (Roques 2010, 2015; Matošević and Pajač Živković 2013; Avtzis et al. 2017). A fast and quite linear increase, with about 10 new species per year, was noticed in Italy since World War II (Inghilesi et al. 2013), and an even higher rate of increase was noted in Croatia since 2007 (Matošević and Pajač Živković 2013). The species newly established during the last three decades tend to spread all over the Mediterranean Basin significantly faster than those that arrived between 1900–1990s (Roques et al. 2016). Such a rapid spread was especially impressive in some species, often relying on multiple introductions in different countries being used as bridgeheads (Rugman-Jones et al. 2013; Kerdelhué et al. 2014; Garnas et al. 2016; Roques et al. 2016; Bras et al. 2019; Lesieur et al. 2019).

Non-indigenous plant pathogenic species have increased exponentially in the last four decades (Santini et al. 2013). Since then, new non-indigenous plant pathogenic species have been introduced mainly from North America, and recently from Asia. Hybrid pathogens also appeared. Countries with a wider range of environments, higher human disturbances or international trade host more non-indigenous species. Rainfall influences the diffusion rates. Environmental conditions of the new and original ranges and systematic and ecological attributes affect pathogen success (Santini et al. 2013).

For plants, the success of introduction in terms of their area of occupancy is larger in species introduced a few centuries ago than species introduced in the 20th century (Lambdon and Hulme 2006), while for birds, establishment success is positively related to time since first introduction (Abellán et al. 2017).

### ***Pathways of introduction (intentional and accidental) of non-indigenous species and pests***

The majority of non-indigenous plants have been introduced into the Mediterranean Basin intentionally, as ornamentals that have escaped from gardens associated with anthropic developments and housing (e.g., touristic urbanizations) but also to embellish infrastructures (Hulme et al. 2008). Furthermore, many non-native trees (e.g., *Acacia*, *Pinus*, *Eucalyptus*) have been planted at large scales as forestry species and also in restoration programs for dune-stabilization, riverine water flux control, soil fertilization or afforestation of agricultural abandoned land. Many plant species have also been introduced unintentionally (accidentally) as “hitchhikers” or seed contaminants.

The main pathway of introduction for vertebrates, for example birds, are accidental escapes from private collections (Abellán et al. 2016). International wildlife trade is one of the main (if not the main) sources of current vertebrate non-indigenous species. When the EU banned the imports of wild birds, there was a rapid trade shift from wild-caught birds to captive-bred birds (which have lower potential to establish populations than wild-caught birds) (Carrete and Tella 2008, 2015; Cabezas et al. 2013) and a sharp decrease in the number of new introduced avian species in the wild (Cardador et al. 2019). However, this positive effect of the EU ban on wild-caught birds coincides with a significant increase in the trade in reptiles (Cardador et al. 2019).

For invertebrates, the vast majority of species introductions have been accidental (Hulme 2009; Roques 2015). A few introductions have been intentional, mostly for biological control between 1950 and 1999 (Rasplus et al. 2010), but such species always represent less than 15% of the total number of non-indigenous species per Mediterranean country, except in Israel (17.4%) (Roll et al. 2007). Since the majority of invertebrates established in the Mediterranean Basin are phytophagous, the major pathway of unintentional introductions appears to be via international trade in live plants (Rabitsch 2010; Inghilesi et al. 2013; Eschen et al. 2015; Roques 2015). Seed trade has also provided a few species and pests (Auger-Rozenberg and Boivin 2016) as well as firewood, logs and fallen timber (Meurisse et al. 2019). The trade of vegetable and fruit commodities also constitute an important pathway for non-indigenous pests (Desneux et al. 2010; Abbes et al. 2012; Cini et al. 2014). Hitchhiking is another significant pathway of pests as stowaways using wood packaging material


(Rassati et al. 2015; Javal et al. 2019; Lesieur et al. 2019), transport infrastructure and vehicles (Javal et al. 2019; Kirichenko et al. 2019) or used tires such as for mosquitoes (Rabitsch 2010). Several other examples are associated with beekeeping (Mutinelli et al. 2014).

For plant pathogens, all the introductions occurred unintentionally. The exact pathway of introduction is almost unknown for most species. However, the most probable is the trade of living plants (57%) or wood (10%). Less than 10% of the introductions occurred through any of the other pathways (Santini et al. 2013). Introductions of even harmless fungi in a new environment give them the opportunity of mating with local or introduced related species giving rise to hybrid progenies. The hybridization process may result in an increase of pathogenicity in one of the species or in the emergence of a completely new plant disease, both of which may threaten the original host plant, and new and naïve host species (Ghelardini et al. 2017).

### ***2.5.2.2 Non-indigenous species as drivers of biodiversity and ecosystem change***

The introduction of non-native plants can decrease local flora and fauna diversity and change the community composition and functional structure of affected ecosystems (Vilà et al. 2006; Zahn et al. 2009; Rascher et al. 2011). Native plants that are most vulnerable to such introductions are those with small population sizes (Lapiedra et al. 2015). At least 12 endemic or critically endangered plant species from the “Top 50 Mediterranean Island Plant” list are threatened by non-native plants (de Montmollin and Strahm 2005). For birds, the impact of non-indigenous species is higher on island species, and in those species with small distribution ranges (Clavero et al. 2009). Destabilized ecosystems, including systems used for food and agricultural production, tend to be more vulnerable to the spread of non-indigenous species (e.g., Marvier et al. 2004; Chytrý et al. 2008). However, there is little evidence to support the hypothesis that highly diverse ecosystems are inherently more resistant to non-indigenous species than less-diverse systems (e.g., Keller et al. 2011).

Changes in ecosystem functioning after the introduction of non-indigenous plants are highly context-dependent and include alterations in decomposition rates, light and water soil availability, and changes in soil carbon and nitrogen pools (Vilà et al. 2006; Castro-Díez et al. 2009; Rascher et al.



**Figure 2.29 | Number of non-indigenous woody plant pathogens per country** (Santini et al. 2013).

2011). Plant introductions can disrupt the positive relationship between native species diversity and multifunctionality (Constán-Nava et al. 2015) and may trigger regime shifts by changing plant succession (Stinca et al. 2015).

The impacts of vertebrates can be dramatic through competition for resources, predation and as vectors of diseases (Genovesi et al. 2009). For example, the introduction of non-native ungulates is a major threat to endangered plants, especially in islands (Pisanu et al. 2012). Some native amphibians have collapsed after the introduction of non-native anurans (Lillo et al. 2011). Non-indigenous parakeets can also interact with native species and have an impact on native populations and communities, largely in the form of harassment, displacement from nest sites and food competition (Hernández-Brito et al. 2014, 2018; Menchetti and Mori 2014; Menchetti et al. 2016; Covas et al. 2017). NIS arthropods can also negatively impact native biodiversity and ecosystem processes by destroying host plant populations, causing disturbances in native genetic resources or, indirectly, affecting the affected communities because of new species assemblages (Kenis et al. 2009; Kenis and Branco 2010; Auger-Rozenberg and Boivin 2016).

Some non-indigenous species can damage productive sectors, most notably agriculture and forestry, with important economic consequences. Non-native insects impact major crops in many countries of the Mediterranean region (Abbes et al. 2012; Abdallah et al. 2012; Mutke et al. 2016). Some non-indigenous bird species such as Monk and ring-necked parakeets can also cause important damages to crops (Senar et al. 2016; Turbé et al. 2017).

Forests as well as urban trees in the Mediterranean Basin can be severely economically impacted by non-indigenous pathogens (Fig. 2.29) (Santini et al. 2013; Ghelardini et al. 2017) and arthropods (Kenis and Branco 2010) some of which can be considered pests (Santini et al. 2013; Rassati et al. 2015; Auger-Rozenberg and Boivin 2016; Branco et al. 2016; Mendel et al. 2016) and can also transmit pathogenic fungi (Montecchio et al. 2014). For example, after World War II, chestnut blight epidemic in the mountains of southern Europe aggravated food shortages for local human populations and increased migration to urban areas (Adua 1999). Besides, canker stain disease of the plane tree was considered a nasty non-indigenous species of urban trees, until it was introduced into Greece, where the Oriental plane is endemic. The disease is presently destroying natural river wood ecosystems in Greece (Ghelardini et al. 2017; Tsopeles et al. 2017) and it is also spreading in neighbouring countries such as Albania (Tsopeles et al. 2017) and Turkey (Lehtijärvi et al. 2018). Currently, the pinewood nematode is massively killing pines, changing the landscape in Portugal (Naves et al. 2016).

Human health can also be affected by non-indigenous species. Of major concern are non-native plants that are allergenic; their advanced flowering phenology enlarges the period of airborne prevalence of allergens (Belmonte and Vilà 2004). Non-indigenous birds of the order Psittaciformes (parrots) are potential reservoirs of *Chlamydophila psittaci*, the etiological agent of human psittacosis, and can transmit other diseases to humans and wildlife (Menchetti and Mori 2014; Turbé et al. 2017). Some non-native invertebrates, mainly insects, can cause distress, and allergic reactions

(e.g., the Asian hornet, *Vespa velutina*) or be vectors of infectious diseases, (e.g., the tiger mosquito, *Aedes albopictus*) (Lounibos 2002; Jucker and Lupi 2011; Monceau et al. 2014; Goubert et al. 2016; Roques et al. 2018; Liroy et al. 2019).

Some impacts of introduced non-indigenous species have attracted worldwide attention, e.g., on cultural heritage in Palermo (Manachini et al. 2013), on the survival of the endangered date palm, *Phoenix theophrasti* in Crete (Avtzis et al. 2017), and the severe attacks of cypress canker disease in southern Tuscany (Italy). Since researchers and policymakers rarely address the connection between non-indigenous species and damage to cultural heritage directly, the cost of these losses is often neglected or underestimated. The Mediterranean Basin has a long history of civilization and it is rich in cultural heritage that can be threatened by non-indigenous species. For example, in southern Tuscany (Italy), severe attacks of the cypress canker disease (caused by the North American fungal pathogen *Seiridium cardinale*) are threatening the survival of trees flanking a monumental avenue (Danti and Della Rocca 2017).

### 2.5.2.3 Further introductions, spread and impacts of non-indigenous species and pests

#### Existing tools for predicting the risk of introduction and research needs

Horizon scanning, prioritization and Pest Risk Analysis (PRA) are essential tools for focusing limited resources to predict the species which can have a high rate of spread, inflict high impacts, and can be cost-effectively managed. PRA are defined by the International Plant Protection Convention as "the process of evaluating biological or other scientific and economic evidence to determine whether a pest should be regulated and the strength of any phytosanitary measures to be taken against it" (FAO 2017b). An important step in the PRA scheme is the "Pest management section" which assesses phytosanitary measures for relevant pathways and their effectiveness in preventing the entry, establishment and spread of non-indigenous species.

Since 2006, European and Mediterranean Plant Protection Organization (EPPO) has formed expert working groups (EWG) to conduct PRA comprised of experts on the pest and cropping systems, mapping and modelling experts, along with experts on EPPO's PRA scheme, risk managers and EPPO PRA Core Members, all which acts to ensure

consistency. EPPO is an international organization responsible for cooperation and harmonization in plant protection within the European and Mediterranean region. One of EPPO's main aims is to provide assistance and guidance to member governments on the administrative, legislative and operational measures necessary to prevent the introduction and spread of non-native plant pests (Smith 1979; Roy et al. 2011). Since 1999, EPPO has maintained an Alert List of plant pests and non-indigenous plants which acts as an early warning for pests, which can present a risk to the EPPO region. EPPO has also developed a prioritization tool for non-indigenous plants that classifies species into one of three lists: minor concern, observation list or list of non-indigenous plant species (EPPO 2012a, 2012b). Those species included in the list of non-indigenous plants are assessed for a PRA, where a higher priority is given to those species with a limited distribution in the EPPO region (EPPO 2012b).

In addition to the EPPO PRA tools, there are also a number of other PRA protocols (Roy et al. 2018) that can be applied to the Mediterranean Basin. To better improve PRA and the risk assessment process, a greater level of transparency and consistency between protocols would be beneficial (Vanderhoeven et al. 2017; González-Moreno et al. 2019). EU Mediterranean member states need to follow the EU Regulation that includes restrictions on keeping, importing, selling, breeding and growing non-indigenous species (European Union 2014).

A quite novel empirical approach to identify potential pests is the use of sentinel plantations of Mediterranean trees, e.g., cork oak, evergreen oak and cypress, in other continents as a priori identification of non-indigenous insect and pathogens capable of colonizing such plants. For example, such plantations in China provided a list of 39 potential non-indigenous insects of which five could be highly damaging (Roques 2015) and several pathogens (Vettraino et al. 2015). The development at potential ports of entry of trapping programs using lures presenting a generic attractiveness for some insect groups is expected to allow early detection of emerging non-indigenous species, even when not listed on quarantine lists (Rassati et al. 2014, 2015; Fan et al. 2019).

#### Non-indigenous species likely to be introduced into the Mediterranean in the next 20-50 years

As temperature increases, current major non-indigenous species are predicted to shift north-

wards at an average pace of 37-55 km decade<sup>-1</sup>, leaving a window of opportunity for new non-indigenous species better adapted to xeric conditions (Gallardo et al. 2017). Regarding non-native plants, gardening practices and ornamental trade will have a major impact on the selection of these future non-indigenous species. The use of non-native drought-tolerant species for gardening and landscaping (i.e., xeriscape) is at its earliest stages in the Mediterranean, but it has already raised concerns in California because of its potential risk as a source of new non-indigenous species (Bradley et al. 2012). Global species niche modelling indicates that xeric shrublands in Mediterranean areas are among the most susceptible ecosystems to introduction by Cactaceae plant species from arid American areas (Novoa et al. 2015). Although some Cactaceae are already largely distributed across the Mediterranean (e.g., *Opuntia ficus-indica*), it is very likely that close relative species with currently restricted distribution or absence in the Mediterranean, such as *Cylindropuntia* spp., would thrive in the next decades aided by new gardening practices (Essl and Kobler 2009). Besides gardening, other relevant terrestrial plant species could be easily introduced and established as contaminants in soil, seeds or containers. For instance, *Parthenium hysterophorus* is a species not currently in the Mediterranean that has been highlighted as of high risk for the region because of its large potential negative impact on agriculture and human health (Kriticos et al. 2015).

While plants have been proportionally the main new non-indigenous species in Europe up until the 19th century, the trend has shifted towards an increasing number of introduced invertebrates and vertebrates in the 20th century (Hulme 2009). This is a pattern that is very likely to continue in the near future by increasing air and maritime cargo, where these taxa can be easily transported as stowaways. The establishment of non-indigenous

invertebrates of tropical origin affecting woody ornamental plants has increased (Eschen et al. 2015), meaning that many ornamental plants, especially palms, fig trees and exotic legumes, are at risk for further introduction as well as *Citrus* and *Eucalyptus* trees (Floris et al. 2018). The recent establishment of ambrosia beetles of tropical origin directly threatening plants of the Mediterranean maquis (Faccoli et al. 2016; Francardi et al. 2017) suggests that such process is going to be amplified with global warming. A list of fruit flies likely to be introduced has been recently proposed (Suffert et al. 2018). Special attention should be paid to major agricultural pests currently not present in the Mediterranean but with the potential to be introduced and cause a major impact. The EPPO A1 quarantine list considers up to 128 species of insects, mites, nematodes and gastropods, currently absent from the EPPO region, recommended for quarantine measurements. For instance, Lepidoptera species such as *Spodoptera* or *Helicoverpa* spp. are polyphagous species that could easily thrive in the Mediterranean if they become established. *Spodoptera frugiperda*, a pest native to the Americas, has quickly spread in Africa causing large yield loss. A recent modelling exercise has identified small pockets of suitable habitats in the Mediterranean area but the potential for permanent populations is still uncertain (Early et al. 2018).

Regarding terrestrial vertebrates, several species have been recently highlighted in a horizon scanning exercise for European non-indigenous species, including the Mediterranean (Roy et al. 2019). Of special relevance is the common myna, *Acridotheres tristis*, a non-indigenous species with very restricted populations in the region, and *Lampropeltis* spp., a family of snakes mainly native to North America and adapted to arid conditions. These species are traded as pets and can easily be introduced due to accidental escapes.

## 2.6 Interaction among drivers

### 2.6.1 Drivers impacting other drivers

The potential for interactions among drivers is a key issue for analyzing their impacts on environment and human societies, and for developing effective conservation policies (Brook et al. 2008). Climate change, pollution, land and sea use change, and non-indigenous species are of-

ten studied and managed in isolation, although it is becoming increasingly clear that a single driver perspective is inadequate when ecosystems are threatened by multiple, co-occurring drivers (Halpern et al. 2008a, 2008b). Conceptually, there are three broad categories of interaction types describing the outcome of multiple stressors, the effects can be additive/cumulative (all the dif-

ferent stresses derived from the implied drivers show up), synergistic (increased stress) or antagonistic (decreased stress) (Folt et al. 1999; Crain et al. 2008). Also, and particularly for the Mediterranean, how different drivers interact could result in alteration, intensification, and even in generation of new impacts (Doblas-Miranda et al. 2017).

In order to facilitate the multi-stressor approach, as a key recognized concept, this section offers two different approximations. First, we describe potential pair interactions within the individual driver classes described before, and second, we provide a few examples of characteristic disturbances of Mediterranean ecosystem that are the result of the combination among multiple drivers.

## 2.6.2 Pairs of interacting drivers

### 2.6.2.1 Climate change effects on pollution

Generally, increases in temperature enhance the toxicity of contaminants and increase concentrations of tropospheric O<sub>3</sub> regionally, but will also likely increase rates of chemical degradation (Lelieveld et al. 2014). In general, climate change coupled with air pollutant exposures may have potentially serious adverse consequences for human health in urban and polluted regions (Noyes et al. 2009).

The increase in the intensity and frequency of storm events linked to climate change can lead to more severe episodes of chemical contamination of water bodies and surrounding watersheds (Noyes et al. 2009). Climate change may also increase the occurrence and the global expansion of harmful algal blooms (Paerl and Paul 2012) (Sections 2.3.3 and 2.3.4).

### 2.6.2.2 Pollution effects on climate change

Many air pollutants that are harmful to human health and ecosystems also contribute to climate change by affecting the amount of incoming sunlight that is reflected or absorbed by the atmosphere, with some pollutants warming and others cooling the Earth. These so-called short-lived climate-forcing pollutants include methane, black carbon, ground-level O<sub>3</sub>, and sulfate aerosols. They have significant impacts on the climate; black carbon and methane in particular are among the top contributors to global warming after CO<sub>2</sub> (Shindell et al. 2009; Stohl et al. 2015). Over the Mediterranean Basin the increase and decrease of anthropogenic aerosols during the second half of the 20th century have had an important role in the dimming-brightening phases, because of their direct action on the incoming solar radiation (Section 2.2.3.1).

Pollution by heavy metals or organic compounds can also affect ecosystem functioning by inhibiting CO<sub>2</sub> fixation performed by photosynthetic organisms, thereby increasing global warming (Rochelle-Newall et al. 2008; Magnusson et al. 2010; Ben Othman et al. 2012) (Sections 2.2.3 and 2.3.2).

### 2.6.2.3 Impact of climate on land and sea use

#### Effects of climate change on land use

Recent accelerated climate change has exacerbated existing environmental problems in the Mediterranean Basin caused by the combination of changes in land use, increasing pollution and biodiversity decline (Cramer et al. 2018). Sea-level rise, combined with land subsidence, may significantly reduce the area available for agriculture. The effects of sea level rise in North Africa, especially

Impacting (column) – Impacted (row)	Climate change	Pollution	Land and sea use changes	Non-indigenous species
Climate change		2.6.2.2	2.6.2.4	?
Pollution	2.6.2.1		2.6.2.7	?
Land and water use changes	2.6.2.3	2.6.2.6		2.6.2.10
Non-indigenous species	2.6.2.5	2.6.2.8	2.6.2.9	

Table 2.3 | Main interactions among drivers

on the coast of the Delta region of Egypt, would impose additional constraints to the agricultural land (Section 3.2.2.1), and also the salinization of coastal aquifers (Section 3.1.2.2).

Similarly, with 42% of the population living in coastal areas (Mediterranean Wetlands Observatory 2018), important direct effects of climate change on coastal settlements include dry-land loss due to erosion and submergence, damage of extreme events (such as wind storms, storm surges, floods, heat extremes, and droughts) on built environments, effects on health (food- and water-borne disease), effects on energy use, effects on water availability and resources, and loss of cultural heritage (Hunt and Watkiss 2011) (Section 2.2.8.2). Coastal industries, their supporting infrastructure including transport (ports, roads, rail, airports), power and water supply, storm water, and sewerage are highly sensitive to a range of extreme weather and climate events including temporary and permanent flooding arising from extreme precipitation, high winds, storm surges, and sea level rise (Horton et al. 2010; Handmer et al. 2012; Hanson and Nicholls 2012; Aerts et al. 2013). The tourism development experienced a comparable pattern, requiring host facilities and corresponding services. In Algeria, for example, construction projects have been carried out among the coastal paleo-dunes despite the existing Littoral Law 02-2002 (coastal protection) and the Law 01-3-2003 related to the Impact Expertise (Senouci and Taibi 2019). A similar situation exists in the industrial sector (e.g., desalination plant and electricity power station built on the beach).

Increases in temperature and decreases in precipitation could alter fire regimes affecting forest cover and could increase the intensity and frequency of drought resulting, in combination with other factors, in desertification (Sections 2.4.1.2 and 2.6.3). Future changes in climate could decrease food production (Section 2.4.1.2) and may alter the use of land all over the Mediterranean (Section 2.4.1.3). Future changes in the quantity and intensity of rain could affect the water cycles and increase the risk of floods (Sections 2.2.6 and 2.6.3).

### **Climate change and variability drives dynamics of marine species**

Climate change and variability has led to concomitant changes in Mediterranean marine ecosystems and resources, with various implications on species diversity and composition, where species with limited locomotive capacity or confined in fragmented habitats seem more likely to be affected (Lejeune

and Chevaldonné 2006; Ledoux et al. 2015). Examples of this changing environment, among others, are the mass mortality events of gorgonians and other sessile metazoans in northwestern Mediterranean (Garrahou et al. 2009; Rivetti et al. 2014) and the continuous decline of *Posidonia* meadows (Marba and Duarte 2010), the increase in the frequency of red tides and of gelatinous carnivore outbreaks (Conversi et al. 2010), the “tropicalization” of marine fauna in favour of the more thermophilic ones (Bianchi 2007), and the increase spread of microbial pathogens associated with water temperature rise (Danovaro et al. 2009) (Section 4.1.1.1).

At the end of 1980s and especially during the mid-1990s the Mediterranean Sea underwent regime shifts (Conversi et al. 2010; Alheit et al. 2019) that inflicted major atmospheric, hydrological and ecosystem changes, also affecting marine resources, mainly fisheries. There have been various studies linking ocean-atmospheric processes such as the Atlantic Multidecadal Oscillation (AMO), the North Atlantic Oscillation (NAO) and the Western Mediterranean Oscillation (WeMO) indices to alterations on the distribution and biomass of pelagic fish, as well as their catch composition (Alheit et al. 2019). Pelagic fish populations, more than other fish species, act as sentinels of these environmental changes. For example, during mid-1990s in the Mediterranean the highly correlated sea surface temperature and AMO index show a sharp increase (Marullo et al. 2011; Macías et al. 2013), whereas the dynamics of many fish species - mainly pelagic - show a conspicuous change around that time.

It is not yet clear how these changes impact pelagic fish population dynamics, combined with the pressures imposed by anthropogenic activity. Fifty-nine taxonomic groups (species or groups of species) showed an abrupt change in their landings in the mid-late 1990s (Tzanatos et al. 2014) with approximately 64% of these changes being correlated with sea surface temperature, mostly inversely correlated. The landings of some species (European sardine *Sardina pilchardus*, squids *Loligo* spp., Norway lobster *Nephrops norvegicus*, and hake *Merluccius merluccius*) decreased conspicuously in the mid-1990s, whereas those of other species (European anchovy *Engraulis encrasicolus* and greater amberjack *Seriola dumerili*) increased (Tzanatos et al. 2014). A study of the fisheries landings of 30 fish and invertebrate taxonomic groups revealed regime shifts at the mid-1990s, concurrent with the sea temperature increase in the eastern and western basins (Vasilakopoulos et al. 2017). The late 1990s



was determined as the turning point for the northward expansion of warm-water species in the Mediterranean (Azzurro et al. 2011). This was confirmed by Raitzos et al. (2010) who showed a clear increase of non-indigenous species entering into the eastern Mediterranean Sea in 1998. Earlier, Pinnegar et al. (2003) reported that the diversity of the western Mediterranean finfish landings increased dramatically after 1995 as a result of new species entering the catch.

Round sardinella (*Sardinella aurita*), a warm-water small pelagic fish species distributed along the southern Mediterranean coastline has been reported to have expanded its distribution to the northern Aegean (Tsikliras 2008), the northern Adriatic (Sinovčić et al. 2004), the Gulf of Lions (Francour et al. 1994), and the northwestern Mediterranean (Sabatés et al. 2006, 2009). A significant positive relationship between round sardinella landings and sea surface temperature anomalies has been reported for the western (Sabatés et al. 2006) and eastern Mediterranean (Tsikliras 2008). The northward distributional shift coincides with the beginning of positive temperature anomalies in the mid-1990s (Tsikliras 2008; Sabatés et al. 2009; Stergiou et al. 2016). Similarly, concomitant with the sea surface temperature change in the western Mediterranean, the landings of bluefish (*Pomatomus saltatrix*) quadrupled due to a northward expansion of the species (Sabatés et al. 2012) and, at the same time, anchovies returned to high biomass, as a result of increasing sea surface temperatures in the Adriatic Sea (Vilibić et al. 2016).

The effect of the AMO and NAO signals across the Mediterranean Sea sub-regions (western, central and eastern) on the small (European sardine *Sardina pilchardus*, European anchovy *Engraulis encrasicolus*, round sardinella *Sardinella aurita* and European sprat *Sprattus sprattus*) and medium (Atlantic mackerel *Scomber scombrus*, Atlantic chub mackerel *Scomber japonicus*, Atlantic horse mackerel *Trachurus trachurus*, Mediterranean horse mackerel *Trachurus mediterraneus*) pelagic fishes have been recently studied in the western, central and eastern Mediterranean Sea (Tsikliras et al. 2019). The pelagic fishes of the central and eastern Mediterranean respond most strongly to AMO variability and those of the central and western Mediterranean also respond to the NAO, while the effect of the NAO on pelagic fishes of the eastern Mediterranean was not significant (Tsikliras et al. 2019). Generally, various indicators revealed that the time of the pelagic fish response to the AMO and NAO signals varied among the

Mediterranean sub-regions (Alheit et al. 2014; Tsikliras et al. 2019).

Finally, the mean temperature of the catch, an indicator that assesses the effect of global warming on the exploited marine communities (Cheung et al. 2013b), has been increasing across the Mediterranean showing that the ratio of thermophilous (warm-water) to psychrophilous (cold-water) marine species has been changing in favour of the former. This is indicative of either an increase in the relative proportion of thermophilous species in the catches or a decrease in the relative proportion of the psychrophilous ones (Tsikliras and Stergiou 2014).

#### 2.6.2.4 Effects of land use on climate change

Changes in crop use (Tribouillois et al. 2018), especially in forest cover, affect the balance between sink and release of CO<sub>2</sub> and the emissions of biogenic volatile organic compounds (BVOCs) in the atmosphere (Doblas-Miranda et al. 2017) (Sections 2.4.1.2 and 3.1.2.1).

Modification of surface albedo by land use changes also entail a highly potential impact on climate change (Benas and Chrysoulakis 2015). Changes in forest or dehesa/montado cover due to reforestation could reduce albedo (Rotenberg and Yakir 2011; Godinho et al. 2016), while fires increase radiations returns to the atmosphere (Sánchez et al. 2015), with contrasting effects on local climate. Agricultural cover may decrease or increase albedo (Giannakopoulou and Toumi 2012; Carrer et al. 2018), while urban sprawl definitely increases the radiation absorption and therefore local temperature (Salvati et al. 2019) (Section 3.1.3.1).

Change of land use and irrigation practices increase evapotranspiration and have a net cooling effect in some areas of the Mediterranean region (Zampieri and Lionello 2011; Thiery et al. 2017; Gormley-Gallagher et al. 2020).

#### 2.6.2.5 Links between trends in non-indigenous species and climate change

##### Impact of climate change on marine non-indigenous species

The introduction of non-indigenous species and global warming interact in complex ways (Stachowicz et al. 2002), and are linked also in the Mediterranean Sea (Occhipinti-Ambrogi 2007).

This connection strongly depends on the species and the mode of its introduction, establishment and colonization. Overall, there is a strong trend of "tropicalization" of temperate areas through the movement of warm-loving (thermophilic) species toward the poles in areas of rapid ocean warming, and with increasingly strong impacts on local communities (Vergés et al. 2014a, 2016). These are not considered as introductions of non-indigenous species per se. But ocean warming may facilitate the establishment and spread of thermophilic non-indigenous species. The success of establishment of an introduced species depends on how suitable the ocean climatic parameters are in the region of introduction. Because successful non-indigenous species are typically generalists with broader climatic tolerances, they are usually considered able to cope better with climate change than native ones (Walther et al. 2009).

There is limited evidence for effects of climate change on the introduction of non-indigenous species. Theoretically, at the trailing "warm" edge of species distributions, the populations of sensitive cold-affinity species should reduce (and eventually extirpate) and that of warm affinity species (including thermophilic species) should increase (Bates et al. 2014). In the Mediterranean Sea, tropicalization evidently occurs (Vergés et al. 2014a), mainly in the Levant, by Lessepsian introductions, and ocean warming was suggested to facilitate the successful establishment of non-indigenous species (Raitsos et al. 2010). Ocean warming probably also helps to spread both native thermophilic species and successful Lessepsian species westward along the basin's temperature gradient, and also northward into the Aegean and Adriatic seas, or even the Ligurian Sea, but there very few direct empirical studies to demonstrate that. Recent analysis of fish trawl data from the southeast Mediterranean (Israel) does strongly suggest that non-indigenous species are indeed promoted by warming while natives are declining (Givan et al. 2017a).

Some studies suggest that habitats degraded by global warming are more likely affected by non-indigenous species than nearly-pristine habitats, envisaging explicitly or not, a cause and effect link between climate warming and the success of introductions (Stachowicz et al. 2002; Bianchi 2007; Galil 2007; Occhipinti-Ambrogi 2007). However, field observations do not support this idea, but reveal instead conflicting results that have provoked intense debate (Boudouresque and Verlaque 2010). For example, in the Mediterranean

Sea, well-structured and conserved habitats (such as coralligenous or *Cystoseira* forests) are able to mitigate and delay the proliferation and spread of the non-indigenous alga *Caulerpa cylindracea*, probably because the complexity of substrata (enhanced by gorgonians or canopy algae presence) is a key factor limiting its colonization and spread (Ceccherelli et al. 2002; Bulleri and Benedetti-Cecchi 2008; Verdura et al. 2019). In contrast, mass mortality of structural native species and subsequent increase of turf-forming species due to an extreme climatic event indirectly promoted the introduction of *C. cylindracea* in a coralligenous habitat (Verdura et al. 2019). However, in other non-indigenous algae such as *Lophocladia lallemandii*, introduction is favoured by more complex and rich communities (Cebrián et al. 2018), and thus simplification derived from climate change effects is expected not to enhance the capacity of *Lophocladia* to establish itself, but prevents its spread.

Using natural laboratories to test the thermal performance curves and sensitivity to acidification of key native and non-indigenous species, as well as the impact of climate change related environmental alteration on species interactions and communities and their ecosystem functions, are critical for better understanding and forecasting of the interactions between climate change and the introduction of non-indigenous species (Rilov et al. 2019a). For example, heat polluted areas and CO<sub>2</sub> vents (Hall-Spencer et al. 2008), as well as laboratory experiments in near-natural mesocosm systems (Wahl et al. 2015). Such recent measurements and experiments in the southeastern Levant have shown that some non-indigenous species (foraminifera) are tolerant to extreme thermal stress (Titelboim et al. 2017), that under warming and acidification conditions most Lessepsian species perform better than native species (Guy-Haim et al. 2016; Guy-Haim 2017), and demonstrated that different thermal performance of two Red Sea foraminifera explain why one species was introduced and the other did not (Titelboim et al. 2019). Furthermore, mesocosm work showed that a *Cystoseira* community becomes more heterotrophic and more dominated by non-indigenous species (but species richness does not change), demonstrating the profound impact of the combination of climate change and the introduction of non-indigenous species on ecosystem function (Guy-Haim et al. 2016; Rilov et al. 2019b).

### **Impacts of climate change on terrestrial non-indigenous species**

There are five non-exclusive consequences of climate change on non-indigenous species: (1) altered transport and introduction mechanisms, (2) establishment of new species, (3) altered impact of existing non-indigenous species, (4) altered distribution of existing non-indigenous species, and (5) altered effectiveness of control strategies (Hellmann et al. 2008) [Section 2.5.1.3].

The influence of climate change on terrestrial non-indigenous species highly depends on species physiological strategy and reproductive adaptations (Bale and Hayward 2010; Antunes et al. 2018). Generalized ecosystem models of plant functional groups applied to Mediterranean islands indicate that climate change might promote the introduction of broadleaved trees (e.g., *Ailanthus altissima*) more than C<sub>4</sub> tropical grasses (e.g., *Amaranthus retroflexus*) (Gritti et al. 2006). Many non-indigenous species from temperate and cold climates might only be able to shift their ranges northward or to expand in altitude because they will be limited by drought and high temperatures (Storkey et al. 2014; Gallardo et al. 2017). While non-indigenous species whose native ranges are drier and warmer than their introduced ranges can be at an advantage to occupy niches at southern latitudes (Gallardo et al. 2017). Therefore, some species might lose and some gain suitable areas for introduction. Regions which will get drier are predicted to lose the highest number of potential non-indigenous species.

For introduced gardening plants, the climatically suitable areas with future climate change are unequally distributed across Europe with more suitable areas in the East than in the West of the Mediterranean Basin (Dullinger et al. 2017). This will be the case for *Cortaderia* which suitable area can increase 69-116% for 2060 (Tarabon et al. 2018) or for *Nassella* that can increase up to 47% for 2018 (Watt et al. 2011).

Similarly, weeds in crops can experience range shifts, niche shifts and trait shifts with climate change that will influence the agronomic practices to reduce their interference to crop production (Peters et al. 2014). Weeds in cereals crops will also advance towards northeastern Europe and remain or contract their distribution in warm areas of the Mediterranean region (Castellanos-Frías et al. 2014).

Climate change can advance the phenology of non-indigenous plant species including their fecundity

(Chuine et al. 2012), pollen production and seed maturation (Leiblein-Wild et al. 2016). Changes in pollen production can exacerbate the problem caused by allergenic non-indigenous plants such as the American *Ambrosia* because the allergenic risk is predicted to increase under all climate scenarios tested (Rasmussen et al. 2017).

Besides the influence of climate change on the establishment and spread of non-indigenous species, a remaining question is whether their impacts on native species increase in combination with climate change. A few greenhouse experiments have explored the interaction between competition of non-indigenous species and drought on the performance of native species (García-Serrano et al. 2007; Matesanz et al. 2008; Werner et al. 2010) and have found a non-synergistic effect. The interaction of climate change and introduction of non-indigenous species is a research area that requires further experimentation for productive systems such as the effect of weeds, pests and pathogens on crops and forestry (Ramesh et al. 2017).

### **2.6.2.6 Impacts of pollution on land and sea use**

One of the major drivers relative to greenhouse gases pollution may be the CO<sub>2</sub> fertilization affecting forests. The balance between faster growth due to the fertilization effect and hydric stress due to most likely warmer and drier conditions have generated a considerable debate in the Mediterranean area (Keenan et al. 2011; Peñuelas et al. 2011). However, the most recent studies mainly corroborate that the effects of CO<sub>2</sub> fertilization will be negligible under the predicted climate conditions for the region (Camarero et al. 2015; Nunes et al. 2015; Gea-Izquierdo et al. 2017), despite some exceptions (Koutavas 2013; Barbata and Peñuelas 2017). The potential effects of nitrogen deposition on Mediterranean forest growth also seem to be low (Ochoa-Hueso et al. 2014).

### **2.6.2.7 Impacts of land and sea use change on pollution**

Intensive farming increases releases of nutrients and pesticides in aquifers while higher releases of methane in the air. The effects of the increase of livestock production on greenhouse gas emissions are assessed in Section 3.2.3.2.

Urban sprawl is associated to higher traffic related emissions [Sections 2.3.3 and 2.4.1.2].

### 2.6.2.8 Pollution effects on non-indigenous species

Pollution can make environmental conditions less tolerable for native species, and provide space and nutrients for opportunists, including non-indigenous species (Crooks et al. 2011).

### 2.6.2.9 Effects of land and water use on non-indigenous species

Habitat destruction causes disturbance, which opens space for non-indigenous species (Hobbs and Huenneke 1992).

### 2.6.2.10 Effects of non-indigenous species on land and sea use

Outbreaks of forest non-indigenous insects could alter forest cover (Section 2.5.2.2).

## 2.6.3 More complex interactions among drivers

### 2.6.3.1 Floods

Floods are an illustrative example of the combination of different drivers such as climate change (extreme precipitation events), land use change (catchment changes on river forests, forest cover, etc.) and even indirect drivers (among them and principally, urban sprawl in risk areas) (Sections 3.1.3.3 and 3.1.4.1).

### 2.6.3.2 Desertification

Puigdefábregas and Mendizabal (1998) analyzed FAO data from Morocco, Algeria and Tunisia during the period 1950-1993, associating desertification to socio-economic boundary conditions and over-exploitation by showing clear increases in population (pressure) and in the use of unsustainable land use practices in the Mediterranean, principally irrigation (Section 6.6).

Desertification is in fact the result of two different factors in origin operating in combination, prolonged drought of climatic origin and land exploitation of human origin (Le Houérou 1996). In Mediterranean arid lands, mainly during the 20th century, short-term planning of agricultural policies and overexploitation, mainly in the form of overgrazing but also fuelwood collection and ground water exploitation, contributed to soil quality decline and massive erosion. Deteriorating conditions have a great impact on the lives of inhabitants of Mediterranean drylands and force

most of them to migrate (Mohamed and Squires 2018) (Sections 3.2.1.4, 4.3.1 and 6.6).

### 2.6.3.3 Wildfires

One relevant consequence of the Mediterranean Climate characterized by dry summers are forest fires. Those can be exacerbated by drought conditions (Turco et al. 2018) but in turn they can affect drastically the flood generation both due to the erosion and the loss of forest mass. Although some forest fires can be provoked or as a result of recklessness, they mostly depend on the state of the vegetation and the climatic and meteorological situation. Consequently, fire regimes will be affected by climate change, if not already affected (Sarris et al. 2014). The Mediterranean is a high fire-risk region, where fires are the cause of severe agricultural, economic and environmental losses and even human casualties (Moreira et al. 2011; Keeley et al. 2012; San-Miguel-Ayanz et al. 2013; Bowman et al. 2017). For instance, the fire seasons in 2017 and 2018 was severe in many regions of Southern Europe, with large wildfires associated with unusually intense droughts and heat-waves (Sánchez-Benítez et al. 2018). In Portugal, the year of 2017 was particularly tragic. An extended and extraordinarily intense fire season yielded a record total burned area of about 500,000 hectares and more than 120 fatalities (Turco et al. 2019). Instead, the summer of 2018 will be remembered by the deadliest fires ever recorded affecting Greece, when a series of wildfires close to Athens killed 99 people, the deadliest in Greece history (AghaKouchak et al. 2018).

However, although several reports, ranging from popular media through to peer-reviewed scientific literature, have led to a shared perception that fires have increased or aggravated in recent years, the quantitative evidence available indicated that fires are decreasing on recent decades in this area (Turco et al. 2016). The increased efforts in fire suppression have probably played an important role in driving the general downward trends described for most of the Mediterranean area (Moreno et al. 2014; Ruffault et al. 2015). In recent decades fire management strategies have improved thanks to new technologies and experience while climate drivers have led to an opposite trend (Amatulli et al. 2013; Batllori et al. 2013; Bedía et al. 2013; Turco et al. 2014; Dupire et al. 2017; Fréjaville and Curt 2017) (Section 4.3.2.1).

FULL NAME	SHORT NAME	THEMATIC FOCUS	NO. OF SCENARIOS	TIME HORIZON	SPECIFIC REGIONAL FOCUS	REFERENCE(S)
<b>GLOBAL SCALE</b>						
Special Report on Emissions Scenarios	SRES	Emission of greenhouse gases	4	2100	-	Nakićenović, 2000
Shared Socioeconomic Pathways	SSPS	Multidisciplinary with a focus on challenges to climate change adaptation and mitigation	5	2100	-	(O'Neill et al. 2014, 2017)
<b>MEDITERRANEAN SCALE</b>						
A sustainable future for the Mediterranean	-	Multidisciplinary and cross-sectoral with an emphasis on sustainable development	2	2025	-	Benoit and Comeau, 2005
Mediterranean scenarios (MedAction project)	-	Multidisciplinary and cross-sectoral with an emphasis on desertification	3	2030	Northern Mediterranean case studies in ES, GR, IT, PT	Kok et al., 2006
EuroMed-2030	-	Multidisciplinary and cross-sectoral with a focus on the Euro-Mediterranean relationship	4	2030	-	EC/DG for Research and Innovation, 2011
Tomorrow, the Mediterranean	-	Multidisciplinary, cross-sectoral, with emphasis on economic development	3	2030	-	IPEMED, 2011
Scenarios for the Mediterranean Region	-	Evolution of regional dynamics and role of the private sector in shaping business and political environments	3	2030	Exclusion of AL, BA, HR, IL, ME, PS, TR	World Economic Forum, 2011
Mediterranean Coastal SSPs	-	Regional and sectoral extension of the global SSPs for Mediterranean coastal regions	5	2100	-	Reimann et al., 2018
<b>EUROPEAN SCALE</b>						
Integrated Visions for a Sustainable Europe	VISIONS	Sustainable development	3	2020, 2050	-	Rotmans et al., 2000
Demographic and Migratory Flows Affecting European Regions and Cities	DEMIFER	Demography and European policies	5	2050	-	Rees et al., 2012
Climate Change Integrated Assessment Methodology for Cross-Sectoral Adaptation and Vulnerability in Europe	CLIMSAVE	Multidisciplinary and cross-sectoral, with emphasis on ecosystem services and provisions	4	2050	-	Gramberger et al., 2013
Territorial Scenarios and Visions for Europe	ET2050	Territorial development and cohesion	4	2050	-	MCRIT, 2015
Demographic Scenarios for the EU	-	Demographic development with a focus on aging, migration and education	3-4	2060	EU	Lutz et al., 2019
European Shared Socioeconomic Pathways	Eur-SSPs	Regional extension of the global SSPs for the European context	4	2040, 2070, 2100	Additional case study in Iberia	Kok et al., 2019

**Table 2.4 | Overview of selected socioeconomic scenarios that cover Mediterranean countries**, partly based on Rohat et al. (2018) and Sanna and Le Tellier (2013). ISO country codes: AL: Albania, BA: Bosnia and Herzegovina, ES: Spain, GR: Greece, HR: Croatia, IL: Israel, IT: Italy, ME: Montenegro, PT: Portugal, PS: State of Palestine, TR: Turkey.

## 2.7 Mediterranean socioeconomic scenarios

Environmental-change-related impacts will be driven not only by changes in climatic conditions, but also by changes in socioeconomic conditions. Prevailing socioeconomic conditions, in particular, determine a society's resilience to climatic hazards. When assessing future risks due to climate change, it is therefore crucial to account for plausible changes in socioeconomic conditions using a range of socioeconomic scenarios (González-Moreno et al. 2013).

A large number of socioeconomic scenarios have been developed in the past decades, focusing on a multitude of disciplines, sectors, and regions. Few of these scenarios were developed specifically for the Mediterranean region and even those usually only cover some of the Mediterranean countries, with a strong bias toward northern Mediterranean countries that are members of the European Union. Table 2.4 provides an overview of a range of

socioeconomic scenarios developed in the last two decades that cover socioeconomic developments




Figure 2.30 | Mediterranean population projections under the different Shared Socioeconomic Pathways (SSPs), A) total Mediterranean population, B) population by geographical region (Kc and Lutz 2017).

in the Mediterranean either fully or partially in terms of geographic coverage.

The most recent socioeconomic scenarios that account for socioeconomic developments in the entire Mediterranean region (as defined in this report) are the state-of-the-art global-scale Shared Socioeconomic Pathways (SSPs). The SSPs explore broad-scale societal trends in the course of the 21st century both qualitatively, in the form of scenario storylines, and quantitatively, in the form of national-level projections of key variables such as population (Kc and Lutz 2017), urbanization (Jiang and O'Neill 2017), and Gross Domestic Product (GDP) (Cuaresma 2017; Dellink et al. 2017; Leimbach et al. 2017). The Mediterranean population is projected to range from 607 million (SSP1) to 659 million (SSP3) in 2050 and from 518 million to 770 million in 2100 (Fig. 2.30a), with considerable differences across regions (Fig. 2.30b). The largest share of the population is projected to live in Egypt under all SSPs, except SSP5 where France

is the most populous country in 2100 due to very high work migration into northern Mediterranean countries (O'Neill et al. 2017).





Figure 2.31 | Mediterranean Gross Domestic Product (GDP) projections under the different Shared Socioeconomic Pathways (SSPs), A) Mediterranean average GDP per capita, B) average GDP per capita by geographical region [Dellink et al. 2017].

The regional average GDP per capita is projected to grow from 16,000 US\$ yr<sup>-1</sup> in 2010 to between 24,000 US\$ yr<sup>-1</sup> (SSP3) and 45,000 US\$ yr<sup>-1</sup> (SSP5) in 2050 and to roughly 35,000 US\$ yr<sup>-1</sup> (SSP3) to 142,000 US\$ yr<sup>-1</sup> (SSP5) in 2100 (Fig. 2.31a). The differences in average GDP per capita are small between (potential) EU candidate countries, countries from the Middle East and the Maghreb region. EU member states have the highest average GDP per capita under all SSPs (Fig. 2.31b).

In order to increase the usefulness of SSPs for impact, adaptation, and vulnerability assessments (van Ruijven et al. 2014), spatially explicit population projections that account for spatial changes in population distribution in the course of the 21st century have been produced, using the national totals as input data. These are available for all Mediterranean countries at a horizontal resolution of 7.5 arc minutes (Jones and O'Neill 2016) and 30 arc seconds (Merkens et al. 2016; Gao 2017). Further, downscaled GDP projections are available for SSPs 1-3 at a resolution of 30 arc minutes (Murakami and Yamagata 2019). All of these projections are based on the underlying global SSP assumptions.

As the global assumptions do not necessarily reflect the socioeconomic developments at the regional scale, extensions of the global-SSPs for the Mediterranean coastal zone have been developed (Reimann et al. 2018). These Mediterranean coastal SSPs account for region-specific developments as well as for changing attractiveness of coastal regions for human settlement across the SSPs, while at the same time ensuring consistency with the global SSPs (Zurek and Henrichs 2007). The Mediterranean coastal SSPs consist of qualitative narratives for each coastal SSP – SSP1 "Green Coast", SSP2 "No Wind of Change", SSP3 "Troubled Waters", SSP4 "Fragmented Coast", and SSP5 'Coast Rush' – differentiating between regional socioeconomic developments in northern versus southern and eastern parts of the region; and of spatially explicit population projections for all Mediterranean riparian countries at a resolution of 30 arc seconds (Fig. 2.32).



**Figure 2.32 | Spatially explicit population projections produced for the Mediterranean Coastal SSPs.** Selected population grids for the base year 2010 and each SSP in 2100 (Reimann et al. 2018), GPW = Gridded Population of the World (Center for International Earth Science Information Network - CIESIN - Columbia University 2016).



## BOX 2.1

### How much has the Mediterranean Basin warmed since the pre-industrial period?

The UNFCCC Paris Agreement of 2015 strengthens the initial goal of the Article 2 in the United Nations Framework Convention on Climate Change (UNFCCC), “to achieve ... stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system”, by “holding the increase in the global average temperature to well below 2°C above pre-industrial levels and pursuing efforts to limit the temperature increase to 1.5°C above pre-industrial levels”. While these temperature goals refer to the global average, it is a natural question to ask, for any region, how much warming has been observed “above pre-industrial levels”.


This is only apparently a simple question. To provide such an estimate it is necessary to clarify the meaning of pre-industrial, the information available on its average temperature in the region, a definition of present average temperature and the method used for estimating it. These issues were at the core of the IPCC special report on global warming of 1.5°C (SR15, IPCC 2018) and specifically considered in its *Chapter 1* (Allen et al. 2018).

The period 1850-1900 has been identified (Allen et al. 2018) as a suitable approximation for the estimate of the pre-industrial average temperature, because it combines typical pre-industrial solar and volcanic forcing, low anthropogenic greenhouse gas concentrations, and sufficient coverage of instrumental temperature observations. The choice of the period 1850-1900 is not completely free of problems, because it is indeed already affected by increasing greenhouse gas concentrations, with a partial compensation caused by aerosols. Further, strong volcanic eruptions occurred in the period 1880-1900. In this box, we follow SR15 but also the indications of the World Meteorological Organization (WMO 2017) and compute average pre-industrial temperature for a 30-years period, the central 1861-1890 period of the 1850-1900 “pre-industrial” period.

The number and distributions of instrumental observation have changed significantly over time, and station density could become critically low in African and Asian areas of the Mediterranean Basin. Further, the collection of observations over sea is systematically more problematic than over land, notably prior to the 20th century. Rather than analyzing station data, we seek consistency with global estimates and we use the HadCRUT4 dataset (Morice et al. 2012) and the CRUTEM4 (Jones et al. 2012) data sets for the land+sea and land only analysis, respectively. These are two widely used gridded global data sets with a resolution of 5 degrees longitude/latitude since 1850 until present. Other data sets at higher spatial resolution are available (such as the recently updated version of CRU TS) (Harris et al. 2020), but they do not reach back far enough into the 19th century for the estimation of pre-industrial conditions. For this analysis, the Mediterranean Basin is defined as the

domain from 10°W to 40°E of longitude and from 30°N to 47.5°N of latitude. For the averaging, an interpolation (based on the closest neighbours) to 1° spatial resolution was undertaken.

Obviously observed temperature values are not yet available for the period 2020-2034. This prevents computing the level of warming in 2020 using a simple 30-year average. Here, we make the conservative assumption that warming in the future will continue at the same rate of the last 50 years and compute the 2020 temperature by extrapolating the linear trend of the 1970-2019 period (Fig. 2.33). In 2020 the Mediterranean Basin is 1.5°C warmer than in the preindustrial with a likely uncertainty range of +0.11°C. Land areas have warmed more than the sea. If only land areas are considered, the 2020 temperature is 1.8°C warmer than pre-industrial with a likely uncertainty range of +0.12°C.



**Figure 2.33 | Time series (grey bars) of the annual average temperature of the Mediterranean Basin** (30°N to 47.5°N and 10°W to 40°E) considering the whole region (panel A) or only its land areas (panel B). The analysis is based on the HadCRUT4 data set for the land+sea analysis (Morice et al. 2012) and on the CRUTEM4 data set for the land analysis (Jones et al. 2012) (Jones et al., 2012). To avoid edge effects, before calculating the averages, the data have been interpolated to 1° spatial resolution. The values are expressed as anomalies from the pre-industrial period (1860-1890). The blue horizontal lines represent the 30-yr averages by steps of 20 years. The red curve is the linear trend linear trend calculated for the 1970-2019 period, extrapolated up to 2020. The black line is the smoother time-series. The blue square represents the likely interval (probability >0.66) for the present warming.

## BOX 2.2

### Representative Concentration Pathways (RCPs)

Representative Concentration Pathways (RCPs) are greenhouse gas concentration pathways, developed by the IPCC in order to explore the physical outcomes of different climate policies, notably regarding the mitigation of greenhouse gas emissions.

**RCP2.6:** The RCP2.6 was developed by the IMAGE modeling team of the Netherlands Environmental Assessment Agency (van Vuuren et al. 2011). The emission pathway is representative for scenarios in the literature leading to very low greenhouse gas concentration levels. It is a so-called “peak” scenario: its radiative forcing level first reaches a value around  $3.1 \text{ W m}^{-2}$  mid-century, returning to  $2.6 \text{ W m}^{-2}$  by 2100. In order to reach such radiative forcing levels, greenhouse gas emissions (and indirectly emissions of air pollutants) are reduced substantially over time.

**RCP4.5:** It was developed by the MiniCAM modeling team at the Pacific Northwest National Laboratory’s Joint Global Change Research Institute (Clarke et al. 2014). It is a stabilization scenario where total radiative forcing is stabilized before 2100 by employment of a range of technologies and strategies for reducing greenhouse gas emissions. It is often considered as an intermediate scenario.

**RCP8.5:** The RCP8.5 was developed by the MESSAGE modeling team and the IIASA Integrated Assessment Framework from the International Institute for Applied Systems Analysis (IIASA), Austria (Riahi et al. 2011). The RCP8.5 is characterized by increasing greenhouse gas emissions over time representative for scenarios in the literature leading to high greenhouse gas concentration levels, reaching  $+8.5 \text{ W m}^{-2}$  additional surface radiative forcing in 2100. It is often considered as a “business-as-usual” scenario.

## References

- Abadie J, Dupouey J-L, Avon C, Rochel X, Tatoni T et al. 2018 Forest recovery since 1860 in a Mediterranean region: drivers and implications for land use and land cover spatial distribution. *Landsc. Ecol.* 33, 289–305. doi: [10.1007/s10980-017-0601-0](https://doi.org/10.1007/s10980-017-0601-0)
- Abbes K, Harbi A, Chermiti B 2012 The tomato leafminer *Tuta absoluta* (Meyrick) in Tunisia: current status and management strategies. *EPPD Bull.* 42, 226–233. doi: [10.1111/epp.2559](https://doi.org/10.1111/epp.2559)
- Abdallah Z, Mezghani-Kkemakhem M, Bouktila D, Makni H, Makni M 2012 Genetic diversity of an invasive pest (*Oryctes agamemnon* Burmeister, Coleoptera: Scarabaeidae) of date palm in Tunisia, inferred from random amplified polymorphic DNA (RAPD) markers. *African J. Agric. Res.* 7, 1170–1176. doi: [10.5897/AJAR11.1580](https://doi.org/10.5897/AJAR11.1580)
- Abdennadher M, Hamza A, Fekih W, Hannachi I, Zouari Bellaaj A et al. 2012 Factors determining the dynamics of toxic blooms of *Alexandrium minutum* during a 10-year study along the shallow southwestern Mediterranean coasts. *Estuar. Coast. Shelf Sci.* 106, 102–111. doi: [10.1016/J.ECSS.2012.04.029](https://doi.org/10.1016/J.ECSS.2012.04.029)
- Abderrahim H, Chellali MR, Hamou A 2016 Forecasting PM10 in Algiers: efficacy of multilayer perceptron networks. *Environ. Sci. Pollut. Res.* 23, 1634–1641. doi: [10.1007/s11356-015-5406-6](https://doi.org/10.1007/s11356-015-5406-6)
- Abella A, Ria M, Mancusi C 2010 Assessment of the status of the coastal groundfish assemblage exploited by the Viareggio fleet (Southern Ligurian Sea). *Sci. Mar.* 74, 793–805. doi: [10.3989/scimar.2010.74n4793](https://doi.org/10.3989/scimar.2010.74n4793)
- Abellán P, Carrete M, Anadón JD, Cardador L, Tella JL 2016 Non-random patterns and temporal trends (1912–2012) in the transport, introduction and establishment of exotic birds in Spain and Portugal. *Divers. Distrib.* 22, 263–273. doi: [10.1111/ddi.12403](https://doi.org/10.1111/ddi.12403)
- Abellán P, Tella JL, Carrete M, Cardador L, Anadón JD 2017 Climate matching drives spread rate but not establishment success in recent unintentional bird introductions. *Proc. Natl. Acad. Sci. U. S. A.* 114, 9385–9390. doi: [10.1073/pnas.1704815114](https://doi.org/10.1073/pnas.1704815114)
- Acácio V, Dias FS, Catry FX, Rocha M, Moreira F 2017 Landscape dynamics in Mediterranean oak forests under global change: understanding the role of anthropogenic and environmental drivers across forest types. *Glob. Chang. Biol.* 23, 1199–1217. doi: [10.1111/gcb.13487](https://doi.org/10.1111/gcb.13487)
- Acri F, Aubry FB, Berton A, Bianchi F, Boldrin A et al. 2004 Plankton communities and nutrients in the Venice Lagoon. Comparison between current and old data. *J. Mar. Syst.* 51, 321–329. doi: [10.1016/j.jmarsys.2004.05.019](https://doi.org/10.1016/j.jmarsys.2004.05.019)
- Adhikari PL, Maiti K, Overton EB 2015 Vertical fluxes of polycyclic aromatic hydrocarbons in the northern Gulf of Mexico. *Mar. Chem.* 168, 60–68. doi: [10.1016/j.marchem.2014.11.001](https://doi.org/10.1016/j.marchem.2014.11.001)
- Adloff F, Jordà G, Somot S, Sevault F, Arsouze T et al. 2018 Improving sea level simulation in Mediterranean regional climate models. *Clim. Dyn.* 51. doi: [10.1007/s00382-017-3842-3](https://doi.org/10.1007/s00382-017-3842-3)
- Adloff F, Somot S, Sevault F, Jordà G, Aznar R et al. 2015 Mediterranean Sea response to climate change in an ensemble of twenty first century scenarios. *Clim. Dyn.* 45, 2775–2802. doi: [10.1007/s00382-015-2507-3](https://doi.org/10.1007/s00382-015-2507-3)
- Adua M 1999 The sweet chestnut throughout history from the Miocene to the Third Millennium. *Acta Hort.*, 29–36. doi: [10.17660/ActaHortic.1999.494.2](https://doi.org/10.17660/ActaHortic.1999.494.2)
- Aerts JCJH, Lin N, Botzen W, Emanuel K, de Moel H 2013 Low-Probability Flood Risk Modeling for New York City. *Risk Anal.* 33, 772–788. doi: [10.1111/risa.12008](https://doi.org/10.1111/risa.12008)
- AghaKouchak A, Huning LS, Chiang F, Sadegh M, Vahedifard F et al. 2018 How do natural hazards cascade to cause disasters? *Nature* 561, 458–460. doi: [10.1038/d41586-018-06783-6](https://doi.org/10.1038/d41586-018-06783-6)
- Aissaoui A, Armi Z, Akrouf F, Ben Hassine OK 2014 Environmental factors and seasonal dynamics of *Proocentrum lima* population in coastal waters of the Gulf of Tunis, South Mediterranean. *Water Environ. Res.* 86, 2256–2270. doi: [10.2175/106143014x13975035526266](https://doi.org/10.2175/106143014x13975035526266)
- Alessandri A, de Felice M, Zeng N, Mariotti A, Pan Y et al. 2015 Robust assessment of the expansion and retreat of Mediterranean climate in the 21st century. *Sci. Rep.* 4, 7211. doi: [10.1038/srep07211](https://doi.org/10.1038/srep07211)
- Alexander MA, Scott JD, Friedland KD, Mills KE, Nye JA et al. 2018 Projected sea surface temperatures over the 21<sup>st</sup> century: Changes in the mean, variability and extremes for large marine ecosystem regions of Northern Oceans. *Elementa* 6, 9. doi: [10.1525/elementa.191](https://doi.org/10.1525/elementa.191)
- Alexandri G, Georgoulas AK, Meleti C, Balis D, Kourtidis KA et al. 2017 A high resolution satellite view of surface solar radiation over the climatically sensitive region of Eastern Mediterranean. *Atmos. Res.* 188, 107–121. doi: [10.1016/J.ATMOSRES.2016.12.015](https://doi.org/10.1016/J.ATMOSRES.2016.12.015)
- Alfaro-Reyna T, Retana J, Martínez-Vilalta J 2018 Is there a substitution of Pinaceae by Fagaceae in temperate forests at the global scale? *Glob. Planet. Change* 166, 41–47. doi: [10.1016/j.gloplacha.2018.04.001](https://doi.org/10.1016/j.gloplacha.2018.04.001)
- Alheit J, Gröger J, Licandro P, McQuinn IH, Pohlmann T et al. 2019 What happened in the mid-1990s? The coupled ocean-atmosphere processes behind climate-induced ecosystem changes in the Northeast Atlantic and the Mediterranean. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 159, 130–142. doi: [10.1016/j.dsr2.2018.11.011](https://doi.org/10.1016/j.dsr2.2018.11.011)
- Alheit J, Licandro P, Coombs S, García A, Giráldez A et al. 2014 Atlantic Multidecadal Oscillation (AMO) modulates dynamics of small pelagic fishes and ecosystem regime shifts in the eastern North and Central Atlantic. *J. Mar. Syst.* 131, 21–35. doi: [10.1016/j.jmarsys.2013.11.002](https://doi.org/10.1016/j.jmarsys.2013.11.002)
- Allen MR, Dube OP, Solecki W, Aragón-Durand F, Cramer W et al. 2018 Framing and Context, in *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, eds. Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J et al. (Cambridge, United Kingdom and New York, USA: Cambridge University Press), 49–91.

- Alpert P, Baldi M, Ilani R, Krichak S, Price C et al. 2006 Chapter 2 Relations between climate variability in the Mediterranean region and the tropics: ENSO, South Asian and African monsoons, hurricanes and Saharan dust, in *Mediterranean*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo RBT-D in E and ES [Elsevier], 149–177.  
doi: [10.1016/S1571-9197\(06\)80005-4](https://doi.org/10.1016/S1571-9197(06)80005-4)
- Alpert P, Neeman BU, Shay-El Y 1990a Climatological analysis of Mediterranean cyclones using ECMWF data. *Tellus A Dyn. Meteorol. Oceanogr.* 42, 65–77.  
doi: [10.3402/tellusa.v42i1.11860](https://doi.org/10.3402/tellusa.v42i1.11860)
- Alpert P, Neeman BU, Shay-El Y 1990b Intermonthly Variability of Cyclone Tracks in the Mediterranean. *J. Clim.* 3, 1474–1478.  
doi: [10.1175/1520-0442\(1990\)003<1474:IVOCTI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1990)003<1474:IVOCTI>2.0.CO;2)
- Alphan H 2018 Analysis of road development and associated agricultural land use change. *Environ. Monit. Assess.* 190, 5.  
doi: [10.1007/s10661-017-6379-3](https://doi.org/10.1007/s10661-017-6379-3)
- Amatulli G, Camia A, San-Miguel-Ayanz J 2013 Estimating future burned areas under changing climate in the EU-Mediterranean countries. *Sci. Total Environ.* 450–451, 209–222.  
doi: [10.1016/j.scitotenv.2013.02.014](https://doi.org/10.1016/j.scitotenv.2013.02.014)
- Améztegui A, Brotons L, Coll L 2010 Land-use changes as major drivers of mountain pine (*Pinus uncinata* Ram.) expansion in the Pyrenees. *Glob. Ecol. Biogeogr.* 19, 632–641.  
doi: [10.1111/j.1466-8238.2010.00550.x](https://doi.org/10.1111/j.1466-8238.2010.00550.x)
- Amici V, Maccherini S, Santi E, Torri D, Vergari F et al. 2017 Long-term patterns of change in a vanishing cultural landscape: A GIS-based assessment. *Ecol. Inform.* 37, 38–51.  
doi: [10.1016/J.ECOINF.2016.11.008](https://doi.org/10.1016/J.ECOINF.2016.11.008)
- Anav A, Mariotti A 2011 Sensitivity of natural vegetation to climate change in the Euro-Mediterranean area. *Clim. Res.* 46, 277–292. doi: [10.3354/cr00993](https://doi.org/10.3354/cr00993)
- Anderson DM, Cembella AD, Hallegraeff GM 2012 Progress in understanding Harmful Algal Blooms: Paradigm shifts and new technologies for research, monitoring, and management. *Ann. Rev. Mar. Sci.* 4, 143–176.  
doi: [10.1146/annurev-marine-120308-081121](https://doi.org/10.1146/annurev-marine-120308-081121)
- Antunes C, Pereira AJ, Fernandes P, Ramos M, Ascensão L et al. 2018 Understanding plant drought resistance in a Mediterranean coastal sand dune ecosystem: differences between native and exotic invasive species. *J. Plant Ecol.* 11, 26–38.  
doi: [10.1093/jpe/rtx014](https://doi.org/10.1093/jpe/rtx014)
- Argenti G, Bottai L, Chiesi M, Maselli F, Stagliano N et al. 2011 Analisi e valutazione di pascoli montani attraverso l'integrazione di dati multispettrali e ausiliari. *Riv. Ital. di Telerilevamento* 43, 45–57. [https://flore.unifi.it/handle/2158/406248#XMN-mJS\\_pPOQ](https://flore.unifi.it/handle/2158/406248#XMN-mJS_pPOQ) [Accessed April 26, 2019].
- Arhami M, Hosseini V, Zare Shahne M, Bigdeli M, Lai A et al. 2017 Seasonal trends, chemical speciation and source apportionment of fine PM in Tehran. *Atmos. Environ.* 153, 70–82. doi: [10.1016/j.atmosenv.2016.12.046](https://doi.org/10.1016/j.atmosenv.2016.12.046)
- Arianoutsou M, Delipetrou P, Celesti-Gradow L, Basnou C, Bazos I et al. 2010 Comparing naturalized alien plants and recipient habitats across an east-west gradient in the Mediterranean Basin. *J. Biogeogr.* 37, 1811–1823.  
doi: [10.1111/j.1365-2699.2010.02324.x](https://doi.org/10.1111/j.1365-2699.2010.02324.x)
- Arianoutsou M, Delipetrou P, Vilà M, Dimitrakopoulos PG, Celesti-Gradow L et al. 2013 Comparative patterns of plant invasions in the Mediterranean biome. *PLoS One* 8, e79174.  
doi: [10.1371/journal.pone.0079174](https://doi.org/10.1371/journal.pone.0079174)
- Armi Z, Milandri A, Turki S, Hajjem B 2011 *Alexandrium catenella* and *Alexandrium tamarense* in the North Lake of Tunis: Bloom characteristics and the occurrence of paralytic shellfish toxin. *African J. Aquat. Sci.* 36, 47–56.  
doi: [10.2989/16085914.2011.559688](https://doi.org/10.2989/16085914.2011.559688)
- Arndt E, Givan O, Edelist D, Sonin O, Belmaker J et al. 2018 Shifts in Eastern Mediterranean fish communities: Abundance changes, trait overlap, and possible competition between native and non-native species. *Fishes* 3, 19.  
doi: [10.3390/fishes3020019](https://doi.org/10.3390/fishes3020019)
- Artale V, Calmanti S, Malanotte-Rizzoli P, Pisacane G, Rupolo V et al. 2006 The Atlantic and Mediterranean Sea as connected systems, in *Developments in Earth and Environmental Sciences* [Elsevier], 282–323.  
doi: [10.1016/S1571-9197\(06\)80008-X](https://doi.org/10.1016/S1571-9197(06)80008-X)
- Asaf D, Perderson D, Peleg M, Matveev V, Luria M 2008 Evaluation of background levels of air pollutants over Israel. *Atmos. Environ.* 42, 8453–8463.  
doi: [10.1016/j.atmosenv.2008.08.011](https://doi.org/10.1016/j.atmosenv.2008.08.011)
- Asia L, Mazouz S, Guiliano M, Dournenq P, Mille G et al. 2009 Occurrence and distribution of hydrocarbons in surface sediments from Marseille Bay (France). *Mar. Pollut. Bull.* 58, 443–451. doi: [10.1016/j.marpolbul.2008.11.022](https://doi.org/10.1016/j.marpolbul.2008.11.022)
- Astraldi M, Conversano F, Civitarese G, Gasparini GP, D'Alcala MR et al. 2002 Water mass properties and chemical signatures in the central Mediterranean region. *J. Mar. Syst.* 33, 155–177. doi: [10.1016/S0924-7963\(02\)00057-X](https://doi.org/10.1016/S0924-7963(02)00057-X)
- Athanasiadis PJ, Wallace JM, Wettstein JJ 2010 Patterns of wintertime jet stream variability and their relation to the storm tracks. *J. Atmos. Sci.* 67, 1361–1381.  
doi: [10.1175/2009JAS3270.1](https://doi.org/10.1175/2009JAS3270.1)
- Auger-Rozenberg M-A, Boivin T 2016 Invasive fruit, cone and seed insects in the Mediterranean Basin, in *Insects and Diseases of Mediterranean Forest Systems* (Cham: Springer International Publishing), 239–259.  
doi: [10.1007/978-3-319-24744-1\\_9](https://doi.org/10.1007/978-3-319-24744-1_9)
- Avtzis DN, Coyle DR, Christopoulos V, Roques A 2017 Biological invasions, national borders, and the current state of non-native insect species in Greece and the neighbouring Balkan countries. *Bull. Insectology* 70, 161–169.  
<http://easin.jrc.ec.europa.eu/> [Accessed April 26, 2019]
- Azorin-Molina C, Vicente-Serrano SM, Mccvicar TR, Jerez S, Sánchez-Lorenzo A et al. 2014 Homogenization and assessment of observed near-surface wind speed trends over Spain and Portugal, 1961–2011. *J. Clim.* 27, 3692–3712.  
doi: [10.1175/JCLI-D-13-00652.1](https://doi.org/10.1175/JCLI-D-13-00652.1)
- Azzurro E, Bariche M 2017 Local knowledge and awareness on the incipient lionfish invasion in the eastern Mediterranean Sea. *Mar. Freshw. Res.* 68, 1950. doi: [10.1071/MF16358](https://doi.org/10.1071/MF16358)
- Azzurro E, Moschella P, Maynou F 2011 Tracking signals of change in Mediterranean fish diversity based on local ecological knowledge. *PLoS One* 6, e24885.  
doi: [10.1371/journal.pone.0024885](https://doi.org/10.1371/journal.pone.0024885)

- Azzurro E, Stancanelli B, Martino V Di, Bariche M 2017 Range expansion of the common lionfish *Pterois miles* (Bennett, 1828) in the Mediterranean Sea: an unwanted new guest for Italian waters. *6*, 95–98. doi: [10.3391/bir.2017.6.2.01](https://doi.org/10.3391/bir.2017.6.2.01)
- Bacer S, Christoudias T, Pozzer A 2016 Projection of North Atlantic Oscillation and its effect on tracer transport. *Atmos. Chem. Phys.* 16, 15581–15592. doi: [10.5194/acp-16-15581-2016](https://doi.org/10.5194/acp-16-15581-2016)
- Baeza A, Rodríguez-Perulero A, Guillén J 2016 Anthropogenic and naturally occurring radionuclide content in near surface air in Cáceres (Spain). *J. Environ. Radioact.* 165, 24–31. doi: [10.1016/j.jenvrad.2016.08.018](https://doi.org/10.1016/j.jenvrad.2016.08.018)
- Balaguer R, Dimastrogiovanni G, García K, González E, Lysimachou A et al. 2018 *Ríos hormonados, Amplia presencia de plaguicidas endocrinos en los ríos españoles*. Ecologista. <https://www.ecologistasenaccion.org/wp-content/uploads/adjuntos-spip/pdf/informe-rios-hormonados.pdf>
- Baldacconi R, Corriero G 2009 Effects of the spread of the alga *Caulerpa racemosa* var. *cylindracea* on the sponge assemblage from coralligenous concretions of the Apulian coast (Ionian Sea, Italy). *Mar. Ecol.* 30, 337–345. doi: [10.1111/j.1439-0485.2009.00282.x](https://doi.org/10.1111/j.1439-0485.2009.00282.x)
- Bale JS, Hayward SAL 2010 Insect overwintering in a changing climate. *J. Exp. Biol.* 213, 980–994. doi: [10.1242/jeb.037911](https://doi.org/10.1242/jeb.037911)
- Balog I, Ruti PM, Tobin I, Armenio V, Vautard R 2016 A numerical approach for planning offshore wind farms from regional to local scales over the Mediterranean. *Renew. Energy* 85, 395–405. doi: [10.1016/j.renene.2015.06.038](https://doi.org/10.1016/j.renene.2015.06.038)
- Ban N, Schmidli J, Schär C 2015 Heavy precipitation in a changing climate: Does short-term summer precipitation increase faster? *Geophys. Res. Lett.* 42, 1165–1172. doi: [10.1002/2014GL025888](https://doi.org/10.1002/2014GL025888) @10.1002/(ISSN)1944-8007.2015 EDHIGHLIGHTS
- Barbeta A, Peñuelas J 2017 Increasing carbon discrimination rates and depth of water uptake favor the growth of Mediterranean evergreen trees in the ecotone with temperate deciduous forests. *Glob. Chang. Biol.* 23, 5054–5068. doi: [10.1111/gcb.13770](https://doi.org/10.1111/gcb.13770)
- Barcikowska MJ, Kapnick SB, Krishnamurty L, Russo S, Cherchi A et al. 2020 Changes in the future summer Mediterranean climate: contribution of teleconnections and local factors. *Earth Syst. Dyn.* 11, 161–181. doi: [10.5194/esd-11-161-2020](https://doi.org/10.5194/esd-11-161-2020)
- Barhoumi B, Castro-Jiménez J, Guigue C, Goutx M, Sempéré R et al. 2018 Levels and risk assessment of hydrocarbons and organochlorines in aerosols from a North African coastal city (Bizerte, Tunisia). *Environ. Pollut.* 240, 422–431. doi: [10.1016/j.envpol.2018.04.109](https://doi.org/10.1016/j.envpol.2018.04.109)
- Bariche M, Torres M, Azzurro E 2013 The presence of the invasive lionfish *Pterois miles* in the Mediterranean Sea. *Mediterr. Mar. Sci.* 14, 292–294. doi: [10.12681/mms.428](https://doi.org/10.12681/mms.428)
- Barkaoui A 2003 *La marine carthaginoise : approche des activités militaires des carthagoins sur mer depuis les origines jusqu'en 146 av. J.-C.* L'Or de temps
- Barnes DKA, Galgani F, Thompson RCC, Barlaz M 2009 Accumulation and fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 1985–1998. doi: [10.1098/rstb.2008.0205](https://doi.org/10.1098/rstb.2008.0205)
- Barnes EA, Polvani L, Barnes EA, Polvani L 2013 Response of the midlatitude jets, and of their variability, to increased greenhouse gases in the CMIP5 models. *J. Clim.* 26, 7117–7135. doi: [10.1175/JCLI-D-12-00536.1](https://doi.org/10.1175/JCLI-D-12-00536.1)
- Barnston AG, Livezey RE 1987 Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. *Mon. Weather Rev.* 115, 1083–1126. doi: [10.1175/1520-0493\(1987\)115<1083:CSAPOL>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<1083:CSAPOL>2.0.CO;2)
- Barreiro-Lostres F, Moreno A, Giralt S, Nadal-Romero E, Valero-Garcés B 2017 Erosion in Mediterranean mountain landscapes during the last millennium: a quantitative approach based on lake sediment sequences (Iberian Range, Spain). *Catena* 149, 782–798. doi: [10.1016/J.CATENA.2016.05.024](https://doi.org/10.1016/J.CATENA.2016.05.024)
- Barrios-Estrada C, de Jesús Rostro-Alanis M, Muñoz-Gutiérrez BD, Iqbal HMN, Kannan S et al. 2018 Emergent contaminants: Endocrine disruptors and their laccase-assisted degradation – A review. *Sci. Total Environ.* 612, 1516–1531. doi: [10.1016/j.scitotenv.2017.09.013](https://doi.org/10.1016/j.scitotenv.2017.09.013)
- Bartók B, Wild M, Folini D, Lüthi D, Kotlarski S et al. 2017 Projected changes in surface solar radiation in CMIP5 global climate models and in EURO-CORDEX regional climate models for Europe. *Clim. Dyn.* 49, 2665–2683. doi: [10.1007/s00382-016-3471-2](https://doi.org/10.1007/s00382-016-3471-2)
- Bartolini G, Morabito M, Crisci A, Grifoni D, Torrigiani T et al. 2008 Recent trends in Tuscany (Italy) summer temperature and indices of extremes. *Int. J. Climatol.* 28, 1751–1760. doi: [10.1002/joc.1673](https://doi.org/10.1002/joc.1673)
- Basnou C, Álvarez E, Bagaria G, Guardiola M, Isern R et al. 2013 Spatial patterns of land use changes across a Mediterranean metropolitan landscape: implications for biodiversity management. *Environ. Manage.* 52, 971–980. doi: [10.1007/s00267-013-0150-5](https://doi.org/10.1007/s00267-013-0150-5)
- Basnou C, Iguzquiza J, Pino J 2015 Examining the role of landscape structure and dynamics in alien plant invasion from urban Mediterranean coastal habitats. *Landsc. Urban Plan.* 136, 156–164. doi: [10.1016/j.landurbplan.2014.12.001](https://doi.org/10.1016/j.landurbplan.2014.12.001)
- Bates AE, Pecl GT, Frusher S, Hobday AJ, Wernberg T et al. 2014 Defining and observing stages of climate-mediated range shifts in marine systems. *Glob. Environ. Chang.* 26, 27–38. doi: [10.1016/J.GLOENVCHA.2014.03.009](https://doi.org/10.1016/J.GLOENVCHA.2014.03.009)
- Battlori E, Parisien M-A, Krawchuk MA, Moritz MA 2013 Climate change-induced shifts in fire for Mediterranean ecosystems. *Glob. Ecol. Biogeogr.* 22, 1118–1129. doi: [10.1111/geb.12065](https://doi.org/10.1111/geb.12065)
- Beaufoy G 2001 EU policies for olive farming-unsustainable on all counts.
- Beaufoy G, Pienkowski M 2000 The environmental impact of olive oil production in the European Union: practical options for improving the environmental impact. <http://ec.europa.eu/environment/agriculture/pdf/oliveoil.pdf> [Accessed April 26, 2019].
- Bedía J, Herrera S, Martín DS, Koutsias N, Gutiérrez JM 2013 Robust projections of Fire Weather Index in the Mediterranean using statistical downscaling. *Clim. Change* 120, 229–247. doi: [10.1007/s10584-013-0787-3](https://doi.org/10.1007/s10584-013-0787-3)
- Belhout D, Kerbachi R, Relvas H, Miranda AI 2018 Air quality assessment in Algiers city. *Air Qual. Atmos. Heal.* 11, 897–906. doi: [10.1007/s11869-018-0589-x](https://doi.org/10.1007/s11869-018-0589-x)

- Belmonte J, Vilà M 2004 Atmospheric invasion of non-native pollen in the Mediterranean region. *Am. J. Bot.* 91, 1243–1250. doi: [10.3732/ajb.91.8.1243](https://doi.org/10.3732/ajb.91.8.1243)
- Belušić Vozila A, Güttler I, Ahrens B, Obermann-Hellhund A, Telišman Prtenjak M 2019 Wind Over the Adriatic Region in CORDEX Climate Change Scenarios. *J. Geophys. Res. Atmos.* 124, 110–130. doi: [10.1029/2018JD028552](https://doi.org/10.1029/2018JD028552)
- Ben-Gai T, Bitan A, Manes A, Alpert P, Kushnir Y 2001 Temperature and surface pressure anomalies in Israel and the North Atlantic Oscillation. *Theor. Appl. Climatol.* 69, 171–177. doi: [10.1007/s007040170023](https://doi.org/10.1007/s007040170023)
- Ben Othman H, Leboulanger C, Le Floc'h E, Mabrouk HH, Sakka Hlaili A 2012 Toxicity of benz(a)anthracene and fluoranthene to marine phytoplankton in culture: Does cell size really matter? *J. Hazard. Mater.* 243, 204–211. doi: [10.1016/j.jhazmat.2012.10.020](https://doi.org/10.1016/j.jhazmat.2012.10.020)
- Benabdelkader A, Taleb A, Probst JL, Belaidi N, Probst A 2018 Anthropogenic contribution and influencing factors on metal features in fluvial sediments from a semi-arid Mediterranean river basin (Tafna River, Algeria): A multi-indices approach. *Sci. Total Environ.* 626, 899–914. doi: [10.1016/j.scitotenv.2018.01.107](https://doi.org/10.1016/j.scitotenv.2018.01.107)
- Benas N, Chrysoulakis N 2015 Estimation of the Land Surface Albedo Changes in the Broader Mediterranean Area, Based on 12 Years of Satellite Observations. *Remote Sens.* 7, 16150–16163. doi: [10.3390/rs71215816](https://doi.org/10.3390/rs71215816)
- Bengtsson L, Hodges KI, Roeckner E 2006 Storm tracks and climate change. *J. Clim.* 19.
- Beniston M, Farinotti D, Stoffel M, Andreassen LM, Coppola E et al. 2018 The European mountain cryosphere: a review of its current state, trends, and future challenges. *Cryosph.* 12, 759–794. doi: [10.5194/tc-12-759-2018](https://doi.org/10.5194/tc-12-759-2018)
- Beniston M, Stephenson DB, Christensen OB, Ferro CAT, Frei C et al. 2007 Future extreme events in European climate: an exploration of regional climate model projections. *Clim. Change* 81, 71–95. doi: [10.1007/s10584-006-9226-z](https://doi.org/10.1007/s10584-006-9226-z)
- Benoit G, Comeau A 2005 *A Sustainable Future for the Mediterranean*. London: Routledge doi: [10.4324/9781849770323](https://doi.org/10.4324/9781849770323)
- Bensoussan N, Chiggiato J, Buongiorno Nardelli B, Pisano A, Garrabou J 2019 Insights on 2017 Marine Heat Waves in the Mediterranean Sea. In: Copernicus Marine Service Ocean State Report, Issue 3. *J. Oper. Oceanogr.* 12, S26–S30. doi: [10.1080/1755876X.2019.1633075](https://doi.org/10.1080/1755876X.2019.1633075)
- Berndtsson R, Jebari S, Hashemi H, Wessels J 2016 Traditional irrigation techniques in MENA with focus on Tunisia. *Hydrol. Sci. J.* 61, 1346–1357. doi: [10.1080/02626667.2016.1165349](https://doi.org/10.1080/02626667.2016.1165349)
- Berrojalbiz N, Dachs J, Ojeda MJ, Valle MC, Castro-Jiménez J et al. 2011 Biogeochemical and physical controls on concentrations of polycyclic aromatic hydrocarbons in water and plankton of the Mediterranean and Black Seas. *Global Biogeochem. Cycles* 25, 1–14. doi: [10.1029/2010GB003775](https://doi.org/10.1029/2010GB003775)
- Berthou S, Kendon EJ, Chan SC, Ban N, Leutwyler D et al. 2018 Pan-European climate at convection-permitting scale: a model intercomparison study. *Clim. Dyn.*, 1–25. doi: [10.1007/s00382-018-4114-6](https://doi.org/10.1007/s00382-018-4114-6)
- Béthoux JP, Gentili B 1999 Functioning of the Mediterranean Sea: past and present changes related to freshwater input and climate changes. *J. Mar. Syst.* 20, 33–47. doi: [10.1016/S0924-7963\(98\)00069-4](https://doi.org/10.1016/S0924-7963(98)00069-4)
- Béthoux JP, Gentili B, Morin P, Nicolas E, Pierre C et al. 1999 The Mediterranean Sea: a miniature ocean for climatic and environmental studies and a key for the climatic functioning of the North Atlantic. *Prog. Oceanogr.* 44, 131–146. doi: [10.1016/S0079-6611\(99\)00023-3](https://doi.org/10.1016/S0079-6611(99)00023-3)
- Béthoux JP, Morin P, Chaumery C, Connan O, Gentili B et al. 1998 Nutrients in the Mediterranean Sea, mass balance and statistical analysis of concentrations with respect to environmental change. *Mar. Chem.* 63, 155–169. doi: [10.1016/S0304-4203\(98\)00059-0](https://doi.org/10.1016/S0304-4203(98)00059-0)
- Bett PE, Thornton HE, Clark RT 2013 European wind variability over 140 yr. *Adv. Sci. Res.* 10, 51–58. doi: [10.5194/asr-10-51-2013](https://doi.org/10.5194/asr-10-51-2013)
- Beuvier J, Sevault F, Herrmann M, Kontoyiannis H, Ludwig W et al. 2010 Modeling the Mediterranean Sea interannual variability during 1961–2000: Focus on the Eastern Mediterranean Transient. *JGR Ocean.* 115, C08017. doi: [10.1029/2009JC005950](https://doi.org/10.1029/2009JC005950)
- Bianchi CN 2007 Biodiversity issues for the forthcoming tropical Mediterranean Sea. in *Biodiversity in Enclosed Seas and Artificial Marine Habitats* Developments in (Hydrobiology)., eds. Relini G, Ryland J (Springer Netherlands), 7–21. doi: [10.1007/s10750-006-0469-5](https://doi.org/10.1007/s10750-006-0469-5)
- Biel-Maeso M, Baena-Nogueras RM, Corada-Fernández C, Lara-Martín PA 2018 Occurrence, distribution and environmental risk of pharmaceutically active compounds (PhACs) in coastal and ocean waters from the Gulf of Cadiz (SW Spain). *Sci. Total Environ.* 612, 649–659. doi: [10.1016/j.scitotenv.2017.08.279](https://doi.org/10.1016/j.scitotenv.2017.08.279)
- Bilal M, Adeel M, Rasheed T, Zhao Y, Iqbal HMN 2019 Emerging contaminants of high concern and their enzyme-assisted biodegradation – A review. *Env. Int* 124, 336–353. doi: [10.1016/j.envint.2019.01.011](https://doi.org/10.1016/j.envint.2019.01.011)
- Bilal M, Rasheed T, Sosa-Hernández JE, Raza A, Nabeel F et al. 2018 Biosorption: An interplay between marine algae and potentially toxic elements-A review. *Mar. Drugs* 16, 65. doi: [10.3390/md16020065](https://doi.org/10.3390/md16020065)
- Bilbao J, Román R, de Miguel A 2019 Temporal and Spatial Variability in Surface Air Temperature and Diurnal Temperature Range in Spain over the Period 1950–2011. *Climate* 7, 16. doi: [10.3390/cli7010016](https://doi.org/10.3390/cli7010016)
- Bladé I, Fortuny D, van Oldenborgh GJ, Liebmann B 2012 The summer North Atlantic Oscillation in CMIP3 models and related uncertainties in projected summer drying in Europe. *JGR Atmos.* 117. doi: [10.1029/2012JD017816](https://doi.org/10.1029/2012JD017816)
- Blondel J 2006 The 'design' of Mediterranean landscapes: a millennial story of humans and ecological systems during the historic period. *Hum. Ecol.* 34, 713–729. doi: [10.1007/s10745-006-9030-4](https://doi.org/10.1007/s10745-006-9030-4)
- Boberg F, Christensen JH 2012 Overestimation of Mediterranean summer temperature projections due to model deficiencies. *Nat. Clim. Chang.* 2, 433–436. doi: [10.1038/nclimate1454](https://doi.org/10.1038/nclimate1454)
- Boé J, Somot S, Corre L, Nabat P 2020 Large discrepancies in summer climate change over Europe as projected by glob-

- al and regional climate models: causes and consequences. *Clim. Dyn.* 54, 2981–3002. doi: [10.1007/s00382-020-05153-1](https://doi.org/10.1007/s00382-020-05153-1)
- Boé J, Terray L 2014 Land–sea contrast, soil–atmosphere and cloud–temperature interactions: interplays and roles in future summer European climate change. *Clim. Dyn.* 42, 683–699. doi: [10.1007/s00382-013-1868-8](https://doi.org/10.1007/s00382-013-1868-8)
- Borghini M, Bryden HL, Schroeder K, Sparnocchia S, Vetrano A 2014 The Mediterranean is becoming saltier. *Ocean Sci.* 10, 693–700. doi: [10.5194/os-10-693-2014](https://doi.org/10.5194/os-10-693-2014)
- Borrego C, Monteiro A, Sá E, Carvalho A, Coelho D et al. 2012 Reducing NO<sub>2</sub> pollution over urban areas: Air quality modelling as a fundamental management tool. *Water, Air, Soil Pollut.* 223, 5307–5320. doi: [10.1007/s11270-012-1281-7](https://doi.org/10.1007/s11270-012-1281-7)
- Bouchouicha-Smida D, Lundholm N, Sahraoui I, Lambert C, Mabrouk HH et al. 2015 Detection of domoic acid in *Mytilus galloprovincialis* and *Ostrea edulis* linked to the presence of *Nitzschia bizertensis* in Bizerte Lagoon (SW Mediterranean). *Estuar. Coast. Shelf Sci.* 165, 270–278. doi: [10.1016/j.ecss.2015.05.029](https://doi.org/10.1016/j.ecss.2015.05.029)
- Boudouresque C-F, Verlaque M 2010 Is global warming involved in the success of seaweed introductions in the Mediterranean Sea?, in *Seaweeds and their role in globally changing environments*, eds. Israel A, Einav R, Seckbach J (Springer Netherlands), 31–50. doi: [10.1007/978-90-481-8569-6\\_3](https://doi.org/10.1007/978-90-481-8569-6_3)
- Boutou D, Peliz Á, Miranda PMA, Soares PMM, Cardoso RM et al. 2014 Inter-annual variability and long term predictability of exchanges through the Strait of Gibraltar. *Glob. Planet. Change* 114, 23–37. doi: [10.1016/j.gloplacha.2013.12.009](https://doi.org/10.1016/j.gloplacha.2013.12.009)
- Bowman DMJS, Williamson GJ, Abatzoglou JT, Kolden CA, Cochrane MA et al. 2017 Human exposure and sensitivity to globally extreme wildfire events. *Nat. Ecol. Evol.* 1, 1–6. doi: [10.1038/s41559-016-0058](https://doi.org/10.1038/s41559-016-0058)
- Boxall ABA, Hardy A, Beulke S, Boucard T, Burgin L et al. 2009 Impacts of Climate Change on Indirect Human Exposure to Pathogens and Chemicals from Agriculture. *Environ. Health Perspect.* 117, 508–514. doi: [10.1289/ehp.0800084](https://doi.org/10.1289/ehp.0800084)
- Bradley BA, Blumenthal DM, Early R, Grosholz ED, Lawler JJ et al. 2012 Global change, global trade, and the next wave of plant invasions. *Front. Ecol. Environ.* 10, 20–28. doi: [10.1890/110145](https://doi.org/10.1890/110145)
- Branco M, Battisti A, Mendel Z 2016 Foliage Feeding Invasive Insects: Defoliators and Gall Makers, in *Insects and Diseases of Mediterranean Forest Systems*, eds. Paine T, Lieutier F (Cham: Springer International Publishing), 211–238. doi: [10.1007/978-3-319-24744-1\\_8](https://doi.org/10.1007/978-3-319-24744-1_8)
- Bras A, Avtzis DN, Kenis M, Li H, Véték G et al. 2019 A complex invasion story underlies the fast spread of the invasive box tree moth (*Cydalima perspectalis*) across Europe. *J. Pest Sci. (2004)*. doi: [10.1007/s10340-019-01111-x](https://doi.org/10.1007/s10340-019-01111-x)
- Bregaglio S, Hossard L, Cappelli G, Resmond R, Bocchi S et al. 2017 Identifying trends and associated uncertainties in potential rice production under climate change in Mediterranean areas. *Agric. For. Meteorol.* 237–238, 219–232. doi: [10.1016/J.AGRFORMET.2017.02.015](https://doi.org/10.1016/J.AGRFORMET.2017.02.015)
- Brilli L, Lugato E, Moriondo M, Gioli B, Toscano P et al. 2019 Carbon sequestration capacity and productivity responses of Mediterranean olive groves under future climates and management options. *Mitig. Adapt. Strateg. Glob. Chang.* 24, 467–491. doi: [10.1007/s11027-018-9824-x](https://doi.org/10.1007/s11027-018-9824-x)
- Brook BW, Sodhi NS, Bradshaw CJA 2008 Synergies among extinction drivers under global change. *Trends Ecol. Evol.* 23, 453–460. doi: [10.1016/J.TREE.2008.03.011](https://doi.org/10.1016/J.TREE.2008.03.011)
- Brown RD, Mote PW 2009 The response of Northern hemisphere snow cover to a changing climate. *J. Clim.* 22, 2124–2145. doi: [10.1175/2008JCLI2665.1](https://doi.org/10.1175/2008JCLI2665.1)
- Brundu G 2013 Invasive Alien Plants in Protected Areas in Mediterranean Islands: Knowledge Gaps and Main Threats, in *Plant Invasions in Protected Areas*, eds. Foxcroft LC, Pyšek P, Richardson DM, Genovesi P (Dordrecht: Springer Netherlands), 395–422. doi: [10.1007/978-94-007-7750-7\\_18](https://doi.org/10.1007/978-94-007-7750-7_18)
- Brunetti M, Maugeri M, Monti F, Nanni T 2006 Temperature and precipitation variability in Italy in the last two centuries from homogenised instrumental time series. *Int. J. Climatol.* 26, 345–381. doi: [10.1002/joc.1251](https://doi.org/10.1002/joc.1251)
- Buccolieri A, Buccolieri G, Cardellicchio N, Dell'Atti A, di Leo A et al. 2006 Heavy metals in marine sediments of Taranto Gulf (Ionian Sea, Southern Italy). *Mar. Chem.* 99, 227–235. doi: [10.1016/j.marchem.2005.09.009](https://doi.org/10.1016/j.marchem.2005.09.009)
- Bulleri F, Benedetti-Cecchi L 2008 Facilitation of the introduced green alga *Caulerpa racemosa* by resident algal turfs: experimental evaluation of underlying mechanisms. *Mar. Ecol. Prog. Ser.* 364, 77–86. doi: [10.3354/meps07484](https://doi.org/10.3354/meps07484)
- Bustamante MA, Helmer EH, Schill S, Belnap J, Brown LK et al. 2018 Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for the Americas*, eds. Rice J, Seixas CS, Zaccagnini ME, Bedoya-Gaitán M, Valderrama N (Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), 335–509.
- Cabezas S, Carrete M, Tella JL, Marchant TA, Bortolotti GR 2013 Differences in acute stress responses between wild-caught and captive-bred birds: a physiological mechanism contributing to current avian invasions? *Biol. Invasions* 15, 521–527. doi: [10.1007/s10530-012-0304-z](https://doi.org/10.1007/s10530-012-0304-z)
- Cabral JPS 2010 Water Microbiology. Bacterial Pathogens and Water. *Int. J. Environ. Res. Public Health* 7, 3657–3703. doi: [10.3390/ijerph7103657](https://doi.org/10.3390/ijerph7103657)
- Calafat FM, Jordà G, Marcos M, Gomis D 2012 Comparison of Mediterranean sea level variability as given by three baroclinic models. *JGR Ocean.* 117. doi: [10.1029/2011JC007277](https://doi.org/10.1029/2011JC007277)
- Caloiero T, Veltri S, Caloiero P, Frustaci F 2018 Drought Analysis in Europe and in the Mediterranean Basin Using the Standardized Precipitation Index. *Water* 10, 1043. doi: [10.3390/w10081043](https://doi.org/10.3390/w10081043)
- Camarero JJ, Gazol A, Tardif JC, Conciatori F 2015 Attributing forest responses to global-change drivers: limited evidence of a CO<sub>2</sub>-fertilization effect in Iberian pine growth. *J. Biogeogr.* 42, 2220–2233. doi: [10.1111/jbi.12590](https://doi.org/10.1111/jbi.12590)
- Camera C, Bruggeman A, Hadjinicolaou P, Michaelides S, Lange MA 2017 Evaluation of a spatial rainfall generator for generating high resolution precipitation projections over orographically complex terrain. *Stoch. Environ. Res. Risk*

- Assess. 31, 757–773. doi: [10.1007/s00477-016-1239-1](https://doi.org/10.1007/s00477-016-1239-1)
- Cameron EK, Vilà M, Cabeza M 2016 Global meta-analysis of the impacts of terrestrial invertebrate invaders on species, communities and ecosystems. *Glob. Ecol. Biogeogr.* 25, 596–606. doi: [10.1111/geb.12436](https://doi.org/10.1111/geb.12436)
- Campins J, Genovés A, Picornell MA, Jansà A 2011 Climatology of Mediterranean cyclones using the ERA-40 dataset. *Int. J. Climatol.* 31, 1596–1614. doi: [10.1002/joc.2183](https://doi.org/10.1002/joc.2183)
- Campo J, Masiá A, Blasco C, Picó Y 2013 Occurrence and removal efficiency of pesticides in sewage treatment plants of four Mediterranean River Basins. *J. Hazard. Mater.* 263, 146–157. doi: [10.1016/j.jhazmat.2013.09.061](https://doi.org/10.1016/j.jhazmat.2013.09.061)
- Camuffo D, Bertolin C, Barriendos M, Domínguez-Castro F, Cocheo C et al. 2010 500-Year temperature reconstruction in the Mediterranean Basin by means of documentary data and instrumental observations. *Clim. Change* 101, 169–199. doi: [10.1007/s10584-010-9815-8](https://doi.org/10.1007/s10584-010-9815-8)
- Cannarsa S, Abete MC, Zanardi M, Squadrone S 2014 Polycyclic aromatic hydrocarbons (PAH) in marine sediment of the northwestern Mediterranean Sea (Italy). <http://blackmed-itjournal.org/wp-content/uploads/137-141-Vol20No2-Cannarsa.pdf> [Accessed April 17, 2019]
- Cardador L, Tella JL, Anadón JD, Abellán P, Carrete M 2019 The European trade ban on wild birds reduced invasion risks. *Conserv. Lett.*, e12631. doi: [10.1111/conl.12631](https://doi.org/10.1111/conl.12631)
- Cardinale M, Osio GC, Scarcella G 2017 Mediterranean Sea: a failure of the European fisheries management system, in *Frontiers in Marine Science* (Frontiers Media SA). doi: [10.3389/fmars.2017.00072](https://doi.org/10.3389/fmars.2017.00072)
- Carillo A, Sannino G, Artale V, Ruti PM, Calmanti S et al. 2012 Steric sea level rise over the Mediterranean Sea: present climate and scenario simulations. *Clim. Dyn.* 39, 2167–2184. doi: [10.1007/s00382-012-1369-1](https://doi.org/10.1007/s00382-012-1369-1)
- Carlton JT 1989 Man's Role in Changing the Face of the Ocean: Biological Invasions and Implications for Conservation of Near-Shore Environments. *Conserv. Biol.* 3, 265–273. doi: [10.2307/2386170](https://doi.org/10.2307/2386170)
- Carlton JT, Chapman JW, Geller JB, Miller JA, Carlton DA et al. 2017 Tsunami-driven rafting: Transoceanic species dispersal and implications for marine biogeography. *Science* (80-.). 357, 1402–1406. doi: [10.1126/science.aao1498](https://doi.org/10.1126/science.aao1498)
- Carranza ML, Drius M, Malavasi M, Frate L, Stanisci A et al. 2018 Assessing land take and its effects on dune carbon pools. An insight into the Mediterranean coastline. *Ecol. Indic.* 85, 951–955. doi: [10.1016/j.ecolind.2017.10.052](https://doi.org/10.1016/j.ecolind.2017.10.052)
- Carrer D, Pique G, Ferlicoq M, Ceamanos X, Ceschia E 2018 What is the potential of cropland albedo management in the fight against global warming? A case study based on the use of cover crops. *Environ. Res. Lett.* 13, 044030. doi: [10.1088/1748-9326/aab650](https://doi.org/10.1088/1748-9326/aab650)
- Carrete M, Tella J 2008 Wild-bird trade and exotic invasions: a new link of conservation concern? *Front. Ecol. Environ.* 6, 207–211. doi: [10.1890/070075](https://doi.org/10.1890/070075)
- Carrete M, Tella JL 2015 Rapid loss of antipredatory behaviour in captive-bred birds is linked to current avian invasions. *Sci. Rep.* 5, e18274. doi: [10.1038/srep18274](https://doi.org/10.1038/srep18274)
- Carta A, Taboada T, Müller J V 2018 Diachronic analysis using aerial photographs across fifty years reveals significant land use and vegetation changes on a Mediterranean island. *Appl. Geogr.* 98, 78–86. doi: [10.1016/j.apgeog.2018.07.010](https://doi.org/10.1016/j.apgeog.2018.07.010)
- Castellanos-Frías E, García de León D, Pujadas-Salva A, Dorado J, Gonzalez-Andujar JL 2014 Potential distribution of *Avena sterilis* L. in Europe under climate change. *Ann. Appl. Biol.* 165, 53–61. doi: [10.1111/aab.12117](https://doi.org/10.1111/aab.12117)
- Castro-Díez P, González-Muñoz N, Alonso AM, Gallardo A, Poorter L 2009 Effects of exotic invasive trees on nitrogen cycling: A case study in Central Spain. *Biol. Invasions* 11, 1973–1986. doi: [10.1007/s10530-008-9374-3](https://doi.org/10.1007/s10530-008-9374-3)
- Castro-Díez Y, Pozo-Vázquez D, Rodrigo FS, Esteban-Parra MJ 2002 NAO and winter temperature variability in southern Europe. *Geophys. Res. Lett.* 29, 1-1-1–4. doi: [10.1029/2001gl014042](https://doi.org/10.1029/2001gl014042)
- Cavicchia L, von Storch H, Gualdi S 2014 Mediterranean Tropical-Like Cyclones in Present and Future Climate. *J. Clim.* 27, 7493–7501. doi: [10.1175/jcli-d-14-00339.1](https://doi.org/10.1175/jcli-d-14-00339.1)
- Cazenave A, WCRP Global Sea Level Budget Group 2018 Global sea-level budget 1993-present. *Earth Syst. Sci. Data* 10, 1551–1590. doi: [10.5194/essd-10-1551-2018](https://doi.org/10.5194/essd-10-1551-2018)
- Ccancapa A, Masiá A, Andreu V, Picó Y 2015 Spatio-temporal patterns of pesticide residues in the Turia and Júcar Rivers (Spain). *Sci. Total Environ.* 540, 200–210. doi: [10.1016/j.scitotenv.2015.06.063](https://doi.org/10.1016/j.scitotenv.2015.06.063)
- CEAM 2019 Mediterranean Sea Surface Temperature report (Summer 2019). doi: [10.13140/RG.2.2.23375.23209](https://doi.org/10.13140/RG.2.2.23375.23209)
- Cebrián E, Tomas F, López-Sendino P, Vilà M, Ballesteros E 2018 Biodiversity influences invasion success of a facultative epiphytic seaweed in a marine forest. *Biol. Invasions* 20, 2839–2848. doi: [10.1007/s10530-018-1736-x](https://doi.org/10.1007/s10530-018-1736-x)
- Ceccherelli G, Piazzini L, Balata D 2002 Spread of introduced *Caulerpa* species in macroalgal habitats. *J. Exp. Mar. Biol. Ecol.* 280, 1–11. doi: [10.1016/s0022-0981\(02\)00336-2](https://doi.org/10.1016/s0022-0981(02)00336-2)
- Ceccherini G, Russo S, Amezttoy I, Marchese AF, Carmo-Moreno C 2017 Heat waves in Africa 1981–2015, observations and reanalysis. *Nat. Hazards Earth Syst. Sci.* 17, 115–125. doi: [10.5194/nhess-2016-90](https://doi.org/10.5194/nhess-2016-90)
- Cecchi P, Garrido M, Collos Y, Pasqualini V 2016 Water flux management and phytoplankton communities in a Mediterranean coastal lagoon. Part II: Mixotrophy of dinoflagellates as an adaptive strategy? *Mar. Pollut. Bull.* 108, 120–133. doi: [10.1016/j.marpolbul.2016.04.041](https://doi.org/10.1016/j.marpolbul.2016.04.041)
- Ceglar A, Zampieri M, Toreti A, Dentener FJ 2019 Observed Northward Migration of Agro-Climatic Zones in Europe Will Further Accelerate Under Climate Change. *Earth's Futur.* 7, 1088–1101. doi: [10.1029/2019ef001178](https://doi.org/10.1029/2019ef001178)
- Celesti-Grapow L, Bassi L, Brundu G, Camarda I, Carli E et al. 2016 Plant invasions on small Mediterranean islands: An overview. *Plant Biosyst. - An Int. J. Deal. with all Asp. Plant Biol.* 150, 1119–1133. doi: [10.1080/11263504.2016.1218974](https://doi.org/10.1080/11263504.2016.1218974)
- Cerdan O, Le Bissonnais Y, Govers G, Lecomte V, van Oost K et al. 2004 Scale effect on runoff from experimental plots to catchments in agricultural areas in Normandy. *J. Hydrol.* 299, 4–14. doi: [10.1016/j.jhydrol.2004.02.017](https://doi.org/10.1016/j.jhydrol.2004.02.017)
- Chang EKM, Guo Y, Xia X 2012 CMIP5 multimodel ensemble



- projection of storm track change under global warming. *JGR Atmos.* 117. doi: [10.1029/2012jd018578](https://doi.org/10.1029/2012jd018578)
- Chatha SAS, Asgher M, Iqbal HMN 2017 Enzyme-based solutions for textile processing and dye contaminant biodegradation—a review. *Environ. Sci. Pollut. Res.* 24, 14005–14018. doi: [10.1007/s11356-017-8998-1](https://doi.org/10.1007/s11356-017-8998-1)
- Cheggour A, Simonneaux V, Roose E 2012 Effets des plantations forestières sur banquettes sur le ruissellement et l'érosion par rapport aux parcours dans les montagnes semi-arides du Haut-Atlas de Marrakech (Maroc), in *Lutte antiérosive: réhabilitation des sols tropicaux et protection contre les pluies exceptionnelles*, eds. Roose E, Duchaufour H (Marseille, France).
- Cheminée A, Sala E, Pastor J, Bodilis P, Thiriet P et al. 2013 Nursery value of *Cystoseira* forests for Mediterranean rocky reef fishes. *J. Exp. Mar. Bio. Ecol.* 442, 70–79. doi: [10.1016/j.jembe.2013.02.003](https://doi.org/10.1016/j.jembe.2013.02.003)
- Chen Y, Paytan A, Chase Z, Measures C, Beck AJ et al. 2008 Sources and fluxes of atmospheric trace elements to the Gulf of Aqaba, Red Sea. *JGR Atmos.* 113. doi: [10.1029/2007JD009110](https://doi.org/10.1029/2007JD009110)
- Cheung WWL, Lam WWY, Sarmiento JL, Kearney K, Watson R et al. 2009 Projecting global marine biodiversity impacts under climate change scenarios. *Fish Fish.* 10, 235–251. doi: [10.1111/j.1467-2979.2008.00315.x](https://doi.org/10.1111/j.1467-2979.2008.00315.x)
- Cheung WWL, Lam WWY, Sarmiento JL, Kearney K, Watson R et al. 2010 Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change. *Glob. Chang. Biol.* 16, 24–35. doi: [10.1111/j.1365-2486.2009.01995.x](https://doi.org/10.1111/j.1365-2486.2009.01995.x)
- Cheung WWL, Sarmiento JL, Dunne J, Frölicher TL, Lam WWY et al. 2013a Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nat. Clim. Chang.* 3, 254–258. doi: [10.1038/nclimate1691](https://doi.org/10.1038/nclimate1691)
- Cheung WWL, Watson R, Pauly D 2013b Signature of ocean warming in global fisheries catch. *Nature* 497, 365–368. doi: [10.1038/nature12156](https://doi.org/10.1038/nature12156)
- Chiacchio M, Wild M 2010 Influence of NAO and clouds on long-term seasonal variations of surface solar radiation in Europe. *JGR Atmos.* 115, D00D22. doi: [10.1029/2009JD012182](https://doi.org/10.1029/2009JD012182)
- Cho S-H, Richmond-Bryant J, Thornburg J, Portzer J, Vanderpool R et al. 2011 A literature review of concentrations and size distributions of ambient airborne Pb-containing particulate matter. *Atmos. Environ.* 45, 5005–5015. doi: [10.1016/j.atmosenv.2011.05.009](https://doi.org/10.1016/j.atmosenv.2011.05.009)
- Christensen JH, Carter TR, Giorgi F 2002 PRUDENCE employs new methods to assess European climate change. *Eos, Trans. Am. Geophys. Union* 83, 147. doi: [10.1029/2002eo000094](https://doi.org/10.1029/2002eo000094)
- Christensen JH, Hewitson B, Busuioc A, Chen A, Gao X et al. 2007 Regional Climate Projections, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Solomon S, Qin D, Manning M, Chen Z, Marquis M et al., 847–940.
- Chu Y, Salles C, Tournoud M-G, Got P, Troussellier M et al. 2011 Faecal bacterial loads during flood events in Northwestern Mediterranean coastal rivers. *J. Hydrol.* 405, 501–511. doi: [10.1016/j.jhydrol.2011.05.047](https://doi.org/10.1016/j.jhydrol.2011.05.047)
- Chuine I, Morin X, Sonié L, Collin C, Fabreguettes J et al. 2012 Climate change might increase the invasion potential of the alien *C<sub>4</sub>* grass *Setaria parviflora* (Poaceae) in the Mediterranean Basin. *Divers. Distrib.* 18, 661–672. doi: [10.1111/j.1472-4642.2011.00880.x](https://doi.org/10.1111/j.1472-4642.2011.00880.x)
- Church JA, Clark PU, Cazenave A, Gregory JM, Jevrejeva S et al. 2013 Sea level change. PM Cambridge University Press
- Chytrý M, Maskell LC, Pino J, Pyšek P, Vilà M et al. 2008 Habitat invasions by alien plants: a quantitative comparison among Mediterranean, subcontinental and oceanic regions of Europe. *J. Appl. Ecol.* 45, 448–458. doi: [10.1111/j.1365-2664.2007.01398.x](https://doi.org/10.1111/j.1365-2664.2007.01398.x)
- Ciminiello P, Dell'Aversano C, Dello Iacovo E, Fattorusso E, Forino M et al. 2014 First finding of *Ostreopsis cf. ovata* toxins in marine aerosols. *Environ. Sci. Technol.* 48, 3532–3540. doi: [10.1021/es405617d](https://doi.org/10.1021/es405617d)
- Cincinelli A, Martellini T, Guerranti C, Scopetani C, Chelazzi D et al. 2019 A potpourri of microplastics in the sea surface and water column of the Mediterranean Sea. *TrAC - Trends Anal. Chem.* 110, 321–326. doi: [10.1016/j.trac.2018.10.026](https://doi.org/10.1016/j.trac.2018.10.026)
- Cini A, Anfora G, Escudero-Colomar LA, Grassi A, Santosuosso U et al. 2014 Tracking the invasion of the alien fruit pest *Drosophila suzukii* in Europe. *J. Pest Sci. (2004)*. 87, 559–566. doi: [10.1007/s10340-014-0617-z](https://doi.org/10.1007/s10340-014-0617-z)
- Clarke L, Jiang K, Akimoto K, Babiker M, Blanford G et al. 2014 Assessing Transformation Pathways, in *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Edenhofer O, Pichs-Madruga R, Sokona Y, Farahani E, Kadner S et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press).
- Clavero M, Brotons L, Pons P, Sol D 2009 Prominent role of invasive species in avian biodiversity loss. *Biol. Conserv.* 142, 2043–2049. doi: [10.1016/j.biocon.2009.03.034](https://doi.org/10.1016/j.biocon.2009.03.034)
- Clotet M, Basnou C, Bagaria G, Pino J 2016 Contrasting historical and current land-use correlation with diverse components of current alien plant invasions in Mediterranean habitats. *Biol. Invasions* 18, 2897–2909. doi: [10.1007/s10530-016-1181-7](https://doi.org/10.1007/s10530-016-1181-7)
- Cofiño AS, San-Martín D, Gutiérrez JM 2007 A web portal for regional projection of weather forecast using GRID middleware, in *Lecture Notes in Computer Science*, 82–89.
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Ben Rais Lasram F et al. 2010 The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS One* 5, e11842. doi: [10.1371/journal.pone.0011842](https://doi.org/10.1371/journal.pone.0011842)
- Collins M, Knutti R, Arblaster J, Dufresne J-L, Fichetef T et al. 2013 Long-term Climate Change: Projections, Commitments and Irreversibility, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. (Cambridge, United Kingdom and New York, NY, USA), 1029–1136. doi: [10.1017/CBO9781107415324.024](https://doi.org/10.1017/CBO9781107415324.024)

- Colloca F, Cardinale M, Maynou F, Giannoulaki M, Scarcella G et al. 2013 Rebuilding Mediterranean fisheries: a new paradigm for ecological sustainability. *Fish Fish.* 14, 89–109. doi: [10.1111/j.1467-2979.2011.00453.x](https://doi.org/10.1111/j.1467-2979.2011.00453.x)
- Colloca F, Scarcella G, Libralato S 2017 Recent Trends and Impacts of Fisheries Exploitation on Mediterranean Stocks and Ecosystems. *Front. Mar. Sci.* 4. doi: [10.3389/fmars.2017.00244](https://doi.org/10.3389/fmars.2017.00244)
- Colucci RR, Guglielmin M 2019 Climate change and rapid ice melt: Suggestions from abrupt permafrost degradation and ice melting in an alpine ice cave. *Prog. Phys. Geogr. Earth Environ.* 43. doi: [10.1177%2F0309133319846056](https://doi.org/10.1177%2F0309133319846056)
- Colucci RR, Žebre M 2016 Late Holocene evolution of glaciers in the southeastern Alps. *J. Maps* 12, 289–299. doi: [10.1080/17445647.2016.1203216](https://doi.org/10.1080/17445647.2016.1203216)
- Compa M, Alomar C, Wilcox C, van Sebille E, Lebreton L et al. 2019 Risk assessment of plastic pollution on marine diversity in the Mediterranean Sea. *Sci. Total Environ.* 678, 188–196. doi: [10.1016/j.scitotenv.2019.04.355](https://doi.org/10.1016/j.scitotenv.2019.04.355)
- Constán-Nava S, Soliveres S, Torices R, Serra L, Bonet A 2015 Direct and indirect effects of invasion by the alien tree *Ailanthus altissima* on riparian plant communities and ecosystem multifunctionality. *Biol. Invasions* 17, 1095–1108. doi: [10.1007/s10530-014-0780-4](https://doi.org/10.1007/s10530-014-0780-4)
- Conte D, Lionello P 2013 Characteristics of large positive and negative surges in the Mediterranean Sea and their attenuation in future climate scenarios. *Glob. Planet. Change* 111, 159–173. doi: [10.1016/j.gloplacha.2013.09.006](https://doi.org/10.1016/j.gloplacha.2013.09.006)
- Conti G, Fagarazzi L 2015 Forest expansion in mountain ecosystems: “environmentalist’s dream” or societal nightmare? *Planum* 11, 1–20.
- Conversi A, Fonda Umani S, Peluso T, Molinero JC, Santojanni A et al. 2010 The Mediterranean Sea regime shift at the end of the 1980s, and intriguing parallels with other European basins. *PLoS One* 5, e10633. doi: [10.1371/journal.pone.0010633](https://doi.org/10.1371/journal.pone.0010633)
- Cook CM, Vardaka E, Lanaras T 2004 Toxic cyanobacteria in Greek freshwaters, 1987–2000: Occurrence, toxicity, and impacts in the Mediterranean region. *Acta Hydrochim. Hydrobiol.* 32, 107–124. doi: [10.1002/aheh.200300523](https://doi.org/10.1002/aheh.200300523)
- Cook KH, Vizy EK 2015 Detection and Analysis of an Amplified Warming of the Sahara Desert. *J. Clim.* 28, 6560–6580. doi: [10.1175/jcli-d-14-00230.1](https://doi.org/10.1175/jcli-d-14-00230.1)
- Coppola E, Sobolowski S, Pichelli E, Raffaele F, Ahrens B et al. 2020 A first-of-its-kind multi-model convection permitting ensemble for investigating convective phenomena over Europe and the Mediterranean. *Clim. Dyn.* 55, 3–34. doi: [10.1007/s00382-018-4521-8](https://doi.org/10.1007/s00382-018-4521-8)
- Corrales X, Coll M, Ofir E, Heymans JJ, Steenbeek J et al. 2018 Future scenarios of marine resources and ecosystem conditions in the Eastern Mediterranean under the impacts of fishing, alien species and sea warming. *Sci. Rep.* 8, 14284. doi: [10.1038/s41598-018-32666-x](https://doi.org/10.1038/s41598-018-32666-x)
- Corte-Real JM, Zhang X, Wang X 1995 Large-scale circulation regimes and surface climatic anomalies over the Mediterranean. *Int. J. Climatol.* 15, 1135–1150. doi: [10.1002/joc.3370151006](https://doi.org/10.1002/joc.3370151006)
- Cosentino SL, Porqueddu C, Copani V, Patané C, Testa G et al. 2014 European grasslands overview: Mediterranean region. *Grassl. Sci. Eur.* 19, 41–56.
- Coumou D, Rahmstorf S 2012 A decade of weather extremes. *Nat. Clim. Chang.* 2, 491–496. doi: [10.1038/nclimate1452](https://doi.org/10.1038/nclimate1452)
- Covas L, Senar JC, Roqué L, Quesada J 2017 Records of fatal attacks by Rose-ringed Parakeets *Psittacula krameri* on native avifauna. [http://ornitologia.org/mm/file/2017\\_06.pdf](http://ornitologia.org/mm/file/2017_06.pdf) [Accessed May 14, 2019].
- Cózar A, Sanz-Martín M, Martí E, Ignacio González-Gordillo J, Ubeda B et al. 2015 Plastic accumulation in the Mediterranean Sea. *PLoS One* 10. doi: [10.1371/journal.pone.0121762](https://doi.org/10.1371/journal.pone.0121762)
- Crain CM, Kroeker KJ, Halpern BS 2008 Interactive and cumulative effects of multiple human stressors in marine systems. *Ecol. Lett.* 11, 1304–1315. doi: [10.1111/j.1461-0248.2008.01253.x](https://doi.org/10.1111/j.1461-0248.2008.01253.x)
- Craine JM, Elmore AJ, Olson KC, Tolleson D 2010 Climate change and cattle nutritional stress. *Glob. Chang. Biol.* 16, 2901–2911. doi: [10.1111/j.1365-2486.2009.02060.x](https://doi.org/10.1111/j.1365-2486.2009.02060.x)
- Cramer W, Guiot J, Fader M, Garrabou J, Gattuso J-P et al. 2018 Climate change and interconnected risks to sustainable development in the Mediterranean. *Nat. Clim. Chang.* 8, 972–980. doi: [10.1038/s41558-018-0299-2](https://doi.org/10.1038/s41558-018-0299-2)
- Crooks JA, Chang AL, Ruiz GM 2011 Aquatic pollution increases the relative success of invasive species. *Biol. Invasions* 13, 165–176. doi: [10.1007/s10530-010-9799-3](https://doi.org/10.1007/s10530-010-9799-3)
- Crooks JA, Suarez A 2006 Hyperconnectivity, invasive species, and the breakdown of barriers to dispersal, in *Connectivity Conservation*, eds. Crooks KR, Sanjayan MA (Cambridge, New York, USA: Cambridge University Press), 451–478.
- Cuaresma JC 2017 Income projections for climate change research: A framework based on human capital dynamics. *Glob. Environ. Chang.* 42, 226–236. doi: [10.1016/j.gloenvcha.2015.02.012](https://doi.org/10.1016/j.gloenvcha.2015.02.012)
- Cudennec C, Leduc C, Koutsoyiannis D 2007 Dryland hydrology in Mediterranean regions—a review. *Hydrol. Sci. J.* 52, 1077–1087. doi: [10.1623/hysj.52.6.1077](https://doi.org/10.1623/hysj.52.6.1077)
- Cullen HM, DeMenocal PB 2000 North Atlantic influence on Tigris–Euphrates streamflow. *Int. J. Climatol.* 20, 853–863. doi: [10.1002/1097-0088\(20000630\)20:8<853::AID-JOC497>3.0.CO;2-M](https://doi.org/10.1002/1097-0088(20000630)20:8<853::AID-JOC497>3.0.CO;2-M)
- Cyrys J, Eeftens M, Heinrich J, Ampe C, Armengaud A et al. 2012 Variation of NO<sub>2</sub> and NO<sub>x</sub> concentrations between and within 36 European study areas: Results from the ESCAPE study. *Atmos. Environ.* 62, 374–390. doi: [10.1016/j.atmosenv.2012.07.080](https://doi.org/10.1016/j.atmosenv.2012.07.080)
- D’Agostino R, Lionello P 2020 The atmospheric moisture budget in the Mediterranean: Mechanisms for seasonal changes in the Last Glacial Maximum and future warming scenario. *Quat. Sci. Rev.* 241, 106392. doi: [10.1016/j.quascirev.2020.106392](https://doi.org/10.1016/j.quascirev.2020.106392)
- D’Agostino R, Lionello P, Adam O, Schneider T 2017 Factors controlling Hadley circulation changes from the Last Glacial Maximum to the end of the 21<sup>st</sup> century. *Geophys. Res. Lett.* 44, 8585–8591. doi: [10.1002/2017gl074533](https://doi.org/10.1002/2017gl074533)
- D’Agostino R, Scambiati AL, Jungclaus J, Lionello P 2020 Poleward Shift of Northern Subtropics in Winter: Time of Emergence of Zonal Versus Regional Signals. *Geophys. Res. Lett.* 47, e2020GL089325. doi: [10.1029/2020GL089325](https://doi.org/10.1029/2020GL089325)

- D'Silva MS, Anil AC, Naik RK, D'Costa PM 2012 Algal blooms: a perspective from the coasts of India. *Nat. Hazards* 63, 1225–1253. doi: [10.1007/s11069-012-0190-9](https://doi.org/10.1007/s11069-012-0190-9)
- Dafka S, Toreti A, Zanis P, Xoplaki E, Luterbacher J 2019 Twenty-first-century changes in the Eastern Mediterranean Etesians and associated midlatitude atmospheric circulation. *JGR Atmos.* 124, 12741–12754. doi: [10.1029/2019JD031203](https://doi.org/10.1029/2019JD031203)
- Dahshan H, Megahed AM, Abd-Elall AM, Abd-El-Kader MA-G, Nabawy E et al. 2016 Monitoring of pesticides water pollution-The Egyptian River Nile. *J. Environ. Heal. Sci. Eng.* 14, 15. doi: [10.1186/s40201-016-0259-6](https://doi.org/10.1186/s40201-016-0259-6)
- Dai A 2013 Increasing drought under global warming in observations and models. *Nat. Clim. Chang.* 3, 52–58. doi: [10.1038/nclimate1633](https://doi.org/10.1038/nclimate1633)
- DAISIE 2009 *Handbook of Alien Species in Europe*. Springer Netherlands doi: [10.1007/978-1-4020-8280-1](https://doi.org/10.1007/978-1-4020-8280-1)
- Danovaro R, Fonda Umani S, Pusceddu A, Umani SF, Pusceddu A et al. 2009 Climate Change and the Potential Spreading of Marine Mucilage and Microbial Pathogens in the Mediterranean Sea. *PLoS One* 4, e7006. doi: [10.1371/journal.pone.0007006](https://doi.org/10.1371/journal.pone.0007006)
- Danti R, Della Rocca G 2017 Epidemiological History of Cypress Canker Disease in Source and Invasion Sites. *Forests* 8, 121. doi: [10.3390/f8040121](https://doi.org/10.3390/f8040121)
- Darmaraki S, Somot S, Sevault F, Nabat P 2019a Past Variability of Mediterranean Sea Marine Heatwaves. *Geophys. Res. Lett.* doi: [10.1029/2019gl082933](https://doi.org/10.1029/2019gl082933)
- Darmaraki S, Somot S, Sevault F, Nabat P, Cabos Narvaez WD et al. 2019b Future evolution of Marine Heatwaves in the Mediterranean Sea. *Clim. Dyn.* 53, 1371–1392. doi: [10.1007/s00382-019-04661-z](https://doi.org/10.1007/s00382-019-04661-z)
- Dayan U, Ricaud P, Zbinden R, Dulac F 2017 Atmospheric pollution over the eastern Mediterranean during summer - A review. *Atmos. Chem. Phys.* 17, 13233–13263. doi: [10.5194/acp-17-13233-2017](https://doi.org/10.5194/acp-17-13233-2017)
- de Giglio O, Caggiano G, Bagordo F, Barbuti G, Brigida S et al. 2017 Enteric viruses and fecal bacteria indicators to assess groundwater quality and suitability for irrigation. *Int. J. Environ. Res. Public Health* 14, 558. doi: [10.3390/ijerph14060558](https://doi.org/10.3390/ijerph14060558)
- de Montmollin B, Strahm W 2005 *The Top 50 Mediterranean Island Plants. Wild Plants At the Brink of Extinction, and What is Needed to Save Them*. IUCN/SSC Mediterranean Islands Plant Specialist Group. IUCN, Gland, Switzerland and Cambridge doi: [10.1017/s0030605306270200](https://doi.org/10.1017/s0030605306270200)
- Debolini M, Marraccini E, Dubeuf JP, Geijzendorffer IR, Guerra CA et al. 2018 Land and farming system dynamics and their drivers in the Mediterranean Basin. *Land use policy* 75, 702–710. doi: [10.1016/j.landusepol.2017.07.010](https://doi.org/10.1016/j.landusepol.2017.07.010)
- Deidun A 2018 Back with a bang – an unexpected massive bloom of *Cassiopea andromeda* (Forskaal, 1775) in the Maltese Islands, nine years after its first appearance. *Bio-Invasions Rec.* 7, 399–404. doi: [10.3391/bir.2018.7.4.07](https://doi.org/10.3391/bir.2018.7.4.07)
- Dell'Aquila A, Mariotti A, Bastin S, Calmanti S, Cavicchia L et al. 2018 Evaluation of simulated decadal variations over the Euro-Mediterranean region from ENSEMBLES to Med-CORDEX. *Clim. Dyn.* 51, 857–876. doi: [10.1007/s00382-016-3143-2](https://doi.org/10.1007/s00382-016-3143-2)
- Dellink R, Chateau J, Lanzi E, Magné B 2017 Long-term economic growth projections in the Shared Socioeconomic Pathways. *Glob. Environ. Chang.* 42, 200–214. doi: [10.1016/j.gloenvcha.2015.06.004](https://doi.org/10.1016/j.gloenvcha.2015.06.004)
- Déqué M 2007 Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: Model results and statistical correction according to observed values. *Glob. Planet. Change* 57, 16–26. doi: [10.1016/j.gloplacha.2006.11.030](https://doi.org/10.1016/j.gloplacha.2006.11.030)
- Déqué M, Somot S, Sánchez-Gomez E, Goodess CM, Jacob D et al. 2012 The spread amongst ENSEMBLES regional scenarios: Regional climate models, driving general circulation models and interannual variability. *Clim. Dyn.* 38, 951–964. doi: [10.1007/s00382-011-1053-x](https://doi.org/10.1007/s00382-011-1053-x)
- Deser C, Hurrell JW, Phillips AS 2017 The role of the North Atlantic Oscillation in European climate projections. *Clim. Dyn.* 49, 3141–3157. doi: [10.1007/s00382-016-3502-z](https://doi.org/10.1007/s00382-016-3502-z)
- Desneux N, Wajnberg E, Wyckhuys KAG, Burgio G, Arpaia S et al. 2010 Biological invasion of European tomato crops by *Tuta absoluta*: Ecology, geographic expansion and prospects for biological control. *J. Pest Sci. (2004)*. 83, 197–215. doi: [10.1007/s10340-010-0321-6](https://doi.org/10.1007/s10340-010-0321-6)
- DG/ACTA 1993 Stratégie nationale pour la conservation des eaux et des sols. Publication de la direction de la conservation des eaux et des sols.
- di Castri F 1991 An ecological overview of the five regions of the world with a mediterranean climate, in *Biogeography of Mediterranean Invasions*, eds. Groves RH, di Castri F (Cambridge University Press), 3–16. doi: [10.1017/cbo9780511525544.002](https://doi.org/10.1017/cbo9780511525544.002)
- Diapouli E, Manousakas M, Vratolis S, Vasilatou V, Maggos T et al. 2017 Evolution of air pollution source contributions over one decade, derived by PM10 and PM2.5 source apportionment in two metropolitan urban areas in Greece. *Atmos. Environ.* 164, 416–430. doi: [10.1016/j.atmosenv.2017.06.016](https://doi.org/10.1016/j.atmosenv.2017.06.016)
- Diappi L 2015 City Size and Urbanization in Mediterranean Cities. *Sci. Reg.*, 129–137. doi: [10.3280/SCRE2015-001011](https://doi.org/10.3280/SCRE2015-001011)
- Diffenbaugh NS, Giorgi F 2012 Climate change hotspots in the CMIP5 global climate model ensemble. *Clim. Change* 114, 813–822. doi: [10.1007/s10584-012-0570-x](https://doi.org/10.1007/s10584-012-0570-x)
- Diffenbaugh NS, Pal JS, Giorgi F, Gao X 2007 Heat stress intensification in the Mediterranean climate change hotspot. *Geophys. Res. Lett.* 34, L11706. doi: [10.1029/2007gl030000](https://doi.org/10.1029/2007gl030000)
- Dimarchopoulou D, Stergiou KI, Tsikliras AC 2017 Gap analysis on the biology of Mediterranean marine fishes. *PLoS One* 12, e0175949. doi: [10.1371/journal.pone.0175949](https://doi.org/10.1371/journal.pone.0175949)
- Doblas-Miranda E, Alonso R, Arnan X, Bermejo V, Brotons L et al. 2017 A review of the combination among global change factors in forests, shrublands and pastures of the Mediterranean Region: Beyond drought effects. *Glob. Planet. Change* 148, 42–54. doi: [10.1016/j.gloplacha.2016.11.012](https://doi.org/10.1016/j.gloplacha.2016.11.012)
- Dobrynin M, Murawsky J, Yang S 2012 Evolution of the global wind wave climate in CMIP5 experiments. *Geophys. Res. Lett.* 39. doi: [10.1029/2012gl052843](https://doi.org/10.1029/2012gl052843)
- Domon G 2011 Landscape as resource: Consequences, challenges and opportunities for rural development. *Landscape Urban Plan.* 100, 338–340. doi: [10.1016/j.landurbplan.2011.02.014](https://doi.org/10.1016/j.landurbplan.2011.02.014)

- Donat MG, Peterson TC, Brunet M, King AD, Almazroui M et al. 2014 Changes in extreme temperature and precipitation in the Arab region: long-term trends and variability related to ENSO and NAO. *Int. J. Climatol.* 34, 581–592. doi: [10.1002/joc.3707](https://doi.org/10.1002/joc.3707)
- Donat MG, Renggli D, Wild S, Alexander L V., Leckebusch GC et al. 2011 Reanalysis suggests long-term upward trends in European storminess since 1871. *Geophys. Res. Lett.* 38, n/a-n/a. doi: [10.1029/2011GL047995](https://doi.org/10.1029/2011GL047995)
- Dorn W, Dethloff K, Rinke A, Roeckner E 2003 Competition of NAO regime changes and increasing greenhouse gases and aerosols with respect to Arctic climate projections. *Clim. Dyn.* 21, 447–458. doi: [10.1007/s00382-003-0344-2](https://doi.org/10.1007/s00382-003-0344-2)
- Dosio A, Fischer EM 2018 Will Half a Degree Make a Difference? Robust Projections of Indices of Mean and Extreme Climate in Europe Under 1.5°C, 2°C, and 3°C Global Warming. *Geophys. Res. Lett.* 45, 935–944. doi: [10.1002/2017gl076222](https://doi.org/10.1002/2017gl076222)
- Douville H, Ribes A, Decharme B, Alkama R, Sheffield J 2013 Anthropogenic influence on multidecadal changes in reconstructed global evapotranspiration. *Nat. Clim. Chang.* 3, 59–62. doi: [10.1038/nclimate1632](https://doi.org/10.1038/nclimate1632)
- Drira Z, Kmiha-Megdiche S, Sahnoun H, Hammami A, Allouche N et al. 2016 Assessment of anthropogenic inputs in the surface waters of the southern coastal area of Sfax during spring (Tunisia, Southern Mediterranean Sea). *Mar. Pollut. Bull.* 104, 355–363. doi: [10.1016/j.marpolbul.2016.01.035](https://doi.org/10.1016/j.marpolbul.2016.01.035)
- Dris R, Gasperi J, Mirande C, Mandin C, Guerrouache M et al. 2017 A first overview of textile fibers, including microplastics, in indoor and outdoor environments. *Environ. Pollut.* 221, 453–458. doi: [10.1016/J.ENVPOL.2016.12.013](https://doi.org/10.1016/J.ENVPOL.2016.12.013)
- Dris R, Gasperi J, Saad M, Mirande C, Tassin B 2016 Synthetic fibers in atmospheric fallout: A source of microplastics in the environment? *Mar. Pollut. Bull.* 104, 290–293. doi: [10.1016/J.MARPOLBUL.2016.01.006](https://doi.org/10.1016/J.MARPOLBUL.2016.01.006)
- Dubois A, Lacouture L, Feuillet C 2010 Les pesticides dans les milieux aquatiques : données 2007. *Etudes Doc. du Commissariat Général au Développement Durable* 26, 50. <https://www.statistiques.developpement-durable.gouv.fr/sites/default/files/2019-01/les-pesticides-dans-les-milieux-aquatiques-etudes-et-documents-26-juillet2010.pdf>
- Dubois C, Somot S, Calmanti S, Carillo A, Déqué M et al. 2012 Future projections of the surface heat and water budgets of the Mediterranean Sea in an ensemble of coupled atmosphere–ocean regional climate models. *Clim. Dyn.* 39, 1859–1884. doi: [10.1007/s00382-011-1261-4](https://doi.org/10.1007/s00382-011-1261-4)
- Dubrovský M, Hayes M, Duce P, Trnka M, Svoboda M et al. 2014 Multi-GCM projections of future drought and climate variability indicators for the Mediterranean region. *Reg. Environ. Chang.* 14, 1907–1919. doi: [10.1007/s10113-013-0562-z](https://doi.org/10.1007/s10113-013-0562-z)
- Dullinger I, Wessely J, Bossdorf O, Dawson W, Essl F et al. 2017 Climate change will increase the naturalization risk from garden plants in Europe. *Glob. Ecol. Biogeogr.* 26, 43–53. doi: [10.1111/geb.12512](https://doi.org/10.1111/geb.12512)
- Dunić N, Vilibić I, Šepić J, Mihanović H, Sevault F et al. 2019 Performance of multi-decadal ocean simulations in the Adriatic Sea. *Ocean Model.* 134, 81–109. doi: [10.1016/j.ocemod.2019.01.006](https://doi.org/10.1016/j.ocemod.2019.01.006)
- Dünkeloh A, Jacobeit J 2003 Circulation dynamics of Mediterranean precipitation variability 1948–98. *Int. J. Climatol.* 23, 1843–1866. doi: [10.1002/joc.973](https://doi.org/10.1002/joc.973)
- Dupire S, Curt T, Bigot S 2017 Spatio-temporal trends in fire weather in the French Alps. *Sci. Total Environ.* 595, 801–817. doi: [10.1016/j.scitotenv.2017.04.027](https://doi.org/10.1016/j.scitotenv.2017.04.027)
- Early R, González-Moreno P, Murphy ST, Day R 2018 Forecasting the global extent of invasion of the cereal pest *Spodoptera frugiperda*, the fall armyworm. *bioRxiv*, 391847. doi: [10.3897/neobiota.40.28165.suppl2](https://doi.org/10.3897/neobiota.40.28165.suppl2)
- EC/DG for Research and Innovation 2011 *EuroMed-2030: Long term challenges for the Mediterranean area; report of an expert group*. Luxembourg: Publ. Office of the European Union.
- Edelist D, Rilov G, Golani D, Carlton JT, Spanier E 2013 Restructuring the Sea: profound shifts in the world's most invaded marine ecosystem. *Divers. Distrib.* 19, 69–77. doi: [10.1111/ddi.12002](https://doi.org/10.1111/ddi.12002)
- EEA 1999 Environmental Indicators: Typology and Overview, Technical Report No. 25. Copenhagen, Denmark <https://www.eea.europa.eu/publications/TEC25>
- Efthymiadis D, Goodess CM, Jones PD 2011 Trends in Mediterranean gridded temperature extremes and large-scale circulation influences. *Nat. Hazards Earth Syst. Sci.* 11, 2199–2214. doi: [10.5194/nhess-11-2199-2011](https://doi.org/10.5194/nhess-11-2199-2011)
- El Ayni F, Manoli E, Cherif S, Jrad A, Assimacopoulos D et al. 2013 Deterioration of a Tunisian coastal aquifer due to agricultural activities and possible approaches for better water management. *Water Environ. J.* 27, 348–361. doi: [10.1111/j.1747-6593.2012.00354.x](https://doi.org/10.1111/j.1747-6593.2012.00354.x)
- El Bakouri H, Ouassini A, Morillo J, Usero J 2008 Pesticides in ground water beneath Loukkos perimeter, Northwest Morocco. *J. Hydrol.* 348, 270–278. doi: [10.1016/j.jhydrol.2007.10.002](https://doi.org/10.1016/j.jhydrol.2007.10.002)
- El Kenawy AM, López Moreno JI, Vicente-Serrano SM 2011 Recent trends in daily temperature extremes over north-eastern Spain (1960–2006). *Nat. Hazards Earth Syst. Sci.* 11, 2583–2603. doi: [10.5194/nhess-11-2583-2011](https://doi.org/10.5194/nhess-11-2583-2011)
- El Kenawy AM, López Moreno JI, Vicente-Serrano SM, Mekld MS 2009 Temperature trends in Libya over the second half of the 20<sup>th</sup> century. *Theor. Appl. Climatol.* 98, 1–8. doi: [10.1007/s00704-008-0089-2](https://doi.org/10.1007/s00704-008-0089-2)
- El Nemr A, Abd-Allah AMA 2003 Contamination of polycyclic aromatic hydrocarbons (PAHs) in microlayer and subsurface waters along Alexandria coast, Egypt. *Chemosphere* 52, 1711–1716. doi: [10.1016/S0045-6535\(03\)00300-X](https://doi.org/10.1016/S0045-6535(03)00300-X)
- Elbakidze M, Hahn T, Zimmermann NE, Cudlin P, Friberg N et al. 2018 Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for Europe and Central Asia*, eds. Rounsevell M, Fischer M, Torre-Marín Rando A, Mader A (Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), 385–568.
- Elguindi N, Somot S, Déqué M, Ludwig W 2011 Climate change evolution of the hydrological balance of the Mediterranean, Black and Caspian Seas: impact of climate model resolution. *Clim. Dyn.* 36, 205–228. doi: [10.1007/s00382-009-0715-4](https://doi.org/10.1007/s00382-009-0715-4)

- Elton CS 1958 *The Ecology of Invasions by Animals and Plants*. Chapman and Hall, New York, US.  
doi: [10.1007/978-1-4899-7214-9](https://doi.org/10.1007/978-1-4899-7214-9)
- Emanuel K 2005 Genesis and maintenance of "Mediterranean hurricanes." *Adv. Geosci.* 2, 217–220.  
doi: [10.5194/adgeo-2-217-2005](https://doi.org/10.5194/adgeo-2-217-2005)
- Emara HI, Said TO, El Naggar NA, Shreadah MA 2008 Aliphatic and polycyclic hydrocarbon compounds as chemical markers for pollution sources in relation to physico-chemical characteristics of the eastern harbour (Egyptian Mediterranean Sea). *Egypt. J. Aquat. Res.* 34, 1–19.
- Engelbrecht JP, Jayanty RKM 2013 Assessing sources of airborne mineral dust and other aerosols, in Iraq. *Aeolian Res.* 9, 153–160. doi: [10.1016/J.AEOLIA.2013.02.003](https://doi.org/10.1016/J.AEOLIA.2013.02.003)
- Enriquez-Alonso A, Sánchez-Lorenzo A, Calbó J, González J-A, Norris JR 2016 Cloud cover climatologies in the Mediterranean obtained from satellites, surface observations, reanalyses, and CMIP5 simulations: validation and future scenarios. *Clim. Dyn.* 47, 249–269. doi: [10.1007/s00382-015-2834-4](https://doi.org/10.1007/s00382-015-2834-4)
- EPP0 2012a Decision-support scheme for an Express Pest Risk Analysis. *EPP0 Bull.* 42, 457–462. doi: [10.1111/epp.2591](https://doi.org/10.1111/epp.2591)
- EPP0 2012b EPP0 prioritization process for invasive alien plants. *EPP0 Bull.* 42, 463–474. doi: [10.1111/epp.2592](https://doi.org/10.1111/epp.2592)
- Eschen R, Grégoire J-C, Hengeveld GM, de Hoop BM, Rigaux L et al. 2015 Trade patterns of the tree nursery industry in Europe and changes following findings of citrus longhorn beetle, *Anoplophora chinensis* Forster. *NeoBiota* 26, 1–20. doi: [10.3897/neobiota.26.8947](https://doi.org/10.3897/neobiota.26.8947)
- Escudero M, Querol X, Ávila A, Cuevas E 2007 Origin of the exceedances of the European daily PM limit value in regional background areas of Spain. *Atmos. Environ.* 41, 730–744. doi: [10.1016/J.ATMOENV.2006.09.014](https://doi.org/10.1016/J.ATMOENV.2006.09.014)
- Essid N, Boufahja F, Beyrem H, Aïssa P, Mahmoudi E 2013 Effects of 17- $\alpha$ -estradiol on a free-living marine nematode community: A microcosm experiment. *African J. Aquat. Sci.* 38, 305–311. doi: [10.2989/16085914.2013.832652](https://doi.org/10.2989/16085914.2013.832652)
- Essl F, Kobler J 2009 Spiny invaders – Patterns and determinants of cacti invasion in Europe. *Flora - Morphol. Distrib. Funct. Ecol. Plants* 204, 485–494. doi: [10.1016/j.flora.2008.06.002](https://doi.org/10.1016/j.flora.2008.06.002)
- Etteieb S, Cherif S, Kawachi A, Han J, Elayni F et al. 2016 Combining Biological and Chemical Screenings to Assess Cytotoxicity of Emerging Contaminants in Discharges into Surface Water. *Water, Air, Soil Pollut.* 227, 341. doi: [10.1007/s11270-016-3049-y](https://doi.org/10.1007/s11270-016-3049-y)
- European Food Safety Authority 2011 The 2009 European Union Report on Pesticide Residues in Food. *EFSA J.* 9, 2430. doi: [10.2903/j.efsa.2011.2430](https://doi.org/10.2903/j.efsa.2011.2430)
- European Union 2014 Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species. *Off. J. Eur. Union* 57. doi: [10.5040/9781509909568.0031](https://doi.org/10.5040/9781509909568.0031)
- Evan AT, Flamant C, Lavaysse C, Kocha C, Saci A 2015 Water Vapor–Forced Greenhouse Warming over the Sahara Desert and the Recent Recovery from the Sahelian Drought. *J. Clim.* 28, 108–123. doi: [10.1175/jcli-d-14-00039.1](https://doi.org/10.1175/jcli-d-14-00039.1)
- Evangelidou N, Balkanski Y, Cozic A, Hao WM, Möller AP 2014 Wildfires in Chernobyl-contaminated forests and risks to the population and the environment: A new nuclear disaster about to happen? *Environ. Int.* 73, 346–358. doi: [10.1016/J.ENVINT.2014.08.012](https://doi.org/10.1016/J.ENVINT.2014.08.012)
- Ezber Y 2018 Assessment of the changes in the Etesians in the EURO-CORDEX regional model projections. *Int. J. Climatol.* 39, 1213–1229. doi: [10.1002/joc.5872](https://doi.org/10.1002/joc.5872)
- Fabbio G, Merlo M, Tosi V 2003 Silvicultural management in maintaining biodiversity and resistance of forests in Europe—the Mediterranean region. *J. Environ. Manage.* 67, 67–76. doi: [10.1016/s0301-4797\(02\)00189-5](https://doi.org/10.1016/s0301-4797(02)00189-5)
- Faccoli M, Campo G, Perrotta G, Rassati D 2016 Two newly introduced tropical bark and ambrosia beetles (Coleoptera: Curculionidae, Scolytinae) damaging figs (*Ficus carica*) in southern Italy. *Zootaxa* 4138, 189. doi: [10.11646/zootaxa.4138.1.10](https://doi.org/10.11646/zootaxa.4138.1.10)
- Fan J, Denux O, Courtin C, Bernard A, Javal M et al. 2019 Multi-component blends for trapping native and exotic longhorn beetles at potential points-of-entry and in forests. *J. Pest Sci. (2004)*. 92, 281–297. doi: [10.1007/s10340-018-0997-6](https://doi.org/10.1007/s10340-018-0997-6)
- Fantini A, Raffaele F, Torma CZ, Bacer S, Coppola E et al. 2018 Assessment of multiple daily precipitation statistics in ERA-Interim driven Med-CORDEX and EURO-CORDEX experiments against high resolution observations. *Clim. Dyn.* 51, 877–900. doi: [10.1007/s00382-016-3453-4](https://doi.org/10.1007/s00382-016-3453-4)
- FAO 2015 Towards a Regional Collaborative Strategy on Sustainable Water Management and Food Security in the Near East and North Africa Region. Cairo, Egypt.
- FAO 2017a Integrated Monitoring Guide for SDG 6. Step-by-step monitoring methodology for indicator 6.4.1 on water-use efficiency.
- FAO 2017b International standards for phytosanitary measures 5. Glossary of phytosanitary terms.
- Faurès J-M, Hoogeveen J, Bruinsma J 2002 The FAO Irrigated Area Forecast for 2030. <http://www.fao.org/tempref/agl/AGLW/docs/fauresetalagadir.pdf>
- Feidas H, Makrogiannis T, Bora-Senta E 2004 Trend analysis of air temperature time series in Greece and their relationship with circulation using surface and satellite data: 1955?2001. *Theor. Appl. Climatol.* 79, 185–208. doi: [10.1007/s00704-004-0064-5](https://doi.org/10.1007/s00704-004-0064-5)
- Feki W, Hamza A, Frossard V, Abdennadher M, Hannachi I et al. 2013 What are the potential drivers of blooms of the toxic dinoflagellate *Karenia selliformis*? A 10-year study in the Gulf of Gabes, Tunisia, southwestern Mediterranean Sea. *Harmful Algae* 23, 8–18. doi: [10.1016/j.hal.2012.12.001](https://doi.org/10.1016/j.hal.2012.12.001)
- Feliks Y, Ghil M, Robertson AW 2010 Oscillatory Climate Modes in the Eastern Mediterranean and Their Synchronization with the North Atlantic Oscillation. *J. Clim.* 23, 4060–4079. doi: [10.1175/2010jcli3181.1](https://doi.org/10.1175/2010jcli3181.1)
- Felis T, Rimbu N 2010 Mediterranean climate variability documented in oxygen isotope records from northern Red Sea corals—A review. *Glob. Planet. Change* 71, 232–241. doi: [10.1016/j.gloplacha.2009.10.006](https://doi.org/10.1016/j.gloplacha.2009.10.006)
- Fenoglio-Marc L, Mariotti A, Sannino G, Meyssignac B, Carillo A et al. 2013 Decadal variability of net water flux at the Med-

- iterranean Sea Gibraltar Strait. *Glob. Planet. Change* 100, 1–10. doi: [10.1016/J.GLOPLACHA.2012.08.007](https://doi.org/10.1016/J.GLOPLACHA.2012.08.007)
- Fernandes PG, Ralph GM, Nieto A, Criado MG, Vasilakopoulos P et al. 2017 Coherent assessments of Europe's marine fishes show regional divergence and megafauna loss. *Nat. Ecol. Evol.* 1, 0170. doi: [10.1038/s41559-017-0170](https://doi.org/10.1038/s41559-017-0170)
- Fernández J, Frías MD, Cabos WD, Cofiño AS, Domínguez M et al. 2019 Consistency of climate change projections from multiple global and regional model intercomparison projects. *Clim. Dyn.* 52, 1139–1156. doi: [10.1007/s00382-018-4181-8](https://doi.org/10.1007/s00382-018-4181-8)
- Filahi S, Tanarhte M, Mouhir L, El Morhit M, Trambly Y 2015 Trends in indices of daily temperature and precipitations extremes in Morocco. *Theor. Appl. Climatol.* 124, 959–972. doi: [10.1007/s00704-015-1472-4](https://doi.org/10.1007/s00704-015-1472-4)
- Fischer-Bruns I, Banse DF, Feichter J 2009 Future impact of anthropogenic sulfate aerosol on North Atlantic climate. *Clim. Dyn.* 32, 511–524. doi: [10.1007/s00382-008-0458-7](https://doi.org/10.1007/s00382-008-0458-7)
- Fischer EM, Schär C 2010 Consistent geographical patterns of changes in high-impact European heatwaves. *Nat. Geosci.* 3, 398–403. doi: [10.1038/ngeo866](https://doi.org/10.1038/ngeo866)
- Flaounas E, Drobinski P, Borga M, Calvet J-C, Delrieu G et al. 2012 Assessment of gridded observations used for climate model validation in the Mediterranean region: the HyMeX and MED-CORDEX framework. *Environ. Res. Lett.* 7, 024017. doi: [10.1088/1748-9326/7/2/024017](https://doi.org/10.1088/1748-9326/7/2/024017)
- Flocas HA, Simmonds I, Kouroutzoglou J, Keay K, Hatzaki M et al. 2010 On Cyclonic Tracks over the Eastern Mediterranean. *J. Clim.* 23, 5243–5257. doi: [10.1175/2010JCLI3426.1](https://doi.org/10.1175/2010JCLI3426.1)
- Floris I, Cocco A, Buffa F, Mannu R, Satta A 2018 Insect pests of *Eucalyptus* plantations in Sardinia (Italy). *Redia*, 61–71. doi: [10.19263/redia-101.18.09](https://doi.org/10.19263/redia-101.18.09)
- Floutsi AA, Korras-Carraca MB, Matsoukas C, Hatzianastassiou N, Biskos G 2016 Climatology and trends of aerosol optical depth over the Mediterranean basin during the last 12 years (2002–2014) based on Collection 006 MODIS-Aqua data. *Sci. Total Environ.* 551–552, 292–303. doi: [10.1016/J.SCITOTENV.2016.01.192](https://doi.org/10.1016/J.SCITOTENV.2016.01.192)
- Foley JA, DeFries RS, Asner GP, Barford C, Bonan G et al. 2005 Global consequences of land use. *Science* (80-. ). 309, 570–574. doi: [10.1126/science.1111772](https://doi.org/10.1126/science.1111772)
- Folini D, Wild M 2011 Aerosol emissions and dimming/brightening in Europe: Sensitivity studies with ECHAM5-HAM. *JGR Atmos.* 116. doi: [10.1029/2011JD016227](https://doi.org/10.1029/2011JD016227)
- Folt CL, Chen CY, Moore M V., Burnaford J 1999 Synergism and antagonism among multiple stressors. *Limnol. Oceanogr.* 44, 864–877. doi: [10.4319/lo.1999.44.3\\_part\\_2.0864](https://doi.org/10.4319/lo.1999.44.3_part_2.0864)
- Fosser G, Khodayar S, Berg P 2015 Benefit of convection permitting climate model simulations in the representation of convective precipitation. *Clim. Dyn.* 44, 45–60. doi: [10.1007/s00382-014-2242-1](https://doi.org/10.1007/s00382-014-2242-1)
- Fourati R, Tedetti M, Guigue C, Goutx M, García N et al. 2018a Sources and spatial distribution of dissolved aliphatic and polycyclic aromatic hydrocarbons in surface coastal waters of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea). *Prog. Oceanogr.* 163, 232–247. doi: [10.1016/j.pocean.2017.02.001](https://doi.org/10.1016/j.pocean.2017.02.001)
- Fourati R, Tedetti M, Guigue C, Goutx M, Zaghden H et al. 2018b Natural and anthropogenic particulate-bound aliphatic and polycyclic aromatic hydrocarbons in surface waters of the Gulf of Gabès (Tunisia, southern Mediterranean Sea). *Environ. Sci. Pollut. Res.* 25, 2476–2494. doi: [10.1007/s11356-017-0641-7](https://doi.org/10.1007/s11356-017-0641-7)
- Francardi V, Noal A, Francescato S, Pinto R, Bruni A et al. 2017 Coexistence of *Xylosandrus crassiusculus* (Motschulsky) and *X. compactus* (Eichhoff) (Coleoptera Curculionidae Scolytinae) in the National Park of Circeo (Lazio, Italy). *Redia* 100, 149–155.
- Francour P, Boudouresque C-F, Harmelin JG, Harmelin-Vivien ML, Quignard JP 1994 Are the Mediterranean waters becoming warmer? Information from biological indicators. *Mar. Pollut. Bull.* 28, 523–526. doi: [10.1016/0025-326x\(94\)90071-x](https://doi.org/10.1016/0025-326x(94)90071-x)
- Fréjaville T, Curt T 2017 Seasonal changes in the human alteration of fire regimes beyond the climate forcing. *Environ. Res. Lett.* 12, 035006. doi: [10.1088/1748-9326/aa5d23](https://doi.org/10.1088/1748-9326/aa5d23)
- Froese R, Kesner-Reyes K 2002 Impact of Fishing on the Abundance of Marine Species. 12.
- Froese R, Winker H, Coro G, Demirel N, Tsikliras AC et al. 2018 Status and rebuilding of European fisheries. *Mar. Policy* 93, 159–170. doi: [10.1016/j.marpol.2018.04.018](https://doi.org/10.1016/j.marpol.2018.04.018)
- Frölicher TL, Fischer EM, Gruber N 2018 Marine heatwaves under global warming. *Nature* 560, 360–364. doi: [10.1038/s41586-018-0383-9](https://doi.org/10.1038/s41586-018-0383-9)
- Fumière Q, Déqué M, Nuissier O, Somot S, Alias A et al. 2019 Extreme rainfall in Mediterranean France during the fall: added value of the CNRM-AROME Convection-Permitting Regional Climate Model. *Clim. Dyn.*, 1–15. doi: [10.1007/s00382-019-04898-8](https://doi.org/10.1007/s00382-019-04898-8)
- Fyfe JC, Boer GJ, Flato GM 1999 The Arctic and Antarctic Oscillations and their projected changes under global warming. *Geophys. Res. Lett.* 26, 1601–1604. doi: [10.1029/1999GL900317](https://doi.org/10.1029/1999GL900317)
- Gačić M, Borzelli GLE, Civitarese G, Cardin V, Yari S 2010 Can internal processes sustain reversals of the ocean upper circulation? The Ionian Sea example. *Geophys. Res. Lett.* 37, n/a-n/a. doi: [10.1029/2010GL043216](https://doi.org/10.1029/2010GL043216)
- Gaertner MÁ, González-Alemán JJ, Romera R, Domínguez M, Gil V et al. 2018 Simulation of medicanes over the Mediterranean Sea in a regional climate model ensemble: impact of ocean-atmosphere coupling and increased resolution. *Clim. Dyn.* 51, 1041–1057. doi: [10.1007/s00382-016-3456-1](https://doi.org/10.1007/s00382-016-3456-1)
- Gaertner MÁ, Jacob D, Gil V, Domínguez M, Padorno E et al. 2007 Tropical cyclones over the Mediterranean Sea in climate change simulations. *Geophys. Res. Lett.* 34, L14711. doi: [10.1029/2007gl029977](https://doi.org/10.1029/2007gl029977)
- Galdies C 2012 Temperature trends in Malta (central Mediterranean) from 1951 to 2010. *Meteorol. Atmos. Phys.* 117, 135–143. doi: [10.1007/s00703-012-0187-7](https://doi.org/10.1007/s00703-012-0187-7)
- Galil BS 2007 Loss or gain? Invasive aliens and biodiversity in the Mediterranean Sea. *Mar. Pollut. Bull.* 55, 314–322. doi: [10.1016/j.marpolbul.2006.11.008](https://doi.org/10.1016/j.marpolbul.2006.11.008)
- Galindo N, Yubero E, Nicolás JF, Varea M, Crespo J 2018 Characterization of metals in PM1 and PM10 and health risk evaluation at an urban site in the western Mediterranean. *Chemosphere* 201, 243–250. doi: [10.1016/j.chemosphere.2018.02.162](https://doi.org/10.1016/j.chemosphere.2018.02.162)

- Gallardo B, Aldridge DC, González-Moreno P, Pergl J, Pizarro M et al. 2017 Protected areas offer refuge from invasive species spreading under climate change. *Glob. Chang. Biol.* 23, 5331–5343. doi: [10.1111/gcb.13798](https://doi.org/10.1111/gcb.13798)
- Gallart F, Delgado JM, Beatson SJ V., Posner H, Llorens P et al. 2011 Analysing the effect of global change on the historical trends of water resources in the headwaters of the Llobregat and Ter river basins (Catalonia, Spain). *Phys. Chem. Earth, Parts A/B/C* 36, 655–661. doi: [10.1016/j.pce.2011.04.009](https://doi.org/10.1016/j.pce.2011.04.009)
- Galloway TS 2015 Micro- and nano-plastics and human health, in *Marine Anthropogenic Litter* (Cham: Springer International Publishing), 343–366. doi: [10.1007/978-3-319-16510-3\\_13](https://doi.org/10.1007/978-3-319-16510-3_13)
- Ganor E, Osetinsky I, Stupp A, Alpert P 2010 Increasing trend of African dust, over 49 years, in the eastern Mediterranean. *JGR Atmos.* 115, D07201. doi: [10.1029/2009JD012500](https://doi.org/10.1029/2009JD012500)
- Gao J 2017 Downscaling Global Spatial Population Projections from 1/8-degree to 1-km Grid Cells. doi: [10.5065/d60z721h](https://doi.org/10.5065/d60z721h)
- Gao X, Pal JS, Giorgi F 2006 Projected changes in mean and extreme precipitation over the Mediterranean region from a high resolution double nested RCM simulation. *Geophys. Res. Lett.* 33, 2–5. doi: [10.1029/2005GL024954](https://doi.org/10.1029/2005GL024954)
- Garcés E, Camp J 2012 Habitat changes in the Mediterranean Sea and the consequences for Harmful Algal Blooms formation, in *Life in the Mediterranean Sea: A Look at Habitat Changes*, ed. Stambler N (Nova Science Publishers, Inc.), 519–541.
- García-Ayllón S 2018 Predictive Diagnosis of Agricultural Peri-urban Areas Based on Territorial Indicators: Comparative Landscape Trends of the So-Called “Orchard of Europe.” *Sustainability* 10, 1820. doi: [10.3390/su10061820](https://doi.org/10.3390/su10061820)
- García-Gómez H, Izquieta-Rojano S, Aguilauame L, González-Fernández I, Valiño F et al. 2018 Joining empirical and modelling approaches to estimate dry deposition of nitrogen in Mediterranean forests. *Environ. Pollut.* 243, 427–436. doi: [10.1016/j.envpol.2018.09.015](https://doi.org/10.1016/j.envpol.2018.09.015)
- García-Serrano H, Sans FX, Escarré J 2007 Interspecific competition between alien and native congeneric species. *Acta Oecologica* 31, 69–78. doi: [10.1016/J.ACTAO.2006.09.005](https://doi.org/10.1016/J.ACTAO.2006.09.005)
- Garnas JR, Auger-Rozenberg M-A, Roques A, Bertelsmeier C, Wingfield MJ et al. 2016 Complex patterns of global spread in invasive insects: eco-evolutionary and management consequences. *Biol. Invasions* 18, 935–952. doi: [10.1007/s10530-016-1082-9](https://doi.org/10.1007/s10530-016-1082-9)
- Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P et al. 2009 Mass mortality in Northwestern Mediterranean rocky benthic communities: Effects of the 2003 heat wave. *Glob. Chang. Biol.* 15, 1090–1103. doi: [10.1111/j.1365-2486.2008.01823.x](https://doi.org/10.1111/j.1365-2486.2008.01823.x)
- Garrido A, Martínez-Santos P, Llamas MR 2006 Groundwater irrigation and its implications for water policy in semiarid countries: the Spanish experience. *Hydrogeol. J.* 14, 340–349. doi: [10.1007/s10040-005-0006-z](https://doi.org/10.1007/s10040-005-0006-z)
- Garrido M, Cecchi P, Collos Y, Agostini S, Pasqualini V 2016 Water flux management and phytoplankton communities in a Mediterranean coastal lagoon. Part I: How to promote dinoflagellate dominance? *Mar. Pollut. Bull.* 104, 139–152. doi: [10.1016/j.marpolbul.2016.01.049](https://doi.org/10.1016/j.marpolbul.2016.01.049)
- Gasperi J, Wright SL, Dris R, Collard F, Mandin C et al. 2018 Microplastics in air: Are we breathing it in? *Curr. Opin. Environ. Sci. Heal.* 1, 1–5. doi: [10.1016/J.COESH.2017.10.002](https://doi.org/10.1016/J.COESH.2017.10.002)
- Gassó N, Pino J, Font X, Vilà M 2012 Regional context affects native and alien plant species richness across habitat types. *Appl. Veg. Sci.* 15, 4–13. doi: [10.1111/j.1654-109x.2011.01159.x](https://doi.org/10.1111/j.1654-109x.2011.01159.x)
- Gattuso J-P, Magnan A, Billé R, Cheung WWL, Howes EL et al. 2015 Contrasting futures for ocean and society from different anthropogenic CO<sub>2</sub> emissions scenarios. *Science (80-. ).* 349. doi: [10.1126/science.aac4722](https://doi.org/10.1126/science.aac4722)
- Gea-Izquierdo G, Nicault A, Battipaglia G, Dorado Liñán I, Gutiérrez E et al. 2017 Risky future for Mediterranean forests unless they undergo extreme carbon fertilization. *Glob. Chang. Biol.*, 1–13. doi: [10.1111/gcb.13597](https://doi.org/10.1111/gcb.13597)
- Geng Q, Sugi M 2003 Possible Change of Extratropical Cyclone Activity due to Enhanced Greenhouse Gases and Sulfate Aerosols—Study with a High-Resolution AGCM. *J. Clim.* 16, 2262–2274. doi: [10.1175/1520-0442\(2003\)16<2262:pcoeca>2.0.co;2](https://doi.org/10.1175/1520-0442(2003)16<2262:pcoeca>2.0.co;2)
- Genovesi P, Bacher S, Kobelt M, Pascal M, Scalera R 2009 Alien mammals of Europe, in *Handbook of alien species in Europe*, ed. DAISIE (Springer Netherlands), 119–128. doi: [10.1007/978-1-4020-8280-1\\_9](https://doi.org/10.1007/978-1-4020-8280-1_9)
- Gerard F, Petit S, Smith G, Thomson A, Brown N et al. 2010 Land cover change in Europe between 1950 and 2000 determined employing aerial photography. *Prog. Phys. Geogr. Earth Environ.* 34, 183–205. doi: [10.1177/0309133309360141](https://doi.org/10.1177/0309133309360141)
- Gerasopoulos E, Kouvarakis G, Babasakalis P, Vrekoussis M, Putaud JP et al. 2006 Origin and variability of particulate matter (PM<sub>10</sub>) mass concentrations over the Eastern Mediterranean. *Atmos. Environ.* 40, 4679–4690. doi: [10.1016/j.atmosenv.2006.04.020](https://doi.org/10.1016/j.atmosenv.2006.04.020)
- Geyer R, Jambeck JR, Law KL 2017 Production, use, and fate of all plastics ever made. *Sci. Adv.* 3, e1700782. doi: [10.1126/sciadv.1700782](https://doi.org/10.1126/sciadv.1700782)
- Ghelardini L, Luchi N, Pecori F, Pepori AL, Danti R et al. 2017 Ecology of invasive forest pathogens. *Biol. Invasions* 19, 3183–3200. doi: [10.1007/s10530-017-1487-0](https://doi.org/10.1007/s10530-017-1487-0)
- Giakoumi S, Pey A, Huseyinoglu MF 2019 Assessing the state of invasive fishes in two Mediterranean marine protected areas and adjacent unprotected areas. in *1st Mediterranean Symposium on Non-Indigenous Species, Antalya, Turkey* (Frontiers Media SA), 53–58. doi: [10.3389/fmars.2017.00049](https://doi.org/10.3389/fmars.2017.00049)
- Giannakopoulou E-M, Toumi R 2012 Impacts of the Nile Delta land-use on the local climate. *Atmos. Sci. Lett.* 13, 208–215. doi: [10.1002/asl.381](https://doi.org/10.1002/asl.381)
- Gil-Tena A, Brotons L, Saura S 2010 Effects of forest landscape change and management on the range expansion of forest bird species in the Mediterranean region. *For. Ecol. Manage.* 259, 1338–1346. doi: [10.1016/j.foreco.2009.10.026](https://doi.org/10.1016/j.foreco.2009.10.026)
- Gillett NP, Fyfe JC, Parker DE 2013 Attribution of observed sea level pressure trends to greenhouse gas, aerosol, and ozone changes. *Geophys. Res. Lett.* 40, 2302–2306. doi: [10.1002/grl.50500](https://doi.org/10.1002/grl.50500)
- Gillett NP, Graf HF, Osborn TJ 2003 Climate change and the North Atlantic oscillation. *Geophys. Monogr. Ser.* 134, 193–209. doi: [10.1029/134GM09](https://doi.org/10.1029/134GM09)

- Giménez Papiol G, Casanova A, Fernández-Tejedor M, de la Iglesia P, Diogène J 2013 Management of domoic acid monitoring in shellfish from the Catalan coast. *Environ. Monit. Assess.* 185, 6653–6666. doi: [10.1007/s10661-012-3054-6](https://doi.org/10.1007/s10661-012-3054-6)
- Gimeno I, Vilà M, Hulme PE 2006 Are islands more susceptible to plant invasion than continents? A test using *Oxalis pes-caprae* L. in the western Mediterranean. *J. Biogeogr.* 33, 1559–1565. doi: [10.1111/j.1365-2699.2006.01525.x](https://doi.org/10.1111/j.1365-2699.2006.01525.x)
- Giorgi F 2006 Climate change hotspots. *Geophys. Res. Lett.* 33, L08707. doi: [10.1029/2006GL025734](https://doi.org/10.1029/2006GL025734)
- Giorgi F, Bi X 2005 Updated regional precipitation and temperature changes for the 21<sup>st</sup> century from ensembles of recent AOGCM simulations. *Geophys. Res. Lett.* 32, L21715. doi: [10.1029/2005GL024288](https://doi.org/10.1029/2005GL024288)
- Giorgi F, Coppola E 2007 European climate-change oscillation (ECO). *Geophys. Res. Lett.* 34, L21703. doi: [10.1029/2007g1031223](https://doi.org/10.1029/2007g1031223)
- Giorgi F, Coppola E 2009 Projections of twenty-first century climate over Europe. *EPJ Web Conf.* 1, 29–46. doi: [10.1140/epjconf/e2009-00908-9](https://doi.org/10.1140/epjconf/e2009-00908-9)
- Giorgi F, Coppola E, Raffaele F 2014 A consistent picture of the hydroclimatic response to global warming from multiple indices: Models and observations. *J. Geophys. Res. Atmos.* 119, 11,611–695,708. doi: [10.1002/2014JD022238](https://doi.org/10.1002/2014JD022238)
- Giorgi F, Im E-S, Coppola E, Diffenbaugh NS, Gao XJ et al. 2011 Higher Hydroclimatic Intensity with Global Warming. *J. Clim.* 24, 5309–5324. doi: [10.1175/2011JCLI3979.1](https://doi.org/10.1175/2011JCLI3979.1)
- Giorgi F, Jones C, Asrar GR 2009 Addressing climate information needs at the regional level: The CORDEX framework. *WMO Bull.* 58, 175.
- Giorgi F, Lionello P 2008 Climate change projections for the Mediterranean region. *Glob. Planet. Change* 63, 90–104. doi: [10.1016/j.gloplacha.2007.09.005](https://doi.org/10.1016/j.gloplacha.2007.09.005)
- Giorgi F, Raffaele F, Coppola E 2019 The response of precipitation characteristics to global warming from climate projections. *Earth Syst. Dynam.* 10, 73–89. doi: [10.5194/esd-10-73-2019](https://doi.org/10.5194/esd-10-73-2019)
- Giorgi F, Torma CZ, Coppola E, Ban N, Schär C et al. 2016 Enhanced summer convective rainfall at Alpine high elevations in response to climate warming. *Nat. Geosci.* 9, 584–589. doi: [10.1038/ngeo2761](https://doi.org/10.1038/ngeo2761)
- Gitay H, Brown S, Easterling W, Jallow B, Antle J et al. 2001 Ecosystems and their goods and services, in *Climate Change 2001: Impacts, Adaptation, and Vulnerability*, eds. McCarthy JJ, Canziani OF, Leary NA, Dokken DJ, White KS (Cambridge, UK and New York, NY, USA), 235–342.
- Givan O, Edelist D, Sonin O, Belmaker J 2017a Thermal affinity as the dominant factor changing Mediterranean fish abundances. *Glob. Chang. Biol.* 24, e80–e89. doi: [10.1111/gcb.13835](https://doi.org/10.1111/gcb.13835)
- Givan O, Parravicini V, Kulbicki M, Belmaker J 2017b Trait structure reveals the processes underlying fish establishment in the Mediterranean. *Glob. Ecol. Biogeogr.* 26, 142–153. doi: [10.1111/geb.12523](https://doi.org/10.1111/geb.12523)
- Godinho S, Gil A, Guiomar N, Costa MJ, Neves N 2016 Assessing the role of Mediterranean evergreen oaks canopy cover in land surface albedo and temperature using a remote sensing-based approach. *Appl. Geogr.* 74, 84–94. doi: [10.1016/j.apgeog.2016.07.004](https://doi.org/10.1016/j.apgeog.2016.07.004)
- Gogou A, Bouloubassi I, Stephanou EG 2000 Marine organic geochemistry of the Eastern Mediterranean: 1. Aliphatic and polyaromatic hydrocarbons in Cretan Sea surficial sediments. *Mar. Chem.* 68, 265–282. doi: [10.1016/S0304-4203\(99\)00082-1](https://doi.org/10.1016/S0304-4203(99)00082-1)
- Gómez-Gutiérrez AI, Jover E, Bodineau L, Albaigés J, Bayona JM 2006 Organic contaminant loads into the Western Mediterranean Sea: Estimate of Ebro River inputs. *Chemosphere* 65, 224–236. doi: [10.1016/j.chemosphere.2006.02.058](https://doi.org/10.1016/j.chemosphere.2006.02.058)
- Gómez F, González N, Echevarría F, García CM 2000 Distribution and fluxes of dissolved nutrients in the Strait of Gibraltar and its relationships to microphytoplankton biomass. *Estuar. Coast. Shelf Sci.* 51, 439–449. doi: [10.1006/ecss.2000.0689](https://doi.org/10.1006/ecss.2000.0689)
- González-Alemán JJ, Pascale S, Gutierrez-Fernandez J, Murakami H, Gaertner MÁ et al. 2019 Potential Increase in Hazard From Mediterranean Hurricane Activity With Global Warming. *Geophys. Res. Lett.* 46, 1754–1764. doi: [10.1029/2018GL081253](https://doi.org/10.1029/2018GL081253)
- González-Moreno P, Lazzaro L, Vilà M, Preda C, Adriaens T et al. 2019 Consistency of impact assessment protocols for non-native species. *NeoBiota* 44, 1–25. doi: [10.3897/neobiota.44.31650](https://doi.org/10.3897/neobiota.44.31650)
- González-Moreno P, Pino J, Gassó N, Vilà M 2013 Landscape context modulates alien plant invasion in Mediterranean forest edges. *Biol. Invasions* 15, 547–557. doi: [10.1007/s10530-012-0306-x](https://doi.org/10.1007/s10530-012-0306-x)
- Gormley-Gallagher AM, Sterl S, Hirsch AL, Seneviratne SI, Davin EL et al. 2020 Agricultural management effects on mean and extreme temperature trends. *Earth Syst. Dynam. Discuss.*, 1–27. doi: [10.5194/esd-2020-35](https://doi.org/10.5194/esd-2020-35)
- Gottipolu RR, Landa ER, Schladweiler MC, McGee JK, Ledbetter AD et al. 2008 Cardiopulmonary Responses of Intratracheally Instilled Tire Particles and Constituent Metal Components. *Inhal. Toxicol.* 20, 473–484. doi: [10.1080/08958370701858427](https://doi.org/10.1080/08958370701858427)
- Goubanova K, Li L 2007 Extremes in temperature and precipitation around the Mediterranean basin in an ensemble of future climate scenario simulations. *Glob. Planet. Change* 57, 27–42. doi: [10.1016/j.gloplacha.2006.11.012](https://doi.org/10.1016/j.gloplacha.2006.11.012)
- Goubert C, Minard G, Vieira C, Boulesteix M 2016 Population genetics of the Asian tiger mosquito *Aedes albopictus*, an invasive vector of human diseases. *Heredity (Edinb.)* 117, 125–134. doi: [10.1038/hdy.2016.35](https://doi.org/10.1038/hdy.2016.35)
- Goyet C, Hassoun AER, Gemayel E, Touratier F, Abboud-Abi Saab M et al. 2016 Thermodynamic forecasts of the Mediterranean Sea acidification. *Mediterr. Mar. Sci.* 17, 508–518. doi: [10.12681/mms.1487](https://doi.org/10.12681/mms.1487)
- Gramberger M, Harrison PA, Jäger J, Kok K, Libbrecht S et al. 2013 Report on the third CLIMSAVE European stakeholder workshop. CLIMSAVE Project.
- Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J et al. 2008 Global change and the ecology of cities. *Science (80-. )* 319, 756–760. doi: [10.1126/science.1150195](https://doi.org/10.1126/science.1150195)
- Grise KM, Davis SM, Simpson IR, Waugh DW, Fu Q et al. 2019



- Recent tropical expansion: natural variability or forced response? *J. Clim.* 32, 1551–1571. doi: [10.1175/jcli-d-18-0444.1](https://doi.org/10.1175/jcli-d-18-0444.1)
- Grisogono B, Belušić D 2009 A review of recent advances in understanding the meso- and microscale properties of the severe Bora wind. *Tellus A Dyn. Meteorol. Oceanogr.* 61 A, 1–16. doi: [10.1111/j.1600-0870.2008.00369.x](https://doi.org/10.1111/j.1600-0870.2008.00369.x)
- Gritti ES, Smith B, Sykes MT 2006 Vulnerability of Mediterranean Basin ecosystems to climate change and invasion by exotic plant species. *J. Biogeogr.* 33, 145–157. doi: [10.1111/j.1365-2699.2005.01377.x](https://doi.org/10.1111/j.1365-2699.2005.01377.x)
- Gros M, Petrović M, Ginebreda A, Barceló D 2010 Removal of pharmaceuticals during wastewater treatment and environmental risk assessment using hazard indexes. *Environ. Int.* 36, 15–26. doi: [10.1016/j.envint.2009.09.002](https://doi.org/10.1016/j.envint.2009.09.002)
- Gualdi S, Somot S, Li L, Artale V, Adani M et al. 2013 The CIRCE Simulations: Regional Climate Change Projections with Realistic Representation of the Mediterranean Sea. *Bull. Am. Meteorol. Soc.* 94. doi: [10.1175/BAMS-D-11-00136.1](https://doi.org/10.1175/BAMS-D-11-00136.1)
- Guichard L, Dedieu F, Jeuffroy M-H, Meynard J-M, Reau R et al. 2017 Le plan Ecophyto de réduction d'usage des pesticides en France : décryptage d'un échec et raisons d'espérer. *Cah. Agric.* 26, 14002. doi: [10.1051/cagri/2017004](https://doi.org/10.1051/cagri/2017004)
- Guigé C, Tedetti M, Dang DH, Mullet J-U, Garnier C et al. 2017 Remobilization of polycyclic aromatic hydrocarbons and organic matter in seawater during sediment resuspension experiments from a polluted coastal environment: Insights from Toulon Bay (France). *Environ. Pollut.* 229, 627–638. doi: [10.1016/j.envpol.2017.06.090](https://doi.org/10.1016/j.envpol.2017.06.090)
- Guigé C, Tedetti M, Ferretto N, García N, Méjanelle L et al. 2014 Spatial and seasonal variabilities of dissolved hydrocarbons in surface waters from the Northwestern Mediterranean Sea: Results from one year intensive sampling. *Sci. Total Environ.* 466–467, 650–662. doi: [10.1016/j.scitotenv.2013.07.082](https://doi.org/10.1016/j.scitotenv.2013.07.082)
- Guigé C, Tedetti M, Giorgi S, Goutx M 2011 Occurrence and distribution of hydrocarbons in the surface microlayer and subsurface water from the urban coastal marine area off Marseilles, Northwestern Mediterranean Sea. *Mar. Pollut. Bull.* 62, 2741–2752. doi: [10.1016/j.marpolbul.2011.09.013](https://doi.org/10.1016/j.marpolbul.2011.09.013)
- Gutiérrez C, Somot S, Nabat P, Mallet M, Corre L et al. 2020 Future evolution of surface solar radiation and photovoltaic potential in Europe: investigating the role of aerosols. *Environ. Res. Lett.* 15, 034035. doi: [10.1088/1748-9326/ab6666](https://doi.org/10.1088/1748-9326/ab6666)
- Guy-Haim T 2017 The impact of ocean warming and acidification on coastal benthic species and communities.
- Guy-Haim T, Silverman J, Raddatz S, Wahl M, Rilov G 2016 Shifted coastal communities and ecosystem functions under predicted warming and acidification. in *41st CIESM Congress, Kiel, Germany*.
- Guzmán GI, González de Molina M, Soto Fernández D, Infante-Amate J, Aguilera E 2018 Spanish agriculture from 1900 to 2008: a long-term perspective on agroecosystem energy from an agroecological approach. *Reg. Environ. Chang.* 18, 995–1008. doi: [10.1007/s10113-017-1136-2](https://doi.org/10.1007/s10113-017-1136-2)
- Haarsma RJ, Selten F, Van den Hurk B, Hazeleger W, Wang X 2009 Drier Mediterranean soils due to greenhouse warming bring easterly winds over summertime central Europe. *Geophys. Res. Lett.* 36. doi: [10.1029/2008gl036617](https://doi.org/10.1029/2008gl036617)
- Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M et al. 2008 Seawater carbonate chemistry in Ischia, Italy, 2008. doi: [10.1594/pangaea.819633](https://doi.org/10.1594/pangaea.819633)
- Hallegraeff GM 2010 Ocean climate change, phytoplankton community responses, and harmful algal blooms: A formidable predictive challenge. *J. Phycol.* 46, 220–235. doi: [10.1111/j.1529-8817.2010.00815.x](https://doi.org/10.1111/j.1529-8817.2010.00815.x)
- Halpern BS, McLeod KL, Rosenberg AA, Crowder LB 2008a Managing for cumulative impacts in ecosystem-based management through ocean zoning. *Ocean Coast. Manag.* 51, 203–211. doi: [10.1016/j.ocecoaman.2007.08.002](https://doi.org/10.1016/j.ocecoaman.2007.08.002)
- Halpern BS, Walbridge S, Selkoe KA, Kappel C V., Micheli F et al. 2008b A global map of human impact on marine ecosystems. *Science (80-. )*. 319, 948–952. doi: [10.1126/science.1149345](https://doi.org/10.1126/science.1149345)
- Handmer J, Honda Y, Kundzewicz ZW, Arnell NW, Benito G et al. 2012 Changes in impacts of climate extremes: human systems and ecosystems, in *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC)*, eds. Field CB, Barros V, Stocker TF, Qin D, Dokken DJ et al. (Cambridge, United Kingdom and New York, NY, USA), 231–290.
- Hannah L, Roehrdanz PR, Ikegami M, Shepard A V., Shaw MR et al. 2013 Climate change, wine, and conservation. *Proc. Natl. Acad. Sci. U. S. A.* 110, 6907–6912. doi: [10.1073/pnas.1210127110](https://doi.org/10.1073/pnas.1210127110)
- Hansen MC, DeFries RS 2004 Detecting Long-term Global Forest Change Using Continuous Fields of Tree-Cover Maps from 8-km Advanced Very High Resolution Radiometer (AVHRR) Data for the Years 1982–99. *Ecosystems* 7, 695–716. doi: [10.1007/s10021-004-0243-3](https://doi.org/10.1007/s10021-004-0243-3)
- Hanson S, Nicholls RJ 2012 Extreme flood events and port cities through the twenty-first century: implications of climate change and other drivers, in *Maritime Transport and the Climate Change Challenge*, eds. Asariotis R, Benamara H (Abingdon, GB: Earthscan from Routledge), 243–265.
- Hanzer F, Förster K, Nemeš J, Strasser U 2018 Projected cryospheric and hydrological impacts of 21<sup>st</sup> century climate change in the Ötztal Alps (Austria) simulated using a physically based approach. *Hydrol. Earth Syst. Sci.* 22, 1593–1614. doi: [10.5194/hess-22-1593-2018](https://doi.org/10.5194/hess-22-1593-2018)
- Hao R, Yu D, Liu Y, Liu Y, Qiao J et al. 2017 Impacts of changes in climate and landscape pattern on ecosystem services. *Sci. Total Environ.* 579, 718–728. doi: [10.1016/j.scitotenv.2016.11.036](https://doi.org/10.1016/j.scitotenv.2016.11.036)
- Harris I, Osborn TJ, Jones PD, Lister D 2020 Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Sci. Data* 7, 109. doi: [10.1038/s41597-020-0453-3](https://doi.org/10.1038/s41597-020-0453-3)
- Harvey BJ, Shaffrey LC, Woollings TJ, Zappa G, Hodges KI 2012 How large are projected 21<sup>st</sup> century storm track changes. *Geophys. Res. Lett.* 39. doi: [10.1029/2012GL052873](https://doi.org/10.1029/2012GL052873)
- Harzallah A, Jordà G, Dubois C, Sannino G, Carillo A et al. 2018 Long term evolution of heat budget in the Mediterranean Sea from Med-CORDEX forced and coupled simulations.

- Clim. Dyn.* 51, 1145–1165. doi: [10.1007/s00382-016-3363-5](https://doi.org/10.1007/s00382-016-3363-5)
- Hasanean HM 2004 Precipitation variability over the Mediterranean and its linkage with El Niño Southern Oscillation (ENSO). *J. Meteorol.* 29, 151–160.
- Hassanien MA, Abdel-Latif NM 2008 Polycyclic aromatic hydrocarbons in road dust over Greater Cairo, Egypt. *J. Hazard. Mater.* 151, 247–254. doi: [10.1016/j.jhazmat.2007.05.079](https://doi.org/10.1016/j.jhazmat.2007.05.079)
- Hatfield JL, Boote KJ, Kimball BA, Ziska LH, Izaurralde RC et al. 2011 Climate Impacts on Agriculture: Implications for Crop Production. *Agron. J.* 103, 351. doi: [10.2134/agronj2010.0303](https://doi.org/10.2134/agronj2010.0303)
- Hatzaki M, Flocas HA, Giannakopoulos C, Maheras P 2009 The impact of the eastern Mediterranean teleconnection pattern on the Mediterranean climate. *J. Clim.* 22, 977–992. doi: [10.1175/2008JCLI2519.1](https://doi.org/10.1175/2008JCLI2519.1)
- Hatzaki M, Flocas HA, Simmonds I, Kouroutzoglou J, Keay K et al. 2014 Seasonal Aspects of an Objective Climatology of Anticyclones Affecting the Mediterranean. *J. Clim.* 27, 9272–9289. doi: [10.1175/JCLI-D-14-00186.1](https://doi.org/10.1175/JCLI-D-14-00186.1)
- Haywood JM, Bellouin N, Jones A, Boucher O, Wild M et al. 2011 The roles of aerosol, water vapor and cloud in future global dimming/brightening. *JGR Atmos.* 116, D20203. doi: [10.1029/2011JD016000](https://doi.org/10.1029/2011JD016000)
- Hellmann JJ, Byers JE, Bierwagen BG, Dukes JS 2008 Five Potential Consequences of Climate Change for Invasive Species. *Conserv. Biol.* 22, 534–543. doi: [10.1111/j.1523-1739.2008.00951.x](https://doi.org/10.1111/j.1523-1739.2008.00951.x)
- Hentgen L, Ban N, Kröner N, Leutwyler D, Schär C 2019 Clouds in Convection-Resolving Climate Simulations Over Europe. *JGR Atmos.* 124, 3849–3870. doi: [10.1029/2018JD030150](https://doi.org/10.1029/2018JD030150)
- Hernández-Brito D, Carrete M, Ibáñez C, Juste J, Tella J 2018 Nest-site competition and killing by invasive parakeets cause the decline of a threatened bat population. *R. Soc. Open Sci.* 5. doi: [10.1098/rsos.172477](https://doi.org/10.1098/rsos.172477)
- Hernández-Brito D, Carrete M, Popa-Lisseanu AG, Ibáñez C, Tella JL 2014 Crowding in the City: Losing and Winning Competitors of an Invasive Bird. *PLoS One* 9, e100593. doi: [10.1371/journal.pone.0100593](https://doi.org/10.1371/journal.pone.0100593)
- Hernandez-Vargas G, Sosa-Hernández JE, Saldarriaga-Hernandez S, Villalba-Rodríguez AM, Parra-Saldívar R et al. 2018 Electrochemical biosensors: A solution to pollution detection with reference to environmental contaminants. *Biosensors* 8, 1–21. doi: [10.3390/bios8020029](https://doi.org/10.3390/bios8020029)
- Hernández A, Miranda M, Arellano EC, Saura S, Ovalle C 2015 Landscape dynamics and their effect on the functional connectivity of a Mediterranean landscape in Chile. *Ecol. Indic.* 48, 198–206. doi: [10.1016/j.ecolind.2014.08.010](https://doi.org/10.1016/j.ecolind.2014.08.010)
- Hernández F, Alonso-Pérez S, Hernández-Armas J, Cuevas E, Karlsson L et al. 2005 Influence of major African dust intrusions on the 137Cs and 40K activities in the lower atmosphere at the Island of Tenerife. *Atmos. Environ.* 39, 4111–4118. doi: [10.1016/j.atmosenv.2005.03.032](https://doi.org/10.1016/j.atmosenv.2005.03.032)
- Herrero-Hernández E, Andrades MS, Álvarez-Martín A, Pose-Juan E, Rodríguez-Cruz MS et al. 2013 Occurrence of pesticides and some of their degradation products in waters in a Spanish wine region. *J. Hydrol.* 486, 234–245. doi: [10.1016/j.jhydrol.2013.01.025](https://doi.org/10.1016/j.jhydrol.2013.01.025)
- Herrmann M, Somot S, Calmanti S, Dubois C, Sevault F 2011 Representation of spatial and temporal variability of daily wind speed and of intense wind events over the Mediterranean Sea using dynamical downscaling: impact of the regional climate model configuration. *Nat. Hazards Earth Syst. Sci.*, 1983–2011. <https://hal.archives-ouvertes.fr/hal-00766447/> [Accessed September 17, 2019]
- Hertig E, Seubert S, Jacobeit J 2010 Temperature extremes in the Mediterranean area: Trends in the past and assessments for the future. *Nat. Hazards Earth Syst. Sci.* 10, 2039–2050. doi: [10.5194/nhess-10-2039-2010](https://doi.org/10.5194/nhess-10-2039-2010)
- Hewitson B, Janetos AC, Carter TR, Giorgi F, Jones RG et al. 2014 Regional context, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 1133–1197.
- Hildebrandt A, Guillaumon M, Lacorte S, Tauler R, Barceló D 2008 Impact of pesticides used in agriculture and vineyards to surface and groundwater quality (North Spain). *Water Res.* 42, 3315–26. doi: [10.1016/j.watres.2008.04.009](https://doi.org/10.1016/j.watres.2008.04.009)
- Hill J, Stellmes M, Udelhoven T, Röder A, Sommer S 2008 Mediterranean desertification and land degradation. *Glob. Planet. Change* 64, 146–157. doi: [10.1016/j.gloplacha.2008.10.005](https://doi.org/10.1016/j.gloplacha.2008.10.005)
- Hirai H, Takada H, Ogata Y, Yamashita R, Mizukawa K et al. 2011 Organic micropollutants in marine plastics debris from the open ocean and remote and urban beaches. *Mar. Pollut. Bull.* 62, 1683–1692. doi: [10.1016/j.marpolbul.2011.06.004](https://doi.org/10.1016/j.marpolbul.2011.06.004)
- Hirche A, Salamani M, Tarhouni M, Nedjraoui D, El Hag M et al. 2018 The Maghreb (North Africa) rangelands evolution over forty years: re-greening or degradation, in *Desertification: Past, Current and Future Trends*, eds. Squires VR, Ariapour A (New York, United States: Nova Science Publishers), 73–106.
- Hirose K, Povinec PP 2015 Sources of plutonium in the atmosphere and stratosphere-troposphere mixing. *Sci. Rep.* 5, 15707. doi: [10.1038/srep15707](https://doi.org/10.1038/srep15707)
- Hoagland P, Anderson DM, Kaoru Y, White AW 2002 The economic effects of harmful algal blooms in the United States: Estimates, assessment issues, and information needs. *Estuaries* 25, 819–837. doi: [10.1007/BF02804908](https://doi.org/10.1007/BF02804908)
- Hobbs RJ, Huenneke LF 1992 Disturbance, Diversity, and Invasion: Implications for Conservation. *Conserv. Biol.* 6, 324–337. doi: [10.1046/j.1523-1739.1992.06030324.x](https://doi.org/10.1046/j.1523-1739.1992.06030324.x)
- Hock R, Bliss A, Marzeion B, Giesen RH, Hirabayashi Y et al. 2019 GlacierMIP - A model intercomparison of global-scale glacier mass-balance models and projections. *J. Glaciol.* 65, 453–467. doi: [10.1017/jog.2019.22](https://doi.org/10.1017/jog.2019.22)
- Hódar JA, Zamora R 2004 Herbivory and climatic warming: a Mediterranean outbreaking caterpillar attacks a relict, boreal pine species. *Biodivers. Conserv.* 13, 493–500. doi: [10.1023/b:bioc.0000009495.95589.a7](https://doi.org/10.1023/b:bioc.0000009495.95589.a7)
- Hoerling M, Eischeid J, Perlwitz J, Quan X, Zhang T et al. 2012 On the Increased Frequency of Mediterranean Drought. *J. Clim.* 25, 2146–2161. doi: [10.1175/JCLI-D-11-00296.1](https://doi.org/10.1175/JCLI-D-11-00296.1)

- Horton R, Rosenzweig C, Gornitz V, Bader D, O'Grady M 2010 Climate risk information: climate change scenarios and implications for NYC infrastructure. New York City Panel on Climate Change. *Ann. N. Y. Acad. Sci.* 1196, 147–228. doi: [10.1111/j.1749-6632.2010.05323.x](https://doi.org/10.1111/j.1749-6632.2010.05323.x)
- Hoskins BJ, Hodges KI 2002 New perspectives on the northern hemisphere winter storm tracks. *J. Atmos. Sci.* 59, 1041–1061. doi: [10.1175/1520-0469\(2002\)059<1041:npotnh>2.0.co;2](https://doi.org/10.1175/1520-0469(2002)059<1041:npotnh>2.0.co;2)
- Hostert P, Röder A, Hill J, Udelhoven T, Tsiourlis G 2003 Retrospective studies of grazing-induced land degradation: A case study in central Crete, Greece. *Int. J. Remote Sens.* 24, 4019–4034. doi: [10.1080/0143116031000103844](https://doi.org/10.1080/0143116031000103844)
- Houpuert L, de Madron XD, Testor P, Bosse A, D'Ortenzio F et al. 2016 Observations of open-ocean deep convection in the northwestern Mediterranean Sea: Seasonal and interannual variability of mixing and deep water masses for the 2007–2013 Period. *JGR Ocean.* 121, 8139–8171. doi: [10.1002/2016JC011857](https://doi.org/10.1002/2016JC011857)
- Hu ZZ, Wu Z 2004 The intensification and shift of the annual North Atlantic Oscillation in a global warming scenario simulation. *Tellus A Dyn. Meteorol. Oceanogr.* 56, 112–124. doi: [10.1111/j.1600-0870.2004.00050.x](https://doi.org/10.1111/j.1600-0870.2004.00050.x)
- Hueging H, Haas R, Born K, Jacob D, Pinto JG 2013 Regional changes in wind energy potential over Europe using regional climate model ensemble projections. *J. Appl. Meteorol. Climatol.* 52, 903–917. doi: [10.1175/JAMC-D-12-086.1](https://doi.org/10.1175/JAMC-D-12-086.1)
- Hughes PD 2018 Little Ice Age glaciers and climate in the Mediterranean mountains: a new analysis. *Geogr. Res. Lett.* 44, 15–45. doi: [10.18172/cig.3362](https://doi.org/10.18172/cig.3362)
- Hulme PE 2009 Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J. Appl. Ecol.* 46, 10–18. doi: [10.1111/j.1365-2664.2008.01600.x](https://doi.org/10.1111/j.1365-2664.2008.01600.x)
- Hulme PE, Brundu G, Camarda I, Dalias P, Lambdon PW et al. 2008 Assessing the risks to Mediterranean islands ecosystems from alien plant introductions, in *Plant invasions: human perception, ecological impacts and management*, eds. Tokarska-Guzik B, Brock JH, Brundu G, Child L, Daehler CC et al. (Buckhuys Publishers, Leiden), 39–56.
- Hunt A, Watkiss P 2011 Climate change impacts and adaptation in cities: a review of the literature. *Clim. Change* 104, 13–49. doi: [10.1007/s10584-010-9975-6](https://doi.org/10.1007/s10584-010-9975-6)
- Hurrell JW 1995 Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and Precipitation. *Science (80-. J.)* 269, 676–679. doi: [10.1126/science.269.5224.676](https://doi.org/10.1126/science.269.5224.676)
- Hurrell JW, Kushnir Y, Ottersen G, Visbeck M 2003 *The North Atlantic Oscillation: Climatic Significance and Environmental Impact*. Washington DC: American Geophysical Union doi: [10.1029/gm134p0vii](https://doi.org/10.1029/gm134p0vii)
- Hurrell JW, Van Loon H 1997 Decadal variations in climate associated with the North Atlantic oscillation. *Clim. Change* 36, 301–326. doi: [10.1023/A:1005314315270](https://doi.org/10.1023/A:1005314315270)
- Hussein RA, Al-Ghanim KA, Abd-El-Atty MM, Mohamed LA 2016 Contamination of red sea shrimp (*Palaemon serratus*) with polycyclic aromatic hydrocarbons: A health risk assessment study. *Polish J. Environ. Stud.* 25, 615–620. doi: [10.15244/pjoes/60767](https://doi.org/10.15244/pjoes/60767)
- Ignatiades L, Gotsis-Skretas O, Pagou K, Krasakopoulou E 2009 Diversification of phytoplankton community structure and related parameters along a large-scale longitudinal east-west transect of the Mediterranean Sea. *J. Plankton Res.* 31, 411–428. doi: [10.1093/plankt/fbn124](https://doi.org/10.1093/plankt/fbn124)
- Iles C, Hegerl GC 2017 Role of the North Atlantic Oscillation in decadal temperature trends. *Environ. Res. Lett.* 12, 114010. doi: [10.1088/1748-9326/aa9152](https://doi.org/10.1088/1748-9326/aa9152)
- Im U 2013 Impact of sea-salt emissions on the model performance and aerosol chemical composition and deposition in the East Mediterranean coastal regions. *Atmos. Environ.* 75, 329–340. doi: [10.1016/J.ATMOSENV.2013.04.034](https://doi.org/10.1016/J.ATMOSENV.2013.04.034)
- Inghilesi AF, Mazza G, Cervo R, Gherardi F, Sposimo P et al. 2013 Alien Insects in Italy: Comparing Patterns from the Regional to European Level. *J. Insect Sci.* 13, 1–13. doi: [10.1673/031.013.7301](https://doi.org/10.1673/031.013.7301)
- IPCC 2007 *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.*, eds. Solomon S, Qin D, Manning M, Chen Z, Marquis M et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- IPCC 2013a *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.*, eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. Cambridge, United Kingdom and New York, NY, USA.
- IPCC 2013b Summary for Policymakers, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press).
- IPCC 2018 *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change.*, eds. Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J et al. In press.
- IPEMED 2011 Tomorrow, the Mediterranean Scenarios and projections for 2030. Paris.
- Ivar do Sul JA, Costa MF 2014 The present and future of microplastic pollution in the marine environment. *Environ. Pollut.* 185, 352–364. doi: [10.1016/j.envpol.2013.10.036](https://doi.org/10.1016/j.envpol.2013.10.036)
- Ivy-Ochs S, Kerschner H, Maisch M, Christl M, Kubik PW et al. 2009 Latest Pleistocene and Holocene glacier variations in the European Alps. *Quat. Sci. Rev.* 28, 2137–2149. doi: [10.1016/j.quascirev.2009.03.009](https://doi.org/10.1016/j.quascirev.2009.03.009)
- Izaurrealde RC, Thomson AM, Morgan JA, Fay PA, Polley HW et al. 2011 Climate Impacts on Agriculture: Implications for Forage and Rangeland Production. *Agron. J.* 103, 371. doi: [10.2134/agronj2010.0304](https://doi.org/10.2134/agronj2010.0304)
- Jacob D, Petersen J, Eggert B, Alias A, Christensen OB et al. 2014 EURO-CORDEX: new high-resolution climate change projections for European impact research. *Reg. Environ. Chang.* 14, 563–578. doi: [10.1007/s10113-013-0499-2](https://doi.org/10.1007/s10113-013-0499-2)
- Jacobit J, Hertig E, Seubert S, Lutz K 2014 Statistical down-

- scaling for climate change projections in the Mediterranean region: methods and results. *Reg. Environ. Chang.* 14, 1891–1906. doi: [10.1007/s10113-014-0605-0](https://doi.org/10.1007/s10113-014-0605-0)
- Jaeger EB, Seneviratne SI 2011 Impact of soil moisture–atmosphere coupling on European climate extremes and trends in a regional climate model. *Clim. Dyn.* 36, 1919–1939. doi: [10.1007/s00382-010-0780-8](https://doi.org/10.1007/s00382-010-0780-8)
- Javal M, Roques A, Haran J, Hérard F, Keena M et al. 2019 Complex invasion history of the Asian long-horned beetle: fifteen years after first detection in Europe. *J. Pest Sci. (2004)*. 92, 173–187. doi: [10.1007/s10340-017-0917-1](https://doi.org/10.1007/s10340-017-0917-1)
- Jeanmonod D, Schlüssel A, Gamisans J 2011 Status and trends in the alien flora of Corsica. *EPPO Bull.* 41, 85–99. doi: [10.1111/j.1365-2338.2011.02440.x](https://doi.org/10.1111/j.1365-2338.2011.02440.x)
- Jebari S 2009 Water erosion modeling using fractal rainfall disaggregation - A study in semi-arid Tunisia.
- Jebari S, Berndtsson R, Bahri A, Boufaroua M 2010 Spatial soil loss risk and reservoir siltation in semi-arid Tunisia. *Hydrol. Sci. J.* 55, 121–137. doi: [10.1080/02626660903529049](https://doi.org/10.1080/02626660903529049)
- Jenkins P 1999 Trade and exotic species introductions, in *Invasive species and biodiversity management*, eds. Sandlund O, Schei P, Viken A (Kluwer, Dordrecht).
- Jeschke JM, Strayer DL 2005 Invasion success of vertebrates in Europe and North America. *Proc. Natl. Acad. Sci. U. S. A.* 102, 7198–7202. doi: [10.1073/pnas.0501271102](https://doi.org/10.1073/pnas.0501271102)
- Jeschke JM, Strayer DL 2006 Determinants of vertebrate invasion success in Europe and North America. *Glob. Chang. Biol.* 12, 1608–1619. doi: [10.1111/j.1365-2486.2006.01213.x](https://doi.org/10.1111/j.1365-2486.2006.01213.x)
- Jiang L, O'Neill BC 2017 Global urbanization projections for the Shared Socioeconomic Pathways. *Glob. Environ. Chang.* 42, 193–199. doi: [10.1016/j.gloenvcha.2015.03.008](https://doi.org/10.1016/j.gloenvcha.2015.03.008)
- Joffre RR, Rambal S 1993 How Tree Cover Influences the Water Balance of Mediterranean Rangelands. *Ecology* 74, 570–582. doi: [10.2307/1939317](https://doi.org/10.2307/1939317)
- Jones B, O'Neill BC 2016 Spatially explicit global population scenarios consistent with the Shared Socioeconomic Pathways. *Environ. Res. Lett.* 11, 84003. doi: [10.1088/1748-9326/11/8/084003](https://doi.org/10.1088/1748-9326/11/8/084003)
- Jones PD, Lister DH, Osborn TJ, Harpham C, Salmon M et al. 2012 Hemispheric and large-scale land surface air temperature variations: an extensive review and an update to 2010. *JGR Atmos.* 117, D05127. doi: [10.1029/2011JD017139](https://doi.org/10.1029/2011JD017139)
- Jordà G, Gomis D, Adloff F 2020 Vulnerability of marginal seas to sea-level rise: The case of the Mediterranean Sea. (*in Rev.*)
- Jordà G, Gomis D, Álvarez-Fanjul E 2012 The VAN2-ERA hind-cast of sea-level residuals: atmospheric forcing of sea-level variability in the Mediterranean Sea (1958–2008). *Sci. Mar.* 76S1, 133–146. doi: [10.3989/scimar.03612.19C](https://doi.org/10.3989/scimar.03612.19C)
- Jucker C, Lupi D 2011 Exotic Insects in Italy: An Overview on Their Environmental Impact. *Importance Biol. Interact. Study Biodivers.* doi: [10.5772/24263](https://doi.org/10.5772/24263)
- Justine J-L, Winsor L, Gey D, Gros P, Thévenot J 2018 Giant worms chez moi! Hammerhead flatworms (Platyhelminthes, Geoplanidae, *Bipalium* spp., *Diversibipalium* spp.) in metropolitan France and overseas French territories. *PeerJ* 6, e4672. doi: [10.7717/peerj.4672](https://doi.org/10.7717/peerj.4672)
- Kairis O, Karavitis C, Salvati L, Kounalaki A, Kosmas K 2015 Exploring the Impact of Overgrazing on Soil Erosion and Land Degradation in a Dry Mediterranean Agro-Forest Landscape (Crete, Greece). *Arid L. Res. Manag.* 29, 360–374. doi: [10.1080/15324982.2014.968691](https://doi.org/10.1080/15324982.2014.968691)
- Kalabokas PD, Thouret V, Cammas J-P, Volz-Thomas A, Boulanger D et al. 2015 The geographical distribution of meteorological parameters associated with high and low summer ozone levels in the lower troposphere and the boundary layer over the eastern Mediterranean (Cairo case). *Tellus B Chem. Phys. Meteorol.* 67, 27853. doi: [10.3402/tellusb.v67.27853](https://doi.org/10.3402/tellusb.v67.27853)
- Kalimeris A, Ranieri E, Founda D, Norrant C 2017 Variability modes of precipitation along a Central Mediterranean area and their relations with ENSO, NAO, and other climatic patterns. *Atmos. Res.* 198, 56–80. doi: [10.1016/j.atmosres.2017.07.031](https://doi.org/10.1016/j.atmosres.2017.07.031)
- Kallos G, Solomos S, Kushta J, Mitsakou C, Spyrou C et al. 2014 Natural and anthropogenic aerosols in the Eastern Mediterranean and Middle East: Possible impacts. *Sci. Total Environ.* 488–489, 389–397. doi: [10.1016/j.scitotenv.2014.02.035](https://doi.org/10.1016/j.scitotenv.2014.02.035)
- Kambezidis HD, Psiloglou BE, Karagiannis D, Dumka UC, Kaskaoutis DG 2016 Recent improvements of the Meteorological Radiation Model for solar irradiance estimates under all-sky conditions. *Renew. Energy* 93, 142–158. doi: [10.1016/j.renene.2016.02.060](https://doi.org/10.1016/j.renene.2016.02.060)
- Kantiani L, Farré M, Asperger D, Rubio F, González S et al. 2008 Triclosan and methyl-triclosan monitoring study in the northeast of Spain using a magnetic particle enzyme immunoassay and confirmatory analysis by gas chromatography-mass spectrometry. *J. Hydrol.* 361, 1–9. doi: [10.1016/j.jhydrol.2008.07.016](https://doi.org/10.1016/j.jhydrol.2008.07.016)
- Karanasiou A, Querol X, Alastuey A, Perez N, Pey J et al. 2014 Particulate matter and gaseous pollutants in the Mediterranean Basin: Results from the MED-PARTICLES project. *Sci. Total Environ.* 488–489, 297–315. doi: [10.1016/j.scitotenv.2014.04.096](https://doi.org/10.1016/j.scitotenv.2014.04.096)
- Karanasiou AA, Siskos PA, Eleftheriadis K 2009 Assessment of source apportionment by Positive Matrix Factorization analysis on fine and coarse urban aerosol size fractions. *Atmos. Environ.* 43, 3385–3395. doi: [10.1016/j.atmosenv.2009.03.051](https://doi.org/10.1016/j.atmosenv.2009.03.051)
- Kark S, Solarz W, Chiron F, Clergeau P, Shirley S 2009 Alien Birds, Amphibians and Reptiles of Europe, in *Handbook of Alien Species in Europe*, ed. DAISIE (Springer Netherlands), 105–118. doi: [10.1007/978-1-4020-8280-1\\_8](https://doi.org/10.1007/978-1-4020-8280-1_8)
- Kaskaoutis DG, Dumka UC, Rashki A, Psiloglou BE, Gavriil A et al. 2019 Analysis of intense dust storms over the eastern Mediterranean in March 2018: Impact on radiative forcing and Athens air quality. *Atmos. Environ.* 209, 23–39. doi: [10.1016/j.atmosenv.2019.04.025](https://doi.org/10.1016/j.atmosenv.2019.04.025)
- Kassomenos PA, Kelessis A, Paschalidou AK, Petrakakis M 2011 Identification of sources and processes affecting particulate pollution in Thessaloniki, Greece. *Atmos. Environ.* 45, 7293–7300. doi: [10.1016/j.atmosenv.2011.08.034](https://doi.org/10.1016/j.atmosenv.2011.08.034)
- Katsanevakis S, Coll M, Piroddi C, Steenbeek J, Ben Rais Lasram F et al. 2014a Invading the Mediterranean Sea: biodi-

- versity patterns shaped by human activities. *Front. Mar. Sci.* 1. doi: [10.3389/fmars.2014.00032](https://doi.org/10.3389/fmars.2014.00032)
- Katsanevakis S, Rilov G, Edelist D 2018 Impacts of marine invasive alien species on European fisheries and aquaculture - plague or boon?, in *Engaging marine scientists and fishers to share knowledge and perceptions - early lessons. CIESM Monograph n° 50*, ed. Briand F (CIESM Publisher, Monaco and Paris), 125–132.
- Katsanevakis S, Tempera F, Teixeira H 2016 Mapping the impact of alien species on marine ecosystems: The Mediterranean Sea case study. *Divers. Distrib.* 22, 694–707. doi: [10.1111/ddi.12429](https://doi.org/10.1111/ddi.12429)
- Katsanevakis S, Wallentinus I, Zenetos A, Leppäkoski E, Çinar ME et al. 2014b Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Aquat. Invasions* 9, 391–423. doi: [10.3391/ai.2014.9.4.01](https://doi.org/10.3391/ai.2014.9.4.01)
- Katsanevakis S, Zenetos A, Belchior C, Cardoso AC 2013 Invading European Seas: Assessing pathways of introduction of marine aliens. *Ocean Coast. Manag.* 76, 64–74. doi: [10.1016/j.ocecoaman.2013.02.024](https://doi.org/10.1016/j.ocecoaman.2013.02.024)
- Kc S, Lutz W 2017 The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100. *Glob. Environ. Chang.* 42, 181–192. doi: [10.1016/j.gloenvcha.2014.06.004](https://doi.org/10.1016/j.gloenvcha.2014.06.004)
- Kchih H, Perrino C, Cherif S 2015 Investigation of Desert Dust Contribution to Source Apportionment of PM10 and PM2.5 from a Southern Mediterranean Coast. *Aerosol Air Qual. Res.* 15, 454–464. doi: [10.4209/aaqr.2014.10.0255](https://doi.org/10.4209/aaqr.2014.10.0255)
- Keeley JE, Bond WJ, Bradstock RA, Pausas JG, Rundel PW 2012 *Fire in Mediterranean ecosystems: Ecology, evolution and management*. Cambridge, UK: Cambridge University Press doi: [10.1017/cbo9781139033091](https://doi.org/10.1017/cbo9781139033091)
- Keenan TF, Maria Serra J, Lloret F, Ninyerola M, Sabate S 2011 Predicting the future of forests in the Mediterranean under climate change, with niche- and process-based models: CO<sub>2</sub> matters! *Glob. Chang. Biol.* 17, 565–579. doi: [10.1111/j.1365-2486.2010.02254.x](https://doi.org/10.1111/j.1365-2486.2010.02254.x)
- Kelepertzis E 2014 Accumulation of heavy metals in agricultural soils of Mediterranean: Insights from Argolida basin, Peloponnese, Greece. *Geoderma* 221–222, 82–90. doi: [10.1016/j.geoderma.2014.01.007](https://doi.org/10.1016/j.geoderma.2014.01.007)
- Keller RP, Geist J, Jeschke JM, Kühn I 2011 Invasive species in Europe: ecology, status, and policy. *Environ. Sci. Eur.* 23, 23. doi: [10.1186/2190-4715-23-23](https://doi.org/10.1186/2190-4715-23-23)
- Kelley C, Ting M, Seager R, Kushnir Y 2012 The relative contributions of radiative forcing and internal climate variability to the late 20<sup>th</sup> century winter drying of the Mediterranean region. *Clim. Dyn.* 38, 2001–2015. doi: [10.1007/s00382-011-1221-z](https://doi.org/10.1007/s00382-011-1221-z)
- Kendon EJ, Roberts NM, Fowler HJ, Roberts MJ, Chan SC et al. 2014 Heavier summer downpours with climate change revealed by weather forecast resolution model. *Nat. Clim. Chang.* 4, 570–576. doi: [10.1038/nclimate2258](https://doi.org/10.1038/nclimate2258)
- Kenis M, Auger-Rozenberg M-A, Roques A, Timms L, Péré C et al. 2009 Ecological effects of invasive alien insects. *Biol. Invasions* 11, 21–45. doi: [10.1007/s10530-008-9318-y](https://doi.org/10.1007/s10530-008-9318-y)
- Kenis M, Branco M 2010 Impact of alien terrestrial arthropods in Europe. Chapter 5. *BioRisk* 4, 51–71. doi: [10.3897/biorisk.4.42](https://doi.org/10.3897/biorisk.4.42)
- Kerdelhué C, Boivin T, Burban C 2014 Contrasted invasion processes imprint the genetic structure of an invasive scale insect across southern Europe. *Heredity (Edinb)*. 113, 390–400. doi: [10.1038/hdy.2014.39](https://doi.org/10.1038/hdy.2014.39)
- Kirichenko N, Augustin S, Kenis M 2019 Invasive leafminers on woody plants: a global review of pathways, impact, and management. *J. Pest Sci. (2004)*. 92, 93–106. doi: [10.1007/s10340-018-1009-6](https://doi.org/10.1007/s10340-018-1009-6)
- Klaper R, Welch L 2011 Emerging contaminant threats and the Great Lakes. [https://c.ymcdn.com/sites/www.productstewardship.us/resource/collection/FFDF28A1-9926-46E7-87F9-70C7BBD95491/Emerging\\_Contaminant\\_Threats\\_and\\_the\\_Great\\_Lakes.pdf](https://c.ymcdn.com/sites/www.productstewardship.us/resource/collection/FFDF28A1-9926-46E7-87F9-70C7BBD95491/Emerging_Contaminant_Threats_and_the_Great_Lakes.pdf)
- Kleitou P, Hall-Spencer J, Rees S, Sfenthourakis S, Demetriou A et al. 2019 Tackling the lionfish invasion in the Mediterranean. The EU-LIFE RELIONMED Project: progress and results. in *Proceedings of the 1st Mediterranean Symposium on the Non-Indigenous Species* (Antalya, Turkey), 65–70.
- Kletou DC, Hall-Spencer JM, Kleitou P 2016 A lionfish (*Pterois miles*) invasion has begun in the Mediterranean Sea. *Mar. Biodivers. Rec.* 9. doi: [10.1186/s41200-016-0065-y](https://doi.org/10.1186/s41200-016-0065-y)
- Klippertz P, Ulbrich U, Marques F, Corte-Real J 2003 Decadal changes in the link between El Niño and springtime North Atlantic oscillation and European–North African rainfall. *Int. J. Climatol.* 23, 1293–1311. doi: [10.1002/joc.944](https://doi.org/10.1002/joc.944)
- Klippertz P, Ulbrich U, Speth P 2000 Changing cyclones and surface wind speeds over the North Atlantic and Europe in a transient GHG experiment. *Clim. Res.* 15, 109–122. doi: [10.3354/cr015109](https://doi.org/10.3354/cr015109)
- Knutson TR, Zeng F, Wittenberg AT, Knutson TR, Zeng F et al. 2013 Multimodel Assessment of Regional Surface Temperature Trends: CMIP3 and CMIP5 Twentieth-Century Simulations. *J. Clim.* 26, 8709–8743. doi: [10.1175/JCLI-D-12-00567.1](https://doi.org/10.1175/JCLI-D-12-00567.1)
- Knutti R, Sedláček J 2012 Robustness and uncertainties in the new CMIP5 climate model projections. *Nat. Clim. Chang.* 3, 369–373. doi: [10.1038/nclimate1716](https://doi.org/10.1038/nclimate1716)
- Knutti R, Sedláček J, Sanderson BM, Lorenz R, Fischer EM et al. 2017 A climate model projection weighting scheme accounting for performance and interdependence. *Geophys. Res. Lett.* 44, 1909–1918.
- Koçak M, Mihalopoulos N, Kubilay NN 2007a Chemical composition of the fine and coarse fraction of aerosols in the north-eastern Mediterranean. *Atmos. Environ.* 41, 7351–7368. doi: [10.1016/j.atmosenv.2007.05.011](https://doi.org/10.1016/j.atmosenv.2007.05.011)
- Koçak M, Mihalopoulos N, Kubilay NN 2007b Contributions of natural sources to high PM10 and PM2.5 events in the eastern Mediterranean. *Atmos. Environ.* 41, 3806–3818. doi: [10.1016/j.atmosenv.2007.01.009](https://doi.org/10.1016/j.atmosenv.2007.01.009)
- Koelmans AA, Besseling E, Foekema E, Kooi M, Mintenig S et al. 2017 Risks of plastic debris: unravelling fact, opinion, perception, and belief. *Environ. Sci. Technol.* 51, 11513–11519. doi: [10.1021/acs.est.7b02219](https://doi.org/10.1021/acs.est.7b02219)
- Kok K, Pedde S, Gramberger M, Harrison PA, Holman IP 2019

- New European socio-economic scenarios for climate change research: Operationalising concepts to extend the shared socio-economic pathways. *Reg. Environ. Chang.* 19, 643–654. doi: [10.1007/s10113-018-1400-0](https://doi.org/10.1007/s10113-018-1400-0)
- Kok K, Rothman DS, Patel M 2006 Multi-scale narratives from an IA perspective: Part I. European and Mediterranean scenario development. *Futures* 38, 261–284. doi: [10.1016/j.futures.2005.07.001](https://doi.org/10.1016/j.futures.2005.07.001)
- Konstantinou IK, Hela DG, Albanis TA 2006 The status of pesticide pollution in surface waters (rivers and lakes) of Greece. Part I. Review on occurrence and levels. *Environ. Pollut.* 141, 555–570. doi: [10.1016/j.envpol.2005.07.024](https://doi.org/10.1016/j.envpol.2005.07.024)
- Kostopoulou E, Giannakopoulos C, Hatzaki M, Karali A, Hadjinicolaou P et al. 2014 Spatio-temporal patterns of recent and future climate extremes in the eastern Mediterranean and Middle East region. *Nat. Hazards Earth Syst. Sci.* 14, 1565–1577. doi: [10.5194/nhess-14-1565-2014](https://doi.org/10.5194/nhess-14-1565-2014)
- Kostopoulou E, Jones PD 2007 Comprehensive analysis of the climate variability in the eastern Mediterranean. Part II: relationships between atmospheric circulation patterns and surface climatic elements. *Int. J. Climatol.* 27, 1351–1371. doi: [10.1002/joc.1466](https://doi.org/10.1002/joc.1466)
- Kotlarski S, Szabó P, Herrera S, Rätty O, Keuler K et al. 2019 Observational uncertainty and regional climate model evaluation: A pan-European perspective. *Int. J. Climatol.* 39, 3730–3749. doi: [10.1002/joc.5249](https://doi.org/10.1002/joc.5249)
- Kouroutzoglou J, Flocas HA, Keay K, Simmonds I, Hatzaki M 2010 Climatological aspects of explosive cyclones in the Mediterranean. *Int. J. Climatol.* 31, 1785–1802. doi: [10.1002/joc.2203](https://doi.org/10.1002/joc.2203)
- Kouroutzoglou J, Flocas HA, Keay K, Simmonds I, Hatzaki M 2011 Climatological aspects of explosive cyclones in the Mediterranean. *Int. J. Climatol.* 31, 1785–1802. doi: [10.1002/joc.2203](https://doi.org/10.1002/joc.2203)
- Koutavas A 2013 CO<sub>2</sub> fertilization and enhanced drought resistance in Greek firs from Cephalonia Island, Greece. *Glob. Chang. Biol.* 19, 529–539. doi: [10.1111/gcb.12053](https://doi.org/10.1111/gcb.12053)
- Koutsodendris A, Papatheodorou G, Kougiourouki O, Georgiadis M 2008 Benthic marine litter in four Gulfs in Greece, Eastern Mediterranean; abundance, composition and source identification. *Estuar. Coast. Shelf Sci.* 77, 501–512. doi: [10.1016/j.ecss.2007.10.011](https://doi.org/10.1016/j.ecss.2007.10.011)
- Kress N, Manca BB, Klein B, Deponte D 2003 Continuing influence of the changed thermohaline circulation in the eastern Mediterranean on the distribution of dissolved oxygen and nutrients: Physical and chemical characterization of the water masses. *JGR Ocean.* 108. doi: [10.1029/2002jc001397](https://doi.org/10.1029/2002jc001397)
- Kriticos DJ, Brunel S, Ota N, Fried G, Oude Lansink AGJM et al. 2015 Downscaling pest risk analyses: Identifying current and future potentially suitable habitats for *Parthenium hysterophorus* with particular reference to Europe and North Africa. *PLoS One* 10, e0132807. doi: [10.1371/journal.pone.0132807](https://doi.org/10.1371/journal.pone.0132807)
- Kuglitsch FG, Toreti A, Xoplaki E, Della-Marta PM, Zerefos CS et al. 2010 Heat wave changes in the eastern Mediterranean since 1960. *Geophys. Res. Lett.* 37, 1–5. doi: [10.1029/2009GL041841](https://doi.org/10.1029/2009GL041841)
- Kuper M, Ameer F, Hammani A 2017 *Unraveling the enduring paradox of increased pressure on groundwater through efficient drip irrigation*. [www.routledge.com/Drip-Irrigation-for-Agriculture-Untold-Stories-of-Efficiency-Innovation/Venot-Kuper-Zwarteveen/p/book/9781138687073](http://www.routledge.com/Drip-Irrigation-for-Agriculture-Untold-Stories-of-Efficiency-Innovation/Venot-Kuper-Zwarteveen/p/book/9781138687073)
- Laabir M, Collos Y, Masseret E, Grzebyk D, Abadie E et al. 2013 Influence of environmental factors on the paralytic shellfish toxin content and profile of *Alexandrium catenella* (Dinophyceae) Isolated from the Mediterranean Sea. *Mar. Drugs* 11, 1583–1601. doi: [10.3390/md11051583](https://doi.org/10.3390/md11051583)
- Lacombe H 1990 A model of the world ocean, water, salt heat, and wind in the Mediterranean. *Oceanus* 33, 26–36.
- Lamb PJ, Pepler RA 1987 North Atlantic Oscillation: Concept and an Application. *Bull. Am. Meteorol. Soc.* 68, 1218–1225. doi: [10.1175/1520-0477\(1987\)068<1218:naocaa>2.0.co;2](https://doi.org/10.1175/1520-0477(1987)068<1218:naocaa>2.0.co;2)
- Lambdon PW, Hulme PE 2006 Predicting the invasion success of Mediterranean alien plants from their introduction characteristics. *Ecography (Cop.)*. 29, 853–865. doi: [10.1111/j.2006.0906-7590.04614.x](https://doi.org/10.1111/j.2006.0906-7590.04614.x)
- Lambdon PW, Pyšek P, Basnou C, Hejda M, Arianoutsou M et al. 2008 Alien flora of Europe: species diversity, temporal trends, geographical patterns and research needs. *Preslia* 80, 101–149.
- Lapiedra O, Sol D, Traveset A, Vilà M 2015 Random processes and phylogenetic loss caused by plant invasions. *Glob. Ecol. Biogeogr.* 24, 774–785. doi: [10.1111/geb.12310](https://doi.org/10.1111/geb.12310)
- Lasanta T, Arnáez J, Pascual N, Ruiz-Flaño P, Errea MP et al. 2017 Space-time process and drivers of land abandonment in Europe. *Catena* 149, 810–823. doi: [10.1016/j.catena.2016.02.024](https://doi.org/10.1016/j.catena.2016.02.024)
- Laza-Martinez A, David H, Riobo P, Miguel I, Orive E et al. 2016 Characterization of a Strain of *Fukuyoa paulensis* (Dinophyceae) from the Western Mediterranean Sea. *J. Eukaryot. Microbiol.* 63, 481–497. doi: [10.1111/jeu.12292](https://doi.org/10.1111/jeu.12292)
- Le Cozannet G, Thieblemont R, Rohmer J, Idir D, Manceau J-C et al. 2019 Low-end probabilistic sea-level projections. *Water* 11, 1507.
- Le Houérou HN 1995 *Bioclimatologie et biogéographie des steppes arides du Nord de l'Afrique: diversité biologique, développement durable et désertisation*. Montpellier: CIHEAM.
- Le Houérou HN 1996 Climate change, drought and desertification. *J. Arid Environ.* 34, 133–185.
- Leckebusch GC, Koffi B, Ulbrich U, Pinto JG, Spanghel T et al. 2006 Analysis of frequency and intensity of European winter storm events from a multi-model perspective, at synoptic and regional scales. *Clim. Res.* 31, 59–74. doi: [10.3354/cr031059](https://doi.org/10.3354/cr031059)
- Ledoux JB, Aurelle D, Bensoussan N, Marschal C, Féral JP et al. 2015 Potential for adaptive evolution at species range margins: Contrasting interactions between red coral populations and their environment in a changing ocean. *Ecol. Evol.* 5, 1178–1192. doi: [10.1002/ece3.1324](https://doi.org/10.1002/ece3.1324)
- Lehtijärvi A, Oskay F, Doğmuş Lehtijärvi HT, Aday Kaya AG, Pecori F et al. 2018 *Ceratocystis platani* is killing plane trees in Istanbul (Turkey). *For. Pathol.* 48, e12375. doi: [10.1111/efp.12375](https://doi.org/10.1111/efp.12375)
- Leiblein-Wild MC, Steinkamp J, Hickler T, Tackenberg O 2016

- Modelling the potential distribution, net primary production and phenology of common ragweed with a physiological model. *J. Biogeogr.* 43, 544–554. doi: [10.1111/jbi.12646](https://doi.org/10.1111/jbi.12646)
- Leimbach M, Kriegler E, Roming N, Schwanitz J 2017 Future growth patterns of world regions – A GDP scenario approach. *Glob. Environ. Chang.* 42, 215–225. doi: [10.1016/j.gloenvcha.2015.02.005](https://doi.org/10.1016/j.gloenvcha.2015.02.005)
- Lejeune C, Chevalloné P 2006 Brooding crustaceans in a highly fragmented habitat: The genetic structure of Mediterranean marine cave-dwelling mysid populations. *Mol. Ecol.* 15, 4123–4140. doi: [10.1111/j.1365-294x.2006.03101.x](https://doi.org/10.1111/j.1365-294x.2006.03101.x)
- Lekkas T, Kolokythas G, Nikolaou A, Kostopoulou M, Kotrikla A et al. 2004 Evaluation of the pollution of the surface waters of Greece from the priority compounds of list II, 76/464/EEC directive, and other toxic compounds. *Environ. Int.* 30, 995–1007. doi: [10.1016/j.envint.2004.04.001](https://doi.org/10.1016/j.envint.2004.04.001)
- Lelieveld J 2002 Global Air Pollution Crossroads over the Mediterranean. *Science (80-. ).* 298, 794–799. doi: [10.1126/science.1075457](https://doi.org/10.1126/science.1075457)
- Lelieveld J, Hadjinicolaou P, Kostopoulou E, Chenoweth J, El Maayar M et al. 2012 Climate change and impacts in the Eastern Mediterranean and the Middle East. *Clim. Change* 114, 667–687. doi: [10.1007/s10584-012-0418-4](https://doi.org/10.1007/s10584-012-0418-4)
- Lelieveld J, Hadjinicolaou P, Kostopoulou E, Giannakopoulos C, Pozzer A et al. 2014 Model projected heat extremes and air pollution in the eastern Mediterranean and Middle East in the twenty-first century. *Reg. Environ. Chang.* 14, 1937–1949. doi: [10.1007/s10113-013-0444-4](https://doi.org/10.1007/s10113-013-0444-4)
- Lelieveld J, Proestos Y, Hadjinicolaou P, Tanarhte M, Tyrlis E et al. 2016 Strongly increasing heat extremes in the Middle East and North Africa (MENA) in the 21<sup>st</sup> century. *Clim. Change* 137, 245–260. doi: [10.1007/s10584-016-1665-6](https://doi.org/10.1007/s10584-016-1665-6)
- Lenderink G, Belušić D, Fowler HJ, Kjellström E, Lind P et al. 2019 Systematic increases in the thermodynamic response of hourly precipitation extremes in an idealized warming experiment with a convection-permitting climate model. *Environ. Res. Lett.* 14, 074012. doi: [10.1088/1748-9326/ab214a](https://doi.org/10.1088/1748-9326/ab214a)
- Lesieur V, Lombaert E, Guillemaud T, Courtial B, Strong W et al. 2019 The rapid spread of *Leptoglossus occidentalis* in Europe: a bridgehead invasion. *J. Pest Sci. (2004).* 92, 189–200. doi: [10.1007/s10340-018-0993-x](https://doi.org/10.1007/s10340-018-0993-x)
- Lévesque B, Gervais M-C, Chevalier P, Gauvin D, Anasour-Laouan-Sidi E et al. 2014 Prospective study of acute health effects in relation to exposure to cyanobacteria. *Sci. Total Environ.* 466–467, 397–403. doi: [10.1016/j.scitotenv.2013.07.045](https://doi.org/10.1016/j.scitotenv.2013.07.045)
- Levin Z, Teller A, Ganor E, Graham B, Andreae MO et al. 2003 Role of aerosol size and composition in nucleation scavenging within clouds in a shallow cold front. *JGR Atmos.* 108, 4700. doi: [10.1029/2003JD003647](https://doi.org/10.1029/2003JD003647)
- Li J, Carlson BE, Dubovnik O, Lacis AA 2014 Recent trends in aerosol optical properties derived from AERONET measurements. *Atmos. Chem. Phys.* 14, 12271–12289. doi: [10.5194/acp-14-12271-2014](https://doi.org/10.5194/acp-14-12271-2014)
- Licciardello F, Antoci ML, Brugaletta L, Cirelli GL 2011 Evaluation of groundwater contamination in a coastal area of south-eastern Sicily. *J. Environ. Sci. Heal. Part. B Pestic. Food Contam. Agric. Wastes* 46, 498–508. doi: [10.1080/03601234.2011.583870](https://doi.org/10.1080/03601234.2011.583870)
- Liebhold AM, Yamanaka T, Roques A, Augustin S, Chown SL et al. 2016 Global compositional variation among native and non-native regional insect assemblages emphasizes the importance of pathways. *Biol. Invasions* 18, 893–905. doi: [10.1007/s10530-016-1079-4](https://doi.org/10.1007/s10530-016-1079-4)
- Liebig MA, Tanaka DL, Wienhold BJ 2004 Tillage and cropping effects on soil quality indicators in the northern Great Plains. *Soil Tillage Res.* 78, 131–141. doi: [10.1016/j.still.2004.02.002](https://doi.org/10.1016/j.still.2004.02.002)
- Lillo F, Faraone FP, Lo Valvo M 2011 Can the introduction of *Xenopus laevis* affect native amphibian populations? Reduction of reproductive occurrence in presence of the invasive species. *Biol. Invasions* 13, 1533–1541. doi: [10.1007/s10530-010-9911-8](https://doi.org/10.1007/s10530-010-9911-8)
- Lionello P 2012 *The Climate of the Mediterranean Region: From the Past to the Future*. Elsevier Science. doi: [10.1016/C2011-0-06210-5](https://doi.org/10.1016/C2011-0-06210-5)
- Lionello P, Abrantes FG, Congedi L, Dulac F, Gačić M et al. 2012a Introduction: Mediterranean climate-background information, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Elsevier Science). doi: [10.1016/B978-0-12-416042-2.00012-4](https://doi.org/10.1016/B978-0-12-416042-2.00012-4)
- Lionello P, Cavaleri L, Nissen KM, Pino C, Raicich F et al. 2012b Severe marine storms in the Northern Adriatic: characteristics and trends. *Phys. Chem. Earth, Parts A/B/C* 40–41, 93–105. doi: [10.1016/j.pce.2010.10.002](https://doi.org/10.1016/j.pce.2010.10.002)
- Lionello P, Cogo S, Galati MB, Sanna A 2008 The Mediterranean surface wave climate inferred from future scenario simulations. *Glob. Planet. Change* 63, 152–162. doi: [10.1016/j.gloplacha.2008.03.004](https://doi.org/10.1016/j.gloplacha.2008.03.004)
- Lionello P, Conte D, Marzo L, Scarascia L 2017 The contrasting effect of increasing mean sea level and decreasing storminess on the maximum water level during storms along the coast of the Mediterranean Sea in the mid 21<sup>st</sup> century. *Glob. Planet. Change* 151, 80–91. doi: [10.1016/j.gloplacha.2016.06.012](https://doi.org/10.1016/j.gloplacha.2016.06.012)
- Lionello P, Dalan F, Elvini E 2002 Cyclones in the Mediterranean region: the present and the doubled CO<sub>2</sub> climate scenarios. *Clim. Res.* 22, 147–159. doi: [10.3354/cr022147](https://doi.org/10.3354/cr022147)
- Lionello P, Galati MB 2008 Links of the significant wave height distribution in the Mediterranean sea with the Northern Hemisphere teleconnection patterns. *Adv. Geosci.* 17, 13–18. doi: [10.5194/adgeo-17-13-2008](https://doi.org/10.5194/adgeo-17-13-2008)
- Lionello P, Giorgi F 2007 Winter precipitation and cyclones in the Mediterranean region: Future climate scenarios in a regional simulation. *Adv. Geosci.* 12, 153–158. doi: [10.5194/adgeo-12-153-2007](https://doi.org/10.5194/adgeo-12-153-2007)
- Lionello P, Malanotte-Rizzoli P, Boscolo R, Alpert P, Artale V et al. 2006 The Mediterranean climate: An overview of the main characteristics and issues, in *Mediterranean climate variability*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo R (Elsevier), 1–26. doi: [10.1016/s1571-9197\(06\)80003-0](https://doi.org/10.1016/s1571-9197(06)80003-0)
- Lionello P, Sanna A 2005 Mediterranean wave climate variability and its links with NAO and Indian Monsoon. *Clim. Dyn.* 25, 611–623. doi: [10.1007/s00382-005-0025-4](https://doi.org/10.1007/s00382-005-0025-4)

- Lionello P, Scarascia L 2018 The relation between climate change in the Mediterranean region and global warming. *Reg. Environ. Chang.* 18, 1481–1493. doi: [10.1007/s10113-018-1290-1](https://doi.org/10.1007/s10113-018-1290-1)
- Lionello P, Scarascia L 2020 The relation of climate extremes with global warming in the Mediterranean region and its north versus south contrast. *Reg. Environ. Chang.* 20, 31. doi: [10.1007/s10113-020-01610-z](https://doi.org/10.1007/s10113-020-01610-z)
- Lionello P, Trigo IF, Gil V, Liberato MLR, Nissen KM et al. 2016 Objective climatology of cyclones in the Mediterranean region: A consensus view among methods with different system identification and tracking criteria. *Tellus A Dyn. Meteorol. Oceanogr.* 68, 29391. doi: [10.3402/tellusa.v68.29391](https://doi.org/10.3402/tellusa.v68.29391)
- Lioy S, Manino A, Porporato M, Laurino D, Romano A et al. 2019 Establishing surveillance areas for tackling the invasion of *Vespa velutina* in outbreaks and over the border of its expanding range. *NeoBiota* 46, 51–69. doi: [10.3897/neobiota.46.33099](https://doi.org/10.3897/neobiota.46.33099)
- Lippmann M, Chen L-C 2009 Health effects of concentrated ambient air particulate matter (CAPs) and its components. *Crit. Rev. Toxicol.* 39, 865–913. doi: [10.3109/10408440903300080](https://doi.org/10.3109/10408440903300080)
- Liubartseva S, Coppini G, Lecci R, Clementi E 2018 Tracking plastics in the Mediterranean: 2D Lagrangian model. *Mar. Pollut. Bull.* 129, 151–162. doi: [10.1016/j.marpolbul.2018.02.019](https://doi.org/10.1016/j.marpolbul.2018.02.019)
- Lloret F, Médail F, Brundu G, Camarda I, Moragues E et al. 2005 Species attributes and invasion success by alien plants on Mediterranean islands. *J. Ecol.* 93, 512–520. doi: [10.1111/j.1365-2745.2005.00979.x](https://doi.org/10.1111/j.1365-2745.2005.00979.x)
- Lloret F, Médail F, Brundu G, Hulme PE 2004a Local and regional abundance of exotic plant species on Mediterranean islands: are species traits important? *Glob. Ecol. Biogeogr.* 13, 37–45. doi: [10.1111/J.1466-882x.2004.00064.X](https://doi.org/10.1111/J.1466-882x.2004.00064.X)
- Lloret J, Palomera I, Salat J, Sole I 2004b Impact of freshwater input and wind on landings of anchovy (*Engraulis encrasicolus*) and sardine (*Sardina pilchardus*) in shelf waters surrounding the Ebro (Ebro) River delta (north-western Mediterranean). *Fish. Oceanogr.* 13, 102–110. doi: [10.1046/j.1365-2419.2003.00279.x](https://doi.org/10.1046/j.1365-2419.2003.00279.x)
- Loepfe L, Martinez-Vilalta J, Oliveres J, Piñol J, Lloret F 2010 Feedbacks between fuel reduction and landscape homogenisation determine fire regimes in three Mediterranean areas. *For. Ecol. Manage.* 259, 2366–2374. doi: [10.1016/j.foreco.2010.03.009](https://doi.org/10.1016/j.foreco.2010.03.009)
- Loepfe L, Martinez-Vilalta J, Piñol J 2011 An integrative model of human-influenced fire regimes and landscape dynamics. *Environ. Model. Softw.* 26, 1028–1040. doi: [10.1016/j.envsoft.2011.02.015](https://doi.org/10.1016/j.envsoft.2011.02.015)
- López-Parages J, Rodríguez-Fonseca B 2012 Multidecadal modulation of El Niño influence on the Euro-Mediterranean rainfall. *Geophys. Res. Lett.* 39. doi: [10.1029/2011GL050049](https://doi.org/10.1029/2011GL050049)
- Lopez B, Ollivier P, Togola A, Baran N, Ghestem J-P 2015 Screening of French groundwater for regulated and emerging contaminants. *Sci. Total Environ.* 518–519, 562–573. doi: [10.1016/j.scitotenv.2015.01.110](https://doi.org/10.1016/j.scitotenv.2015.01.110)
- Löptien U, Zolina O, Gulev S, Latif M, Soloviev V 2008 Cyclone life cycle characteristics over the Northern Hemisphere in coupled GCMs. *Clim. Dyn.* 31, 507–532. doi: [10.1007/s00382-007-0355-5](https://doi.org/10.1007/s00382-007-0355-5)
- Louchart X, Voltz M, Andrieux P, Moussa R 2001 Herbicide transport to surface waters at field and watershed scales in a Mediterranean vineyard area. *J. Environ. Qual.* 30, 982–991. <http://www.ncbi.nlm.nih.gov/pubmed/11401289>
- Loumou A, Giourga C 2003 Olive groves: “The life and identity of the Mediterranean.” *Agric. Human Values* 20, 87–95. doi: [10.1023/a:1022444005336](https://doi.org/10.1023/a:1022444005336)
- Lounibos LP 2002 Invasions by Insect Vectors of Human Disease. *Annu. Rev. Entomol.* 47, 233–266. doi: [10.1146/annurev.ento.47.091201.145206](https://doi.org/10.1146/annurev.ento.47.091201.145206)
- Lowenthal DH, Gertler AW, Labib MW 2014 Particulate matter source apportionment in Cairo: recent measurements and comparison with previous studies. *Int. J. Environ. Sci. Technol.* 11, 657–670. doi: [10.1007/s13762-013-0272-6](https://doi.org/10.1007/s13762-013-0272-6)
- Lu J, Vecchi GA, Reichler T 2007a Correction to “Expansion of the Hadley cell under global warming.” *Geophys. Res. Lett.* 34. doi: [10.1029/2007gl030931](https://doi.org/10.1029/2007gl030931)
- Lu J, Vecchi GA, Reichler T 2007b Expansion of the Hadley cell under global warming. *Geophys. Res. Lett.* 34, L06805. doi: [10.1029/2006GL028443](https://doi.org/10.1029/2006GL028443)
- Ludwig W, Dumont E, Meybeck M, Heussner S 2009 River discharges of water and nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during past and future decades? *Prog. Oceanogr.* 80, 199–217. doi: [10.1016/j.pocean.2009.02.001](https://doi.org/10.1016/j.pocean.2009.02.001)
- Luterbacher J, Werner JP, Smerdon JE 2016 European summer temperatures since Roman times. *Environ. Res. Lett.* 11, 2. doi: [10.1088/1748-9326/11/2/024001](https://doi.org/10.1088/1748-9326/11/2/024001)
- Luterbacher J, Xoplaki E 2003 500-year Winter Temperature and Precipitation Variability over the Mediterranean area and its Connection to the Large-scale Atmospheric Circulation, in *Mediterranean Climate. Variability and Trends*, ed. Bolle HJ (Berlin: Springer Verlag), 133–153.
- Lutz W, Amran G, Bélanger A, Conte A, Gailey N et al. 2019 Demographic Scenarios for the EU - Migration, Population and Education. EUR 29739 EN, Publications Office, Luxembourg doi: [10.2760/590301](https://doi.org/10.2760/590301)
- M'Rabet C, Kéfi-Daly YO, Couet D, Gueroun SKM, Pringault O 2019 Consequences of a contaminant mixture of bisphenol A (BPA) and di-[2-ethylhexyl] phthalate (DEHP), two plastic-derived chemicals, on the diversity of coastal phytoplankton. *Mar. Pollut. Bull.* 138, 385–396. doi: [10.1016/j.marpolbul.2018.11.035](https://doi.org/10.1016/j.marpolbul.2018.11.035)
- Macías DM, García-Gorriç E, Stips A 2013 Understanding the causes of recent warming of Mediterranean waters. How much could be attributed to climate change? *PLoS One* 8, e81591. doi: [10.1371/journal.pone.0081591](https://doi.org/10.1371/journal.pone.0081591)
- Macías DM, García-Gorriç E, Stips A 2018 Deep winter convection and phytoplankton dynamics in the NW Mediterranean Sea under present climate and future (horizon 2030) scenarios. *Sci. Rep.* 8, 6626. doi: [10.1038/s41598-018-24965-0](https://doi.org/10.1038/s41598-018-24965-0)
- Maggi V, Colucci RR, Scoto F, Giudice G, Randazzo L 2018 Chapter 19 - Ice Caves in Italy, in, eds. Perçoiu A, Lauritzen S-EBT-IC (Elsevier), 399–423.



- doi: [10.1016/B978-0-12-811739-2.00019-X](https://doi.org/10.1016/B978-0-12-811739-2.00019-X)
- Magnusson M, Heimann K, Quayle P, Negri AP 2010 Additive toxicity of herbicide mixtures and comparative sensitivity of tropical benthic microalgae. *Mar. Pollut. Bull.* 60, 1978–1987. doi: [10.1016/j.marpolbul.2010.07.031](https://doi.org/10.1016/j.marpolbul.2010.07.031)
- Maheras P, Flocas HA, Patrikas I, Anagnostopoulou C 2001 A 40 year objective climatology of surface cyclones in the Mediterranean region: spatial and temporal distribution. *Int. J. Climatol.* 21, 109–130. doi: [10.1002/joc.599](https://doi.org/10.1002/joc.599)
- Mahmoud KF, Alfaro SC, Favez O, Abdel Wahab MM, Sciare J 2008 Origin of black carbon concentration peaks in Cairo (Egypt). *Atmos. Res.* 89, 161–169. doi: [10.1016/J.ATMOSRES.2008.01.004](https://doi.org/10.1016/J.ATMOSRES.2008.01.004)
- Malagó A, Bouraoui F, Grizzetti B, De Roo A 2019 Modelling nutrient fluxes into the Mediterranean Sea. *J. Hydrol. Reg. Stud.* 22, 100592. doi: [10.1016/j.ejrh.2019.01.004](https://doi.org/10.1016/j.ejrh.2019.01.004)
- Malek Ž, Verburg PH 2018 Adaptation of land management in the Mediterranean under scenarios of irrigation water use and availability. *Mitig. Adapt. Strateg. Glob. Chang.* 23, 821–837. doi: [10.1007/s11027-017-9761-0](https://doi.org/10.1007/s11027-017-9761-0)
- Manachini B, Billeci N, Palla F 2013 Exotic insect pests: The impact of the Red Palm Weevil on natural and cultural heritage in Palermo (Italy). *J. Cult. Herit.* 14, e177–e182. doi: [10.1016/j.culher.2012.11.028](https://doi.org/10.1016/j.culher.2012.11.028)
- Manara V, Brunetti M, Celozzi M, Maugeri M, Sánchez-Lorenzo A et al. 2016 Detection of dimming / brightening in Italy from homogenized all-sky and clear-sky surface solar radiation records and underlying causes (1959–2013). *Atmos. Chem. Phys.* 16, 11145–11161. doi: [10.5194/acp-16-11145-2016](https://doi.org/10.5194/acp-16-11145-2016)
- Mandarić L, Kalogianni E, Skoulikidis N, Petrovic M, Sabater S 2019 Contamination patterns and attenuation of pharmaceuticals in a temporary Mediterranean river. *Sci. Total Environ.* 647, 561–569. doi: [10.1016/j.scitotenv.2018.07.308](https://doi.org/10.1016/j.scitotenv.2018.07.308)
- Manodori L, Gambaro A, Piazza R, Ferrari S, Stortini AM et al. 2006 PCBs and PAHs in sea-surface microlayer and sub-surface water samples of the Venice Lagoon (Italy). *Mar. Pollut. Bull.* 52, 184–192. doi: [10.1016/j.marpolbul.2005.08.017](https://doi.org/10.1016/j.marpolbul.2005.08.017)
- Mansui J, Molcard A, Ourmières Y 2015 Modelling the transport and accumulation of floating marine debris in the Mediterranean basin. *Mar. Pollut. Bull.* 91, 249–257. doi: [10.1016/j.marpolbul.2014.11.037](https://doi.org/10.1016/j.marpolbul.2014.11.037)
- Manzano S, Carrión JS, García-Murillo P, López-Merino L 2019 When dynamism is the baseline: long-term ecology of a Mediterranean seasonal wetland in the Doñana National Park (Southwestern Europe). *Biodivers. Conserv.* 28, 501–522. doi: [10.1007/s10531-018-1674-z](https://doi.org/10.1007/s10531-018-1674-z)
- Maraun D, Widmann M, Gutiérrez JM 2019 Statistical downscaling skill under present climate conditions: A synthesis of the VALUE perfect predictor experiment. *Int. J. Climatol.* 39, 3692–3703. doi: [10.1002/joc.5877](https://doi.org/10.1002/joc.5877)
- Marba N, Duarte CM 2010 Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality. *Glob. Chang. Biol.* 16, 2366–2375.
- Marcelli M, Dayan AR, Langeneck J 2017 Finding Dory: first record of *Paracanthurus hepatus* (Perciformes: Acanthuriidae) in the Mediterranean Sea. *Mar. Biodivers.* 47, 599–602. doi: [10.1007/s12526-016-0573-3](https://doi.org/10.1007/s12526-016-0573-3)
- Marcos M, Calafat FM, Berihuete A, Dangendorf S 2015 Long-term variations in global sea level extremes. *JGR Ocean.* 120, 8115–8134. doi: [10.1002/2015JC011173](https://doi.org/10.1002/2015JC011173)
- Marcos M, Jordà G, Gomis D, Pérez Gómez B 2011 Changes in storm surges in southern Europe from a regional model under climate change scenarios. *Glob. Planet. Change* 77, 116–128. <https://www.sciencedirect.com/science/article/pii/S0921818111000555> [Accessed September 17, 2019]
- Marcos M, Tsimplis MN 2008 Comparison of results of AOGCMs in the Mediterranean Sea during the 21<sup>st</sup> century. *JGR Ocean.* 113. doi: [10.1029/2008JC004820](https://doi.org/10.1029/2008JC004820)
- Marcos M, Tsimplis MN, Shaw AGP 2009 Sea level extremes in southern Europe. *JGR Ocean.* 114, C01007. doi: [10.1029/2008JC004912](https://doi.org/10.1029/2008JC004912)
- Mariotti A 2010 Recent changes in the Mediterranean water cycle: A pathway toward long-term regional hydroclimatic change? *J. Clim.* 23, 1513–1525. doi: [10.1175/2009JCLI3251.1](https://doi.org/10.1175/2009JCLI3251.1)
- Mariotti A, Ballabrera-Poy J, Zeng N 2005 Tropical influence on Euro-Asian autumn rainfall variability. *Clim. Dyn.* 24, 511–521.
- Mariotti A, Dell'Aquila A 2012 Decadal climate variability in the Mediterranean region: roles of large-scale forcings and regional processes. *Clim. Dyn.* 38, 1129–1145. doi: [10.1007/s00382-011-1056-7](https://doi.org/10.1007/s00382-011-1056-7)
- Mariotti A, Pan Y, Zeng N, Alessandri A 2015 Long-term climate change in the Mediterranean region in the midst of decadal variability. *Clim. Dyn.* 44, 1437–1456. doi: [10.1007/s00382-015-2487-3](https://doi.org/10.1007/s00382-015-2487-3)
- Mariotti A, Struglia MV, Zeng N, Lau K-M 2002a The Hydrological Cycle in the Mediterranean Region and Implications for the Water Budget of the Mediterranean Sea. *J. Clim.* 15, 1674–1690. doi: [10.1175/1520-0442\(2002\)015<1674:THCITM>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1674:THCITM>2.0.CO;2)
- Mariotti A, Zeng N, Lau KM 2002b Euro-Mediterranean rainfall and ENSO—a seasonally varying relationship. *Geophys. Res. Lett.* 29, 1621. doi: [10.1029/2001GL014248](https://doi.org/10.1029/2001GL014248)
- Mariotti A, Zeng N, Yoon J-H, Artale V, Navarra A et al. 2008 Mediterranean water cycle changes: transition to drier 21<sup>st</sup> century conditions in observations and CMIP3 simulations. *Environ. Res. Lett.* 3, 044001. doi: [10.1088/1748-9326/3/4/044001](https://doi.org/10.1088/1748-9326/3/4/044001)
- Marmy A, Rajczak J, Delaloye R, Hilbich C, Hoelzle M et al. 2016 Semi-automated calibration method for modelling of mountain permafrost evolution in Switzerland. *Cryosph.* 10, 2693–2719. doi: [10.5194/tc-10-2693-2016](https://doi.org/10.5194/tc-10-2693-2016)
- Marty J-C, Chiavérini J 2010 Hydrological changes in the Ligurian Sea (NW Mediterranean, DYFAMED site) during 1995–2007 and biogeochemical consequences. *Biogeosciences* 7, 2117–2128. doi: [10.5194/bg-7-2117-2010](https://doi.org/10.5194/bg-7-2117-2010)
- Marullo S, Artale V, Santoleri R 2011 The SST Multidecadal Variability in the Atlantic–Mediterranean Region and Its Relation to AMO. *J. Clim.* 24, 4385–4401. doi: [10.1175/2011JCLI3884.1](https://doi.org/10.1175/2011JCLI3884.1)
- Marullo S, Santoleri R, Banzon V, Evans RH, Guarracino M 2010 A diurnal-cycle resolving sea surface temperature product for the tropical Atlantic. *JGR Atmos.* 115, C05011.

- doi: [10.1029/2009JC005466](https://doi.org/10.1029/2009JC005466)
- Marvier M, Kareiva P, Neubert MG 2004 Habitat Destruction, Fragmentation, and Disturbance Promote Invasion by Habitat Generalists in a Multispecies Metapopulation. *Risk Anal.* 24, 869–878. doi: [10.1111/j.0272-4332.2004.00485.x](https://doi.org/10.1111/j.0272-4332.2004.00485.x)
- Matesanz S, Escudero A, Valladares F 2008 Additive effects of a potentially invasive grass and water stress on the performance of seedlings of gypsum specialists. *Appl. Veg. Sci.* 11, 287–296. doi: [10.3170/2008-7-18425](https://doi.org/10.3170/2008-7-18425)
- Matošević D, Pajač Živković I 2013 Alien phytophagous insect and mite species on woody plants in Croatia (in croatian). *Šumarski List*, 191–200.
- Matzrafi M 2019 Climate change exacerbates pest damage through reduced pesticide efficacy. *Pest Manag. Sci.* 75, 9–13. doi: [10.1002/ps.5121](https://doi.org/10.1002/ps.5121)
- McInnes KL, Erwin TA, Bathols JM 2011 Global Climate Model projected changes in 10 m wind speed and direction due to anthropogenic climate change. *Atmos. Sci. Lett.* 12, 325–333. doi: [10.1002/asl.341](https://doi.org/10.1002/asl.341)
- MCRIT 2015 ET2050. Territorial Scenarios and Visions for Europe. Volume 1 - Approach to Scenario Building and Storylines. [https://www.espon.eu/sites/default/files/attachments/ET2050\\_FR-03\\_Volume\\_1\\_-\\_Approach\\_to\\_Scenario\\_Building\\_and\\_Storylines.pdf](https://www.espon.eu/sites/default/files/attachments/ET2050_FR-03_Volume_1_-_Approach_to_Scenario_Building_and_Storylines.pdf)
- MEA 2005 *Ecosystems and human well-being: synthesis*. Island Press, Washington, DC <https://www.cifor.org/library/1888/>
- Mediterranean Wetlands Observatory 2018 *Mediterranean Wetland Outlook 2: Solutions for sustainable Mediterranean Wetlands*. , eds. Geijzendorffer IR, Chazée L, Gaget E, Galewski T, Guelmami A et al. Arles, France <https://tourduvalat.org/en/actions/les-zones-humides-mediterraneennes-en-jeux-et-perspectives-2-solutions-pour-des-zones-humides-mediterraneennes-durables/>
- Meffe R, de Bustamante I 2014 Emerging organic contaminants in surface water and groundwater: A first overview of the situation in Italy. *Sci. Total Environ.* 481, 280–295. doi: [10.1016/j.scitotenv.2014.02.053](https://doi.org/10.1016/j.scitotenv.2014.02.053)
- Menchetti M, Mori E 2014 Worldwide impact of alien parrots (Aves Psittaciformes) on native biodiversity and environment: a review. *Ethol. Ecol. Evol.* 26, 172–194. doi: [10.1080/03949370.2014.905981](https://doi.org/10.1080/03949370.2014.905981)
- Menchetti M, Mori E, Angelici FM 2016 Effects of the recent world invasion by ring-necked parakeets *Psittacula krameri*, in *Problematic Wildlife*, ed. Angelici FM (Springer International Publishing), 253–266. doi: [10.1007/978-3-319-22246-2\\_12](https://doi.org/10.1007/978-3-319-22246-2_12)
- Mendel Z, Branco M, Battisti A 2016 Invasive Sap-Sucker Insects in the Mediterranean Basin. *Insects Dis. Mediterr. For. Syst.*, 261–291. doi: [10.1007/978-3-319-24744-1\\_10](https://doi.org/10.1007/978-3-319-24744-1_10)
- Menemenlis D, Fukumori I, Lee T 2007 Atlantic to Mediterranean Sea Level Difference Driven by Winds near Gibraltar Strait. *J. Phys. Oceanogr.* 37, 359–376. doi: [10.1175/JPO3015.1](https://doi.org/10.1175/JPO3015.1)
- Menendez M, García-Díez M, Fita L, Fernández J, Méndez FJ et al. 2014 High-resolution sea wind hindcasts over the Mediterranean area. *Clim. Dyn.* 42, 1857–1872. doi: [10.1007/s00382-013-1912-8](https://doi.org/10.1007/s00382-013-1912-8)
- Merhaby D, Net S, Halwani J, Ouddane B 2015 Organic pollution in surficial sediments of Tripoli harbour, Lebanon. *Mar. Pollut. Bull.* 93, 284–293. doi: [10.1016/j.marpolbul.2015.01.004](https://doi.org/10.1016/j.marpolbul.2015.01.004)
- Merhaby D, Ouddane B, Net S, Halwani J 2018 Assessment of trace metals contamination in surficial sediments along Lebanese Coastal Zone. *Mar. Pollut. Bull.* 133, 881–890. doi: [10.1016/j.marpolbul.2018.06.031](https://doi.org/10.1016/j.marpolbul.2018.06.031)
- Merkens J-L, Reimann L, Hinkel J, Vafeidis AT 2016 Gridded population projections for the coastal zone under the Shared Socioeconomic Pathways. *Glob. Planet. Change* 145, 57–66. doi: [10.1016/j.gloplacha.2016.08.009](https://doi.org/10.1016/j.gloplacha.2016.08.009)
- Meurisse N, Rassati D, Hurley BP, Brockerhoff EG, Haack RA 2019 Common pathways by which non-native forest insects move internationally and domestically. *J. Pest Sci. (2004)*. 92, 13–27. doi: [10.1007/s10340-018-0990-0](https://doi.org/10.1007/s10340-018-0990-0)
- Miglietta MM 2019 Mediterranean Tropical-Like Cyclones (Medicanes). *Atmosphere (Basel)*. 10, 206. doi: [10.3390/atmos10040206](https://doi.org/10.3390/atmos10040206)
- Miglietta MM, Laviola S, Malvaldi A, Conte D, Levizzani V et al. 2013 Analysis of tropical-like cyclones over the Mediterranean Sea through a combined modeling and satellite approach. *Geophys. Res. Lett.* 40, 2400–2405. doi: [10.1002/grl.50432](https://doi.org/10.1002/grl.50432)
- Miglietta MM, Mastrangelo E, Conte D 2015 Influence of physics parameterization schemes on the simulation of a tropical-like cyclone in the Mediterranean Sea. *Atmos. Res.* 153, 360–375. doi: [10.1016/j.atmosres.2014.09.008](https://doi.org/10.1016/j.atmosres.2014.09.008)
- Millán MM, Mantilla E, Salvador R, Carratalá A, Sanz MJ et al. 2000 Ozone Cycles in the Western Mediterranean Basin: Interpretation of Monitoring Data in Complex Coastal Terrain. *J. Appl. Meteorol.* 39, 487–508. doi: [10.1175/1520-0450\(2000\)039<0487:OCITWM>2.0.CO;2](https://doi.org/10.1175/1520-0450(2000)039<0487:OCITWM>2.0.CO;2)
- Millot C, Candela J, Fuda J-L, Tber Y 2006 Large warming and salinification of the Mediterranean outflow due to changes in its composition. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 53, 656–666. doi: [10.1016/J.DSR.2005.12.017](https://doi.org/10.1016/J.DSR.2005.12.017)
- Mitsakou C, Kallos G, Papantoniou N, Spyrou C, Solomos S et al. 2008 Saharan dust levels in Greece and received inhalation doses. *Atmos. Chem. Phys.* 8, 7181–7192. doi: [10.5194/acp-8-7181-2008](https://doi.org/10.5194/acp-8-7181-2008)
- Moemken J, Reyers M, Feldmann H, Pinto JG 2018 Future Changes of Wind Speed and Wind Energy Potentials in EURO-CORDEX Ensemble Simulations. *JGR Atmos.* 123, 6373–6389. doi: [10.1029/2018JD028473](https://doi.org/10.1029/2018JD028473)
- Mohamed AH, Squires VR 2018 Drylands of the Mediterranean Basin: Challenges, Problems and Prospects, in *Climate Variability Impacts on Land Use and Livelihoods in Drylands*, eds. Gaur M, Squires V (Springer, Cham).
- Molle F, Tanouti O 2017 Squaring the circle: Agricultural intensification vs. Water conservation in Morocco. *Agric. Water Manag.* 192, 170–179. doi: [10.1016/j.agwat.2017.07.009](https://doi.org/10.1016/j.agwat.2017.07.009)
- Monceau K, Bonnard O, Thiéry D 2014 *Vespa velutina*: a new invasive predator of honeybees in Europe. *J. Pest Sci. (2004)*. 87, 1–16. doi: [10.1007/s10340-013-0537-3](https://doi.org/10.1007/s10340-013-0537-3)
- Moncer M, Hamza A, Feki-Sahnoun W, Mabrouk L, Hassen MB 2017 Variability patterns of epibenthic microalgae in eastern Tunisian coasts. *Sci. Mar.* 81, 487–498. doi: [10.3989/scimar.04651.17A](https://doi.org/10.3989/scimar.04651.17A)
- Montecchio L, Fanchin G, Simonato M, Faccoli M 2014 First

- record of thousand cankers disease fungal pathogen *Geosmithia morbida* and Walnut Twig Beetle *Pityophthorus juglandis* on *Juglans regia* in Europe. *Plant Dis.* 98, 1445. doi: [10.1094/pdis-07-14-0719-pdn](https://doi.org/10.1094/pdis-07-14-0719-pdn)
- Monteiro A, Russo M, Gama C, Borrego C 2018a How important are maritime emissions for the air quality: At European and national scale. *Environ. Pollut.* 242, 565–575. doi: [10.1016/j.envpol.2018.07.011](https://doi.org/10.1016/j.envpol.2018.07.011)
- Monteiro A, Sá E, Fernandes AP, Gama C, Sorte S et al. 2018b How healthy will be the air quality in 2050? *Air Qual. Atmos. Heal.* 11, 353–362. doi: [10.1007/s11869-017-0466-z](https://doi.org/10.1007/s11869-017-0466-z)
- Mooney HA, Hobbs RJ 2000 *Invasive Species in a Changing World*. Washington DC, USA: Island Press
- Moreira F, Viedma O, Arianoutsou M, Curt T, Koutsias N et al. 2011 Landscape – wildfire interactions in southern Europe: Implications for landscape management. *J. Environ. Manage.* 92, 2389–2402. doi: [10.1016/j.jenvman.2011.06.028](https://doi.org/10.1016/j.jenvman.2011.06.028)
- Moreno-González R, Rodriguez-Mozaz S, Gros M, Barceló D, León VM 2015 Seasonal distribution of pharmaceuticals in marine water and sediment from a mediterranean coastal lagoon (SE Spain). *Environ. Res.* 138, 326–344. doi: [10.1016/j.envres.2015.02.016](https://doi.org/10.1016/j.envres.2015.02.016)
- Moreno J, Arianoutsou M, González-Cabán A, Mouillot F, Oechel W et al. 2014 Forest fires under climate, social and economic changes in Europe, the Mediterranean and other fire-affected areas of the world.
- Morice CP, Kennedy JJ, Rayner NA, Jones PD 2012 Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 data set. *JGR Atmos.* 117, D08101. doi: [10.1029/2011JD017187](https://doi.org/10.1029/2011JD017187)
- Moriondo M, Trombi G, Ferrise R, Brandani G, Dibari C et al. 2013 Olive trees as bio-indicators of climate evolution in the Mediterranean Basin. *Glob. Ecol. Biogeogr.* 22. doi: [10.1111/qeb.12061](https://doi.org/10.1111/qeb.12061)
- Moullec F, Barrier N, Drira S, Guilhaumon F, Marsaleix P et al. 2019 An End-to-End Model Reveals Losers and Winners in a Warming Mediterranean Sea. *Front. Mar. Sci.* 6. doi: [10.3389/fmars.2019.00345](https://doi.org/10.3389/fmars.2019.00345)
- Müller R-J, Kleeberg I, Deckwer W-D 2001 Biodegradation of polyesters containing aromatic constituents. *J. Biotechnol.* 86, 87–95. doi: [10.1016/S0168-1656\(00\)00407-7](https://doi.org/10.1016/S0168-1656(00)00407-7)
- Murakami D, Yamagata Y 2019 Estimation of Gridded Population and GDP Scenarios with Spatially Explicit Statistical Downscaling. *Sustainability* 11, 2106. doi: [10.3390/su11072106](https://doi.org/10.3390/su11072106)
- Mutinelli F, Montarsi F, Federico G, Granato A, Ponti AM et al. 2014 Detection of *Aethina tumida* Murray (Coleoptera: Nitidulidae) in Italy: outbreaks and early reaction measures. *J. Apic. Res.* 53, 569–575. doi: [10.3896/ibra.1.53.5.13](https://doi.org/10.3896/ibra.1.53.5.13)
- Mutke S, Calama R, Neaymeh EN, Roques A 2016 Impact of the dry cone syndrome on commercial kernel yield of stone pine cones. *Options Méditerranéennes* 122, 78–84.
- Nabat P, Somot S, Cassou C, Mallet M, Michou M et al. 2020 Modulation of radiative aerosols effects by atmospheric circulation over the Euro-Mediterranean region. *Atmos. Chem. Phys.* 20, 8315–8349.
- Nabat P, Somot S, Mallet M, Chiappello I, Morcrette JJ et al. 2013 A 4-D climatology (1979–2009) of the monthly tropospheric aerosol optical depth distribution over the Mediterranean region from a comparative evaluation and blending of remote sensing and model products. *Atmos. Meas. Tech.* 6. doi: [10.5194/amt-6-1287-2013](https://doi.org/10.5194/amt-6-1287-2013)
- Nabat P, Somot S, Mallet M, Michou M, Sevault F et al. 2015a Dust aerosol radiative effects during summer 2012 simulated with a coupled regional aerosol–atmosphere–ocean model over the Mediterranean. *Atmos. Chem. Phys.* 15, 3303–3326. doi: [10.5194/acp-15-3303-2015](https://doi.org/10.5194/acp-15-3303-2015)
- Nabat P, Somot S, Mallet M, Sánchez-Lorenzo A, Wild M 2014 Contribution of anthropogenic sulfate aerosols to the changing Euro-Mediterranean climate since 1980. *Geophys. Res. Lett.* 41, 5605–5611. doi: [10.1002/2014GL060798](https://doi.org/10.1002/2014GL060798)
- Nabat P, Somot S, Mallet M, Sevault F, Chiacchio M et al. 2015b Direct and semi-direct aerosol radiative effect on the Mediterranean climate variability using a coupled regional climate system model. *Clim. Dyn.* 44, 1127–1155. doi: [10.1007/s00382-014-2205-6](https://doi.org/10.1007/s00382-014-2205-6)
- Naidja L, Ali-Khodja H, Khaldi S 2018 Sources and levels of particulate matter in North African and Sub-Saharan cities: a literature review. *Environ. Sci. Pollut. Res.* 25, 12303–12328. doi: [10.1007/s11356-018-1715-x](https://doi.org/10.1007/s11356-018-1715-x)
- Nakićenović N 2000 *Special report on emissions scenarios: A special report of Working Group III of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press
- Naranjo C, García-Lafuente J, Sammartino S, Sánchez-Garrido JC, Sánchez-Leal R et al. 2017 Recent changes (2004–2016) of temperature and salinity in the Mediterranean outflow. *Geophys. Res. Lett.* 44, 5665–5672. doi: [10.1002/2017GL072615](https://doi.org/10.1002/2017GL072615)
- Nashwan MS, Shahid S, Abd Rahim N 2018 Unidirectional trends in annual and seasonal climate and extremes in Egypt. *Theor. Appl. Climatol.* 136, 457–473. doi: [10.1007/s00704-018-2498-1](https://doi.org/10.1007/s00704-018-2498-1)
- Nasrallah HA, Balling RC 1993 Spatial and temporal analysis of Middle Eastern temperature changes. *Clim. Change* 25, 153–161. doi: [10.1007/bf01661203](https://doi.org/10.1007/bf01661203)
- Nastos PT, Karavana Papadimou K, Matsangouras IT 2018 Mediterranean tropical-like cyclones: Impacts and composite daily means and anomalies of synoptic patterns. *Atmos. Res.* 208, 156–166. doi: [10.1016/J.ATMOSRES.2017.10.023](https://doi.org/10.1016/J.ATMOSRES.2017.10.023)
- Naumann G, Alfieri L, Wyser K, Mentaschi L, Betts RA et al. 2018 Global changes in drought conditions under different levels of warming. *Geophys. Res. Lett.* 45, 3285–3296. doi: [10.1002/2017GL076521](https://doi.org/10.1002/2017GL076521)
- Navarro-Pedreño J, Gómez I, Almendro-Candel MB, Meléndez-Pastor I 2008 Heavy metals in Mediterranean soils, in *Soil contamination research trends* (Nova Science Publishers Inc), 161–176. [https://www.novapublishers.com/catalog/product\\_info.php?products\\_id=6860](https://www.novapublishers.com/catalog/product_info.php?products_id=6860)
- Naves P, Bonifácio L, de Sousa E 2016 The Pine Wood Nematode and Its Local Vectors in the Mediterranean Basin. *Insects Dis. Mediterr. For. Syst.*, 329–378. doi: [10.1007/978-3-319-24744-1\\_12](https://doi.org/10.1007/978-3-319-24744-1_12)
- Nehme N, Haydar C, Koubaissy B, Fakihi M, Awad S et al. 2014 The distribution of heavy metals in the Lower River Basin,

- Lebanon. *Phys. Procedia* 55, 456–463. doi: [10.1016/j.phpro.2014.07.066](https://doi.org/10.1016/j.phpro.2014.07.066)
- Nelson GC, Bennett EM, Berhe AA, Cassman KG, DeFries RS et al. 2006 Anthropogenic drivers of ecosystem change: An overview. *Ecol. Soc.* 11, 29. doi: [10.5751/ES-01826-110229](https://doi.org/10.5751/ES-01826-110229)
- Nepstad DC, Verssimo A, Alencar A, Nobre CA, Lima E et al. 1999 Large-scale impoverishment of Amazonian forests by logging and fire. *Nature* 398, 505–508. doi: [10.1038/19066](https://doi.org/10.1038/19066)
- Nieto R, Gimeno L, Trigo RM 2006 A Lagrangian identification of major sources of Sahel moisture. *Geophys. Res. Lett.* 33. doi: [10.1029/2006GL027232](https://doi.org/10.1029/2006GL027232)
- Nikulin G, Lennard C, Dosio A, Kjellström E, Chen Y et al. 2018 The effects of 1.5 and 2 degrees of global warming on Africa in the CORDEX ensemble. *Environ. Res. Lett.* 13, 65003. doi: [10.1088/1748-9326/aab1b1](https://doi.org/10.1088/1748-9326/aab1b1)
- Nissen KM, Leckebusch GC, Pinto JG, Renggli D, Ulbrich S et al. 2010 Cyclones causing wind storms in the Mediterranean: characteristics, trends and links to large-scale patterns. *Nat. Hazards Earth Syst. Sci.* 10, 1379–1391. doi: [10.5194/nhess-10-1379-2010](https://doi.org/10.5194/nhess-10-1379-2010)
- Nissen KM, Leckebusch GC, Pinto JG, Ulbrich U 2014 Mediterranean cyclones and windstorms in a changing climate. *Reg. Environ. Chang.* 14, 1873–1890. doi: [10.1007/s10113-012-0400-8](https://doi.org/10.1007/s10113-012-0400-8)
- Nödler K, Voutsas D, Licha T 2014 Polar organic micropollutants in the coastal environment of different marine systems. *Mar. Pollut. Bull.* 85, 50–59. doi: [10.1016/j.marpolbul.2014.06.024](https://doi.org/10.1016/j.marpolbul.2014.06.024)
- Nori M 2018 Migration, agriculture and rural territories in the Mediterranean, in *Migration and Inclusive Rural Development in the Mediterranean*, ed. MediTERRA C/ (Presses de Sciences Po).
- Novoa A, Le Roux JJ, Robertson MP, Wilson JRU, Richardson DM 2015 Introduced and invasive cactus species: a global review. *AoB Plants* 7. doi: [10.1093/aobpla/plu078](https://doi.org/10.1093/aobpla/plu078)
- Noyes PD, McElwee MK, Miller HD, Clark BW, Van Tiem LA et al. 2009 The toxicology of climate change: Environmental contaminants in a warming world. *Environ. Int.* 35, 971–986. doi: [10.1016/j.envint.2009.02.006](https://doi.org/10.1016/j.envint.2009.02.006)
- Nunes L, Gower ST, Peckham SD, Magalhães M, Lopes D et al. 2015 Estimation of productivity in pine and oak forests in northern Portugal using Biome-BGC. *For. An Int. J. For. Res.* 88, 200–212. doi: [10.1093/forestry/cpu044](https://doi.org/10.1093/forestry/cpu044)
- Nyingi W, Oguge N, Dziba L, Chandipo R, Didier TA et al. 2018 Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for Africa*, eds. Archer E, Dziba L, Mulongoy KJ, Maoela MA, Walters M (Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), 207–296.
- Nykjaer L 2009 Mediterranean Sea surface warming 1985–2006. *Clim. Res.* 39, 11–17. doi: [10.3354/cr00794](https://doi.org/10.3354/cr00794)
- O'Neill BC, Kriegler E, Ebi KL, Kemp-Benedict E, Riahi K et al. 2017 The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21<sup>st</sup> century. *Glob. Environ. Chang.* 42, 169–180. doi: [10.1016/j.gloenvcha.2015.01.004](https://doi.org/10.1016/j.gloenvcha.2015.01.004)
- O'Neill BC, Kriegler E, Riahi K, Ebi KL, Hallegatte S et al. 2014 A new scenario framework for climate change research: The concept of shared socioeconomic pathways. *Clim. Change* 122, 387–400. doi: [10.1007/s10584-013-0905-2](https://doi.org/10.1007/s10584-013-0905-2)
- Obermann-Hellhund A, Conte D, Somot S, Torma CZ, Ahrens B 2018 Mistral and Tramontane wind systems in climate simulations from 1950 to 2100. *Clim. Dyn.* 50, 693–703. doi: [10.1007/s00382-017-3635-8](https://doi.org/10.1007/s00382-017-3635-8)
- Obermann A, Bastin S, Belamari S, Conte D, Gaertner MÁ et al. 2018 Mistral and Tramontane wind speed and wind direction patterns in regional climate simulations. *Clim. Dyn.* 51, 1059–1076. doi: [10.1007/s00382-016-3053-3](https://doi.org/10.1007/s00382-016-3053-3)
- Occhipinti-Ambrogi A 2007 Global change and marine communities: Alien species and climate change. *Mar. Pollut. Bull.* 55, 342–352. doi: [10.1016/j.marpolbul.2006.11.014](https://doi.org/10.1016/j.marpolbul.2006.11.014)
- Ochoa-Hueso R, Bell MD, Manrique E 2014 Impacts of increased nitrogen deposition and altered precipitation regimes on soil fertility and functioning in semiarid Mediterranean shrublands. *J. Arid Environ.* 104, 106–115. doi: [10.1016/j.jaridenv.2014.01.020](https://doi.org/10.1016/j.jaridenv.2014.01.020)
- Ochoa-Hueso R, Munzi S, Alonso R, Arróniz-Crespo M, Avila A et al. 2017 Ecological impacts of atmospheric pollution and interactions with climate change in terrestrial ecosystems of the Mediterranean Basin: Current research and future directions. Elsevier <https://www.sciencedirect.com/science/article/pii/S0269749116320760?via%3Dihub> [Accessed March 30, 2019]
- Olenycz M, Sokotowski A, Niewińska A, Wołowicz M, Namieśnik J et al. 2015 Comparison of PCBs and PAHs levels in European coastal waters using mussels from the *Mytilus edulis* complex as biomonitors. *Oceanologia* 57, 196–211. doi: [10.1016/j.oceano.2014.12.001](https://doi.org/10.1016/j.oceano.2014.12.001)
- Oliva M, Gómez-Ortiz A, Salvador-Franch F, Salvà-Catarineu M, Palacios D et al. 2016 Inexistence of permafrost at the top of the Veleta peak (Sierra Nevada, Spain). *Sci. Total Environ.* 550, 484–494. doi: [10.1016/j.scitotenv.2016.01.150](https://doi.org/10.1016/j.scitotenv.2016.01.150)
- Oliva M, Žebre M, Guglielmin M, Hughes PD, Çiner A et al. 2018 Permafrost conditions in the Mediterranean region since the Last Glaciation. *Earth-Science Rev.* 185, 397–436. doi: [10.1016/j.earscirev.2018.06.018](https://doi.org/10.1016/j.earscirev.2018.06.018)
- Oliveira MA, Tomlinson SJ, Carnell EJ, Dore AJ, Serrano HC et al. 2020 Nitrogen and sulfur deposition over a region in SW Europe based on a regional atmospheric chemical transport model. *Atmos. Environ.* 223, 117290. doi: [10.1016/j.atmosenv.2020.117290](https://doi.org/10.1016/j.atmosenv.2020.117290)
- Oliver ECJ, Donat MG, Burrows MT, Moore PJ, Smale DA et al. 2018 Longer and more frequent marine heatwaves over the past century. *Nat. Commun.* 9, 1324. doi: [10.1038/s41467-018-03732-9](https://doi.org/10.1038/s41467-018-03732-9)
- Onea F, Deleanu L, Rusu L, Georgescu C 2016 Evaluation of the wind energy potential along the Mediterranean Sea coasts. *Energy Explor. Exploit.* 34, 766–792. doi: [10.1177/0144598716659592](https://doi.org/10.1177/0144598716659592)
- Onorati G, di Meo T, Bussettini M, Fabiani C, Farrace MGG et al. 2006 Groundwater quality monitoring in Italy for the implementation of the EU water framework directive. *Phys. Chem. Earth, Parts A/B/C* 31, 1004–1014. doi: [10.1016/j.pce.2006.07.001](https://doi.org/10.1016/j.pce.2006.07.001)

- Oppenheimer M, Glavovic BC, Hinkel J, van de Wal R, Magnan AK et al. 2019 Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities, in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, eds. Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M et al. (in press).
- Osborn TJ, Briffa KR, Tett SFB, Jones PD, Trigo RM 1999 Evaluation of the North Atlantic Oscillation as simulated by a coupled climate model. *Clim. Dyn.* 15, 685–702. doi: [10.1007/s003820050310](https://doi.org/10.1007/s003820050310)
- Osio GC, Orio A, Millar CP 2015 Assessing the vulnerability of Mediterranean demersal stocks and predicting exploitation status of un-assessed stocks. *Fish. Res.* 171, 110–121. doi: [10.1016/j.fishres.2015.02.005](https://doi.org/10.1016/j.fishres.2015.02.005)
- Ouzeau G, Soubeyroux J-M, Schneider M, Vautard R, Planton S 2016 Heat waves analysis over France in present and future climate: Application of a new method on the EURO-CORDEX ensemble. *Clim. Serv.* 4, 1–12. doi: [10.1016/j.cliser.2016.09.002](https://doi.org/10.1016/j.cliser.2016.09.002)
- Ozer T, Gertman I, Kress N, Silverman J, Herut B 2017 Interannual thermohaline (1979–2014) and nutrient (2002–2014) dynamics in the Levantine surface and intermediate water masses, SE Mediterranean Sea. *Glob. Planet. Change* 151, 60–67. doi: [10.1016/j.gloplacha.2016.04.001](https://doi.org/10.1016/j.gloplacha.2016.04.001)
- Ozturk T, Ceber ZP, Türkeş M, Kurnaz ML 2015 Projections of climate change in the Mediterranean Basin by using down-scaled global climate model outputs. *Int. J. Climatol.* 35, 4276–4292. doi: [10.1002/joc.4285](https://doi.org/10.1002/joc.4285)
- Paerl HW, Paul VJ 2012 Climate change: Links to global expansion of harmful cyanobacteria. *Water Res.* 46, 1349–1363. doi: [10.1016/j.watres.2011.08.002](https://doi.org/10.1016/j.watres.2011.08.002)
- Palanques A, Lopez L, Guillén J, Puig P, Masqué P 2017 Decline of trace metal pollution in the bottom sediments of the Barcelona City continental shelf (NW Mediterranean). *Sci. Total Environ.* 579, 755–767. doi: [10.1016/j.scitotenv.2016.11.031](https://doi.org/10.1016/j.scitotenv.2016.11.031)
- Palmiéri J, Orr JC, Dutay J-C, Béranger K, Schneider A et al. 2015 Simulated anthropogenic CO<sub>2</sub> storage and acidification of the Mediterranean Sea. *Biogeosciences* 12, 781–802. doi: [10.5194/bg-12-781-2015](https://doi.org/10.5194/bg-12-781-2015)
- Paluselli A, Aminot Y, Galgani F, Net S, Sempéré R 2018a Occurrence of phthalate acid esters (PAEs) in the northwestern Mediterranean Sea and the Rhône River. *Prog. Oceanogr.* 163, 221–231. doi: [10.1016/j.pocean.2017.06.002](https://doi.org/10.1016/j.pocean.2017.06.002)
- Paluselli A, Fauvelle V, Schmidt N, Galgani F, Net S et al. 2018b Distribution of phthalates in Marseille Bay (NW Mediterranean Sea). *Sci. Total Environ.* 621, 578–587. doi: [10.1016/j.scitotenv.2017.11.306](https://doi.org/10.1016/j.scitotenv.2017.11.306)
- Pant P, Harrison RM 2013 Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements: A review. *Atmos. Environ.* 77, 78–97. doi: [10.1016/j.atmosenv.2013.04.028](https://doi.org/10.1016/j.atmosenv.2013.04.028)
- Papaconstantinou C, Farrugio H 2000 Fisheries in the Mediterranean. *Mediterr. Mar. Sci.* 1, 5. doi: [10.12681/mms.2](https://doi.org/10.12681/mms.2)
- Papanastasis VP 2004 Vegetation degradation and land use changes in agrosilvopastoral systems, in *Sustainability of Agrosilvopastoral Systems*, eds. Schnabel S, Ferreira A [Reiskirchen, Germany: Catena Verlag], 1–12.
- Papanastasis VP, Kazaklis A 1998 Land use changes and conflicts in the Mediterranean-Type ecosystems of Western Crete. *Ecol. Stud.*, 141–154. doi: [10.1007/978-3-662-03543-6\\_8](https://doi.org/10.1007/978-3-662-03543-6_8)
- Parravicini V, Mangialajo L, Mousseau L, Peirano A, Morri C et al. 2015 Climate change and warm-water species at the north-western boundary of the Mediterranean Sea. *Mar. Ecol.* 36, 897–909. doi: [10.1111/maec.12277](https://doi.org/10.1111/maec.12277)
- Pascual A, Vidal-Vijande E, Ruiz S, Somot S, Papadopoulos V 2014 Spatio-temporal variability of the surface circulation in the western Mediterranean: A comparative study using altimetry and modeling, in *The Mediterranean Sea: Temporal Variability and Spatial Patterns*, eds. Borzelli GLE, Gacic M, Lionello P, Malanotte-Rizzoli P, 5–24. <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/9781118847572#page=13>
- Pastor F, Valiente JA, Palau JL 2018 Sea surface temperature in the Mediterranean: trends and spatial patterns (1982–2016). *Pure Appl. Geophys.* 175, 4017–4029. doi: [10.1007/s00024-017-1739-z](https://doi.org/10.1007/s00024-017-1739-z)
- Pauly D, Zeller D 2016 Catch reconstructions reveal that global marine fisheries catches are higher than reported and declining. *Nat. Commun.* 7, 10244. doi: [10.1038/ncomms10244](https://doi.org/10.1038/ncomms10244)
- Pausas JG, Fernández-Muñoz S 2012 Fire regime changes in the Western Mediterranean Basin: from fuel-limited to drought-driven fire regime. *Clim. Change* 110, 215–226. doi: [10.1007/s10584-011-0060-6](https://doi.org/10.1007/s10584-011-0060-6)
- Pellet V, Aires F, Munier S, Fernández Prieto D, Jordà G et al. 2019 Integrating multiple satellite observations into a coherent dataset to monitor the full water cycle – application to the Mediterranean region. *Hydrol. Earth Syst. Sci.* 23, 465–491. doi: [10.5194/hess-23-465-2019](https://doi.org/10.5194/hess-23-465-2019)
- Peña-Angulo D, Vicente-Serrano SM, Domínguez-Castro F, Murphy C, Reig F et al. 2020 Long-term precipitation in Southwestern Europe reveals no clear trend attributable to anthropogenic forcing. *Environ. Res. Lett.* 15, 094070.
- Penna A, Bertozzini E, Battocchi C, Galluzzi L, Giacobbe MG et al. 2007 Monitoring of HAB species in the Mediterranean Sea through molecular methods. *J. Plankton Res.* 29, 19–38. doi: [10.1093/plankt/fbl053](https://doi.org/10.1093/plankt/fbl053)
- Peñuelas J, Canadell JG, Ogaya R 2011 Increased water-use efficiency during the 20<sup>th</sup> century did not translate into enhanced tree growth. *Glob. Ecol. Biogeogr.* 20, 597–608. doi: [10.1111/j.1466-8238.2010.00608.x](https://doi.org/10.1111/j.1466-8238.2010.00608.x)
- Perez J, Menendez M, Camus P, Mendez Fernando J, Losada JJ 2015 Statistical multi-model climate projections of surface ocean waves in Europe. *Ocean Model.* 96, 161–170. doi: [10.1016/j.ocemod.2015.06.001](https://doi.org/10.1016/j.ocemod.2015.06.001)
- Perović V, Jakšić D, Jaramaz D, Koković N, Čakmak D et al. 2018 Spatio-temporal analysis of land use/land cover change and its effects on soil erosion (Case study in the Oplenac wine-producing area, Serbia). *Environ. Monit. Assess.* 190. doi: [10.1007/s10661-018-7025-4](https://doi.org/10.1007/s10661-018-7025-4)
- Perry AL, Low PJ, Ellis JR, Reynolds JD 2005 Climate Change and Distribution Shifts in Marine Fishes. *Science* (80-. ). 308, 1912–1915. doi: [10.1126/science.1111322](https://doi.org/10.1126/science.1111322)
- Petanidou T, Kizos T, Soulakellis N 2008 Socioeconomic Di-

- mensions of Changes in the Agricultural Landscape of the Mediterranean Basin: A Case Study of the Abandonment of Cultivation Terraces on Nisyros Island, Greece. *Environ. Manage.* 41, 250–266. doi: [10.1007/s00267-007-9054-6](https://doi.org/10.1007/s00267-007-9054-6)
- Peters K, Breitsamer L, Gerowitt B 2014 Impact of climate change on weeds in agriculture: a review. *Agron. Sustain. Dev.* 34, 707–721. doi: [10.1007/s13593-014-0245-2](https://doi.org/10.1007/s13593-014-0245-2)
- Pettersen S 1956 *Weather Analysis and Forecasting*. McGraw-Hill
- Pey J, Querol X, Alastuey A, Forastiere F, Stafoggia M 2013 African dust outbreaks over the Mediterranean Basin during 2001–2011: PM10 concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology. *Atmos. Chem. Phys.* 13, 1395–1410. doi: [10.5194/acp-13-1395-2013](https://doi.org/10.5194/acp-13-1395-2013)
- Pfeifroth U, Sánchez-Lorenzo A, Manara V, Trentmann J, Hollmann R 2018 Trends and variability of surface solar radiation in Europe based on surface- and satellite-based data records. *JGR Atmos.* 123, 1735–1754. doi: [10.1002/2017JD027418](https://doi.org/10.1002/2017JD027418)
- Pham MK, Chamizo E, Mas Balbuena JL, Miquel J-CC, Martín J et al. 2017 Impact of Saharan dust events on radionuclide levels in Monaco air and in the water column of the north-west Mediterranean Sea. *J. Environ. Radioact.* 166, 2–9. doi: [10.1016/j.jenvrad.2016.04.014](https://doi.org/10.1016/j.jenvrad.2016.04.014)
- Phelps LN, Kaplan JO 2017 Land use for animal production in global change studies: Defining and characterizing a framework. *Glob. Chang. Biol.* 23, 4457–4471. doi: [10.1111/gcb.13732](https://doi.org/10.1111/gcb.13732)
- Piazza L, Balata D, Bulleri F, Gennaro P, Ceccherelli G 2016 The invasion of *Caulerpa cylindracea* in the Mediterranean: the known, the unknown and the knowable. *Mar. Biol.* 163. doi: [10.1007/s00227-016-2937-4](https://doi.org/10.1007/s00227-016-2937-4)
- Piazza L, Ceccherelli G 2006 Persistence of biological invasion effects: Recovery of macroalgal assemblages after removal of *Caulerpa racemosa* var. *cylindracea*. *Estuar. Coast. Shelf Sci.* 68, 455–461. doi: [10.1016/j.ecss.2006.02.011](https://doi.org/10.1016/j.ecss.2006.02.011)
- Piazza L, Meinesz A, Verlaque M, Akçali B, Antolic B et al. 2005 Invasion of *Caulerpa racemosa* var. *cylindracea* (Caulerpaceae, Chlorophyta) in the Mediterranean Sea: an assessment of the spread. *Cryptogam. Algal.* 26, 189–202.
- Pikitch EK, Santora C, Babcock EA, Bakun A, Bonfil R et al. 2004 Ecosystem-Based Fishery Management. *Science (80- )*. 305, 346–347. doi: [10.1126/science.1098222](https://doi.org/10.1126/science.1098222)
- Pinilla V, Ayuda MI, Saez LA 2008 Rural depopulation and the migration turnaround in Mediterranean Western Europe: a case study of Aragon. *J. Rural Community Dev.* 3.
- Pinnegar JK, Polunin NVC, Badalamenti F 2003 Long-term changes in the trophic level of western Mediterranean fishery and aquaculture landings. *Can. J. Fish. Aquat. Sci.* 60, 222–235. doi: [10.1139/f03-016](https://doi.org/10.1139/f03-016)
- Pinto-Correia T, Vos W 2004 Multifunctionality in Mediterranean landscapes - past and future. *New Dimens. Eur. Landsc.*, 135–164. doi: [10.1007/978-1-4020-2911-0\\_10](https://doi.org/10.1007/978-1-4020-2911-0_10)
- Pinto JG, Raible CC 2012 Past and recent changes in the North Atlantic oscillation. *Wiley Interdiscip. Rev. Clim. Chang.* 3, 79–90. doi: [10.1002/wcc.150](https://doi.org/10.1002/wcc.150)
- Pinto JG, Ulbrich U, Leckebusch GC, Spangehl T, Reyers M et al. 2007 Changes in storm track and cyclone activity in three SRES ensemble experiments with the ECHAM5/MPI-OM1 GCM. *Clim. Dyn.* 29, 195–210. doi: [10.1007/s00382-007-0230-4](https://doi.org/10.1007/s00382-007-0230-4)
- Pirazzoli PA, Tomasin A 2003 Recent near-surface wind changes in the central Mediterranean and Adriatic areas. *Int. J. Climatol.* 23, 963–973. doi: [10.1002/joc.925](https://doi.org/10.1002/joc.925)
- Pisano A, Nardelli BB, Tronconi C, Santoleri R 2016 The new Mediterranean optimally interpolated pathfinder AVHRR SST data set. *Remote Sens. Environ.* 176, 107–116. doi: [10.1016/j.rse.2016.01.019](https://doi.org/10.1016/j.rse.2016.01.019)
- Pisanu S, Farris E, Filigheddu R, García MB 2012 Demographic effects of large, introduced herbivores on a long-lived endemic plant. *Plant Ecol.* 213, 1543–1553. doi: [10.1007/s11258-012-0110-9](https://doi.org/10.1007/s11258-012-0110-9)
- Planton S, Lionello P, Artale V, Aznar R, Carrillo A et al. 2012 The climate of the Mediterranean region in future climate projections, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Oxford: Elsevier), 449–502. doi: [10.1016/B978-0-12-416042-2.00008-2](https://doi.org/10.1016/B978-0-12-416042-2.00008-2)
- Poli JX 1791 Testacea utriusque Siciliae eorumque historia et anatome tabulis aeneis illustrata. *Ex Reg. Typogr. Parmae* 1, i–lxxiii, [1–6].
- Pomaro A, Cavaleri L, Lionello P 2017 Climatology and trends of the Adriatic Sea wind waves: Analysis of a 37-year long instrumental data set. *Int. J. Climatol.* 37, 4237–4250. doi: [10.1002/joc.5066](https://doi.org/10.1002/joc.5066)
- Por FD 1978 *Lessepsian migration: the influx of Red Sea biota into the Mediterranean by way of the Suez Canal*. Springer-Verlag, Berlin doi: [10.1016/0160-9327\(79\)90119-4](https://doi.org/10.1016/0160-9327(79)90119-4)
- Povinec PP, Holý K, Chudý M, Šivo A, Sýkora I et al. 2012 Long-term variations of <sup>14</sup>C and <sup>137</sup>Cs in the Bratislava air – implications of different atmospheric transport processes. *J. Environ. Radioact.* 108, 33–40. doi: [10.1016/J.JENVRAD.2011.08.004](https://doi.org/10.1016/J.JENVRAD.2011.08.004)
- Pozo-Vázquez D, Esteban-Parra MJ, Rodrigo FS, Castro-Díez Y 2001 The Association between ENSO and Winter Atmospheric Circulation and Temperature in the North Atlantic Region. *J. Clim.* 14, 3408–3420. doi: [10.1175/1520-0442\(2001\)014<3408:tabeaw>2.0.co;2](https://doi.org/10.1175/1520-0442(2001)014<3408:tabeaw>2.0.co;2)
- Prein AF, Gobiet A 2017 Impacts of uncertainties in European gridded precipitation observations on regional climate analysis. *Int. J. Climatol.* 37, 305–327. doi: [10.1002/joc.4706](https://doi.org/10.1002/joc.4706)
- Price C, Michaelides S, Pashiardis S, Alpert P 1999 Long term changes in diurnal temperature range in Cyprus. *Atmos. Res.* 51, 85–98. doi: [10.1016/S0169-8095\(99\)00022-8](https://doi.org/10.1016/S0169-8095(99)00022-8)
- Proia L, Lupini G, Osorio V, Pérez S, Barceló D et al. 2013 Response of biofilm bacterial communities to antibiotic pollutants in a Mediterranean river. *Chemosphere* 92, 1126–1135. doi: [10.1016/j.chemosphere.2013.01.063](https://doi.org/10.1016/j.chemosphere.2013.01.063)
- Puigdefábregas J, Mendizabal T 1998 Perspectives on desertification: western Mediterranean. *J. Arid Environ.* 39, 209–224. doi: [10.1006/jare.1998.0401](https://doi.org/10.1006/jare.1998.0401)
- Pujo-Pay M, Conan P, Oriol L, Cornet-Barthaux V, Falco C et al. 2011 Integrated survey of elemental stoichiometry (C, N, P) from the western to eastern Mediterranean Sea. *Biogeosciences* 8, 883–899. doi: [10.5194/bg-8-883-2011](https://doi.org/10.5194/bg-8-883-2011)

- Putaud J-P, Van Dingenen R, Alastuey A, Bauer H, Birmili W et al. 2010 A European aerosol phenomenology – 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe. *Atmos. Environ.* 44, 1308–1320. doi: [10.1016/j.atmosenv.2009.12.011](https://doi.org/10.1016/j.atmosenv.2009.12.011)
- Pyrina M, Hatzianastassiou N, Matsoukas C, Fotiadi A, Papadimas CD et al. 2015 Cloud effects on the solar and thermal radiation budgets of the Mediterranean basin. *Atmos. Res.* 152, 14–28. doi: [10.1016/j.atmosres.2013.11.009](https://doi.org/10.1016/j.atmosres.2013.11.009)
- Quadrelli R, Pavan V, Molteni F 2001 Wintertime variability of Mediterranean precipitation and its links with large-scale circulation anomalies. *Clim. Dyn.* 17, 457–466. doi: [10.1007/s003820000121](https://doi.org/10.1007/s003820000121)
- Quan X-W, Hoerling MP, Perlwitz J, Diaz HF 2018 On the Time of Emergence of Tropical Width Change. *J. Clim.* 31, 7225–7236. doi: [10.1175/jcli-d-18-0068.1](https://doi.org/10.1175/jcli-d-18-0068.1)
- Querol X, Alastuey A, Gangoiti G, Perez N, Lee HK et al. 2018 Phenomenology of summer ozone episodes over the Madrid Metropolitan Area, central Spain. *Atmos. Chem. Phys.* 18, 6511–6533. doi: [10.5194/acp-18-6511-2018](https://doi.org/10.5194/acp-18-6511-2018)
- Querol X, Alastuey A, Viana M-M, Rodríguez S, Artíñano B et al. 2004 Speciation and origin of PM10 and PM2.5 in Spain. *J. Aerosol Sci.* 35, 1151–1172. doi: [10.1016/j.jaerosci.2004.04.002](https://doi.org/10.1016/j.jaerosci.2004.04.002)
- Querol X, Pey J, Pandolfi M, Alastuey A, Cusack M et al. 2009 African dust contributions to mean ambient PM10 mass-levels across the Mediterranean Basin. *Atmos. Environ.* 43, 4266–4277. doi: [10.1016/j.atmosenv.2009.06.013](https://doi.org/10.1016/j.atmosenv.2009.06.013)
- Quesada B, Vautard R, Yiou P, Hirschi M, Seneviratne SI 2012 Asymmetric European summer heat predictability from wet and dry southern winters and springs. *Nat. Clim. Chang.* 2, 736–741. doi: [10.1038/nclimate1536](https://doi.org/10.1038/nclimate1536)
- Quintana-Seguí P, Martin E, Sánchez E, Zribi M, Vennetier M et al. 2016 Drought: Observed trends, future projections, in *The Mediterranean Region under Climate Change. A Scientific Update*, eds. Thiébaud S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 123–132.
- Rabatel A, Letréguilly A, Dedieu J-P, Eckert N 2013 Changes in glacier equilibrium-line altitude in the western Alps from 1984 to 2010: evaluation by remote sensing and modeling of the morpho-topographic and climate controls. *Cryosph.* 7, 1455–1471. doi: [10.5194/tc-7-1455-2013](https://doi.org/10.5194/tc-7-1455-2013)
- Rabitsch W 2010 True Bugs (Hemiptera, Heteroptera). *BioRisk* 4, 407–433. doi: [10.3897/biorisk.4.44](https://doi.org/10.3897/biorisk.4.44)
- Raclot D, Le Bissonnais Y, Annabi M, Sabir M, Smetanova A 2018 Main issues for preserving Mediterranean soil resources from water erosion under global change. *L. Degrad. Dev.* 29, 789–799. doi: [10.1002/ldr.2774](https://doi.org/10.1002/ldr.2774)
- Rafael S, Martins H, Borrego C, Lopes M 2015 Urban vulnerability and resilience to climate change. *WIT Trans. Ecol. Environ.* 198, 379–390. doi: [10.2495/AIR150331](https://doi.org/10.2495/AIR150331)
- Raible CC, Saaroni H, Wild M 2010 Winter synoptic-scale variability over the Mediterranean Basin under future climate conditions as simulated by the ECHAM5. *Clim. Dyn.* 35, 473–488. doi: [10.1007/s00382-009-0678-5](https://doi.org/10.1007/s00382-009-0678-5)
- Räisänen J 2002 CO<sub>2</sub>-induced changes in interannual temperature and precipitation variability in 19 CMIP2 experiments. *Bull. Am. Meteorol. Soc.* 15, 2395–2411. doi: [10.1175/1520-0442\(2002\)015%3C2395:CICIIT%3E2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015%3C2395:CICIIT%3E2.0.CO;2)
- Raitsos DE, Beaugrand G, Georgopoulos D, Zenetos A, Pancucci-Papadopoulou AM et al. 2010 Global climate change amplifies the entry of tropical species into the eastern Mediterranean Sea. *Limnol. Oceanogr.* 55, 1478–1484. doi: [10.4319/lo.2010.55.4.1478](https://doi.org/10.4319/lo.2010.55.4.1478)
- Ramesh K, Matloob A, Aslam F, Florentine SK, Chauhan BS 2017 Weeds in a Changing Climate: Vulnerabilities, Consequences, and Implications for Future Weed Management. *Front. Plant Sci.* 8, 95. doi: [10.3389/fpls.2017.00095](https://doi.org/10.3389/fpls.2017.00095)
- Rascher KG, Große-Stoltenberg A, Máguas C, Meira-Neto JAA, Werner C 2011 *Acacia longifolia* invasion impacts vegetation structure and regeneration dynamics in open dunes and pine forests. *Biol. Invasions* 13, 1099–1113. doi: [10.1007/s10530-011-9949-2](https://doi.org/10.1007/s10530-011-9949-2)
- Rasheed T, Bilal M, Nabeel F, Adeel M, Iqbal HMN 2019 Environmentally-related contaminants of high concern: Potential sources and analytical modalities for detection, quantification, and treatment. *Env. Int.* 122, 52–66. doi: [10.1016/j.envint.2018.11.038](https://doi.org/10.1016/j.envint.2018.11.038)
- Rasmussen E, Zick C 1987 A subsynoptic vortex over the Mediterranean with some resemblance to polar lows. *Tellus A Dyn. Meteorol. Oceanogr.* 39 A, 408–425. doi: [10.1111/j.1600-0870.1987.tb00318.x](https://doi.org/10.1111/j.1600-0870.1987.tb00318.x)
- Rasmussen K, Thyrring J, Muscarella R, Borchsenius F 2017 Climate-change-induced range shifts of three allergenic ragweeds (*Ambrosia* L.) in Europe and their potential impact on human health. *PeerJ* 5, e3104. doi: [10.7717/peerj.3104](https://doi.org/10.7717/peerj.3104)
- Rasplus J-Y, Villemant C, Rosa Paiva M, Delvare G, Roques A 2010 Hymenoptera. *BioRisk* 4, 669–776. doi: [10.3897/biorisk.4.55](https://doi.org/10.3897/biorisk.4.55)
- Rassati D, Faccoli M, Petrucco Toffolo E, Battisti A, Marini L 2015 Improving the early detection of alien wood-boring beetles in ports and surrounding forests. *J. Appl. Ecol.* 52, 50–58. doi: [10.1111/1365-2664.12347](https://doi.org/10.1111/1365-2664.12347)
- Rassati D, Petrucco Toffolo E, Roques A, Battisti A, Faccoli M 2014 Trapping wood boring beetles in Italian ports: a pilot study. *J. Pest Sci. (2004)*. 87, 61–69. doi: [10.1007/s10340-013-0499-5](https://doi.org/10.1007/s10340-013-0499-5)
- Ratola N, Cincinelli A, Alves A, Katsoyiannis A 2012 Occurrence of organic microcontaminants in the wastewater treatment process. A mini review. *J. Hazard. Mater.* 239–240, 1–18. doi: [10.1016/j.jhazmat.2012.05.040](https://doi.org/10.1016/j.jhazmat.2012.05.040)
- Rauthe M, Hense A, Paeth H 2004 A model intercomparison study of climate change-signals in extratropical circulation. *Int. J. Climatol.* 24, 643–662. doi: [10.1002/joc.1025](https://doi.org/10.1002/joc.1025)
- Reale M, Liberato MLR, Lionello P, Pinto JG, Salon S et al. 2019 A Global Climatology of Explosive Cyclones using a Multi-Tracking Approach. *Tellus A Dyn. Meteorol. Oceanogr.* 71, 1611340. doi: [10.1080/16000870.2019.1611340](https://doi.org/10.1080/16000870.2019.1611340)
- Reale M, Lionello P 2013 Synoptic climatology of winter intense precipitation events along the Mediterranean coasts. *Nat. Hazards Earth Syst. Sci.* 13, 1707–1722. doi: [10.5194/nhess-13-1707-2013](https://doi.org/10.5194/nhess-13-1707-2013)
- Reale O, Atlas R 2001 Tropical Cyclone-Like Vortices in the Extratropics: Observational Evidence and Synoptic Analysis. *Weather Forecast.* 16, 7–34. doi: [10.1175/1520-0434\(2001\)016<0007:tcclvit>2.0.co;2](https://doi.org/10.1175/1520-0434(2001)016<0007:tcclvit>2.0.co;2)

- Rees P, van der Gaag N, de Beer J, Heins F 2012 European Regional Populations: Current Trends, Future Pathways, and Policy Options. *Eur. J. Popul.* 28, 385–416. doi: [10.1007/s10680-012-9268-z](https://doi.org/10.1007/s10680-012-9268-z)
- Reichard SH, Hamilton CW 1997 Predicting invasions of woody plants introduced into North America. *Conserv. Biol.* 11, 193–203. doi: [10.1046/j.1523-1739.1997.95473.x](https://doi.org/10.1046/j.1523-1739.1997.95473.x)
- Reimann L, Merckens J-L, Vafeidis AT 2018 Regionalized Shared Socioeconomic Pathways: Narratives and spatial population projections for the Mediterranean coastal zone. *Reg. Environ. Chang.* 18, 235–245. doi: [10.1007/s10113-017-1189-2](https://doi.org/10.1007/s10113-017-1189-2)
- Rejda M 2004 Forest cover changes in the Maghreb countries with special reference to Morocco, in *Recent Dynamics of the Mediterranean Vegetation and Landscape*, eds. Mazzoleni S, Pasquale G Di, Mulligan M, Di Martino P, Rego F (John Wiley & Sons, Ltd), 21–31. doi: [10.1002/0470093714.ch3](https://doi.org/10.1002/0470093714.ch3)
- Rezg R, El-Fazaa S, Gharbi N, Mornagui B 2014 Bisphenol A and human chronic diseases: Current evidences, possible mechanisms, and future perspectives. *Environ. Int.* 64, 83–90. doi: [10.1016/j.envint.2013.12.007](https://doi.org/10.1016/j.envint.2013.12.007)
- Riahi K, Rao S, Krey V, Cho C, Chirkov V et al. 2011 RCP 8.5-A scenario of comparatively high greenhouse gas emissions. *Clim. Change* 109, 33–57. doi: [10.1007/s10584-011-0149-y](https://doi.org/10.1007/s10584-011-0149-y)
- Ribes A, Thao S, Vautard R, Dubuisson B, Somot S et al. 2019 Observed increase in extreme daily rainfall in the French Mediterranean. *Clim. Dyn.* 52, 1095–1114. <https://link.springer.com/article/10.1007/s00382-018-4179-2> [Accessed September 17, 2019].
- Ricart M, Guasch H, Barceló D, Brix R, Conceição MH et al. 2010 Primary and complex stressors in polluted mediterranean rivers: Pesticide effects on biological communities. *J. Hydrol.* 383, 52–61. doi: [10.1016/j.jhydrol.2009.08.014](https://doi.org/10.1016/j.jhydrol.2009.08.014)
- Rilov G 2016 Multi-species collapses at the warm edge of a warming sea. *Sci. Rep.* 6, 36897. doi: [10.1038/srep36897](https://doi.org/10.1038/srep36897)
- Rilov G, Crooks JA 2009 *Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives*. Springer, Heidelberg, Germany.
- Rilov G, Galil BS 2009 Marine Bioinvasions in the Mediterranean Sea – History, Distribution and Ecology, in *Biological Invasions in Marine Ecosystems*, eds. Rilov G, Crooks JA (Springer-Verlag, Heidelberg, Germany.), 549–575. doi: [10.1007/978-3-540-79236-9\\_31](https://doi.org/10.1007/978-3-540-79236-9_31)
- Rilov G, Mazaris AD, Stelzenmüller V, Helmuth B, Wahl M et al. 2019a Adaptive marine conservation planning in the face of climate change: What can we learn from physiological, ecological and genetic studies? *Glob. Ecol. Conserv.* 17, e00566. doi: [10.1016/j.gecco.2019.e00566](https://doi.org/10.1016/j.gecco.2019.e00566)
- Rilov G, Peleg O, Guy-Haim T 2019b The Restructuring of Levant Reefs by Aliens, Ocean Warming and Overfishing: Implications to Species Interactions and Ecosystem Functions, in *Interactions in the Marine Benthos – Global Patterns and Processes*, eds. Hawkins SJ, Bohn K, Firth LB, Williams GA (Cambridge University Press, Cambridge).
- Rivetti I, Boero F, Fraschetti S, Zambianchi E, Lionello P 2017 Anomalies of the upper water column in the Mediterranean Sea. *Glob. Planet. Change* 151, 68–79. doi: [10.1016/j.gloplacha.2016.03.001](https://doi.org/10.1016/j.gloplacha.2016.03.001)
- Rivetti I, Fraschetti S, Lionello P, Zambianchi E, Boero F 2014 Global Warming and Mass Mortalities of Benthic Invertebrates in the Mediterranean Sea. *PLoS One* 9, e115655. doi: [10.1371/journal.pone.0115655](https://doi.org/10.1371/journal.pone.0115655)
- Rixen M, Beckers J-M, Levitus S, Antonov J, Boyer T et al. 2005 The Western Mediterranean Deep Water: A proxy for climate change. *Geophys. Res. Lett.* 32, n/a-n/a. doi: [10.1029/2005GL022702](https://doi.org/10.1029/2005GL022702)
- Roberts CM, O'Leary BC, McCauley DJ, Cury PM, Duarte CM et al. 2017 Marine reserves can mitigate and promote adaptation to climate change. *Proc. Natl. Acad. Sci. U. S. A.* 114, 6167–6175. doi: [10.1073/pnas.1701262114](https://doi.org/10.1073/pnas.1701262114)
- Robles-Molina J, Gilbert-López B, García-Reyes JF, Molina-Díaz A 2014 Monitoring of selected priority and emerging contaminants in the Guadalquivir River and other related surface waters in the province of Jaén, South East Spain. *Sci. Total Environ.* 479–480, 247–257. doi: [10.1016/j.scitotenv.2014.01.121](https://doi.org/10.1016/j.scitotenv.2014.01.121)
- Rochelle-Newall EJ, Delesalle B, Mari X, Rouchon C, Torrétón JP et al. 2008 Zinc induces shifts in microbial carbon flux in tropical coastal environments. *Aquat. Microb. Ecol.* 52, 57–68. doi: [10.3354/ame01212](https://doi.org/10.3354/ame01212)
- Rodger HD, Henry L, Mitchell SO 2011 Non-infectious gill disorders of marine salmonid fish. *Rev. Fish Biol. Fish.* 21, 423–440. doi: [10.1007/s11160-010-9182-6](https://doi.org/10.1007/s11160-010-9182-6)
- Rodó X 2001 Reversal of three global atmospheric fields linking changes in SST anomalies in the Pacific, Atlantic and Indian oceans at tropical latitudes and midlatitudes. *Clim. Dyn.* 18, 203–217. doi: [10.1007/s003820100171](https://doi.org/10.1007/s003820100171)
- Rodó X, Baert E, Comin FA 1997 Variations in seasonal rainfall in Southern Europe during the present century: relationships with the North Atlantic Oscillation and the El Niño-Southern Oscillation. *Clim. Dyn.* 13, 275–284.
- Rodríguez-Fonseca B, de Castro M 2002 On the connection between winter anomalous precipitation in the Iberian Peninsula and North West Africa and the summer subtropical Atlantic sea surface temperature. *Geophys. Res. Lett.* 29, 10-1-10-4. doi: [10.1029/2001gl014421](https://doi.org/10.1029/2001gl014421)
- Rodríguez-Lazaro D, Cook N, Ruggeri FM, Sellwood J, Nasser A et al. 2012 Virus hazards from food, water and other contaminated environments. *FEMS Microbiol. Rev.* 36, 786–814. doi: [10.1111/j.1574-6976.2011.00306.x](https://doi.org/10.1111/j.1574-6976.2011.00306.x)
- Rodríguez S, Querol X, Alastuey A, de la Rosa J 2007 Atmospheric particulate matter and air quality in the Mediterranean: a review. *Environ. Chem. Lett.* 5, 1–7. doi: [10.1007/s10311-006-0071-0](https://doi.org/10.1007/s10311-006-0071-0)
- Rodríguez S, Querol X, Alastuey A, Viana M-M, Mantilla E 2003 Events Affecting Levels and Seasonal Evolution of Airborne Particulate Matter Concentrations in the Western Mediterranean. *Environ. Sci. Technol.* 37, 216–222. doi: [10.1021/es020106p](https://doi.org/10.1021/es020106p)
- Rodwell MJ, Hoskins BJ 1996 Monsoons and the dynamics of deserts. *Q. J. R. Meteorol. Soc.* 122, 1385–1404. doi: [10.1002/qj.49712253408](https://doi.org/10.1002/qj.49712253408)
- Roether W, Klein B, Manca BB, Theocharis A, Kioroglou S 2007 Transient Eastern Mediterranean deep waters in response to the massive dense-water output of the Aegean Sea in the 1990s. *Prog. Oceanogr.* 74, 540–571.



- doi: [10.1016/j.pocean.2007.03.001](https://doi.org/10.1016/j.pocean.2007.03.001)
- Rohat G, Flacke J, Dao H, van Maarseveen M 2018 Co-use of existing scenario sets to extend and quantify the shared socioeconomic pathways. *Clim. Change* 151, 619–636. doi: [10.1007/s10584-018-2318-8](https://doi.org/10.1007/s10584-018-2318-8)
- Rohling EJ, Bryden HL 1992 Man-induced salinity and temperature increases in Western Mediterranean deep water. *JGR Atmos.* 97, 11191–11198. doi: [10.1029/92JC00767](https://doi.org/10.1029/92JC00767)
- Roig N, Sierra J, Rovira J, Schuhmacher M, Domingo JL et al. 2013 In vitro tests to assess toxic effects of airborne PM10 samples. Correlation with metals and chlorinated dioxins and furans. *Sci. Total Environ.* 443, 791–797. doi: [10.1016/j.scitotenv.2012.11.022](https://doi.org/10.1016/j.scitotenv.2012.11.022)
- Rojas M, Li LZ, Kanakidou M, Hatzianastassiou N, Seze G et al. 2013 Winter weather regimes over the Mediterranean region: their role for the regional climate and projected changes in the twenty-first century. *Clim. Dyn.* 41, 551–571.
- Roll U, Dayan T, Simberloff D, Goren M 2007 Characteristics of the introduced fish fauna of Israel. *Biol. Invasions* 9, 813–824. doi: [10.1007/s10530-006-9083-8](https://doi.org/10.1007/s10530-006-9083-8)
- Romera R, Gaertner MÁ, Sánchez E, Domínguez M, González-Alemán JJ et al. 2017 Climate change projections of medicanes with a large multi-model ensemble of regional climate models. *Glob. Planet. Change* 151, 134–143. doi: [10.1016/j.gloplacha.2016.10.008](https://doi.org/10.1016/j.gloplacha.2016.10.008)
- Romero R, Emanuel K 2013 Medicanes risk in a changing climate. *JGR Atmos.* 118, 5992–6001. doi: [10.1002/jgrd.50475](https://doi.org/10.1002/jgrd.50475)
- Romero R, Emanuel K 2017 Climate Change and Hurricane-Like Extratropical Cyclones: Projections for North Atlantic Polar Lows and Medicanes Based on CMIP5 Models. *J. Clim.* 30, 279–299. doi: [10.1175/jcli-d-16-0255.1](https://doi.org/10.1175/jcli-d-16-0255.1)
- Romo S, Soria J, Fernandez F, Ouahid Y, Baron-Sola A 2013 Water residence time and the dynamics of toxic cyanobacteria. *Freshw. Biol.* 58, 513–522. doi: [10.1111/j.1365-2427.2012.02734.x](https://doi.org/10.1111/j.1365-2427.2012.02734.x)
- Roques A 2010 Taxonomy, time and geographic patterns. Chapter 2. *BioRisk* 4, 11–26. doi: [10.3897/biorisk.4.70](https://doi.org/10.3897/biorisk.4.70)
- Roques A 2015 Drivers and pathways of forest insect invasions in Europe, can we predict the next arrivals? *Atti Accad. Naz. Ital. di Entomol.* 53, 145–150.
- Roques A, Auger-Rozenberg M-A, Blackburn TM, Garnas J, Pyšek P et al. 2016 Temporal and interspecific variation in rates of spread for insect species invading Europe during the last 200 years. *Biol. Invasions* 18, 907–920. doi: [10.1007/s10530-016-1080-y](https://doi.org/10.1007/s10530-016-1080-y)
- Roques A, Kenis M, Lees D, Lopez-Vaamonde C, Rabitsch W et al. 2010 Alien terrestrial arthropods of Europe. *BIORISK*, 1028. doi: [10.1007/978-1-4020-8280-1\\_5](https://doi.org/10.1007/978-1-4020-8280-1_5)
- Roques A, Preda C, Augustin S, Auger-Rozenberg M-A 2018 Bugs, ants, wasps, moths and other insect species, in *Invasive species and human health*, eds. Maza G, Tricarico E (CABI, Wallingford, UK), 63–75. doi: [10.1079/9781786390981.0063](https://doi.org/10.1079/9781786390981.0063)
- Rotenberg E, Yakir D 2011 Distinct patterns of changes in surface energy budget associated with forestation in the semi-arid region. *Glob. Chang. Biol.* 17, 1536–1548. doi: [10.1111/j.1365-2486.2010.02320.x](https://doi.org/10.1111/j.1365-2486.2010.02320.x)
- Rotmans J, van Asselt M, Anastasi C, Greeuw S, Mellors J et al. 2000 Visions for a sustainable Europe. *Futures* 32, 809–831. doi: [10.1016/s0016-3287\(00\)00033-1](https://doi.org/10.1016/s0016-3287(00)00033-1)
- Rousi E, Anagnostopoulou C, Tolika K, Maheras P 2015 Representing teleconnection patterns over Europe: A comparison of SOM and PCA methods. *Atmos. Res.* 152, 123–137. doi: [10.1016/j.atmosres.2013.11.010](https://doi.org/10.1016/j.atmosres.2013.11.010)
- Rousis NI, Bade R, Bijlsma L, Zuccato E, Sancho J V. et al. 2017 Monitoring a large number of pesticides and transformation products in water samples from Spain and Italy. *Environ. Res.* 156, 31–38. doi: [10.1016/j.envres.2017.03.013](https://doi.org/10.1016/j.envres.2017.03.013)
- Roy HE, Bacher S, Essl F, Adriaens T, Aldridge DC et al. 2019 Developing a list of invasive alien species likely to threaten biodiversity and ecosystems in the European Union. *Glob. Chang. Biol.* 25, gcb.14527. doi: [10.1111/gcb.14527](https://doi.org/10.1111/gcb.14527)
- Roy HE, Rabitsch W, Scalera R, Stewart A, Gallardo B et al. 2018 Developing a framework of minimum standards for the risk assessment of alien species. *J. Appl. Ecol.* 55, 526–538. doi: [10.1111/1365-2664.13025](https://doi.org/10.1111/1365-2664.13025)
- Roy HE, Rhule E, Harding S, Lawson Handley L-J, Poland RL et al. 2011 Living with the enemy: Parasites and pathogens of the ladybird *Harmonia axyridis*. *BioControl* 56, 663–679. doi: [10.1007/s10526-011-9387-1](https://doi.org/10.1007/s10526-011-9387-1)
- Ruffault J, Mouillot F, Peters DPC 2015 How a new fire-suppression policy can abruptly reshape the fire-weather relationship. *Ecosphere* 6, 1–19. doi: [10.1890/ES15-00182.1](https://doi.org/10.1890/ES15-00182.1)
- Rugman-Jones PF, Hoddle CD, Hoddle MS, Stouthamer R 2013 The Lesser of Two Weevils: Molecular genetics of pest palm weevil populations confirm *Rhynchophorus vulneratus* (Panzer 1798) as a valid species distinct from *R. ferrugineus* (Olivier 1790), and reveal the global extent of both. *PLoS One* 8, e78379. doi: [10.1371/journal.pone.0078379](https://doi.org/10.1371/journal.pone.0078379)
- Ruiz-Ramos M, Ferrise R, Rodríguez A, Lorite IJ, Bindi M et al. 2018 Adaptation response surfaces for managing wheat under perturbed climate and CO<sub>2</sub> in a Mediterranean environment. *Agric. Syst.* 159, 260–274. doi: [10.1016/j.agsy.2017.01.009](https://doi.org/10.1016/j.agsy.2017.01.009)
- Rusiñol M, Fernandez-Cassi X, Timoneda N, Carratalà A, Abril JF et al. 2015 Evidence of viral dissemination and seasonality in a Mediterranean river catchment: Implications for water pollution management. *J. Environ. Manage.* 159, 58–67. doi: [10.1016/j.jenvman.2015.05.019](https://doi.org/10.1016/j.jenvman.2015.05.019)
- Russo MA, Leitão J, Gama C, Ferreira J, Monteiro A 2018 Shipping emissions over Europe: A state-of-the-art and comparative analysis. *Atmos. Environ.* 177, 187–194. doi: [10.1016/j.atmosenv.2018.01.025](https://doi.org/10.1016/j.atmosenv.2018.01.025)
- Russo S, Dosio A, Graversen RG, Sillmann J, Carrao H et al. 2014 Magnitude of extreme heat waves in present climate and their projection in a warming world. *JGR Atmos.* 119, 12,500–12,512. doi: [10.1002/2014jd022098](https://doi.org/10.1002/2014jd022098)
- Ruti PM, Somot S, Giorgi F, Dubois C, Flaouanas E et al. 2016 Med-CORDEX initiative for Mediterranean climate studies. *Bull. Am. Meteorol. Soc.* 97, 1187–1208. doi: [10.1175/BAMS-D-14-00176.1](https://doi.org/10.1175/BAMS-D-14-00176.1)
- Ryan PG, Moore CJ, van Franeker JA, Moloney CL 2009 Monitoring the abundance of plastic debris in the marine environment. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 1999–2012. doi: [10.1098/rstb.2008.0207](https://doi.org/10.1098/rstb.2008.0207)

- Saadi S, Todorovic M, Tanasijevic L, Pereira LS, Pizzigalli C et al. 2015 Climate change and Mediterranean agriculture: Impacts on winter wheat and tomato crop evapotranspiration, irrigation requirements and yield. *Agric. Water Manag.* 147, 103–115. doi: [10.1016/J.AGWAT.2014.05.008](https://doi.org/10.1016/J.AGWAT.2014.05.008)
- Saaroni H, Ziv B, Edelson J, Alpert P 2003 Long-term variations in summer temperatures over the Eastern Mediterranean. *Geophys. Res. Lett.* 30. doi: [10.1029/2003gl017742](https://doi.org/10.1029/2003gl017742)
- Sabatés A, Martín P, Lloret J, Raya V 2006 Sea warming and fish distribution: The case of the small pelagic fish, *Sardinella aurita*, in the western Mediterranean. *Glob. Chang. Biol.* 12, 2209–2219. doi: [10.1111/j.1365-2486.2006.01246.x](https://doi.org/10.1111/j.1365-2486.2006.01246.x)
- Sabatés A, Martín P, Raya V, Sabatés A, Martín P et al. 2012 Changes in life-history traits in relation to climate change: bluefish (*Pomatomus saltatrix*) in the northwestern Mediterranean. *ICES J. Mar. Sci.* 69, 1000–1009. doi: [10.1093/icesjms/fss053](https://doi.org/10.1093/icesjms/fss053)
- Sabatés A, Salat J, Raya V, Emelianov M, Segura-Noguera M 2009 Spawning environmental conditions of *Sardinella aurita* at the northern limit of its distribution range, the western Mediterranean. *Mar. Ecol. Prog. Ser.* 385, 227–236. doi: [10.3354/meps08058](https://doi.org/10.3354/meps08058)
- Sabine CL, Feely RA, Gruber N, Key RM, Lee K et al. 2004 The Oceanic Sink for Anthropogenic CO<sub>2</sub>. *Science* (80-. ). 305, 367–371. doi: [10.1126/science.1097403](https://doi.org/10.1126/science.1097403)
- Sáenz J, Rodríguez-Puebla C, Fernández J, Zubillaga J 2001 Interpretation of interannual winter temperature variations over southwestern Europe. *JGR Atmos.* 106, 20641–20651. doi: [10.1029/2001JD900247](https://doi.org/10.1029/2001JD900247)
- Saffioti C, Fischer EM, Scherrer SC, Knutti R 2016 Reconciling observed and modeled temperature and precipitation trends over Europe by adjusting for circulation variability. *Geophys. Res. Lett.* 43, 8189–8198. doi: [10.1002/2016GL069802](https://doi.org/10.1002/2016GL069802)
- Sahraoui I, Grami B, Bates SS, Bouchouicha D, Chikhaoui MA et al. 2012 Response of potentially toxic Pseudo-nitzschia (Bacillariophyceae) populations and domoic acid to environmental conditions in a eutrophied, SW Mediterranean coastal lagoon (Tunisia). *Estuar. Coast. Shelf Sci.* 102, 95–104. doi: [10.1016/j.ecss.2012.03.018](https://doi.org/10.1016/j.ecss.2012.03.018)
- Sahu LK, Saxena P 2015 High time and mass resolved PTR-TOF-MS measurements of VOCs at an urban site of India during winter: Role of anthropogenic, biomass burning, biogenic and photochemical sources. *Atmos. Res.* 164–165, 84–94. doi: [10.1016/J.ATMOSRES.2015.04.021](https://doi.org/10.1016/J.ATMOSRES.2015.04.021)
- Sahu LK, Yadav R, Pal D 2016 Source identification of VOCs at an urban site of western India: Effect of marathon events and anthropogenic emissions. *JGR Atmos.* 121, 2416–2433. doi: [10.1002/2015JD024454](https://doi.org/10.1002/2015JD024454)
- Sakka Hlaili A, Chikhaoui M-A, El Grami B, Mabrouk HH 2006 Effects of N and P supply on phytoplankton in Bizerte Lagoon (western Mediterranean). *J. Exp. Mar. Bio. Ecol.* 333, 79–96. doi: [10.1016/j.jembe.2005.12.049](https://doi.org/10.1016/j.jembe.2005.12.049)
- Sakka Hlaili A, Sahraoui I, Bouchouicha D, Meliti Garali S, Ksouri J et al. 2016 Toxic and potentially toxic diatoms blooms in SW Mediterranean waters: review of ten years investigations, in *Advances in Environmental Research*, ed. Daniels JA (Nova Science Publishers, Inc.).
- Sala E, Kizilkaya Z, Yildirim D, Ballesteros E 2011 Alien Marine Fishes Deplete Algal Biomass in the Eastern Mediterranean. *PLoS One* 6, e17356. doi: [10.1371/journal.pone.0017356](https://doi.org/10.1371/journal.pone.0017356)
- Saliba NA, Massoud R 2010 A Comparative Review of PM Levels, Sources, and Their Likely Fates in the Eastern Mediterranean Region, in *Environmental Science and Engineering (Subseries: Environmental Science)*, eds. Zereini F, Wiseman C (Springer, Berlin, Heidelberg), 3–17. doi: [10.1007/978-3-642-12278-1\\_1](https://doi.org/10.1007/978-3-642-12278-1_1)
- Saliba NA, Moussa S, Salame H, El-Fadel M 2006 Variation of selected air quality indicators over the city of Beirut, Lebanon: Assessment of emission sources. *Atmos. Environ.* 40, 3263–3268. doi: [10.1016/j.atmosenv.2006.01.054](https://doi.org/10.1016/j.atmosenv.2006.01.054)
- Sallustio L, di Cristofaro M, Hashmi M, Vizzarri M, Sitzia T et al. 2018 Evaluating the Contribution of Trees outside Forests and Small Open Areas to the Italian Landscape Diversification during the Last Decades. *Forests* 9, 701. doi: [10.3390/f9110701](https://doi.org/10.3390/f9110701)
- Salvador P, Artñano B, Viana M-M, Alastuey A, Querol X 2012 Evaluation of the changes in the Madrid metropolitan area influencing air quality: Analysis of 1999–2008 temporal trend of particulate matter. *Atmos. Environ.* 57, 175–185. doi: [10.1016/J.ATMOSENV.2012.04.026](https://doi.org/10.1016/J.ATMOSENV.2012.04.026)
- Salvati A, Monti P, Coch Roura H, Cecere C 2019 Climatic performance of urban textures: Analysis tools for a Mediterranean urban context. *Energy Build.* 185, 162–179. doi: [10.1016/j.enbuild.2018.12.024](https://doi.org/10.1016/j.enbuild.2018.12.024)
- Samaniego L, Thober S, Kumar R, Wanders N, Rakovec O et al. 2018 Anthropogenic warming exacerbates European soil moisture droughts. *Nat. Clim. Chang.* 8, 421–426. doi: [10.1038/s41558-018-0138-5](https://doi.org/10.1038/s41558-018-0138-5)
- San-Miguel-Ayanz J, Moreno JM, Camia A 2013 Analysis of large fires in European Mediterranean landscapes: Lessons learned and perspectives. *For. Ecol. Manage.* 294, 11–22. doi: [10.1016/j.foreco.2012.10.050](https://doi.org/10.1016/j.foreco.2012.10.050)
- Sánchez-Benítez A, García-Herrera R, Barriopedro D, Sousa PM, Trigo RM 2018 June 2017: The Earliest European Summer Mega-heatwave of Reanalysis Period. *Geophys. Res. Lett.* 45, 1955–1962. doi: [10.1002/2018GL077253](https://doi.org/10.1002/2018GL077253)
- Sánchez-Gomez E, Somot S, Déqué M 2009 Ability of an ensemble of regional climate models to reproduce weather regimes over Europe–Atlantic during the period 1961–2000. *Clim. Dyn.* 33, 723–736. doi: [10.1007/s00382-008-0502-7](https://doi.org/10.1007/s00382-008-0502-7)
- Sánchez-Lorenzo A, Brunetti M, Calbó J, Martín-Vide J 2007 Recent spatial and temporal variability and trends of sunshine duration over the Iberian Peninsula from a homogenized data set. *JGR Atmos.* 112, D20115. doi: [10.1029/2007JD008677](https://doi.org/10.1029/2007JD008677)
- Sánchez-Lorenzo A, Calbó J, Wild M 2013 Global and diffuse solar radiation in Spain: Building a homogeneous dataset and assessing their trends. *Glob. Planet. Change* 100, 343–352. doi: [10.1016/J.GLOPLACHA.2012.11.010](https://doi.org/10.1016/J.GLOPLACHA.2012.11.010)
- Sánchez-Lorenzo A, Enriquez-Alonso A, Calbó J, González J-A, Wild M et al. 2017 Fewer clouds in the Mediterranean: consistency of observations and climate simulations. *Sci. Rep.* 7, 41475. doi: [10.1038/srep41475](https://doi.org/10.1038/srep41475)

- Sánchez J, Bisquert M, Rubio E, Caselles V 2015 Impact of Land Cover Change Induced by a Fire Event on the Surface Energy Fluxes Derived from Remote Sensing. *Remote Sens.* 7, 14899–14915. doi: [10.3390/rs71114899](https://doi.org/10.3390/rs71114899)
- Sanna S, Le Tellier J 2013 Building on the Mediterranean scenario experiences: Cross-cutting approaches between regional foresight analysis and participatory prospective. Valbonne: Plan Bleu.
- Santi Delia A, Caruso G, Melcarne L, Caruso G, Parisi S et al. 2015 Biological Toxins from Marine and Freshwater Microalgae, in *Microbial Toxins and Related Contamination in the Food Industry*, eds. Caruso G, Caruso G, Laganà P, Santi Delia A, Parisi S et al. (Springer International Publishing).
- Santini A, Ghelardini L, de Pace C, Desprez-Loustau ML, Capretti P et al. 2013 Biogeographical patterns and determinants of invasion by forest pathogens in Europe. *New Phytol.* 197, 238–250. doi: [10.1111/j.1469-8137.2012.04364.x](https://doi.org/10.1111/j.1469-8137.2012.04364.x)
- Sarris D, Christopoulou A, Angelonidi E, Koutsias N, Fulé PZ et al. 2014 Increasing extremes of heat and drought associated with recent severe wildfires in southern Greece. *Reg. Environ. Chang.* 14, 1257–1268. doi: [10.1007/s10113-013-0568-6](https://doi.org/10.1007/s10113-013-0568-6)
- Saura S, Estreguil C, Mouton C, Rodriguez-Freire M 2011 Network analysis to assess landscape connectivity trends: application to European forests (1990–2000). *Ecol. Indic.* 11, 407–416. doi: [10.1016/j.ecolind.2010.06.011](https://doi.org/10.1016/j.ecolind.2010.06.011)
- Sauvé S, Desrosiers M 2014 A review of what is an emerging contaminant. *Chem. Cent. J.* 8. doi: [10.1186/1752-153X-8-15](https://doi.org/10.1186/1752-153X-8-15)
- Schär C, Vidale PL, Lüthi D, Frei C, Häberli C et al. 2004 The role of increasing temperature variability in European summer heatwaves. *Nature* 427, 332–336. doi: [10.1038/nature02300](https://doi.org/10.1038/nature02300)
- Schembari C, Cavalli F, Cuccia E, Hjorth J, Calzolari G et al. 2012 Impact of a European directive on ship emissions on air quality in Mediterranean harbours. *Atmos. Environ.* 61, 661–669. doi: [10.1016/j.atmosenv.2012.06.047](https://doi.org/10.1016/j.atmosenv.2012.06.047)
- Schneider A, Tanhua T, Körtzinger A, Wallace DWR 2010 High anthropogenic carbon content in the eastern Mediterranean. *J. Geophys. Res. Ocean.* 115. doi: [10.1029/2010JC006171](https://doi.org/10.1029/2010JC006171)
- Schroeder K, Chiggiato J, Bryden HL, Borghini M, Ben Ismail S 2016 Abrupt climate shift in the Western Mediterranean Sea. *Sci. Rep.* 6. doi: [10.1038/srep23009](https://doi.org/10.1038/srep23009)
- Schroeder K, Chiggiato J, Josey SA, Borghini M, Aracri S et al. 2017 Rapid response to climate change in a marginal sea. *Sci. Rep.* 7, 4065. doi: [10.1038/s41598-017-04455-5](https://doi.org/10.1038/s41598-017-04455-5)
- Schroeder K, García Lafuente J, Josey SA, Artale V, Buongiorno Nardelli B et al. 2012 Circulation of the Mediterranean Sea and its variability, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Elsevier). doi: [10.1016/B978-0-12-416042-2.00012-4](https://doi.org/10.1016/B978-0-12-416042-2.00012-4)
- Scyphers SB, Powers SP, Akins JL, Drymon JM, Martin CW et al. 2015 The role of citizens in detecting and responding to a rapid marine invasion. *Conserv. Lett.* 8, 242–250. doi: [10.1111/conl.12127](https://doi.org/10.1111/conl.12127)
- Seager R, Liu H, Henderson N, Simpson IR, Kelley C et al. 2014 Causes of increasing aridification of the Mediterranean region in response to rising greenhouse gases. *J. Clim.* 27, 4655–4676. doi: [10.1175/jcli-d-13-00446.1](https://doi.org/10.1175/jcli-d-13-00446.1)
- Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE et al. 2017 No saturation in the accumulation of alien species worldwide. *Nat. Commun.* 8, 14435. doi: [10.1038/ncomms14435](https://doi.org/10.1038/ncomms14435)
- Seebens H, Gastner MT, Blasius B 2013 The risk of marine bio-invasion caused by global shipping. *Ecol. Lett.* 16, 782–790. doi: [10.1111/ele.12111](https://doi.org/10.1111/ele.12111)
- Seinfeld JH, Pandis SN 2006 *Atmospheric chemistry and physics: from air pollution to climate change*. 2nd ed. Hoboken, N.J.: Wiley.
- Seljak G 2013 Dinamika vnosa tujerodnih fitofagnih žuželk in pršic v Slovenijo. *Acta Entomol. Slov.* 21, 85–122.
- Sellner KG, Doucette GJ, Kirkpatrick GJ 2003 Harmful algal blooms: Causes, impacts and detection. *J. Ind. Microbiol. Biotechnol.* 30, 383–406. doi: [10.1007/s10295-003-0074-9](https://doi.org/10.1007/s10295-003-0074-9)
- Sen OL, Ezber Y, Bozkurt D 2019 Euro-Mediterranean climate variability in boreal winter: A potential role of the East Asian trough. *Clim. Dyn.* 52, 7071–7084. doi: [10.1007/s00382-018-4573-9](https://doi.org/10.1007/s00382-018-4573-9)
- Senar JC, Domènech J, Arroyo L, Torre I, Gordo D 2016 An evaluation of monk parakeet damage to crops in the metropolitan area of Barcelona. *Anim. Biodivers. Conserv.* 39, 141–145. doi: [10.32800/abc.2016.39.0141](https://doi.org/10.32800/abc.2016.39.0141)
- Seneviratne SI, Lüthi D, Litschi M, Schär C 2006 Land-atmosphere coupling and climate change in Europe. *Nature* 443, 205–209. doi: [10.1038/nature05095](https://doi.org/10.1038/nature05095)
- Senouci R, Taibi N-E 2019 Impact of the urbanization on coastal dune: Case of Kharrouba, West of Algeria. *J. Sediment. Environ.* 4, 90–98. doi: [10.12957/jse.2019.39951](https://doi.org/10.12957/jse.2019.39951)
- Sevaut F, Somot S, Alias A, Dubois C, Lebeaupin-Brossier C et al. 2014 A fully coupled Mediterranean regional climate system model: design and evaluation of the ocean component for the 1980–2012 period. *Tellus A Dyn. Meteorol. Oceanogr.* 66, 23967. doi: [10.3402/tellusa.v66.23967](https://doi.org/10.3402/tellusa.v66.23967)
- Severin T, Conan P, de Madron XD, Houpert L, Oliver MJ et al. 2014 Impact of open-ocean convection on nutrients, phytoplankton biomass and activity. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 94, 62–71. doi: [10.1016/j.dsr.2014.07.015](https://doi.org/10.1016/j.dsr.2014.07.015)
- Shaltout M, Omstedt A 2014 Recent sea surface temperature trends and future scenarios for the Mediterranean Sea. *Oceanologia* 56, 411–443. doi: [10.5697/oc.56-3.411](https://doi.org/10.5697/oc.56-3.411)
- Shaw TA, Baldwin M, Barnes EA, Caballero R, Garfinkel CI et al. 2016 Storm track processes and the opposing influences of climate change. *Nat. Geosci.* 9, 656–664. doi: [10.1038/ngeo2783](https://doi.org/10.1038/ngeo2783)
- Sheffield J, Wood EF 2008 Global Trends and Variability in Soil Moisture and Drought Characteristics, 1950–2000, from Observation-Driven Simulations of the Terrestrial Hydrologic Cycle. *J. Clim.* 21, 432–458. doi: [10.1175/2007JCLI1822.1](https://doi.org/10.1175/2007JCLI1822.1)
- Shepherd TG 2014 Atmospheric circulation as a source of uncertainty in climate change projections. *Nat. Geosci.* 7, 703–708. doi: [10.1038/NNGEO2253](https://doi.org/10.1038/NNGEO2253)
- Shindell DT, Faluvegi G, Koch DM, Schmidt GA, Unger N et al. 2009 Improved Attribution of Climate Forcing to Emissions. *Science (80-. )*. 326, 716–718. doi: [10.1126/science.1174760](https://doi.org/10.1126/science.1174760)
- Shindell DT, Lamarque J-F, Schulz M, Flanner M, Jiao C et al. 2013 Radiative forcing in the ACCMIP historical and future

- climate simulations. *Atmos. Chem. Phys.* 13, 2939–2974. doi: [10.5194/acp-13-2939-2013](https://doi.org/10.5194/acp-13-2939-2013)
- Shomar BH, Müller G, Yahya A 2006 Occurrence of Pesticides in Groundwater and Topsoil of the Gaza Strip. *Water, Air, Soil Pollut.* 171, 237–251. doi: [10.1007/s11270-005-9038-1](https://doi.org/10.1007/s11270-005-9038-1)
- Sicard P, de Marco A, Troussier F, Renou C, Vas N et al. 2013 Decrease in surface ozone concentrations at Mediterranean remote sites and increase in the cities. *Atmos. Environ.* 79, 705–715. doi: [10.1016/j.atmosenv.2013.07.042](https://doi.org/10.1016/j.atmosenv.2013.07.042)
- Sillmann J, Kharin V V., Zwiers FW, Zhang X, Bronaugh D 2013 Climate extremes indices in the CMIP5 multimodel ensemble: Part 2. Future climate projections. *JGR Atmos.* 118, 2473–2493. doi: [10.1002/jgrd.50188](https://doi.org/10.1002/jgrd.50188)
- Simpson IR, Seager R, Shaw TA, Ting M 2015 Mediterranean summer climate and the importance of middle East topography. *J. Clim.* 28, 1977–1996. doi: [10.1175/jcli-d-14-00298.1](https://doi.org/10.1175/jcli-d-14-00298.1)
- Singh AN, Kumar P 2017 Emerging Contaminants – An Concise Overview.
- Sinovičić G, Franičević M, Zorica B, Čikeš-Keč V 2004 Short Communication Length-weight and length-length relationships for 10 pelagic fish species from the Adriatic Sea (Croatia). *J. Appl. Ichthyol.* 20, 156–158. doi: [10.1046/j.1439-0426.2003.00519.x](https://doi.org/10.1046/j.1439-0426.2003.00519.x)
- Sire A, Amouroux I 2016 Détermination de Valeurs Guides Environnementales (VGE) mollusques alternatives aux Normes de Qualité Environnementale (NQE) définies dans la DCE. <https://archimer.ifremer.fr/doc/00333/44378/> [Accessed April 17, 2019].
- Sisma-Ventura G, Yam R, Shemesh A 2014 Recent unprecedented warming and oligotrophy of the eastern Mediterranean Sea within the last millennium. *Geophys. Res. Lett.* 41, 5158–5166. doi: [10.1002/2014GL060393](https://doi.org/10.1002/2014GL060393)
- Skejic S, Car A, Marasovic I, Jozic S, Buzancic M et al. 2017 Morphology and ecology of the poorly known dinoflagellate *Prorocentrum arcuatum* (Dinophyceae) from the Medulin Bay (eastern Adriatic Sea). *Acta Adriat.* 58, 41–52.
- Skliris N, Zika JD, Herold L, Josey SA, Marsh R 2018 Mediterranean sea water budget long-term trend inferred from salinity observations. *Clim. Dyn.* 51, 2857–2876. doi: [10.1007/s00382-017-4053-7](https://doi.org/10.1007/s00382-017-4053-7)
- Slangen ABA, Adloff F, Jevrejeva S, Leclercq PW, Marzeion B et al. 2017 A review of recent updates of sea-level projections at global and regional scales. *Surv. Geophys.* 38, 395–416. doi: [10.1007/978-3-319-56490-6\\_17](https://doi.org/10.1007/978-3-319-56490-6_17)
- Slimani H, Aidoud A 2004 Desertification in the Maghreb: A case study of an Algerian high-plain steppe, in *Environmental Challenges in the Mediterranean 2000-2050*, ed. Marquina A [Dordrecht, Netherlands: Springer].
- Smith IM 1979 EPP0: the work of a regional plant protection organization, with particular reference to phytosanitary regulations, in *Plant Health: The Scientific Basis for Administrative Control of Plant Diseases and Pests*, eds. Ebbels DL, King JE [Oxford, UK: Blackwell Scientific Publications].
- Smouni A, Ater M, Auguy F, Laplaze L, El Mzibri M et al. 2010 Assessment of contamination by metallic trace elements in a mining area of eastern Morocco. *Cah. Agric.* 19, 273–279. doi: [10.1684/agr.2010.0413](https://doi.org/10.1684/agr.2010.0413)
- Sobrinho E, Sanz-Elorza M, Dana ED, González-Moreno A 2002 Invasibility of a coastal strip in NE Spain by alien plants. *J. Veg. Sci.* 13, 585–594. doi: [10.1111/j.1654-1103.2002.tb02085.x](https://doi.org/10.1111/j.1654-1103.2002.tb02085.x)
- Somot S, Houpert L, Sevault F, Testor P, Bosse A et al. 2018 Characterizing, modelling and understanding the climate variability of the deep water formation in the North-Western Mediterranean Sea. *Clim. Dyn.* 51, 1179–1210.
- Somot S, Sevault F, Déqué M 2006 Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model. *Clim. Dyn.* 27, 851–879. doi: [10.1007/s00382-006-0167-z](https://doi.org/10.1007/s00382-006-0167-z)
- Somot S, Sevault F, Déqué M, Crépon M 2008 21<sup>st</sup> century climate change scenario for the Mediterranean using a coupled atmosphere-ocean regional climate model. *Glob. Planet. Change* 63, 112–126. doi: [10.1016/j.gloplacha.2007.10.003](https://doi.org/10.1016/j.gloplacha.2007.10.003)
- Soto-Navarro J, Jordà G, Amores A, Cabos W, Somot S et al. 2020 Evolution of Mediterranean Sea water properties under climate change scenarios in the Med-CORDEX ensemble. *Clim. Dyn.* 54, 2135–2165. doi: [10.1007/s00382-019-05105-4](https://doi.org/10.1007/s00382-019-05105-4)
- Soto-Navarro J, Somot S, Sevault F, Beuvier J, Criado-Aldeanueva F et al. 2015 Evaluation of regional ocean circulation models for the Mediterranean Sea at the Strait of Gibraltar: volume transport and thermohaline properties of the outflow. *Clim. Dyn.* 44, 1277–1292. doi: [10.1007/s00382-014-2179-4](https://doi.org/10.1007/s00382-014-2179-4)
- Spinoni J, Naumann G, Vogt J, Barbosa P 2015 European drought climatologies and trends based on a multi-indicator approach. *Glob. Planet. Change* 127, 50–57. doi: [10.1016/j.gloplacha.2015.01.012](https://doi.org/10.1016/j.gloplacha.2015.01.012)
- Spinoni J, Vogt J V., Naumann G, Barbosa P, Dosio A 2018 Will drought events become more frequent and severe in Europe? *Int. J. Climatol.* 38, 1718–1736. doi: [10.1002/joc.5291](https://doi.org/10.1002/joc.5291)
- Squadrone S, Brizio P, Stella C, Prearo M, Pastorino P et al. 2016 Presence of trace metals in aquaculture marine ecosystems of the northwestern Mediterranean Sea (Italy). *Environ. Pollut.* 215, 77–83. doi: [10.1016/j.envpol.2016.04.096](https://doi.org/10.1016/j.envpol.2016.04.096)
- Stachowicz JJ, Terwin JR, Whitlatch RB, Osman RW 2002 Linking climate change and biological invasions: Ocean warming facilitates nonindigenous species invasions. *Proc. Natl. Acad. Sci. U. S. A.* 99, 15497–15500. doi: [10.1073/pnas.242437499](https://doi.org/10.1073/pnas.242437499)
- Stagge JH, Kohn I, Tallaksen LM, Stahl K 2015 Modeling drought impact occurrence based on meteorological drought indices in Europe. *J. Hydrol.* 530, 37–50. doi: [10.1016/j.jhydrol.2015.09.039](https://doi.org/10.1016/j.jhydrol.2015.09.039)
- Stamatis N, Hela D, Triantafyllidis V, Konstantinou I 2013 Spatiotemporal variation and risk assessment of pesticides in water of the lower catchment basin of Acheloos River, Western Greece. *ScientificWorldJournal.* 2013, 231610. doi: [10.1155/2013/231610](https://doi.org/10.1155/2013/231610)
- Statuto D, Cillis G, Picuno P 2016 Analysis of the effects of agricultural land use change on rural environment and

- landscape through historical cartography and GIS tools. *J. Agric. Eng.* 47, 468.
- Stehle S, Schulz R 2015 Agricultural insecticides threaten surface waters at the global scale. *Proc. Natl. Acad. Sci. U. S. A.* 112, 5750–5755. doi: [10.1073/pnas.1500232112](https://doi.org/10.1073/pnas.1500232112)
- Stephenson DB, Pavan V, Collins M, Junge MM, Quadrelli R 2006 North Atlantic Oscillation response to transient greenhouse gas forcing and the impact on European winter climate: A CMIP2 multi-model assessment. *Clim. Dyn.* 27, 401–420. doi: [10.1007/s00382-006-0140-x](https://doi.org/10.1007/s00382-006-0140-x)
- Stergiou KI, Somarakis S, Triantafyllou G, Tsiaras KP, Giannoulaki M et al. 2016 Trends in productivity and biomass yields in the Mediterranean Sea Large Marine Ecosystem during climate change. *Environ. Dev.* 17, 57–74. doi: [10.1016/j.envdev.2015.09.001](https://doi.org/10.1016/j.envdev.2015.09.001)
- Stergiou KI, Tsikliras AC 2011 Fishing down, fishing through and fishing up: fundamental process versus technical details. *Mar. Ecol. Prog. Ser.* 441, 295–301. doi: [10.3354/meps09377](https://doi.org/10.3354/meps09377)
- Stinca A, Chirico GB, Incerti G, Bonanomi G 2015 Regime shift by an exotic nitrogen-fixing shrub mediates plant facilitation in primary succession. *PLoS One* 10, e0123128. doi: [10.1371/journal.pone.0123128](https://doi.org/10.1371/journal.pone.0123128)
- Stohl A, Aamaas B, Amann M, Baker LH, Bellouin N et al. 2015 Evaluating the climate and air quality impacts of short-lived pollutants. *Atmos. Chem. Phys.* 15, 10529–10566. doi: [10.5194/acp-15-10529-2015](https://doi.org/10.5194/acp-15-10529-2015)
- Storkey J, Stratonovitch P, Chapman DS, Vidotto F, Semenov MA 2014 A Process-Based Approach to Predicting the Effect of Climate Change on the Distribution of an Invasive Allergenic Plant in Europe. *PLoS One* 9, e88156. doi: [10.1371/journal.pone.0088156](https://doi.org/10.1371/journal.pone.0088156)
- Strode SA, Ott LE, Pawson S, Bowyer TW 2012 Emission and transport of cesium-137 from boreal biomass burning in the summer of 2010. *JGR Atmos.* 117, n/a-n/a. doi: [10.1029/2011JD017382](https://doi.org/10.1029/2011JD017382)
- Sturaro E, Thiene M, Cocca G, Mrad M, Tempesta T et al. 2013 Factors influencing summer farms management in the Alps. *Ital. J. Anim. Sci.* 12, e25. doi: [10.4081/ijas.2013.e25](https://doi.org/10.4081/ijas.2013.e25)
- Suffert M, Wilstermann A, Petter F, Schrader G, Grousset F 2018 Identification of new pests likely to be introduced into Europe with the fruit trade. *Bull. OEPP/EPPO* 48, 144–154.
- Sumaila UR, Cheung WWL, Lam WVY, Pauly D, Herrick S 2011 Climate change impacts on the biophysics and economics of world fisheries. *Nat. Clim. Chang.* 1, 449–456. doi: [10.1038/nclimate1301](https://doi.org/10.1038/nclimate1301)
- Sumaila UR, Tai TC, Lam WVY, Cheung WWL, Bailey M et al. 2019 Benefits of the Paris Agreement to ocean life, economies, and people. *Sci. Adv.* 5, eaau3855. doi: [10.1126/sciadv.aau3855](https://doi.org/10.1126/sciadv.aau3855)
- Sun X, Ren G, You Q, Ren Y, Xu W et al. 2019 Global diurnal temperature range (DTR) changes since 1901. *Clim. Dyn.* 52, 3343–3356. doi: [10.1007/s00382-018-4329-6](https://doi.org/10.1007/s00382-018-4329-6)
- Taillefumier F, Piégay H 2003 Contemporary land use changes in Prealpine Mediterranean mountains. A multivariate GIS-based approach applied to two municipalities in the Southern French Prealps. *Catena* 51, 267–296. doi: [10.1016/S0341-8162\(02\)00168-6](https://doi.org/10.1016/S0341-8162(02)00168-6)
- Talbi A, Kerchich Y, Kerbachi R, Boughedaoui M 2018 Assessment of annual air pollution levels with PM1, PM2.5, PM10 and associated heavy metals in Algiers, Algeria. *Environ. Pollut.* 232, 252–263. doi: [10.1016/j.envpol.2017.09.041](https://doi.org/10.1016/j.envpol.2017.09.041)
- Tanarhte M, Hadjinicolaou P, Lelieveld J 2012 Intercomparison of temperature and precipitation data sets based on observations in the Mediterranean and the Middle East. *JGR Atmos.* 117, n/a-n/a. doi: [10.1029/2011jd017293](https://doi.org/10.1029/2011jd017293)
- Tanasijevic L, Todorovic M, Pereira LS, Pizzigalli C, Lionello P 2014 Impacts of climate change on olive crop evapotranspiration and irrigation requirements in the Mediterranean region. *Agric. Water Manag.* 144, 54–68. doi: [10.1016/J.AGWAT.2014.05.019](https://doi.org/10.1016/j.agwat.2014.05.019)
- Tanhua T, Hainbucher D, Schroeder K, Cardin V, Alvarez M et al. 2013 The Mediterranean Sea system: a review and an introduction to the special issue. *Ocean Sci.* 9, 789–803. doi: [10.5194/os-9-789-2013](https://doi.org/10.5194/os-9-789-2013)
- Tarabon S, Bertrand R, Lavoie C, Vigouroux T, Isselin-Nondedeu F 2018 The effects of climate warming and urbanised areas on the future distribution of *Cortaderia seloana*, pampas grass, in France. *Weed Res.* 58, 413–423. doi: [10.1111/wre.12330](https://doi.org/10.1111/wre.12330)
- Taylor KE, Stouffer RJ, Meehl GA 2012 An Overview of CMIP5 and the Experiment Design. *Bull. Am. Meteorol. Soc.* 93, 485–498. doi: [10.1175/BAMS-D-11-00094.1](https://doi.org/10.1175/BAMS-D-11-00094.1)
- Taylor PD, Fahrig L, Henein K, Merriam G 1993 Connectivity is a vital element of landscape structure. *Oikos* 68, 571–573.
- Tchounwou PB, Yedjou CG, Patlolla AK, Sutton DJ 2012 Heavy metal toxicity and the environment. *Exp. Suppl.* 101, 133–164. doi: [10.1007/978-3-7643-8340-4\\_6](https://doi.org/10.1007/978-3-7643-8340-4_6)
- Testor P, Bosse A, Houpert L, Margirier F, Mortier L et al. 2018 Multiscale Observations of Deep Convection in the Northwestern Mediterranean Sea During Winter 2012–2013 Using Multiple Platforms. *J. Geophys. Res. Ocean.* 123, 1745–1776. doi: [10.1002/2016JC012671](https://doi.org/10.1002/2016JC012671)
- Testor P, de Madron XD, Mortier L, D'Ortenzio F, Legoff H et al. 2019 LION observatory data. *SEANOE*. doi: [10.17882/44411](https://doi.org/10.17882/44411)
- Thiéblemont R, Le Cozannet G, Toimil A, Meyssignac B, Losada IJ 2019 Likely and High-End Impacts of Regional Sea-Level Rise on the Shoreline Change of European Sandy Coasts Under a High Greenhouse Gas Emissions Scenario. *Water* 11, 2607.
- Thierry W, Davin EL, Lawrence DM, Hirsch AL, Hauser M et al. 2017 Present-day irrigation mitigates heat extremes. *J. Geophys. Res. Atmos.* 122, 1403–1422. doi: [10.1002/2016JD025740](https://doi.org/10.1002/2016JD025740)
- Thornes JB 2009 Land degradation, in *The Physical Geography of the Mediterranean*, ed. Woodward J (New York: Oxford University Press), 563–581.
- Thorpe A, Harrison RM 2008 Sources and properties of non-exhaust particulate matter from road traffic: A review. *Sci. Total Environ.* 400, 270–282. doi: [10.1016/J.SCITOTENV.2008.06.007](https://doi.org/10.1016/J.SCITOTENV.2008.06.007)
- Thorpe RB, Bigg GR 2000 Modelling the sensitivity of Mediterranean outflow to anthropogenically forced climate change. *Clim. Dyn.* 16, 355–368. doi: [10.1007/s003820050333](https://doi.org/10.1007/s003820050333)
- Titelboim D, Almogi-Labin A, Herut B, Kucera M, Askenazi-Polivoda S et al. 2019 Thermal tolerance and range

- expansion of invasive foraminifera under climate changes. *Sci. Rep.* 9, 1–5. doi: [10.1038/s41598-019-40944-5](https://doi.org/10.1038/s41598-019-40944-5)
- Titelboim D, Sadekov A, Almogi-Labin A, Herut B, Kucera M et al. 2017 Geochemical signatures of benthic foraminiferal shells from a heat-polluted shallow marine environment provide field evidence for growth and calcification under extreme warmth. *Glob. Chang. Biol.* 23, 4346–4353. doi: [10.1111/gcb.13729](https://doi.org/10.1111/gcb.13729)
- Tiwari D, Kamble J, Chilgunde S, Patil P, Maru G et al. 2012 Clastogenic and mutagenic effects of bisphenol A: An endocrine disruptor. *Mutat. Res. - Genet. Toxicol. Environ. Mutagen.* 743, 83–90. doi: [10.1016/j.mrgentox.2011.12.023](https://doi.org/10.1016/j.mrgentox.2011.12.023)
- Tobin I, Vautard R, Balog I, Bréon F-M, Jerez S et al. 2015 Assessing climate change impacts on European wind energy from ENSEMBLES high-resolution climate projections. *Clim. Change* 128, 99–112. doi: [10.1007/s10584-014-1291-0](https://doi.org/10.1007/s10584-014-1291-0)
- Tolika K 2019 Assessing Heat Waves over Greece Using the Excess Heat Factor (EHF). *Climate* 7, 9. doi: [10.3390/cli7010009](https://doi.org/10.3390/cli7010009)
- Toreti A, Xoplaki E, Maraun D, Kuglitsch FG, Wanner H et al. 2010 Characterisation of extreme winter precipitation in Mediterranean coastal sites and associated anomalous atmospheric circulation patterns. *Nat. Hazards Earth Syst. Sci.* 10, 1037–1050. doi: [10.5194/nhess-10-1037-2010](https://doi.org/10.5194/nhess-10-1037-2010)
- Torma CZ 2019 Detailed validation of EURO-CORDEX and Med-CORDEX regional climate model ensembles over the Carpathian Region. *Q. J. Hungarian Meteorol. Serv.* 123, 217–240.
- Torma CZ, Giorgi F, Coppola E 2015 Added value of regional climate modeling over areas characterized by complex terrain-Precipitation over the Alps. *JGR Atmos.* 120, 3957–3972. doi: [10.1002/2014JD022781](https://doi.org/10.1002/2014JD022781)
- Touaylia S, Ghannem S, Toumi H, Bejaoui M 2016 Assessment of heavy metals status in northern Tunisia using contamination indices: Case of the Ichkeul steams system. 3, 209–217.
- Touratier F, Goyet C 2011 Impact of the Eastern Mediterranean Transient on the distribution of anthropogenic CO<sub>2</sub> and first estimate of acidification for the Mediterranean Sea. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 58, 1–15. doi: [10.1016/j.dsr.2010.10.002](https://doi.org/10.1016/j.dsr.2010.10.002)
- Tous M, Romero R 2013 Meteorological environments associated with medicane development. *Int. J. Climatol.* 33, 1–14. doi: [10.1002/joc.3428](https://doi.org/10.1002/joc.3428)
- Tous M, Zappa G, Romero R, Shaffrey L, Vidale PL 2016 Projected changes in medicanes in the HadGEM3 N512 high-resolution global climate model. *Clim. Dyn.* 47, 1913–1924. doi: [10.1007/s00382-015-2941-2](https://doi.org/10.1007/s00382-015-2941-2)
- Tovar-Sánchez A, Basterretxea G, Ben Omar M, Jordi A, Sánchez-Quiles D et al. 2016 Nutrients, trace metals and B-vitamin composition of the Moulouya River: A major North African river discharging into the Mediterranean Sea. *Estuar. Coast. Shelf Sci.* 176, 47–57. doi: [10.1016/j.ecss.2016.04.006](https://doi.org/10.1016/j.ecss.2016.04.006)
- Traina A, Bono G, Bonsignore M, Falco F, Giuga M et al. 2019 Heavy metals concentrations in some commercially key species from Sicilian coasts (Mediterranean Sea): Potential human health risk estimation. *Ecotoxicol. Environ. Saf.* 168, 466–478. doi: [10.1016/j.ecoenv.2018.10.056](https://doi.org/10.1016/j.ecoenv.2018.10.056)
- Tramblay Y, Feki H, Quintana-Seguí P, Guijarro JA 2019 The SAFRAN daily gridded precipitation product in Tunisia (1979–2015). *Int. J. Climatol.*, joc.6181. doi: [10.1002/joc.6181](https://doi.org/10.1002/joc.6181)
- Tribouillois H, Constantin J, Justes E 2018 Cover crops mitigate direct greenhouse gases balance but reduce drainage under climate change scenarios in temperate climate with dry summers. *Glob. Chang. Biol.* 24, 2513–2529. doi: [10.1111/gcb.14091](https://doi.org/10.1111/gcb.14091)
- Trigo IF, Bigg GR, Davies TD 2002a Climatology of Cyclogenesis Mechanisms in the Mediterranean. *Mon. Weather Rev.* 130, 549–569. doi: [10.1175/1520-0493\(2002\)130<0549:COCMIT>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<0549:COCMIT>2.0.CO;2)
- Trigo IF, Davies TD, Bigg GR 1999 Objective Climatology of Cyclones in the Mediterranean Region. *J. Clim.* 12, 1685–1696. doi: [10.1175/1520-0442\(1999\)012<1685:OCOCIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1685:OCOCIT>2.0.CO;2)
- Trigo IF, Davies TD, Bigg GR 2000 Decline in Mediterranean rainfall caused by weakening of Mediterranean cyclones. *Geophys. Res. Lett.* 27, 2913–2916. doi: [10.1029/2000GL011526](https://doi.org/10.1029/2000GL011526)
- Trigo R, Xoplaki E, Zorita E, Luterbacher J, Krichak SO et al. 2006 Chapter 3 Relations between variability in the Mediterranean region and mid-latitude variability, in *Mediterranean*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo RBT-D in E and ES (Elsevier), 179–226. doi: [10.1016/S1571-9197\(06\)80006-6](https://doi.org/10.1016/S1571-9197(06)80006-6)
- Trigo RM, Osborn TJ, Corte-Real JM 2002b The North Atlantic Oscillation influence on Europe: climate impacts and associated physical mechanisms. *Clim. Res.* 20, 9–17. doi: [10.3354/cr020009](https://doi.org/10.3354/cr020009)
- Trigo RM, Pozo-Vázquez D, Osborn TJ, Castro-Díez Y, Gámiz-Fortis S et al. 2004 North Atlantic oscillation influence on precipitation, river flow and water resources in the Iberian Peninsula. *Int. J. Climatol.* 24, 925–944. doi: [10.1002/joc.1048](https://doi.org/10.1002/joc.1048)
- Trouet V 2014 A Tree-Ring Based Late Summer Temperature Reconstruction (AD 1675–1980) for the Northeastern Mediterranean. *Radiocarbon* 56, S69–S78. doi: [10.1017/s003822200050372](https://doi.org/10.1017/s003822200050372)
- Tsikliras AC 2008 Climate-related geographic shift and sudden population increase of a small pelagic fish (*Sardinella aurita*) in the eastern Mediterranean Sea. *Mar. Biol. Res.* 4, 477–481. doi: [10.1080/17451000802291292](https://doi.org/10.1080/17451000802291292)
- Tsikliras AC 2014 Fisheries Mismanagement in the Mediterranean: A Greek Tragedy. *Fish. Aquac. J.* 05. doi: [10.4172/2150-3508.1000e113](https://doi.org/10.4172/2150-3508.1000e113)
- Tsikliras AC, Dinouli A, Tsalkou E 2013a Exploitation trends of the Mediterranean and Black Sea fisheries. *Acta Adriat.* 54, 273–282.
- Tsikliras AC, Dinouli A, Tsiros V-Z, Tsalkou E 2015 The Mediterranean and Black Sea Fisheries at Risk from Overexploitation. *PLoS One* 10, e0121188. doi: [10.1371/journal.pone.0121188](https://doi.org/10.1371/journal.pone.0121188)
- Tsikliras AC, Licandro P, Pardalou A, McQuinn IH, Gröger JP et al. 2019 Synchronization of Mediterranean pelagic fish populations with the North Atlantic climate variability. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 159, 143–151.

- doi: [10.1016/j.dsr2.2018.07.005](https://doi.org/10.1016/j.dsr2.2018.07.005)
- Tsikliras AC, Stergiou KI 2014 Mean temperature of the catch increases quickly in the Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 515, 281–284. doi: [10.3354/meps11005](https://doi.org/10.3354/meps11005)
- Tsikliras AC, Tsiros VZ, Stergiou KI 2013b Assessing the state of Greek marine fisheries resources. *Fish. Manag. Ecol.* 20, 34–41. doi: [10.1111/j.1365-2400.2012.00863.x](https://doi.org/10.1111/j.1365-2400.2012.00863.x)
- Tsimplis MN, Álvarez-Fanjul E, Gomis D, Fenoglio-Marc L, Pérez B 2005 Mediterranean Sea level trends: Atmospheric pressure and wind contribution. *Geophys. Res. Lett.* 32, L20602. doi: [10.1029/2005GL023867](https://doi.org/10.1029/2005GL023867)
- Tsimplis MN, Zervakis V, Josey S, Peneva E, Struglia MV et al. 2006 Variability of the Mediterranean sea level and oceanic circulation and their relation to climate patterns, in *Mediterranean climate variability*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo R [Elsevier].
- Tsopelas P, Santini A, Wingfield MJ, de Beer ZW 2017 Canker stain: A lethal disease destroying iconic plane trees. *Plant Dis.* 101, 645–658. doi: [10.1094/PDIS-09-16-1235-FE](https://doi.org/10.1094/PDIS-09-16-1235-FE)
- Turbé A, Strubbe D, Mori E, Carrete M, Chiron F et al. 2017 Assessing the assessments: evaluation of four impact assessment protocols for invasive alien species. *Divers. Distrib.* 23, 297–307. doi: [10.1111/ddi.12528](https://doi.org/10.1111/ddi.12528)
- Turco M, Bedía J, di Liberto F, Fiorucci P, Von Hardenberg J et al. 2016 Decreasing fires in mediterranean Europe. *PLoS One* 11. doi: [10.1371/journal.pone.0150663](https://doi.org/10.1371/journal.pone.0150663)
- Turco M, Jerez S, Augusto S, Tarín-Carrasco P, Ratola N et al. 2019 Climate drivers of the 2017 devastating fires in Portugal. *Sci. Rep.* 9, 13886. doi: [10.1038/s41598-019-50281-2](https://doi.org/10.1038/s41598-019-50281-2)
- Turco M, Llasat MC, von Hardenberg J, Provenzale A 2014 Climate change impacts on wildfires in a Mediterranean environment. *Clim. Change* 125, 369–380. doi: [10.1007/s10584-014-1183-3](https://doi.org/10.1007/s10584-014-1183-3)
- Turco M, Rosa-Cánovas JJ, Bedía J, Jerez S, Montávez JP et al. 2018 Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models. *Nat. Commun.* 9, 3821. doi: [10.1038/s41467-018-06358-z](https://doi.org/10.1038/s41467-018-06358-z)
- Türkeş M, Ertat E 2003 Precipitation changes and variability in Turkey linked to the North Atlantic oscillation during the period 1930–2000. *Int. J. Climatol.* 23, 1771–1796. doi: [10.1002/joc.962](https://doi.org/10.1002/joc.962)
- Tyrlis E, Hoskins BJ 2008 Aspects of a northern hemisphere atmospheric blocking climatology. *J. Atmos. Sci.* 65, 1638–1652. doi: [10.1175/2007JAS2337.1](https://doi.org/10.1175/2007JAS2337.1)
- Tyrlis E, Lelieveld J 2013 Climatology and Dynamics of the Summer Etesian Winds over the Eastern Mediterranean\*. *J. Atmos. Sci.* 70, 3374–3396. doi: [10.1175/jas-d-13-035.1](https://doi.org/10.1175/jas-d-13-035.1)
- Tzanatos E, Raitzos DE, Triantafyllou G, Somarakis S, Tsonis AA 2014 Indications of a climate effect on Mediterranean fisheries. *Clim. Change* 122, 41–54. doi: [10.1007/s10584-013-0972-4](https://doi.org/10.1007/s10584-013-0972-4)
- Ulbrich U, Christoph M 1999 A shift of the NAO and increasing storm track activity over Europe due to anthropogenic greenhouse gas forcing. *Clim. Dyn.* 15, 551–559. doi: [10.1007/s003820050299](https://doi.org/10.1007/s003820050299)
- Ulbrich U, Leckebusch GC, Pinto JG 2009 Extra-tropical cyclones in the present and future climate: A review. *Theor. Appl. Climatol.* 96, 117–131. doi: [10.1007/s00704-008-0083-8](https://doi.org/10.1007/s00704-008-0083-8)
- Ulbrich U, Lionello P, Belušić D, Jacobeit J, Knippertz P et al. 2012 Climate of the Mediterranean: Synoptic patterns, temperature, precipitation, winds and their extremes, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P [Elsevier], 1–61.
- Ulbrich U, Pinto JG, Kupfer H, Leckebusch GC, Spanghel T et al. 2008 Changing Northern Hemisphere Storm Tracks in an Ensemble of IPCC Climate Change Simulations. *J. Clim.* 21, 1669–1679. doi: [10.1175/2007jcli1992.1](https://doi.org/10.1175/2007jcli1992.1)
- Ullah S, Zuberi A, Alagawany M, Farag MR, Dadar M et al. 2018 Cypermethrin induced toxicities in fish and adverse health outcomes: Its prevention and control measure adaptation. *J. Environ. Manage.* 206, 863–871. doi: [10.1016/j.jenvman.2017.11.076](https://doi.org/10.1016/j.jenvman.2017.11.076)
- UN-Water 2014 *The United Nations World Water Development Report 3*. doi: [10.4324/9781849773355](https://doi.org/10.4324/9781849773355)
- UN 2015 Transforming our world: the 2030 Agenda for Sustainable Development. New York [https://www.un.org/ga/search/view\\_doc.asp?symbol=A/RES/70/1&Lang=E](https://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E)
- Underwood EC, Viers JH, Klausmeyer KR, Cox RL, Shaw MR 2009 Threats and biodiversity in the mediterranean biome. *Divers. Distrib.* 15, 188–197. doi: [10.1111/j.1472-4642.2008.00518.x](https://doi.org/10.1111/j.1472-4642.2008.00518.x)
- UNEP-GRID Arendal 2013 State of the Mediterranean Marine and Coastal Environment.
- UNEP/MAP 2012a Initial integrated assessment of the Mediterranean sea: fulfilling step 3 of the ecosystem approach process.
- UNEP/MAP 2012b State of the Mediterranean Marine and Coastal Environment. UNEP/MAP – Barcelona Convention, Athens.
- UNEP/MAP 2017 2017 Mediterranean Quality Status Report.
- Valladares F, Matesanz S, Guilhaumon F, Araújo MB, Balaguer L et al. 2014 The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecol. Lett.* 17, 1351–1364. doi: [10.1111/ele.12348](https://doi.org/10.1111/ele.12348)
- Van der Sluis T, Pedrolí B, Kristensen SBP, Cosor GL, Pavlis E 2016 Changing land use intensity in Europe—Recent processes in selected case studies. *Land use policy* 57, 777–785. doi: [10.1016/j.landusepol.2014.12.005](https://doi.org/10.1016/j.landusepol.2014.12.005)
- Van Dingenen R, Raes F, Putaud J-P, Baltensperger U, Charron A et al. 2004 A European aerosol phenomenology—1: physical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe. *Atmos. Environ.* 38, 2561–2577. doi: [10.1016/j.atmosenv.2004.01.040](https://doi.org/10.1016/j.atmosenv.2004.01.040)
- Van Kleunen M, Essl F, Pergl J, Brundu G, Carboni M et al. 2018 The changing role of ornamental horticulture in alien plant invasions. *Biol. Rev.* 93, 1421–1437. doi: [10.1111/brv.12402](https://doi.org/10.1111/brv.12402)
- van Ruijven BJ, Levy MA, Agrawal A, Biermann F, Birkmann J et al. 2014 Enhancing the relevance of Shared Socioeconomic Pathways for climate change impacts, adaptation and vulnerability research. *Clim. Change* 122, 481–494. doi: [10.1007/s10584-013-0931-0](https://doi.org/10.1007/s10584-013-0931-0)
- van Vuuren DP, Edmonds JE, Kainuma M, Riahi K, Thomson A

- et al. 2011 The representative concentration pathways: An overview. *Clim. Change* 109, 5–31. doi: [10.1007/s10584-011-0148-z](https://doi.org/10.1007/s10584-011-0148-z)
- Vanderhoeven S, Branquart E, Casaer J, D'hondt B, Hulme PE et al. 2017 Beyond protocols: improving the reliability of expert-based risk analysis underpinning invasive species policies. *Biol. Invasions* 19, 2507–2517. doi: [10.1007/s10530-017-1434-0](https://doi.org/10.1007/s10530-017-1434-0)
- Vargas-Yáñez M, García-Martínez MC, Moya F, Balbín R, López-Jurado JL et al. 2017 Updating temperature and salinity mean values and trends in the Western Mediterranean: The RADMED project. *Prog. Oceanogr.* 157, 27–46. doi: [10.1016/j.poccean.2017.09.004](https://doi.org/10.1016/j.poccean.2017.09.004)
- Vargas-Yáñez M, Zunino P, Benali A, Delpy M, Pastre F et al. 2010 How much is the western Mediterranean really warming and salting? *JGR Atmos.* 115, C04001. doi: [10.1029/2009jc005816](https://doi.org/10.1029/2009jc005816)
- Vasilakopoulos P, Maravelias CD, Tserpes G 2014 The alarming decline of mediterranean fish stocks. *Curr. Biol.* 24, 1643–1648. doi: [10.1016/j.cub.2014.05.070](https://doi.org/10.1016/j.cub.2014.05.070)
- Vasilakopoulos P, Raitsois DE, Tzanatos E, Maravelias CD 2017 Resilience and regime shifts in a marine biodiversity hotspot. *Sci. Rep.* 7. doi: [10.1038/s41598-017-13852-9](https://doi.org/10.1038/s41598-017-13852-9)
- Vautard R, Cattiaux J, Yiou P, Thépaut J-N, Ciais P 2010 Northern Hemisphere atmospheric stilling partly attributed to an increase in surface roughness. *Nat. Geosci.* 3, 756–761. doi: [10.1038/ngeo979](https://doi.org/10.1038/ngeo979)
- Vautard R, Gobiet A, Sobolowski SP, Kjellström E, Stegehuis A et al. 2014 The European climate under a 2°C global warming. *Environ. Res. Lett.* 9, 34006. doi: [10.1088/1748-9326/9/3/034006](https://doi.org/10.1088/1748-9326/9/3/034006)
- Vayreda J, Martínez-Vilalta J, Gracia M, Retana J 2012 Recent climate changes interact with stand structure and management to determine changes in tree carbon stocks in Spanish forests. *Glob. Chang. Biol.* 18, 1028–1041. doi: [10.1111/j.1365-2486.2011.02606.x](https://doi.org/10.1111/j.1365-2486.2011.02606.x)
- Velaoras D, Kassis D, Perivoliotis L, Pagonis P, Hondronasios A et al. 2013 Temperature and salinity variability in the Greek Seas based on POSEIDON stations time series: preliminary results. *Mediterr. Mar. Sci.* 14, 5–18. doi: [10.12681/mms.446](https://doi.org/10.12681/mms.446)
- Verdura J, Linares C, Ballesteros E, Coma R, Uriz MJ et al. 2019 Biodiversity loss in a Mediterranean ecosystem due to an extreme warming event unveils the role of an engineering gorgonian species. *Sci. Rep.* 9. doi: [10.1038/s41598-019-41929-0](https://doi.org/10.1038/s41598-019-41929-0)
- Verfaillie D, Lafaysse M, Déqué M, Eckert N, Lejeune Y et al. 2018 Multi-component ensembles of future meteorological and natural snow conditions for 1500 m altitude in the Chartreuse mountain range, Northern French Alps. *Cryosph.* 12, 1249–1271. doi: [10.5194/tc-12-1249-2018](https://doi.org/10.5194/tc-12-1249-2018)
- Vergés A, Doropoulos C, Malcolm HA, Skye M, García-Pizá M et al. 2016 Long-term empirical evidence of ocean warming leading to tropicalization of fish communities, increased herbivory, and loss of kelp. *Proc. Natl. Acad. Sci. U. S. A.* 113, 13791–13796. doi: [10.1073/pnas.1610725113](https://doi.org/10.1073/pnas.1610725113)
- Vergés A, Steinberg PD, Hay ME, Poore AGB, Campbell AH et al. 2014a The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B Biol. Sci.* 281, 20140846. doi: [10.1098/rspb.2014.0846](https://doi.org/10.1098/rspb.2014.0846)
- Vergés A, Tomas F, Cebrián E, Ballesteros E, Kizilkaya Z et al. 2014b Tropical rabbitfish and the deforestation of a warming temperate sea. *J. Ecol.* 102, 1518–1527. doi: [10.1111/1365-2745.12324](https://doi.org/10.1111/1365-2745.12324)
- Verheyen J, Stoks R 2019 Current and future daily temperature fluctuations make a pesticide more toxic: Contrasting effects on life history and physiology. *Environ. Pollut.* 248, 209–218. doi: [10.1016/j.envpol.2019.02.022](https://doi.org/10.1016/j.envpol.2019.02.022)
- Verkerk PJ, Sánchez A, Libbrecht S, Broekman A, Bruggeman A et al. 2017 A participatory approach for adapting river basins to climate change. *Water* 9, 958. doi: [10.3390/w9120958](https://doi.org/10.3390/w9120958)
- Vettraino AM, Roques A, Yart A, Fan J, Sun J et al. 2015 Sentinel trees as a tool to forecast invasions of alien plant pathogens. *PLoS One* 10, e0120571. doi: [10.1371/journal.pone.0120571](https://doi.org/10.1371/journal.pone.0120571)
- Vicente-Serrano SM, Domínguez-Castro F, Murphy C, Hannaford J, Reig F et al. 2020 Long-term variability and trends in meteorological droughts in Western Europe (1851–2018). *Int. J. Climatol.* n/a. doi: [10.1002/joc.6719](https://doi.org/10.1002/joc.6719)
- Vicente-Serrano SM, López Moreno JI, Beguería S, Lorenzo-Lacruz J, Sánchez-Lorenzo A et al. 2014 Evidence of increasing drought severity caused by temperature rise in southern Europe. *Environ. Res. Lett.* 9. doi: [10.1088/1748-9326/9/4/044001](https://doi.org/10.1088/1748-9326/9/4/044001)
- Vicente-Serrano SM, López Moreno JI, Gimeno L, Nieto R, Morán-Tejeda E et al. 2011 A multiscale global evaluation of the impact of ENSO on droughts. *JGR Atmos.* 116, D20109. doi: [10.1029/2011JD016039](https://doi.org/10.1029/2011JD016039)
- Vicente-Serrano SM, Van der Schrier G, Beguería S, Azorin-Molina C, López Moreno JI 2015 Contribution of precipitation and reference evapotranspiration to drought indices under different climates. *J. Hydrol.* 526, 42–54. doi: [10.1016/j.jhydrol.2014.11.025](https://doi.org/10.1016/j.jhydrol.2014.11.025)
- Vieira G, Mora C, Faleh A 2017 New observations indicate the possible presence of permafrost in North Africa (Djebel Toubkal, High Atlas, Morocco). *Cryosph.* 11, 1691–1705. doi: [10.5194/tc-11-1691-2017](https://doi.org/10.5194/tc-11-1691-2017)
- Vilà-Cabrera A, Coll L, Martínez-Vilalta J, Retana J 2018 Forest management for adaptation to climate change in the Mediterranean basin: A synthesis of evidence. *For. Ecol. Manage.* 407, 16–22. doi: [10.1016/j.foreco.2017.10.021](https://doi.org/10.1016/j.foreco.2017.10.021)
- Vila M, Camp J, Garces E, Maso M, Delgado M 2001 High resolution spatio-temporal detection of potentially harmful dinoflagellates in confined waters of the NW Mediterranean. *J. Plankton Res.* 23, 497–514. doi: [10.1093/plankt/23.5.497](https://doi.org/10.1093/plankt/23.5.497)
- Vilà M, Espinar JL, Hejda M, Hulme PE, Jarošík V et al. 2011 Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol. Lett.* 14, 702–708. doi: [10.1111/j.1461-0248.2011.01628.x](https://doi.org/10.1111/j.1461-0248.2011.01628.x)
- Vilà M, Hulme PE 2017 *Impact of Biological Invasions on Ecosystem Services*. Springer
- Vilà M, Pino J, Font X 2007 Regional assessment of plant inva-



- sions across different habitat types. *J. Veg. Sci.* 18, 35–42. doi: [10.1111/j.1654-1103.2007.tb02513.x](https://doi.org/10.1111/j.1654-1103.2007.tb02513.x)
- Vilà M, Pino J, Montero A, Font X 2010 Are island plant communities more invaded than their mainland counterparts? *J. Veg. Sci.* 21, 438–446. <https://www.jstor.org/stable/40925501>
- Vilà M, Pujadas J 2001 Land-use and socio-economic correlates of plant invasions in European and North African countries. *Biol. Conserv.* 100, 397–401. doi: [10.1016/s0006-3207\(01\)00047-7](https://doi.org/10.1016/s0006-3207(01)00047-7)
- Vilà M, Tessier M, Suehs CM, Brundu G, Carta L et al. 2006 Local and regional assessments of the impacts of plant invaders on vegetation structure and soil properties of Mediterranean islands. *J. Biogeogr.* 33, 853–861. doi: [10.1111/j.1365-2699.2005.01430.x](https://doi.org/10.1111/j.1365-2699.2005.01430.x)
- Vilibić I, Čikeš Keč V, Zorica B, Šepić J, Matijević S et al. 2016 Hydrographic conditions driving sardine and anchovy populations in a land-locked sea. *Mediterr. Mar. Sci.* 17, 1–12. doi: [10.12681/mms.1120](https://doi.org/10.12681/mms.1120)
- von Schuckmann K, Le Traon P-Y, Smith N, Pascual A, Brousseau P et al. 2018 Copernicus Marine Service Ocean State Report. *J. Oper. Oceanogr.* 11, S1–S142. doi: [10.1080/1755876X.2018.1489208](https://doi.org/10.1080/1755876X.2018.1489208)
- Vousdoukas MI, Mentaschi L, Voukouvalas E, Verlaan M, Feyen L 2017 Extreme sea levels on the rise along Europe's coasts. *Earth's Futur.* 5, 304–323. doi: [10.1002/2016EF000505](https://doi.org/10.1002/2016EF000505)
- Wahl M, Buchholz B, Winde V, Golomb D, Guy-Haim T et al. 2015 A mesocosm concept for the simulation of near-natural shallow underwater climates: The Kiel Outdoor Benthocosms (KOB). *Limnol. Oceanogr. Methods* 13, 651–663. doi: [10.1002/lom3.10055](https://doi.org/10.1002/lom3.10055)
- Waldman R, Somot S, Herrmann M, Sevault F, Isachsen PE 2018 On the chaotic variability of deep convection in the Mediterranean Sea. *Geophys. Res. Lett.* 45, 2433–2443. doi: [10.1002/2017GL076319](https://doi.org/10.1002/2017GL076319)
- Walsh K, Giorgi F, Coppola E 2014 Mediterranean warm-core cyclones in a warmer world. *Clim. Dyn.* 42, 1053–1066. doi: [10.1007/s00382-013-1723-y](https://doi.org/10.1007/s00382-013-1723-y)
- Walther GR, Roques A, Hulme PE, Sykes MT, Pyšek P et al. 2009 Alien species in a warmer world: risks and opportunities. *Trends Ecol. Evol.* 24, 686–693. doi: [10.1016/j.tree.2009.06.008](https://doi.org/10.1016/j.tree.2009.06.008)
- Wang F, Polcher J 2019 Assessing the freshwater flux from the continents to the Mediterranean Sea. *Sci. Rep.* 9, 8024. doi: [10.1038/s41598-019-44293-1](https://doi.org/10.1038/s41598-019-44293-1)
- Watt MS, Kriticos DJ, Lamoureux SL, Bourdôt GW 2011 Climate change and the potential global distribution of serrated tussock (*Nassella trichotoma*). *Weed Sci.* 59, 538–545. doi: [10.1614/WS-D-11-00032.1](https://doi.org/10.1614/WS-D-11-00032.1)
- Werner C, Zumkier U, Beyschlag W, Máguas C 2010 High competitiveness of a resource demanding invasive *Acacia* under low resource supply. *Plant Ecol.* 206, 83–96. doi: [10.1007/s11258-009-9625-0](https://doi.org/10.1007/s11258-009-9625-0)
- Wernli H, Schwierz C 2006 Surface Cyclones in the ERA-40 Dataset (1958–2001). Part I: Novel Identification Method and Global Climatology. *J. Atmos. Sci.* 63, 2486–2507. doi: [10.1175/jas3766.1](https://doi.org/10.1175/jas3766.1)
- Wibig J 1999 Precipitation in Europe in relation to circulation patterns at the 500 hPa level. *Int. J. Climatol.* 19, 253–269. doi: [10.1002/\(SICI\)1097-0088\(19990315\)19:3<253::AID-JOC366>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1097-0088(19990315)19:3<253::AID-JOC366>3.0.CO;2-0)
- Wild M 2009 Global dimming and brightening: A review. *JGR Atmos.* 114, D00D16. doi: [10.1029/2008JD011470](https://doi.org/10.1029/2008JD011470)
- Wild M 2012 Enlightening Global Dimming and Brightening. *Bull. Am. Meteorol. Soc.* 93, 27–37. doi: [10.1175/BAMS-D-11-00074.1](https://doi.org/10.1175/BAMS-D-11-00074.1)
- Wild RJ, Dubé WP, Aikin KC, Eilerman SJ, Neuman JA et al. 2017 On-road measurements of vehicle NO<sub>z</sub>/NO<sub>x</sub> emission ratios in Denver, Colorado, USA. *Atmos. Environ.* 148, 182–189. doi: [10.1016/j.atmosenv.2016.10.039](https://doi.org/10.1016/j.atmosenv.2016.10.039)
- Wilks DS 2016 “The stippling shows statistically significant grid points”: How research results are routinely overstated and overinterpreted, and what to do about it. *Bull. Am. Meteorol. Soc.* 97, 2263–2273. doi: [10.1175/BAMS-D-15-00267.1](https://doi.org/10.1175/BAMS-D-15-00267.1)
- Witkowski F, Andral B, Derolez V, Tomasino C 2017 Campagne de surveillance DCE 2015 en Méditerranée française Districts « Rhône et Cotiers Méditerranéens » et « Corse ».
- WMO 2017 WMO Guidelines on the Calculation of Climate Normals - 2017 edition. Geneva, Switzerland.
- Wood S, Sebastian K, Scherr SJ 2000 Pilot analysis of global ecosystems: Agroecosystems.
- Woollings T, Blackburn M 2012 The North Atlantic Jet Stream under Climate Change and Its Relation to the NAO and EA Patterns. *J. Clim.* 25, 886–902. doi: [10.1175/jcli-d-11-00087.1](https://doi.org/10.1175/jcli-d-11-00087.1)
- Wöppelmann G, Marcos M 2012 Coastal sea level rise in Southern Europe and the nonclimate contribution of vertical land motion. *JGR Ocean.* 117. doi: [10.1029/2011JC007469](https://doi.org/10.1029/2011JC007469)
- World Economic Forum 2011 Scenarios for the Mediterranean Region. 40.
- Wu J, Zha J, Zhao D, Yang Q 2018a Changes in terrestrial near-surface wind speed and their possible causes: an overview. *Clim. Dyn.* 51, 2039–2078. doi: [10.1007/s00382-017-3997-y](https://doi.org/10.1007/s00382-017-3997-y)
- Wu N, Wang C, Ausseil AG, Alhafedh Y, Broadhurst L et al. 2018b Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for Asia and the Pacific*, eds. Karki M, Senaratna Sellamuttu S, Okayasu S, Suzuki W (Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), 265–370.
- Wurl O, Obbard JP 2004 A review of pollutants in the sea-surface microlayer (SML): a unique habitat for marine organisms. *Mar. Pollut. Bull.* 48, 1016–1030. doi: [10.1016/j.marpolbul.2004.03.016](https://doi.org/10.1016/j.marpolbul.2004.03.016)
- Xoplaki E, González-Rouco JF, Gyalistras D, Luterbacher J, Rickli R et al. 2003a Interannual summer air temperature variability over Greece and its connection to the large-scale atmospheric circulation and Mediterranean SSTs 1950–1999. *Clim. Dyn.* 20, 537–554. doi: [10.1007/s00382-002-0291-3](https://doi.org/10.1007/s00382-002-0291-3)
- Xoplaki E, Gonzalez-Rouco JF, Luterbacher J, Wanner H 2004 Wet season Mediterranean precipitation variability: influence of large-scale dynamics and trends. *Clim. Dyn.* 23,

- 63–78. doi: [10.1007/s00382-004-0422-0](https://doi.org/10.1007/s00382-004-0422-0)
- Xoplaki E, González-Rouco JF, Luterbacher J, Wanner H 2003b Mediterranean summer air temperature variability and its connection to the large-scale atmospheric circulation and SSTs. *Clim. Dyn.* 20, 723–739. doi: [10.1007/s00382-003-0304-x](https://doi.org/10.1007/s00382-003-0304-x)
- Yeruham E, Rilov G, Shpigel M, Abelson A 2015 Collapse of the echinoid *Paracentrotus lividus* populations in the Eastern Mediterranean - result of climate change? *Sci. Rep.* 5, 13479. doi: [10.1038/srep13479](https://doi.org/10.1038/srep13479)
- Yin JH 2005 A consistent poleward shift of the storm tracks in simulations of 21<sup>st</sup> century climate. *Geophys. Res. Lett.* 32, {L18}701. doi: [10.1029/2005GL023684](https://doi.org/10.1029/2005GL023684)
- Youssef L, Younes G, Kouzayha A, Jaber F 2015 Occurrence and levels of pesticides in South Lebanon water. *Chem. Speciat. Bioavailab.* 27, 62–70. doi: [10.1080/09542299.2015.1023092](https://doi.org/10.1080/09542299.2015.1023092)
- Yurtkuran Z, Saygı Y 2013 Assessment of Pesticide Residues in Karaboğaz Lake from Kızılırmak Delta, Turkey. *Bull. Environ. Contam. Toxicol.* 91, 165–170. doi: [10.1007/s00128-013-1037-0](https://doi.org/10.1007/s00128-013-1037-0)
- Zaghden H, Kallel M, Louati A, Elleuch B, Oudot J et al. 2005 Hydrocarbons in surface sediments from the Sfax coastal zone, (Tunisia) Mediterranean Sea. *Mar. Pollut. Bull.* 11, 1287–1294. doi: [10.1016/j.marpolbul.2005.04.045](https://doi.org/10.1016/j.marpolbul.2005.04.045)
- Zahn A, Rainho A, Rodrigues L, Palmeirim JM 2009 Low macro-arthropod abundance in exotic Eucalyptus plantations in the Mediterranean. *Appl. Ecol. Environ. Res.* 7, 297–301.
- Zalidis GC, Stamatiadis S, Takavagoglou V, Eskridge K, Misopolinos N 2002 Impacts of agricultural practices on soil and water quality in the Mediterranean region and proposed assessment methodology. *Agric. Ecosyst. Environ.* 88, 137–146.
- Zampieri M, Lionello P 2011 Anthropogenic land use causes summer cooling in Central Europe. *Clim. Res.* 46, 255–268. doi: [10.3354/cr00981](https://doi.org/10.3354/cr00981)
- Zanis P, Hadjinicolaou P, Pozzer A, Tyrlis E, Dafka S et al. 2014 Summertime free-tropospheric ozone pool over the eastern Mediterranean/Middle East. *Atmos. Chem. Phys.* 14, 115–132. doi: [10.5194/acp-14-115-2014](https://doi.org/10.5194/acp-14-115-2014)
- Zappa G, Hoskins BJ, Shepherd TG 2015 The dependence of wintertime Mediterranean precipitation on the atmospheric circulation response to climate change. *Environ. Res. Lett.* 10, 104012.
- Zappa G, Shaffrey LC, Hodges KI, Sansom PG, Stephenson DB et al. 2013 A Multimodel Assessment of Future Projections of North Atlantic and European Extratropical Cyclones in the CMIP5 Climate Models. *J. Clim.* 26, 5846–5862. doi: [10.1175/JCLI-D-12-00573.1](https://doi.org/10.1175/JCLI-D-12-00573.1)
- Zemp M, Paul F, Hoelzle M, Haeberli W 2008 Glacier fluctuations in the European Alps, 1850–2000, in *Darkening Peaks: Glacier Retreat, Science, and Society*, eds. Orlove B, Wiegandt E, Luckman BH (Berkeley, Los Angeles, London: University of California Press), 152–167.
- Zenetos A 2019 Mediterranean Sea: 30 Years of Biological Invasions (1988–2017). in *Proceedings of the 1st Mediterranean Symposium on the Non-Indigenous Species* (Tunis, Antalya: SPA/RAC Publications), 13–19.
- Zenetos A, Çinar ME, Crocetta F, Golani D, Rosso A et al. 2017 Uncertainties and validation of alien species catalogues: The Mediterranean as an example. *Estuar. Coast. Shelf Sci.* 191, 171–187. doi: [10.1016/j.ecss.2017.03.031](https://doi.org/10.1016/j.ecss.2017.03.031)
- Zenetos A, Mačić V, Jaklin A, Lipej L, Poursanidis D et al. 2016 Adriatic ‘opisthobranchs’ (Gastropoda, Heterobranchia): shedding light on biodiversity issues. *Mar. Ecol.* 37, 1239–1255. doi: [10.1111/maec.12306](https://doi.org/10.1111/maec.12306)
- Zerefos CS, Eleftheratos K, Meleti C, Kazadzis S, Romanou A et al. 2009 Solar dimming and brightening over Thessaloniki, Greece, and Beijing, China. *Tellus B Chem. Phys. Meteorol.* 61, 657–665. doi: [10.1111/j.1600-0889.2009.00425.x](https://doi.org/10.1111/j.1600-0889.2009.00425.x)
- Zhou JL, Maskouki K 2003 Distribution of polycyclic aromatic hydrocarbons in water and surface sediments from Daya Bay, China. *Environ. Pollut.* 121, 269–281. doi: [10.1016/S0269-7491\(02\)00215-4](https://doi.org/10.1016/S0269-7491(02)00215-4)
- Zittis G 2018 Observed rainfall trends and precipitation uncertainty in the vicinity of the Mediterranean, Middle East and North Africa. *Theor. Appl. Climatol.* 134, 1207–1230. doi: [10.1007/s00704-017-2333-0](https://doi.org/10.1007/s00704-017-2333-0)
- Zittis G, Hadjinicolaou P 2017 The effect of radiation parameterization schemes on surface temperature in regional climate simulations over the MENA-CORDEX domain. *Int. J. Climatol.* 37, 3847–3862. doi: [10.1002/joc.4959](https://doi.org/10.1002/joc.4959)
- Zittis G, Hadjinicolaou P, Fnais M, Lelieveld J 2016 Projected changes in heat wave characteristics in the eastern Mediterranean and the Middle East. *Reg. Environ. Chang.* 16, 1863–1876. doi: [10.1007/s10113-014-0753-2](https://doi.org/10.1007/s10113-014-0753-2)
- Zittis G, Hadjinicolaou P, Klangidou M, Proestos Y, Lelieveld J 2019 A multi-model, multi-scenario, and multi-domain analysis of regional climate projections for the Mediterranean. *Reg. Environ. Chang.* 19, 2621–2635. doi: [10.1007/s10113-019-01565-w](https://doi.org/10.1007/s10113-019-01565-w)
- Zittis G, Hadjinicolaou P, Lelieveld J 2014 Role of soil moisture in the amplification of climate warming in the eastern Mediterranean and the Middle East. *Clim. Res.* 59, 27–37. doi: [10.3354/cr01205](https://doi.org/10.3354/cr01205)
- Zubler EM, Folini D, Lohmann U, Lüthi D, Schär C et al. 2011 Simulation of dimming and brightening in Europe from 1958 to 2001 using a regional climate model. *JGR Atmos.* 116, D18205. doi: [10.1029/2010JD015396](https://doi.org/10.1029/2010JD015396)
- Zurek MB, Henrichs T 2007 Linking scenarios across geographical scales in international environmental assessments. *Technol. Forecast. Soc. Change* 74, 1282–1295. doi: [10.1016/j.techfore.2006.11.005](https://doi.org/10.1016/j.techfore.2006.11.005)

## Information about authors

### Coordinating Lead Authors

Semia Cherif:

*Tunis El Manar University, Higher Institute for Applied Biological Sciences of Tunis (UTM-ISSBAT), Tunis, Tunisia*

Enrique Doblas-Miranda:

*CREAF, Bellaterra (Barcelona), Spain & Universitat Autònoma de Barcelona, Bellaterra (Barcelona), Spain*

Piero Lionello:

*University of Salento, Lecce, Italy*

### Lead Authors

Carlos Borrego:

*Department of Environment and Planning (DAO), Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal*

Filippo Giorgi:

*Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste, Italy*

Ana Iglesias:

*Universidad Politécnica de Madrid (UPM), Madrid, Spain*

Sihem Jebari:

*Tunisian National Research Institute for Rural Engineering, Water and Forestry (INRGREF), Tunis, Tunisia*

Ezzeddine Mahmoudi:

*University of Carthage, Bizerte, Tunisia*

Marco Moriondo:

*Institute of BioEconomy, Italian National Research Council (CNR-IBE), Florence, Italy*

Olivier Pringault:

*Mediterranean Institute of Oceanography (MIO), Institute of Research for Development (IRD), Marseille, France*

Gil Rilov:

*National Institute of Oceanography, Israel Oceanographic and Limnological Research (IOLR), Haifa, Israel*

Samuel Somot:

*Centre National de Recherches Météorologiques (CNRM, Université de Toulouse, Météo-France, CNRS), Toulouse, France*

Athanossios Tsikliras:

*Aristotle University of Thessaloniki, Thessaloniki, Greece*

Monsterrat Vila:

*Spanish National Research Council, Doñana Biological Station (EBD-CSIC), Sevilla, Spain*

Georgios Zittis:

*The Cyprus Institute, Nicosia, Cyprus*

### Contributing Authors

Giovanni Argenti:

*University of Florence, Florence, Italy*

Marie-Anne Auger-Rozenberg:

*Orléans Forest Zoology research unit, French National Research Institute for Agriculture, Food and Environment (INRAE), Orléans, France*

Ernesto Azzurro:

*Institute for Marine Biological Resources and Biotech-*

*nology of the National Research Council (CNR-IRBIM), Ancona, Italy*

Corina Basnou:

*CREAF, Bellaterra (Barcelona), Spain*

Sophie Bastin:

*Laboratoire ATmospheres Milieux Observations Spatiales/Institut Pierre Simon Laplace (LATMOS/IPSL), UVSQ Université Paris-Saclay, SU, CNRS, Guyancourt, France*

Mustapha Béjaoui:

*University of Carthage, Bizerte, Tunisia*

Lorenzo Brilli:

*Institute of BioEconomy, Italian National Research Council (CNR-IBE), Florence, Italy*

Martina Carrete:

*University Pablo de Olavide, Sevilla, Spain*

Emma Cebria:

*Blanes Centre for Advanced Studies (CEAB), Spanish National Research Council (CSIC), Blanes, Girona, Spain*

Hanene Chaabane :

*Laboratory of Pest control and Integrated Protection in Agriculture / LMI NAILA, National Institute of Agronomy of Tunisia (INAT), University of Carthage, Tunis, Tunisia*

Sílvia Coelho:

*Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal*

Renato Colucci:

*Institute of Polar Sciences, Marine Science Institute, National Research Council (CNR-ISP), Trieste, Italy*

Styliani Dafka:

*Justus-Liebig University, Giessen, Germany*

Sofia Darmaraki:

*Dalhousie University, Canada*

Camilla Dibari:

*University of Florence, Florence, Italy*

Donna Dimarchopoulou:

*Aristotle University of Thessaloniki, Thessaloniki, Greece*

Jean-Claude Dutay:

*Laboratory of Climate and Environmental Sciences (LSCE), Pierre Simon Laplace Institute (IPSL), Gif-sur-Yvette, France*

Monia El Bour:

*National Institute of Marine Sciences and Technology (INSTM), Carthage - Salammbô, Tunisia*

Antonietta Elia:

*University of Santiago de Compostela, Santiago de Compostela, Spain*

Elena Georgopoulou:

*National Observatory of Athens, Athens, Greece*

Sylvaine Giakoumi:

*Zoological Station Anton Dohrn, Naples, Italy & School of Biological Sciences, University of Queensland, Australia*

Juan Jesus Gonzalez Aleman:

*Complutense University of Madrid, Madrid, Spain*

Pablo González-Moreno:

*CABI, Egham, United Kingdom*

Madeleine Goutx:

*Mediterranean Institute of Oceanography (MIO), CNRS, Marseille, France*

- Olivier Grunberger:  
*National Agricultural Institute of Tunisia (INAT), French National Research Institute for Sustainable Development (IRD), Tunis, Tunisia*
- Ivan Güttler:  
*Croatian Meteorological and Hydrological Service, Zagreb, Croatia*
- Nathalie Hilmi:  
*Monaco Scientific Centre (MSC), Monaco*
- Gabriel Jorda:  
*Spanish Institute of Oceanography, Balearic Islands Oceanographic Centre (COB-IEO), Palma, Spain*
- Stelios Katsanevakis:  
*University of the Aegean, Mitilini, Greece*
- Mehdi Lahlou:  
*National Institute of Statistics and Applied Economics (INSEA), Rabat, Morocco*
- Manfred A. Lange:  
*The Cyprus Institute, Nicosia, Cyprus*
- Luisa Leolini:  
*University of Florence, Florence, Italy*
- Myriam Lopes:  
*Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal*
- Annarita Mariotti:  
*National Oceanic and Atmospheric Administration (NOAA), Silver Spring, United States of America*
- Ana Isabel Miranda:  
*Centre for Environmental and Marine Studies (CESAM), Department of Environment and Planning, University of Aveiro, Aveiro, Portugal*
- Meryem Mojtahid:  
*University of Angers, Angers, France*
- Alexandra Monteiro:  
*Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal*
- Samuel Morin:  
*Université Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Études de la Neige, Grenoble, France*
- Pierre Nabat:  
*Centre National de Recherches Météorologiques (CNRM, Université de Toulouse, Météo-France, CNRS), Toulouse, France*
- Anika Obermann-Hellhund:  
*Goethe University, Frankfurt am Main, Germany*
- Tuğba Öztürk:  
*Işık University, Istanbul, Turkey*
- Androniki Pardalou:  
*Aristotle University of Thessaloniki, Thessaloniki, Greece*
- Sandra Rafael:  
*Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal*
- Francesca Raffaele:  
*Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste, Italy*
- Lena Reimann:  
*Christian-Albrechts University, Kiel, Germany*
- Alain Roques:  
*Orléans Forest Zoology research unit, French National Research Institute for Agriculture, Food and Environment (INRAE), Orléans, France*
- Asma Sakka Hlaili:  
*University of Carthage, Bizerte, Tunisia*
- Alberto Santini:  
*Institute for Sustainable Plant Protection, Italian National Research Council (CNR), Sesto Fiorentino, Italy*
- Giuseppe Scarcella:  
*National Research Council, Institute of Marine Biological Resources and Biotechnologies (CNR-IRBIM), Ancona, Italy*
- Katrin Schroeder:  
*Italian National Research Council, Institute of Marine Science (CNR-ISMAR), Venice, Italy*
- Isla Simpson:  
*Climate and Global Dynamics Laboratory of the National Center for Atmospheric Research in Boulder, Colorado, United States of America*
- Nicolina Staglianò:  
*University of Florence, Florence, Italy*
- Meryem Tanarhte:  
*Hassan II Mohammedia University, Casablanca, Morocco*
- Rob Tanner :  
*European and Mediterranean Plant Protection Organisation (EPPO), Paris, France*
- Rémi Thiéblemont:  
*French Geological Survey, Orléans, France*
- Yves Tramblay:  
*HydroSciences Montpellier (University Montpellier, CNRS, IRD), Montpellier, France*
- Marco Turco:  
*University of Murcia, Murcia, Spain*
- Athanasios Vafeidis:  
*Christian-Albrechts University, Kiel, Germany*
- Martin Wild:  
*Institute of Atmospheric and Climate Science, ETH Zürich, Zürich, Switzerland*
- Elena Xoplaki:  
*Department of Geography & Center for international Development and Environmental Research, Justus Liebig University, Giessen, Germany*
- Argyro Zenetos:  
*Institute of Marine Biological Resources and Inland Waters, Hellenic Centre for Marine Research (HCMR), Argiroupoli, Greece*



# 3 RESOURCES 1-WATER

**Coordinating Lead Authors:**

Marianela Fader (Germany), Carlo Giupponi (Italy)


**Lead Authors:**

Selmin Burak (Turkey), Hamouda Dakhlaoui (Tunisia), Aristeidis Koutroulis (Greece), Manfred A. Lange (Cyprus), María Carmen Llasat (Spain), David Pulido-Velazquez (Spain), Alberto Sanz-Cobeña (Spain)

**Contributing Authors:**

Manolis Grillakis (Greece), Rachid Mrabet (Morocco), David Sauri Pujol (Spain), Robert Savé (Spain), Mladen Todorovic (Italy), Yves Trambly (France), Veronika Zwirgmaier (Germany)

*This chapter should be cited as: Fader M, Giupponi C, Burak S, Dakhlaoui H, Koutroulis A, Lange MA, Llasat MC, Pulido-Velazquez D, Sanz-Cobeña A 2020 Water. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 181-236.*



# Table of contents

<b>3.1 Water</b>	<b>184</b>
Executive summary	<b>184</b>
3.1.1 Water resources in the Mediterranean Basin	<b>184</b>
3.1.1.1 Water availability	<b>184</b>
3.1.1.2 Rivers	<b>185</b>
3.1.1.3 Groundwater	<b>186</b>
3.1.1.4 Lakes and reservoirs	<b>187</b>
3.1.1.5 Country-level water availability	<b>187</b>
3.1.2 Water use per sector	<b>188</b>
3.1.2.1 Overview across economic sectors	<b>188</b>
3.1.2.2 Agriculture	<b>188</b>
3.1.2.3 Tourism	<b>190</b>
3.1.2.4 Industry and energy	<b>190</b>
3.1.2.5 Municipal water withdrawal	<b>191</b>
3.1.3 Past changes in hydrological variables	<b>192</b>
3.1.3.1 Evapotranspiration and soil moisture	<b>192</b>
3.1.3.2 Runoff and water resources	<b>194</b>
3.1.3.3 Extreme events	<b>194</b>
Floods	<b>194</b>
Droughts	<b>195</b>
3.1.3.4 Groundwater	<b>195</b>
3.1.3.5 Water quality	<b>196</b>
3.1.4 Projections, vulnerabilities and risks	<b>198</b>
3.1.4.1 Impacts of 1.5-2 °C global warming and associated socio-economic pathways on water	<b>198</b>
Evapotranspiration and soil moisture	<b>198</b>
Runoff	<b>198</b>
Extreme events	<b>199</b>
- Floods	<b>199</b>
- Droughts	<b>199</b>
Groundwater resources	<b>200</b>
Water quality	<b>200</b>
3.1.4.2 Impacts of higher end global warming on water	<b>201</b>
Soil moisture	<b>201</b>
Runoff	<b>201</b>
Extreme events	<b>204</b>
Floods	<b>204</b>
Droughts	<b>204</b>
Groundwater	<b>204</b>
Water quality	<b>204</b>
Vulnerabilities and risks in the water-food-energy nexus	<b>204</b>
3.1.5 Water management and adaptation	<b>205</b>
3.1.5.1 Integrated Water Resources Management (IWRM)	<b>205</b>
Definition, components and link to climate change adaptation	<b>205</b>
3.1.5.2 Adaptation measures	<b>206</b>
Supply-side adaptation measures	<b>207</b>
- Desalination	<b>207</b>
- Wastewater treatment and reuse	<b>208</b>
- Artificial recharge of groundwater	<b>208</b>
- Inter-basin transfers	<b>210</b>
- Dams	<b>211</b>
- Virtual water trade	<b>213</b>
Demand-side adaptation measures	<b>214</b>
- Efficient water use in households and economic sectors	<b>214</b>
- Agricultural management for water conservation	<b>216</b>
- Reduction of water losses	<b>217</b>
<b>Box 3.1.1 Impacts of structural aging and climate change on water infrastructure</b>	<b>218</b>
Dams	<b>218</b>
Pipelines	<b>218</b>
<b>Box 3.1.2 Water use and the specific Mediterranean diet</b>	<b>219</b>
<b>Supplementary information</b>	<b>220</b>
<b>References</b>	<b>221</b>
<b>Information about authors</b>	<b>236</b>

## 3.1 Water

### Executive summary

Water resources in the Mediterranean are scarce. They are limited, unevenly distributed and often mismatching human and environmental needs. Three quarters of the resource are located in the northern Mediterranean while three quarters of the needs are in the south and east. As a consequence, approx. 180 million people in the southern and eastern Mediterranean countries suffer from water scarcity (<math>1,000 \text{ m}^3 \text{ capita}^{-1} \text{ yr}^{-1}</math>). The main water user is agriculture, in particular on the southern and eastern rim. The percentage of irrigated land of the total cultivated area is 25% for the Mediterranean Basin and is currently increasing, likely with higher rates under even drier climate conditions in the future. Water demand for both tourism and agriculture peak in summer, potentially enhancing tensions and conflicts in the future. Municipal water use is particularly constrained in the south and will likely be exacerbated in the future by demographic and migration phenomena. In parallel, northern countries face additional risks in flood prone areas where population and urban settlements are rapidly increasing.

Climate change, in interaction with other drivers (mainly demographic and socio-economic developments), has mainly negative consequences for the water cycle in the Mediterranean Basin, including reduced runoff and groundwater recharge, increased crop water requirements, increased conflicts among users, and increased risk of over-exploitation and degradation. These impacts will be much more important for global warming higher than 2°C.

Strategies and policies for water management and climate change adaptation are strongly interconnected with all other sectors (e.g., the Water-Energy and Food Nexus). Technical solutions are available for improving water use efficiency and increasing reuse. Seawater desalination is increasingly used as adaptation measure to reduce (potable) water scarcity in arid and semi-arid Mediterranean countries, despite known drawbacks in terms of environmental impacts and energy requirements. Promising solar technologies are under development, potentially reducing emissions and costs. Reuse of wastewater is a solution for agriculture and industrial activities but also recharge of aquifers. Inter-basin transfers may lead to controversies and conflicts. Construction of dams

contributes to combat water and energy scarcities, but with trade-offs in terms of social and environmental impacts. Overall, water demand management, which increases water use efficiency and reduces water losses, particularly in urban environments, is crucial for a sustainable development. Maintaining Mediterranean diet or coming back to it on the basis of locally produced food and reducing food wastes may save water but also carbon emissions while having nutritional benefits.

### 3.1.1 Water resources in the Mediterranean Basin

#### 3.1.1.1 Water availability

The total renewable freshwater resources of the countries belonging to the Mediterranean Basin are estimated to between 1,212 km<sup>3</sup> yr<sup>-1</sup> and 1,452 km<sup>3</sup> yr<sup>-1</sup> (Ferragina 2010; FAO 2016a), distributed unevenly. Northern Mediterranean countries hold approx. 72 to 74% of the resources, while the eastern Mediterranean (including Turkey) and the southern Mediterranean countries (including Egypt and the Nile) share the remaining approx. 26 to 28% (Ferragina 2010; FAO 2016a). Besides the heterogeneous distribution of total freshwater resources, the partitioning of surface and groundwater differs as well. In northern Mediterranean countries, 96% of the renewable water is surface water, whereof 25% are contributing as base flow to river discharges after percolating to the aquifer. The 25% are referred to as shared surface/groundwater resource. Only 4% of the total water is recharging the groundwater (FAO 2016a). In the southern Mediterranean, the share of renewable groundwater resources is 11% of its total renewable freshwater. In eastern Mediterranean countries it even amounts to 20% (FAO 2016a). Especially in southern and eastern Mediterranean countries non-renewable "fossil" groundwater resources account for almost 66% of the total groundwater (MED-EUWI 2007; Lezzaik and Milewski 2018).

As aquifers and rivers are often situated across political borders, the dependency among countries concerning freshwater resources is common (Ganoulis 2006; Iglesias et al. 2007, 2011). In the southern and eastern Mediterranean, more than 60% of the surface water is transboundary and all Middle East and North Africa countries share at least one aquifer (World Bank 2018). Expressed as a dependency ratio, i.e., percentage of renewable



freshwater resources originating in another country, the mean dependency of the northern Mediterranean countries is 22%, the eastern 27% and the southern 18% (FAO 2016a).

The total human population of Mediterranean countries is rising and is expected to increase from 466 million people in 2010 to 529 million people in 2025 (UNEP/MAP 2016). Thus, while only covering 2.6% of the freshwater resources, 7.4% of the world's population has to be supplied with water (MED-EUWI 2007). Contrary to the total population development of the Mediterranean region, some single country projections show a decrease in population of 1% to 5% until 2025 and even 16% to 62% until 2100. Most of the countries with a negative population growth rate are in the northern Mediterranean region (Albania, Bosnia and Herzegovina, Greece, Italy, Malta, Montenegro, Macedonia, Portugal) except for Lebanon, which belongs to the eastern part (UN 2019). Comparing available freshwater resources to the population of the Mediterranean regions, the northern part has 36% of the population and 72% to 74% of the renewable freshwater, the east 24% and 19.5% to 21%

and southern Mediterranean 40% and 5% to 8.5% respectively (FAO 2016a). As a result, 180 million people in the southern and eastern Mediterranean suffer from water scarcity (<1,000 m<sup>3</sup> capita<sup>-1</sup> yr<sup>-1</sup>) and 80 million people from extreme water shortage (<500 m<sup>3</sup> capita<sup>-1</sup> yr<sup>-1</sup>) (Ferragina 2010). In the northern Mediterranean however, an average water availability of 1,700 m<sup>3</sup> capita<sup>-1</sup> yr<sup>-1</sup> is given, in some Balkan states even a supply of 10,000 m<sup>3</sup> capita<sup>-1</sup> yr<sup>-1</sup> (Milano et al. 2013).

### 3.1.1.2 Rivers

River basins draining into the Mediterranean Sea cover an area of over 5 million km<sup>2</sup> including the entire Nile river basin but not the rivers draining Portugal into the Atlantic Ocean (Ludwig et al. 2009; Lionello et al. 2012). Portugal is considered a Mediterranean country and three large-scale river basins are shared between Spain and Portugal, i.e., Duero with 96,200 km<sup>2</sup>, Tejo with 69,900 km<sup>2</sup> and Guadiana with 65,200 km<sup>2</sup> (Wolf et al. 1999). Besides a few major river basins (>80,000 km<sup>2</sup>, Fig. 3.1), most catchments are medium to small-scale (Lionello et al. 2012).




Figure 3.1 | Major river basins draining into the Mediterranean (Struglia et al. 2004).

In terms of discharge the ten largest rivers are the Rhône, Po, Drin-Buna, Nile, Neretva, Ebro, Tiber, Adige, Seyhan and Ceyhan rivers (Ludwig et al. 2009). Seven of these rivers are located in the northern Mediterranean countries, two in the eastern Mediterranean (Turkey) and one (the Nile) in the southern Mediterranean. Consequently, 71% of the mean annual discharge into the Mediterranean Sea originates from the northern part, whereas the eastern countries are contributing 12% and the southern 17% (Struglia et al. 2004). The large share of the southern countries comes mostly from the Nile, while 25% of the discharge in the northern countries is discharged by the Rhône and the Po River (Struglia et al. 2004; PERSEUS – UNEP/MAP 2015). Estimates of the total annual freshwater flux into the Mediterranean and Black Sea range from 305 to 737 km<sup>3</sup> yr<sup>-1</sup> (Struglia et al. 2004; Ludwig et al. 2009).

The seasonal distribution of discharge is highly variable, depending on the climatic and geographical features of the river basins. Due to the Mediterranean climate, precipitation is mostly available for river discharge during autumn, winter and spring. Some Mediterranean rivers have an ephemeral or intermittent character (Argyroudi et al. 2009). In the mountain ranges of the Mediterranean region, precipitation mostly falls in form of snow in winter and is stored until late spring. During snowmelt in late spring this freshwater is contributing to the river discharge (Nogués-Bravo et al. 2008; García-Ruiz et al. 2011; Lionello et al. 2012). Most mountain ranges are more humid than lowland regions in the Mediterranean and therefore a source of water throughout the year (López-Moreno et al. 2008). A number of mountain ranges are almost entirely located in the Mediterranean Basin (Pyrenees, Apennines, Dinaric Alps, the Taurus and Pinthos mountain ranges and the Atlas Mountains), but also the main Alps contribute to the discharge into the Mediterranean Sea (e.g., through the Rhone, Adige and Po) (Lutz et al. 2016).

Many Mediterranean river basins are transboundary. The Nile River crosses ten countries before entering Egypt and then the Mediterranean Sea. Only 9% of the total basin area belongs to Egypt (Wolf et al. 1999). The Jordan is another important transboundary river, subjected to great water scarcity and political tensions between riparian states (Hoff et al. 2011). The largest northern transboundary river basins flowing into the Mediterranean Sea are the Ebro Basin in Spain, the Po Basin (shared by Italy, Switzerland and France) and the Rhone Basin (shared by France and Switzerland) (Wolf et al. 1999).

### 3.1.1.3 Groundwater

Groundwater resources are the main source of the water supply in many Mediterranean countries (e.g., Libya, Palestine and Israel) (FAO 2003; Leduc et al. 2017). Of the total abstracted 60,000 km<sup>3</sup> yr<sup>-1</sup>, 54% are supplying the northern Mediterranean, 18% the eastern Mediterranean and 28% the southern Mediterranean countries (MED-EUWI 2007). Accessibility to the groundwater resources depends on several factors, for example the aquifer type. Three aquifer types are most common in the Mediterranean region: The karstic carbonated aquifer is the most common aquifer type. It is mainly recharged by surface water drainage, springs or adjacent aquifers. The levels of the groundwater and the volumes of karstic aquifers are highly diverse. Nevertheless, they are frequently used for water abstraction. Alluvial aquifers emerge in valleys or deltas of large rivers, providing a distinct layer of interaction between surface water and groundwater, often with a water table close to the surface. The two major Mediterranean alluvial aquifers are located in Italy (Po delta) and Egypt (Nile delta). The third aquifer type originates from sedimentary formations. Usually it comprises a large volume at a great depth and is not renewable. Connections to surface water fluxes are therefore not common. Considering the fact that no recharge is given, the groundwater of this aquifer type is referred to as "fossil". The spatial distribution of fossil groundwater resources is mainly concentrated in the southern Mediterranean countries (e.g., the Nubian sandstone aquifer in southern Mediterranean) (Aureli et al. 2008).

The recharging of groundwater is spatially variable. In the total Mediterranean region, 92% of the total recharge is contributing to northern, 3% to the eastern and 5% to the southern countries. In the southern countries the abstraction of renewable groundwater resources is exceeded by 24% and so an overexploitation of mainly fossil groundwater is necessary to meet the demand. In northern as well as in eastern Mediterranean countries on average 31% and 92% of their renewable groundwater respectively is abstracted (Aureli et al. 2008). The situation in those regions also differs among countries. For example, overexploitation of renewable water resources is found in Palestine and Jordan, leading to depletion of the aquifers (Saghir et al. 2000).

Potential groundwater resources in the Mediterranean are not only subjected to pressures resulting from unequal distribution and accessibility but also quality issues. Agricultural activities, leakage

from urban areas or saltwater intrusion are the main sources of groundwater pollution, which can lead the resource to become unusable (Garrido and Iglesias 2006; Ferragina 2010).

Further, aquifers are often crossing political borders making an integrated management difficult. 274 underground water fields (aquifers) are known in the Mediterranean (Ferragina 2010). One of the largest aquifers, the Nubian Sandstone Aquifer, is located in the southern Mediterranean region and is shared by four countries from which two are bordering to the Mediterranean Sea (Libya and Egypt). Approximately 37% of the water is located in Egypt and 34% in Libya, which obtains 90% of its water supply from groundwater (Margat and van der Gun 2013; Leduc et al. 2017).

### 3.1.1.4 Lakes and reservoirs

“Large dams” are all dams higher than 15 m from their lowest foundation to crest and also dams between 5 m and 15 m impounding more than

3 million m<sup>3</sup> and in the Mediterranean the countries with the highest numbers are Spain (1,064), Turkey (974), France (720) and Italy (541) (ICOLD 2019). The two biggest dams in the European Mediterranean area are the Kremasta dam in the Aspropótamos River in Greece and the Alqueva dam in the Guadiana River in Portugal, whose capacity are 4.75 and 4.15 km<sup>3</sup>, respectively. Although during the last two centuries the size and number of large storage capacity reservoirs have increased, it is now growing very slowly, due to the low availability of unused suitable places and the increase of environmental concerns (EEA 2018). The largest natural freshwater lake in Southern Europe is the Lake Skadar shared by Albania and Montenegro, which volume is 1.9 km<sup>3</sup> (Lasca et al. 1981).

### 3.1.1.5 Country-level water availability

Available and exploitable water resources of the Mediterranean region per country are listed in *Table 3.1*. It is important to differentiate availability

	Population (x1,000)	Renewable water resources (km <sup>3</sup> yr <sup>-1</sup> )	Exploitable water resources (km <sup>3</sup> yr <sup>-1</sup> )	Renewable water resources per capita (m <sup>3</sup> yr <sup>-1</sup> )	Exploitable water resources per capita (m <sup>3</sup> yr <sup>-1</sup> )
Albania	2,930	30.2	13	10,307.2	4,436.9
Algeria	41,318	11.67	7.9	282.4	191.2
Bosnia and Herzegovina	3,507	37.5	-	10,692.9	-
Croatia	4,189	105.5	-	25,185.0	-
Cyprus	1,180	0.78	0.54	661.0	457.6
Egypt	97,553	57.5	49.7	589.4	509.5
France	64,980	211	100	3,247.2	1,538.9
Greece	11,160	68.4	29	6,129.0	2,598.6
Israel	8,322	1.78	1.636	2,13.9	196.6
Italy	59,360	191.3	123	3,222.7	2,072.1
Jordan	9,702	0.937	-	96.6	-
Lebanon	6,082	4.503	2.08	740.4	342.0
Libya	6,375	0.7	0.635	109.8	99.6
Malta	430.8	0.0505	0.015	117.2	34.8
Monaco	38.7	-	-	-	-
Montenegro	629	-	-	-	-
Morocco	35,740	29	20	811.4	559.6
North Macedonia	2,083	6.4	3	3,072.5	1,440.2
Palestine	4,921	0.837	0.715	170.1	145.3
Portugal	10,330	77.4	13	7,492.7	1,258.5
Serbia	8,791	162.2	-	18,450.7	-
Slovenia	2,080	31.87	-	15,322.1	-
Spain	46,354	111.5	46.3	2,405.4	998.8
Syrian Arab Republic	18,270	16.8	20.6	919.5	1,127.5
Tunisia	11,532	4.615	3.625	400.2	314.3
Turkey	80,745	211.6	112	2,620.6	1,387.1

**Table 3.1 | Available and exploitable water resources in the Mediterranean region per country** [Data Source: FAO, 2003, 2016].

and exploitability to assess the water situation of a country. Not all water can be used due to technical or environmental limitations, like a minimum required flow or uneconomical groundwater pumping (FAO 2003). The availability as well as the potential usable water vary among countries and so does ratio of water that can be exploited to the total available water.

### 3.1.2 Water use per sector

#### 3.1.2.1 Overview across economic sectors

In the southern Mediterranean countries most water is used for agriculture (76%) whereas the industrial consumption and the public amount only to 4% and 20%, respectively, of the total abstracted water (Hamdy et al. 1995; FAO 2016a). In the eastern part, agriculture uses 79% of the abstracted water, whereas the industrial and public sector have a relatively small share with 6% and 13%, respectively (FAO 2016a). The northern Mediterranean countries have also the largest water usage in agriculture (36%) (FAO 2016a). Industrial (incl. cooling 48%) and public use (16%) are much higher than the rest of the Mediterranean regions (Hamdy et al. 1995; FAO 2016a). Looking at country values, agricultural use dominates generally water demand with some prominent exceptions, for example Slovenia and France having predominant industrial water demand (Fig. 3.2) (Burak and Margat 2016).

The part of overall water abstraction for different uses from surface water and groundwater varies between countries, from 100% of groundwater resources in Malta to approx. 20% in France (Leduc et al. 2017). In most of the Mediterranean, water demand is satisfied by freshwater withdrawal but in northern Africa the proportion of demand covered with fossil groundwater is high, as are the use of treated municipal water in Israel and desalinated water in Cyprus and Malta (Fig. 3.3, Table S3.1). Water demand in Northern Africa is thus met increasingly by non-renewable water resources, estimated at 16 km<sup>3</sup> yr<sup>-1</sup>, of which more than 60% is withdrawn from fossil resources and more than 30% is due to overexploitation of renewable groundwater (WWC 2009).

#### 3.1.2.2 Agriculture

In the Mediterranean countries, water withdrawal for the agricultural sector is about 193 km<sup>3</sup> yr<sup>-1</sup>, 64-69% of total water withdrawal (FAO 2016a; Malek and Verburg 2018). These amounts depend mainly on climate, from very low levels in some Balkan countries to more than 80% in the countries with arid and semi-arid climate. The quality of water used for irrigation is a matter of concern, as low water quality may cause water-borne diseases and crop damage which would reduce agricultural production (Etteieb et al. 2017). In some countries (e.g., Egypt), non-conventional, water is used in the agricultural sector from brackish water collected from drainage canals and municipal wastewater. Using municipal waste water, even




Figure 3.2 | Water demand per sectoral use as percentage of total water demand (Burak and Margat 2016).




Figure 3.3 | Sources of water supply as percentage of total water supply (Burak and Margat 2016).

after conventional treatment, while it is beneficial in regions suffering of water scarcity, should be applied only on selected plants and carefully monitored, because of the nutrient content, as well as bacteriological pollution, not to create sanitary issues (El Ayni et al. 2011).

The total area of the Mediterranean currently equipped for irrigation is about 27 million ha (FAO 2016a). Its percentage of agricultural land is the largest in Egypt (almost 100%), very high in Israel (76.3%), Turkey (71%), Lebanon (78.8%), Greece (70.9%), Cyprus, (65.6%) and Italy (60.7%). In a given year, only a part of the equipped area is actually irrigated (about 86% or 23.2 million ha) due to lack of water for irrigation, inadequate maintenance, operation and governance, obsolete irrigation systems, etc. Average water consumption for irrigation of the agricultural sector in the Mediterranean countries is estimated to about 8,340 m<sup>3</sup> ha<sup>-1</sup>. It goes from a few thousand m<sup>3</sup> ha<sup>-1</sup> in the Balkan area (Albania, Montenegro, Bosnia and Herzegovina, Croatia and Slovenia) to much higher values, such as in Portugal (20,800 m<sup>3</sup> ha<sup>-1</sup>), Egypt (18,000 m<sup>3</sup> ha<sup>-1</sup>), Syrian Arab Republic (12,000 m<sup>3</sup> ha<sup>-1</sup>), Lebanon (8,700 m<sup>3</sup> ha<sup>-1</sup>), Malta (7,900 m<sup>3</sup> ha<sup>-1</sup>), Jordan (7,500 m<sup>3</sup> ha<sup>-1</sup>), Greece (6,800 m<sup>3</sup> ha<sup>-1</sup>), Tunisia (6,500 m<sup>3</sup> ha<sup>-1</sup>), Turkey (6,400 m<sup>3</sup> ha<sup>-1</sup>) and Morocco (6,300 m<sup>3</sup> ha<sup>-1</sup>). These differences are, besides the specific climatic conditions, due to different crop-

ping pattern, irrigation methods, and overall efficiency of water withdrawal, storage, conveyance, distribution and application. Many Mediterranean countries widely use surface irrigation, such as Turkey (87.8% of irrigated area), Syrian Arab Republic (77.8%), Egypt (75.6%), Morocco (71.6%) etc.

There is a trend in several Mediterranean countries towards the substitution of surface irrigation with more efficient localized irrigation (Rodríguez-Díaz et al. 2011), e.g., in southern Spain and the Maghreb oases (Ibáñez et al. 2008), it is also reflected in the National Strategy for Irrigation Water Saving launched by the Moroccan government within the overall Green Morocco Development Plan. The trend towards more efficient irrigation systems may not have led to absolute water savings due to simultaneous changes towards water-demanding, more profitable crops (e.g., vegetables) and/or expansion of irrigated areas (Ward and Pulido-Velazquez 2008). Yet, the implementation of water-saving irrigation systems has led to higher water productivity in terms of tons and revenues produced per unit of water applied (Rodríguez-Díaz et al. 2011; Shah 2014). The implementation of pressurized systems has also led to higher energy requirements and, thus, higher greenhouse gas emissions. Daccache et al. (2014) state that irrigation modernization in the Mediterranean could save 8 km<sup>3</sup> of water per year, but it would

also increase CO<sub>2</sub> emissions by 2.42 Gt CO<sub>2e</sub> (+135%). The new development of solar pumps in drylands and desertic environments has created a substantial decrease of the fossil water table and increased the risk of salinization (Zammouri et al. 2007; Gonçalves et al. 2013).

Changes in irrigation systems affect key variables of the water cycle such as soil evaporation, infiltration and percolation, water storage in soils, groundwater recharge, runoff and return flow. These changes affect the availability of water resources. For example, implementing drippers instead of flooding irrigation reduces in most cases soil evaporation, surface runoff, groundwater recharge and return flows (Cooley et al. 2009), potentially causing water scarcity in downstream areas. This transformation can also generate significant environmental issues in groundwater dependent ecosystems influencing the biodiversity and functioning of aquatic and terrestrial ecosystems (Kløve et al. 2011), in coastal aquifer vulnerable to seawater intrusion (Kouzana et al. 2009; Mazi et al. 2014), and in terms of soil salinization (Clemmens et al. 2008).

In some regions, there is still conversion of natural ecosystems to croplands, to non-natural grasslands/grazing areas and, especially, a widespread conversion of all uses to urban areas. This affects hydrological variables, such as soil evaporation, plant transpiration, infiltration, percolation, water storage in soils, groundwater recharge, runoff and return flow. For example, deforestation was found to reduce spring and summer evaporation by more than 1 mm day<sup>-1</sup> and decrease precipitation in the western Mediterranean (Gaertner et al. 2001).

### 3.1.2.3 Tourism

Most tourist modalities are highly dependent on sufficient water resources and at the same time a major actor in water use that may contribute to the overexploitation of existing supplies and degradation of freshwater ecosystems (de Stefano 2004). In 2017, the Mediterranean received 289 million visitors (76% of which in the countries of the North of the basin including Turkey) reaffirming the position of this area as the largest single tourist destination in the world (UNTWO 2018). In the Mediterranean, tourist activity is at its highest in summer coinciding with peak demands by irrigated agriculture which may create tensions regarding water availability likely to be exacerbated in the future due to climate change (Toth et al. 2018).

Although usually higher than that of permanent residents, water consumption by tourism is strongly influenced by the tourist modality as well as location. Gössling et al. (2012) estimated wide variations of consumption, ranging from 84 l person<sup>-1</sup> day<sup>-1</sup> for campsites in Spain to close more than 2,000 l person<sup>-1</sup> day<sup>-1</sup> in hotels in Thailand. A correspondence between hotel category and water consumption has been found with establishments in upper categories consuming more water than establishments in lower categories (Gössling et al. 2015; Rico et al. 2020) but hotel-based tourism shows also less consumption per capita than residential tourism based on house rentals (Rico-Amoros et al. 2009). High water use is related to the presence of outdoor amenities such as lawns, swimming pools or golf courses (Gössling et al. 2015). In the Mediterranean, small insular states dedicate a significant part of their total water supply to tourism (5% in Cyprus and more than 7% in Malta) while in the large countries, tourism represents at the most 1% of total water use at the country level but sometimes 5% or more of domestic use (Gössling et al. 2012).

Overall, tourism-related water consumption appears to decrease, at least in the developed mass tourism destinations of the northern part of the basin (Rico et al. 2020), due to increasing efficiencies and also to the use of non-conventional resources such as recycled water (Gabarda-Mallorquí et al. 2017), or due to the exchange between agriculture and tourism of water flows of different qualities (Rico-Amoros et al. 2013). These options respond to increasing episodes of water stress linked to climate change in the region which may also increase coastal erosion and jeopardize beaches and natural and cultural heritage sites, especially in the southern and eastern countries (Bocci and Murciano 2018).

### 3.1.2.4 Industry and energy

Water use in the industrial sector of Mediterranean countries is estimated at 59.6 km<sup>3</sup> yr<sup>-1</sup>. Additionally, 38 km<sup>3</sup> yr<sup>-1</sup> are used for the cooling of thermal power plants (Burak and Margat 2016). The two figures combined would represent around 30% of water use in the Mediterranean Basin. Most of this consumption occurs in the large developed countries of the North (France, Italy and Spain) which concentrate 80% of water used in the industrial sector and 87.5% of water used for cooling purposes (France alone concentrates more than 60% of water used for cooling). In the East and South, Turkey represents 7% of industrial

water use and Egypt 2% (Förster 2014; Burak and Margat 2016).

In most countries, chemical and especially petrochemical facilities are the main industrial users of water. More than 200 petrochemical plants and basic chemical plants are located along the Mediterranean coast and in adjacent river basins, including at least 40 major oil refineries with important concentrations in Spain, France, Italy, and some Northern African countries such as Algeria and Egypt (IDAEA 2015). Mining and manufacturing of basic metals is the main water-using industry in Serbia and Turkey while water use for food processing is present in most countries although in small quantities (Förster 2014).

The abstraction of water for industrial activities decreased in most of the developed North of the basin during the first decade of the 21st century and overall demand for water from the energy and industrial sectors is projected to decline in the following decades in these areas mainly as a result of improving efficiencies. Treated wastewater is increasingly used in some industrial sectors. For example, the petrochemical complex of Tarragona, Spain, the largest in the Mediterranean, will cover around 80% of the  $27 \times 10^6 \text{ m}^3$  of water used annually with reclaimed water in 2020 (Molist et al. 2011). In contrast, demand in the industrial sector of the South and the East is projected to increase significantly and could account for over 7% of the total water demand by 2025 (Verdier and Viollet 2015).

Hydropower constitutes a large part of the 228 GW (38%) of the installed capacity for electricity production in the Mediterranean. Around 80% of this capacity is located in France, Italy and Spain, although possibilities for further development in these countries are severely limited. *Table 3.2* indicates the percentage of electricity produced from hydropower in several countries. Of these, the only country where electricity production from hydropower is expanding is Turkey, especially in the Tigris and Euphrates river basins. Hydropower is very sensible to climate change (*Section 3.3.3.5*). For example, the average flow of the Rhône River (supplying 25% of hydroelectricity in France) could be reduced to a third of its current flow by 2100 (European Water Movement 2018).

### 3.1.2.5 Municipal water withdrawal

Municipal water withdrawal refers primarily to the direct use of water by the population, including renewable and non-renewable sources, treated, desalinated and drainage water. It is usually computed as the

Country	Electricity generated from hydropower (%)
France	12.5
Italy	25
Spain	20
Turkey	31.2
Egypt	9

**Table 3.2 | Percentage of electricity generated from hydropower in selected Mediterranean countries** (Bocchiola and Rosso 2014; France Hydroelectricité 2018; OME 2018; Estado de los embalses, pantanos y presas de España 2019; TSKB Ekonomik Araştırmalar 2020).

total water withdrawn by the public distribution network. *Table 3.3* shows the municipal water withdrawal for Mediterranean countries. A fraction of 30%, on average, of total water withdrawal is consumed for municipal use in the Mediterranean. In absolute terms, Egypt and Italy have the largest municipal water withdrawal, while when computed as percentage of total withdrawal, Bosnia and Monaco have the largest values. However, values per capita may give a more accurate picture of the situation, since population numbers differ largely from country to country (*Table 3.3*, right column). However, national statistics about drinking and sanitation water use may differ from these numbers. According to the last national study on the supply of drinking water and sanitation in Spain, for 2017 (AEAS-AGA 2018), the total water use in Spanish households (drinking, washing, cooking, toilet, shower, cleaning, etc.) was, on average,  $132 \text{ l capita}^{-1} \text{ day}^{-1}$  [i.e.,  $48 \text{ m}^3 \text{ yr}^{-1} \text{ capita}^{-1}$ , differing from the  $105.5 \text{ m}^3 \text{ yr}^{-1} \text{ capita}^{-1}$  shown in *Table 3.3*].

Domestic water consumption in the Mediterranean depends on regional socioeconomic and socio-demographic circumstances with large differences from place to place. *Fig. 3.4* shows the temporal evolution in municipal water withdrawal for selected Mediterranean countries according to FAO Aquastat database (FAO 2016a). Increasing withdrawal is reported for Cyprus, Jordan and Algeria during the last decades in the absolute values of municipal water, as well as in the fraction of municipal compared to total water withdrawal. On the other hand, the fraction of municipal water for France, Monaco and Spain shows a decreasing trend despite the relatively constant absolute values. Decreasing trends in municipal water use has also been reported in specific urban areas during the recent past, as for example in the city

	Population (x1,000)	Total water withdrawal (10 <sup>9</sup> m <sup>3</sup> yr <sup>-1</sup> )	Municipal water withdrawal (10 <sup>9</sup> m <sup>3</sup> yr <sup>-1</sup> )	Municipal water withdrawal as % of total withdrawal (%)	Municipal water withdrawal per capita (m <sup>3</sup> yr <sup>-1</sup> )
Albania	2,930	1.311	0.283	21.6	96.6
Algeria	41,318	10.46	3.600	34.4	87.1
Bosnia and Herzegovina	3,507	-	0.361	-	102.9
Croatia	4,189	0.715	0.455	63.6	108.6
Cyprus	1,180	0.311	0.110	35.4	93.2
Egypt	97,553	77.5	10.750	13.9	110.2
France	64,980	26.44	5.175	19.6	79.6
Greece	11,160	11.24	1.991	17.7	178.4
Israel	8,322	2.304	0.983	42.7	118.1
Italy	59,360	34.19	9.488	27.8	159.8
Jordan	9,702	1.044	0.457	43.8	47.1
Lebanon	6,082	1.84	0.240	13.0	39.5
Libya	6,375	5.83	0.700	12.0	109.8
Malta	430.8	0.0638	0.037	58.6	86.8
Monaco	38.7	0.005	0.005	100.0	129.2
Montenegro	629	0.1609	0.096	59.9	153.3
Morocco	35,740	10.43	1.063	10.2	29.7
North Macedonia	2,083	0.5235	0.278	53.0	133.2
Palestine	4,921	0.3752	0.181	48.3	36.8
Portugal	10,330	9.151	0.914	10.0	88.5
Serbia	8,791	5.377	0.660	12.3	75.0
Slovenia	2,080	0.9314	0.170	18.2	81.5
Spain	46,354	31.22	4.890	15.7	105.5
Syrian Arab Republic	18,270	16.76	1.475	8.8	80.7
Tunisia	11,532	4.875	0.137	2.8	11.9
Turkey	80,745	58.79	5.839	9.9	72.3

**Table 3.3 | Municipal water withdrawal in absolute values, in percentage of total withdrawal and per capita.**  
The values shown are the most recent values present in FAO AQUASTAT Database from the period 2003-2017.

of Alicante from a combination of water saving as a response to water pricing and increased environmental awareness, as well as water reuse (Morote et al. 2016).

Municipal water distribution systems of many Mediterranean countries are old and as a result, losses and leaks are estimated of the order of 35% of the total water demand (UNEP/MAP and Plan Bleu 2020). Several Mediterranean countries have set specific targets for improving water use efficiency in the context of the Mediterranean Strategy for Sustainable Development.

### 3.1.3 Past changes in hydrological variables


In order to isolate the impacts of climatic conditions on groundwater and surface water resources it is necessary to remove anthropogenic detractions from the results of monitoring. In many cases, this is performed by applying hydrological models (Escriva-Bou et al. 2017; Trichakis et al. 2017) that

simulate first the link between the climatic driving forces and the hydrological variables in natural conditions for calibration, and then compare simulations with observations.

#### 3.1.3.1 Evapotranspiration and soil moisture

The increasing Mediterranean temperatures translate directly into higher evaporative demand. However, observations denote a recent (1998 onwards) decline of land evapotranspiration in a global context that could be attributed to limitations in moisture supply (Jung et al. 2010). Evapotranspiration is controlled by water demand and supply limitation conditions, which are highly variable depending on the region and the season (Wang et al. 2010). Weather variables affecting evapotranspiration in arid and semi-arid climates range over a large interval making difficult the evaluation of actual evapotranspiration (Rana and Katerji 2000).





**Figure 3.4 | Trends in municipal water withdrawal.** Absolute (left y-axis) withdrawal values and in percentage of total withdrawal (right y-axis) for selected Mediterranean countries according to (FAO 2016a).

In several Mediterranean regions small trend changes of  $\pm 0.1 \text{ mm yr}^{-1}$  in annual evapotranspiration have been detected from 1982 to 2008, with large regional variations. Positive multi-decadal evapotranspiration trends in Mediterranean have been found by several authors (Miralles et al. 2014; Zhang et al. 2016, 2019), as a consequence of increases in transpiration and interception components, counterbalanced by decreasing soil evaporation (Zhang et al. 2016).

Soil moisture is one of the most important water resources for agriculture, especially during the dry season, and it also affects temperature variability (D'Andrea et al. 2006). Mediterranean ecosystems respond to soil moisture shortage by directly reducing gross primary productivity (Piayda et al. 2014; Meza et al. 2018), hence

soil moisture variability affects also long-term terrestrial carbon storage (Green et al. 2019). Sparse and uneven observations make it difficult to assess the past trends in soil moisture across the Mediterranean. Assessments mostly rely on hydrological and water accounting models driven by observed climate data. Such studies indicate a historical decrease in soil moisture in most of the Mediterranean region, particularly in southeastern Europe, southwestern Europe and southern France, as well as a substantial increase over western Turkey (Sheffield and Wood 2008; Kurnik et al. 2015). A progressive decrease in total soil moisture of the Mediterranean region during the twentieth century of about 2-3% that continues at an increased pace until today has been estimated through simulations (based on CMIP5 simulations) (Mariotti et al. 2015).

Spatially distributed soil moisture detection can be derived from remote sensed products. Feng (2016) analyzed the temporal trends of global soil moisture during 1982 to 2013 on European Space Agency's Climate Change Initiative soil moisture data. They found no significant trend in soil moisture in the coastal regions of south Mediterranean countries, except for Egypt that exhibits a marginally negative trend. Soil moisture in southwest Turkey decreased, but in increased in Southern Italy. Similar results were obtained by Dorigo et al. (2012) from microwave surface soil moisture measurements. These findings have to be interpreted with caution, as the depth of the soil moisture that can be detected with these equipments is limited to the first few centimeters.

### 3.1.3.2 Runoff and water resources

Several studies indicate an important reduction of streamflow in basins of the Mediterranean region during recent decades (Lutz et al. 2016; Suárez-Almiñana et al. 2017). For example, in the Jucar Basin (East Spain) streamflow in natural regime has experienced a reduction close to 40% since the 1980s (Suárez-Almiñana et al. 2017). Decreasing long-term flow trends are also detectable for a large part of the Mediterranean rivers (Su et al. 2018). The strong significant runoff decrease in the Mediterranean has also been identified by Gudmundsson et al. (2017) and is likely attributed to anthropogenic forces.

Overall, Mediterranean catchments are prone to drier climate and declining water resources apart from the alpine catchments in the north of the Mediterranean region, as for example in the Adige Region (Lutz et al. 2016). This reduction affects surface and groundwater resources.

### 3.1.3.3 Extreme events

#### Floods


Floods are the most frequent and among the costliest and deadliest natural disasters in the Mediterranean area (Swiss Re 2015; UNISDR 2015), where flooding has produced more than 85 billion euros of damages since 1900 (a 42.5% of total damages related to various disasters, EM-DAT)<sup>9</sup>. Floods and droughts present significant and increasing risks for water stress (OECD 2016) and can significantly erode poor people's assets and further undermine their livelihoods in terms

of labor productivity, housing, infrastructure, and social networks (Olsson et al. 2014).

In recent decades, a mixed signal of increasing and decreasing trends in flood occurrence has been reported from many local studies over the European Mediterranean (Hall et al. 2014). In Spain and southern France, generally decreasing trends in annual maximum floods have been found (Renard et al. 2008; Stahl et al. 2010; Giuntoli et al. 2012; Mediero et al. 2014). Blöschl et al. (2019) show a common negative change in mean annual flood discharge (between -5% and -24%) in the northern and eastern Mediterranean Basin for the period 1960-2010. Although flood trend attribution is uncertain, it is often possible identify the likely key drivers of this negative trend (Merz et al. 2012). As there is not a common negative trend in precipitation, neither in maximum precipitation in those regions, causes may be related to other changes in rainfall-runoff processes at the catchment scale, such as changes in water tables caused by either overexploitation or recharge of aquifers, or land use changes, such as deforestation or forestation, urbanization, wildfires and agricultural use changes (Mediero et al. 2014). As an example, the abandonment of agricultural activities in Catalonia (northeastern Spain) has led to an increase of the forest density in the region from 30% to 70% in less than 100 years (Boada and Gómez 2011). Finally, structural flood protection measures like flood-control reservoirs have led to a decrease in flood probability although in some cases have reduced preparedness. This is known as the "levee effect" (di Baldassarre et al. 2018).

For the Po River (Italy) there is no clear trend in annual maximum floods (Montanari 2012). In Greece, around Athens, an increase in flood frequency has been observed in recent decades (Diakakis 2014). For the largest rivers in Mediterranean basins, Blöschl et al. (2017) indicate later winter floods and Mangini et al. (2018) noted a tendency towards increasing flood magnitude and decreasing flood frequency. Studies of historical flood series for more than 500 years show the great dependence of floods on land use changes and increased exposure in Mediterranean flood prone areas since the 18th century (Barriendos et al. 2003; Wilhelm et al. 2012). Flood-rich periods associated to climate anomalies (e.g., the Little Ice Age), can often be explained by natural climate variability (Glaser et al. 2010; Barrera-Escoda and Llasat 2015).

<sup>9</sup> <http://www.emdat.be/>



**Figure 3.5 | Potentially renewable water resources** (positive values) and their official exploitation (negative values) for surface water (SW) and groundwater (GW) in the Mediterranean countries in 10<sup>9</sup> m<sup>3</sup> yr<sup>-1</sup>. Source: Leduc et al. (2017), based on AQUASTAT database (FAO 2016a). Country codes are: FR France, TR Turkey, IT Italy, ES Spain, HR Croatia, GR Greece, EG Egypt, SI Slovenia, AL Albania, MA Morocco, SY Syria, DZ Algeria; TN Tunisia, LB Lebanon, IL Israel, JO Jordan, PS Palestine, CY Cyprus, LY Libya, MT Malta.

Disastrous flash floods are much more frequent in some parts of the Mediterranean region than in the rest of Europe, affecting mainly the coastal areas, where population and urban settlements are rapidly increasing in flood-prone areas (Gaume et al. 2016). Flash floods and minor floods have increased since 1981 in regions of Italy, France and Spain (Llasat et al. 2013). This increase is mainly associated to non-climatic factors such as increasingly sealed surfaces in urban areas and suboptimal storm-water management systems (Gaume et al. 2016). In the eastern Iberian Peninsula, observations points to an increase in convective and heavy precipitation concentrated in fewer days, consistent with climate change expected for this part of the basin (Llasat et al. 2016) and that could explain the positive trend found in flash floods.

**Droughts**

In the Mediterranean region the frequency and intensity of drought has increased since 1950 (Seneviratne et al. 2012), but more severe droughts have also been recorded in the past (Quintana-Seguí et al. 2016). Gudmundsson and Seneviratne (2016) and Gudmundsson et al. (2017) suggest that anthropogenic climate change has substantially increased the probability of drought years in the Mediterranean region and conclude that there is medium confidence that enhanced greenhouse forcing contributed to increased

drying in the entire Mediterranean region. This is coherent with the increasing length of dry spells observed in the Mediterranean region (Turco and Llasat 2011; Turco et al. 2017; Hoegh-Guldberg et al. 2018).

**3.1.3.4 Groundwater**

In the Mediterranean area there is a wide range of hydrogeological contexts, aquifer recharge conditions and groundwater exploitation rates. With changing climate and growing scarcity of water, groundwater could act as a buffer during shortages of surface water supply, as aquifers have a high storage capacity and respond with a certain time lag to climatic changes. However, increases in population, rise of living standards, development of irrigated agriculture, and new activities, especially tourism, have drastically increased groundwater depletion in many countries of the regions, and the overall very high rates of withdrawal of groundwater (FAO 2015; GEF 2015). The growing exploitation, favored by many technical innovations, is the most important driver of the change in Mediterranean groundwater resources that have been reduced significantly during the last 50 years, mainly to satisfy agricultural demand, tourism and coastal cities (Leduc et al. 2017). Declining freshwater availability due to groundwater overexploitation over the southern Mediterranean Basin is detectable from large-scale satellite gravity data (GRACE) (Gonçalvès et al. 2013; Rodell et al. 2018).

An additional factor affecting trends is that the intensification of groundwater use for irrigation has occurred without governmental control during decades, affecting both the quantity and quality of the resources (Llamas et al. 2015). In the EU countries the implementation of the Water Framework Directive (WFD 2000) and the specific Directive for protection of groundwater (Directive 2006/118/EC) have helped to strengthen the governmental control for sustainable management of water resources, including groundwater and dependent ecosystem issues (Garrido et al. 2006; de Stefano et al. 2014). Nevertheless, there are frequent cases of extreme overexploitation in which to recover a sustainable use will be difficult (Leduc et al. 2017). In southeastern Spain drawdowns up to several hundred meters have been observed (Custodio et al. 2016). Particularly severe examples are known from Libya, where a very significant drop in water levels due to pumping volumes exceeds renewal resources by nearly one order of magnitude (Wada et al. 2012). Fig. 3.5 shows the available information about renewable water resources (positive values) and their exploitation in the Mediterranean countries. Despite significant uncertainties (Leduc et al. 2017), there are huge differences between big northern countries and

southern countries (Libya and Tunisia). At a country scale, the figure only shows overexploitation for Libya and Tunisia, mainly since the late 1970s (Gonçalvès et al. 2020), although there are aquifers with significant overexploitation problems also in other Mediterranean countries (e.g., Spain).

### 3.1.3.5 Water quality

Inland water pollution and seawater pollution have different characteristics (Section 2.3.1). Continental water discharging in the sea carries the most dangerous pollutants such as heavy metals, Polychlorinated Biphenyls (PCBs), aromatic hydrocarbons etc. (EPA 2001). Here, the reasons and sources of land-based pollution will be assessed, bearing in mind that 75-80% of the sea water pollution in the Mediterranean Basin is land-based generated (EPA 2001; Civili 2010) (Fig. 2.21).

There are two types of pollution sources, point source and diffuse source. Point sources can be enumerated as untreated municipal wastewater and industrial discharges whereas diffuse sources are generated by irrigated agriculture, with river discharges carrying both point and diffuse sources.

	Urban effluent	Urban solid waste	Industrial effluent	Oil effluent	Chemical toxic product	Coastal eutrophication	Coastal urbanization
Albania	+	+	-	-	+	+/-	+/-
Algeria	+	+	+	+	-	+/-	+
Bosnia and Herzegovina	+	+	-	-	+/-	-	+
Croatia	+	+	-	+ (expected)	-	+	+
Cyprus	+/-	-	+	-	-	-	+/-
Egypt	+	+	+	+/-	-	+	+
Spain	+	-	+	-	-	+/-	+
France	+	-	+	-	-	+/-	+/-
Greece	+	+	+	-	-	+/-	+/-
Israel	+	-	+	+/-	-	+/-	+/-
Italy	+	-	+	+	-	+	+
Lebanon	+	+	+/-	-	-	-	+
Libya	+	+	+	+/-	-	-	-
Malta	+	+/-	+/-	+/-	-	-	+
Morocco	+	+	+	+	+/-	+/-	+
Gaza	+	+	+	-	-	+/-	+
Monaco	-	-	-	-	-	-	+
Slovenia	+	-	+	-	-	+/-	+
Syrian Arab Republic	+	+	+	+	-	+/-	+/-

**Table 3.4 | Major environmental problems for water quality along the coastal zone of Mediterranean countries** (+: Important problem; +/-: Medium problem; -: Minor problem). Source: EEA (2006).



Table 3.4 summarizes existing major environmental problems for water quality along the coastal zone of Mediterranean countries and shows the spatial heterogeneity of water quality problems. It can be observed that urban effluents are an important problem for all Mediterranean countries with the exception of Monaco.

Inland waters such as lakes and rivers have high importance for drinking water supply. Therefore, monitoring of inland water quality is done with more stringent standards with respect to some parameters that are a constraint for human consumption (e.g., pesticides) (WFD 2006). The European Commission has launched in 2005 an initiative in order to control the most important polluting sources in the Mediterranean (i.e., industrial discharges, urban solid wastes, and urban wastewater), while reinforcing the capacity of non-EU neighboring countries with regard to pollution abatement actions (MSFD 2008/2008/56/EC).

Transboundary pollution is a severe concern with regard to persistent organic pollutants (POPs) as their transmission can be long distances away from their sources since these are not biodegradable in water but in fatty acids of living organisms and can, thus, enter the food chain (Section 2.3.3.4). In the Mediterranean region, PCBs have been used throughout urban and industrial areas (Pozo et al. 2016). In Italy, for example, PCBs were widely used as insulating fluids in electrical equipment and for many other uses (Breivik et al. 2007). In Europe, lindane usage has been estimated at 287,160 tons between 1950 and 2000 representing 63% of the total world consumption (Vijgen et al. 2011).

In the Mediterranean Sea inputs through rivers and wadis can be relevant during flash flood events, which may represent a significant portion of the yearly input of organic pollutants (Velasco et al. 2006). The total input of polyaromatic hydrocarbons (PAHs), organochlorinated phenyls (OCPs) and PCBs during two flash flood events in the coastal lagoon Mar Menor (Spain) was estimated at 0.98 kg, 1.32 kg and 0.34 kg respectively (León et al. 2017). Emerging POPs contamination has also been studied in the Albufera Natural Park (Spain), a recognized Ramsar site after 1989, where the largest contribution is via the Turia and Júcar Rivers, and also from some major irrigation channels. Emerging POPs, such as Perfluoroalkyl Substances (PFASs) and Organophosphate Flame Retardants were found in multiple environmental compartments of the Albufera wetland introduced mainly from point sources like wastewater treatment plants (WWTPs) and diffuse sources

conveyed by the two rivers and irrigation channels (Lorenzo et al. 2019).

Tourism activities lead to water pollution problems as the infrastructure facilities have to comply with an increase of polluting load by 5-fold in many cases, during the summer season (Burak et al. 2004). In several coastal settlements of eastern and southern Mediterranean countries, this issue is a big challenge for the municipal management in the sense that sudden increase in population must receive the corresponding services in good quality in order to sustain touristic activities, which constitute the major income in such cities.

Eutrophication is the result of the enrichment of water bodies with nutrients such as nitrogen and phosphorus compounds which exist mainly in domestic wastewater and industrial wastewater generated by e.g., fertilizer industries and non-point sources generated from agricultural irrigation waters that carry fertilizers rich in nitrogen and phosphorus compounds (see Section 2.3.4). The problem emerges when overfeeding of aquatic ecosystems depletes the dissolved oxygen in water during their decomposition (decay) phase. When water becomes eutrophic, change in the initial (baseline) conditions of water quality is perceived to be detrimental and harmful for the ecosystem. Eutrophication causes the degradation of the water quality, which results in negative impacts on living and non-living environment of the receiving water body. This becomes increasingly a threat for receiving water especially in semi-enclosed bays and estuaries, coastal lagoons and deltas having high productivity. Coastal eutrophication is of medium or important significance in 13 Mediterranean countries (Table 3.4, Sections 2.3.3.1 and 4.2.2.1).

Bacteriological contamination of bathing water in particular is a threat to human health, therefore sea-outfalls have to be designed and operated in order to ensure that there is no adverse impact of pathogen microorganisms on human health. Monitoring of bathing water and the EU Directive EU 76/160/EEC on this subject has been a significant achievement for Mediterranean countries (EEA 2017), either member-state or non-member state since the quality of bathing water is a prerequisite for sustainable tourism, a major income source for all the coastal cities of the Mediterranean region.

Spreading of marine mucilage, which is an aggregate of mucus-like organic matter found in the Mediterranean Sea, is linked to climate-driven sea surface warming. The presence of

mucilage makes the seawater unsuitable for bathing because of its smell and its adherence on the skins of the bathers. The mucilage can act as a controlling factor of microbial diversity across oceanic regions and could have the potential to act as a carrier of specific microorganisms, thus increasing the spreading of pathogen bacteria (Danovaro et al. 2009).

### 3.1.4 Projections, vulnerabilities and risks

#### 3.1.4.1 Impacts of 1.5-2 °C global warming and associated socio-economic pathways on water

##### Evapotranspiration and soil moisture

Evapotranspiration is an important part of the water balance at the catchment scale, especially for the Mediterranean region where around 90% of the annual rainfall can be lost through evapotranspiration (Wilcox et al. 2003) (Section 2.2.5.3). In the Maghreb region, Tramblay et al. (2018) reported that under RCP4.5, potential evapotranspiration (PET) is projected to increase (+6% to +11%) for 2036–2065 period and (+7% to +14%) for 2066–2095 period compared to historical period (1976–2005), in most areas. The relative potential evapotranspiration increase is the most important during the winter and spring months. Similar projections comparing the Temperature-based PET formula and Penman Monteith equations were reported, which indicate that the main driver of change is the temperature increase. In contrast, the projected changes in actual evapotranspiration in the Maghreb region are negative from -10% up to -35%, under RCP4.5. The strongest decline is observed in spring. This change in actual evapotranspiration is correlated to the decrease in precipitation (Tramblay et al. 2018) (Section 2.2.5.3).

Overall, soil moisture is expected to decrease by the end of this century, with a significantly lower risk at 1.5°C warming as compared to higher levels (Stocker et al. 2013; Lehner et al. 2017). Under RCP2.6 and RCP 6.0 scenarios and global warming by roughly 1.3°C and 2.4°C degrees relatively to the recent past, the European Mediterranean region is expected to exhibit increase in area affected by soil moisture drought by 14.1% and 16.3%, respectively. Most of affected areas are expected to be in Greece, the southern Iberian region (Grillakis 2019) and Mediterranean area of Iberian Peninsula (Savé et al. 2012). Likewise limited to the European Mediterranean (Portugal to Greece)

and warming of 1.5°C and 2°C, an increase in soil moisture drought area by 34% and 38% is expected (Samaniego et al. 2018).

A general decline of mean soil water availability is expected at the beginning of the growing season in Sicily, due to the expected reduction of winter rainfall. Higher water stress is likely to reduce the optimal rooting depth, possibly favouring a transition toward shrubs at the expense of forests (Viola et al. 2008). Bioclimatic and evapotranspiration projections for 2070 in Malta, under a RCP 6.0 scenario, show that arable lands of the country would need at least an additional 6 m<sup>3</sup> ha<sup>-1</sup> day<sup>-1</sup> of water to make up for the expected increased water loss. The already existent scarcity of surface water supply through reservoirs and ground water is likely to limit the future potential for irrigation, which has critical implications for future crop production (Galdies and Vella 2019).

##### Runoff

Several studies show that future reduced precipitation, associated with increased evaporation will lead to a decline of runoff in the Mediterranean region (Droogers et al. 2012; Mariotti et al. 2015; Marx et al. 2018; Thober et al. 2018) (Section 2.2.5.3). The median reduction in annual runoff is projected to almost double from about 9% (likely range 4.5–15.5%) at 1.5°C to 17% (likely range 8–25%) at 2°C (Schleussner et al. 2016; Donnelly et al. 2017) and yet higher levels corresponding to stronger warming (Döll et al. 2018; Thober et al. 2018). Overall, these projections are considered robust, since all models of the multi-model ensemble agree on the same decreasing trend (Tramblay et al. 2016).

Marx et al. (2018) found that the Alpine region shows the strongest low flow increase, from 22% at 1.5°C to 30% at 2°C, because of the relatively large snowmelt contribution. For the Mediterranean region, Thober et al. (2018) project significant decreases in high flows of -11% and -13% at 1.5°C and 2°C, respectively, mainly resulting from reduced precipitation.

Several studies have shown a future potential decrease in water resources for the southern Mediterranean region (Tramblay et al. 2013b; Ruelland et al. 2015; Seif-Ennasr et al. 2016; Marchane et al. 2017; Dakhlaoui et al. 2019a, 2019b). The projected decline in surface water in the Maghreb region is significant in winter and spring (Tramblay et al. 2016). In snow-dominated catchments in the Atlas Mountains (Morocco) a stronger climate change signal points to a major

decrease in spring runoff associated with reduced snow cover (Marchane et al. 2017). This could have serious consequences since these arid regions depend to a large extent on the water resources provided by the mountain ranges (Tramblay et al. 2016).

### Extreme events

#### Floods

Flood risk, associated with extreme rainfall events, are likely to increase due to climate change, but also due to non-climatic factors such as increasingly sealed surfaces in urban areas and ill-conceived storm water management systems and major exposure and vulnerability in flood-prone regions (Alfieri et al. 2015). Floods also affect the supply of drinking water, because in circumstances of very high flows, sewage treatment plants cannot operate and, usually, pollutants are discharged into watercourses or directly to the sea. In countries such as Spain, where hydroelectric production dams are also used for flood mitigation, the forecast of heavy rains and floods obliges partially to evacuate part of the dammed water, decreasing the energy resource.

There may be important local effects beyond the effects of land use. In a study on the impacts of climate change on floods in central Italy, basins with permeable soils have been found under greater flood risk (Camici et al. 2017). In Sardinia impermeable and flat sub-basins are predicted to experience more intense flood events in future scenarios, while more permeable and steep sub-catchments will have an opposite tendency (Piras et al. 2014). The timing of floods is changing. High flows are expected to occur up to 14 days earlier per decade in the north of Italy, the south of France and eastern Greece, or later (1 day per decade) near the north-eastern Adriatic coast, eastern Spain, the south of Italy and Greece (Blöschl et al. 2017).

There are systematic differences between projections of changes in flood hazard in the south of Europe (Italy, Greece and Iberian Peninsula) in most European and global studies using large-scale hydrological models (Kundzewicz et al. 2017). Flood events with occurrence interval larger than the return period of present flood protections are projected to increase in all continents under all considered warming levels (1.5°C, 2°C and 4°C), leading to widespread increment in flood hazard (Alfieri et al. 2017). A future increase in floods corresponding to a 10-year return level in southern French basins has been projected using the ISBA land surface scheme with different downscaling

methods, but with different magnitudes depending on the basins (Quintana-Seguí et al. 2011). Other studies suggest a decrease (Donnelly et al. 2017; Thober et al. 2018). This is due to different climate model types, scenarios and downscaling approaches (Section 2.2.1.2 and Box 2.2), but also the use of large-scale hydrological models often not calibrated and validated for all basins. This type of global (or large scale) hydrological models (LISFLOOD, VIC, HYPE...) is not well adapted to small river basins (<500 km<sup>2</sup>) which is the typical catchment size found in the Mediterranean region.

In the western Mediterranean, the lower Rhône Basin and the Po catchment, the 100-year flood is projected to mainly increase in height (Dankers and Feyen 2009; Rojas et al. 2012; Dumas et al. 2013). For the upper Soca River in Slovenia, increasing high-flow magnitudes have also been projected as well (Janža 2011). For 2°C warming, river flood magnitudes are expected to increase significantly in Mediterranean Europe except for Bulgaria and southern Spain (Roudier et al. 2016). In contrast, Thober et al. (2018) has identified significant decreases of -11% (-13%) in high flows in the Mediterranean Region at 1.5°C (2°C) scenario, mainly resulting from reduced precipitation.

#### Droughts

Drought affects both the quantity and the quality of water resources. Enhanced evapotranspiration and reduced rainfall (4% decrease of precipitation per degree of global warming, Section 2.2.5.2) both reduce water availability (Baouab and Cherif 2015). In the Mediterranean, water availability could be reduced by 2-15% for 2°C warming, among the largest decreases in the world (Schleussner et al. 2016). Regional climate simulations project (medium confidence) an increase in duration and intensity of droughts in the Mediterranean, by the end of the 21st century, for different kind of drought such as streamflow droughts (Feyen and Dankers 2009; Forzieri et al. 2014; Prudhomme et al. 2014; Giuntoli et al. 2015; Quintana-Seguí et al. 2016), meteorological droughts (Koutroulis et al. 2011) or generally low water availability (Tsanis et al. 2011). Decreased low-flow was also projected by Marx et al. (2018) using a combination of three hydrological models with five climate models and three scenarios (RCP2.6, RCP6.0, RCP8.5). They found a decrease for Euro-Mediterranean areas (France, Spain, Italy, Balkans and Greece) ranging from -12% with +1.5°C warming up to -35% with 3°C warming. Liu et al. (2018) suggest that more urban populations will be exposed to severe droughts in the Mediterranean, and the

number of the affected people will escalate further the larger will be the temperature increase.

Basin-scale studies arrive to similar results. Summer low flows are reduced between -15% and -25% for the Jordan River Basin (Smiatek et al. 2014). The intermittent flow regime of the Guadiana River (south of the Iberian Peninsula) may intensify in climate change simulations, according to the JULES land surface model with an ensemble of Euro-CORDEX simulations under RCP8.5 (Papadimitriou et al. 2016). Overall, most studies conducted with hydrological models forced by climate models in different basins, found in future projections an increase of the low flow period during summer, an increased frequency of no-flow events in France (Lespinas et al. 2014), Italy (Senatore et al. 2011; Fiseha et al. 2014; Piras et al. 2014), Spain (Majone et al. 2012), Portugal (Mourato et al. 2015), Morocco and Tunisia (Tramblay et al. 2013a, 2016; Marchane et al. 2017).

Future scenarios are most extreme when both climate and human drivers are considered. For the Durance River in southern France, regulated by large reservoirs, decrease of mean annual renewable water resources has been demonstrated, with a decrease in summer low flows, associated with a greater pressure on water demand (Andrew and Sauquet 2017). For the Mediterranean Basin in southern Europe an increase in discharge intermittency is likely to be exacerbated in the future since large amounts of water are already withdrawn for irrigation purposes (Schneider et al. 2013). For the Ebro (Spain) and Herault (France) basins an integrated modelling framework considering both hydrological processes and water demand has been applied. According to the scenarios built from nine GCM under RCP8.5 it has been found that a future increase in human activities (tourism, agriculture etc.) may have more impact on water demand than climatic changes (Grouillet et al. 2015). To conclude, water demand is already large and may severely increase in the future, in particular in North Africa, and impact water resources, and subsequently low flows (Droogers et al. 2012; Milano et al. 2013).

Projected frequency and magnitude of floods and droughts at the global scale are smaller under a 1.5°C versus 2°C of warming (*medium confidence*) (Hoegh-Guldberg et al. 2018). There is medium confidence that a global warming of 2°C (1.5°C) would lead to an increase of the area at global scale with significant increases in runoff as well as an increase in flood hazard in some regions, as compared to conditions at 1.5°C global warming (present-day condition) (Hoegh-Guldberg et al. 2018). Human exposure to increased flooding is

projected to be substantially lower at 1.5°C as compared to 2°C of global warming, although projected changes create regionally differentiated risks (*medium confidence*) (Hoegh-Guldberg et al. 2018). The risks (with current adaptation) related to water deficit in the Mediterranean are high for a global warming of 2°C, but can be substantially reduced if global warming is limited to 1.5°C (Guiot and Cramer 2016; Schleussner et al. 2016; Donnelly et al. 2017).

### Groundwater resources

Aquifer recharge is also likely to be affected by climate change. In the semi-arid zone of the Mediterranean, several regions show important reductions in future potential recharge for most of the considered projections. For example, decreases of net aquifer recharge by 12% on average over continental Spain in the horizon 2011-2045 under the highest emission scenario (RCP8.5) are expected (Pulido-Velazquez et al. 2018a). The standard deviation of annual mean recharge is expected to increase by 8% on average in the future, and the spatial distribution of the reduction is quite heterogeneous. Approx. 6.6% of the territory would have reductions of more than 20%, 52.3% of the area would suffer reductions between 10% and 20%, and the reduction would be between 0% and 10% over 40.9% of continental Spain. For some climate models, the simulations predict total recharge increases over the historical values, even though climate change would produce a reduction in the mean rainfall and an increased mean temperature (Pulido-Velazquez et al. 2015). Overexploitation of groundwater is often the most important factor in lowering of groundwater levels as compared to climate change. In Tunisia, groundwater depletion is projected to reach -28% by 2050 (Requier-Desjardins 2010). Reductions in groundwater recharge and levels, independently of the drivers, might produce significant hydrological impacts, especially in the aquifers with higher vulnerability, as for example coastal aquifers where the salt-water intrusion could be exacerbated (Pulido-Velazquez et al. 2018a).

### Water quality

Important challenges to groundwater quality in coastal areas will probably arise from salt-water intrusion driven by enhanced extraction of coastal groundwater aquifers and sea-level rise, as well as from increasing water pollution in the southern and eastern Mediterranean (Ludwig et al. 2010). Serpa et al. (2017), evaluating the impacts of climate change on nutrient and copper exports from a



humid Portuguese Mediterranean catchment (São Lourenço), found that climate change could lead to a decline in annual total nitrogen, total phosphorus exports mostly due to a decrease in runoff and erosion induced by a reduction in rainfall, but it hardly affected copper (Cu) exports largely due to its strong immobilization in soils. The changes in water quality varied markedly according to the scenarios considered. A substantial decrease in total nitrogen, total phosphorus and Cu exports was simulated under intermediate scenario A1B (rapid economy growth with balanced energy sources). Under lower emission scenario B1, however, total phosphorus exports decreased much less while total nitrogen exports hardly changed, Cu exports also remained the same (Sections 2.3.2 and 2.3.3).

### 3.1.4.2 Impacts of higher end global warming on water

A number of recent studies of potential hydrological impacts of climate change are focusing on the ambitious warming targets of the Paris Agreement (UNFCCC 2016). Given the current trends in greenhouse gas emissions, the remaining challenges for mitigation and the risk of crossing planetary stabilization thresholds (Steffen et al. 2018), the target of limiting global warming to 1.5°C (and 2°C) is becoming increasingly more difficult to achieve (Mitchell et al. 2018). Higher levels of global warming are associated with significantly increased risks and vulnerabilities in the Mediterranean freshwater resources. The present section deals with the impacts of higher end global warming levels on water variables. Since the majority of climate-change impacts assessments have tended to be framed in terms of future time horizons, rather policy relevant warming level studies (Betts et al. 2018), the context of this assessment is also framed with impact studies using time horizons. These studies consider the high emission scenario RCP8.5 and examine hydrological impacts in time horizons close to the end of the 21st century, given that 80% of the CMIP5 models are crossing the 4°C global level above the pre-industrial period before 2100 under RCP8.5 [see Section 2.2.4.2].

#### Soil moisture

There is high agreement in the degree of change of the soil moisture in the Mediterranean region (Table 3.5). High warming scenario RCP8.5 projections for the end of the century (2070–2099) show an overall soil moisture drying pattern, more pronounced in western Mediterranean and mainly in the Iberian part but also in the Aegean and


Region	Soil moisture change (%)
Iberian Peninsula	-6 / -14
Italy	0 / -6
Aegean	-4 / -10
Major emerging changes	-4 / -8
Egypt	9

**Table 3.5 | Projected changes (%) of soil moisture for the different Mediterranean regions** as indicated in Berg et al. (2017), Ruosteenoja et al. (2018).

Eastern Mediterranean regions. The already dry regions of Tunisia, Libya and Egypt are projected to be less impacted (Berg et al. 2017). A scenario for 2°C warming relative to the pre-industrial for Mediterranean Europe shows an increase in drought areas with 38.4% of surface area affected and 3.7 months of drought conditions per year (Samaniego et al. 2018).

#### Runoff

There is a high level of agreement for decreased discharge of the order of -10% to -50% over the Mediterranean region during the 21st century (Jiménez Cisneros et al. 2014; Schewe et al. 2014). Such reductions in mean discharge have also been found by Koirala et al. (2014) who applied a high-resolution routing scheme on the runoff output from 11 CMIP5 GCMs. In the same study, a significant decrease in high flows (Q5, i.e., flows exceeding 5% of time within a year) and more exaggerated in low flows (Q95, exceeded 95% of time within a year) is foreseen under high-end climate change. Assessments of higher resolution hydrological projections have been made by Betts et al. (2018), Koutroulis et al. (2018) and Papadimitriou et al. (2016) in order to assess hydrological changes at different levels of global warming (1.5°C, 2°C and 4°C relative to pre-industrial), under high-end climate change (RCP8.5). A set of high-resolution AGCM projections has been used to drive a land surface model (Papadimitriou et al. 2017) and simulate regional transient hydrologic responses (Wyser et al. 2016). Fig. 3.6 shows regional patterns of changes in multi-model mean simulated annual runoff production at different levels of global warming, relative to the 1981–2010 mean runoff states, Table 3.5 contains the corresponding



**Figure 3.6 | Regional patterns of changes in multi-model mean simulated annual runoff** (in millimetres of rain equivalent – left panel, and relative values [%] – right panel) at different warming levels (1.5°C, 2°C and 4°C relative to pre-industrial) relative to the 1981-2010 mean runoff states. The corresponding degree of agreement towards drier conditions in a set of high-resolution climate projections is shown in the middle panel (Papadimitriou et al. 2016; Wyser et al. 2016; Betts et al. 2018; Koutroulis et al. 2018).

spatially averaged values over the Mediterranean SREX domain.


With global warming level of 4°C above pre-industrial conditions, these high-resolution projections project precipitation to be reduced by a median of 10.4% relative to 1981-2010 (-6.0% to +21.1% between ensemble members). With these precipitation changes, and combined with rising temperatures and thus higher evaporation demand, runoff is expected to be 7.4% less

(-4.4% to -21.1% between ensemble members) (Table 3.6). There are large local uncertainties in hydrological impacts, but in most locations, hydrological response indicates drier conditions at 4°C, with an increasing level of agreement between ensemble models.

The severe drying is particularly apparent over the southern Mediterranean, southern and western Iberian Peninsula and France, Italy and south Greece and the Levant, with relative changes

	PRECIPITATION CHANGE						RUNOFF CHANGE					
	[mm]			[%]			[mm]			[%]		
	Median	Max	Min	Median	Max	Min	Median	Max	Min	Median	Max	Min
<b>1.5°C</b>	-2.9	26.4	-37.80	-0.6%	5.5%	-7.8%	1.6	13.4	-11.1	1.3%	10.4%	-8.6%
<b>2°C</b>	-20.1	14.3	-44.69	-4.2%	3.0%	-9.3%	-4.6	11.5	-12.5	-3.6%	8.9%	-9.7%
<b>4°C</b>	-50.1	-28.9	-101.56	-10.4%	-6.0%	-21.1%	-9.8	5.6	-27.3	-7.6%	4.4%	-21.1%

**Table 3.6 | Simulated changes in spatially averaged multi-model mean annual runoff** (in millimeters of rain equivalent and relative values) at different warming levels (1.5°C, 2°C and 4°C relative to pre-industrial) relative to the 1981-2010 mean precipitation and runoff states. Percent changes are calculated based on the spatially averaged values over the Mediterranean SREX domain (Papadimitriou et al. 2016; Wyser et al. 2016; Betts et al. 2018; Koutroulis et al. 2018).



**Figure 3.7 | Same as Fig. 3.6 for the relative changes in multi-model mean simulated annual runoff at 4°C above pre-industrial** with the locations of selected basin scale assessments. Source: see Table 3.7.

in mean annual runoff up to -50%. Global and regional scale studies show consistent patterns toward a drier Mediterranean, even if different in magnitude. Gaps in the spatial scale of these assessments are covered by a wealth of local and watershed scale studies on the simulated impacts of climate change on runoff and streamflow. Gosling et al. (2017) compared global hydrological simulations with catchment level models for the Tagus Basin and found consistency in the median values and spread of mean runoff between the

two ensembles. For 3°C global warming, mean runoff is projected, by both ensembles, to decline by approx. 40% relative to 1980-2010. For high flows (Q5) the projected median decrease is also similar, 28% and 32% between the global and the basin scale multi-model ensembles, respectively. On the other hand, for low flows (Q95) the median decrease is considerably lower (35% vs 50%) as projected by the catchment hydrological model ensemble compared to the global hydrological model ensemble.

Ref. N° Fig. 3.7	Country	Watershed /Region	Size (km <sup>2</sup> )	Future/baseline period of reported changes	No of climate models	Relative changes			Reference
						Mean	Max	Min	
1	Israel	Lake Kinneret watershed	800	2050-2079/ 1979-2005	15	-35%	-9%	-51%	[Givati et al. 2019]
2	Greece	Crete	8,320	2047-2076/ 1990-2011	5	-27%	-37%	-3%	[Koutroulis et al. 2016]
3	Italy	Imera Meridionale river basin	1,782	2080-2100/ 1990-2010	32	-50%	-25%	-80%	[Viola et al. 2016]
4	Spain	Tagus	80,000	2071-2100/ 1971-2000	5	-60%	-50%	-75%	[Lobanova et al. 2016]
5	Morocco	Rheraya catchment (high Atlas)	225	1979-2005/ 2049-2065	5	-50%	-35%	-65%	[Marchane et al. 2017]
6	Tunisia	North Tunisia (5 catchments)	81-315	1970-2000/ 2070-2100	8	-50%	-37%	-57%	[Dakhlouli et al. 2019a, 2019b, 2020]
7	France	Rhône at Beaucaire	98,000	1970-2000/ 2070-2100	8	-17%	-30%	-5%	[Dayon et al. 2018]

**Table 3.7 | Characteristics and relative changes in runoff and discharge under high-end climate change** as reported by a number of recent basin scale assessments. The ref. no is a cross-reference with Fig. 3.7.

Basin scale assessments include local scale information and thus can provide detailed impact projections not only in terms of spatial scale but also on plausible developments of local socioeconomic and environmental conditions (i.e., land use changes and human management). A number of recent such studies, listed in *Table 3.7*, project runoff reduction across the Mediterranean, with regionally variable magnitude. The mean and the range of the relative projected changes (*Table 3.7*) are largely comparable with the regional changes simulated by the higher resolution hydrological projections (*Fig. 3.6* and *3.7*).

### Extreme events

#### Floods

Global projections of river flood risk at a 4°C warming indicate that countries representing 70% of the world population and GDP will likely face an increase in flood impact above 500% (Alfieri et al. 2017). Countries of the northern Mediterranean like Italy, France and Portugal belong to this list. For southern Mediterranean countries projections rather indicate an average decrease in impacted population and expected damage at a 4°C above preindustrial levels. In a pan-European study based on Lisflood model simulations of Euro-CORDEX projections, Alfieri et al. (2015) found a general increase in 100-year daily peak flow and in average frequency of peak flow events for the majority of the northern Mediterranean river network, but the projected changes had large uncertainties under high-end climate change. An opposite (decreasing) signal was found for southern Spain caused by an overall reduction in the components contributing to river runoff. Using both socioeconomic and heavy precipitation scenarios for 1.5°C, 2°C and 3°C, Cortès et al. (2019) have demonstrated an increase in the probability of damaging events due to flash floods in the eastern part of the Spanish Mediterranean region that can arrive to be above 60% for an increase of 3°C.

#### Droughts

Regarding the evolution of drought occurrence, progressively drier conditions may be expected, based on outputs from a variety of studies, from the catchment to the global scale (Orlowsky and Seneviratne 2013; Prudhomme et al. 2014). For the RCP8.5 emission scenario, a significant increase in frequency of droughts is projected by the end of the 21st century for the Mediterranean Basin, where droughts are projected to happen 5 to 10 times more frequently not only for a global warming of 3°C, especially in Northern Africa (Naumann et al. 2018).

### Groundwater

The assessment of changes in rainfall recharge in the more pessimistic emission scenarios shows reductions even higher than 55%. The heterogeneity described for the 1.5-2°C global warming scenarios, is expected to increase in higher end warming scenarios (Pulido-Velazquez et al. 2015).

### Water quality

Climate change may affect water quality, through changing precipitation, temperature variability, frequency and occurrence of extreme events. For example, floods may result in the contamination of water sources (receiving media) with wastewater and solid waste leachate. Droughts can also affect water quality because lower water flows reduce dilution of pollutants (e.g., organic matter, heavy metals) and increase contamination of remaining water sources (Wilk and Wittgren 2009). Floods, for example, may magnify the risk of contamination in case sewerage network is composed of combined sewers collecting also rainwater. These systems are designed generally with overflow chambers to provide the security of the sewerage network by discharging the surplus water mixed with sewage into the receiving media (e.g., river, lake, sea). Leachate generated at solid waste dumping areas may contaminate water resources with hazardous pollutants disposed in such areas.

Surface waters are threatened by various kinds of point source pollution including municipal sewage discharges, industrial wastewater loads, and nonpoint source pollution from agriculture, inducing a metallic, nutrient and organic pollution, particularly cytotoxic emerging micropollutants, in river waters that can even be used for drinking purposes at a large scale (Etteieb et al. 2016; Khaled-Khodja et al. 2018) (*Section 2.3.3*).

### Vulnerabilities and risks in the water-food-energy nexus

Global sustainability is intertwined with freshwater security. The combined dynamics of climate and socio-economic changes suggest that although there is an important potential for adaptation to reduce freshwater vulnerability, climate change exposure cannot be totally and uniformly counterbalanced. In many regions, socio-economic developments will have greater impact on water availability compared to climate-induced changes. However, under a global warming level of 4°C, freshwater vulnerability in the Mediterranean is expected to increase,

regardless of the level of adaptation potential as formulated by the different Shared Socio-Economic Pathways (SSPs) (Section 2.7).

Changes in hydrological variables affect the functioning of all economic sectors, especially the food and the energy sector (Fader et al. 2018). For example, reduced river flows lead to large (>15%) declines in hydropower potential as projected for southeastern Europe (Balkan countries) (van Vliet et al. 2015). This, combined with strong increases in water temperature, makes the use of water for cooling purposes more difficult and challenging (Section 3.3).

The agricultural sector will also be severely affected by reduced water availability and increased drought under high-end climate change (van Vliet et al. 2015). Agricultural expansion in the Mediterranean region will be limited by the generally lower levels of productivity and water resources. More frequent and prolonged droughts in combination with heat stress is estimated to be the major limiting factor in crop yields, causing increased crop stress and failure in parts of central and southern Europe, especially in the European Mediterranean (Berry et al. 2017). Policy support will be increasingly important to maintain rural agricultural employment in southern Europe as increasing water scarcity and decreasing land suitability impact production and profitability (van Vliet et al. 2015). The water scarcity pressures are not homogeneous across Mediterranean and local management at the basin level is of crucial importance, but the potential benefits depend on the appropriate multi-institutional and multi-stakeholder coordination (Iglesias et al. 2007) (Section 3.2.3).

### 3.1.5 Water management and adaptation

Risk, vulnerability and impacts of climate change and other anthropogenic interventions on water resources are not static variables depending only on the strength and characteristics of human interventions. Robust design, construction and operation of infrastructure can alleviate climate-driven hazards (e.g., appropriate location of landfill sites equipped with liner and well-operating on-site leachate drainage system can reduce possible flood-induced contamination of water resources as explained in the above paragraph). This approach can be a no-regret measure for climate change adaptation.

Regulatory frameworks for water quality management vary between and within countries, also

in degrees of efficiency. Few legal and regulatory texts directly consider the impacts on water quality (Cross and Latorre 2015). Understanding that different uses require different water qualities provides an opportunity to increase water use efficiency (WUE) by developing an integrated framework regulating water qualities 'fit for purpose', drawing from the wide range of water quality standards and guidelines currently available (UN-Water 2015). Through (water, landscape, land use, etc.) management and adaptation measures, impacts, vulnerabilities and risks may be potentially reduced. This section shortly analyses strategies for management and adaptation in the water domain, divided into two subsections: (i) Integrated Water Resources Management and (ii) adaptation measures (supply and demand-side).

#### 3.1.5.1 Integrated Water Resources Management (IWRM)

##### **Definition, components and link to climate change adaptation**

Integrated Water Resources Management (IWRM) has been defined as a "process which promotes the coordinated development and management of water, land and related resources in order to maximize economic and social welfare in an equitable manner without compromising the sustainability of vital ecosystems and the environment" (Global Water Partnership 2011). The three main principles of IWRM are economic efficiency, equity and environmental sustainability. Based on them, three pillars should be developed: developing management instruments for institutions and stakeholders, establishing an enabling environment that supports IWRM implementation, and putting in place an institutional framework needed for the implementation of policies, strategies and legislation (Hassing et al. 2009). Through management of the resource at the most adequate level, the organization of participation in management practices and policy development, and assuring that the most vulnerable groups are considered, IWRM instruments directly assist communities to cope with climate variability.

There are similarities and differences between IWRM and adaptation to climate change. The main difference between the two is the focus on current and historic issues of IWRM compared to the (long-term) future focus of adaptation. Water management systems design has been based on historical climate and hydrological data assuming stationarity of systems behavior (Ludwig et al. 2014). However, future changes in the climate

system no longer allow for such assumptions and historical data are no longer sufficient as the only source of information to plan for variability and extremes (Milly et al. 2008; Ludwig et al. 2009). Thus, climate change impacts will require new approaches to guarantee sufficient water resources and also to ensure that current investments will not become obsolete. In the Mediterranean Basin, where water distribution is uneven in time and in space and some regions suffer from structural drought, appropriate planning and management considering climate change impacts is the key issue. IWRM is increasingly viewed as comprising the best available framework for building the resilience needed to adapt to climate change. Any deficiency in pertinent decision-making process may result in severe shortfalls in the water management system, which may have adverse impacts on resource availability, including water supply.

Uncertainty management is crucial for the water sector given the marked inertia, which prevails as a result of the predominance of the long-term sectoral planning timeframes and in the lifespan of investments (de Perthuis et al. 2010). Many decisions relating to water, including adaptation, are sensitive to uncertainty and imply a particularly high risk of “maladaptation” when certain solutions are excessively structured or proven rather rigid (Plan Bleu 2011). However, the uncertainty surrounding impact and risk assessment should not be seen as hampering action. On the contrary, it should encourage the emergence of a dual approach: “no regret” actions and adaptation.

A challenge for implementing IWRM or adaptation measures is assuring the coverage of investment costs and long-term funding for functioning. Public-private partnerships (PPPs) can be an adequate approach for the implementation of some of the measures. This financial approach can be applied for example for local and targeted projects (IPEMED 2018), and for the construction and operation of “centralized” wastewater treatment plants (WWTP), that require a significant technical and financial support (e.g., the As-Samra WWTP that was built according to the Build-Operate-Transfer model over 25 years in Morocco). Although PPPs are developing, they remain marginal in medium-sized cities and almost non-existent in peri-urban and rural areas, especially in southeastern Mediterranean countries.

### 3.1.5.2 Adaptation measures

The existence of uncertainties in the evaluation of future climate change impacts (Pulido-Velazquez

et al. 2018b) should not be an excuse for delay or inaction in the analysis and implementation of adaptation measures, especially in the Mediterranean region, which has been identified as one of the most vulnerable areas (Milano et al. 2013). However, due to these uncertainties, adaptation must be flexible, and adopt a comprehensive approach, considering not only climate change, but also other potential socioeconomic and environmental changes (UN 2009). The impacts will affect the private (for example irrigation communities), and the public (e.g., environmental impact, quality, and supply reliability) context. For this reason, the market for technologies for adaptation to climate change grows rapidly, given that “the cost of repairing damages is estimated to be 6 times greater than adaptation costs” (H2020WATER-2014/2015, Part 12 - Page 23 of 76).

Different approaches are applied to define adaptation scenarios. In a “top-down” approach, adaptation scenarios are developed based on expert criteria that considering the assessment of potential physical vulnerability obtained by simulating/propagating future potential scenarios within a modeling framework. Examples of application of this procedure can be found in many Mediterranean systems. Pulido-Velazquez et al. (2011) and Escrivá-Bou et al. (2017) show that the systems are vulnerable to future climate change scenarios and suggest different adaptation strategies, for example, demand management alternatives or the introduction of complementary resources (additional pumping or water transfer), which can save important quantity of money (3-65 million € yr<sup>-1</sup> in the Jucar Basin). “Bottom-up” approaches include definition of scenarios through participatory processes assessing social vulnerability (Culley et al. 2016). In this case, seminars are designed to involve the main stakeholders in the process of defining the adaptation scenarios. There are also combinations of both approaches (Brown et al. 2012; Girard et al. 2015), integrating the advantages of both of them (Serra-Llobet et al. 2016).

Adaptation measures can also be classified in measures on the demand side and on the supply side of water resources. The first group has the aim of control water demand and use through for example efficiency management, modernization in irrigation (Sanchis-Ibor et al. 2017), and application of economic instruments (prices policies, markets and subsidies) to reduce demand. In the group of measures on the supply side, we observe measures oriented to obtain complementary resources (water reuse, desalination, water transfers, etc.), measures to improve allocation and availability of

water resources (for example building new small dams or channels), and conjunctive strategies, including Management Aquifer Recharge techniques.

### **Supply-side adaptation measures**

In this section we include a short introduction to desalination, wastewater treatment and reuse, artificial recharge of groundwater, inter-basin transfer, dams and virtual water trade.

#### **Desalination**

The conversion of seawater or saline groundwater into drinking water increasingly provides a source of potable water in almost all Mediterranean countries, particularly in the eastern basin, the Arabian Peninsula and North Africa. Of the currently almost 16,000 operational desalination plants that are found in 177 countries, about half are located in the Middle East and North Africa region (Jones et al. 2019). In the Mediterranean Basin, desalination capacity has increased over the last few decades and the production of desalinated seawater in the MENA region is projected to be thirteen times higher in 2040 than 2014, the most advanced countries being presently Algeria, Egypt, Israel, Italy and Spain (UNEP/MAP and Plan Bleu 2020). Given the anticipated increase in demand as a result of growing population pressures in most Mediterranean countries on the one hand and diminishing supply resulting from precipitation decreases due to climate change, seawater desalination as an alternative source of (drinking) water will grow in importance for the region.

Desalination technologies fall into two basic groups and involve either (Younos and Tulou 2005):

- a phase change process of the water-salt mixture through the boiling of feed water; the evolving steam is subsequently cooled and condensed, leaving salts, minerals and pollutants in a highly enriched brine solution, which is separated from the clean condensed water; or
- the employment of semi-permeable membranes to separate the solvent or solutes from the water by including pressure, electric potential, and concentration to overcome natural osmotic pressures and effectively force water through the membrane, leaving all substances other than water behind.

For each group a number of different technologies have been developed (Miller 2003; Younos and Tulou 2005; Khawaji et al. 2008). Common to all of these technologies are a number of challenges. Most

of them are relatively energy intensive, which is mainly due to the need for extensive pretreatment and post-treatment steps, implying a strong correlation between electricity prices and the price for the water produced (Semiat 2008; Elimelech and Phillip 2011). Utilizing conventional, hydrocarbon sources for electricity production results in the emission of air pollutants and greenhouse gases that further exacerbate climate change (Elimelech and Phillip 2011). The impingement and entrainment of marine organisms associated with the seawater intake of a desalination plant represents a further disadvantage (Elimelech and Phillip 2011). The discharge of high-salinity brines as well as of the chemicals used in the pretreatment and membrane-cleaning protocols into the sea adjacent to a desalination plant adversely affects near coastal marine ecosystems and represents an environmental problem that is increasingly recognized (Lattemann and Höpner 2008; Elimelech and Phillip 2011; Missimer and Maliva 2018; Jones et al. 2019).

Addressing particularly the first two challenges, there have been numerous efforts to improve existing technologies (Khawaji et al. 2008; Shannon et al. 2008; Elimelech and Phillip 2011; Subramani and Jacangelo 2015). More experiments and field monitoring are needed to assess adverse impacts of brine discharge from desalination into the ocean (Elimelech and Phillip 2011).

New solutions have been proposed, particularly with regard to the high demand for energy for desalination (Papanicolas 2010; Lange 2013, 2019; Georgiou et al. 2016). In this regard, utilizing renewable energies for desalination appear to be particularly promising. Given the environmental conditions in the Mediterranean Basin, solar energy appears to be the most suitable alternative related to other renewables (Li et al. 2013). Solar desalination can be achieved either directly by coupling a solar collector with a distilling mechanism through a one-stage cycle (García-Rodríguez 2003; Kalogirou 2004; Qiblawey and Banat 2008) or indirectly by connecting a conventional distillation plant to a solar thermal system (Eltawil et al. 2009; Li et al. 2013). It is also possible to combine electricity production with seawater desalination by utilizing concentrated solar power (CSP) (El-Nashar 2001; Trieb and Müller-Steinhagen 2008; Papanicolas et al. 2016). While these technologies offer the advantage of providing “clean” electricity and potable water from one plant by utilizing cost-free solar energy in regions where solar radiation is plentiful and water availability is scarce, there are a number of significant challenges including

the requirement to improve existing technologies, the relatively high capital cost to build such plants, the need to build CSP plants close to the sea, where prices for land are usually particularly high, adding to the aforementioned capital cost, and the risk of enhanced corrosion of the plant's technical installations through sea spray and relatively high dust loads that reduce the efficiency of the CSP mirrors.

Despite these challenges, solar technologies in general and the co-generation of electricity and potable water in integrated CSP plants, in particular, appear as a viable alternative to conventionally driven seawater desalination (Lange 2013; Georgiou et al. 2016; Papanicolas et al. 2016; Bonanos et al. 2017). Seawater desalination thus clearly represents an adaptation measure to reduce (potable) water scarcity in arid and semi-arid Mediterranean countries. Desalination capacity in the Mediterranean is increasing. While promising new (solar) technologies are being developed, they still have their drawbacks and need to prove their economic feasibility. Importantly, operators will have to deal with the environmental repercussions of desalination and significant adverse impacts on near-coastal marine ecosystems (Missimer and Maliva 2018; Jones et al. 2019).

### **Wastewater treatment and reuse**

The volume of wastewater produced in southern and eastern Mediterranean countries (SEMCs) was estimated at 8,134 km<sup>3</sup> (with the exception of Israel), which makes it a valuable source with regard to its quantity (IPEMED 2018). In order to reuse wastewater, the first requirement is to have access to sewerage network connected to wastewater treatment plants (WWTPs) and network for reuse complying with the corresponding standards (IPEMED 2018). Not all effluent qualities match with the required reuse. According to the World Bank (2019)<sup>10</sup>, 90 and 97% of Mediterranean populations (south-east and European Mediterranean, respectively) had improved access to sanitation services in 2015. However, these figures do not mean that there is available treated effluent for reuse. The situation is complex with regard to efficient interception of the sewage and treatment. Although in coastal urban areas, sewerage network is satisfactory, in general, in inland areas, less developed settlements have poor sanitation networks, with often leaking septic tanks, combined sewer system with overflow structure, illegal connection to streams etc. (EPA 2001; IPEMED 2018).

Fig. 3.8 shows the sectors in which treated wastewater can be reused (Lautze et al. 2014). Agricultural, industrial and watering activities present together approx. 70% water reuse potential. The heterogeneity of goals connected to reuse of wastewater shows that the quality of the treated water may differ for end-users accordingly. Most importantly, reclaimed water use practices are finding more users as a reliable alternative and low-cost resource in line with improved treatment technologies and standards in parallel with awareness raising campaigns with regard to cultural and social acceptance. Israel and Jordan have a leading role in SEMCs with a reuse rate of over 85% of their collected wastewater. Cyprus and Malta have high levels with 90% and 60%, respectively, of their treated wastewater re-used.

Wastewater reuse should be considered not only as a reduction of losses, but also as an improvement of water quality and a change in water fluxes inside a watershed, in accordance with the principles of circular economy. For example, grey water reuse, as partial recycling inside the buildings comprises flushing water for toilets from recirculated wastewater that has been treated. Introducing this system is recommended in newly constructed, smart buildings in Istanbul (Turkey). Some research work is being carried out in residences in order to work out the conversion of grey water into water source for flushing in some new buildings at planning level.


### **Artificial recharge of groundwater**

Groundwater, which underlies most of earth's surface, represents one of the most important sources of freshwater. It is protected from evaporation by the overlying soil cover and is naturally replenished/recharged by percolation of surface water and during precipitation events. Percolation through the soil reduces impurities and thus improves water quality (Racz et al. 2012). Being protected from evaporation, groundwater resources are also less sensitive to annual and inter-annual rainfall fluctuations than surface water (Giordano 2009). For these reasons, aquifers with high mean residence time can play a significant role as buffer values to reduce the impacts of meteorological droughts (Foster et al. 2017).

Many groundwater aquifers in the Mediterranean, particularly in the eastern basin, are overexploited by groundwater extraction that exceeds surface water extraction (FAO 2016a; Jódar-Abellán et al. 2017). Satellite observations confirm these trends

<sup>10</sup> <https://donnees.banquemondiale.org/indicateur/SH.STA.BASS.ZS>





**Figure 3.8 | Potential wastewater reuse per sector** (Lautze et al. 2014).

(Voss et al. 2013; Rodell et al. 2018). In order to mitigate groundwater depletion, artificial recharge of aquifers has been used worldwide since the 1960s and 1970s (de Giglio et al. 2018). For this purpose, surface water has been pumped underground in order to re-fill the aquifer. More recently, more efficient techniques have been developed, referred to as Managed Aquifer Recharge (MAR) (de Giglio et al. 2018). These techniques include the use of treated waste water or saline water for aquifer storage and recovery (Foster and Chilton 2004; Koussis et al. 2010; Maliva et al. 2011; Djuma et al. 2016), building underground dams (Nilsson 1988; Onder and Yilmaz 2005; Chezgi et al. 2016), using groundwater in combination with other sources to minimize its usage and implementing water saving technologies (Giordano 2009), and groundwater recharge by check dams (Hashemi et al. 2015; Stein et al. 2016).

The recharge of groundwater aquifers with treated wastewater is often seen critical because of potential water quality problems. Application of this technique is therefore often restricted by regulatory authorities and lacks public acceptance (Kazner et al. 2012). Nevertheless, it is used to counter seawater intrusion in order to maintain heavily exploited coastal aquifers (Koussis et al. 2010). For example, in Tunisia a Treated Waste Water (TWW) recharge in the aquifer in order to counter its salinization due to seawater intrusion and pollution due to agricultural activities has shown a reduc-

tion in groundwater salinity. Contamination by nitrate and bacteria remained a major problem of the aquifer (Cherif et al. 2013). In Israel a tertiary treated wastewater was used for the recharge of an aquifer during a 300 days experiment. The resulting water met irrigation standards with unrestricted use since no bacteriological contamination was found in the aquifer (Idelovitch 1978). This technique not only allows the exploitation of a high amount of non-conventional water resource that is TWW, but also enables the remediation of over-exploited aquifers by increasing the water table level, and its relative quality.

A less frequently applied technology is the use of underground or subsurface dams, which are claimed to enable management of groundwater in a more sustainable manner (Onder and Yilmaz 2005). Underground dams represent subsurface barriers across a stream and can be compared to check dams or sand-storage dams (Nilsson 1988; Onder and Yilmaz 2005). The construction of subsurface dams restricts the natural groundwater flow of the system and enables the storage of water below the surface. Such dams can contribute to meet demands during droughts or heavy irrigation periods (Nilsson 1988). This is often used in near-coastal situations, where groundwater would otherwise be discharged into the sea and lost for utilization. Similarly, subsurface dams are being employed in restricting saltwater intrusion into coastal aquifers.

Recharge check- or sand-storage dams represent barriers that are placed across a river or channel to slow the movement of water, encouraging groundwater recharge (Djuma et al. 2017a). Various materials have been used to build the barrier (Onder and Yilmaz 2005). Recharge behind the check dam depends on the build-up of sediment. More specifically, a growing layer of sediment reduces the volume of stored water that eventually recharges underground aquifers. Sediment accumulation is a result of riverbank erosion or erosion in the upstream watershed area. This is affected by land use, climate, topography and soils (Abedini et al. 2012; García Lorenzo et al. 2013; Djuma et al. 2017a, 2017b). Only few studies quantify the groundwater recharge efficiency of check dams. Djuma et al. (2017b) applied a water-balance approach for the Peristerona, an ephemeral river located on the northeastern hill slopes of the Troodos Mountains, Cyprus. They found that check dams can be valuable structures for increasing groundwater resources in semi-arid regions.

### Inter-basin transfers

The movement of water through artificial conveyance schemes between river basins is called inter-basin transfer (IBT). IBT is mainly employed in order to ease water shortages in the receiving basin and can be traced back to ancient times (Shiklomanov 1999; Gupta and van der Zaag 2008; Pittock et al. 2009; Boddu et al. 2011). Ever since dams have been built during the last half of the 1900s more than 364 large-scale inter-basin water transfer schemes have been established (Pittock et al. 2009). These IBTs transfer around 400 km<sup>3</sup> of water per year (Shiklomanov 1999) and are considered viable solutions to meeting escalating water demands in water scarce regions. Pittock et al. (2009) estimate that the total number of large-scale water transfer schemes may rise to between 760 and 1240 by 2020 and will transfer up to 800 km<sup>3</sup> of water per year (Shiklomanov 1999).

While potentially solving water supply issues in regions of water shortage, IBTs have significant social and environmental costs usually for both the river basin providing and the river basin receiving the water (Pittock et al. 2009). The large scale of most IBTs usually renders them expensive and thus economically risky. From an environmental point of view, IBTs interrupt the connectivity of river systems and therefore disrupt fish spawning and migration. Natural flow regimes are usually altered, sometimes with great ecological cost to threatened aquatic species or protected areas. IBTs often also modify river morphology and contribute to salinization. Finally, IBTs may also enable the transfer of invasive alien species between river basins. Short, medium and longer-term impacts of moving water from one community (the donor basin) and providing it to another (the recipient basin) are often overlooked in IBT development (Pittock et al. 2009). This may lead to controversies and conflict.

In the Mediterranean Basin, Spain has a long history of water transfers and one of the largest systems of IBTs. Despite general agreement among the main water decision-makers and stakeholders on projects and plans regarding water distribution and management, several factors have thrown this old system into crisis (Hernández-Mora et al. 2014). The Ebro inter-basin transfer, which was the main project of the Spanish National Hydrological Plan, was initiated because of pervasive pressures, scarcity, and degradation of southeastern basins in Spain (Albiac et al. 2006). The project caused heated political debates, and ultimately failed due to difficulties in achieving a sustainable management of water resources, which was caused by conflict-

ing interests of stakeholders and regions. Hernández-Mora et al. (2014) conclude that currently no technical, territorial, political, or social agreement exists on how to allocate water in Spain despite significant public and private investments in water supply infrastructure. These challenges, while depicted for Spain, are of a more general nature and can be seen in other countries of the Mediterranean Basin as well (Donta et al. 2008). They include increasing interregional conflicts and water allocation demands, the appearance of new water users who challenge the long-term privileges of large historic water holders, and a lack of understanding of water scarcity as a risk to be managed, not as a geophysical imbalance or a structural hydrological deficit (Hernández-Mora et al. 2014).

Water transfer projects have also played a major role in Turkey (Karakaya et al. 2014). The 25 main watersheds of Turkey have distinct characteristics regarding their water potential, their economy, culture, and demography. Since some of them do meet growing, but also conflicting water demands, inter-basin water transfer projects have been planned and implemented. IBTs in Turkey primarily supply water to watersheds that contain big cities, major industries, and significant agricultural activities (Karakaya et al. 2014). While water resources in Turkey are considered state property, their utilization is guaranteed for any user. Conflicts nevertheless arise between different donor/source and receiver/user basins as well as between various water consumer groups. Economic costs in the source basins have partly been met through financial compensations, and/or transfer of wealth associated with use of water resources from the user basins to the source basins. In order to address short- and long-term socio-economic implications of inter-basin water transfers, integrated assessments and specific studies are needed (Karakaya et al. 2014).

In Cyprus, groundwater was the main source of water supply for both drinking and irrigation until the 1970s. This resulted in almost all aquifers being significantly depleted because of overpumping. Seawater intrusion was observed in most of the coastal aquifers. Population increase, as well as rising numbers of tourist arrivals on the island exacerbated this problem. Already in the early 1960s, Cyprus engaged in a program to build dams to enable the collection of rainwater in surface reservoirs (see next subsection). While this somewhat eased the supply shortage of irrigation water to the agricultural sector, the gradient in precipitation values from relatively copious amounts in the central and western part of the island versus



**Figure 3.9 | Example for a large-scale IBT project is the Southern Conveyor Project on the island of Cyprus** (Water Development Department Cyprus 2000).

the eastern regions of Cyprus required additional measures (Nikolakis 2008). This led to the implementation of the Southern Conveyor Project, which was seen as a necessity and a basic prerequisite for the further agricultural and economic development of the island (Water Development Department Cyprus 2000). The Southern Conveyor Project is the largest water development project ever undertaken by the Government of Cyprus (Fig. 3.9). Its main objective is to collect and store surplus water flowing to the sea and convey it to areas of demand. Major components of the project, aside from the pipeline transporting the water to the east, are the Kouris Dam in south-central Cyprus, river diversions and underground tunnels. The project aims at the agricultural development of the coastal region between Limassol and Famagusta, as well as to meet the domestic water demand of Limassol, Larnaca, Famagusta, Nicosia, a number of villages and the tourist and industrial demand of the southern, eastern and central areas of the island (Water Development Department Cyprus 2009). The project area extends along the southern coast, between the Dhiarizos River in the west and the Kokkinokhoria irrigation area in the east (Fig. 3.9).

Large-scale IBTs are often seen as a technical solution to restore perceived imbalances in water distribution between neighboring basins. However, the disadvantages and pitfalls that often accompany such infrastructures cast doubts on the ultimate

usefulness of IBT (Pittock et al. 2009). While providing irrigation water to agriculture, IBT can also be considered to promote unsustainable and subsidized cropping practices. In planning, implementing and constructing IBTs, alternatives to the IBT that may mean delaying, deferring or avoiding the costs (in every sense) of an IBT are often overlooked or omitted. In addition, poor to non-existent consultation with affected stakeholders frequently characterizes IBT development. Finally, sufficient and adequate consideration to the environmental, social and cultural impacts of the IBT, in both the donor and recipient basins are often neglected.

### Dams

Freshwater flowing into the sea is “water lost” to arid and semi-arid countries. Reservoirs and dams play a crucial role in water resources management, but also in flood abatement, mitigating the adverse effects downriver from these structures (Sordo-Ward et al. 2012, 2013). Dams are thus built to store water, but also to divert rivers so that the bulk of their water can be used by various consumers. Dams also serve as major elements of hydro-power generation in several large Mediterranean rivers.

The first evidence for dam building dates back to the early and middle Bronze Age (2,500–1,600 BC). One of the oldest records of dam building are found in the ruins of the Saad el Kafara dam near Cairo

indicating that it was built in around 3,000 BC (Water Development Department Cyprus 2009). Roman dams were the first that were used to create reservoirs of fresh water to secure a permanent drinking water supply for urban settlements over the dry season, and also the first to introduce dam-construction types that are being used until today (Schnitter 1978, 1987; Hodge 2000). Throughout the following millennia, relatively little progress was made and it was not until the 19th century when engineering skills and construction materials available were capable of building the first large-scale arch dams. The era of building large dams was initiated by the construction of the Aswan Low Dam in Egypt in 1902 by the British.

The abatement capacity of a dam depends on the hydrologic load, the dam and reservoir characteristics, the existing operational rules, the volume to abate and other foreseen uses related with socioeconomic activities, which in some cases may lead to conflicts. Those conflicts are particularly important in those dams that are multipurpose, usually involving flood control and other purposes such as hydropower, ecological discharges, water supply or irrigation, which may be in conflict (Labadie 2004; Dittmann et al. 2009; Bianucci et al. 2013). To reduce conflicts, the decision-making process can be improved by applying a combined approach, including simulation of predefined rules (modeling without considering any inflow forecast) and optimization programming (i.e., from stochastically generated floods or flood forecasting), taking into account the different purposes through indices such as minimizing the expected deficit of water availability (for a certain purpose), or maximizing the reliability of satisfying downstream requirements (Bianucci et al. 2015). In all cases, to minimize the conflict between consumptive demands and flood abatement, the participation of users is crucial (Martín Carrasco et al. 2007). A final consideration refers to the use of meteorological forecasting to improve dam management in Mediterranean context. As the flood events are relatively short (fast response basin) and are usually due to heavy precipitation, short-term forecasting facilitated by the combination of mesoscale models and radar imagery is needed, while to manage droughts or water resources for irrigation, seasonal forecast gives substantial added value (Marcos et al. 2017).

Large dams are defined to be of at least 15 m in height, impounding more than 3 million m<sup>3</sup> of water (ICOLD 2019). In the Mediterranean Basin, the World Register on Dams lists 5,731 dams (ICOLD 2018), both single- and multi-purpose dams. In

both, the single- and the multi-purpose dams, irrigation stands out as the most frequent purpose for dam building (50% and 24%, respectively), followed by hydropower (Fig. 3.10).

Many North African countries, particularly Algeria, Egypt, Libya and Morocco, but also other Mediterranean countries including Cyprus, rely on dams and reservoirs to provide irrigation water (AQUASTAT Programme 2007; Water Development Department Cyprus 2009). While the importance and benefits of dams for the provision of water and hydroelectric power for many of the Mediterranean countries is obvious, there are a number of adverse impacts that need to be considered (Scudder 2006; Tortajada et al. 2012).




Figure 3.10 | Number of large dams in Mediterranean countries (ICOLD 2018).

The reservoirs created by a dam affect many ecological aspects of a river. The impacts of large dams on ecosystems, biodiversity and downstream livelihoods have been debated for many year and include the loss of forests and wildlife species and habitats, due to inundation, and the loss of aquatic biodiversity of upstream and downstream fisheries, amongst others (World Commission on Dams 2000).

On balance, the ecosystem impacts of practically all dams are considered more negative than positive, but enhancements of ecosystem values



through the creation of new wetland habitat and the fishing and recreational opportunities provided by new reservoirs have also been observed. Most efforts to counter the ecosystem impacts of large dams have had limited success. This has led to increased attention to legislation aimed to avoid or minimize ecological impacts. This includes setting aside particular river segments or basins in their natural state and the selection of alternative projects, sites or designs (World Commission on Dams 2000).

The impacts of dam building on people are also significant (Scudder 2006; Tortajada et al. 2012). In many cases, dam construction requires the state to displace individual households or entire communities in the name of the common good, leading to hardships and conflicts. In some cases, these negative effects have not been assessed nor accounted for by the relevant authorities. In addition, large dams frequently cause significant adverse effects on cultural heritage through the loss of cultural resources of local communities and the submergence and degradation of plant and animal remains, burial sites and archaeological monuments. The World Commission on Dams (2000) concludes that the poor, other vulnerable groups and future generations are likely to bear a disproportionate share of the social and environmental costs of large dam projects without gaining a commensurate share of the economic benefits. Nevertheless, if this report is cited in the context of negative impacts, Schulz and Adams (2019) conclude that neither the impacts nor the controversy over large dams have ended.

### Virtual water trade


Water (or any other resource) scarcity produced through an imbalance between demand and supply, leads to price increases or negative consequences for the sectors and stakeholders that need that resource. This, in turn, typically induces several societal adaptation mechanisms, such as increased efficiency through technological development, and the opportunity to import water-intensive products from other markets. More difficult is the import of the large water volumes required for local food production due to its weight and bulkiness. The strategy of trading the commodities that would be produced with the lacking water is called “virtual water trade” and can be considered an adaptation option. “Virtual water” (VW), as defined by Allan (1998), is the volume of water used to produce a good in the various steps of the production chain. Agricultural commodities require large amounts

of water from rainfall (green water) or from freshwater resources (rivers, lakes, reservoirs, canals, etc.) (blue water) (Fader et al. 2011). The third main component of the anthropogenic water cycle is grey water, defined as water released from those activities, generated in households or office buildings from streams without fecal contamination, i.e., all streams except for the wastewater from toilets affected by the consequent degree of pollution. The trade of any commodity, but in particular of agricultural ones, is associated with a virtual transport of the green and blue water used and the grey water generated for their production (Hoekstra and Chapagain 2008). Water footprints (WFs) are the other side of the medal, indicating how much water (virtual or real) a country needs to produce the products consumed by its population.

Virtual water trade, even if widely disputed as it neglects fundamental strategic and national security issues (Fader et al. 2013), supports global food security (Merrett 2003). Not only the trade of water embedded in agricultural products has gained the attention of scholars and media, but also the phenomena of “appropriation” of resources across the globe has emerged under the label of “land grabbing”, i.e., large-scale acquisition of farmland in developing countries by international investors (privates or sovereign funds). Scarcity of water, food and biofuels may partially drive the international trade of commodities, or otherwise stimulate the direct acquisition of resources where they are with a phenomenon, which is currently in the order of tens of million hectares (Land Matrix Initiative)<sup>11</sup>. It has been argued that this phenomenon should be better named as water grabbing, since water is the resource that lacks and drives the acquisitions, more than land (Johansson et al. 2016).

Hoekstra and Mekonnen (2012) calculated and mapped the green, blue and grey water footprints. They assessed both national footprints and the international virtual water flows deriving from trade of agricultural and industrial commodities. According to their analysis, most countries of the Mediterranean Basin and the Middle East are hotspots of virtual water imports (*Fig. 3.11*). Countries such as Portugal, Spain, Italy, Greece, Israel and Turkey are among those with the highest WF of national consumption (above 2,000 m<sup>3</sup> yr<sup>-1</sup> capita<sup>-1</sup>) (not shown). Hoekstra and Mekonnen (2012) also found that cereal products have the largest contribution to the WF of the average consumer (27%), followed by meat (22%) and milk products (7%).

<sup>11</sup> <https://landmatrix.org/>



**Figure 3.11 | Net virtual water imports of countries**, after Fig. 2 of Hoekstra and Mekonnen (2012).

Antonelli et al. (2012) focus on 11 Mediterranean countries, critique earlier approaches for virtual water “flow” calculations, and propose an input-output approach to account for both direct and indirect (e.g., irrigation schemes providing water to households and livestock) consumption of blue and green water. In their calculations, consideration of indirect water consumptions increases the values calculated for national WF, with remarkable differences in results for countries like France (higher WF and higher estimated VW imports), ending up with consideration that focus should be on blue water and on the economic potential for re-allocating water from agriculture (low marginal value) to other uses (households, industry), where the marginal value is higher. They also affirm that, since “green water cannot be moved” there should be no interest in saving it. In their analyses countries like Morocco and Tunisia appear to be much less blue water intensive than Egypt, with the latter showing very high potential for blue water saving. Antonelli and Tamea (2015) calculated the average VW imports of the MENA countries as  $601 \text{ m}^3 \text{ capita}^{-1} \text{ yr}^{-1}$ . Sebri (2017) suggests that the Maghreb countries are already relying on non-conventional water sources, such as waste water reuse and desalination and that they should invest more and more on strategies focused on increasing virtual water trade and enhancing water desalination technologies with the use of renewable energy as a means for abating energy costs. The author affirms that those countries do not benefit enough from virtual water trade, but that it should be considered as an important policy instrument with great care, given the relevant


strategic issues related to dependency on foreign countries for basic population needs.

#### **Demand-side adaptation measures**

Water demand management (WDM), seen as any method that saves water, or at least saves water of higher quality, is central to the reduction of water losses. WDM incorporates, (i) improving the efficiency of water used to achieve a specific task; (ii) adjusting the nature of the task or the way it is accomplished with less water or with lower quality; (iii) minimizing the loss in water quantity or quality as it flows from source through use to disposal; (iv) shifting the timing of use from peak to off-peak periods and (v) increasing the ability of the water system to continue to supply water to the users at times when water is in short supply (Brooks 2006). This embodies technical, economic, administrative, financial and/or social measures. This section assesses some demand-side measures aiming at reducing the demand for water, such as efficient water use in households and economic sector, agricultural management for water conservation, reduction of water losses, and returning or maintaining the Mediterranean diet.

#### **Efficient water use in households and economic sectors**

Water plays an important role in the context of the UN Sustainable Development Goals (SDGs, UN 2015) and is key in several SDG targets. While water for households is dealt by Target 6.1 (Achieve safe and affordable drinking water) and 6.2 (Achieve access to sanitation and hygiene and



**Figure 3.12 | Water-use efficiency per country** (UN-Water 2018).

end open defecation), consideration of efficient use of water by different sectors means focusing on Target 6.4 (Increase water-use efficiency and ensure freshwater supplies), and in particular on Indicator 6.4.1 measuring water use efficiency (WUE) to address the economic component.

A recent report produced by UN Water (UN-Water 2018) provides a comprehensive assessment of the state of SDG 6 targets across countries. Concerning Target 6.1, the proportion of population using at least basic drinking water services in 2015 appears above 90% for all Mediterranean countries, with the exception of Morocco (75-90%), and almost the same happens for 6.2, regarding the proportion of population using at least basic sanitation services, with Morocco and Algeria being between 75 and 90%, with all the other countries being above 90%. Nevertheless, there is ample room for improvement of the current situation, by adopting both innovative technologies complemented by targeted education strategies.

On the technological side of possible solutions, Campisano et al. (2017) examine opportunities for improving household water management, with focus on the potential domestic Rain Water Harvesting (RWH) systems in multi-story buildings, demonstrating that in the case study of the old town of Lipari (Aeolian Islands) there is potential for yearly water savings between 30% and 50%. Regarding household behavior, Gul et al. (2017) point out that water consumption habits are quite similar across countries and that awareness campaigns and marketing policies could both contribute to more conscious tap water use in households.

The analysis of the situation of the productive sectors is more complicated, as it requires the quantification of the various components of WUE. As suggested by FAO (2017), the WUE is defined as the value added per unit of water withdrawn over time (showing the trend in water use efficiency over time) and is calculated in US\$ m<sup>-3</sup> of abstracted water as the sum of the three main sectors (agriculture, industry and services), weighted according to the proportion of water withdrawn by each sector over the total withdrawals. UN-Water (2018) provides a global map of WUE per country. Mediterranean countries appear grouped into three classes of WUE (Fig. 3.12). Morocco, Syria, Egypt and Albania show WUE below 10 US\$ m<sup>-3</sup>. On the contrary, France and Croatia have a WUE between 40 and 80 US\$ m<sup>-3</sup>, while all other countries are between 10 and 40 US\$ m<sup>-3</sup>.

Maximizing WUE means optimizing water allocation, in order to use the scarce resource for those uses that generate the highest value added. Wimmer et al. (2014) studied future scenarios of water allocation in Europe. Their results indicate that significant physical water shortages may result from climate and socio-economic change in many regions of Europe, particularly in the Mediterranean. Therefore, specific policies will be necessary in order to prevent conflicts among users and negative economic and social, but also environmental consequences. They also point out that cross-sectoral impacts can be limited if higher priority is assigned to the domestic or industry sectors, instead of to agriculture.

For Mediterranean tourism, Hadjikakou et al. (2013) examined five cases of holiday destinations in semi-arid eastern Mediterranean, and found that food tourists' consumption is by far the most significant contribution to the sector's water footprint, but they recommend also considering the links with energy use (*Section 3.3.4.1*). Moresi (2014) explored environmental impacts of the food industry, with focus on the carbon footprint, pointing out that agricultural production appears as the hotspot in the life cycle of food products, but also that Mediterranean-type diet may have positive effects on both the environment and health. Hence, prioritizing increases water use efficiencies in the tourism and food sector may contribute substantially to the adaptation potential of the region.

#### **Agricultural management for water conservation**

Water saving in agriculture includes a set of different actions (technical, socio-economic, environmental and institutional) that should be governed and adopted at each specific location according to the effective needs, priorities and probability of success. Therefore, the solutions differ for different regions and consider both rainfed and irrigated agricultural systems. In marginal rural areas, which are usually among those most vulnerable to climate change, the overall objective beside durable improvement of the agricultural water management is the stabilization of yield and a broader socio-economic development. In this context, there is a need to integrate the traditional knowledge of cultivation with the modern technical achievements and application of new technologies. For example, the adoption of minimum-tillage in suitable soils and crops can be accompanied with a series of synchronized activities that include residue cover during the off-season, appropriate crop rotation program, adequate sowing machines, use of varieties tolerant to abiotic stresses, proper sowing, planting time and density, optimized water, nutrient inputs, weed and plant disease control, harvesting, yield storage and economic evaluation of the products.

In the ACLIMAS project ([www.aclimas.eu](http://www.aclimas.eu)), selected crop cultivars and best management practices were implemented in five Mediterranean countries (Lebanon, Jordan, Tunisia, Algeria and Morocco) at 109 farms over a total area of about 287 ha (*Section 3.2.3.1*). Overall, yields increased by 19-33% compared to traditional cultivation and water saving and water rose by 20-50%. These results are in line with the findings obtained in small Medi-

terranean basins<sup>12</sup>, where it has been shown that the increase of water use efficiency at basin level provide more water for all involved stakeholders, and also important economic, social and environmental benefits.

Traditional water harvesting techniques are already widely applied in the Mediterranean and they include interventions in micro and macro catchments, i.e., floodwater diversion to agricultural fields and construction of storage reservoirs, tanks, ponds and cisterns (FAO 2016b). Traditional techniques are assessed and designed by the application of modern technologies (GIS, digital land cover data, elevation models and satellite images) and implemented on the ground by new technologies for land preparation (Grum et al. 2016). Water saving and increase of water productivity can be achieved manipulating the microclimate of growing conditions by the application of different types of shelters (Ilić et al. 2012; Tanny 2013) and windbreaks (Lasco et al. 2014).

In the case of irrigated agriculture, the use of modern technologies, including remote sensing, for monitoring of crop water status and optimization of irrigation scheduling can contribute to more efficient use of water, nutrients and energy (El Ayni et al. 2012; Abi Saab et al. 2019) (*Section 3.2.3.1*). Water conservation and water productivity enhancement can be achieved applying supplementary irrigation and deficit irrigation strategies as regulated deficit irrigation and partial root drying (Kang et al. 2017). Other water conservation solutions, still under investigation, include plant conditioners (Boari et al. 2015; Ćosić et al. 2015; Cantore et al. 2016; AbdAllah et al. 2018) like anti-transpirants, bio-stimulants and plant growth regulators, which regulate crop transpiration and mitigate the effects of abiotic stresses, and soil conditioners (Guilherme et al. 2015), which aim to improve soil physical properties.

Some techniques of sustainable intensification, such as mulching, zero tillage, etc. increase the water retention capacity of soils making them more capable of coping with dry spells and increasing the water amount accessible to plants (Kassam et al. 2012) (*Section 3.2.3.1* and 6.4). Also, more efficient irrigation systems, shifts towards drought tolerant crops, adaptation of sowing dates, application of deficit irrigation schemes, land reclamation, and land management for carbon sequestration (Almagro et al. 2016; Funes et

<sup>12</sup> <http://medacc-life.eu/>



al. 2019) may reduce water needs for agriculture and increase water use efficiency in terms of m<sup>3</sup> per tons. For example, the yearly water withdrawal for irrigation in the Mediterranean region amounts to ~223 km<sup>3</sup> (Fader et al. 2015), but there is a water saving potential of 35% through implementation of efficient irrigation systems (Fader et al. 2016). This would, however, increase the energy costs of farmers substantially (Rodríguez-Díaz et al. 2011), driving among others changes towards more profitable but more water-intensive crops such as citrus (Fernández García et al. 2014), and potentially increasing carbon emissions of energy generation. Also, the water saving effect of efficient irrigation systems may be counterbalanced by expansion of irrigated areas.

Another important factor under Mediterranean conditions relates to soil management (*Section 3.2.3.2*). Water scarcity, soil disturbance and nutrient deficiencies limit net primary productivity in agriculture and consequently reduce soil organic carbon (SOC) stocks, since carbon inputs, such as litter, roots or crop residues, are limited. Soil carbon sequestration occurs if the balance between carbon inputs and outputs (through emissions from respiration and mineralization) is positive and finally leads to increased SOC stocks. Future increases in temperatures linked with a decrease in available soil water content, and the corresponding decrease in yields (Waha et al. 2017) may, hence, decrease soil carbon inputs. However, although it is widely known that warming increases microbial activity, soil moisture could act as the main driver of soil biomes in Mediterranean environments, limiting SOC losses by microbial mineralization. Also, agricultural management practices can significantly affect soil hydraulic properties and processes in space and time. These responses are coupled with the processes of infiltration, runoff, erosion, chemical movement, and crop growth (Green et al. 2003). All of them promote low soil water availability for crops. In all cases, water management (irrigation or soil water harvesting and storage) is critical to the feasibility of the agricultural sector in Mediterranean regions and the avoidance of SOC losses, since available water for crops increases biomass productivity, turnover of organic matter timing and humus formation (Funes et al. 2019). More information improved water management may be found in *Section 3.2.3.2*.

#### **Reduction of water losses**

Reduction of water losses in all sectors of water use is crucial for sustainable management and adaptation strategies by alleviating pressure on

freshwater supplies and protecting quality. Increasing water efficiency by reducing physical losses, is the basic principle in urban water use. This measure requires adequate monitoring between supplied and consumed water in order to minimize the non-revenue water (NRW) ratio. NRW is composed of water produced but not consumed i.e., not metered, not billed (wasted), and non-physical portion (consumed) not metered, not billed which is unauthorized consumption (*Table 3.8*) (Alegre et al. 2006). This indicates the presence of illegal connections to the municipal water network.

The water use efficiency index indicates how to measure progress in water savings through demand management, by reducing losses and wasteful use during its transmission and distribution. It covers total and sectoral efficiency in domestic (municipal), agricultural and industrial water use (Blinda 2012). The municipal water use efficiency index is defined as the ratio of the 'total drinking water volume billed' to the 'total volume supplied (abstracted/treated and distributed)' to customers by the municipalities.

A good information basis about the sources of non-revenue water (NRW) is important for water demand management, avoiding both physical (real) and commercial/non-physical (apparent) losses. In Turkey, where municipal water use has significant NRW, comprehensive rehabilitation encompassing both physical/technical and administrative improvement has decreased NRW considerably. NRW losses have been reduced by measures including the installation of bulk water meters at source to precisely measure the volume of water supplied to the city; water balance calculations by reading source, bulk and customer meters regularly; preventing reservoir overflows; synchronizing district water supply and district meter readings (establishing controlled supply zones); conducting regular leak detection studies; replacing outdated pipes and repairing leaking house connections; detecting, correcting and preventing illegal connections; and others (Burak and Mat 2010). Eliminating illegal connections in itself will not directly conserve water because consumers will still require the water they previously acquired illegally. However, once legally connected, the consumer will be subject to tariffs, which in turn should reduce the previously unmetered levels of consumption. A further benefit of legalizing these connections will be that they are properly made: illegal connections are often sub-standard and lead to high losses.

System input volume (corrected for known errors) (Water Produced + Water Imported)	Authorized consumption	Billed authorized consumption	Billed metered consumption (including water exported)	Revenue water
			Billed unmetered consumption	
		Unbilled authorized consumption	Unbilled metered consumption	Non-Revenue Water (NRW)
			Unbilled unmetered consumption	
	Water losses	Apparent losses	Unauthorized consumption	
			Customer metering / billing inaccuracies	
		Real losses	Leakage on transmission and/or distribution mains	
			Leakage and overflows at utility's storage tanks	
Leakage on service connections up to point of customer metering				

Table 3.8 | Water loss definitions and classifications (Alegre et al. 2006).

Water losses that could be recovered losses by improved network efficiency for drinking water and irrigation have been estimated to be 56 km<sup>3</sup> for the whole Mediterranean region covering the northern, eastern and southern rims in 2005 (Margat and Blinda 2005). This estimate is based on improve-

ment of drinking water (municipal) network efficiency raised to 85%, end-user (customer connection) efficiency to 90%, irrigation network efficiency increased to 90%, and plot efficiency increased to 80%. Although particularly the targeted irrigation efficiency seems to be ambitious, the correspond-

**BOX 3.1.1**

**Impacts of structural aging and climate change on water infrastructure**

Climate change and structural aging poses challenges for the functioning and security of water infrastructure, sometimes reducing water availability and quality. This subsection shortly summarizes this aspect with respect to dams and pipelines.

**Dams**

In the Mediterranean region, dams are important structures for the storage capacity of water for municipal and industrial use, irrigation purpose and energy production. Although they do not have environmental acceptance in recent years, these structures are also very important for water management in the Mediterranean Basin where available water quantity does not exist where and when required. Therefore, they are also key water structures for flood control and for maintaining water readily available for inter-basin transfer projects that have been widely implemented in several water-scarce regions in recent years (Gohari et al. 2013).

The security of dams in the face of climate change impacts is very important with respect their structural security due to increase in extreme conditions (heavy storms and flooding), changing runoff conditions (Alcocer-Yamanaka and Murillo-Fernandez 2016), and also any changes in storage capacity due to siltation (Burak and Margat 2016). It is estimated that actual capacities of dam reservoirs in Maghreb will decrease by 50% by

2100 due to siltation (Burak and Margat 2016). Permanent flow from upstream riparian countries (e.g., Turkey, Sudan) may not be ensured due to drought conditions (Margat 2011).

**Pipelines**

Pipelines are the closest infrastructure to the users; therefore, robust and well-operating water network pipelines are very important. It is quite common that water supply utilities face operational difficulties within their distribution network. Significant challenges are encountered for both rehabilitating and replacing aging infrastructure in response to growing population and new development patterns and/or shifting population (Grayman et al. 2009). In old systems, it is possible that asbestos cement pipelines (ACPs) exist even at present in some parts of the region. In Turkey, for instance, this material has been replaced in several municipal networks with ductile iron and/or high-density polyethylene (HDPE) pipes (e.g., Istanbul, Bursa and Adana) because they are low-standard pipes and because of their possible carcinogenic effect in the water network, even though there is no proven studies as stipulated by some researchers and by the WHO guidelines (Polissar et al. 1984; WHO 2003). Also, starting in 1990 in Istanbul and in other cities in the following years, investments for rehabilitation of existing water network in the new service area have been implemented in order to reduce physical losses (World Bank 2016). However, with regard to possible health risks generated by the use of ACP, practices vary from one country to another (Polissar et al. 1984).



## BOX 3.1.2

**Water use and the specific Mediterranean diet<sup>13</sup>**

The choice of diets influences the amount of water needed to produce and process the corresponding food (Section 3.2.1.2). Similarly, food waste is, at the same time, a waste of the water that was used to produce that food. Hence, influencing diet choices can be regarded as an adaptation option.

Countries like Spain are making significant efforts to reduce food loss and waste, reverse growing obesity trends, and promote the adoption of healthier food habits like the recommended and traditional Mediterranean diet. This is recognized as a key strategy to improve the population's health with locally grown, traditional, and seasonal products like fruits, vegetables, olive oil, and fish. Nevertheless, current Spanish consumption patterns (especially among younger generations, and urban and/or low-income citizens) appear to be shifting towards unhealthier diets. The largest share of the WF of current Spanish diet, as occurring with for example North American diet, is always linked to green water, which implies that the largest impact of dietary shifts is also linked to land use. Grey water in the US is 67% higher than in Spain. Only few products account for the largest share of the total WF of the two dietary options in both countries, being meat, fats, oil, and dairy products the food items with the largest WFs.

For the year 2014, the total WF of current consumption in Spain was equivalent to around 3,302 l per capita per day (of which 2,555 are green, and 400 blue WF). The products that account for the largest share in the total WF are once again meat, animal

fats, and dairy products. Likewise, roughly 41% of the total WF linked to household diets is foreign, i.e., imported Virtual Water, and the main countries of origin are Tunisia, Portugal, and France. The Total WF of food waste at households' level is estimated at 131 l per capita per day (of which 97 are green and 19 blue WF), equivalent to 4% of the Total WF of current consumption. In addition, regarding nutritional analysis, the nutrients wasted (because of food waste) per capita year were 40,385 kcal, almost 7.5 kg of macronutrients (proteins, fats, and carbohydrates), 483 grams of fiber and almost 160 grams of micronutrients (vitamins and minerals).

Current Spanish household diet is shifting away from the recommended Mediterranean towards alternative diet containing three times more meat, dairy and sugar products, and 1/3 fewer fruits, vegetables and cereals. The Mediterranean diet is also less caloric, as it contains lesser amounts of proteins and fats, and is richer in fiber and micronutrients. Due to the high water content embedded in animal products, a shift towards a Mediterranean diet would reduce the consumptive water use by about 753 l per capita per day (of which 34 are blue WF). In addition, the Mediterranean diet has higher water-nutritional efficiency than current consumption: more energy, fiber, and macro- and micro-nutrients are made available per liter of consumptive water used. In conclusion, a shift back to a locally produced Mediterranean diet (in which fruits, fish and vegetables account for a larger share of the food intake) and lessening food waste, would deliver large water savings (753 and 116 liters of consumptive water per capita per day, respectively) and nutritional benefits.

ing saved water volume appears to deserve almost any affordable effort. An overall water use efficiency index of 74% was adopted for 2015 to be one of the desirable goals by the Mediterranean countries as part of the Mediterranean Water Strategy. In that spirit, Turkey government has decided that municipalities and water administrations had to reduce the rate of water loss, averaging 25% by 2023<sup>14</sup>. Tariff structures should seek to cover the operation and investment costs whilst at the same time trying to strike a balance with what is considered fair and socially acceptable.

<sup>13</sup> This box is based on the PhD thesis and related research papers by Alejandro Blas (Blas et al. 2016, 2018).

<sup>14</sup> [https://sustainabledevelopment.un.org/content/documents/23862Turkey\\_VNR\\_110719.pdf](https://sustainabledevelopment.un.org/content/documents/23862Turkey_VNR_110719.pdf) - page 75

## Supplementary information

Country	Fresh surface water withdrawal	Fresh groundwater withdrawal	Total freshwater withdrawal	Direct use of treated municipal wastewater	Direct use of agricultural drainage water	Desalinated water produced
<b>(10<sup>9</sup> m<sup>3</sup> yr<sup>-1</sup>)</b>						
Albania	1.081	0.107	1.188	0.050		
Algeria	1.699	8.103	9.802			0.631
Bosnia and Herzegovina	0.107	0.294	0.401			
Croatia	0.248	0.425	0.673	0.015		
Cyprus	0.061	0.155	0.216	1.200		0.065
Egypt		6.500	64.400	0.411	11.9	0.2
France	20.930	5.506	26.440	0.104		0.0117
Greece	4.386	6.854	11.240	0.520		0.01
Israel	0.358	0.840	1.198	0.045		0.586
Italy			34.190	0.004		0.0973
Jordan	0.289	0.615	0.904	0.002		0.1363
Lebanon	0.396	0.700	1.812		0.028	0.0473
Libya	0.170	5.550	5.760	0.001		0.07
Malta	0.003	0.040	0.043			0.0202
Monaco			0.005			
Montenegro			0.161	0.070		
Morocco	8.251	2.322	10.350			0.007
North Macedonia	0.885	0.162	0.524	0.013		
Palestine	0.024	0.265	0.288	0.003		0.004
Portugal	4.352	4.794	9.146			0.0016
Serbia	4.917	0.460	5.377			
Slovenia	0.741	0.190	0.931			
Spain	24.830	6.394	31.220	0.592		0.364
Syrian Arab Republic			14.140	0.550	2.246	
Tunisia	1.151	2.066	4.768	0.042	0.01	0.055
Turkey	44.550	15.460	58.760	0.025		0.0082

Table S3.1 | Sources of water supply (FAO 2016a).

## References

- AbdAllah AM, Burkey KO, Mashaheet AM 2018 Reduction of plant water consumption through anti-transpirants foliar application in tomato plants (*Solanum lycopersicum* L.). *Sci. Hortic. (Amsterdam)* 235, 373–381.  
doi: [10.1016/j.scienta.2018.03.005](https://doi.org/10.1016/j.scienta.2018.03.005)
- Abedini M, Md Said MA, Ahmad F 2012 Effectiveness of check dam to control soil erosion in a tropical catchment (The Ulu Kinta Basin). *Catena* 97, 63–70.  
doi: [10.1016/j.catena.2012.05.003](https://doi.org/10.1016/j.catena.2012.05.003)
- Abi Saab M, Jomaa I, Skaf S, Fahed S, Todorovic M 2019 Assessment of a Smartphone Application for Real-Time Irrigation Scheduling in Mediterranean Environments. *Water* 11, 252.  
doi: [10.3390/w11020252](https://doi.org/10.3390/w11020252)
- AEAS-AGA 2018 XV Estudio Nacional de Suministro de Agua Potable y Saneamiento en España 2018.
- Albiac J, Hanemann M, Calatrava J, Uche J, Tapia J 2006 The rise and fall of the Ebro water transfer. *Nat. Resour. J.* 46, 727–757.
- Alcocer-Yamanaka VH, Murillo-Fernandez R 2016 Adaptation and Mitigation Measures for High-Risk Dams, Considering Changes in Their Climate and Basin, in, 179–204.  
doi: [10.1007/978-981-10-1914-2\\_9](https://doi.org/10.1007/978-981-10-1914-2_9)
- Alegre H, Baptista JM, Cabrera E, Cubillo F, Duarte P et al. 2006 *Performance Indicators for Water Supply Services*. London: IWA Publishing.  
doi: [10.2166/9781780406336](https://doi.org/10.2166/9781780406336)
- Alfieri L, Bisselink B, Dottori F, Naumann G, de Roo A et al. 2017 Global projections of river flood risk in a warmer world. *Earth's Futur.* 5, 171–182.  
doi: [10.1002/2016EF000485](https://doi.org/10.1002/2016EF000485)
- Alfieri L, Burek P, Feyen L, Forzieri G 2015 Global warming increases the frequency of river floods in Europe. *Hydrol. Earth Syst. Sci.* 19, 2247–2260.  
doi: [10.5194/hess-19-2247-2015](https://doi.org/10.5194/hess-19-2247-2015)
- Allan JA 1998 Virtual water: A strategic resource global solutions to regional deficits. *Groundwater* 36, 545–546. doi: [10.1111/j.1745-6584.1998.tb02825.x](https://doi.org/10.1111/j.1745-6584.1998.tb02825.x)
- Almagro M, de Vente J, Boix-Fayos C, García-Franco N, Melgares de Aguilar J et al. 2016 Sustainable land management practices as providers of several ecosystem services under rainfed Mediterranean agroecosystems. *Mitig. Adapt. Strateg. Glob. Chang.* 21, 1029–1043. doi: [10.1007/s11027-013-9535-2](https://doi.org/10.1007/s11027-013-9535-2)
- Andrew J, Sauquet E 2017 Climate Change Impacts and Water Management Adaptation in Two Mediterranean-Climate Watersheds: Learning from the Durance and Sacramento Rivers. *Water* 9, 126.  
doi: [10.3390/w9020126](https://doi.org/10.3390/w9020126)
- Antonelli M, Roson R, Sartori M 2012 Systemic Input-Output Computation of Green and Blue Virtual Water 'Flows' with an Illustration for the Mediterranean Region. *Water Resour. Manag.* 26, 4133–4146.  
doi: [10.1007/s11269-012-0135-9](https://doi.org/10.1007/s11269-012-0135-9)
- Antonelli M, Tamea S 2015 Food-water security and virtual water trade in the Middle East and North Africa. *Int. J. Water Resour. Dev.* 31, 326–342.  
doi: [10.1080/07900627.2015.1030496](https://doi.org/10.1080/07900627.2015.1030496)
- AQUASTAT Programme 2007 Dams and Agriculture in Africa.
- Argyroudi A, Chatzinikolaou Y, Poirazidis K, Lazaridou M 2009 Do intermittent and ephemeral Mediterranean rivers belong to the same river type? *Aquat. Ecol.* 43, 465–476. doi: [10.1007/s10452-008-9176-9](https://doi.org/10.1007/s10452-008-9176-9)
- Aureli A, Ganoulis J, Margat J 2008 Water in the Mediterranean. Groundwater resources in the Mediterranean region: importance, uses and sharing, in *IEMed Yearbook 2008*, 96–105.
- Baouab MH, Cherif S 2015 Changement climatique et ressources en eau : tendances, fluctuations et projections pour un cas d'étude de l'eau potable en Tunisie. *La Houille Blanche*, 99–107.  
doi: [10.1051/lhb/20150061](https://doi.org/10.1051/lhb/20150061)
- Barrera-Escoda A, Llasat MC 2015 Evolving flood patterns in a Mediterranean region (1301–2012) and climatic factors - the case of Catalonia. *Hydrol. Earth Syst. Sci.* 19, 465–483.  
doi: [10.5194/hess-19-465-2015](https://doi.org/10.5194/hess-19-465-2015)
- Barriandos M, Coeur D, Lang M, Llasat MC, Naulet R et al. 2003 Stationarity analysis of historical flood series in France and Spain (14<sup>th</sup>–20<sup>th</sup> centuries). *Nat. Hazards Earth Syst. Sci.* 3, 583–592.  
doi: [10.5194/nhess-3-583-2003](https://doi.org/10.5194/nhess-3-583-2003)
- Berg AM, Sheffield J, Milly PCD 2017 Divergent surface and total soil moisture projections under global warming. *Geophys. Res. Lett.* 44, 236–244.  
doi: [10.1002/2016gl071921](https://doi.org/10.1002/2016gl071921)
- Berry PM, Betts RA, Harrison PA, Sánchez-Arcilla A 2017 *High-end climate change in Europe: impacts, vulnerability and adaptation*. Sofia: Pensoft Publishers <https://upcommons.upc.edu/handle/2117/106769> [Accessed December 21, 2019]
- Betts RA, Alfieri L, Bradshaw C, Caesar J, Feyen L et al. 2018 Changes in climate extremes, fresh water availability and vulnerability to food insecurity projected at 1.5°C and 2°C global warming with a higher-resolution global climate model. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376.  
doi: [10.1098/rsta.2016.0452](https://doi.org/10.1098/rsta.2016.0452)
- Bianucci P, Sordo-Ward Á, Moralo J, Garrote L 2015 Probabilistic-multiobjective comparison of user-defined operating rules. Case study: hydropower dam in Spain. *Water* 7, 956–974.  
doi: [10.3390/w7030956](https://doi.org/10.3390/w7030956)
- Bianucci P, Sordo-Ward Á, Pérez JI, García-Palacios J, Mediero L et al. 2013 Risk-based methodology for parameter calibration of a reservoir flood control model. *Nat. Hazards Earth Syst. Sci.* 13, 965–981.  
doi: [10.5194/nhess-13-965-2013](https://doi.org/10.5194/nhess-13-965-2013)
- Blas A, Garrido A, Willaarts B 2016 Evaluating the Wa-

- ter Footprint of the Mediterranean and American Diets. *Water* 8, 448. doi: [10.3390/w8100448](https://doi.org/10.3390/w8100448)
- Blas A, Garrido A, Willaarts B 2018 Food consumption and waste in Spanish households: Water implications within and beyond national borders. *Ecol. Indic.* 89, 290–300. doi: [10.1016/j.ecolind.2018.01.057](https://doi.org/10.1016/j.ecolind.2018.01.057)
- Blinda M 2012 More efficient water use in the Mediterranean. Valbonne, France
- Blöschl G, Hall J, Parajka J, Perdigão RAP, Merz B et al. 2017 Changing climate shifts timing of European floods. *Science (80-. )*. 357, 588–590. doi: [10.1126/science.aan2506](https://doi.org/10.1126/science.aan2506)
- Blöschl G, Hall J, Viglione A, Perdigão RAP, Parajka J et al. 2019 Changing climate both increases and decreases European river floods. *Nature* 573, 108–111. doi: [10.1038/s41586-019-1495-6](https://doi.org/10.1038/s41586-019-1495-6)
- Boada M, Gómez FJ 2011 *Forests of Catalonia (in Catalan language)*. Ed. Lumberg
- Boari F, Donadio A, Schiattone MI, Cantore V 2015 Particle film technology: A supplemental tool to save water. *Agric. Water Manag.* 147, 154–162. doi: [10.1016/j.agwat.2014.07.014](https://doi.org/10.1016/j.agwat.2014.07.014)
- Bocchiola D, Rosso R 2014 Safety of Italian dams in the face of flood hazard. *Adv. Water Resour.* 71, 23–31. doi: [10.1016/j.advwatres.2014.05.006](https://doi.org/10.1016/j.advwatres.2014.05.006)
- Bocci M, Murciano C 2018 Climate Change Impact on the Tourism Sector in the Southern Mediterranean. Foreseen Developments and Policy Measures. Final Report.
- Boddu M, Gaayam T, Annamdas VGM 2011 A Review on Inter Basin Transfer of Water. in *IPWE 2011, Proceedings of 4th International Perspective on Water Resources & the Environment, January 4-6, 2011* (National University of Singapore [NUS], Singapore).
- Bonanos AM, Georgiou MC, Guillen E, Papanicolas CN 2017 CSP+D: The case study at the PROTEAS facility. in *AIP Conference Proceedings, 1850, 170001* doi: [10.1063/1.4984564](https://doi.org/10.1063/1.4984564)
- Breivik K, Sweetman A, Pacyna JM, Jones KC 2007 Towards a global historical emission inventory for selected PCB congeners - A mass balance approach. 3. An update. *Sci. Total Environ.* 377, 296–307. doi: [10.1016/j.scitotenv.2007.02.026](https://doi.org/10.1016/j.scitotenv.2007.02.026)
- Brooks DB 2006 An operational definition of water demand management. *Int. J. Water Resour. Dev.* 22, 521–528. doi: [10.1080/07900620600779699](https://doi.org/10.1080/07900620600779699)
- Brown C, Ghile Y, Lavery M, Li K 2012 Decision scaling: Linking bottom-up vulnerability analysis with climate projections in the water sector. *Water Resour. Res.* 48. doi: [10.1029/2011wr011212](https://doi.org/10.1029/2011wr011212)
- Burak S, Dogan E, Gazioglu C 2004 Impact of urbanization and tourism on coastal environment. *Ocean Coast. Manag.* 47, 515–527. doi: [10.1016/J.OCECOAMAN.2004.07.007](https://doi.org/10.1016/J.OCECOAMAN.2004.07.007)
- Burak S, Margat J 2016 Water Management in the Mediterranean Region: Concepts and Policies. *Water Resour. Manag.* 30, 5779–5797. doi: [10.1007/s11269-016-1389-4](https://doi.org/10.1007/s11269-016-1389-4)
- Burak S, Mat H 2010 Municipal water demand and efficiency analysis: Case studies in Turkey. *Water Policy* 12, 695–706. doi: [10.2166/wp.2009.209](https://doi.org/10.2166/wp.2009.209)
- Camici S, Brocca L, Moramarco T 2017 Accuracy versus variability of climate projections for flood assessment in central Italy. *Clim. Change* 141, 273–286. doi: [10.1007/s10584-016-1876-x](https://doi.org/10.1007/s10584-016-1876-x)
- Campisano A, D'Amico G, Modica C 2017 Water Saving and Cost Analysis of Large-Scale Implementation of Domestic Rain Water Harvesting in Minor Mediterranean Islands. *Water* 9, 916. doi: [10.3390/w9120916](https://doi.org/10.3390/w9120916)
- Cantore V, Lechkar O, Karabulut E, Sellami MH, Albrizio R et al. 2016 Combined effect of deficit irrigation and strobilurin application on yield, fruit quality and water use efficiency of “cherry” tomato (*Solanum lycopersicum* L.). *Agric. Water Manag.* 167, 53–61. doi: [10.1016/j.agwat.2015.12.024](https://doi.org/10.1016/j.agwat.2015.12.024)
- Cherif S, El Ayni F, Jrad AT, Trabelsi-Ayadi M 2013 Aquifer Recharge by Treated Wastewaters: Korba case study (Tunisia). *Sustain. Sanit. Pract.* 14, 41–48.
- Chezgi J, Pourghasemi HR, Naghibi SA, Moradi HR, Kheirkhah Zarkesh M 2016 Assessment of a spatial multi-criteria evaluation to site selection underground dams in the Alborz Province, Iran. *Geocarto Int.* 31, 628–646. doi: [10.1080/10106049.2015.1073366](https://doi.org/10.1080/10106049.2015.1073366)
- Civili FS 2010 The Land-Based Pollution of the Mediterranean Sea: Present State and Prospects. *Econ. Territ. - Sustain. Dev.* 34, 241–245.
- Clemmens AJ, Allen RG, Burt CM 2008 Technical concepts related to conservation of irrigation and rainwater in agricultural systems. *Water Resour. Res.* 44. doi: [10.1029/2007wr006095](https://doi.org/10.1029/2007wr006095)
- Cooley H, Christian-Smith J, Gleick PH, Allen L, Cohen M 2009 Understanding and reducing the risks of climate change for transboundary waters. Oakland, California [http://www.pacinst.org/reports/trans-boundary\\_waters/transboundary\\_water\\_and\\_climate\\_report.pdf](http://www.pacinst.org/reports/trans-boundary_waters/transboundary_water_and_climate_report.pdf)
- Cortès M, Turco M, Ward P, Sánchez-Espigares JA, Alfieri L et al. 2019 Changes in flood damage with global warming on the eastern coast of Spain. *Nat. Hazards Earth Syst. Sci.* 19, 2855–2877. doi: [10.5194/nhess-19-2855-2019](https://doi.org/10.5194/nhess-19-2855-2019)
- Ćosić M, Djurović N, Todorović M, Maletić R, Zečević B et al. 2015 Effect of irrigation regime and application of kaolin on yield, quality and water use efficiency of sweet pepper. *Agric. Water Manag.* 159, 139–147. doi: [10.1016/j.agwat.2015.05.014](https://doi.org/10.1016/j.agwat.2015.05.014)
- Cross K, Latorre C 2015 Which water for which use? Exploring water quality instruments in the context of a changing climate. *Aquat. Procedia* 5, 104–110. doi: [10.1016/j.aqpro.2015.10.012](https://doi.org/10.1016/j.aqpro.2015.10.012)
- Culley S, Noble S, Yates A, Timbs M, Westra S et al. 2016 A bottom-up approach to identifying the

- maximum operational adaptive capacity of water resource systems to a changing climate. *Water Resour. Res.* 52, 6751–6768. doi: [10.1002/2015wr018253](https://doi.org/10.1002/2015wr018253)
- Custodio E, Andreu-Rodes JM, Aragón R, Estrela T, Ferrer J et al. 2016 Groundwater intensive use and mining in south-eastern peninsular Spain: Hydrogeological, economic and social aspects. *Sci. Total Environ.* 559, 302–316. doi: [10.1016/j.scitotenv.2016.02.107](https://doi.org/10.1016/j.scitotenv.2016.02.107)
- D'Andrea F, Provenzale A, Vautard R, de Noblet-Ducoudré N 2006 Hot and cool summers: Multiple equilibria of the continental water cycle. *Geophys. Res. Lett.* 33. doi: [10.1029/2006GL027972](https://doi.org/10.1029/2006GL027972)
- Daccache A, Ciurana JS, Rodríguez-Díaz JA, Knox JW 2014 Water and energy footprint of irrigated agriculture in the Mediterranean region. *Environ. Res. Lett.* 9, 124014. doi: [10.1088/1748-9326/9/12/124014](https://doi.org/10.1088/1748-9326/9/12/124014)
- Dakhlaoui H, Ruelland D, Trambly Y 2019a A bootstrap-based differential split-sample test to assess the transferability of conceptual rainfall-runoff models under past and future climate variability. *J. Hydrol.* 575, 470–486. doi: [10.1016/j.jhydrol.2019.05.056](https://doi.org/10.1016/j.jhydrol.2019.05.056)
- Dakhlaoui H, Seibert J, Hakala K 2019b Hydrological Impacts of Climate Change in Northern Tunisia, in *Advances in Sustainable and Environmental Hydrology, Hydrogeology, Hydrochemistry and Water Resources. Proceedings of the 1st Springer Conference of the Arabian Journal of Geosciences (CAJG-1), Tunisia 2018*, eds. Chaminé HI, Barbieri M, Kisi O, Chen M, Merkel BJ (Springer International Publishing), 301–303. doi: [10.1007/978-3-030-01572-5\\_71](https://doi.org/10.1007/978-3-030-01572-5_71)
- Dakhlaoui H, Seibert J, Hakala K 2020 Sensitivity of discharge projections to potential evapotranspiration estimation in Northern Tunisia. *Reg. Environ. Chang.* 20, 1–12. doi: [10.1007/s10113-020-01615-8](https://doi.org/10.1007/s10113-020-01615-8)
- Dankers R, Feyen L 2009 Flood hazard in Europe in an ensemble of regional climate scenarios. *JGR Atmos.* 114. doi: [10.1029/2008jd011523](https://doi.org/10.1029/2008jd011523)
- Danovaro R, Fonda Umani S, Pusceddu A, Umani SF, Pusceddu A et al. 2009 Climate Change and the Potential Spreading of Marine Mucilage and Microbial Pathogens in the Mediterranean Sea. *PLoS One* 4, e7006. doi: [10.1371/journal.pone.0007006](https://doi.org/10.1371/journal.pone.0007006)
- Dayon G, Boé J, Martin É, Gailhard J 2018 Impacts of climate change on the hydrological cycle over France and associated uncertainties. *Comptes Rendus Geosci.* 350, 141–153. doi: [10.1016/J.CRTE.2018.03.001](https://doi.org/10.1016/J.CRTE.2018.03.001)
- de Giglio O, Caggiano G, Apollonio F, Marzella A, Brigida S et al. 2018 The aquifer recharge: An overview of the legislative and planning aspect. *Ann. di Ig.* 30, 34–43. doi: [10.7416/ai.2018.2193](https://doi.org/10.7416/ai.2018.2193)
- de Perthuis C, Hallegatte S, Lecocq F 2010 Économie de l'adaptation au changement climatique. de Stefano L 2004 Freshwater and Tourism in the Mediterranean. Rome, Italy
- de Stefano L, Fornés JM, López-Geta JA, Villarroya F 2014 Groundwater use in Spain: an overview in light of the EU Water Framework Directive. *Int. J. Water Resour. Dev.* 31, 640–656. doi: [10.1080/07900627.2014.938260](https://doi.org/10.1080/07900627.2014.938260)
- di Baldassarre G, Kreibich H, Vorogushyn S, Aerts JCJH, Arnbjerg-Nielsen K et al. 2018 HESS Opinions: An interdisciplinary research agenda to explore the unintended consequences of structural flood protection. *Hydrol. Earth Syst. Sci.* 22, 5629–5637. doi: [10.5194/hess-22-5629-2018](https://doi.org/10.5194/hess-22-5629-2018)
- Diakakis M 2014 An inventory of flood events in Athens, Greece, during the last 130 years. Seasonality and spatial distribution. *J. Flood Risk Manag.* 7, 332–343. doi: [10.1111/jfr3.12053](https://doi.org/10.1111/jfr3.12053)
- Dittmann R, Froehlich F, Pohl R, Ostrowski M 2009 Optimum multi-objective reservoir operation with emphasis on flood control and ecology. *Nat. Hazards Earth Syst. Sci.* 9, 1973–1980. doi: [10.5194/nhess-9-1973-2009](https://doi.org/10.5194/nhess-9-1973-2009)
- Djuma H, Bruggeman A, Camera C, Eliades M, Kostarelos K 2017a The Impact of a Check Dam on Groundwater Recharge and Sedimentation in an Ephemeral Stream. *Water* 9, 813. doi: [10.3390/w9100813](https://doi.org/10.3390/w9100813)
- Djuma H, Bruggeman A, Camera C, Zoumidis C 2017b Combining Qualitative and Quantitative Methods for Soil Erosion Assessments: An Application in a Sloping Mediterranean Watershed, Cyprus. *L. Degrad. Dev.* 28, 243–254. doi: [10.1002/ldr.2571](https://doi.org/10.1002/ldr.2571)
- Djuma H, Bruggeman A, Eliades M, Lange MA 2016 Non-conventional water resources research in semi-arid countries of the Middle East. *Desalin. Water Treat.* 57, 2290–2303. doi: [10.1080/19443994.2014.984930](https://doi.org/10.1080/19443994.2014.984930)
- Döll P, Trautmann T, Gerten D, Müller Schmied H, Ostberg S et al. 2018 Risks for the global freshwater system at 1.5 °C and 2 °C global warming. *Environ. Res. Lett.* 13, 044038. doi: [10.1088/1748-9326/aab792](https://doi.org/10.1088/1748-9326/aab792)
- Donnelly C, Greuell W, Andersson J, Gerten D, Pisacane G et al. 2017 Impacts of climate change on European hydrology at 1.5, 2 and 3 degrees mean global warming above preindustrial level. *Clim. Change* 143, 13–26. doi: [10.1007/s10584-017-1971-7](https://doi.org/10.1007/s10584-017-1971-7)
- Donta AA, Lange MA, MEDIS Consortium 2008 Water Management on Mediterranean Islands: Pressure, Recommended Policy and Management Options, in *Coping with Water Deficiency. Environment & Policy, vol 48* (Springer, Dordrecht), 11–44. doi: [10.1007/978-1-4020-6615-3\\_2](https://doi.org/10.1007/978-1-4020-6615-3_2)
- Dorigo WA, de Jeu RAM, Chung D, Parinussa RM, Liu Y et al. 2012 Evaluating global trends (1988–2010) in harmonized multi-satellite surface soil moisture. *Geophys. Res. Lett.* 39. doi: [10.1029/2012gl052988](https://doi.org/10.1029/2012gl052988)
- Droogers P, Immerzeel WW, Terink W, Hoogeveen J,

- Bierkens MFP et al. 2012 Water resources trends in Middle East and North Africa towards 2050. *Hydrol. Earth Syst. Sci.* 16, 3101–3114. doi: [10.5194/hess-16-3101-2012](https://doi.org/10.5194/hess-16-3101-2012)
- Dumas P, Hallegatte S, Quintana-Seguí P, Martin E 2013 The influence of climate change on flood risks in France – first estimates and uncertainty analysis. *Nat. Hazards Earth Syst. Sci.* 13, 809–821. doi: [10.5194/nhess-13-809-2013](https://doi.org/10.5194/nhess-13-809-2013)
- EEA 2006 *Priority issues in the Mediterranean environment*. Copenhagen, Denmark
- EEA 2017 *Qualité des eaux de baignade européenne en 2016*.
- EEA 2018 *Reservoirs and dams*.
- El-Nashar AM 2001 Cogeneration for power and desalination - state of the art review. *Desalination* 134, 7–28. doi: [10.1016/s0011-9164\(01\)00111-4](https://doi.org/10.1016/s0011-9164(01)00111-4)
- El Ayni F, Cherif S, Jrad A, Trabelsi-Ayadi M 2011 Impact of Treated Wastewater Reuse on Agriculture and Aquifer Recharge in a Coastal Area: Korba Case Study. *Water Resour. Manag.* 25, 2251–2265. doi: [10.1007/s11269-011-9805-2](https://doi.org/10.1007/s11269-011-9805-2)
- El Ayni F, Cherif S, Jrad A, Trabelsi-Ayadi M 2012 A New Approach for the Assessment of Groundwater Quality and Its Suitability for Irrigation: A Case Study of the Korba Coastal Aquifer (Tunisia, Africa). *Water Environ. Res.* 84, 673–681. doi: [10.2175/106143012x13378023685673](https://doi.org/10.2175/106143012x13378023685673)
- Elimelech M, Phillip WA 2011 The Future of Seawater Desalination: Energy, Technology, and the Environment. *Science (80-. )*. 333, 712–717. doi: [10.1126/science.1200488](https://doi.org/10.1126/science.1200488)
- Eltawil MA, Zhengming Z, Yuan L 2009 A review of renewable energy technologies integrated with desalination systems. *Renew. Sustain. Energy Rev.* 13, 2245–2262. doi: [10.1016/j.rser.2009.06.011](https://doi.org/10.1016/j.rser.2009.06.011)
- EPA 2001 *Parameters Of Water Quality - Interpretation and Standards*. Wexford, Ireland
- Escriba-Bou A, Pulido-Velazquez M, Pulido-Velazquez D 2017 Economic Value of Climate Change Adaptation Strategies for Water Management in Spain's Jucar Basin. *J. Water Resour. Plan. Manag.* 143, 04017005. doi: [10.1061/\(ASCE\)WR.1943-5452.0000735](https://doi.org/10.1061/(ASCE)WR.1943-5452.0000735)
- Estado de los embalses, pantanos y presas de España 2019 <https://www.embalses.net/> [Accessed January 10, 2020]
- Etteieb S, Cherif S, Kawachi A, Han J, Elayni F et al. 2016 Combining Biological and Chemical Screenings to Assess Cytotoxicity of Emerging Contaminants in Discharges into Surface Water. *Water, Air, Soil Pollut.* 227, 341. doi: [10.1007/s11270-016-3049-y](https://doi.org/10.1007/s11270-016-3049-y)
- Etteieb S, Cherif S, Tarhouni J 2017 Hydrochemical assessment of water quality for irrigation: a case study of the Medjerda River in Tunisia. *Appl. Water Sci.* 7, 469–480. doi: [10.1007/s13201-015-0265-3](https://doi.org/10.1007/s13201-015-0265-3)
- European Water Movement 2018 *Barrages et hydroélectricité du Rhône*. <http://europeanwater.org/fr/ressources/rapports-et-publications/746-barrages-et-hydroelectricite-du-rhone> [Accessed January 10, 2020]
- Fader M, Cranmer C, Lawford R, Engel-Cox J 2018 Toward an Understanding of Synergies and Trade-Offs Between Water, Energy, and Food SDG Targets. *Front. Environ. Sci.* 6, 1–11. doi: [10.3389/fenvs.2018.00112](https://doi.org/10.3389/fenvs.2018.00112)
- Fader M, Gerten D, Krause M, Lucht W, Cramer W 2013 Spatial decoupling of agricultural production and consumption: quantifying dependences of countries on food imports due to domestic land and water constraints. *Environ. Res. Lett.* 8. doi: [10.1088/1748-9326/8/1/014046](https://doi.org/10.1088/1748-9326/8/1/014046)
- Fader M, Gerten D, Thammer M, Heinke J, Lotze-Campen H et al. 2011 Internal and external green-blue agricultural water footprints of nations, and related water and land savings through trade. *Hydrol. Earth Syst. Sci.* 15, 1641–1660. doi: [10.5194/hess-15-1641-2011](https://doi.org/10.5194/hess-15-1641-2011)
- Fader M, Shi S, Von Bloh W, Bondeau A, Cramer W 2016 Mediterranean irrigation under climate change: More efficient irrigation needed to compensate for increases in irrigation water requirements. *Hydrol. Earth Syst. Sci.* 20, 953–973. doi: [10.5194/hess-20-953-2016](https://doi.org/10.5194/hess-20-953-2016)
- Fader M, Von Bloh W, Shi S, Bondeau A, Cramer W 2015 Modelling Mediterranean agro-ecosystems by including agricultural trees in the LPJmL model. *Geosci. Model Dev.* 8, 3545–3561. doi: [10.5194/gmd-8-3545-2015](https://doi.org/10.5194/gmd-8-3545-2015)
- FAO 2003 *Review of World Water Resources by Country*. Water Reports 23. Rome
- FAO 2015 *Towards a Regional Collaborative Strategy on Sustainable Water Management and Food Security in the Near East and North Africa Region*. Cairo, Egypt.
- FAO 2016a AQUASTAT Main Database, Food and Agriculture Organization of the United Nations (FAO). <http://www.fao.org/nr/water/aquastat/data/query/index.html?lang=en> [Accessed July 23, 2019]
- FAO 2016b *Assessment of the water harvesting sector in Jordan*. Final Report. Rome, Italy
- FAO 2017 *Integrated Monitoring Guide for SDG 6*. Step-by-step monitoring methodology for indicator 6.4.1 on water-use efficiency.
- Feng H 2016 Individual contributions of climate and vegetation change to soil moisture trends across multiple spatial scales. *Sci. Rep.* 6, 32782. doi: [10.1038/srep32782](https://doi.org/10.1038/srep32782)
- Fernández García I, Rodríguez Díaz JA, Camacho Poyato E, Montesinos P, Berbel J 2014 Effects of modernization and medium term perspectives on water and energy use in irrigation districts. *Agric. Syst.* 131, 56–63. doi: [10.1016/j.agry.2014.08.002](https://doi.org/10.1016/j.agry.2014.08.002)
- Ferragina E 2010 *The Water Issue in the Mediterranean*.



- [https://www.iemed.org/observatori-es/arees-danalisi/arxiu-adjunts/10-papers-for-barcelona-2010/8-environmental-and-sustainable-development-in-the-mediterranean/ferragina\\_8.pdf](https://www.iemed.org/observatori-es/arees-danalisi/arxiu-adjunts/10-papers-for-barcelona-2010/8-environmental-and-sustainable-development-in-the-mediterranean/ferragina_8.pdf)
- Feyen L, Dankers R 2009 Impact of global warming on streamflow drought in Europe. *JGR Atmos.* 114. doi: [10.1029/2008jd011438](https://doi.org/10.1029/2008jd011438)
- Fiseha BM, Setegn SG, Melesse AM, Volpi E, Fiori A 2014 Impact of Climate Change on the Hydrology of Upper Tiber River Basin Using Bias Corrected Regional Climate Model. *Water Resour. Manag.* 28, 1327–1343. doi: [10.1007/s11269-014-0546-x](https://doi.org/10.1007/s11269-014-0546-x)
- Förster J 2014 Cooling for electricity production dominates water use in industry. *Stat. Focus. Eurostat* 14/2014. [https://ec.europa.eu/eurostat/statistics-explained/index.php/Archive:Water\\_use\\_in\\_industry](https://ec.europa.eu/eurostat/statistics-explained/index.php/Archive:Water_use_in_industry)
- Forzieri G, Feyen L, Rojas R, Flörke M, Wimmer F et al. 2014 Ensemble projections of future streamflow droughts in Europe. *Hydrol. Earth Syst. Sci.* 18, 85–108. doi: [10.5194/hess-18-85-2014](https://doi.org/10.5194/hess-18-85-2014)
- Foster SSD, Chilton PJ 2004 Downstream of downtown: urban wastewater as groundwater recharge. *Hydrogeol. J.* 12, 115–120. doi: [10.1007/s10040-003-0296-y](https://doi.org/10.1007/s10040-003-0296-y)
- Foster T, Brozović N, Butler AP 2017 Effects of initial aquifer conditions on economic benefits from groundwater conservation. *Water Resour. Res.* 53, 744–762. doi: [10.1002/2016WR019365](https://doi.org/10.1002/2016WR019365)
- France Hydroélectricité 2018 L'énergie Hydraulique: Chiffres clés 2017. <http://www.france-hydro-electricite.fr/lenergie-hydraulique/chiffres-cles> [Accessed November 20, 2019]
- Funes I, Savé R, Rovira P, Molowny-Horas R, Alcañiz JM et al. 2019 Agricultural soil organic carbon stocks in the north-eastern Iberian Peninsula: Drivers and spatial variability. *Sci. Total Environ.* 668, 283–294. doi: [10.1016/j.scitotenv.2019.02.317](https://doi.org/10.1016/j.scitotenv.2019.02.317)
- Gabarda-Mallorquí A, García X, Ribas A 2017 Mass tourism and water efficiency in the hotel industry: A case study. *Int. J. Hosp. Manag.* 61, 82–93. doi: [10.1016/j.ijhm.2016.11.006](https://doi.org/10.1016/j.ijhm.2016.11.006)
- Gaertner MÁ, Christensen OB, Prego JA, Polcher J, Gallardo C et al. 2001 The impact of deforestation on the hydrological cycle in the western Mediterranean: an ensemble study with two regional climate models. *Clim. Dyn.* 17, 857–873. doi: [10.1007/s003820100151](https://doi.org/10.1007/s003820100151)
- Galdies C, Vella K 2019 Future Climate Change Impacts on Malta's Agriculture, Based on Multi-model Results from WCRP's CMIP5. *Clim. Chang. Manag.*, 137–156. doi: [10.1007/978-3-319-75004-0\\_8](https://doi.org/10.1007/978-3-319-75004-0_8)
- Ganoulis J 2006 Water resources management and environmental security in Mediterranean transboundary river basins, in *Environmental Security and Environmental Management: The Role of Risk Assessment*, eds. Morel B, Linkov I (Springer Netherlands), 49–58. doi: [10.1007/1-4020-3893-3](https://doi.org/10.1007/1-4020-3893-3)
- García-Rodríguez L 2003 Renewable energy applications in desalination: state of the art. *Sol. Energy* 75, 381–393. doi: [10.1016/j.solener.2003.08.005](https://doi.org/10.1016/j.solener.2003.08.005)
- García-Ruiz JM, López Moreno JI, Vicente-Serrano SM, Lasanta-Martínez T, Beguería S 2011 Mediterranean water resources in a global change scenario. *Earth-Science Rev.* 105, 121–139. doi: [10.1016/J.EARSCIREV.2011.01.006](https://doi.org/10.1016/J.EARSCIREV.2011.01.006)
- García Lorenzo R, Conesa García C, Martínez Salvador A 2013 Assessing Soil Erosion in Semi-Arid Check Dam Watersheds Using GeoWEPP (South-East Spain), in *Geomorphology: Processes Taxonomy and Applications*, ed. Miguel, H.S. and DCP (Hauppauge, NY, USA: Nova Science), 115–146.
- Garrido A, Iglesias A 2006 Groundwater's role in managing water scarcity in the Mediterranean Region. in *International Symposium on Groundwater Sustainability*, 113–138.
- Garrido A, Martínez-Santos P, Llamas MR 2006 Groundwater irrigation and its implications for water policy in semiarid countries: the Spanish experience. *Hydrogeol. J.* 14, 340–349. doi: [10.1007/s10040-005-0006-z](https://doi.org/10.1007/s10040-005-0006-z)
- Gaume E, Borga M, Llasat MC, Maouche S, Lang M et al. 2016 Mediterranean extreme floods and flash floods, in *The Mediterranean Region under Climate Change. A Scientific Update* Coll. Synthèses., eds. Thiébaud S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 133–144. <https://hal.archives-ouvertes.fr/hal-01465740>
- GEF 2015 Global Diagnostic on Groundwater Governance. Rome
- Georgiou MC, Bonanos AM, Georgiadis JG 2016 Evaluation of a solar powered distillation unit as a mitigation to water scarcity and climate change in Cyprus. *Desalin. Water Treat.* 57, 2325–2335. doi: [10.1080/19443994.2014.989637](https://doi.org/10.1080/19443994.2014.989637)
- Giordano M 2009 Global Groundwater? Issues and Solutions. *Annu. Rev. Environ. Resour.* 34, 153–178. doi: [10.1146/annurev.enviro.030308.100251](https://doi.org/10.1146/annurev.enviro.030308.100251)
- Girard C, Pulido-Velazquez M, Rinaudo J-D, Pagé C, Caballero Y 2015 Integrating top-down and bottom-up approaches to design global change adaptation at the river basin scale. *Glob. Environ. Chang.* 34, 132–146. doi: [10.1016/j.gloenvcha.2015.07.002](https://doi.org/10.1016/j.gloenvcha.2015.07.002)
- Giuntoli I, Renard B, Lang M 2012 Floods in France, in *Changes in Flood Risk in Europe*, ed. Kundzewicz ZW (CRC Press), 199–211. doi: [10.1201/b12348-10](https://doi.org/10.1201/b12348-10)
- Giuntoli I, Vidal J-P, Prudhomme C, Hannah DM 2015 Future hydrological extremes: the uncertainty from multiple global climate and global hydrological models. *Earth Syst. Dyn.* 6, 267–285. doi: [10.5194/esd-6-267-2015](https://doi.org/10.5194/esd-6-267-2015)
- Givati A, Thirel G, Rosenfeld D, Paz D 2019 Climate change impacts on streamflow at the upper Jordan River based on an ensemble of regional climate models. *J. Hydrol. Reg. Stud.* 21, 92–109.

- doi: [10.1016/J.EJRH.2018.12.004](https://doi.org/10.1016/J.EJRH.2018.12.004)
- Glaser R, Riemann D, Schönbein J, Barriendos M, Brázdil R et al. 2010 The variability of European floods since AD 1500. *Clim. Change* 101, 235–256. doi: [10.1007/s10584-010-9816-7](https://doi.org/10.1007/s10584-010-9816-7)
- Global Water Partnership 2011 What is IWRM? <https://www.gwp.org/en/GWP-CEE/about/why/what-is-iwrm/> [Accessed September 9, 2019]
- Gohari A, Eslamian S, Mirchi A, Abedi-Koupaei J, Masah Bavani A et al. 2013 Water transfer as a solution to water shortage: A fix that can Backfire. *J. Hydrol.* 491, 23–39. doi: [10.1016/J.JHYDROL.2013.03.021](https://doi.org/10.1016/J.JHYDROL.2013.03.021)
- Gonçalvès J, Deschamps P, Hamelin B, Vallet-Coulomb C, Petersen J et al. 2020 Revisiting recharge and sustainability of the North-Western Sahara aquifers. *Reg. Environ. Chang.* 20. doi: [10.1007/s10113-020-01627-4](https://doi.org/10.1007/s10113-020-01627-4)
- Gonçalvès J, Petersen J, Deschamps P, Hamelin B, Baba-Sy O 2013 Quantifying the modern recharge of the “fossil” Sahara aquifers. *Geophys. Res. Lett.* 40, 2673–2678. doi: [10.1002/grl.50478](https://doi.org/10.1002/grl.50478)
- Gosling SN, Zaherpour J, Mount NJ, Hattermann FF, Dankers R et al. 2017 A comparison of changes in river runoff from multiple global and catchment-scale hydrological models under global warming scenarios of 1 °C, 2 °C and 3 °C. *Clim. Change* 141, 577–595. doi: [10.1007/s10584-016-1773-3](https://doi.org/10.1007/s10584-016-1773-3)
- Gössling S, Hall CM, Scott D 2015 *Tourism and Water*. Channel View Publications. doi: [10.21832/9781845415006](https://doi.org/10.21832/9781845415006)
- Gössling S, Peeters P, Hall CM, Ceron J-P, Dubois G et al. 2012 Tourism and water use: Supply, demand, and security. An international review. *Tour. Manag.* 33, 1–15. doi: [10.1016/j.tourman.2011.03.015](https://doi.org/10.1016/j.tourman.2011.03.015)
- Grayman WM, Murray R, Savic DA 2009 Effects of Redesign of Water Systems for Security and Water Quality Factors. *World Environ. Water Resour. Congr. 2009* 41036, 1–11. doi: [10.1061/41036\(342\)49](https://doi.org/10.1061/41036(342)49)
- Green JK, Seneviratne SI, Berg AM, Findell KL, Hagemann S et al. 2019 Large influence of soil moisture on long-term terrestrial carbon uptake. *Nature* 565, 476–479. doi: [10.1038/s41586-018-0848-x](https://doi.org/10.1038/s41586-018-0848-x)
- Green TR, Ahuja LR, Benjamin JG 2003 Advances and challenges in predicting agricultural management effects on soil hydraulic properties. *Geoderma* 116, 3–27. doi: [10.1016/s0016-7061\(03\)00091-0](https://doi.org/10.1016/s0016-7061(03)00091-0)
- Grillakis MG 2019 Increase in severe and extreme soil moisture droughts for Europe under climate change. *Sci. Total Environ.* 660, 1245–1255. doi: [10.1016/J.SCITOTENV.2019.01.001](https://doi.org/10.1016/J.SCITOTENV.2019.01.001)
- Grouillet B, Fabre J, Ruelland D, Dezetter A 2015 Historical reconstruction and 2050 projections of water demand under anthropogenic and climate changes in two contrasted Mediterranean catchments. *J. Hydrol.* 522, 684–696. doi: [10.1016/j.jhydrol.2015.01.029](https://doi.org/10.1016/j.jhydrol.2015.01.029)
- Grum B, Hessel R, Kessler A, Woldearegay K, Yazew E et al. 2016 A decision support approach for the selection and implementation of water harvesting techniques in arid and semi-arid regions. *Agric. Water Manag.* 173, 35–47. doi: [10.1016/j.agwat.2016.04.018](https://doi.org/10.1016/j.agwat.2016.04.018)
- Gudmundsson L, Seneviratne SI 2016 Anthropogenic climate change affects meteorological drought risk in Europe. *Environ. Res. Lett.* 11, 044005. doi: [10.1088/1748-9326/11/4/044005](https://doi.org/10.1088/1748-9326/11/4/044005)
- Gudmundsson L, Seneviratne SI, Zhang X 2017 Anthropogenic climate change detected in European renewable freshwater resources. *Nat. Clim. Chang.* 7, 813–816. doi: [10.1038/nclimate3416](https://doi.org/10.1038/nclimate3416)
- Guilherme MR, Aouada FA, Fajardo AR, Martins AF, Paulino AT et al. 2015 Superabsorbent hydrogels based on polysaccharides for application in agriculture as soil conditioner and nutrient carrier: A review. *Eur. Polym. J.* 72, 365–385. doi: [10.1016/j.eurpolymj.2015.04.017](https://doi.org/10.1016/j.eurpolymj.2015.04.017)
- Guiot J, Cramer W 2016 Climate change: The 2015 Paris Agreement thresholds and Mediterranean basin ecosystems. *Science (80-. )*. 354, 4528–4532. doi: [10.1126/science.aah5015](https://doi.org/10.1126/science.aah5015)
- Gul M, Akpınar MG, Ceylan RF 2017 Water use efficiency of urban households in the Mediterranean region of Turkey. *Desalin. Water Treat.* 76, 364–368. doi: [10.5004/dwt.2017.20445](https://doi.org/10.5004/dwt.2017.20445)
- Gupta J, van der Zaag P 2008 Interbasin water transfers and integrated water resources management: Where engineering, science and politics interlock. *Phys. Chem. Earth, Parts A/B/C* 33, 28–40. doi: [10.1016/j.pce.2007.04.003](https://doi.org/10.1016/j.pce.2007.04.003)
- Hadjikakou M, Chenoweth J, Miller G 2013 Estimating the direct and indirect water use of tourism in the eastern Mediterranean. *J. Environ. Manage.* 114, 548–556. doi: [10.1016/j.jenvman.2012.11.002](https://doi.org/10.1016/j.jenvman.2012.11.002)
- Hall J, Arheimer B, Borga M, Brázdil R, Claps P et al. 2014 Understanding flood regime changes in Europe: a state-of-the-art assessment. *Hydrol. Earth Syst. Sci.* 18, 2735–2772. doi: [10.5194/hess-18-2735-2014](https://doi.org/10.5194/hess-18-2735-2014)
- Hamdy A, Abu-Zeid M, Lacirignola C 1995 Water Crisis in the Mediterranean: Agricultural Water Demand Management. *Water Int.* 20, 176–187. doi: [10.1080/02508069508686473](https://doi.org/10.1080/02508069508686473)
- Hashemi H, Berndtsson R, Persson M 2015 Artificial recharge by floodwater spreading estimated by water balances and groundwater modelling in arid Iran. *Hydrol. Sci. J.* 60, 336–350. doi: [10.1080/02626667.2014.881485](https://doi.org/10.1080/02626667.2014.881485)
- Hassing J, Ipsen N, Jonch Clausen T, Larsen H, Lindgaard-Jorgensen P 2009 Integrated water resources management in action. <https://unesdoc.unesco.org/ark:/48223/pf0000181891>
- Hernández-Mora N, del Moral Ituarte L, La-Roca F, La

- Calle A, Schmidt G 2014 Interbasin Water Transfers in Spain: Interregional Conflicts and Governance Responses. *Glob. Water*, 175–194. doi: [10.1007/978-94-007-7323-3\\_13](https://doi.org/10.1007/978-94-007-7323-3_13)
- Hodge AT 2000 Reservoirs and dams, in *Handbook of Ancient Water Technology*, ed. Wikander Ö (Leiden, The Netherlands), 331–342.
- Hoegh-Guldberg O, Jacob D, Taylor M, Bindi M, Brown S et al. 2018 Impacts of 1.5°C of global warming on natural and human systems, in *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, eds. Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J et al. (Cambridge, United Kingdom and New York, NY, USA: In press), 175–311.
- Hoekstra AY, Chapagain AK. 2008 *Globalization of Water: Sharing the Planet's Freshwater Resources*. Wiley-Blackwell
- Hoekstra AY, Mekonnen MM 2012 The water footprint of humanity. *Proc. Natl. Acad. Sci. U. S. A.* 109, 3232–3237. doi: [10.1073/pnas.1109936109](https://doi.org/10.1073/pnas.1109936109)
- Hoff H, Bonzi C, Joyce B, Tielbörger K 2011 A Water Resources Planning Tool for the Jordan River Basin. *Water* 3, 718–736. doi: [10.3390/w3030718](https://doi.org/10.3390/w3030718)
- Ibáñez J, Valderrama JM, Puigdefábregas J 2008 Assessing overexploitation in Mediterranean aquifers using system stability condition analysis. *Ecol. Model.* 218, 260–266. doi: [10.1016/j.ecolmodel.2008.07.004](https://doi.org/10.1016/j.ecolmodel.2008.07.004)
- ICOLD 2018 Number of Dams by Country Members. *Int. Comm. Large Dams*. [https://www.icold-cigb.org/article/GB/world\\_register/general\\_synthesis/number-of-dams-by-country-members](https://www.icold-cigb.org/article/GB/world_register/general_synthesis/number-of-dams-by-country-members) [Accessed March 9, 2019]
- ICOLD 2019 Definition of a Large Dam, International Committee on Large Dams. [https://www.icold-cigb.org/GB/dams/definition\\_of\\_a\\_large\\_dam.asp](https://www.icold-cigb.org/GB/dams/definition_of_a_large_dam.asp) [Accessed May 4, 2019]
- IDAEA 2015 Water Strategy in the Western Mediterranean. <https://www.idaea.csic.es/sites/default/files/5%2B5-Water-strategy-in-the-western-Mediterranean.pdf>
- Idelovitch E 1978 Wastewater reuse by biological-chemical treatment and groundwater recharge. *J. Water Pollut. Control Fed.* 50, 2723–2740.
- Iglesias A, Garrote L, Flores F, Moneo M 2007 Challenges to Manage the Risk of Water Scarcity and Climate Change in the Mediterranean. *Water Resour. Manag.* 21, 775–788. doi: [10.1007/s11269-006-9111-6](https://doi.org/10.1007/s11269-006-9111-6)
- Iglesias A, Mougou R, Moneo M, Quiroga S 2011 Towards adaptation of agriculture to climate change in the Mediterranean. *Reg. Environ. Chang.* 11, 159–166. doi: [10.1007/s10113-010-0187-4](https://doi.org/10.1007/s10113-010-0187-4)
- Ilić ZS, Milenković L, Stanojević L, Cvetković D, Fallik E 2012 Effects of the modification of light intensity by color shade nets on yield and quality of tomato fruits. *Sci. Hortic. (Amsterdam)*. 139, 90–95. doi: [10.1016/j.scienta.2012.03.009](https://doi.org/10.1016/j.scienta.2012.03.009)
- IPEMED 2018 Reuse of treated waste water in the Mediterranean and impacts on territories. [http://www.ipemed.coop/adminIpemed/media/fich\\_articulo/1521051262\\_palimpseste-n19-en.pdf](http://www.ipemed.coop/adminIpemed/media/fich_articulo/1521051262_palimpseste-n19-en.pdf)
- Janža M 2011 Impact assessment of projected climate change on the hydrological regime in the SE Alps, Upper Soča River basin, Slovenia. *Nat. Hazards* 67, 1025–1043. doi: [10.1007/s11069-011-9892-7](https://doi.org/10.1007/s11069-011-9892-7)
- Jiménez Cisneros BE, Oki T, Arnell NW, Benito G, Cogley JG et al. 2014 Freshwater Resources, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 229–269. [https://www.ipcc.ch/pdf/assessment-report/ar5/wg2/WGIAR5-Chap3\\_FINAL.pdf](https://www.ipcc.ch/pdf/assessment-report/ar5/wg2/WGIAR5-Chap3_FINAL.pdf) [Accessed March 11, 2017]
- Jódar-Abellán A, Albaladejo-García JA, Prats-Rico D 2017 Artificial groundwater recharge. Review of the current knowledge of the technique. *Rev. la Soc. Geológica España* 30, 85–96.
- Johansson EL, Fader M, Seaquist JW, Nicholas KA 2016 Green and blue water demand from large-scale land acquisitions in Africa. *Proc. Natl. Acad. Sci. U. S. A.* 113, 11471–11476. doi: [10.1073/pnas.1524741113](https://doi.org/10.1073/pnas.1524741113)
- Jones E, Qadir M, van Vliet MTH, Smakhtin V, Kang S 2019 The state of desalination and brine production: A global outlook. *Sci. Total Environ.* 657, 1343–1356. doi: [10.1016/j.scitotenv.2018.12.076](https://doi.org/10.1016/j.scitotenv.2018.12.076)
- Jung M, Reichstein M, Ciais P, Seneviratne SI, Sheffield J et al. 2010 Recent decline in the global land evapotranspiration trend due to limited moisture supply. *Nature* 467, 951–954. doi: [10.1038/nature09396](https://doi.org/10.1038/nature09396)
- Kalogirou SA 2004 Solar thermal collectors and applications. *Prog. Energy Combust. Sci.* 30, 231–295. doi: [10.1016/j.peccs.2004.02.001](https://doi.org/10.1016/j.peccs.2004.02.001)
- Kang S, Hao X, Du T, Tong L, Su X et al. 2017 Improving agricultural water productivity to ensure food security in China under changing environment: From research to practice. *Agric. Water Manag.* 179, 5–17. doi: [10.1016/j.agwat.2016.05.007](https://doi.org/10.1016/j.agwat.2016.05.007)
- Karakaya N, Evrendilek F, Gonenc E 2014 Interbasin Water Transfer Practices in Turkey. *J. Ecosyst. Ecography* 04, 149–153. doi: [10.4172/2157-7625.1000149](https://doi.org/10.4172/2157-7625.1000149)
- Kassam A, Friedrich T, Derpsch R, Lahmar R, Mrabet R et al. 2012 Conservation agriculture in the dry Mediterranean climate. *F. Crop. Res.* 132, 7–17.

- doi: [10.1016/j.fcr.2012.02.023](https://doi.org/10.1016/j.fcr.2012.02.023)
- Kazner C, Wintgens T, Dillon P 2012 Water Reclamation Technologies for Safe Managed Aquifer Recharge, in *Water Reclamation Technologies for Safe Managed Aquifer Recharge*, eds. Kazner C, Wintgens T, Dillon P (IWA Publishing), 460. doi: [10.2166/9781780400648](https://doi.org/10.2166/9781780400648)
- Khaled-Khodja S, Cherif S, Durand G 2018 Seasonal assessment of metal trace element contamination by PCA in Seybouse wadi (Algeria). *Water Supply* 18, 1897–1905. doi: [10.2166/ws.2018.010](https://doi.org/10.2166/ws.2018.010)
- Khawaji AD, Kutubkhanah IK, Wie J-M 2008 Advances in seawater desalination technologies. *Desalination* 221, 47–69. doi: [10.1016/j.desal.2007.01.067](https://doi.org/10.1016/j.desal.2007.01.067)
- Kløve B, Allan A, Bertrand G, Druzyńska E, Ertürk A et al. 2011 Groundwater dependent ecosystems. Part II. Ecosystem services and management in Europe under risk of climate change and land use intensification. *Environ. Sci. Policy* 14, 782–793. doi: [10.1016/j.envsci.2011.04.005](https://doi.org/10.1016/j.envsci.2011.04.005)
- Koirala S, Hirabayashi Y, Mahendran R, Kanae S 2014 Global assessment of agreement among stream-flow projections using CMIP5 model outputs. *Environ. Res. Lett.* 9, 64017. doi: [10.1088/1748-9326/9/6/064017](https://doi.org/10.1088/1748-9326/9/6/064017)
- Koussis AD, Georgopoulou E, Kotronarou A, Mazi K, Restrepo P et al. 2010 Cost-efficient management of coastal aquifers via recharge with treated wastewater and desalination of brackish groundwater: application to the Akrotiri basin and aquifer, Cyprus. *Hydrol. Sci. J.* 55, 1234–1245. doi: [10.1080/02626667.2010.512469](https://doi.org/10.1080/02626667.2010.512469)
- Koutroulis AG, Grillakis MG, Daliakopoulos IN, Tsanis IK, Jacob D 2016 Cross sectoral impacts on water availability at +2 degrees C and +3 degrees C for east Mediterranean island states: The case of Crete. *J. Hydrol.* 532, 16–28. doi: [10.1016/j.jhydrol.2015.11.015](https://doi.org/10.1016/j.jhydrol.2015.11.015)
- Koutroulis AG, Papadimitriou L V., Grillakis MG, Tsanis IK, Wyser K et al. 2018 Simulating Hydrological Impacts under Climate Change: Implications from Methodological Differences of a Pan European Assessment. *Water* 10, 1331. doi: [10.3390/w10101331](https://doi.org/10.3390/w10101331)
- Koutroulis AG, Vrohidou A-EK, Tsanis IK 2011 Spatio-temporal Characteristics of Meteorological Drought for the Island of Crete. *J. Hydrometeorol.* 12, 206–226. doi: [10.1175/2010jhm1252.1](https://doi.org/10.1175/2010jhm1252.1)
- Kouzana L, Mammou A Ben, Felfoul MS 2009 Seawater intrusion and associated processes: Case of the Korba aquifer (Cap-Bon, Tunisia). *Comptes Rendus Geosci.* 341, 21–35. doi: [10.1016/j.crte.2008.09.008](https://doi.org/10.1016/j.crte.2008.09.008)
- Kundzewicz ZW, Krysanova V, Dankers R, Hirabayashi Y, Kanae S et al. 2017 Differences in flood hazard projections in Europe – their causes and consequences for decision making. *Hydrol. Sci. J.* 62, 1–14. doi: [10.1080/02626667.2016.1241398](https://doi.org/10.1080/02626667.2016.1241398)
- Kurnik B, Kajfež-Bogataj L, Horion S 2015 An assessment of actual evapotranspiration and soil water deficit in agricultural regions in Europe. *Int. J. Climatol.* 35, 2451–2471. doi: [10.1002/joc.4154](https://doi.org/10.1002/joc.4154)
- Labadie JW 2004 Optimal Operation of Multireservoir Systems: State-of-the-Art Review. *J. Water Resour. Plan. Manag.* 130, 93–111. doi: [10.1061/\(asce\)0733-9496\(2004\)130:2\(93\)](https://doi.org/10.1061/(asce)0733-9496(2004)130:2(93))
- Lange MA 2013 Renewable Energy and Water Resources, in *Climate Vulnerability*, eds. Roger A, Pielke S (San Diego, USA: Elsevier), 149–166. doi: [10.1016/b978-0-12-384703-4.00320-8](https://doi.org/10.1016/b978-0-12-384703-4.00320-8)
- Lange MA 2019 Impacts of Climate Change on the Eastern Mediterranean and the Middle East and North Africa Region and the Water–Energy Nexus. *Atmosphere (Basel)*. 10, 455. doi: [10.3390/atmos10080455](https://doi.org/10.3390/atmos10080455)
- Lasca NP, Radulović V, Ristić RJ, Cherkauer DS 1981 Geology, hydrology, climate and bathymetry of Lake Skadar, in *The biota and limnology of Lake Skadar*, eds. Beeton A, Karaman G (Montenegro, Yugoslavia: University Veljko Vlahović, Institute of Biological and Medicine Research Titograd,), 17–38.
- Lasco RD, Delfino RJP, Espaldon MLO 2014 Agroforestry systems: helping smallholders adapt to climate risks while mitigating climate change. *Wiley Interdiscip. Rev. Clim. Chang.* 5, 825–833. doi: [10.1002/wcc.301](https://doi.org/10.1002/wcc.301)
- Lattemann S, Höpner T 2008 Environmental impact and impact assessment of seawater desalination. *Desalination* 220, 1–15. doi: [10.1016/j.desal.2007.03.009](https://doi.org/10.1016/j.desal.2007.03.009)
- Lautze J, Stander E, Drechsel P, Da Silva AK, Keraita B 2014 Global experiences in water reuse. *CGIAR Res. Progr. Water, L. Ecosyst. (WLE). Int. Water Manag. Inst. (IWMI), Colombo, Sri Lanka* 31.
- Leduc C, Pulido-Bosch A, Remini B 2017 Anthropization of groundwater resources in the Mediterranean region: processes and challenges. *Hydrogeol. J.* 25, 1529–1547. doi: [10.1007/s10040-017-1572-6](https://doi.org/10.1007/s10040-017-1572-6)
- Lehner F, Coats S, Stocker TF, Pendergrass AG, Sanderson BM et al. 2017 Projected drought risk in 1.5°C and 2°C warmer climates. *Geophys. Res. Lett.* 44, 7419–7428. doi: [10.1002/2017GL074117](https://doi.org/10.1002/2017GL074117)
- León VM, Moreno-González R, García V, Campillo JA 2017 Impact of flash flood events on the distribution of organic pollutants in surface sediments from a Mediterranean coastal lagoon (Mar Menor, SE Spain). *Environ. Sci. Pollut. Res.* 24, 4284–4300. doi: [10.1007/s11356-015-4628-y](https://doi.org/10.1007/s11356-015-4628-y)
- Lespinas F, Ludwig W, Heussner S 2014 Hydrological and climatic uncertainties associated with modeling the impact of climate change on water resources of small Mediterranean coastal rivers. *J. Hydrol.* 511, 403–422. doi: [10.1016/j.jhydrol.2014.01.033](https://doi.org/10.1016/j.jhydrol.2014.01.033)
- Lezzaik K, Milewski A 2018 A quantitative assessment of groundwater resources in the Middle East and North Africa region. *Hydrogeol. J.* 26, 251–266. doi: [10.1007/s10040-017-1646-5](https://doi.org/10.1007/s10040-017-1646-5)

- Li C, Goswami Y, Stefanakos E 2013 Solar assisted sea water desalination: A review. *Renew. Sustain. Energy Rev.* 19, 136–163. doi: [10.1016/j.rser.2012.04.059](https://doi.org/10.1016/j.rser.2012.04.059)
- Lionello P, Abrantes FG, Congedi L, Dulac F, Gačić M et al. 2012 Introduction: Mediterranean climate-background information, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Elsevier Science). doi: [10.1016/B978-0-12-416042-2.00012-4](https://doi.org/10.1016/B978-0-12-416042-2.00012-4)
- Liu W, Sun F, Lim WH, Zhang J, Wang H et al. 2018 Global drought and severe drought-affected populations in 1.5 and 2°C warmer worlds. *Earth Syst. Dyn.* 9, 267–283. doi: [10.5194/esd-9-267-2018](https://doi.org/10.5194/esd-9-267-2018)
- Llamas MR, Custodio E, de la Hera A, Fornés JM 2015 Groundwater in Spain: increasing role, evolution, present and future. *Environ. Earth Sci.* 73, 2567–2578. doi: [10.1007/s12665-014-4004-0](https://doi.org/10.1007/s12665-014-4004-0)
- Llasat MC, Llasat-Botija M, Petrucci O, Pasqua AA, Rosselló J et al. 2013 Towards a database on societal impact of Mediterranean floods within the framework of the HYMEX project. *Nat. Hazards Earth Syst. Sci.* 13, 1337–1350. doi: [10.5194/nhess-13-1337-2013](https://doi.org/10.5194/nhess-13-1337-2013)
- Llasat MC, Marcos R, Turco M, Gilabert J, Llasat-Botija M 2016 Trends in flash flood events versus convective precipitation in the Mediterranean region: The case of Catalonia. *J. Hydrol.* 541. doi: [10.1016/j.jhydrol.2016.05.040](https://doi.org/10.1016/j.jhydrol.2016.05.040)
- Lobanova A, Koch H, Liersch S, Hattermann FF, Krysanova V 2016 Impacts of changing climate on the hydrology and hydropower production of the Tagus River basin. *Hydrol. Process.* 30, 5039–5052. doi: [10.1002/hyp.10966](https://doi.org/10.1002/hyp.10966)
- López-Moreno JI, Beniston M, García-Ruiz JM 2008 Environmental change and water management in the Pyrenees: Facts and future perspectives for Mediterranean mountains. *Glob. Planet. Change* 61, 300–312. doi: [10.1016/j.gloplacha.2007.10.004](https://doi.org/10.1016/j.gloplacha.2007.10.004)
- Lorenzo M, Campo J, Morales Suárez-Varela M, Picó Y 2019 Occurrence, distribution and behavior of emerging persistent organic pollutants (POPs) in a Mediterranean wetland protected area. *Sci. Total Environ.* 646, 1009–1020. doi: [10.1016/j.scitotenv.2018.07.304](https://doi.org/10.1016/j.scitotenv.2018.07.304)
- Ludwig F, van Slobbe E, Cofino W 2014 Climate change adaptation and Integrated Water Resource Management in the water sector. *J. Hydrol.* 518, 235–242. doi: [10.1016/J.JHYDROL.2013.08.010](https://doi.org/10.1016/J.JHYDROL.2013.08.010)
- Ludwig W, Bouwman AF, Dumont E, Lespinas F 2010 Water and nutrient fluxes from major Mediterranean and Black Sea rivers: Past and future trends and their implications for the basin-scale budgets. *Global Biogeochem. Cycles* 24, n/a-n/a. doi: [10.1029/2009gb003594](https://doi.org/10.1029/2009gb003594)
- Ludwig W, Dumont E, Meybeck M, Heussner S 2009 River discharges of water and nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during past and future decades? *Prog. Oceanogr.* 80, 199–217. doi: [10.1016/j.pocean.2009.02.001](https://doi.org/10.1016/j.pocean.2009.02.001)
- Lutz SR, Mallucci S, Diamantini E, Majone B, Bellin A et al. 2016 Hydroclimatic and water quality trends across three Mediterranean river basins. *Sci. Total Environ.* 571, 1392–1406. doi: [10.1016/j.scitotenv.2016.07.102](https://doi.org/10.1016/j.scitotenv.2016.07.102)
- Majone B, Bovolo CI, Bellin A, Blenkinsop S, Fowler HJ 2012 Modeling the impacts of future climate change on water resources for the Gállego river basin (Spain). *Water Resour. Res.* 48. doi: [10.1029/2011wr010985](https://doi.org/10.1029/2011wr010985)
- Malek Ž, Verburg PH 2018 Adaptation of land management in the Mediterranean under scenarios of irrigation water use and availability. *Mitig. Adapt. Strateg. Glob. Chang.* 23, 821–837. doi: [10.1007/s11027-017-9761-0](https://doi.org/10.1007/s11027-017-9761-0)
- Maliva RG, Missimer TM, Winslow FP, Herrmann R 2011 Aquifer Storage and Recovery of Treated Sewage Effluent in the Middle East. *Arab. J. Sci. Eng.* 36, 63–74. doi: [10.1007/s13369-010-0011-y](https://doi.org/10.1007/s13369-010-0011-y)
- Mangini W, Viglione A, Hall J, Hundecha Y, Ceola S et al. 2018 Detection of trends in magnitude and frequency of flood peaks across Europe. *Hydrol. Sci. J.* 63, 493–512. doi: [10.1080/02626667.2018.1444766](https://doi.org/10.1080/02626667.2018.1444766)
- Marchane A, Trambly Y, Hanich L, Ruelland D, Jarlan L 2017 Climate change impacts on surface water resources in the Rheraya catchment (High Atlas, Morocco). *Hydrol. Sci. J.* 62, 979–995. doi: [10.1080/02626667.2017.1283042](https://doi.org/10.1080/02626667.2017.1283042)
- Marcos R, Llasat MC, Quintana-Seguí P, Turco M 2017 Seasonal predictability of water resources in a Mediterranean freshwater reservoir and assessment of its utility for end-users. *Sci. Total Environ.* 575. doi: [10.1016/j.scitotenv.2016.09.080](https://doi.org/10.1016/j.scitotenv.2016.09.080)
- Margat J 2011 Sources d’approvisionnement en eau durables et non durables au Maghreb. in *4ème Colloque International Ressources en Eau et Développement Durable (CIRED 4)* (Alger).
- Margat J, Blinda M 2005 L’avenir de l’eau en Méditerranée. Problèmes et solutions: nouvelle prospective 2025 du Plan Bleu. in *International Conference on Water, land and Food Security in Arid and Semi-arid Regions*, 44–63. doi: [10.1163/q3\\_SIM\\_00374](https://doi.org/10.1163/q3_SIM_00374)
- Margat J, van der Gun J 2013 *Groundwater around the World*. CRC Press <http://dx.doi.org/10.1201/b13977>
- Mariotti A, Pan Y, Zeng N, Alessandri A 2015 Long-term climate change in the Mediterranean region in the midst of decadal variability. *Clim. Dyn.* 44, 1437–1456. doi: [10.1007/s00382-015-2487-3](https://doi.org/10.1007/s00382-015-2487-3)
- Martín Carrasco FJ, Garrote de Marcos LM, Mediero Orduña L 2007 Gestión de conflictos de compatibilidad de usos en embalses multipropósito. *Tecnol. y ciencias del agua* 22, 81–90.
- Marx A, Kumar R, Thober S, Rakovec O, Wanders N et al. 2018 Climate change alters low flows in Europe

- under global warming of 1.5, 2, and 3 °C. *Hydrol. Earth Syst. Sci.* 22, 1017–1032. doi: [10.5194/hess-22-1017-2018](https://doi.org/10.5194/hess-22-1017-2018)
- Mazi K, Koussis AD, Destouni G 2014 Intensively exploited Mediterranean aquifers: resilience to seawater intrusion and proximity to critical thresholds. *Hydrol. Earth Syst. Sci.* 18, 1663–1677. doi: [10.5194/hess-18-1663-2014](https://doi.org/10.5194/hess-18-1663-2014)
- MED-EUWI 2007 *Mediterranean Groundwater Report - Technical report on groundwater management in the Mediterranean and the Water Framework Directive.*
- Mediero L, Santillán D, Garrote L, Granados A 2014 Detection and attribution of trends in magnitude, frequency and timing of floods in Spain. *J. Hydrol.* 517, 1072–1088. doi: [10.1016/j.jhydrol.2014.06.040](https://doi.org/10.1016/j.jhydrol.2014.06.040)
- Merett S 2003 Virtual Water and Occam's Razor. *Water Int.* 28, 103–105. doi: [10.1080/02508060.2003.9724811](https://doi.org/10.1080/02508060.2003.9724811)
- Merz B, Vorogushyn S, Uhlemann S, Delgado JM, Hundscha Y 2012 More efforts and scientific rigour are needed to attribute trends in flood time series. *Hydrol. Earth Syst. Sci.* 16, 1379–1387. doi: [10.5194/hess-16-1379-2012](https://doi.org/10.5194/hess-16-1379-2012)
- Meza FJ, Montes C, Bravo-Martínez F, Serrano-Ortiz P, Kowalski AS 2018 Soil water content effects on net ecosystem CO<sub>2</sub> exchange and actual evapotranspiration in a Mediterranean semiarid savanna of Central Chile. *Sci. Rep.* 8, 8570. doi: [10.1038/s41598-018-26934-z](https://doi.org/10.1038/s41598-018-26934-z)
- Milano M, Ruelland D, Fernandez S, Dezetter A, Fabre J et al. 2013 Current state of Mediterranean water resources and future trends under climatic and anthropogenic changes. *Hydrol. Sci. J.* 58, 498–518. doi: [10.1080/02626667.2013.774458](https://doi.org/10.1080/02626667.2013.774458)
- Miller JE 2003 Review of Water Resources and Desalination Technologies. doi: [10.2172/809106](https://doi.org/10.2172/809106)
- Milly PCD, Betancourt J, Falkenmark M, Hirsch RM, Kundzewicz ZW et al. 2008 Stationarity Is Dead: Whither Water Management? *Science* (80-. ). 319, 573–574. doi: [10.1126/science.1151915](https://doi.org/10.1126/science.1151915)
- Miralles DG, van den Berg MJ, Gash JH, Parinussa RM, de Jeu RAM et al. 2014 El Niño–La Niña cycle and recent trends in continental evaporation. *Nat. Clim. Chang.* 4, 122–126. doi: [10.1038/nclimate2068](https://doi.org/10.1038/nclimate2068)
- Missimer TM, Maliva RG 2018 Environmental issues in seawater reverse osmosis desalination: Intakes and outfalls. *Desalination* 434, 198–215. doi: [10.1016/j.desal.2017.07.012](https://doi.org/10.1016/j.desal.2017.07.012)
- Mitchell D, Allen MR, Hall JW, Muller B, Rajamani L et al. 2018 The myriad challenges of the Paris Agreement. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376, 20180066. doi: [10.1098/rsta.2018.0066](https://doi.org/10.1098/rsta.2018.0066)
- Molist J, Gómez JM, Sanz J 2011 Water reclamation for industrial reuse in Tarragona. in *International Water Association (IWA) Conference* [Barcelona, Spain].
- Montanari A 2012 Hydrology of the Po River: looking for changing patterns in river discharge. *Hydrol. Earth Syst. Sci.* 16, 3739–3747. doi: [10.5194/hess-16-3739-2012](https://doi.org/10.5194/hess-16-3739-2012)
- Moresi M 2014 Assessment of the life cycle greenhouse gas emissions in the food industry. *Agro Food Ind. Hi-Tech.* 25(3), 53–62.
- Morote Á-F, Hernández M, Rico A-M 2016 Causes of Domestic Water Consumption Trends in the City of Alicante: Exploring the Links between the Housing Bubble, the Types of Housing and the Socio-Economic Factors. *Water* 8, 374. doi: [10.3390/w8090374](https://doi.org/10.3390/w8090374)
- Mourato S, Moreira M, Corte-Real JM 2015 Water Resources Impact Assessment Under Climate Change Scenarios in Mediterranean Watersheds. *Water Resour. Manag.* 29, 2377–2391. doi: [10.1007/s11269-015-0947-5](https://doi.org/10.1007/s11269-015-0947-5)
- MSFD 2008 Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive). <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX-32008L0056&from=EN>
- Naumann G, Alfieri L, Wyser K, Mentaschi L, Betts RA et al. 2018 Global changes in drought conditions under different levels of warming. *Geophys. Res. Lett.* 45, 3285–3296. doi: [10.1002/2017GL076521](https://doi.org/10.1002/2017GL076521)
- Nikolakis D 2008 A statistical study of precipitation in Cyprus. *Hell. J. Geosci.* 43, 67–74.
- Nilsson Å 1988 *Groundwater Dams for Small-Scale Water Supply.* Practical Action Publishing. doi: [10.3362/9781780442297](https://doi.org/10.3362/9781780442297)
- Nogués-Bravo D, Araújo MB, Lasanta T, Moreno JLL 2008 Climate change in Mediterranean mountains during the 21<sup>st</sup> century. *Ambio* 37, 280–285. doi: [10.1579/0044-7447\(2008\)37\[280:ccimmd\]2.0.co;2](https://doi.org/10.1579/0044-7447(2008)37[280:ccimmd]2.0.co;2)
- OECD 2016 *Mitigating Droughts and Floods in Agriculture: Policy Lessons and Approaches.* Paris. doi: [10.1787/9789264246744-en](https://doi.org/10.1787/9789264246744-en)
- Olsson P, Galaz V, Boonstra WJ 2014 Sustainability transformations: a resilience perspective. *Ecol. Soc.* 19. doi: [10.5751/es-06799-190401](https://doi.org/10.5751/es-06799-190401)
- OME 2018 *Mediterranean Energy Perspectives 2018.* Paris.
- Onder H, Yilmaz M 2005 Underground Dams. A Tool of Sustainable Development and Management of Groundwater Resources. *Eur. Water* 11, 35–45.
- Orlowsky B, Seneviratne SI 2013 Elusive drought: uncertainty in observed trends and short- and long-term CMIP5 projections. *Hydrol. Earth Syst. Sci.* 17, 1765–1781. doi: [10.5194/hess-17-1765-2013](https://doi.org/10.5194/hess-17-1765-2013)
- Papadimitriou L V., Koutroulis AG, Grillakis MG, Tsanis IK 2016 High-end climate change impact on European runoff and low flows-exploring the effects of forcing biases. *Hydrol. Earth Syst. Sci.* 20, 1785–1808. doi: [10.5194/hess-20-1785-2016](https://doi.org/10.5194/hess-20-1785-2016)
- Papadimitriou L V., Koutroulis AG, Grillakis MG, Tsanis

- IK 2017 The effect of GCM biases on global runoff simulations of a land surface model. *Hydrol. Earth Syst. Sci.* 21, 4379–4401. doi: [10.5194/hess-21-4379-2017](https://doi.org/10.5194/hess-21-4379-2017)
- Papanicolas C 2010 Research and development study for a concentrated solar power-desalination of sea water (CSP-DSW) project.
- Papanicolas CN, Bonanos AM, Georgiou MC, Guillen E, Jarraud N et al. 2016 CSP cogeneration of electricity and desalinated water at the Pentakomo field facility. doi: [10.1063/1.4949196](https://doi.org/10.1063/1.4949196)
- PERSEUS – UNEP/MAP 2015 *Atlas of Riverine Inputs to the Mediterranean Sea*. <http://www.perseus-net.eu/assets/media/PDF/5567.pdf>
- Piayda A, Dubbert M, Rebmann C, Kolle O, Costa e Silva F et al. 2014 Drought impact on carbon and water cycling in a Mediterranean *Quercus suber* L. woodland during the extreme drought event in 2012. *Biogeosciences* 11, 7159–7178. doi: [10.5194/bg-11-7159-2014](https://doi.org/10.5194/bg-11-7159-2014)
- Piras M, Mascaro G, Deidda R, Vivoni ER 2014 Quantification of hydrologic impacts of climate change in a Mediterranean basin in Sardinia, Italy, through high-resolution simulations. *Hydrol. Earth Syst. Sci.* 18, 5201–5217. doi: [10.5194/hess-18-5201-2014](https://doi.org/10.5194/hess-18-5201-2014)
- Pittock J, Meng J, Geiger M 2009 Interbasin water transfers and water scarcity in a changing world—a solution or a pipedream. Frankfurt am Main, Germany.): WWF Germany.
- Plan Bleu 2011 Adapting to climate change in the water sector in the Mediterranean: situation and prospects. Valbonne OneSearch.
- Polissar L, Severson RK, Boatman ES 1984 A Case-Control Study of Asbestos in Drinking Water And Cancer Risk. *Am. J. Epidemiol.* 119, 456–471. doi: [10.1093/oxfordjournals.aje.a113763](https://doi.org/10.1093/oxfordjournals.aje.a113763)
- Pozo K, Palmeri M, Palmeri V, Estellano VH, Mulder MD et al. 2016 Assessing persistent organic pollutants (POPS) in the sicily island atmosphere, mediterranean, using PUF disk passive air samplers. *Environ. Sci. Pollut. Res.* 23, 20796–20804. doi: [10.1007/s11356-016-7131-1](https://doi.org/10.1007/s11356-016-7131-1)
- Prudhomme C, Giuntoli I, Robinson EL, Clark DB, Arnell NW et al. 2014 Hydrological droughts in the 21<sup>st</sup> century, hotspots and uncertainties from a global multimodel ensemble experiment. *Proc. Natl. Acad. Sci. U. S. A.* 111, 3262–7. doi: [10.1073/pnas.1222473110](https://doi.org/10.1073/pnas.1222473110)
- Pulido-Velazquez D, Collados-Lara A-J, Alcalá FJ 2018a Assessing impacts of future potential climate change scenarios on aquifer recharge in continental Spain. *J. Hydrol.* 567, 803–819. doi: [10.1016/j.jhydrol.2017.10.077](https://doi.org/10.1016/j.jhydrol.2017.10.077)
- Pulido-Velazquez D, García-Aróstegui JL, Molina J-L, Pulido-Velazquez M 2015 Assessment of future groundwater recharge in semi-arid regions under climate change scenarios (Serral-Salinas aquifer, SE Spain). Could increased rainfall variability increase the recharge rate? *Hydrol. Process.* 29, 828–844. doi: [10.1002/hyp.10191](https://doi.org/10.1002/hyp.10191)
- Pulido-Velazquez D, Garrote L, Andreu J, Martin-Carrasco F-J, Iglesias A 2011 A methodology to diagnose the effect of climate change and to identify adaptive strategies to reduce its impacts in conjunctive-use systems at basin scale. *J. Hydrol.* 405, 110–122. doi: [10.1016/j.jhydrol.2011.05.014](https://doi.org/10.1016/j.jhydrol.2011.05.014)
- Pulido-Velazquez D, Renau-Pruñonosa A, Llopis-Albert C, Morell I, Collados-Lara A-J et al. 2018b Integrated assessment of future potential global change scenarios and their hydrological impacts in coastal aquifers – a new tool to analyse management alternatives in the Plana Oropesa-Torreblanca aquifer. *Hydrol. Earth Syst. Sci.* 22, 3053–3074. doi: [10.5194/hess-22-3053-2018](https://doi.org/10.5194/hess-22-3053-2018)
- Qiblawey HM, Banat F 2008 Solar thermal desalination technologies. *Desalination* 220, 633–644. doi: [10.1016/j.desal.2007.01.059](https://doi.org/10.1016/j.desal.2007.01.059)
- Quintana-Seguí P, Habets F, Martin E 2011 Comparison of past and future Mediterranean high and low extremes of precipitation and river flow projected using different statistical downscaling methods. *Nat. Hazards Earth Syst. Sci.* 11, 1411–1432. doi: [10.5194/nhess-11-1411-2011](https://doi.org/10.5194/nhess-11-1411-2011)
- Quintana-Seguí P, Martin E, Sánchez E, Zribi M, Vennetier M et al. 2016 Drought: Observed trends, future projections, in *The Mediterranean Region under Climate Change. A Scientific Update*, eds. Thiébaud S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 123–132.
- Racz AJ, Fisher AT, Schmidt CM, Lockwood BS, Huertos ML 2012 Spatial and Temporal Infiltration Dynamics During Managed Aquifer Recharge. *Groundwater* 50, 562–570. doi: [10.1111/j.1745-6584.2011.00875.x](https://doi.org/10.1111/j.1745-6584.2011.00875.x)
- Rana G, Katerji N 2000 Measurement and estimation of actual evapotranspiration in the field under Mediterranean climate: a review. *Eur. J. Agron.* 13, 125–153. doi: [10.1016/s1161-0301\(00\)00070-8](https://doi.org/10.1016/s1161-0301(00)00070-8)
- Renard B, Lang M, Bois P, Dupeyrat A, Mestre O et al. 2008 Regional methods for trend detection: Assessing field significance and regional consistency. *Water Resour. Res.* 44. doi: [10.1029/2007wr006268](https://doi.org/10.1029/2007wr006268)
- Requier-Desjardins M 2010 Impacts des changements climatiques sur l'agriculture au Maroc et en Tunisie et priorités d'adaptation. Notes d'Analyse du CI-HEAM (56).
- Rico-Amoros AM, Olcina-Cantos J, Sauri D 2009 Tourist land use patterns and water demand: Evidence from the Western Mediterranean. *Land use policy* 26, 493–501. doi: [10.1016/j.landusepol.2008.07.002](https://doi.org/10.1016/j.landusepol.2008.07.002)
- Rico-Amoros AM, Sauri D, Olcina-Cantos J, Vera-Rebollo JF 2013 Beyond Megaprojects? Water Alternatives for Mass Tourism in Coastal Mediterranean Spain. *Water Resour. Manag.* 27, 553–565. doi: [10.1007/s11269-012-0201-3](https://doi.org/10.1007/s11269-012-0201-3)

- Rico A, Olcina J, Baños C, García X, Sauri D 2020 Declining water consumption in the hotel industry of mass tourism resorts: contrasting evidence for Benidorm, Spain. *Curr. Issues Tour.* 23, 770–783. doi: [10.1080/13683500.2019.1589431](https://doi.org/10.1080/13683500.2019.1589431)
- Rodell M, Famiglietti JS, Wiese DN, Reager JT, Beaudoin HK et al. 2018 Emerging trends in global freshwater availability. *Nature* 557, 651–659. doi: [10.1038/s41586-018-0123-1](https://doi.org/10.1038/s41586-018-0123-1)
- Rodríguez-Díaz JA, Pérez-Urrestarazu L, Camacho-Poyato E, Montesinos P 2011 The paradox of irrigation scheme modernization: more efficient water use linked to higher energy demand. *Spanish J. Agric. Res.* 4, 1000–1008. doi: [10.5424/sjar/20110904-492-10](https://doi.org/10.5424/sjar/20110904-492-10)
- Rojas R, Feyen L, Bianchi A, Dosio A 2012 Assessment of future flood hazard in Europe using a large ensemble of bias-corrected regional climate simulations. *JGR Atmos.* 117, n/a-n/a. doi: [10.1029/2012jd017461](https://doi.org/10.1029/2012jd017461)
- Roudier P, Andersson JCM, Donnelly C, Feyen L, Greuell W et al. 2016 Projections of future floods and hydrological droughts in Europe under a +2°C global warming. *Clim. Change* 135, 341–355. doi: [10.1007/s10584-015-1570-4](https://doi.org/10.1007/s10584-015-1570-4)
- Ruelland D, Hublart P, Tramblay Y 2015 Assessing uncertainties in climate change impacts on runoff in Western Mediterranean basins. *Proc. Int. Assoc. Hydrol. Sci.* 371, 75–81. doi: [10.5194/piahs-371-75-2015](https://doi.org/10.5194/piahs-371-75-2015)
- Ruosteenoja K, Markkanen T, Venäläinen A, Räisänen P, Peltola H 2018 Seasonal soil moisture and drought occurrence in Europe in CMIP5 projections for the 21<sup>st</sup> century. *Clim. Dyn.* 50, 1177–1192. doi: [10.1007/s00382-017-3671-4](https://doi.org/10.1007/s00382-017-3671-4)
- Saghir J, Schiffler M, Woldu M 2000 Urban water and sanitation in the Middle East and North Africa Region: The way forward.
- Samaniego L, Thober S, Kumar R, Wanders N, Rakovec O et al. 2018 Anthropogenic warming exacerbates European soil moisture droughts. *Nat. Clim. Chang.* 8, 421–426. doi: [10.1038/s41558-018-0138-5](https://doi.org/10.1038/s41558-018-0138-5)
- Sanchis-Ibor C, García-Mollá M, Avellà-Reus L 2017 Effects of drip irrigation promotion policies on water use and irrigation costs in Valencia, Spain. *Water Policy* 19, 165–180. doi: [10.2166/wp.2016.025](https://doi.org/10.2166/wp.2016.025)
- Savé R, de Herralde F, Aranda X, Pla E, Pascual D et al. 2012 Potential changes in irrigation requirements and phenology of maize, apple trees and alfalfa under global change conditions in Fluvia watershed during XXI<sup>st</sup> century: Results from a modeling approximation to watershed-level water balance. *Agric. Water Manag.* 114, 78–87. doi: [10.1016/j.agwat.2012.07.006](https://doi.org/10.1016/j.agwat.2012.07.006)
- Schewe J, Heinke J, Gerten D, Haddeland I, Arnell NW et al. 2014 Multimodel assessment of water scarcity under climate change. *Proc. Natl. Acad. Sci. U. S. A.* 111, 3245–3250. doi: [10.1073/pnas.1222460110](https://doi.org/10.1073/pnas.1222460110)
- Schleussner C-F, Lissner TK, Fischer EM, Wohland J, Perrette M et al. 2016 Differential climate impacts for policy-relevant limits to global warming: The case of 1.5 °C and 2 °C. *Earth Syst. Dyn.* 7, 327–351. doi: [10.5194/esd-7-327-2016](https://doi.org/10.5194/esd-7-327-2016)
- Schneider C, Laizé CLR, Acreman MC, Flörke M 2013 How will climate change modify river flow regimes in Europe? *Hydrol. Earth Syst. Sci.* 17, 325–339. doi: [10.5194/hess-17-325-2013](https://doi.org/10.5194/hess-17-325-2013)
- Schnitter N 1978 Römische Talsperren. *Antike Welt* 8, 25–32.
- Schnitter N 1987 Die Entwicklungsgeschichte der Pfeilerstaumauer, in *Historische Talsperren*, ed. , Garbrecht G (ed. . (Stuttgart, Germany), 57–74.
- Schulz C, Adams WM 2019 Debating dams: The World Commission on Dams 20 years on. *Wiley Interdiscip. Rev. Water* 6. doi: [10.1002/wat2.1369](https://doi.org/10.1002/wat2.1369)
- Scudder TT 2006 *The future of large dams: dealing with social, environmental, institutional and political costs*. Routledge doi: [10.4324/9781849773904](https://doi.org/10.4324/9781849773904)
- Sebri M 2017 Bridging the Maghreb's water gap: from rationalizing the virtual water trade to enhancing the renewable energy desalination. *Environ. Dev. Sustain.* 19, 1673–1684. doi: [10.1007/s10668-016-9820-9](https://doi.org/10.1007/s10668-016-9820-9)
- Seif-Ennasr M, Zaaboul R, Hirich A, Caroletti GN, Bouchaou L et al. 2016 Climate change and adaptive water management measures in Chtouka Aït Baha region (Morocco). *Sci. Total Environ.* 573, 862–875. doi: [10.1016/j.scitotenv.2016.08.170](https://doi.org/10.1016/j.scitotenv.2016.08.170)
- Semiati R 2008 Energy Issues in Desalination Processes. *Environ. Sci. Technol.* 42, 8193–8201. doi: [10.1021/es801330u](https://doi.org/10.1021/es801330u)
- Senatore A, Mendicino G, Smiatek G, Kunstmann H 2011 Regional climate change projections and hydrological impact analysis for a Mediterranean basin in Southern Italy. *J. Hydrol.* 399, 70–92. doi: [10.1016/j.jhydrol.2010.12.035](https://doi.org/10.1016/j.jhydrol.2010.12.035)
- Seneviratne SI, Nicholls N, Easterling DR, Goodess CM, Kanae S et al. 2012 Changes in Climate Extremes and their Impacts on the Natural Physical Environment, in *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, eds. Field CB, Barros V, Stocker TF, Qin D, Dokken DJ et al. (Cambridge, United Kingdom and New York, NY, USA: A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC). Cambridge University Press), 109–230. doi: [10.1017/cbo9781139177245.006](https://doi.org/10.1017/cbo9781139177245.006)
- Serpa D, Nunes JP, Keizer JJ, Abrantes N 2017 Impacts of climate and land use changes on the water quality of a small Mediterranean catchment with intensive viticulture. *Environ. Pollut.* 224, 454–465. doi: [10.1016/j.envpol.2017.02.026](https://doi.org/10.1016/j.envpol.2017.02.026)
- Serra-Llobet A, Conrad E, Schaefer K 2016 Governing for Integrated Water and Flood Risk Management:



- Comparing Top-Down and Bottom-Up Approaches in Spain and California. *Water* 8, 445. doi: [10.3390/w8100445](https://doi.org/10.3390/w8100445)
- Shah T 2014 Groundwater governance and irrigated agriculture. [https://www.gwp.org/globalassets/global/toolbox/publications/background-papers/gwp\\_tec\\_19\\_web.pdf](https://www.gwp.org/globalassets/global/toolbox/publications/background-papers/gwp_tec_19_web.pdf)
- Shannon MA, Bohn PW, Elimelech M, Georgiadis JG, Mariñas BJ et al. 2008 Science and technology for water purification in the coming decades. *Nature* 452, 301–310. doi: [10.1038/nature06599](https://doi.org/10.1038/nature06599)
- Sheffield J, Wood EF 2008 Global Trends and Variability in Soil Moisture and Drought Characteristics, 1950–2000, from Observation-Driven Simulations of the Terrestrial Hydrologic Cycle. *J. Clim.* 21, 432–458. doi: [10.1175/2007JCLI1822.1](https://doi.org/10.1175/2007JCLI1822.1)
- Shiklomanov IA 1999 Water transfer as one of the most important ways to eliminate water resources deficits and to solve water management problems. in *International Workshop on Interbasin Water Transfer*, ed. UNESCO (Paris), 203–210.
- Smiatek G, Kunstmann H, Heckl A 2014 High-Resolution Climate Change Impact Analysis on Expected Future Water Availability in the Upper Jordan Catchment and the Middle East. *J. Hydrometeorol.* 15, 1517–1531. doi: [10.1175/jhm-d-13-0153.1](https://doi.org/10.1175/jhm-d-13-0153.1)
- Sordo-Ward Á, Garrote L, Bejarano MD, Castillo LG 2013 Extreme flood abatement in large dams with gate-controlled spillways. *J. Hydrol.* 498, 113–123. doi: [10.1016/j.jhydrol.2013.06.010](https://doi.org/10.1016/j.jhydrol.2013.06.010)
- Sordo-Ward Á, Garrote L, Martín-Carrasco F, Bejarano MD 2012 Extreme flood abatement in large dams with fixed-crest spillways. *J. Hydrol.* 466–467, 60–72. doi: [10.1016/j.jhydrol.2012.08.009](https://doi.org/10.1016/j.jhydrol.2012.08.009)
- Stahl K, Hisdal H, Hannaford J, Tallaksen LM, van Lanen HAJ et al. 2010 Streamflow trends in Europe: evidence from a dataset of near-natural catchments. *Hydrol. Earth Syst. Sci.* 14, 2367–2382. doi: [10.5194/hess-14-2367-2010](https://doi.org/10.5194/hess-14-2367-2010)
- Steffen W, Rockström J, Richardson K, Lenton TM, Folke C et al. 2018 Trajectories of the Earth System in the Anthropocene. *Proc. Natl. Acad. Sci. U. S. A.* 115, 8252–8259. doi: [10.1073/pnas.1810141115](https://doi.org/10.1073/pnas.1810141115)
- Steinel A, Schelkes K, Subah A, Himmelsbach T 2016 Spatial multi-criteria analysis for selecting potential sites for aquifer recharge via harvesting and infiltration of surface runoff in north Jordan. *Hydrogeol. J.* 24, 1753–1774. doi: [10.1007/s10040-016-1427-6](https://doi.org/10.1007/s10040-016-1427-6)
- Stocker TF, Qin D, Plattner G-K, Alexander L V., Allen SK et al. 2013 Technical Summary, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 33–115. doi: [10.1017/CBO9781107415324.005](https://doi.org/10.1017/CBO9781107415324.005)
- Struglia MV, Mariotti A, Filograsso A 2004 River Discharge into the Mediterranean Sea: Climatology and Aspects of the Observed Variability. *J. Clim.* 17, 4740–4751. doi: [10.1175/jcli-3225.1](https://doi.org/10.1175/jcli-3225.1)
- Su L, Miao C, Kong D, Duan Q, Lei X et al. 2018 Long-term trends in global river flow and the causal relationships between river flow and ocean signals. *J. Hydrol.* 563, 818–833. doi: [10.1016/j.jhydrol.2018.06.058](https://doi.org/10.1016/j.jhydrol.2018.06.058)
- Suárez-Almiñana S, Pedro-Monzonís M, Paredes-Arquiola J, Andreu J, Solera A 2017 Linking Pan-European data to the local scale for decision making for global change and water scarcity within water resources planning and management. *Sci. Total Environ.* 603–604, 126–139. doi: [10.1016/j.scitotenv.2017.05.259](https://doi.org/10.1016/j.scitotenv.2017.05.259)
- Subramani A, Jacangelo JG 2015 Emerging desalination technologies for water treatment: A critical review. *Water Res.* 75, 164–187. doi: [10.1016/j.watres.2015.02.032](https://doi.org/10.1016/j.watres.2015.02.032)
- Swiss Re 2015 Sigma 02/2015: Natural catastrophes and man-made disasters in 2014: convective and winter storms generate most losses. [www.swiss-re.com/institute/research/sigma-research/sigma-2015-02.html](http://www.swiss-re.com/institute/research/sigma-research/sigma-2015-02.html) [Accessed August 9, 2019]
- Tanny J 2013 Microclimate and evapotranspiration of crops covered by agricultural screens: A review. *Biosyst. Eng.* 114, 26–43. doi: [10.1016/j.biosystemseng.2012.10.008](https://doi.org/10.1016/j.biosystemseng.2012.10.008)
- Thober S, Kumar R, Wanders N, Marx A, Pan M et al. 2018 Multi-model ensemble projections of European river floods and high flows at 1.5, 2, and 3 degrees global warming. *Environ. Res. Lett.* 13, 14003. doi: [10.1088/1748-9326/aa9e35](https://doi.org/10.1088/1748-9326/aa9e35)
- Tortajada C, Altinbilek D, Biswas AK 2012 *Impacts of Large Dams: A Global Assessment*. Springer, doi: [10.1007/978-3-642-23571-9](https://doi.org/10.1007/978-3-642-23571-9)
- Toth E, Bragalli C, Neri M 2018 Assessing the significance of tourism and climate on residential water demand: Panel-data analysis and non-linear modelling of monthly water consumptions. *Environ. Model. Softw.* 103, 52–61. doi: [10.1016/j.envsoft.2018.01.011](https://doi.org/10.1016/j.envsoft.2018.01.011)
- Tramblay Y, El Adlouni S, Servat E 2013a Trends and variability in extreme precipitation indices over Maghreb countries. *Nat. Hazards Earth Syst. Sci.* 13, 3235–3248. doi: [10.5194/nhess-13-3235-2013](https://doi.org/10.5194/nhess-13-3235-2013)
- Tramblay Y, Jarlan L, Hanich L, Somot S 2018 Future Scenarios of Surface Water Resources Availability in North African Dams. *Water Resour. Manag.* 32, 1291–1306. doi: [10.1007/s11269-017-1870-8](https://doi.org/10.1007/s11269-017-1870-8)
- Tramblay Y, Ruelland D, Hanich L, Dakhlouli H 2016 Hydrological impacts of climate change in North African countries, in *The Mediterranean Region under Climate Change. A Scientific Update*, eds. Thiébault

- S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 295–302.  
doi: [10.4000/books.irdeditions.23496](https://doi.org/10.4000/books.irdeditions.23496)
- Trambly Y, Ruelland D, Somot S, Bouaicha R, Servat E 2013b High-resolution Med-CORDEX regional climate model simulations for hydrological impact studies: a first evaluation of the ALADIN-Climate model in Morocco. *Hydrol. Earth Syst. Sci.* 17, 3721–3739. doi: [10.5194/hess-17-3721-2013](https://doi.org/10.5194/hess-17-3721-2013)
- Trichakis I, Burek P, de Roo A, Pistocchi A 2017 Towards a Pan-European Integrated Groundwater and Surface Water Model: Development and Applications. *Environ. Process.* 4, 81–93.  
doi: [10.1007/s40710-017-0216-0](https://doi.org/10.1007/s40710-017-0216-0)
- Trieb F, Müller-Steinhagen H 2008 Concentrating solar power for seawater desalination in the Middle East and North Africa. *Desalination* 220, 165–183.  
doi: [10.1016/j.desal.2007.01.030](https://doi.org/10.1016/j.desal.2007.01.030)
- Tsanis IK, Koutroulis AG, Daliakopoulos IN, Jacob D 2011 Severe climate-induced water shortage and extremes in Crete. *Clim. Change* 106, 667–677.  
doi: [10.1007/s10584-011-0048-2](https://doi.org/10.1007/s10584-011-0048-2)
- TSKB Ekonomik Araştırmalar 2020 Aylık Enerji Bülteni.
- Turco M, Llasat MC 2011 Trends in indices of daily precipitation extremes in Catalonia (NE Spain), 1951–2003. *Nat. Hazards Earth Syst. Sci.* 11, 3213–3226.  
doi: [10.5194/nhess-11-3213-2011](https://doi.org/10.5194/nhess-11-3213-2011)
- Turco M, Llasat MC, Herrera S, Gutiérrez JM 2017 Bias correction and downscaling of future RCM precipitation projections using a MOS-Analog technique. *JGR Atmos.* 122. doi: [10.1002/2016JD025724](https://doi.org/10.1002/2016JD025724)
- UN-Water 2015 Compendium of Water Quality Regulatory Frameworks: Which Water for Which Use?
- UN-Water 2018 Sustainable Development Goal 6. Synthesis Report on Water and Sanitation. Geneva, Switzerland.
- UN 2009 Guidance on Water and Adaptation to Climate Change. Geneva.
- UN 2015 Transforming our world: the 2030 Agenda for Sustainable Development. New York [https://www.un.org/ga/search/view\\_doc.asp?symbol=A/RES/70/1&Lang=E](https://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E)
- UN 2019 World Population Prospects 2019, Online Edition. <https://population.un.org/wpp/DefinitionOfProjectionVariants> [Accessed August 1, 2019]
- UNEP/MAP 2016 Mediterranean Strategy for Sustainable Development 2016–2025. Valbonne.
- UNEP/MAP, Plan Bleu 2020 State of the Environment and Development in the Mediterranean. Nairobi
- UNFCCC 2016 Report of the Conference of the Parties on its twenty-first session, held in Paris from 30 November to 13 December 2015.
- UNISDR 2015 *Global Assessment Report on Disaster Risk Reduction (GAR)*. United Nations.
- UNTWO 2018 UNTWO Tourist Highlights 2017. *World Tour. Organ.* <https://www.e-unwto.org/doi/pdf/10.18111/9789284419876> [Accessed December 11, 2019]
- van Vliet MTH, Donnelly C, Strömbäck L, Capell R, Ludwig F 2015 European scale climate information services for water use sectors. *J. Hydrol.* 528, 503–513. doi: [10.1016/J.JHYDROL.2015.06.060](https://doi.org/10.1016/J.JHYDROL.2015.06.060)
- Velasco J, Lloret J, Millan A, Marin A, Barahona J et al. 2006 Nutrient and particulate inputs into the Mar Menor lagoon (Se Spain) from an intensive agricultural watershed. *Water. Air. Soil Pollut.* 176, 37–56.  
doi: [10.1007/s11270-006-2859-8](https://doi.org/10.1007/s11270-006-2859-8)
- Verdier J, Viollet P-L 2015 Water Tensions in Europe and in the Mediterranean: water crisis by 2050? *La Houille Blanche*, 102–107.  
doi: [10.1051/lhb/20150075](https://doi.org/10.1051/lhb/20150075)
- Vijgen J, Abhilash PC, Li YF, Lal R, Forter M et al. 2011 Hexachlorocyclohexane (HCH) as new Stockholm Convention POPs—a global perspective on the management of Lindane and its waste isomers. *Environ. Sci. Pollut. Res.* 18, 152–162.  
doi: [10.1007/s11356-010-0417-9](https://doi.org/10.1007/s11356-010-0417-9)
- Viola F, Daly E, Vico G, Cannarozzo M, Porporato A 2008 Transient soil-moisture dynamics and climate change in Mediterranean ecosystems. *Water Resour. Res.* 44. doi: [10.1029/2007wr006371](https://doi.org/10.1029/2007wr006371)
- Viola F, Francipane A, Caracciolo D, Pumo D, La Loggia G et al. 2016 Co-evolution of hydrological components under climate change scenarios in the Mediterranean area. *Sci. Total Environ.* 544, 515–524.  
doi: [10.1016/J.SCITOTENV.2015.12.004](https://doi.org/10.1016/J.SCITOTENV.2015.12.004)
- Voss KA, Famiglietti JS, Lo M, Linage C, Rodell M et al. 2013 Groundwater depletion in the Middle East from GRACE with implications for transboundary water management in the Tigris-Euphrates-Western Iran region. *Water Resour. Res.* 49, 904–914.  
doi: [10.1002/wrcr.20078](https://doi.org/10.1002/wrcr.20078)
- Wada Y, van Beek LPH, Bierkens MFP 2012 Nonsustainable groundwater sustaining irrigation: A global assessment. *Water Resour. Res.* 48.  
doi: [10.1029/2011wr010562](https://doi.org/10.1029/2011wr010562)
- Waha K, Krümmenauer L, Adams S, Aich V, Baarsch F et al. 2017 Climate change impacts in the Middle East and Northern Africa (MENA) region and their implications for vulnerable population groups. *Reg. Environ. Chang.* 17, 1623–1638.  
doi: [10.1007/s10113-017-1144-2](https://doi.org/10.1007/s10113-017-1144-2)
- Wang K, Dickinson RE, Wild M, Liang S 2010 Evidence for decadal variation in global terrestrial evapotranspiration between 1982 and 2002: 2. Results. *JGR Atmos.* 115. doi: [10.1029/2010jd013847](https://doi.org/10.1029/2010jd013847)
- Ward FA, Pulido-Velazquez M 2008 Water conservation in irrigation can increase water use. *Proc. Natl. Acad. Sci. U. S. A.* 105, 18215–18220.  
doi: [10.1073/pnas.0805554105](https://doi.org/10.1073/pnas.0805554105)
- Water Development Department Cyprus 2000 Southern Conveyor Project. Nicosia, Cyprus.
- Water Development Department Cyprus 2009 Dams of Cyprus. Nicosia, Cyprus.
- WFD 2000 Directive 2000/60/EC of the European Parlia-

- ment and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy (OJ L 327 22.12.2000). *Doc. Eur. Community Environ. Law*, 879–969. doi: [10.1017/cbo9780511610851.056](https://doi.org/10.1017/cbo9780511610851.056)
- WFD 2006 2006/118/EC Groundwater Daughter Directive to WFD. <https://rod.eionet.europa.eu/instruments/625> [Accessed September 3, 2019]
- WHO 2003 Asbestos in drinking-water. Background document for development of WHO guidelines for drinking-water quality.
- Wilcox BP, Seyfried MS, Breshears DD 2003 The water balance on rangelands, in *Encyclopedia of Water Science*, eds. Stewart BA, Howell TA (New York: Marcel Dekker), 791–794.
- Wilhelm B, Arnaud F, Sabatier P, Crouzet C, Brisset E et al. 2012 1400 years of extreme precipitation patterns over the Mediterranean French Alps and possible forcing mechanisms. *Quat. Res.* 78, 1–12. doi: [10.1016/j.yqres.2012.03.003](https://doi.org/10.1016/j.yqres.2012.03.003)
- Wilk J, Wittgren HBB 2009 Adapting Water Management to Climate Change. Swedish Water House
- Wimmer F, Audsley E, Malsy M, Savin C, Dunford R et al. 2014 Modelling the effects of cross-sectoral water allocation schemes in Europe. *Clim. Change* 128, 229–244. doi: [10.1007/s10584-014-1161-9](https://doi.org/10.1007/s10584-014-1161-9)
- Wolf AT, Natharius JA, Danielson JJ, Ward BS, Pender JK 1999 International River Basins of the World. *Int. J. Water Resour. Dev.* 15, 387–427. doi: [10.1080/07900629948682](https://doi.org/10.1080/07900629948682)
- World Bank 2016 High and Dry: Climate Change, Water, and the Economy. Washington, DC.
- World Bank 2018 Beyond Scarcity: Water Security in the Middle East and North Africa. Washington, DC. doi: [10.1596/978-1-4648-1144-9](https://doi.org/10.1596/978-1-4648-1144-9)
- World Bank 2019 People using at least basic sanitation services (% of population). <https://donnees.banquemondiale.org/indicateur/SH.STA.BASS.ZS> [Accessed June 5, 2019]
- World Commission on Dams 2000 Dams and Development. A New Framework for Decision-Making. London and Sterling, VA, Ltd, E.P. doi: [10.4324/9781315541518](https://doi.org/10.4324/9781315541518)
- WWC 2009 World Water Forum Mediterranean Session Regional Document. Istanbul.
- Wyser K, Strandberg G, Caesar J, Gohar L 2016 Documentation of changes in climate variability and extremes simulated by the HELIX AGCMs at the 3 SWLs and comparison in equivalent SST/SIC low-resolution CMIP5 projections.
- Younos T, Tulou KE 2005 Overview of Desalination Techniques. *J. Contemp. Water Res. Educ.* 132, 3–10. doi: [10.1111/j.1936-704x.2005.mp132001002.x](https://doi.org/10.1111/j.1936-704x.2005.mp132001002.x)
- Zammouri M, Siegfried T, El-Fahem T, Kriâa S, Kinzelbach W 2007 Salinization of groundwater in the Nefzawa oases region, Tunisia: results of a regional-scale hydrogeologic approach. *Hydrogeol. J.* 15, 1357–1375. doi: [10.1007/s10040-007-0185-x](https://doi.org/10.1007/s10040-007-0185-x)
- Zhang Y, Kong D, Gan R, Chiew FHS, McVicar TR et al. 2019 Coupled estimation of 500 m and 8-day resolution global evapotranspiration and gross primary production in 2002–2017. *Remote Sens. Environ.* 222, 165–182. doi: [10.1016/j.rse.2018.12.031](https://doi.org/10.1016/j.rse.2018.12.031)
- Zhang Y, Peña-Arancibia JL, McVicar TR, Chiew FHS, Vaze J et al. 2016 Multi-decadal trends in global terrestrial evapotranspiration and its components. *Sci. Rep.* 6, 19124. doi: [10.1038/srep19124](https://doi.org/10.1038/srep19124)

## Information about authors

### Coordinating Lead Authors

Marianela Fader:

*International Centre for Water Resources and Global Change (UNESCO), Federal Institute of Hydrology, Koblenz, Germany*

Carlo Giupponi:

*Ca' Foscari University and Venice International University, Venice, Italy*

### Lead Authors

Selmin Burak:

*Istanbul University-Institute of Marine Sciences and Management (IU-IMSM), Istanbul, Turkey*

Hamouda Dakhlaoui:

*LMHE, Ecole Nationale d'Ingénieurs de Tunis, Université Tunis El Manar, Tunis Le Belvédère/ Ecole Nationale d'Architecture et d'Urbanisme, Université de Carthage, Sidi Bou Said, Tunisia*

Aristeidis Koutroulis:

*Technical University of Crete, Chania, Greece*

Manfred A. Lange:

*The Cyprus Institute, Nicosia, Cyprus*

Maria Carmen Llasat:

*University of Barcelona, Barcelona, Spain*

David Pulido-Velazquez:

*Spanish Geological Survey, Granada, Spain*

Alberto Sanz-Cobeña:

*CEIGRAM, Universidad Politécnica de Madrid (UPM), Madrid, Spain*

### Contributing Authors

Manolis Grillakis:

*Institute of Mediterranean Studies (Institute of Technology & Research – Hellas), Rethymno, Crete, Greece*

Rachid Mrabet :

*National Institute for Agricultural Research (INRA Morocco), Rabat, Morocco*

David Sauri Pujol:

*Universitat Autònoma de Barcelona, Barcelona, Spain*

Robert Savé:

*Institute of Agrifood Research and Technology (IRTA), Caldes de Montbui (Barcelona), Spain*

Mladen Todorović:

*Centre International de Hautes Études Agronomiques Méditerranéennes (CIHEAM), Mediterranean Agronomic Institute of Bari (IAMB), Bari, Italy*

Yves Trambly:

*HydroSciences Montpellier (University Montpellier, CNRS, IRD), Montpellier, France*

Veronika Zwirgmaier:

*Ludwig Maximilian University of Munich, Munich, Germany*



# RESOURCES 2-FOOD

**Coordinating Lead Authors:**

Rachid Mrabet (Morocco), Robert Savé (Spain)


**Lead Authors:**

Andrea Toreti (Italy), Nuno Caiola (Spain), Mouad Chentouf (Morocco), Maria Carmen Llasat (Spain), Assem Abdelmonem Ahmed Mohamed (Egypt), Fabio G. Santeramo (Italy), Alberto Sanz-Cobena (Spain), Athanassios Tsikliras (Greece)

**Contributing Authors:**

Eduardo Aguilera (Spain), Luis Asin (Spain), Andrej Cegljar (Italy), Alejandro de Blas (Spain), Donna Dimarchopou-lou (Greece), Elena Georgopoulou (Greece), Luis Lassaletta (Spain), Androniki Pardalou (Greece), Giuseppe Scarcella (Italy), Marco Turco (Spain), Matteo Zampieri (Italy)

*This chapter should be cited as: Mrabet R, Savé R, Toreti A, Caiola N, Chentouf M, Llasat MC, Mohamed AAA, Santeramo FG, Sanz-Cobena A, Tsikliras A 2020 Food. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 237-264.*



## Table of contents

<b>3.2 Food</b> .....	<b>240</b>
Executive summary.....	<b>240</b>
3.2.1 Past trends and current situation.....	<b>241</b>
3.2.1.1 <i>Demand for agricultural products, consumption and trade</i> .....	<b>242</b>
3.2.1.2 <i>Marine food resources</i> .....	<b>245</b>
3.2.1.3 <i>Observed impacts of extreme weather and climate events on food production</i> .....	<b>246</b>
3.2.1.4 <i>Food policy and economics</i> .....	<b>246</b>
3.2.2 Projections, vulnerabilities and risks.....	<b>247</b>
3.2.2.1 <i>Agricultural resources</i> .....	<b>247</b>
3.2.2.2 <i>Marine food resources</i> .....	<b>249</b>
3.2.3 Adaptation and mitigation.....	<b>249</b>
3.2.3.1 <i>Adaptation of the food system to environmental change</i> .....	<b>249</b>
3.2.3.2 <i>Mitigation of climate change drivers</i> .....	<b>251</b>
<i>Nitrogen fertilization optimization</i> .....	<b>251</b>
<i>Improved water management</i> .....	<b>252</b>
<i>Soil improvement</i> .....	<b>252</b>
3.2.3.3 <i>Synergies and trade-offs between adaptation and mitigation</i> .....	<b>253</b>
<b>References</b> .....	<b>255</b>
<b>Information about authors</b> .....	<b>264</b>

## 3.2 Food

### Executive summary

Food production in the Mediterranean Basin, from both land and the sea, is impacted by climate change, more frequent and intense extreme events, jointly with land degradation, overfishing, ocean acidification and salinization of coastal soils. Climate extremes pose a threat to the entire agriculture sector. Extremes, such as heat stress, droughts but also floods, can cause crop yield losses/failures, crop quality reduction and impacts on livestock. Perturbations in the global agricultural markets may exacerbate the local impacts of climate change, especially because most Mediterranean countries are net importers of cereals and fodder/feeding products. Mostly due to unsustainable fishing, total fish landings in the Mediterranean Sea have declined by 28% from 1994 to 2017.

Climate projections show a decrease of water availability and an intensification of extremes in the Mediterranean region, and thus a higher risk for the agriculture sector. Crop yield reductions are projected for the next decades in most current areas of production and for most crops. The cultivation of some water demanding crops like maize or vegetables could become impossible in many Mediterranean regions if there will be no enough water for irrigation. This will potentially be worsened by emerging pests and pathogens, and

perturbations in the global food markets due to environmental crises elsewhere.

Sea level rise will also negatively affect the agriculture sector by its direct impact on agricultural areas and associated increasing soil salinity, which could be multiplied by three. Rice production in Egypt and Spain could be the most affected.

Climate change is projected to heavily affect marine resources in the next decades. Warmer temperatures, acidification and water pollution will likely reduce marine productivity, affect species distribution and trigger local extinction of more than 20% of exploited fish and marine invertebrates around 2050.

In agriculture, there are large possibilities for adaptation consisting mostly in changing farming practices and application of more sustainable methods, including agroecological strategies. Successful strategies for sustainable development and enhanced resilience to environmental change are based on combining different approaches, i.e., reduced tillage, varieties, rotational patterns, and crop diversity or diversification of income. Sectorial co-designed climate services will represent a key asset to reduce the risk linked to unfavorable climate conditions and extremes.



**Figure 3.13 | Total agricultural land in the Mediterranean countries** in 2016 [% with respect to the total land at country scale]. Source: World Bank (accessed February 2020).





**Figure 3.14 | Total irrigated land in the Mediterranean countries** (latest reported value in % with respect to the agricultural land at the country scale). Source of data: World Bank Data (accessed February 2020).

Sustainable intensification of farming systems offers greenhouse gas mitigation options by nitrogen fertilization optimization, improved water management, higher storage of soil organic carbon and carbon sequestration both in annual and perennial cropping systems, management of crop residues and agroindustry by-products.

### 3.2.1 Past trends and current situation

The ensemble of the Mediterranean countries has approximately 877 million ha of land, of which about 28% is devoted to agriculture (Fig. 3.13). There is a pronounced spatial heterogeneity in the share of agricultural land, from 4% of Egypt to almost 76% of Syria (Fig. 3.13). The agriculture sector is characterized by a variety of different farm structures and agro-management practices combined with pronounced differences in environmental conditions, rendering substantial variation in agricultural inputs (e.g., nutrients, pesticides, water for irrigation) and outputs (e.g., crop yields). Irrigation is practised only on 8% of the Mediterranean agricultural land area (Fig. 3.14), however uncertainties characterize this value as data for several countries are neither available nor updated. Israel has the highest portion of agricultural land being irrigated (approx. 33%, Fig. 3.14).

The Mediterranean agriculture production is characterized by high spatial variability and differences (Table 3.9). Annual crops include cereals (e.g., wheat, maize, barley and rice), and

vegetables (e.g., potatoes and tomatoes). Together, wheat, maize, barley and rice cover, for almost all Mediterranean countries, more than 90% of the entire cereal production, with rice having a significant share (>3%) only in Egypt, Greece, North Macedonia, Portugal, Spain and Italy. Permanent crops consist of fruit, olives, grapes and dates. For cereals, France, Turkey, Egypt, Spain and Italy produce (2014-2018 average) about 66, 35, 23, 21, and 18 million t, respectively (Table 3.9). As for fruit and vegetable production, the highest values (15-22 million t for fruit, and 13-24 million t for vegetables) come from Egypt, Italy, Spain and Turkey (Table 3.9).

Although productivity has increased in recent decades, there are still large differences in the region, with for instance wheat yield ranging from approx. 1 to almost 7 t ha<sup>-1</sup> (FAO 2017). These differences are also reflected in the estimated yield gap for wheat, maize and barley (e.g., Mueller et al. 2012; Schils et al. 2018). Improved agro-management practices can contribute to close the gap in regions where large differences exist between potential and farm yield. As an example, Pala et al. (2011) found that wheat yields can be increased 1.6–2.5 times in Morocco, 1.7–2.0 times in Syria and 1.5–3.0 times in Turkey.

Large spatial differences also characterize the livestock subsector, with meat (beef and buffalo) production varying from 0.1 to 143.9x10<sup>4</sup> t; while milk production varies from 0.4 to 262.7x10<sup>5</sup> t (Table 3.9). Milk productivity also spans a vast range from

	Cereal	Fruit	Vegetables	Meat (beef & buffalo)	Milk
Albania	6.9	7.8	8.0	0.39	11.4
Algeria	40.4	67.6	63.4	1.57	35.8
Bosnia and Herzegovina	13.6	3.3	7.7	0.15	7.0
Bulgaria	92.5	4.9	4.6	0.18	11.3
Croatia	30.6	3.2	12.4	0.43	6.8
Cyprus	0.4	1.8	0.7	0.05	2.5
Egypt	229.7	150.8	158.2	7.89	51.6
France	662.6	92.2	52	14.39	262.7
Greece	38.5	40.5	25.2	0.43	19.4
Israel	2.9	13.8	14.9	1.29	15.4
Italy	175.8	175.3	125.9	7.75	119.3
Jordan	1.0	5.4	16.2	0.27	3.5
Lebanon	1.7	8.0	8.2	0.45	2.6
Libya	2.7	6.8	6.8	0.09	2.3
Malta	0.1	0.1	0.8	0.01	0.4
Montenegro	0.1	0.8	0.2	0.04	1.7
Morocco	84.7	57.3	40.3	2.61	23.9
North Macedonia	5.6	5.8	6.9	0.05	4.5
Palestine	0.5	1.2	6.4	0.08	1.6
Portugal	11.9	19.6	24.4	0.89	20.8
Serbia	95.2	16.5	8.5	0.69	16.0
Slovenia	6.2	2.1	0.9	0.34	6.4
Spain	211.6	192.7	128	6.32	80.2
Syria	31.4	25.1	17.9	0.70	22.3
Tunisia	17.8	20.9	30.3	0.59	13.8
Turkey	354.3	217.5	239.8	9.90	197.2
Kosovo	95.2	16.5	8.5	0.69	16.0

**Table 3.9 | Production of cereals, fruit, vegetables, meat and milk in the Mediterranean countries, 2014-2018 average, 10<sup>5</sup> tonnes.** Data source: FAOSTAT (accessed February 2020).

800 kg animal<sup>-1</sup> in Libya to 5,500 kg animal<sup>-1</sup> in Slovenia. Overall, Mediterranean countries of the MENA region have an average milk productivity of 700 kg animal<sup>-1</sup>, compared to 1,800 kg animal<sup>-1</sup> of the other countries and 2,300 kg animal<sup>-1</sup> for the EU countries in the region.

### 3.2.1.1 Demand for agricultural products, consumption and trade

Agricultural demand in the Mediterranean region is influenced by changing dietary patterns and by the socio-economic and political situation of each country, including population growth and import/export flows. In 2013, the Mediterranean diet has been recognized by UNESCO as intangible cultural heritage of humanity, involving not only food production, processing and consumption but


also social behaviour and community identity. Its low environmental impact and its importance for a sustainable future have been highlighted in many studies (Sofi et al. 2010; Capone et al. 2012; CIHEAM and FAO 2015). Recent studies indicate a diet transition affecting the Mediterranean countries and posing a threat for the preservation and enhancement of the Mediterranean diet (Bonaccio et al. 2012, 2014; CIHEAM and FAO 2015). These changes may further affect nutritional issues and human health in Mediterranean countries, where already malnutrition (characterized by the presence of significant percentage of overweight and underweight population) takes place.

The food system of the Mediterranean also contributes to the ecological deficit as estimated by Galli et al. (2015) and updated in the National Footprint Accounts 2019<sup>15</sup>. The amount and the

<sup>15</sup> Global Footprint Network, <http://data.footprintnetwork.org/>

type of contribution are country dependent, and in some cases (e.g., Portugal, Greece, Spain, Malta, Croatia and Italy) characterized by a relevant component of meat, dairy and fish (Galli et al. 2017). Changes during the last decade (2000-2013) in meat consumption are not homogeneous in the Mediterranean region, with twelve countries show-

ing a significant increase, six having characterized by a significant decrease and the others having a stationary pattern. For fish, fourteen countries show a significant increase, only two a significant decrease and all others have a stationary pattern (Fig. 3.15).



**Figure 3.15 | Meat and fish consumption (kg capita<sup>-1</sup> yr<sup>-1</sup>) in Mediterranean countries** from 2000 to 2013 (FAO 2017).


Trade patterns play a key role in the Mediterranean region, with most of the countries being net importer of cereal and fodder/feeding products (Fig. 3.16 and 3.17). Concerning cereal, four countries (Italy, Spain, Egypt, and Algeria) import 12-19 million t of cereal, while France exports about 29 million t (Fig. 3.16). As for fodder and feeding products, five Mediterranean countries import more than one million t, with Turkey reaching about 4.3 million t (Fig. 3.17). The current trade patterns have been reached by a profound transformation of the agricultural systems that has occurred in the last decades, often characterized by a decoupling of the crop and livestock producing systems (Lassaletta et al. 2014).

Overall, the Mediterranean region, in terms of nitrogen (N) import has moved towards a more unbalanced situation with most of the countries being net larger importer (Fig. 3.18) (Lassaletta et al. 2014; Sanz-Cobena et al. 2017). The decoupling of the crop and livestock producing systems caused a lower nutrient efficiency and issues associated with the lack of manure in cropping area and


excessive manure in livestock farms (Lassaletta et al. 2012; Sanz-Cobena et al. 2017). The excessive manure production is difficult to manage, and over-application in areas close to high-density livestock systems can severely affect the environment. As a consequence, a high risk of catchment pollution has been estimated and reported in some studies (Lassaletta et al. 2012; Romero et al. 2016).

The Mediterranean is among the oldest examples of strongly coupled human-environment system that has undergone very profound land/landscape changes driven by activities such as the agriculture and by the human-water interaction (Barton et al. 2010, 2016) (Section 4.3.1.1). Many factors have contributed to these changes in the Mediterranean, e.g., people’s mobility towards the coast and the urban areas, tourism expansion, industrialization, agriculture intensification (Bajocco et al. 2012; Niedertscheider and Erb 2014).


These changes have contributed to alter land quality, productivity and to degradation. Changes have not been homogeneous as very different



**Figure 3.16 | Cereal trade patterns** (average 2014-2017 values in tonnes x 10<sup>7</sup>) in the Mediterranean countries: import (deep blue) and export (light blue) contribution for each Mediterranean country (identified by the ISO 3166-1 alpha-2 code) (FAO 2017).



**Figure 3.17 | Trade patterns in fodder and feeding products** (average 2014-2017 values in tonnes x 10<sup>6</sup>) in the Mediterranean countries: import (deep blue) and export (light blue) contribution for each Mediterranean country (FAO 2017).



**Figure 3.18 | Net protein fluxes** (Gg N) of food and feed imported to the Mediterranean regions from the other countries in 2009. Green countries are net N exporters to the Mediterranean. Yellow/red countries are net N importing from the Mediterranean. Fluxes below 50 Gg N are not represented (adapted from Sanz-Cobena et al. 2017).

socio-economic conditions characterize the region as well as behavioural patterns in farming. For instance, in some areas of the western Mediterranean the abandonment of dryland farming, of farming activities in mountainous and/or re-mote regions, and the consequent afforestation modified the ecosystems and the services provided (Kauppi et al. 2006; Falcucci et al. 2007; Padilla et al. 2010). Abandonment of agricultural terraces in mountainous regions has in some cases also favored erosion processes and loss of fertile soil (Arnaez et al. 2011). Land competition has also played a key role in some regions of southern Mediterranean, e.g., Morocco (Debolini et al. 2015). Mobility towards urban areas, evolving economic conditions, modified productivity in agricultural areas of Mediterranean countries also contributed to shifts in the cultivated crops. In Crete (Greece), for instance, a transition from cereal production towards olive cultivation characterized the 20th century (Karamesouti et al. 2015). In some countries, urbanization has forced agricultural expansion towards marginal areas requiring higher management levels in terms of irrigation and fertilization (Abd-Elmabod et al. 2019).


### 3.2.1.2 Marine food resources

Mediterranean total fishery landings have been declining in the last years (Fig. 3.19) (Tsikliras et al. 2015; FAO 2018) and so are the reconstructed catches that included discarded, illegal and unreported and recreational fisheries catch (Pauly and Zeller 2016). Total landings of the entire Mediterranean Basin exceeded 1.16 million t in 1994 and declined to around 842,000 t in 2017, i.e., a decrease of 28% (Fig. 3.19).

While the peaks occurred relatively early in the central Mediterranean (~1985) and eastern Mediterranean (~1994), the landings peaked much later in the western Mediterranean (~2006) (Fig. 3.20). In 2017, the landings were relatively low: from 161,000 t in eastern Mediterranean to 325,000 t in central Mediterranean (Fig. 3.20). In 2017, the highest contribution to the total landings in the Mediterranean came from Italy (22%), followed by Algeria (12%), Tunisia (12%), and Spain (10%)<sup>16</sup>.

Small pelagic fishes constitute the vast majority of landings across the entire Mediterranean Sea, with European anchovy (*Engraulis encrasicolus*) and

<sup>16</sup> Data from FAO-GFCM, accessed March 2020. <http://www.fao.org/gfcm/en/>




**Figure 3.19 | Total fish landings (tonnes) in the Mediterranean Sea** from 1970 to 2017. Data source: FAO-GFCM<sup>16</sup>, accessed in March 2020.

European sardine (*Sardina pilchardus*) being the species with the most landings accounting for 34% of the 2014-2016 average annual landings (FAO 2018; Tsikliras et al. 2019). The main pelagic fish species landed in the western Mediterranean are European anchovy, European sardine and sardinella nei (*Sardinella* spp.) accounting for 46% of the total landings (FAO 2018). In the Ionian part of central Mediterranean, European sardine, sardinella nei, jack and horse mackerel nei (*Trachurus* spp.), and common Pandora (*Pagellus erythrinus*) account for 30% of the total landings (FAO 2018). While in the Adriatic part of Central Mediterranean, landings of European sardine and anchovy reach 61% (FAO 2018). Finally, in the eastern Mediterranean, European sardine, anchovy and European sardine are again the main species landed accounting for 33% of the total landings (FAO 2018) [Sections 2.4.2, 4.1 and 4.2].

### 3.2.1.3 Observed impacts of extreme weather and climate events on food production

Extreme weather and climate events (such as floods, droughts, storms, heat waves and cold spells) pose a threat for agricultural production (Lesk et al. 2016; Zampieri et al. 2017; FAO 2018). The impacts of heat stress occurring in critical phenological phases can induce serious crop yield losses and quality reduction. In Italy, for instance, early heat waves have been associated to durum wheat yield losses occurred in the last decades (Fontana et al. 2015; Zampieri et al. 2017). In Greece, recent trends in extreme temperatures reduced cereal yields by 1.8-7.1% per degree increase in maximum temperatures (Mavromatis 2015). Also milk production and quality are affected by heat stress (Bernabucci et al. 2010, 2015; Gantner et al. 2017) as well as livestock fertility (de Rensis et al. 2015). Temperature changes during important



**Figure 3.20 | Total Landings (t)** from 1970 to 2017 in the Mediterranean Sea. Data source: FAO-GFCM<sup>16</sup>, accessed in March 2020.

phenological stages such as blossoming may affect yields of maize, alfalfa, apples, almonds and other crops (Savé et al. 2012; Funes et al. 2016; Díez-Palet et al. 2019).

The entire agriculture sector in the region is also heavily affected by drought events (Blauhut et al. 2015; Zampieri et al. 2017). Severe socio-economic impacts triggered by drought events were reported on Moroccan agriculture (Verner et al. 2018), with the events of 2007 (Schilling et al. 2012) and 2015-16 that caused heavy losses on wheat, citrus and olive production, posing a threat for the livestock sector. Severe droughts can also modify the rural landscape, preventing the adoption of new crops and ultimately forcing farmers to emigrate (Ruauudel and Morrison-Métois 2017). The costs and the risks associated with climate extremes are not only related to direct losses, such as crop failures, but also to a wide range of indirect effects triggered by market reactions to events occurring in other producing regions of the world (Chatzopoulos et al. 2019).

The Mediterranean is also a high fire-risk region, where fires are the cause of severe agricultural, economic and environmental losses and even human casualties (San-Miguel-Ayanz et al. 2013; Moritz et al. 2014; Bowman et al. 2017). The abandonment of agricultural land leads to an increased risk of forest fires due to the occupation of what were agricultural lands by forest and the bushes, and increasing the biomass available for burning as well as its spatial continuity. Conversely, some forest fires may be triggered for the creation of more pastures for livestock or farmland.

### 3.2.1.4 Food policy and economics

Agriculture and the entire food system are generally influenced by socio-economic conditions, also in Mediterranean countries. The strong

fluctuations in food markets are partly due to the characteristics of agricultural production itself (perishable products, climatic and health risks, seasonal production cycles, size of farms, distance from markets etc.), which, together with aspects of the overall economy, even of a geopolitical nature, can modify the food supply-demand balances (Reguant and Savé 2016). These conditions also endanger the capacity of Mediterranean countries to guarantee food security (Santeramo 2015). Among the Mediterranean countries, some in the southern and eastern shores have also suffered political instabilities and conflicts that have posed a challenge to the maintenance and development of the agriculture system (Tanyeri-Abur 2015; Petit and Le Grusse 2018).

The food system of the Mediterranean region in all its aspects (production, trade patterns, etc.) is under strong influence from the policies of high-income countries and, in particular, the European Union (Caracciolo et al. 2014). The tight links among Mediterranean countries imply that changes in the EU Common Agricultural Policy and in trade agreements may have important impacts on national agri-food sectors also outside the EU. For instance, the Euro-Mediterranean trade partnership between the EU and the southern and eastern Mediterranean non-EU countries (except for Syria and Libya, entered into force to promote trade and investments in the region) tends to influence market fundamentals in all Mediterranean countries. Furthermore, food quality standards and entry price mechanisms are very important for trade patterns (Cioffi et al. 2011; Santeramo and Cioffi 2012; Marquez-Ramos and Martinez-Gomez 2016; Bureau and Swinnen 2018).

Trade has prioritized the export of fruits and vegetables and has widened the production-consumption gap of cereals, which are the main food of the most vulnerable segments of the population in the southern and eastern parts of the region (Larson et al. 2002; Cioffi and Dell'Aquila 2004; García Martínez and Poole 2004). The vulnerability of the cereal sector has enhanced the impact of food price fluctuation on food security, which may have severe impacts (e.g., in terms of income level and income distribution) depending on the capacity of the countries to be self-sufficient (Caracciolo and Santeramo 2013).

Pasture based systems are becoming less competitive, due to the high labor costs (on-farm resources are being substituted with external inputs) promoting intensive livestock systems near urban areas (Malek et al. 2018). This has resulted in the increase in landless livestock systems in the Mediterranean region.

Mediterranean countries are vulnerable to price fluctuations on international markets due to their dependence on imports of basic foodstuffs. Worldwide phenomena (e.g., food crisis) have accentuated the structural weakness of the agricultural production model adopted by these countries, increasing social and political frustrations (Reguant and Savé 2016). In this context, it is worth to mention initiatives such as the Mediterranean Agricultural Market Information Network (MED-Amin)<sup>17</sup> and the MedAgri platform<sup>18</sup> from FAO, the European Bank and the World Bank. The drivers of price volatility are numerous and complex (Santeramo et al. 2018b), but it seems there is a consensus that arbitrage, and price discovery mechanisms tend to have a positive impact on price stabilization (Santeramo and Lamonaca 2019).

Access to agricultural technology is unequally distributed across countries of the Mediterranean area and is usually more accessible to farmers in the north-western part. It is also true that most developed countries tend to provide higher subsidies (at least in nominal terms) to their agricultural sector. The adoption of risk management strategies, and the access to credit are also very unequally distributed and generally lower in the developing countries of the Mediterranean area (Santeramo et al. 2014). These peculiarities allow to conclude that the less developed countries, among the Mediterranean ones are likely to be the most vulnerable to food security issues. On the other hand, investments on technology, on policies to promote the agri-food sector, and in particular to promote risk management strategies may prove effective mechanisms to enhance resilience to food security.

### 3.2.2 Projections, vulnerabilities and risks

#### 3.2.2.1 Agricultural resources

Climate projections indicate significant warming and drying in the Mediterranean Basin, together with intensification of climate extremes such as


<sup>17</sup> [www.med-amin.org](http://www.med-amin.org)

<sup>18</sup> [www.medagri.org](http://www.medagri.org)

drought and heat waves (Sections 2.2.5.2 and 2.2.4). Thus, severe impacts on the agriculture sector are to be expected if no adaptation and mitigation will take place. These impacts include changes in phenology and growing cycle of many crops (Trnka et al. 2011; Funes et al. 2016), combined with higher water demands due to enhanced evaporation (Savé et al. 2012; Girard et al. 2015; Saadi et al. 2015; Valverde et al. 2015; Phogat et al. 2018). The wheat growing period in Tunisia is expected to be shortened by 16 days for 2.5°C and by 30 days for 4°C (Mougou et al. 2011). Additional constraints include water scarcity (Section 3.1.4.1) (Vicente-Serrano et al. 2017, 2018) and soil salinity (Lagacherie et al. 2018; Phogat et al. 2018).

As a consequence, potential yields of crops and livestock yields are projected to decline in many areas due to climatic and other stress factors without adaptation. Several regions of the Mediterranean might entirely lose their suitability for growing specific crops (Ceglar et al. 2019). Crop yields in MENA countries are expected to decline by approx. 30% with 1.5-2°C warming in Jordan (Al-Bakri et al. 2011) and similarly in North Africa (Drine 2011) and up to 60% with 3-4°C warming (Schilling et al. 2012). Maize is projected to be among the most affected crops (Webber et al. 2018; Zampieri et al. 2019; Feyen et al. 2020), with significant yield decline of, e.g., 10-17% in Italy, Bulgaria, and Greece by the mid-century (2021-2050, under the business as usual RCP8.5 scenario and assuming the current agro-management will still be in place). Wheat yield losses are projected for some European countries in the Mediterranean region (5%-22% in 2021-2050 under the RCP8.5 scenario with no adaptation) (Feyen et al. 2020) associated with higher inter-annual variability and decrease resilience of the production (Zampieri et al. 2020). Reductions in wheat yield in case of no adaptation have been also reported for Algeria (Chourghal et al. 2016). However, reductions in water availability for maize irrigation could bring much bigger losses. Based on a meta-analysis of 16 studies available at the time, Waha et al. (2017) conclude that climate change constitutes a significant risk for crop yields across the MENA region (Fig. 3.21).

Soil and agro-management influence on the projected changes has also been reported for wheat yield in an Italian region, showing moderate yield increase as well as heavy decrease (63% in 2040-2070 under the A1B scenario, which is close to the relatively high scenario RCP6.0) (Bird et al. 2016) according to the soil type. While a strong dependence on water availability has been pointed out for tomato yield in Tunisia, where a 10% reduction in



**Figure 3.21 | Crop yield changes in the MENA region** based on a meta-analysis of 16 different studies (Waha et al. 2017).

water for irrigation could make some productions not feasible (Bird et al. 2016). In Tunisia, wheat yields may increase in some producing areas (Annabi et al. 2018). However, recurrent drought events may induce losses of approximately -50% in olive production, and the increase in floods could lead to a decrease of -13% in rainfed cereal production (Requier-Desjardins 2010). The effects of changes in precipitation regimes on olive production in the entire Mediterranean region were investigated by Tanasijevic et al. (2014) that pointed to the likely absolute need of irrigation in future olive cultivation. Large climate impacts have been also estimated (assuming no adaptation) for agriculture in Egypt increasing over time and triggering exceptional food price increases (McCarl et al. 2015). Sea level rise will also pose a threat to agriculture in Egypt leading to area losses and affecting, for instance, rice production (Chen et al. 2012; Sušnik et al. 2015). Similarly, heavy impacts of sea level rise and associated increased soil salinity (estimated to be three times the current one) on rice production have been estimated at the end of the century in Spain (Genua-Olmedo et al. 2016).

The increase in the atmospheric concentration of CO<sub>2</sub> could bring beneficial effects in terms of yield under optimal growing conditions (especially for C<sub>3</sub> crops such as wheat and barley) and could buffer some days more under drought conditions (Kimball 2016) but may also bring new nutritional challenges (Uddling et al. 2018; Asseng et al. 2019). Results also highlight the limited beneficial contribution of elevated atmospheric CO<sub>2</sub> concentration under pronounced water-stress conditions. Reductions in wheat protein yield by 5-10% have been estimat-



ed at some south-western locations of the basin (2040-2069 under the RCP8.5 scenario) (Asseng et al. 2019).

Concurrent and recurrent extremes, not fully considered in the current impact assessments, may well pose the main threat to the stability and the resilience of the Mediterranean production systems. Climate extremes occurring in other regions of the world may also trigger impacts through increased market volatility and price spikes (Chatzopoulos et al. 2019; Toreti et al. 2019). New and re-emerging pest and pathogens, usually not fully considered in impact assessments, may contribute to larger than estimated losses (Bebber et al. 2013, 2014). Another threat to food security and quality may be represented by mycotoxigenic fungal pathogens and higher level of contamination (Medina et al. 2017). Agriculture in the region may be also affected by increased risk of large fires, with a 34 to 140% rise according to the location and the scenario (Section 4.3.2.1).

### 3.2.2.2 Marine food resources

Projected climate change is also expected to heavily impact marine food resources, which are over-exploited already. Ocean warming, acidification, water pollution and constrained migration possibilities to cooler areas (due to sea enclosure) may lead to local extinction of up to 50% of exploited fish and marine invertebrates around 2050, affecting also endemic fishes, including commercial ones (e.g., Cheung et al. 2016) (Section 4.1.2.1). Pollution from anthropogenic activities also affects fish population, notably in the Nile delta, with potentially serious consequences on human food security.

Besides warming, marine ecosystems are also sensitive to increasing atmospheric CO<sub>2</sub> concentration due to its rapid dissolution in seawater, which causes alterations in the chemistry of inorganic carbon with a lower pH and higher concentration of carbonic ions (Doney et al. 2009). The carbonic ion is an essential element for organisms that depend on the deposition of calcium carbonate (CaCO<sub>3</sub>) through biomineralization for the formation of calcareous structures, such as the mollusk shells. If the biomineralization process does not occur properly, organisms reduce their growth rate and may present morphological anomalies that cause, for example, the loss of capacity of fixation to the substrate and diminishes feeding activity.

The joint effects of ocean warming and acidification may also include a number of biological alterations such as decreased ocean productivity

(Behrenfeld et al. 2006), reduced growth and survival of calcifying organisms (Hoegh-Guldberg et al. 2007), changes in species distributions, altered food dynamics (Vergés et al. 2014), and altered incidence of disease (Burge et al. 2014). These effects can be translated into a diminution of the abundance and, therefore, of the fisheries production. The affected activities would be both extractive fishing (shellfish, in this case) and aquaculture that is extensively used in coastal areas (Gazeau et al. 2014; Prado et al. 2016). The example of mollusks is perhaps the most quoted but we should not forget that other organisms are subject to biomineralization processes for the formation of the skeleton (e.g., fish), and thus, can also be negatively affected. The effects on habitats can also be quite important because at lower pH some plants can be affected directly or indirectly and even calcareous substrates of biological or mineral origin. All this has to be added to the benthic organisms with calcareous structures that are not commercial species but are the base of the food web for upper trophic groups.

To fully understand the potential effects of global change, it is important to focus on population bottlenecks, which are usually early developmental and reproductive stages (Thorson 1950). Survival of adult and juvenile bivalves shows little dependence on pCO<sub>2</sub> (Berge et al. 2006; Hendriks et al. 2010; Range et al. 2012), although increasing temperature may result in increasing mortality and metabolic rates (Basso et al. 2015). In contrast, gametes, embryos, and larvae are generally more sensitive to both temperature and elevated pCO<sub>2</sub> stress (Havenhand et al. 2008; Parker et al. 2009, 2010) because the deposition of CaCO<sub>3</sub> shells and skeletons begins in these stages (Kurihara et al. 2007). Yet, there is also a wide natural variability in pH ranges of seawater (7.5-8.5, with even lower values possible in semiconfined waters; Flecha et al. 2015) which may partly account for observed differences in the responses of calcifying organisms to acidification and complicates the generalization of patterns across species and ecosystems (Kurihara 2008).

## 3.2.3 Adaptation and mitigation

### 3.2.3.1 Adaptation of the food system to environmental change

The assessment of how climate change will affect crops is essential for policymakers, planners, farmers and all the other actors of the agriculture sector to develop, propose and implement adaptation and mitigation strategies at the local/regional

scale to make agriculture more resilient to changes (Liebig et al. 2016). For instance, future water availability and water demands put the current management model in question, so adaptation choices have to be necessarily developed (Iglesias and Garrote 2015; Ronco et al. 2017). The projected water scarcity and increase of drought events will limit adaptation actions based on irrigation. Under some scenarios combining climate change and population growth, half of the Mediterranean countries (mainly in the southern and eastern shores) are projected to be unable to cover irrigation water demands by the end of the century (Fader et al. 2016) (Section 3.1.5.2).

Crop distribution, diversity, varieties, rotation patterns, and agro-management represent key elements of adaptation strategies at the farm scale (Valverde et al. 2015; EEA 2019). Breeding and sowing new varieties water and heat stress tolerant (del Pozo et al. 2016, 2019; Hatfield and Dold 2019), adapting the crop calendar (Ronco et al. 2017), using optimal crop diversification (Lin 2011) could be all used as adaptation strategies. The inclusion or reintroduction of wild food plants, neglected and underutilized crops, also add to diversifying the agricultural portfolio of crops with potential resilience against climate change. North-south differences were estimated for the adaptive capacity in agriculture (Grasso and Feola 2012), mainly associated with soft factors (e.g., information) rather than with other more structural ones (such as technological and infrastructural perspective). Looking at the implemented adaptation strategies in the Mediterranean (Harmanny and Malek 2019), the most common ones are farming practices (diversify and change crop types, adjust crop rotation), water management (modify irrigation practices), and farm management (diversify source of income). The main drivers of such adaptation actions are water scarcity, environmental factors (climate change, soil degradation and erosion, sustainability), and socio-economic factors (Harmanny and Malek 2019).

Combining several actions can also lead to better results in terms of crop yield. Higher wheat yield under different water conditions in Lebanon were achieved by using a drought-tolerant variety, conservation tillage and precise irrigation during grain filling<sup>19</sup>. A higher degree of diversification, more varieties of the main crops, earlier sowing,

and hedgerow planting were also identified as actions to increase the resilience in a pilot farm project in southern France<sup>20</sup>. Crop productivity (vines, corn, apples, lucerne) was increased using different agronomical practices to increase water availability by plants without increase water from irrigation in Spain<sup>21</sup>. Some strategies have also additional indirect benefits, such as soil organic carbon accrual due to agroforestry (Chatterjee et al. 2018), cover crops (Aguilera et al. 2013b; Vicente-Vicente et al. 2016), and local crop varieties with a higher residue and root biomass production (Carranza-Gallego et al. 2018) (Section 6.4). Improved soil erosion control, increasing soil fertility, retaining soil moisture and resource efficiency are the dominant drivers for conservation agriculture, organic farming and agroforestry (Lagacherie et al. 2018). Conservation agriculture represents a relatively widely adopted management system that aims to sustain long-term crop productivity and system's sustainability (Kassam et al. 2012). The environmental and economic benefits of no-till implemented as its core principle combined with other practices have been pointed out in several studies (Peigné et al. 2007, 2015; Cooper et al. 2016; Vincent-Caboud et al. 2017). The use of sectorial climate services (Buontempo et al. 2020; Ceglar et al. 2020) at different spatio-temporal scales will also be a key adaptation measure to reduce the risks and alleviate the impacts of extreme events.

Advanced agricultural technologies may also influence the ability of the region to produce food (Asseng et al. 2019) and adapt to the changing climate and environment. Precision agriculture will make possible a targeted monitoring of plant growth and thus a more efficient use of resources (water, pesticides, nutrients) by combining technologies for data collection (e.g., in-field sensors, weather stations, imaging) with analytical tools, computer vision and artificial intelligence technologies (Bhakta et al. 2019). Precision agriculture has been already applied by some Mediterranean countries (e.g., Israel, Italy, Spain), in viticulture and other crops, and holds a significant technological innovation potential. At the same time, bio and nanotechnologies may help to ensure food security and increase productivity (King et al. 2018; Santeramo et al. 2018a). Cultured, plant-based and insect-based meat are emerging technologies for producing alternatives to meat-

<sup>19</sup> Results from the SWIM-project ACLIMAS, [www.aclimas.eu](http://www.aclimas.eu)

<sup>20</sup> Results from the LIFE-project AGRI-ADAPT, [www.agriadapt.eu](http://www.agriadapt.eu)

<sup>21</sup> Results from the LIFE-project MEDACC, <http://medacc-life.eu/>

derived proteins whose demand is growing. High costs and consumer reluctance appear to be major obstacles to the implementation of these techniques (Santeramo et al. 2018a; Gómez-Luciano et al. 2019).

### 3.2.3.2 Mitigation of climate change drivers

Mediterranean climatic conditions host two main crop production systems, rain-fed and irrigated, largely differing in terms of management and, consequently, emissions of N<sub>2</sub>O, a potent greenhouse gas. Rain-fed systems are usually characterized by periods with low soil moisture and cold temperatures, thus with decreased soil micro-biological activity and N<sub>2</sub>O fluxes. Recent reviews have shown that N<sub>2</sub>O emission factors (EF) from rain-fed Mediterranean cropping systems are much lower than the IPCC-default EF threshold of 1% (Aguilera et al. 2013b; Cayuela et al. 2017). Rain-fed crops in Mediterranean regions have lower EFs (EF: 0.27%) than irrigated crops (EF: 0.63%). Irrigated systems usually receive large amounts of water and nitrogen inputs, which create favorable soil conditions for N<sub>2</sub>O emissions. Emission factors in these systems fluctuate greatly according to water management and the type and amount of fertilizer used (e.g., synthetic, solid or liquid manures). Sprinkler irrigated crops lead to N<sub>2</sub>O emission factor of 0.91%; whereas, drip irrigated systems emit at a lower rate (EF:0.51%) (Cayuela et al. 2017).

Among the most relevant mitigation strategies, there are: nitrogen fertilization optimization; improved water management; better storage of soil organic carbon and carbon sequestration in soil and perennial wood structures of woody crops; management of crop residues and agroindustry by-products.

#### **Nitrogen fertilization optimization**

Optimized nitrogen fertilizer application (in terms of input rate and time of application), as well as the careful selection of the type of fertilizer used are crucial to improve crop productivity while reducing N<sub>2</sub>O emissions (Sanz-Cobena et al. 2017). An additional effect could be achieved by applying already existing nitrogen (organic fertilizer) when possible or with the use of nitrification and urease inhibitors. Reduction of nitrogen application rates according to soil nitrogen availability and crop yield potential may decrease nitrogen surpluses and subsequent direct and indirect N<sub>2</sub>O emissions, while saving energy and abating other greenhouse gas emissions (e.g.,

associated to manufacturing synthetic fertilizers). Significant effects of nitrogen application timing on N<sub>2</sub>O emissions have been reported from cereal crops in Mediterranean countries such as Spain (Abalos et al. 2016). The estimated N<sub>2</sub>O mitigation potential, through adjusted fertilization (rate and timing) in Mediterranean agro-ecosystems ranges between 30% and 50% compared to non-adjusted practices. Replacing mineral nitrogen with organic fertilization provides not only nitrogen, phosphorus, potassium (NPK) and micronutrients to the soil and crop, but also organic carbon when using solid fertilizers (i.e., solid manure, composts, etc.), which is highly beneficial in Mediterranean soils with low organic carbon contents (Aguilera et al. 2013a; Funes et al. 2019).

In areas where croplands co-exist with livestock farms, using a farm sub-product allows the reuse/recovery of farm products, thus decreasing the volume of waste that needs to be managed, and then avoiding the emission of greenhouse gases both in the management of such wastes and in the manufacturing of new synthetic fertilizers. In Mediterranean areas, the efficient use of manure of fertilizer should be encouraged, and this could be facilitated by increased cooperation between farmers' unions. The use of organic sources of fertilizers may also decrease the need to import synthetic sources thus decreasing greenhouse gas emissions from the production and transport stages. Unfortunately, current intensive livestock production systems are often decoupled from agricultural systems (Sanz-Cobena et al. 2017).

The N<sub>2</sub>O emission reduction at plot scale depends on the form of manure used. Solid manures have proved to significantly decrease N<sub>2</sub>O emissions (ca. 23%) in Mediterranean systems (Aguilera et al. 2013b), although there is some contradictory information in the scientific literature (Webb et al. 2004; Thorman et al. 2007). For liquid manures (i.e., slurries), no significant differences have been observed when these substitute synthetic nitrogen sources. This seems to be a consequence of the strong similarities between available nitrogen, in the form of NH<sub>4</sub><sup>+</sup>, in both fertilizer types (Meijide et al. 2009; Plaza-Bonilla et al. 2014).

Trade-offs in the form of NH<sub>3</sub> emissions, odors, enhanced denitrification rates due to coexistence of high soil water contents and organic carbon suitable for denitrifiers, must be considered together with the application technology used to fertilize with liquid manures. Nitrification and urease inhibitors (NI) are used in a wide range of agro-climatic regions (Akiyama et al. 2010; Gilsanz

et al. 2016). In Mediterranean soils, NIs have shown high mitigation efficiency in rain-fed and irrigated fields, with a likely indirect effect on denitrification in the latter systems (Meijide et al. 2010). Soil texture may regulate mitigation efficiency (Barth et al. 2008) but to a limited extent, since soil texture has been shown to have a small influence on the inhibition of nitrification (Gilsanz et al. 2016).

### **Improved water management**

The different soil conditions between irrigated and rain-fed crops affect soil microbial processes, which control the fluxes of carbon (carbon dioxide, CO<sub>2</sub>; methane, CH<sub>4</sub>; organic carbon) and nitrogen (nitrous oxide, N<sub>2</sub>O; molecular nitrogen, N<sub>2</sub>; nitrate, NO<sub>3</sub>; ammonia, NH<sub>3</sub>). Soil moisture is a key factor affecting N<sub>2</sub>O losses (del Prado et al. 2006; García-Marco et al. 2014), hence the potential for N<sub>2</sub>O mitigation linked to irrigation technologies is high, even above 50% (Sánchez-Martín et al. 2008, 2010; Guardia et al. 2016). The lower amounts of water applied in subsurface drip irrigation (SDI) or normal/superficial drip irrigation (DI) through more frequent irrigation events, generate “dry” and “wet” areas in the soil, lowering the overall soil moisture and favoring nitrification over denitrification (Sánchez-Martín et al. 2010), thus reducing N<sub>2</sub>O emissions. Drip irrigation systems have shown an N<sub>2</sub>O emission factor of only 0.18%, compared to 1% in sprinkler systems (SI), showing the mitigation potential of irrigation technologies in the Mediterranean region (Cayuela et al. 2017). Optimized irrigation techniques to decrease greenhouse gas emissions from Mediterranean regions are particularly used in perennial crops and intensive vegetable cropping systems and in paddy soils (water table management). Other strategies which have been shown to be effective increasing nitrogen use efficiency and reducing N<sub>2</sub>O emissions are fertigation and sub-surface drip irrigation (Ayars et al. 2015).

### **Soil improvement**

Most Mediterranean agricultural landscapes are subject to soil organic matter depletion, particularly in the southern and eastern parts of the basin (Ryan et al. 2006). In the northern part of the basin, the issue of low soil organic matter (SOM) is of particular concern for perennial systems such as orchards and vineyards (Meersmans et al. 2012). Maetens et al. (2012) showed that bare soils, vineyards and orchards in Europe are prone to high mean soil losses (10-20 t ha<sup>-1</sup> yr<sup>-1</sup>), while cropland and fallow show smaller values (6.5 and 5.8 t ha<sup>-1</sup> yr<sup>-1</sup>) largely because the latter occupy

land exhibiting little or no slope. SOM in the Mediterranean countries is somewhat affected by climate change, with land use types such as permanent pasture and cropland being more sensitive than forests (Fantappié et al. 2011). Large losses of SOM may also be caused by erosion caused by the torrential storms that frequently occur in Mediterranean regions (Lagacherie et al. 2018). Likewise, rainfall shortage limits net primary productivity and, in turn, soil carbon buildup. Low carbon inputs driven by limited soil moisture availability are exacerbated by the adoption of certain management practices. Crop residues competition for livestock feeding or the introduction of long fallowing in the crop rotation are two examples of typical management practices in the Mediterranean region that have contributed to the reduction of carbon inputs returned into the soil.

Besides decreases in carbon inputs, agricultural management may also cause soil organic carbon (SOC) losses. Reduction or complete cessation of tillage decreases the direct incorporation of fresh organic debris into deeper soil layers. The absence of tillage (NT) slows down aggregate turnover and increases the physical stabilization of SOC within soil aggregates (Álvaro-Fuentes et al. 2008; Mrabet 2008; Plaza-Bonilla et al. 2010). When tillage is avoided, an approximate annual increase of 1% in SOC can be observed in Mediterranean croplands (Aguilera et al. 2013a). These estimates are highly dependent on soil depth, since vertical SOC distribution in no tillage (NT) and conventional tillage (CT) systems are different (Cantero-Martínez et al. 2007). Further, the assumption of a steady and linear C sequestration may not hold true, because the annual carbon accumulation rate tends to decrease in the long-term (Álvaro-Fuentes et al. 2014).

Long crop rotations have been proposed in rain-fed Mediterranean systems to enhance carbon sequestration and restore soil fertility and structure (Benhabib et al. 2014). The effect of crop rotations on carbon sequestration is highly dependent on time with no significant effect reported in short-term studies (Saber and Mrabet 2002; López-Bellido et al. 2010). Positive effects in long-term experiments (>15 years) could appear if crop biomass is properly managed after harvest (Masri and Ryan 2006; López-Bellido et al. 2010; Martiniello and Teixeira da Silva 2011). The effect of crop rotations on SOC stocks is also dependent on the type of crops included in the rotation (Triberti et al. 2016) and the management of crop residues. The introduction of perennial crops has

shown benefits (di Bene et al. 2011; Pellegrino et al. 2011). The substitution of bare fallows by any crop has been associated with SOC stabilization in NT systems (Álvaro-Fuentes et al. 2009), and to reduced soil erosion (Boellstorff and Benito 2005). The effect of inclusion of grain legumes in rain-fed yearly rotations on carbon sequestration is uncertain, due to their low biomass production, although their conversion to stabilized soil organic matter could be more efficient than that of cereals (Carranca et al. 2009). Consequently, the highest potential of fallow and legumes for mitigating greenhouse gases from these types of cropping systems comes from the avoidance of fertilizer production emissions.

Estimating the greenhouse gas mitigation potential of using crop residues and organic by-products from agroindustry in Mediterranean areas implies accounting for: (i) soil amendments to improve SOM and enhance SOC sequestration (Aguilera et al. 2013a), (ii) feedstock for bioenergy production (di Giacomo and Taglieri 2009; Spinelli and Picchi 2010), (iii) co-substrate for composting (Santos et al. 2016), (iv) feed for livestock (Molina-Alcaide and Yáñez-Ruiz 2008) or (v) construction materials (e.g., animal beds, buildings). The potential to increase SOC levels by using agroindustry by-products, as in crop residues, depends on their composition and degradability. However, agroindustry by-products vary widely in their chemical composition and therefore in their degradation rates. For example, olive and mill waste as they have very low degradation rate in the soil have been found to be good amendments to increase SOC when applied to the soil (Saviozzi et al. 2001).

Besides the potential direct greenhouse gas reduction that any strategy involving the return of the crop residues and agroindustry by-products to the soil may cause (Kassam et al. 2012; Plaza-Bonilla et al. 2014), applying these materials, treated or untreated, as soil amendments can also deliver environmental co-benefits. These benefits include erosion reduction when raw products are used for mulching (Blavet et al. 2009; Jordán et al. 2010) or, in general, the closing the nutrient cycles, with associated potential reductions of fertilizer use and reductions in draught force and fuel consumption for soil tillage (Peltre et al. 2015). Trade-offs may occur with some of the strategies that may result in larger greenhouse gas mitigation potential. For example, the use of crop residues on the soil surface might pose a risk of fire in some Mediterranean areas and, sanitary, pollution and

legal constraints may apply, especially if the by-product is applied to crops e.g., fresh vegetables without pre-treatment.

### **3.2.3.3 Synergies and trade-offs between adaptation and mitigation**

Developing sectorial adaptation strategies requires considering also the mitigation needs and efforts (Sanz-Cobena et al. 2017). For N<sub>2</sub>O mitigation and its links with adaptation measures, the pedoclimatic conditions for soil processes in Mediterranean cropping systems imply different N<sub>2</sub>O emission patterns as compared to temperate soils (Aguilera et al. 2013b). Nitrification and nitrifier-denitrification, and not denitrification, are very often the main pathways leading to emissions of nitrogen oxides in rain-fed Mediterranean cropping systems (Sánchez-Martín et al. 2008; Kool et al. 2011; Aguilera et al. 2013b; Norton and Ouyang 2019). These two processes are favored by conditions of soil water content (i.e., water filled pore space, WFPS) under saturation (i.e., 40–60% WFPS). Denitrification may play a predominant role in anaerobic soil microsites in intensively managed and irrigated systems (Sanz-Cobena et al. 2012, 2014). Consequently, different cumulative N<sub>2</sub>O emissions have been proposed for rainfed crops (0.7 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) and for e.g., sprinkler irrigated crops in Mediterranean areas (4.4 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) (Cayuela et al. 2017).

The importance and potential for N<sub>2</sub>O mitigation and the best mitigation strategy differ greatly depending on the cropping system, being highly affected by adaptation strategies regarding to soil organic carbon and water management in cropping systems. In this sense, increasing the generally low carbon content of Mediterranean soils is an important greenhouse gas mitigation strategy (Robertson et al. 2000), and is also a priority for preventing erosion and improving soil quality. Soil organic carbon content of Mediterranean soils is typically lower than in temperate areas (Chiti et al. 2012), and degradation processes are present in many areas (Lahmar and Ruellan 2007), a trend that is expected to be exacerbated by climate change in the coming decades (Al-Adamat et al. 2007). However, SOC in Mediterranean croplands is also highly responsive to management changes such as organic amendments, cover crops and tillage reductions, and there is a high potential for SOC storage through land restoration. Significant carbon sequestration rates have been observed after the application of recommended management practices and organic management

in Mediterranean cropping systems (Aguilera et al. 2013a). This high responsiveness is reflected in SOC storage rates nearly one order of magnitude higher than the 0.4% annual SOC increase proposed by the “4 per 1,000” initiative, as reported in a recent meta-analysis (Chabbi et al. 2017; Minasny et al. 2017) . This meta-analysis underlined the differences between herbaceous and woody crops regarding the carbon sequestration potential and

the practices to be applied in each system. Thus, organic fertilizers, tillage reduction and residue retention are effective practices in herbaceous systems. Woody systems, in which the storage potential is higher, would greatly benefit from maintaining a soil cover and making use of agro-industry byproducts, such as composted olive mill waste, as a source of organic matter (Vicente-Vicente et al. 2016).

## References

- Abalos D, Jeffery S, Drury CF, Wagner-Riddle C 2016 Improving fertilizer management in the U.S. and Canada for N<sub>2</sub>O mitigation: Understanding potential positive and negative side-effects on corn yields. *Agric. Ecosyst. Environ.* 221, 214–221. doi: [10.1016/j.agee.2016.01.044](https://doi.org/10.1016/j.agee.2016.01.044)
- Abd-Elmabod SK, Fitch AC, Zhang Z, Ali RR, Jones L 2019 Rapid urbanisation threatens fertile agricultural land and soil carbon in the Nile delta. *J. Environ. Manage.* 252, 109668. doi: [10.1016/j.jenvman.2019.109668](https://doi.org/10.1016/j.jenvman.2019.109668)
- Aguilera E, Lassaletta L, Gattinger A, Gimeno BS 2013a Managing soil carbon for climate change mitigation and adaptation in Mediterranean cropping systems: A meta-analysis. *Agric. Ecosyst. Environ.* 168, 25–36. doi: [10.1016/j.agee.2013.02.003](https://doi.org/10.1016/j.agee.2013.02.003)
- Aguilera E, Lassaletta L, Sanz-Cobena A, Garnier J, Vallejo A 2013b The potential of organic fertilizers and water management to reduce N<sub>2</sub>O emissions in Mediterranean climate cropping systems. A review. *Agric. Ecosyst. Environ.* 164, 32–52. doi: [10.1016/j.agee.2012.09.006](https://doi.org/10.1016/j.agee.2012.09.006)
- Akiyama H, Yan X, Yagi K 2010 Evaluation of effectiveness of enhanced-efficiency fertilizers as mitigation options for N<sub>2</sub>O and NO emissions from agricultural soils: meta-analysis. *Glob. Chang. Biol.* 16, 1837–1846. doi: [10.1111/j.1365-2486.2009.02031.x](https://doi.org/10.1111/j.1365-2486.2009.02031.x)
- Al-Adamat R, Rawajfih Z, Easter M, Paustian K, Coleman K et al. 2007 Predicted soil organic carbon stocks and changes in Jordan between 2000 and 2030 made using the GEFSOC Modelling System. *Agric. Ecosyst. Environ.* 122, 35–45. doi: [10.1016/j.agee.2007.01.006](https://doi.org/10.1016/j.agee.2007.01.006)
- Al-Bakri J, Suleiman A, Abdulla F, Ayad J 2011 Potential impact of climate change on rainfed agriculture of a semi-arid basin in Jordan. *Phys. Chem. Earth* 36, 125–134. doi: [10.1016/j.pce.2010.06.001](https://doi.org/10.1016/j.pce.2010.06.001)
- Álvaro-Fuentes J, Cantero-Martínez C, López M V., Paustian K, Deneff K et al. 2009 Soil aggregation and soil organic carbon stabilization: effects of management in semiarid Mediterranean agroecosystems. *Soil Sci. Soc. Am. J.* 73, 1519–1529. doi: [10.2136/sssaj2008.0333](https://doi.org/10.2136/sssaj2008.0333)
- Álvaro-Fuentes J, López M V., Cantero-Martínez C, Arrúe JL 2008 Tillage effects on soil organic carbon fractions in Mediterranean dryland agroecosystems. *Soil Sci. Soc. Am. J.* 72, 541. doi: [10.2136/sssaj2007.0164](https://doi.org/10.2136/sssaj2007.0164)
- Álvaro-Fuentes J, Plaza-Bonilla D, Arrúe JL, Lampurlanés J, Cantero-Martínez C 2014 Soil organic carbon storage in a no-tillage chronosequence under Mediterranean conditions. *Plant Soil* 376, 31–41. doi: [10.1007/s11104-012-1167-x](https://doi.org/10.1007/s11104-012-1167-x)
- Annabi M, Bahri H, M'hamed HC 2018 Integrating future climate change, CO<sub>2</sub> increase and technology progress on wheat production in Northern Tunisia, in *Recent Advances in Environmental Science from the Euro-Mediterranean and Surrounding Regions. Proceedings of Euro-Mediterranean Conference for Environmental Integration (EMCEI-1), Tunisia 2017*, eds. Kallel A, Ksibi M, Ben Dhia H, Khélifi N (Springer International Publishing), 69–70. doi: [10.1007/978-3-319-70548-4](https://doi.org/10.1007/978-3-319-70548-4)
- Arnaez J, Lasanta T, Errea MP, Ortigosa L 2011 Land abandonment, landscape evolution, and soil erosion in a Spanish Mediterranean mountain region: The case of Camero Viejo. *L. Degrad. Dev.* 22, 537–550. doi: [10.1002/ldr.1032](https://doi.org/10.1002/ldr.1032)
- Asseng S, Martre P, Maiorano A, Rötter RP, O'Leary GJ et al. 2019 Climate change impact and adaptation for wheat protein. *Glob. Chang. Biol.* 25, 155–173. doi: [10.1111/gcb.14481](https://doi.org/10.1111/gcb.14481)
- Ayars JE, Fulton A, Taylor B 2015 Subsurface drip irrigation in California—Here to stay? *Agric. Water Manag.* 157, 39–47. doi: [10.1016/j.agwat.2015.01.001](https://doi.org/10.1016/j.agwat.2015.01.001)
- Bajocco S, de Angelis A, Perini L, Ferrara A, Salvati L 2012 The impact of Land Use/Land Cover Changes on land degradation dynamics: A Mediterranean case study. *Environ. Manage.* 49, 980–989. doi: [10.1007/s00267-012-9831-8](https://doi.org/10.1007/s00267-012-9831-8)
- Barth G, Von Tucher S, Schmidhalter U 2008 Effectiveness of 3,4-Dimethylpyrazole Phosphate as Nitrification Inhibitor in Soil as Influenced by Inhibitor Concentration, Application Form, and Soil Matric Potential. *Pedosphere* 18, 378–385. doi: [10.1016/s1002-0160\(08\)60028-4](https://doi.org/10.1016/s1002-0160(08)60028-4)
- Barton CM, Ullah IIT, Bergin SM 2010 Land use, water and Mediterranean landscapes: Modelling long-term dynamics of complex socio-ecological systems. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 368, 5275–5297. doi: [10.1098/rsta.2010.0193](https://doi.org/10.1098/rsta.2010.0193)
- Barton CM, Ullah IIT, Bergin SM, Sarjoughian HS, Mayer GR et al. 2016 Experimental socioecology: Integrative science for anthropocene landscape dynamics. *Anthropocene* 13, 34–45. doi: [10.1016/j.ancene.2015.12.004](https://doi.org/10.1016/j.ancene.2015.12.004)
- Basso L, Hendriks IE, Duarte CM 2015 Juvenile pen shells (*Pinna nobilis*) tolerate acidification but are vulnerable to warming. *Estuaries and Coasts* 38, 1976–1985. doi: [10.1007/s12237-015-9948-0](https://doi.org/10.1007/s12237-015-9948-0)
- Bebber DP, Holmes T, Gurr SJ 2014 The global spread of crop pests and pathogens. *Glob. Ecol. Biogeogr.* 23, 1398–1407. doi: [10.1111/geb.12214](https://doi.org/10.1111/geb.12214)
- Bebber DP, Ramotowski MAT, Gurr SJ 2013 Crop pests and pathogens move polewards in a warming world. *Nat. Clim. Chang.* 3, 985–988. doi: [10.1038/nclimate1990](https://doi.org/10.1038/nclimate1990)
- Behrenfeld MJ, O'Malley RT, Siegel DA, McClain CR, Sarmiento JL et al. 2006 Climate-driven trends in contemporary ocean productivity. *Nature* 444,

- 752–755. doi: [10.1038/nature05317](https://doi.org/10.1038/nature05317)
- Benhabib O, Yazar A, Qadir M, Lourenço E, Jacobsen S-E 2014 How Can We Improve Mediterranean Cropping Systems? *J. Agron. Crop Sci.* 200, 325–332. doi: [10.1111/jac.12066](https://doi.org/10.1111/jac.12066)
- Berge JA, Bjerkeng B, Pettersen O, Schaanning MT, Øxnevad S 2006 Effects of increased sea water concentrations of CO<sub>2</sub> on growth of the bivalve *Mytilus edulis* L. *Chemosphere* 62, 681–687. doi: [10.1016/j.chemosphere.2005.04.111](https://doi.org/10.1016/j.chemosphere.2005.04.111)
- Bernabucci U, Basiricò L, Morera P, Dipasquale D, Vitali A et al. 2015 Effect of summer season on milk protein fractions in Holstein cows. *J. Dairy Sci.* 98, 1815–1827. doi: [10.3168/jds.2014-8788](https://doi.org/10.3168/jds.2014-8788)
- Bernabucci U, Lacetera N, Baumgard LH, Rhoads RP, Ronchi B et al. 2010 Metabolic and hormonal acclimation to heat stress in domesticated ruminants. *Animal* 4, 1167–1183. doi: [10.1017/S175173111000090X](https://doi.org/10.1017/S175173111000090X)
- Bhakta I, Phadikar S, Majumder K 2019 State-of-the-art technologies in precision agriculture: a systematic review. *J. Sci. Food Agric.* 99, 4878–4888. doi: [10.1002/jsfa.9693](https://doi.org/10.1002/jsfa.9693)
- Bird DN, Benabdallah S, Gouda N, Hummel F, Koebel J et al. 2016 Modelling climate change impacts on and adaptation strategies for agriculture in Sardinia and Tunisia using AquaCrop and value-at-risk. *Sci. Total Environ.* 543, 1019–1027. doi: [10.1016/j.scitotenv.2015.07.035](https://doi.org/10.1016/j.scitotenv.2015.07.035)
- Blauhut V, Gudmundsson L, Stahl K 2015 Towards pan-European drought risk maps: quantifying the link between drought indices and reported drought impacts. *Environ. Res. Lett.* 10, 14008. doi: [10.1088/1748-9326/10/1/014008](https://doi.org/10.1088/1748-9326/10/1/014008)
- Blavet D, de Noni G, le Bissonnais Y, Leonard M, Maillo L et al. 2009 Effect of land use and management on the early stages of soil water erosion in French Mediterranean vineyards. *Soil Tillage Res.* 106, 124–136. doi: [10.1016/j.still.2009.04.010](https://doi.org/10.1016/j.still.2009.04.010)
- Boellstorff D, Benito G 2005 Impacts of set-aside policy on the risk of soil erosion in central Spain. *Agric. Ecosyst. Environ.* 107, 231–243. doi: [10.1016/j.agee.2004.11.002](https://doi.org/10.1016/j.agee.2004.11.002)
- Bonaccio M, di Castelnuovo A, Bonanni A, Costanzo S, de Lucia F et al. 2014 Decline of the Mediterranean diet at a time of economic crisis. Results from the Moli-sani study. *Nutr. Metab. Cardiovasc. Dis.* 24, 853–860. doi: [10.1016/j.numecd.2014.02.014](https://doi.org/10.1016/j.numecd.2014.02.014)
- Bonaccio M, Iacoviello L, de Gaetano G 2012 The Mediterranean diet: The reasons for a success. *Thromb. Res.* 129, 401–404. doi: [10.1016/j.thromres.2011.10.018](https://doi.org/10.1016/j.thromres.2011.10.018)
- Bowman DMJS, Williamson GJ, Abatzoglou JT, Kolden CA, Cochrane MA et al. 2017 Human exposure and sensitivity to globally extreme wildfire events. *Nat. Ecol. Evol.* 1, 1–6. doi: [10.1038/s41559-016-0058](https://doi.org/10.1038/s41559-016-0058)
- Buontempo C, Hutjes R, Beavis P, Berckmans J, Cagnazzo C et al. 2020 Fostering the development of climate services through Copernicus Climate Change Service (C3S) for agriculture applications. *Weather Clim. Extrem.* 27, 100226. doi: [10.1016/j.wace.2019.100226](https://doi.org/10.1016/j.wace.2019.100226)
- Bureau JC, Swinnen J 2018 EU policies and global food security. *Glob. Food Sec.* 16, 106–115. doi: [10.1016/j.gfs.2017.12.001](https://doi.org/10.1016/j.gfs.2017.12.001)
- Burge CA, Eakin CM, Friedman CS, Froelich B, Hershberger PK et al. 2014 Climate change influences on marine infectious diseases: implications for management and society. *Ann. Rev. Mar. Sci.* 6, 249–277. doi: [10.1146/annurev-marine-010213-135029](https://doi.org/10.1146/annurev-marine-010213-135029)
- Cantero-Martínez C, Angás P, Lampurlanés J 2007 Long-term yield and water use efficiency under various tillage systems in Mediterranean rainfed conditions. *Ann. Appl. Biol.* 150, 293–305. doi: [10.1111/j.1744-7348.2007.00142.x](https://doi.org/10.1111/j.1744-7348.2007.00142.x)
- Capone R, El Bilali H, Elferchichi A, Lamaddalena N, Lamberti L 2012 Natural resources and food in the Mediterranean, in *MediTERRA 2012: The mediterranean diet for sustainable regional development*, ed. CIHEAM (Paris: Presses de Sciences Po), 171–193.
- Caracciolo F, Gotor E, Santeramo FG 2014 European Common Agricultural Policy Impacts On Developing Countries Commodities Prices. *Reg. Sect. Econ. Stud.* 14, 2.
- Caracciolo F, Santeramo FG 2013 Price Trends and Income Inequalities: Will Sub-Saharan Africa Reduce the Gap? *African Dev. Rev.* 25, 42–54. doi: [10.1111/j.1467-8268.2013.12012.x](https://doi.org/10.1111/j.1467-8268.2013.12012.x)
- Carranca C, Oliveira A, Pampulha E, Torres MO 2009 Temporal dynamics of soil nitrogen, carbon and microbial activity in conservative and disturbed fields amended with mature white lupine and oat residues. *Geoderma* 151, 50–59. doi: [10.1016/j.geoderma.2009.03.012](https://doi.org/10.1016/j.geoderma.2009.03.012)
- Carranza-Gallego G, Guzmán GI, García-Ruiz R, González de Molina M, Aguilera E 2018 Contribution of old wheat varieties to climate change mitigation under contrasting managements and rainfed Mediterranean conditions. *J. Clean. Prod.* 195, 111–121. doi: [10.1016/j.jclepro.2018.05.188](https://doi.org/10.1016/j.jclepro.2018.05.188)
- Cayuela ML, Aguilera E, Sanz-Cobena A, Adams DC, Abalos D et al. 2017 Direct nitrous oxide emissions in Mediterranean climate cropping systems: Emission factors based on a meta-analysis of available measurement data. *Agric. Ecosyst. Environ.* 238, 25–35. doi: [10.1016/j.agee.2016.10.006](https://doi.org/10.1016/j.agee.2016.10.006)
- Ceglar A, Toreti A, Zampieri M, Manstretta V, Bettati T et al. 2020 Clisagri: An R package for agro-climate services. *Clim. Serv.* 20, 100197. doi: [10.1016/j.cliser.2020.100197](https://doi.org/10.1016/j.cliser.2020.100197)
- Ceglar A, Zampieri M, Toreti A, Dentener FJ 2019 Observed Northward Migration of Agro-Climatic Zones in Europe Will Further Accelerate Under Climate Change. *Earth's Futur.* 7, 1088–1101. doi: [10.1029/2019ef001178](https://doi.org/10.1029/2019ef001178)



- Chabbi A, Lehmann J, Ciais P, Loescher HW, Cotrufo MF et al. 2017 Aligning agriculture and climate policy. *Nat. Clim. Chang.* 7, 307–309. doi: [10.1038/nclimate3286](https://doi.org/10.1038/nclimate3286)
- Chatterjee N, Nair PKR, Chakraborty S, Nair VD 2018 Changes in soil carbon stocks across the Forest-Agroforest-Agriculture/Pasture continuum in various agroecological regions: A meta-analysis. *Agric. Ecosyst. Environ.* 266, 55–67. doi: [10.1016/j.agee.2018.07.014](https://doi.org/10.1016/j.agee.2018.07.014)
- Chatzopoulos T, Pérez Domínguez I, Zampieri M, Toreti A 2019 Climate extremes and agricultural commodity markets: A global economic analysis of regionally simulated events. *Weather Clim. Extrem.*, 100193. doi: [10.1016/j.wace.2019.100193](https://doi.org/10.1016/j.wace.2019.100193)
- Chen C-C, McCarl B, Chang C-C 2012 Climate change, sea level rise and rice: Global market implications. *Clim. Change* 110, 543–560. doi: [10.1007/s10584-011-0074-0](https://doi.org/10.1007/s10584-011-0074-0)
- Cheung WWL, Frölicher TL, Asch RG, Jones MC, Pinsky ML et al. 2016 Building confidence in projections of the responses of living marine resources to climate change. *ICES J. Mar. Sci. J. du Cons.* 73, 1283–1296. doi: [10.1093/icesjms/fsv250](https://doi.org/10.1093/icesjms/fsv250)
- Chiti T, Gardin L, Perugini L, Quarantino R, Vaccari FP et al. 2012 Soil organic carbon stock assessment for the different cropland land uses in Italy. *Biol. Fertil. Soils* 48, 9–17. doi: [10.1007/s00374-011-0599-4](https://doi.org/10.1007/s00374-011-0599-4)
- Chourghal N, Lhomme JP, Huard F, Aidaoui A 2016 Climate change in Algeria and its impact on durum wheat. *Reg. Environ. Chang.* 16, 1623–1634. doi: [10.1007/s10113-015-0889-8](https://doi.org/10.1007/s10113-015-0889-8)
- CIHEAM, FAO 2015 Mediterranean food consumption patterns: diet, environment, society, economy and health. A White Paper.
- Cioffi A, Dell'Aquila C 2004 The effects of trade policies for fresh fruit and vegetables of the European Union. *Food Policy* 29, 169–185. doi: [10.1016/s0306-9192\(04\)00010-7](https://doi.org/10.1016/s0306-9192(04)00010-7)
- Cioffi A, Santeramo FG, Vitale CD 2011 The price stabilization effects of the EU entry price scheme for fruit and vegetables. *Agric. Econ.* 42, 405–418. doi: [10.1111/j.1574-0862.2010.00526.x](https://doi.org/10.1111/j.1574-0862.2010.00526.x)
- Cooper J, Baranski M, Stewart G, Nobel-de Lange M, Bàrberi P et al. 2016 Shallow non-inversion tillage in organic farming maintains crop yields and increases soil C stocks: a meta-analysis. *Agron. Sustain. Dev.* 36, 22. doi: [10.1007/s13593-016-0354-1](https://doi.org/10.1007/s13593-016-0354-1)
- de Rensis F, García-Ispierto I, López-Gatius F 2015 Seasonal heat stress: Clinical implications and hormone treatments for the fertility of dairy cows. *Theriogenology* 84, 659–666. doi: [10.1016/j.theriogenology.2015.04.021](https://doi.org/10.1016/j.theriogenology.2015.04.021)
- Debolini M, Valette E, François M, Chéry JP 2015 Mapping land use competition in the rural-urban fringe and future perspectives on land policies: A case study of Meknès (Morocco). *Land use policy* 47, 373–381. doi: [10.1016/j.landusepol.2015.01.035](https://doi.org/10.1016/j.landusepol.2015.01.035)
- del Pozo A, Brunel-Saldias N, Engler A, Ortega-Farías S, Acevedo-Opazo C et al. 2019 Climate change impacts and adaptation strategies of agriculture in Mediterranean-climate regions (MCRs). *Sustain.* 11, 2769. doi: [10.3390/su11102769](https://doi.org/10.3390/su11102769)
- del Pozo A, Yáñez A, Matus IA, Tapia G, Castillo D et al. 2016 Physiological Traits Associated with Wheat Yield Potential and Performance under Water-Stress in a Mediterranean Environment. *Front. Plant Sci.* 7, 987. doi: [10.3389/fpls.2016.00987](https://doi.org/10.3389/fpls.2016.00987)
- del Prado A, Merino P, Estavillo JM, Pinto M, González-Murua C 2006 N<sub>2</sub>O and NO emissions from different N sources and under a range of soil water contents. *Nutr. Cycl. Agroecosystems* 74, 229–243. doi: [10.1007/s10705-006-9001-6](https://doi.org/10.1007/s10705-006-9001-6)
- di Bene C, Tavarini S, Mazzoncini M, Angelini LG 2011 Changes in soil chemical parameters and organic matter balance after 13 years of ramie [*Boehmeria nivea* (L.) Gaud.] cultivation in the Mediterranean region. *Eur. J. Agron.* 35, 154–163. doi: [10.1016/j.eja.2011.05.007](https://doi.org/10.1016/j.eja.2011.05.007)
- di Giacomo G, Taglieri L 2009 Renewable energy benefits with conversion of woody residues to pellets. *Energy* 34, 724–731. doi: [10.1016/j.energy.2008.08.010](https://doi.org/10.1016/j.energy.2008.08.010)
- Díez-Palet I, Funes I, Savé R, Biel C, de Herralde F et al. 2019 Blooming under Mediterranean Climate: Estimating Cultivar-Specific Chill and Heat Requirements of Almond and Apple Trees Using a Statistical Approach. *Agronomy* 9, 760. doi: [10.3390/agronomy9110760](https://doi.org/10.3390/agronomy9110760)
- Doney SC, Fabry VJ, Feely RA, Kleypas JA 2009 Ocean acidification: The other CO<sub>2</sub> problem. *Ann. Rev. Mar. Sci.* 1, 169–192. doi: [10.1146/annurev.marine.010908.163834](https://doi.org/10.1146/annurev.marine.010908.163834)
- Drine I 2011 Climate Change Compounding Risks in North Africa.
- EEA 2019 Climate change adaptation in the agriculture sector in Europe.
- Fader M, Shi S, Von Bloh W, Bondeau A, Cramer W 2016 Mediterranean irrigation under climate change: More efficient irrigation needed to compensate for increases in irrigation water requirements. *Hydrol. Earth Syst. Sci.* 20, 953–973. doi: [10.5194/hess-20-953-2016](https://doi.org/10.5194/hess-20-953-2016)
- Falcucci A, Maiorano L, Boitani L 2007 Changes in land-use/land-cover patterns in Italy and their implications for biodiversity conservation. *Landsc. Ecol.* 22, 617–631. doi: [10.1007/s10980-006-9056-4](https://doi.org/10.1007/s10980-006-9056-4)
- Fantappiè M, L'Abate G, Costantini EAC 2011 The influence of climate change on the soil organic carbon content in Italy from 1961 to 2008. *Geomorphology* 135, 343–352. doi: [10.1016/j.geomorph.2011.02.006](https://doi.org/10.1016/j.geomorph.2011.02.006)
- FAO 2017 FAOSTAT.
- FAO 2018 The State of Mediterranean and Black Sea Fisheries. Rome.

- Feyen L, Ciscar JC, Gosling S, Ibarreta D, Soria A 2020 Climate change impacts and adaptation in Europe. JRC PESETA IV final report. Luxembourg. doi: [10.2760/171121](https://doi.org/10.2760/171121)
- Flecha S, Pérez FF, García-Lafuente J, Sammartino S, Ríos AF et al. 2015 Trends of pH decrease in the Mediterranean Sea through high frequency observational data: indication of ocean acidification in the basin. *Sci. Rep.* 5, 16770. doi: [10.1038/srep16770](https://doi.org/10.1038/srep16770)
- Fontana G, Toreti A, Ceglar A, de Sanctis G 2015 Early heat waves over Italy and their impacts on durum wheat yields. *Nat. Hazards Earth Syst. Sci.* 15, 1631–1637. doi: [10.5194/nhess-15-1631-2015](https://doi.org/10.5194/nhess-15-1631-2015)
- Funes I, Aranda X, Biel C, Carbó J, Camps F et al. 2016 Future climate change impacts on apple flowering date in a Mediterranean subbasin. *Agric. Water Manag.* 164, 19–27. doi: [10.1016/j.agwat.2015.06.013](https://doi.org/10.1016/j.agwat.2015.06.013)
- Funes I, Savé R, Rovira P, Molowny-Horas R, Alcañiz JM et al. 2019 Agricultural soil organic carbon stocks in the north-eastern Iberian Peninsula: Drivers and spatial variability. *Sci. Total Environ.* 668, 283–294. doi: [10.1016/j.scitotenv.2019.02.317](https://doi.org/10.1016/j.scitotenv.2019.02.317)
- Galli A, Halle M, Grunewald N 2015 Physical limits to resource access and utilisation and their economic implications in Mediterranean economies. *Environ. Sci. Policy* 51, 125–136. doi: [10.1016/j.envsci.2015.04.002](https://doi.org/10.1016/j.envsci.2015.04.002)
- Galli A, Iha K, Halle M, El Bilali H, Grunewald N et al. 2017 Mediterranean countries' food consumption and sourcing patterns: An Ecological Footprint viewpoint. *Sci. Total Environ.* 578, 383–391. doi: [10.1016/j.scitotenv.2016.10.191](https://doi.org/10.1016/j.scitotenv.2016.10.191)
- Gantner V, Bobic T, Gantner R, Gregic M, Kuterovac K et al. 2017 Differences in response to heat stress due to production level and breed of dairy cows. *Int. J. Biometeorol.* 61, 1675–1685. doi: [10.1007/s00484-017-1348-7](https://doi.org/10.1007/s00484-017-1348-7)
- García-Marco S, Ravella SR, Chadwick DR, Vallejo A, Gregory AS et al. 2014 Ranking factors affecting emissions of GHG from incubated agricultural soils. *Eur. J. Soil Sci.* 65, 573–583. doi: [10.1111/ejss.12143](https://doi.org/10.1111/ejss.12143)
- García Martínez M, Poole N 2004 The development of private fresh produce safety standards: Implications for developing Mediterranean exporting countries. *Food Policy* 29, 229–255. doi: [10.1016/j.foodpol.2004.04.002](https://doi.org/10.1016/j.foodpol.2004.04.002)
- Gazeau F, Alliouane S, Bock C, Bramanti L, López Correa M et al. 2014 Impact of ocean acidification and warming on the Mediterranean mussel (*Mytilus galloprovincialis*). *Front. Mar. Sci.* 1, 62. doi: [10.3389/fmars.2014.00062](https://doi.org/10.3389/fmars.2014.00062)
- Genua-Olmedo A, Alcaraz C, Caiola N, Ibáñez C 2016 Sea level rise impacts on rice production: The Ebro Delta as an example. *Sci. Total Environ.* 571, 1200–1210. doi: [10.1016/j.scitotenv.2016.07.136](https://doi.org/10.1016/j.scitotenv.2016.07.136)
- Gilsanz C, Báez D, Misselbrook TH, Dhanoa MS, Cárdenas LM 2016 Development of emission factors and efficiency of two nitrification inhibitors, DCD and DMPP. *Agric. Ecosyst. Environ.* 216, 1–8. doi: [10.1016/j.agee.2015.09.030](https://doi.org/10.1016/j.agee.2015.09.030)
- Girard C, Pulido-Velazquez M, Rinaudo J-D, Pagé C, Caballero Y 2015 Integrating top-down and bottom-up approaches to design global change adaptation at the river basin scale. *Glob. Environ. Chang.* 34, 132–146. doi: [10.1016/j.gloenvcha.2015.07.002](https://doi.org/10.1016/j.gloenvcha.2015.07.002)
- Gómez-Luciano CA, de Aguiar LK, Vriesekoop F, Urbano B 2019 Consumers' willingness to purchase three alternatives to meat proteins in the United Kingdom, Spain, Brazil and the Dominican Republic. *Food Qual. Prefer.* 78, 103732. doi: [10.1016/j.foodqual.2019.103732](https://doi.org/10.1016/j.foodqual.2019.103732)
- Grasso M, Feola G 2012 Mediterranean agriculture under climate change: Adaptive capacity, adaptation, and ethics. *Reg. Environ. Chang.* 12, 607–618. doi: [10.1007/s10113-011-0274-1](https://doi.org/10.1007/s10113-011-0274-1)
- Guardia G, Tellez-Rio A, García-Marco S, Martin-Lammerding D, Tenorio JL et al. 2016 Effect of tillage and crop (cereal versus legume) on greenhouse gas emissions and Global Warming Potential in a non-irrigated Mediterranean field. *Agric. Ecosyst. Environ.* 221, 187–197. doi: [10.1016/j.agee.2016.01.047](https://doi.org/10.1016/j.agee.2016.01.047)
- Harmanny KS, Malek Ž 2019 Adaptations in irrigated agriculture in the Mediterranean region: an overview and spatial analysis of implemented strategies. *Reg. Environ. Chang.* 19, 1401–1416. doi: [10.1007/s10113-019-01494-8](https://doi.org/10.1007/s10113-019-01494-8)
- Hatfield JL, Dold C 2019 Water-use efficiency: Advances and challenges in a changing climate. *Front. Plant Sci.* 10, 103. doi: [10.3389/fpls.2019.00103](https://doi.org/10.3389/fpls.2019.00103)
- Havenhand JN, Buttler F-R, Thorndyke MC, Williamson JE 2008 Near-future levels of ocean acidification reduce fertilization success in a sea urchin. *Curr. Biol.* 18, R651–R652. doi: [10.1016/j.cub.2008.06.015](https://doi.org/10.1016/j.cub.2008.06.015)
- Hendriks IE, Duarte CM, Álvarez M 2010 Vulnerability of marine biodiversity to ocean acidification: A meta-analysis. *Estuar. Coast. Shelf Sci.* 86, 157–164. doi: [10.1016/j.ecss.2009.11.022](https://doi.org/10.1016/j.ecss.2009.11.022)
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P et al. 2007 Coral reefs under rapid climate change and ocean acidification. *Science (80-. J.)* 318, 1737–1742. doi: [10.1126/science.1152509](https://doi.org/10.1126/science.1152509)
- Iglesias A, Garrote L 2015 Adaptation strategies for agricultural water management under climate change in Europe. *Agric. Water Manag.* 155, 113–124. doi: [10.1016/j.agwat.2015.03.014](https://doi.org/10.1016/j.agwat.2015.03.014)
- Jordán A, Zavala LM, Gil J 2010 Effects of mulching on soil physical properties and runoff under semi-arid conditions in southern Spain. *Catena* 81, 77–85. doi: [10.1016/j.catena.2010.01.007](https://doi.org/10.1016/j.catena.2010.01.007)
- Karamesouti M, Detsis V, Kounalaki A, Vasilioi P,

- Salvati L et al. 2015 Land-use and land degradation processes affecting soil resources: Evidence from a traditional Mediterranean cropland (Greece). *Catena* 132, 45–55. doi: [10.1016/j.catena.2015.04.010](https://doi.org/10.1016/j.catena.2015.04.010)
- Kassam A, Friedrich T, Derpsch R, Lahmar R, Mrabet R et al. 2012 Conservation agriculture in the dry Mediterranean climate. *F. Crop. Res.* 132, 7–17. doi: [10.1016/j.fcr.2012.02.023](https://doi.org/10.1016/j.fcr.2012.02.023)
- Kauppi PE, Ausubel JH, Fang J, Mather AS, Sedjo RA et al. 2006 Returning forests analyzed with the forest identity. *Proc. Natl. Acad. Sci. U. S. A.* 103, 17574–17579. doi: [10.1073/pnas.0608343103](https://doi.org/10.1073/pnas.0608343103)
- Kimball BA 2016 Crop responses to elevated CO<sub>2</sub> and interactions with H<sub>2</sub>O, N, and temperature. *Curr. Opin. Plant Biol.* 31, 36–43. doi: [10.1016/j.pbi.2016.03.006](https://doi.org/10.1016/j.pbi.2016.03.006)
- King T, Osmond-McLeod MJ, Duffy LL 2018 Nanotechnology in the food sector and potential applications for the poultry industry. *Trends Food Sci. Technol.* 72, 62–73. doi: [10.1016/j.tifs.2017.11.015](https://doi.org/10.1016/j.tifs.2017.11.015)
- Kool DM, Dolfing J, Wrage N, Van Groenigen JW 2011 Nitrifier denitrification as a distinct and significant source of nitrous oxide from soil. *Soil Biol. Biochem.* 43, 174–178. doi: [10.1016/j.soilbio.2010.09.030](https://doi.org/10.1016/j.soilbio.2010.09.030)
- Kurihara H 2008 Effects of CO<sub>2</sub>-driven ocean acidification on the early developmental stages of invertebrates. *Mar. Ecol. Prog. Ser.* 373, 275–284. doi: [10.3354/meps07802](https://doi.org/10.3354/meps07802)
- Kurihara H, Kato S, Ishimatsu A 2007 Effects of increased seawater pCO<sub>2</sub> on early development of the oyster *Crassostrea gigas*. *Aquat. Biol.* 1, 91–98. doi: [10.3354/ab00009](https://doi.org/10.3354/ab00009)
- Lagacherie P, Álvaro-Fuentes J, Annabi M, Bernoux M, Bouarfa S et al. 2018 Managing Mediterranean soil resources under global change: expected trends and mitigation strategies. *Reg. Environ. Chang.* 18, 663–675. doi: [10.1007/s10113-017-1239-9](https://doi.org/10.1007/s10113-017-1239-9)
- Lahmar R, Ruellan A 2007 Soil degradation in the Mediterranean region and cooperative strategies. *Cah. Agric.* 16, 318–323. doi: [10.1684/agr.2007.0119](https://doi.org/10.1684/agr.2007.0119)
- Larson BA, Nicolaidis E, Al Zu'bi B, Sukkar N, Laraki K et al. 2002 The Impact of Environmental Regulations on Exports: Case Study Results from Cyprus, Jordan, Morocco, Syria, Tunisia, and Turkey. *World Dev.* 30, 1057–1072. doi: [10.1016/s0305-750x\(02\)00023-2](https://doi.org/10.1016/s0305-750x(02)00023-2)
- Lassaletta L, Billen G, Grizzetti B, Garnier J, Leach AM et al. 2014 Food and feed trade as a driver in the global nitrogen cycle: 50-year trends. *Biogeochemistry* 118, 225–241. doi: [10.1007/s10533-013-9923-4](https://doi.org/10.1007/s10533-013-9923-4)
- Lassaletta L, Romero E, Billen G, Garnier J, García-Gómez H et al. 2012 Spatialized N budgets in a large agricultural Mediterranean watershed: High loading and low transfer. *Biogeosciences* 9, 57–70. doi: [10.5194/bg-9-57-2012](https://doi.org/10.5194/bg-9-57-2012)
- Lesk C, Rowhani P, Ramankutty N 2016 Influence of extreme weather disasters on global crop production. *Nature* 529, 84–87. doi: [10.1038/nature16467](https://doi.org/10.1038/nature16467)
- Liebig MA, Franzluebbers AJ, Alvarez C, Chiesa TD, Lewczuk N et al. 2016 MAGNet: An international network to foster mitigation of agricultural greenhouse gases. *Carbon Manag.* 7, 243–248. doi: [10.1080/17583004.2016.1180586](https://doi.org/10.1080/17583004.2016.1180586)
- Lin BB 2011 Resilience in Agriculture through Crop Diversification: Adaptive Management for Environmental Change. *Bioscience* 61, 183–193. doi: [10.1525/bio.2011.61.3.4](https://doi.org/10.1525/bio.2011.61.3.4)
- López-Bellido RJ, Fontán JM, López-Bellido FJ, López-Bellido L 2010 Carbon Sequestration by Tillage, Rotation, and Nitrogen Fertilization in a Mediterranean Vertisol. *Agron. J.* 102, 310. doi: [10.2134/agronj2009.0165](https://doi.org/10.2134/agronj2009.0165)
- Maetens W, Vanmaercke M, Poesen J, Jankauskas B, Jankauskiene G et al. 2012 Effects of land use on annual runoff and soil loss in Europe and the Mediterranean. *Prog. Phys. Geogr. Earth Environ.* 36, 599–653. doi: [10.1177/0309133312451303](https://doi.org/10.1177/0309133312451303)
- Malek Ž, Verburg PH, Geijzendorffer IR, Bondeau A, Cramer W 2018 Global change effects on land management in the Mediterranean region. *Glob. Environ. Chang.* 50, 238–254. doi: [10.1016/j.gloenvcha.2018.04.007](https://doi.org/10.1016/j.gloenvcha.2018.04.007)
- Marquez-Ramos L, Martinez-Gomez VD 2016 On the effect of EU trade preferences: evidence for monthly exports of fruit and vegetables from Morocco. *New Medit* 15, 14–21.
- Martiniello P, Teixeira da Silva JA 2011 Physiological and Bioagronomical Aspects Involved in Growth and Yield Components of Cultivated Forage Species in Mediterranean Environments: A Review. *Eur. J. Plant Sci. Biotechnol.* 5, 64–98.
- Masri Z, Ryan J 2006 Soil organic matter and related physical properties in a Mediterranean wheat-based rotation trial. *Soil Tillage Res.* 87, 146–154. doi: [10.1016/j.still.2005.03.003](https://doi.org/10.1016/j.still.2005.03.003)
- Mavromatis T 2015 Crop-climate relationships of cereals in Greece and the impacts of recent climate trends. *Theor. Appl. Climatol.* 120, 417–432. doi: [10.1007/s00704-014-1179-y](https://doi.org/10.1007/s00704-014-1179-y)
- McCarl BA, Musumba M, Smith JB, Kirshen P, Jones R et al. 2015 Climate change vulnerability and adaptation strategies in Egypt's agricultural sector. *Mitig. Adapt. Strateg. Glob. Chang.* 20, 1097–1109. doi: [10.1007/s11027-013-9520-9](https://doi.org/10.1007/s11027-013-9520-9)
- Medina A, Akbar A, Baazeem A, Rodriguez A, Magan N 2017 Climate change, food security and mycotoxins: Do we know enough? *Fungal Biol. Rev.* 31, 143–154. doi: [10.1016/j.fbr.2017.04.002](https://doi.org/10.1016/j.fbr.2017.04.002)
- Meersmans J, Martin MP, Lacarce E, de Baets S, Jolivet C et al. 2012 A high resolution map of French soil organic carbon. *Agron. Sustain. Dev.* 32, 841–851.

- doi: [10.1007/s13593-012-0086-9](https://doi.org/10.1007/s13593-012-0086-9)
- Meijide A, Cárdenas LM, Sánchez-Martín L, Vallejo A 2010 Carbon dioxide and methane fluxes from a barley field amended with organic fertilizers under Mediterranean climatic conditions. *Plant Soil* 328, 353–367. doi: [10.1007/s11104-009-0114-y](https://doi.org/10.1007/s11104-009-0114-y)
- Meijide A, García-Torres L, Arce A, Vallejo A 2009 Nitrogen oxide emissions affected by organic fertilization in a non-irrigated Mediterranean barley field. *Agric. Ecosyst. Environ.* 132, 106–115. doi: [10.1016/j.agee.2009.03.005](https://doi.org/10.1016/j.agee.2009.03.005)
- Minasny B, Malone BP, McBratney AB, Angers DA, Arrouays D et al. 2017 Soil carbon 4 per mille. *Geoderma* 292, 59–86. doi: [10.1016/j.geoderma.2017.01.002](https://doi.org/10.1016/j.geoderma.2017.01.002)
- Molina-Alcaide E, Yáñez-Ruiz DR 2008 Potential use of olive by-products in ruminant feeding: A review. *Anim. Feed Sci. Technol.* 147, 247–264. doi: [10.1016/j.anifeedsci.2007.09.021](https://doi.org/10.1016/j.anifeedsci.2007.09.021)
- Moritz MA, Battlori E, Bradstock RA, Gill AM, Handmer J et al. 2014 Learning to coexist with wildfire. *Nature* 515, 58–66. doi: [10.1038/nature13946](https://doi.org/10.1038/nature13946)
- Mougou R, Mansour M, Iglesias A, Chebbi RZ, Battagliani A 2011 Climate change and agricultural vulnerability: A case study of rain-fed wheat in Kairouan, Central Tunisia. *Reg. Environ. Chang.* 11, 137–142. doi: [10.1007/s10113-010-0179-4](https://doi.org/10.1007/s10113-010-0179-4)
- Mrabet R 2008 *No-Tillage systems for sustainable dryland agriculture in Morocco*. INRA Publication, Fanigraph Edition.
- Mueller ND, Gerber JS, Johnston M, Ray DK, Ramankutty N et al. 2012 Closing yield gaps through nutrient and water management. *Nature* 490, 254–257. doi: [10.1038/nature11420](https://doi.org/10.1038/nature11420)
- Niedertscheider M, Erb K 2014 Land system change in Italy from 1884 to 2007: Analysing the North-South divergence on the basis of an integrated indicator framework. *Land use policy* 39, 366–375. doi: [10.1016/j.landusepol.2014.01.015](https://doi.org/10.1016/j.landusepol.2014.01.015)
- Norton J, Ouyang Y 2019 Controls and adaptive management of nitrification in agricultural soils. *Front. Microbiol.* 10. doi: [10.3389/fmicb.2019.01931](https://doi.org/10.3389/fmicb.2019.01931)
- Padilla FM, Vidal B, Sánchez J, Pugnaire FI 2010 Land-use changes and carbon sequestration through the twentieth century in a Mediterranean mountain ecosystem: implications for land management. *J. Environ. Manage.* 91, 2688–2695. doi: [10.1016/j.jenvman.2010.07.031](https://doi.org/10.1016/j.jenvman.2010.07.031)
- Pala M, Oweis T, Benli B, de Pauw E, El Mourid M et al. 2011 Assessment of wheat yield gap in the Mediterranean: case studies from Morocco, Syria and Turkey. *ICARDA Rep.*, 36.
- Parker LM, Ross PM, O'Connor WA 2009 The effect of ocean acidification and temperature on the fertilization and embryonic development of the Sydney rock oyster *Saccostrea glomerata* (Gould 1850). *Glob. Chang. Biol.* 15, 2123–2136. doi: [10.1111/j.1365-2486.2009.01895.x](https://doi.org/10.1111/j.1365-2486.2009.01895.x)
- Parker LM, Ross PM, O'Connor WA 2010 Comparing the effect of elevated pCO<sub>2</sub> and temperature on the fertilization and early development of two species of oysters. *Mar. Biol.* 157, 2435–2452. doi: [10.1007/s00227-010-1508-3](https://doi.org/10.1007/s00227-010-1508-3)
- Pauly D, Zeller D 2016 Catch reconstructions reveal that global marine fisheries catches are higher than reported and declining. *Nat. Commun.* 7, 10244. doi: [10.1038/ncomms10244](https://doi.org/10.1038/ncomms10244)
- Peigné J, Ball BC, Roger-Estrade J, David C 2007 Is conservation tillage suitable for organic farming? A review. *Soil Use Manag.* 23, 129–144. doi: [10.1111/j.1475-2743.2006.00082.x](https://doi.org/10.1111/j.1475-2743.2006.00082.x)
- Peigné J, Lefevre V, Vian JF, Fleury P 2015 Conservation agriculture in organic farming: experiences, challenges and opportunities in Europe, in *Conservation Agriculture* (Switzerland: Springer International Publishing), 559–578. doi: [10.1007/978-3-319-11620-4\\_21](https://doi.org/10.1007/978-3-319-11620-4_21)
- Pellegrino E, di Bene C, Tozzini C, Bonari E 2011 Impact on soil quality of a 10-year-old short-rotation coppice poplar stand compared with intensive agricultural and uncultivated systems in a Mediterranean area. *Agric. Ecosyst. Environ.* 140, 245–254. doi: [10.1016/j.agee.2010.12.011](https://doi.org/10.1016/j.agee.2010.12.011)
- Peltre C, Nyord T, Bruun S, Jensen LS, Magid J 2015 Repeated soil application of organic waste amendments reduces draught force and fuel consumption for soil tillage. *Agric. Ecosyst. Environ.* 211, 94–101. doi: [10.1016/j.agee.2015.06.004](https://doi.org/10.1016/j.agee.2015.06.004)
- Petit M, Le Grusse P 2018 Food and water security in the Mediterranean Basin, in *The Oxford handbook of food, water and society*, eds. Allan T, Bromwich B, Keulertz M, Colman A (Oxford, Royaume-Uni: Oxford University Press), 1–16.
- Phogat V, Pitt T, Cox JW, Šimůnek J, Skewes MA 2018 Soil water and salinity dynamics under sprinkler irrigated almond exposed to a varied salinity stress at different growth stages. *Agric. Water Manag.* 201, 70–82. doi: [10.1016/j.agwat.2018.01.018](https://doi.org/10.1016/j.agwat.2018.01.018)
- Plaza-Bonilla D, Álvaro-Fuentes J, Arrúe JL, Cantero-Martínez C 2014 Tillage and nitrogen fertilization effects on nitrous oxide yield-scaled emissions in a rainfed Mediterranean area. *Agric. Ecosyst. Environ.* 189, 43–52. doi: [10.1016/j.agee.2014.03.023](https://doi.org/10.1016/j.agee.2014.03.023)
- Plaza-Bonilla D, Cantero-Martínez C, Álvaro-Fuentes J 2010 Tillage effects on soil aggregation and soil organic carbon profile distribution under Mediterranean semi-arid conditions. *Soil Use Manag.* 26, 445–474. doi: [10.1111/j.1475-2743.2010.00298.x](https://doi.org/10.1111/j.1475-2743.2010.00298.x)
- Prado P, Roque A, Pérez J, Ibáñez C, Alcaraz C et al. 2016 Warming and acidification-mediated resilience to bacterial infection determine mortality of early *Ostrea edulis* life stages. *Mar. Ecol. Prog. Ser.* 545, 189–202. doi: [10.3354/meps11618](https://doi.org/10.3354/meps11618)
- Range P, Piló D, Ben-Hamadou R, Chícharo MA, Matias

- D et al. 2012 Seawater acidification by CO<sub>2</sub> in a coastal lagoon environment: Effects on life history traits of juvenile mussels *Mytilus galloprovincialis*. *J. Exp. Mar. Bio. Ecol.* 424–425, 89–98. doi: [10.1016/j.jembe.2012.05.010](https://doi.org/10.1016/j.jembe.2012.05.010)
- Reguant F, Savé R 2016 Disponibilidad alimentaria y desarrollo global sostenible, in *El sistema alimentario. Globalización sostenibilidad seguridad y cultura alimentaria*, eds. Colomer-Xena Y, Clotet-Ballús R, González-Vaqué L (Thomson Reuters Proview Aranzadi).
- Requier-Desjardins M 2010 Impacts des changements climatiques sur l'agriculture au Maroc et en Tunisie et priorités d'adaptation. Notes d'Analyse du CIHEAM (56).
- Robertson GP, Paul EA, Harwood RR 2000 Greenhouse gases in intensive agriculture: Contributions of individual gases to the radiative forcing of the atmosphere. *Science (80-. )*. 289, 1922–1925. doi: [10.1126/science.289.5486.1922](https://doi.org/10.1126/science.289.5486.1922)
- Romero E, Garnier J, Billen G, Peters F, Lassaletta L 2016 Water management practices exacerbate nitrogen retention in Mediterranean catchments. *Sci. Total Environ.* 573, 420–432. doi: [10.1016/j.scitotenv.2016.08.007](https://doi.org/10.1016/j.scitotenv.2016.08.007)
- Ronco P, Zennaro F, Torresan S, Critto A, Santini M et al. 2017 A risk assessment framework for irrigated agriculture under climate change. *Adv. Water Resour.* 110, 562–578. doi: [10.1016/j.advwatres.2017.08.003](https://doi.org/10.1016/j.advwatres.2017.08.003)
- Ruauudel H, Morrison-Métois S 2017 Responding to Refugee Crises in Developing Countries: What Can We Learn From Evaluations? Paris. doi: [10.1787/ae4362bd-en](https://doi.org/10.1787/ae4362bd-en)
- Ryan J, de Pauw E, Gomez H, Mrabet R 2006 Drylands of the Mediterranean Zone: Biophysical Resources and Cropping Systems, in *Dryland Agriculture*, eds. Peterson G, Unger P, Payne W, 577–624.
- Saadi S, Todorovic M, Tanasijevic L, Pereira LS, Pizigalli C et al. 2015 Climate change and Mediterranean agriculture: Impacts on winter wheat and tomato crop evapotranspiration, irrigation requirements and yield. *Agric. Water Manag.* 147, 103–115. doi: [10.1016/J.AGWAT.2014.05.008](https://doi.org/10.1016/J.AGWAT.2014.05.008)
- Saber N, Mrabet R 2002 Impact of no tillage and crop sequence on selected soil quality attributes of a vertic calcixeroll soil in Morocco. *Agronomie* 22, 451–459. doi: [10.1051/agro:2002026](https://doi.org/10.1051/agro:2002026)
- San-Miguel-Ayanz J, Moreno JM, Camia A 2013 Analysis of large fires in European Mediterranean landscapes: Lessons learned and perspectives. *For. Ecol. Manage.* 294, 11–22. doi: [10.1016/j.foreco.2012.10.050](https://doi.org/10.1016/j.foreco.2012.10.050)
- Sánchez-Martín L, Meijide A, García-Torres L, Vallejo A 2010 Combination of drip irrigation and organic fertilizer for mitigating emissions of nitrogen oxides in semiarid climate. *Agric. Ecosyst. Environ.* 137, 99–107. doi: [10.1016/j.agee.2010.01.006](https://doi.org/10.1016/j.agee.2010.01.006)
- Sánchez-Martín L, Vallejo A, Dick J, Skiba UM 2008 The influence of soluble carbon and fertilizer nitrogen on nitric oxide and nitrous oxide emissions from two contrasting agricultural soils. *Soil Biol. Biochem.* 40, 142–151. doi: [10.1016/j.soilbio.2007.07.016](https://doi.org/10.1016/j.soilbio.2007.07.016)
- Santeramo FG 2015 On the composite indicators for food security: Decisions matter! *Food Rev. Int.* 31, 63–73. doi: [10.1080/87559129.2014.961076](https://doi.org/10.1080/87559129.2014.961076)
- Santeramo FG, Capitanio F, Adinolfi F 2014 Integrating agricultural risks management strategies in selected EU partner countries: Syria, Tunisia, Turkey. *Rom. J. Eur. Aff.* 14, 22.
- Santeramo FG, Carlucci D, de Devitiis B, Seccia A, Stasi A et al. 2018a Emerging trends in European food, diets and food industry. *Food Res. Int.* 104, 39–47. doi: [10.1016/j.foodres.2017.10.039](https://doi.org/10.1016/j.foodres.2017.10.039)
- Santeramo FG, Cioffi A 2012 The entry price threshold in EU agriculture: Deterrent or barrier? *J. Policy Model.* 34, 691–704. doi: [10.1016/j.jpolmod.2012.02.001](https://doi.org/10.1016/j.jpolmod.2012.02.001)
- Santeramo FG, Lamonaca E 2019 On the drivers of global grain price volatility: an empirical investigation. *Agric. Econ.* 65, 31–42. doi: [10.17221/76/2018-AGRICECON](https://doi.org/10.17221/76/2018-AGRICECON)
- Santeramo FG, Lamonaca E, Contò F, Nardone G, Stasi A 2018b Drivers of grain price volatility: a cursory critical review. *Agric. Econ.* 64, 347–356. doi: [10.17221/55/2017-AGRICECON](https://doi.org/10.17221/55/2017-AGRICECON)
- Santos A, Bustamante MA, Moral R, Bernal MP 2016 Carbon conservation strategy for the management of pig slurry by composting: Initial study of the bulking agent influence. *Mitig. Adapt. Strateg. Glob. Chang.* 21, 1093–1105. doi: [10.1007/s11027-014-9593-0](https://doi.org/10.1007/s11027-014-9593-0)
- Sanz-Cobena A, Lassaletta L, Aguilera E, del Prado A, Garnier J et al. 2017 Strategies for greenhouse gas emissions mitigation in Mediterranean agriculture: A review. *Agric. Ecosyst. Environ.* 238, 5–24. doi: [10.1016/j.agee.2016.09.038](https://doi.org/10.1016/j.agee.2016.09.038)
- Sanz-Cobena A, Lassaletta L, Estellés F, del Prado A, Guardia G et al. 2014 Yield-scaled mitigation of ammonia emission from N fertilization: the Spanish case. *Environ. Res. Lett.* 9, 125005. doi: [10.1088/1748-9326/9/12/125005](https://doi.org/10.1088/1748-9326/9/12/125005)
- Sanz-Cobena A, Sánchez-Martín L, García-Torres L, Vallejo A 2012 Gaseous emissions of N<sub>2</sub>O and NO and NO<sub>3</sub>-leaching from urea applied with urease and nitrification inhibitors to a maize (*Zea mays*) crop. *Agric. Ecosyst. Environ.* 149, 64–73. doi: [10.1016/j.agee.2011.12.016](https://doi.org/10.1016/j.agee.2011.12.016)
- Savé R, de Herralde F, Aranda X, Pla E, Pascual D et al. 2012 Potential changes in irrigation requirements and phenology of maize, apple trees and alfalfa under global change conditions in Fluvia watershed during XXIst century: Results from a

- modeling approximation to watershed-level water balance. *Agric. Water Manag.* 114, 78–87. doi: [10.1016/j.agwat.2012.07.006](https://doi.org/10.1016/j.agwat.2012.07.006)
- Saviozzi A, Levi-Minzi R, Cardelli R, Biasci A, Riffaldi R 2001 Suitability of Moist Olive Pomace as Soil Amendment. *Water, Air, Soil Pollut.* 128, 13–22. doi: [10.1023/a:1010361807181](https://doi.org/10.1023/a:1010361807181)
- Schilling J, Freier KP, Hertig E, Scheffran J 2012 Climate change, vulnerability and adaptation in North Africa with focus on Morocco. *Agric. Ecosyst. Environ.* 156, 12–26. doi: [10.1016/j.agee.2012.04.021](https://doi.org/10.1016/j.agee.2012.04.021)
- Schils R, Olesen JE, Kersebaum KC, Rijk B, Oberforster M et al. 2018 Cereal yield gaps across Europe. *Eur. J. Agron.* 101, 109–120. doi: [10.1016/j.eja.2018.09.003](https://doi.org/10.1016/j.eja.2018.09.003)
- Sofi F, Abbate R, Gensini GF, Casini A 2010 Accruing evidence on benefits of adherence to the Mediterranean diet on health: an updated systematic review and meta-analysis. *Am. J. Clin. Nutr.* 92, 1189–1196. doi: [10.3945/ajcn.2010.29673](https://doi.org/10.3945/ajcn.2010.29673)
- Spinelli R, Picchi G 2010 Industrial harvesting of olive tree pruning residue for energy biomass. *Bioresour. Technol.* 101, 730–735. doi: [10.1016/j.biortech.2009.08.039](https://doi.org/10.1016/j.biortech.2009.08.039)
- Sušnik J, Vamvakieridou-Lyroudia LS, Baumert N, Kloos J, Renaud FG et al. 2015 Interdisciplinary assessment of sea-level rise and climate change impacts on the lower Nile delta, Egypt. *Sci. Total Environ.* 503–504, 279–288. doi: [10.1016/j.scitotenv.2014.06.111](https://doi.org/10.1016/j.scitotenv.2014.06.111)
- Tanasijevic L, Todorovic M, Pereira LS, Pizzigalli C, Lionello P 2014 Impacts of climate change on olive crop evapotranspiration and irrigation requirements in the Mediterranean region. *Agric. Water Manag.* 144, 54–68. doi: [10.1016/j.agwat.2014.05.019](https://doi.org/10.1016/j.agwat.2014.05.019)
- Tanyeri-Abur A 2015 Food Security in the southern Mediterranean/North Africa, in *The sustainability of agro-food and natural resource systems in the Mediterranean Basin.*, ed. Vastola A (Springer International Publishing), 3–14. doi: [10.1007/978-3-319-16357-4](https://doi.org/10.1007/978-3-319-16357-4)
- Thorman RE, Chadwick DR, Harrison R, Boyles LO, Matthews R 2007 The effect on N<sub>2</sub>O emissions of storage conditions and rapid incorporation of pig and cattle farmyard manure into tillage land. *Bio-syst. Eng.* 97, 501–511. doi: [10.1016/j.biosystemseng.2007.03.039](https://doi.org/10.1016/j.biosystemseng.2007.03.039)
- Thorson G 1950 Reproductive and larval ecology of marine bottom invertebrates. *Biol. Rev. Camb. Philos. Soc.* 25, 1–45. doi: [10.1111/j.1469-185x.1950.tb00585.x](https://doi.org/10.1111/j.1469-185x.1950.tb00585.x)
- Toreti A, Cronie O, Zampieri M 2019 Concurrent climate extremes in the key wheat producing regions of the world. *Sci. Rep.* 9, 5493. doi: [10.1038/s41598-019-41932-5](https://doi.org/10.1038/s41598-019-41932-5)
- Triberti L, Nastri A, Baldoni G 2016 Long-term effects of crop rotation, manure and mineral fertilisation on carbon sequestration and soil fertility. *Eur. J. Agron.* 74, 47–55. doi: [10.1016/j.eja.2015.11.024](https://doi.org/10.1016/j.eja.2015.11.024)
- Trnka M, Olesen JE, Kersebaum KC, Skjelvåg AO, Eitzinger J et al. 2011 Agroclimatic conditions in Europe under climate change. *Glob. Chang. Biol.* 17, 2298–2318. doi: [10.1111/j.1365-2486.2011.02396.x](https://doi.org/10.1111/j.1365-2486.2011.02396.x)
- Tsikliras AC, Dinouli A, Tsiros V-Z, Tsalkou E 2015 The Mediterranean and Black Sea Fisheries at Risk from Overexploitation. *PLoS One* 10, e0121188. doi: [10.1371/journal.pone.0121188](https://doi.org/10.1371/journal.pone.0121188)
- Tsikliras AC, Licandro P, Pardalou A, McQuinn IH, Gröger JP et al. 2019 Synchronization of Mediterranean pelagic fish populations with the North Atlantic climate variability. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 159, 143–151. doi: [10.1016/j.dsr2.2018.07.005](https://doi.org/10.1016/j.dsr2.2018.07.005)
- Uddling J, Broberg MC, Feng Z, Pleijel H 2018 Crop quality under rising atmospheric CO<sub>2</sub>. *Curr. Opin. Plant Biol.* 45, 262–267. doi: [10.1016/j.pbi.2018.06.001](https://doi.org/10.1016/j.pbi.2018.06.001)
- Valverde P, Serralheiro R, de Carvalho M, Maia R, Oliveira B et al. 2015 Climate change impacts on irrigated agriculture in the Guadiana river basin (Portugal). *Agric. Water Manag.* 152, 17–30. doi: [10.1016/j.agwat.2014.12.012](https://doi.org/10.1016/j.agwat.2014.12.012)
- Vergés A, Steinberg PD, Hay ME, Poore AGB, Campbell AH et al. 2014 The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B Biol. Sci.* 281, 20140846. doi: [10.1098/rspb.2014.0846](https://doi.org/10.1098/rspb.2014.0846)
- Verner D, Treguer DO, Redwood J, Christensen JH, McDonnell R et al. 2018 Climate variability, drought, and drought management in Morocco's agricultural sector. Washington, D.C.
- Vicente-Serrano SM, Nieto R, Gimeno L, Azorin-Molina C, Drumond A et al. 2018 Recent changes of relative humidity: regional connections with land and ocean processes. *Earth Syst. Dyn.* 9, 915–937. doi: [10.5194/esd-9-915-2018](https://doi.org/10.5194/esd-9-915-2018)
- Vicente-Serrano SM, Tomas-Burguera M, Beguería S, Reig F, Latorre B et al. 2017 A High Resolution Dataset of Drought Indices for Spain. *Data* 2, 22. doi: [10.3390/data2030022](https://doi.org/10.3390/data2030022)
- Vicente-Vicente JL, García-Ruiz R, Francaviglia R, Aguilera E, Smith P 2016 Soil carbon sequestration rates under Mediterranean woody crops using recommended management practices: A meta-analysis. *Agric. Ecosyst. Environ.* 235, 204–214. doi: [10.1016/j.agee.2016.10.024](https://doi.org/10.1016/j.agee.2016.10.024)
- Vincent-Cabou L, Peigné J, Casagrande M, Silva E 2017 Overview of Organic Cover Crop-Based No-Tillage Technique in Europe: Farmers' Practices and Research Challenges. *Agriculture* 7, 42. doi: [10.3390/agriculture7050042](https://doi.org/10.3390/agriculture7050042)
- Waha K, Krümmenauer L, Adams S, Aich V, Baarsch F

- et al. 2017 Climate change impacts in the Middle East and Northern Africa (MENA) region and their implications for vulnerable population groups. *Reg. Environ. Chang.* 17, 1623–1638. doi: [10.1007/s10113-017-1144-2](https://doi.org/10.1007/s10113-017-1144-2)
- Webb J, Chadwick DR, Ellis S 2004 Emissions of ammonia and nitrous oxide following incorporation into the soil of farmyard manures stored at different densities. *Nutr. Cycl. Agroecosystems* 70, 67–76. doi: [10.1023/b:fres.0000045985.32440.27](https://doi.org/10.1023/b:fres.0000045985.32440.27)
- Webber H, Ewert F, Olesen JE, Müller C, Fronzek S et al. 2018 Diverging importance of drought stress for maize and winter wheat in Europe. *Nat. Commun.* 9, 4249. doi: [10.1038/s41467-018-06525-2](https://doi.org/10.1038/s41467-018-06525-2)
- Zampieri M, Ceglar A, Dentener FJ, Dosio A, Naumann G et al. 2019 When will current climate extremes affecting maize production become the norm? *Earth's Futur.* 7, 113–122. doi: [10.1029/2018ef000995](https://doi.org/10.1029/2018ef000995)
- Zampieri M, Ceglar A, Dentener FJ, Toreti A 2017 Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales. *Environ. Res. Lett.* 12, 064008. doi: [10.1088/1748-9326/aa723b](https://doi.org/10.1088/1748-9326/aa723b)
- Zampieri M, Toreti A, Ceglar A, Naumann G, Turco M et al. 2020 Climate resilience of the top ten wheat producers in the Mediterranean and the Middle East. *Reg. Environ. Chang.* 20, 41. doi: [10.1007/s10113-020-01622-9](https://doi.org/10.1007/s10113-020-01622-9)

## Information about authors

### Coordinating Lead Authors

Rachid Mrabet:

*National Institute for Agricultural Research (INRA Morocco), Rabat, Morocco*

Robert Savé:

*Institute of Agrifood Research and Technology (IRTA), Caldes de Montbui (Barcelona), Spain*

### Lead Authors

Nuno Caiola:

*Climate Change department, Technology Centre of Catalonia, EURECAT, Spain*

Mouad Chentouf:

*National Institute for Agricultural Research (INRA Morocco), Tangier, Morocco*

Llasat Maria Carmen:

*University of Barcelona, Barcelona, Spain*

Assem Abdelmonem Ahmed Mohamed:

*Agricultural Research Center (ARC), Central Laboratory for Agricultural Climate (CLAC), Giza, Egypt*

Fabio G. Santeramo:

*University of Foggia, Foggia, Italy*

Alberto Sanz-Cobeña:

*CEIGRAM, Universidad Politécnica de Madrid (UPM), Madrid, Spain*

Andrea Toreti:

*European Commission, Joint Research Centre, Ispra, Italy*

Athanosios Tsikliras:

*Aristotle University of Thessaloniki, Thessaloniki, Greece*

### Contributing Authors

Eduardo Aguilera:

*CEIGRAM, Universidad Politécnica de Madrid (UPM), Madrid, Spain*

Luis Asin:

*Institute of Agrifood Research and Technology (IRTA), Lleida, Spain*

Alejandro Blas:

*CEIGRAM, Universidad Politécnica de Madrid (UPM), Madrid, Spain*

Andrej Ceglar:

*European Commission, Joint Research Centre, Ispra, Italy*

Donna Dimarchopoulou:

*University of Rhode Island, United States of America*

Elena Georgopoulou:

*National Observatory of Athens, Athens, Greece*

Luis Lassaletta:

*CEIGRAM, Universidad Politécnica de Madrid (UPM), Madrid, Spain*

Androniki Pardalou:

*Aristotle University of Thessaloniki, Thessaloniki, Greece*

Giuseppe Scarcella:

*National Research Council, Institute of Marine Biological Resources and Biotechnologies (CNR-IRBIM), Ancona, Italy*


Marco Turco:

*University of Barcelona, Barcelona, Spain*

Matteo Zampieri:

*European Commission, Joint Research Centre, Ispra, Italy*





# 3

## RESOURCES

### 3-ENERGY TRANSITION

#### Coordinating Lead Authors:

Philippe Drobinski (France), Brian Azzopardi (Malta)


#### Lead Authors:

Houda Ben Janet Allal (Tunisia/France), Vincent Bouchet (France), Edouard Civel (France), Anna Creti (France), Neven Duic (Croatia), Nestor Fylaktos (Cyprus), Joseph Mutale (United Kingdom), Silvia Pariente-David (France), Joe Ravetz (United Kingdom), Constantinos Taliotis (Cyprus), Robert Vautard (France)

#### Contributing Authors:

Kaouther Ben Nasr (Tunisia), Thierry Brunelle (France), Mikaël Cugnet (France), Paola de Joanna (Italy), Sokol Dervishi (Albania), Juan Fernandez-Manjarrés (France), Dora Francese (Italy), Benoit Gabrielle (France), Lisa Guarrera (France), Victor Homar Santaner (Spain), Boutaina Ismaili Idrissi (Morocco), Rémy Lapère (France), Aina Maimo-Far (Spain), Emanuela Menichetti (France), Lina Murauskaitė (Lithuania), Federico Pontoni (Italy), Gianmaria Sannino (Italy), Roxane Sansilvestri (Spain), Alexis Tantet (France), Michelle Van Vliet (The Netherlands)

*This chapter should be cited as: Drobinski P, Azzopardi B, Ben Janet Allal H, Bouchet V, Civel E, Creti A, Duic N, Fylaktos N, Mutale J, Pariente-David S, Ravetz J, Taliotis C, Vautard R 2020 Energy transition in the Mediterranean. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 265-322.*



# Table of contents

<b>3.3 Energy transition in the Mediterranean</b> .....	<b>268</b>
Executive summary.....	<b>268</b>
3.3.1 Introduction.....	<b>268</b>
3.3.2 Past trends and current situation.....	<b>269</b>
3.3.2.1 <i>Mediterranean energy situation</i> .....	<b>269</b>
3.3.2.2 <i>Renewable energy resources</i> .....	<b>273</b>
<i>Wind power</i> .....	<b>273</b>
<i>Solar power</i> .....	<b>274</b>
<i>Hydropower and thermoelectric power</i> .....	<b>274</b>
<i>Marine energy</i> .....	<b>275</b>
<i>Bioenergy</i> .....	<b>277</b>
3.3.2.3 <i>Energy system vulnerability to climate extremes</i> .....	<b>281</b>
3.3.3 Projections, vulnerabilities and risks.....	<b>282</b>
3.3.3.1 <i>Energy transition scenarios</i> .....	<b>283</b>
3.3.3.2 <i>Energy demand</i> .....	<b>283</b>
3.3.3.3 <i>Energy supply</i> .....	<b>284</b>
3.3.3.4 <i>An NDC-based scenario for the Mediterranean region</i> .....	<b>286</b>
3.3.3.5 <i>Impact of climate change on energy resources</i> .....	<b>287</b>
3.3.3.6 <i>Impact of climate change on energy demand</i> .....	<b>291</b>
3.3.4 Adaptation and mitigation.....	<b>293</b>
3.3.4.1 <i>Adaptation of energy systems to water constraints</i> .....	<b>293</b>
3.3.4.2 <i>Integration of renewables into the energy mix</i> .....	<b>295</b>
<i>Incentive effects of support mechanisms for renewable energy</i> .....	<b>295</b>
<i>The specific case of bioenergy: barriers and opportunities</i> .....	<b>296</b>
3.3.4.3 <i>Energy access in developing countries of the Mediterranean region</i> .....	<b>297</b>
<i>Financial, behavioral, institutional and regulatory impediments to energy access</i> .....	<b>297</b>
<i>Regulatory framework and business models to reach universal access</i> .....	<b>298</b>
<i>The Green Transition in North Africa and the Middle East</i> .....	<b>298</b>
3.3.4.4 <i>Financing the energy transition</i> .....	<b>299</b>
<i>Financial needs by sector and region</i> .....	<b>300</b>
<i>Financial actors</i> .....	<b>301</b>
<i>Financial instruments</i> .....	<b>302</b>
3.3.4.5 <i>Supra-regional cooperation for energy transition</i> .....	<b>302</b>
<i>Mediterranean energy market integration</i> .....	<b>303</b>
<i>The need for supra-regional interconnections</i> .....	<b>305</b>
<i>Strategy of transnational associations of regulators</i> .....	<b>307</b>
<b>Box 3.3.1 Climate variability and energy planning</b> .....	<b>308</b>
<b>Box 3.3.2 Energy issues for Mediterranean islands</b> .....	<b>310</b>
<i>Energy production and demand in the Mediterranean islands</i> .....	<b>310</b>
<i>Interlocking challenges of energy security and climate resilience</i> .....	<b>310</b>
<b>References</b> .....	<b>312</b>
<b>Information about authors</b> .....	<b>322</b>

### 3.3 Energy transition in the Mediterranean

#### Executive summary

Current Mediterranean greenhouse gas emissions amount to a relatively low level of 6% of the global emissions, a proportion close to its proportion of the world population (7.4%). The expected impacts of climate and environmental changes necessitate an accelerated energy transition in the countries of this region to enable a secure, sustainable and inclusive development. The primary energy consumption in the Mediterranean Basin from 1980 to 2016 has steadily increased by approximate 1.7% annually. This trend is mostly related to a steady increase in the consumption of oil, gas, nuclear and renewables and is caused by changes in demographic, socioeconomic (lifestyle and consumption) and climatic conditions in the region. While the northern rim countries advance in gradually diversifying their energy mix, improving energy efficiency and increasing the fraction of renewable energy sources, the eastern and southern rim countries (SEMCs) still lack behind in these developments. The Mediterranean Basin, especially the SEMCs, has large potential for renewable energy, terrestrial as well as marine, including wind, solar, hydro, geothermal, bioenergy, waves and currents. With the increase of the share of renewables, the electricity transmission system will be more exposed to weather variations and may be threatened by specific weather conditions that are usually not considered as extremes.

The projected energy demand trajectories for the next few decades in the Mediterranean Basin are significantly different for the northern versus the eastern and southern rim countries. The energy demand in the North has decreased by 4% since 2010, due to a moderate population growth and a decreasing gross domestic product, and expected to continue to decrease until 2040. The SEMCs have experienced sustained economic and population growth over the past years, which resulted in a growth in a 6% energy demand since 2010. Towards 2040, the energy demand is expected to continue to increase. Although fossil fuels are currently expected to remain the dominant component of the energy mix until 2040, renewables will become the second most used energy source in the Mediterranean Basin and triple until 2040.

A significant gap between energy supply and demand is expected, particularly in SEMCs. It is, therefore, more than necessary to move rapidly towards a restructuring of the energy sector,


particularly the more pronounced integration of renewable energies. Mitigation of greenhouse gas emissions and adaptation to climate change will require investments from households, companies and governments. Regional energy market integration and cooperation are crucial to unleashing cost-effective climate change mitigation.

#### 3.3.1 Introduction

Despite a relatively low share in global greenhouse gas emissions (6%), close to its proportion of the world population (7.4%), the Mediterranean region is severely hit by impacts of climate change (*Section 2.2*). The nature and magnitude of current and future impacts of climate and environmental change in the Mediterranean region, and the associated vulnerability of people, are important imperatives to accelerate the energy transition in all countries of this region in order to enable them to secure a sustainable and inclusive development trajectory.

While the energy transition raises common issues for the region as a whole, the nature and extent of these issues are expressed differently between both shores of the Mediterranean. Northern countries have achieved at least two decades of reforms that enabled them to gradually diversify their energy mix and control to some extent their energy demand due to the deceleration of their demographic growth and the gains in terms of energy efficiency. With respect to the relatively high level of CO<sub>2</sub> emissions from these countries, their energy transition is bound by a rationale that is different from that of Southern and Eastern Mediterranean countries (SEMCs).

The energy transition issue in the SEMCs is closely linked to the sustainability of their development model. These countries face multiple challenges related among others to the fast population growth (607 to 659 million inhabitants, depending on the scenario, by 2050 against 534 million in 2015, *Section 2.7*) which would put additional pressure on energy demand to accompany the surge of urbanization and increasing needs of various sectors of the economy. Therefore, the energy demand in the south-east Mediterranean countries is expected to rise nearly 118% by 2040. The polarization of SEMCs' energy mix on fossil fuels, mainly oil and natural gas, is another important challenge, especially for the countries



**Figure 3.22 | Evolution of Primary Energy Consumption** across the Mediterranean for the period 1980-2016, in exajoules EJ ( $10^{18}$  joules) RES - Renewable Energy Sources (EIA 2019).


where hydrocarbon export revenues play a central role in the macro-financial balance.

### 3.3.2 Past trends and current situation


#### 3.3.2.1 Mediterranean energy situation

The Mediterranean region holds 5% of the world oil and gas reserves, of which 98% are in the countries of the southern rim (UNEP/MAP 2007;


Plan Bleu 2008; UNEP/MAP-Plan Bleu 2009). There is also a significant potential of renewable energies, in particular, solar and wind energies. The Mediterranean region as a whole has been experiencing a steady increase in primary energy consumption, from about 26 exajoules (EJ) in 1980 to 34 EJ in 1995 to 43 EJ in 2016, representing an approximate annual growth rate of 1.7% as in Fig. 3.22. This trend concerns oil, gas, nuclear and renewables, combined with a small decline in the use of coal. In 2005 the combination of



**Figure 3.23 | Evolution of per capita Energy Intensity Index** (energy divided by GDP) in selected Mediterranean countries; 1995 used as a year of reference (EIA 2019).



**Figure 3.24 | Evolution of primary energy production** across the Mediterranean between 1995 (A) and 2016 (B) [Data from the EIA 2019].



**Figure 3.25 | Evolution of primary energy consumption** across the Mediterranean between 1995 (A) and 2016 (B) [Data from the EIA 2019].

mechanisms that include reduced energy intensity by some large consumers, economic crises and political instability, and improved energy efficiency halted the increase in consumption.

Mediterranean countries differ in their patterns of energy use. The level of industrialization, sectoral energy profiles, prevailing climatic conditions and the level of economic growth is among the crucial aspects affecting the energy intensity of each economy. *Fig. 3.23* illustrates the evolution of the per capita energy intensity, expressed as units of energy per unit of GDP, in a selection of Mediterranean economies. Egypt, Morocco and Turkey show the highest increase in energy

intensity during the period 1995-2016, driven by the economic growth. Partly hampered by the financial crisis and partly driven by energy efficiency measures, the French and Italian economies show the highest decline in energy intensity. Others, such as Greece and Spain, had experienced an increase until the mid-2000s but the subsequent financial crisis brought energy intensity back down and has stabilized in the period 2012-2016.

In terms of the energy mix, *Fig. 3.24* and *3.25* illustrate the recent evolution of primary energy production and consumption across the Mediterranean for the period 1995-2016 (EIA 2019). In absolute terms, during this period, the




Figure 3.26 | Energy weight in trade balance (2017-2018) (MEF 2019).

(Country code: Appendix D)

contribution of oil has remained stable at 17 EJ, while consumption of coal has experienced a mild gradual decrease from 4.1 to 3.7 EJ during the two reference years. Primary energy consumption of natural gas has doubled from 6 EJ in 1995 to 12 EJ in 2016, while nuclear and renewable energy sources contribution has risen from 7 to 10 EJ between 1995 and 2016.

On the national level, France has the largest energy-consuming economy in the Mediterranean. Its primary energy consumption has risen from 9.1 EJ in 1980 to 10.7 EJ in 1995, at which level it has returned by 2016 after peaking at 12.1 EJ in 2006. In both 1995 and 2016, roughly 50% of the primary energy consumed, was produced locally (EIA 2019); the vast majority of this relating to the large share of nuclear power in the country's energy mix (UN DESA 2016).


In 1995, the vast majority of Mediterranean nations were mostly dependent on oil, primarily imported, except for Algeria, Egypt, Libya, Tunisia and Syria (Fig. 3.25). The importance of fossil fuels in the energy mix of most of the Mediterranean countries affects their trade balance negatively (Fig. 3.26) (MEF 2019). The SEMCs suffer more from this negative impact because most of them are net importers, and the energy imports can amount to, for some countries, up to 20% in the trade balance.

By 2016, oil dependence was reduced to a certain extent, as the share of natural gas and low carbon technologies has increased. Despite the increased

shares of less carbon-intensive or carbon-neutral energy sources, increasing energy demand has continued to drive greenhouse gas emissions upwards in the region. An attempt to diversify and decarbonize the energy mix is underway, driven mainly by the deployment of renewable energy technologies, as well as substitution of carbon-intensive coal and oil with natural gas. The energy mix is still heavily dependent on fossil fuels.

The shift in energy carriers is a consequence of costs, constraints, regulations and new technologies, among others. The services that make use of these energy sources are, again, disparate and in various states of growth or decline depending on the country. Fig. 3.27 shows the change in energy use in various sectors of the economy for selected Mediterranean countries. The selected countries provide a good representation of the situation in the Mediterranean as a whole, as they are the region's most significant energy users.

There are some clear-cut sectoral trends worthy of note, and some which are not easily explainable by raw data alone. All examined countries exhibit consistent growth in the transportation sector. This is in line with increases in both passenger-km and increased economic activity. This trend is mostly following an increased uptake of private vehicles in the South such as in Turkey, Israel, Egypt, Algeria and Morocco, which is also partially the case for countries in the North Mediterranean such as France, Italy, Spain and Greece, but for those countries, final consumption



**Figure 3.27 | Energy consumption** in selected Mediterranean countries 1990-2016. Data for France, Italy, Spain, Greece, Turkey, Israel, Egypt, Algeria and Morocco (IEA 2018).

in transportation has been declining since 2010. This decline is attributable to a modal shift, policies and regulations that target energy savings in transportation, economic recession, increased use of more efficient vehicles and gradual electrification of transport. These characteristics are not yet strongly apparent in the countries of the South. There is evidence, that energy consumption in transportation will fall, with wider adoption of some (or all) of the above measures across the Mediterranean (OME/MEDENER 2016).

Residential energy use follows a similar upward trajectory in both North and South Mediterranean countries up to 2010, after which final energy consumption in the North is decreasing. The partial decline shown in *Fig. 3.27* is attributable to lower consumption in the four EU countries of this set. There are numerous challenges and opportunities for reducing energy demand. Improved energy efficiency measures and technologies will allow buildings to lower their energy demand per m<sup>2</sup>, but the proliferation of air-conditioning due to higher living standards and increased demand due to climate change-induced higher ambient temperatures may also intensify demand in the region (OME/MEDENER 2016).

Energy use in industry differs significantly between northern and southern Mediterranean countries (*Fig. 3.27*). While the countries of the

north experience a decline due to structural change in the economy, with a shift to services and transformation industry that are less carbon-intensive (from primary industries that are more carbon-intensive), fuel switching and technical efficiency measures, the countries in the south exhibit the reverse, mainly due to new investments in industrial infrastructure and growing economies. Commercial and public service energy use is growing across all the countries examined, but at a much higher rate in the south, where the sector's registered energy use was limited in 1990. Growth in this sector's energy use reflects the changing dynamics in the economies of all countries involved.

Overall the energy consumption in the Mediterranean is increasing but tapering-off of overall consumption. There are multiple sectoral dynamics and a distinct North-South differentiation in the energy use patterns. While the total use seems to be levelling off, there are significant challenges on the horizon about increased demand for industry and services, especially in the south. The projected increase in cooling of living and working places, as caused by climate-change-induced temperature increases, represents the most significant increase in projected energy consumption in the region. The rapid growth in electricity demand is contributing most to the energy demand increase. Between 1971 and 2006 the Mediterranean region saw a fourfold increase in electricity consumption.



Transport continues to be the primary consumer compared with the other sectors in the Mediterranean. Transport electrification is set to shift the resource mix towards low carbon forms of energy but does not reduce overall energy consumption.

### 3.3.2.2 Renewable energy resources

The Mediterranean Basin benefits from a temperate climate with mild winters and warm and sunny summers, with a large potential for energy production from terrestrial renewable energies, as well as for the development of marine energy (Soukissian et al. 2017). These include energies drawn from wind, solar, hydro, bioenergy (crops and forests), waves and currents. Geothermal is an additional key renewable energy source in Europe that can provide low-carbon base-load power. Capacity factors of new geothermal power plants can reach 95% (Chamorro et al. 2014). In the early 2010s, a resurgent interest in geothermal power was observed after nearly a decade of only small development. A substantial number of projects have been developed throughout Europe, and geothermal energy is on its way to becoming a key player in the European energy market (Bertani 2017).

The assessment of resources for wind and solar is generally based on in-situ measurements. Long-term datasets including seasonal, yearly and decadal variability are required. These are usually available from in-situ or remote sensed measurements and gridded re-analyses. These tools have various limitations. In-situ measurements are local and may not have large measurement footprints, and homogeneity of long time series is not granted. At regional scales re-analyses are generally used but have biases, which require specific bias correction methods (Staffell and Pfenninger 2016). Remote-sensed data sets have large spatial coverage but are often short, even though several datasets are currently used for solar radiation estimate (e.g., Müller et al. 2015). Renewable energy resources should be assessed using a variety of tools combined together (Pfenninger and Staffell 2016).

#### Wind power

Wind power is essentially affected by wind speed in the lower part of the atmosphere, at the altitude of the hubs of wind turbines (from 50 to 150 m). Wind power production is a highly nonlinear function of hub-height wind speed, with no

electricity generation below wind speeds of a few  $\text{m s}^{-1}$ , a rapid growth and a saturation at nominal wind speed (typically  $10\text{--}15 \text{ m s}^{-1}$ ). To protect turbines, production is usually cut beyond a threshold of about  $25 \text{ m s}^{-1}$ . Therefore, production is extremely sensitive to low wind-speed changes as well as extreme stormy winds. Wind power production is also marginally sensitive to air density, the denser the air, the larger the production. It is sensitive to turbulence as it decreases efficiency.

A number of studies have currently assessed both the wind power potential in Europe and in Mediterranean areas, as part of the enhanced effort to develop prospective energy mix scenarios including intensive share of renewable energy sources. Beyond national assessments of wind resources in many countries, a New European Wind Atlas (NEWA)<sup>22</sup> (Petersen et al. 2014) is currently being developed, combining wind observations with model results. The offshore component of this atlas includes Mediterranean areas and presents regional climate model results calibrated with satellite scatterometer observations (Karagali et al. 2018).

Over the Mediterranean Sea, larger wind potentials are found in the northwestern part (the Gulf of Lions), in the Alboran Sea and in the Aegean Sea, as indicated by satellite datasets and regional climate modeling (Balog et al. 2016; Onea et al. 2016; Omrani et al. 2017; Soukissian et al. 2017; Rusu and Rusu 2019) (Section 2.2.2.4). Offshore installations can theoretically extract much more kinetic energy from the lower atmosphere than onshore installation in large-scale wind farms (Possner and Caldeira 2017). Currently most installed power lies onshore, with reported installed power provided in Table 3.10. More than 80 gigawatts (GW) are currently installed in Mediterranean countries, but the potential is much higher. Near the shorelines, wind power also benefits from regular and generally predictable land/sea breezes.

Observed near-surface winds have long-term trends (i.e., multidecadal). In Europe in general, winds have been declining for several decades (McVicar et al. 2012), a more general phenomenon called “wind stilling” (Vautard et al. 2010). In the Mediterranean region this trend was less clear. Recent observations show that the wind-stilling trend is recovering on a global scale (Zeng et al. 2019).

<sup>22</sup> <http://www.neweuropeanwindatlas.eu/>

	Installed wind power (MW)	Installed solar PV power (MW <sub>peak</sub> ) 2018
Albania	150	
Algeria	10	
Bulgaria	644	1,036 <sup>(1)</sup>
Croatia	529	61 <sup>(1)</sup>
Cyprus	188	113 <sup>(1)</sup>
Egypt	1,375	1,800 <sup>(2)</sup>
Greece	8,256	2,652 <sup>(1)</sup>
France	19,668	9,466 <sup>(4)</sup>
Israel	123	1,450 <sup>(5)</sup>
Italy	11,175	20,107 <sup>(1)</sup>
Libya	20	
Malta		131 <sup>(6)</sup>
Montenegro	118	
Morocco	1,343	
North Macedonia	37	
Portugal	5,567	671 <sup>(1)</sup>
Spain	24,664	4,751 <sup>(1)</sup>
Tunisia	242	
Turkey	9,384	5,063 <sup>(3)</sup>
<b>Total</b>	<b>83,165</b>	<b>47,170</b>

**Table 3.10 | Reported installed wind and solar (photovoltaic, PV) power in Mediterranean countries in 2019.** References are given in the footnote.

For wind power, see:

<https://www.thewindpower.net/>

For solar power, see:

(1) [https://en.wikipedia.org/wiki/Solar\\_energy\\_in\\_the\\_European\\_Union](https://en.wikipedia.org/wiki/Solar_energy_in_the_European_Union)

(2) <https://spectrum.ieee.org/energywise/energy/renewables/egypts-massive-18gw-benban-solar-park-nears-completion>

(3) [https://en.wikipedia.org/wiki/Solar\\_power\\_in\\_Turkey](https://en.wikipedia.org/wiki/Solar_power_in_Turkey)

(4) [https://en.wikipedia.org/wiki/Solar\\_power\\_in\\_France](https://en.wikipedia.org/wiki/Solar_power_in_France)

(5) [https://en.wikipedia.org/wiki/Solar\\_power\\_in\\_Israel](https://en.wikipedia.org/wiki/Solar_power_in_Israel)

(6) <https://solarfeeds.com/wiki/solar-energy-in-eu/>

### Solar power

Solar power production (concentrated solar power, CSP, or photovoltaic, PV) is mostly influenced by surface solar radiation (*Section 2.2.3.1*), whose variations depend mostly on atmospheric composition (aerosols, water vapor) and clouds. The importance of aerosols has been noted in several studies for Mediterranean areas (e.g., Gutiérrez et al. 2018). Solar production is extremely sensitive to clouds and cloud types. For PV, panels efficiency also largely depends on cell temperature, which itself depends on air temperature, radiation

and near-surface wind speed. Solar panels may also be sensitive to the presence of snow and ice cover or particulate matter potentially covering panels. Solar resources are of particular interest in Mediterranean countries due to the high mean solar irradiance in the region (Hadjipanayi et al. 2016). Solar radiation increases from North to South in the Mediterranean Basin, with typical yearly mean values of 150-250 W m<sup>-2</sup>, and 1,300 to 2,000 kWh m<sup>-2</sup> yr<sup>-1</sup>.<sup>23</sup> On the European side weather disturbances make the resource variability higher than on the southern side.

Solar radiation has undergone varying trends in past decades, due to cloud changes and the “dimming and brightening” phenomena linked to changing aerosols atmospheric composition (Wild et al. 2005) (*Section 2.2.3.1*). In Europe an increase of solar radiation of 2 W m<sup>-2</sup> decade<sup>-1</sup> was observed from 1983 to 2010 (Sánchez-Lorenzo et al. 2017), with higher values in the Mediterranean regions found in a set of ground stations (about 5 W m<sup>-2</sup> decade<sup>-1</sup>) (Pfeifroth et al. 2018). These trends are probably mostly attributable to changes in cloudiness but aerosol variations also affect mean solar radiation in a significant manner (Philipona et al. 2009; Nabat et al. 2015) (*Fig. 2.4*). Such changes are also likely to affect summer temperatures. Dong et al. (2017) found that aerosols decline explain about half of the rapid rise of summertime extreme temperatures.

### Hydropower and thermoelectric power

Hydropower relies on the availability of water in large reservoirs, or the streamflow intensity for run-off-the-river production. Production is sensitive to precipitation and snowpack melt, allowing to feed the reservoirs. Droughts and associated low flows are limiting the production. Bioenergy, just as agriculture and forestry, is largely dependent on climate conditions in many ways (seasonality of temperature, radiation, precipitation). Marine energies depend on currents, which have a low frequency variability, and on waves, themselves influenced by wind speed conditions.

Addressing impacts of climate variability and change on water resources, electricity supply and energy infrastructure vulnerability and resilience relies on a global hydrological-electricity coupled modelling framework. It consists of a physically based hydrological (Liang et al. 1994) and water temperature model (Yearsley 2009; van Vliet et al.

<sup>23</sup> <https://globalsolaratlas.info/>


2012a), which are linked to hydropower and thermoelectric power models (Koch and Vögele 2009; van Vliet et al. 2012b). Fig. 3.28 shows simulated hydropower utilization rates and utilisation rates of thermoelectric power over the period 1981–2010 in the form of boxplots with the distributions of utilisation rates of all plants for the Mediterranean (van Vliet et al. 2016a). The utilization rates are fairly constant, but they may be reduced in severe drought years. During such years, hydropower utilisation rates were on average reduced by 5.2%, and thermoelectric power by 3.8% (worldwide average). This corresponds to severe streamflow drought years for hydropower and to streamflow drought and high water temperature for thermoelectric power. Overall utilisation rates of thermoelectric plants are lower than for hydropower, since usable thermoelectric power capacity may be limited by more factors (i.e., streamflow drought and high water temperature) and benefits less from storage of water in reservoirs during low flow conditions than conventional hydropower plants (van Vliet et al. 2016a).

### Marine energy


The energy resource in the ocean comes from five distinct sources, each with different origins and requiring different technologies for conversion, including, (1) tidal currents that extract kinetic energy from tidal flow, (2) tidal range which captures the potential energy created by the difference in sea level between high and low tides, (3) waves which convert kinetic energy transmitted by the wind to the upper surface of the ocean, (4)

ocean thermal energy conversion which exploits the temperature difference between deep and surface ocean layers, and (5) salinity gradients which exploit the chemical potential due to salinity gradients in water bodies. These resources are not uniformly distributed on the globe. Also, the degree of maturity of the technology necessary to their exploitation is different. In the Mediterranean Sea, the two ocean energy sources with the highest potential are tidal currents and waves.


The Mediterranean coastal areas experience two high tides and two low tides per day. Tidal currents are generated by horizontal movements of water, modified by seabed bathymetry, particularly near coasts or other constrictions (e.g., islands). Tidal current flows result from the rise and fall of the tide; although short-term weather fluctuations can slightly influence these flows, their timing and magnitude are highly predictable and largely insensitive to climate change influences (Lewis et al. 2011). In the Mediterranean Basin, there is no commercial development of the tidal energy sector. As tidal turbines need a stream speed of at least  $1.5\text{--}2\text{ m s}^{-1}$  to operate effectively, the tidal energy potential of the basin sets specific constraints. Given the minimal flow needs provided above, some Mediterranean sites could be of particular interest. The Straits of Gibraltar, and particularly the Strait of Messina (where tidal stream energy resource presents its highest values in the Mediterranean) have been under consideration (Soukissian et al. 2017). The Strait of Messina is characterized by high-energy tidal currents with maximum velocities at spring peak tides ranging



**Figure 3.28 | Impacts of streamflow drought and high water temperature on utilisation rates of hydropower (A) and thermoelectric power (B) for 1981–2010.** Boxplots with distributions of utilisation rates of hydropower and for thermoelectric power are presented with the largest number of plants and installed capacity. Values of 1 indicate that a power plant works at full capacity (no constraints) while for instance a value of 0.8 indicate that the plant works at 80% of the maximum capacity. Highlighted years indicate that utilisation rates were reduced significantly compared to the average over 1981–2010 (van Vliet et al. 2016a).



**Figure 3.29 | Distribution of wave energy flux** in  $\text{kW m}^{-2}$  averaged over the period 2001–2010 in the Mediterranean Sea. The energy resource was evaluated through of a numerical simulation performed using an ocean wave model. The model was forced with six-hourly wind fields obtained from European Center for Medium-Range Weather Forecast (ECMWF) operational analysis at  $1/4^\circ$  spatial resolution (Liberti et al. 2013).



**Figure 3.30 | Potential locations for offshore wind farms** (Piante and Ody 2015).

from  $1.8 \text{ m s}^{-1}$  to more than  $3 \text{ m s}^{-1}$ , proving the suitability of the site for tidal energy harnessing (El-Geziry et al. 2009; Coiro et al. 2013). An estimation of the marine current energy fluxes in the Gibraltar Strait has been provided in Calero Quesada et al. (2014), revealing the suitability of two main sills (Camarinall in the middle of the Strait, and

Espartell at the wester entrance of the Strait) for a power plant installation, with computed averaged fluxes in these areas that can exceed  $1.8 \text{ kW m}^{-2}$ .

Ocean wave energy is energy that has been transferred from the wind to the ocean. As the wind blows over the ocean, air-sea interaction transfers

some of the wind energy to the water, forming waves, which store this energy as potential and kinetic energy. The size and period of the resulting waves depend on the amount of transferred energy, which is a function of the wind speed, the length of time the wind blows (order of days) and the length of ocean over which the wind blows (fetch). Energy availability is certainly a major factor affecting wave energy production but high energy potential usually implies exceptional wave conditions during extreme events. Such conditions pose serious engineering challenges to the design and deployment of wave energy converters increasing the costs of development, production, installation, maintenance and insurance of these devices. On the other hand, in calmer and semi-enclosed seas such as the Mediterranean, where lower amounts of wave energy are available, many technical issues related to extreme sea climate could be more easily solved, possibly making wave energy production still economically viable. From this point of view, wave energy production in the Mediterranean is particularly appealing (Fig. 3.29) (Liberti et al. 2013).

Offshore wind is likely the aspect of the energy transition of the Mediterranean region with the most important development potential, particularly in SEMCs. It has been introduced in other parts of the world, as it is less environmentally disturbing than on-shore alternatives (Piante and Ody 2015). So far, there is no offshore wind farm in the Mediterranean, although offshore wind production could be highly profitable (Fig. 3.30) (Gaudiosi and Borri 2010).

### Bioenergy


Bioenergy is an important source of renewable energy in the Mediterranean, with an annual output from solid biofuels that dominates by far the production of electricity from solar or wind sources (Table 3.10 and 3.11; Bryden et al. 2013; IEA Bioenergy 2016). Traditional biomass remains a major source of renewable energy on the south side, given the low development of renewable energy in this part of the region (Table 3.11). Biomass is the least promising sector for electric energy production and it is rather reserved for the production of heat or fuel. This stems from the fact that bioenergy encompasses a broad range of value-chains and end-uses, providing heat, electricity, and transportation fuels from a variety of biomass sources and conversion pathways (Sansilvestri et al. 2020).

In the Mediterranean region, the importance of bioenergy is highly variable across countries as

	Installed wind power (MW)	Installed solar PV power (MW <sub>peak</sub> ) 2018
Albania	150	
Algeria	10	
Bulgaria	644	1,036 <sup>(1)</sup>
Croatia	529	61 <sup>(1)</sup>
Cyprus	188	113 <sup>(1)</sup>
Egypt	1,375	1,800 <sup>(2)</sup>
Greece	8,256	2,652 <sup>(1)</sup>
France	19,668	9,466 <sup>(4)</sup>
Israel	123	1,450 <sup>(5)</sup>
Italy	11,175	20,107 <sup>(1)</sup>
Libya	20	
Malta		131 <sup>(6)</sup>
Montenegro	118	
Morocco	1,343	
North Macedonia	37	
Portugal	5,567	671 <sup>(1)</sup>
Spain	24,664	4,751 <sup>(1)</sup>
Tunisia	242	
Turkey	9,384	5,063 <sup>(3)</sup>
<b>Total</b>	<b>83,165</b>	<b>47,170</b>

**Table 3.11 | Levels of primary solid biofuels** (in terajoules ( $10^{12}$  joules, TJ) from domestic supply in Mediterranean countries for which information is available for 2017. Data downloaded from <https://www.iea.org/data-and-statistics>.

it depends on the available biomass from forests, agriculture and organic waste. Bioenergy is difficult to characterize as it also has the advantage of producing fertilizers after the organic matter has been digested. In terms of technology, it is common to separate traditional biomass, which predominates in developing countries and involves the burning of wood fuels and agricultural residues for heating and cooking, from modern forms of bioenergy production relying on somewhat more complex biomass processing systems (Chum et al. 2011). Those include liquid biofuels for transport (e.g., bioethanol from sugar crops), pellets from forest residues or agricultural biomass, or electricity generated by dedicated power plants. Anaerobic digestion of organic waste (from cattle, agro-industry or municipal sources) to produce biogas has been growing lately throughout the Mediterranean area. Traditional biomass and modern uses of biomass are both present in the Mediterranean region, although the latter is more difficult to quantify due to a lack of statistics.




**Figure 3.31 | Firewood production** in the southern and northern parts of the Mediterranean Basin (FAO 2017).

Overall production of energy from solid biomass in the Mediterranean region amounts to at least 1.56 PW according to the statistics published for 2017 by the International Energy Agency<sup>24</sup>. Energy, including heat production from biomass, varies significantly from one country to another, being concentrated in northern Mediterranean countries (Fig. 3.31). The consumption of wood, as reported by FAOSTAT includes traditional and modern usages of the biomass, presumably with a large proportion of the former when compared to other

statistics available on modern value-chains relying on this feedstock. The production of firewood has increased by about 90% in North Africa over the last 60 years while it is back to its 1960's level in southern Europe after a large dip from 1973 to 2009 (Fig. 3.32). The increased demand for firewood in North Africa arises from rising demographic pressure, especially in the rural areas (Schilling et al. 2012). The pressure on wood resources may be alleviated by improving the efficiency of cook stoves (thereby reduce health damages associated with open-hearth indoor fires), or switching to alternative renewable energy sources (Chum et al. 2011). In northern Europe, the competition with other end-uses for wood (e.g., for building, furniture-making or pulp and paper) explains the temporary decrease in firewood consumption.

The availability of biomass from forests is highly asymmetric between northern and southern Mediterranean countries (Fig. 3.32). Considering its large area of forests, the Mediterranean Basin represents a good candidate to develop wood biomass energy for the renewable energy sector development (Gómez et al. 2010). Wood biomass




**Figure 3.32 | Forest area and production forests available for industrial use including biomass for energy purposes.** Values correspond to 1,000 hectares. Data obtained from the 2015 FAO Global forest Assessment<sup>25</sup>. Data for France obtained from official reports<sup>26</sup>, whose classification of production forests may vary with respect to FAO guidelines.

<sup>24</sup> <https://www.iea.org/data-and-statistics>

<sup>25</sup> <http://www.fao.org/forest-resources-assessment/en/>

<sup>26</sup> <https://inventaire-forestier.ign.fr/>



**Figure 3.33 | Electricity production from biogas and solid biomass** in Turkey, Israel and Spain (IEA 2018).

has low energy density and it is highly spread, two problems that increase harvesting and transportation costs (Caputo et al. 2005; Yoon et al. 2012). While forest surface may be increasing on northern Mediterranean countries, a continuous decline in northern African countries is currently occurring (see FAO forest Global Assessment<sup>27</sup>). Forest fragility is exacerbated in the Mediterranean area where forests have low productivity and agriculture is difficult considering regional climate conditions (González et al. 2015).

Large bioenergy facilities may constitute threats to the biodiversity and forest cover in any region, this also in the Mediterranean. Bioenergy must be ecologically sustainable, environmentally acceptable for the public, and the delivery costs need to be lower than for fossil fuels (Bilgen et al. 2015), but this is often not the case. An example can be found in the Mediterranean zone of France where the conversion of a carbon-based electric central to wood biomass-fueled power plant in 2016 caused conflict between citizens, the forestry sector and regional authorities (Sansilvestri et al. 2020). In contrast, small sized boilers using wastewood resources do not raise the same levels of concern to citizens and are more accepted.

Important efforts exist in Algeria and Morocco to scale up renewable energy sources. For instance, the Renewable Energy Development Centre in Algeria<sup>28</sup> has programs on the whole array of renewable energy including biomass and solar. Clearly, North African countries and other Mediterranean countries will rely more on non-forest biomass as long as there are agricultural or domestic waste as the forest biofuel supply is limited in this region.

<sup>27</sup> <http://www.fao.org/3/a-i4808e.pdf>

<sup>28</sup> <https://www.cder.dz/>

Spain uses mainly biofuels for transports and more recently solid biomass for heating network installations (Paredes-Sánchez et al. 2016) In France, the number of heat boilers and networks increased from 30 in 2003 to 284 in 2016 (Neumuller 2015; OFME 2015). The main bioenergy potential in Portugal is domestic wastes (Ferreira et al. 2017). For Morocco wood biomass represents a real economic market with the heat demand for hammam and domestic cooking, causing continuous loss of forest surface (Zouiri and Elmessaoudi 2018).

While firewood is mostly used for heating and cooking, the recent development of more refined bioenergy systems may be captured by statistics on biogas production for the co-generation of heat and electricity in Turkey, Israel or Spain (Fig. 3.33). Power plants running on solid biomass (from forestry or agriculture) are operating in Spain, Italy and Portugal, with similar outputs (2,600 - 43,000 GWh range in 2018) (IEA 2018). No estimates were found for northern Africa, but some programme targeting small-scale household digestion systems have been reported in Morocco and Tunisia (Mulinda et al. 2013). Biogas and bioelectricity production use mostly residues as feedstocks, but statistics on the amount of biomass hereby mobilized are not available. Overall, the contribution of biomass to the national energy mixes is variable in the transport sector (from 0% in the SEMCs to 9% in France), usually larger in the heating sector, and small but growing in the electricity sector.

The production of liquid biofuels (which currently relies on food crops as feedstocks) has only been reported in three southern Europe countries (France, Italy and Spain), as a result of the changes in the Common Agricultural Policy of the EU in the 1990's, and of a series of policies such as the renewable energy directive of the EU. The latter mentioned a 10% target for the share of renewables in the transport sector in 2020, most of which would be achieved with biofuels. Specific rules will be applicable for bioenergy produced from food and feed crops with a target of no more than 7%. The contribution of biofuels with a high risk of indirect land use change (i.e., mainly imported biomass such as palm oil) will be gradually reduced to 0% by 2030.

Unlike other renewables, biomass and biofuels in general may be traded across countries and con-


tinents. In 2017 for instance, Italy imported large amounts of wood pellets from North America (totalling 1.8 M tons or about 32.4 PJ of energy content). This means that the lack of local biomass production potential in the Mediterranean Basin may be compensated for by imports, although its consequences in terms of overall pressure on land resources should be carefully assessed (Searchinger et al. 2008).

In general, agriculture and forestry in the Mediterranean countries are faced with a growing array of challenges (IPCC 2014): the net primary production of managed ecosystems is constrained by limited water resources, the low availability of land on which to grow crops, and soil degradation (in general from erosion, salinity or desertification in the southern Mediterranean) (Olsson et al. 2019). This clearly hinders the development of purpose-grown plants (whether biofuel crops, lignocellulosic plants such as miscanthus, or short rotation coppice), which make up a large part of biomass supply in bioenergy expansion scenarios, e.g., in the 1.5°C warming scenarios of the IPCC SR1.5 (IPCC 2018). Also, regarding agricultural land management, the emphasis in the

SEMCs is primarily on food production since all countries rely on imports to meet the needs of their inhabitants. The food constraint is such that in the study of biomass potentials by Beringer et al. (2011), no bioenergy plantations are projected in the SEMCs area by 2050, and to a very limited extent in the northern part of the basin (Fig. 3.34).

Since biomass availability is the primary constraint to the development of bioenergy and is limited by a range of physical and economic factors (e.g., land availability, competition with other uses, productivity), many studies have attempted an assessment of “biomass potentials” given a set of assumptions and limitations. Sustainability has also become a major issue with bioenergy systems, arising in particular from the consequences of developing biofuels on the use of land worldwide, and the possibility that these may negate the climate benefits of substituting biofuels (Searchinger et al. 2008; ElAkkari et al. 2018). In principle, “sustainable biomass potentials” would provide the most useful guidance, but are not easy to determine because the multifaceted nature of sustainability is difficult to factor in (Chum et al. 2011). The trade-offs with soil quality when exporting agricultural or forest residues for bioenergy purposes can be evaluated (Saffih-Hdadi and Mary 2008). It results in a drastic reduction of removal rates (by two thirds in France, for instance).

Regarding the use of land for bioenergy plantations, an option to mitigate the competition between food production and bioenergy markets would consist in growing these plants on marginal land, i.e., land which is unsuitable for other purposes (Fritsche et al. 2017). There is still considerable debate as to the actual amount of such land worldwide, and whether it would be economically feasible to grow biomass plants on these lands. Current estimates range from 350 to 6,000 Mha worldwide (compared to a global cropland area of 1,700 Mha), and a recent study estimated an area of 69 Mha in Europe, among which 43 Mha are located in the northern Mediterranean area (Elbersen et al. 2018). Assuming an energy yield of about 60 GJ ha<sup>-1</sup> on marginal land (Gelfand et al. 2013), this would translate as an output of 2,600 PJ yr<sup>-1</sup> for the northern Mediterranean. This is a large amount compared to the current use of biomass in this area (for instance biomass produces 73 PJ of heat yr<sup>-1</sup> in France and Italy (IEA 2018), with wood as the main feedstock). Aside from this potential opportunity on marginal land, avenues to increase biomass outputs in Europe include an intensification of forestry, the development of purpose-grown plants, which only occupy a marginal fraction of



**Figure 3.34 | Projected constraints to the establishment of bioenergy plantations by 2050.** The “LUC emission” constraint corresponds to a loss of soil C upon conversion to bioenergy crops, which could not be paid back in less than 10 years. Adapted from Beringer et al. (2011).



land so far (Don et al. 2011), and an increased valorization of residues and waste streams (for biogas, heat and power). Regarding energy crops, a study factoring in sustainability constraints estimated that the Northern Mediterranean countries could produce 630 PJ yr<sup>-1</sup> in 2030 from less than 10 PJ yr<sup>-1</sup> in 2010 (Don et al. 2011; Elbersen et al. 2012).

Regarding the SEMCs, Stecher et al. (2013) reviewed the biomass potential studies and their numbers for the African continent, concluding that the potential for energy crops could range from 0-13,900 PJ yr<sup>-1</sup>, 0-5,400 PJ yr<sup>-1</sup> for forestry biomass and 10-5,254 PJ yr<sup>-1</sup> for residues and waste by 2020. While those numbers could not be disaggregated across regions in Africa, they point to significant potential for all three feedstock categories. From a sustainability perspective, as suggested in Chum et al. (2011), residues are particularly efficient at reducing GHG emissions. Their use was prioritized in the strategic energy plan of Morocco for 2030, which emphasizes the use of organic waste, agricultural residues, and algae – a medium-term technology unlikely to be commercialized before 2040 (Chum et al. 2011; Royaume du Maroc 2017).


### 3.3.2.3 Energy system vulnerability to climate extremes

There are a number of ways extreme weather events affect energy resources and energy systems. Individual events may threaten localized installations but large-scale events such as cold spells may threaten the electricity load balancing at a large scale. Also, with the increase of the share of renewables, the electricity transmission system

will be more exposed to weather variations and may be threatened by specific weather conditions that are usually not considered as extreme.

Heat waves increase energy demand for cooling and increase rivers temperature. Long heat waves are generally associated with low flows (see below). Increased river temperatures reduce the permissible temperature increase in power plants, thereby inhibiting cooling the cooling water capacity and inducing plant production (Koch and Vögele 2009; van Vliet et al. 2016a). Permissible temperature increase is in particular bounded by critical biological threshold for freshwater species and strongly regulated. Excessive heat also affects power lines capacity to dissipate heat and reduces transmission capacity (Bartos et al. 2016). Heat also reduces the efficiency of solar panels and long-term thermal stress deteriorates PV cells (Chow 2010).

Long-lasting droughts may induce low flows in rivers, reducing production of run-of-the-river hydropower and stock in reservoirs for hydropower (van Vliet et al. 2016a). Droughts also increase the sensitivity of river temperature to air temperature, so that low flows combined with a heat wave can lead to a large warming of river temperatures with consequences on thermal production. The 2003 heatwave and associated drought impacted production in 30 nuclear power plants (Schewe et al. 2019). During the drought, warm year of 2003 simulated hydropower utilisation in Europe was significantly reduced by 6.6% and thermoelectric power by 4.7% compared to the average of 1981–2010 with a smaller impact in the Mediterranean than in northern Europe as Mediterranean power plants are hampered by water constraints on a more



**Figure 3.35 | Mediterranean patterns with number of days in the given year with streamflow drought** (left), **high water temperature** (middle), **and that both events coincide** (right). Results are presented for year 2003 with both streamflow drought and high water temperature. Scale for right figure panels (streamflow drought and high water temperature) differs from the scale of the left (streamflow drought) and middle (high water temperature) panels (van Vliet et al. 2016a).


regular basis, resulting in lower absolute values of plant utilisation rates and smaller relative changes in usable capacity for 2003 compared to the long-term average for 1981–2010. In 2003, northern Europe and the Mediterranean region display large regions with streamflow drought for more than 100 days and high water temperature for more than 50 days (Fig. 3.35; van Vliet et al. 2016a).

Cold spells are exacerbating energy demand, due to increased heating (Thornton et al. 2016). Cold spells generally cover a large geographical extent (typically 1,000-2,000 km<sup>2</sup>) and may induce challenging conditions for the transmission system. Floods threaten all infrastructures and power devices. Storms and wind gusts also threaten infrastructures, in particular power lines, which may be damaged by falling trees. Solar panels

are sensitive to several sorts of extreme weather events (floods, storms), which may threaten the infrastructure. Storms also induce halting of wind turbines for their protection. Storm surges threaten coastal infrastructures. By contrast, low winds induce a loss in wind power production. There are also a number of other hazards, which affect energy systems such as icing on wind turbines, freezing rain and heavy snowfalls endangering power lines, landslides affecting infrastructures.

### 3.3.3 Projections, vulnerabilities and risks

By improving energy efficiency and deploying renewables on a large scale, the Mediterranean region would enhance energy security for all countries, improve export potential for exporting ones and reduce energy costs and environmental



**Figure 3.36 | Voluntary commitments of SEMC** (Nationally Determined Contribution – NDC – and funding). The upper part shows the installed power capacity of non-hydro renewable energy technologies in 2018, and the extra effort needed to achieve the targets established by the SEMCs at the horizon 2020-2030. The lower part shows the level of funding needed to implement the measures included in the NDCs including the share covered by local vs. international climate funding (unconditional vs. conditional). Sources: OME/MEDENER (2018) - bottom part: UNFCCC<sup>29</sup>

<sup>29</sup> <https://www4.unfccc.int/sites/submissions/INDC/>

damages for the whole region. Embarking on an energy transition path will also help improve social welfare in the region and contribute to job creation, among other positive externalities.

Nationally Determined Contributions (NDCs) are at the heart of the Paris Agreement and the achievement of the long-term goals to keep temperatures below 2°C. They are voluntary commitments to greenhouse gas emission reduction in all sectors, among which the energy sector, which is the main responsible for greenhouse gas emissions at global scale. The NDCs of the Mediterranean countries and their implementation have been analyzed in depth by several organisations like IPEMED (Robin 2015), OME/MEDENER (2016), the UfM (Fernández and Hertz 2019) or the European Union project ClimaSouth (Rizzo and Maro 2018) and have served for elaborating energy transition scenarios. *Fig. 3.36* shows the voluntary commitments of SEMC to the required energy transition.

### 3.3.3.1 Energy transition scenarios

Since 2008, OME regularly issues a Mediterranean prospective analysis to 2040, the “Mediterranean Energy Perspectives” series (MEP). The MEP analyses the trends for energy demand in the different use sectors and the implications in terms of security of supply, CO<sub>2</sub> emissions, and environmental impacts. Within this framework, in 2015 MEDENER and OME published a joint Mediterranean Energy Transition Scenario (TS), an ambitious scenario that goes beyond the plans and targets announced by governments and policymakers. In all scenarios, the OME follows a structural econometric approach that combines economic theory and statistical methods to produce a system of equations establishing causal relationships between energy demand and activity variables (such as GDP, population etc.). This system of equations is then used to generate medium and long-term forecasts of future energy demand. The descriptions of the model and of the energy scenarios assumptions are available in the Appendix of the OME/MEDENER report (OME/MEDENER 2016).

The Mediterranean Energy Transition Scenario (TS) assumes the implementation of those measures that are currently the most technically, economically, and politically mature for large-scale rollout of energy efficiency and renewable energies. This scenario assumes no major technology breakthrough, but the deployment of existing technologies and sound energy efficiency policies and measures across all Mediterranean countries.


According to this analysis, under a Business-As-Usual or so-called “Conservative” Scenario (CS) the situation would evolve critically on all counts over the next 25 years: doubling of energy demand and tripling of electricity consumption, soaring infrastructure and import bills (+443 GW to be installed and doubling of the fossil-fuel imports) and a critical rise in carbon emissions (+45%). Such a scenario, based essentially on fossil fuels, would put further strain on the climate and exacerbate geopolitical tensions in the region. A change of energy trajectory is therefore necessary for all Mediterranean countries to curb the trends through increasing energy efficiency and renewable energy deployment.

Based on this exercise, in 2018, a new edition of the Mediterranean Energy Perspectives was released by OME, which includes two scenarios: i) the Reference Scenario (RS) which considers past trends, current policies and ongoing projects and incorporates the unconditional targets of Nationally Determined Contributions (NDCs); in other terms the RS assumes that international financing and other aids will not be forthcoming. ii) The Proactive Scenario (PS) is based on the implementation of strong energy efficiency programmes and increased diversification in the energy mix based on the NDCs submitted by each country and assumes that international financing will be made readily available and that all targets of the NDCs will be met in full.

### 3.3.3.2 Energy demand

In the Mediterranean region, energy demand increased from 711 Mtoe in 1990 to 978 Mtoe in 2015, an average growth of 1.3% yr<sup>-1</sup>. The largest share of regional energy demand is in the North Mediterranean countries, which account for over 63%. Expected trajectories for energy demand in the region are very contrasted across the two shores of the Mediterranean (*Fig. 3.37*).

The northern countries are ahead in terms of a transition path with substantial levels of renewables and effective demand-side management. The energy demand in the North has decreased by 8% since 2010. This decrease is not only due to energy efficiency efforts but should also be seen in the light of a very moderate population growth (+0.5%) and decelerating gross domestic product growth, especially after the 2008 financial crisis (-2%). In both scenarios, by 2040, energy demand in the North Mediterranean would continue to decrease. In 2040, North Mediterranean energy demand would be 10% and 23% lower than 2015



**Figure 3.37 | Primary energy demand by region, in megatons of oil equivalent (Mtoe).** RS = Reference Scenario; PS = Proactive Scenario (adapted from OME 2018), see Section 3.3.3.1 for definition of the scenarios.

levels, in the Reference Scenario (RS) and Proactive Scenario (PS), respectively.

The South and East Mediterranean, on the other hand, have experienced sustained economic and population growth over the past years (+6% and +5% respectively), translating into growth in energy demand by +6% since 2010. In all scenarios, energy demand continues to increase by 118% and 72% from 2015 levels, for the Reference Scenario (RS) and Proactive Scenario (PS), respectively. Energy savings would be of 21% in the Proactive Scenario (PS) compared to the Reference Scenario energy demand forecasts.

The South and East Mediterranean would account for 61% of the energy savings with 2025 Mtoe, six times the 2015 primary energy demand of all south and East Mediterranean countries. Cumulative potential energy savings in the North Mediterranean, while less substantial than in the South and East, would still be considerable at around 1315 Mtoe over the same period – more than double current North Mediterranean primary energy demand.


The potential for energy efficiency is substantial in the Mediterranean region, particularly in the South and East. Despite some improvements, energy efficiency is still, at present, in its infancy stage in the region. Overall, energy intensity is decreasing in the region, largely related to shifts in the buildings, industry and transport sector. Globally, 45% of energy consumption (and a similar share

of greenhouse gas emissions) are attributable to buildings (Butler 2008). New constructions in developing economies are numerous (Hui 2000). Building regulations aimed at energy efficiency are indispensable, especially in these developing economies, where the energy market alone, does not allow for the activation of incentive mechanisms. Substantial energy savings can be reached in the building sector (residential and tertiary sectors), especially in the South Mediterranean where over 50 million new dwellings are expected to be built over the next decades (OME/MEDENER 2015; OME, 2018). Barriers exist, including the high cost of efficient equipment, the difficulty of changing habits, the lack of adequate technology and also the ignorance of energy and climate issues by some architects in these countries (Iwaro and Mwashia 2010). Jaber and Ajib (2011) propose an assessment of the best orientation of the building, the optimal window size, the optimal thickness of thermal insulation from an energy, economic and environmental point of view for a typical residential building in the Mediterranean region. They suggest that about 28% of annual energy consumption can be saved by choosing the best orientation, optimal window size and optimal insulation thickness. The choice of new materials can also contribute to energy savings (Zabalza Bribián et al. 2011; Buoninconti and Filagrossi Ambrosino 2015).

Industry can substantially improve its efficiency of electricity consumption, but there is less scope to decrease its fossil fuel use in heavy industries, especially in the South and East Mediterranean countries. For all SEMCs, except Tunisia, the share of industry in the final energy consumption is declining. The share of industry in final energy demand is already low in Lebanon (12%) because the activity of this sector is traditionally less developed while it is high in Tunisia (35%). It fell sharply in Lebanon (-10%), after the war with Israel in 2005-2006, with the destruction of many industrial infrastructures, and to a lesser extent in Italy, France and Greece due to the economic crisis and the increasing trend of the service sector (MEDENER 2014). The transport sector would witness strong efficiency gains over the outlook period (18%) as they offer great scope for efficiency improvements in areas such as improved engines, and modal transport expansion.

### 3.3.3.3 Energy supply


In both OME scenarios, fossil fuels remain the dominant component of the mix. While oil alone is expected to remain the dominant fuel of the energy mix, mainly for its demand in the transport



**Figure 3.38 | Primary energy resources in the Mediterranean energy mix in megatonnes of oil equivalent.** RS = Reference Scenario; PS = Proactive Scenario (OME 2018).

sector, a complete reshape is occurring in the electricity mix, where renewable energy sources have overtaken natural gas capacity and become the main fuel in the Mediterranean electricity mix. The Mediterranean region has abundant renewable energy resources. Yet, today renewables still account for a limited share of the region’s primary energy supply (11% in 2015, Fig. 3.38). Traditionally the most exploited renewable energy sources have been biomass and hydropower. Geothermal energy contributes in a few countries mainly Italy, Turkey

and, to a lesser extent, France, Spain and Portugal. In recent years, wind and solar, both for electricity and heat production have entered the energy mix. In 2040, the share of renewables would reach 13% in the Reference Scenario (RS) and 24% in the Proactive Scenario (PS) (OME 2018). Most of the increase is expected to come from wind and solar. Among the various renewable energy technologies, solar is expected to grow at the fastest pace in both sub-regions. End usage of solar thermal energy, in particular solar water heaters, offers great poten-



**Figure 3.39 | Electricity generation mix by fuel type, in terawatt-hours (TWh).** RS = Reference Scenario; PS = Proactive Scenario, RES = Renewable Energy Sources. (Adapted from OME 2018).

tial in the South and is efficient with good return on investment. Solar water heating and solar cooling demand will also increase by 2040.

The most significant change ahead is a substantial increase in the contribution of renewables to power generation (*Fig. 3.39*). With 124 GW and 104 GW respectively, hydro and non-hydro renewable technologies covered 38% of the cumulative power capacity in the Mediterranean region in 2015. If current trends continue, renewable energy technology will dominate the Mediterranean electricity market in the next years in terms of net generation capacity additions. In 2015, the net renewable energy capacity added was almost the half of the one of natural gas, which historically represented the first-generation source in the Mediterranean electricity mix (10 GW against 5.6 GW). In particular, the net additions of non-hydro renewable electricity capacity in were larger than 8 GW yr<sup>-1</sup> on average during the last 10 years.

The important growth in terms of new renewable electricity capacity expected will lead to a drastic restructuring of the power generation infrastructure. By 2040, renewables would in fact account for about 70% of total installed capacity and more than 50% of electricity generation in the PS. North Mediterranean countries are expected to add about 9 GW of new renewable capacity per year to reach a total of 410 GW by 2040 (thus more than doubling current power installed capacity). South and East Mediterranean countries will contribute some 6 GW yr<sup>-1</sup>, to reach 181 GW by 2040, a five times growth in the PS compared to current levels. This would completely change the electricity market supply and demand structure in South Mediterranean countries.

In terms of electricity generation, renewables will generate 1,137 TWh in 2040 in the RS, or 34% of total electricity generation in the Mediterranean. This implies an average annual growth rate of 1.3% for hydro, which would generate 357 TWh, and 5.1% for non-hydro renewables (780 TWh) over the period 2015-2040. In the Proactive Scenario, electricity generated from renewables is expected to reach over 1,438 TWh, around 52% of total production in 2040. This trend is influenced by 20% less growth in electricity generation in the Proactive Scenario than in the reference case, the progressive phase out of oil and coal-fired electricity production plants and a further boost to renewable energy technologies, both in North Mediterranean countries, where non-hydro renewables would experience a compounded average annual growth rate (CAAGR) of over 5.5%,


and in South and East Mediterranean countries, where non-hydro renewables are expected to grow at 11% CAAGR.

In terms of the outlook for the three fastest growing renewable energy technologies in the Mediterranean region, North West Mediterranean countries maintain their regional leadership in both scenarios for wind generation, with a projected electricity output of 267 TWh in the RS and 325 TWh in the PS by 2040, around 70% of wind electricity generation in the Mediterranean. The Proactive Scenario foresees a more accelerated rate of growth of wind in the South and East Mediterranean region, which would supply about one-third of the total wind-generated electricity by 2040 (over 148 TWh). The South and East Mediterranean should produce about 32% of the total solar CSP-based electricity in the region (15 TWh) by 2040 in the RS, and 43% (27 TWh) in the PS.

### 3.3.3.4 An NDC-based scenario for the Mediterranean region

While the energy transition raises common challenges for the entire region, the nature and scale of the challenges are different between the northern countries and SEMC. *Fig. 3.40* shows the demographic and energy projection in the Reference Scenario (RS) of OME MEP2018 which will be used as a reference in *Section 3.3.4.4* to discuss energy transition financing objectives. The countries on the northern shore have more than two decades of reforms that have gradually diversified their energy mix and controlled the growth of their energy demand, in a context of demographic stability. SEMCs are facing rapid population growth, which should lead to a significant increase in their energy demand. Their energy mix is widely relying on fossil fuels and fossil fuel export revenues which play a central role in the macro-financial balance of some countries (Algeria, Egypt, Tunisia) (MEF 2019).

Finally, compared to the RS by 2040, *Fig. 3.41* shows that the Proactive Scenario would lead to significant benefits, in terms of reduced energy dependence, energy efficiency, renewable energy growth and climate mitigation. More specifically, compared to the RS, the Proactive Scenario (PS) would reduce the energy dependency of the region by 45% (from 43% to 24%); as well energy demand and electricity generation would be reduced by 20%, each; the share of renewables in electricity production would be 57% higher in the PS compared to the RS. Finally, the PS would see a reduction of CO<sub>2</sub> emissions of 30%.




**Figure 3.40 | Demographic and energy projections in the Mediterranean in 2040.** Source: MEF (2019), based on OME (2018) Reference Scenario (RS).

### 3.3.3.5 Impact of climate change on energy resources

Although solar photovoltaics and wind power are growing rapidly, several scenario studies show that thermoelectric (fossil, nuclear, geothermal, biomass-fueled) power, together with hydropower, will most likely remain the dominant power-generating technologies during the whole of the twenty-first century (IEA 2018). Overall reduction in total power generation is projected under global


warming as highlighted in a study investigating wind, solar, hydropower and thermoelectric power generation evolution in Europe (Tobin et al. 2018).

Several studies using regional or global climate projections show that the western Mediterranean Basin is likely to undergo slightly decreasing winds in future decades (Hueging et al. 2013; Tobin et al. 2016) due to the poleward shift of the Hadley cell (Section 2.2.2.1). Surface wind speed declines remain moderate by the mid-century (generally of the order of a few percent). Wind speeds usually undergo small expected changes (Solaun and Cerdá 2020). By contrast, wind speeds are consistently projected to increase in the Aegean Sea (Tobin et al. 2015, 2016) where more persistent episodes of stable wind mill production regime are found (Section 2.2.2.4) (Weber et al. 2018). In any case wind resource is not threatened by climate change. Overall, the magnitude of change is small (< 5%) for all countries under a 1.5°C and 2°C global warming (Fig. 3.42; panel A).



**Figure 3.41 | Benefits of the implementation of the Proactive Scenario (PS) compared to the Reference Scenario (RS) at the horizon 2040** (OME 2018).

For a 3°C warming, most countries undergo changes with a magnitude also below 5% except for Portugal, Ireland and Cyprus where decreases in magnitude are expected to exceed 5%, approaching 10% for Cyprus. A 2°C warming does not systematically lead to higher change magnitudes than 1.5°C, while 3°C warming leads to stronger changes in most cases. In terms of individual cli-



(Country code: Appendix D)

**Figure 3.42 | Future changes in national wind power (A), solar PV power (B), hydropower (C) and thermo-electric power (D) production under +1.5°C global warming (yellow bars), 2°C (orange bars) and 3°C (red bars).** Changes are relative to the reference period 1971–2000. Colored bars correspond to the ensemble mean. The black thin error bars represent ensemble-mean confidence intervals (95% level based on the Wilcoxon-Mann-Whitney test). Adapted from Tobin et al. (2018).

mate model signals, the spread among the models is limited.


Climate change also impacts solar radiation and thereby solar energy resource. For solar PV generation (Fig. 3.42; panel B), projected ensemble mean impacts show moderate reductions for most countries in Europe, except for the Mediterranean countries, i.e., Portugal, Spain, Greece and Cyprus where changes are very small (but the magnitude and direction of changes are robust since most models agree). The overall magnitude of the signals is correlated with the amount of warming. Crook et al. (2011) and Wild et al. (2015, 2017) studied the global changes on PV and CSP outputs as a result from climate change, combining solar radiation with other factors such as cell temperature and wind. They found that solar PV output is generally declining in future scenarios worldwide except

over a few areas in Europe and the Mediterranean region where the expected increase in solar radiation is expected to overpass the temperature effect.

Over Mediterranean areas there is agreement among studies that radiation will increase and this will lead to an increase in solar PV output despite the temperature increase (Jerez et al. 2015; Gil et al. 2019). There is large spread in estimated amplitudes and also significant differences between regional and global climate modeling results (Bartók et al. 2017), which may be at least partly explained by the lack of aerosol changes in most regional climate models.

CSP potential is also expected to increase with climate change. North-West Africa (Morocco and North of Algeria) appears to be one of the areas of Africa where solar potential is increasing in





**Figure 3.43 | Impacts of climate change on annual mean streamflow (A) and water temperature (B) for RCP8.5 for 2040–2069 (2050s) relative to 1971–2000 (adapted from van Vliet et al. 2016b).**

climate change scenarios (Soares et al. 2019), but over the whole of North Africa projections indicate a small decrease in resource (Bichet et al. 2019).

Climate variability and the likelihood of heat waves and droughts are expected to increase in the Mediterranean (section 2.2.4.2; Raymond et al. 2019). This may have important impacts on water resources available for hydropower (e.g., Hamududu and Killingtveit 2012) and thermoelectric power generation (van Vliet et al. 2012b, 2016b) in Europe and Africa. Consistent decreases in streamflow are projected for the Mediterranean region (up to -30% to -50% in south and east according to scenarios, Fig. 3.7). Water temperatures continue to increase during the twenty-first century for RCP8.5 (+1.0°C to +2.0°C for 2050s; van Vliet et al. 2016b; Fig. 3.43).

Spatial patterns of changes in hydropower usable capacities strongly correspond with the projected impacts on streamflow, showing overall decreases in hydropower usable capacity the Mediterranean, with reductions in the annual hydropower capacities of 2.5–7.0% for the 2050s [RCP2.6–RCP8.5; van Vliet et al. 2016b] (Fig. 3.43; panel A). Thermoelectric power usable capacities are projected to decrease for more than 60% (RCP8.5) of the power plants (Fig. 3.43; panel B). Thermoelectric power plants in the Mediterranean are situated in areas with expected declines in mean annual streamflow (Section 3.1.4.1) combined with strong water temperature increases (Section 2.2.4.2), which both amplify restrictions on cooling water use. Fig. 3.43 (panel B) shows considerable reductions in thermoelectric power usable capacity in the Mediterranean region of 10–15% [for the 2050s, RCP2.6–RCP8.5].

Impacts of climate change on gross hydropower potential have also been studied for Europe. Gross hydropower potential refers to “the annual energy

potentially available when all natural runoff in a country is harnessed down to sea level (or to the border line of the country) without any energy losses” (Eurelectric 1997). Mean gross hydropower potential is projected to increase in northern, eastern and western Europe and to decrease in southern Europe (Fig. 3.42; panel C). Overall, higher warming results in stronger changes. Results for the individual regional climate model projections show that many individual signals are not significant and the spread amongst models is substantial. The most negatively impacted countries will be Greece, Portugal, and Spain. Impacts in these southern European countries can be reduced by limiting global warming. A warming of 3°C reduces hydropower potential by 15–20% while limiting to 2°C warming would keep decreases below 10%.

The usable capacity of thermoelectric power plants using river water for cooling is expected to reduce in all European countries due to a combination of higher water temperatures and reduced summer river flows (Fig. 3.42; panel D). The magnitudes of the decrease are about 5% for 1.5°C, 10% for 2°C and -15% for 3°C for most countries. Bulgaria, Greece and Spain will be the most strongly impacted (15–20% decrease). Results based on output of various climate models project significant changes and agree on the direction of changes as the spread among signals is limited. Robust and significant negative climate change effects are found, with a magnitude higher than for other power generating technologies in Europe (Tobin et al. 2018) (Fig. 3.44).


Regarding bioenergy, in the long term (>2040), biomass is projected to be a key option to meet the most stringent climate targets (Clarke et al. 2014; Rose et al. 2014; Rogelj et al. 2015), because of its ability to produce negative emissions in association with carbon capture and storage (Section 3.2.3.2).

Integrated assessment models involved in the IPCC 1.5°C report project that biomass will take a growing share in primary energy use in most regions of the world during the 21st century (Fig. 3.45). The main driver behind this result is the need to decarbonize the energy sector (Bauer et al. 2018) and the availability of other sources of renewable energy.


Future environmental and climatic conditions of the Mediterranean Basin reinforce uncertainties related to bioenergy sector development. The Mediterranean zone is expected to have huge shifts in the distribution of agricultural and forest areas with climate change making the region vulnerable (Fernández-Manjarrés et al. 2018). The current low primary production and the fragmentation of private lands could limit the development of the bioenergy sector. Furthermore, the future increase of dry conditions and extreme natural events, reinforce tough environmental conditions for the biomass sources production especially forestry, agricultural and energy crops. Fires can be favored with increased dry conditions making many forest biomass reserves vulnerable, therefore needing adequate management (Fernández-Manjarrés et al. 2018). Also, the less pregnant decarbonisation constraint and the availability of solar energy explain why the share of biomass is not changing much in SEMCs.

Agriculture in the Mediterranean area is particularly vulnerable to climate change due to its reliance on rainfed systems and the increased water shortages, which are anticipated in this region (Section 3.2.2.1). These are very likely to have a negative impact on crop yields, in particular for wheat in North Africa. A decrease in crop yields would directly affect the amount of land available to grow energy crops, which are scarce to start with, especially in the SEMCs. Herbaceous bioenergy crops (*Miscanthus*, *Panicum virgatum*, cardoon, *Arundo donax*) produced on marginal or abandoned land offer promising prospects in southern Europe with positive effects in terms of sustainability criteria (Pulighe et al. 2019). To ensure the feasibility of production with regard to environmental criteria, the choice of cultivars most resilient to water stress and agronomic management will play a critical role.

While there is a large body of work on the combined effects of CO<sub>2</sub> concentration rises and climate change on staple food crops (IPCC 2014), only few studies exist on lignocellulosic plants (Chum et al. 2011). One study on miscanthus (a perennial grass) concluded that this crop would no longer be suitable in the Mediterranean area by 2050 due to drought kill (Hastings et al. 2009). Conversely, a simulation study involving another perennial crop (*Arundo donax*) in northern Italy concluded that its



**Figure 3.44 | Impacts of climate and water resources change on annual mean usable capacity of current hydropower and thermoelectric power plants.** Relative changes in annual mean usable capacity of hydropower plants (A) and thermoelectric power plants (B) for RCP8.5 for 2040–2069 (2050s) relative to the control period 1971–2000 (van Vliet et al. 2016b).



**Figure 3.45 | Regional share of biomass in the primary energy use in the IPCC 1.5°C scenarios.** Source: data available from the IAMC 1.5°C Scenario Explorer hosted by IIASA available at <https://data.ene.iiasa.ac.at/iamc-1.5c-explorer/#/login?redirect=%2Fworkspaces>. ASIA: Asian countries except Japan, LAM: Latin America, REF: Countries from Reforming Economies of the Former Soviet Union, MAF: Countries of the Middle East and Africa, OECD90+EU: OECD90 and EU (and EU candidate) countries.

yield could increase by 20% within 2050 due to an increase in water-use efficiency permitted by higher ambient CO<sub>2</sub> concentrations (Cappelli et al. 2015). For any bioenergy project, the selection of the most appropriate feedstocks should clearly be based on the local conditions and contexts, but while there is an increasing body of data and technical knowledge available on energy crops (Laurent et al. 2015), there is a scientific gap as far as the impact of climate change is concerned, in particular in the Mediterranean. Finally, climate change should also reduce the amount of agricultural residues available for bioenergy, which projections usually consider stable. The same applies to forest residues or waste from the agri-food sector, although the latter may import raw material from other world regions to compensate for a decrease in local supply.


Biomass production in the Mediterranean coastal regions should eventually, be limited by the lack of available land suitable for bioenergy production (Daioglou et al. 2019) and by water constraints. In a scenario projecting a warming limited to 2°C, Near East and North Africa will become major importers of biomass with, by 2100, a level of biomass imports in monetary values higher than their current fossil fuel exports (Fig. 3.46; Muratori et al. 2016).

Factoring in food security and bioenergy production is usually seen as a dilemma in projections and

foresight studies (Tilman et al. 2009). Integrated assessment models show that the agricultural and forestry sectors can meet an increasing demand both for food and bioenergy products by various adjustments (increase in crop yields via higher input rates, expansion of cropland area) (IPCC 2018), but in the Mediterranean area it is likely that there is less room for such adjustments given the environmental and economic constraints. Overall the highest potential should come from residue and waste streams, for which there are unfortunately few estimates available, and the use of marginal land – pending a proof of concept that this strategy can be implemented at a large enough scale beyond simulation studies (Gelfand et al. 2013). In the rural areas of the southern Mediterranean Basin where the use of traditional biomass predominates, the generalization of more efficient bioenergy technologies (improved cookstoves in particular) would also reduce the pressure on biomass resources and make biomass available for more refined uses (for power and transportation fuels) (Chum et al. 2011; Mulinda et al. 2013).

### 3.3.3.6 Impact of climate change on energy demand

Climate change in the Mediterranean is expected to impact the energy consumption mainly as a decrease (respectively increase) in space heating



**Figure 3.46 | Global fossil fuels, biomass, and agricultural products financial flows in 2010 and in 2100** (Muratori et al. 2016).


(respectively cooling) demand, all other factors remaining fixed. These results from a number of studies (see below) are derived from climate change projections from general or regional climate models reviewed by IPCC’s 4th and 5th assessment reports, in combination with econometric analyses or with bottom-up energy models. The net change in the yearly energy demand for heating and cooling in the 21st century associated with climate change depends on location via both the distribution of temperatures and the thermal sensitivity of energy consumption (Wenz et al. 2017). Thermal sensitivity may be defined as the change in mean energy consumption or in peak load associated with a unit change in temperature. Thermal sensitivity is usually negative for low temperatures, due to increased energy consumption associated with heating and positive for high temperatures, due to increased energy consumption from cooling. Thus, thermal sensitivity depends on local efficiency of appliances and isolation and on consumer behavior.

Due to heating, countries historically experiencing relatively low temperatures, such as in northern

Europe, are expected to see a net decrease in energy demand (Damm et al. 2017; Wenz et al. 2017) as opposed to countries which already or which will experience high temperatures, such as most Mediterranean countries (Giannakopoulos et al. 2009; Eskeland and Mideksa 2010), which are expected to see a net increase in the energy demand due to cooling. This is illustrated by the maps of Fig. 3.47 which represent changes in average daily peak electric load at the end of the 21st century relative to the beginning of the century estimated by Wenz et al. (2017) for the RCP4.5 (A) and RCP8.5 (B) scenarios. Differences between countries may not only be due to differences in temperature distributions, but also to varying thermal sensitivities between countries. Annual demand values do not reflect the seasonality of these changes, which may significantly impact the seasonal planning of energy resources.

Using regional climate models or statistical downscaling, case studies for specific countries such as Algeria (Ghedamsi et al. 2016), Cyprus (Zachariadis and Hadjinicolaou 2014), Slovenia (Dolinar et al. 2010), or Spain (Pérez-Andreu et al. 2018), offers





**Figure 3.47 | Percentage change in average daily peak electric load from 2006–2012 to 2080–2099 for projected daily maximum temperatures under RCP4.5 (A) and RCP8.5 (B) climate change scenarios.**

While daily peak load decreases in northern European countries, it increases in southern and western European countries. This trend is most pronounced for a scenario of unabated climate change (RCP-8.5, B) but still holds for a scenario of mitigated climate change (RCP-4.5, A) (Wenz et al. 2017).

a more detailed picture showing that the expected distribution in demand changes may depend on local climates and that extremes in both heating and cooling demand may intensify even if the demand is reduced on average.

Energy demand may also be indirectly affected by climate change as impacts on the agricultural sector associated with reduced precipitation in North Africa may motivate replacing biomass as traditional energy source in rural areas (Schilling et al. 2012).


### 3.3.4 Adaptation and mitigation

#### 3.3.4.1 Adaptation of energy systems to water constraints

Adaptation measures to mitigate the vulnerability of the electricity sector to future water constraints under changing climate have been investigated in the form of six options (van Vliet et al. 2016b). Options 1 and 2 assume increases in efficiencies of hydropower plants and thermoelectric power plants. Option 3 assumes replacement of fuel sources of thermoelectric power plants (coal- and oil-fired plants replaced by gas-fired plants). Option 4 assumes replacement of once-through cooling systems by recirculation (wet tower) cooling systems. Option 5 assumes switch to seawater cooling for thermoelectric power plants close to the coast (<100 km), and option 6 assumes decoupling from freshwater resources by switch to seawater and dry (air) cooling for 10% of the thermoelectric power plants that are most vulnerable to water constraints under climate change (Fig. 3.48).

Increasing total efficiencies of hydropower plants up to 10% is able to completely offset the mean annual impacts of increased water constraints under changing climate for most regions, including the Mediterranean area (Europe and Africa) (Fig. 3.48; panel A). For thermoelectric power, increased power plant efficiencies also positively contribute in reducing water demands and decreasing the vulnerability to water constraints under climate change (Fig. 3.48; panel B). A strong increase in power plant efficiencies up to 20% is for most regions still insufficient to mitigate overall reductions in cooling water use potential under changing climate. This is the case for Europe, but not necessarily for Africa.

Changes in sources of fuel and switches in cooling system types (from once-through to recirculation with wet cooling towers or dry cooling) are for most regions more effective in reducing plant vulnerabilities to water constraints (Macknick et



**Figure 3.48 | Impacts of adaptation options on power-generation vulnerability to water constraints under climate change.** Relative changes for the baseline settings and for various adaptation options of hydropower (A) and thermoelectric power (B). The GCM-ensemble mean changes are presented by the bars. In addition, changes for the five individual GCM experiments for RCP8.5 (2050s) are presented to show the range between the five different GCM experiments (van Vliet et al. 2016b).

al. 2012; van Vliet et al. 2016b). On average, fuel switching to higher efficiency gas-fired plants with lower cooling water demands can be sufficient to mitigate plant vulnerability to water constraints for the 2020s and 2050s under RCP2.6 scenario. This adaptation option will be insufficient for Europe under RCP8.5 scenario in the 2050s. The strongest positive impacts relative to the baseline settings are found for Africa, where the relative number of coal-fired plants that can be substituted by gas-fired plants is high.

A switch to recirculation (wet tower) cooling will decrease water withdrawals and reduce plant

vulnerabilities to water constraints. A switch from freshwater cooling to seawater cooling for plants nearby the coast also reduces vulnerabilities to freshwater constraints. Decoupling of cooling water systems from freshwater resources (by switching to dry cooling or sea water cooling) for the 10% most severely impacted plants is a more effective adaptation option.

On top of these options, a higher share in non-water dependent power generating technologies, such as solar PV and wind power, will strongly reduce the dependency on freshwater resources. The uncertainty range is considerable and strongly

depends on the relative share of different technologies within the energy transition (e.g., Mouratiadou et al. 2018).

### 3.3.4.2 Integration of renewables into the energy mix

In view of the current dynamics of energy supply and demand, a large gap is expected between anticipated demand and supply, particularly in SEMCs (Hawila et al. 2012). In order to address both the depletion of fossil fuels and climate change issues, it is necessary to improve rapidly the penetration of renewable energies. To this end, measures have already been taken to shift the energy mix in the SEMCs from fossil fuels to renewables. Large-scale deployment of renewable energy technologies and a necessary profound transformation of the energy sector are still lagging behind in some countries in the region (Vidican 2016).

#### *Incentive effects of support mechanisms for renewable energy*

SEMCs already consume a lot of energy and their energy demand is expected to continue to grow in the coming decades (El-Katiri 2014). As a result, energy plays an increasing role in national development policies in most of these countries. Given their climatic advantage, particularly in solar energy, the countries of the region therefore want to seize their comparative advantage. To achieve effective integration of renewable energy in these countries, a key solution would be to reform energy pricing mechanisms in the region and promote the participation of private actors in order to address the distortions of the energy market in the region. If this cannot be achieved for any reason, an alternative solution could be the introduction of tax and regulatory incentives aimed at reducing remaining cost disadvantages of renewable energies compared to fossil fuels (El-Katiri 2014). Since energy prices for the final consumer are heavily subsidized in most SEMCs renewable energy sources could also receive subsidies to make renewable energy production viable. Poudineh et al. (2018) argue that renewable energy support policy for SEMCs should meet criteria such as compatibility with the structure of the region's electricity system, harmonization with existing institutions, suitability at the project scale, coverage of economic risks and provision of efficiency.

Large-scale deployment of renewable energy could have significant positive impact on SEMCs. By examining the implications of renewable energy deployment in the SEMCs through the LEAP (Long

Range Energy Alternatives Planning System) forecasting model, El Fadel et al. (2013) find that on a regional scale: (i) per capita greenhouse gas emissions could be drastically reduced, (ii) the return on investment in renewable energies promises up to 54% savings excluding positive externalities and (iii) the establishment of a CO<sub>2</sub> emissions trading market would provide economic incentives that make investment in renewable energies more attractive. An assessment by Timmerberg et al. (2019) of national renewable energy production targets and a diagnosis of the current energy sector in some SEMCs shows that the plans currently in place to expand renewable energy in these countries will be able to roughly meet the expected future growth in national electricity demand. Therefore, they estimate that if the targets set by the states for 2030 are met, CO<sub>2</sub> emissions per kWh, for example, could drop drastically to 341-514 g CO<sub>2e</sub> kWh<sup>-1</sup> compared to 396-682 g CO<sub>2e</sub> kWh<sup>-1</sup> in 2017.

Using a cost-minimizing electricity market model, Brand (2013) explores the option of optimized infrastructure for the integration of renewables into the interconnected North African electricity grids until 2030. The results show that the five countries, Morocco, Algeria, Tunisia, Libya and Egypt, could together achieve significant economic benefits of up to 3.4 billion € if they increase electricity grid integration, build interconnections and cooperate in the joint use of their power generation assets. The challenge remains to eliminate political obstacles to cooperation. Brand and Blok (2015) also assessed several studies (including Haller et al. 2012; Fragkos et al. 2013) using economic models of electricity supply and demand to assess the possible development paths of electricity systems in the North African region from today to 2030 (or even 2050). All studies agree that additional costs associated with the expansion of renewable energy in SEMCs could in most cases be offset by avoided fuel costs and avoided investments in fossil fuel power plants. Even electricity exports to Europe could become viable.

Despite the remarkable political objectives announced by SEMCs leaders to increase the deployment of renewable energy to meet growing energy demand and mitigate climate change in the region, there are still some inconsistencies between the approaches and the objectives. The deployment of renewable energy faces many obstacles in the region, namely, the state monopoly on the electricity sector, persistent fossil fuel subsidies, and weak regulatory institutions and bureaucratic issues (Lilliestam and Patt 2015). Al-Asaad (2009) notes mainly technical problems. For example, it points

out that much of the region's network infrastructure (e.g., sub-stations and power lines) and network codes have long since become obsolete and therefore need to be upgraded to effectively meet the objectives of diversifying energy sources.

Finally, regarding renewable energy costs, Krupa and Poudineh (2017), using simulations, show that the levelized cost of energy (LCOE) of renewables is more sensitive to the discount rate compared to fossil energy sources in SEMCs. This is certainly due to the fact that the investment costs are very high whereas the production costs of fossils are more evenly spread over the life of the project.

### ***The specific case of bioenergy: barriers and opportunities***

There are four main barriers to develop bioenergy in Mediterranean area: financial, technological, social and institutional. Technical barriers include investment costs and lack of experience in biomass conversion technologies, and development of sustainable management of biomass sources. The regulation barrier requires new policies for bioenergy promotion, and socio-economic and environmental issues analysis. Financial decisions by funding agency adds the problem of competitiveness with other industry and energy markets. In fact, the development of bioenergy represents a new competitive sector because of the use of sources already used in industrial sectors as the paper industry or agriculture (Ferreira et al. 2017). For now, cultivation of energy crops is limited because of farmers' reluctance considering uncertainty of productivity and economic markets, and less attractive prices compared to cereal crops (Pulighe et al. 2019). In the same way, wood-biomass energy still remains low owing to the weak economic power of the forestry sector and the hard biodiversity regulation (Cavicchi et al. 2017).

The development of bioenergy implies increasing the production of local biomass sources, which raises ecological, economic and socio-political issues. Increasing biomass sources can be effective through tree plantations, forest management improvement, energy crops implementation, agricultural sources plantations, among others. Yet, these new biomass sources at large-scale production could increase greenhouse gas emissions considering the land-use changes associated (Fargione et al. 2008; Gibbs et al. 2008; Searchinger et al. 2008). In the Mediterranean Basin, tree plantations focus on *Eucalyptus* spp., *Populus* spp. and *Pinus* spp., but these species have high ecological impacts during the clearcutting and plantation operations, and soil

preparation (Rodríguez-Loinaz et al. 2013). The development of agricultural energy crops would be competitive if some of the following situations happen: distinctive quality allowing competitive prices, new market opportunities that cover production costs, providing subsidies to these cultures by environmental reasons (Ferreira et al. 2017).

The development of bioenergy represents a larger opportunity than simple renewable and carbon neutral fuel source. The implementation of a holistic bioenergy sector can play a crucial role in the transition by improving Mediterranean forests management, promoting new socio-economic models and rethinking the relation between human and nature (Paredes-Sánchez et al. 2019; Sansilvestri et al. 2020). The bioenergy sector in Mediterranean Basin is an opportunity for land restoration and abandonment issues because bioenergy crops can be implanted on abandoned or polluted lands without sacrificing pastures, arable or protected lands (Pulighe et al. 2019).

It has been proposed that the evolution of the renewable energy based on biomass depends on two interacting variables: the strength of policy for centralization and the development of competitive markets for biomass-based alternatives. These two interacting axes will determine, by cascade effects, how much centralized policies will favor large industrial developments in contrast to a more widespread biomass conversion where sustainability of the resource is paramount. The development of biomass-based solutions also depends on how competitive solar and wind base solutions are (Section 3.3.2.2). In any case, northern Mediterranean countries have more margin of strategy evolution than southern ones, as biomass production is inherently much more restricted in the southern region.

Bioenergy has a place in a new economical organization with a clear position, rules and limits. Bioenergy can valorize residual productions from other industries, which can be upgraded in a bioeconomy sector. Pulighe et al. (2019) suggest that future support schemes and business models for mobilizing financing and attracting investors should be more aligned with greenhouse gas emissions, ecosystem services and sustainability indicators, avoiding criticism raised for the biogas sector regarding tradeoffs on land use pressure for biomass. For the time being, bioenergy is mostly based on an incentive economy, tax credits and fiscal exemptions, but to increase engagement of industry, foresters and farmers, it needs clear economic possibilities with a long-term visibility.



### 3.3.4.3 Energy access in developing countries of the Mediterranean region

Unlike the sub-Saharan African zone (only 45.6% electrification rate), SEMCs do not yet have major difficulties in providing electricity to their populations. The energy costs supported by households (very often subsidized) are largely lower compared to the world average. According to World Bank data, the electrification rate was almost 100% in 2017 for Algeria, Morocco, Tunisia, Egypt, Israel, Jordan and Lebanon. Libya is the only country with a lower rate (70.1%), due to the political instability (it had an electrification rate of 99.8% in 2000). Most analyses find that the energy dynamics of these countries are unsustainable in the long term, for several reasons. First, the region is experiencing strong demographic growth and prospects for strong economic growth are also expected. Therefore, energy demand is expected to grow strongly in the future for the net energy exporting countries (Algeria, Libya, Syria, Iran etc.) and for the net importing countries (Morocco, Tunisia, Egypt, Jordan and Lebanon) of the area (difficult combination of future depletion of reserves and energy subsidies). Second, the region is one of the most affected by the effects of global warming. Countries are therefore compelled to make an energy transition to renewable energy sources as quickly as possible and gradually move away from fossil energy sources. Finally, price control policies such as energy subsidies are quite widespread in the region. These policies are still costly to state budgets and quite ineffective. Energy subsidies frequently distort price signals and lead to systemic misallocation of resources (Fattouh and El-Katiri 2013). The main concerns are therefore how to produce enough renewable energy to meet the anticipated high future energy demand in the region, still considering potential political and economic barriers.

#### **Financial, behavioral, institutional and regulatory impediments to energy access**

The SEMCs are known for their oil and gas reserves, but also for their immense potential in renewable energies, particularly solar and wind energy. According to estimates, with only 0.2% of the land suitable for concentrating solar power, plants in seven countries of the region (Morocco, Tunisia, Algeria, Libya, Egypt, Jordan and Saudi Arabia) could cover 15% of the electricity demand expected in Europe in 2050 (Trieb et al. 2012). Thus, there are economic as well as environmental reasons to switch to renewable energy sources. But for those countries in the region that are highly

dependent on fossil revenues (oil, natural gas), it would be very difficult to switch entirely to green forms of energy (El-Katiri 2014). The economies of these countries largely depend on the rents from hydrocarbons in the 1960s and 1970s. Also, the social contract between the government and citizens is based on the fact that energy is virtually free, making the investments into the transition difficult.

For the reasons mentioned above, the countries of the area need to make their energy transition quickly. By the end of 2013, some of them had set targets for the share of renewables in total electricity generation, in total installed power generation capacity or in total electrical and thermal energy (El-Katiri 2014). Algeria has set a target of 15% of electricity generation by 2020 and 40% by 2030, Morocco (42% of installed power generation capacity by 2020), Tunisia (25% of electricity generation by 2030), Egypt (20% of electricity generation by 2022) and Lebanon (12% of electrical and thermal energy by 2020). Despite high penetration of solar water heaters in Palestinian households (56%), solar thermal energy represents only a small fraction of the Palestinian Energy mix (8%). Transitions to renewable energy sources require huge investments, and therefore large-scale regional and international projects such as joint projects with the European Union. The realisation of large-scale energy projects between Europe and the region, such as the Mediterranean Solar Plan or Desertec, faces several barriers. Fritzsche et al. (2011) mainly refers to political constraints, such as the lack of subsidies, incentives and liberalization of the renewable energy market in the region. Authors including de Souza et al. (2018) rather refer to cultural barriers to renewable energy trade projects in the Mediterranean (between the EU and southern countries), alluding to the failures of energy cooperation projects between Europe and its former colonies.

While voices are being raised in the North of the shore to denounce what should increase Europe's energy dependence, and therefore additional diplomatic pressure (Lilliestam and Ellenbeck 2011), in the South, there is talk of a new form of resource exploitation, and therefore of the notion of solar colonialism (Marktanner and Salman 2011; Rowlinson 2015). There are three main risks to their implementation, risks that affect the cost of capital in particular: (1) the complexity and instability of national regulations, (2) low political stability and (3) terrorist threats in the region (Komendantova et al. 2011). The World Bank has repeatedly pointed out that among the constraints to investment in

the renewable energy sector in the region are the narrowness of local markets (which discourages private investment in particular) and the need to consider a regional energy trading market, the weakness of public finance and the competition of energy investments with other public service and public infrastructure investments. Far from the reality of the major oil exporting countries of the region, Morocco depends heavily on its energy imports, i.e., 95% of its consumption in 2007 (Fritzsche et al. 2011), despite a rather remarkable potential in renewable energy. This energy deficit partly explains the strong pressure on traditional energy sources (wood, coal, etc.). According to Zejli et Bennouna (2009), between 30,000 and 50,000 hectares of forest disappear every year. Aware of this and aiming in particular at reducing its dependence and energy bill, the country is widely involved in projects and initiatives aimed at promoting renewable energy. With this in mind, in November 2009 the country launched the pharaonic Noor project (I, II and III) in Ouarzazate at an estimated cost of 2.5 billion US\$ (9 billion US\$ for the full Moroccan Solar Plan), a project that should eventually provide electricity to more than 2 million people in the country.

### **Regulatory framework and business models to reach universal access**

According to de Souza et al. (2018), to facilitate the successful implementation of renewable energy projects, it is particularly important that the countries of the South are sufficiently integrated into the value chains of production, from distribution to transportation of energy (acceptability), put in place a plan to mitigate the environmental impacts of site operations and facilitate technology transfer. Fritzsche et al. (2011) point out that specialized agencies should be created to overcome political constraints to facilitate the establishment of renewable energy facilities in the area and multiply tax incentives to help attract foreign direct investment. They also point out that funding and technical assistance from international and regional donors and development banks are essential for the implementation of these types of projects. They also argue that technology transfer from North to South should be promoted, R&D should be encouraged and EU-SEMCs and PPP (public-private partnership) type partnerships should be implemented. Blimpo and Cosgrove-Davies (2019) argue that it would be viable to establish a regional energy exchange market between the countries in the area, since energy trade would halve the cost of kWh. As private investors are more attracted to large markets, establishing a common regional market will

also allow small economies and countries that are not abundant in energy resources to benefit from cheaper electricity. As an example of an integration model, Blimpo and Cosgrove-Davies (2019) mention the example of the West African Power Pool (EEOA), which brings together 14 countries of the zone and 27 companies. This system would have enabled many countries in the West African zone to benefit from an affordable, reliable and sustainable electricity supply (typical case of the Manantali dam on the Senegal River, which benefited from a lower investment cost to the benefit of Mali, Senegal and Mauritania). El-Katiri (2014) emphasizes the need to reform the system governing the energy price mechanism (local market prices strictly controlled by the State to its liking) at the national and regional levels and to implement a fiscal policy aimed essentially at reducing the cost disadvantage of renewable energies compared to fossil fuels.

### **The Green Transition in North Africa and the Middle East**

With the exception of Libya, the other six states in the North African region have ratified the Paris Agreement of 2015. For countries in North Africa, the energy transition requires a profound structural transformation of the energy sector and all other sectors of the economy, but one of the major difficulties of this structural transformation is to effectively find a compromise between long-term and short-term political and economic objectives (Pye and Bataille 2016).

Another major difficulty remains the question of financing, as green projects are unfortunately still quite expensive and require large investments. To illustrate, estimates have shown that for some 28 countries on the African continent, for example, meeting national commitments for energy transition would require an overall investment of more than 240 billion US\$. On the other hand, these investments would be quite beneficial for the economies of North Africa, particularly in view of their potential for renewable energy. Brand and Zingerle (2011) show that for every € spent on renewable energy, 0.15, 0.16 and 0.27 € of savings could be made in the national electricity system in Tunisia, Algeria and Morocco respectively. Also, according to Alnaser and Alnaser (2011), the costs of generating electricity from photovoltaic systems and wind turbines have been on a downward trend for several years. In some SEMCs, it is already possible to produce green power much cheaper than electricity production from fossil sources (IRENA 2018). The competitiveness of renewable energy

sources depends not only on the technology used for energy production and the availability of energy resources, but depends especially on the cost of fossil fuels (substitutes for renewable energy). The costs of fossil fuels are much lower for most SEMCs compared to other regions of the world. This does not promote the competitiveness of renewable energies.

Timmerberg et al. (2019), based on an assessment of national renewable energy production targets and a diagnosis of the current energy sector in selected SEMCs (Algeria, Egypt, Morocco, Saudi Arabia and Tunisia), show that current plans for renewable energy expansion in these countries will be able to approximately meet the expected future growth in national electricity demand. They estimate that if the targets set by the States for 2030 are met, CO<sub>2</sub> emissions per kWh, for example, could drop drastically to 341-514 g CO<sub>2</sub>e kWh<sup>-1</sup> (compared with 396-682 g CO<sub>2</sub>e kWh<sup>-1</sup> in 2017).

### 3.3.4.4 Financing the energy transition

As discussed in the previous sections, the energy transition requires a significant transformation of the energy and economic model in the Mediterranean region. Mitigation and adaptation will require

investments from households, companies and governments, which in turn require sources of financing. Policymakers in Mediterranean countries need to focus policies and public investment to achieve three main objectives: i) to bring the benefits of the new energy technologies to citizens via accelerated uptake of distributed generation, ii) incentivize the deployment of utility-scale PV parks, preferably including storage, and iii) to ease the opening of new energy technology factories in their countries. Today's low-cost solar energy and electricity storage technology, make the first objective achievable thanks to new legislation supporting distributed generation for example updating obsolete construction regulation making long and costly the permit process to functionalize buildings with solar modules. The growth potential is huge as less than 5% of the world's buildings, and even a lesser fraction of the large Mediterranean built environment, are functionalized with solar collectors. Achieving the second objective in industrialized countries such as Spain, Italy, and France is feasible by allowing renewable energy generation companies to take part not only to the day-ahead electricity market but also to the dispatching and energy services markets, so far opened to thermal production units only. Sunnier SEMCs will preferably continue with tenders to purchase clean electricity produced via utility-scale PV plants at prices





Figure 3.49 | Economic and financial contexts around the Mediterranean (MEF 2019).



**Figure 3.50 | Change in infrastructure spending for a 2°C scenario in the Mediterranean region, percentage change in expenditure over 2015-2030 compared to Business-as-usual.** (MEF 2019).

that in the case of Egypt have already reached very low levels. Finally, the third goal is to make their countries home to new industrial plants of the solar economy, requires to concentrate the financial resources on establishing new partnerships with the leading renewable energy and energy storage technology manufacturers, which are not based in Mediterranean countries. There are many ways to incentivize foreign companies to invest, as shown by Morocco with their new electric vehicle plant; or by Algeria, requiring renewable energy companies to use components made in Algeria for the utility-scale PV plants awarded the Public-Private Alliances.


**Financial needs by sector and region**

Beyond the economic contexts, public finance capacities of SEMCs are subject to strong tensions, with a high weight of the treasury’s debt in terms of GDP. SEMCs financial systems are also characterized by a domination of banking, a weak development of financial markets and a low degree of international market openness (Fig. 3.49). In the Mediterranean region, financing needs for SEMCs

NDC are estimated at more than 170 billion US\$, of which 81% are related to conditional NDC (MEF 2019).

While there is no specific figure for the Mediterranean region, Western Europe benefited from 106 billion US\$, while Middle East and North Africa benefited from 13 billion US\$ during the period 2017/2018 (CBI 2019). Focusing on international public climate finance, Egypt, Turkey, and Tunisia were the top-3 recipients in 2017, altogether representing 5.6 billion US\$ (of 8.1 billion US\$ for the SEMCs) (UfM 2019). International public climate finance captured by the SEMCs has shown an encouraging trend since 2012, since the amounts committed have increased by an annual average of 21.5% over the 2012-2017 period (MEF 2019). In the Mediterranean region, 135 billion US\$ per year in the energy sector will be needed until 2040 (approx. 1% of each country’s GDP) and will have to be redirected from fossil fuels to renewables and energy efficiency (IEA 2018; IPCC 2018). Natural gas and renewables will dominate future investments in power generation across the region and close to 44 billion € per year will be needed to reach the





**Figure 3.51 | International public climate finance by sector in SEMCs, 2017** (UfM 2019).

energy demand levels of the Reference Scenario (40% increase in demand) while energy efficiency investments will need to reach 61 billion € per year (Fig. 3.50). About 55% of these energy investments will be required on the north shore, 25% in North Africa and 20% in the South East (OME 2018).

In the 2017/2018 period, the following financial flows in SEMCs went to mitigation, and specifically to renewable energy generation (1.6 billion US\$), water supply and sanitation (1.4 billion US\$), transport and storage (1.2 billion US\$) and agriculture (0.6 billion US\$) (UfM 2019; Fig. 3.51).

Currently, adaptation flows account for a small percentage of total climate finance, illustrating what Abadie et al. (2013) call the 'mitigation bias'. This bias is also present in the SEMCs (Fig. 3.51). Beyond investment in hard projects for mitigation and adaptation of infrastructures, financial flows are needed to push R&D capabilities in both northern and SEMCs.

### Financial actors

Different actors are expected to finance climate change mitigation and adaptation. In the 2017/18

period private actors represented 326 billion US\$ while public actors represented 253 billion US\$ of annual flows (CBI 2019). While 75% of climate finance is invested in the same country in which it is sourced, developed countries have pledged to raise 100 billion US\$ per year by 2020 for global climate action to developing countries (UNFCCC 2018), including both public and private sources.

Climate-related multilateral funds, such as the Green Climate Fund (GCF) and the Adaptation Fund (AF), remain less important (0.39 billion US\$). Major bilateral donors are member countries of the Development Assistance Committee, including France (0.88 billion US\$), Germany (0.75 billion US\$), EU institutions (0.62 billion US\$) and Japan (0.43 billion US\$ in 2017 after 1 billion US\$ in 2016) (UfM 2019). Since 2013, the contribution of multilateral development banks has been steadily increasing, in line with their commitment taken at COP24 in December 2018 to align their activities with the goals of the Paris Agreement. On the northern shore, EU countries committed to dedicate at least 20% of the EU budget from 2014-2020 to climate-related actions (representing around 180 billion €, threefold increase from the 6-8% share in 2007-2013<sup>30</sup>).

<sup>30</sup> Source: [https://ec.europa.eu/clima/policies/international/finance\\_en](https://ec.europa.eu/clima/policies/international/finance_en)




Figure 3.52 | International public climate financial instruments by SEMCs recipient countries, 2017 (UfM 2019).

### Financial instruments

Worldwide, debt is the main financial instrument of climate finance. SEMCs benefited mostly from debt instruments from international public finance (Fig. 3.52). Financing the energy transition goes beyond debt, equity and grants. Above the various mitigation mechanism, putting a price on carbon with tax or quotas can encourage to consume and invest in goods and equipment that emit less greenhouse gases and generates a complementary source of financing for the energy transition. In the Mediterranean region, carbon price mechanisms are only present on the north shore. The EU Emission Trading System (EU-ETS), launched in 2005, is the world’s biggest emissions trading system, accounting for 45% of total EU greenhouse gas emissions. Different countries have also applied carbon taxes (e.g., France, Portugal, Slovenia in the Mediterranean region).

Another tax measure to accelerate the energy transition and provide financing is to reduce fossil fuel subsidies. Subsidies were estimated to 130 billion US\$ for SEMCs in 2015 (OECD 2018). These subsidies allocated to fossil fuels reach significantly high levels in countries such as Egypt, Lebanon and Algeria while Tunisia, Jordan or Morocco have relatively more moderate levels (Fig. 3.53). In Egypt, a program was adopted in 2014 for gradually subsidy phase out taking into consideration the social impacts, with a foreseen deadline in 2021.




Figure 3.53 | Energy subsidies in SEMCs (World Bank 2018).

Among the new instruments of the climate finance, green bonds are bonds issued by companies, governments or local authorities on the financial markets to finance a project or activity with an environmental benefit and are subject to standards (Green Bonds Principles). Worldwide, \$168 billion were issued in 2018 (40 billion US\$ in 2014). Since 2007, Europe accounts for more than 36% of green bonds issued while Africa accounts for less than 0.4% (CBI 2019).

### 3.3.4.5 Supra-regional cooperation for energy transition

Faced with this economic, social and environmental situation, energy and climate challenges are of

major importance. Accelerating the energy transition in the Mediterranean would help to control the growing demand for energy, to promote renewable resources and finally to optimize the use of fossil resources. The optimisation of the regional energy system, would pass through a better integration of markets, increased interconnection and intelligent management of networks, including the facilitation of access for renewable energy and demand side management.

Energy should be considered as part of the larger process of political shifting to a more inclusive, democratic and sustainable development paradigm (new social contract) that concentrates on a fair split of resources, opportunities and the results of growth all the while ensuring the right of all people to equitably participate in decisions that shape the future of their societies. This approach mirrors the dominant one that demands more decisive investment in renewable energies, but mainly as a tool for socio-economic development. For investments in renewable energies to have concrete value for the larger sustainable development goals of a country, projects should include technology transfers and capacity building for the population. In that regard, new initiatives for promoting renewable energies in the Mediterranean should avoid the 'Eurocentric focus' of the Mediterranean Solar Plan, which was seen as an instrument to support mostly the interests of European companies. Studies of the implementation of renewable energy projects in the SEMCs have also alerted on their consequences in terms of reinforcing the private sector and central government at the expense of local populations. In brief, if Euro-Mediterranean energy cooperation wants to thrive, it is necessary to reconsider who matters in energy security, from market and state actors to society at large, and consider particularly marginalized sectors of the population.

The Euro-Mediterranean dialogue, under the chair of the European Commission and the Kingdom of Jordan, at the Rome conference in November 2014, decided to establish three platforms for exchange and cooperation. The ultimate goal for these platforms is to operate as permanent consultation forum on strategic objectives and measures to be implemented under the auspices of the Union for the Mediterranean. The three platforms cover: (i) the gas sector managed by the Mediterranean Observatory of Energy (OME), (ii) the electricity market with support provided by the Association of Mediterranean Regulators (MEDREG) and the Association of Mediterranean Transmission System Operators (MEDTSO) and finally (iii) the renewable

energy and energy efficiency with support of MEDENER and RCREEE.

Many studies have shown that energy cooperation projects with North Africa aimed at large-scale energy production and exchange (mainly solar and wind power) would be an ideal instrument to achieve greenhouse gas reduction (Komendantova et al. 2012). Other studies have also analysed the barriers that constrain these energy projects, which Flyvbjerg et al. (2003) describe as "mega-projects" because they require huge costs. All of these studies are based on a perception of risk at two levels: regulatory and political risks, and force majeure or security risks (including, in particular, the terrorist threat in the region).


### *Mediterranean energy market integration*

Achieving the objectives of the Paris Agreement requires a massive scale-up of renewable energy sources and regional energy market integration facilitates the large-scale development of renewable energy, as it increases the area over which electricity supply and demand must be balanced in real time, making it less likely that the resource will not be available when needed. Regional energy market integration offers numerous benefits to the power systems and the economies of participating countries: enhanced energy security and power system reliability, reduced need for back-up capacity thanks to reserve sharing, supply mix diversification, more efficient use of power plants, lower power system costs (both investment and operating) and therefore lower consumer prices (World Bank 2010; UK DECC 2013). With more ambitious climate mitigation objectives since the December 2015 Paris Agreement, the climate benefits of regional integration are increasingly acknowledged as being as important if not more than the energy and economy benefits. Some of the climate benefits result from the increased efficiency of the power system, but most of them are derived from the fact that regional integration increases power system flexibility, and therefore facilitates renewable energy scale-up.

Although several studies have estimated the costs and benefits of the integration of electricity markets, particularly in Europe (Bockers et al. 2013; Newbery et al. 2016), there is less research on the preconditions and the required policies for establishing a successfully integrated market for electricity and a truly seamless transmission system (Oseni and Pollitt 2016; Roques and Verhaeghe 2016). Newbery et al. (2016) estimate the benefits of integrating the European Union (EU)

# Physical energy flows across Europe

GWh average over the year



<sup>1</sup> Consolidated yearly values might differ from detailed flow data from the ENTSO-E database due to ex-post consolidation taking into account national statistical resources.

<sup>2</sup> All data with the country code GB represents monthly statistical data as sum of England, Northern Ireland, Scotland and Wales.

	Sum of imports	Sum of exports	Balance (imp-exp)		Sum of imports	Sum of exports	Balance (imp-exp)
<b>AL</b>	1,771	2,683	-912	<b>IT</b>	47,169	3,268	43,902
<b>AT</b>	29,393	19,057	10,336	<b>LT</b>	12,850	3,219	9,631
<b>BA</b>	3,091	7,796	-4,605	<b>LU</b>	7,514	1,349	6,166
<b>BE</b>	21,650	4,313	17,338	<b>LV</b>	5,179	4,272	907
<b>BG</b>	2,220	10,029	-7,809	<b>ME</b>	2,760	3,011	-251
<b>CH</b>	30,420	31,693	-1,274	<b>MK</b>	4,144	2,224	1,921
<b>CZ</b>	11,562	25,453	-13,891	<b>NL</b>	26,818	18,596	8,223
<b>DE</b>	31,542	82,673	-51,131	<b>NO</b>	8,085	17,954	-9,869
<b>DK</b>	15,606	10,413	5,193	<b>PL</b>	13,839	8,121	5,718
<b>EE</b>	3,514	5,364	-1,850	<b>PT</b>	5,669	8,324	-2,655
<b>ES</b>	24,014	12,910	11,104	<b>RO</b>	2,829	5,370	-2,541
<b>FI</b>	23,397	3,459	19,938	<b>RS</b>	7,300	6,703	597
<b>FR</b>	13,466	76,020	-62,554	<b>SE</b>	14,234	31,561	-17,328
<b>GB</b>	22,662	2,189	20,473	<b>SI</b>	8,928	9,320	-392
<b>GR</b>	8,552	2,265	6,288	<b>SK</b>	12,544	8,747	3,797
<b>HR</b>	12,692	6,533	6,160	<b>TR</b>	2,638	3,046	-408
<b>HU</b>	18,613	4,265	14,348	<b>ENTSO-E</b>	<b>458,274</b>	<b>443,734</b>	<b>-14,540</b>
<b>IE</b>	1,614	1,643	-29	Physical flow values in GWh			

(Country code: Appendix D)

Figure 3.54 | Cross-border physical electricity flows, in gigawatt-hours (GWh) (ENTSO-E 2019).



markets at 13-40 billion € per year for the EU as a whole, depending on assumptions on fuel and carbon prices, renewable energy costs and penetration. If the market integration were broadened to include the whole Mediterranean region, then the savings would be even larger possibly delivering additional savings of 30 billion € per year according to studies conducted by Desert industrial energy initiative (Dii) (Zickfeld and Wieland 2012).

An integrated energy market and a cooperative approach would reduce the cost of meeting the ambitious EU CO<sub>2</sub> reduction and renewable energy targets (Caldés et al. 2015; Szabó et al. 2015). Associating the southern shore of the Mediterranean would further reduce the cost of decarbonization of the EU power sector, as the region is endowed with a massive renewable energy potential and a vast stock of unused land where solar panels and other renewable plants can be sited without creating a nuisance for nearby population. The Sahara Desert is a prime location for solar power generation and could potentially produce several times the level of demand for carbonless electricity in Europe, while also covering demand in the SEMCs.

The two Mediterranean shores are already inter-connected, in the West by a submarine cable of 1,400 MW under the Strait of Gibraltar connecting Spain and Morocco, and in the East with Turkey connected to Bulgaria through two 400 kV lines (for a total capacity of 2,500 MW) and to Greece through a 400-kV line with a capacity of 500 MW. Other interconnections between Europe and the southern shore of the Mediterranean are being studied, in particular, the Italy-Tunisia connection to the centre of the Mediterranean (Fig. 3.54). Contrary to Europe, the network in the SEMCs is sparser, and interrupted between Tunisia and Libya. In this region, renewable energy integration often needs complementary developments in generation and transmission, due to more isolated systems requiring back-up capacity.

The idea of exporting solar electricity from the Sahara is not new and dates back to the 1940s (Escibano et al. 2019), but the concept gained momentum when the first EU renewable energy Directive (EU 2009) was being prepared. The period 2008-2012 saw a flurry of initiatives such as Mediterranean Solar Plan, Desertec, Dii and MEDGRID to connect the southern and northern shores of the Mediterranean when it seemed that many EU countries would face difficulties in meeting their commitments under the Directive. Most of these initiatives went into hibernation as EU countries did not express much interest in cooperation mechanisms with southern Mediterranean countries, either because

they could meet their 2020 targets from their own renewable energy resources (in part because of stagnant demand), or because they preferred to use other mechanisms such as statistical transfers or cooperation mechanisms with other EU countries. Caldés et al. (2018) and Lilliestam et al. (2016) identified the following reasons for the failures of these ambitious Mediterranean integration initiatives: (1) underdeveloped legal and regulatory frameworks, (2) weak grid infrastructures (in particular lack of interconnection between the two shores), (3) lower than expected socio-economic benefits, (4) high upfront costs and lack of financing mechanisms, (5) high fossil fuel subsidies in SEMCs and (6) energy policy giving priority to domestic renewable energy production over electricity imports and electricity surpluses in EU countries.

The EU has now set more stringent targets for 2030 and is aiming for carbon neutrality in 2050 under the European Green Deal proposed by the Commission in December 2019. Furthermore, the benefits of regional market integration in terms of investment and operating cost savings, and facilitated renewable energy grid integration, are increasingly acknowledged. Regional electricity market integration appears to be recognized as one of the most cost-effective options to increase power system flexibility.

### ***The need for supra-regional interconnections***

The integration of a high-level of renewable energy requires that power systems have flexibility to cope with the stress resulting from sudden and unpredictable variations in the availability of renewable energy. Power system flexibility is defined as the ability to reliably and cost-effectively manage the variability and uncertainty of demand and supply across all relevant timescales, from ensuring instantaneous stability of the power system to support the long-term security of supply (Taibi et al. 2018; Mohandes et al. 2019). Flexibility can be provided by dispatchable power plants, demand-side response, storage and network infrastructure, in particular, the one that supports regional market integration (Baritaud and Volk 2014) and requires a combination of regulatory, operational and investment measures (Hirth and Ziegenhagen 2015). Greater transmission interconnectivity yields substantial economic and environmental benefits and requires the strengthening of interconnectors, the hardware of regional electricity market integration (Crisan and Kuhn 2017). Ensuring that interconnectors are efficiently used and properly remunerated for their flexibility services both reduces the short-run cost of integrating renewables and increases the attractiveness of investing in additional interconnection (Newbery et

al. 2016; Newbery 2017). Regional market integration through interconnectors expands the region's access to carbonless sources of energy, such as hydro reserves in the North or more predictable plentiful solar power in North Africa and reduces renewable energy sources curtailment.

The degree of interconnectivity will affect the need for new power plants and their location. Power generation and transmission planning, therefore, needs to be integrated. Successful renewable energy sources integration requires integration of power generation and transmission planning, of operational and investment decisions and of national markets into regional markets (Pariante-David 2014). A holistic approach is needed. Reaching high renewable energy shares entails "integration costs" elsewhere in the power system such as balancing services and firm back-up capacity on standby that are not reflected in traditional planning and economics approach based on the LCOE. A system-wide approach is needed that integrates all the costs and derived effects of renewable energy sources integration to determine the optimal mix of power generation plants and transmission lines to meet electricity demand at the lowest cost while satisfying the climate change and other policy objectives. This is the "total system cost" approach which focuses on the total cost of the power system, rather than trying to allocate some of the cost components to specific technologies, or part of the power system, to compare the technologies on the basis of LCOE (Pariante-David 2016).

After a period of lull, there is a revival of interest in electricity exchanges across the Mediterranean. Designed by Med-TSO between 2015 and 2018 in

the framework of its Mediterranean project "the Mediterranean Master Plan plays a key role for consolidating a secure and sustainable electricity infrastructure through the development of interconnections, while facilitating the integration of Renewable Energy Sources in the Mediterranean Region" (Illiceto and Ferrante 2018). The Mediterranean Master Plan identified 14 clusters of projects for the interconnection of Mediterranean electricity grids according to a 2030 horizon scenario (Fig. 3.55). The Plan targets 18 GW of new interconnection capacity, corresponding to 2,200 km of new transmission lines, requiring about 16 G billion € of additional investments.

The circumstances seem to be auspicious in 2019 to relaunch the Mediterranean Energy Union process. The EU "Clean Energy for All Europeans" (CE4ALL) Package (European Commission 2016) was fully adopted in May 2019, as the implementing instrument of the 2015 Energy Union strategy, which acknowledges that regional market integration is key in achieving EU climate change objectives at least cost and that strongly interconnected networks are required to support unhampered electricity trade and sharing of ancillary services. Cooperation, coordination (both on national policies and power system operations) as well as and regional market integration are a central part of the CE4ALL Package. Although the Package primarily aims at the European internal energy market, it recognises that the EU is not isolated and includes several cooperation mechanisms and financing instruments for joint projects with third-party countries (Held et al. 2019), so that the EU can tap into the best resources to achieve power sector decarbonisation.




Figure 3.55 | Mediterranean interconnection projects (Illiceto and Ferrante 2018).

In summary, regional energy market integration and cooperation are crucial to unleashing the region's solar potential for cost-effective climate change mitigation in the Mediterranean. Many obstacles remain, including lack of the necessary transmission/interconnection infrastructure, difficulty in siting renewable energy plants in the best locations because of high up-front costs that makes financing risky, geopolitical considerations, insufficient coordination of power systems and lack of alignment of market rules. It is necessary to ensure interoperability of wholesale markets, value properly flexibility services and allocate interconnection costs to reflect the benefits

accrued to the different stakeholders. This will extract value from regional market integration. The process can take years, happens in stages and requires appropriate national and regional institutions (Oseni and Pollitt 2016; Pariente-David and Jannet Allal 2019).

**Strategy of transnational associations of regulators**

Regulations regarding energy transmission systems and operations vary significantly among Mediterranean countries (MEDREG 2018). Transnational partnerships thus require forming a common set of




Figure 3.56 | Strategy of Mediterranean regulators for regional integration (Illiceto and Ferrante 2018).

## BOX 3.3.1


## Climate variability and energy planning

The variability of the renewable energies production, such as from wind and PV energy, adds to that of the energy demand and may pose challenges to the adequation and stability of networks which can translate into a decreased quality of services or into an increase in system costs (Ueckerdt et al. 2015) due to additional flexibility requirements. The variability of renewable energies production becomes critical when their penetration reaches a level, which is higher than what existing flexibility mechanisms allow the energy system to cope with (Creti and Fontini 2019, chap. 26).

Not only the variability of meteorological conditions at single locations is relevant to energy planning, but also the relations between electricity demand and capacity factors, between capacity factors at different locations, and between capacity factors for different technologies. These correlations influence the potential smoothing of production once aggregated by interconnections, illustrated in Fig. 3.57 for the case of wind and PV energy in Italy analyzed by Tantet et al. (2019). In panel a, points in the curves represent optimal distributions of wind and PV capacities among the six Italian electric regions for a varying trade-off between maximizing the mean and minimizing the standard deviation of the wind and PV penetration. Due to the variability of the wind and PV production, changing the weight put on each objective results in mixes with different characteristics, a higher mean penetration also leading to an increased variance. In addition,


ignoring capacity-factor correlations between regions leads to sub-optimal mixes as shown by the weaker slope of the dashed lines (which corresponds to a weaker increase of the mean penetration as the variance is allowed to increase). The blue curve represents the optimal frontier with the addition of the constraint that the total wind-PV capacity be the same as in 2015. The blue dot (resp. blue diamond) on this frontier represents the optimal mix that has the same renewable energies penetration variance (resp. mean) as the actual 2015 mix (represented by the gray dot). The geographical wind-PV distribution of the capacities corresponding to the blue dot and the blue diamond are represented in panels b and c, respectively. Depending on whether more weight is put on the mean or on the variance of the renewable energy penetration, different optimal mixes are obtained.

Regarding locations for renewable energy development, the European case is studied by Pryor et al. (2006), specifically for wind energy, by Pfenninger and Staffell (2016) for photovoltaic energy and by Rodriguez et al. (2014) for both wind and PV energy with network constraints; the case of northern Spain by Marcos et al. (2012); and the case of Italy by Tantet et al. (2019). Concerning technologies, the complementarity of wind and solar energy in Europe is analyzed by Buttler et al. (2016) and Miglietta et al. (2017), in Italy by Monforti et al. (2014) and Tantet et al. (2019), and in Spain by Santos-Alamillos et al. (2012, 2017); and the case of run-of-the river hydropower and wind and PV energy with countries including France, Greece, Italy, Spain and Tunisia by François et al. (2016).



**Figure 3.57 | Panel A: Illustration of the optimal, or Pareto, frontiers for two objectives – maximizing the mean (y-axis, in % of the total demand) and minimizing the standard deviation (x axis) of the total wind-PV penetration** obtained by distributing wind and PV capacities among the six bidding zones (electric regions) of Italy. The gray area represents infeasible mixes. The thick line is the frontier when taking all capacity-factor correlations between regions and technologies into account in the bi-objective optimization, while the dashed line (resp. the point-dashed line) represents the frontier when only capacity-factor correlations between wind and PV energy in the same region (resp. when no correlations) are considered. The thick blue line is like the thick black line, but with the constraint that the total wind-PV capacity be the same as in 2015. The blue dot (resp. blue diamond) represents the optimal mix with the same renewable energy penetration variance (resp. mean) as in 2015. The gray dot represents the actual 2015 mix.

To our knowledge, there are no studies analyzing how past changes in climate variability has impacted energy systems and their planning, let alone in the Mediterranean. With the development of various renewable energy types, as well as with existing or new hydroelectric capacities, changes in climate variability in the Mediterranean potentially affect the variability of the energy production (Widén et al. 2015) on time scales ranging from seconds (Apt 2007) to years or more (Pryor et al. 2006; Pozo-Vazquez et al. 2011; Pfenninger and Staffell 2016; Collins et al. 2018). Only few studies analyze past changes in the variability of the wind, solar and hydroelectric resources. No clear trend is found in European wind variability over the last 140 years or so (Bett et al. 2013), but there is significant variability between decades, depending on location and specific conditions. For instance, only small changes in the occurrence of Mistral and Tramontane winds are found in regional climate simulations forced by a reanalysis over the 1950-2010 period (Obermann et al. 2018).



**Panel B: PV (blue) and wind (orange) optimal capacity distribution resulting in the same variable renewable energies penetration variance as the actual 2015 mix.**

**Panel C: same as panel b for the same variable renewable energy penetration mean as the actual 2015 mix.** (Tantet et al. 2019).

regulations so as to enable enhanced connectivity of energy markets. Setting such conditions for a future Mediterranean energy community is the aim of the Association of Mediterranean Energy Regulators (MEDREG). Gathering 27 energy regulators from 22 countries spanning the EU, the Balkans and the SEMCs, MEDREG targets the establishment of a level playing field for all Mediterranean energy actors through an adapted legal and regulatory framework.

The stability and reliability of the regulatory framework is key to provide clear rules for investors to develop their confidence and ensure technical standards compatibility, which is a prerequisite for interconnecting markets. As a result, regulators have a crucial mission in implementing a good investment climate ensuring that network developments are delivered in due time, in providing guidance to Transmission System Operators about how to use interconnections and regulatory compatibility, in articulating a sound regulatory framework and a clear strategy, in ensuring an effective coordination between regulated networks and private/competitive activities, in improving the investment planning capacity, with long term assessment of energy needs and financial charges, and in ensuring a high level of transparency and education. MEDREG recognizes the importance of developing inter-operable electricity systems at sub-regional level. This requires first working on assessing the usage and problems of current interconnections and at a second stage evaluate what added value new interconnections could bring.

In this framework, the development of renewable energy sources requires a specific focus on network regulation. At the national level, it implies the connection and integration of renewable energy sources. Cross-border regulations require convergence of national regulations to allow interconnections to work effectively. Investment regulation requires the design and develop infrastructure that will be needed for promoting international complementarities.

The framework of EU Projects of Common Interest is an example of reflection that aims to build a shared vision to ensure the security of supply and facilitate renewable energy development in a coordinated way. Fig. 3.56 highlights the three strategic corridors regulators aim at to ensure a full integration of the Mediterranean (Illiceto and Ferrante 2018).

## BOX 3.3.2

**Energy issues for Mediterranean islands**

Islands are physically isolated territories, a characteristic that sets specific threats, challenges and opportunities in the context of global change and energy transition. The European Union recognizes that “insular regions suffer from structural handicaps linked to their island status, the permanence of which impairs their economic and social development” (Treaty of Amsterdam 1999). Geographical and socioeconomic singularities of Mediterranean Islands put additional pressure on water and energy, leading to resource depletion and degraded environment, threatening sustainable development (Gold and Webber 2015). More than 11 million people live in Mediterranean islands (Sen Nag 2017). Except for Sicily, Sardinia and Cyprus, all Mediterranean islands are below the million permanent inhabitants, with notable cases, such as Majorca, which frequently double its population during high touristic season. Mediterranean islands suffer strong limitations due to the limited range of their accessible resources, the inability to achieve economies of scale, the strong seasonal population variation, higher infrastructure costs and particular climatic conditions (Erdinc et al. 2015).

Characteristic aspects of Mediterranean climate such as large seasonal temperature and irradiance variations, occurrence of strong winds, heavy precipitations and the impacts of a range of cyclone, interact with the islands, rendering unique climates, even at local scales (Homar et al. 2010). They also enhance Mediterranean islands vulnerability, especially in the context of global climate change. In addition to the Mediterranean specificities, the IPCC 4th and 5th assessment reports state with high confidence that, globally, coasts are undergoing adverse consequences from climate change, such as sea level rise, inundation, erosion, and ecosystem loss. The reports also state that coasts are highly vulnerable to extreme events such as cyclones, extreme waves, storm surges, altered rainfall and runoff patterns, and ocean acidification. Therefore, Mediterranean islands are essentially isolated coastal territories with double penalty from climate change impacts.

**Energy production and demand in the Mediterranean islands**

Climate effects on the energy transition process in Mediterranean islands are of capital importance. On the one hand, the water-energy nexus is a central aspect in islands, as pressure on water resources is exceptional and highly anticorrelated with the seasonal precipitations and the availability of fresh water depends on techniques such as dwelling, impoundment and desalination. This severe fresh water scarcity is projected to aggravate in the future, since a drying of the region is expected throughout the 21st century (Dubrovský et al. 2014) and dry spells are projected to increase in duration and increasingly affect the wet season (Raymond et al. 2019). This climate and the projected scenarios not only severely hamper the adoption of hydropower electric sources in Mediterranean islands, but also increases the projected electricity demand derived from increased use of desalination strategies. On the other hand, Mediterranean

islands have an important wind power climatic resource originating from various climatic features. Larger islands develop sea breeze very in a consistent way, especially during summer, in phase with the highest seasonal electricity demand. In contrast, smaller -and usually flatter- islands do not develop relevant sea breezes, but are more exposed to the weakly dragged maritime winds. The Mediterranean is well known to have the highest concentration of cyclones across the globe (Petterssen 1956). An additional climatic resource for Mediterranean islands, that has raised some attention in recent years, is the wave energy (Franzitta and Curto 2017) and maritime underwater flows (*Section 3.3.2.2*). Despite wave energy converters are becoming more efficient, the spatial and temporal variability of this maritime resource hampers its general implementation. Wave energy in the Mediterranean Sea is larger during the cyclogenetic cold season and over the area of influence of the most frequent cyclogenetic regions (Ponce de León et al. 2016). This resource is significantly weaker during the summer season, when energy demand steered by tourist activity is higher. This characteristic favors its use in energy transition planning as a complementary electricity generation technology to photovoltaic, which has an opposite seasonal phase (Curto and Trapanese 2018; Curto et al. 2019). Climate projections for wave energy in the western Mediterranean show that this resource will remain reliable with a reduced temporal variability and slight reduction of the annual and seasonal wave power (Sierra et al. 2017).

Regarding the demand side, islands are expected to follow the Mediterranean mainland projections of increases of 6% in demand by 2050 (Zachariadis and Hadjinicolaou 2014), even possibly amplified due to larger tourist activity in currently underexploited environments. The non-linear relation between consumed energy and total population could be explained by differences in existing economic activities, geographical site and cultural aspects (Neves et al. 2014).


**Interlocking challenges of energy security and climate resilience**

In addition to the effects of climate and climate change as an energy resource to consider in sustainable energy transition planning for the Mediterranean islands, climate extremes pose both energy extraction opportunities (i.e., for wave energy) but more importantly engineering and protection challenges. Strong winds and heavy precipitation are projected to decrease in frequency but become more intense (*Subchapter 2.2*), threatening renewable energy infrastructure both in centralized plants and distributed generation topologies.

These climatic conditions add upon other economic, sociocultural and differential characteristics of Mediterranean insular territories compared to mainland, a fact that is also reflected to their power system structures and the energy transition objectives (Erdinc et al. 2015). Challenges such as the geographical limitations, protection of natural and cultural values, or the technical limitations of small size grid structures with low inertia, are all barriers to overcome in order to seize the opportunities of the energy transition for ensuring a sustainable insular power system (Andaloro et al. 2012). Nowadays,

in most non-interconnected islands the electricity generation cost is extremely high due to the utilization of outdated autonomous power stations based on oil-fuel imports and diesel-electric generators, and the most frequent energy mix proposed considers solar, wind and sea wave renewable sources (Franzitta and Curto 2017; Curto et al. 2019). Interconnection with mainland is frequent although not for all cases profitable (Lobato et al. 2017). This solution may alleviate the reduced inertia challenge and externalize the dependency on fossil fuel, but does not contribute to achieve the near zero energy system proposed by some insular communities (Sanseverino et al. 2014).

For stand-alone power systems, the management of renewable energy surplus is an important concern. Hydrogen generation, commercial batteries and the deferrable load of desalinated water-production are proposed as two effective renewable energy buffering strategies for Mediterranean islands (Corsini et al. 2009; Kaldellis et al. 2012). In the absence of a single solve-all solution, hybrid solutions are hypothesized to lead to a remarkable reduction in power generation costs, although more efforts are needed for making battery/hydrogen systems technically and economically viable (Corsini et al. 2009; Beccali et al. 2018; Wang et al. 2020). Besides generation-side measures and energy storage for reserve provision, demand-side measures have also some specificities in Mediterranean insular areas which can foster their transition to a sustainable and autonomous energy system. Outdated distribution grids, impact of the lack of economy of scale on the reduced budgets for new investments and the low penetration of automation in domestic utilities prevent the widespread implementation of solutions for the automation of the end-user's electrical installations, in order to offer to the utility flexibility to be used for the improvement of the generation and distribution efficiency (Zizzo et al. 2017). In this regard, electric vehicles have ranges suitable for the great majority of Mediterranean islands sizes, also offering new grid-to-vehicle and vehicle-to-grid managing alternatives which can catalyze the solutions for the inherent reduced grid inertia problems (Groppi et al. 2019).



## References

- Abadie LM, Galarraga I, Rübhelke D 2013 An analysis of the causes of the mitigation bias in international climate finance. *Mitig. Adapt. Strateg. Glob. Chang.* 18, 943–955. doi: [10.1007/s11027-012-9401-7](https://doi.org/10.1007/s11027-012-9401-7)
- Al-Asaad HK 2009 Electricity Power Sector Reform in the GCC Region. *Electr. J.* 22, 58–64. doi: [10.1016/j.tej.2009.08.013](https://doi.org/10.1016/j.tej.2009.08.013)
- Alnaser WE, Alnaser NW 2011 The status of renewable energy in the GCC countries. *Renew. Sustain. Energy Rev.* 15, 3074–3098. doi: [10.1016/j.rser.2011.03.021](https://doi.org/10.1016/j.rser.2011.03.021)
- Andaloro APF, Salomone R, Andaloro L, Briguglio N, Sparacia S 2012 Alternative energy scenarios for small islands: A case study from Salina Island (Aeolian Islands, Southern Italy). *Renew. Energy* 47, 135–146. doi: [10.1016/j.renene.2012.04.021](https://doi.org/10.1016/j.renene.2012.04.021)
- Apt J 2007 The spectrum of power from wind turbines. *J. Power Sources* 169, 369–374. doi: [10.1016/j.jpowsour.2007.02.077](https://doi.org/10.1016/j.jpowsour.2007.02.077)
- Balog I, Ruti PM, Tobin I, Armenio V, Vautard R 2016 A numerical approach for planning offshore wind farms from regional to local scales over the Mediterranean. *Renew. Energy* 85, 395–405. doi: [10.1016/j.renene.2015.06.038](https://doi.org/10.1016/j.renene.2015.06.038)
- Baritaud M, Volk D 2014 Seamless Power Markets: Regional Integration of Electricity Markets in IEA Member Countries. Paris. <https://webstore.iea.org/seamless-power-markets>
- Bartók B, Wild M, Folini D, Lüthi D, Kotlarski S et al. 2017 Projected changes in surface solar radiation in CMIP5 global climate models and in EURO-CORDEX regional climate models for Europe. *Clim. Dyn.* 49, 2665–2683. doi: [10.1007/s00382-016-3471-2](https://doi.org/10.1007/s00382-016-3471-2)
- Bartos M, Chester M V., Johnson N, Gorman B, Eisenberg D et al. 2016 Impacts of rising air temperatures on electric transmission ampacity and peak electricity load in the United States. *Environ. Res. Lett.* 11, 114008. doi: [10.1088/1748-9326/11/11/114008](https://doi.org/10.1088/1748-9326/11/11/114008)
- Bauer N, Rose SK, Fujimori S, van Vuuren DP, Weyant J et al. 2018 Global energy sector emission reductions and bioenergy use: overview of the bioenergy demand phase of the EMF-33 model comparison. *Clim. Change*, 1–16. doi: [10.1007/s10584-018-2226-y](https://doi.org/10.1007/s10584-018-2226-y)
- Beccali M, Finocchiaro P, Ippolito MG, Leone G, Panno D et al. 2018 Analysis of some renewable energy uses and demand side measures for hotels on small Mediterranean islands: A case study. *Energy* 157, 106–114. doi: [10.1016/j.energy.2018.05.139](https://doi.org/10.1016/j.energy.2018.05.139)
- Beringer T, Lucht W, Schaphoff S 2011 Bioenergy production potential of global biomass plantations under environmental and agricultural constraints. *GCB Bioenergy* 3, 299–312. doi: [10.1111/j.1757-1707.2010.01088.x](https://doi.org/10.1111/j.1757-1707.2010.01088.x)
- Bertani R 2017 *Perspectives for Geothermal Energy in Europe*. World Scientific Publishing doi: 10.1142/q0069
- Bett PE, Thornton HE, Clark RT 2013 European wind variability over 140 yr. *Adv. Sci. Res.* 10, 51–58. doi: [10.5194/asr-10-51-2013](https://doi.org/10.5194/asr-10-51-2013)
- Bichet A, Hingray B, Evin G, Diedhiou A, Kebe CMF et al. 2019 Potential impact of climate change on solar resource in Africa for photovoltaic energy: analyses from CORDEX-AFRICA climate experiments. *Environ. Res. Lett.* 14, 124039. doi: [10.1088/1748-9326/ab500a](https://doi.org/10.1088/1748-9326/ab500a)
- Bilgen S, Keleş S, Sarikaya I, Kaygusuz K 2015 A perspective for potential and technology of bioenergy in Turkey: Present case and future view. *Renew. Sustain. Energy Rev.* 48, 228–239. doi: [10.1016/j.rser.2015.03.096](https://doi.org/10.1016/j.rser.2015.03.096)
- Blimpo MP, Cosgrove-Davies M 2019 Electricity Access in Sub-Saharan Africa: Uptake, Reliability, and Complementary Factors for Economic Impact. World Bank Group. doi: [10.1596/978-1-4648-1361-0](https://doi.org/10.1596/978-1-4648-1361-0)
- Bockers V, Haucap J, Heimeshoff U 2013 Cost of Non-Europe in the Single Market for Energy: ANEX IV Benefits of an Integrated European Electricity Market. Brussels [http://publications.europa.eu/resource/cellar/99d4fd94-7619-44f4-9f4b-5541235b90d1.0001.04/DOC\\_1](http://publications.europa.eu/resource/cellar/99d4fd94-7619-44f4-9f4b-5541235b90d1.0001.04/DOC_1)
- Brand B 2013 Transmission topologies for the integration of renewable power into the electricity systems of North Africa. *Energy Policy* 60, 155–166. doi: [10.1016/j.enpol.2013.04.071](https://doi.org/10.1016/j.enpol.2013.04.071)
- Brand B, Blok K 2015 Renewable energy perspectives for the North African electricity systems: A comparative analysis of model-based scenario studies. *Energy Strateg. Rev.* 6, 1–11. doi: [10.1016/j.esr.2014.11.002](https://doi.org/10.1016/j.esr.2014.11.002)
- Brand B, Zingerle J 2011 The renewable energy targets of the Maghreb countries: Impact on electricity supply and conventional power markets. *Energy Policy* 39, 4411–4419. doi: [10.1016/j.enpol.2010.10.010](https://doi.org/10.1016/j.enpol.2010.10.010)
- Bryden J, Riahi L, Zissler R 2013 MENA Renewables Status Report. Paris, France
- Buoninconti L, Filagrossi Ambrosino C 2015 Water saving assessment in residential buildings. *Sustain. Mediterr. Constr.* [http://www.sustainablemediterraneanconstruction.eu/SMC/The\\_Magazine\\_n.2\\_files/Smc\\_N.2\\_pap\\_10.pdf](http://www.sustainablemediterraneanconstruction.eu/SMC/The_Magazine_n.2_files/Smc_N.2_pap_10.pdf) [Accessed June 30, 2019]
- Butler D 2008 Architects of a low-energy future. *Nature* 452, 520–523. doi: [10.1038/452520a](https://doi.org/10.1038/452520a)
- Buttler A, Dinkel F, Franz S, Spliethoff H 2016 Variability of wind and solar power – An assessment of the current situation in the European Union based on the year 2014. *Energy* 106, 147–161. doi: [10.1016/j.energy.2016.03.041](https://doi.org/10.1016/j.energy.2016.03.041)
- Caldés N, de la Rúa C, Lechón Y, Rodríguez I, Trieb F et al. 2015 Bringing Europe and Third countries closer together through renewable energies (BETTER): Summary Report. [http://better-project.net/sites/default/files/BETTER\\_Summary\\_Report\\_0.pdf](http://better-project.net/sites/default/files/BETTER_Summary_Report_0.pdf) [Accessed June 29, 2019]



- Caldés N, del Río P, Lechón Y, Gerbeti A 2018 Renewable Energy Cooperation in Europe: What Next? Drivers and Barriers to the Use of Cooperation Mechanisms. *Energies* 12, 70. doi: [10.3390/en12010070](https://doi.org/10.3390/en12010070)
- Calero Quesada MC, García Lafuente J, Sánchez Garrido JC, Sammartino S, Delgado J 2014 Energy of marine currents in the Strait of Gibraltar and its potential as a renewable energy resource. *Renew. Sustain. Energy Rev.* 34, 98–109. doi: [10.1016/j.rser.2014.02.038](https://doi.org/10.1016/j.rser.2014.02.038)
- Cappelli G, Yamaç SS, Stella T, Francone C, Paleari L et al. 2015 Are advantages from the partial replacement of corn with second-generation energy crops undermined by climate change? A case study for giant reed in northern Italy. *Biomass and Bioenergy* 80, 85–93. doi: [10.1016/j.biombioe.2015.04.038](https://doi.org/10.1016/j.biombioe.2015.04.038)
- Caputo AC, Palumbo M, Pelagagge PM, Scacchia F 2005 Economics of biomass energy utilization in combustion and gasification plants: effects of logistic variables. *Biomass and Bioenergy* 28, 35–51. doi: [10.1016/j.biombioe.2004.04.009](https://doi.org/10.1016/j.biombioe.2004.04.009)
- Cavicchi B, Palmieri S, Odaldi M 2017 The Influence of Local Governance: Effects on the Sustainability of Bioenergy Innovation. *Sustainability* 9, 406. doi: [10.3390/su9030406](https://doi.org/10.3390/su9030406)
- CBI 2019 Green Bonds: The State of the Market 2018. Climate Bonds Initiative. [https://www.climatebonds.net/files/reports/cbi\\_gbm\\_final\\_032019\\_web.pdf](https://www.climatebonds.net/files/reports/cbi_gbm_final_032019_web.pdf)
- Chamorro CR, García-Cuesta JL, Mondéjar ME, Pérez-Madrado A 2014 Enhanced geothermal systems in Europe: An estimation and comparison of the technical and sustainable potentials. *Energy* 65, 250–263. doi: [10.1016/j.energy.2013.11.078](https://doi.org/10.1016/j.energy.2013.11.078)
- Chow TT 2010 A review on photovoltaic/thermal hybrid solar technology. *Appl. Energy* 87, 365–379. doi: [10.1016/j.apenergy.2009.06.037](https://doi.org/10.1016/j.apenergy.2009.06.037)
- Chum H, Faaij A, Moreira J, Berndes G, Dhamija P et al. 2011 Bioenergy, in *Renewable Energy Sources and Climate Change Mitigation - Special Report of the Intergovernmental Panel on Climate Change*, eds. Edenhofer O, Pichs-Madruga R, Sokona Y, Seyboth K, Matschoss P et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 209–332. doi: [10.1017/cbo9781139151153.006](https://doi.org/10.1017/cbo9781139151153.006)
- Clarke L, Jiang K, Akimoto K, Babiker M, Blanford G et al. 2014 Assessing Transformation Pathways, in *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Edenhofer O, Pichs-Madruga R, Sokona Y, Farahani E, Kadner S et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press).
- Coiro DP, Troise G, Ciuffardi T, Sannino G 2013 Tidal current energy resource assessment: The Strait of Messina test case. in *International Conference on Clean Electrical Power (ICCEP), Alghero, Italy (IEEE)*. doi: [10.1109/iccep.2013.6586992](https://doi.org/10.1109/iccep.2013.6586992)
- Collins S, Deane P, Ó Gallachóir B, Pfenninger S, Staffell I 2018 Impacts of Inter-annual Wind and Solar Variations on the European Power System. *Joule* 2, 2076–2090. doi: [10.1016/j.joule.2018.06.020](https://doi.org/10.1016/j.joule.2018.06.020)
- Corsini A, Rispoli F, Gamberale M, Tortora E 2009 Assessment of H<sub>2</sub>- and H<sub>2</sub>O-based renewable energy-buffering systems in minor islands. *Renew. Energy* 34, 279–288. doi: [10.1016/j.renene.2008.03.005](https://doi.org/10.1016/j.renene.2008.03.005)
- Creti A, Fontini F 2019 *Economics of Electricity: Markets, Competition and Rules*. Cambridge University Press doi: [10.1017/9781316884614](https://doi.org/10.1017/9781316884614)
- Crisan A, Kuhn M 2017 The Energy Network: Infrastructure as the Hardware of the Energy Union, in *Energy Union: Europe's New Liberal Mercantilism? International Political Economy Series.*, eds. Andersen SS, Goldthau A, Sitter N (London: Palgrave Macmillan UK), 165–182. doi: [10.1057/978-1-137-59104-3\\_10](https://doi.org/10.1057/978-1-137-59104-3_10)
- Crook JA, Jones LA, Forster PM, Crook R 2011 Climate change impacts on future photovoltaic and concentrated solar power energy output. *Energy Environ. Sci.* 4, 3101–3109. doi: [10.1039/c1ee01495a](https://doi.org/10.1039/c1ee01495a)
- Curto D, Franzitta V, Viola A, Cirrincione M, Mohammadi A et al. 2019 A renewable energy mix to supply small islands. A comparative study applied to Balearic Islands and Fiji. *J. Clean. Prod.* 241, 118356. doi: [10.1016/j.jclepro.2019.118356](https://doi.org/10.1016/j.jclepro.2019.118356)
- Curto D, Trapanese M 2018 A Renewable Energy mix to Supply the Balearic Islands: Sea Wave, Wind and Solar. in *2018 IEEE International Conference on Environment and Electrical Engineering and 2018 IEEE Industrial and Commercial Power Systems Europe (EEEIC / I&CPS Europe)* (IEEE). doi: [10.1109/eeeic.2018.8493876](https://doi.org/10.1109/eeeic.2018.8493876)
- Daioglou V, Doelman JC, Wicke B, Faaij A, van Vuuren DP 2019 Integrated assessment of biomass supply and demand in climate change mitigation scenarios. *Glob. Environ. Chang.* 54, 88–101. doi: [10.1016/j.gloenvcha.2018.11.012](https://doi.org/10.1016/j.gloenvcha.2018.11.012)
- Damm A, Köberl J, Prettenhaler F, Rogler N, Töglhofer C 2017 Impacts of +2 °C global warming on electricity demand in Europe. *Clim. Serv.* 7, 12–30. doi: [10.1016/j.cliser.2016.07.001](https://doi.org/10.1016/j.cliser.2016.07.001)
- de Souza LEV, Bosco EMGRL, Cavalcante AG, Da Costa Ferreira L 2018 Postcolonial theories meet energy studies: “Institutional orientalism” as a barrier for renewable electricity trade in the Mediterranean region. *Energy Res. Soc. Sci.* 40, 91–100. doi: [10.1016/j.erss.2017.12.001](https://doi.org/10.1016/j.erss.2017.12.001)
- Dolarin M, Vidrih B, Kajfež-Bogataj L, Medved S 2010 Predicted changes in energy demands for heating and cooling due to climate change. *Phys. Chem. Earth, Parts A/B/C* 35, 100–106. doi: [10.1016/j.pce.2010.03.003](https://doi.org/10.1016/j.pce.2010.03.003)
- Don A, Osborne B, Hastings A, Skiba U, Carter MS et al. 2011 Land-use change to bioenergy production in



- François B, Borga M, Creutin J-D, Hingray B, Raynaud D et al. 2016 Complementarity between solar and hydro power: Sensitivity study to climate characteristics in Northern-Italy. *Renew. Energy* 86, 543–553. doi: [10.1016/j.renene.2015.08.044](https://doi.org/10.1016/j.renene.2015.08.044)
- Franzitta V, Curto D 2017 Sustainability of the Renewable Energy Extraction Close to the Mediterranean Islands. *Energies* 10, 283. doi: [10.3390/en10030283](https://doi.org/10.3390/en10030283)
- Fritsche UR, Berndes G, Cowie AL, Dale VH, Kline KL et al. 2017 Energy and Land-Use. <https://knowledge.unccd.int/sites/default/files/2018-06/2.Fritsche%2Bet%2Bal%2B%282017%29%2BEnergy%2Band%2BLand%2BUse%2B-%2BGL0%2Bpaper-corr.pdf>
- Fritzsche K, Zejli D, Tänzler D 2011 The relevance of global energy governance for Arab countries: The case of Morocco. *Energy Policy* 39, 4497–4506. doi: [10.1016/j.enpol.2010.11.042](https://doi.org/10.1016/j.enpol.2010.11.042)
- Gaudiosi G, Borri C 2010 Offshore wind energy in the mediterranean countries. *Rev. des Energies Renouvelables SMEE'10 Bou Ismail Tipaza*, 173 – 188. [https://www.cder.dz/download/smee2010\\_19.pdf](https://www.cder.dz/download/smee2010_19.pdf)
- Gelfand I, Sahajpal R, Zhang X, Izaurrealde RC, Gross KL et al. 2013 Sustainable bioenergy production from marginal lands in the US Midwest. *Nature* 493, 514–517. doi: [10.1038/nature11811](https://doi.org/10.1038/nature11811)
- Ghedamsi R, Settou N, Gouareh A, Khamouli A, Saifi N et al. 2016 Modeling and forecasting energy consumption for residential buildings in Algeria using bottom-up approach. *Energy Build.* 121, 309–317. doi: [10.1016/j.enbuild.2015.12.030](https://doi.org/10.1016/j.enbuild.2015.12.030)
- Giannakopoulos C, Hadjinicolaou P, Zerefos CS, Demosthenous G 2009 Changing energy requirements in the Mediterranean under changing climatic conditions. *Energies* 2, 805–815. doi: [10.3390/en20400805](https://doi.org/10.3390/en20400805)
- Gibbs HK, Johnston M, Foley JA, Holloway T, Monfreda C et al. 2008 Carbon payback times for crop-based biofuel expansion in the tropics: the effects of changing yield and technology. *Environ. Res. Lett.* 3, 34001. doi: [10.1088/1748-9326/3/3/034001](https://doi.org/10.1088/1748-9326/3/3/034001)
- Gil V, Gaertner MA, Gutiérrez C, Losada T 2019 Impact of climate change on solar irradiation and variability over the Iberian Peninsula using regional climate models. *Int. J. Climatol.* 39, 1733–1747. doi: [10.1002/joc.5916](https://doi.org/10.1002/joc.5916)
- Gold G, Webber M 2015 The Energy-Water Nexus: An Analysis and Comparison of Various Configurations Integrating Desalination with Renewable Power. *Resources* 4, 227–276. doi: [10.3390/resources4020227](https://doi.org/10.3390/resources4020227)
- Gómez A, Rodrigues M, Montañés C, Dopazo C, Fueyo N 2010 The potential for electricity generation from crop and forestry residues in Spain. *Biomass and Bioenergy* 34, 703–719. doi: [10.1016/j.biombioe.2010.01.013](https://doi.org/10.1016/j.biombioe.2010.01.013)
- González A, Riba JR, Puig R, Navarro P 2015 Review of micro- and small-scale technologies to produce electricity and heat from Mediterranean forests' wood chips. *Renew. Sustain. Energy Rev.* 43, 143–155. doi: [10.1016/j.rser.2014.11.013](https://doi.org/10.1016/j.rser.2014.11.013)
- Groppi D, Astiaso Garcia D, Lo Basso G, de Santoli L 2019 Synergy between smart energy systems simulation tools for greening small Mediterranean islands. *Renew. Energy* 135, 515–524. doi: [10.1016/j.renene.2018.12.043](https://doi.org/10.1016/j.renene.2018.12.043)
- Gutiérrez C, Somot S, Nabat P, Mallet M, Gaertner M et al. 2018 Impact of aerosols on the spatiotemporal variability of photovoltaic energy production in the Euro-Mediterranean area. *Sol. Energy* 174, 1142–1152. doi: [10.1016/j.solener.2018.09.085](https://doi.org/10.1016/j.solener.2018.09.085)
- Hadjipanayi M, Koumparou I, Philippou N, Paraskeva V, Phinikarides A et al. 2016 Prospects of photovoltaics in southern European, Mediterranean and Middle East regions. *Renew. Energy* 92, 58–74. doi: [10.1016/j.renene.2016.01.096](https://doi.org/10.1016/j.renene.2016.01.096)
- Haller M, Ludig S, Bauer N 2012 Decarbonization scenarios for the EU and MENA power system: Considering spatial distribution and short term dynamics of renewable generation. *Energy Policy* 47, 282–290. doi: [10.1016/j.enpol.2012.04.069](https://doi.org/10.1016/j.enpol.2012.04.069)
- Hamududu B, Killingtveit A 2012 Assessing climate change impacts on global hydropower. *Energies* 5, 305–322. doi: [10.3390/en5020305](https://doi.org/10.3390/en5020305)
- Hastings A, Clifton-Brown J, Wattenbach M, Mitchell CP, Stampfl P et al. 2009 Future energy potential of *Miscanthus* in Europe. *GCB Bioenergy* 1, 180–196. doi: [10.1111/j.1757-1707.2009.01012.x](https://doi.org/10.1111/j.1757-1707.2009.01012.x)
- Hawila D, Mezher T, Kennedy SW, Mondal A 2012 Renewable energy readiness assessment for North African countries. *Proc. PICMET '12 Technol. Manag. Emerg. Technol.*, 2970–2982. doi: [10.1016/j.rser.2014.01.066](https://doi.org/10.1016/j.rser.2014.01.066)
- Held A, Ragwitz M, Winkler J 2019 “Clean energy for all Europeans” package: Implications and opportunities for the Mediterranean. [https://www.cmimarseille.org/sites/default/files/newsite/english\\_version\\_online.pdf](https://www.cmimarseille.org/sites/default/files/newsite/english_version_online.pdf)
- Hirth L, Ziegenhagen I 2015 Balancing power and variable renewables: Three links. *Renew. Sustain. Energy Rev.* 50, 1035–1051. doi: [10.1016/j.rser.2015.04.180](https://doi.org/10.1016/j.rser.2015.04.180)
- Homar V, Ramis C, Romero R, Alonso S 2010 Recent trends in temperature and precipitation over the Balearic Islands (Spain). *Clim. Change* 98, 199. doi: [10.1007/s10584-009-9664-5](https://doi.org/10.1007/s10584-009-9664-5)
- Hueging H, Haas R, Born K, Jacob D, Pinto JG 2013 Regional changes in wind energy potential over Europe using regional climate model ensemble projections. *J. Appl. Meteorol. Climatol.* 52, 903–917. doi: [10.1175/JAMC-D-12-086.1](https://doi.org/10.1175/JAMC-D-12-086.1)
- Hui SCM 2000 Building energy efficiency standards in Hong Kong and mainland China. in *2000 ACEEE Summer Study on Energy Efficiency in Buildings, 20-25 August 2000, Pacific Grove, California*.

- IEA 2018 Data and Statistics. *Int. Energy Agency*. [https://www.iea.org/data-and-statistics?country=WORLD&fuel=Energy supply&indicator=Total primary energy supply](https://www.iea.org/data-and-statistics?country=WORLD&fuel=Energy%20supply&indicator=Total%20primary%20energy%20supply) (TPES) by source [Accessed June 30, 2019].
- IEA Bioenergy 2016 IEA Bioenergy Countries' Report: Bioenergy policies and status of implementation.
- Iliceto A, Ferrante A 2018 Consolidating a Secure and Sustainable Electricity Infrastructure in the Mediterranean Region : The Mediterranean Project of Med-TSO. in *2018 AEIT International Annual Conference* (IEEE). doi: [10.23919/AEIT.2018.8577419](https://doi.org/10.23919/AEIT.2018.8577419)
- IPCC 2014 *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* , eds. Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press. doi: [10.1017/CBO9781107415386](https://doi.org/10.1017/CBO9781107415386)
- IPCC 2018 *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change.* , eds. Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J et al. In press
- IRENA 2018 Renewable power generation costs in 2014. [https://www.irena.org/documentdownloads/publications/irena\\_re\\_power\\_costs\\_2014\\_report.pdf](https://www.irena.org/documentdownloads/publications/irena_re_power_costs_2014_report.pdf)
- Iwano J, Mwasha A 2010 A review of building energy regulation and policy for energy conservation in developing countries. *Energy Policy* 38, 7744–7755. doi: [10.1016/j.enpol.2010.08.027](https://doi.org/10.1016/j.enpol.2010.08.027)
- Jaber S, Ajib S 2011 Optimum, technical and energy efficiency design of residential building in Mediterranean region. *Energy Build.* 43, 1829–1834. doi: [10.1016/j.enbuild.2011.03.024](https://doi.org/10.1016/j.enbuild.2011.03.024)
- Jerez S, Tobin I, Vautard R, Montávez JP, López-Romero JM et al. 2015 The impact of climate change on photovoltaic power generation in Europe. *Nat. Commun.* 6, 10014. doi: [10.1038/ncomms10014](https://doi.org/10.1038/ncomms10014)
- Kaldellis JK, Gkikaki A, Kaldelli E, Kapsali M 2012 Investigating the energy autonomy of very small non-interconnected islands. *Energy Sustain. Dev.* 16, 476–485. doi: [10.1016/j.esd.2012.08.002](https://doi.org/10.1016/j.esd.2012.08.002)
- Karagali I, Mann J, Dellwik E, Vasiljević N 2018 New European Wind Atlas: The Østerild balconies experiment. *J. Phys. Conf. Ser.* 1037, 52029. doi: [10.1088/1742-6596/1037/5/052029](https://doi.org/10.1088/1742-6596/1037/5/052029)
- Koch H, Vögele S 2009 Dynamic modelling of water demand, water availability and adaptation strategies for power plants to global change. *Ecol. Econ.* 68, 2031–2039. doi: [10.1016/j.ecolecon.2009.02.015](https://doi.org/10.1016/j.ecolecon.2009.02.015)
- Komendantova N, Patt A, Barras L, Battaglini A 2012 Perception of risks in renewable energy projects: The case of concentrated solar power in North Africa. *Energy Policy* 40, 103–109. doi: [10.1016/j.enpol.2009.12.008](https://doi.org/10.1016/j.enpol.2009.12.008)
- Komendantova N, Patt A, Williges K 2011 Solar power investment in North Africa: Reducing perceived risks. *Renew. Sustain. Energy Rev.* 15, 4829–4835. doi: [10.1016/j.rser.2011.07.068](https://doi.org/10.1016/j.rser.2011.07.068)
- Krupa J, Poudineh R 2017 Financing renewable electricity in the resource-rich countries of the Middle East and North Africa: a review. Oxford Institute for Energy Studies. doi: [10.26889/9781784670788](https://doi.org/10.26889/9781784670788)
- Laurent A, Pelzer E, Loyce C, Makowski D 2015 Ranking yields of energy crops: A meta-analysis using direct and indirect comparisons. *Renew. Sustain. Energy Rev.* 46, 41–50. doi: [10.1016/j.rser.2015.02.023](https://doi.org/10.1016/j.rser.2015.02.023)
- Lewis A, Estefen S, Huckerby J, Musial W, Pontes T et al. 2011 Ocean Energy, in *IPCC Special Report on Renewable Energy Sources and Climate Change Mitigation*, eds. Edenhofer O, Pichs-Madruga R, Sokona Y, Seyboth K, Matschoss P et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press).
- Liang X, Lettenmaier DP, Wood EF, Burges SJ 1994 A simple hydrologically based model of land surface water and energy fluxes for general circulation models. *J. Geophys. Res.* 99, 14415–14428. doi: [10.1029/94jd00483](https://doi.org/10.1029/94jd00483)
- Liberti L, Carillo A, Sannino G 2013 Wave energy resource assessment in the Mediterranean, the Italian perspective. *Renew. Energy* 50, 938–949. doi: [10.1016/j.renene.2012.08.023](https://doi.org/10.1016/j.renene.2012.08.023)
- Lilliestam J, Ellenbeck S 2011 Energy security and renewable electricity trade—Will Desertec make Europe vulnerable to the “energy weapon”? *Energy Policy* 39, 3380–3391. doi: [10.1016/j.enpol.2011.03.035](https://doi.org/10.1016/j.enpol.2011.03.035)
- Lilliestam J, Ellenbeck S, Karakosta C, Caldés N 2016 Understanding the absence of renewable electricity imports to the European Union. *Int. J. Energy Sect. Manag.* 10, 291–311. doi: [10.1108/IJESM-10-2014-0002](https://doi.org/10.1108/IJESM-10-2014-0002)
- Lilliestam J, Patt A 2015 Barriers, risks and policies for renewables in the Gulf States. *Energies* 8, 8263–8285. doi: [10.3390/en8088263](https://doi.org/10.3390/en8088263)
- Lobato E, Sigrist L, Rouco L 2017 Value of electric interconnection links in remote island power systems: The Spanish Canary and Balearic archipelago cases. *Int. J. Electr. Power Energy Syst.* 91, 192–200. doi: [10.1016/j.ijepes.2017.03.014](https://doi.org/10.1016/j.ijepes.2017.03.014)
- Macknick J, Newmark R, Heath G, Hallett KC 2012 Operational water consumption and withdrawal factors for electricity generating technologies: a review of existing literature. *Environ. Res. Lett.* 7, 45802. doi: [10.1088/1748-9326/7/4/045802](https://doi.org/10.1088/1748-9326/7/4/045802)
- Marcos J, Marroyo L, Lorenzo E, García M 2012 Smoothing of PV power fluctuations by geographical dispersion. *Prog. Photovoltaics Res. Appl.* 20, 226–237. doi: [10.1002/pip.1127](https://doi.org/10.1002/pip.1127)

- Marktanner M, Salman L 2011 Economic and geopolitical dimensions of renewable vs. nuclear energy in North Africa. *Energy Policy* 39, 4479–4489. doi: [10.1016/j.enpol.2010.12.047](https://doi.org/10.1016/j.enpol.2010.12.047)
- McVicar TR, Roderick ML, Donohue RJ, Li LT, Van Niel TG et al. 2012 Global review and synthesis of trends in observed terrestrial near-surface wind speeds: Implications for evaporation. *J. Hydrol.* 416–417, 182–205. doi: [10.1016/j.jhydrol.2011.10.024](https://doi.org/10.1016/j.jhydrol.2011.10.024)
- MEDENER 2014 Trends in energy efficiency in countries of the Mediterranean rim.
- MEDREG 2018 Mediterranean Energy Regulatory Outlook 2017.
- MEF 2019 Financing energy transition in the Mediterranean: Issues, challenges and key responses.
- Miglietta MM, Huld T, Monforti-Ferrario F 2017 Local Complementarity of Wind and Solar Energy Resources over Europe: An Assessment Study from a Meteorological Perspective. *J. Appl. Meteorol. Climatol.* 56, 217–234. doi: [10.1175/jamc-d-16-0031.1](https://doi.org/10.1175/jamc-d-16-0031.1)
- Mohandes B, Moursi MS El, Hatzigiorgiou N, Khatib S El 2019 A Review of Power System Flexibility With High Penetration of Renewables. *IEEE Trans. Power Syst.* 34, 3140–3155. doi: [10.1109/TPWRS.2019.2897727](https://doi.org/10.1109/TPWRS.2019.2897727)
- Monforti F, Huld T, Bódis K, Vitali L, D'Isidoro M et al. 2014 Assessing complementarity of wind and solar resources for energy production in Italy. A Monte Carlo approach. *Renew. Energy* 63, 576–586. doi: [10.1016/j.renene.2013.10.028](https://doi.org/10.1016/j.renene.2013.10.028)
- Mouratiadou I, Bevione M, Bijl DL, Drouet L, Hejazi M et al. 2018 Water demand for electricity in deep decarbonisation scenarios: a multi-model assessment. *Clim. Change* 147, 91–106. doi: [10.1007/s10584-017-2117-7](https://doi.org/10.1007/s10584-017-2117-7)
- Mulinda C, Hu Q, Pan K 2013 Dissemination and Problems of African Biogas Technology. *Energy Power Eng.* 05, 506–512. doi: [10.4236/epe.2013.58055](https://doi.org/10.4236/epe.2013.58055)
- Müller R, Pfeifroth U, Träger-Chatterjee C, Cremer R, Trentmann J et al. 2015 Surface Solar Radiation Data Set - Heliosat (SARAH) - Edition 1, Satellite Application Facility on Climate Monitoring. doi: [10.5676/EUM\\_SAF\\_CM/SARAH/V001](https://doi.org/10.5676/EUM_SAF_CM/SARAH/V001)
- Muratori M, Calvin K V., Wise M, Kyle P, Edmonds JE 2016 Global economic consequences of deploying bioenergy with carbon capture and storage (BECCS). *Environ. Res. Lett.* 11, 95004. doi: [10.1088/1748-9326/11/9/095004](https://doi.org/10.1088/1748-9326/11/9/095004)
- Nabat P, Somot S, Mallet M, Sevault F, Chiacchio M et al. 2015 Direct and semi-direct aerosol radiative effect on the Mediterranean climate variability using a coupled regional climate system model. *Clim. Dyn.* 44, 1127–1155. doi: [10.1007/s00382-014-2205-6](https://doi.org/10.1007/s00382-014-2205-6)
- Neumuller M 2015 Retour à l'équilibre du marché des bois provençaux. *Econostrum*.
- Neves D, Silva CA, Connors S 2014 Design and implementation of hybrid renewable energy systems on micro-communities: A review on case studies. *Renew. Sustain. Energy Rev.* 31, 935–946. doi: [10.1016/j.rser.2013.12.047](https://doi.org/10.1016/j.rser.2013.12.047)
- Newbery D 2017 Tales of two islands – Lessons for EU energy policy from electricity market reforms in Britain and Ireland. *Energy Policy* 105, 597–607. doi: [10.1016/j.enpol.2016.10.015](https://doi.org/10.1016/j.enpol.2016.10.015)
- Newbery D, Strbac G, Viehoff I 2016 The benefits of integrating European electricity markets. *Energy Policy* 94, 253–263. doi: [10.1016/j.enpol.2016.03.047](https://doi.org/10.1016/j.enpol.2016.03.047)
- Obermann A, Bastin S, Belamari S, Conte D, Gaertner MÁ et al. 2018 Mistral and Tramontane wind speed and wind direction patterns in regional climate simulations. *Clim. Dyn.* 51, 1059–1076. doi: [10.1007/s00382-016-3053-3](https://doi.org/10.1007/s00382-016-3053-3)
- OECD 2018 Climate-related Development Finance Data. *OECD, Dev. Assist. Comm.* <https://www.oecd.org/dac/financing-sustainable-development/development-finance-topics/Climate-related-development-finance-in-2018.pdf>
- OFME 2015 Données et chiffres clés de la forêt Méditerranéenne. Marseille [https://www.ofme.org/documents/Chiffres-cles/Chiffres-cles-2015\\_pl\\_web.pdf](https://www.ofme.org/documents/Chiffres-cles/Chiffres-cles-2015_pl_web.pdf)
- Olsson L, Barbosa H, Bhadwal S, Cowie AL, Delusca K et al. 2019 Land Degradation, in *Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems*, eds. Shukla PR, Skea J, Buendia EC, Masson-Delmotte V, Pörtner H-O et al. (In press).
- OME/MEDENER 2016 Executive Summary. Mediterranean Energy Transition: 2040 Scenario. Paris <https://www.medener.org/wp-content/uploads/2015/05/Transition-énergétique-en-Méditerranée-scénario-2040-en.pdf> [Accessed June 30, 2019].
- OME/MEDENER 2018 Les Energies Renouvelables en Méditerranée : Tendances, Perspectives et Bonnes Pratiques. Paris.
- OME 2018 Mediterranean Energy Perspectives 2018. Paris.
- Omrani H, Drobinski P, Arsouze T, Bastin S, Lebeaupin-Brossier C et al. 2017 Spatial and temporal variability of wind energy resource and production over the North Western Mediterranean Sea: Sensitivity to air-sea interactions. *Renew. Energy* 101, 680–689. doi: [10.1016/j.renene.2016.09.028](https://doi.org/10.1016/j.renene.2016.09.028)
- Onea F, Deleanu L, Rusu L, Georgescu C 2016 Evaluation of the wind energy potential along the Mediterranean Sea coasts. *Energy Explor. Exploit.* 34, 766–792. doi: [10.1177/0144598716659592](https://doi.org/10.1177/0144598716659592)
- Oseni MO, Pollitt MG 2016 The promotion of regional integration of electricity markets: Lessons for developing countries. *Energy Policy* 88, 628–638. doi: [10.1016/j.enpol.2015.09.007](https://doi.org/10.1016/j.enpol.2015.09.007)
- Paredes-Sánchez JP, López-Ochoa LM, López-González LM, Las-Heras-Casas J, Xiberta-Bernat J 2019

- Evolution and perspectives of the bioenergy applications in Spain. *J. Clean. Prod.* 213, 553–568. doi: [10.1016/j.jclepro.2018.12.112](https://doi.org/10.1016/j.jclepro.2018.12.112)
- Paredes-Sánchez JP, López-Ochoa LM, López-González LM, Xiberta-Bernat J 2016 Bioenergy for District Bioheating System (DBS) from eucalyptus residues in a European coal-producing region. *Energy Convers. Manag.* 126, 960–970. doi: [10.1016/j.enconman.2016.08.084](https://doi.org/10.1016/j.enconman.2016.08.084)
- Pariente-David S 2014 Successful Grid Integration of Renewable Energy: Integration is the Name of the Game. *Int. Assoc. Energy Econ.*, 29–30. <https://www.iaee.org/newsletter/issue/29>
- Pariente-David S 2016 The Cost and Value of Renewable Energy: Revisiting Electricity Economics. *Int. Assoc. Energy Econ. IAAE Energy Forum*, 21–23.
- Pariente-David S, Jannet Allal H Ben 2019 North Africa. An energy bridge between the African and European Continents, in *Empowering Africa: Access to power in the African continent*, eds. Colantoni L, Montesano G, Sartori N (Peter Lang), 199–231. doi: [10.3726/b15292/20](https://doi.org/10.3726/b15292/20)
- Pérez-Andreu V, Aparicio-Fernández C, Martínez-Iberón A, Vivancos J-L 2018 Impact of climate change on heating and cooling energy demand in a residential building in a Mediterranean climate. *Energy* 165, 63–74. doi: [10.1016/j.energy.2018.09.015](https://doi.org/10.1016/j.energy.2018.09.015)
- Petersen LE, Troen I, Ejsing Jørgensen H, Mann J 2014 The new European wind atlas. *Energy Bull.* 17, 34–39.
- Petterssen S 1956 *Weather Analysis and Forecasting*. McGraw-Hill.
- Pfeifroth U, Sánchez-Lorenzo A, Manara V, Trentmann J, Hollmann R 2018 Trends and variability of surface solar radiation in Europe based on surface- and satellite-based data records. *JGR Atmos.* 123, 1735–1754. doi: [10.1002/2017JD027418](https://doi.org/10.1002/2017JD027418)
- Pfenninger S, Staffell I 2016 Long-term patterns of European PV output using 30 years of validated hourly reanalysis and satellite data. *Energy* 114, 1251–1265. doi: [10.1016/j.energy.2016.08.060](https://doi.org/10.1016/j.energy.2016.08.060)
- Philipona R, Behrens K, Ruckstuhl C 2009 How declining aerosols and rising greenhouse gases forced rapid warming in Europe since the 1980s. *Geophys. Res. Lett.* 36. doi: [10.1029/2008gl036350](https://doi.org/10.1029/2008gl036350)
- Piante C, Ody D 2015 Blue Growth in the Mediterranean Sea: The Challenge of Good Environmental Status. <http://www.developpement-durable.gouv.fr/>
- Plan Bleu 2008 Climate Change and Energy in the Mediterranean. Sophia Antipolis [http://www.eib.org/attachments/country/climate\\_change\\_energy\\_mediterranean\\_en.pdf](http://www.eib.org/attachments/country/climate_change_energy_mediterranean_en.pdf).
- Ponce de León S, Orfila A, Simarro G 2016 Wave energy in the Balearic Sea. Evolution from a 29 year spectral wave hindcast. *Renew. Energy* 85, 1192–1200. doi: [10.1016/j.renene.2015.07.076](https://doi.org/10.1016/j.renene.2015.07.076)
- Possner A, Caldeira K 2017 Geophysical potential for wind energy over the open oceans. *Proc. Natl. Acad. Sci. U. S. A.* 114, 11338–11343. doi: [10.1073/pnas.1705710114](https://doi.org/10.1073/pnas.1705710114)
- Poudineh R, Sen A, Fattouh B 2018 Advancing renewable energy in resource-rich economies of the MENA. *Renew. Energy* 123, 135–149. doi: [10.1016/j.renene.2018.02.015](https://doi.org/10.1016/j.renene.2018.02.015)
- Pozo-Vazquez D, Santos-Alamillos FJ, Lara-Fanego V, Ruiz-Arias JA, Tovar-Pescador J 2011 The Impact of the NAO on the Solar and Wind Energy Resources in the Mediterranean Area, in *Hydrological Socio-economic and Ecological Impacts of the North Atlantic Oscillation in the Mediterranean Region, Advances in Global Change Research*, eds. Vicente-Serrano SM, Trigo RM (Springer Netherlands), 213–231. doi: [10.1007/978-94-007-1372-7\\_15](https://doi.org/10.1007/978-94-007-1372-7_15)
- Pryor SC, Barthelmie RJ, Schoof JT 2006 Inter-annual variability of wind indices across Europe. *Wind Energy* 9, 27–38. doi: [10.1002/we.178](https://doi.org/10.1002/we.178)
- Pulighe G, Bonati G, Colangeli M, Morese MM, Traverso L et al. 2019 Ongoing and emerging issues for sustainable bioenergy production on marginal lands in the Mediterranean regions. *Renew. Sustain. Energy Rev.* 103, 58–70. doi: [10.1016/J.RSER.2018.12.043](https://doi.org/10.1016/J.RSER.2018.12.043)
- Pye S, Bataille C 2016 Improving deep decarbonization modelling capacity for developed and developing country contexts. *Clim. Policy* 16, S27–S46. doi: [10.1080/14693062.2016.1173004](https://doi.org/10.1080/14693062.2016.1173004)
- Raymond F, Ullmann A, Trambly Y, Drobinski P, Camberlin P 2019 Evolution of Mediterranean extreme dry spells during the wet season under climate change. *Reg. Environ. Chang.* 19, 2339–2351. doi: [10.1007/s10113-019-01526-3](https://doi.org/10.1007/s10113-019-01526-3)
- Rizzo A, Maro P 2018 Implementing Nationally Determined Contributions (NDCs) in the South Mediterranean region. Perspectives on climate action from eight countries.
- Robin K 2015 Climate - What did Mediterranean countries commit to in view of the COP21?
- Rodríguez-Loinaz G, Amezcaga I, Onaindia M 2013 Use of native species to improve carbon sequestration and contribute towards solving the environmental problems of the timberlands in Biscay, northern Spain. *J. Environ. Manage.* 120, 18–26. doi: [10.1016/j.jenvman.2013.01.032](https://doi.org/10.1016/j.jenvman.2013.01.032)
- Rogelj J, Luderer G, Pietzcker RC, Kriegler E, Schaeffer M et al. 2015 Energy system transformations for limiting end-of-century warming to below 1.5 °C. *Nat. Clim. Chang.* 5, 519–527. doi: [10.1038/nclimate2572](https://doi.org/10.1038/nclimate2572)
- Roques F, Verhaeghe C 2016 Options for the future of Power System Regional Coordination. Paris [https://docstore.entsoe.eu/Documents/Publications/Position\\_papers\\_and\\_reports/entsoe\\_fti\\_161207.pdf](https://docstore.entsoe.eu/Documents/Publications/Position_papers_and_reports/entsoe_fti_161207.pdf)
- Rose SK, Kriegler E, Bibas R, Calvin K V., Popp A et al. 2014 Bioenergy in energy transformation and climate management. *Clim. Change* 123, 477–493.

- doi: [10.1007/s10584-013-0965-3](https://doi.org/10.1007/s10584-013-0965-3)
- Rowlinson ET 2015 À Qui le Soleil: How Morocco's Developing Solar Capacities Have Altered Urban Infrastructural Provisions. <https://repositories.lib.utexas.edu/bitstream/handle/2152/35485/ROWLINSON-THESIS-2015.pdf?sequence=1&isAllowed=y>
- Royaume du Maroc 2017 Stratégie nationale énergétique – horizon 2030.
- Rusu E, Rusu L 2019 Evaluation of the wind power potential in the European nearshore of the Mediterranean Sea. *E3S Web Conf.* 103, 1003. doi: [10.1051/e3sconf/201910301003](https://doi.org/10.1051/e3sconf/201910301003)
- Saffih-Hdadi K, Mary B 2008 Modeling consequences of straw residues export on soil organic carbon. *Soil Biol. Biochem.* 40, 594–607. doi: [10.1016/j.soilbio.2007.08.022](https://doi.org/10.1016/j.soilbio.2007.08.022)
- Sánchez-Lorenzo A, Enriquez-Alonso A, Wild M, Trentmann J, Vicente-Serrano SM et al. 2017 Trends in downward surface solar radiation from satellites and ground observations over Europe during 1983–2010. *Remote Sens. Environ.* 189, 108–117. doi: [10.1016/J.RSE.2016.11.018](https://doi.org/10.1016/J.RSE.2016.11.018)
- Sanseverino ER, Sanseverino RR, Favuzza S, Vaccaro V 2014 Near zero energy islands in the Mediterranean: Supporting policies and local obstacles. *Energy Policy* 66, 592–602. doi: [10.1016/j.enpol.2013.11.007](https://doi.org/10.1016/j.enpol.2013.11.007)
- Sansilvestri R, Cuccarollo M, Frascaria-Lacoste N, Benito-Garzon M, Fernandez-Manjarrés J 2020 Evaluating climate change adaptation pathways through capital assessment: five case studies of forest social-ecological systems in France. *Sustain. Sci.* 15, 539–553. doi: [10.1007/s11625-019-00731-7](https://doi.org/10.1007/s11625-019-00731-7)
- Santos-Alamillos FJ, Pozo-Vázquez D, Ruiz-Arias JA, Lara-Fanego V, Tovar-Pescador J 2012 Analysis of Spatiotemporal Balancing between Wind and Solar Energy Resources in the Southern Iberian Peninsula. *J. Appl. Meteorol. Climatol.* 51, 2005–2024. doi: [10.1175/jamc-d-11-0189.1](https://doi.org/10.1175/jamc-d-11-0189.1)
- Santos-Alamillos FJ, Thomaidis NS, Usaola-García J, Ruiz-Arias JA, Pozo-Vázquez D 2017 Exploring the mean-variance portfolio optimization approach for planning wind repowering actions in Spain. *Renew. Energy* 106, 335–342. doi: [10.1016/j.renene.2017.01.041](https://doi.org/10.1016/j.renene.2017.01.041)
- Schewe J, Gosling SN, Reyer C, Zhao F, Ciais P et al. 2019 State-of-the-art global models underestimate impacts from climate extremes. *Nat. Commun.* 10, 1005. doi: [10.1038/s41467-019-08745-6](https://doi.org/10.1038/s41467-019-08745-6)
- Schilling J, Freier KP, Hertig E, Scheffran J 2012 Climate change, vulnerability and adaptation in North Africa with focus on Morocco. *Agric. Ecosyst. Environ.* 156, 12–26. doi: [10.1016/J.AGEE.2012.04.021](https://doi.org/10.1016/J.AGEE.2012.04.021)
- Searchinger T, Heimlich R, Houghton RA, Dong F, Elobeid A et al. 2008 Use of U.S. Croplands for Biofuels Increases Greenhouse Gases Through Emissions from Land-Use Change. *Science (80-. )*. 319, 1238–1240. doi: [10.1126/science.1151861](https://doi.org/10.1126/science.1151861)
- Sen Nag O 2017 The Most Populated Islands In The Mediterranean Sea. <https://www.worldatlas.com/articles/the-most-populated-islands-in-the-mediterranean-sea.html>
- Sierra JP, Casas-Prat M, Campins E 2017 Impact of climate change on wave energy resource: The case of Menorca (Spain). *Renew. Energy* 101, 275–285. doi: [10.1016/j.renene.2016.08.060](https://doi.org/10.1016/j.renene.2016.08.060)
- Soares PMM, Brito MC, Careto JAM 2019 Persistence of the high solar potential in Africa in a changing climate. *Environ. Res. Lett.* 14, 124036. doi: [10.1088/1748-9326/ab51a1](https://doi.org/10.1088/1748-9326/ab51a1)
- Solaun K, Cerdá E 2020 Impacts of climate change on wind energy power – Four wind farms in Spain. *Renew. Energy* 145, 1306–1316. doi: [10.1016/j.renene.2019.06.129](https://doi.org/10.1016/j.renene.2019.06.129)
- Soukissian TH, Denaxa D, Karathanasi F, Prospathopoulos A, Sarantakos K et al. 2017 Marine renewable energy in the Mediterranean Sea: Status and perspectives. *Energies* 10, 1512. doi: [10.3390/en10101512](https://doi.org/10.3390/en10101512)
- Staffell I, Pfenninger S 2016 Using bias-corrected reanalysis to simulate current and future wind power output. *Energy* 114, 1224–1239. doi: [10.1016/j.energy.2016.08.068](https://doi.org/10.1016/j.energy.2016.08.068)
- Stecher K, Brosowski A, Thrän D 2013 Biomass potential in Africa. Abu Dhabi.
- Szabó L, Mezősi A, Törőcsik Á, Kotek P, Kácsor E et al. 2015 Dialogue on a RES policy Framework for 2030. D3.1a Renewable Based District Heating in Europe - Policy Assessment of Selected Member States.
- Taibi E, Nikolakakis T, Gutierrez L, Fernandez C, Kiviluoma J et al. 2018 Power system flexibility for the energy transition. Part 1: Overview for policy makers. Abu Dhabi [https://irena.org/-/media/Files/IRENA/Agency/Publication/2018/Nov/IRENA\\_Power\\_system\\_flexibility\\_1\\_2018.pdf?la=en&hash=72EC26336F127C7D51DF798CE19F-477557CE9A82](https://irena.org/-/media/Files/IRENA/Agency/Publication/2018/Nov/IRENA_Power_system_flexibility_1_2018.pdf?la=en&hash=72EC26336F127C7D51DF798CE19F-477557CE9A82)
- Tantet A, Stéfanon M, Drobinski P, Badosa J, Concettini S et al. 2019 E4Clim 1.0: The Energy for a Climate Integrated Model: Description and Application to Italy. *Energies* 12, 4299. doi: [10.3390/en12224299](https://doi.org/10.3390/en12224299)
- Thornton HE, Hoskins BJ, Scaife AA 2016 The role of temperature in the variability and extremes of electricity and gas demand in Great Britain. *Environ. Res. Lett.* 11, 114015. doi: [10.1088/1748-9326/11/11/114015](https://doi.org/10.1088/1748-9326/11/11/114015)
- Tilman D, Socolow R, Foley JA, Hill J, Larson E et al. 2009 Beneficial Biofuels--The Food, Energy, and Environment Trilemma. *Science (80-. )*. 325, 270–271. doi: [10.1126/science.1177970](https://doi.org/10.1126/science.1177970)
- Timmerberg S, Sanna A, Kaltschmitt M, Finkbeiner M 2019 Renewable electricity targets in selected MENA countries – Assessment of available resources, generation costs and GHG emissions. *Energy Re-*

- ports 5, 1470–1487. doi: [10.1016/j.egy.2019.10.003](https://doi.org/10.1016/j.egy.2019.10.003)
- Tobin I, Greuell W, Jerez S, Ludwig F, Vautard R et al. 2018 Vulnerabilities and resilience of European power generation to 1.5°C, 2°C and 3°C warming. *Environ. Res. Lett.* 13, 44024. doi: [10.1088/1748-9326/aab211](https://doi.org/10.1088/1748-9326/aab211)
- Tobin I, Jerez S, Vautard R, Thais F, van Meijgaard E et al. 2016 Climate change impacts on the power generation potential of a European mid-century wind farms scenario. *Environ. Res. Lett.* 11, 34013. doi: [10.1088/1748-9326/11/3/034013](https://doi.org/10.1088/1748-9326/11/3/034013)
- Tobin I, Vautard R, Balog I, Bréon F-M, Jerez S et al. 2015 Assessing climate change impacts on European wind energy from ENSEMBLES high-resolution climate projections. *Clim. Change* 128, 99–112. doi: [10.1007/s10584-014-1291-0](https://doi.org/10.1007/s10584-014-1291-0)
- Trieb F, Schillings C, Pregger T, O'Sullivan M 2012 Solar electricity imports from the Middle East and North Africa to Europe. *Energy Policy* 42, 341–353. doi: [10.1016/j.enpol.2011.11.091](https://doi.org/10.1016/j.enpol.2011.11.091)
- Ueckerdt F, Brecha R, Luderer G 2015 Analyzing major challenges of wind and solar variability in power systems. *Renew. Energy* 81, 1–10. doi: [10.1016/j.renene.2015.03.002](https://doi.org/10.1016/j.renene.2015.03.002)
- UfM 2019 Tracking and enhancing international private climate finance in the Southern-Mediterranean Region. <https://ufmsecretariat.org/wp-content/uploads/2019/09/Private-Climate-Finance-Tracking-and-enhancing-international-private-climate-finance-in-the-Southern-Mediterranean-Region.pdf>
- UK DECC 2013 More interconnection: improving energy security and lowering bills. [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/266460/More\\_interconnection\\_-\\_improving\\_energy\\_security\\_and\\_lowering\\_bills.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/266460/More_interconnection_-_improving_energy_security_and_lowering_bills.pdf) [Accessed June 29, 2019]
- UN DESA 2016 Energy Balances. <https://unstats.un.org/unsd/energy/balance/default.htm>
- UNEP/MAP-Plan Bleu 2009 State of the Environment and Development in the Mediterranean. Athens, Greece.
- UNEP/MAP 2007 MAP Mediterranean Action Plan. Energy and sustainable development in the Mediterranean: in *Proceedings of the Regional Workshop, Monaco* (Monaco). <https://wedocs.unep.org/bitstream/handle/20.500.11822/516/mts167.pdf?sequence=2&is-Allowed=y> [Accessed June 30, 2019].
- UNFCCC 2018 Biennial Assessment and Overview of Climate Finance Flows. Technical Report. [https://unfccc.int/sites/default/files/resource/2018\\_BA\\_Technical\\_Report\\_Final\\_Feb\\_2019.pdf](https://unfccc.int/sites/default/files/resource/2018_BA_Technical_Report_Final_Feb_2019.pdf)
- van Vliet MTH, Sheffield J, Wiberg D, Wood EF 2016a Impacts of recent drought and warm years on water resources and electricity supply worldwide. *Environ. Res. Lett.* 11, 124021. doi: [10.1088/1748-9326/11/12/124021](https://doi.org/10.1088/1748-9326/11/12/124021)
- van Vliet MTH, Wiberg D, Leduc S, Riahi K 2016b Power-generation system vulnerability and adaptation to changes in climate and water resources. *Nat. Clim. Chang.* 6, 375–380. doi: [10.1038/nclimate2903](https://doi.org/10.1038/nclimate2903)
- van Vliet MTH, Yearsley JR, Franssen WHP, Ludwig F, Haddeland I et al. 2012a Coupled daily streamflow and water temperature modelling in large river basins. *Hydrol. Earth Syst. Sci.* 16, 4303–4321. doi: [10.5194/hess-16-4303-2012](https://doi.org/10.5194/hess-16-4303-2012)
- van Vliet MTH, Yearsley JR, Ludwig F, Vögele S, Lettenmaier DP et al. 2012b Vulnerability of US and European electricity supply to climate change. *Nat. Clim. Chang.* 2, 676–681. doi: [10.1038/nclimate1546](https://doi.org/10.1038/nclimate1546)
- Vautard R, Cattiaux J, Yiou P, Thépaut J-N, Ciais P 2010 Northern Hemisphere atmospheric stilling partly attributed to an increase in surface roughness. *Nat. Geosci.* 3, 756–761. doi: [10.1038/ngeo979](https://doi.org/10.1038/ngeo979)
- Vidican G 2016 Scaling Up Renewable Energy Deployment in North Africa. *Regul. Investments Energy Mark.*, 73–87. doi: [10.1016/b978-0-12-804436-0.00004-7](https://doi.org/10.1016/b978-0-12-804436-0.00004-7)
- Wang Z, Lin X, Tong N, Li Z, Sun S et al. 2020 Optimal planning of a 100% renewable energy island supply system based on the integration of a concentrating solar power plant and desalination units. *Int. J. Electr. Power Energy Syst.* 117, 105707. doi: [10.1016/j.ijepes.2019.105707](https://doi.org/10.1016/j.ijepes.2019.105707)
- Weber J, Gotzens F, Witthaut D 2018 Impact of strong climate change on the statistics of wind power generation in Europe. *Energy Procedia* 153, 22–28. doi: [10.1016/j.egypro.2018.10.004](https://doi.org/10.1016/j.egypro.2018.10.004)
- Wenz L, Levermann A, Auffhammer M 2017 North-south polarization of European electricity consumption under future warming. *Proc. Natl. Acad. Sci. U. S. A.* 114, E7910–E7918. doi: [10.1073/pnas.1704339114](https://doi.org/10.1073/pnas.1704339114)
- Widén J, Carpman N, Castellucci V, Lingfors D, Olauson J et al. 2015 Variability assessment and forecasting of renewables: A review for solar, wind, wave and tidal resources. *Renew. Sustain. Energy Rev.* 44, 356–375. doi: [10.1016/j.rser.2014.12.019](https://doi.org/10.1016/j.rser.2014.12.019)
- Wild M, Folini D, Henschel F 2017 Impact of climate change on future concentrated solar power (CSP) production. *AIP Conf. Proc.* 1810, 100007. doi: [10.1063/1.4975562](https://doi.org/10.1063/1.4975562)
- Wild M, Folini D, Henschel F, Fischer N, Müller B 2015 Projections of long-term changes in solar radiation based on CMIP5 climate models and their influence on energy yields of photovoltaic systems. *Sol. Energy* 116, 12–24. doi: [10.1016/j.solener.2015.03.039](https://doi.org/10.1016/j.solener.2015.03.039)
- Wild M, Gilgen H, Roesch A, Ohmura A, Long CN et al. 2005 From dimming to brightening: Decadal changes in solar radiation at Earth's surface. *Science (80-. J.)* 308, 847–850. doi: [10.1126/science.1103215](https://doi.org/10.1126/science.1103215)
- World Bank 2010 2010 Annual Report. Washington, DC. <http://documents.worldbank.org/curated/en/408911468331735129/2010-annual-report>
- World Bank 2018 Energy Subsidies in Mediterranean



- developing countries and their reform. [https://www.cape4financeministry.org/sites/cape/files/in-line-files/Session 1-3. Thomas Flochel\\_M](https://www.cape4financeministry.org/sites/cape/files/in-line-files/Session%201-3.Thomas%20Flochel_M)
- Yearsley JR 2009 A semi-Lagrangian water temperature model for advection-dominated river systems. *Water Resour. Res.* 45. doi: [10.1029/2008wr007629](https://doi.org/10.1029/2008wr007629)
- Yoon SJ, Son Y-I, Kim Y-K, Lee J-G 2012 Gasification and power generation characteristics of rice husk and rice husk pellet using a downdraft fixed-bed gasifier. *Renew. Energy* 42, 163–167. doi: [10.1016/j.renene.2011.08.028](https://doi.org/10.1016/j.renene.2011.08.028)
- Zabalza Bribián I, Valero Capilla A, Aranda Usón A 2011 Life cycle assessment of building materials: Comparative analysis of energy and environmental impacts and evaluation of the eco-efficiency improvement potential. *Build. Environ.* 46, 1133–1140. doi: [10.1016/j.buildenv.2010.12.002](https://doi.org/10.1016/j.buildenv.2010.12.002)
- Zachariadis T, Hadjinicolaou P 2014 The effect of climate change on electricity needs - A case study from Mediterranean Europe. *Energy* 76, 899–910. doi: [10.1016/j.energy.2014.09.001](https://doi.org/10.1016/j.energy.2014.09.001)
- Zejli D, Bennouna A 2009 Wind Energy in Morocco: Which Strategy for Which Development?, in (Springer, Dordrecht), 151–173. doi: [10.1007/978-1-4020-9892-5\\_9](https://doi.org/10.1007/978-1-4020-9892-5_9)
- Zeng Z, Ziegler AD, Searchinger T, Yang L, Chen A et al. 2019 A reversal in global terrestrial stilling and its implications for wind energy production. *Nat. Clim. Chang.* 9, 979–985. doi: [10.1038/s41558-019-0622-6](https://doi.org/10.1038/s41558-019-0622-6)
- Zickfeld F, Wieland A 2012 Desert Power 2050: Perspectives on a Sustainable Power System for EUMENA. Germany [http://mait.camins.cat/ET2050\\_library/docs/tech/energy/2012\\_2050\\_Desert\\_Power.pdf](http://mait.camins.cat/ET2050_library/docs/tech/energy/2012_2050_Desert_Power.pdf) [Accessed June 29, 2019]
- Zizzo G, Beccali M, Bonomolo M, di Pietra B, Ippolito MG et al. 2017 A feasibility study of some DSM enabling solutions in small islands: The case of Lampedusa. *Energy* 140, 1030–1046. doi: [10.1016/j.energy.2017.09.069](https://doi.org/10.1016/j.energy.2017.09.069)
- Zouiri H, Elmessaoudi H 2018 Energies renouvelables et développement durable au Maroc. *Rev. Estud. Front. del Estrecho Gibraltar REFEG* 6/20, 1–29.



## Information about authors

### Coordinating Lead Authors

Brian Azzopardi:

*Malta College of Arts, Science and Technology (MCAST), Paola, Malta*

Philippe Drobinski:

*Laboratoire de Météorologie Dynamique / Institut Pierre Simon Laplace (LMD/IPSL), École Polytechnique, IP Paris, Sorbonne Université, ENS, PSL Université, CNRS, Palaiseau, France*

### Lead Authors

Houda Allal:

*Mediterranean Energy Observatory (OME), Paris, France*

Vincent Bouchet:

*École polytechnique (i3-CRG), Palaiseau, France*

Eduard Civel:

*Ecole Polytechnique, Palaiseau, France*

Anna Creti:

*Climate Economics Chair, Economics of Gas Chair, Paris Dauphine University, Paris, France*

Neven Duic:

*Faculty of Mechanical Engineering and Naval Architecture, University of Zagreb, Croatia*

Nestor Fylaktos:

*The Cyprus Institute, Nicosia, Cyprus*

Joseph Mutale:

*University of Manchester, Manchester, United Kingdom*

Silvia Pariente-David:

*International Energy Consultant, Marseille, France*

Joe Ravetz:

*Manchester Urban Institute, Manchester, United Kingdom*

Constantinos Taliotis:

*The Cyprus Institute, Nicosia, Cyprus*

Robert Vautard:

*Laboratory of Climate and Environmental Sciences (LSCE), Pierre Simon Laplace Institute (IPSL), France*

### Contributing Authors

Kaouther Ben Nasr:

*University of Carthage, Research and Technology Centre of Energy (CRTE), Hammam-Lif, Tunisia*

Thierry Brunelle:

*International Research Center on Environment and Development (CIRED), Montpellier, France*

Mikaël Cugnet:

*French Commission for Atomic and Alternative Energies (CEA), Grenoble, France*

Paola de Joanna:

*University of Naples Federico II, Naples, Italy*

Sokol Dervishi:

*Epoka University, Tirana, Albania*

Juan Fernandez-Manjarrés:

*University Paris-Saclay, Paris, France*

Dora Francese:

*University of Naples Federico II, Naples, Italy*

Benoit Gabrielle:

*Paris Institute of Life and Environmental Sciences (AgroParisTech), Paris, France*

Lisa Guarrera:

*Mediterranean Energy Observatory (OME), Paris, France*

Victor Homar Santaner:

*University of the Balearic Islands, Palma, Spain*

Boutaina Ismaili Idrissi:

*Agdal -University Mohammed V, Rabat, Morocco*

Rémy Lapère:

*Laboratoire de Météorologie Dynamique / Institut Pierre Simon Laplace (LMD/IPSL), École Polytechnique, IP Paris, Sorbonne Université, ENS, PSL Université, CNRS, Palaiseau, France*

Aina Maimo-Far:

*University of the Balearic Islands, Palma, Spain*

Emanuela Menichetti:

*Mediterranean Energy Observatory (OME), Paris, France*

Lina Murauskaite:

*Lithuanian Energy Institute, Kaunas, Lithuania*

Federico Pontoni:

*Centre for Research on Geography, Resources, Environment, Energy & Networks (GREEN), Bocconi University, Milan, Italy*

Gianmaria Sannino:

*Climate Modelling Laboratory and Impacts, National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Rome, Italy*

Roxane Sansilvestri:

*Campus de la transition, France*

Alexis Tantet:

*Laboratoire de Météorologie Dynamique / Institut Pierre Simon Laplace (LMD/IPSL), École Polytechnique, IP Paris, Sorbonne Université, ENS, PSL Université, CNRS, Palaiseau, France*

Michelle Van Vliet:

*Utrecht University, Utrecht, Netherlands*



# 4 ECOSYSTEMS

**Coordinating Lead Authors:**

Mario V. Balzan (Malta), Abed El Rahman Hassoun (Lebanon)


**Lead Authors:**

Najet Aroua (Algeria), Virginie Baldy (France), Magda Bou Dagher (Lebanon), Cristina Branquinho (Portugal), Jean-Claude Dutay (France), Monia El Bour (Tunisia), Frédéric Médail (France), Meryem Mojtahid (Morocco/France), Alejandra Morán-Ordóñez (Spain), Pier Paolo Roggero (Italy), Sergio Rossi Heras (Italy), Bertrand Schatz (France), Ioannis N. Vogiatzakis (Cyprus), George N. Zaimis (Greece), Patrizia Ziveri (Spain)

**Contributing Authors:**

Marie Abboud-Abi Saab (Lebanon), Aitor Ameztegui (Spain), Margaretha Breil (Italy), Thierry Gauquelin (France), Ulse R. Geijzendorffer (France), Aristeidis Koutroulis (Greece), Jürg Luterbacher (Germany), Mohammad Merheb (Lebanon), Cesar Terrer Moreno (Spain), Marco Turco (Spain), Elena Xoplaki (Germany)

*This chapter should be cited as: Balzan MV, Hassoun AER, Aroua N, Baldy V, Bou Dagher M, Branquinho C, Dutay J-C, El Bour M, Médail F, Mojtahid M, Morán-Ordóñez A, Roggero PP, Rossi Heras S, Schatz B, Vogiatzakis IN, Zaimis GN, Ziveri P 2020 Ecosystems. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 323-468.*



# Table of contents

<b>4 Ecosystems</b>	<b>328</b>
Executive summary	328
<b>4.1 Marine ecosystems</b>	<b>329</b>
4.1.1 Current condition and past trends	329
4.1.1.1 Observed changes	329
Key habitats undergoing change	331
- Coralligenous	331
- Deep sea ecosystems	332
- Planktonic ecosystems	333
- Large vertebrates	335
Changes in biodiversity	335
4.1.1.2 Past changes	339
Response of marine ecosystems to past temperature changes	339
Response of marine ecosystems to past changes in stratification and ventilation	340
Response of marine ecosystems to past changes in productivity	341
Response of marine ecosystems to past changes in pH	341
4.1.2 Projected vulnerabilities and risks	342
4.1.2.1 Projected impacts and risks	342
Projected impacts on microbes	342
Projected impacts on primary and secondary production	342
Projected impacts on macrobenthic and pelagic species	344
- Corals	344
- Seagrass	345
- Mussels	346
- Jellyfish	346
Winners and losers	346
4.1.2.2 Vulnerabilities	347
Climate-related vulnerabilities	347
Anthropogenic vulnerabilities	348
4.1.3 Adaptation	348
4.1.3.1 Long-term monitoring and adaptation strategies	348
4.1.3.2 The role of Marine Protected Areas (MPAs) for adaptation	348
4.1.3.3 Management of fisheries and adaptation	349
4.1.3.4 Adaptation strategies for ocean warming and ocean acidification in the Mediterranean Sea	350
4.1.3.5 Regional observation networks as a tool for adaptation	351
<b>4.2 Coastal ecosystems</b>	<b>352</b>
4.2.1 Current condition and past trends	352
4.2.1.1 Observed changes	352
Natural Mediterranean habitats under severe degradation	352
- Sandy beaches and sand dunes	353
- Rocky coasts	354
- Coastal wetlands	355
- Seagrass meadows	355
- Coastal lagoons and deltas	356
- Salt marshes	357

- Coastal aquifers .....	358
Risks from non-indigenous species .....	358
- Phytoplankton .....	358
- Jellyfish .....	359
- Fish .....	359
- Plants .....	360
- Other non-indigenous species .....	360
4.2.1.2 Past changes .....	360
Response of coastal ecosystems to past changes in sea level .....	360
Response of coastal ecosystems to past climate variability .....	362
4.2.2 Projected vulnerabilities and risks .....	362
4.2.2.1 Projections and risks based on biological groups .....	362
Phytoplankton .....	362
Fish .....	363
Seaweed .....	364
Corals .....	364
4.2.2.2 Projections and risks based on key natural habitats .....	365
Sandy beaches/dunes .....	365
Rocky shores .....	366
Coastal wetlands .....	366
Seagrass meadows .....	367
Coastal lagoons .....	367
Deltas .....	367
Coastal aquifers .....	368
4.2.2.3 Vulnerabilities .....	369
Coastal urbanization .....	369
Sea level rise .....	371
4.2.3 Adaptation .....	371
4.2.3.1 Adaptation of different coastal systems .....	371
4.2.3.2 Harmful algal bloom monitoring .....	372
4.2.3.3 Early detection of potentially dangerous species .....	373
4.2.3.4 Adaptation management strategies for the jellyfish <i>Pelagia noctiluca</i> .....	373
4.2.3.5 Ecosystem-based adaptation management .....	374
4.2.3.6 The role of institutions/actors and local communities: recommendations .....	374
<b>4.3 Terrestrial and freshwater ecosystems .....</b>	<b>375</b>
4.3.1 Current conditions and past trends .....	375
4.3.1.1 Past climate variability and its impact on terrestrial ecosystems .....	375
4.3.1.2 Direct human impacts on ecosystems in the past .....	376
Forests .....	377
- The human footprint in Mediterranean forests .....	378
- Ecosystem services provision by Mediterranean forests .....	379
Mountains .....	380
- Land use changes in mountain regions .....	381
- Mountain biodiversity changes .....	382
Drylands and shrublands .....	382
Agroecosystems .....	383
- Agroecosystem development in different regions .....	384
- Perennial crops .....	385
- Vegetables .....	386
- Winter cereals .....	386
- Grasslands and grazing systems .....	387
- Ecosystem services related to pollination .....	387
Freshwater ecosystems .....	388
- River regulation .....	388

- Groundwater depletion .....	388
- Hydrologic regimes .....	389
- Land-use changes, reduction of wetlands and riparian areas .....	389
- Water quality .....	390
- Freshwater species .....	391
- Protected areas (Natura 2000 network and Ramsar Convention) .....	391
<b>4.3.2 Projected vulnerabilities and risks .....</b>	<b>392</b>
4.3.2.1 Forests .....	392
Changes in forest ecosystem health and ecosystem services provision .....	393
Changes in species range, abundance and extinction .....	394
Fire activity and burnt areas across the Mediterranean .....	395
4.3.2.2 Mountains .....	395
4.3.2.3 Drylands and shrublands .....	396
Droughts .....	398
4.3.2.4 Agriculture and pasturelands .....	398
Cropping systems .....	399
Grasslands and grazing systems .....	401
4.3.2.5 Freshwater ecosystems .....	401
Rivers and streams .....	401
Wetlands .....	402
Freshwater biodiversity .....	402
<b>4.3.3 Adaptation .....</b>	<b>402</b>
4.3.3.1 Forests .....	403
Biological adaptation .....	403
Limits to adaptation .....	404
Measures to promote adaptation .....	405
4.3.3.2 Mountain ecosystems .....	406
4.3.3.3 Drylands and shrublands .....	406
4.3.3.4 Agriculture and pasturelands .....	407
4.3.3.5 Freshwater ecosystems .....	410
<b>Box 4.1 Bio-indicators for the assessment of changes in Mediterranean marine ecosystems .....</b>	<b>413</b>
<b>Box 4.2 Urban biodiversity in the Mediterranean Region .....</b>	<b>414</b>
Consequences for biodiversity and ecosystem services in urban areas .....	414
Urban biodiversity and ecosystem services .....	415
<b>Box 4.3 Nitrogen deposition and ecosystems .....</b>	<b>414</b>
<b>Box 4.4 Mediterranean islands .....</b>	<b>416</b>
Islands as laboratories .....	416
Recent evidence of change .....	416
Climate change projections and islands .....	416
Vulnerability/resilience .....	416
Conservation and adaptation .....	416
<b>References .....</b>	<b>418</b>
<b>Information about authors .....</b>	<b>468</b>

## 4 Ecosystems

### Executive summary

#### Marine ecosystems

Despite covering only 0.82% of the ocean's surface, the Mediterranean Sea supports up to 18% of all known marine species, with 21% being listed as vulnerable and 11% as endangered. The accelerated spread of tropical non-indigenous species is leading to the "tropicalization" of Mediterranean fauna and flora as a result of warming and extreme heat waves since the 1990s. The acidification rate in the Mediterranean waters has ranged between 0.055 and 0.156 pH units since the pre-industrial period, affecting the marine trophic chain, from its primary producers (i.e., coccolithophores and foraminifera) to corals and coralline red algae.

Projections for high emission scenarios show that endemic assemblages will be modified with numerous species becoming extinct in the mid 21st century and changes to the natural habitats of commercially valuable species, which would have many repercussions on marine ecosystem services such as tourism, fisheries, climate regulation, and ultimately on human health.

Adaptation strategies to reduce environmental change impacts need effective mitigation policies and actions. They require anticipatory planning to enable them to tackle problems while they are still manageable. Given the diversity of each Mediterranean sub-basin, wider monitoring coverage is needed to strengthen our knowledge about the different adaptation processes that characterize and best suit each geographical zone. Adaptation implies the implementation of more sustainable fishing practices as well as reducing pollution from agricultural activity, sustainable tourism or developing more effective waste management. Marine protected areas can potentially have an insurance role if they are established in locations not particularly vulnerable to ocean acidification and climate change.

#### Coastal ecosystems

The coastal zone, i.e., the area in which the interaction between marine systems and the land dominate ecological and resource systems, is a hotspot of risks, especially in the south-eastern Mediterranean region. Alterations to coastal ecosystems (lagoons, deltas, salt marshes, etc.) due to climate change and human activities affect

the flow of nutrients to the sea, the magnitude, timing and composition of potentially harmful/toxic plankton blooms. They also significantly increase the number and frequency of jellyfish outbreaks, and could have negative impacts on fisheries. 1.2 to 5% of seagrass meadows in the Mediterranean Sea, which represent 5 to 17% of the worldwide seagrass habitat, are lost each year. Among them, almost half of the surveyed *Posidonia oceanica* sites have suffered net density losses of over 20% in 10 years. As for fish, non-indigenous species and climate change cause local extinction.

Projected temperature increases combined with a decrease in nutrient replenishment and ocean acidification, are expected to cause changes in plankton communities, negative impacts on fish, corals, seagrass meadows and propagation of non-indigenous species. Projected sea level rise will impact coastal wetlands deltas and lagoons. Extensive urbanization added to climate change is also expected to threaten coastal ecosystems, human health and well-being.

A nexus approach is required when trying to establish adaptation methods for the entire Mediterranean, while taking into account ecosystem-based management, synergies and conflicts, integrating local knowledge and institutions. Suitable adaptation policies include reducing pollution runoff, both from agriculture and industry and waste management, and policies to limit or prevent acidification. Conservation planning and management should focus on cross-cutting approaches and building resilience between structural and functional connectivities of various fields.

#### Terrestrial ecosystems

Biodiversity changes in the Mediterranean over the past 40 years have occurred more quickly and been more significant than in other regions of the world. Urbanization and the loss of grasslands are key factors of ecosystem degradation across the region. Since 1990, agricultural abandonment has led to a general increase in forest areas in the northern Mediterranean, while in the southern Mediterranean, ecosystems are still at risk of fragmentation or disappearance due to human pressure from clearing and cultivation, overexploitation of firewood and overgrazing. Drylands have significant biodiversity value, with many of the plants and animals highly adapted to water-limited conditions. They are undergoing



an overall increase in response to climate change and extensive land abandonment. 48% of Mediterranean wetlands were lost between 1970 and 2013, with 36% of wetland-dependent animals in the Mediterranean threatened with extinction. Because of the reduction in river flows, 40% of fish species in Mediterranean rivers are endangered. Projections for the 21st century indicate drier climate and increased human pressure, with negative impacts on terrestrial biodiversity, forest productivity, burned areas, freshwater ecosystems and agrosystems. Future projections indicate that burnt areas can increase across the region by up to 40% in a 1.5°C warming scenario and up to 100% from current levels for 3°C warming at the end of the century. Mediterranean drylands will become drier and their extent is expected to increase across the region. Projections suggest decreased hydrological connectivity, increased concentration of pollutants during droughts, changes in biological communities as a result of harsher environmental

conditions, and a decrease in biological processes such as nutrient uptake, primary production, and decomposition.

Promotion of "climate-wise connectivity" through permeability of the landscape matrix, dispersal corridors and habitat networks are key to facilitating upward the migration of lowland species to mountains in order to adapt to new climate change conditions. Promotion of mixed-species forest stands and silvicultural practices such as thinning, and management of understory can promote the adaptation of Mediterranean forests to climate change. Promotion of the spatial heterogeneity of the landscape matrix can help reduce fire impacts. The preservation of the natural flow variability of Mediterranean rivers and streams and wide riparian areas, along with reductions in water demand are key to the adaptation of freshwater ecosystems to future climate change.

## 4.1 Marine ecosystems

### 4.1.1 Current condition and past trends

#### 4.1.1.1 Observed changes

Despite only covering 0.82% of the ocean surface, the Mediterranean Sea supports a high level of biodiversity, including about 18% of all known marine species (~17,000) (Bianchi and Morri 2000; UNEP/MAP-RAC/SPA 2009; Coll et al. 2010). The Mediterranean Sea is biologically diverse because it is a warm sea at temperate latitudes, and is thus home to both temperate and subtropical species, and has been further diversified by its complex geological history (Bianchi and Morri 2000; Merheb et al. 2016). As a result, the present marine biota of the Mediterranean is composed of species belonging to: (1) temperate Atlantic-Mediterranean species; (2) cosmopolitan species; (3) endemic elements, comprising both paleoendemic (Tethyan origin) and neoendemic species (Pliocenic origin); (4) subtropical Atlantic species (interglacial remnants); (5) boreal Atlantic species (ice-age remnants); (6) Red Sea migrants (especially into the Levantine Basin); (7) eastern Atlantic migrants (especially into the Alboran Sea) (Bianchi and Morri 2000).

In marine ecosystems, specific drivers of environmental change include: i) the increasing tem-

perature and salinity of surface waters (Coma et al. 2009; Conversi et al. 2010; Calvo et al. 2011) and the deep-sea (>400 m) (Béthoux et al. 1990; Rixen et al. 2005; Vargas-Yáñez et al. 2010; Skliris et al. 2014; Schroeder et al. 2016), ii) enhanced thermal stratification (Powley et al. 2016), which can increase eutrophication and O<sub>2</sub> consumption due to increasing dissolved organic carbon (DOC) concentrations in the mixed layer (Ferreira et al. 2011; Santinelli et al. 2013; Ngatia et al. 2019), and iii) decreasing ocean pH fundamentally changing ocean carbonate chemistry (Calvo et al. 2011; The MerMex Group et al. 2011; Flecha et al. 2015; Hassoun et al. 2015, 2019; Merlivat et al. 2018). Detailed information about these drivers, namely temperature and salinity changes, Mediterranean hydrology and ocean acidification can be found in *Sections 2.2.4, 2.2.7.2 and 2.2.9*. Risks and vulnerabilities caused by these drivers are also affected by non-climate related anthropogenic stressors, such as industrialization, urbanization and agriculture, fishing, maritime traffic, harbor activities, tourism (Macías et al. 2014; Thiébault et al. 2016) and floating plastics and other polymers (Fossi et al. 2012, 2018; Suaria et al. 2016). These non-climate drivers are thoroughly described in *Chapters 2 and 3.1 (Section 3.1.2.3)* and can be classified as pollution (*Section 2.3*) and land and sea-use changes (*Section 2.4*).

The interconnected effects of climate change and several non-climate related drivers, covered in *Chapter 2, Section 2.6* affect the way the Mediterranean marine ecosystem functions at all levels, from primary producers to upper trophic-levels (The MerMex Group et al. 2011; Doney et al. 2012; IPCC 2014) (*Fig. 4.1*). Consequences include enhanced mortality of key marine habitat species, e.g., coralligenous outcrops, maërl beds (Paireud et al. 2014; Molina et al. 2016) and the bivalve *Pinna nobilis* (Vázquez-Luis et al. 2017), as well as the increased establishment of new communities and disease outbreaks (Rubio-Portillo et al. 2018; Berzak et al. 2019). Impacts of warming on marine biota not only result from the direct impact of increasing temperature on organism physiology, but also from the effect of warming on other biological (e.g., microbial activity, metabolic rates) and abiotic (e.g., oxygen solubility) components of ecosystem functions (Vaquer-Sunyer and Duarte 2013).

Since the mid-1980s, regime shifts in the Mediterranean Sea have impacted different ecosystem components (e.g., diversity and abundance of zooplankton, abundance of anchovy stocks, frequency of harmful algal blooms, mucilage outbreaks), possibly due to regional effects of climate modes (*Section 2.2.2*), such as a positive state of the North Atlantic Oscillation (NAO) that affects the physical properties of the water column (Conversi et al. 2010; Barausse et al. 2011). The recent study by Fortibuoni et al. (2017), while confirming the existence of some regime shifts, does not support the hypothesis of climatic change as a main driver for these, and rather points to the impact of local pressures, i.e., overexploitation and nutrient loads.

Increasing temperatures are driving the northward spread of warm-water species (Sabatés et al. 2006; Tsikliras 2008; Bianchi et al. 2018), and have contributed to the spread of the non-indigenous Atlantic coral *Oculina patagonia* (Serrano et al. 2013). The recent spread of warm-water species that have entered from Red Sea and Atlantic Ocean into cooler northern areas is leading to the “tropicalization” of Mediterranean fauna (Vergés et al. 2014; Bianchi et al. 2018; Galil et al. 2018). Non-indigenous species are extensively detailed as a driver in *Section 2.5*. Species that need certain temperature ranges cannot migrate further, as the different areas in which they usually live and span are becoming more and more restricted, e.g., the anchovy *Engraulis engrasicolus* (Sabatés et al. 2006). Warming water may also have strong effects on deep Mediterranean areas of the two zones where cold water is formed, as increasing

temperature may slow the potential downwelling and the provision of oxygen both in the Gulf of Lions and in the Adriatic Sea, leaving the cold-water coral communities exposed to a certain degree of hypoxia (Taviani et al. 2016).

In addition to the general warming patterns, periods of extreme temperatures have had large-scale and negative consequences for Mediterranean marine ecosystems (*Sections 2.2.1* and *2.2.2*). A link between positive thermal anomalies and observed invertebrate mass mortalities has been observed in the Mediterranean Sea (Rivetti et al. 2014). Also, unprecedented mass mortality events, which affected at least 25 prominent sessile metazoans, occurred during the summers of 1999, 2003, and 2006 across hundreds of kilometers of coastline in the northwest Mediterranean Sea (Cerrano et al. 2000; Calvo et al. 2011). These events coincided with either short periods (2 to 5 days: 2003, 2006) of high sea temperatures (27°C) or longer periods (30 to 40 days) of less extreme temperatures (24°C: 1999) (Crisci et al. 2011). Impacts of these events on marine organisms have particularly been reported between 0 and 35 m depths, such as gorgonian coral mortality (Coma et al. 2009) or shoot mortality and flowering of seagrasses (Díaz-Almela et al. 2007; Marba and Duarte 2010). A collaborative database for tracking mass mortality events in the Mediterranean Sea has been recently launched to support the analysis of relationships between thermal conditions and/or other environmental drivers (Garrabou et al. 2019), and can be helpful for better detecting changes across the Mediterranean Basin.

In addition, ocean acidification is an emerging human health issue, that also threatens the marine realm (Falkenberg et al. 2020) (*Section 2.2.9*). Studies of the consequences of ocean acidification on marine Mediterranean ecosystems report diverse responses (Martin and Gattuso 2009; Rodolfo-Metalpa et al. 2010; Movilla et al. 2012; Bramanti et al. 2013; Gazeau et al. 2014; Lacoue-Labarthe et al. 2016). Insights have been gained by studying natural CO<sub>2</sub> seeps at Mediterranean sites such as Ischia and Vulcano in Italy, where biodiversity decreases with decreasing pH toward the vents, with a notable decline in calcifiers (Hall-Spencer et al. 2008; Prada et al. 2017). Transplants of corals, mollusks, and bryozoans along the acidification gradients around seeps reveal a low level of vulnerability to CO<sub>2</sub> levels expected over the next 100 years (Rodolfo-Metalpa et al. 2010, 2011). However, periods of high temperature increase vulnerability to ocean acidification, thereby increasing the long-term risk posed to

Mediterranean organisms and ecosystems as temperatures rise (Gazeau et al. 2014; Lacoue-Labarthe et al. 2016). Ocean acidification seems to have a slower but unstoppable effect on several organisms, the increase of temperature being a more immediate stress factor in most species (Lejeusne et al. 2010). A recent overview (Gao et al. 2020) showed that the combination of ocean acidification and warming may affect food webs from different directions; ocean acidification is more likely to follow bottom-up controls (resource driven), while temperature drives top-down controls (consumer driven).

### **Key habitats undergoing change**

Rapid warming of the Mediterranean Sea, in synergy with other climate and non-climate related drivers (see *Chapter 2*), threatens marine biodiversity, and particularly some key ecosystems that have high vulnerability to such pressures, as presented below.

#### **Coralligenous**

The coralligenous is a typical Mediterranean underwater seascape, present on hard bottoms from ~15 to 120 m depths and is mainly produced by the accumulation of calcareous encrusting algae (*Lithophyllum*, *Lithothamnion*, *Mesophyllum* and *Peyssonnelia*) growing in dim light conditions and relatively calm waters (Ballesteros 2006; Boudouresque et al. 2015). These outcrops foster one of the richest assemblages found in the Mediterranean, harboring approximately 10% of Mediterranean marine species (Ros et al. 1985; Boudouresque 2004; Ballesteros 2006; Casas-Güell et al. 2016), most of which are long-lived algae and sessile invertebrates (sponges, corals, bryozoans and tunicates) (Garrabou et al. 2002; Ballesteros 2006). The different habitats that make up these biogenic formations are mainly determined by light exposure, so that some coralligenous habitats can be dominated by calcareous algae and others completely dominated by macroinvertebrates with almost no algae (Gili et al. 2014; Casas-Güell et al. 2016). Red coral, *Corallium rubrum*, is one of the habitat-forming species that plays a key role in the functioning of coralligenous habitats because of its trophic activity, biomass and perennial biogenic structure, like other Mediterranean gorgonian species (Gili et al. 2014; Ponti et al. 2014b, 2016, 2018). Red coral is a slow-growing, long-lived species that grows in dim light habitats (e.g., caves, vertical cliffs and overhangs) between 10 and 200m depths. Despite its essential ecosystemic role, little is known about the geographical distribution of red coral up to 400km offshore the coastline due to its large

bathymetric range and afferent constraints (Casas-Güell et al. 2015, 2016), and the major studies focus on the phytobenthic component (Piazzi et al. 2009, 2012; Boudouresque et al. 2015). Studies at an intermediate scale (tens of km) have been conducted with key species, pinpointing the fact that their distribution may be very heterogeneous depending on the environmental factors (Gori et al. 2012; Coppari et al. 2014, 2016). Due to this lack of baseline data, the structure of coralligenous outcrops is still poorly understood, preventing a proper assessment of its current state of biodiversity and the potential impacts of harvesting, and other disturbances related to global change, on red coral assemblages. A recent study (Mallo et al. 2019) based on historical red coral data from the north western Mediterranean Sea, documented the halt in the *C. rubrum* decrease and the first recovery response due to effective protection measures in some areas.

Coralligenous outcrops are affected by several consequences of global change such as nutrient enrichment, non-indigenous species, increased sedimentation, mechanical impacts, mainly from fishing activities, e.g., mechanical injuries and sediment re-suspension (Cebrián et al. 2012; Piazzi et al. 2012; Gatti et al. 2015), as well as sea warming (e.g., massive mortalities related to temperature anomalies) and the potential effects of ocean acidification (Bramanti et al. 2013; Cerrano et al. 2013; Gili et al. 2014). Recently, potential synergies between these stressors have been hypothesized (*Section 2.6*), especially in shallow areas where heat waves may have a large impact on several organisms (Galli et al. 2017), resulting in a fragmentation of the habitat that can open new space for non-indigenous species (Vezzulli et al. 2013). It has also been demonstrated that a decrease in the abundance of coralligenous habitat-forming species leads to a rapid fragmentation in community structure and a loss of species benefiting from the structural complexity these species provide (Ponti et al. 2014b; di Camillo and Cerrano 2015; Valls et al. 2015).

In addition to marine heat waves (Garrabou et al. 2001, 2009), one of the main past threats for the red coral *Corallium rubrum* has been intensive harvesting (see *Section 2.4*), which has caused an overall shift in population structure, resulting in a decrease in both biomass and colony size (Tsounis et al. 2010; Bramanti et al. 2014; Montero-Serra et al. 2015). Moreover, its Mg-calcite skeleton makes it vulnerable to ocean acidification (Bramanti et al. 2015). Bramanti et al. (2013) experimentally

evaluated the effects of low pH on *C. rubrum* over a 314-day period under two pH levels (8.10 and 7.81). This study concludes that exposure to lower pH conditions negatively affected skeletal growth and spicule morphology (i.e., abnormal shapes).

Mediterranean gorgonian “forests” (e.g., *Paramuricea clavata*, *Eunicella cavolinii*) are threatened by several human activities and are affected by climatic anomalies that have led to mass mortality events in recent decades (Ponti et al. 2014b, 2018; Verdura et al. 2019). Observed mortality events have been linked to temperature-dependent bacterial pathogens (Bally and Garrabou 2007). Also, diverse responses to thermal stress have been shown in gorgonians (Pivotto et al. 2015; Crisci et al. 2017). This may condition the future response of these species to climate change.

The ecological role of these habitats and the possible consequence of their loss are still poorly understood. The experimental study of Ponti et al. (2014b) reports a significant effect of gorgonians (*E. cavolinii*, and *P. clavata*) on the recruitment of epibenthic organisms and their presence mainly limits the growth of erect algae and enhances the abundance of encrusting algae and sessile invertebrates. This effect could be due to microscale modification of hydrodynamics and sediment deposition rate by i) a shading effect that reduces light intensity, ii) intercepting settling propagules, iii) competing for food with filter-feeders and/or iv) competing for space by producing allelochemicals. Although the biological interaction between gorgonians and other species deserves further study, changes to the edaphic conditions caused by gorgonian forests influences the larval settlement and recruitment processes of benthic assemblages (Ponti et al. 2014b, 2018).

In addition to the long-term effects of global change and its consequences on the Mediterranean coralligenous, short-term extreme events may be even more devastating than heat waves. Teixidó et al. (2013) show how an extreme storm event affected the dynamics of benthic coralligenous outcrops in the northwestern Mediterranean Sea using data acquired before (2006–2008) and after the impact (2009–2010) of a major storm. The most exposed and impacted site experienced a major shift immediately after the storm and over the following year. This impact consists of changes in the species richness and diversity of benthic species such as calcareous algae, sponges, anthozoans, bryozoans and tunicates. In this site, benthic species recorded a 22% to 58% loss of cover on average, with those with fragile forms

showing cover losses up to 50 to 100%. Small patches survived after the storm and began to grow slightly during the following year, and the sheltered sites showed no significant changes in all the studied parameters, indicating no variations due to the storm (Teixidó et al. 2013).

### Deep sea ecosystems

Although poorly known, deep seafloor ecosystems provide services that are vitally important to the entire ocean and biosphere, and play a particularly major role in climate change mitigation. For instance, by storing a large amount of anthropogenic CO<sub>2</sub> and by absorbing heat accumulated from the greenhouse effect, the deep sea Mediterranean waters and ecosystems capture large quantities of carbon and, as such, slow down the warming of surface waters and land (Luna et al. 2012; Palmiéri et al. 2015) (Sections 2.2.7 and 2.2.9). Rising atmospheric greenhouse gases are affecting water column oxygenation, temperature, pH and food supply (Section 3.2), with similar impacts on deep-sea ecosystems (Sweetman et al. 2017). As they are deprived of light, deep-sea ecosystems are greatly dependent on surface primary production: “marine snow” (Gambi et al. 2014). Surface water also oxygenates deep-sea environments when they sink to form deep and intermediate water masses. When surface water is warmer, it does not mix well with deep water (Section 2.2.7).

In the Mediterranean, the deep sea covers about 79% of the Mediterranean Basin, including habitats potentially able to deliver multiple ecosystem services and numerous resources of high economic value (Manea et al. 2020). Despite this fact, very few studies address the response of deep-sea ecosystems to ongoing climate change in this sea. In contrast with most oceans where the flux of particulate organic matter to the seafloor is likely to decline significantly in response to climate change (Sweetman et al. 2017), a study from the eastern Mediterranean shows that climate change has caused an immediate accumulation of organic matter on the deep-sea floor in recent decades (Danovaro et al. 2001). This led, together with deep-sea warming, to alteration of carbon and nitrogen cycles and has had negative effects on deep-sea bacteria and benthic fauna (Danovaro et al. 2001, 2004). For instance, the observed salinity and temperature changes in eastern Mediterranean deep and bottom waters from 1987 to 1994 (Roether et al. 1996; Theocharis et al. 2002) led to the uplift of these water masses by several hundred meters, reaching shallower depths (100–150 m; i.e., close to the euphotic zone) under the influence of cyclonic circulation. This resulted

in increased biological production and therefore enhanced flux of organic carbon to the deep sea, thereby significantly and quickly changing the way deep-sea ecosystems function (Psarra et al. 2000; Danovaro et al. 2001). The review of Yasuhara and Danovaro (2016) on temperature impacts on deep-sea Mediterranean biodiversity shows that minor temperature shifts of around 0.1°C or less are sufficient to cause significant changes in biodiversity and the community structure of deep-sea nematode assemblages.

### Planktonic ecosystems


Several studies have addressed the possible impact of climate change on marine phytoplankton diversity and distribution in the Mediterranean Sea, highlighting highly contrasting regional patterns (Duarte et al. 2000; Goffart et al. 2002; Marty et al. 2002; Bosc et al. 2004; Ribera d'Alcalà et al. 2004; Marty and Chiavérini 2010; Herrmann et al. 2014; Oviedo et al. 2015; D'Amario et al. 2017). Some studies from the northwestern Mediterranean have reported a positive trend in phytoplankton biomass in response to the expansion of the summer stratification. This trend was accompanied by an increase in picoplankton and nanoflagellates (i.e., small-sized phytoplankton) and a decline in diatoms, which are responsible for new production (Goffart et al. 2002; Marty et al. 2002; Mena et al. 2019; Ramírez-Romero et al. 2020). However, other studies report that the spring bloom in many Mediterranean regions tends to occur earlier in the year, possibly in relation to earlier water warming and high irradiance, in contrast with the autumn bloom that tends to disappear because of a longer stratification period (Bosc et al. 2004). Bosc et al. (2004) also reveal significant interannual variations in biomass and primary production, not only in the northwestern basin (e.g., the exceptional bloom in spring 1999), but also, and more surprisingly, in the oligotrophic waters of the eastern basin (e.g., the 9% decrease in primary production from 2000 to 2001). In this latter basin, phytoplankton shifts seem to be concurrent with rising winter precipitation and sea surface temperature (Mena et al. 2019) (Section 2.2.4, 2.2.5 and 2.2.7).

In some Mediterranean settings, such as the central Ligurian Sea, increased deep-water convection (as deep as 2,000 m) has been attributed to greater surface salinity causing increased nutrient supply near the surface, and thus more primary production (Marty and Chiavérini 2010). In contrast, in the productive northwestern Mediterranean Sea, deep convection could significantly decrease under the influence of climate change (Herrmann et al. 2014), impacting

pelagic planktonic ecosystem, which are strongly influenced by these hydrodynamics. The weakening of deep convection and surface warming modifies the pelagic planktonic ecosystem and associated carbon cycle indirectly only: the spring bloom occurs one month earlier, and the bottom up control of phytoplankton development and bacteria growth by nitrogen and phosphorus availability strengthens, and the microbial loop intensifies as the small-sized plankton biomass increases (Herrmann et al. 2014). Net carbon fixation and deep export do not change significantly. In the Tyrrhenian Sea, Ribera d'Alcalà et al. (2004) explain the significant changes in the long-term patterns of rare copepod species as a symptom of large-scale meteorological phenomena of the North Atlantic sector.

In the NW Mediterranean Sea, decadal climatic oscillations linked to the NAO forcing of the precipitation regime led to an increase in the upper salinity in the 1980s and in the late 1990s and early 2000s (Chapter 2, Section 2.2.7). In saline years, the annual abundance of zooplankton is higher than otherwise (Fernández de Puellas and Molinero 2007). According to Molinero et al. (2008), large-scale climate forcing has altered the local environment and the pelagic food-web dynamics in the NW Mediterranean Sea through changes in biological interactions, competition and predation. The authors also suggest that warming, the dominance of small phytoplankton and predation pressure by jellyfish negatively affected copepod populations (recruitment, life-history traits and physiological thresholds) in the early 1990s, whereas chaetognaths were surpassed by jellyplankton as the most frequent copepod prey. A more recent study from the same Ligurian time-series updated with ten more years (up to 2003) revealed that the zooplankton, mainly copepods, recovered their initial concentrations after 2000, suggesting a quasi-decadal cycle (Coma et al. 2009). This illustrates the difficulty in identifying long-term changes from decadal oscillation in short time-series in plankton. However, surface salinity appears to be a common physical indicator of changes in the pelagic ecosystem of the NW Mediterranean Sea for jellyfish (Buecher et al. 1997), crustaceans (García-Comas et al. 2011) and phytoplankton (Marty and Chiavérini 2010).

Gallisai et al. (2014) report that aerosol deposition from the Sahara may explain 1 to 10% of seasonally detrended chlorophyll variability in the nutrient-low Mediterranean with main effects in spring over the eastern and central Mediterranean, corresponding to dust events fueling needed nutrients for the



**Figure 4.1 | Summary of interactions between large marine vertebrates and marine litter** (Galgani et al. 2014). Fluxes of litter in the life cycle and intensity of its effects on large marine vertebrates, (a: entanglement; b: ingestion), depending on various factors such as ingestion mechanisms (predation, active or passive filter feeding), development stage (benthic or pelagic phases for sea turtles), behavior and foraging strategy (feeding on the sea floor, in the water column or on the surface, selectivity according to color, shape etc., ecological plasticity in diet and habitat), types of litter (micro/macro litter) and types of fishing gear (nets, hooks and lines). The thicker arrows indicate key processes. Although trophic transfer from one level to another has been demonstrated in vitro for microplastics in plankton, it remains controversial in situ, as most ingested litter is excreted in feces.

planktonic community (Ternon et al. 2011). The areas showing negative effects on chlorophyll from dust deposition are regions under significant influence from European aerosols. Anthropogenic aerosol deposition of nitrate and phosphate largely influence primary production in the northern Mediterranean Sea (Richon et al. 2018a, 2018b) (Section 2.2.3). This response of chlorophyll dynamics to dust deposition is important when knowing that future scenarios predict increased aridity and shallowing of the mixed layer (Gallisai et al. 2014) (Section 2.3.2).

From around the island of Lampedusa (central Mediterranean), the multi-year evolution of biogenic dimethylsulfide (DMS) production in the marine surface layer and the resulting methanesulfonate on the atmosphere are mainly attributed to phytoplankton physiology (Becagli et al. 2013). High phytoplankton productivity can also be the expression of stressed cells, especially during summer when high irradiance and the shallow depth of the upper mixed layer prevails. This therefore leads to higher methanesulfonate concentrations in the atmosphere. These dynamics can be further controlled by the North Atlantic Oscillation, and related oceanic and atmospheric processes (Becagli et al. 2013).

### Large vertebrates

One of the biggest threats to large marine vertebrates is litter debris, such as fishing gear or other large items (Galgani et al. 2014) (Fig. 4.1). Regularly, in the Mediterranean Sea and worldwide, large vertebrates such as sea birds (van Franeker et al. 2011), cetaceans (de Stephanis et al. 2013; Notarbartolo di Sciara 2014) and marine turtles (Lazar and Gračan 2011; Campani et al. 2013; Camedda et al. 2014) accidentally swallow micro and macro-plastic debris that is often found in their digestive tracts. The plastic debris (Section 2.3.2.3) affects the marine biota of the Mediterranean at macro, micro- and nano-levels.


Sperm whales (*Physeter macrocephalus*) in the Mediterranean Sea, which are believed to be fewer than 2,500 mature individuals, are endangered world-wide (Notarbartolo di Sciara 2014). A decline in sperm whales in the Mediterranean has been observed over the last half-century. In addition to ingestion of solid debris, other anthropogenic activities at sea are suspected to have caused the decline of this species and continue to threaten its survival in various ways: bycatch, collisions with vessels, debilitation by chemical pollution, anthropogenic noise, disturbance from irresponsible whale watching and most likely

climate change, and prey depletion (Notarbartolo di Sciara 2014). Regarding specifically ingested debris, ingestion rates are as high as 31% in some marine mammal populations, and sub-lethal effects could result in impacts at the population level (Baulch and Perry 2014). Campani et al. (2013) and Camedda et al. (2014) investigated the interaction between loggerhead sea turtles (*Caretta caretta*) and marine litter in the northern Tyrrhenian Sea and around Sardinia, respectively. In thirty-one *C. caretta* individuals found stranded or accidentally bycaught in northern Tyrrhenian Sea, marine debris, mainly plastics, were present in 71% of specimens (Campani et al. 2013). In Sardinia, only 14% of the 121 monitored turtles had debris in their digestive tracts but plastic was the main physical category (Fossi et al. 2013; Camedda et al. 2014).

Sharks and rays are also seriously threatened by anthropogenic pressures, mainly as a result of overfishing (Dulvy et al. 2014) (Fig. 4.2), as described in Section 2.4.2 in the context of the increasing sea use changes. Some sharks live in narrow climatic ranges (Chin et al. 2010), putting them at risk in a climate change hotspot such as the Mediterranean (Ben Rais Lasram et al. 2010). Microplastic (<5 mm) ingestion has been recorded in 16.8% of the analyzed specimens of the blackmouth catshark *Galeus melastomus* around the Balearic Islands, with higher quantities of filament-type microplastics (Alomar and Deudero 2017). In three striped dolphin populations living in the Pelagos Sanctuary (bordered by western Italy, southern France and northern Sardinia), the highest toxicological stress was from PBT (persistent, bioaccumulative and toxic substances) chemical levels, combined with correlated biomarker responses (Fossi et al. 2013). More on chemical pollution is covered in Section 2.3.

### Changes in biodiversity

To date, changes in Mediterranean marine biodiversity are essentially driven by human activities (Mannino et al. 2017), i.e., pollution (Section 2.3), sea use changes (Section 2.4.2), the introduction of non-indigenous species (Section 2.5), together with climate change (Section 2.2) (Lejeune et al. 2010; Zenetos et al. 2012; Katsanevakis et al. 2013, 2014b). In general, the Mediterranean Sea represents the highest proportion of threatened marine habitats in Europe (32%, 15 habitats) with 21% being listed as vulnerable and 11% as endangered (see review in Mannino et al. 2017). This threat includes several valuable and unique habitats (e.g., seagrasses and coralligenous), supporting an



**Figure 4.2 | The trajectory and spatial pattern of chondrichthyan (cartilaginous fishes that include sharks, skates, rays and chimaeras) fisheries catch landings and fin exports.** (A) The landed catch of chondrichthyans reported to the United Nations Food and Agriculture Organization from 1950 to 2009 up to the peak in 2003 (dark blue) and subsequent decline (orange). (B) The rising contribution of rays to the taxonomically-differentiated global reported landed catch: shark landings (blue), ray landings (dark blue), log ratio [rays/sharks], (orange). Log ratios >0 occur when more rays are landed than sharks. The peak catch of taxonomically-differentiated rays peaks at 289,353 tonnes in 2003. (C) The main shark and ray fishing nations are gray-shaded according to their percentage share of the total average annual chondrichthyan landings reported to the FAO from 1999 to 2009. The relative share of shark and ray fin trade exports to Hong Kong in 2010 are represented by fin size. The taxonomically-differentiated proportion excludes the 'nei' (not elsewhere included) and generic 'sharks, rays, and chimaeras' category (adapted from Dulvy et al. 2014).

extensive repository of biodiversity (Gubbay et al. 2016).

The shallow depth (on average 1,450 m) of the Mediterranean Sea and the relatively fast deep-water turnover in comparison to the open ocean, coupled with a high degree of endemism (about 20% of Mediterranean marine species; Coll et al. [2010])

point to a potential amplification of climate change impacts. These are expected to cause earlier changes in biodiversity in comparison with other seas, thus making this system a model for investigating biodiversity response to direct and indirect effects of temperature changes and other climate-related and non-related drivers (Chapter 2).



Species with low dispersal ability are particularly affected by climate change, which may also lead to local extinctions, greatly contributing to biodiversity loss (Mannino et al. 2017). Any change in biodiversity may affect ecosystem functioning, even in the case of the establishment of a single species and may lead to important consequences both for nature as well as for society. However, the extreme richness of microclimates in the Mediterranean (ranging from climate conditions similar to those of the Northern Sea in the Adriatic to an almost tropical condition in the eastern basin) makes prediction at large spatial scales difficult. Most effects of climate change (or climate anomalies) on marine biodiversity have been so far identified at regional scales (Philippart et al. 2011).


During recent decades, Mediterranean marine communities have shown significant changes in taxa composition and distribution. In the western Mediterranean, climate change is influencing the boundaries of biogeographic regions and thus warm water marine species are extending their ranges and colonizing new regions where they were previously absent (Katsanevakis et al. 2014a). For instance, mucilages have appeared more frequently (associated with a malfunctioning of the microbial loop) in the Adriatic Sea, where it was documented for the first time, and in several regions beyond, in recent decades, concomitantly with a significant increase in sea surface temperature (Danovaro et al. 2009). Mucilage is not closely associated with the presence of eutrophic conditions, as several mucilage outbreaks have been recently observed in oligotrophic seas, such as the Aegean Sea (Danovaro et al. 2009). The Ligurian Sea, one of the coldest areas of the Mediterranean Sea, displays a low number of subtropical species and a higher abundance of cold-temperate water species. However, the recent warming of Ligurian seawater has favored the penetration of warm-water species (e.g., *Thalassoma pavo*), which from 1985 onward, established large and stable populations (Parravicini et al. 2015).

Temperature anomalies, even of short duration, can dramatically change Mediterranean faunal diversity. The largest mass-mortality event recorded in the Mediterranean Sea so far occurred in 1999 along the French and Italian coasts (Cerrano et al. 2000; Perez et al. 2000; Garrabou et al. 2001). That year was characterized by a summer with a positive thermal anomaly that extended the thermocline down to a depth of 40 m (Romano et al. 2000) and resulted in the extensive mortality of 28 epibenthic invertebrate species (Fig. 4.3) (Perez et al. 2000; Rivetti et al. 2014). Among benthic organ-

isms, sponges and gorgonians were most severely affected (Cerrano et al. 2000; Perez et al. 2000; Romano et al. 2000; Garrabou et al. 2001; Rivetti et al. 2014). The shortage of food over several weeks is a common phenomenon in the Mediterranean Sea due to summer water stratification, but very long periods with high temperatures may explain such mass mortalities (Rossi et al. 2017a).

In the eastern Mediterranean, the rise of seawater temperatures may also be partly responsible for the entrance of non-indigenous species (Section 2.5), mostly from the tropical Indo-Pacific (Galil 2000; Por 2009; Zenetos et al. 2012; Rilov 2016). The increased introduction and spread of non-indigenous species may be a supplementary stress factor for native species already weakened by climate variations resulting in the dislocation of indigenous species' niches and possibly cascade effects on the food webs (Rilov 2016; Corrales et al. 2018). Non-indigenous species are a recognized threat to diversity and the abundance of native species as well as a threat to the ecological stability of the infested ecosystems. Despite the overall tendency towards ocean warming, the eastern Mediterranean also experiences occasional climate anomalies, for example between 1992 and 1994, when temperatures dropped by about 0.4°C (Danovaro et al. 2001). This caused a drastic decrease in nematode abundance and overall faunal diversity (e.g., a roughly 50% decrease in nematode diversity, Danovaro et al. 2004). After 1994, when the temperature gradually recovered, biodiversity started to reverse to previous conditions but had not recovered fully in 1998 (Danovaro et al. 2004).

Sea warming may also have effects on the virulence of pathogens, favoring the frequency of epidemiological events, as most pathogens are temperature sensitive (Bally and Garrabou 2007; Vezzulli et al. 2013). Mass mortalities of the gorgonian *Paramunicea clavata*, scleractinian corals, zoanthids, and sponges observed in 1999 in the Ligurian Sea were indeed promoted by a temperature shift, in conjunction with the growth of opportunistic pathogens (including some fungi) (Cerrano et al. 2000). Increased surface temperatures and altered circulation and precipitation regimes have been evoked to explain the increased frequency of bottom water hypoxia or anoxia in coastal areas of the northern Adriatic. These phenomena, often associated with mass mortalities of fish and benthic fauna, alter food webs and might have important cascade effects on biodiversity (Coll et al. 2010). The Adriatic Sea can undergo dramatic change in the lower part of its temperature ranges. In winter 2001, the



**Figure 4.3 | Temperature trends across the Mediterranean Basin.** (Temperature trends at 0–10 m (A), 11–30 m (B), 31–50 m (C) depth layers for the period 1945–2011 in July–November. Linear regressions have been calculated on grids of 1° latitude by 1° longitude and tested for statistical significance at the 90% level. Significant increased/decreased temperature trends are reported as colored cells, non-significant increased/decreased temperature trends are reported as grey areas. Dots show the locations of documented mass mortalities for a depth layer, each color represents a single event. The asterisks in the legend of mass mortalities (MM) events refer to the taxa affected: \* stands for sponges, \*\* for cnidarians, \*\*\* for bryozoans, \*\*\*\* for ascidians, \*\*\*\*\* for bivalves (Rivetti et al. 2014).

Adriatic Sea experienced a period of abnormally low surface temperatures (from 9°C to freezing) that led to mass mortalities of sardines (*Sardinella aurita*) (Guidetti et al. 2002), with resulting alteration of the food webs. The Adriatic Basin is also the site for deep-water formation, as a result of the bora winds associated with decreased temperatures, but recent studies have reported the shift of

this water formation site towards the Aegean Sea by a phenomenon known as eastern-Mediterranean Transient (EMT), related mainly to climatic sea and atmosphere conditions (Hassoun et al. 2015). EMTs change the salinity distribution with surface water freshening linked to enhanced deep-water production and in turn to strengthened Mediterranean thermohaline circulation (Incarbona et al.

2016). This phenomenon can thus affect the marine biodiversity not only in the Adriatic and Ionian Seas but much further, as documented by Ouba et al. (2016), who have correlated the salinity variations and increase in total zooplankton abundance in Lebanese waters to the activation of the Aegean Sea as a major source of dense water formation as part of an “eastern Mediterranean Transient-like” event (see *Section 2.2.7* for more details about Mediterranean circulation changes).

In response to ocean acidification, calcifying organisms (planktonic and benthic) such as corals, foraminifera, coccolithophores and coralline red algae, important contributors of marine calcium carbonate production, may be greatly affected (Langer et al. 2009; Moy et al. 2009; Bramanti et al. 2013; Cerrano et al. 2013; Kroeker et al. 2013). Based on experiments, the impact of ocean acidification on Mediterranean corals was examined and a significant decrease in calcification rates in most tested species was reported (Movilla et al. 2012, 2014). In the latter study, there was a heterogeneous effect of low pH on the skeletal growth rate of the organisms depending on their initial weight, suggesting that those specimens with high calcification rates may be the most susceptible to the negative effects of acidification. Also, a significant effect on benthic foraminiferal communities of low-pH seawaters around the island of Ischia (Italy) has been demonstrated as a result of volcanic gas vents with significant changes in distribution, diversity and nature of the fauna (Dias et al. 2010).

Coccolithophores, which are the primary calcifying phytoplankton group, and especially the most abundant species, *Emiliana huxleyi*, have shown a reduction of calcification at increased CO<sub>2</sub> concentrations for the majority of strains tested in culture experiments (Meyer and Riebesell 2015). Meier et al. (2014) analyzed in situ *E. huxleyi* coccolith weight from the NW Mediterranean Sea in a 12-year sediment trap series, and surface sediment and sediment core samples. Their findings clearly show a continuous decrease in the average coccolith weight of *E. huxleyi* from 1993 to 2005, reaching levels below pre-industrial (Holocene) and industrial (20th century) values recorded in the sedimentary record, as most likely a result of the changes in the surface ocean carbonate system. Also, a drastic decrease in production, species diversity and anomalous calcification in coccolithophores has been shown along a natural pH gradient caused by marine CO<sub>2</sub> seeps off Vulcano Island (Italy) (Ziveri et al. 2014).

To conclude, (1) Mediterranean fauna is highly vulnerable to human activities and climate change; (2) both structural and functional biodiversity of continental margins are significantly affected by very small temperature changes; and (3) the impact of human activities and climate change on marine biodiversity might be non-reversible. Since there are close interactions between deep and shallow systems, the vulnerability of deep-sea ecosystems to climate change might also have important implications on the biodiversity and functioning of continental shelves.

The extent of changes caused by climate and non-climate drivers, the responses of Mediterranean marine biota to these changes and their local-regional consequences are yet to be investigated, as slow but significant transformations that may modify the neritic, pelagic, and benthic zones are still ongoing.

#### 4.1.1.2 Past changes

Understanding the degree to which changes in Mediterranean marine ecosystems point to a directional trend driven by global warming remains a challenge for marine ecology (Bertolino et al. 2017a). Reconstructing the temporal variability of Mediterranean marine ecosystems on time scales longer than a few centuries beyond the instrumental records, crossing relevant climate variations and historical periods, can be critical for interpreting these changes.

Climate forcings of Mediterranean marine ecosystems over the past thousand years have occurred on different time scales (Abrantes et al. 2005; Hennekam et al. 2014; Xoplaki et al. 2018). During the Holocene, rapid warming and cooling events have occurred which can, to some degree, provide analogues for the projected changes for the coming centuries (Blois et al. 2013; Benito-Garzón et al. 2014; Raji et al. 2015). In the Mediterranean, these past climate changes impacted the marine physico-chemical parameters of surface and deep waters (e.g., salinity, temperature, oxygenation, pH) which in turn affected marine ecosystems (Frigola et al. 2008; Schmiedl et al. 2010a; Mojtahid et al. 2015; Bertolino et al. 2017b).

#### **Response of marine ecosystems to past temperature changes**

In the Mediterranean region, the most abundant Holocene temperature proxy data, especially for the Common Era (the last 2,000 years) are alkenone-derived records (Abrantes et al. 2012; Jalali

et al. 2016; Sicre et al. 2016). These studies document natural long-term trends superimposed on a multidecadal variability in response to external (e.g., solar) and internal forcings (e.g., NAO) which might explain some recently observed sea surface temperature trends (Versteegh et al. 2007). These studies also reveal a strong regional component. For example, a high resolution study from the Gulf of Lion shows an overall sea surface temperature cooling trend since the mid-holocene followed by a rapid warming from ~1850 AD onwards that may parallel recent climate change (Jalali et al. 2016). In contrast, south of Sicily and in the eastern Levantine Basin, sea surface temperature records show progressive warming since the early Holocene without a clear signature of the recent anthropogenic change (Castañeda et al. 2010; Luterbacher et al. 2012; Jalali et al. 2017). The planktonic ecosystem in the Siculo-Tunisian Strait responded to this progressive warming of the sea surface temperature by increasing the abundance of warm dinocyst species (*Spiniferites mirabilis* and *Impagidinium aculeatum*) and planktonic foraminifera (*Globorotalia inflata* and *Globigerinoides ruber*) (Rouis-Zargouni et al. 2010).

The Holocene was interrupted by at least four brief cooling events at ~9.2 ka, ~8 ka, ~7 ka and ~2.2 ka cal. BP, which may be correlated to climate events recorded elsewhere, including in Greenland ice cores and in Atlantic Ocean sediments. Investigations on cetacean bones from the Grotta dell'Uzzo in northwestern Sicily (Italy) show that the rapid climate change around 8 ka coincided with increased strandings in the Mediterranean Sea (Mannino et al. 2015). Also, the diversity of sponge species living in coralligenous habitats from the Ionian and Ligurian was strongly affected by Holocene warming episodes with a significant loss of their biodiversity in recent decades (Bertolino et al. 2017b, 2019).

In the eastern Mediterranean, multiproxy records derived from sediments from the southeastern Levantine (Schilman et al. 2001b; Mojtahid et al. 2015) and the Adriatic Sea (Piva et al. 2008) reveal complex paleo-oceanographic changes during the late Holocene, with pronounced anomalies during the Medieval Warm Period (MWP) (ca. AD 1150) and the Little Ice Age (ca. AD 1730). These temperature anomalies were accompanied in the eastern Levantine Basin by a drastic change in planktonic foraminiferal successions indicating periods oscillating between cold and warm surface waters in

opposite phase with the western Mediterranean records (Mojtahid et al. 2015). This east-west contrast in the climate signals has been confirmed by other proxy data (Jalali et al. 2016, 2017).

These findings imply that long-term and short-term climate-driven environmental changes, caused by global warming, will likely impact the entire food chain from planktonic ecosystems to large mammals (e.g., cetaceans) in the near future.

### **Response of marine ecosystems to past changes in stratification and ventilation**

Throughout the Pleistocene, the eastern Mediterranean experienced numerous anoxic events recorded by the cyclical deposition of organic-rich layers or sapropels (Rossignol-Strick et al. 1982; Rohling 1994), the most recent being Sapropel S1 from ~10 to 6 cal ka BP. Maximum insolation due to the Earth's orbital precession minimum significantly intensified the northeast African monsoon, leading to enhanced discharge of fresh and nutrient-rich Nile River water into the eastern Mediterranean (Rossignol-Strick et al. 1982; Emeis et al. 2000). In the Levantine Basin, sea surface salinity during S1 dropped by about 2.0 to 4.0 units compared to present values (Kallel et al. 1997; Myers et al. 1998). This led to severe water column stratification and organic enrichment from the Nile river water. In the Ionian Sea, the correspondence of recent sapropel layers with peaks of the lower photic zone coccolithophore species *Florisphaera profunda* indicated the development of a deep chlorophyll maximum, due to the pycnocline/nutricline shallowing in the lower part of the photic zone (Incarbona et al. 2011). In the SE Levantine Basin, a severe drop in planktonic foraminiferal diversity was recorded in response to the water column stratification and expressed by the near exclusive presence of the euryhaline tropical-subtropical species *Globigerinoides ruber* and the disappearance of deep-dwelling species (Mojtahid et al. 2015).

The combination of higher organic matter remineralization and decreased ventilation resulted in widespread bottom water anoxia (Rohling 1994; Hennekam et al. 2014). In the Southern Aegean and Levantine Seas, there was a gradual increase in deep-water residence times, preceding S1 formation by approximately 1–1.5 kyr. Once oxygen levels fell below a critical threshold, the benthic ecosystems collapsed almost synchronously with the onset of S1 deposition. The recovery of benthic ecosystems during the terminal phase of S1

formation is controlled by subsequently deeper convection and re-ventilation over a period of approximately 1500 years. After the re-ventilation of the various sub-basins during the middle and late Holocene, deep-water renewal was more or less similar to recent rates (Schmiedl et al. 2010b). Several species of deep-water ostracods that are still common in the western Mediterranean became extinct in the eastern Mediterranean Basin at the onset of early Holocene S1 sapropel deposition and the related anoxia (Van Harten 1987). The deep-water ostracode *Bythocypris obtusata* apparently survived the oxygen crisis in the eastern basin itself. This suggests that full oxygen depletion may not have affected the bottom of all deep sub-basins and supports a midwater oxygen-minimum model for these sub-basins (Van Harten 1987; Schmiedl et al. 2010b).

These paleoclimatic findings suggest that eastern Mediterranean pelagic and benthic marine ecosystems are capable of abrupt transitions in response to gradual forcing. This is crucial for the projection of whether an increase in oceanic moisture availability under current and future warming could trigger a sudden intensification of monsoon rainfall further inland from today's core monsoon region (Schewe and Levermann 2017).

### **Response of marine ecosystems to past changes in productivity**

In the western Mediterranean, productivity has shown an overall decreasing trend since the early Holocene with a marked fall in productivity after the 8.2 ky BP dry-cold event (Ciampo 2004; Jiménez-Espejo et al. 2007; Melki et al. 2009). Superimposed on this long-term pattern, some studies show millennial-centennial time scale variability linked with weakening and strengthening of upwelling conditions that have been simultaneous to changes in Western Mediterranean Deep Water (WMDW) formation in the Gulf of Lions and by extent to the NAO over the past 7.7 ka (Ausín et al. 2015). These changes were accompanied by re-organization in coccolithophore assemblages showing in particular, several high-amplitude oscillations of the productivity indicator species *F. profunda* (Ausín et al. 2015).

In the eastern Mediterranean, several proxy data support overall increased productivity during Sapropel S1 in a high-nutrient stratified environment (Gennari et al. 2009; Castañeda et al. 2010; Mojtahid et al. 2015). This period is characterized by the highest accumulation rates of planktonic foraminifera together with the productivity indica-

tor coccolithophore species *F. profunda* (Incarbona et al. 2011; Mojtahid et al. 2015). After Sapropel 1, a progressive decrease in surface water productivity was recorded and surface and deep-sea ecosystems were driven by short-term changes in food quantity and quality as well as in seasonality, all of which are linked to millennial-scale changes in river runoff and associated nutrient input (Kuhnt et al. 2008; Schmiedl et al. 2010b). Particularly, the last 2.9 ka encompassed a succession of three ecosystem states characterized by nutrient-limiting surface waters from 2.9 to 1.1 ka, and during the Little Ice Age, and by nutrient-rich waters from 1.1 to 0.54 ka (Medieval Climate anomaly) (Mojtahid et al. 2015). These conditions were linked to periods of low and high Nile River runoff respectively, in line with arid and humid climate conditions in the Levant and Nile headwaters.

These findings imply that surface productivity in the overall oligotrophic Mediterranean Sea responds rapidly to short and long-term changes in nutrient input, either via rivers, winds or upwelling activity, modifying the benthic-pelagic ecosystems by extending into the entire food chain (Marino and Ziveri 2013), ultimately increasing eutrophication.

### **Response of marine ecosystems to past changes in pH**

Holocene reconstructions of paleo-pH have yet to be undertaken in the Mediterranean. There is a promising raw data record of planktonic foraminiferal (*Neogloboquadrina incompta*)  $\delta^{11}\text{B}$  and B/Ca. These geochemical proxies can be used for paleo-pH and show an overall decreasing trend in both sub-basins of the Mediterranean Sea during the last deglacial episode of glacial-interglacial  $\text{CO}_2$  rise (Grelaud et al. 2012; Marino and Ziveri 2013). The response of marine calcifiers to this trend can be estimated via planktonic foraminifera shell weight that shows overall decreasing planktonic calcification in response to this variability. In addition to this general trend, periods of changing seawater carbonate chemistry can be observed, which could be linked to low/high primary production activity such as the anomaly observed during Sapropel 1 period, which can be linked to enhanced mineralization of organic matter.

These first studies show that Mediterranean marine calcifiers responded to past changes in surface seawater carbonate chemistry conditions. The extent to which this affects marine ecosystems needs to be analyzed in the context of the current acidification in the Mediterranean's surface and deep seawaters.

### 4.1.2 Projected vulnerabilities and risks

#### 4.1.2.1 Projected impacts and risks

As already discussed in *Section 2.2.4.1*, annual mean temperatures in the Mediterranean are now 1.5°C above late 19th century levels with magnitudes that vary locally depending on the period of analysis, the region and the type of dataset. The diurnal temperature range has also changed in some parts of the Mediterranean (*Section 2.2.4.1*). In absolute terms, the warmest parts are the southern and eastern Mediterranean and the major impact in these parts is the immigration of Indo-Pacific species (around a thousand species), which has accelerated in recent years, mainly for thermophilic species, due to rapid warming conditions (more than 50% of Mediterranean non-indigenous species are in the eastern Mediterranean) (Azzurro et al. 2011; Marbà et al. 2015; Kletou et al. 2016; Bariche et al. 2017). All Mediterranean waters, even the deepest, are affected by ocean acidification driven by Mediterranean Sea uptake of atmospheric CO<sub>2</sub> (Flecha et al. 2015; Hassoun et al. 2015; Palmiéri et al. 2015; Ingrassio et al. 2017) (*Section 6.11*). In addition, the effects of climate change are amplified by other major non-climate-related anthropogenic forcings, as the Mediterranean has one of the most populated coastlines with a long human history of exploitation of marine resources (with presently one of the world's most intense coastal and maritime tourism areas), habitat degradation and plastic pollution (Cózar et al. 2015; Compa et al. 2019). More information about sea use changes and pollution are covered in *Chapter 2*.

The combination of various ongoing climate change processes (e.g., sea warming, ocean acidification, and sea level rise; *Section 2.6*) has caused detectable effects on marine organisms at individual, population, and ecosystem scales (*Fig. 4.4*). Future risks of sea level rise, marine heat waves, and ocean acidification are also highlighted in *Sections 6.9, 6.10 and 6.11* respectively. In fact, sponges, gorgonians, bryozoans, molluscs, and seagrasses are all affected by these drivers (Cerrano et al. 2006; Garrabou et al. 2009; Bensoussan et al. 2010; Marbà and Duarte 2010), but primary producers, mainly calcifiers such as coccolithophores, are among the most vulnerable organisms (Meier et al. 2014). The impacts are expected to affect endemic and iconic ecosystems including major reorganizations of the biota distribution, species loss, marine productivity, increases in non-indigenous species, and potential species extinction (Malcolm et al. 2006; Ramírez et al. 2018; Gao et al. 2020).

### Projected impacts on microbes

Sea warming may have effects on the virulence of pathogens (viruses, parasites, etc.), favoring the frequency of epidemiological events, as most pathogens are temperature sensitive (Vezzulli et al. 2013) [see *Section 4.1.1* and *Section 2.3.4* in *Chapter 2* for more information about biological pollutants], as observed for *Vibrio shiloi*, responsible for the whitening of the coral *Oculina patagonica* in the eastern Mediterranean (Kushmaro et al. 1998). This warming is also responsible for the expansion of harmful and/or toxic microalgae, mainly dinobionts such as *Ostreopsis ovata*, which produces palytoxins, a serious public health hazard (Accoroni et al. 2016; Vila et al. 2016). Temperature anomalies also seem to negatively affect the chemical defenses of marine organisms (Thomas et al. 2007), allowing pathogens to act undisturbed. Given the predicted rise in temperatures over the coming decades, a better understanding of the factors and mechanisms that affect the disease process will be of critical importance in predicting future threats to temperate gorgonian communities (Bally and Garrabou 2007), and other affected species in the Mediterranean Sea.

In deep waters, a recent study has shown that deep-sea benthic Archaea can be more sensitive to temperature shifts than their bacterial counterparts. Changes in deep-water temperature may thus alter the relative importance of Archaea in benthic ecosystem processes (Danovaro et al. 2016). With rising deep-water temperatures, the predicted positive response of prokaryotic metabolism to temperature increases may accelerate oxygen depletion in deep Mediterranean waters, with domino effects on carbon cycling and biogeochemical processes across the entire deep basin (Luna et al. 2012). Along canyon-cut margins (e.g., the western Mediterranean), warming may additionally reduce density-driven domino effects, leading to decreased organic matter transport to the seafloor (Canals et al. 2006), though this very process is also likely to reduce physical disturbance on the seafloor and therefore affect deep-sea ecosystems.

### Projected impacts on primary and secondary production

Climate change affects the functioning of the biological components of ecosystems, from the basis of the food webs (plankton) to the higher trophic levels (e.g., predator fish). Phytoplankton constitutes the autotrophic primary producers in the pelagic food chains in marine waters and their annu-

al cycle is affected by many physical features that in turn control nutrient levels. These include large horizontal gradients in temperature (Izrael 1991). Due to their rapid turnover and fast responses to environmental changes, plankton is considered a suitable proxy to highlight either environmental changes circumscribed in space and/or time or wider climatic variations. Warming, for example, is responsible for the expansion of harmful and/or toxic microalgae, mainly the dinobionts such as *Ostreopsis ovata*, which produces palytoxins, a serious public health hazard (Accoroni et al. 2016; Vila et al. 2016). A new study in the Eastern Mediterranean has shown the occurrence of important concentrations of biotoxins (domoic acid, gymnodimines and spirolides) in various marine organisms sampled from the Lebanese shores (Hassoun et al. 2021). These concentrations were correlated with the abundance of biotoxins' producers such as *Pseudo-nitzschia*, *Prorocentrum*, *Alexandrium*, and other species that could be occurring more frequently due to climate change (Hassoun et al. 2021).

Moreover, phytoplankton species responsible for bloom at late winter and at the beginning of spring (like *Skeletonema costatum*, *Nitzschia* spp., *Leptocylindrus danicus* and *L. minimus* and others) could start earlier, because features of temperate marine planktonic ecosystems are not only sensitive to annual variations in weather, but also any trends that might result from greenhouse warming or other factors that affect the climate system and both the density and timing of spring blooms will be altered in some regions (Townsend et al. 1994).

The taxonomic compositions of phyto- and zooplankton may change under the influence of changes in ocean structure (Kawasaki 1991; Berlin et al. 2012; Howes et al. 2015) (Section 2.2.7). A thermophilic phytoplankton species could proliferate especially in some enriched areas and could be ichthyotoxic or even toxic for humans (Abboud-Abi Saab 2008, 2009; Accoroni et al. 2016; Abboud-Abi Saab and Hassoun 2017). Some examples can explain such variations. In the Mediterranean Sea, phytoplankton biomass abundance and sea surface thermal stratification show a strong inverse relationship at seasonal and sub-basin scales. At inter-annual and sub-basin scales, a gradual decline of the phytoplankton biomass across the entire central Mediterranean occurs with a delay of one year (Volpe et al. 2012). In the Adriatic Sea, during the past decade, the community structure and seasonality of phytoplankton have changed significantly. The phytoplankton annual cycle has become more irregular with sudden diatom

blooms, reflecting the variability of meteorological events in recent years (Totti et al. 2019).

Only a few regional studies have investigated the sensitivity of the oligotrophic Mediterranean Sea to future climate change. The first investigations considered only the changes in circulation. For instance, a regional model of the northwestern Mediterranean domain found that the effect of local stratification due to climate change would have no drastic effect on the pelagic ecosystem (Herrmann et al. 2014). However, one study investigated the overall effects of a moderate climate change scenario (A1B SRES) on Mediterranean biological productivity and plankton communities and found an overall decrease in phytoplankton biomass in response to the stratification simulated in their dynamic climate change scenario (Lazzari et al. 2014). A simulation was carried out for an increase in integrated primary productivity across the eastern Mediterranean Basin as a result of changes in density (decreased stratification) (Macías et al. 2015). However, conclusions from these studies remain limited by the fact that they are based on non-transient simulations and present-day nutrient inputs.

A new study has investigated the influence of both changes in circulation and biogeochemical forcings (rivers and input at Gibraltar) (Richon et al. 2019). It suggests that climate change and nutrient inputs from river sources and fluxes through the Strait of Gibraltar have contrasting influences on Mediterranean Sea productivity (Section 2.3.3). Increased future stratification globally reduces surface productivity in the eastern basin, but the biogeochemistry in the western basin is strongly controlled by nutrient input across the Strait of Gibraltar, while the eastern basin is more sensitive to vertical mixing and river inputs. In the near future, longer water stratification and warmer conditions may be essential clues to understanding local stress and species mortalities, especially because of the changes in primary productivity and lack of enough resources to face starvation periods (Rossi et al. 2017a). Once some species are eradicated, other species, pre-adapted to the new conditions, can replace locally extinct species, thus hampering ecosystem resilience to pre-impact conditions.

Unfortunately, the future evolution of nutrient atmospheric deposition has not been considered in modelling studies so far, despite the important role of nutrients for marine primary production (Richon et al. 2018b, 2018a). Using aerosol sampling and microcosm experiments performed during the TransMed BOUM cruise (June–July 2008), Ternon

et al. (2011) showed that primary production significantly increased at all tested stations after aerosol addition collected on-board and after Saharan dust analog addition, indicating that both additions relieved on-going co-limitations, whereas a decline in the future of primary production is predicted and associated with trophic amplification toward phytoplankton and zooplankton (Richon et al. 2019).

Abrupt community shifts are expected in plankton communities coinciding with climate changes that alter local thermal regimes, which in turn interact with the thermal niche of species to trigger long-term and sometimes abrupt shifts at the community level (Beaugrand et al. 2019). Planktonic calcifying organisms such as foraminifera, pteropods and coccolithophores are expected to be particularly affected by ocean acidification and climate change. A recent study in the Mediterranean found foraminifera to be highly susceptible to temperature-induced surface water stratification and food availability. In correlation with these results, rapid warming increased surface ocean stratification impacting food availability and changes in trophic conditions could be the causes of reduced foraminiferal abundance, diversity, and species-specific changes in planktonic foraminiferal calcification (Mallo et al. 2017). Coccolithophores, an abundant unicellular calcifying phytoplankton, are known to have a haplo-diploid life cycle with environmental affinities. This dimorphic life cycle might provide the ability to adapt to the "tropicalization" of Mediterranean environments under climate change, in conditions characterized by surface water with a relatively high calcite saturation state, high temperature, stratification and nutrient limitation (D'Amario et al. 2017).

### **Projected impacts on macrobenthic and pelagic species**

Sea warming, ocean acidification, sea level rise and changes in circulation patterns will likely change Mediterranean benthic and pelagic ecosystems, as shown in Fig. 4.4, where potential impacts related to climate change are presented (Rossi et al. 2019). Repercussions will be different depending on the region and on the dominance of certain benthic organisms, species interactions, metabolic constraints, dispersion capability, and the presence of alien species that may take advantage of the new physical, chemical, and biological conditions in the future oceans. Changes in trophic relationships will likely change biodiversity, both in the water column and in the different benthic communities. The nursery effect (i.e., sheltering and feeding grounds for juveniles and larvae) may


be compromised if massive mortalities or deoxygenation affect the three-dimensional live structures (i.e., Posidonia meadows, gorgonian forests, pluriannual macroalgae, cold water corals, etc.). Also, changes in river runoff due to different rain frequency/quantities will also have effects on coastal nutrient and pollutant inputs. Nutrient and pollutant equilibria will also have direct or indirect effects on pelagic and benthic communities (Rossi et al. 2019).

### **Corals**

Environmental change-driven modifications to the environment of gorgonian forests influence the larval settlement and recruitment processes of the benthic assemblages (Ponti et al. 2014b). In experiments, red coral (*Corallium rubrum*) showed a 59% decrease in its calcification rate at lowered pH based on experimental studies (Bramanti et al. 2013). Disturbances such as harvesting pressure could act in synergy with ocean acidification bringing local populations to extinction. The changes in the likelihood of occurrence obtained by differences between present conditions and future scenarios show that the projected extent of potential mortality zones is higher than in the current climate for red coral and that coralligenous formations along the Mediterranean sites are less likely, mostly due to acidification increase. However, the adverse impact is localized to certain regions: an increase in likelihood is also reported for the presence of coralligenous in the North Aegean and northern Adriatic Seas (MEDSEA 2015; Gómez-Gras et al. 2019).

The effects of in situ exposure to different pH levels (7.4–8.1) and temperatures (15.5–25.6°C) on mortality and net calcification rates have been assessed for Mediterranean scleractinian corals transplanted near a volcanic CO<sub>2</sub> vent off Panarea Island (Prada et al. 2017). Results of this *in situ* study show a synergistic adverse effect on mortality rates for all corals (up to 60%), suggesting that high seawater temperatures may have increased their metabolic rates which, in conjunction with decreasing pH, could have led to rapid deterioration of cellular processes and performance. The same study (Prada et al. 2017) suggests that symbiotic corals may be more tolerant to increasing warming and acidifying conditions compared to asymbiotic corals. Laboratory cultures of coralline algae under conditions of elevated temperature and pCO<sub>2</sub> revealed effects on photosynthesis, growth and calcification. Crustose coralline algae (*Neogoniolithon brassica-florida*) sensitivity to ocean acidification examined in CO<sub>2</sub> seeps confirmed that calcifying algae are likely to be threatened





**Figure 4.4 | Different drivers potentially affecting marine pelagics and benthos in the Mediterranean Sea** (original diagram in Rossi et al. 2019, artwork by A. Gennari).

by ocean acidification, especially species living near their thermal limit. Further in situ and laboratory experiments indicate that *N. brassica-florida* may not be able to contribute to reef accretion under the levels of seawater warming and ocean acidification projected by the end of this century (Fine et al. 2017).

### Seagrass

Warming can induce declines in abundance through increased shoot mortality in Mediterranean *Posidonia oceanica* meadows. Younger life stages (i.e., seedlings) of *P. oceanica* may be particularly vulnerable to climate change. Insights into acidification effects on seagrasses have come from CO<sub>2</sub> vent surveys showing consistent loss of crustose coralline algal epiphytes on seagrass leaves, and greater seagrass density close to seeps with a lower pH (Hendriks et al. 2017). Lower epiphyte loads can have positive impacts for seagrass as it reduced shading and nutrient uptake by the epiphytes.

In the case of seagrass such as *P. oceanica*, projections show a negative impact due to the

effects of global warming over the next century and to significant climate change challenges posed to an endemic system that is already suffering losses from anthropogenic impacts (Jordà et al. 2012). The trajectory of *P. oceanica* meadows under the warming expected in the western Mediterranean and warming seems to likely lead to the functional extinction of *P. oceanica* meadows by the middle of this century (year 2049 ± 10) even under a relatively mild greenhouse-gas emissions scenario (Jordà et al. 2012). Similarly, the distribution of two seagrass species under different scenarios was forecasted (Chefaoui et al. 2018), and the results found that, in the worst-case scenario (RCP 8.5 scenario), *P. oceanica* might lose 75% of suitable habitat by 2050 and is at risk of functional extinction by 2100, whereas *Cymodocea nodosa* would lose around 46.5% of suitable habitat by 2050. The same study (Chefaoui et al. 2018) also predicts that erosion of present genetic diversity and vicariant processes can happen, as all Mediterranean genetic regions could decrease considerably in extension in future warming scenarios. The functional extinction of *P. oceanica* would have important ecological

impacts and may also lead to the release of the massive carbon stocks these ecosystems stored over millennia.

### Mussels

The Mediterranean mussel *Mytilus galloprovincialis* is particularly sensitive to warming. A significant decrease in growth (total weight, shell length, shell weight) has been found in warmer conditions as well as clear dissolutions of the shells exposed to low pH conditions (-0.3 compared to ambient). Mussels exposed to low pH showed a clear loss in the organic layer covering the shell in summer, explaining the clear dissolution signal measured on these mussels (Gazeau et al. 2014). Non-climate drivers, such as chemical pollution (Section 2.3.3), are exacerbating the pressures on this important commercial mussel (*M. galloprovincialis*).

### Jellyfish

The sensitivity and specificity of the Mediterranean Sea to climate change and other human-related stressors have been unequivocally documented in recent years by a significant increase in the number and frequency of jellyfish outbreaks (Coll et al. 2010; Canepa et al. 2014). Ocean warming and acidification may favor the dissemination of the non-indigenous *Cassiopea andromeda* that seems to benefit from the changed conditions (Fuentes et al. 2018). Also, seawater temperature, together with the quantity and quality of available food resources, are known as major drivers of gonadal outputs (Harland et al. 1992; Ben-David-Zaslow and Benayahu 1999). Some studies showed that elevated temperature by itself or in combination with high feeding frequency (due to raised zooplankton prey abundance) increased the budding rate and bud size in *Aurelia* polyps populations worldwide (Hočvar et al. 2018). Thus, more food and warmer waters may be the key to understanding proliferation of jellyfish in general and non-indigenous tropical species in particular. In a sea highly impacted by the alteration of the trophic chains due to overfishing, seawater warming favors the successful dispersion and growth of jellyfish.

### Winners and losers

The western basin is acidifying faster than the eastern basin (Goyet et al. 2016). A first tipping point has already been reached, since anthropogenic CO<sub>2</sub> is already over 82 μmol kg<sup>-1</sup> in many Mediterranean areas (Hassoun et al. 2015). The exact timing of the tipping points (Section 2.2.9.2) will strongly depend on the policies controlling human activities, which will impact both global

warming and the anthropogenic CO<sub>2</sub> increase both in the atmosphere and into the ocean. The results of these projections raise concerns about how marine organisms will respond in the context of each scenario, after reaching every tipping point.

Mean warming, acidification and associated non-climatic stressors will have varying impact across the Mediterranean marine ecosystems, and result in both potential winners and losers. Impacts of ocean acidification and warming may extend to several Mediterranean marine and coastal ecosystem services, food provision, recreational activities, carbon absorption, climate regulation, coastal protection, and ultimately affecting human health (Falkenberg et al. 2020). Marine areas with economic activities directly depending on marine resources may face serious impacts on employment and benefits in sectors like aquaculture, open sea fisheries and tourism, which is relevant to many Mediterranean countries. Tourism may be affected by acidification and warming through the impact of degraded marine ecosystems (loss of iconic coralligenous species, such as gorgonians - soft coral (Bramanti et al. 2013) from diving experiences and through jellyfish outbreaks). Sensitivity of shell-forming species such as bivalve mollusks to changes in temperature and acidity presents a threat to the aquaculture sector (Rodrigues et al. 2015).

A reduction in primary production linked to an increase in sea surface temperature could have negative impacts on fisheries catches and could exacerbate current overfishing trends (Cheung et al. 2010). Projections of biomass and fisheries catches across the Mediterranean under the high emission scenario RCP8.5 suggest a 5 and 22% increase in total fish and macroinvertebrate biomass, and a 0.3 and 7% in fisheries catches by 2021–2050 and 2071–2100, respectively, combined with changes in primary and secondary production (Moullec et al. 2019). Winners were mainly small pelagic species, thermophilic and/or exotic species, of smaller size and of low trophic levels. Loser species are generally large-sized, some of which are of great commercial interest. The bulk of the increase is expected in the southeastern part of the basin whereas significant decreases are most likely in the western Mediterranean Sea. Read more about projections for marine food resources and fisheries in Sections 2.4.2.3 and 3.2.2.2.

Since temperature seems to be a main environmental parameter driving the cnidarian community composition, abundance and spatial distribution patterns in the Mediterranean Sea, jellyfish

are considered a possible group of winners under warming (Guerrero et al. 2018). The structure and phenology of the Mediterranean hydrozoan community displayed significant changes in species composition, bathymetric distribution, and reproductive timing over the last decades. When the Scyphozoa group is considered, *Pelagia noctiluca* (among the most abundant jellyfish in the Mediterranean Sea and Eastern Atlantic waters) has increasingly frequent massive outbreaks associated with warmer winters (Milisenda et al. 2018). Swarms of the Portuguese Man-of-War (*Physalia physalis*), in summer 2010 were the result of an unusual combination of meteorological and oceanographic conditions during the previous winter and not a permanent invasion favored by climate changes (Prieto et al. 2015). However, many studies have attributed the increase in *Pelagia noctiluca* outbreaks to the alteration of the trophic structure of ecosystems due to overfishing and/or eutrophication on the one hand, and by sea warming and changes in surface hydrography on the other (Licandro et al. 2010; Canepa et al. 2014). Water temperature affects sexual reproduction through changes in the energy storage and gonad development cycles and it is still expected that the species composition and biogeography of jellyfish communities will change under global warming.

#### 4.1.2.2 Vulnerabilities

##### Climate-related vulnerabilities

It is expected that the ocean's primary production will, in general, be reduced with environmental change. As a result, production zones may be redistributed and the natural habitat of commercially valuable species of fish may change (Izrael 1991). On the other hand, climate change can also lead to changes in the composition of the bottom of marine food webs. The rise in water temperature has already increased jellyfish population outbreaks in the Mediterranean Sea (Section 4.1.2.1), such as *Pelagia noctiluca*, a planktonic predator of fish larvae and of their zooplankton prey. The outbreaks of this species, along with other jellyfish species, may become more frequent in the Mediterranean Basin in the future and extend over a longer period of the year than previously, causing changes to the pelagic food web and thereby reducing fishery production (Licandro et al. 2010). Rising seawater temperatures might also trigger the increased spread of pathogens throughout the Mediterranean in the future, affecting both marine organisms, and human health (Danovaro et al. 2009) [see Section 5.2.3 on heat-related impacts].

Temperature has a major direct impact on the physiology, growth, reproduction, recruitment and behavior of marine organisms such as fish. Warming associated with climate change already affect the Mediterranean ecosystem for some benthic and pelagic species (Marbà et al. 2015). Warming combined with a decline in oxygen and resource availability reduces fish body size, with the average maximum body weight of fish expected to shrink by 4% to 49% from 2000 to 2050 (Cheung et al. 2013). Also, fish tend to adapt to local environmental temperatures. Therefore, among the most perceptible large-scale consequences of climate change is the shift in spatial distribution range of marine organisms, which will make some Mediterranean sub-basins more vulnerable to drivers than the others. Seawater warming will induce a loss of climatically suitable habitats for various organisms, causing distribution shifts, as well as species extinction. The diversity of fish assemblages is predicted to be severely affected due to their loss of suitable climatic niches. Demersal species will suffer regional impacts associated with the expected changes in primary production, thermohaline circulation, and the severity of winter weather (Section 2.2.2). Warming and the expected increase in Atlantic water entering into the Mediterranean will likely affect migrations and spawning behavior in large pelagic fish (Barange et al. 2018). In recent decades, several mass mortality events of invertebrates have occurred in the Mediterranean which have been linked to the documented rise in seawater temperatures (Rivetti et al. 2014).

Projections for the global warming scenario (SRES A2) for the potential future distribution of 75 Mediterranean endemic fish species have shown that by 2041-2060, 31 species are projected to extend their geographic range, whereas the geographic range of 44 species is projected to reduce (Ben Rais Lasram et al. 2010). Also, 25% of the Mediterranean continental shelf is predicted to undergo an overall change to endemic assemblages by the end of the 21st century, where the survival of 25 species is threatened and six species would become extinct (for example, starry sturgeon *Acipenser stellatus* and European sturgeon *Huso huso*). For "narrow" endemic species found strictly in the Mediterranean Sea that do not reach the neighboring Atlantic Ocean and Black Sea, their extinction would be irreversible. By the middle of the 21st century, the coldest areas of the Mediterranean Sea, namely the Adriatic Sea and the Gulf of Lion, would act as a refuge for cold-water species, but by the end of the century, those areas are projected to become a "cul-de-sac" that would drive those species towards extinction (Ben Rais Lasram et al. 2010). Another study concludes that 54 species

will have lost their climatically suitable habitat at the end of the century and that species richness will decrease across 70.4% of the continental shelf area (Albouy et al. 2013). Information about the trends of fisheries in the Mediterranean is detailed in *Sections 2.4.2.1 and 3.2.1.2*.

### **Anthropogenic vulnerabilities**

Ship collisions and harmful fishing practices are among the non-climate drivers exacerbating the vulnerability of marine cetaceans in the changing Mediterranean. The previous analysis of ship collision records for the relatively isolated population of fin whales in the Mediterranean Sea from 1972 to 2001, indicated that the fatal collision rate increased from 1 to 1.7 whales per year during this period mainly in the Pelagos Sanctuary (the largest marine protected area created for marine mammals in the Mediterranean; see *Section 4.1.1*) due to high levels of traffic and whale concentrations (Panigada et al. 2006). Studies by Pennino et al. (2016, 2017) assessed the risk exposure for high intensity vessel traffic areas for the three most abundant cetacean species (*Stenella coeruleoalba*, *Tursiops truncatus* and *Balaenoptera physalus*) in the southern area of the Pelagos Sanctuary. They modeled both the occurrence of three cetacean species and marine traffic intensity, and identified two main hotspots of high intensity marine traffic in the area, which partially overlap with the area where the studied species are present. International shipping, although considered as an environment-friendly form of transportation, directly and indirectly impacts cetaceans in many ways, particularly in the Mediterranean Sea, one of the world's busiest waterways (Bray et al. 2016; Coomber et al. 2016). More recent data about maritime traffic in relation to cetaceans, investigated through direct observations (July 2013–June 2015) and along three fixed transects in western Mediterranean areas, showed seasonal maritime traffic intensity with the highest vessel abundance impacts on cetaceans in most offshore sub-areas in the spring and summer, especially for the species *B. physalus* and *S. coeruleoalba* (Campana et al. 2017).

### **4.1.3 Adaptation**

#### **4.1.3.1 Long-term monitoring and adaptation strategies**

Temperature significantly affects eukaryotic phytoplankton metabolism, increasing the demand for nitrogen with consequences for the marine carbon cycle due to shifts towards N-limitation

(Toseland et al. 2013). Experiments reveal that some taxa of marine phytoplankton may adapt to ocean acidification, and there are also strong indications from studies of variation and structure in natural populations that selection on standing genetic variation is likely (Collins et al. 2014).

To better evaluate the adaptation strategies of plankton communities to the diverse climate and non-climate related drivers, it is necessary to assess the vulnerabilities of the pelagic ecosystem, including both plankton and nekton communities, to the impacts of climate change, including temperature and pH variations. Taking into consideration the diversity of plankton communities' responses to climate change and other stressors in every Mediterranean sub-basin (Crise et al. 1999; Psarra et al. 2005; Vadrucchi et al. 2008; Calvo et al. 2011; Marić et al. 2012; Ouba et al. 2016; Danovaro et al. 2017; Benedetti et al. 2018), wider monitoring coverage is needed to improve our knowledge about the different adaptation processes that characterize and best suit each geographical zone. Since each Mediterranean Basin represents a unique set of interrelated physical, biological and human components and processes, the extent and nature of impacts in one basin will differ between sub-basins as well.

Also, an assessment of the implications of accelerated temperature increase in the Mediterranean Sea and the identification of the types of problems that a marine area will face is crucial in order to anticipate the need for action (Lacoue-Labarthe et al. 2016). In their review about the biodiversity and ecosystem functioning, Danovaro and Pusceddu (2007) have recommended enhanced strategies for protecting the Mediterranean Sea, such as monitoring environmental quality, grey and black lists of chemicals, utilizing the best available technologies once they have been tested for their eco-sustainability, applying precautionary principles (e.g., reducing pollution emissions), monitoring biodiversity and long-term temporal changes in community structure. The suggestion has been made to assess not only the apparent changes, but also the potential biodiversity (cyst banks) while paying particular attention to species replacement in relation to functional biodiversity (Danovaro and Pusceddu 2007).

#### **4.1.3.2 The role of Marine Protected Areas (MPAs) for adaptation**

Marine Protected Areas cannot halt climate change and impacts such as ocean acidification,

but they are an important tool for enhancing the resilience and adaptive capacity of ecosystems. A topical example in this sense are Mediterranean coralligenous reefs, critical for life-supporting and ecological functions and providing a natural capital like biomass production, erosion control, historical, economic and intrinsic value. Marine protected areas provide refuge and replenishment zones for this heavily exploited and vulnerable species. Moreover, they protect, aid recovery and are home to a large number of species (Rodríguez-Rodríguez et al. 2015; Pascual et al. 2016). Few MPA designs account for official MPA boundaries regardless of boundary overlaps and their ecological implications (Gabrié et al. 2012), whereas others account for MPA boundary overlaps and thus foster a more ecologically meaningful, functional spatial approach through “protected polygons” (Foster et al. 2014).

The effectiveness of MPAs can be improved if they form part of a system of protected areas geared towards ensuring ecological representativeness and creating networks. Nevertheless, subdividing an area into zones with varying intensities of use (zoning), ranging from total protection (marine reserves where extractive use is prohibited) to areas serving primarily to uphold sustainable and/or traditional use of marine resources, and areas that are closed to fishing activities, is increasingly recognized as a useful instrument for sustainable, ecosystem-based fisheries management, particularly artisanal fisheries (Pascual et al. 2016). MPA ecological effectiveness, defined as species self-replenishment and colonization through dispersal, depends, among other variables, on MPA design factors such as size, shape, spacing and location (Shanks et al. 2003; OSPAR Commission 2007; Roberts et al. 2010; Sciberras et al. 2013). Different categories of MPAs often sit side by side with core areas under strict protection and peripheral zones with fewer restrictions relating to use.

An example of these MPAs is the Mediterranean Marine Protected Area of Medes Islands, in the northwestern Mediterranean Sea. This area was recently the subject of an economic assessment, using monetary valuation, of changes in the quality of highly biodiverse coralligenous systems (Rodrigues et al. 2015). The Adriatic Sea, for example, has the largest number of MPAs and also the smallest, least spaced and least compact designated and functional MPAs. The design pattern in this ecoregion seems to have randomly followed a design approach of “several small” MPAs (Rodríguez-Rodríguez et al. 2015). The

establishment of a transboundary Large Marine Protected Area (LMPA) and Fisheries Restricted Area (FRA) can reverse ecological and socio-economic losses in the Adriatic, one of the most exploited areas of the Mediterranean as shown in a study that presents current opportunities and expected benefits of LMPAs (Bastari et al. 2016). This demonstrates that the establishment of MPAs should take into consideration the structural and functional links between key organisms within the ecosystem and between ecosystems to guarantee a sustainable adaptation strategy.

The majority of MPAs are located along the basin’s northern shores, highlighting the lack of MPAs in the south and east coasts (Abdulla et al. 2008). Coll et al. (2012) studied the interaction between marine biodiversity and threats (including climate change) across the Mediterranean and assessed their spatial overlap with current marine protected areas. They identified areas of conservation concern where future protection activities should be targeted through spatial prioritization. Spatial prioritization in conservation is commonly employed to direct limited resources to where actions are most urgently needed and most likely to produce effective conservation outcomes. Resilience is increasing through building MPA networks and setting priorities at the regional level. Examples for the conservation of three key Mediterranean habitats, i.e., seagrass *Posidonia oceanica* meadows, coralligenous formations, and marine caves, were determined through a systematic planning approach (Giakoumi et al. 2013).

#### 4.1.3.3 Management of fisheries and adaptation

Fisheries is one of the main sectors related to resource-based growth activities in the Mediterranean area (Section 5.1.1.3). Sustainable development of fishing activities and the management of their impacts require better constructive collaboration between scientists, industry and government agencies. The analysis of all available stock assessment and effort data for the most important commercial species and fleets in the Mediterranean Sea since 2003, demonstrated a significant decline for red mullet and giant red shrimp stocks (Cardinale et al. 2017). This latter study concluded that the European Common Fisheries Policies have failed to achieve the maximum sustainable yield before 2015 for the Mediterranean Sea and will face large difficulties to reach maximum sustainable yield and Marine Strategy Framework Directive targets before 2020 under the current

management system, due to many factors, such as the ineffectiveness of the putative effort reductions to control fishing mortalities, the continuous failure to follow scientific advice, and the existence of ineffective national management plans as a primary management measure (Cardinale et al. 2017).

The establishment and implementation of management plans that could efficiently help fisheries to adapt relies on knowing each species properties and demographic features. For example, the local and isolated groups of the European anchovy (*Engraulis encrasicolus*) may have unique demographic properties and should be managed separately since they may react independently to exploitation (Jemaa et al. 2015a). The same conclusion has been highlighted for pelagic species (such as the European sardine, *Sardina pilchardus*) with high gene flow to ensure sustainable fishery benefits and efficient conservation as they also may have unique demographic properties and responses to exploitation (Jemaa et al. 2015b).

The current knowledge on Mediterranean fisheries and ecosystems is limited. In fact, the effect of poorly regulated fisheries, in combination with ongoing climate forcing and the rapid expansion of non-indigenous species, are rapidly changing the structure and functioning of ecosystems with unpredictable effects on the goods and services provided (Colloca et al. 2017). Although this would call for urgent conservation actions, the management system implemented in the region appears too slow and probably inadequate to protect biodiversity and secure fisheries resources for future generations. This is why some studies are encouraging the adoption of other management approaches such as the establishment of a transboundary Large Marine Protected Area (LMPA), specifically a no-trawl area LMPA or Fisheries Restricted Area (FRA), which is a promising and feasible approach for reversing ecological and socio-economic losses in some Mediterranean sub-basins such as the Adriatic (Bastari et al. 2016), as mentioned earlier in Section 4.1.3.2.

Maintaining ecosystem services (through efficient fisheries management, sustainable and ecofriendly aquaculture industry (Section 3.2) is crucial for the food security, economic growth and well-being of neighboring populations (Section 5.1.1.3). Developing practical management actions that take into consideration the uniqueness of each species and their responses towards different drivers is crucial

to increasing their resilience and plasticity in the context of climate change.

#### 4.1.3.4 Adaptation strategies for ocean warming and ocean acidification in the Mediterranean Sea

Studies on adaptation to climate change in the Mediterranean Sea are still very limited. Actions considered are mostly supply-side oriented, aimed at restoring or protecting the production of marine goods and services harmed by ocean warming and acidification for example (Ziveri et al. 2017). The demand-side dimension can ultimately produce economic consequences of the same or greater magnitude than adaptation through supply-side strategies and actions.

The Mediterranean Sea is a marine biodiversity hotspot (Coll et al. 2010) and ecosystems with high biodiversity and/or redundancy of functional groups (for example, several species fill the role of algal grazers) tend to be more resilient and recover more quickly following disturbance. This implies that biodiversity preservation and improvement are logical methods for sustaining ecosystems responding to rapid environmental stressors. In practice this means, on the one hand, exploiting the acclimation potential of many calcifying species of the Mediterranean and, on the other hand, protecting other species. The recommendation of a drastic reduction of local drivers is strictly connected with mitigation and adaptation strategies, since reducing local stressors – such as land-based pollution, coastal development and overharvesting – is the most common strategy for improving or maintaining ecosystem resilience (read more on economic vulnerabilities/risks and the adaptation measures in Sections 5.1, 6.10.2 and 6.11.2).

In the Mediterranean, commercial fisheries are economically important on a regional and local scale or for some specific communities and user groups. Although there is still limited knowledge on the combined direct impacts of ocean acidification and warming on fish, there are indications of the physiological and behavioral effects of CO<sub>2</sub> on fish (Nilsson et al. 2012; Milazzo et al. 2016). However, the two phenomena could indirectly impact fisheries affecting phytoplankton community structures at the bottom of the food web (Nagelkerken et al. 2016). In general, when fishing activity is more "sustainable", it tends also to be more resilient to negative shocks. All the measures working in this direction are thus

also suitable for mitigating the adverse economic effects of ocean acidification.

Aquaculture is a key economic sector of fisheries affected by ocean acidification and warming. In the Mediterranean, detrimental effects on bivalve mollusk species might arise from the associated increase in sea surface temperature, ocean acidification and possible synergies with other non-climate drivers (Gazeau et al. 2014). A study suggests that the increase in frequency and duration of summer heatwaves are perceived as the highest threat, having been observed in a majority of the studied production sites in past years, with effects on seed (spat), adult mortality and byssus attachment (Rodrigues et al. 2015). Ocean acidification knowledge transfer and monitoring programs are essential for the development of appropriate strategies to counteract the effects of these phenomena, which are still poorly known by stakeholders. Adaptation in this sector tends to be particularly expensive since it usually requires costly investment in new machinery or in modifying existing machinery so as to reduce negative environmental impacts. A recent study emphasizes that the cost of management and mitigation strategies and actions will be dependent upon the socio-economic context. Specifically, costs will likely be greater for socio-economically disadvantaged populations, exacerbating the current inequitable distribution of environmental and human health challenges (Falkenberg et al. 2020).

Some species are proven to improve the resilience of their habitat to various drivers. For example, *Paramuricea clavata* forests may enhance bioconstruction processes and increase the resistance and resilience of benthic assemblages in Mediterranean coralligenous habitats (Ponti et al. 2014b, 2018). The lack of available food, rising temperature and decreasing pH trends will be essential to understanding future population dynamics. Bioengineering as a possible adaptation strategy includes techniques to mitigate chemical effects of increased atmospheric CO<sub>2</sub> concentrations on the oceans. These chemical changes may have a variety of important biological consequences, including some potentially negative impacts, which are controversial and surely require further consideration. These ideas have never been tested *in situ* (Ziveri et al. 2017).

In conclusion, any kind of action that improves marine ecosystem health, resilience or biodiversity could delay and reduce the adverse effects of climate drivers. This includes the implementation

of more sustainable fishing practices as well as reducing pollution from agricultural activity, sustainable tourism and developing more effective waste management. Marine protected areas can potentially have an insurance role if they are placed in locations not particularly vulnerable to ocean acidification and climate change. However, the detrimental effects of these global phenomena on certain habitats in vulnerable regions do not make MPAs easily effective in improving species resilience to environmental change when considering long-term strategies. Developing specific adaptation options, for example new practices in aquaculture or improving marine and coastal protection against storm surges and coastal erosion, could be effective but particularly costly (see Sections 3.3.2.2 and 3.3.2.3 for more information about the use of marine energy as a renewable energy resource and the vulnerability of coastal energy systems to climate extremes).

Adaptation strategies must have medium- to long-term effectiveness. They thus require careful and anticipatory planning to enjoy their benefits reasonably soon, and especially to enable them to tackle problems while they are still manageable. Overall, adaptation strategies are a necessary response to ongoing and expected Mediterranean environmental changes. However, the necessary strategy for reducing climate change impacts needs effective mitigation policies and actions to be implemented.

#### 4.1.3.5 Regional observation networks as a tool for adaptation

Another aspect that might improve the effectiveness of managed adaptation strategies is the establishment of active regional and local observation networks. Local observation programs and regional networks that include scientists from different Mediterranean countries/sub-basins create a solid platform for peers to collaborate in monitoring climate change drivers and impacts, enhance data sharing policies and accessibility, and improve capacity-building among the members of their scientific community.

Furthermore, long-term active hubs would definitely help in deriving more robust findings about the different environmental trends in the Mediterranean, which will provide more comprehensive and conclusive results for decision makers. Within the Global Ocean Acidification-Observing Network (GOA-ON), a regional Mediterranean hub has been recently established, called the Ocean Acid-

ification Mediterranean Hub<sup>31</sup> (OA Med-Hub). This hub could be an important platform for providing robust ocean acidification-related results for the scientific community, the general public and decisions-makers, which would help to create relevant future adaptation actions in Mediterranean countries.

Moreover, these regional networks could be an effective tool for improving public awareness and enhancing capacity-building among scientists who are not able or do not have the appropriate tools/equipment to monitor specific phenomenon. Thus, such hubs could unify the methodologies and tools adopted to measure and monitor short-

and long-term climate change trends, and their effects on local and regional marine resources and ecosystems. They could also help obtain funds for laboratories that do not have the capacities to properly survey climate change drivers and impacts. For example, GOA-ON published recently an implementation strategy document<sup>32</sup> to provide guidance that will harmonize sampling and analysis procedures, to compare results and trends. Creating similar networks that could target other phenomena, such as deoxygenation, warming, etc., and good communication between these hubs will be crucial for developing suitable and holistic key messages that could be provided to policymakers.

## 4.2 Coastal ecosystems

### 4.2.1 Current condition and past trends

#### 4.2.1.1 Observed changes

The coastal zone refers to the area in which the interaction between marine systems and the land dominate ecological and resource systems. These rather complex systems consist of coexisting biotic and abiotic components which interact with human communities and their socio-economic activities (UNEP/MAP/PAP 2008). In sensitivity studies concerning expected sea level rise, the term "Low-Elevation Coastal Zone" (LECZ) has been used for the specific area up to an elevation of 10 m (Vafeidis et al. 2011). Ecosystems in this zone are referred to hereafter as "coastal ecosystems". The natural coastal systems include distinct coastal features and ecosystems such as rocky coasts, coral reefs, beaches, barriers and sand dunes, estuaries, lagoons, and generally the land located at the lower end of drainage basins, where stream and river systems meet the sea and are mixed by currents and tides (i.e., deltas, river mouths, wetlands) (Convertino et al. 2013).

To evaluate the risks that could affect Mediterranean coastal systems, their natural habitats, or particular events that could occur on its shores, many indexes have been developed, such as the Environmental Sensitivity Indices (ESI) developed for the evaluation of oil spill risk in Mediterranean

coasts (Gugliermetti et al. 2007), the participatory multicriteria analysis (MCA) for a multidimensional assessment of coastal erosion risks (Roca et al. 2008), and the coastal dune vulnerability index for Mediterranean ecosystems (Ciccarelli et al. 2017). Also, a Coastal Risk Index (CRI-Med) has been developed to assess coastal vulnerabilities associated with the physical and socio-economic impacts of climate change in all Mediterranean coastal zones (Satta et al. 2017; *Fig. 4.5*). By applying the CRI-Med on 21 Mediterranean countries, coastal risk hotspots are found to be predominantly located in the South-Eastern Mediterranean region. Countries with the highest percentage of extremely high-risk values are Syria (30.5%), Lebanon (22.1%), Egypt (20.7%), and Palestine (13.7%). Coastal hotspots are designated to support the prioritization of policies and resources for adaptation and Integrated Coastal Zone Management (ICZM).


#### **Natural Mediterranean habitats under severe degradation**

Climate change affects marine biodiversity, especially in coastal habitats (Gatti et al. 2015; Bertolino et al. 2016, 2017a; Betti et al. 2017; Longobardi et al. 2017). Mediterranean shores have unique and highly diversified landscapes, which harbor a high level of diversity in fauna and flora (*Section 2.4*). Directly or indirectly, sea level rise (*Section 2.2.8*), global warming (*Section 2.2.4*) and changes in

<sup>31</sup> [http://goa-on.org/regional\\_hubs/mediterranean/about/introduction.php](http://goa-on.org/regional_hubs/mediterranean/about/introduction.php)

<sup>32</sup> [http://goa-on.org/documents/general/GOA-ON\\_Implementation\\_Strategy.pdf](http://goa-on.org/documents/general/GOA-ON_Implementation_Strategy.pdf)





**Figure 4.5 | Coastal Risk Index (CRI-MED) map of the Mediterranean**, indicating five levels of vulnerability (Satta et al. 2017).

rainfall patterns (*Section 2.2.5*) would greatly modify coastal ecosystems and rivers with significant impacts on the way they function. For instance, in response to relative sea level rise, coastal wetlands and river mouths would be affected, while reduced precipitation and prolonged droughts will reduce the water discharge of Mediterranean rivers and catchments (Merheb et al. 2016). Dynamic coasts are likely to retreat or disappear because of the effects of erosion due to the accelerated rise in sea level. With the accelerated rise in sea level, the least mobile species will likely be the most severely impacted in contrast to mobile organisms such as fish and water-birds which are more able to adapt.

Coastal ecosystems function in a way that maintains the preservation of energy and matter transfer and plays an important role in global cycles (such as carbon and nitrogen global cycles) between the continental and marine realms. Furthermore, in addition of hosting a wide diversity of wild faunal and floral species, coastal ecosystems are also often used as aquaculture platforms (i.e., fish, shellfish cultures, etc.), and the pressures on them may have significant impacts on their uses.

#### **Sandy beaches and sand dunes**

Escalating pressures caused by the combined effects of population growth, demographic shifts, economic development and global climate change

pose unprecedented threats to sandy beach ecosystems across the world (Velegrakis et al. 2016; Vousdoukas et al. 2016). From the sub- to the supra-littoral, sandy habitats are important in preventing coastal erosion and flooding, but their value may be enhanced by the many biological processes that complement or even increase their role in coastal defense. For example, in addition to their role in nourishing other sandy systems, shallow, sub-tidal sands also support seagrass beds, a habitat increasingly recognized as important for coastal protection due to their ability to stabilize and accumulate sediment, and attenuate and dissipate waves (Christianen et al. 2013; Ondiviela et al. 2014).

In addition to direct anthropogenic impacts (e.g., urban and agricultural development, *Sections 2.3 and 2.4*), climate change is predicted to have dramatic, widespread and long-lasting consequences on sandy coastal ecosystems. For instance, sea-level rise increases the phenomenon of "coastal squeeze", but increased storm intensity and frequency are likely to be the major challenges faced (Feagin et al. 2005; Harley et al. 2006; Velegrakis et al. 2016; Vousdoukas et al. 2016).

As ecosystems, sandy beaches play varying important roles, and interact closely with coastal dunes both physically and biologically. The typical

coastal dune system is composed mainly of three components: the submerged beach, the emerged beach and the dune (Fig. 4.6). For thousands of years, human activities have been impacting the sandy beaches and coastal sand dunes of the Mediterranean Basin through agriculture, husbandry and the deliberate use of fire (Lavorel et al. 1998; Cori 1999; Falcucci et al. 2007). In recent decades, tourism has caused important damages as the main driver of coastal urbanization, the increase of summer visitors, and the introduction of non-indigenous or exotic species (Tzatzanis et al. 2003). The pedestrian and motorized pathways all over dunes lead to vegetation destruction and therefore enhanced weathering and erosion. Waste and non-indigenous species introduction are also destruction factors among many other drivers highlighted in Sections 2.3 and 2.5.

The impact of human pressure on landscape patterns and plant species richness in Mediterranean coastal dunes was assessed and a general simplification was observed in the natural dune spatial pattern with a decline in plant richness where human pressure is important (Malavasi et al. 2018). Assessing the conservation status of coastal dune systems in Tuscany (Italy), Ciccarelli (2014) showed that the general spatial pattern of vegetation there was close to the natural zonation (Acosta et al. 2007; Forey et al. 2008; Miller et al. 2009; Fenu et al. 2013), with a variable sequence of coastal dune plant communities, ranging from the disappearance of the foredune habitats to the presence of the complete sequence. Vegetation of the driftline disappeared in this study's transects

where erosion was significant, while embryonic shifting dunes with *Elymus farctus* were well represented in the area. However, embryonic dune habitat showed a decrease in coverage in transects belonging to coastal tracts affected by erosion. Although vegetation of the driftline and embryonic shifting dunes have few taxa with low coverage, they represent an important element that must be considered in conservation programs (Ciccarelli 2014).

**Rocky coasts**

Rocky intertidal shores play an important role for marine habitat functioning (Sugden et al. 2009). They are a destination for leisure, exploration and relaxation (Hall et al. 2002; Sugden et al. 2009), and are subject to extractive activities for food, bait or ornamental purposes (Murray et al. 1999; Hall et al. 2002). Hence, the benefits provided by the Mediterranean intertidal area, and the associated economic assets, are relevant in the perspective of integrated coastal zone management and under the current climate change scenario. Rocky shores are biologically rich and diverse habitats (Benedetti-Cecchi et al. 2003; Ceccherelli et al. 2005; Schembri et al. 2005; Mangialajo et al. 2008a). Their diversity is multiplied from the local scale up to the basin-wide level by the interplay of biotic and abiotic factors such as interaction among species, exposure, microtopography, island or mainland location and latitude. The presence of ecosystem engineering species also plays a crucial role here, where their disappearance is typically associated with radical changes in the structure of the assemblages. The genus *Cystoseira*, for example,




Figure 4.6 | Cross-section of a sandy dune system<sup>33</sup>

<sup>33</sup> [http://www.marinespecies.org/traits/wiki/File:Sand\\_dunes.JPG](http://www.marinespecies.org/traits/wiki/File:Sand_dunes.JPG)

includes low-shore canopy algae responsible for habitat formation for sessile invertebrates requiring high humidity and low light levels, such as sponges and tunicates (Benedetti-Cecchi et al. 2001; Ceccherelli et al. 2005; Mangialajo et al. 2008b). The loss of *Cystoseira* canopies results in disruptions to habitat complexity and species diversity patterns, and in the development of algal turfs associated with low-abundance invertebrate assemblages (Benedetti-Cecchi et al. 2001; Mangialajo et al. 2008b).

*Mytilus galloprovincialis* is an edible mussel, widespread in intertidal and shallow subtidal areas throughout the Mediterranean Sea, where it is also aquacultured. The loss or reduction of natural *M. galloprovincialis* populations (Sarà et al. 2011) and of another autochthonous element of bivalve fauna, *Mytilaster minimus* (Sarà and de Pirro 2011), has been shown to produce changes in the associated assemblages, enhanced when the contemporary reduction in canopy-forming algae occurs (Maggi et al. 2009). Although not an endangered species, its presence and abundance might be altered by the establishment of non-indigenous species like the mussel *Brachidontes pharaonis* (Sarà et al. 2008).

Mediterranean vermetid reefs are found where the temperature of surface coastal waters is no lower than 14°C (in winter) (Chemello and Silenzi 2011). A typical vermetid reef is the outcome of complex synergistic building activity by the vermetid mollusc *Dendropoma (Novastoa) petraeum* and the encrusting red algae *Neogoniolithon brassica-florida* (Chemello and Silenzi 2011). Other species, such as the red algae *Lithophyllum* spp., support the process of bioconstruction. Vermetid reefs develop in the lower mesolittoral and upper infralittoral, on rocky coasts only, and precipitate an aragonite shell (Silenzi et al. 2004; Sisma-Ventura et al. 2009). In addition to temperature and the type of substrate, the hydrodynamism of shallow water layers influences the distribution and size of these structures on a small scale, because vermetid platforms are rare along sheltered coasts (Chemello and Silenzi 2011). As for ocean acidification (Section 2.2.9), although *D. petraeum* were able to reproduce and brood at high levels of CO<sub>2</sub>, recruitment success was found to be adversely affected (Milazzo et al. 2014).

Experimental work on the Mediterranean subtidal red alga *Peyssonnelia squamaria* shows that this species may benefit from ocean acidification, as its own nitrogen metabolism will be regulated (Yildiz 2018). Among the engineering species, although

vermetids are resilient to near-future pCO<sub>2</sub> levels, it is likely that their reefs will not be able to withstand levels of acidification predicted for the end of this century, and the associated community will change as a result (Section 2.2.9).

Mediterranean subtidal rocky ecosystems have not been well studied. An initial current baseline and gradient of ecosystem structure was established for nearshore subtidal rocky reefs on a Mediterranean scale, at 8 to 12 m water depths (Sala et al. 2012). This baseline study showed remarkable variation in the structure of rocky reef ecosystems and suggested that the healthiest shallow rocky reef ecosystems in the Mediterranean have both large fish and algal biomass. Protection level and primary production were the only variables significantly correlated to community biomass structure. Fish biomass was significantly larger in well-enforced no-take marine reserves, but there were no significant differences between multi-use marine protected areas (which allow some fishing) and open access areas at the regional scale (Sala et al. 2012).

Overall, intertidal systems are poorly represented in the socio-economic literature, and there appears to be low awareness of their value among stakeholders. Subsequently, conservation efforts for intertidal communities are minimal.

### Coastal wetlands

Mediterranean coastal wetlands include a wide variety of natural habitats such as river deltas, coastal lagoons and salt marshes, intertidal wetlands, and coastal aquifers. Global warming and direct anthropogenic impacts, such as water extraction (more on land and sea use changes and practices in Sections 2.4 and 3.1.5.2), largely affect water budgets in Mediterranean wetlands, thereby increasing wetland salinities and isolation, and decreasing water depths, water quality and hydroperiods (duration of the flooding period) (Mediterranean Wetlands Observatory 2018). These wetland features are key elements that structure waterbird communities for instance (Ramírez et al. 2018). However, the ultimate and net consequences of these dynamic conditions on key species assemblages are largely unknown, although recent work indicates that waterbird communities, for example, tend to adapt to climate conditions (Gaget et al. 2018).

### Seagrass meadows

Seagrass meadows in the Mediterranean Sea cover 1.35 to 5 million hectares, between 5 and 17% of the worldwide seagrass habitat. The rate

of loss of seagrass is above 1.2% each year and up to 5% in the Mediterranean (Marba and Duarte 2010). The Mediterranean Sea is home to four seagrass genera (*Cymodocea*, *Halophila*, *Posidonia* and *Zostera*) encompassing four indigenous species (*C. nodosa*, *P. oceanica*, *Z. marina* and *Z. noltii*) and one non-indigenous species (*H. stipulacea*) (Sghaier et al. 2011). The largest Mediterranean sublittoral area occupied by seagrasses is dominated by the endemic *P. oceanica* (neptune seagrass). It clonally reproduces in meadows and can spread up to 15 km while being hundreds to thousands of years old. In certain areas its populations are significantly declining due to a combination of human-induced factors (Vohník et al. 2016). *Posidonia oceanica* is the most threatened seagrass habitat: almost half of the surveyed sites in the Mediterranean have suffered net density losses of over 20% in 10 years. These *P. oceanica* losses are directly linked to human activities: eutrophication from nutrient pollution, alteration of coastal sediment balance and physical disturbances from trawling, anchoring of yachts, dredging, and other activities highlighted in Sections 2.3 and 2.4. Other non-human impacts include rising sea temperature (Section 2.2.8) and non-indigenous species (Section 2.5) (Claudet and Fraschetti 2010; Crooks et al. 2011).

The non-indigenous variety of *Caulerpa racemosa*, currently spreading in the Mediterranean Sea, was first discovered in the early 1990s near Tripoli Harbour in Libya (Nizamuddin 1991). It spreads rapidly, and is now found off the coasts of multiple countries around the Mediterranean Sea (Lebanon, Tunisia, Libya, Egypt, Cyprus, Turkey, Greece, Malta, Croatia, Italy, France and Spain), and has reached the Canary Islands in the Atlantic Ocean (Verlaque et al. 2004; Bitar et al. 2017). Recent work has shown that this non-indigenous variety is *C. racemosa* var. *cylindracea*, which was introduced from southwestern Australia (Verlaque et al. 2003). Long-range dispersal of the algae seems to be a result of human activities (e.g., dispersal by anchors, fishing). *C. racemosa* can inhabit a wide range of subtidal substrata (sand, mud, rocks, dead mat of seagrass, from 0 to 50 m depths), and has the potential to expand its range around the entire coastline of the Mediterranean Sea. *C. racemosa* modifies the density and diversity of benthic communities (Capiomont et al. 2005).

Early studies on spatial distribution and expansion of *H. stipulacea* had been focused on the northern Mediterranean - the spreading of *H. stipulacea* along the southern and southeastern Mediterranean coasts was only noticed later (Lipkin 1975;


Bitar et al. 2000, 2017; Abboud-Abi Saab et al. 2003; Lakkis and Novel-Lakkis 2007; Verlaque et al. 2015).

### Coastal lagoons and deltas

The small range of tides associated with low-speed currents has encouraged the establishment of lagoon or endogenic systems around much of the Mediterranean. Lagoons and deltas provide support for rich biodiversity, including vital habitats for bivalves, crustaceans, fish and birds. They provide a physical refuge from predation and are used as nursery and feeding areas for some endangered species (Franco et al. 2006; Le Pape et al. 2013; Escalas et al. 2015; Isnard et al. 2015). There are more than 100 coastal lagoons in the Mediterranean, holding an important ecological role, and also providing essential goods and services for humans (read how the effects of climate change on coastal ecosystems could affect livelihood, culture and human rights in Section 5.3). Over 621 macrophyte species and 199 fish species are present in Atlantic-Mediterranean lagoons (Pérez-Ruzafa et al. 2011).

Due to their location between land and open sea, coastal lagoons are subject to strong anthropogenic pressures in parallel with climate change. Habitat destruction, pollution, water withdrawal, overexploitation and non-indigenous species are the main causes of their degradation (Newton et al. 2018). These pressures are responsible for major ecosystem alterations such as eutrophication, bacterial contamination, algal blooms (toxic, harmful or not), anoxia and fish killings. Furthermore, additional problems arise from coastal erosion, subsidence and effects related to extreme meteorological events, typical for the Mediterranean (Aliaume et al. 2007).

In the Mediterranean, the largest coastal wetlands are found in delta areas like that of the Po (Italy), Nile (Egypt), Rhône (France) and Ebro (Spain) rivers (Section 3.1.1.3). Like most wetlands, deltas are diverse and ecologically important ecosystems (UNEP-MAP-RAC/SPA 2010). Deltas absorb runoff from both floods (from rivers) and storms (from lakes or the ocean). Deltas also filter water as it slowly makes its way through the delta's tributary network. This can reduce the impact of pollution flowing from upstream. Plants such as lilies and hibiscus grow in deltas, as well as herbs such as wort, which are used in traditional medicines. Many animals are indigenous to the shallow, shifting waters of a delta. Fish, crustaceans such as oysters, birds, insects are also part of a delta's ecosystem (UNEP-MAP-RAC/SPA 2010; Medi-



**Figure 4.7 | Typical salt marsh zonation** (modified from Bertness et al. 2002). Species along the tidal elevation gradient are adapted to the inundation frequency, including extreme flooding and storm events. MLW: Mean Low Water; MHWN: Mean High Water of Neap tides; MHWS: Mean High Water Spring tide; HAT: Highest Astronomical Tide.

terranean Wetlands Observatory 2012). In deltaic areas and low-lying coastal plains climate change, particularly sea-level rise, is already considered as an important issue (Nicholls and Hoozemans 1996), that significantly decreases the return periods of maximum water levels due to storm surges (Section 2.2.8 and Box 3.1.1). This has been discussed by Sánchez-Arcilla et al. (2008) for the Ebro delta where other phenomena are affecting deltaic behavior such as increases in inundation/flooding, coastal erosion, salinity intrusion, and changes in wave climate (wave height, direction, and storminess).

### Salt marshes

Salt marshes cover low-energy, intertidal shorelines worldwide and are among the most abundant and productive coastal ecosystems. Salt-marsh ecosystems provide a wide array of benefits to coastal populations, including shoreline protection, fishery support, water quality improvement, wildlife habitat provision, and carbon sequestration (Hansen and Reiss 2015). These are specialized habitats, characterized by high primary productivity and species diversity, which support a wide variety of native flora and fauna, and also constitute important areas for wintering aquatic birds (Simas et al. 2001). Salt marshes protect lowlands from marine flooding by damping storms and waves and by slowing flows pushing inland (Allen 2000).

In the Mediterranean Basin, coastal salt marshes (Fig. 4.7) include various Mediterranean plant communities of the classes *Juncetea maritimi* and *Salicornietea fruticosae* which are under the influence of saline seawater. The vegetation is dominated by perennial and shrubby halophytes growing on the extreme upper shores of low sedimentary coasts, sheltered from the mechanical action of waves. Their habitat is especially diverse in the Iberian Peninsula and in southern Italy (Sicily, Apulia, Calabria) (Molina et al. 2003; Cutini et al. 2010; Sciandrello and Tomaselli 2014). In these parts, the habitat forms a mosaic of tall rushes mixed with shrubby and other herbaceous species, often with succulent stems and/or leaves, forming halophytic shrublands and thickets. In soils with brackish water, beds of reed and other tall helophytes grow. The habitat further includes Mediterranean halo-psammophile meadows (*Plantaginion crassifoliae*), humid halophilous moors with the shrubby stratum dominated by *Artemisia coerulescens*, halo-nitrophilous shrubby seablite thickets of *Suaeda vera* rarely inundated, shrub communities of *Limonium* spp., and communities in the Dalmatian coastal region, in somewhat drier habitats with less salt, which are not directly affected by waves and tides. On intertidal muds, cord grasses (*Spartinion maritimae*) may grow, but these are relatively rare in the Mediterranean (Molina et al. 2003; Cutini et al. 2010; Sciandrello and Tomaselli

2014). Saltmarshes include a wide diversity not only of vegetation (Ayyad and El-Ghareeh 1982), but also of plankton, crustaceans and fish species. The zooplankton structure and dynamics in permanent and temporary Mediterranean salt marshes were investigated, and the results found that although temporary and permanent basins show no differences in the most abundant zooplankton species, they differ in their zooplankton diversity, temporal pattern and size structure (Brucet et al. 2005). The same study also found that the presence of a stable fish population in the permanent salt marshes may explain the high values of zooplankton diversity and the low densities of large zooplankton (Brucet et al. 2005).

In an assessment of the threat level of non-indigenous species on salt marshes of the southeastern Iberian Peninsula, the status of its habitats has been categorized as "particularly critical" as many of them were destroyed in the past, due to their transformation into cropland or by desiccation for fear of malaria (Al Hassan et al. 2016). In the Valencia region (Spain), the coastline supports virtually all farming, much of the region's industrial activity, and shelters large population centers. This, along with huge pressure from tourism (i.e., pressures related to water consumption, *Section 3.1.2.3*), have highly impacted salt marshes. These ecosystems house a specific flora of halophytes and their high specialization contributes to their vulnerability (Pétillon et al. 2005).

### Coastal aquifers

In Mediterranean coastal systems, seawater intrusion is an important consequence of climate change and human action in coastal aquifers (Sherif and Singh 1999). Salinization alters the fundamental physicochemical nature of the soil-water environment, increasing ionic concentrations and altering chemical equilibria and mineral solubility (Herbert et al. 2015). Increased concentrations of solutes, especially sulfate, alter the biogeochemical cycling of major elements including carbon, nitrogen, phosphorus, sulfur, iron, and silica (Herbert et al. 2015), which has negative impacts on photosynthetic pigments and global biomass (Parihar et al. 2015). More about seawater intrusion in coastal aquifers can be found in *Section 3.1.2.2*. Three realistic scenarios (no change, sea rise of 0.5 m, land side lowered by 0.5 m by water pumping) were considered by Sherif and Singh (1999), who found that the Nile Delta aquifer is vulnerable to climate change and sea level rise. However, salinization tolerance can be found in some soil inhabitants like the spider *Arctosa fulvolineate* and the beetle *Merizodus soledadinus*, which survived

salinity levels up to 70‰ (Pereira et al. 2019). Also, some littoral and terrestrial amphipod species can survive salinity levels up to 900 mOsm external concentration (Morritt 1988).

### Risks from non-indigenous species

#### Phytoplankton

The Mediterranean Sea has experienced particularly strong algal bloom events over the past 50 years, mostly near the coast, in bays, lagoons, ports, beaches and estuaries, leading to deterioration in water quality, increasing the mortality of fish and risks to human health due to specific toxins that could be released into the marine environment and bio-accumulated through the marine trophic chain (Aligizaki 2009; Vlamis and Katikou 2015; Griffith and Gobler 2020). Climate-induced changes in water temperature, stratification and other physical properties appear to strongly impact the physiology and behavior of harmful algae bloom species, in terms of occurrence, physiology and toxin production (*Section 2.3.4*) in Mediterranean coastal areas where already more frequent Harmful Algal Blooms "HABs", and "novel" nuisance species have been recorded (Legrand and Casotti 2009).

The potential impact of climate-induced changes to phytoplankton, and especially HABs, has raised attention in the scientific communities and directed their research in this field, mainly driven by human health concerns due to the potency of some algal toxins that are transferred through the marine food web (Turki et al. 2006; Drira et al. 2008; Mabrouk et al. 2012; Estevez et al. 2019). In monitoring southern Mediterranean countries, more than 64 dinoflagellate species were identified with a remarkable increase in spring and summer (Feki et al. 2016). Dinoflagellate abundance between tidal periods was variable and the highest abundance was detected in the slack period in the Gulf of Gabès (southern coast of Tunisia), suffering from the pressure of high urbanization and industrialization rates, as well as rapidly increasing population growth rates. The dinoflagellates represent a major part of the eukaryotic primary production in marine ecosystems and the ability of several strains to cause shellfish poisoning and/or to form resting cysts has led to considerable attention being paid to the diversity and distribution of planktonic dinoflagellates in relation to environmental factors, hydrodynamism, nutrients and microalgae/biotoxins (Monti et al. 2007).

*Ostreopsis* (a dinoflagellate) blooms have become common in temperate areas as well, and regularly

occur in the Mediterranean Sea in the summer and autumn (Vila et al. 2001; Selina et al. 2014). In these areas, *Ostreopsis* was well-known as its blooms were often associated with harmful effects on the health of both humans and benthic marine organisms (Vila et al. 2001; Aligizaki and Nikolaidis 2006). Additionally, *Ostreopsis* often appeared in association with other toxic or potentially toxic benthic dinoflagellates such as *Prorocentrum* spp., *Amphidinium* spp. and *Coolia monotis* in several Mediterranean areas (Monti et al. 2007; Mabrouk et al. 2012; Selina et al. 2014). The genus *Ostreopsis* includes several species producing various palytoxin-like compounds with harmful effects on humans and marine fauna (Scalco et al. 2012). Species of this genus are regular members of the epiphytic community in tropical seas but their geographic range has shown an apparent expansion towards temperate regions of the Mediterranean Sea.

### Jellyfish

*Cassiopea andromeda* is a non-indigenous jellyfish species that possibly takes advantage of the warming tendency in the Mediterranean Sea. Recently detected in Malta, Sicily and other areas beyond the Eastern Mediterranean Sea, this benthic jellyfish seems to be well adapted to mesotrophic waters near harbors and closed bays in where water has low hydrodynamism (Yokeş et al. 2018). Due to the fact that many coastal areas all over the Mediterranean are no longer oligotrophic, the dispersion of this species may accelerate its path. The huge biomass reached in certain zones and its fast-growing features (Deidun 2018) may be a problem for fisheries, coastal tourism and management.

Other non-indigenous jellyfishes are also increasing their abundance in the warming Mediterranean Sea. The dreaded cubomedusa is typical of tropical seas, such as around the Australian Great Barrier Reef or the Philippines. Cubozoans, or "box jellyfish", are considered to be the cnidarian group's most dangerous, with an extremely painful sting that changes the lives of dozens of Australians every year. During the summer of 2008, a great abundance of a cubomedusa occurred along the Spanish coast, spotted off the beaches of Denia, Alicante, particularly *Carybdea marsupialis*, a small jellyfish species. It is rare in the Mediterranean Sea, so had never been considered to be a species that would form a major proliferation, yet during the summer of 2008 the Red Cross reported a high number of stinging incidents in this area due, no doubt, to this almost imperceptible, transparent and seemingly harmless jellyfish forming dense swarms in the breakers (Kingsford et al. 2018).

It is possible that *C. marsupialis* appeared because of the changing conditions throughout the water column, but the information about its distribution or what factors influence its life cycle is still limited (Canepa et al. 2017). Sea warming seems to be one of the key factors explaining its acute proliferation, and is likely the reason of the already changing trophic interaction map of the Mediterranean Sea due, in part, to climate change. Water temperature, together with quantity and quality of available food resources, are known as major drivers of gonadal outputs (Harland et al. 1992; Ben-David-Zaslow and Benayahu 1999). In general, jellyfish sexual and asexual reproduction is known to be influenced by warming. Some studies have shown that elevated temperature by itself or in combination with high feeding frequency (due to raised zooplankton prey abundance) increased budding rate and bud size in *Aurelia* polyps populations worldwide (Hočvar et al. 2018).

### Fish

The establishment of the lionfish (*Pterois* sp.) and the blowfish (*Lagocephalus* sp.) in Mediterranean waters can be envisaged as a paradigm of how climate change helps the dispersion of tropical species in a warm temperate sea (Section 2.5.1). The lionfish, for example, is a predator that has almost no controlling species (other fishes, sharks, etc.) and is a generalist, living in all shallow and mesophotic zones. The species has high reproductive and dispersal capacities, a massive production of well-protected eggs all year long and a fast spread and high larval survival rates (Betancur et al. 2011). In the Eastern Mediterranean Sea, these non-indigenous species have been introduced from the Red Sea, and have been documented in many areas (Bariche et al. 2013, 2017; Kletou et al. 2016). The blowfish (*Lagocephalus sceleratus*) is another example, with dispersion apparently faster in certain zones (Boustany et al. 2015; Kara et al. 2015). One of the first records indicate fast mobility from the original source (the Suez Canal) (Akyol et al. 2005; Kara et al. 2015), and since its first recorded sighting, it has been detected even in coastal waters in southern Italy (Azzurro et al. 2014). All the non-indigenous vagile fauna would have a very restricted dispersion if the water column temperature conditions were stable, with a clear marked low temperature during the autumn and winter, but changing conditions may be favoring its definitive establishment in Mediterranean waters (Bianchi and Morri 2003). The rapid spread of some of these species will be a serious problem for fisheries and trophic relationships in coastal areas, causing the likely local extinction of some species that may be preys of these generalist fish species (Coro et al. 2018).

### Plants

Among numerous threats, the pressure of non-indigenous plants has strong effects in these fragmented and linear ecosystems. *Dittrichia viscosa* is a perennial, 40-130 cm high plant, very common in the western Mediterranean but also found in its eastern part (Al Hassan et al. 2016). The species shows a remarkable pioneering character, and in recent decades largely expanded its range in Mediterranean countries, possibly due to increased human disturbances (Wacquart 1990; Mateo et al. 2013). Its capability to colonize new habitats and threaten biodiversity has been well-documented (Wacquart 1990) and related to characteristics such as its phenotypic plasticity (Wacquart and Bouab 1983), high stress tolerance (Curadi et al. 2005) and resistance to chemical pollution (Murciago et al. 2007; Fernández et al. 2013), as well as to its allelopathic effects (Omezzine et al. 2011). In the last 50 years, *D. viscosa* has become an invader in the NW Mediterranean region, since it increased its ecological range under disturbance pressure and is colonizing new habitats (Wacquart 1990; Boonne et al. 1992; Wacquart and Picard 1992; Mateo et al. 2013). The species' recent expansion in the Iberian Peninsula has also been correlated to temperature increases due to accelerated global warming (Vesperinas et al. 2001). Although *D. viscosa* cannot directly compete with true halophytes in highly saline environments, it is nevertheless quite stress tolerant and therefore presents a threat to the vegetation located on salt marsh borders, where several endemic and threatened species are found in the area of study conducted by Al Hassan et al. (2016).

### Other non-indigenous species

Some non-indigenous eco-engineering species are also favored by sea warming in the Mediterranean Sea (Section 2.5). Sea forests are living three-dimensional structures, similar to terrestrial forests but basically comprised of seaweeds, seagrasses, sponges, cnidarians, bryozoans, ascidians and other sessile organisms in the ocean benthos (Rossi et al. 2017a). These forests are dominated by ecosystem engineering species, organisms which directly or indirectly modulate the availability of resources to other species, causing changes to the physical condition of biotic or abiotic materials (Jones et al. 1994). In the case of the non-indigenous species, we can highlight two different case studies of non-indigenous eco-engineering species that may be already changing the benthic seascape in many areas of the Mediterranean Sea due to the suitable conditions for their expansion. The first case is represented by the seaweed *Caulerpa cylindracea*. This chlorophyte has been identified

as one of the most successful bioinvaders (Montefalcone et al. 2015). This species has been much more successful with respect to *Caulerpa taxifolia*, the so-called killer algae. In several areas, it is replacing other algae, phanerogams and sessile animals creating a new seascape in which the biodiversity and biomass are rapidly changing (Alomar et al. 2016).

A different case study is that of myxotrophic scleractinians that are more present toward the northern and western Mediterranean Sea. *Oculina patagonica* has been recorded in many Mediterranean areas but was not present in northern Spanish coastal areas until recently (Leydet et al. 2018). This species seems to be rapidly adapting to new temperature trends, a factor that drives new populations to quickly move north. Originating from the south, this species is now an invader that proliferates, replacing other native species in shallow waters, where sea urchin barrens may be essential to understand their capability to cope with new spaces (Coma et al. 2011). The presence of this species in northern areas seems to be correlated with sea warming, but has clear photobiological limits due to temperature factors that have to be considered (Rodolfo-Metalpa et al. 2014).

#### 4.2.1.2 Past changes

Although human activity is considered as a major driving force affecting the distribution and dynamics of Mediterranean ecosystems, the full consequences of projected climate variability and relative sea-level changes on fragile coastal ecosystems for the next century are still unknown. It is unclear how these waterfront ecosystems, as well as the services they provide, can be sustained, when relative sea-level rise and global warming are expected to exert even greater pressures in the near future (drought, habitat degradation and accelerated shoreline retreat) (Kaniewski et al. 2014). The most suitable archives for such paleoreconstructions are located in coastal wetlands that are highly vulnerable to global warming and the rapid rise in sea level, as they are highly exposed to processes such as flooding, subsidence, sediment scarcity and coastline erosion (Anthony et al. 2014; Wong et al. 2014) (Chapter 2).

#### Response of coastal ecosystems to past changes in sea level

During the Holocene period, sea level worldwide exhibited significant fluctuations, mostly responding to the advance and retreat of the continental



ice sheets (Lambeck et al. 2010). In the Mediterranean, regional sea level has risen continuously for the whole Holocene with a sudden slowdown at ~7.5 ka BP and a further deceleration during the last ~4.0 ka BP, after which time observed regional sea-level changes have mainly related to variability in isostatic adjustment (Galili et al. 2005; Vacchi et al. 2016). This mid-Holocene sea level stabilization had a major influence on fluvial sedimentation in coastal regions, and the formation of many coastal wetlands such as flood plains, coastal lagoons and deltas can be traced back to this period (Pavlopoulos et al. 2006; Fouache et al. 2008; Carmona et al. 2016). Although the stabilization of sea level and availability of sediment are important variables in this process, the palaeogeography of the lagoon environment and specific fluvial and marine sedimentary dynamics are also important variables in each case study (Huntley 2001).

Coastal wetland-based archives have an exceptional scientific potential to observe past ecosystem changes and identify key thresholds for particular ecosystems facing sea-level rise. In northeastern Spain, a reconstruction of past ecosystems was presented for the Castelló lagoon using ostracods, diatoms, and pollen and non-pollen palynomorphs as bio-indicators of marine vs freshwater influences (Ejarque et al. 2016). This study pointed out the strong link between the lagoon ecosystem, changing sea-level and the historical anthropogenic pressure. From the late Neolithic to the medieval period, dynamics of the lagoon ecosystem were mainly driven by changing marine influence. From ~1,550 and ~150 cal BC maximum marine flooding hampered agropastoral activities. After the Medieval period, societies actively controlled lagoon dynamics and became a major agent of landscape transformation. The removal of littoral woodlands after the 8th century was followed by the expansion of agrarian and industrial activities. The expansion of the milling industry and of agricultural lands led to the channelization of the river Muga into the lagoon after ~1,250 cal AD, which caused its transformation into a freshwater lake ecosystem (Ejarque et al. 2016).

In the Corsican back-barrier wetlands, a study of the fossil Coleoptera in two sediment cores supplemented by pollen and geochemical data show that 60% of past wetland beetle fauna at the Grecu pond became locally extinct because of the increase in salinity caused by marine intrusions (Poher et al. 2018). Most of this diversity loss occurred 3,700 years ago, when relative sea-level reported in the region rose. Regarding the Cannuta marsh, results of the same study show evolution

from a brackish lagoon to a freshwater environment marked by diversification of wetland beetle fauna 1,000 years ago, which is possibly due to relative sea-level stability and floodplain progradation (Poher et al. 2018).

In the Eastern Mediterranean, the rapid response of the Alikes Lagoon of Zakynthos Island ecosystem has been attributed from 8,540 and 3,400 BP to climate-related events such as storms or tsunamis where marine characteristics dominate (Avramidis et al. 2013). Furthermore, the study of Kaniewski et al. (2014) shows that Haifa Bay underwent a landward sea invasion, with a maximum sea penetration 4,000 years ago. The main consequences of the sea invasion were a retreat of the coastal forest, a loss of resilience and disappearance of the initial local biogeographic zonation. The forest replacement by a thorny shrub-steppe and then by an open-steppe appeared to follow, rather than cause, failure of tree regeneration. Due to the intrusion of the saline water table in freshwater streams, the freshwater wetland associated with the Na'aman River was deeply impacted after 4,000 cal. years, with a fall in hygrophilous-hydrophilous herbs. The subsequent coastal progradation that started at 3,400-3,300 cal yr BP left an eroded sandy-salty area, colonized by a steppe vegetation that became dominant until the end of the shoreline retreat (2,900 cal yr BP). A similar process was observed in the Salt Lake of Larnaca, Cyprus, where a shift from sheltered marine to lagoon environments produced an ecological change with a strong increase in xerophytic vegetation-types colonizing the shores that were no longer washed by seawater (Kaniewski et al. 2013b). The 3.2 K yr BP drought event (Schilman et al. 2001a; Kaniewski et al. 2013a) caused a dramatic demise in wooded ecosystems unrecovered until after 2,850 cal yr BP. Due to human-induced modification during the last ~200 years, it retreated by 15 m (Zviely et al. 2009). This new phase of sea-level invasion is well attested by high values of steppe vegetation and a renewed drop in hygrophilous-hydrophilous herbs. This suggests that similar stresses generate analogous biological processes, whatever the period under consideration.

Mediterranean coastal ecosystems, including fauna and flora components, are very sensitive to sea-level changes. The mid- to late Holocene changes in sea-level caused strong biological stresses and major ecological alterations providing a foreshadowing of potential future diversity and community changes along Mediterranean coastal wetland ecosystems and a model to understand the consequences of sea-level rise for

the next century (Marcos and Tsimplis 2008). This retrospective approach encompassing historical anthropogenic pressure also clearly indicates that for efficient management of coastal habitats, anthropogenic pressures linked to urban development (Section 5.1.1.3) must be reduced in order to mitigate the predicted effects of global change.

### **Response of coastal ecosystems to past climate variability**

In addition to eustatic sea level curves and coastal geomorphological dynamics, recent research has revealed that the climate variability of the Holocene impacted the Mediterranean coastal ecosystems particularly through storms and floods (Carmona et al. 2016). Storm surges, leading to catastrophic coastal flooding, are amongst the most feared natural hazards due to the high population densities and economic importance of coastal areas. Strong evidence has been provided for enhanced periods of storms that caused coastal flooding over the last 4,500 years as a result of solar activity, acting on cycles of around 2,200-yr and 230-yr (Kaniewski et al. 2016). These storm surges were characterized by inland intrusion of ostracods and dinoflagellate cysts, while the intrusion of saline water into the freshwater-fed plains greatly affected terrestrial ecosystems leading to land fragmentation by salt encroachment (Kaniewski et al. 2016). An important part of paleoenvironmental Holocene research has focused on identifying phases of high frequency and magnitude of floods related to climate variability (Thorndyraft and Benito 2006; Benito et al. 2008, 2015). In the Western Mediterranean region, periods with more frequent heavy-rain flooding events coincide with transitions to cooler and wetter climates, while flood frequency in northern Africa is linked to drier climate and in the eastern Mediterranean, to wetter conditions (Benito et al. 2015). Some studies hypothesized that these have been driven by internal modes of atmospheric and oceanic changes such as the East Atlantic pattern, the NAO, and thermohaline circulation (Degeai et al. 2017). However, the effect of these flood episodes on coastal ecosystems is much less documented and needs to be further addressed in high time-resolution studies.

Based on these findings, short-term climate events, which are driven by high frequency external and internal forcing factors, are superimposed on anthropogenic-driven factors. This complex interaction may either accentuate or attenuate the effect of current and future global warming. For instance, climate models are predicting a decrease in Mediterranean storms in the second half of the 21st century while the study by Kaniewski et al.

(2016) suggests that a decrease in solar activity will increase and intensify the risk of frequent flooding in coastal areas.


## **4.2.2 Projected vulnerabilities and risks**

### **4.2.2.1 Projections and risks based on biological groups**

#### **Phytoplankton**

Climate change consequences, particularly increasing temperatures (Section 2.2.4), decreasing nutrient replenishment (Section 2.3), and ocean acidification (Section 2.2.9), are expected to cause changes in plankton communities at different levels, from phenology and biomass to community structure. For example, a shift in phytoplankton community, dominance of smaller species (picophytoplankton and nanoflagellates) and a decrease in diatoms, with an expected decrease in the biomass of calcifying organisms such as coccolithophorids are some of the expected outcomes (Dias et al. 2010; The MerMex Group et al. 2011). There are still many uncertainties when it comes to the impact of sea warming and acidification on primary production in the Mediterranean, but it is clear that physico-chemical changes will affect the magnitude, timing and composition of phytoplankton blooms, with associated changes in the seasonal distribution of zooplankton (Moullec et al.) (Section 4.1.2). Ocean acidification combined with warming and deoxygenation, has been shown to cause negative effects on marine animals and to stimulate the production of primary producers, particularly in coastal waters that do not experience stratification or nutrient limitation (Gao et al. 2020). The associated decreased predatory pressure has the potential to further increase primary production. The increased primary production will stimulate the respiration of bacteria and thus intensify the hypoxia and low pH zone (Gao et al. 2020).

To understand how climate variation controls phyto- and zooplankton dynamics and possibly affects artisanal and small-scale fisheries exploiting areas near the coast, Goffart et al. (2017) used a unique long-term (1979-2014) time series obtained from a Mediterranean coastal area unbiased by local anthropogenic pressure in the Bay of Calvi, Corsica. They identified threshold values of physical variables below and above which they strongly impact nutrient availability, phyto- and zooplankton bloom characteristics and seasonality succession of plankton functional groups, stressing the importance of winter conditions in determining the state of Mediterranean pelagic



**Figure 4.8 | Expected geographical species shifts for the 288 coastal Mediterranean fish species** for both periods (a: 2040–2059; b: 2080–2099). Shifts were calculated from geographic range centroids and are expressed in degrees (NW: North West, NE: North East, SE: South East, SW: South West). Triangles represent the overall displacement of the fish assemblage calculated as the average centroid geographical shift (Albouy et al. 2012).

ecosystems, and highlighting that the thresholds obtained from their long-term time series provide key information for improving model scenarios of the impact of climate change on Mediterranean ecosystems.

The functional traits and geographic distribution of 106 copepod species were used to estimate the zooplankton functional diversity of Mediterranean surface assemblages for the 1965–1994 and 2069–2098 periods (Benedetti et al. 2018). Multiple environmental niche models were trained at the global scale to project species habitat suitability in the Mediterranean Sea and assess their sensitivity to climate change predicted by several scenarios. A relatively low decrease in species richness is predicted for 97% of the Mediterranean Basin, with higher losses in the eastern regions (Benedetti et al. 2018). The results of the same study show that climate change is not expected to alter copepod functional traits distribution in the Mediterranean Sea, as the most and the least sensitive species are functionally redundant. Such redundancy should buffer the loss of ecosystem functions in Mediterranean zooplankton assemblages induced by climate change. Since the most negatively impacted species are affiliated with temperate regimes and share Atlantic biogeographic origins, the results of Benedetti et al. (2018) are in line with the hypothesis of increasingly more tropical Mediterranean communities (Section 2.6.2.3).

### Fish

In the Mediterranean Sea, a reduction in primary production linked to an increase in sea surface

temperature (see previous sub-sections in this chapter and Section 3.2.2.2) could have negative impacts on fisheries catch and could exacerbate current trends of overfishing. Projected changes in primary and secondary productions suggest that trophic mismatches between fish pre-recruits and their prey could increase in the future, with negative consequences for recruitment success, sustainable fisheries and conservation of biodiversity (Lejeune et al. 2010; Stergiou et al. 2016). Also, jellyfish outbreaks (e.g., *P. noctiluca*) may become more frequent in the Mediterranean Basin (see previous sub-sections) and may extend over a longer period of the year than previously, causing alteration of the pelagic food web and thereby reducing fishery production (Licandro et al. 2010).

Using Bioclimatic Envelope Models (BEMs), the potential future climatic niches of 288 coastal Mediterranean fish species were projected based on a global warming scenario, then the species-level projections were geographically aggregated to analyze the projected changes in species richness and composition (Albouy et al. 2012). The results show that projected changes in assemblage composition are caused by different processes (species replacement vs. nestedness) in several areas of the Mediterranean Sea, and that the coastal fish fauna in several regions of the Mediterranean Sea could experience a “cul-de-sac” effect if exposed to climate warming (Albouy et al. 2012) (Section 4.1.2.2).

Fish species ranges are expected to move northwards and eastwards, and most of the Gulf of Lion

as well as the Adriatic and Aegean sub-basins are projected to experience a net decrease in species richness resulting from a loss of thermal niches for numerous fish that are not balanced by the arrival of other species from the south by the end of the 21st century (Fig. 4.8). In a best-case scenario, remaining or arriving species in local assemblages after global change impact would share combinations of functional traits with lost species, thereby maintaining ecosystem functioning, while in the worst-case scenario, lost species would have functional traits distinct from those of remaining or arriving species, hence the functions they support would go extinct and would imperil ecosystem functioning. Overall, the joint exploration of changes in species richness and composition coupled with the distinction between species replacement and nestedness bears important information for understanding the nature of climate change impacts on biodiversity (Albouy et al. 2012).

Lionfish (*Pterois* sp.) and blowfish (*Lagocephalus* sp.) in Mediterranean waters reflect how climate change helps the dispersion of tropical species in a warm temperate sea (Section 4.2.1.1). The question is when these species will migrate to the Western Mediterranean Sea. Johnston and Purkis (2014) argue that this is unlikely to happen. The connectivity between different areas is not the same as in the Caribbean for example, and the shifts in temperature need to be much more dramatic to make this invasion possible during the coming decades. However, the generalist behavior of this species may be one of the essential points to understanding its future success if a shift of 1 to 1.5 °C occurs in surface waters in some areas. This alien species from warm tropical waters may present a problem for other species because of the sea warming tendency (Section 2.2.8). It is not only a problem of the alien species per se, but a problem of shift from warm temperate to tropical trophic chain structures. The rapid spread of some of these species will be a serious problem for fisheries and trophic relationships in coastal areas, as the local extinction of some species that may be preys of these generalist fishes is very likely (Coro et al. 2018).

All the above-mentioned projections highlight the pressures that could increase the risk to fish and their habitats, namely for commercially valuable euryhaline coastal fish species (i.e., sole, seabass, seabream, mullet, eel) in coastal nursery sites such as lagoons, estuaries, and deltas. These pressures vary from rising temperatures that could exacerbate the occurrence of HABs (i.e., ciguatoxins, produced by dinoflagellates) and

thus the distribution of biotoxins and pathogens (i.e., *Vibrio* bacteria) (Lloret et al. 2016), to the depletion of oxygen that may cause suffocation, which kills fish, to plastic pollution (Barange et al. 2018). A likely decrease in connectivity between neighboring ecosystems within the Mediterranean is expected because of a decrease in the size of spawning areas and an increase in larval retention on smaller areas of the continental shelf (Barange et al. 2018). In addition to warming and ocean acidification, changes to fisheries' structures will contribute to the disappearance and modification of fragile and long-lived species that create biogenic structures or seagrass meadows, which provide important ecosystem services as well (Jordà et al. 2012).

### Seaweed

Other non-indigenous species are also favored by sea warming in the Mediterranean Sea. For the non-indigenous seaweed *Caulerpa cylindracea* (Section 4.2.1.1), the effects on the quality and quantity of available seston may be positive, also changing biogeochemical cycles and benthic-pelagic coupling relationships (Rizzo et al. 2017). Warming and acidification may be a perfect match for these fleshy algae (Comeau and Cornwall 2017), promoting its proliferation, in part, by the resistance to sedimentation processes, increased in many areas due to direct human impact (Alomar et al. 2016). Several algae species' dispersal may be thus enhanced not only by direct climate change effects but also by direct human impacts in coastal areas.

### Corals

As for myxotrophic scleractinians, such as *Oculina patagonica*, projected global warming is likely to cause a gradual contraction in their distribution zones, where temperatures are too high due to their temperature constraints (Rodolfo-Metalpa et al. 2014). Higher transparency of water and rising temperatures in surface waters may also be the key to understanding its successful proliferation. Myxotrophy needs a photosynthetic component that is not present in Mediterranean waters as it is in tropical waters: light (water transparency) and temperature, combined, are the key to understanding high photosynthetic performance (Schubert et al. 2017). Some of these species from tropical or subtropical waters, may be favored by the new conditions of the water column in the Mediterranean Sea, but not all of them. It is possible that Indo-Pacific species (Lessepsian species), adapted to higher temperatures, have more potential of spreading over Mediterranean shallow areas over the coming decades.

The reproductive features and the larval release strategy of octocorals species make them highly sensitive to global climate change (especially in shallow areas, above 40 m depths). Recent models highlight that water stratification in the Mediterranean Sea may last for longer periods and warmer waters may stress non-mobile organisms (Galli et al. 2017). Whether the food availability for benthic suspension feeders would be affected by elevated temperatures is not clear, but it has been shown that under anomalous warming episodes in shallow water adults of *P. clavate*, *E. singularis* and *C. rubrum* suffer from partial or total tissue loss (Garrabou et al. 2001; Linares et al. 2005, 2008; Rossi and Tsounis 2007). Also, the new recruit mortality in shallow populations is very high when compared to deeper populations (Bramanti et al. 2005; Linares et al. 2008; Coma et al. 2009). Furthermore, mother care (i.e., the energy invested by mother gorgonian colonies to the offspring) will be crucial to understanding the potential survival in a warmer and less productive ocean (Viladrich et al. 2016, 2017). Within the context of global change, there is a risk that the period of trophic crisis might be significantly prolonged to the point that the capacity of energy reserves in lecithotrophic larvae would not last until the arrival of favorable feeding conditions in early autumn. This situation could be even worse if the spawning of these species is triggered earlier by the increase in temperature. Asexual reproduction may enable some individuals to survive catastrophic mortality events such as warming episodes and then expand following the disturbance (Lasker and Coffroth 1999). However, chronic stress that reduces recruitment will have less obvious effects on these clonal taxa and may be the key to understanding future composition of benthic communities. Climate change could lead to partial recruitment failure in the affected species, with major changes in the population structure and dynamics, and a drastic change in ecosystem functioning. These combined factors may be crucial to understanding how seascapes will change in shallow Mediterranean benthic communities.

Non-indigenous species in the Mediterranean Sea may be invasive or simply immigrant species (Section 2.5). The new suitable conditions are key to understanding the transition observed in coastal and offshore areas. Higher temperatures that may be bad for native species (adapted to clear seasonal trends and certain limits of temperature and light), may be positive for the incoming species that are stressed by the same rising temperature phenomenon in their native areas.

#### 4.2.2.2 Projections and risks based on key natural habitats

##### Sandy beaches/dunes

The impacts from reshaped coastlines as a result of sea level rise and changes in wave climate were assessed via regional climate models, indicating that beaches of the Balearic Islands (western Mediterranean) would suffer a coastal retreat of 7 to 50 m, equivalent to half of the present-day aerial beach surface, under the RCP4.5 and RCP8.5 climate scenarios (Enríquez et al. 2017). Also, beach erosion due to sea level rise in the Aegean archipelago (eastern Mediterranean) was evaluated: under a mean sea level rise of 0.5 m (RCP4.5), a storm-induced sea level rise of 0.6 m is projected to result in complete erosion of 31 to 88% of all beaches (29 to 87% of beaches are currently fronting coastal infrastructure and assets), at least temporarily (Monioudi et al. 2017). The projections of the same study suggest a very considerable risk, which will require significant effort, financial resources and policies/regulation in order to protect/maintain the critical economic resources of the Aegean archipelago (Monioudi et al. 2017). Biodiversity loss will be the outcome of the negative pressures driven by climate change consequences, which would hamper beach ecosystem resilience (Scapini et al. 2019). The specificity of sandy beaches as narrow ecotones between sea and land may be lost under climate change pressure, adversely affecting fine-tuned macrofaunal adaptations and therefore ecosystem functioning (Scapini et al. 2019). In comparing two coastal plant communities, one in Montenegro and another in Albania, it is demonstrated that the less disturbed beach had zonation very similar to potential vegetation, while plant communities of the touristic beach were fragmented or even substituted by replacement communities (Šilc et al. 2016).

The way habitat distribution will be altered under the effects of two climate change scenarios were analyzed, and the efficiency of the current Italian network of protected areas in the future after distribution shifts was evaluated in Prisco et al. (2013). According to this latter study the range of habitats is currently sufficiently covered by protected areas, achieving the conservation target. However, according to their predictions, protection levels for mobile and fixed dune habitats is predicted to drop drastically under climate change (Prisco et al. 2013).

After combining a digital terrain model with 5 years of nest survey data describing location and clutch

depth, Varela et al. (2019) identified (a) regions with the highest nest densities for the loggerhead (*Caretta caretta*) and green turtle (*Chelonia mydas*), (b) nest elevation by species and beach, and (c) the estimated proportion of nests inundated under each sea-level rise scenario. On average, green turtles nested at higher elevations than loggerheads. However, because green turtles dig deeper nests than loggerheads, these were at similar risk of inundation. For a sea-level rise of 1.2 m, a loss of 67.3% for loggerhead turtle nests and 59.1% for green turtle nests were estimated (Varela et al. 2019). Existing natural and artificial barriers may affect the ability of these nesting habitats to remain suitable for nesting through beach migration.

### Rocky shores

The prolonged desiccation events on the southeastern rocky shores of the Mediterranean were characterized, and their potential ecological impacts on the unique intertidal Mediterranean Sea ecosystem of vermetid reefs were examined (from 2012 to 2014) by Zamir et al. (2018). This study shows that desiccation stress has already increased on southeastern Mediterranean vermetid reef ecological communities, and if this trend continues, further increases in aerial exposure and desiccation stress could be expected, which could have long-term impacts on this fragile ecosystem (Zamir et al. 2018). For the vermetid *Dendropoma petraeum*, long-term exposure to acidified conditions predicted for the year 2100 and beyond caused shell dissolution and a significant increase in shell Mg content. Unless CO<sub>2</sub> emissions are reduced and conservation measures taken, these reefs are in danger of extinction within this century (Zamir et al. 2018), with significant ecological and socio-economic ramifications for coastal systems (Milazzo et al. 2014).

The narrow range of the intertidal in the Mediterranean has particular implications for its resilience to climate change and sea level rise. For example, in the Mediterranean, the potential harsh effects of tidal aerial exposure on the ecological responses of intertidal organisms is, in fact, usually buffered by wave splashing (Sarà et al. 2011). This might help limit the otherwise detrimental impacts of increasing aerial temperature and dryness on organisms. However, the limited amplitude of the Mediterranean intertidal area implies a very small optimal range of environmental features. As such, some species may be able to adapt and migrate as sea levels rise, but others will not. This knowledge is not definitive as historical exploitation of the Mediterranean Sea and the absence of rigorous baselines makes it difficult to evaluate the current

health of these ecosystems and the efficacy of conservation actions at the ecosystem level (Sala et al. 2012).

### Coastal wetlands

The extent of wetland salinization and thus its effect on Mediterranean wetlands are still poorly known. Typically, increased salt and sulfide concentrations induce physiological stress in wetland biota and ultimately can result in significant shifts in wetland communities and their associated ecosystem functions (Herbert et al. 2015). In a large-scale outdoor mesocosm experiment, the effects of salinity on successional patterns, diversity, and relative abundance of Camargue (southern France) temporary pool crustaceans were studied (Waterkeyn et al. 2010). Salinity significantly altered crustacean communities hatching from the resting egg bank through a number of direct and indirect effects. Salinity had a significant negative effect on the establishment of large branchiopods and copepods. Both the diversity and density of cladocerans, especially chydorids, were positively related to salinity, possibly due to the absence of biotic interactions with large branchiopods at the highest salinity values (Waterkeyn et al. 2010). In the same study, the authors hypothesize that the salinity-mediated presence of the large branchiopod keystone group can shift the whole wetland regime from a zooplankton-rich clear-water state to a zooplankton-poor turbid state. Crustacean succession was significantly altered by salinity, by slowed development rates, population growth or maturation rates of some species. This suggests that in addition to salinity changes, any alteration of wetland hydroperiod (e.g., through aridification or poor water management) could have a synergistic effect on community structure and the diversity of invertebrate communities, including some keystone species.

Based on Multi-Criteria Decision Analysis techniques, it is documented that wetlands and terrestrial ecosystems have the highest relative risk scores in the Tunisian coastal zone of the Gulf of Gabes (Rizzi et al. 2016). A combination was made for regular sampling of waterbird presence through one annual cycle with in-situ data on relevant environmental predictors of waterbird distribution to model habitat selection for 69 species in a typical Mediterranean wetland network in southwestern Spain (Ramírez et al. 2018). Species associations with environmental features were subsequently used to predict changes in habitat suitability for each species under three climate change scenarios (encompassing changes in environmental predictors that ranged from 10% to 50% change as

predicted by regional climate models). Waterbirds distributed themselves unevenly throughout environmental gradients and water salinity was the most important gradient structuring the distribution of the community. Environmental suitability for the guilds of diving birds and vegetation gleaners will decline in future climate scenarios, while many small wading birds will benefit from changing conditions. Resident species and those that breed in this wetland network will also be more negatively impacted than those using this area for wintering or stopover (Ramírez et al. 2018).

### Seagrass meadows

Considering the rapid loss of seagrass habitat in the Mediterranean Basin and its capacity to capture carbon, preventing seagrass degradation by developing blue carbon projects is a major opportunity for financing seagrass protection. The avoided degradation of *Posidonia oceanica*, for instance, could provide an extra source of CO<sub>2</sub> capture of 4 tons per hectare per year (Sifleet et al. 2011). More importantly, avoiding their destruction would also prevent the washing away of carbon stored in the sediments under the seagrass beds and thus avoid the release of more than 500 tons of CO<sub>2</sub> per hectare, stored over millennia (Chefaoui et al. 2018). Temperate seagrass ecosystems as the thermal regime of the Mediterranean Sea, are sensitive to ocean warming and will exceed the upper thermal limit of the endemic *P. oceanica* in some areas (Marba and Duarte 2010; Jordà et al. 2012).

Using *Cymodocea nodosa* as a model species, Ontoria et al. (2019) assessed the joint effects of warming (at 20°C, 30°C and 35°C) with two potential outcomes of eutrophication. They found that in addition to the possibility of the persistence of *C. nodosa* being directly jeopardized by temperature increase, the joint effects of warming and eutrophication may further curtail its survival (see projected impacts on seagrasses in Section 4.1.2.1, whereas the drivers “Eutrophication” and “Warming” are detailed in Sections 2.2.7 and 2.3.2).

### Coastal lagoons

Coastal lagoons are sentinel systems that are highly vulnerable to potential impacts associated with climate change, particularly, as these systems have a key role in regulating the fluxes of water, nutrients and organisms between land, rivers and the ocean (Newton et al. 2018).

The effects of seawater acidification were assessed on a number of biological responses for one mussel and two clam species, including

growth and calcification, at two locations, namely a coastal lagoon in southern Portugal and in the Northern Adriatic Sea (Range et al. 2014). In this study, the CO<sub>2</sub> perturbation experiments produce contrasting responses depending on the species and location. Whereas the effects of acidification on growth and calcification in water of the Adriatic Sea were significant, in the coastal lagoon, these effects were much less evident, probably buffered by a high carbonate content in the water (Range et al. 2014). The same study reveals major variations in macroinvertebrate response to the imposed changes (in temperature and pH, respectively), underpinning the need for species-specific and location-specific adaptation measures. A reflection on threats to integrated management of the Thau coastal lagoon (France) due to climate change and the multi-scalar water scarcity adaptation strategy underlines that although water uses are currently secured thanks to the regional transfer of water, they are not coherent with local water management and create new vulnerabilities in the context of climate change (La Jeunesse et al. 2016). Climate change scenarios predict intensified terrestrial storm runoff, providing coastal ecosystems with large nutrient pulses and increased turbidity, with unknown consequences for the phytoplankton community. In the same lagoon (Thau), a 12-day mesocosm experiment shows that pulsed terrestrial runoff can cause rapid, low quality (high carbon: nutrient) diatom blooms (Deininger et al. 2016). However, bloom duration may be short and reduced in magnitude by fish. Thus, climate change may shift shallow coastal ecosystems towards famine or feast dynamics.

Covering most of the bottom of the Mar Menor lagoon (southeastern Spain), *Caulerpa prolifera* has probably increased the resistance of the lagoon to eutrophication processes through the high uptake of nutrients from the water column and their retention in the sediments, avoiding high phytoplankton densities (Lloret et al. 2008). Nevertheless, if climate change predictions prove true, the current status of the lagoon is likely to collapse, since future environmental conditions could make *C. prolifera* unable to reach values of net photosynthesis greater than zero, and eutrophication processes are expected to appear (Lloret et al. 2008).

### Deltas

A comprehensive overview of the status and sustainability of the Ebro, Rhône, and Po Deltas and Venice Lagoon has been published by Day et al. (2019), showing that all of these systems have been strongly modified by human activities. However, each system has a unique combination of


impacts that informs management and restoration approaches (see the interaction among drivers in Section 2.6).

As an example, the Ebro Delta is a diverse area in terms of wetland habitat types and has high ecological (e.g., it is the second most important "Special Protection Area" for birds in Spain) and economic value (e.g., third largest producer of rice in Europe) (Fatorić and Chelleri 2012). In the last 150 years, the Ebro delta has been largely transformed into rice fields, which now cover 70% of the total area (Cardoch et al. 2002) and have both direct and indirect effects on the ecology of the area, such as salt infiltration in ground water. The main impact is the destruction of natural habitats, but even the remaining deltaic ecosystems have been affected by rice production, through alteration of the natural hydrological cycle as a result of freshwater inputs during the rice growing season (April to September). In addition, large amounts of nutrients and pesticides are delivered for the fertilization and care of the rice paddies (Forès 1992). The planned construction of 49 new reservoirs mainly for irrigation purposes and withdrawal of water upstream from the delta are forecasted to have drastic ecological consequences such as reductions in sediment and freshwater inputs into the delta and detrimental side effects on deltaic ecosystems fauna and flora (Prat and Ibáñez 1995). Overall, because of its morphology, relative sea-level rise will become the most important climate-induced potential hazard for the Ebro delta (Sánchez-Arcilla et al. 2008).

Sea level rise may severely threaten many key coastal ecosystems such as the Nile delta and may cause the loss of important habitats such as the loggerhead (*Caretta caretta*) nesting beaches (UNEP/MAP-RAC/SPA 2009). Projection of averaged sea-level rise trends by El Sayed Frihy et al. (2010) indicates that the coastal plain of the Nile Delta and Alexandria is vulnerable to accelerated sea-level rise but not at the same level due to wide variability of the land topography, which includes low-lying areas, high-elevated coastal ridges and sand dunes, accretionary beaches, and artificially protective structures. Similarly, based on Earth System model simulations, the sea-level variation along the Egyptian coasts is significantly affected by other factors such as sea-level variation West of the Gibraltar Strait, steric sea level, and sea-surface temperature (Shaltout et al. 2015).

**Coastal aquifers**

Several recent studies have considered the possible impacts of climate change and seawater level rise on seawater intrusion in coastal aquifers (Sefelnasr and Sherif 2014). All have revealed the severity of the problem and the significance of the landward movement of the dispersion zone under seawater level rise. Most of the studies did not consider the possible effects of seawater rise on the inland movement of the shoreline and the associated changes in the boundary conditions at the seaside and the domain geometry. Such effects become more evident in flat, lowland, coastal alluvial plains where large areas might be submerged with seawater under a relatively small increase in seawater



**Figure 4.9 | Nile Delta, Egypt.** A) Submerged land in the coastal zone under 0.5 m seawater rise. B) Submerged land in the coastal zone under 1.0 m sea-level rise (Sefelnasr and Sherif 2014). Red line indicates the border of the basin.



level. None of the studies combined the effect of increased groundwater pumping, due to the possible decline in precipitation and shortage in surface water resources, with the expected landward shift of the shoreline. Using a two-dimensional horizontal model, the study of Sefelnasr and Sherif (2014) investigated the possible effects of seawater level rise in the Mediterranean Sea on the seawater intrusion problem in the Nile Delta Aquifer. They concluded that large areas in the coastal zone of the Nile Delta will be submerged by seawater, and the coastline will shift landward by several kilometers on the eastern and western sides of the Delta (Fig. 4.9). By using an equivalent porous continuous medium to represent a karstic Apulian aquifer (southern Italy), an evident piezometric drop was confirmed for the past period (until 1999) and a likely similar dramatic drop in the future was projected (Romanazzi et al. 2015). All phenomena considered in this study's models (e.g., sea level and sea salinity) showed non-negligible effects on coastal groundwater (Romanazzi et al. 2015).

The effects of salinization on coastal aquifers' biogeochemistry typically include decreased inorganic nitrogen removal (with implications for water quality and climate regulation), decreased carbon storage (with implications for climate regulation and wetland accretion), and increased generation of toxic sulfides (with implications for nutrient cycling and the health/functioning of wetland biota) (Herbert et al. 2015). In agriculture, studies on the salinization effects on soil organisms are scarce, but negative effects of saline conditions on survival and reproduction of soil invertebrate species (Owojori et al. 2008, 2014) or on avoidance behavior of earthworms (Bencherif et al. 2015) have been reported. Deleterious effects of soil salinization on diverse life stages of agriculture plants have also been described (Wichern et al. 2006), including decreased and/or delayed germination and/or effects on seedling physiognomic state, deficient growth, as well as a decrease in photosynthetic pigments, and global biomass (Parihar et al. 2015). However, some littoral and terrestrial amphipod species can survive salinity levels of up to 900 mOsm external concentration (Morritt 1988), as along with other spiders mentioned earlier (Pereira et al. 2019).

#### 4.2.2.3 Vulnerabilities

##### Coastal urbanization

The Mediterranean bioregion is currently suffering severe disturbance due to intensive urbanization

and climate change effects (Adloff et al. 2015)<sup>34</sup>. The situation is expected to worsen as land availability decreases (due to the global warming and infrastructure impairment), while demographic growth and migration flows are likely to pursue (Burak and Margat 2016) (read more on land and sea use changes in *Section 2.4*, water management and infrastructure in *Box 3.1.1*, and the vulnerability of coastal energy systems to climate extremes in *Section 3.3.2.3*). In the future, hydroclimatic hazards, probably more frequent and intense, will have adverse impacts on ecological balances and human health and well-being, particularly in coastal Mediterranean cities where almost one-third of the population lives (Hallegatte et al. 2009; Magnan et al. 2009; Adloff et al. 2015; Im et al. 2018). However, Mediterranean coastal cities seem to lack a long-term vision (i.e., establishing smart cities, green cities, etc.) for planning future urban development and valuable policies and social-economic resources for establishing participative governance (Mazurek 2018) (*Section 5.1.3.1*).

Social-economic contexts and urban growth rates, trends and phases are quite variable from North to East and South and even across each country (Im et al. 2018). While the size of these urban settlements varies from North to South, most have a historic urban center developed around a harbor near the sea, which makes the different biological species living in or near these areas highly vulnerable not only to human stressors, but also to global phenomena like climate change (*Chapter 2*).

The survey of juvenile fish populations across various infrastructures and natural sites along a 100 km shoreline of the French Mediterranean coast demonstrated that anthropogenic structures can play an important role as potential juvenile fish habitats, particularly in harbors where highly variable densities were found, with densities on ripraps or jetties that were equivalent to those of natural sites (Mercader et al. 2018). This is the case of the herbivorous fish *Siganus rivulatus* in Lebanon-Eastern Mediterranean where it settled in protected shallow areas offering hard substrates and algal communities such as muddy harbors (Bariche et al. 2004).

One of the most frequently documented negative impacts of the high density of harbors and boats, are the collisions and disturbance of large mammals (dolphins, whales, sea turtles). In particular, the destructive impact of fishing practices on dolphin populations has reached international news

<sup>34</sup> <http://www.medqsr.org/>

headlines [see *Section 4.1.2.2* for more information about ship collision effects on cetaceans]. Another threat for marine mammals in harbor areas is marine dredging with high concern and impact on the marine life of cetaceans, pinnipeds, and sirenians with effects largely unknown (Todd et al. 2015). The leisure activities, habitat degradation, noise, contaminant remobilization, suspended sediments, and sedimentation may impact marine mammals indirectly through changes to prey. Dredging has the potential to impact marine mammals with specific species and location effects, depending on the type of dredging equipment. In harbor areas, marine mammals continue to be impacted by many anthropogenic activities and almost all marine mammal species have been reported to face at least one threat in the Mediterranean (Laran et al. 2017; Avila et al. 2018).

Reduced primary production (*Section 4.1.2.2*) and the possible increase of local blooms of toxic algae in some urbanized coastal areas could have many repercussions on marine ecosystem services such as tourism and fisheries, and ultimately on human health (read how the effects of climate change on coastal ecosystems could affect livelihood, culture and human rights in *Chapter 5.3*). The impact of eutrophication is largely observed in many Mediterranean systems, namely in harbors. For example, in Punic harbors of Carthage, the oldest and most well-preserved in the Mediterranean Basin (Gulf of Tunis, South Mediterranean Sea), the harmful blooms of *Dinophysis sacculus*, *D. acuminata*, *Alexandrium* spp., *Gymnodinium aureolum*, *Gymnodinium impudicum*, *Akashiwo sanguinea*, *Scrippsiella* spp. and *Prorocentrum gracile* were identified in correlation with water temperature and orthophosphate concentrations (Aissaoui et al. 2014) (*Section 2.3.3*).

The Mediterranean Basin is particularly exposed to biological invasions through shipping from maritime traffic and the high number of harbors constitute large areas for the extension of several non-indigenous species (Izquierdo-Muñoz et al. 2009). The species *Pseudonereis anomala* (Gravier 1900) (Polychaeta, Nereididae) first recorded in Alexandria (Egypt) by (Fauvel 1937), was recorded in several harbor areas in the Mediterranean within the period (2003 and 2005) (Kambouroglou and Nicolaidou 2006), indicating shipping transfer of benthic species (read more on biological pollutants in *Section 2.3.4*, on non-indigenous species in *Section 2.5*, and on future risks associated with non-indigenous species in *Section 6.12*).

In harbor systems, where macrofauna is scarce and difficult to sample, the study of meiofaunal

assemblages is proposed as the most suitable instrument for monitoring purposes since, ports, ranging from large commercial harbors to small tourist marinas, are the main link between anthropized and natural coastal ecosystems, and should be taken as primary sources of coastal disturbances (Sedano Vera et al. 2014). Other Mediterranean species associated with marine fouling harbors have been described by Khedhri et al. (2016). The brachyuran decapod is associated with marine fouling in Egyptian Mediterranean harbors and nine species of 9 genera affiliated with 5 families have been recorded so far.

The spatial and temporal changes in climate attractiveness in the Mediterranean could have major impacts on the sustainability of tourism development as suggested by Amelung and Viner (2006), who used a Tourism Climate Index based on future climate change scenarios for the Mediterranean region. This intense tourism activity is harming the Mediterranean shores. Based on the type of garbage on 13 Mediterranean beaches, there are indications that most Mediterranean coastal litter is land-based (Gabrielides et al. 1991). In fact, based on beach cleanups organized over the summers of 2016 and 2017 in Cyprus, Loizidou et al. (2018) suggest that although these initiatives are quite successful at collecting large pieces of marine litter, small pieces of litter (such as cigarette butts and small pieces of plastic items related to recreational activities) remain, accumulating or buried over time, with some items becoming a nuisance to beach goers and a potential source of marine litter. This issue is already influencing coastal organisms such as loggerhead sea turtles (*Caretta caretta*) where, according to a survey by Tomas et al. (2002), the most frequent type of debris in their gastrointestinal tract is plastics (75.9%). Furthermore, the environmental impacts of sunscreen chemicals are likely to be exacerbated in the Mediterranean waters due to the massive influx of tourists and its densely populated coasts, the basin's limited exchanges with the ocean, the high residence time of surface waters, and its oligotrophic waters, which raises significant concerns about its toxicity on marine biota and its bioaccumulation in the marine trophic chain (Tovar-Sánchez et al. 2019) (*Section 2.3*).

Coastal cities with a sandy ground are often exposed to massive sand extraction. An investigation on the short-term effects of sand extraction on macrozoobenthic communities before and after beach dredging along the Emilia-Romagna coast (northern Adriatic Sea) showed no significant settlement of opportunistic species (Simonini et al.

2005). The same study suggested that the limited impact of sand extraction operations on the physical characteristics of sediment and hydrological-sedimentary characteristics in the relict sand area should aid its rapid recovery and the restoration of the original community in a short period of time (2–4 years after dredging). The same pattern was documented in the coastal ecosystems of the Bay of Blanes on the Catalan coast (Sardá et al. 2000) where recolonization in these dredged habitats was fast, and no changes in seasonal trends were detected after dredging. However, this latter study documented that the filter-feeder *Callista chione* and the carnivorous polychaetes *Protodorvillea kefersteini* and *Glycera* spp., were still significantly reduced after two years, suggesting that a longer period is needed to restructure dredged bottoms to their initial situation (Sardá et al. 2000).

### Sea level rise

Similar to the impacts of sea level rise elsewhere (Bernstein et al. 2019; Mullin et al. 2019; Murfin and Spiegel 2020), many Mediterranean regions will be increasingly exposed to a major risk of submersion and erosion, affecting several parts of the coast (with extreme cases being Venice, Kerkennah archipelago in Tunisia, Alexandria and the Nile delta) (UNEP/MAP/PAP 2015; UNEP/MAP 2016). Future risks associated with sea level rise are detailed in Section 6.9. The main consequences on coastal ecosystems include more frequent and/or intensive flooding along low-lying coasts, particularly in delta areas, lagoon coasts, tideland and some islands (Sections 2.2.8.1 and 6.9.1). Slight increases in mean sea level will lead to relatively quick inundation, deterioration and displacement of significant areas of wetland vegetation. Severe losses of coastal wetlands are expected in the Mediterranean (McFadden et al. 2007). Apart from the actual loss of land area, these wetlands support rare and localized habitats containing highly specialized organisms, the degradation or loss of which will in turn impact migratory bird populations, particularly along main migratory routes (Cyprus, Malta, Palm Islands Nature Reserve in Lebanon). Mediterranean waterbird communities already show changes in community composition based on the recent changes in temperature and whether or not they have a strict protection status, greatly improves the adaptability of species and communities (Gaget et al. 2018). Future breeding suitability maps indicate that the little tern (*Sterna albifrons*) and the common tern (*Sterna hirundo*) could potentially face a drastic decrease in suitable breeding grounds even in protected areas (Ivajnić et al. 2017).

A modelling study from the island of Zakynthos-Greece for the loggerhead sea turtle (*Caretta caretta*) suggests that even under the most conservative 0.2 m sea-level rise scenario, about 38% (range: 31 to 48%) of total nesting beach area would be lost, while an average of 13% (range: 7 to 17%) of current nesting beach area would be lost (Katselidis et al. 2014). For a sea-level rise of 1.2 m, they estimated a loss of 67.3% for loggerhead turtle nests and 59.1% for green turtle nests although suitability of nesting sites for future migration will also be dependent on existing natural and artificial barriers (Varela et al. 2019).

Accelerated cliff and beach erosion will result in habitat and species loss. For example many cliffs host chasmophytic endemics while many coastal habitats of priority importance at the European/global level (i.e., coastal Junipers, *Posidonia* meadows) (Gubbay et al. 2016; Janssen et al. 2016). In the western Mediterranean, seagrass could reach functional extinction under warming scenarios (Jordà et al. 2012; Telesca et al. 2015). The effects of sea-level rise on competition and the subsequent plant diversity decrease in Mediterranean-climate marshes (Noto and Shurin 2017). Increased salinization in the estuaries will result in species changes/structure, function, and occurrence of eutrophication (EEA 2004; Bernes 2005; Robinson et al. 2005; Smayda 2006).

## 4.2.3 Adaptation

### 4.2.3.1 Adaptation of different coastal systems

When it comes to adaptation strategies for coastal systems to environmental changes, different zones require specific actions. For example, shorelines are mainly affected by deterioration of engineering species such as corals, and vermetids forming reefs that protect coasts from erosion, regulate sediment transport and accumulation and provide habitat for other species. Estuaries are particularly vulnerable to pollution, including plastic from nearby human settlements and require different adaptation strategies. Thus, suitable adaptation policies include (i) reducing pollution runoff, both from agriculture and industry and waste management, (ii) policies to limit or prevent acidification and (iii) moving aquaculture operations to areas protected from critical acidification levels (Sections 3.1.5 and 6.11).

Keystone Mediterranean benthic species are vulnerable to ocean acidification and warming (Rodolfo-Metalpa et al. 2011, 2014; Milazzo et al. 2014;

Zunino et al. 2017; Verdura et al. 2019). Therefore, spatial planning for these areas should include plans for coastal protection and different urbanization schemes (infrastructure sector), plans to enhance the attractiveness of these zones for tourists, and different regulations for recreational boats. The most likely mechanisms by which ocean acidification refugia (OAR) can mitigate ocean acidification impacts are reducing exposure to harmful conditions or enhancing adaptive capacity (Kapsenberg and Cyronak 2019). While local management options, such as creating OAR, can help coastal ecosystems to adapt, they present unique challenges, and reducing global anthropogenic CO<sub>2</sub> emissions remains a priority. Given the scale of ocean acidification impacts on human health and well-being, recognizing and researching these complexities may allow the adaptation of management such that both the harms to human health are reduced and the benefits enhanced (Falkenberg et al. 2020).

Deep waters are mainly impacted by changes in wild harvests so adaptation measures should focus on fisheries indirectly impacted by changes in phytoplankton production at the surface as well as ocean warming. Shallow coastal zones are exposed to changes in availability of fish and shellfish. Hence, here the most effective adaptive measures involve the management of both fisheries and aquaculture, and the wise use of coastal habitats. The resilience of socio-ecological systems to sea level rise, storms and flooding can be enhanced when coastal habitats are used as natural infrastructure since they provide similar services and added benefits that support short- and long-term biological, cultural, social, and economic goals (Powell et al. 2019). Better integration across policy and planning instruments is needed to enhance adaptive capacity at the interface of climate change adaptation, marine and aquaculture planning and management. This requires holistic and cooperative management tools, such as aquaculture management areas, that could support adaptation across wider spatial scales (Greenhill et al. 2020). This could be enabled by establishing links between existing and proposed collaborative groups to enhance development of adaptation responses and through co-ordination of monitoring and review processes to promote learning across scales (Kapsenberg and Cyronak 2019; Powell et al. 2019; Greenhill et al. 2020). Economic and

financial tools to promote environmental management are detailed in *Section 5.1.3.2*.

#### 4.2.3.2 Harmful algal bloom monitoring

In the last two decades, Harmful Algal Bloom (HAB) events have increased, with many species suddenly emerging in regions previously free from such toxic or potentially harmful algae. Along the Mediterranean coastline, several phytoplankton toxic networks have been established such as the French REPHY network<sup>35</sup>. The recent observations are quite atypical for phytoplankton blooms, and may be partially explained by exceptionally favorable new environmental conditions related to climate change (Draredja et al. 2019; Jenhani et al. 2019; Ninčević Gladan et al. 2020). Coastal HABs appear to have increased on a global scale and several reasons have been suggested: better knowledge of toxic species, better monitoring and alerting systems, the transport of algal cysts in ballast waters, the development of aquaculture, the stimulating effect of urban and industrial activities and/or atypical climate conditions (Glibert et al. 2005). The same trend has been observed in the Mediterranean (see *Sections 2.3.4 and 4.2.1.1*). Thus, national and regional water quality assessment efforts and routine coastal monitoring programs intended to detect species, and the study their toxicities have increased worldwide and in the Mediterranean area as well (Nastasi 2010).

Similar to programs elsewhere around the Mediterranean, the Tunisian national monitoring network of phytoplankton and phycotoxins, has been implemented since 1995 to ensure public safety by establishing tools for early warning of bloom events. Also, a regional project "Risk-Monitoring, Modelling and Mitigation (M3-HABs) of benthic microalgal blooms across the Mediterranean regions" found that better awareness of the risks associated with the *Ostreopsis* blooms could be achieved, including appropriate diffusion of cautionary measures, the production of common monitoring protocols, the development of new technologies for species-specific identification, species counting, and the build-up of prediction models in order to prevent and reduce risk factors for the environment, human health and economic activities. Despite the efforts in management and monitoring work, predicting the impact of climate

<sup>35</sup> [http://www.ifremer.fr/envlit/surveillance/phytoplancton\\_phycotoxines](http://www.ifremer.fr/envlit/surveillance/phytoplancton_phycotoxines)

change on HABs in the Mediterranean is very challenging as it involves many combining factors. Thus, the calls for increased awareness in both HAB and seafood monitoring programs remain a high priority in the Mediterranean region (Turki et al. 2014; Visciano et al. 2016; Estevez et al. 2019; Ordoñez et al. 2019).

#### 4.2.3.3 Early detection of potentially dangerous species

An updated list of introduced alien species in the Mediterranean (November 2018) shows that the non-indigenous species (NIS) number is close to 1,000 species (Section 2.5.1.1). Adaptation to NIS requires protecting the coastal population against the possible risks associated with the establishment of NIS (Section 2.5). In fact, in the past two decades research interest in NIS has increased, mostly stimulated by evidence about their ecological and socio-economic impacts in the Mediterranean region. This has also raised the urgency of innovative approaches to forecast, track and manage these species (Corrales et al. 2018) (Section 6.12). For example, the Early Detection and Rapid Response (EDRR) has been recognized as a key aspect for NIS management and acknowledged by the European Commission, and has been included in the new European regulation (EU) No 1143/2014 on the prevention and management of the introduction and spread of NIS. Efficient public awareness campaigns disseminating information to local communities, also through “specific alerts”, was adopted as the key driver to quickly detect unwanted NIS and are still used in the last few years with several theoretical frameworks developed through formalized early warning systems (Azzurro et al. 2014).

The silver-cheeked toadfish *Lagocephalus sceleratus* (Gmelin, 1789) (Tetraodontidae) has expanded rapidly through the Western Mediterranean (Stefantaris and Zenetos 2006; Jribi and Bradai 2012; Kara et al. 2015). Due to its toxicity, many Mediterranean countries have quickly responded by informing the general public about the risks associated to the consumption of this species. These awareness initiatives, necessary to limit the impacts of this invasion (Nader et al. 2012), have been carried out in countries such as Egypt, Turkey, Lebanon, Cyprus, Greece and Tunisia (Ben Souissi et al. 2014). The same strategy is being adopted for the common lionfish *Pterois miles* (Section 2.5.1.3). Hence, early detection and continuous monitoring of these species is a successful example of positive interaction between citizens, researchers, and policymakers (Azzurro et al. 2016).

The current list of NIS provides a reliable updated database and basis to continue monitoring the arrival and spread of NIS in the Mediterranean, as well as to provide counsel to governmental agencies with respect to management and control. The current geographical, taxonomical and impact data gaps can be reduced only by instituting harmonized standards and methodologies for monitoring alien populations in all countries bordering the Mediterranean Sea.

#### 4.2.3.4 Adaptation management strategies for the jellyfish *Pelagia noctiluca*

Long-term climate fluctuations have been correlated with jellyfish abundance in Mediterranean waters as revealed by (Molinero et al. 2005, 2008). The *Pelagia noctiluca* represents the most important jellyfish species in the Mediterranean Sea (an oceanic scyphozoan that has become very abundant along the coasts) with negative interaction and toxicity (Condon et al. 2013). In order to better monitor and track the dispersion of jellyfish in the Mediterranean and raise awareness about these species, many networks have been established. The CIESM JellyWatch Program was set up in 2009 to gather baseline data on the frequency and extent of jellyfish outbreaks across the Mediterranean Sea (CIESM 2009b, 2009a). The Medusa Project set up in Catalonia set out to understand the spatio-temporal dynamics of the jellyfish populations in the NW Mediterranean Sea by carrying out daily sampling during summer (May to September) of 243 beaches, covering more than 500 points. The recommendations of Medusa were to enhance similar sampling programs for all Mediterranean coasts to better understand changes in the distribution, abundance, and blooming patterns of dangerous jellyfish species (Canepa et al. 2014). The MED-JELLYRISK project “towards an early warning system to detect jellyfish swarms”, started with a campaign to better understand the movement of jellyfish blooms. Three sea drifters were deployed off the coast of Mellieha Bay. Based on satellite tracking, the information gathered by the sea drifters - including sea surface currents (direction and strength) and temperature allowed scientists to validate numerical models that can simulate the dispersion of jellyfish blooms and predict their incidence on coastal areas. These jellyfish dispersion models constituted the basic element of a prototype system intended to act as an early warning of jellyfish swarms impacting Mediterranean beaches. The mission of the Italian, Maltese, Spanish and Tunisian scientists behind the MED-JELLYRISK project (2014-2015) was

making public authorities, local businesses and beachgoers ready to live together with jellyfish and adapting solutions to address the growth of such fascinating creatures (Lucas et al. 2014). The possible relationships between climate change and blooms of *P. noctiluca* have been studied and it has been proposed that *P. noctiluca* may be an indicator of climate variability in the Mediterranean Sea (Daly Yahia et al. 2010; Condon et al. 2013; Rosa et al. 2013).

Despite these studies and monitoring surveys, additional studies and long-term surveys are needed to improve knowledge of the eco-physiology of the marine species, which will help to better manage and maybe take advantage of NIS, harmful microalgal and jellyfish biomasses and/or their bioactive molecules as a resource for biotechnological applications, from biofuels to pharmacology, cosmetics, health products, food for humans and feed for livestock or aquaculture farms (de Domenico et al. 2019).

#### 4.2.3.5 Ecosystem-based adaptation management

Given the already poor conditions of exploited resources, there is a need for fisheries management to adapt to future changes and to incorporate climate change impacts into future management strategy assessment (Moullec et al. 2019) (Section 3.2). Ecosystem-based adaptation is gaining attention as a cost-effective method for protecting human and ecological communities against the impacts of climate change. This approach has been supported by many studies for various Mediterranean habitats to improve their resilience against the consequences of various drivers. For example, an initial assessment of vulnerability to sea-level rise to help decision makers, and other relevant stakeholders, to develop appropriate public policies and land-use planning measures has been provided (Demirkesen et al. 2008). Also, potential strategies to ameliorate the impact of seawater inundation have been proposed, such as: wetland preservation, beach nourishment at tourist resorts and the afforestation of dunes (Snoussi et al. 2008). Protected areas can play an important role in safeguarding coastal dune plant communities against land-use transformations (Prisco et al. 2016). In this context, functional traits can guide conservation planning, helping to identify groups of species most at risk of population declines. Future conservation interventions need to be mindful to ensure that the natural disturbance regime of dune ecosystems is not disrupted. At the water body management level, scientific cooperation is

necessary to deal with the conceptual and ecological difficulties derived from inter and intra-lagoon variability in hydrology and biological assemblages, which are inherent factors in the functioning of these complex ecosystems (Pérez-Ruzafa et al. 2011).

The multiple levels of land-sea interactions (Fang et al. 2018) require a new approach to Integrated Coastal Zone Management (ICZM) and marine spatial planning. The Mediterranean includes hotspots of global priority for land-sea integration (Halpern et al. 2009) and there are emerging Mediterranean case studies which embrace such an approach (Ramieri et al. 2019). Conservation planning and management should focus on cross-realm processes and building resilience between realms. In this respect, connectivity between processes and structural elements is of the utmost importance. Single realm connectivity is inadequate (Fang et al. 2018) since it cannot account for cross-system threats (Beger et al. 2010) and multi-realm species (Giakoumi et al. 2019). Interaction between realms should be translated into structural connectivity (see for example the framework proposed by (Beger et al. 2010) or functional connectivity (Magris et al. 2018) where multi-realm species are taken into consideration. In addition, integrating connectivity and climate change (Magris et al. 2014; Keeley et al. 2018) can be used in various spatio-temporal scales and could be fully applied to maintain and restore land-sea processes. Case studies and innovative approaches are highlighted in Sections 6.9.3 and 6.9.4.

#### 4.2.3.6 The role of institutions/actors and local communities: recommendations

Adaptation efforts often focus on one species, or species group (e.g., jellyfish, algae) or on a specific land-sea ecosystem (e.g., sandy beaches). A more holistic approach is required when trying to establish adaptation methods for the entire Mediterranean, which could include:

- Ecosystem Based Management (EBM) of coastal areas: ecosystem-based approaches (the integrated management of land, water and living resources) to climate change adaptation and mitigation.
- Identify adaptation and mitigation interactions (synergies and conflicts) and assumptions related to adaptation/mitigation.
- Building institutional capacity (governance - adaptive management/monitoring) to improve governance over land/sea natural resources and climate

change adaptation. A Nexus approach has been proposed by the IUCN Commission on Ecosystem Management - coastal ecosystem group (CEM/CEG) based on three complementary approaches: (i) sectoral adaptation, (ii) cross-sectoral governance and (iii) territorial strategic planning (Krchnak et al. 2011; Ozment et al. 2015).

- Adaptation and coastal community resilience: Impacts will directly affect coastal communities. A participatory research approach may assist in assessing vulnerability of socio-economic groups, their current adaptation strategies and their adaptive capacity to cope with the impacts of climate variability and extremes and sea-level rise. Such an approach may identify barriers and opportunities for community response to climate change and place emphasis on any existing knowledge/practices promoting adaptation.
- Integrating local knowledge and institutions with

respect to risk management and adaptation is part of an ecosystem-based approach (Colls et al. 2009).

- At a practical/management level, adaptation to coastal risks can be achieved via hard structures but also soft protection including beach nourishments and dune rehabilitation, in addition to improved spatial planning regulations, and anticipatory and reactive adaptation.

Overall, information, data, adaptation techniques and networks do exist, but collaboration can be enhanced. Also, public involvement in the development and implementation of adaptation strategies for these ecosystems can be improved. Awareness campaigns on the importance of adaptation measures for these ecosystems may help policymakers to make the necessary steps to ensure their protection and conservation against potential climate change impacts.

## 4.3 Terrestrial and freshwater ecosystems

### 4.3.1 Current conditions and past trends

The Mediterranean region is recognized as a global biodiversity hotspot (Myers et al. 2000; Mittermeier et al. 2005), representing one of the Earth's most geologically, biologically, and culturally rich and complex regions (Blondel and Aronson 1999; Blondel et al. 2010; Visconti et al. 2018). More than 50 plant refuges during recent ice ages (Médail and Diadema 2009) have allowed plant diversity to be exceptionally high (Mittermeier et al. 2005), with about 25,000 plant species today (Myers et al. 2000) and 60% endemism (Thompson 2005). There are 290 tree species (Noce et al. 2016), of which 200 are endemic (Quézel and Médail 2003; Gauquelin et al. 2018). Two thirds of Mediterranean amphibian species, 48% of reptiles, a quarter of mammals, 14% of dragonflies, and 3% of birds are endemic (Mittermeier et al. 2005; Lefèvre and Fady 2016; Paine and Lieutier 2016). However, the diversity of several taxa has not been analyzed in detail under the taxonomic and ecological angles (Azam et al. 2016).

#### 4.3.1.1 Past climate variability and its impact on terrestrial ecosystems

Regional temperatures in the Mediterranean Basin are now ~1.4°C higher than during the 1880-1920 period, higher than the increase in global temper-

ature of 0.85°C (Chapter 2). During the Holocene (especially in the second half of this period), periods of precipitation deficits have occurred, but in contrast to the 21st-century situation, temperatures did not rise above the present average. These periods of precipitation deficits (~6 to ~5.2, ~4.2 to ~4, and ~3.1 to ~2.9 thousand yr BP) have been identified as possible causes of declines or collapses in civilization in the eastern Mediterranean region (Guiot and Kaniewski 2015).

Information from tree rings from different tree species growing at high elevation can provide annually-resolved, absolutely dated climate information across the Mediterranean covering the past centuries. Reconstructions from locations across the Mediterranean reflect different climate conditions during different times of the year. Recent tree ring-based climate reconstructions reflect not only seasonal temperatures but also parts of the hydrological cycle including drought stress. Only few tree-ring based climate reconstructions go beyond the past 600 years and this limits our understanding of drought variability, the magnitude and timing of long-term trends and centennial-scale variability across the Mediterranean back to medieval times.

The most detailed depiction of Mediterranean drought variability over the last 900 years is presented in the Old World Drought Atlas (OWDA), a tree-ring-based field reconstruction of warm-season drought severity




Figure 4.10 | Mediterranean land systems (Malek et al. 2018).

(Cook et al. 2015). Summer drought reconstructions from high-elevation sites from Mount Smolikias in the Pindus Mountains (Northern Greece) go back to AD 730 (Konter et al. 2017; Klippel et al. 2018). Drought variability displays significant East-West coherence between the western (Spain, Morocco, Algeria, and Tunisia) and eastern (Balkans, Greece, and Turkey) Mediterranean Basin on multi-decadal to centennial timescales (Cook et al. 2016). There appears to be a north-south contrast in the eastern Mediterranean, with a tendency for wet anomalies in Greece, Anatolia, and the Balkans while Libya, the southern Levant, and the Middle East are dry and vice versa associated with North Atlantic Oscillation (NAO) and other atmospheric circulation dynamics (Cook et al. 2016). The recent droughts in north-western Africa (Morocco) and the Levant are unusual in the context of the past 900 years (Esper et al. 2007; Cook et al. 2016). In the Pyrenees, the Alps, the northern Apennines, the Balkans, the north-western and southern Carpathians long tree ring width formation are mostly controlled by summer temperature (Buntgen et al. 2007; Büntgen et al. 2009, 2017; Popa and Kern 2009; Panayotov et al. 2010). Reconstructions from the Iberian Peninsula and northern Africa for the past 900 years reflect overall warmer conditions around 1200 and 1400, and again after around 1850. Cooler conditions are reconstructed for the mid 13th century and between the 15th and 18th century.


The only winter/spring  $\delta^{13}C$  from tree ring-derived temperature reconstruction from southwestern Turkey indicates warmer conditions during the early 12th century and the late 15th century and lower temperatures from the early 16th century to the late 19th century (Heinrich et al. 2013). Tree ring-based climate reconstructions account for a maximum of 35% explained variance and thus are associated with large uncertainties.

Pollen-based reconstructions for the entire Holocene confirm the picture that significant switches between drier and wetter conditions have occurred around the Mediterranean Basin, even if temperatures have never reached current levels. Vegetation has switched between major biome categories in up to 10% of the land area from one century to another, with only slightly higher values during the particular shifts identified above (Guiot and Cramer 2016).

#### 4.3.1.2 Direct human impacts on ecosystems in the past

The Mediterranean Region is also one of the regions with the longest and most intense human occupation in the world (Underwood et al. 2009), and its diversity is the result of co-evolution between human societies and their environment, characterized by constantly evolving land use practices over at least the past 300 generations of human occupation (Blondel 2006). The





**Figure 4.11 | Map of Global Dryland Assessment (GDA) plots** showing main changes in land use over the years 2000 to 2015. Non-forest land uses are shown in green. Land use shifting from forests to other uses is shown in red. Changes from other land to cropland and settlements are shown in yellow. Plots that did not change are shown in black (Martín-Ortega et al. 2018).

presence of many endemic species is closely related to extensive use of Mediterranean landscapes, particularly agro-silvo-pastoral mosaic systems (Médail and Quézel 1999) and wetlands (Cuttelod et al. 2009). Reconstructing landscapes and ecosystems over the course of time remains difficult. Knowledge of the human history of the region is therefore still limited. Most changes in land cover appear to have been due to change in human activities, often inducing diversity through changes in different taxonomic groups and their interactions (Sirami et al. 2010). Throughout the Holocene, Mediterranean ecosystems appear to have been rather resilient to perturbations (Blondel 2006; Underwood et al. 2009). Together with urbanization and agriculture intensification (Myers et al. 2000), land abandonment and the decrease in open habitats are key trends in several countries of the northwestern part of the Mediterranean Basin (Portugal, Spain, France and Italy) (Mazzoleni et al. 2004).

Since about 1980, biodiversity changes are faster and greater across different Mediterranean taxonomic groups and habitats (Blondel et al. 2010; Vogt-Schilb et al. 2016; Delpon et al. 2018). Species loss is marked by a general trend of homogenization (loss of vulnerable and rare species) recorded in several taxonomic groups and by a general simplification of biotic interactions (loss of specialized relationship) (Blondel et al. 2010; Visconti et al. 2018).

The most detailed land use map of the Mediterranean indicates a highly heterogeneous spatial structure of land use systems (Fig. 4.10) (Malek and Verburg

2017; Malek et al. 2018). In a coarser reconstruction of land use change during recent decades, cropland was found to be the dominant land use (35.2%), grassland was the second most common land cover (26% of plots), followed by forest (20.7%) and other lands (13.4%). Settlement and wetlands accounted for the smallest number of plots, with 3.3% and 1.4% respectively (Martín-Ortega et al. 2018).

During the period 2000-2015, human activities have intensified in the Mediterranean region, particularly in Spain, France, Turkey and most North African countries (Fig. 4.11), where an increase in cropland was recorded. An intensification of agricultural activities in the region is associated with marked transitions from non-irrigated or heterogeneous cropland to permanently irrigated cropland (Ruiz-Benito et al. 2012), leading to an increased use of freshwater resources, with similar projections being made for the future (Malek et al. 2018). New areas containing settlements occurred concurrently with this regional expansion in cropland because of urbanization and tourism, indicating an important trend of urbanization across the region (Martín-Ortega et al. 2018), impacting landscape character, resources use and ecosystem services capacity (Martínez-Fernández et al. 2009; Ruiz-Benito et al. 2012; Balzan et al. 2018).

### Forests

In the Mediterranean region, the term “forest” comprises a variety of vegetation types interleaved with one another in complex patterns created by

variations in soil, topography, climate, human use and fire history, among other factors. This includes dense stands with a closed canopy, as well as pre-forest or pre-steppe structures with lower tree density and tree height (Gauquelin et al. 2018) such as the human-shaped Mediterranean mosaic landscapes, including *dehesas* or *montados*, and multifunctional agro-silvopastoral systems of pastures with scattered oaks, typical of the Iberian Peninsula. The forests of the Mediterranean Basin cover more than 48.2 million ha of which 35 million are in southern Europe, 8.8 in the Middle East and 4.4 in North Africa (Quézel and Médail 2003; Fady and Médail 2004). However, based on the FAO definition of forests (“Land spanning more than 0.5 ha with trees higher than 5 m and a canopy cover of more than 10%”), there were an estimated 88 million ha of forest area in Mediterranean countries in 2015, representing 2.2% of the world’s total forest area (FAO and Plan Bleu 2018).

Despite the small extent of the Mediterranean forest area compared to rest of the world, it is a floristic global hotspot with ca. 25,000 flowering plant and fern species (4.5% of the world’s endemics), which represent approx. 10% of the world’s flowering plants (Myers et al. 2000). It is also the world’s second highest region in terms of in plant endemism, with 50–60% of the plants being found nowhere else, including emblematic species such as cork oak (*Quercus suber*), argan (*Argania spinosa*), cypresses such as *Tetraclinis articulata* or *Juniperus thurifera*, or fir species such as *Abies pinsapo*, *A. marocana*, *A. nebrodensis*, many of them endemic to the different mountain ranges across the Mediterranean (Thompson 2005; Blondel et al. 2010). The Mediterranean Basin hosts 290 indigenous woody species and subspecies (in comparison to 135 for non-Mediterranean Europe), 201 of which are endemic (Fady-Welterlen 2005). Intra-region variability in climate, soil and human factors result in a mosaic of forest types (Masiero et al. 2013). The relatively harsh climate conditions in arid zones prevent the existence of tall forests and lead to the formation of *maquis* and *garriga* shrublands, dominated by evergreen shrubs such as *Pistacia lentiscus*, *Quercus coccifera*, *Q. calliprinos* and *Cistus sp.* The semiarid zones are dominated by *Pinus halepensis* in the western part and *Pinus brutia* in the eastern areas. Sub-humid areas are the typical habitat for evergreen oaks such as *Quercus ilex* or *Q. suber*, but also *Pinus pinea* and other accompanying species such as *Arbutus unedo* or *Erica arborea*. Deciduous and marcescent oaks appear in the sub-humid to humid Mediterranean areas, with oak species such as *Quercus pubescens*, *Q. cerris*, *Q. pyrenaica*, *Q. faginea* or *Q. macrolepis*,

among others, accompanied by conifers such as *Cedrus sp.* or by Mediterranean firs (*Abies pinsapo* or *A. cephalonica*). In mountain areas pines become the dominant species including *P. nigra* and *P. sylvestris*, and it is also possible to find islands of oceanic climate with *Q. robur*, *Q. petraea*, *Fagus sylvatica* or *Abies alba*. Along the rivers, forests of *Fraxinus sp.*, *Populus alba* and *P. nigra* can prosper (FAO and Plan Bleu 2013). The wildlife diversity associated with this variety of forest environments is also high: 786 of 1,601 vertebrate Mediterranean species live in forest habitats and 792 of 1,184 terrestrial insects assessed by the IUCN Red List (as in 2018) are recorded as living in forests, 364 of which are endemic to the Mediterranean region (FAO and Plan Bleu 2018).

### The human footprint in Mediterranean forests

The current composition, structure, dynamics and biological diversity of Mediterranean forests cannot be understood without considering the long history of uses and changes induced by human activities, which have contributed to shaping the Mediterranean landscapes as we know them today (Blondel 2006). Human influence in the Mediterranean dates back several thousand years, to the point that some authors argue that a “coevolution” has shaped the interactions between these ecosystems and the human societies that inhabited them (Blondel 2006).

Transformation into agricultural fields, over-exploitation, the prevalence of livestock grazing within forests, and the repeated occurrence of natural and human-caused fires led to a progressive reduction and fragmentation of vegetation cover, and forests are mainly confined into the less fertile slopes and occupy less than 15% of their potential area (Quézel and Médail 2003). In some areas, the loss of forest canopy on slopes and their associated understory after fire events has led to important soil erosion (Cerdà and Mataix-Solera 2009; Shakesby 2011). Nevertheless, most soil degradation in forests of the Mediterranean Basin is associated with overgrazing and trampling of the forest understory (Le Houérou 1990; FAO 2016; FAO and Plan Bleu 2018).

In many areas, however, the combination of forests, pastures and fields, together with the high variability in climate, relief and soil resulted in a mosaic-type landscape that greatly contributed to maintaining the biological diversity of Mediterranean landscapes. These landscapes are highly dependent on human stewardship to maintain their resilience to disturbances (e.g., by reducing fire risk through browsing the forest understory (Blondel 2006). The population increase and industrial

development from the 18th century onwards led to an increase in the pressure on forests for wood and charcoal on the northern shore of the Mediterranean Basin (Nocentini and Coll 2013). Forests were intensively cut and transformed into coppices, some species were overexploited, and many forests were transformed into uniform, even-aged systems managed through clearcutting or uniform shelter-wood (Puettmann et al. 2008).

At the beginning of the 20th century, extensive reforestation plans were implemented in many European countries to reverse the trend. For example, 3.3 million ha were reforested between 1938 and 1984 in Spain, 460,000 ha in Portugal and around 1.3 million ha in Italy during the 20th century (Pemán and Serrada 2017). These large national reforestation programs mainly used conifers (mostly pines) due to their ability to grow in degraded soils and harsh environments. Many reforested areas contributed to a general improvement of environmental conditions, but the use of a single species over vast areas, together with the lack of subsequent management led to very homogeneous forests, often at excessive densities, with associated expansion of pests and a high risk of wildfires (Nocentini and Coll 2013; Guijarro et al. 2017; Martín-Alcón et al. 2017).

Since 1990, overall forest area has increased by  $0.67\% \text{ yr}^{-1}$  across the Mediterranean Basin (FAO and Plan Bleu 2018). Despite this generally increasing trend, forest loss and degradation still prevail around most of the Mediterranean Basin, especially in coastal areas, due to population increase and urban expansion (FAO and Plan Bleu 2018). Sharp differences can be observed between sub-regions. Almost all countries in the North experienced a huge increase in forest area, with rates around  $1\% \text{ yr}^{-1}$  in Italy, France and Spain (Masiero et al. 2013), to which afforestation only contributes  $0.23\% \text{ yr}^{-1}$ . The major part of this trend is due to the decline of agriculture and grazing and the consequent abandonment of marginal lands that are colonized by forests, a process that has been stimulated by European Common Agricultural Policy subsidies (FAO and Plan Bleu 2013, 2018). In contrast, on the southern Mediterranean shore, forest ecosystems are still at risk of fragmentation or disappearance due to human pressure from clearing and cultivation, overexploitation of firewood and overgrazing (Gauquelin et al. 1999; Croitoru 2007; Palahi et al. 2008; Djema and Messaoudene 2009; Masiero et al. 2013; FAO and Plan Bleu 2018). For example, Algerian forests decreased at a rate of  $0.5\%$  from 1990 to 2010 (FAO and Plan Bleu 2013) and a decrease rate of  $\sim 126,000 \text{ ha yr}^{-1}$

across North Africa has been estimated over the last 25 years (Keenan et al. 2015). This degradation continues despite forest representing 22% of the protected land area in North African countries (FAO and Plan Bleu 2018). However, many of these protected areas generally lack management plans or the resources to implement them (IPBES 2018).

### *Ecosystem services provision by Mediterranean forests*

Mediterranean forests are complex and biodiversity-rich socio-ecological systems, resulting from the coevolution of plants and societies through millennia of human perturbations and management (Blondel 2006; Doblás-Miranda et al. 2015; Gauquelin et al. 2018). Currently, Mediterranean forests play a key role in the livelihoods of diverse communities across the Mediterranean by providing people with ecosystem services, food and products for home consumption and income generation. In particular, the provision of non-wood forest products (NWFPs) and other services (e.g., watershed protection, soil erosion mitigation) (Merlo and Croitoru 2005; Croitoru 2007) stands out from the provision of wood forest products (WFPs) (FAO and Plan Bleu 2018). Removal of WFPs represents about 20 to 40% of the estimated total economy value in most northern countries, but less than 15% in most southern and eastern countries (Croitoru 2007). Northern Mediterranean countries dominate all areas of wood production, especially roundwood, pulpwood and derived products, with countries in eastern Mediterranean making a significant contribution to fiberboard production (Turkey produces 50% of the fiberboard products in the region) (FAO and Plan Bleu 2013). Production is low in the southern Mediterranean countries except for wood fuel, which constitutes one-third of the total production in the sub-region (FAO and Plan Bleu 2013), with firewood reaching 80 to 100% of total removals in Tunisia, Morocco and Lebanon (Croitoru 2007). In any case, the overall production of WFP in the Mediterranean is insufficient to meet regional demands, making the region a net importer of wood and wood forest products (FAO and Plan Bleu 2013).

The main NWFPs of Mediterranean forests include cork, pine cones and pine nuts, mushrooms, chestnuts, honey, truffles, berries, acorns, carob, myrtle, rosemary, and other products. Most of these NWFPs are generally harvested, stored and consumed by local communities or constitute their main source of income (FAO and Plan Bleu 2018). There is a significant geographical variation in production and consumption of NWFPs, largely dependent on the tree species available

in each country. Portugal is the main producer of cork (50 percent of total production), followed by Spain (30%), Morocco (6%), Algeria (5%), Tunisia (4%), France (3%) and Italy (3%) (APCOR 2015). In 2016, the estimated annual export value of cork by Mediterranean countries was €1,295.8 million (APCOR 2015). Cork production is mostly concentrated in cork-oak savannas (also called *dehesas* or *montados*) that result from an intentionally induced simplification (both in terms of structure and species diversity) of the Mediterranean forest: human intervention reduces tree density, removes shrub cover and fosters the growth of grass. These are considered “biodiversity-based product systems” by the Convention of Biological Diversity and have a multi-functional character, contributing to the provision of other services such as fuelwood, acorns (to feed animals), carbon storage and pasture, while supporting important habitats for biodiversity (Bugalho et al. 2011).

Pine nut extraction (from stone pine *Pinus pinea*) generates an income of about €50–60 ha<sup>-1</sup> yr<sup>-1</sup> (for a cone yield of 200 kg ha<sup>-1</sup> yr<sup>-1</sup>), which is higher than the revenue from timber (€20–30 ha<sup>-1</sup> yr<sup>-1</sup>), fuelwood, and other products or uses (FAO and Plan Bleu 2013), where the most productive areas are Portugal and Lebanon, producing 4–7 t ha<sup>-1</sup> yr<sup>-1</sup>. In Mediterranean Europe, mushroom picking is a recreational activity and mushrooms are marketed according to origin (e.g., France, Italy and Spain) rather than used solely as a direct food source for local communities (FAO and Plan Bleu 2018). The saffron milk cap (*Lactarius deliciosus*) and porcino (*Boletus edulis*) are the most important species commercially traded, along with the highly-valued black truffle (*Tuber melanosporum*) which is increasingly artificially inoculated on purposefully cultivated oak trees. In Turkey, where 90% of the total pine honey is produced (*Pinus brutia* and *P. halepensis*), the beekeeping sector is the main source of income for nearly 10,000 families in the region (Croitoru and Liagre 2013). In Morocco, for example, argan forests contribute to 7% of regional GDP and ensure subsistence for 14% of the rural population (Croitoru and Liagre 2013). Grazing in forested areas remains the main source of subsistence for local populations in Algeria, Morocco, Tunisia, Lebanon and Turkey (Daly Hassen 2016). One of the most recent marketed values of Mediterranean forests is their attraction for tourism, sometimes to the detriment of other forest services (García-Nieto et al. 2013) or to the forest itself (Kuvan 2010).

Mediterranean forests also hold many important non-marketed values, mostly regulating services.

Among these, one of the most relevant is watershed protection: forests regulate watershed hydrological regimes and protect against erosion and extreme flooding events while filtering and purifying water for its local consumption (Palahi et al. 2008; Guerra et al. 2016). Watershed protection is the single most valuable benefit from forests in Syria, for example, accounting for more than 50% of the total economic value of forests (US\$100 ha<sup>-1</sup> yr<sup>-1</sup>). In the Maghreb countries, it is second in value only to grazing, varying within US\$26–32 ha<sup>-1</sup> yr<sup>-1</sup> (Croitoru and Liagre 2013). Mediterranean forests play an important role in regulating micro-climatic conditions, atmospheric composition, water and biochemical cycles (Peñuelas et al. 2017). They also represent a net carbon sink, helping mitigate climate change impacts (Section 4.3.3). In 2005, the economic value of carbon storage in Mediterranean forests ranged between US\$ 37 billion and US\$ 63 billion, i.e., 13% of the forests’ total economic value, when assuming the SRES IPCC scenarios A1 and B1, respectively, for the 2050 horizon (Ding et al. 2010). However, carbon storage capacity by Mediterranean forests is strongly modulated by management (Seidl et al. 2014; Bravo et al. 2017). In fact, forests can act as carbon sources if disturbed, poorly managed, overexploited or burnt (Ding et al. 2010; Peñuelas et al. 2017).

Other societal values of Mediterranean forests include their cultural, spiritual and religious importance (especially for the few remaining ancient forests) (Mansourian et al. 2013) and their attractiveness (aesthetics) for recreational activities and tourism (FAO and Plan Bleu 2013; Bernetti et al. 2019; Raviv et al. 2020). Recreational uses of Mediterranean forests can lead to trade-offs with other services: for example, in Tunisia, the number of visitors to parks demanding recreational services from forests increased from 93,000 to 110,000 between 1998 and 2014 (Daly-Hassen et al. 2017). In these areas, limiting the access to the public also limits soil erosion, one of the main explicit concerns of Tunisian forests (Daly-Hassen et al. 2017).

### Mountains

According to the UNEP definition, Mediterranean mountains cover some 1.7 million km<sup>2</sup>. Seven Mediterranean countries are among the top 20 mountainous countries in the world, and half of the countries in the region have at least 50% of their land classified as mountain areas (Regato and Salman 2008). Mediterranean mountains exhibit similarities in their biotic, ecological, physical and

environmental characteristics but also significant differences (floristically, human colonization patterns, historic land uses and current anthropogenic pressures). Mediterranean mountains host many regional and local endemic species, some of which are relicts of past biogeographical patterns. Médail and Diadema (2009) identified 33 mountainous areas within 52 refugia in the Mediterranean Basin. Some of these mountains had already been identified as regional biodiversity hotspots (Médail and Quézel 1999) and global centers of plant diversity (Davis et al. 1994).

Historically, Mediterranean mountain forests have been crucial for the development of all civilizations and countries in the region. Most of the prehistoric populations of the Near East originated in Mediterranean mountain areas with very high plant and animal diversity, year-round water, shelter and suitable conditions for survival. Early mountain farmer-herders in the eastern Mediterranean and North African mountains changed pastoral and cropping patterns leading to the domestication of major livestock and domestic species but also important crops, including barley and wheat.

#### **Land use changes in mountain regions**

The long history of human intervention has modified land cover and resulted in numerous land use changes over time. Although Grove and Rackham (2003) support the resilience of modern Mediterranean landscape to changes since ancient times, McNeill (1992) argues that for most of the mountains, the changes that destroyed the environment and left behind skeletal landscapes are comparatively recent (past 200 years). Major land uses included woodcutting, pastoralism, agriculture and mining. Drivers of land use changes (north vs. south) included socio-political, economic, environmental expansion vs. population decline, urbanization and industrialization. The beginning of the 20th century, marked the start of rural emigration from the mountains of northern Mediterranean countries that peaked after the Second World War, resulting in land abandonment and the remarkable increase of forest cover seen in recent years. In southern Mediterranean countries, in contrast, an opposite trend is recorded due to the substantial increase in the rural population and consequent pressure on the mountains for arable and grazing land.

Recent land use changes have disrupted the traditional agro-silvopastoral equilibrium of the Mediterranean mountains. The increased forest cover in the northern Mediterranean has resulted in a decline in species and especially of habitat diver-

sity and an increase in natural hazards, especially forest fires, but increased carbon sequestration and decreased soil erosion and sediment transport to the lowlands. The decrease in forest cover in the southern Mediterranean, on the other hand, has led to severe soil erosion.

The intensity and therefore impact of the principal human activities, i.e., agriculture, grazing and tourism, vary significantly from north to south of the basin. For example, agriculture was historically more important in the Sierra Nevada, the Lucanian Apennines of Italy and the Rif mountains than in Taurus mountains or the Pindos mountains (McNeill 1992). In addition, in Morocco the extensive deforestation of mountains has been reported as a result of an increasing rural population, intensive grazing and the end of traditional pastoral nomadic migrations (Rejdali 2004). Regional differences have also been demonstrated in the case of Lefka Ori and the Psiloritis mountains in Crete where the opposite pattern was observed with abandonment mainly due to different grazing practices and number of stock density (Papanastasis 2012). This reflects the impacts of cultural practices (tradition) even within the same region (Regato and Salman 2008; Papanastasis 2012).

Land use change is still considered a more imminent threat in the short to medium term for mountain areas compared to climate changes (Tasser et al. 2017). Abandonment of agricultural activities (including grazing) emerges as a common trend that can be identified in most of the Euro-Mediterranean mountains as a result of rural depopulation which started in the 1950s, with plenty of examples from Spain, Italy and Greece (Papanastasis 2012). Typical countryside structures such as terraces and stone walls are collapsing, soil erosion is increasing while secondary succession is taking place in addition to reforestation activities carried out to mitigate the effects of torrential floods within rivers and ravines and to reduce the siltation of reservoirs within the valley bottoms (López-Moreno et al. 2008). Abandonment continues to date with farmlands abandoned as a consequence of migration to urban settlements in lowland areas, resulting in significant reduction in cultivation in the northern Mediterranean (Vicente-Serrano et al. 2004; Lasanta-Martínez et al. 2005). However, reports from Turkey support the argument that 'people are still in the mountains' since the decline in forest cover is small compared to other areas in the Mediterranean (Kadioğullari and Başkent 2008; Günlü et al. 2009) while in the Taurus, livestock husbandry is still very active (Kaniewski et al. 2007).

### Mountain biodiversity changes

Mediterranean mountains located on the borders of different biogeographical regions, and three continents, are biodiversity hotspots with a flora which comprises different phytogeographical elements ranging from Euro-Siberian to Arctic-Alpine, and Irano-Turanian in the eastern Mediterranean. This is particularly demonstrated in the mountain flora of Crete, Cyprus and Turkey. In the Mediterranean Basin, with its long history of human activity, mountains are considered to be some of the last remaining wilderness areas with high landscape and biodiversity value providing a wide range of ecosystem services within and beyond their boundaries. Geology, tectonic activity, isolation and limited human activity explain the current biogeographical patterns occurring in Mediterranean mountains. These patterns conform to theory, with mountains displaying low species richness but high endemism along altitudinal gradients. In recognition of their importance, a high number of protected areas and many mountainous areas are part of the UNESCO World Network of Biosphere Reserves (Sierra Nevada, Mount Olympus and Lefka Ori-Crete).

Community composition changes have been recorded both because of land use as well as climate change. The increased forest cover in the northern Mediterranean has resulted in a decline in species and especially of habitat diversity (Papanastasis 2012). Altitudinal shifts have been already reported from the Montseny mountains (Peñuelas and Boada 2003), as well as changes in the abundance of endemic species (Fernández Calzado et al. 2012). Community-level studies in Mediterranean mountain ranges indicate that there will be colonization of high altitudes by subalpine species (Stanisci et al. 2005; Kazakis et al. 2007) or what Gottfried et al. (2012) have termed "thermophilization". There is already evidence in Mediterranean mountain areas of an increase in the frequency of extreme events, a direct result of climate change, manifesting itself as droughts and, sediment transfer (Maas and Macklin 2002).

### Drylands and shrublands


Drylands are characterized by low precipitation levels which do not compensate for the evaporative demands imposed by high temperatures and solar radiation, thereby exhibiting high aridity levels (MEA 2005; Reynolds et al. 2007). Dryland ecosystems have low productivity, which is often exacerbated by the highly irregular, low predictable pulses of rain, resulting in a long period during which soil moisture depletion with no restoration

prevails. Low productivity generates relatively low plant biomass, which produces only small amounts of plant litter and leads to low soil organic contents (Safriel 2006).

Drylands are classified using the aridity index (AI related to the average ratio of annual precipitation over potential evapotranspiration) developed by the United Nations (Middleton and Thomas 1997) into four classes: hyper-arid ( $AI < 0.05$ ), arid ( $0.05 < AI < 0.20$ ), semi-arid ( $0.20 < AI < 0.50$ ), dry sub-humid ( $0.50 < AI < 0.65$ ). In the Mediterranean, drylands represent almost 80% of its area (Fig. 4.15a) from which 12.3% are classified as hyper-arid, 16.5% arid, 36.7% semi-arid and 14.5% dry-sub-humid (considering the boundaries of the Mediterranean SREX region defined in the 5th IPCC Assessment Report). Climatic constraints limit the productivity of drylands, increasing their susceptibility to wind and water erosion. These climatic limitations, coupled with intense human activity (e.g. agriculture, grazing and deforestation) has been leading to desertification and land degradation particularly in the Mediterranean (Olsson et al. 2019) (Fig. 4.12).

The Mediterranean domain has undergone an overall increase in arid area of almost 15% (from 64% to 78%) at the cost of the more humid aridity classes (Daliakopoulos et al. 2017; Elsen et al. 2017). Changes in land cover in drylands have both human-driven and climate variability as underlying causes and have resulted in extensive land abandonment especially after 1960 (Moreira et al. 2011; Stellmes et al. 2013). The depopulation of marginal areas includes the abandonment of extensively used agricultural areas, the discontinuation of traditional forms of land use, e.g., dehesas (wooded pastureland) and a decrease in livestock grazing (Delgado et al. 2010; Rescia et al. 2010). Moreover, extensive active afforestation measures resulted in an increase in forested areas (Valbuena-Carabaña et al. 2010).

Despite their relative levels of aridity, drylands contain a great variety of biodiversity, much of which is highly adapted to water-limited conditions. As a result, there are many animal and plant species and habitats found only in drylands: some semi-arid and dry sub-humid areas are among the most biodiverse regions in the world (Gudka et al. 2014). Diversity is also high in drylands, for example between ecotones, areas of different aridity, temperature or altitude. Species have adapted to these factors in many unique ways, creating a variety of habitats that are essential to the survival of species as well as to the livelihoods of people.



**Figure 4.12 | Change in the percentage of land-use change from 1961 to present in relation to desertification and land degradation.** Dryland areas were estimated using TerraClimate precipitation and potential evapotranspiration (1980–2015) to identify areas where the Aridity Index is below 0.65. Population data are from the HYDE3.2 database. Areas in drought are based on the 12-month accumulation Global Precipitation Climatology Centre Drought Index. The inland wetland extent (including peatlands) is based on aggregated data from more than 2,000 time series that report changes in local wetland area over time (IPCC 2019).

Steppes occupy important areas (630,000 km<sup>2</sup>) of the arid zones of the Mediterranean Basin, from the Red Sea to southeastern Spain (Maestre and Cortina 2005). Subjected to strong human impact for millennia, they have been greatly modified. Moreover, some of them, especially the graminean steppe of *Stipa tenacissima*, are the consequence of the degradation of the former open forests. In the past, particularly at the end of the last glacial episode and during the following glacial-interglacial transition, steppes were important biomes representing a significant part of the global continental carbon reservoir.

Shrubland ecosystems account for a substantial part of total land cover and are particularly relevant in arid, semi-arid, and dry-subhumid areas (Reynolds et al. 2007). Shrub-dominated ecosystems are increasing worldwide, a process with important implications for the structure and functioning of terrestrial ecosystems (Van Auken 2000; Berlow et al. 2002; Anthelme et al. 2007). In shrublands, the dominant maquis has many local names reflecting indigenous and local knowl-

edge, such as macchia in Italy, matorral in Spain, phryganae in Greece or bartha in Israel. It is characterized by hard-leaved shrubby evergreen species of genera *Cistus*, *Erica*, *Genista*, *Juniperus*, *Myrtus*, *Phillyrea* and *Pistacia*. The term “garrigue” is restricted to the limestone, semi-arid, lowland and coastal regions of the basin and is maintained by grazing and fires.

Biological soil crusts are complex topsoil microbial assemblages composed of eukaryotic algae, cyanobacteria, mosses, liverworts, fungi and lichens (Velasco Ayuso et al. 2017). They cover the uppermost mm of the soil surface in most arid and semi-arid ecosystems throughout the globe and are one of the most conspicuous and important biotic components of these areas (Belnap and Lange 2013). They exert a strong influence on key ecosystem processes such as runoff (Alexander and Calvo 1990; Belnap 2006), soil respiration (Maestre and Cortina 2003), nitrogen fixation and transformations (Belnap 2002; Castillo-Monroy et al. 2010), establishment and performance of vascular plants (Defalco et al. 2001; Escudero et al. 2007) and act as habitats for a dependent food web of arthropods, fungi, bacteria, and other soil organisms (Belnap and Lange 2013).

Dryland biodiversity also provides significant global economic values through the provision of ecosystem services and biodiversity products. Many cultivated plants and livestock breeds originate in drylands, providing a genetic reservoir whose importance is increasing as climate change drives the demand for new adaptations and extinctions of wild breeds. These services, such as cultural identity and spirituality are central to dryland cultures and can be integral to the protection of dryland ecosystems. There has been an observable correlation between land degradation and cultural degradation in drylands demonstrating their interconnectedness (Davies et al. 2012).

### Agroecosystems

Agroecosystems support high levels of biodiversity and then a rich diversity of habitats and landscapes because of traditional, low-intensity and diverse agricultural systems (Levers et al. 2016). However, this biodiversity has declined dramatically since the early 1950s due to the intensification of agriculture, leading to an increase in highly modified agroecosystems and simplified and agricultural landscapes (Poláková et al. 2011). The common farmland bird index indicates a reduction in agricultural biodiversity by 34% over the time period

1989-2016<sup>36</sup>. In agricultural landscapes, intensification of agricultural systems has generally induced decreased crop diversity, decreased coverage of natural and semi-natural areas (hedgerows, isolated trees, ponds, permanent grasslands) and lower connectivity between the remaining natural and semi-natural habitats (Stoate et al. 2001, 2009).

Agroecosystems provide important ecosystem services to society, but these are threatened by agricultural abandonment and intensification of agricultural practices. These threaten multifunctional landscapes and erode the capacity to deliver ecosystem services, particularly regulating, and cultural ecosystem services (Nieto-Romero et al. 2014; Balzan et al. 2020). Despite the increasing availability of literature about the topic, there are disparities in the availability of research about agroecosystem services within the Mediterranean region, in particular north-south trends. The limited availability of social research on the topic was identified in a review of Nieto-Romero et al. (2014). Most scientific studies focus on provisioning ecosystem services from intensely managed agroecosystems, whilst regulating and cultural ecosystem services were primarily studied in extensive agroecosystems (Nieto-Romero et al. 2014). These results are supported by recent literature, and a parallel can be drawn with the land-sparing vs. land-sharing debate (Phalan et al. 2011). Intensive agricultural districts, characterized by high landscape homogeneity, were shown to provide food products but are relatively poor in terms of capacity to deliver other services in Barcelona (Baró et al. 2017). Extensive agriculture and semi-natural habitats in Malta and in Sardinia were associated with ecosystem service synergies indicating high landscape multifunctionality (Bagella et al. 2013; Balzan et al. 2018).

Approaches that maintain farmland biodiversity have been linked with an improvement in the delivery of regulation ecosystem services in Mediterranean climates. There is evidence that habitat management through the provision of non-crop plant resources (e.g., floral) and conditions can contribute to increased abundance and diversity of natural enemies, biological control and suppression of crop pests when compared to lower biodiversity controls (Shackelford et al. 2017).

#### **Agroecosystem development in different regions**

The ongoing changes in Mediterranean agricultural ecosystems are driven by the dynamics of

the global market of food, energy and technology (e.g., seeds, feeds, fertilizers and agrochemicals) and by regional societal changes (Debolini et al. 2018). A summary of these dynamics is necessary to understand and frame the ongoing changes of Mediterranean terrestrial ecosystems associated with agricultural systems (*Section 3.2.1.1*).

Agricultural systems of northern and southern Mediterranean countries face contrasting challenges in relation to their diverse historical backgrounds and ecological constraints. On one side, particularly in southern Europe, farm abandonment in marginal land is associated with the bias introduced by the implementation of the Common Agricultural Policy (CAP) and post-socialism dynamics (Lasanta et al. 2017). Both external (migration, socio-economic model, public policies) and internal (local factors and characteristics of the agricultural holdings) factors trigger and control the land abandonment process, respectively. Biophysical and socio-economic drivers are interlinked, and the outcomes are therefore very site-specific, depending on local contexts. Abandonment of agricultural activities has many landscape, ecological and socio-economic implications particularly during the transition process, which is still under way in many mountainous areas of Mediterranean Europe (Sirami et al. 2010; Alary et al. 2019). In Mediterranean Europe, the agricultural subsidies related to the CAP and the agro-environmental measures had contradictory effects on land abandonment over the years, which in the long run resulted in a sharp decrease in the number of farms and the increase in average farm size (Papadopoulos 2015; Lowder et al. 2016). In contrast, between 1960 and 2000, average farm size in North Africa decreased, becoming less than 2 ha in around 70% of the farms operating less than 10% of farmland, as 50% of the land is farmed by holdings above 10 ha in size. This is consistent with a global trend of farm size reduction in low- and middle-income countries (Lowder et al. 2016).

The contrasting evolution of agriculture between the northern and southern Mediterranean shores indicates the profound differences in socio-economic and biophysical conditions driving agroecosystem change. While in northern Mediterranean countries abandonment of mountain and marginal land and intensification of lowland and coastal areas is clear, in southern countries there is still high pressure from agricultural and grazing systems on lands that are vulnerable to land degradation and desertification. In northern countries

<sup>36</sup> European wild bird indicators, 2018 update: <https://pecbms.info/european-wild-bird-indicators-2018-update/>



the role of grazing livestock is strategic to mitigate the negative impacts of abandoned farmland (e.g., wildfire prevention). In southern countries, overgrazing is still a core issue (Lasanta et al. 2015) with important impacts on biodiversity (Plieninger et al. 2013) and related ecosystem services (Hurni et al. 2015).

In many marginal agricultural areas of Mediterranean countries, particularly in the north west, the abandonment of agriculture and livestock activities and the consequent forest transition is leading to the rapid expansion of wild fauna (e.g., wild boars, wolves, wild dogs) which is in turn negatively impacting farming (e.g., increased production costs and lower competitiveness of the agricultural business) (Otero et al. 2015), and soil degradation (Mauri et al. 2019). The conservation of biodiversity and ecosystem services can emerge from the implementation of adaptive management approaches, including monitoring of population dynamics and related environmental indices (Katona and Coetsee 2019). Abandonment is also generating a loss of plant biodiversity and cultural landscapes associated with grasslands and farmland fields (Malavasi et al. 2018) and the loss of valuable plant and animal germplasm selected over centuries for their adaptive capacity to these marginal lands (Bullitta et al. 2017), that is not of interest for intensive farmland and hence is at risk of loss. The encroached abandoned croplands and grasslands become particularly vulnerable to wildfires, particularly during the transition from grassland to forest, which in Mediterranean countries is represented by pyrophilous shrubby vegetation, particularly in oligotrophic soils (López-Poma et al. 2014; Bagella et al. 2017). In southern Mediterranean countries, overgrazing still prevails with impacts on soil degradation (Martínez-Valderrama et al. 2018) that are compensated by the increasing import of feed for animal food supplementation, which has doubled in Northern African countries in the past two decades (FAO 2017).

A key issue related to the ongoing changes to Mediterranean agriculture is the impact of these changes on ecosystem water resources and the related hydrological cycle (Milano et al. 2013; Martínez-Valderrama et al. 2018). In silvopastoral ecosystems, the transition from grass to woody vegetation exacerbates the negative effects of increasingly frequent drought events and extreme heatwaves associated with ongoing climate changes (Rolo and Moreno 2019). Deep-rooted tall evergreen trees increase actual evapotranspiration beyond the expected increase of reference evapotranspiration due to increased temperatures. Land use abandonment

therefore results in the loss of ground and surface water resources, which is expected just when more water is needed both for civil and agricultural uses (García-Ruiz and Lana-Renault 2011).

Agriculture absorbs 80% and 60% of total water demand in African and European countries surrounding the Mediterranean Sea, respectively. Under business-as-usual trends, this demand is expected to rise as a consequence of temperature rise and higher drought frequency, resulting in higher evapotranspiration, while at the same time, groundwater recharge and runoff are expected to be reduced as a consequence of the altered water balance due the above-mentioned land use changes in northern Mediterranean countries (García et al. 2017).

The intensification of agricultural activities in lowland and coastal lands is also impacting biodiversity and ecosystem services as is the abandonment of marginal land. The impact of such intensification processes goes beyond provisioning services and the impact on agricultural biodiversity and multiple regulating and cultural ecosystem services is one of the main focuses of the CAP reform debate in Europe (Nieto-Romero et al. 2014). In the following paragraphs we describe the dynamics and drivers of different Mediterranean agroecosystems to understand the implications for biodiversity and ecosystem services.

### **Perennial crops**

In 2017, over 80% of the 10 Mha of olive harvested area in Mediterranean countries was located in Spain (25%), Tunisia (17%), Italy (13%), Morocco (10%), Greece (9%), and Turkey (8%). The harvested area is steadily increasing at a rate of some 140 kha yr<sup>-1</sup> because of the area increments in the MENA and North African countries, where many new plantations are increasing their productivity under introduced irrigation. However, the sustainability of such irrigated croplands is sometimes questioned in the arid lands of North Africa by the use of non-renewable deep groundwater and the high cost of non-conventional treated wastewater or seawater desalinization (Mualla 2018). Crop yield of perennial crops is instead stable and relatively high in European Mediterranean countries (Tanasijevec et al. 2014). In the case of grapes, the harvested area has declined from 200 kha to 150 kha in the past 2 to 3 decades, but production is stable as crop yield increased from 7.5 t ha<sup>-1</sup> in the 1990s to some 8.0-8.5 t ha<sup>-1</sup> in recent years, again as a consequence of the improvement of agronomic techniques and the use of irrigation (data from FAOSTAT).

Olives and vineyards are a fundamental part of the agricultural landscape and cultural heritage of Mediterranean croplands. In the traditional cropping systems, often based on some sort of agroforestry systems or, sometimes, agro-silvo-pastoral systems, are designed for a mix of provisioning services (food, wine, cork etc.), but they also provide unique habitats for agrobiodiversity and contribute to multiple ecosystem services (Cohen et al. 2015; Brambilla et al. 2017; Assandri et al. 2018). Almonds and other traditional fruit Mediterranean orchards, and agroforestry systems in many cases represent a traditional and cultural landscape (Moreno et al. 2018).

### Vegetables

The production of fresh vegetables is increasing in some Northern African and Western Asian countries, particularly Egypt, Algeria, Israel and Turkey. In Egypt the area harvested has doubled in the last 10 years and now represents over 20% of the total harvestable area of fresh vegetables in the Mediterranean area, with just over 140 kha, slightly higher than Italy, traditionally the first country in the Mediterranean. In all other southern European countries, the harvestable area of fresh vegetables has remained stable during the past three decades. The cultivation of vegetables is related to a wide range of farming systems, ranging from very small family farms for subsistence, mainly in the northern African and Near East countries, to very well-organized industrial horticulture value chains. An extreme example of industrial vegetable production is that of Almeria, in Southern Spain, where some 30-40 kha of greenhouses for vegetables and ornamental plants in a very arid area (200 mm yr<sup>-1</sup> rain) are producing a gross value of some €1.5-2.0 billion, 75% is generated through the export of fresh vegetables, primarily to northern Europe. These systems were developed relatively recently (the first greenhouse in Almeria was built in the 1960s) and rely on groundwater (80%), with potential overexploitation and salinization of aquifers under way (Custodio et al. 2016). However, a novel bioeconomy model is being developed in Almeria, to increase its sustainability (Egea et al. 2018). Such systems are increasingly growing in other countries, pushed by the demand for out-of-seasons vegetables across Europe, which is sometimes considered more sustainable than domestic production (Tobarra et al. 2018).

Intensive vegetable cropping systems increase the supply of provisioning ecosystem services but impact biodiversity and may lead to trade-offs with other regulating and cultural ecosystem

services (Balzan et al. 2020). For example, the introduction of irrigation in arid and semi-arid agroecosystems generates a deep transformation of habitat, species composition and related ecosystem services. The mismanagement of irrigation can lead to soil salinization and impacts on agricultural biodiversity (De Frutos et al. 2015; Juárez-Escario et al. 2017). Intensive production sometimes includes the intensive use of agrochemicals with almost total control of weeds, pests and diseases. Furthermore, the industry includes investments in the development of new varieties characterized by tolerance or resistance to biotic and abiotic stress, reduction of harvesting costs, adaptation to long shelf-life and post harvest packaging, which in practice are reducing the diversity of varieties being grown. On the other hand, old varieties are often more suitable for organic farming systems and can provide valuable germplasm for future needs. Small holders therefore still represent a residual source of valuable germplasm that is at risk of extinction and deserves political attention. This is particularly true for Mediterranean germplasm that had been selected by farmers over centuries. Such farming systems therefore provide a valuable ecosystem service in terms of germplasm in situ conservation which is often linked to the cultural values of the traditional rural societies of the Mediterranean Basin.

### Winter cereals

In Mediterranean countries, winter cereals often cover more than 50% of the arable land. Their impact on agroecosystems and ecosystem services is therefore very relevant. However, the winter cereals harvested area in the Mediterranean is generally declining, particularly in southern Europe (e.g., Italy) and is increasing in MENA countries (e.g., Egypt). In contrast, the grain yield is steadily increasing almost everywhere at an average rate ranging from less than +20 kg ha<sup>-1</sup> yr<sup>-1</sup> in North African countries to +40 to 60 kg ha<sup>-1</sup> yr<sup>-1</sup> in southern and eastern Europe (FAOSTAT). Schils et al. (2018) have shown that the yield gap between actual and water limited yield potential for wheat is relatively low in central western Europe and is increasing in Mediterranean countries and eastern Europe, where crop stresses other than just water are still limiting actual yield. This decline in harvested area and increase in yield indicates that, in the past, winter cereals were grown on marginal land. This is certainly the case of EU countries where the CAP subsidies were coupled with winter cereal crops until the CAP reform in early 2000, with farmers also “growing the subsidies” in unsuitable areas (Balkhausen et al. 2007).

In Mediterranean agroecosystems, winter cereals are sown between early and late autumn and harvested in early summer. This guarantees soil cover and protection from erosion in winter and spring but as most winter cereal fields are tilled, soils are exposed to water erosion during the early stages of the crop, corresponding to the heavy rains that are frequent in the Mediterranean climate at the start of the season. Furthermore, with intensive crops the capacity of cereal seedlings in the early growth stages to uptake nitrate nitrogen is low, hence either nitrates derived from the natural mineralization of organic matter or from mineral fertilizers distributed before seeding can contribute to the contamination of groundwater. These processes are expected to increase due to increased temperatures leading to higher mineralization rates, and the higher frequency of heavy storms.

### **Grasslands and grazing systems**

Mediterranean grasslands of the “old world” cover over 1 billion ha of land, mostly in the MENA regions. They host some 240 million dairy and meat sheep, 100 million dairy goats and 95 million beef cattle, mostly based on livestock-cereal, agricultural and agro-silvopastoral systems (Porqueddu et al. 2016). Large-scale grazing systems in southern Europe have almost completely abandoned the traditional transhumant system and only few short-distance vertical movements between lowland and upland pastures are maintained (Caballero et al. 2011). Large-scale grazing systems in rangelands and common grasslands are always associated with specific grazing institutions, regulating grazing management and different arrangements between landowners and pastoralists. Such arrangements shape the cohesion of the local rural society, thus generating complex relationships between biophysical and socio-economic processes leading to more or less desirable outcomes at environmental and social scales (Caballero et al. 2011). In MENA countries, the grazing systems are still shaped by such dynamics and overgrazing is among the main drivers impacting land degradation and desertification, with site-specific issues that call for the development of integrated policy implementation frameworks (Middleton 2018).

In Mediterranean countries, livestock grazing systems are often well integrated with winter cereal cropping systems. Grazing can stimulate tillering and hence contribute to increasing the number of cereal heads per unit area. This practice is coupled with early seeding, which can contribute to preventing soil erosion and nitrate leaching. Graz-

ing is suspended before heading and resumed after grain harvest, when grains losses and straw greatly contribute to animal feeding.

### **Ecosystem services related to pollination**

The decline of pollinators is largely seen across Europe (Biesmeijer et al. 2006; Potts et al. 2010; IPBES 2016), but it strongly contrasts with the steadily growing demand for pollination in crop production (Klein et al. 2006; Aizen and Harder 2009; Garibaldi et al. 2013; Breeze et al. 2014; IPBES 2016). Over the last five decades, agriculture has become increasingly pollinator-dependent, with a three-fold increase in the number of crops requiring the intervention of pollinators (Aizen and Harder 2009). The recommended number of honeybees and hives required to provide crop pollination (by considering the natural presence and action of wild pollinators) across 41 European countries rose 4.9 times faster than honeybee stocks between 2005 and 2010 (Garibaldi et al. 2013). As a result, 90% of the demand for honeybee stocks is not met in 22 out of the 41 countries studied (Breeze et al. 2014). The Mediterranean climate zone has the highest bee species richness in Europe, with the Iberian, Italian and Balkan peninsulas being the most important areas of species richness. Southern Europe also has the highest concentration of endemism, and threatened species. The high diversity of bees in the Mediterranean region is a consequence of the climate of the region and the associated resource heterogeneity, which provide optimal conditions for bee diversity. Petanidou et al. (2008) provide evidence of high temporal plasticity in species composition and interaction identity, indicating that even flower visitation networks show high temporal variation. The main threat to European bees is habitat loss as a result of agriculture intensification and urban development, increased frequency of fires and climate change (Nieto et al. 2015). Fires considerably change vegetation and land cover conditions, and can therefore have an important effect on pollinators and plant pollination (IPBES 2016). For example, fires in Mediterranean oak-pine forests lead to an initial reduction of bee diversity in recently burnt areas. However, these areas recover in the following years, and this recovery is highly correlated to floral diversity (Potts et al. 2003). In Europe, 179 non-threatened species and two threatened species are regarded as under threat from an increased susceptibility to fire, whilst 113 non-threatened species and 23 threatened species are regarded as threatened by climate change (Nieto et al. 2015).

Traditional and non-intensive agricultural practices have a positive impact on agricultural biodiver-

sity in the Mediterranean region (Sokos et al. 2013; Balzan et al. 2020). Similarly, several studies indicate a positive effect of diversified farming systems and organic management related to conventional monocultures (Kennedy et al. 2013). In the meta-analysis by Kennedy et al. (2013), Mediterranean organic fields were estimated to harbor 68% and 56% higher bee abundance and species richness respectively when compared to conventional fields. This study also recorded a significant positive effect of landscape composition, with average increases of 129% and 41% in bee abundance and richness, respectively, for each 0.1 unit increase in the Lonsdorf Landscape Index (an ecologically scaled index of landscape composition) (Kennedy et al. 2013). The Middle East and Mediterranean Europe recorded higher monetary benefits in crop production that is directly linked with pollination services in comparison to other regions. This is mainly due to the cultivation of a variety of fruit and seed crops (IPBES 2016).

### Freshwater ecosystems

Freshwater ecosystems, including streams, rivers, lakes, riparian areas and terrestrial wetlands, offer many important ecosystem services such as water supply for drinking, agriculture and industries (Brauman et al. 2007), water purification, erosion control (MEA 2005; de Groot et al. 2010), recreation, tourism and flood mitigation (Mediterranean Wetlands Observatory 2018). Humans have used these services for thousands of years, and in the process, have severely degraded these ecosystems (Zaimes and Emmanouloudis 2012; Geijzendorffer et al. 2019b). This is particularly true for the Mediterranean region that has been inhabited for thousands of years. The region is characterized by limited water resources and strong population growth (+70% increase in population since 1970 (UN 2013) and +30% in the last 20 years (Abis 2006). Furthermore, substantial increases in seasonal tourism are forecasted in many Mediterranean countries (Burak et al. 2004; Gober 2010), particularly in coastal regions, which can triple in population during the summer (Abis 2006; Collet et al. 2014). Overall, water demand has doubled in the second half of the 20th century in the Mediterranean (Blinda and Thivet 2009; Collet et al. 2013). Accelerated population growth, tourism and globalization are expected to further exacerbate agriculture, urbanization and subsequent pressures leading to an increase in water demand and to significant changes in water use patterns, thus affecting surface waters in the decades to come (Sala et al. 2000; Ferreira et al. 2019; Mack et al. 2019).


### River regulation

The highly irregular rainfall patterns and strong seasonal and annual variability of the flow regimes of Mediterranean rivers and streams (Garofano-Gomez et al. 2011), along with the high topographic relief of many of its river (Grantham et al. 2013) have led to the building of more than 3,500 dams in rivers during the 20th century (Cuttelod et al. 2009; Lobera et al. 2016). Large dams regulate river flow hydrology and influence water chemistry, sediment dynamics, channel form and biotic communities, act as barriers to sediments, fish migration and vegetation propagules (Brierley and Fryirs 2005; Charlton 2008), change the thermal regime, water quality and biogeochemical fluxes, thus impacting habitat availability and connectivity along the fluvial continuum (Van Steeter and Pitlick 1998; Gasith and Resh 1999; Brierley and Fryirs 2005; Nilsson et al. 2005; Garde 2006; Garofano-Gomez et al. 2011; Bernal et al. 2013; Bonada and Resh 2013; Mediterranean Wetlands Observatory 2018). Reservoirs can reduce the sediment load up to 90% and change the flow from a flashy Mediterranean river to a more constant flow regime below the dam. Downstream, the main consequences of water with reduced sediment supply from upstream include: i) river channel degradation (e.g., bed incision), coarsening of the surface layer and channel narrowing; ii) ecological degradation, damaging the availability and quality of habitat for both the aquatic and riparian biota; and iii) reduction of the sediment supply to the development of the river delta and hence accelerated coastal erosion (Kondolf 1997; Liébault and Piégay 2001; Simon and Rinaldi 2006; Vericat and Batalla 2006; Gendaszek et al. 2012; Lobera et al. 2016). The decline in river sediment inputs can be the result of human activities such as the stabilization of mountain slopes because of rural agriculture decline, rural exodus, reforestation and engineered torrent control (Provansal et al. 2014).

When considering all rivers, the total quantity of freshwater discharged into the Mediterranean each year (not including precipitation) has declined by about 45% during the 20th century (*Section 3.1.3*). The reduction in river flows is a probable cause of the very unfavorable conservation status of the biodiversity dependent on rivers: 40% of the fish species found in Mediterranean wetlands are endangered (Mediterranean Wetlands Observatory 2018; *Fig. 4.13*).

### Groundwater depletion

The close connection between streams and aquifers is responsible for base flow during periods of scarce recharge, controlling stream discharge as



**Figure 4.13 | Reduction in freshwater discharge flows into the Mediterranean for all rivers** (Mediterranean Wetlands Observatory, 2018).

well as other hydromorphological characteristics (Winter 1999; Woessner 2000; Menció and Mas-Pla 2010). Human activities, such as groundwater withdrawal or major changes in land cover primarily for agriculture, can result in a reduction of subsurface inflow to streams, changes in groundwater dynamics, and thus, loss of biological quality (Benejam et al. 2008; Menció and Mas-Pla 2010). The needs for water abstraction have risen and it has become difficult to meet water needs (Qadir et al. 2007; Collet et al. 2013; Mediterranean Wetlands Observatory 2018), which is also disrupting the functioning of surface irrigation (EEA 2018; Mack et al. 2019). This is particularly true for Mediterranean islands that are largely dependent on groundwater resources (MED-EUWI 2007; Koutroulis et al. 2013). Exploitation of water resources causes changes in the water balance which leads to low or zero flows, especially during the summer, but also decreases surface water quality (Baldock et al. 2000; Moustadraf et al. 2008; Menció and Mas-Pla 2010).

### Hydrologic regimes

Rivers and streams in Mediterranean areas are subject to naturally occurring high variability in their flow, with predictable seasonal disturbances such as floods and droughts (Menció and Mas-Pla 2010; Zaimes et al. 2010) (Section 3.1.3). They can experience wet winters and consequent floods to severe droughts in the summer, when intermittency in otherwise perennial systems can occur (Cid et al. 2017) (Section 2.2.5.3). In addition, Mediterranean regions are often rugged, marked by a notable altitudinal gradient between the headwaters and the outlet (Emmanouloudis et al. 2011). Mediterranean streams, located in

high elevated areas experience annual rainfall exceeding 1,000 mm, and are characterized by low temperatures in winter with the chance of snow accumulation. This creates a typically bimodal pattern in the flow regime, with the highest discharge following the onset of rain and following snowmelt in spring (Sabater et al. 1992), but maintaining a permanent flow throughout the year. In contrast, rivers located in semi-arid areas (low land areas), with mean annual precipitation ranging from 200 to 500 mm, show a less permanent flow regime (many are intermittent and ephemeral) (Lobera et al. 2016).

Human competition for water enhances the natural deficit in water resources of the region, due to mean annual precipitation lower than the mean potential evapotranspiration (Gasith and Resh 1999). Additionally, water diversion, damming, flow regulation, increased salinity, pollution and introduced species have also severely impacted Mediterranean freshwater ecosystems over time (Moyle 1995; Gasith and Resh 1999; Aguiar and Ferreira 2005; Hooke 2006). Overall, the discharge has decreased almost by half in many Mediterranean catchments in the second half of the 20th century (García-Ruiz et al. 2011; Mediterranean Wetlands Observatory 2018). Mediterranean river ecosystems also have a highly endangered biodiversity, with 40% of the fish species being endangered (Mediterranean Wetlands Observatory 2018), which cannot be dissociated from the long history of human disturbances (Zeder 2008; Feio et al. 2014).

Finally, while we are seeing decreases in water availability and runoff, urbanization and infrastructure have encroached river floodplains leading to higher exposure of both people and capital risk to flood damage (Geijzendorffer et al. 2019b). With climate change leading to more unpredictable flash floods, especially in intermittent and ephemeral torrents, the risk and potential damage of river floods has significantly increased in the Mediterranean Basin (Section 3.1.3.3).

### Land-use changes, reduction of wetlands and riparian areas

In recent decades, coastal Mediterranean wetlands have suffered considerable pressures from land use change, intensification of urban growth, increasing tourism infrastructure and intensification of agricultural practices (Sanchez et al. 2015). The recent Ramsar Global Wetland Outlook (Ramsar Convention on Wetlands 2018) highlights that the Mediterranean region, where water shortages coincide with demography, had total wetland losses

of 48% between 1970 and 2013 (significantly higher than other regions). In addition, 36% of assessed wetland-dependent animals in the Mediterranean are threatened with global extinction (Mediterranean Wetlands Observatory 2018). Special attention should be given to temporary aquatic habitats that are characteristics of the Mediterranean region that provide flood control, groundwater recharge, toxin removal and recycling of nutrients (Balzan et al. 2019). Some temporary wetlands in the Mediterranean region are a priority habitat under the Natura 2000 Network (Natura code 3170, 92 / 43 / CEE, 21 May 1992) (Waterkeyn et al. 2010). Land use intensification in/and adjacent to rivers and streams has eliminated or simplified riparian the structural diversity of ecosystems (Robinson et al. 2002; Corbacho et al. 2003; Kingsford and Thomas 2004). These ecosystems have substantial fragmentation in the lowlands of Mediterranean areas primarily due to agriculture, compared to the mountainous areas that can have detrimental effects on their functionality (Zaimes et al. 2011). The maintenance and re-establishment of riparian ecosystems is a difficult but also an important task in southern Europe where most riparian ecosystems have experienced an extensive history of intensive land-use changes and other human disturbances (Corbacho et al. 2003; Zaimes et al. 2010).

The most important parameters for riparian vegetation were the distance from dams, the sea and rivers (Zaimes et al. 2019). Overall, the riparian vegetation of a Mediterranean Basin decreased with increasing drought, flow regulation and agriculture (González et al. 2010; Bruno et al. 2016; Aguiar et al. 2018). Agriculture is the most important stressor for riparian functionality in the Mediterranean. Agricultural land use and hydro-morphological alteration intensification increases in Mediterranean and semi-arid areas (Nilsson and Berggren 2000; Allan 2004; Bruno et al. 2014a) led to a general decrease in both richness and ecological condition. Agricultural land use was the main pressure explaining riparian richness and quality, whereas the responses of aquatic communities were highly related to hydromorphological alteration. These basin-wide variables had a greater effect than variables operating on a local scale (Bruno et al. 2014b).

The riparian sites with the worst quality were near the river mouth and were characterized by an artificial and highly variable flow regime (Zaimes et al. 2011). This artificial flow variability as well as the presence of lateral structures in the river channel and geomorphological characteristics

were the main factors driving the hydromorphological and floristic pattern in the regulated river. This flow-biota interaction is remarkable in Mediterranean rivers (Prenda et al. 2006) due to their high biological diversity and extremely variable flow regimes (Blondel and Aronson 1999; Naiman et al. 2008; Zaimes et al. 2010; Garófano Gómez 2013). Many native species of riparian vegetation exhibit life cycles adapted to seasonal peak flows, the loss of which may hinder the regeneration of these riparian communities, reducing their growth rates or favoring the invasion of alien species (Poff et al. 1997). Lateral connectivity is also altered by the reduction of the frequency, magnitude and duration of events that periodically flood banks and floodplains (Charlton 2008), causing loss of native riparian vegetation (Burch et al. 1987; Garófano-Gomez et al. 2011; Zaimes et al. 2019).

In conclusion, the loss of natural wetlands is a major concern, since their loss is nearly irreversible and leads to significant impacts on wetland biodiversity and ecosystem services. Restoration initiatives exist but have a low rate of success when it comes to re-establishing the same richness and stability that can be found in natural wetlands.

#### **Water quality**

Based on the Water Framework Directive (WFD; Directive 2000/60/EC), European Union (EU) countries are obligated to assess the ecological status of their freshwater ecosystems using biological indicators, as well as chemical, hydrochemical, and hydro-morphological parameters and to achieve good qualitative and quantitative status of all ground and surface water bodies (Van den Broeck et al. 2015). Additionally, the new Groundwater Directive (GD; Directive 2006/118/EC) considers groundwater as a valuable natural resource that should be protected from deterioration and chemical pollution (Menció and Mas-Pla 2010). However, a concern is that the WFD programs do not incorporate assessment techniques for temporary wetlands (Van den Broeck et al. 2015) whilst the links between water quality and ecosystem functions and services, and the implications of water management on ecosystem services are either implicit or overlooked (Acreman et al. 2017). Outside of the EU, data on water quantity and quality are sparse and often biased. This is a real problem, where countries can affect both water quality and quantity flowing downstream to another country, as is the case for some rivers in the eastern part of the Mediterranean Basin.

The global Sustainable Development of Agenda has included water as an important priority, and SDG6

emphasizes safe access to water and sanitation. Water quality is considered as a major environmental problem across the Mediterranean region (Table 3.4, Section 3.1.3.5), with recent assessment indicating that the WFD has improved water quality in the European countries of the Mediterranean Basin while water quality has degraded further in North Africa and the Middle East (Mediterranean Wetlands Observatory 2018).

### Freshwater species

Declines in the Living Planet Index (LPI) of Mediterranean wetlands was continuously observed between 1990 and 2008, after which it increased, but with varying results depending on the group and sub-region. Between 1990 and 2013 waterbird numbers show a positive trend and increased in Western Europe (+101%) and Northern Africa. However, more moderate increases were observed in the eastern Mediterranean (+27%) while declines have been observed in the Middle East since 2008. Contrastingly, declines in amphibians, reptiles, mammals and fish have been observed since 1990 (Mediterranean Wetlands Observatory 2018).

Freshwater communities of the Mediterranean region have adapted to the natural variability in water flows through shorter life spans, mechanisms to resist or avoid desiccation, and higher colonization rates (Lytle and Poff 2004; Bonada et al. 2007; Stromberg et al. 2008; Santos 2010). These Mediterranean communities are, therefore, different from those of temperate rivers, showing interannual fluctuations in richness and composition and in trophic structure (Ferreira et al. 2001, 2002; Bonada et al. 2007; Feio et al. 2010, 2014). During dry seasons (predictable and periodical seasonal droughts), groundwater that flows towards streams is highly significant as it represents a unique input for water discharge, leading to stream/river reaches that are permanent, intermittent or ephemeral (Uys and O'Keeffe 1997; Argyroudi et al. 2009). Seasonal droughts can cause habitat loss, poor water quality and biotic interactions, but in severe droughts (longer, unpredictable, seasonal or supra-seasonal droughts) as expected due to climate change, major ecological effects will be observed, stressing and depleting both fauna and flora (Boulton 2003; Lake 2003; Bond et al. 2008; Menció and Mas-Pla 2010).

Mediterranean rivers present rich and dynamic riparian plant communities, which are highly interconnected with lateral and vertical ecotones and have multi-scaled biotic drivers that act in both space and time (Ferreira et al. 2019; Kontsiotis et al. 2019). Natural and human disturbances are

entwined forces that shape riparian plant communities, to the point that undisturbed plant communities are difficult to find or characterize. Though there are few truly aquatic species, Mediterranean riparian plants nonetheless play an important role in stream functions (Zaimes et al. 2010; Magdaleno and Martinez 2014). The protection of these species should be a priority in the region, and many riparian areas are included in the Natura 2000 Network and the Ramsar Convention (Zaimes et al. 2010; Ferreira et al. 2019).

Freshwater ecosystems are under threat from the effects of multiple stressors, including non-indigenous species (Navarro-Ortega et al. 2015). Non-indigenous species are considered in the top five causes of biodiversity loss (Bruno et al. 2019) and result in the accelerated impairment of aquatic and riparian habitats and their ecosystem services worldwide (Saunders et al. 2002; Dudgeon et al. 2006; Van den Broeck et al. 2015; Rouissi et al. 2018; Fraixedas et al. 2019). Their increase and expansion are due to the alteration of their hydrologic regimes, and biological and morphological functionality due to agriculture on the floodplain, channel diversions and dams, and increased pollution (Jiménez-Ruiz and Santín-Montanyá 2016). Non-indigenous species in many cases can tolerate and adapt easier to the new conditions. Exotic species often thrive in Mediterranean rivers altered by human activity, further homogenizing river communities worldwide (Cooper et al. 2013). Alteration of the vegetative structure, competitive displacement of native riparian vegetation, reduction of arthropod and avian diversities and abundances are some of the major impacts of non-indigenous species (Saunders et al. 2002; Herrera and Dudley 2003; Dudgeon et al. 2006; Bruno et al. 2019). Examples of non-indigenous species that are serious threats and problems in Mediterranean riparian areas are the *Robinia pseudoacacia*, *Ailanthus altissima* and *Arundo donax* (Constán-Nava et al. 2015; Bruno et al. 2019; Nadal-Sala et al. 2019). Examples of non-indigenous species that are serious threats and problems in Mediterranean wetlands include *Myriophyllum aquaticum*, *Carpobrotus edulis* and *Cortaderia seloana* (Lastrucci et al. 2018; Chefaoui and Chozas 2019; Company et al. 2019).

### Protected areas (Natura 2000 network and Ramsar Convention)

In the European Union, the importance of conserving and protecting freshwater ecosystems is recognized through the many that have been designated as Natura 2000 sites (Iakovoglou et al. 2013). Wetland protection is also officially a priority

for the 159 nations (as of 2009) that have ratified the Ramsar Convention, although wetlands still continue to be under threat of being drained and reclaimed (Ramsar Convention on Wetlands 2018; Geijzendorffer et al. 2019a). Degradation is closely related to the rapid increase in human population, and the increased input of nutrients, pollutants, and sediments, due to increases in urban development, industry, agricultural activities, and water abstraction. The most obvious effect is the loss of biodiversity as a consequence of a reduction in area and the deterioration in conditions, especially in arid and semiarid regions (Brinson and Malvárez 2002).


### 4.3.2 Projected vulnerabilities and risks

#### 4.3.2.1 Forests

The interactions between different drivers of climate change (CO<sub>2</sub>, warming, reduced rainfall, increase in drought frequency and intensity) are

predicted to have multiple, and sometimes antagonistic effects on the future condition of Mediterranean ecosystems (Bussotti et al. 2014) (Fig. 4.14). While increasing atmospheric CO<sub>2</sub> concentrations might directly promote forest productivity and growth (Sabaté and Gracia 2002; Keenan et al. 2011), this effect will likely be strongly modulated by increasing temperatures and drought conditions (Peñuelas et al. 2011; Bussotti et al. 2014; Doblas-Miranda et al. 2017; Lo et al. 2019). For pine and oak-dominated Mediterranean woodlands in Israel, Helman et al. (2017) projected that warming scenarios of 1 and 2°C could lead to 16% and 31% reductions of annual gross ecosystem productivity, respectively, despite the increase in atmospheric CO<sub>2</sub>.

Warmer and drier conditions also alter plant phenology (i.e., leaf unfolding, flowering and fruiting), usually lengthening the growing season (Peñuelas et al. 2004), with direct consequences on forest productivity and growth (Kramer et al. 2000). Al-



**Figure 4.14 | Tree- and ecosystem-level responses in Mediterranean forests** to environmental changes associated with climate change and their impacts on ecosystem service (ES) provision. Adapted from Figure 1 in Bussotti et al. (2014) and based on the reviews by Doblas-Miranda et al. (2017) and Peñuelas et al. (2017, 2018).



though a longer growing period may be positive for forest growth, advanced spring phenology may also cause higher risk of frost damage (Mutke et al. 2005), as well as increased transpiration. Moreover, global warming also changes bird migrations and dates of insect outbreaks, leading to a decoupling of species interactions (e.g., decoupling of predator-insect interactions reduces the effectiveness of pine processionary moth control by birds; [Barbaro and Battisti 2011]). All factors considered, we can expect a general reduction of site productivity in the mid- and long-term, particularly for species or populations growing in water-limited environments, which constitute the majority of Mediterranean forests (Sabaté and Gracia 2002; Bravo-Oviedo et al. 2010).

### **Changes in forest ecosystem health and ecosystem services provision**

There is evidence that Mediterranean forests and woodlands now experience climate-driven declines in growth and die-back episodes from drought and heat stress (Allen et al. 2010; Lindner et al. 2010; Anderegg et al. 2013; Gentilesca et al. 2017; Klein et al. 2019), similar to shrublands (Lloret et al. 2016; Sapes et al. 2017). Increasing crown defoliation and soil respiration may reduce net primary production and ultimately limit growth in Mediterranean forests, associated with higher mortality and dieback, especially if warming is combined with drought (Peñuelas et al. 2018). The combination of reduced water availability and increased respiration rates of tissues due to rising temperatures can result in hydraulic failure, the exhaustion of reserve carbohydrates and a general weakening of the trees, also making them more vulnerable to pests and pathogens (Rennenberg et al. 2006). Drought has been linked to the general dieback of *Quercus ilex* and *Q. suber* in southwestern Spain, where, known as “seca”, the weakened trees are more susceptible to the attack of the fungus *Phytophthora* (Sánchez-Salguero et al. 2013). Although even drought-adapted ecosystems are influenced by growth reductions, these phenomena are expected to become particularly frequent in the trailing-edge of species distribution (Jump et al. 2006; Sarris et al. 2011), or for species found in the Mediterranean Basin, the southern limit of their European distribution (Linares et al. 2010; Dorman et al. 2013), which are particularly vulnerable (e.g., *Pinus pinaster*, *P. nigra*, *P. halepensis*, *P. sylvestris*, *Quercus ilex*, *Q. suber*, *Fagus sylvatica*, *Abies alba*, *A. pinsapo*, *Juniperus phoenicea*, *Cedrus atlantica*), and especially in dense, unmanaged forests (Lindner and Calama 2013) or in sites with shallow soils (Lloret et al. 2004).

This additional climate stress may lead to important changes in biotic interactions, affecting forest composition and species distribution (see next section). For example, in southern Spain, in a mixed *Abies pinsapo*–*Pinus halepensis* forest, *A. pinsapo* showed sudden growth reductions under drier conditions, while pine trees were able to maintain almost constant growth values and lower water costs under increasing long-term water stress (Linares et al. 2011). Similarly, Sarris et al. (2011) reported that where mixed *Abies cephalonica*–*Pinus halepensis* forests exist in southern Greece, *Pinus* did not experience any mortality at this altitude (800 a.s.l.) after drought events, unlike *Abies*. Prolonged droughts and hot spells will aggravate the risk of forest fires, which can further induce problems of soil erosion and fertility. In fact, fire-drought interactions can be complex and trigger vegetation transitions, disrupting ecosystem resilience and even leading to non-forest states (Battlori et al. 2019). In the driest areas, desertification can advance and become a major problem (Karavani et al. 2018a).

Climate change is also expected to affect host plant-pest interactions, favoring the establishment of new ones (Lindner and Calama 2013). Warming already causes changes in the distribution areas of pests, mostly upward and northward. An example is the pine processionary moth (*Thaumetopoea pityocampa*), which is expanding upwards in several mountain ranges due to milder winter temperatures, affecting tree populations that had previously never been exposed to this insect (Hódar and Zamora 2004; Battisti et al. 2005; Roques et al. 2015). The succession of several years of mild winters has favored unprecedented outbreaks in northeastern Spain (Roques et al. 2015). Opportunistic fungi and insects such as *Armillaria* or *Ips* spp. are also being favored by warmer temperatures, which induces better conditions for survival, allowing them to complete more than one generation in one year (Lindner and Calama 2013). The greatest impacts are expected to arise from the establishment of alien pests and diseases, i.e., those that are exogenous to a given environment. The number of alien pests is expected to increase under warmer and drier conditions, as has been the case with the pine nematode (*Bursaphelenchus xylophilus*), native from North America, and with the potential to spread across Europe (de la Fuente et al. 2018) and cause massive wilt and mortality in pine species (Vicente et al. 2012).

All these changes may ultimately lead to profound changes in ecosystem function and associated ecosystem services (Seidl et al. 2014; Peñuelas

et al. 2017). Changes in carbon storage and water availability are especially important for their implications in all forest services, because they are the basis of the primary production that supports the services (e.g., timber production) and because of the effects they have on climate change (Peñuelas et al. 2017; Ruiz-Peinado et al. 2017). Increased plant evapotranspiration will decrease the movement of water from forest to downstream ecosystems (Peñuelas et al. 2018), compromising supporting services (e.g., water cycle), provision of habitat for aquatic species and water availability for consumptive uses. Severe summer droughts can reduce the yields of economically relevant NWFP such as cork (Oliveira et al. 2016) and pine nuts (Mutke et al. 2005).

The response of some forest ecosystem services to climate change drivers is still under debate. For example, despite the fact that some studies have highlighted that mushroom productivity in Mediterranean ecosystems may be experiencing a sharp drought-induced decrease (Boddy et al. 2014; Ágreda et al. 2015) due to delayed phenology in the autumn season under warmer and drier conditions (Kausarud et al. 2012; Büntgen et al. 2015), simulations by Karavani et al. (2018b) rather point towards an increase in production of edible and marketable species under climate change scenarios as a consequence of the longer mushroom season. The leisure use of Mediterranean pine forests (for walking, mountain biking hunting, etc.) will probably be negatively affected by the increasing incidence of pest outbreaks of the pine processionary moth (Morán-Ordóñez et al. 2019), as this species is responsible of strong allergic reactions in humans (Battisti et al. 2017). However, simulation studies in Mediterranean forests (Mina et al. 2017) suggest that forest management (i.e., silvicultural interventions) might have a prevailing role over climate in determining the future condition of forests and the provision of their associated ecosystem services. This has also been reported in other forest systems across the globe (Albrich et al. 2018; Schwaiger et al. 2019).

Besides the direct impacts of climate change drivers on the condition of tree species and ecosystem services provision, climate change drivers might push Mediterranean forests past critical thresholds (e.g., changes in community composition, loss of ecosystem functions), which could hamper their capacity to recover from disturbances in the future (Anderson-Teixeira et al. 2013). For example, Mediterranean water-stressed forests are likely to become more vulnerable to pests and pathogens (Lindner and Calama 2013; Gauquelin

et al. 2018), as well as to other disturbances such as fire. Post-fire regeneration might be limited under water-limited conditions, ultimately leading to deforestation or transition from oak and pine forest to shrublands (Karavani et al. 2018a), thereby decreasing the overall capacity of the region to sequester atmospheric CO<sub>2</sub> and potentially losing the recreational value of affected areas (Peñuelas et al. 2017).

### **Changes in species range, abundance and extinction**

Climate change is predicted to induce changes in the geographic ranges for many terrestrial species across the Mediterranean Basin (expansion, shrinkage, geographic shifts), with studies showing contrasting predictions depending on the modelling approach, the drivers and the scenarios considered, even when predictions are made for the same species and the same region. For example, on the basis of a process-based model, Keenan et al. (2011), predicted that around 40% of the current suitable stand locations of *Quercus ilex* in Spain will become unsuitable for the species during 2050-2080 under a non-Paris agreement compliance warming scenario (3.1°C) whereas Lloret et al. (2013), predicted an increase in climatic suitability for the same species, region, scenario and time horizon on the basis of a correlative model.

The EU Mediterranean biome was predicted to be the most vulnerable region to plant species loss and turnover in a study by Thuiller et al. (2005), who simulated climatically determined geographic range loss of 1,350 European plant species under seven climate change scenarios (IPCC AR4 SRES scenarios predicting temperature increases ranging from 1.8 to 3.6°C), with climate-related range contractions already reported in Mediterranean mountains (Pauli et al. 2012). Consistent patterns have been forecasted for other taxonomic groups. For example, using bioclimatic envelope models and ensemble forecasting of SRES scenarios, Levinsky et al. (2007) and Barbet-Massin et al. (2012), predicted losses up to 100% and 30% of current potential species richness of mammals and bird species in EU Mediterranean, respectively, for the end of the century. In general terms, species at the rear edge of their distribution in the Mediterranean (e.g., deciduous temperate species like *Quercus petraea*) and mountain species (e.g., *Pinus sylvestris*, *Abies alba*) will be the species most threatened by climate change (with ranges potentially shrinking), whereas the most xeric Mediterranean species, which are better adapted to drought, are those expected to encounter fewer problems for survival

and range expansion under future climate change (Ruiz-Labourdette et al. 2012; Lindner and Calama 2013; Bussotti et al. 2014).

Projections of species range losses due to climate change across the Mediterranean cannot be taken as precise forecasts given the uncertainties in climate change scenarios. Only a few forecasting studies have assessed the interactions of climate change with other drivers (Morán-Ordóñez et al. 2019), there is therefore a risk that the vulnerability of species to other important disturbances, such as land use change, fires and their synergistic effects, is underestimated (IPBES 2019). An additional caveat for studies projecting changes to climatic range is that generally these do not incorporate the role of interactions between species or the effects of extreme weather events, the latter of which is of great relevance in the context of Mediterranean forest systems.

### **Fire activity and burnt areas across the Mediterranean**

The Mediterranean Basin can be considered as a hotspot under future climate conditions conducive to extreme wildfire events, with significant potential impacts for human well-being (Bowman et al. 2017). How exactly climate change will influence future fire regimes is still under debate. While a warmer and drier climate will upsurge fire activity by increasing water demand and decreasing fuel moisture, increasing temperatures may also negatively affect ecosystem productivity and lead to an overall reduction of fuel biomass, which can counteract warming effects on fire activity (Batllori et al. 2013). Drought increases terpene emissions from Mediterranean plants, which are compounds that play a key role in the flammability of forests (Peñuelas et al. 2018). Warming conditions also increase emissions of other volatile organic compounds (VOCs) besides terpenes, with multiple physiological and ecological functions (e.g., plant defense, communication with other organisms) that, in a cascade effect, can affect communities of organisms, ecosystems, atmospheric chemistry and even meteorological conditions, even potentially generating feedbacks to warming (Doblas-Miranda et al. 2017; Peñuelas et al. 2017) (Fig. 4.14).

The increase in exposure to large wildfires in recent years (Bowman et al. 2017), along with the effects of climate change, might still overcome current fire prevention efforts. More and different fire management approaches must therefore be considered in order to increase our resilience

towards future Mediterranean forest fires (Moritz et al. 2014; Turco et al. 2018a). Projections indicate an increase of burned areas across the Mediterranean in the future, but it is difficult to compare estimates given the variation between scenarios, future periods and models used. For example, Amatulli et al. (2013) estimated increases of up to 66 and 140% in burnt area in EU-Mediterranean countries in 2071–2100 relative to 1985–2004 under the IPCC SRES scenarios B2 and A2, respectively, and Migliavacca et al. (2013) estimated a 34% increase in burnt area in southern Europe, in 2070–2100 relative to 1960–1990 under the A1B scenario. Turco et al. (2018b) projected future summer burned area in Mediterranean Europe under 1.5, 2, and 3°C global warming scenarios, concluding that the higher the warming level, the greater the increase in burned area, ranging from a ~40% (1.5°C scenario) to ~100% (3°C scenario) increase from current levels across the scenarios. Although the future total burnt area could be smaller if a stationary relationship between drought and fires is assumed, in all the cases the burned area is still expected to increase with warming. Significant benefits (regarding burnt area reductions) would be obtained if warming were limited to well below 2°C (Turco et al. 2018b). These benefits extend beyond plant cover protection or human safety. A reduction of burned areas also reduces risks of soil erosion and desertification, especially in very dry areas (Shakesby 2011).

#### **4.3.2.2 Mountains**

Many of the key observed and projected climate changes identified for southern Europe by EEA (2017) apply in the case of Mediterranean mountains, including: (i) significant increase in heat extremes, (ii) decrease in precipitation and river flow, (iii) increasing risk of droughts, (iv) increasing risk of biodiversity loss and (v) increasing risk of forest fires. For Mediterranean mountains, projections indicate warming between 1.4°C and 5.1°C for 2055 (1.6°C and 8.3°C for 2085) and a decrease in precipitation, mainly during spring (-17% under A1fi and -4.8% under B1 for 2085) (Nogués-Bravo et al. 2008).

Mediterranean mountain environments seem to be accelerating towards uncertain ecological states because of changes associated with climate and land use changes (Nogués-Bravo et al. 2008). For the 21st century, projected warming and reduced rainfall are likely to affect: (i) snow pack and glaciers, which provide key habitats for alpine specialist species and, (ii) water availability and river discharge and therefore aquatic and wetland

habitats and species. Beyond these indirect effects of climate change in biodiversity, climate change would, (iii) reduce habitat availability of alpine and sub-alpine belts, increasing the risk of extinction for endemic species or range-restricted species and may well disrupt the biological networks that ultimately support ecosystem functioning (Nogués-Bravo et al. 2008).

Mediterranean mountains are susceptible to forest fires and are vulnerable to hydro-geological risks (floods, landslides, infrastructure damage). Most of these hazards will be increased by the predicted rise in temperature and changes in precipitation patterns. Mediterranean mountains provide basic water-based ecosystem services. Therefore, water management and quality-assurance policies need to consider the specific features of mountains and predicted climate change trends. In mountain environments, changes in precipitation (amount and pattern) will be influenced by local geomorphology and therefore predictions are subject to high uncertainty and variation at local and regional scales. Local climate change adaptation strategies require careful consideration in order to counteract specific pressures.

The homogenization of Mediterranean mountain landscapes due to the abandonment of agropastoralism has negative impacts on biodiversity, water resources, soils and natural hazards (Vogiatzakis 2012). For many species in Mediterranean high-altitude zones, including cedars in Cyprus and Lebanon, migrating upwards is not an option (Fernández Calzado et al. 2012). For other species, such as junipers, climate change and increased fires have adverse effects on regeneration, which is already limited by the environment (Vogiatzakis 2012). Fire events are likely to increase in number and intensity in Mediterranean mountain forests and will be associated with elevational shifts of dominant tree species (Fyllas and Troumbis 2009; Pausas et al. 2009). Reduced depth and persistence of snow cover will also affect high mountain vegetation (García-Romero et al. 2010).


#### 4.3.2.3 Drylands and shrublands

There is high confidence in observed drought increases in the Mediterranean and West Africa and medium confidence that anthropogenic climate change has contributed to increased drying in the Mediterranean region and that this tendency will continue to increase under higher levels of global warming (Koutroulis 2019). According to global warming levels of 1.5°C, 2°C and 4°C above pre-industrial temperatures, Mediterranean land

will shift to drier types by 11.6%, 20.1% and 41.3% respectively (Fig.4.15b-d). With a 2°C global temperature rise, annual warming over the world's drylands is expected to reach 3.2°C–4.0°C, implying about 44% more warming over drylands than elsewhere (Huang et al. 2016), thus potentially aggravating water scarcity issues through increased evaporative demand. The Mediterranean, North Africa and the eastern Mediterranean will be particularly vulnerable to water shortages, and expansion of desert terrain and vegetation is predicted to occur in the Mediterranean biome, an unparalleled change in the last 10,000 years (*medium confidence*) (Guiot and Cramer 2016). At 2.5°C–3.5°C, risks are expected to become very high with migration from some drylands resulting as the only adaptation option (*medium confidence*). Scarcity of water for irrigation is expected to increase, in particular in Mediterranean regions, with limited possibilities for adaptation (Haddeland et al. 2014; Malek and Verburg 2018).

Mediterranean drylands are relatively resilient systems with a certain capacity to recover from various forms of disturbance that have occurred for millennia, such as fires, overgrazing and drought (García-Romero et al. 2010). Nevertheless, these ecosystems face critical thresholds with potential catastrophic shifts that may trigger biological diversity losses and modifications in ecosystem functioning and services (Daliakopoulos and Tsanis 2014). The resilience of Mediterranean drylands is currently under pressure from various factors, such as high permanent and seasonal population density, abandonment of traditional practices, continued habitat conversion (2.5% of Mediterranean habitat was lost between 1950 and 1990) and loss of the typically high spatial natural heterogeneity (Fahrig 2003). Although vegetation recovery both from shrubs and forest can reduce soil erosion and enhance carbon sinks, it might change the cultural landscapes frequently developed from initial mosaics of different land-use types towards homogenized states with dense shrubs (Stellmes et al. 2013). Increases in woody biomass and the loss of gaps and breaks as well as edges between different fuel types make these landscapes more vulnerable to fires compared to disconnected patches of forest, shrub and cultivated fields or grasslands (Puigdefábregas and Mendizabal 1998; Viedma et al. 2006; Röder et al. 2008), in particular during drought (Viegas 1998; Duguy et al. 2007).

Since 1960, wildfire occurrence in many Mediterranean drylands has increased because of changes in land use. The level of soil degradation due to these fires depends on fire recurrence, topography, the



**Figure 4.15 | (A) Distribution of drylands and their subtypes based on observations** (reanalysis combined with station and satellite observations) for the 1981-2010 baseline period. Areal coverage (percentage) of drylands per subtype is presented in the form of pie charts and is calculated within the boundaries of the Mediterranean SREX region (dashed line). **(B, C, D) Distribution of projected dryland transitions according to RCP8.5 for three Global Warming Levels** (GWLs: +1.5°C, +2°C and +4°C from preindustrial levels), relative to the baseline (1981-2010) period, using the high-resolution atmosphere-only version of the HadGEM3A model. Grey shaded areas in (b), (c) and (d) are drylands of the baseline period. Chord diagrams denote the areal extent (fraction of MED SREX land area) of projected transitions in each dryland subtype under the three GWLs. The size of the chord diagrams is proportional to the total areal extent of the MED land changing to drier types. Figure adapted from (Koutroulis 2019).

intensity of soil erosion processes and post-fire plant cover regeneration rate (Caon et al. 2014). To promote the accumulation and retention of nutrients in soil after a fire, it is important to stabilize the burnt site by applying post-fire measures that limit soil erosion, surface runoff and loss of the ash due to wind. Depending on the plant species and the time that elapses between consecutive wildfires, fire is responsible for the transition from forests to shrublands, which are poorer in soil nutrient status. High fire frequency may cause the eradication of keystone species, which has consequences for soil nutrient pool recovery (Caon et al. 2014). In addition to the increased risk of soil erosion, frequent wildfires also induce changes in the water cycle by altering the infiltration capacity of soil and increasing soil hydrophobicity (Vallejo and Alloza 1998; Fernández et al. 2012; Carreiras et al. 2014).

### Droughts

Mediterranean shrublands are experiencing important episodes of drought-induced die-back explained by the decrease in climate suitability for plants during the drought event (Lloret et al. 2016; Sapes et al. 2017). In anticipation of a drier climate and to project future changes in dryland dynamics, it is imperative to understand species-specific differences in drought resistance (Väänänen et al. 2020). In the long rainless eastern Mediterranean summer, it was found that the physiological traits of species exhibiting different levels of mortality and co-existing in the same habitat (*Phillyrea latifolia*, *Pistacia lentiscus* and *Quercus calliprinos*) were more associated with drought resistance strategies rather than actual drought stress experienced by the plants (Väänänen et al. 2020). The intensity of drought effects on shrub performance is thus species-specific, and plant species combination-dependent. High shrub richness levels modulate the negative impacts of aggravated drought conditions. Results point to a probable shift in interspecific relationships in response to water shortage. As drought impacts are not mediated in low-diversity communities, species-specific responses to drier conditions could lead to shifts in plant community composition favoring the most drought-resistant species such as oaks and rosemary. Maintaining high diversity appears critical to mediate drought effects for less resistant species (e.g., *Cistus* and *Ulex*) (Rodríguez-Ramírez et al. 2017).

#### 4.3.2.4 Agriculture and pasturelands

The ongoing changes to agricultural systems in the Mediterranean Basin are the outcomes of a

combination of climate and other drivers that influence farmers' perceptions and choices (Fahrig 2003; Nguyen et al. 2016), which in turn trigger changes in agroecosystems and their service provision at different scales. These processes have a strong site-specific component with a common background of multiple pressures including global and climate changes (Kummu et al. 2017).

Mediterranean agriculture is generally vulnerable to shocks in the flow of agricultural commodities, particularly in southern countries, because of the heavy dependence on imports (Capone et al. 2014). This is coupled with expectations of future adverse climate impacts leading to the decrease of water-limited crop yields (Schils et al. 2018), increase in irrigated crop water demand, increasing risks in livestock production and mortality associated with heatwaves, expansion of habitats for southern disease vectors and increases in multiple climate hazards (EEA 2017). These impacts hamper the profitability and the ecosystem service provisioning of agricultural activities, particularly in the most vulnerable situations, such as those of farm enterprises relying on natural and semi-natural resources in marginal land or where farmers have made large investments and are therefore exposed to high capital risks (Dono et al. 2014). The latter is the case, for instance of intensive dairy farming, where lower animal productions caused by an increase in the frequency of heatwaves are not counterbalanced by potential benefits in terms of irrigated forage crop yield that can be expected as a consequence of higher temperatures in winter and/or CO<sub>2</sub> concentration rise (Dono et al. 2016).

Water represents a key factor for risk, vulnerability and the resilience of agriculture at farming system and landscape scales (Iglesias et al. 2007). Extreme drought and flooding are two side effects of the same climate pressure (Iglesias et al. 2007; Quintana-Seguí et al. 2016) and there is evidence that the Mediterranean Basin is highly vulnerable to reductions (up to -49%) in provisioning and regulating ecosystem services associated with agricultural water use and management (Jorda-Capdevila et al. 2018). Dramatic changes in the water balance of Mediterranean watersheds are ongoing, as a consequence of a combination of changes in rainfall regime, temperature rise and the increase in vegetation cover from the abandonment of agricultural and pastoral activities, particularly in northern countries (Serpa et al. 2015; Krause et al. 2016; Zeng et al. 2018), which can also result into increased soil erosion (Bussi et al. 2014).

These dynamics are affecting grassland-based systems (e.g., reduction of water pools for animal drinking) and large catchments based on mountain water resources and hence the availability and stability of groundwater and reservoir stocks to be potentially used for irrigation (Rolo and Moreno 2019). There is also evidence of increased vulnerability of water stocks potentially affecting agricultural systems in southern Europe and northern Africa, more than in eastern countries of the Mediterranean Basin, caused by both ongoing climate and socio-economic dynamics, which in most cases, cannot be totally eliminated by adaptation strategies (Koutroulis et al. 2019).

The expected impacts of climate change on croplands and grasslands are often assessed by modelling under current and future climate scenarios (Moriondo et al. 2010; Koutroulis et al. 2019), sometimes integrated with economic (Dono et al. 2016) and policy change analyses (Cortignani and Dono 2018). Unfortunately, only a few studies consider the combined effects of the other drivers of ongoing changes such as those related to technology development, consumer behavior, energy production etc. (Alexander et al. 2015; Doblaz-Miranda et al. 2017). Differences in geographic, political and environmental context characterize agriculture and pastureland ecosystems of each sub-region of the Mediterranean Basin, resulting in diverse spatial distribution of vulnerabilities and risks (Prosperi et al. 2014).

In southern European countries the impacts of climate pressures (e.g., increased drought) are coupled with the ongoing transformation agroecosystems (e.g., abandonment in marginal lands and intensification of coastland agriculture). Forest wildfires, landslides and depopulation of marginal rural areas cause additional change (Nainggolan et al. 2012), just as water and air pollution in intensively cultivated areas. Overall, production (food or forage) is given priority over regulating or cultural services (Aguilera et al. 2013; Maes et al. 2018), except for the urbanization of agricultural fertile soils (Ceccarelli et al. 2014).

In the Middle East and northern African countries, multiple stressors include climate, groundwater overexploitation, seawater intrusion in coastal areas, water pollution, land degradation and desertification (Sowers et al. 2011; Schilling et al. 2012; Fouchy et al. 2019), impacting social and political stability (de Châtel 2014). Future trends in agricultural and pastoral land use are very much context-sensitive, as predictions of future dynamics are the consequence of multiple drivers beyond climate, which are much more uncertain in north-

ern Africa than for southern European countries (Prestele et al. 2016).

Focusing on an agricultural and pastoral district (some 54,000 ha) characterized by a mosaic of many different crops and land uses in southern Europe, located in the mid-west Mediterranean Basin, Dono et al. (2016) have shown that the same climate and socio-economic pressures can generate a mosaic of different impacts on diverse farming systems even within the same production system. Several factors, including economic farm size, the dependence on external inputs and the availability of water, contributed to a range of situations: rice and vegetable farms were the “winners”, as long as water is not a constraint in the near future, while “losers” were livestock farmers whose feeding system was based on rain-fed grasslands suffering from the increased frequency of extreme drought, hampering hay stock production (e.g. dairy sheep and beef cattle), or those heavily dependent on external inputs, such as the dairy cattle system. Net impacts were associated mainly with the increased frequency of heat waves with a high temperature-humidity index (Bernabucci et al. 2014).

### ***Cropping systems***

The potential higher resilience of irrigated cropping systems to increased drought must be managed in the context of +4 to +18% increased crop water requirements due to climate forcing under different scenarios, involving water resource availability in quantity and quality and water use efficiency, particularly in southern and eastern Mediterranean countries (Fader et al. 2016; Malek and Verburg 2018). At present, the Mediterranean region could save some 35% of irrigation water by implementing more efficient irrigation and water management systems, but southern and eastern sub-regions would need around 35% more water than today in the future, even after the implementation of some degree of modernization of irrigation and conveyance systems, taking into account increased CO<sub>2</sub> fertilization effects (Fader et al. 2016) and the need for supplemental irrigation for winter cereals (Saadi et al. 2015). Mediterranean irrigated croplands include a wide range of vegetable crops, including potato, orchards and grapes, forage crops and, in southern countries, sugar cane and cotton. Most C<sub>3</sub> irrigated crops (e.g., many vegetables and rice) would benefit from increased CO<sub>2</sub> fertilization effects and some C<sub>4</sub> from the increased temperature (e.g., sugar cane and maize) but others might be negatively affected (e.g., olives) (Makowski et al. 2020).

Studies on climate change impact on vegetable production are scarce. Bisbis et al. (2018) have shown that climate change may threaten vegetable crop yield and quality in response to rising CO<sub>2</sub> and O<sub>3</sub> concentrations as well as extreme events. Heat stress reduces fruit set of fruiting vegetables and accelerates the development of some crops, thus reducing assimilation, resulting in lower quality and higher product waste. Vernalization of some crops such as cauliflower can also be threatened by cool season temperature increase. Fruit crops such as apples may suffer significant delays to flowering dates due to temperature rise and the difficult achievement of chill requirement fulfillment in milder Mediterranean climates, which might threaten the cultivation of sensitive varieties in currently vacated areas (Funes et al. 2016). Most of these impact assessments are made without considering the threats related to increased incidence of pest, diseases and weeds (Bindi and Olesen 2011; Pautasso et al. 2012; Hulme 2017) and those related to extreme events such as flooding (Erol and Randhir 2012).

In the Mediterranean area, rain-fed croplands include mainly winter cereals, forage crops and other autumn-spring herbaceous crops, and perennials such as grapes and olives. Schils et al. (2018) showed that water-limited cereal yield gaps are still relevant, particularly in eastern European countries, as they are substantially higher than for irrigated crops. Unlocking the potential for production growth requires a substantial increase of crop N uptake and/or N use efficiency. Filling these gaps requires ecological or sustainable intensification of agricultural systems, and has many implications on innovation of cropping systems, their ecosystem services and impacts on GHG emissions and soil functions (Cassman 1999; Serpa et al. 2015; Hamidov et al. 2018; Serraj and Pingali 2019). Currently, the climate resilience of European wheat crops is declining because of the decline in the response diversity that is emerging both from farmers' fields and plot experiments also in southern European countries (Kahiluoto et al. 2019). This suggests that current breeding programs and cultivar selection practices do not sufficiently prepare for climate uncertainty and variability and calls for more coordinated assessment and communication of response diversity among plant breeders, the recovery of old varieties that had been abandoned by seed producers to be considered and the need for domestication to broaden the germplasm pool (Langridge 2019).

Climate change will impact olive crop evapotranspiration (+8%) and irrigation requirements

(+18.5%) and crop phenology, up to reducing the possibility of rain-fed cultivation (Tanasišević et al. 2014). Furthermore, climate change will also impact the interaction of olive and the obligate olive fruit fly (*Bactrocera oleae*) and alter the economics of olive crop across the basin. Climate warming will affect olive yield and fly infestation levels resulting in economic winners and losers at the local and regional scales, that overall result in threatened biodiversity and soil conservation (Ponti et al. 2014a).

Wine grape production provides a good test case for measuring indirect impacts mediated by changes in agriculture, because viticulture is sensitive to climate and is concentrated in Mediterranean climate regions. At the global scale, the impacts of climate change on viticulture are expected to be substantial, leading to possible conservation conflicts in land use and freshwater ecosystems. The area suitable for viticulture is expected to drop up to 73% in major wine producing regions by 2050 in the worst scenario (RCP8.5), which could be partially compensated by upland or northward cultivation, or by irrigated crops, possibly resulting in land or water degradation (Hannah et al. 2013). The projected increasing temperatures will result in a general acceleration and shortening of the phenological stages compared to the present period. Accordingly, the reduction in time for biomass accumulation negatively affects the final yield. In the cooler subregions of the Mediterranean Basin such as southern France and western Balkans, climate conditions are not limiting and the crop benefited from enhanced atmospheric concentration of carbon dioxide (Schils et al. 2018). Impacts are also expected on grape composition and hence wine quality, in particular with respect to aroma compounds. Furthermore, the frequency of extreme climate events such as hail and flooding is likely to increase vulnerability and risks in some areas (van Leeuwen and Darriet 2016).

In the warmer areas, increasing temperature can have detrimental impacts on grape yield due to increased asynchrony between the larvae-resistant growth stages of the grapevine and the larvae of the grapevine moth. On the other hand, the increase in pest pressure due to the increased number of generations might not be as severe as expected, because of the advance in harvest dates limiting damages from late-season generations. Furthermore, powdery mildew is expected to decrease in disease severity, especially in years with a later onset of the disease symptoms and under the most extreme warming scenarios (Caffarra et al. 2012).



### Grasslands and grazing systems

Mediterranean pastoral systems in drylands are expected to be severely impacted from climate change, mainly because of altered rainfall regime and grassland ecosystem water balance. Pastoral mobility, where possible, can mitigate the effects in terms of livelihood but not necessarily in economic terms (Martin et al. 2014). These impacts are expected to be site-specific as they are related mainly to the change in precipitation variability, which appears as the main determinant of degradation in terms of losses in fodder and livestock production in drylands. Perennial forage plants adapted to Mediterranean conditions are a fundamental resource (Lelièvre et al. 2011), providing that sufficient rest is allowed between two subsequent grazing periods. However, projected change is expected to outrange the adaptive capacity of pastoralists. Similar conclusions about the dependence of climate change effects on land use and subregions in grassland systems were achieved by Bütöf et al. (2012), who showed how single plant species respond in many different ways to climate pressures because of the complex interactions of climate change with land use practices. More assessments are expected from the use of grassland modelling well-calibrated to Mediterranean-type ecosystems (Pulina et al. 2017; Langridge 2019).

Wooded pastures such as dehesa-type habitats, are a typical high nature value (Bernués et al. 2016) agro-silvopastoral vegetation of many Mediterranean countries, particularly in the western basin (Bagella et al. 2013; Torralba et al. 2016; Seddaiu et al. 2018). These types of ecosystems are already threatened by current management systems under present climate conditions (Rossetti et al. 2015) and by increased drought risks in relation to stocking rates and grazing management, leading to potential higher economic losses with high stocking rates (Iglesias et al. 2016). Tree survival in such ecosystems depends on deep water reserves throughout late spring and summer, which helps to avoid competition for water with herbaceous vegetation (Cubera and Moreno 2007).

Few studies have explored how climate change and grazing interactively affect the biodiversity, primary productivity and ecosystem stability of grassland ecosystems. A recent meta-analysis indicates that the effects of climate change on biodiversity and ecosystem functioning were largely dependent on grazing history within same climate conditions. However, more field studies are needed to test how different climate scenarios affect the biodiversity, functioning, structure and stability of grassland

ecosystems, to address sustainable grassland management in different environmental and climate contexts (Kairis et al. 2015; Li et al. 2018).

Vulnerability and risks are mainly associated with the increased frequency of heat stress in summer, leading to heavy impacts on animal health and welfare, i.e., increased incidence of diseases and mortality or lower fertility (Lacetera 2019). Indirect effects of climate pressures increase vulnerability and risks associated with new vector-borne infections such as bluetongue (driven by *Culicoides imicola*), or other direct parasites, whose spread can be facilitated by a milder winter climate in northern Mediterranean countries (Bosco et al. 2015). Other indirect effects can be related to the increased incidence of mycotoxins in fodders due to a higher incidence of pests and diseases in forage crops favored by increased temperature (Bernabucci et al. 2011).

#### 4.3.2.5 Freshwater ecosystems

##### Rivers and streams

In most of the Mediterranean region average river discharge is predicted to decrease while both water temperature and the frequency of large floods are likely to increase (Calbó 2010). The projected decrease in rainfall and increase in temperatures will result in a 10 to 30% decrease in river discharge by the end of the 21st century and a significant reduction in the availability of freshwater (Allen and Ingram 2002; Milly et al. 2005; Lelieveld et al. 2012).

In the eastern Mediterranean, many authors have detected negative trends in runoff. This was the case for rivers located in Greece (Giakoumakis and Baloutsos 1997) and the Balkans (Genev 2003; Rivas and Koleva-Lizama 2005; Frantar and Hrvatin 2006), Lebanon (Shaban 2009) and Turkey (Kahya and Kalayci 2004). In the western basin, the Duero Basin in the Iberian Peninsula is the most obvious example. Since 1960, Duero River discharges have decreased by 20 to 50% (Ceballos-Barbancho et al. 2008; Morán-Tejeda et al. 2010). Most Mediterranean catchment headwaters are in mountainous areas and are snow-fed. In various regions across the Mediterranean, snow-fed high mountainous springs are the only source of runoff during the long dry summer of the Mediterranean climate. Hence, an increase in temperatures cause less snow accumulation and an irregular and rapid snowmelt, which will result in turn in higher winter and lower spring discharges and decreasing summer low flows. These impacts are observed in various part of the Mediterranean such as the Pyrenees (López-Moreno and García-Ruiz

2004; López-Moreno 2005) and in Lebanon (Shaban 2009).

Overall, projections suggest decreased hydrological connectivity, increased concentration of pollutants during droughts, changes in biological communities as a result of harsher environmental conditions, and a decrease in biological processes like nutrient uptake, primary production, or decomposition. Furthermore, the increased pressure on shrinking water resources will compound the impacts on river ecosystems (Navarro-Ortega et al. 2015).

### **Wetlands**

Mediterranean wetland water depths and hydroperiods (meaning the water inundation period) along with the increase in their salinity levels and isolation and fragmentation are affected by multiple human activities (e.g., water extraction) (Ramírez et al. 2018). These activities are altering the water budgets of wetlands and reducing their ecosystem services. Reed beds in the region have expanded by 89.3% and are the predominant aquatic plant of the all wetlands in the region, which is a major change. In contrast, open water areas and wet meadows have decreased by 53.7 and 96.5% respectively (Papastergiadou et al. 2007). The loss of these key wetland features (e.g., open waters and wet meadows) are impacting the structure of waterbird communities. The future conditions of climate change scenarios will further reduce the environmental suitability of Mediterranean wetlands for the guilds of diving birds and vegetation gleaners (Ramírez et al. 2018).

### **Freshwater biodiversity**

The high intensity and large-scale water management alterations on rivers and streams of the region have had a particularly strong impact on these ecosystems, possibly the highest in the world (Grantham et al. 2013). A similar trend in fish biodiversity loss, also associated with water management pressures, has been reported for rivers in the Iberian Peninsula (Aparicio et al. 2000; Benejam et al. 2008; Clavero et al. 2010). The establishment of alien species in these ecosystems, which can alter natural processes and adversely affect native biota, has also been associated with numerous anthropogenic hydrologic infrastructure in the region (Elvira and Almodovar 2001; Clavero et al. 2004; Light and Marchetti 2007; Grantham et al. 2013).

The Mediterranean-climate freshwater ecosystems host fauna that have evolved and are adapt-

ed to the stresses of its streams and rivers. With climate change predicting longer or more extreme drying events (Lawrence et al. 2010; Filipe et al. 2013), their populations and communities will be highly stressed during dry years, thus reducing the resilience capacity of Mediterranean rivers and streams and compromising the survival of their biota (Magalhães et al. 2007). Under this situation, these new conditions will lead to irreversible, and undesirable, "regime shifts" in Mediterranean rivers (Cid et al. 2017).

Due to climate induced changes, stream biota tend to move towards higher elevations and upper latitudes, while the communities change and homogenize their composition (Filipe et al. 2013). Some life-history traits provide biota with resilience and resistance to adapt to the new conditions although it appears that in many cases, current and future environmental changes are exceeding the biota survival boundaries. The difficulty of distinguishing disturbances due to natural hydrologic variability from the effects of climate change in the region make adaptation forecasts even more challenging. Long-term studies are needed to improve knowledge regarding stream biota ecological responses due to climate change (Filipe et al. 2013).

The reduction of subsurface inflow to streams and the changes in groundwater dynamics that have degraded of their biological quality have already made these ecosystems highly vulnerable (Benejam et al. 2008). In addition, wastewater inflow (whether treated or not) into streams will further exacerbate the pressures on fluvial ecosystems, even though initially the induced drought impacts can be partially offset by these industrial discharges. The hydrological benefits of these discharges are compromised by declines in water quality and habitat quality. Moreover, the capacity of aquatic ecosystems to cope with droughts has been lost or significantly reduced in many regions (Andersen et al. 2004; Bond et al. 2008; Rault et al. 2019).

### **4.3.3 Adaptation**

Communities continue to be significantly dependent on ecosystem services for their livelihoods and therefore the preservation of the livelihood and culture of communities together with its biodiversity, in these areas is considered as important to promoting sustainable development and adaptation to climate stresses within the region. The integration of humans, and human actions within the landscapes and seascapes of the Mediterranean region also embraces the IUCN Category IV - Protected Landscapes (Dudley 2008), which

provides the flexibility to offer protection to entire landscapes, as shaped through the interaction of human actions and nature, as well as specific protection for specially defined purposes (e.g., habitats/species). Appropriate (integrated) landscape management can be used to promote heterogeneity compensating for the loss of habitat diversity:

- the protection of traditional food systems, conservation of species and functional agricultural biodiversity, and improvement in cropping and irrigation systems to adapt to a changing climate,
- sustainable urban development that promotes the uptake of nature-based solutions that are suitable for a Mediterranean climate in urban areas to provide benefits to biodiversity, contribute to ecosystem services and increase resilience to climate change (Box 4.2),
- semi-natural ecosystems: adaptive management includes the implementation of habitat management, restoration and afforestation actions to provide benefits to biodiversity and human well-being, whilst using species adapted to expected future conditions,
- managing changing disturbance regimes: preventive (e.g., pest monitoring) and remedial (e.g., sanitation felling, pest control).

This section provides an overview of the opportunities for adaptation of ecosystems, through incremental (capacity-building) actions and impact-based actions, whilst considering the limitations to the adaptability of Mediterranean social-ecological systems and the impacts of these actions on biodiversity and the ecosystem services and benefits to human well-being. In this analysis, the role of human influences and inputs on ecosystem structure and functions is critical to promoting (or limiting) adaptation of Mediterranean social-ecological systems.

#### 4.3.3.1 Forests

Mediterranean forests will need to adapt to a warmer and drier climate, which entails extended drought periods, long heat waves, increasing fire risk and exposure to increased intensity and frequency of biotic disturbances (e.g., pests). Mediterranean forests, as any other type of ecosystem, have an inherent adaptive capacity as a result of the co-evolution of plants with environmental conditions that have always changed (Valladares et al. 2014a). However, the speed of current environmental change is unprecedented and poses doubts concerning the ability of Mediterranean species to cope with the change to come, and in some cases

might make it advisable to adopt planned adaptive measures.

#### Biological adaptation

The inherent adaptive capacity of forests includes in situ adjustments to new environmental conditions via phenotypic plasticity or natural selection, and migration to more suitable habitats (Mate-sanz and Valladares 2014). Climate envelopes are shifting polewards and upwards, and the easiest response to climate change may be a geographic shift in distribution into climatically suitable areas (Christmas et al. 2016). There is already evidence of some species responding to increasingly warmer and arid conditions through altitudinal or latitudinal migration. For example, Peñuelas et al. (2007) and Peñuelas and Boada (2003) showed a gradual upward shift of the temperate *Fagus sylvatica* species in northeastern Spain and their gradual replacement by the xeric *Quercus ilex* in the mid- and low- altitudes. Similarly, Sanz-Elorza et al. (2003) reported an encroachment of sub-alpine grasslands by Mediterranean woody species characteristic of lower altitudes during the second half of the 20th century in mountain systems of central Spain. Upward migration of forest species has also been reported in the Italian Apennines (Palombo et al. 2013), the Spanish Pyrenees (Améztegui et al. 2010, 2016) or southeastern France (Bodin et al. 2013). However, although these movements coincide with an increase in temperatures, changes in land use (agricultural abandonment and reduction of anthropic pressure on forests) seem to play a preponderant role in forest expansion (Améztegui et al. 2010). Most of the species altitudinal displacements have occurred via the colonization of open areas after their abandonment. Replacement of a given tree species by their low-altitude neighbors is only possible when there is a retraction in its trailing-edge distribution, such as in the case of the Montseny mountains studied by Peñuelas and Boada (2003). However, altitudinal range retractions have received much less attention than lead-edge expansions, particularly in Mediterranean mountains, although they seem to be occurring in many mountain areas worldwide (Jump et al. 2009).

Phenological observations since the 1950s show a fairly consistent response of Mediterranean vegetation to rising temperatures. Between 80 and 96% of the species studied advanced their leaf unfolding, delayed the leaf fall, or both, which resulted in an average extension of 30 days in the growing season between 1952 and 2000 (Peñuelas et al. 2002). The lengthening of the growing season could

trigger increases in growth but can also cause higher frost damage risk and increased water transpiration. There is less consensus about the physiological plasticity of Mediterranean forest species to environmental stressors. The current available information reveals the potential for some Mediterranean plant species for significant plasticity and rapid evolutionary change, and epigenetic responses have also been documented (Madlung and Comai 2004). However, this information is fragmentary and suggests large differences among species, with some of them being quite vulnerable to fast rates of environmental change (Matesanz and Valladares 2014). Some studies reveal contrasting functional responses to disturbances among tree species and forest biomes. For instance, evergreen gymnosperms growing in drought-prone areas showed lower resistance but faster recovery after drought events than plants dominating in temperate or wet regions, which suggests different physiological strategies to cope with drought (Gazol et al. 2018). This may be of great importance in a changing future, as the response of vegetation may be different as droughts become more intense, more frequent, or both. Tree species with wide a distribution range also seem to display contrasting responses across their entire range (Benito-Garzón et al. 2011) that have been related to intraspecific plasticity and genetic differentiation among tree populations as a result of differences in the intensity of the environmental stresses (Benito-Garzón et al. 2013). In a drier environment, interactions between species may also be altered. According to the stress gradient hypothesis (Maestre et al. 2009), facilitative effects may become more frequent. In fact, the role of shrubs as nurse vegetation for pine seedlings has already been documented in semi-arid and arid Mediterranean regions (Castro et al. 2004; Gómez-Aparicio et al. 2008), and this role could become even more important in the future.

The degree to which physiological responses lead to vegetation shifts (i.e., changes in the composition of the vegetation) is fundamentally unresolved. When two or more species coexist and are differently affected by directional changes in climate and/or by disturbance events, demographic responses become fundamental to project the fate of woody plant communities (Martínez-Vilalta and Lloret 2016). Mortality and regeneration thus become the key processes, since a vegetation shift will only occur if the initially affected species is not able to regenerate and dominate again (Martínez-Vilalta and Lloret 2016). In a drier environment, interactions between species may

also be altered. According to the stress gradient hypothesis (Maestre et al. 2009), facilitative effects may become more frequent. In fact, the role of shrubs as nurse vegetation for pine seedlings has already been documented in semi-arid and arid Mediterranean regions (Castro et al. 2004; Gómez-Aparicio et al. 2008), and this role could become even more important in the future.

### **Limits to adaptation**

Forests are particularly sensitive to climate change, because the long life-span of trees does not allow for rapid adaptation to environmental changes (Lindner et al. 2010). Moreover, the Mediterranean region is likely to experience more adverse effects of climate change when compared to Europe, while being the least prepared to cope with such drastic changes (Lindner et al. 2010; Lindner and Calama 2013). On the one hand, the strong human impact on Mediterranean forests has led to high levels of fragmentation, which alter population genetics and species ecology, and affect the ability of populations to respond to environmental changes. Furthermore, adaptive capacity is usually the lowest at the rear edge of species ranges, where plants are growing close to their physiological limits, as is the case for many species dwelling in Mediterranean forests. These forests cannot benefit from gene flow from better adapted populations, so only short-term adaptation and plasticity are available to cope with the extinction risk (Lindner and Calama 2013).

Fragmented populations suffer from greater genetic drift, homozygosity and inbreeding within populations, and are less likely to benefit from the positive effects of gene flow (Valladares et al. 2014b; Christmas et al. 2016). Both the adaptive capacity to new environmental conditions and the ability to migrate are hampered by fragmentation. Shifts in species or population ranges to track optimal climate conditions can be limited by fragmentation, which acts as a barrier for the colonization of many species. Indeed, migrations are not as common as could be expected (Harsch et al. 2009; Zhu et al. 2012), potentially putting populations at higher risk of becoming increasingly maladapted over time (Christmas et al. 2016), particularly at the rear end of species distributions, where populations are deprived of gene flow from better adapted populations (Lindner et al. 2010). Under the projected rates of future climate change, migration will rely on the evolution of very long dispersal distances in order to enable species to reach suitable new habitats. Moreover, global warming will also lead to a decoupling of species interactions (pollinators,

predator-prey, etc.) that can further complicate the migration of several forest species.

If species fail to migrate, then *in situ* adaptation will be the only strategy for persistence (Christmas et al. 2016). In this regard, fragmented populations also show lower capacity of adapting and responding to changing conditions. Forests with a greater diversity of response traits (i.e., traits that confer the organisms the ability to cope with disturbances), are indeed more resistant and resilient to disturbances, increasing the likelihood that such communities may persist under future conditions (Sánchez-Pinillos et al. 2016). Moreover, the effects of fragmentation on the genetic diversity of organisms can last for centuries, and some populations still hold legacies from the effects of human actions centuries ago, as is the case of *Fagus sylvatica* in northeastern Spain (Jump and Peñuelas 2005). In the absence of migration or evolutionary adaptation, the ability of populations to persist in the new environment will depend exclusively on their phenotypic plasticity, i.e., their ability to alter their phenotype with environmental conditions. Although Mediterranean forests have shown relatively high levels of plasticity, being plastic has an important metabolic cost, and there are universal physicochemical constraints that prevent the ability of a species to simultaneously tolerate several stresses (Laanisto and Niinemets 2015). In this sense, the succession of disturbances can cause an important limitation to the adaptation of the species. For example, the regeneration of *Pinus nigra* after wildfires depends both on the existence of nearby unburned vegetation patches and on climate conditions in the years following the fire (Martín-Alcón and Coll 2016; Sánchez-Pinillos et al. 2018). Distinct sequences of disturbance events can cause vegetation transitions, with non-linear responses and tipping points, even if the recurrence of individual disturbances is moderate (Batllori et al. 2019). Therefore, the succession of fires and droughts could trigger massive failures in regeneration, leading to a change in the ecosystem towards a greater dominance of oaks. In the driest areas, the combined effects of several disturbances is likely to exceed the response capacity of organisms, leading to the extinction of some species and even triggering shifts in ecosystem state (from forest to non-forest) (Batllori et al. 2019), which entails a high risk of soil erosion, degradation, and desertification.

### Measures to promote adaptation

Whenever the inherent adaptive capacity of species is not sufficient, or too slow, planned adap-

tation measures can be implemented to decrease the known risks, increase forest resistance, or promote its recovery capacity (Lindner and Calama 2013). Adaptation measures in the Mediterranean commonly seek to address the two main disturbances in the region: drought and fire, and can be classified into five categories (Vilà-Cabrera et al. 2018).

**Reducing tree density through thinning** has the triple effect of increasing the growth and value of the remaining trees while also improving their water status and reducing fire risks. In a climate change context, thinning can diminish interception losses and reduce stand transpiration, increasing the amount of available water, which is apportioned among fewer trees (Sohn et al. 2016b). Some studies report a direct reduction of drought-induced mortality of Scots pine for high thinning intensities (Giuggiola et al. 2013), and an increase in the resistance and recovery of growth following drought events (Martín-Benito et al. 2010; Sohn et al. 2013, 2016a), which may be particularly important in dry areas or under severe climate change scenarios (Ameztegui et al. 2017; del Río et al. 2017). However, there is also evidence to suggest that, under extremely dry conditions, tree mortality risk may be density-independent, as all the available soil moisture can be lost to evapotranspiration before it can be harnessed by trees (Dorman et al. 2015).

**The reduction of the understory cover** has mainly been applied with the aim of reducing the risk of fire propagation by breaking the vertical and horizontal fuel continuity. Reduction of understory cover can be achieved either through mechanical treatments, prescribed burning or by promoting understory grazing in forest areas (Vilà-Cabrera et al. 2018).

The promotion of **mixed-species stands** (at the species or genotype levels) can increase resistance and recovery capacity to extreme droughts (Pretzsch et al. 2013), higher temporal stability (Jucker et al. 2014; Sánchez-Pinillos et al. 2016; del Río et al. 2017), and reduce the risk of biotic and abiotic disturbances (Guyot et al. 2016; Jactel et al. 2017) and the maintenance of ecosystem service provision (Gamfeldt et al. 2013).

**The change in species or genetic composition** seeks to replace the maladapted species or populations with species or genotypes better adapted to the forecasted climate conditions, and can include (i) assisted population migration (i.e., the active relocation of well-adapted populations of a given species within its current range); (ii) assist-

ed range expansion (relocation of a species to an area adjacent to its current range); and (iii) assisted species migration (i.e., the displacement of a species beyond its current range, where the future climate is expected to be suitable for its development) (Williams and Dumroese 2013). Although the effectiveness of these practices seems apparent, there are still many doubts about their risks and consequences on the host environment, especially in the latter sense. At present, there is no consensus on their suitability, and they generate significant rejection both by a large part of the scientific community and by the general population (Lawler and Olden 2011), and in the Mediterranean they have not yet been applied beyond small-scale scientific experiments (Martín-Alcón et al. 2016).

The promotion of **the spatial heterogeneity of the landscape matrix** has mostly been advocated as a way to reduce the impacts of fire by slowing or preventing its expansion and allowing for greater effectiveness in firefighting (e.g., minimizing total burnt area) (Loepfe et al. 2012; Regos et al. 2016). A heterogeneous landscape also allows for the coexistence of different habitats (forests, open areas, etc.) each with different goals and providing different services. Moreover, greater heterogeneity can also contribute to enhancing gene flow and natural species migration, provided that enough corridors are available (Saura et al. 2018; Vilà-Cabrera et al. 2018). Fire risk management can also be achieved through the promotion of particular land covers/uses that reduce the risk of intense crown fires (e.g., fagaceae vs. conifers: Moriondo et al. 2006).

However, the socio-economic adaptive capacity of the Mediterranean forest sector also has to face several constraints derived from the low economic incomes of many Mediterranean forests, mainly due to low fertility and water limitation, the lack of a developed road network and the limited implementation of technological advances (e.g., harvesting machinery), which results in a large part of Mediterranean forests not being managed at all (Lindner et al. 2010; Lindner and Calama 2013), limiting the capacity of forests to adapt to climate change. Moreover, managing forests to increase adaptive capacity can lead to trade-offs with other ecosystem functions and with biodiversity (Vilà-Cabrera et al. 2018).

#### 4.3.3.2 Mountain ecosystems

Implementation of effective adaptation measures depends on the availability of human resources and expertise. However, the knowledge base about Mediterranean mountains varies significantly.

Enhancing connectivity is a key measure to facilitate expected range shifts (Keeley et al. 2018) which in mountain areas may be achieved by “building” linear and latitudinal corridors and taking advantage of the river network. In addition, due to interconnected risks, wider spatial frameworks are necessary, for instance at the watershed level, since upstream changes influence downstream. There are still pristine areas in many Mediterranean mountains that sustain a diversity of plant and animal species. However, this role is impeded by ongoing human activities and most importantly, climate change has pointed to the need to design a flexible reserve system along with conventional ex situ conservation measures. Such as reserve system may place emphasis on the permeability of the intervening landscape matrix, dispersal corridors and habitat networks (Jongman and Pungetti 2004; Watts and Handley 2010). Mountains have played a refuge role in geological history and to a certain extent they retain this role today, with many of their endemic species surviving in places located in such refugia (Vogiatzakis and Griffiths 2008; Vogiatzakis 2012). This is at the core of climate-wise connectivity as proposed by recent studies (Keeley et al. 2018). Mountain ecosystems may prove more resilient since upward migration of lower zone species will be conditioned by topography and geomorphology i.e., habitat suitability (Kazakis et al. 2007).

#### 4.3.3.3 Drylands and shrublands

Plants exhibit a variety of mechanisms to avoid (e.g., annual life-cycle) or to tolerate drought (e.g., perennial shrubs), and to deal with disturbance pressures, such as fire and herbivory (Noy-Meir 1973; Davies et al. 2012). Dryland biodiversity interacts with abiotic factors to determine ecosystem functioning (e.g., productivity, nutrient fluxes) and resilience (i.e., the ability to return to a previous state after disturbance), both of which are critical to ensuring the provision of ecosystem services (MEA 2005). Climate change predictions point to an overall increase in aridity and in the variability of precipitation distribution in drylands (Dai 2013). Climate change is therefore expected to further reduce productivity over time.

Topography creates contrasting microclimates, especially between northern and southern slopes in drylands, which result in clear differences in tree cover patterns in the landscape. These local-scale differences in tree cover patterns may result from limitations occurring at different plant development stages, such as seed germination, seedling establishment, tree growth rate and survival,

all crucial for the maintenance and expansion of plant populations through natural regeneration. Leaf physiological performance was similar under contrasting microclimatic conditions. However, in areas with higher Potential Solar Radiation, tree age and density were significantly reduced. These results suggest that microclimatic differences on southern slopes with high Potential Solar Radiation are limiting for germination and sapling establishment. Thus, forest regeneration, restoration and nature conservation practices aiming at increasing forest resilience in Mediterranean dryland climates should account for the importance of microclimate in defining the niche of seedlings and adult trees (Príncipe et al. 2019).

Shrub encroachment was largely predicted by topo-edaphic factors in Mediterranean dryland ecosystems subject to conventional low-intensity land use composed of savanna-like holm oak woodlands, along with a regional climate gradient (Nunes et al. 2019). Management strategies to reduce encroachment therefore need to take these drivers into account for efficient forecasting and higher cost-effectiveness. Climate had a stronger effect on a set of functional traits involved to a limited extent in shrub encroachment, related to flowering and dispersal strategies. These results suggest that climate change might not greatly impact shrub encroachment in the Mediterranean Basin, but may affect the functional structure and reduce the functional diversity of plant communities, thus affecting ecosystem functioning (Nunes et al. 2019).

Drylands are very susceptible to the effects of climate change due to water stress. One possible climate change adaptation measure is the construction of lakes to increase water availability for drinking and irrigation (food production) and decrease fire risk. These lakes can also increase local biodiversity and human well-being. However, other non-target services such as carbon (C) storage, water purification, and sediment retention might also change.

An evaluation of the trade-offs on non-targeted ecosystem services due to lake construction in drylands was carried out by Santos et al. (2018). This was done using the Integrated Valuation of Ecosystem Services and Tradeoffs (InVEST) modelling tools, comparing a Mediterranean area located in southwestern Europe, with and without artificial lakes. Results showed that the construction of artificial lakes caused an increase of 9.4% in carbon storage. However, the resulting increase in agricultural area decreased water purification

and sediment retention services. This could diminish the lifespan of the lakes, changing the initial beneficial cost-benefit analysis on lakes as adaptation measures to climate change. As a global measure for mitigation and adaptation to climate change strategy, we consider lake construction in drylands to be positive since it can store carbon in sediments and reduces the vulnerability to water scarcity. However, as a general recommendation, and when built to support or increase agriculture in semi-arid landscapes, we consider that lakes should be supplemented with additional measures to reduce soil erosion and nutrient leaching, such as (i) locating agricultural areas outside the lake water basin, (ii) afforestation surrounding the lakes, and (iii) adopting the best local agriculture practices to prevent and control soil erosion and nutrient leaching.

#### 4.3.3.4 Agriculture and pasturelands

Transformational adaptation in agriculture has been described as a redistribution of at least a third of the production factors and/or production output in a 25-year timeframe, which mostly involves qualitative changes in inputs and outputs (Vermeulen et al. 2013). Transformational adaptation can result into a radical change to the area of production, to the main crops or production types (e.g., shift from animal to crop productions, abandonment of a specific type of farming, reclaim of abandoned lands, shift from rain-fed to irrigated agriculture, from nomadic to settled grazing systems, from conventional to organic farming systems). Vermeulen et al. (2018) also pointed out that the success factors and drivers of positive transformational adaptation include changes in governance in favor of disadvantaged stakeholders. They also showed that the capacity of producers, processors and consumers to adapt is highly context-sensitive and depends on public policy, market drivers and cultural values.

Farmers' long-term responses and investments in adaptation are constrained by barriers: (i) climate change signals can be biased by the perceptions of farmers (Nguyen et al. 2016), (ii) the projections are uncertain, (iii) climate change communication is difficult and often does not result in behavioral change (Wise et al. 2014), (iv) timescales for farm planning are relatively short and other priorities take precedence, (v) there are expectations of technology being able to cope with the negative effects of climate change, (vi) scientific knowledge and tools developed in agriculture rarely support long-term strategic decisions (Robertson and Murray-Prior 2016). Also, short-sighted state policies can increase the vulnerability of farmers

to climate change and constrain their adaptive capacities (Turhan et al. 2015).

Opening new spaces for learning and generating enabling contexts through translating systems thinking into practice are fundamental steps for raising awareness about adaptation actions in a climate change world (Ison 2010). Vermeulen et al. (2018) examined several case studies on transformational adaptation in agriculture and showed how in practice it often emerges from a disorganized combination of responses from multiple actors to external pressures where climate change can only be an indirect driver. However, farmers rely on their experience to plan their practices, which will become obsolete if climate change occurs too quickly. They design their activities on the basis of a perceived probability distribution of their known drivers and performance indicators but, under climate change, probability distributions are shifted in mean, variance or both and are unknown, thus resulting in increased frequency of unexpected events. Dono et al. (2016) showed how these shifts between actual and expected probabilities can result in winners and losers in an affected area, depending on the type of farming system under the same expectations of climate pressures in a Mediterranean context.

Adaptation in agriculture requires customized support to the choices of any specific farm type in a given environmental situation or context, which is a challenge both for science and policy. The variety of environmental and socio-economic contexts and agricultural systems across the sub-regions of the Mediterranean Basin and across farming types within sub-regions, generate a great diversity of needs, adaptation strategies and have site-specific implications on biodiversity and the ecosystem services of agroecosystems.

Adaptation strategies and plans are being adopted at the continental, country or local level under the Cancun Adaptation Framework of the UNFCCC<sup>37</sup>. However, by December 2018, among the nine Mediterranean countries of the EU, only three (Cyprus, France and Spain) had already adopted a national adaptation plan, while eight had adopted a national strategy. Such plans should create the enabling environment for “last-mile” adaptation to occur, but moving from planning to implementation is a challenge because of the difficulties found in addressing capacity constraints, securing adequate financing and measuring the success of actions

(Mullan et al. 2015). The following cases provide some examples of the specific adaptation needs of agriculture and pastureland systems in the Mediterranean context.

At present, cereal production is well below potential in southern Europe. Schils et al. (2018) demonstrated that the yield gap between actual and potential production would require good agronomy for sustainable intensification and thus increase the self-sufficiency food production of the entire Mediterranean area (Vermeulen et al. 2013). The self-sufficiency ratio in northern Africa is lower today than it was in the past as a consequence of demographic expansion, and is also resulting in low stability, not only in low GDP countries (Luan et al. 2013). Under pronounced drying trends documented by recent assessments, particularly in northwest Africa, a strategic objective is to move from maximized to stabilized production (Schilling et al. 2012). This can also be achieved through improved climate-proof agronomic practices such as the incorporation of crop residues combined with supplementary irrigation, where available (Benlhabib et al. 2014; Jacobsen 2014).

Adapting dryland agriculture to climate change in the Mediterranean requires substantial investments in plant breeding for heat and water stress tolerance and to increase yield and quality under conditions of high CO<sub>2</sub> concentration (Asseng and Pannell 2013). The same authors suggest investments in new species and cultivars of perennial plants. Such investments should be coupled with improved seasonal forecasting, which would enable farmers to make timely decisions about agronomic practices, thus improving resource use and crop yield. However, climate resilience is currently not receiving the necessary attention from breeders, seed and wheat traders, and farmers, while there are clear signals of declining resilience, at least for durum wheat, also in Mediterranean countries, including Spain (Kahiluoto et al. 2019). This latter assessment revealed that current breeding programs and cultivar selection practices do not sufficiently prepare for climate uncertainty and variability by applying a variety of responses to the same climate pressures by different wheat cultivars. In the case of barley, a pivotal crop in the Mediterranean area, Cammaron et al. (2019) demonstrated that, besides plant breeding, shifting sowing dates and improving soil organic carbon are viable adaptation strategies to mitigate the expected negative impacts of a future drier and

<sup>37</sup> <https://unfccc.int/topics/adaptation-and-resilience/workstreams/national-adaptation-plans>



warmer Mediterranean climate on barley grain yield.

Wheat protein yield gains are expected to be lower and more variable in most rain-fed low-input cropping regions, where nitrogen availability limits growth stimulus from elevated CO<sub>2</sub> (Asseng et al. 2019). This is particularly true for North African countries, where food demand is increasing due to population increase (Schils et al. 2018). Introducing wheat genotypes adapted to warmer climate may not result in increased protein production. Therefore climate adaptations leading to stabilized grain yield could not always be positive in terms of grain quality (Asseng et al. 2019). Adaptive pathways for cereal productions also have impacts on integrated governance aiming to yield stability, such as the implementation of national action plans and policies to regulate and provide incentives for increasing diversity in crop responses to climate uncertainties (Kahiluoto et al. 2019).

Legumes represent a strategic resource for sustainable intensification of agricultural systems and climate change adaptation in the Mediterranean Basin. In addition to serving as a fundamental source of high quality food and feed, legumes contribute to net nitrogen inputs in cropping systems at low N<sub>2</sub>O emissions and contribute to net soil carbon sequestration (Volpi et al. 2016; Stagnari et al. 2017). The environmental services provided by legume cultivation are still undervalued, while new opportunities for yield improvement are arising from the ongoing development of cost-efficient genome-enabled selection procedures, enhanced adaptation to specific cropping conditions and more thorough exploitation of global genetic resources (Annicchiarico 2017).

The cropping systems for the production of bio-energy and biomaterials are assumed to occupy part of the residual agricultural land abandoned in the past 50 years, but this will not be sufficient to meet the increased bioenergy demand associated with climate change energy policies in European countries (Cosentino et al. 2012). These crops may also find a strategic position in Mediterranean cropping systems to reclaim polluted arable land from industrial or mining wastelands (Fagnano and Fiorentino 2018).

Increased irrigation water efficiency and the design of climate-friendly agro-ecosystems are key adaptation strategies for Mediterranean agriculture, in particular for countries such as Algeria, Libya, Israel, Jordan, Lebanon, Syria, Serbia, Morocco,

Tunisia and Spain, which are at high risk of not being able to meet future irrigation needs (Fader et al. 2016). A range of adaptation strategies are being studied or put in place either to store more water in hot and arid environments (e.g., with managed aquifer recharge) (Salameh et al. 2019), use and recycle non-conventional water sources (Ait-Mouheb et al. 2018; Elkiran et al. 2019), desalinate seawater (Stanhill et al. 2015) or improve irrigation efficiency (Tarjuelo et al. 2015; El Jaouhari et al. 2018). All these strategies have some potential side effects in terms of energy requirements (Rodríguez-Díaz et al. 2011), GHG emissions, high capital investments and social acceptance (Daccache et al. 2014; Chartzoulakis and Bertaki 2015). However, in arid zones, these are often the only alternatives to achieve sustainable agricultural intensification. Some solutions, like desalination, should be considered only where there is evidence that the natural recharge available in surface and underground storage might become limiting considering the economic and environmental dimensions of sustainability (Stanhill et al. 2015).

Supplementary irrigation of rain-fed crops is also crucial for increasing the productivity of traditional Mediterranean rain-fed cropping systems, including winter cereals or perennial crops such as olive and vineyards (Fraga et al. 2012; Tanasijevic et al. 2014). However, the introduction of new technologies for irrigation on traditional rain-fed or irrigated cropping systems has many systemic implications in the environmental, socio-cultural, institutional and economic domains (Ortega-Reig et al. 2017).

Increased water and soil salinity is also a threat for future Mediterranean cropping systems, particularly in coastal areas (Maggio et al. 2011; Pittalis et al. 2016). Adaptation strategies include the introduction of salt tolerant crop species for which there is increasing consumer demand. For example, there is increasing interest in quinoa germplasm and production in the Mediterranean Basin, in relation to its tolerance to salinity and water stress, high water use efficiency and the increasing demand of gluten-free food (Hirich et al. 2014; Lavini et al. 2014; Mahmoud 2017; Noulas et al. 2017).

Combined agro-ecological approaches to climate change adaptation in organic horticulture is suggested by Diacono et al. (2016), following long-term field experiments that showed that such cropping systems can sustain the yield of cash crops in rotation, in spite of changes in temperature and rainfall.

Pasturelands and rangelands will face multiple threats from expected drier and warmer climate in Mediterranean countries. These farming systems rely heavily on natural resources on marginal land, often characterized by shallow and oligotrophic soils, with low water holding capacity. Highland pastures and their biodiversity is being threatened by loss in biodiversity due to climate change (Dibari et al. 2015). Dono et al. (2016) showed that near future climate change will result in losses for rain-fed grazing systems, mainly because of the shift in the probability distribution of rain-fed pasture and hay crop production due to drier springs caused by reduced rainfall and higher evapotranspiration, resulting in higher costs for purchasing external feeds or renting more land. Adaptation strategies range from incremental to transformational strategies in this case. Silanikove and Koluman (2015) project an overall negative impact, but a positive role of dairy goats in adaptation to global warming when compared to dairy cows, given their higher tolerance to heat stress.

The savanna-type pastoral vegetation of the dehesa in Spain, montado in Portugal or pastures with scattered cork-oak trees in Sardinia and elsewhere in northern African countries are considered a multifunctional resource that can support adaptive responses to climate change and the provisioning of multiple ecosystem services (den Herder et al. 2017; Castro and Castro 2019). Mediterranean agro-silvopastoral systems generate unique habitats for plant and microbial diversity, resulting in a wide range of services such as forage, wood and non-wood products, soil organic carbon sequestration and landscape cultural values (Bagella et al. 2013; Seddaiu et al. 2013, 2018; Rossetti et al. 2015; Tardy et al. 2015; Torralba et al. 2016; Garrido et al. 2017). Adaptation strategies in pastoral systems based on wooded pastures include actions that can prevent the threats of degradation due to abandonment (e.g., wildfires, loss of cultural landscape and heritage, increased drought stress) or intensification (e.g., lack of tree regeneration) (Garrido et al. 2017; Rolo and Moreno 2019). Given the complexity of the factors driving the sustainability of agro-silvopastoral systems, adaptation strategies should be designed and implemented through systemic and integrated approaches and not by just targeting a specific service or pastoral activity (Hernández-Morcillo et al. 2018). However, more attention should be devoted to these agro-silvopastoral systems, as they are currently overlooked by rural development policies in Europe, while agroforestry systems can effectively contribute to maximizing the productivity of marginal land (Mosquera-Losada et al. 2018).

Agro-silvopastoral systems in the Mediterranean area are under threat because the income of farmers that contribute to their maintenance does not acknowledge the many ecosystem services they provide (Fagerholm et al. 2016; Rodríguez-Ortega et al. 2018).

The adaptive capacity of grazing systems in the Mediterranean depends on local contexts, with contrasting trends in northern and southern countries, rain-fed or irrigated conditions. A general trend towards increased specialization and related environmental risks is occurring almost everywhere, which is in contrast with the need for increased resilience to climate pressures and reduced environmental impacts (Rodríguez-Ortega et al. 2017). Adaptive development strategies include enhancing the spatial dimension of grazing systems through increased animal mobility, increased feeding self-sufficiency and integration of crop-livestock integration at the regional and sub-regional levels (Alary et al. 2019). These livestock farming systems would also respond to the ongoing change in human dietary recommendations, which is one of the drivers of the meat crisis (D'Silva and Webster 2017). However, this may result in different environmental impacts in terms of greenhouse gas emissions, eutrophication and land use in different regions, depending on the income level (Behrens et al. 2017).

Adaptation strategies are more complex for farmers who have made large long-term investments following market pressures and productivity objectives. Dairy cattle farming systems rely mostly on irrigation water and are threatened by the increasing frequency of heatwaves, to which highly productive cows are very sensitive (Lacetera 2019). These farming systems are facing uncertainties caused by fluctuating world feed prices, climate, market and environmental normative pressures (e.g., nitrate vulnerable zones), which are gradually squeezing their marginal net returns (Dono et al. 2016). Adaptation strategies are constrained by multiple pressures and are often based on crop and animal diversification combined with improved animal feeding and genetics (Rojas-Downing et al. 2017; Henry et al. 2018).

#### 4.3.3.5 Freshwater ecosystems

Successful adaptation measures need to follow a large-scale hydrological approach to determine the origin of variations, which are usually related to human pressures, and to provide further strategies for environmental management (Menció and Mas-Pla 2010). Conservation and restoration ef-

forts traditionally carried out at a local scale need to be accompanied by land use and hydrological planning at a basin-wide scale in order to maintain stream ecosystem integrity and biodiversity (Bruno et al. 2014a, 2014b). In this sense, nature-based solutions can play an important role in maintaining freshwater biodiversity, and because of their multifunctionality, in providing critical ecosystem services (e.g., food provisioning, erosion regulation and cultural ecosystem services) (Balzan et al. 2019). Nature-based solutions (NbS) are defined by IUCN as “actions to protect, sustainably manage, and restore natural or modified ecosystems that address societal challenges effectively and adaptively, simultaneously providing human well-being and biodiversity benefits”<sup>38</sup>.

The Mediterranean Strategy for Sustainable Development 2016-2025 prioritizes national action to ensure that natural water resources are extracted sustainably. However, reliable methods to assess water demand are needed. Such methods assess the dynamics and determine the main drivers of each type of water demand, and project future water uses (Charlton and Arnell 2011; Collet et al. 2013; Griffin et al. 2013; Reynard et al. 2014), with interdisciplinary approaches that combine physical and human features and incorporate climate change impacts at the local scale (Grouillet et al. 2015). Local water management planning and adaptation strategies need to be improved and updated in order to attain future water security (Koutroulis et al. 2013) (more on human security related to water is highlighted in *Chapter 5.3*). Achieving integrated and sustainable water management will also require enhanced awareness of climate change effects and public demands for water-use efficiency and improved environmental quality (Grantham et al. 2013). Awareness of the practical implications of plausible hydro-climatic and socio-economic future scenarios will shift perceptions and preference towards a more sustainable model (Koutroulis et al. 2010).

The success of mitigation and adaptation policies to restore sustainability depends on implementation efficiency at local level, where awareness and perception often pose barriers (Betzold 2015; La Jeunesse et al. 2015). Communicating relevant and targeted climate change information to stakeholders and decision makers is crucial for gaining commitment in the field. The projected water scarcity in the region highlights the important role for development and deployment of water con-

servation technologies and practices (Hejazi et al. 2014) and the need for strategic resource planning from global to regional and local scales (Koutroulis et al. 2013). Stakeholders, the beneficiaries of ecosystem services from river bodies and landscapes, play a key role in interpreting the impact of climate change on water resources and usage. A mixed methodology based on a transdisciplinary approach and the involvement of academia, policymakers, and local experts is suggested. Many physical models on the impacts of climate change and on water scarcity exist but approaches that are transdisciplinary with input from local stakeholders and interpretation of intermediary results are limited (Rault et al. 2019). Improving the understanding of ecosystem responses to multiple stressors and defining measures to improve the ecological status of water bodies are needed and sought by the WFD (Menció and Mas-Pla 2010).

Preserving the natural flow variability of rivers and streams is key in sustainable environmental management plans in the Mediterranean (Menció and Mas-Pla 2010) and critical to the long-term conservation of their unique biodiversity (Cid et al. 2017). The high variation in hydrological regimes in the region, however, tends to exacerbate the magnitude of negative responses to anthropogenic and climate impacts. For example, land use changes promote longer dry season flows, concentrating contaminants, allowing the accumulation of waste, algae, and plants, and fostering higher temperatures and lower dissolved oxygen levels, all of which may extirpate sensitive native species. Exotic species often thrive in rivers altered by human activity, further homogenizing river communities worldwide. Future research should rigorously evaluate the effects of management and restoration practices on river ecosystems, determine the cause-effect pathways leading from human disturbances to stream biological communities, and incorporate analyses of the effects of scale, land use heterogeneity, and high temporal hydrological variability on stream communities (Cooper et al. 2013).

The surface water-groundwater relationship is of major interest in the characterization of human pressures on stream hydrological dynamics and the ecological quality of Mediterranean reaches (Menció and Mas-Pla 2010). The ecological status of streams depends on an equilibrium between hydrological processes and biological dynamics. Water discharge is the main requisite for a rich riparian habitat and impacts upon aquifer water storage and

<sup>38</sup> <https://www.iucn.org/commissions/commission-ecosystem-management/our-work/nature-based-solutions>

on base flow generation have a significant effect on stream biology. Tools for managing these systems, such as those related to biomonitoring, climate change, and conservation, must be tailored to the seasonal and inter-annual variability of these systems.

Much of the ecological surface water monitoring under the WFD focuses on the assessment of biological structure, hydromorphological elements and chemical and physicochemical elements, all of which represent important information about ecosystem condition (Balzan et al., 2019). It is assumed that good ecological and chemical status have a positive impact on the capacity of ecosystems to provide ecosystem services and benefits to human well-being (Grizzetti et al. 2019). The link between waterbody condition, and ecosystem function and services has seldom been explored in detail, with the implication of water management for ecosystem services being either implicit or overlooked (Vlachopoulou et al., 2014). Thus, tools that work at the ecosystem function level, connecting ecosystem condition to services, are required especially in climate change scenarios where ecosystem condition is expected to change with the climate and other interacting drivers (*Chapter 2*). There are a number of studies from the Mediterranean Basin that demonstrate that such links can be made in practice, as demonstrated by the study by Acreman et al. (2017), which shows how the implementation of beach restoration is associated with ecological recovery and recreational ecosystem services.

Wetland management and conservation in semi-arid Mediterranean areas is necessary because they have been highly impacted by agriculture. This can be done with pressure and state indicators at landscape and wetland scales that reflect the status, condition, and trends of wetland ecosystems. Ortega et al. (2004) developed an ecological integrity index with 12 indicators (5 at the catchment scale and 7 at the wetland scale) based on the relationship between pressures from anthropogenic activities and the ecological state of wetlands and their catchments, integrating environmental, biological, economic, and social issues. Overall, a wide riparian zone acts as a buffer for wetlands, diminishing the effects of intensive agriculture. Provisioning services are more relevant in normal and wet years, while regulating service water purification provides higher benefits in dry years, when threats to water quality are increased because of a decreased dilution capacity (Terrado et al. 2014). Protecting water towers in semi-arid regions expected to experience dramatic changes is essential to ensuring water provisioning in dry years. However, the protection

of water resources is not sufficient if consumption rates continue or increase in the future. Actions should be planned to enhance the provision of regulating services (Terrado et al. 2014). Overall, in semi-arid basins under continuous human impact, hydrological ecosystem services are very sensitive to climate extremes, and service supply and demand areas are usually spatially and temporally decoupled. Both aspects are relevant and need to be considered in basin management in semi-arid regions (Terrado et al. 2014).

Studies that consider the respective influences of climate, land cover (forest cover dynamics in hydrological processes) and water withdrawals on water availability (e.g., Chauvelon et al. 2003; Varela-Ortega et al. 2011) are required for proper adaptation measures but are still scarce, since the required data are often unavailable or not easily accessible over long periods (Sivapalan et al. 2003). Databases at large spatial and temporal scales are key to understanding the variability of hydrological systems, and in providing water managers with science-based decision-making support information (Hannah et al. 2011). However, such databases are still too scarce in the Mediterranean, despite efforts to maintain and develop data networks at the regional and global levels. At this scale, the physical and human characteristics of catchments, especially in the Mediterranean, are extremely heterogeneous since they encompass extreme contrasts in terms of climatic, topographic and geological characteristics, population distribution and water uses and are therefore difficult to define and grasp (Collet et al. 2014). The climate variability of Mediterranean river basins also makes it difficult to describe general patterns which explain and predict the relationships between runoff, erosion and sediment transport (López-Tarazón et al. 2010; de Vente et al. 2011) and this is even more complicated in rivers affected by regulation. Time-series on sediment transport (Batalla and Vericat 2009) and on lake ecosystems, especially those predating anthropogenic influences, are very scarce (Papastergiadou et al. 2007). Long-term data sets are also important for understanding the interactions among native species and introduced species. These are particularly valuable in understanding the influence of extreme events such as drought and floods (Magalhães et al. 2007; Bêche et al. 2009). Such studies, particularly in regulated systems, help guide flow recommendations to benefit native species but have only begun, although their value is already clear (Kiernan et al. 2012; Resh et al. 2013). Long-term data sets that can reveal trends need to exceed the sub-decade scale for sustainable management of Mediterranean-climate streams and rivers (Cid et al. 2017).

## BOX 4.1

### Bio-indicators for the assessment of changes in Mediterranean marine ecosystems

Awareness of recent changes in ecological conditions in many seas has fostered a need to assess increasing anthropogenic pressures and their consequences on sediment and water quality, and to suggest measures to reverse this trend. In this context, the European Commission has implemented the Water Framework Directive (WFD, Directive 2000/60/EC) with the aim to obtain (or to maintain) a “good status” for all European waters by 2015. To support this, a large number of monitoring tools have been developed, including several bio-indicators such as phytoplankton, macro-algae, seagrass, angiosperms, fish faunas and soft substrate benthic invertebrate fauna, which are benthic foraminifera.

Phytoplankton and zooplankton are ecologically important groups in most aquatic ecosystems and have been an important component of biological monitoring programs in the Mediterranean (Abboud-Abi Saab et al. 2008, 2012; Tunin-Ley et al. 2009; Gharib et al. 2011; Tunin-Ley and Lemée 2013; Abo-Taleb et al. 2016; Ouba et al. 2016; Abboud-Abi Saab and Hassoun 2017). The genus *Neoceratium* (planktonic dinoflagellates) in the NW Mediterranean is known to be particularly sensitive to water temperature, and is responsive to global warming (Tunin-Ley et al. 2009; Tunin-Ley and Lemée 2013). Moreover, the WFD mandates the use of biological quality element (BQE) phytoplankton to assess the ecological status of coastal and transitional water bodies. Alternatively, Camp et al. (2016) propose a methodology to assess water-quality based on the use of chlorophyll-a (Chl-a), as a proxy of phytoplankton biomass.

For soft-bottom marine habitats, macrofauna is traditionally used as a bio-indicator, and a wide range of different biotic indices have been developed (Borja et al. 2016). The use of meiofauna, occurring in higher densities, is less developed. Among these, benthic foraminifera appear particularly suitable for bio-monitoring in the Mediterranean (Barras et al. 2014; Jorissen et al. 2018). The abundant and diverse benthic foraminifera faunas in the Mediterranean react rapidly to environmental changes such as organic pollution, eutrophication and oxygen depletion. These characteristics led to the development of a standardized biotic index based on foraminifera (Jorissen et al. 2018).

For the coralligenous, several indices of its health status have been suggested. One is the INDEX-COR approach, based on long time series of photographic sampling, standardized and used as a large spatial comparison tool (Sartoretto et al. 2017). This type of index integrates the sensitivity of different coralligenous taxa to organic matter and sediment deposition, the observable taxonomic richness and the structural complexity of the benthic assemblages. With these approaches, the health status of this complex ecosystem can be assessed without invasive or directly impacting methods. When deeper areas are considered in the coralligenous assemblages (e.g., from 30–40 to 200 m depth), Remotely Operated Vehicle (ROV) approaches may be useful (Rossi et al. 2008). The Mesophotic

Assemblages Ecological Status (MAES) has been suggested as a tool for conservation and management procedures (Cánovas-Molina et al. 2016). The MAES index is based on community structure, condition of the erect species and visible human impacts (Cánovas-Molina et al. 2016). A combined biomarker index can also be considered as a medium-long term monitoring approach. In selected populations of representative sessile species of the coralligenous, activity (e.g., polyp expansion), growth, reproductive output, stable isotopes, biochemical balance (protein-carbohydrate-lipids), fatty acids and C/N ratio may be used to estimate the nutritional condition and health status of populations or entire communities, considering the biology and ecology of each species (Rossi et al. 2017b).

For large vertebrates, cetaceans and seabirds are widely regarded as reliable indicators of the health of marine ecosystems due to their position near the top of the marine food web, conspicuous nature, and reliance on marine resources (Durant et al. 2009; Bossart 2011; Schwacke et al. 2013; Fossi and Panti 2017; Fossi et al. 2018). Some cetaceans and seabirds are reported as sentinels or indicators for the state of marine ecosystems because they are globally subject to multiple stress factors, such as the bioaccumulation of contaminants, infectious diseases, non-indigenous species, food depletion, and climate change (UNEP/MAP 2012; Poloczanska et al. 2013). The advantage of using cetaceans as sentinels is that they have physiology and/or diets similar to those of humans, so they can indicate earlier potential adverse health effects (Schwacke et al. 2013). For fish, red mullets (*Mullus barbatus* and *M. surmuletus*) have been widely used as quantitative bio-indicators of chemical contamination (Porte et al. 2002; Storelli and Marcotrigiano 2005; Martínez-Gómez et al. 2017). For instance, the recent study by Cresson et al. (2014) confirmed that red mullets are efficient bio-indicators of Mercury (Hg), one of the main chemicals currently altering Mediterranean ecosystems.

## BOX 4.2

### Urban biodiversity in the Mediterranean Region

Further to being a hotspot of biodiversity, the Mediterranean region is also one of the most densely urbanized areas in the world (FAO and Plan Bleu 2018). The overall population in the region grew by 190 million people between 1970 and 2010, while the urban population increased by 163 million, with more 74% of population growth concentrated in the countries of the southern part of the Mediterranean (UNEP/MAP 2012). Despite substantially lower growth rates of urban populations in European countries of the Mediterranean, surfaces occupied by urban settlements have also increased considerably due to tourism and decentralization of population from high density core cities towards low density residential areas and along the coastlines, as for instance in Barcelona (Domene and Sauri 2007) or Athens (Cecchini et al. 2019).

Also, patterns of land use changes differ between southern and northern parts of the region. In European countries, urbanization has been accompanied by abandonment of agriculture since the post war period and extended mainly onto different types of cultivated areas, and onto shrubland and/or herbaceous areas (García-Nieto et al. 2018), leading to substantial losses of biodiversity and of agricultural area, while shrubland and forest land increased (Domene and Sauri 2007; FAO and Plan Bleu 2018; García-Nieto et al. 2018).

The growth of north African peri-urban areas in that period occurred in parallel with an increase in irrigated arable land, permanent crops, complex cultivation patterns and shrublands and/or herbaceous areas and pastures, at the expense of non-irrigated arable land and forest, both around peri-urban areas, as well as at the national level. In southern Mediterranean countries, environmental change contributed, for example, to a rural exodus in Morocco

between 1980 and 1990, and in Algeria and Tunisia in 1999 (García-Nieto et al. 2018).

#### *Consequences for biodiversity and ecosystem services in urban areas*

Consequences for biodiversity and ecosystem services available for urban areas differ accordingly. In most cases, urban areas replace former agricultural land, but, while in the southern part of the Mediterranean this coincides with an extension of cultivated areas outside urbanized areas, in most European countries the contemporary abandonment of agricultural areas leads to an increase in shrubland and uncultivated areas and contributes to increasing vulnerability of surfaces, for instance with the threat of wildfires along the wildland-urban interface (San-Miguel-Ayanz et al. 2013; Laforteza et al. 2015; Xanthopoulos 2015) and desertification (Salvati et al. 2015). Abandonment of agricultural lands also leads to the loss of cultural landscape management practices. For example, traditional Mediterranean agricultural landscapes are in many places characterized by terraces and dry walls which represent small scale practices of erosion prevention (Cecchini et al. 2019). Increasing soil sealing in urban areas and connected infrastructure and abandonment of historic techniques of landscape management lead to increasing risks from flooding and landslides (Salvati et al. 2015; García-Nieto et al. 2018).

The remaining ecosystems are increasingly under threat with particular risks for rare plants depending on small patch ecosystems with highly localized distributions even though they may be protected (Vimal et al. 2012). Peri-urban areas nevertheless provide relevant services for urban populations, for instance the increase in areas of natural or semi-natural

## BOX 4.3

### Nitrogen deposition and ecosystems

Climate change contributes to an increase in dry deposition of nitrogen and increases the negative impacts of excess atmospheric nitrogen on biodiversity (Oliveira et al. 2020). Reactive nitrogen (Nr) impacts vegetation through direct foliar damage, eutrophication, acidification, and susceptibility to secondary stress depending on the nitrogen form and concentration (Krupa 2003). Grassland, heathland and forest ecosystems are recognized as habitats vulnerable to Nr in Europe (Dise et al. 2011). In Spain, natural grasslands, particularly in the northern alpine area, were found to be the most threatened habitat followed by mountain ecosystems (García-Gómez et al. 2014). At least 14% of the Natura 2000 sites in western Iberia are at risk of eutrophication (Oliveira et al. 2020).

It is not yet clear if different wet-deposited forms of Nr (e.g., nitrate,  $\text{NO}_3^-$  – versus ammonium,  $\text{NH}_4^+$ ) have different effects on biodiversity. However, gaseous ammonia ( $\text{NH}_3$ ) can be particularly harmful to vegetation. The highest relative risk of biodiversity change in Natura 2000 sites due to  $\text{NH}_3$

pollution in Portugal was found to be in peats, mires, bogs, and similar acidic and oligotrophic habitats (most located in the northern mountains), whereas in the Atlantic and Mediterranean climate zone (coastal, tidal, and scrubland habitats) they were deemed the least sensitive in Portugal (Pinho et al. 2018).

Exceedance of critical loads for nitrogen is linked to reduced plant species richness in a broad range of European ecosystems (Dise et al. 2011). Experimental evidence shows that species richness and abundance resulted in larger declines with greater amounts of annual N addition including in semi-arid areas (Midolo et al. 2019). Reductions in the abundance of individual species were greater for N-sensitive plant life-form types (legumes and non-vascular plants) (Ochoa-Hueso et al. 2014, 2017; Midolo et al. 2019).

Several conservation plants (e.g., orchids and carnivorous) and cryptogams are naturally adapted to low environment N supply. Thus, increasing Nr alters the natural ecological balance. This results in the loss

## BOX 4.2

vegetation such as ecosystem fragments, reserves, nature parks, forests, and river banks, which house varying amounts of native species, potentially provide cultural services. Formal and informal forms of urban agriculture and horticulture (Domene and Saurí 2007; Cecchini et al. 2019) present an opportunity for integration into diets for urban residents and provide both provisioning and cultural services, as these areas are also used for leisure activities (Domene and Saurí 2007; Cecchini et al. 2019; Palau-Salvador et al. 2019).

The expansion of peri-urban agriculture, for instance, olive cultivation for self-consumption or small-scale economic production, have counteracted land use change in the peri-urban areas of several cities in the Mediterranean Basin (García-Nieto et al. 2018; Cecchini et al. 2019). Olive landscapes have a high tolerance to pests and are characterized by a stable trend of economic production, and abundant insect fauna contributing to biodiversity conservation. They furthermore show a higher resistance to wildfires than other Mediterranean vegetation. The economic stability of olive oil production in small groves, like other small-scale agricultural areas and the status of protected natural areas (e.g., coastal woods in the case of Rome) in the green belt around cities contributes to the ability of these areas to form an efficient barrier against urban dispersion and reduce the impacts of soil sealing on the hydrological cycle and on ecosystem services loss (Salvati et al. 2015; Cecchini et al. 2019).

#### **Urban biodiversity and ecosystem services**

Despite the important provisioning and cultural services and their increasing popularity in many cities, urban gardening seems to be not as effective in protecting peri-urban areas from land use changes as

in the case of peri-urban agriculture observed in Athens (Domene and Saurí 2007; Heywood 2017; Cecchini et al. 2019). This may be due to less stable legal position of such areas which are often the result of squatting on private or, more often, public land (Domene and Saurí 2007) and their less consolidated economic status. In Rome, managed spaces with cultivated vegetation such as parks and gardens represent a lower level of resilience against transformation despite their importance for cultural and regulating services (e.g., leisure and heat mitigation) for urban residents (Salvati et al. 2015).

In many urban areas of the Mediterranean, street trees provide important regulating services for human well-being by offering shade and reducing heat impacts during summer due to their evaporation rates and the albedo created by foliage (Rana and Ferrara 2019). They also provide important cultural services as characterizing elements of Mediterranean urban landscapes (Heywood 2017).

Furthermore, urban wastelands and shrublands which are mainly colonized by weeds, ruderal plants and non-indigenous species, in many cases house considerable numbers of native plants and are potential places for valuable biodiversity (Heywood 2017). The importance of green spaces in urban areas is increasingly recognized by Mediterranean cities, which are increasingly engaging in urban green infrastructure projects, preserving remnants of biodiversity and natural areas within cities. These are expected to provide important regulating, cultural and provisioning services but there is a general lack of data on urban biodiversity in urban and peri-urban areas of the Mediterranean (Heywood 2017).

## BOX 4.3

of the most sensitive species, which are often a priority for protection, and their replacement by non-indigenous or other opportunistic species that prefer high rates of nitrogen supply (Bobbink et al. 2010). Lichens and bryophytes are among the most sensitive organisms to N pollution at the ecosystem level (Cape et al. 2009), having a different response depending on their functional response group (Pinho et al. 2008, 2009, 2011, 2012b, 2012a; Jovan et al. 2012).

There are some clear examples of reductions in faunal diversity that can be linked to Nr deposition, but overall, our knowledge of faunal effects is still limited (Dise et al. 2011). Changes to above-ground faunal communities probably occur primarily through changes in vegetation diversity, composition or structure (Murray et al. 2006). The evidence strongly suggests that ecological communities respond to the accumulated pool of plant-available N in the soil. Thus, the cumulative load of enhanced Nr impacting an ecosystem is probably important (Stevens et al. 2011). Because of this response to cumulative inputs, it is likely that biodiversity has been in decline in Europe for many decades due

to enhanced Nr deposition (Bobbink et al. 2010). Equally, full recovery in response to reduced Nr deposition is likely to be slow, especially in highly impacted ecosystems. In some cases, recovery may require management intervention.



## BOX 4.4

## Mediterranean islands

*Islands as laboratories*

The high concentration of islands (> 10,000) is one of the features which contributes to the Mediterranean's unique character, placing the region within the richest in the world in terms of islands and archipelagos (Médail 2017). In addition to being biological laboratories, the largest of the islands are also the centers of many of the world's ancient civilizations (Patton 1996). On these islands, the diversity of biogeography, geology and human settlement has produced exceptionally high numbers of biodiversity and endemism, earning them a place in the global biodiversity hotspots list (Médail and Quézel 1997; Vogiatzakis et al. 2016; Médail 2017). In addition to the role as tertiary and glacial refuges, islands have also contributed to more recent plant diversification (Médail 2017). Biodiversity on islands display an insular syndrome due to abiotic conditions (nature of isolation, particular climate) and their own biogeographical history, and are characterized by unique specific assembly (with several endemics) and biotic interactions (de Montmollin and Strahm 2005; Blondel et al. 2010; Médail 2013, 2017; Schatz 2017).

*Recent evidence of change*

Despite their relatively small contribution to greenhouse gas emissions, Mediterranean islands are likely to be adversely affected by climate change, in synergy with ongoing land use changes.

- **Land-use change:** The landscapes of Mediterranean islands have evolved as the result of similar pressures to the mainland generated by socio-economic and political factors but amplified on a "matrix" of limited space. Land use changes and associated impacts differ significantly depending on the size of the island and therefore demographics, as well as its popularity as tourist destination (Vogiatzakis et al. 2008). Therefore, to date there is no consensus on the trends of changes since islands seem to respond/ behave individually (Vogiatzakis et al. 2008; Harris 2012). Recent land uses are associated with temporal and spatial shifts in land-use systems with polarization of land-use intensity, particularly on small/medium size islands (Tzanopoulos and Vogiatzakis 2011; Balzan et al. 2018).
- **Climate evidence:** Observed trends for winter (Nov–Feb) precipitation (mm/50 years) and summer (Jun–Sep) temperature (°C/50 years) for Mediterranean island regions during the second half of the twentieth century do not show a consistent climate pattern (Vogiatzakis et al. 2016).
- **Biological/ecosystems evidence:** Documented evidence on plant and animal phenology changes (Peñuelas et al. 2002; Gordo and Sanz 2010), range shifts (Lenoir et al. 2008) and changes in the function, structure and dynamics of ecosystems e.g., temporal mismatches among mutualistic partners (Visser et al. 2004), species loss and changes (+/-) in species richness (Kazakis et al. 2007). Plant communities are steadily changing, such as orchids in Corsica (Vogt-Schilb et al. 2016), as well as specialized plant-insect

interactions (pollination, seed dispersal) (Traveset and Riera 2005; Blondel et al. 2010; Stefanaki et al. 2015).

*Climate change projections and islands*

- **Current scenarios:** Projected trends from various climate models agree as far as the direction of change in precipitation and temperature regimes are concerned (Table 2.1 in Section 2.2) (Vogiatzakis et al. 2016).
- **Sea level rise:** For the Aegean archipelagos, Monioudi et al. (2017) assume mean sea-level rise of 0.5 m for RCP4.5 and predict that a storm-induced sea level rise of 0.6 m would result in complete erosion of between 31 and 88% of all beaches, at least temporarily.
- **Island representation and model resolution:** In most modeling studies (niche models or GCMs), islands are simply a subset of the Mediterranean (Araújo et al. 2006; Settele et al. 2008). As a result, neither distribution nor climate data have sufficient resolution to allow climate envelope models for most endemic island taxa (Henle et al. 2010).
- **Synergies with land cover changes:** Changes in land use (Settele et al. 2005) coupled with climate models predict modifications to species climate space (Settele et al. 2008) and islands are no different. Documented land cover changes related to urban/tourism development and increasing linear infrastructure are already having an impact on island biodiversity (Zomeni and Vogiatzakis 2014).

*Vulnerability/resilience*

The vulnerability of Mediterranean island systems to past and recent extinctions has been well documented (de Montmollin and Strahm 2005; Foufopoulos et al. 2011). A recent assessment of global imminent extinctions includes two Mediterranean Islands (Ricketts et al. 2005). Compared to the rest of the Mediterranean Basin, islands have always been more vulnerable to invasion by exotic species (Hulme 2004). Human activity will be the limiting factor which will determine the future of island flora. Islands are representative examples in the Mediterranean of the co-evolution of social-ecological systems intensified by the element of insularity and which are currently under threat and more susceptible to externalities due to (i) limited resources/space, (ii) administrative/political leverage, (iii) institutional capacities. While many of the islands have experienced demographic losses, concerning their permanent inhabitants, they have become principal tourist destinations (Ioannides et al. 2001) and islands are being faced with a key challenge of balancing economic benefits from ecosystem services delivery (tourism, agriculture) with environmental pressures.

*Conservation and adaptation*

On islands, opportunities for (human assisted) adaptation are limited. The lack of available space for wildlife to shift presents significant barriers to the natural adaptation of species and habitats. It also leads to more intense land-use conflicts, therefore increasing size of protected areas, and connections might be problematic. In an



attempt to promote "climate-wise connectivity" (Keeley et al. 2018), areas not (significantly) affected by climate change could act as refugia to species in the future. Many of the island endemic species of today have survived past climate changes in places in such refugia (Vogiatzakis and Griffiths 2008; Vogiatzakis 2012). In the case of island clusters, the suitable climate space might be a neighboring island, although there are many examples, particularly in the Aegean, where neighboring islands have different floras (Kallimanis et al. 2010). Therefore, biogeography may be more important than climate *per se* in interpreting species distribution patterns (Whittaker and Fernandez-Palacios 2007). Building a coherent "network" of protected areas across islands (e.g., in the Aegean) might provide solutions to safeguarding common biotic elements (species or habitats). At the government level, adaptation should include increased institutional capacity for innovation, the increase of monitoring activities, adaptive management, and promotion of inter-island collaboration (Kark et al. 2009). In the case of managing island ecosystems and their services, the key priorities for the future must be to:

- identify ecosystem service capacity hotspots and how they can be affected under climate change scenarios;
- identify ecosystem service demand and flows in hotspots and manage green infrastructure and co-created nature-based solutions to provide synergies for biodiversity conservation and ecosystem services for human well-being;
- manage protected areas for ecosystem service provision given the fact that lack of space is also problematic;
- evaluate tradeoffs and thresholds of ecosystem service provision and assess the impacts from demographics and tourism, as well as land use and climate change projections.

## References

- Abbound-Abi Saab M 2008 *Tintinnids of the Lebanese coastal waters (Eastern Mediterranean)*. CNRS-Lebanon/ UNEP/MAP/RAC/SPA, 192pp.
- Abbound-Abi Saab M 2009 Studies and changes of phytoplankton populations on the Lebanese coastal waters - a brief overview. in *CIESM Workshop Monographs n°40, Phytoplankton Response to Mediterranean Environmental Change*, 79–82.
- Abbound-Abi Saab M, Bitar G, Harmelin-Vivien M, Harmelin JG, Romano JC et al. 2003 Environnement côtier et biodiversité marine sur les côtes libanaises; inventaire et mise en place d'un ensemble matériel et humain d'observation et d'analyse de leur évolution, degré d'altération des communautés benthiques littorales. Rapport final Franco-Lebanese Cooperation Program CEDRE (1999-2002).
- Abbound-Abi Saab M, Fakhri M, Hassoun AER, Tilbian M, Kassab M-T et al. 2012 Effects of continental input on marine environment in the Lebanese coastal waters. in *INOC-CNRS, International Conference on "Land-Sea Interactions in the Coastal Zone", 06-08 November 2012 (Lebanon)*, 370–382. doi: [10.13140/2.1.4400.9767](https://doi.org/10.13140/2.1.4400.9767) 06-08 November–2012
- Abbound-Abi Saab M, Fakhri M, Kassab M-T, Matar N 2008 Développement exceptionnel des eaux colorées au printemps 2007 dans la zone côtière libanaise entre Zouk-Naher el kelb. *Leban. Sci. J.* 9, 61–70.
- Abbound-Abi Saab M, Hassoun AER 2017 Effects of organic pollution on environmental conditions and the phytoplankton community in the central Lebanese coastal waters with special attention to toxic algae. *Reg. Stud. Mar. Sci.* 10, 38–51. doi: [10.1016/j.rsma.2017.01.003](https://doi.org/10.1016/j.rsma.2017.01.003)
- Abdulla A, Gomei M, Hyrenbach D, Notarbartolo di Sciarra G, Agardy T 2008 Challenges facing a network of representative marine protected areas in the Mediterranean: prioritizing the protection of underrepresented habitats. *ICES J. Mar. Sci.* 66, 22–28. doi: [10.1093/icesjms/fsn164](https://doi.org/10.1093/icesjms/fsn164)
- Abis S 2006 Les dynamiques démographiques en Méditerranée. *Les Notes Danalyse Du CIHEAM* 11, 27.
- Abo-Taleb HA, Abdel Aziz NE, Aboul Ezz SM, El Raey M, Abou Zaid MM 2016 Study of Chromista and Protozoa in a hotspot area at the Mediterranean coast with special reference to the potentiality to use it as bio-indicators. *Int. J. Mar. Sci.* 6. <http://biopublisher.ca/index.php/ijms/article/view/2911> [Accessed March 26, 2019]
- Abrantes FG, Lebreiro S, Rodrigues T, Gil I, Bartels-Jónsdóttir HB et al. 2005 Shallow-marine sediment cores record climate variability and earthquake activity off Lisbon (Portugal) for the last 2000 years. *Quat. Sci. Rev.* 24, 2477–2494. doi: [10.1016/j.quascirev.2004.04.009](https://doi.org/10.1016/j.quascirev.2004.04.009)
- Abrantes FG, Voelker AH, Sierro FJ, Naughton F, Rodrigues T et al. 2012 Paleoclimate Variability in the Mediterranean Region, in *The Climate of the Mediterranean Region*, ed. Lionello P (Oxford: Elsevier), 1–86. <http://www.sciencedirect.com/science/article/pii/B978012416042200001X>
- Accoroni S, Romagnoli T, Penna A, Capellacci S, Ciminiello P et al. 2016 *Ostreopsis fattorussoi* sp. nov. (Dinophyceae), a new benthic toxic *Ostreopsis* species from the eastern Mediterranean Sea. *J. Phycol.* 52, 1064–1084. doi: [10.1111/jpy.12464](https://doi.org/10.1111/jpy.12464)
- Acosta ATR, Ercole S, Stanisci A, Pillar VDP, Blasi C 2007 Coastal vegetation zonation and dune morphology in some Mediterranean ecosystems. *J. Coast. Res.* 23, 1518–1524. doi: [10.2112/05-0589.1](https://doi.org/10.2112/05-0589.1)
- Acreman M, Ludi E, Oates NEM, Parker H, Moncrieff CR et al. 2017 Managing Rivers for Multiple Benefits—A Coherent Approach to Research, Policy and Planning. *Front. Environ. Sci.* 5, 1–8. doi: [10.3389/fenvs.2017.00004](https://doi.org/10.3389/fenvs.2017.00004)
- Adloff F, Somot S, Sevault F, Jordà G, Aznar R et al. 2015 Mediterranean Sea response to climate change in an ensemble of twenty first century scenarios. *Clim. Dyn.* 45, 2775–2802. doi: [10.1007/s00382-015-2507-3](https://doi.org/10.1007/s00382-015-2507-3)
- Ágreda T, Águeda B, Olano JM, Vicente-Serrano SM, Fernández-Toirán M 2015 Increased evapotranspiration demand in a Mediterranean climate might cause a decline in fungal yields under global warming. *Glob. Chang. Biol.* 21, 3499–3510. doi: [10.1111/gcb.12960](https://doi.org/10.1111/gcb.12960)
- Aguiar FC, Ferreira MT 2005 Human-disturbed landscapes: effects on composition and integrity of riparian woody vegetation in the Tagus River basin, Portugal. *Environ. Conserv.* 32, 30–41. doi: [10.1017/s0376892905001992](https://doi.org/10.1017/s0376892905001992)
- Aguiar FC, Segurado P, Martins MJ, Bejarano MD, Nilsson C et al. 2018 The abundance and distribution of guilds of riparian woody plants change in response to land use and flow regulation. *J. Appl. Ecol.* 55, 2227–2240. doi: [10.1111/1365-2664.13110](https://doi.org/10.1111/1365-2664.13110)
- Aguilera E, Lassaletta L, Gattinger A, Gimeno BS 2013 Managing soil carbon for climate change mitigation and adaptation in Mediterranean cropping systems: A meta-analysis. *Agric. Ecosyst. Environ.* 168, 25–36. doi: [10.1016/j.agee.2013.02.003](https://doi.org/10.1016/j.agee.2013.02.003)
- Aissaoui A, Armi Z, Akrouf F, Ben Hassine OK 2014 Environmental factors and seasonal dynamics of *Prorocentrum lima* population in coastal waters of the Gulf of Tunis, South Mediterranean. *Water Environ. Res.* 86, 2256–2270. doi: [10.2175/106143014x13975035526266](https://doi.org/10.2175/106143014x13975035526266)
- Ait-Mouheb N, Bahri A, Thayer B Ben, Benyahia B, Bourrié G et al. 2018 The reuse of reclaimed water for irrigation around the Mediterranean Rim: a step towards a more virtuous cycle? *Reg. Environ. Chang.* 18, 693–705. doi: [10.1007/s10113-018-1292-z](https://doi.org/10.1007/s10113-018-1292-z)
- Aizen MA, Harder LD 2009 The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Curr. Biol.* 19, 915–918. doi: [10.1016/j.cub.2009.03.071](https://doi.org/10.1016/j.cub.2009.03.071)
- Akyol O, Unal V, Ceyhan T, Bilecenoglu M 2005 First con-

- firmed record of *Lagocephalus sceleratus* (Gmelin, 1789) in the Mediterranean Sea. *J. Fish Biol.* 66, 1183–1186. doi: [10.1111/j.0022-1112.2005.00667.x](https://doi.org/10.1111/j.0022-1112.2005.00667.x)
- Al Hassan M, Chaura J, López-Gresa MP, Borsari O, Daniso E et al. 2016 Native-Invasive Plants vs. Halophytes in Mediterranean Salt Marshes: Stress Tolerance Mechanisms in Two Related Species. *Front. Plant Sci.* 7, 473. doi: [10.3389/fpls.2016.00473](https://doi.org/10.3389/fpls.2016.00473)
- Alary V, Moulin C-H, Lasseur J, Aboulnaga A, Srairi T 2019 The dynamic of crop-livestock systems in the Mediterranean and future prospective at local level: A comparative analysis for South and North Mediterranean systems. *Livest. Sci.* 224, 40–49. doi: [10.1016/J.LIVSCI.2019.03.017](https://doi.org/10.1016/J.LIVSCI.2019.03.017)
- Albouy C, Guilhaumon F, Araújo MB, Mouillot D, Leprieux F 2012 Combining projected changes in species richness and composition reveals climate change impacts on coastal Mediterranean fish assemblages. *Glob. Chang. Biol.* 18, 2995–3003. doi: [10.1111/j.1365-2486.2012.02772.x](https://doi.org/10.1111/j.1365-2486.2012.02772.x)
- Albouy C, Guilhaumon F, Leprieux F, Lasram FBR, Somot S et al. 2013 Projected climate change and the changing biogeography of coastal Mediterranean fishes. *J. Biogeogr.* 40, 534–547. doi: [10.1111/jbi.12013](https://doi.org/10.1111/jbi.12013)
- Albrich K, Rammer W, Thom D, Seidl R 2018 Trade-offs between temporal stability and level of forest ecosystem services provisioning under climate change. *Ecol. Appl.* 28, 1884–1896. doi: [10.1002/eap.1785](https://doi.org/10.1002/eap.1785)
- Alexander P, Rounsevell MDA, Dislich C, Dodson JR, Engström K et al. 2015 Drivers for global agricultural land use change: The nexus of diet, population, yield and bioenergy. *Glob. Environ. Chang.* 35, 138–147. doi: [10.1016/J.GLOENVCHA.2015.08.011](https://doi.org/10.1016/J.GLOENVCHA.2015.08.011)
- Alexander RW, Calvo A 1990 The influence of lichens on slope processes in some Spanish badlands, in *Vegetation and Erosion*, ed. Thornes JB (New York: John Wiley), 385–398.
- Aliaume C, Do Chi T, Viaroli P, Zaldívar JM 2007 Coastal lagoons of Southern Europe: recent changes and future scenarios. *Transitional Waters Monogr.* 1, 1–12. doi: [10.1285/i18252273v1n1p1](https://doi.org/10.1285/i18252273v1n1p1)
- Aligizaki K 2009 Spread of potentially toxic benthic dinoflagellates in the Mediterranean Sea: A response to climate change ? in *Phytoplankton Response to Mediterranean Environmental Change, CIESM Workshop Monographs n°40- Tunis, 7 - 10 October 2009.*, ed. Briand F (Paris, France), 57–61.
- Aligizaki K, Nikolaidis G 2006 The presence of the potentially toxic genera *Ostreopsis* and *Coolia* (Dinophyceae) in the North Aegean Sea, Greece. *Harmful Algae* 5, 717–730. doi: [10.1016/j.hal.2006.02.005](https://doi.org/10.1016/j.hal.2006.02.005)
- Allan JD 2004 Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems. *Annu. Rev. Ecol. Syst.* 35, 257–284. doi: [10.1146/annurev.ecolsys.35.120202.110122](https://doi.org/10.1146/annurev.ecolsys.35.120202.110122)
- Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N et al. 2010 A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *For. Ecol. Manage.* 259, 660–684. doi: [10.1016/j.foreco.2009.09.001](https://doi.org/10.1016/j.foreco.2009.09.001)
- Allen JRL 2000 Morphodynamics of Holocene salt marshes: A review sketch from the Atlantic and Southern North Sea coasts of Europe. *Quat. Sci. Rev.* 19, 1155–1231. doi: [10.1016/S0277-3791\(99\)00034-7](https://doi.org/10.1016/S0277-3791(99)00034-7)
- Allen MR, Ingram WJ 2002 Constraints on future changes in climate and the hydrologic cycle. *Nature* 419, 228–232. doi: [10.1038/nature01092](https://doi.org/10.1038/nature01092)
- Alomar C, Deudero S 2017 Evidence of microplastic ingestion in the shark *Galeus melastomus* Rafinesque, 1810 in the continental shelf off the western Mediterranean Sea. *Environ. Pollut.* 223, 223–229. doi: [10.1016/j.envpol.2017.01.015](https://doi.org/10.1016/j.envpol.2017.01.015)
- Alomar C, Deudero S, Andaloro F, Castriota L, Consoli P et al. 2016 *Caulerpa cylindracea* Sonder invasion modifies trophic niche in infralittoral rocky benthic community. *Mar. Environ. Res.* 120, 86–92. doi: [10.1016/j.marenvres.2016.07.010](https://doi.org/10.1016/j.marenvres.2016.07.010)
- Amatulli G, Camia A, San-Miguel-Ayanz J 2013 Estimating future burned areas under changing climate in the EU-Mediterranean countries. *Sci. Total Environ.* 450–451, 209–222. doi: [10.1016/j.scitotenv.2013.02.014](https://doi.org/10.1016/j.scitotenv.2013.02.014)
- Amelung B, Viner D 2006 Mediterranean tourism: Exploring the future with the tourism climatic index. *J. Sustain. Tour.* 14, 349–366. doi: [10.2167/jost549.0](https://doi.org/10.2167/jost549.0)
- Améztegui A, Brotons L, Coll L 2010 Land-use changes as major drivers of mountain pine (*Pinus uncinata* Ram.) expansion in the Pyrenees. *Glob. Ecol. Biogeogr.* 19, 632–641. doi: [10.1111/j.1466-8238.2010.00550.x](https://doi.org/10.1111/j.1466-8238.2010.00550.x)
- Améztegui A, Cabon A, de Cáceres M, Coll L 2017 Managing stand density to enhance the adaptability of Scots pine stands to climate change: A modelling approach. *Ecol. Modell.* 356, 141–150. doi: [10.1016/j.ecolmodel.2017.04.006](https://doi.org/10.1016/j.ecolmodel.2017.04.006)
- Améztegui A, Coll L, Brotons L, Ninot JM 2016 Land-use legacies rather than climate change are driving the recent upward shift of the mountain tree line in the Pyrenees. *Glob. Ecol. Biogeogr.* 25, 263–273. doi: [10.1111/geb.12407](https://doi.org/10.1111/geb.12407)
- Anderegg WRL, Kane JM, Anderegg LDL 2013 Consequences of widespread tree mortality triggered by drought and temperature stress. *Nat. Clim. Chang.* 3, 30–36. doi: [10.1038/nclimate1635](https://doi.org/10.1038/nclimate1635)
- Andersen CB, Lewis GP, Sargent KA 2004 Influence of wastewater-treatment effluent on concentrations and fluxes of solutes in the Bush River, South Carolina, during extreme drought conditions. *Environ. Geosci.* 11, 28–41. doi: [10.1306/eg.10200303017](https://doi.org/10.1306/eg.10200303017)
- Anderson-Teixeira KJ, Miller AD, Mohan JE, Hudiburg TW, Duval BD et al. 2013 Altered dynamics of forest recovery under a changing climate. *Glob. Chang. Biol.* 19, 2001–2021. doi: [10.1111/gcb.12194](https://doi.org/10.1111/gcb.12194)
- Annicchiarico P 2017 Feed legumes for truly sustainable crop-animal systems. *Ital. J. Agron.* 12. doi: [10.4081/ija.2017.880](https://doi.org/10.4081/ija.2017.880)

- Anthelme F, Villaret J, Brun J 2007 Shrub encroachment in the Alps gives rise to the convergence of sub-alpine communities on a regional scale. *J. Veg. Sci.* 18, 355–362. doi: [10.1111/j.1654-1103.2007.tb02547.x](https://doi.org/10.1111/j.1654-1103.2007.tb02547.x)
- Anthony EJ, Marriner N, Morhange C 2014 Human influence and the changing geomorphology of Mediterranean deltas and coasts over the last 6000 years: From progradation to destruction phase? *Earth-Science Rev.* 139, 336–361. doi: [10.1016/j.earscirev.2014.10.003](https://doi.org/10.1016/j.earscirev.2014.10.003)
- Aparicio E, Vargas MJ, Olmo JM, de Sostoa A 2000 Decline of native freshwater fishes in a Mediterranean watershed on the Iberian Peninsula: A quantitative assessment. *Environ. Biol. Fishes* 59, 11–19. doi: [10.1023/A:1007618517557](https://doi.org/10.1023/A:1007618517557)
- APCOR 2015 *APCOR year book*. APCOR [http://www.apcor.pt/wp-content/uploads/2017/12/Boletim\\_Estatistico\\_APCOR\\_17\\_18.pdf](http://www.apcor.pt/wp-content/uploads/2017/12/Boletim_Estatistico_APCOR_17_18.pdf).
- Araújo MB, Thuiller W, Pearson RG 2006 Climate warming and the decline of amphibians and reptiles in Europe. *J. Biogeogr.* 33, 1712–1728. doi: [10.1111/j.1365-2699.2006.01482.x](https://doi.org/10.1111/j.1365-2699.2006.01482.x)
- Argyroudi A, Chatzinikolaou Y, Poirazidis K, Lazaridou M 2009 Do intermittent and ephemeral Mediterranean rivers belong to the same river type? *Aquat. Ecol.* 43, 465–476. doi: [10.1007/s10452-008-9176-9](https://doi.org/10.1007/s10452-008-9176-9)
- Assandri G, Bogliani G, Pedrini P, Brambilla M 2018 Beautiful agricultural landscapes promote cultural ecosystem services and biodiversity conservation. *Agric. Ecosyst. Environ.* 256, 200–210. doi: [10.1016/j.agee.2018.01.012](https://doi.org/10.1016/j.agee.2018.01.012)
- Asseng S, Martre P, Maiorano A, Rötter RP, O'Leary GJ et al. 2019 Climate change impact and adaptation for wheat protein. *Glob. Chang. Biol.* 25, 155–173. doi: [10.1111/gcb.14481](https://doi.org/10.1111/gcb.14481)
- Asseng S, Pannell DJ 2013 Adapting dryland agriculture to climate change: Farming implications and research and development needs in Western Australia. *Clim. Change* 118, 167–181. doi: [10.1007/s10584-012-0623-1](https://doi.org/10.1007/s10584-012-0623-1)
- Ausín B, Flores JA, Sierro FJ, Cacho I, Hernández-Almeida I et al. 2015 Atmospheric patterns driving Holocene productivity in the Alboran Sea (Western Mediterranean): A multiproxy approach. doi: <http://dx.doi.org/10.13039/501100004837>
- Avila IC, Kaschner K, Dormann CF 2018 Current global risks to marine mammals: Taking stock of the threats. *Biol. Conserv.* 221, 44–58. doi: [10.1016/j.biocon.2018.02.021](https://doi.org/10.1016/j.biocon.2018.02.021)
- Avramidis P, Geraga M, Lazarova M, Kontopoulos N 2013 Holocene record of environmental changes and palaeoclimatic implications in Alykes Lagoon, Zakynthos Island, western Greece, Mediterranean Sea. *Quat. Int.* 293, 184–195. doi: [10.1016/j.quaint.2012.04.026](https://doi.org/10.1016/j.quaint.2012.04.026)
- Ayyad MA, El-Ghareh REM 1982 Salt marsh vegetation of the Western Mediterranean desert of Egypt. *Vegetatio* 49, 3–19. doi: [10.1007/bf00051557](https://doi.org/10.1007/bf00051557)
- Azam CS, Gigot G, Witte I, Schatz B 2016 National and subnational Red Lists in European and Mediterranean countries: Current state and use for conservation. *Endanger. Species Res.* 30, 255–266. doi: [10.3354/esr00740](https://doi.org/10.3354/esr00740)
- Azzurro E, Castriota L, Falautano M, Bariche M, Broglio E et al. 2016 New records of the silver-cheeked toadfish *Lagocephalus sceleratus* (Gmelin, 1789) in the Tyrrhenian and Ionian Seas: early detection and participatory monitoring in practice. *Biol. Invasions Rec.* 5, 295–299. doi: [10.3391/bir.2016.5.4.16](https://doi.org/10.3391/bir.2016.5.4.16)
- Azzurro E, Castriota L, Falautano M, Giardina F, Andaloro F 2014 The silver-cheeked toadfish *Lagocephalus sceleratus* (Gmelin, 1789) reaches Italian waters. *J. Appl. Ichthyol.* 30, 1050–1052. doi: [10.1111/jai.12471](https://doi.org/10.1111/jai.12471)
- Azzurro E, Moschella P, Maynou F 2011 Tracking signals of change in Mediterranean fish diversity based on local ecological knowledge. *PLoS One* 6, e24885. doi: [10.1371/journal.pone.0024885](https://doi.org/10.1371/journal.pone.0024885)
- Bagella S, Salis L, Marrosu GM, Rossetti I, Fanni S et al. 2013 Effects of long-term management practices on grassland plant assemblages in Mediterranean cork oak silvo-pastoral systems. *Plant Ecol.* 214, 621–631. doi: [10.1007/s11258-013-0194-x](https://doi.org/10.1007/s11258-013-0194-x)
- Bagella S, Sitzia M, Roggero PP 2017 Soil fertilisation contributes to mitigating forest fire hazard associated with *Cistus monspeliensis* L. (rock rose) shrublands. *Int. J. Wildl. Fire* 26, 156. doi: [10.1071/WF16114](https://doi.org/10.1071/WF16114)
- Baldock D, Caraveli H, Dwyer J, Einschütz S, Petersen JE et al. 2000 The environmental impacts of irrigation in the European Union. Brussels, Belgium.
- Balkhausen O, Banse M, Grethe H 2007 Modelling CAP Decoupling in the EU: A Comparison of Selected Simulation Models and Results. *J. Agric. Econ.* 0, 071003055534004-???. doi: [10.1111/j.1477-9552.2007.00135.x](https://doi.org/10.1111/j.1477-9552.2007.00135.x)
- Ballesteros E 2006 *Mediterranean coralligenous assemblages: A synthesis of present knowledge*. Taylor & Francis <https://digital.csic.es/handle/10261/132026>
- Bally M, Garrabou J 2007 Thermodependent bacterial pathogens and mass mortalities in temperate benthic communities: a new case of emerging disease linked to climate change. *Glob. Chang. Biol.* 13, 2078–2088.
- Balzan M V., Pinheiro AM, Mascarenhas A, Morán-Ordóñez A, Ruiz-Frau A et al. 2019 Improving ecosystem assessments in Mediterranean social-ecological systems: a DPSIR analysis. *Ecosyst. People* 15, 136–155. doi: [10.1080/26395916.2019.1598499](https://doi.org/10.1080/26395916.2019.1598499)
- Balzan M V., Sadula R, Scalvenzi L 2020 Assessing ecosystem services supplied by agroecosystems in Mediterranean Europe: A literature review. *Land* 9, 245. doi: [10.3390/LAND9080245](https://doi.org/10.3390/LAND9080245)
- Balzan M V., Caruana J, Zammit A 2018 Assessing the capacity and flow of ecosystem services in multifunctional landscapes: Evidence of a rural-urban gradient in a Mediterranean small island state. *Land use policy* 75, 711–725. doi: [10.1016/j.landusepol.2017.08.025](https://doi.org/10.1016/j.landusepol.2017.08.025)
- Barange M, Bahri T, Beveridge MCM, Cochrane KL, Funge-Smith S et al. 2018 Impacts of Climate Change on

- Fisheries and Aquaculture: Synthesis of Current Knowledge. Adaptation and Mitigation Options. Rome.
- Barausse A, Michieli A, Riginella E, Palmeri L, Mazzoldi C 2011 Long-term changes in community composition and life-history traits in a highly exploited basin (northern Adriatic Sea): the role of environment and anthropogenic pressures. *J. Fish Biol.* 79, 1453–1486. doi: [10.1111/j.1095-8649.2011.03139.x](https://doi.org/10.1111/j.1095-8649.2011.03139.x)
- Barbaro L, Battisti A 2011 Birds as predators of the pine processionary moth (Lepidoptera: Notodontidae). *Biol. Control* 56, 107–114. doi: [10.1016/j.biocontrol.2010.10.009](https://doi.org/10.1016/j.biocontrol.2010.10.009)
- Barbet-Massin M, Thuiller W, Jiguet F 2012 The fate of European breeding birds under climate, land-use and dispersal scenarios. *Glob. Chang. Biol.* 18, 881–890. doi: [10.1111/j.1365-2486.2011.02552.x](https://doi.org/10.1111/j.1365-2486.2011.02552.x)
- Bariche M, Kleitou P, Kalogirou S, Bernardi G 2017 Genetics reveal the identity and origin of the lionfish invasion in the Mediterranean Sea. *Sci. Rep.* 7, 1–6. doi: [10.1038/s41598-017-07326-1](https://doi.org/10.1038/s41598-017-07326-1)
- Bariche M, Letourneur Y, Harmelin-Vivien M 2004 Temporal fluctuations and settlement patterns of native and Lessepsian herbivorous fishes on the Lebanese coast (eastern Mediterranean). *Environ. Biol. Fishes* 70, 81–90. doi: [10.1023/B:EBFI.0000022928.15148.75](https://doi.org/10.1023/B:EBFI.0000022928.15148.75)
- Bariche M, Torres M, Azzurro E 2013 The presence of the invasive lionfish *Pterois miles* in the Mediterranean Sea. *Mediterr. Mar. Sci.* 14, 292–294. doi: [10.12681/mms.428](https://doi.org/10.12681/mms.428)
- Baró F, Gómez-Baggethun E, Haase D 2017 Ecosystem service bundles along the urban-rural gradient: Insights for landscape planning and management. *Ecosyst. Serv.* 24, 147–159. doi: [10.1016/j.ecoser.2017.02.021](https://doi.org/10.1016/j.ecoser.2017.02.021)
- Barras C, Jorissen FJ, Labrune C, Andral B, Boissery P 2014 Live benthic foraminiferal faunas from the French Mediterranean Coast: Towards a new biotic index of environmental quality. *Ecol. Indic.* 36, 719–743. doi: [10.1016/j.ecolind.2013.09.028](https://doi.org/10.1016/j.ecolind.2013.09.028)
- Bastari A, Micheli F, Ferretti F, Pusceddu A, Cerrano C 2016 Large marine protected areas (LMPAs) in the Mediterranean Sea: The opportunity of the Adriatic Sea. *Mar. Policy* 68, 165–177. doi: [10.1016/j.marpol.2016.03.010](https://doi.org/10.1016/j.marpol.2016.03.010)
- Batalla RJ, Vericat D 2009 Hydrological and sediment transport dynamics of flushing flows: implications for management in large Mediterranean Rivers. *River Res. Appl.* 25, 297–314. doi: [10.1002/rra.1160](https://doi.org/10.1002/rra.1160)
- Batllori E, de Cáceres M, Brotons L, Ackerly DD, Moritz MA et al. 2019 Compound fire-drought regimes promote ecosystem transitions in Mediterranean ecosystems. *J. Ecol.* 107, 1187–1198. doi: [10.1111/1365-2745.13115](https://doi.org/10.1111/1365-2745.13115)
- Batllori E, Parisien M-A, Krawchuk MA, Moritz MA 2013 Climate change-induced shifts in fire for Mediterranean ecosystems. *Glob. Ecol. Biogeogr.* 22, 1118–1129. doi: [10.1111/geb.12065](https://doi.org/10.1111/geb.12065)
- Battisti A, Larsson S, Roques A 2017 Processionary Moths and Associated Urtication Risk: Global Change-Driven Effects. *Annu. Rev. Entomol.* 62, 323–342. doi: [10.1146/annurev-ento-031616-034918](https://doi.org/10.1146/annurev-ento-031616-034918)
- Battisti A, Stastny M, Netherer S, Robinet C, Schopf A et al. 2005 Expansion of geographic range in the pine processionary moth caused by increased winter temperatures. *Ecol. Appl.* 15, 2084–2096. doi: [10.1890/04-1903](https://doi.org/10.1890/04-1903)
- Baulch S, Perry C 2014 Evaluating the impacts of marine debris on cetaceans. *Mar. Pollut. Bull.* 80, 210–221. doi: [10.1016/j.marpolbul.2013.12.050](https://doi.org/10.1016/j.marpolbul.2013.12.050)
- Beaugrand G, Conversi A, Atkinson A, Cloern JE, Chiba S et al. 2019 Prediction of unprecedented biological shifts in the global ocean. *Nat. Clim. Chang.* 9, 237–243. doi: [10.1038/s41558-019-0420-1](https://doi.org/10.1038/s41558-019-0420-1)
- Becagli S, Lazzara L, Fani F, Marchese C, Traversi R et al. 2013 Relationship between methanesulfonate (MS<sup>-</sup>) in atmospheric particulate and remotely sensed phytoplankton activity in oligo-mesotrophic central Mediterranean Sea. *Atmos. Environ.* 79, 681–688. doi: [10.1016/j.atmosenv.2013.07.032](https://doi.org/10.1016/j.atmosenv.2013.07.032)
- Bêche LA, Connors PG, Resh VH, Merenlender AM 2009 Resilience of fishes and invertebrates to prolonged drought in two California streams. *Ecography (Cop.)* 32, 778–788. doi: [10.1111/j.1600-0587.2009.05612.x](https://doi.org/10.1111/j.1600-0587.2009.05612.x)
- Beger M, Grantham HS, Pressey RL, Wilson KA, Peterson EL et al. 2010 Conservation planning for connectivity across marine, freshwater, and terrestrial realms. *Biol. Conserv.* 143, 565–575. doi: [10.1016/j.biocon.2009.11.006](https://doi.org/10.1016/j.biocon.2009.11.006)
- Behrens P, Kieffe-de Jong JC, Bosker T, Rodrigues JFD, de Koning A et al. 2017 Evaluating the environmental impacts of dietary recommendations. *Proc. Natl. Acad. Sci. U. S. A.* 114, 13412–13417. doi: [10.1073/pnas.1711889114](https://doi.org/10.1073/pnas.1711889114)
- Belnap J 2002 Nitrogen fixation in biological soil crusts from southeast Utah, USA. *Biol. Fertil. Soils* 35, 128–135. doi: [10.1007/s00374-002-0452-x](https://doi.org/10.1007/s00374-002-0452-x)
- Belnap J 2006 The potential roles of biological soil crusts in dryland hydrologic cycles. *Hydrol. Process.* 20, 3159–3178. doi: [10.1002/hyp.6325](https://doi.org/10.1002/hyp.6325)
- Belnap J, Lange OL 2013 *Biological soil crusts: structure, function, and management*. Berlin Heidelberg: Springer-Verlag doi: [10.1007/978-3-642-56475-8](https://doi.org/10.1007/978-3-642-56475-8)
- Ben-David-Zaslow R, Benayahu Y 1999 Temporal variation in lipid, protein and carbohydrate content in the Red Sea soft coral *Heteroxenia fuscescens*. *J. Mar. Biol. Assoc. United Kingdom* 79, 1001–1006. doi: [10.1017/s002531549900123x](https://doi.org/10.1017/s002531549900123x)
- Ben Rais Lasram F, Guilhaumon F, Albouy C, Somot S, Thuiller W et al. 2010 The Mediterranean Sea as a “cul-de-sac” for endemic fishes facing climate change. *Glob. Chang. Biol.* 16, 3223–3245. <http://www.documentation.ird.fr/hor/fdi:010054015> [Accessed March 25, 2019].
- Ben Souissi J, Rifi M, Ghanem R, Ghazzi L, Boughedir W et al. 2014 *Lagocephalus sceleratus* (Gmelin, 1789) expands through the African coasts towards the Western

- Mediterranean Sea: A call for awareness. *Manag. Biol. Invasions* 5, 357–362. doi: [10.3391/mbi.2014.5.4.06](https://doi.org/10.3391/mbi.2014.5.4.06)
- Bencherif K, Boutekrabt A, Fontaine J, Laruelle F, Dalpé Y et al. 2015 Impact of soil salinity on arbuscular mycorrhizal fungi biodiversity and microflora biomass associated with *Tamarix articulata* Vahl rhizosphere in arid and semi-arid Algerian areas. *Sci. Total Environ.* 533, 488–494. doi: [10.1016/j.scitotenv.2015.07.007](https://doi.org/10.1016/j.scitotenv.2015.07.007)
- Benedetti-Cecchi L, Maggi E, Bertocci I, Vaselli S, Micheli F et al. 2003 Variation in rocky shore assemblages in the northwestern Mediterranean: contrasts between islands and the mainland. *J. Exp. Mar. Bio. Ecol.* 293, 193–215. doi: [10.1016/S0022-0981\(03\)00220-X](https://doi.org/10.1016/S0022-0981(03)00220-X)
- Benedetti-Cecchi L, Pannaciuoli F, Bulleri F, Moschella PS, Airoidi L et al. 2001 Predicting the consequences of anthropogenic disturbance: large-scale effects of loss of canopy algae on rocky shores. *Mar. Ecol. Prog. Ser.* 214, 137–150. doi: [10.3354/meps214137](https://doi.org/10.3354/meps214137)
- Benedetti F, Guilhaumon F, Adloff F, Ayata S 2018 Investigating uncertainties in zooplankton composition shifts under climate change scenarios in the Mediterranean Sea. *Ecography (Cop.)* 41, 345–360. doi: [10.1111/ecog.02434](https://doi.org/10.1111/ecog.02434)
- Benejam L, Aparicio E, Vargas MJ, Vila-Gispert A, García-Berthou E 2008 Assessing fish metrics and biotic indices in a Mediterranean stream: effects of uncertain native status of fish. *Hydrobiologia* 603, 197–210. doi: [10.1007/s10750-007-9272-1](https://doi.org/10.1007/s10750-007-9272-1)
- Benito-Garzón M, Alía R, Robson TM, Zavala MA 2011 Intra-specific variability and plasticity influence potential tree species distributions under climate change. *Glob. Ecol. Biogeogr.* 20, 766–778. doi: [10.1111/j.1466-8238.2010.00646.x](https://doi.org/10.1111/j.1466-8238.2010.00646.x)
- Benito-Garzón M, Leadley PW, Fernández-Manjarrés JF 2014 Assessing global biome exposure to climate change through the Holocene–Anthropocene transition. *Glob. Ecol. Biogeogr.* 23, 235–244. doi: [10.1111/geb.12097](https://doi.org/10.1111/geb.12097)
- Benito-Garzón M, Ruiz-Benito P, Zavala MA 2013 Interspecific differences in tree growth and mortality responses to environmental drivers determine potential species distributional limits in Iberian forests. *Glob. Ecol. Biogeogr.* 22, 1141–1151. doi: [10.1111/geb.12075](https://doi.org/10.1111/geb.12075)
- Benito G, Thorndyraft VR, Rico M, Sánchez-Moya Y, Sopena A 2008 Palaeoflood and floodplain records from Spain: Evidence for long-term climate variability and environmental changes. *Geomorphology* 101, 68–77. doi: [10.1016/j.geomorph.2008.05.020](https://doi.org/10.1016/j.geomorph.2008.05.020)
- Benito X, Trobajo R, Ibáñez C 2015 Benthic diatoms in a Mediterranean delta: ecological indicators and a conductivity transfer function for paleoenvironmental studies. *J. Paleolimnol.* 54, 171–188. doi: [10.1007/s10933-015-9845-3](https://doi.org/10.1007/s10933-015-9845-3)
- Benlhabib O, Yazar A, Qadir M, Lourenço E, Jacobsen S-E 2014 How Can We Improve Mediterranean Cropping Systems? *J. Agron. Crop Sci.* 200, 325–332. doi: [10.1111/jac.12066](https://doi.org/10.1111/jac.12066)
- Bensoussan N, Romano JC, Harmelin JG, Garrabou J 2010 High resolution characterization of northwest Mediterranean coastal waters thermal regimes: To better understand responses of benthic communities to climate change. *Estuar. Coast. Shelf Sci.* 87, 431–441. doi: [10.1016/j.ecss.2010.01.008](https://doi.org/10.1016/j.ecss.2010.01.008)
- Berline L, Siokou-Frangou I, Marasović I, Vidjak O, Fernández de Puelles ML et al. 2012 Intercomparison of six Mediterranean zooplankton time series. *Prog. Oceanogr.* 97–100, 76–91. doi: [10.1016/j.pocean.2011.11.011](https://doi.org/10.1016/j.pocean.2011.11.011)
- Berlow EL, D'Antonio CM, Reynolds SA 2002 Shrub expansion in montane meadows: The interaction of local-scale disturbance and site aridity. *Ecol. Appl.* 12, 1103–1118. doi: [10.1890/1051-0761\(2002\)012\[1103:SEIMMT\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2002)012[1103:SEIMMT]2.0.CO;2)
- Bernabucci U, Biffani S, Buggiotti L, Vitali A, Lacetera N et al. 2014 The effects of heat stress in Italian Holstein dairy cattle. *J. Dairy Sci.* 97, 471–486. doi: [10.3168/JDS.2013-6611](https://doi.org/10.3168/JDS.2013-6611)
- Bernabucci U, Colavecchia L, Danieli PP, Basiricò L, Lacetera N et al. 2011 Aflatoxin B1 and fumonisin B1 affect the oxidative status of bovine peripheral blood mononuclear cells. *Toxicol. Vitro.* 25, 684–691. doi: [10.1016/J.TIV.2011.01.009](https://doi.org/10.1016/J.TIV.2011.01.009)
- Bernal S, von Schiller D, Sabater F, Martí E 2013 Hydrological extremes modulate nutrient dynamics in Mediterranean climate streams across different spatial scales. *Hydrobiologia* 719, 31–42. doi: [10.1007/s10750-012-1246-2](https://doi.org/10.1007/s10750-012-1246-2)
- Bernes C 2005 Change Beneath the Surface: An In-Depth Look at Sweden's Marine Environment.
- Bernetti I, Chirici G, Sacchelli S 2019 Big data and evaluation of cultural ecosystem services: An analysis based on geotagged photographs from social media in tuscan forest (Italy). *iForest - Biogeosciences For.* 12, 98–105. doi: [10.3832/ifer2821-011](https://doi.org/10.3832/ifer2821-011)
- Bernstein A, Gustafson MT, Lewis R 2019 Disaster on the horizon: The price effect of sea level rise. *J. financ. econ.* 134, 253–272. doi: [10.1016/j.jfineco.2019.03.013](https://doi.org/10.1016/j.jfineco.2019.03.013)
- Bernués A, Tello-García E, Rodríguez-Ortega T, Ripoll-Bosch R, Casasús I 2016 Agricultural practices, ecosystem services and sustainability in High Nature Value farmland: Unraveling the perceptions of farmers and nonfarmers. *Land use policy* 59, 130–142. doi: [10.1016/J.LANDUSEPOL.2016.08.033](https://doi.org/10.1016/J.LANDUSEPOL.2016.08.033)
- Bertness MD, Ewanchuk PJ, Silliman BR 2002 Anthropogenic modification of New England salt marsh landscapes. *Proc. Natl. Acad. Sci. U. S. A.* 99, 1395–1398. doi: [10.1073/pnas.022447299](https://doi.org/10.1073/pnas.022447299)
- Bertolino M, Betti F, Bo M, Cattaneo-Vietti R, Pansini M et al. 2016 Changes and stability of a Mediterranean hard bottom benthic community over 25 years. *J. Mar. Biol. Assoc. United Kingdom* 96, 341–350. doi: [10.1017/S0025315415001186](https://doi.org/10.1017/S0025315415001186)
- Bertolino M, Cattaneo-Vietti R, Costa G, Pansini M, Fraschetti S et al. 2017a Have climate changes driven the diversity of a Mediterranean coralligenous sponge

- assemblage on a millennial timescale? *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 487, 355–363. doi: [10.1016/J.PALAEO.2017.09.020](https://doi.org/10.1016/J.PALAEO.2017.09.020)
- Bertolino M, Costa G, Carella M, Cattaneo-Vietti R, Cer-rano C et al. 2017b The dynamics of a Mediterranean coralligenous sponge assemblage at decennial and millennial temporal scales. *PLoS One* 12. doi: [10.1371/journal.pone.0177945](https://doi.org/10.1371/journal.pone.0177945)
- Bertolino M, Costa G, Cattaneo-Vietti R, Pansini M, Quarta G et al. 2019 Ancient and recent sponge assemblages from the Tyrrhenian coralligenous over millennia (Mediterranean Sea). *Facies* 65, 30. <https://doi.org/10.1007/s10347-019-0573-4>
- Berzak R, Scheinin A, Davidovitch N, Regev Y, Diga R et al. 2019 Prevalence of nervous necrosis virus (NNV) and *Streptococcus* species in wild marine fish and crustaceans from the Levantine Basin, Mediterranean Sea. *Dis. Aquat. Organ.* 133, 7–17. doi: [10.3354/dao03339](https://doi.org/10.3354/dao03339)
- Betancur RR, Hines A, Acero P. A, Ortí G, Wilbur AE et al. 2011 Reconstructing the lionfish invasion: insights into Greater Caribbean biogeography. *J. Biogeogr.* 38, 1281–1293. doi: [10.1111/j.1365-2699.2011.02496.x](https://doi.org/10.1111/j.1365-2699.2011.02496.x)
- Béthoux JP, Gentili B, Raunet J, Tailliez D 1990 Warming trend in the western Mediterranean deep water. *Nature* 347, 660. doi: [10.1038/347660a0](https://doi.org/10.1038/347660a0)
- Betti F, Bavestrello G, Bo M, Asnaghi V, Chiantore M et al. 2017 Over 10 years of variation in Mediterranean reef benthic communities. *Mar. Ecol.* 38, e12439. doi: [10.1111/maec.12439](https://doi.org/10.1111/maec.12439)
- Betzold C 2015 Adapting to climate change in small island developing states. *Clim. Change* 133, 481–489. doi: [10.1007/s10584-015-1408-0](https://doi.org/10.1007/s10584-015-1408-0)
- Bianchi CN, Caroli F, Guidetti P, Morri C 2018 Seawater warming at the northern reach for southern species: Gulf of Genoa, NW Mediterranean. *J. Mar. Biol. Assoc. United Kingdom* 98, 1–12. doi: [10.1017/S0025315417000819](https://doi.org/10.1017/S0025315417000819)
- Bianchi CN, Morri C 2000 Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research. *Mar. Pollut. Bull.* 40, 367–376. doi: [10.1016/S0025-326X\(00\)00027-8](https://doi.org/10.1016/S0025-326X(00)00027-8)
- Bianchi CN, Morri C 2003 Global sea warming and “tropicalization” of the Mediterranean Sea: biogeographic and ecological aspects. *Biogeogr. – J. Integr. Biogeogr.* 24, 319–328. doi: [10.21426/b6110129](https://doi.org/10.21426/b6110129)
- Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M et al. 2006 Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain and the Netherlands. *Science (80-. )*. 313, 351–354. doi: [10.1126/science.1127863](https://doi.org/10.1126/science.1127863)
- Bindi M, Olesen JE 2011 The responses of agriculture in Europe to climate change. *Reg. Environ. Chang.* 11, 151–158. doi: [10.1007/s10113-010-0173-x](https://doi.org/10.1007/s10113-010-0173-x)
- Bisbis MB, Gruda N, Blanke M 2018 Potential impacts of climate change on vegetable production and product quality – A review. *J. Clean. Prod.* 170, 1602–1620. doi: [10.1016/J.JCLEPRO.2017.09.224](https://doi.org/10.1016/J.JCLEPRO.2017.09.224)
- Bitar G, Harmelin JG, Verlaque M, Zibrowius H 2000 Sur la flore marine benthique supposée Lessepsienne de la côte libanaise. Cas particulier de *Stytopodium schimperii*. in *Proceedings of the First Mediterranean Symposium on Marine Vegetation, Ajaccio, 3-4 Oct. 2000*, ed. RAC/SPA (RAC/SPA, PNUE), 97–100.
- Bitar G, Ramos-Esplá A, Ocaña O, Sghaier YR, Forcada A et al. 2017 Introduced marine macroflora of Lebanon and its distribution on the Levantine coast. *Mediterr. Mar. Sci.* 18, 138–155. doi: [10.12681/mms.1993](https://doi.org/10.12681/mms.1993)
- Blinda M, Thivet G 2009 Ressources et demandes en eau en Méditerranée : situation et perspectives. *Sci. Chang. planétaires / Sécheresse* 20, 9–16. doi: [10.1684/SEC.2009.0162](https://doi.org/10.1684/SEC.2009.0162)
- Blois JL, Zarnetske PL, Fitzpatrick MC, Finnegan S 2013 Climate Change and the Past, Present, and Future of Biotic Interactions. *Science (80-. )*. 341, 499–504. doi: [10.1126/science.1237184](https://doi.org/10.1126/science.1237184)
- Blondel J 2006 The ‘design’ of Mediterranean landscapes: a millennial story of humans and ecological systems during the historic period. *Hum. Ecol.* 34, 713–729. doi: [10.1007/s10745-006-9030-4](https://doi.org/10.1007/s10745-006-9030-4)
- Blondel J, Aronson J 1999 *Biology and wildlife of the Mediterranean region*. USA: Oxford University Press. [http://llrc.mcast.edu.mt/digitalversion/Table\\_of\\_Contents\\_13104.pdf](http://llrc.mcast.edu.mt/digitalversion/Table_of_Contents_13104.pdf)
- Blondel J, Aronson J, Bodiou JY, Boeuf G 2010 *The Mediterranean region: biological diversity in space and time*. Oxford University Press.
- Bobbink R, Hicks K, Galloway J, Spranger T, Alkemade R et al. 2010 Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecol. Appl.* 20, 30–59. doi: [10.1890/08-1140.1](https://doi.org/10.1890/08-1140.1)
- Boddy L, Büntgen U, Egli S, Gange AC, Heegaard E et al. 2014 Climate variation effects on fungal fruiting. *Fungal Ecol.* 10, 20–33. doi: [10.1016/j.funeco.2013.10.006](https://doi.org/10.1016/j.funeco.2013.10.006)
- Bodin J, Badeau V, Bruno E, Cluzeau C, Moisselin JM et al. 2013 Shifts of forest species along an elevational gradient in Southeast France: Climate change or stand maturation? *J. Veg. Sci.* 24, 269–283. doi: [10.1111/j.1654-1103.2012.01456.x](https://doi.org/10.1111/j.1654-1103.2012.01456.x)
- Bonada N, Dolédec S, Stutzner B 2007 Taxonomic and biological trait differences of stream macroinvertebrate communities between mediterranean and temperate regions: implications for future climatic scenarios. *Glob. Chang. Biol.* 13, 1658–1671. doi: [10.1111/j.1365-2486.2007.01375.x](https://doi.org/10.1111/j.1365-2486.2007.01375.x)
- Bonada N, Resh VH 2013 Mediterranean-climate streams and rivers: geographically separated but ecologically comparable freshwater systems. *Hydrobiologia* 719, 1–29. doi: [10.1007/s10750-013-1634-2](https://doi.org/10.1007/s10750-013-1634-2)
- Bond NR, Lake PS, Arthington AH 2008 The impacts of drought on freshwater ecosystems: an Australian perspective. *Hydrobiologia* 600, 3–16. doi: [10.1007/s10750-008-9326-z](https://doi.org/10.1007/s10750-008-9326-z)
- Boonne C, Wacquant JP, Jonard R 1992 In-vitro cloning of *Dittrichia viscosa* for screening nutritional ecotypes.

- Plant Soil* 142, 323–328. doi: [10.1007/BF00010978](https://doi.org/10.1007/BF00010978)
- Borja Á, Elliott M, Andersen JH, Berg T, Carstensen J et al. 2016 Overview of Integrative Assessment of Marine Systems: The Ecosystem Approach in Practice. *Front. Mar. Sci.* 3. doi: [10.3389/fmars.2016.00020](https://doi.org/10.3389/fmars.2016.00020)
- Bosc E, Bricaud A, Antoine D 2004 Seasonal and interannual variability in algal biomass and primary production in the Mediterranean Sea, as derived from 4 years of SeaWiFS observations. *Global Biogeochem. Cycles* 18. doi: [10.1029/2003GB002034](https://doi.org/10.1029/2003GB002034)
- Bosco A, Rinaldi L, Musella V, Amadesi A, Cringoli G 2015 Outbreak of acute fasciolosis in sheep farms in a Mediterranean area arising as a possible consequence of climate change. *Geospat. Health* 9, 319. doi: [10.4081/gh.2015.354](https://doi.org/10.4081/gh.2015.354)
- Bossart GD 2011 Marine mammals as sentinel species for oceans and human health. *Vet. Pathol.* 48, 676–690. doi: [10.1177/0300985810388525](https://doi.org/10.1177/0300985810388525)
- Boudouresque C-F 2004 Marine biodiversity in the Mediterranean: status of species, populations and communities, in *Scientific Reports of Port-Cros National Park 20*, 97–146.
- Boudouresque C-F, Blanfuné A, Harmelin-Vivien M, Personnic S, Ruitton S et al. 2015 Where seaweed forests meet animal forests: the examples of macroalgae in coral reefs and the Mediterranean coralligenous ecosystem, in *Marine Animal Forests: The Ecology of Benthic Biodiversity Hotspots*, eds. Rossi S, Bramanti L, Gori A, Orejas Saco del Valle C (Springer International Publishing, Cham), 1–28. doi: [10.1007/978-3-319-17001-5\\_48-1](https://doi.org/10.1007/978-3-319-17001-5_48-1)
- Boulton AJ 2003 Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages. *Freshw. Biol.* 48, 1173–1185. doi: [10.1046/j.1365-2427.2003.01084.x](https://doi.org/10.1046/j.1365-2427.2003.01084.x)
- Boustany L, El Indary S, Nader M 2015 Biological characteristics of the Lessepsian pufferfish *Lagocephalus sceleratus* (Gmelin, 1789) off Lebanon. *Cah. Biol. Mar.* 56, 137–142.
- Bowman DMJS, Williamson GJ, Abatzoglou JT, Kolden CA, Cochrane MA et al. 2017 Human exposure and sensitivity to globally extreme wildfire events. *Nat. Ecol. Evol.* 1, 1–6. doi: [10.1038/s41559-016-0058](https://doi.org/10.1038/s41559-016-0058)
- Bramanti L, Magagnini G, de Maio L, Santangelo G 2005 Recruitment, early survival and growth of the Mediterranean red coral *Corallium rubrum* (L. 1758), a 4-year study. *J. Exp. Mar. Bio. Ecol.* 314, 69–78. doi: [10.1016/j.jembe.2004.08.029](https://doi.org/10.1016/j.jembe.2004.08.029)
- Bramanti L, Movilla J, Calvo EM, Gori A, Domínguez-Carrión C et al. 2015 Effects of ocean acidification on the precious Mediterranean red coral (*Corallium rubrum*). in <https://pdfs.semanticscholar.org/777c/81157a7f-771134ba6a1c7c0d77325a8d1683.pdf>
- Bramanti L, Movilla J, Guron M, Calvo EM, Gori A et al. 2013 Detrimental effects of ocean acidification on the economically important Mediterranean red coral (*Corallium rubrum*). *Glob. Chang. Biol.* 19, 1897–1908. doi: [10.1111/gcb.12171](https://doi.org/10.1111/gcb.12171)
- Bramanti L, Vielmini I, Rossi S, Tsounis G, Iannelli M et al. 2014 Demographic parameters of two populations of red coral (*Corallium rubrum* L. 1758) in the North Western Mediterranean. *Mar. Biol.* 161, 1015–1026. doi: [10.1007/s00227-013-2383-5](https://doi.org/10.1007/s00227-013-2383-5)
- Brambilla M, Ilahiane L, Assandri G, Ronchi S, Bogliani G 2017 Combining habitat requirements of endemic bird species and other ecosystem services may synergistically enhance conservation efforts. *Sci. Total Environ.* 586, 206–214. doi: [10.1016/j.scitotenv.2017.01.203](https://doi.org/10.1016/j.scitotenv.2017.01.203)
- Brauman KA, Daily GC, Duarte TK, Mooney HA 2007 The Nature and Value of Ecosystem Services: An Overview Highlighting Hydrologic Services. *Annu. Rev. Environ. Resour.* 32, 67–98. doi: [10.1146/annurev.energy.32.031306.102758](https://doi.org/10.1146/annurev.energy.32.031306.102758)
- Bravo-Oviedo A, Gallardo-Andrés C, del Río M, Montero G 2010 Regional changes of *Pinus pinaster* site index in Spain using a climate-based dominant height model. *Can. J. For. Res.* 40, 2036–2048. doi: [10.1139/x10-143](https://doi.org/10.1139/x10-143)
- Bravo F, del Río M, Bravo-Oviedo A, Ruiz-Peinado R, del Peso C et al. 2017 Forest Carbon Sequestration: The Impact of Forest Management, in *Managing forest ecosystems: the challenge of climate change*, eds. Bravo F, Le May V, Jandl R (Springer International Publishing), 251–275. doi: [10.1007/978-3-319-28250-3](https://doi.org/10.1007/978-3-319-28250-3)
- Bray L, Reizopoulou S, Voukouvalas E, Soukissian T, Alovermar C et al. 2016 Expected Effects of Offshore Wind Farms on Mediterranean Marine Life. *J. Mar. Sci. Eng.* 4, 18. doi: [10.3390/jmse4010018](https://doi.org/10.3390/jmse4010018)
- Breeze TD, Vaissière BE, Bommarco R, Petanidou T, Seraphides N et al. 2014 Agricultural Policies Exacerbate Honeybee Pollination Service Supply-Demand Mismatches Across Europe. *PLoS One* 9, e82996. doi: [10.1371/journal.pone.0082996](https://doi.org/10.1371/journal.pone.0082996)
- Brierley GJ, Fryirs KA 2005 *Geomorphology and River Management: Applications of the River Styles Framework*. Oxford, UK: Blackwell Publishing. doi: [10.1002/9780470751367](https://doi.org/10.1002/9780470751367)
- Brinson MM, Malvárez AI 2002 Temperate freshwater wetlands: Types, status, and threats. *Environ. Conserv.* 29, 115–133. doi: [10.1017/S0376892902000085](https://doi.org/10.1017/S0376892902000085)
- Bruce S, Boix D, López-Flores R, Badosa A, Moreno-Amich R et al. 2005 Zooplankton structure and dynamics in permanent and temporary Mediterranean salt marshes: taxon-based and size-based approaches. *Arch. für Hydrobiol.* 162, 535–555. doi: [10.1127/0003-9136/2005/0162-0535](https://doi.org/10.1127/0003-9136/2005/0162-0535)
- Bruno D, Belmar O, Sánchez-Fernández D, Guareschi S, Millán A et al. 2014a Responses of Mediterranean aquatic and riparian communities to human pressures at different spatial scales. *Ecol. Indic.* 45, 456–464. doi: [10.1016/j.ecolind.2014.04.051](https://doi.org/10.1016/j.ecolind.2014.04.051)
- Bruno D, Belmar O, Sánchez-Fernández D, Velasco J 2014b Environmental determinants of woody and herbaceous riparian vegetation patterns in a semi-arid mediterranean basin. *Hydrobiologia* 730, 45–57.



- doi: [10.1007/s10750-014-1822-8](https://doi.org/10.1007/s10750-014-1822-8)
- Bruno D, Gutiérrez-Cánovas C, Sánchez-Fernández D, Velasco J, Nilsson C 2016 Impacts of environmental filters on functional redundancy in riparian vegetation. *J. Appl. Ecol.* 53, 846–855. doi: [10.1111/1365-2664.12619](https://doi.org/10.1111/1365-2664.12619)
- Bruno D, Zapata V, Guareschi S, Picazo F, Dettori E et al. 2019 Short-Term Responses of Aquatic and Terrestrial Biodiversity to Riparian Restoration Measures Designed to Control the Invasive *Arundo donax* L. *Water* 11, 2551. doi: [10.3390/w11122551](https://doi.org/10.3390/w11122551)
- Buecher E, Goy J, Planque B, Etienne M, Dallot S 1997 Long-term fluctuations of *Liriope tetraphylla* in Villefranche bay between 1966 and 1993 compared to *Pelagia noctiluca* populations. *Ocean. Acta* 20, 145–157. <https://archimer.ifremer.fr/doc/00093/20387/> [Accessed March 25, 2019]
- Bugalho MN, Caldeira MC, Pereira JS, Aronson J, Pausas JG 2011 Mediterranean cork oak savannas require human use to sustain biodiversity and ecosystem services. *Front. Ecol. Environ.* 9, 278–286. doi: [10.1890/100084](https://doi.org/10.1890/100084)
- Bullitta S, Piluzza G, Re GA, Sulas L 2017 Mediterranean grassland species from traditional to multiple uses, in *Grassland science in Europe, vol. 22 - Grassland resources for extensive farming systems in marginal lands.*, 461–474.
- Büntgen U, Egli S, Galván JD, Diez JM, Aldea J et al. 2015 Drought-induced changes in the phenology, productivity and diversity of Spanish fungi. *Fungal Ecol.* 16, 6–18. doi: [10.1016/j.funeco.2015.03.008](https://doi.org/10.1016/j.funeco.2015.03.008)
- Büntgen U, Frank D, Liebhold A, Johnson D, Carrer M et al. 2009 Three centuries of insect outbreaks across the European Alps. *New Phytol.* 182, 929–941. doi: [10.1111/j.1469-8137.2009.02825.x](https://doi.org/10.1111/j.1469-8137.2009.02825.x)
- Büntgen U, Frank DC, Kaczka RJ, Verstege A, Zwi-jacz-Kozica T et al. 2007 Growth responses to climate in a multi-species tree-ring network in the Western Carpathian Tatra Mountains, Poland and Slovakia. *Tree Physiol.* 27, 689–702. doi: [10.1093/treephys/27.5.689](https://doi.org/10.1093/treephys/27.5.689)
- Büntgen U, Krusic PJ, Verstege A, Sangüesa-Barreda G, Wagner S et al. 2017 New Tree-Ring Evidence from the Pyrenees Reveals Western Mediterranean Climate Variability since Medieval Times. *J. Clim.* 30, 5295–5318. doi: [10.1175/jcli-d-16-0526.1](https://doi.org/10.1175/jcli-d-16-0526.1)
- Burak S, Dogan E, Gazioglu C 2004 Impact of urbanization and tourism on coastal environment. *Ocean Coast. Manag.* 47, 515–527. doi: [10.1016/j.ocecoaman.2004.07.007](https://doi.org/10.1016/j.ocecoaman.2004.07.007)
- Burak S, Margat J 2016 Water Management in the Mediterranean Region: Concepts and Policies. *Water Resour. Manag.* 30, 5779–5797. doi: [10.1007/s11269-016-1389-4](https://doi.org/10.1007/s11269-016-1389-4)
- Burch GJ, Bath RK, Moore ID, O'Loughlin EM 1987 Comparative hydrological behaviour of forested and cleared catchments in southeastern Australia. *J. Hydrol.* 90, 19–42. doi: [10.1016/0022-1694\(87\)90171-5](https://doi.org/10.1016/0022-1694(87)90171-5)
- Bussi G, Francés F, Horel E, López-Tarazón JA, Batalla RJ 2014 Modelling the impact of climate change on sediment yield in a highly erodible Mediterranean catchment. *J. Soils Sediments* 14, 1921–1937. doi: [10.1007/s11368-014-0956-7](https://doi.org/10.1007/s11368-014-0956-7)
- Bussotti F, Ferrini F, Pollastrini M, Fini A 2014 The challenge of Mediterranean sclerophyllous vegetation under climate change: From acclimation to adaptation. *Environ. Exp. Bot.* 103, 80–98. doi: [10.1016/j.envexpbot.2013.09.013](https://doi.org/10.1016/j.envexpbot.2013.09.013)
- Bütöf A, von Riedmatten LR, Dormann CF, Scherer-Lorenzen M, Welk E et al. 2012 The responses of grassland plants to experimentally simulated climate change depend on land use and region. *Glob. Chang. Biol.* 18, 127–137. doi: [10.1111/j.1365-2486.2011.02539.x](https://doi.org/10.1111/j.1365-2486.2011.02539.x)
- Caballero R, Fernández-González F, Pérez Badia R, Molle G, Roggero PP et al. 2011 Grazing systems and biodiversity in Mediterranean areas: Spain, Italy and Greece. *Pastos* 39, 9–152. <http://polired.upm.es/index.php/pastos/article/view/1712>
- Caffarra A, Rinaldi M, Eccel E, Rossi V, Pertot I 2012 Modelling the impact of climate change on the interaction between grapevine and its pests and pathogens: European grapevine moth and powdery mildew. *Agric. Ecosyst. Environ.* 148, 89–101. doi: [10.1016/j.AGEE.2011.11.017](https://doi.org/10.1016/j.AGEE.2011.11.017)
- Calbó J 2010 Possible Climate Change Scenarios with Specific Reference to Mediterranean Regions, in *Water Scarcity in the Mediterranean. Perspectives Under Global Change*, eds. Sabater S, Barceló D (Berlin Heidelberg: Springer-Verlag), 1–13. doi: [10.1007/978-3-642-03971-3](https://doi.org/10.1007/978-3-642-03971-3)
- Calvo EM, Simó R, Coma R, Ribes M, Pascual J et al. 2011 Effects of climate change on Mediterranean marine ecosystems: the case of the Catalan Sea. *Clim. Res.* 50, 1–29. doi: [10.3354/cr01040](https://doi.org/10.3354/cr01040)
- Camedda A, Marra S, Matiddi M, Massaro G, Coppa S et al. 2014 Interaction between loggerhead sea turtles (*Caretta caretta*) and marine litter in Sardinia (Western Mediterranean Sea). *Mar. Environ. Res.* 100, 25–32. doi: [10.1016/j.marenvres.2013.12.004](https://doi.org/10.1016/j.marenvres.2013.12.004)
- Cammarano D, Ceccarelli S, Grando S, Romagosa I, Benbelkacem A et al. 2019 The impact of climate change on barley yield in the Mediterranean basin. *Eur. J. Agron.* 106, 1–11. doi: [10.1016/j.eja.2019.03.002](https://doi.org/10.1016/j.eja.2019.03.002)
- Camp J, Flo E, Vila M, Arin L, Reñé A et al. 2016 Pros and cons of biological quality element phytoplankton as a water-quality indicator in the NW Mediterranean Sea, in *Experiences from Ground, Coastal and Transitional Water Quality Monitoring. The Handbook of Environmental Chemistry, vol 43.*, eds. Munné A, Ginebreda A, Prat N (Cham: Springer International Publishing), 135–160. doi: [10.1007/698\\_2015\\_392](https://doi.org/10.1007/698_2015_392)
- Campana I, Angeletti D, Crosti R, Luperini C, Ruvoletto A et al. 2017 Seasonal characterisation of maritime traffic and the relationship with cetacean presence in the

- Western Mediterranean Sea. *Mar. Pollut. Bull.* 115, 282–291. doi: [10.1016/j.marpolbul.2016.12.008](https://doi.org/10.1016/j.marpolbul.2016.12.008)
- Campani T, Bains M, Giannetti M, Cancelli F, Mancusi C et al. 2013 Presence of plastic debris in loggerhead turtle stranded along the Tuscany coasts of the Pelagos Sanctuary for Mediterranean Marine Mammals (Italy). *Mar. Pollut. Bull.* 74, 225–230. doi: [10.1016/j.marpolbul.2013.06.053](https://doi.org/10.1016/j.marpolbul.2013.06.053)
- Canals M, Puig P, de Madron XD, Heussner S, Palanques A et al. 2006 Flushing submarine canyons. *Nature* 444, 354–357. doi: [10.1038/nature05271](https://doi.org/10.1038/nature05271)
- Canepa A, Fuentes V, Bosch-Belmar M, Acevedo M, Toledo-Guedes K et al. 2017 Environmental factors influencing the spatio-temporal distribution of *Carybdea marsupialis* (Lineo, 1978, Cubozoa) in South-Western Mediterranean coasts. *PLoS One* 12, e0181611. doi: [10.1371/journal.pone.0181611](https://doi.org/10.1371/journal.pone.0181611)
- Canepa A, Fuentes V, Sabatés A, Piraino F, Boero F et al. 2014 *Pelagia noctiluca* in the Mediterranean Sea, in *Jellyfish Blooms*, eds. Pitt KA, Lucas CH, 237–266.
- Cánovas-Molina A, Montefalcone M, Bavestrello G, Cau A, Bianchi CN et al. 2016 A new ecological index for the status of mesophotic megabenthic assemblages in the mediterranean based on ROV photography and video footage. *Cont. Shelf Res.* 121, 13–20. doi: [10.1016/j.csr.2016.01.008](https://doi.org/10.1016/j.csr.2016.01.008)
- Caon L, Vallejo VR, Coen RJ, Geissen V 2014 Effects of wildfire on soil nutrients in Mediterranean ecosystems. *Earth-Science Rev.* 139, 47–58. doi: [10.1016/j.earscirev.2014.09.001](https://doi.org/10.1016/j.earscirev.2014.09.001)
- Cape JN, van der Eerden LJ, Sheppard LJ, Leith ID, Sutton MA 2009 Evidence for changing the critical level for ammonia. *Environ. Pollut.* 157, 1033–1037. doi: [10.1016/j.envpol.2008.09.049](https://doi.org/10.1016/j.envpol.2008.09.049)
- Capiomont A, Breugnot E, den Haan M, Meinesz A 2005 Phenology of a deep-water population of *Caulerpa racemosa* var. *cylindracea* in the northwestern Mediterranean Sea. *Bot. Mar.* 48, 80–83. doi: [10.1515/bot.2005.006](https://doi.org/10.1515/bot.2005.006)
- Capone R, El Bilali H, Cardone G, Debs P, Driouech N 2014 Food economic accessibility and affordability in the Mediterranean region: an exploratory assessment at micro and macro levels. *J. Food Secur.* 2, 1–12. doi: [10.12691/jfs-2-1-1](https://doi.org/10.12691/jfs-2-1-1)
- Cardinale M, Osio GC, Scarcella G 2017 Mediterranean Sea: a failure of the European fisheries management system, in *Frontiers in Marine Science* (Frontiers Media SA). doi: [10.3389/fmars.2017.00072](https://doi.org/10.3389/fmars.2017.00072)
- Cardoch L, Day JW, Ibáñez C 2002 Net primary productivity as an indicator of sustainability in the Ebro and Mississippi deltas. *Ecol. Appl.* 12, 1044–1055. doi: [10.1890/1051-0761\(2002\)012\[1044:NPPAAI\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2002)012[1044:NPPAAI]2.0.CO;2)
- Carmona P, Ruiz-Pérez J-M, Blázquez A-M, López-Belzunce M, Riera S et al. 2016 Environmental evolution and mid-late Holocene climate events in the Valencia lagoon (Mediterranean coast of Spain). *The Holocene* 26, 1750–1765. doi: [10.1177/0959683616645940](https://doi.org/10.1177/0959683616645940)
- Carreiras M, Ferreira AJD, Valente S, Fleskens L, Gonzales-Pelayo Ó et al. 2014 Comparative analysis of policies to deal with wildfire risk. *L. Degrad. Dev.* 25, 92–103. doi: [10.1002/ldr.2271](https://doi.org/10.1002/ldr.2271)
- Casas-Güell E, Cebrián E, Garrabou J, Ledoux J-B, Linares C et al. 2016 Structure and biodiversity of coralligenous assemblages dominated by the precious red coral *Corallium rubrum* over broad spatial scales. *Sci. Rep.* 6, 36535. doi: [10.1038/srep36535](https://doi.org/10.1038/srep36535)
- Casas-Güell E, Teixidó N, Garrabou J, Cebrián E 2015 Structure and biodiversity of coralligenous assemblages over broad spatial and temporal scales. *Mar. Biol.* 162, 901–912. doi: [10.1007/s00227-015-2635-7](https://doi.org/10.1007/s00227-015-2635-7)
- Cassman KG 1999 Ecological intensification of cereal production systems: yield potential, soil quality, and precision agriculture. *Proc. Natl. Acad. Sci. U. S. A.* 96, 5952–9. doi: [10.1073/PNAS.96.11.5952](https://doi.org/10.1073/PNAS.96.11.5952)
- Castañeda IS, Schefuß E, Pätzold J, Sinninghe Damsté JS, Weldeab S et al. 2010 Millennial-scale sea surface temperature changes in the eastern Mediterranean (Nile River Delta region) over the last 27,000 years. *Paleoceanography* 25, PA1208. doi: [10.1029/2009PA001740](https://doi.org/10.1029/2009PA001740)
- Castillo-Monroy AP, Maestre FT, Delgado-Baquerizo M, Gallardo A 2010 Biological soil crusts modulate nitrogen availability in semi-arid ecosystems: insights from a Mediterranean grassland. *Plant Soil* 333, 21–34. doi: [10.1007/s11104-009-0276-7](https://doi.org/10.1007/s11104-009-0276-7)
- Castro H, Castro P 2019 Mediterranean Marginal Lands in Face of Climate Change: Biodiversity and Ecosystem Services, in (Springer, Cham), 175–187. doi: [10.1007/978-3-319-75004-0\\_10](https://doi.org/10.1007/978-3-319-75004-0_10)
- Castro J, Zamora R, Hódar JA, Gómez JM, Gómez-Aparicio L 2004 Benefits of using shrubs as nurse plants for reforestation in Mediterranean mountains: A 4-year study. *Restor. Ecol.* 12, 352–358. doi: [10.1111/j.1061-2971.2004.0316.x](https://doi.org/10.1111/j.1061-2971.2004.0316.x)
- Ceballos-Barbancho A, Morán-Tejeda E, Luengo-Ugidos MÁ, Llorente-Pinto JM 2008 Water resources and environmental change in a Mediterranean environment: The south-west sector of the Duero river basin (Spain). *J. Hydrol.* 351, 126–138. doi: [10.1016/j.jhydrol.2007.12.004](https://doi.org/10.1016/j.jhydrol.2007.12.004)
- Cebrián E, Linares C, Marschal C, Garrabou J 2012 Exploring the effects of invasive algae on the persistence of gorgonian populations. *Biol. Invasions* 14, 2647–2656. doi: [10.1007/s10530-012-0261-6](https://doi.org/10.1007/s10530-012-0261-6)
- Ceccarelli T, Bajocco S, Luigi Perini L, Luca Salvati L 2014 Urbanisation and Land Take of High Quality Agricultural Soils - Exploring Long-term Land Use Changes and Land Capability in Northern Italy. *Int. J. Environ. Res.* 8, 181–192. doi: [10.22059/IJER.2014.707](https://doi.org/10.22059/IJER.2014.707)
- Ceccherelli G, Casu D, Sechi N 2005 Spatial variation of intertidal assemblages at Tavolara-Capo Coda Cavallo MPA (NE Sardinia): geographical vs. protection effect. *Mar. Environ. Res.* 59, 533–546. doi: [10.1016/j.marenvres.2004.09.002](https://doi.org/10.1016/j.marenvres.2004.09.002)

- Cecchini M, Zamboni I, Pontrandolfi A, Turco R, Colantoni A et al. 2019 Urban sprawl and the 'olive' landscape: sustainable land management for 'crisis' cities. *Geo-Journal* 84, 237–255. doi: [10.1007/s10708-018-9848-5](https://doi.org/10.1007/s10708-018-9848-5)
- Cerdà A, Mataix-Solera J 2009 *Efectos de los incendios forestales sobre los suelos en España. El estado de la cuestión visto por los científicos españoles.*, eds. Cerdà A, Mataix-Solera J Valencia: Cátedra de Divulgación de la Ciencia. Universitat de Valencia.
- Cerrano C, Bavestrello G, Bianchi CN, Cattaneo-Vietti R, Bava S et al. 2000 A catastrophic mass-mortality episode of gorgonians and other organisms in the Ligurian Sea (North-western Mediterranean), summer 1999. *Ecol. Lett.* 3, 284–293. doi: [10.1046/j.1461-0248.2000.00152.x](https://doi.org/10.1046/j.1461-0248.2000.00152.x)
- Cerrano C, Cardini U, Bianchelli S, Corinaldesi C, Puscedu A et al. 2013 Red coral extinction risk enhanced by ocean acidification. *Sci. Rep.* 3, 1457. doi: [10.1038/srep01457](https://doi.org/10.1038/srep01457)
- Cerrano C, Totti C, Sponga F, Bavestrello G 2006 Summer disease in *Parazoanthus axinellae* (Schmidt, 1862) (Cnidaria, Zoanthidea). *Ital. J. Zool.* 73, 355–361. doi: [10.1080/11250000600911675](https://doi.org/10.1080/11250000600911675)
- Charlton MB, Arnell NW 2011 Adapting to climate change impacts on water resources in England—An assessment of draft Water Resources Management Plans. *Glob. Environ. Chang.* 21, 238–248. doi: [10.1016/j.gloenvcha.2010.07.012](https://doi.org/10.1016/j.gloenvcha.2010.07.012)
- Charlton R 2008 *Fundamentals of fluvial geomorphology*. Abingdon: Routledge.
- Chartzoulakis KS, Bertaki M 2015 Sustainable Water Management in Agriculture under Climate Change. *Agric. Agric. Sci. Procedia* 4, 88–98. doi: [10.1016/j.aaspro.2015.03.011](https://doi.org/10.1016/j.aaspro.2015.03.011)
- Chauvelon P, Tournoud MG, Sandoz A 2003 Integrated hydrological modelling of a managed coastal Mediterranean wetland (Rhône delta, France): initial calibration. *Hydrol. Earth Syst. Sci. Discuss. Eur. Geosci. Union* 7, 123–132. <https://hal.archives-ouvertes.fr/hal-00330847> [Accessed October 13, 2020]
- Chefaoui RM, Chozas S 2019 Abandonment of traditional saltworks facilitates degradation of halophytic plant communities and *Carpobrotus edulis* invasion. *Appl. Veg. Sci.* 22, 444–453. doi: [10.1111/avsc.12436](https://doi.org/10.1111/avsc.12436)
- Chefaoui RM, Duarte CM, Serrão EA 2018 Dramatic loss of seagrass habitat under projected climate change in the Mediterranean Sea. *Glob. Chang. Biol.* 24, 4919–4928. doi: [10.1111/gcb.14401](https://doi.org/10.1111/gcb.14401)
- Chemello R, Silenzi S 2011 Vermetid reefs in the Mediterranean Sea as archives of sea-level and surface temperature changes. *Chem. Ecol.* 27, 121–127. doi: [10.1080/02757540.2011.554405](https://doi.org/10.1080/02757540.2011.554405)
- Cheung WWL, Lam VWY, Sarmiento JL, Kearney K, Watson R et al. 2010 Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change. *Glob. Chang. Biol.* 16, 24–35. doi: [10.1111/j.1365-2486.2009.01995.x](https://doi.org/10.1111/j.1365-2486.2009.01995.x)
- Cheung WWL, Watson R, Pauly D 2013 Signature of ocean warming in global fisheries catch. *Nature* 497, 365–368. doi: [10.1038/nature12156](https://doi.org/10.1038/nature12156)
- Chin A, Kyne PM, Walker TI, McAuley RB 2010 An integrated risk assessment for climate change: analysing the vulnerability of sharks and rays on Australia's Great Barrier Reef. *Glob. Chang. Biol.* 16, 1936–1953. doi: [10.1111/j.1365-2486.2009.02128.x](https://doi.org/10.1111/j.1365-2486.2009.02128.x)
- Christianen MJA, van Belzen J, Herman PMJ, van Katwijk MM, Lamers LPM et al. 2013 Low-canopy seagrass beds still provide important coastal protection services. *PLoS One* 8, e62413. doi: [10.1371/journal.pone.0062413](https://doi.org/10.1371/journal.pone.0062413)
- Christmas MJ, Breed MF, Lowe AJ 2016 Constraints to and conservation implications for climate change adaptation in plants. *Conserv. Genet.* 17, 305–320. doi: [10.1007/s10592-015-0782-5](https://doi.org/10.1007/s10592-015-0782-5)
- Ciampo G 2004 Ostracods as palaeoenvironmental indicators in the last 30 ky from the Tyrrhenian continental shelf. *Glob. Planet. Change* 40, 151–157. doi: [10.1016/S0921-8181\(03\)00105-X](https://doi.org/10.1016/S0921-8181(03)00105-X)
- Ciccarelli D 2014 Mediterranean coastal sand dune vegetation: influence of natural and anthropogenic factors. *Environ. Manage.* 54, 194–204. doi: [10.1007/s00267-014-0290-2](https://doi.org/10.1007/s00267-014-0290-2)
- Ciccarelli D, Pinna MS, Alquini F, Cogoni D, Ruocco M et al. 2017 Development of a coastal dune vulnerability index for Mediterranean ecosystems: A useful tool for coastal managers? *Estuar. Coast. Shelf Sci.* 187, 84–95. doi: [10.1016/j.ecss.2016.12.008](https://doi.org/10.1016/j.ecss.2016.12.008)
- Cid N, Bonada N, Carlson SM, Grantham TE, Gasith A et al. 2017 High Variability Is a Defining Component of Mediterranean-Climate Rivers and Their Biota. *Water* 9, 52. doi: [10.3390/w9010052](https://doi.org/10.3390/w9010052)
- CIESM 2009a Dynamics of Mediterranean deep waters. in *Malta, 27 - 30 May 2009*, 132 p.
- CIESM 2009b Phytoplankton responses to Mediterranean environmental changes. in *Tunis (Tunisia), 7 - 10 October 2009*, 120 p.
- Claudet S, Frascchetti J 2010 Human-driven impacts on marine habitats: A regional meta-analysis in the Mediterranean Sea. *Biol. Conserv.* 143, 2195–2206. doi: [10.1016/j.biocon.2010.06.004](https://doi.org/10.1016/j.biocon.2010.06.004)
- Clavero M, Blanco-Garrido F, Prenda J 2004 Fish fauna in Iberian Mediterranean river basins: Biodiversity, introduced species and damming impacts. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 14, 575–585. doi: [10.1002/aqc.636](https://doi.org/10.1002/aqc.636)
- Clavero M, Hermoso V, Levin N, Kark S 2010 BIODIVERSITY RESEARCH: Geographical linkages between threats and imperilment in freshwater fish in the Mediterranean Basin. *Divers. Distrib.* 16, 744–754. doi: [10.1111/j.1472-4642.2010.00680.x](https://doi.org/10.1111/j.1472-4642.2010.00680.x)
- Cohen M, Bilodeau C, Alexandre F, Godron M, Andrieu J et al. 2015 What is the plant biodiversity in a cultural landscape? A comparative, multi-scale and interdisciplinary

- plinary study in olive groves and vineyards (Mediterranean France). *Agric. Ecosyst. Environ.* 212, 175–186. doi: [10.1016/j.agee.2015.06.023](https://doi.org/10.1016/j.agee.2015.06.023)
- Coll M, Piroddi C, Albouy C, Ben Rais Lasram F, Cheung WWL et al. 2012 The Mediterranean Sea under siege: Spatial overlap between marine biodiversity, cumulative threats and marine reserves. *Glob. Ecol. Biogeogr.* 21, 465–480. doi: [10.1111/j.1466-8238.2011.00697.x](https://doi.org/10.1111/j.1466-8238.2011.00697.x)
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Ben Rais Lasram F et al. 2010 The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS One* 5, e11842. doi: [10.1371/journal.pone.0011842](https://doi.org/10.1371/journal.pone.0011842)
- Collet L, Ruelland D, Borrell-Estupina V, Dezetter A, Servat E 2013 Integrated modelling to assess long-term water supply capacity of a meso-scale Mediterranean catchment. *Sci. Total Environ.* 461–462, 528–540. doi: [10.1016/j.scitotenv.2013.05.036](https://doi.org/10.1016/j.scitotenv.2013.05.036)
- Collet L, Ruelland D, Borrell-Estupina V, Servat E 2014 Assessing the long-term impact of climatic variability and human activities on the water resources of a meso-scale Mediterranean catchment. *Hydrol. Sci. J.* 59, 1457–1469. doi: [10.1080/02626667.2013.842073](https://doi.org/10.1080/02626667.2013.842073)
- Collins S, Rost B, Rynearson TA 2014 Evolutionary potential of marine phytoplankton under ocean acidification. *Evol. Appl.* 7, 140–155. doi: [10.1111/eva.12120](https://doi.org/10.1111/eva.12120)
- Colloca F, Scarcella G, Libralato S 2017 Recent Trends and Impacts of Fisheries Exploitation on Mediterranean Stocks and Ecosystems. *Front. Mar. Sci.* 4. doi: [10.3389/fmars.2017.00244](https://doi.org/10.3389/fmars.2017.00244)
- Colls A, Ash N, Ikkala N 2009 Ecosystem-based adaptation: a natural response to climate change. Gland, Switzerland.
- Coma R, Ribes M, Serrano E, Jiménez E, Salat J et al. 2009 Global warming-enhanced stratification and mass mortality events in the Mediterranean. *Proc. Natl. Acad. Sci. U. S. A.* 106, 6176–6181. doi: [10.1073/pnas.0805801106](https://doi.org/10.1073/pnas.0805801106)
- Coma R, Serrano E, Linares C, Ribes M, Díaz D et al. 2011 Sea urchins predation facilitates coral invasion in a marine reserve. *PLoS One* 6, e22017. doi: [10.1371/journal.pone.0022017](https://doi.org/10.1371/journal.pone.0022017)
- Comeau S, Cornwall CE 2017 Contrasting Effects of Ocean Acidification on Coral Reef “Animal Forests” Versus Seaweed “Kelp Forests.” *Mar. Anim. For.*, 1083–1107. doi: [10.1007/978-3-319-21012-4\\_29](https://doi.org/10.1007/978-3-319-21012-4_29)
- Compa M, Alomar C, Wilcox C, van Sebille E, Lebreton L et al. 2019 Risk assessment of plastic pollution on marine diversity in the Mediterranean Sea. *Sci. Total Environ.* 678, 188–196. doi: [10.1016/j.scitotenv.2019.04.355](https://doi.org/10.1016/j.scitotenv.2019.04.355)
- Company T, Soriano P, Estrelles E, Mayoral O 2019 Seed bank longevity and germination ecology of invasive and native grass species from Mediterranean wetlands. *Folia Geobot.* 54, 151–161. doi: [10.1007/s12224-019-09350-7](https://doi.org/10.1007/s12224-019-09350-7)
- Condon RH, Duarte CM, Pitt KA, Robinson KL, Lucas CH et al. 2013 Recurrent jellyfish blooms are a consequence of global oscillations. *Proc. Natl. Acad. Sci. U. S. A.* 110, 1000–1005. doi: [10.1073/pnas.1210920110](https://doi.org/10.1073/pnas.1210920110)
- Constán-Nava S, Soliveres S, Torices R, Serra L, Bonet A 2015 Direct and indirect effects of invasion by the alien tree *Ailanthus altissima* on riparian plant communities and ecosystem multifunctionality. *Biol. Invasions* 17, 1095–1108. doi: [10.1007/s10530-014-0780-4](https://doi.org/10.1007/s10530-014-0780-4)
- Conversi A, Fonda Umani S, Peluso T, Molinero JC, Santojanni A et al. 2010 The Mediterranean Sea regime shift at the end of the 1980s, and intriguing parallelisms with other European basins. *PLoS One* 5, e10633. doi: [10.1371/journal.pone.0010633](https://doi.org/10.1371/journal.pone.0010633)
- Convertino M, Nardi F, Kiker GA, Munoz-Carpena R, Troccoli A et al. 2013 Epitomes of Bottom-Up Hydro-Geo-Climatological Analysis to Face Sea Level Rise in Complex Coastal Ecosystems, in *Climate Vulnerability: Understanding and Addressing Threats to Essential Resources* (Elsevier Inc.), 267–282. doi: [10.1016/B978-0-12-384703-4.00502-5](https://doi.org/10.1016/B978-0-12-384703-4.00502-5)
- Cook BI, Anchukaitis KJ, Touchan R, Meko DM, Cook ER 2016 Spatiotemporal drought variability in the Mediterranean over the last 900 years. *JGR Atmos.* 121, 1–15. doi: [10.1002/2015JD023929](https://doi.org/10.1002/2015JD023929). Received
- Cook ER, Seager R, Kushnir Y, Briffa KR, Büntgen U et al. 2015 Old World megadroughts and pluvials during the Common Era. *Sci. Adv.* 1, e1500561–e1500561. doi: [10.1126/sciadv.1500561](https://doi.org/10.1126/sciadv.1500561)
- Coomber FG, D’Inca M, Rosso M, Tepsich P, Notarbartolo di Sciarra G et al. 2016 Description of the vessel traffic within the north Pelagos Sanctuary: Inputs for marine spatial planning and management implications within an existing international Marine Protected Area. *Mar. Policy* 69, 102–113. doi: [10.1016/j.marpol.2016.04.013](https://doi.org/10.1016/j.marpol.2016.04.013)
- Cooper SD, Lake PS, Sabater S, Melack JM, Sabo JL 2013 The effects of land use changes on streams and rivers in Mediterranean climates. *Hydrobiologia* 719, 383–425. doi: [10.1007/s10750-012-1333-4](https://doi.org/10.1007/s10750-012-1333-4)
- Coppari M, Gori A, Rossi S 2014 Size, spatial, and bathymetrical distribution of the ascidian *Halocynthia papillosa* in Mediterranean coastal bottoms: Benthic-pelagic coupling implications. *Mar. Biol.* 161, 2079–2095. doi: [10.1007/s00227-014-2488-5](https://doi.org/10.1007/s00227-014-2488-5)
- Coppari M, Gori A, Viladrich N, Saponari L, Canepa A et al. 2016 The role of Mediterranean sponges in benthic-pelagic coupling processes: *Aplysina aerophoba* and *Axinella polypoides* case studies. *J. Exp. Mar. Bio. Ecol.* 477, 57–68. doi: [10.1016/j.jembe.2016.01.004](https://doi.org/10.1016/j.jembe.2016.01.004)
- Corbacho C, Sánchez JM, Costillo E 2003 Patterns of structural complexity and human disturbance of riparian vegetation in agricultural landscapes of a Mediterranean area. *Agric. Ecosyst. Environ.* 95, 495–507. doi: [10.1016/s0167-8809\(02\)00218-9](https://doi.org/10.1016/s0167-8809(02)00218-9)
- Cori B 1999 Spatial dynamics of Mediterranean coastal regions. *J. Coast. Conserv.* 5, 105–112. doi: [10.1007/BF02802747](https://doi.org/10.1007/BF02802747)
- Coro G, Vilas LG, Magliozzi C, Ellenbroek A, Scarponi P et al. 2018 Forecasting the ongoing invasion of *Lagocephalus sceleratus* in the Mediterranean Sea. *Ecol. Modell.*

- 371, 37–49. doi: [10.1016/j.ecolmodel.2018.01.007](https://doi.org/10.1016/j.ecolmodel.2018.01.007)
- Corrales X, Coll M, Ofir E, Heymans JJ, Steenbeek J et al. 2018 Future scenarios of marine resources and ecosystem conditions in the Eastern Mediterranean under the impacts of fishing, alien species and sea warming. *Sci. Rep.* 8, 14284. doi: [10.1038/s41598-018-32666-x](https://doi.org/10.1038/s41598-018-32666-x)
- Cortignani R, Dono G 2018 Agricultural policy and climate change: An integrated assessment of the impacts on an agricultural area of Southern Italy. doi: [10.1016/j.envsci.2017.12.003](https://doi.org/10.1016/j.envsci.2017.12.003)
- Cosentino SL, Testa G, Scordia D, Alexopoulou E 2012 Future yields assessment of bioenergy crops in relation to climate change and technological development in Europe. *Ital. J. Agron.* 7, 22. doi: [10.4081/ija.2012.e22](https://doi.org/10.4081/ija.2012.e22)
- Cózar A, Sanz-Martín M, Martí E, Ignacio González-Gordillo J, Ubeda B et al. 2015 Plastic accumulation in the Mediterranean Sea. *PLoS One* 10. doi: [10.1371/journal.pone.0121762](https://doi.org/10.1371/journal.pone.0121762)
- Cresson P, Fabri MC, Bouchoucha M, Brach Papa C, Chavanon F et al. 2014 Mercury in organisms from the Northwestern Mediterranean slope: Importance of food sources. *Sci. Total Environ.* 497–498, 229–238. doi: [10.1016/j.scitotenv.2014.07.069](https://doi.org/10.1016/j.scitotenv.2014.07.069)
- Crisci C, Bensoussan N, Romano J-C, Garrabou J 2011 Temperature anomalies and mortality events in marine communities: insights on factors behind differential mortality impacts in the NW Mediterranean. *PLoS One* 6, e23814. doi: [10.1371/journal.pone.0023814](https://doi.org/10.1371/journal.pone.0023814)
- Crisci C, Ledoux J-B, Mokhtar-Jamai K, Bally M, Bensoussan N et al. 2017 Regional and local environmental conditions do not shape the response to warming of a marine habitat-forming species. *Sci. Rep.* 7, 1–13. doi: [10.1038/s41598-017-05220-4](https://doi.org/10.1038/s41598-017-05220-4)
- Crise A, Allen JI, Baretta J, Crispi G, Mosetti R et al. 1999 The Mediterranean pelagic ecosystem response to physical forcing. in *Progress in Oceanography* (Pergamon), 219–243. doi: [10.1016/S0079-6611\(99\)00027-0](https://doi.org/10.1016/S0079-6611(99)00027-0)
- Croituru L 2007 How much are Mediterranean forests worth? *For. Policy Econ.* 9, 536–545. doi: [10.1016/j.forpol.2006.04.001](https://doi.org/10.1016/j.forpol.2006.04.001)
- Croituru L, Liagre L 2013 Contribution of Forests to a Green Economy in the Middle East and North Africa Evidence, drivers and policy orientations. 32.
- Crooks JA, Chang AL, Ruiz GM 2011 Aquatic pollution increases the relative success of invasive species. *Biol. Invasions* 13, 165–176. doi: [10.1007/s10530-010-9799-3](https://doi.org/10.1007/s10530-010-9799-3)
- Cubera E, Moreno G 2007 Effect of single *Quercus ilex* trees upon spatial and seasonal changes in soil water content in dehesas of central western Spain. *Ann. For. Sci.* 64, 355–364. doi: [10.1051/forest:2007012](https://doi.org/10.1051/forest:2007012)
- Curadi M, Picciarelli P, Lorenzi R, Graifenberg A, Geccarelli N 2005 Antioxidant activity and phenolic compounds in the edible parts of early and late Italian artichoke (*Cynara scolymus* L.) varieties. *Ital. J. Food Sci.* 17, 33–44.
- Custodio E, Andreu-Rodes JM, Aragón R, Estrela T, Ferrer J et al. 2016 Groundwater intensive use and mining in south-eastern peninsular Spain: Hydrogeological, economic and social aspects. *Sci. Total Environ.* 559, 302–316. doi: [10.1016/J.SCITOTENV.2016.02.107](https://doi.org/10.1016/J.SCITOTENV.2016.02.107)
- Cutini M, Agostinelli E, Acosta ATR, Molina JA 2010 Coastal salt-marsh zonation in Tyrrhenian central Italy and its relationship with other Mediterranean wetlands. *Plant Biosyst. - An Int. J. Deal. with all Asp. Plant Biol.* 144, 1–11. doi: [10.1080/11263500903178117](https://doi.org/10.1080/11263500903178117)
- Cuttelod A, García N, Abdul Malak D, Temple H, Katariya VA 2009 The Mediterranean: a biodiversity hotspot under threat. IUCN Gland, Switzerland: J.-C. Vié, C. Hilton-Taylor and S.N. Stuart
- D'Amario B, Ziveri P, Grelaud M, Oviedo A, Kralj M 2017 Coccolithophore haploid and diploid distribution patterns in the Mediterranean Sea: can a haplo-diploid life cycle be advantageous under climate change? *J. Plankton Res.* 39, 781–794. doi: [10.1093/plankt/fbx044](https://doi.org/10.1093/plankt/fbx044)
- D'Silva J, Webster J 2017 *The Meat Crisis.*, eds. D'Silva J, Webster J Abingdon, Oxon ; New York, NY : Routledge, 2017.: Routledge doi: [10.4324/9781315562032](https://doi.org/10.4324/9781315562032)
- Daccache A, Ciurana JS, Rodríguez-Díaz JA, Knox JW 2014 Water and energy footprint of irrigated agriculture in the Mediterranean region. *Environ. Res. Lett.* 9, 124014. doi: [10.1088/1748-9326/9/12/124014](https://doi.org/10.1088/1748-9326/9/12/124014)
- Dai A 2013 Increasing drought under global warming in observations and models. *Nat. Clim. Chang.* 3, 52–58. doi: [10.1038/nclimate1633](https://doi.org/10.1038/nclimate1633)
- Daliakopoulos IN, Panagea IS, Tsanis IK, Grillakis MG, Koutroulis AG et al. 2017 Yield Response of Mediterranean Rangelands under a Changing Climate. *L. Degrad. Dev.* 28, 1962–1972. doi: [10.1002/ldr.2717](https://doi.org/10.1002/ldr.2717)
- Daliakopoulos IN, Tsanis IK 2014 Climate-induced catastrophic shifts in pastoralism systems managed under the maximum sustainable yield model. *Oper. Res.* 14, 177–188. doi: [10.1007/s12351-014-0156-7](https://doi.org/10.1007/s12351-014-0156-7)
- Daly-Hassen H, Riera P, Mavsar R, Gammoudi A, García D 2017 Valuing trade-offs between local forest uses and environmental services in Tunisia. *J. Environ. Econ. Policy* 6, 268–282. doi: [10.1080/21606544.2017.1293566](https://doi.org/10.1080/21606544.2017.1293566)
- Daly Hassen H 2016 Assessment of the socio-economic value of the goods and services provided by Mediterranean forest ecosystems: critical and comparative analysis of studies conducted in Algeria, Lebanon, Morocco, Tunisia and Turkey. Valbonne.
- Daly Yahia MN, Batistic M, Lucic D, Fernández de Puelles ML, Licandro P et al. 2010 Are the outbreaks of *Pelagia noctiluca* (Forsskål, 1775) more frequent in the Mediterranean basin? in *Proceedings of the Joint ICES/CIESM Workshop to Compare Zooplankton Ecology and Methodologies between the Mediterranean and the North Atlantic (WKZEM)*, eds. Gislason A, Gorsky G, 8–14.
- Danovaro R, Carugati L, Berzano M, Cahill AE, Carvalho S et al. 2016 Implementing and Innovating Marine Monitoring Approaches for Assessing Marine Environmental Status. *Front. Mar. Sci.* 3. doi: [10.3389/fmars.2016.00213](https://doi.org/10.3389/fmars.2016.00213)

- Danovaro R, Carugati L, Boldrin A, Calafat A, Canals M et al. 2017 Deep-water zooplankton in the Mediterranean Sea: Results from a continuous, synchronous sampling over different regions using sediment traps. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 126, 103–114. doi: [10.1016/j.dsr.2017.06.002](https://doi.org/10.1016/j.dsr.2017.06.002)
- Danovaro R, Dell'Anno A, Fabiano M, Pusceddu A, Tselepidis A 2001 Deep-sea ecosystem response to climate changes: the eastern Mediterranean case study. *Trends Ecol. Evol.* 16, 505–510. doi: [10.1016/S0169-5347\(01\)02215-7](https://doi.org/10.1016/S0169-5347(01)02215-7)
- Danovaro R, Dell'Anno A, Pusceddu A 2004 Biodiversity response to climate change in a warm deep sea. *Ecol. Lett.* 7, 821–828. doi: [10.1111/j.1461-0248.2004.00634.x](https://doi.org/10.1111/j.1461-0248.2004.00634.x)
- Danovaro R, Fonda Umani S, Pusceddu A, Umani SF, Pusceddu A et al. 2009 Climate Change and the Potential Spreading of Marine Mucilage and Microbial Pathogens in the Mediterranean Sea. *PLoS One* 4, e7006. doi: [10.1371/journal.pone.0007006](https://doi.org/10.1371/journal.pone.0007006)
- Danovaro R, Pusceddu A 2007 Ecomanagement of biodiversity and ecosystem functioning in the Mediterranean Sea: concerns and strategies. *Chem. Ecol.* 23, 347–360. doi: [10.1080/02757540701653384](https://doi.org/10.1080/02757540701653384)
- Davies J, Poulsen L, Schulte-Herbrüggen B, Mackinnon K, Crawhall N et al. 2012 Conserving Dryland Biodiversity. Nairobi, Kenya.
- Davis SD, Heywood VH, Hamilton AC 1994 Centres of plant diversity. A guide and strategy for their conservation.
- Day JW, Ibáñez C, Pont D, Scarton F 2019 Status and sustainability of Mediterranean deltas: The case of the Ebro, Rhône, and Po Deltas and Venice Lagoon, in *Coasts and Estuaries*, eds. Wolanski E, Day JW, Elliott M, Ramachandran R (Elsevier), 237–249. doi: [10.1016/b978-0-12-814003-1.00014-9](https://doi.org/10.1016/b978-0-12-814003-1.00014-9)
- de Châtel F 2014 The Role of Drought and Climate Change in the Syrian Uprising: Untangling the Triggers of the Revolution. *Middle East. Stud.* 50, 521–535. doi: [10.1080/00263206.2013.850076](https://doi.org/10.1080/00263206.2013.850076)
- de Domenico S, de Rinaldis G, Paulmery M, Piraino S, Leone A 2019 Barrel jellyfish (*Rhizostoma pulmo*) as source of antioxidant peptides. *Mar. Drugs* 17, 134. doi: [10.3390/md17020134](https://doi.org/10.3390/md17020134)
- De Frutos A, Olea PP, Mateo-Tomás P 2015 Responses of medium- and large-sized bird diversity to irrigation in dry cereal agroecosystems across spatial scales. *Agric. Ecosyst. Environ.* 207, 141–152. doi: [10.1016/j.agee.2015.04.009](https://doi.org/10.1016/j.agee.2015.04.009)
- de Groot RS, Alkemade R, Braat L, Hein L, Willemen L 2010 Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecol. Complex.* 7, 260–272. doi: [10.1016/j.ecocom.2009.10.006](https://doi.org/10.1016/j.ecocom.2009.10.006)
- de la Fuente B, Saura S, Beck PSA 2018 Predicting the spread of an invasive tree pest: The pine wood nematode in Southern Europe. *J. Appl. Ecol.* 55, 2374–2385. doi: [10.1111/1365-2664.13177](https://doi.org/10.1111/1365-2664.13177)
- de Montmollin B, Strahm W 2005 *The Top 50 Mediterranean Island Plants. Wild Plants At the Brink of Extinction, and What is Needed to Save Them*. IUCN/SSC Mediterranean Islands Plant Specialist Group. IUCN, Gland, Switzerland and Cambridge. doi: [10.1017/s0030605306270200](https://doi.org/10.1017/s0030605306270200)
- de Stephanis R, Giménez J, Carpinelli E, Gutierrez-Exposito C, Cañadas A 2013 As main meal for sperm whales: plastics debris. *Mar. Pollut. Bull.* 69, 206–214. doi: [0.1016/j.marpolbul.2013.01.033](https://doi.org/10.1016/j.marpolbul.2013.01.033)
- de Vente J, Verduyn R, Verstraeten G, Vanmaercke M, Poesen J 2011 Factors controlling sediment yield at the catchment scale in NW Mediterranean geosystems. *J. Soils Sediments* 11, 690–707. doi: [10.1007/s11368-011-0346-3](https://doi.org/10.1007/s11368-011-0346-3)
- Debolini M, Marraccini E, Dubeuf JP, Geizendorffer IR, Guerra CA et al. 2018 Land and farming system dynamics and their drivers in the Mediterranean Basin. *Land use policy* 75, 702–710. doi: [10.1016/j.landusepol.2017.07.010](https://doi.org/10.1016/j.landusepol.2017.07.010)
- Defalco LA, Detling JK, Tracy CR, Warren SD 2001 Physiological variation among native and exotic winter annual plants associated with microbiotic crusts in the Mojave Desert. *Plant Soil* 234, 1–14. doi: [10.1023/A:1010323001006](https://doi.org/10.1023/A:1010323001006)
- Degeai JP, Devillers B, Blanchemanche P, Dezileau L, Oueslati H et al. 2017 Fluvial response to the last Holocene rapid climate change in the Northwestern Mediterranean coastlands. *Glob. Planet. Change* 152, 176–186. doi: [10.1016/j.gloplacha.2017.03.008](https://doi.org/10.1016/j.gloplacha.2017.03.008)
- Deidun A 2018 Back with a bang – an unexpected massive bloom of *Cassiopea andromeda* (Forskaal, 1775) in the Maltese Islands, nine years after its first appearance. *BiolInvasions Rec.* 7, 399–404. doi: [10.3391/bir.2018.7.4.07](https://doi.org/10.3391/bir.2018.7.4.07)
- Deininger A, Faithfull CL, Lange K, Bayer T, Vidussi F et al. 2016 Simulated terrestrial runoff triggered a phytoplankton succession and changed seston stoichiometry in coastal lagoon mesocosms. *Mar. Environ. Res.* 119, 40–50. doi: [10.1016/j.marenvres.2016.05.001](https://doi.org/10.1016/j.marenvres.2016.05.001)
- del Río M, Bravo-Oviedo A, Pretzsch H, Löf M, Ruiz-Peinado R 2017 A review of thinning effects on Scots pine stands: from growth and yield to new challenges under global change. *For. Syst.* 26, 9. doi: [10.1007/bf02158077](https://doi.org/10.1007/bf02158077)
- Delgado JA, Del Grosso SJ, Ogle SM 2010 15N isotopic crop residue cycling studies and modeling suggest that IPCC methodologies to assess residue contributions to N<sub>2</sub>O-N emissions should be reevaluated. *Nutr. Cycl. Agroecosystems* 86, 383–390. doi: [10.1007/s10705-009-9300-9](https://doi.org/10.1007/s10705-009-9300-9)
- Delpont G, Vogt-Schilb H, Munoz F, Richard F, Schatz B 2018 Diachronic variations in the distribution of butterflies and dragonflies linked to recent habitat changes in Western Europe. *Insect Conserv. Divers.* 12, 49–68. doi: [10.1111/icad.12309](https://doi.org/10.1111/icad.12309)
- Demirkesen AC, Evrendilek F, Berberoglu S 2008 Quan-

- tifying coastal inundation vulnerability of Turkey to sea-level rise. *Environ. Monit. Assess.* 138, 101–106. doi: [10.1007/s10661-007-9746-7](https://doi.org/10.1007/s10661-007-9746-7)
- den Herder M, Moreno G, Mosquera-Losada RM, Palma JHN, Sidiropoulou A et al. 2017 Current extent and stratification of agroforestry in the European Union. *Agric. Ecosyst. Environ.* 241, 121–132. doi: [10.1016/j.agee.2017.03.005](https://doi.org/10.1016/j.agee.2017.03.005)
- di Camillo CG, Cerrano C 2015 Mass mortality events in the NW Adriatic Sea: phase shift from slow- to fast-growing organisms. *PLoS One* 10, e0126689. doi: [10.1371/journal.pone.0126689](https://doi.org/10.1371/journal.pone.0126689)
- Diacono M, Fiore A, Farina R, Canali S, di Bene C et al. 2016 Combined agro-ecological strategies for adaptation of organic horticultural systems to climate change in Mediterranean environment. *Ital. J. Agron.* 11, 85. doi: [10.4081/ija.2016.730](https://doi.org/10.4081/ija.2016.730)
- Dias BB, Hart MB, Smart CW, Hall-Spencer JM 2010 Modern seawater acidification: the response of foraminifera to high-CO<sub>2</sub> conditions in the Mediterranean Sea. *J. Geol. Soc. London.* 167, 843–846. doi: [10.1144/0016-76492010-050](https://doi.org/10.1144/0016-76492010-050)
- Díaz-Almela E, Marba N, Duarte CM 2007 Consequences of Mediterranean warming events in seagrass (*Posidonia oceanica*) flowering records. *Glob. Chang. Biol.* 13, 224–235. doi: [10.1111/j.1365-2486.2006.01260.x](https://doi.org/10.1111/j.1365-2486.2006.01260.x)
- Dibari C, Argenti G, Catolfi F, Moriondo M, Staglianò N et al. 2015 Pastoral suitability driven by future climate change along the Apennines. *Ital. J. Agron.* 10, 109. doi: [10.4081/ija.2015.659](https://doi.org/10.4081/ija.2015.659)
- Ding H, Nunes PALD, Teelucksingh S 2010 European forests and carbon sequestration services: An economic assessment of climate change impacts. *FEEM Work. Pap.* 10.2010. doi: [10.2139/ssrn.1557689](https://doi.org/10.2139/ssrn.1557689)
- Dise NB, Ashmore M, Belyazid S, Bleeker A, Bobbink R et al. 2011 Nitrogen as a threat to European terrestrial biodiversity, in *The European Nitrogen Assessment* (Cambridge University Press), 463–494. doi: [10.1017/cbo9780511976988.023](https://doi.org/10.1017/cbo9780511976988.023)
- Djema A, Messaoudene M 2009 The Algerian forest: Current situation and prospect. in *Modelling, Valuing and Managing Mediterranean Forest Ecosystems for Non-Timber Goods and Services*, eds. Palahi M, Birot Y, Bravo F, Gorris E (Thessaloniki, Greece: European Forest Institute), 17–28. <https://www.efi.int/publications-bank/modelling-valuing-and-managing-mediterranean-forest-ecosystems-non-timber-goods>
- Doblas-Miranda E, Alonso R, Arnan X, Bermejo V, Brotons L et al. 2017 A review of the combination among global change factors in forests, shrublands and pastures of the Mediterranean Region: Beyond drought effects. *Glob. Planet. Change* 148, 42–54. doi: [10.1016/j.gloplacha.2016.11.012](https://doi.org/10.1016/j.gloplacha.2016.11.012)
- Doblas-Miranda E, Martínez-Vilalta J, Lloret F, Alvarez A, Avila A et al. 2015 Reassessing global change research priorities in mediterranean terrestrial ecosystems: How far have we come and where do we go from here? *Glob. Ecol. Biogeogr.* 24, 25–43. doi: [10.1111/geb.12224](https://doi.org/10.1111/geb.12224)
- Domene E, Saurí D 2007 Urbanization and class-produced natures: Vegetable gardens in the Barcelona Metropolitan Region. *Geoforum* 38, 287–298. doi: [10.1016/j.geoforum.2006.03.004](https://doi.org/10.1016/j.geoforum.2006.03.004)
- Doney SC, Ruckelshaus M, Emmett Duffy J, Barry JP, Chan F et al. 2012 Climate Change Impacts on Marine Ecosystems. *Ann. Rev. Mar. Sci.* 4, 11–37. doi: [10.1146/annurev-marine-041911-111611](https://doi.org/10.1146/annurev-marine-041911-111611)
- Dono G, Cortignani R, Dell'Unto D, Deligios PA, Doro L et al. 2016 Winners and losers from climate change in agriculture: Insights from a case study in the Mediterranean basin. *Agric. Syst.* 147, 65–75. doi: [10.1016/j.agsy.2016.05.013](https://doi.org/10.1016/j.agsy.2016.05.013)
- Dono G, Cortignani R, Giraldo L, Pasqui M, Roggero PP 2014 Income Impacts of Climate Change: Irrigated Farming in the Mediterranean and Expected Changes in Probability of Favorable and Adverse Weather Conditions. *Ger. J. Agric. Econ.* 63, 177–186. doi: [issn 0002-1121](https://doi.org/10.1007/s00442-015-3229-2)
- Dorman M, Perevolotsky A, Sarris D, Svoray T 2015 The effect of rainfall and competition intensity on forest response to drought: lessons learned from a dry extreme. *Oecologia* 177, 1025–1038. doi: [10.1007/s00442-015-3229-2](https://doi.org/10.1007/s00442-015-3229-2)
- Dorman M, Svoray T, Perevolotsky A, Sarris D 2013 Forest performance during two consecutive drought periods: Diverging long-term trends and short-term responses along a climatic gradient. *For. Ecol. Manage.* 310, 1–9. doi: [10.1016/j.foreco.2013.08.009](https://doi.org/10.1016/j.foreco.2013.08.009)
- Draredja MA, Frihi H, Boualleg C, Gofart A, Abadie E et al. 2019 Seasonal variations of phytoplankton community in relation to environmental factors in a protected meso-oligotrophic southern Mediterranean marine ecosystem (Mellah lagoon, Algeria) with an emphasis of HAB species. *Environ. Monit. Assess.* 191, 603. doi: [10.1007/s10661-019-7708-5](https://doi.org/10.1007/s10661-019-7708-5)
- Drira Z, Hamza A, Belhassen M, Ayadi H, Bouain A et al. 2008 Dynamics of dinoflagellates and environmental factors during the summer in the Gulf of Gabes (Tunisia, Eastern Mediterranean Sea). *Sci. Mar.* 72. doi: [10.3989/scimar.2008.72n159](https://doi.org/10.3989/scimar.2008.72n159)
- Duarte CM, Agustí S, Agawin NSR 2000 Response of a Mediterranean phytoplankton community to increased nutrient inputs: a mesocosm experiment. *Mar. Ecol. Prog. Ser.* 195, 61–70. <https://digital.csic.es/handle/10261/54164>
- Dudgeon D, Arthington AH, Gessner MO, Kawabata Z-I, Knowler DJ et al. 2006 Freshwater biodiversity: importance, threats, status and conservation challenges. *Biol. Rev.* 81, 163–182. doi: [10.1017/s1464793105006950](https://doi.org/10.1017/s1464793105006950)
- Dudley N 2008 *Guidelines for applying protected area management categories*. Gland, Switzerland: IUCN doi: [10.2305/iucn.ch.2008.paps.2.en](https://doi.org/10.2305/iucn.ch.2008.paps.2.en)
- Duguy B, Alloza JA, Roder A, Vallejo R, Pastor F 2007 Mod-

- elling the effects of landscape fuel treatments on fire growth and behaviour in a Mediterranean landscape (eastern Spain). *Int. J. Wildl. Fire* 16, 619–632. doi: [10.1071/WF06101](https://doi.org/10.1071/WF06101)
- Dulvy NK, Fowler SL, Musick JA, Cavanagh RD, Kyne PM et al. 2014 Extinction risk and conservation of the world's sharks and rays. *Elife* 3, e00590. doi: [10.7554/eLife.00590](https://doi.org/10.7554/eLife.00590)
- Durant JM, Hjermmann DØ, Frederiksen M, Charrassin J-B, Le Maho Y et al. 2009 Pros and cons of using seabirds as ecological indicators. *Clim. Res.* 39, 115–129. doi: [10.3354/cr00798](https://doi.org/10.3354/cr00798)
- EEA 2004 Agriculture and the environment in the EU accession countries. Implications of applying the EU common agricultural policy. Copenhagen, Denmark
- EEA 2017 Climate change, impacts and vulnerability in Europe 2016. An indicator-based report. doi: [10.2800/534806](https://doi.org/10.2800/534806)
- EEA 2018 European Waters - Assessment of Status and Pressures. Copenhagen. doi: [10.2800/303664](https://doi.org/10.2800/303664)
- Egea FJ, Torrente RG, Aguilar A 2018 An efficient agro-industrial complex in Almería (Spain): Towards an integrated and sustainable bioeconomy model. doi: [10.1016/j.nbt.2017.06.009](https://doi.org/10.1016/j.nbt.2017.06.009)
- Ejarque A, Julià R, Reed JM, Mesquita-Joanes F, Marco-Barba J et al. 2016 Coastal evolution in a Mediterranean microtidal zone: Mid to late Holocene natural dynamics and human management of the Castelló Lagoon, NE Spain. *PLoS One* 11, e0155446. doi: [10.1371/journal.pone.0155446](https://doi.org/10.1371/journal.pone.0155446)
- El Jaouhari N, Abouabdillah A, Bouabid R, Bourioum M, Aleya L et al. 2018 Assessment of sustainable deficit irrigation in a Moroccan apple orchard as a climate change adaptation strategy. *Sci. Total Environ.* 642, 574–581. doi: [10.1016/J.SCITOTENV.2018.06.108](https://doi.org/10.1016/J.SCITOTENV.2018.06.108)
- El Sayed Frihy O, Deabes EA, Shereet SM, Abdalla FA 2010 Alexandria-Nile Delta coast, Egypt: Update and future projection of relative sea-level rise. *Environ. Earth Sci.* 61, 253–273. doi: [10.1007/s12665-009-0340-x](https://doi.org/10.1007/s12665-009-0340-x)
- Elkiran G, Aslanova F, Hiziroglu S, Elkiran G, Aslanova F et al. 2019 Effluent Water Reuse Possibilities in Northern Cyprus. *Water* 11, 191. doi: [10.3390/w11020191](https://doi.org/10.3390/w11020191)
- Elsen HGM, Hessel R, Stringer LC, Daliakopoulos IN, Tsanis I et al. 2017 *CASCADE Catastrophic shifts in drylands: how can we prevent ecosystem degradation? Final Publishable Summary: EU - FP7 project Grant Agreement Number 283068*. Wageningen: Wageningen Environmental Research (Alterra) <http://edepot.wur.nl/441608>
- Elvira B, Almodovar A 2001 Freshwater fish introductions in Spain: facts and figures at the beginning of the 21st century. *J. Fish Biol.* 59, 323–331. doi: [10.1111/j.1095-8649.2001.tb01393.x](https://doi.org/10.1111/j.1095-8649.2001.tb01393.x)
- Emeis K-C, Struck U, Schulz H-M, Rosenberg R, Bernasconi SM et al. 2000 Temperature and salinity variations of Mediterranean Sea surface waters over the last 16,000 years from records of planktonic stable oxygen isotopes and alkenone unsaturation ratios. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 158, 259–280. doi: [10.1016/S0031-0182\(00\)00053-5](https://doi.org/10.1016/S0031-0182(00)00053-5)
- Emmanouloudis D, Garcia Rodriguez J, Zaimis G, Gimenez Suare M, Filippidis E 2011 Euro-Mediterranean torrents: case studies on tools that can improve their management., in *Mountain Ecosystems: dynamics, Management and Conservation.*, ed. Richards KE (Nova Science Publishers, Hauppauge), 1–44.
- Enríquez AR, Marcos M, Álvarez-Ellacuría A, Orfila A, Gomis D 2017 Changes in beach shoreline due to sea level rise and waves under climate change scenarios: application to the Balearic Islands (western Mediterranean). *Nat. Hazards Earth Syst. Sci.* 17, 1075–1089. doi: [10.5194/nhess-17-1075-2017](https://doi.org/10.5194/nhess-17-1075-2017)
- Erol A, Randhir TO 2012 Climatic change impacts on the ecohydrology of Mediterranean watersheds. *Clim. Change* 114, 319–341. doi: [10.1007/s10584-012-0406-8](https://doi.org/10.1007/s10584-012-0406-8)
- Escalas A, Ferraton F, Paillon C, Vidy G, Carcaillet F et al. 2015 Spatial variations in dietary organic matter sources modulate the size and condition of fish juveniles in temperate lagoon nursery sites. *Estuar. Coast. Shelf Sci.* 152, 78–90. doi: [10.1016/j.ecss.2014.11.021](https://doi.org/10.1016/j.ecss.2014.11.021)
- Escudero A, Martínez I, de la Cruz A, Otálora MAG, Maestre FT 2007 Soil lichens have species-specific effects on the seedling emergence of three gypsophile plant species. *J. Arid Environ.* 70, 18–28. doi: [10.1016/j.jaridenv.2006.12.019](https://doi.org/10.1016/j.jaridenv.2006.12.019)
- Esper J, Frank D, Büntgen U, Verstege A, Luterbacher J et al. 2007 Long-term drought severity variations in Morocco. *Geophys. Res. Lett.* 34, 1–5. doi: [10.1029/2007GL030844](https://doi.org/10.1029/2007GL030844)
- Estevez P, Castro D, Pequeño-Valtierra A, Giraldez J, Gago-Martinez A 2019 Emerging Marine Biotoxins in Seafood from European Coasts: Incidence and Analytical Challenges. *Foods* 8, 149. doi: [10.3390/foods8050149](https://doi.org/10.3390/foods8050149)
- Fader M, Shi S, Von Bloh W, Bondeau A, Cramer W 2016 Mediterranean irrigation under climate change: More efficient irrigation needed to compensate for increases in irrigation water requirements. *Hydrol. Earth Syst. Sci.* 20, 953–973. doi: [10.5194/hess-20-953-2016](https://doi.org/10.5194/hess-20-953-2016)
- Fady-Welterlen B 2005 Is there really more biodiversity in Mediterranean forest ecosystems? *Taxon* 54, 905–910.
- Fady B, Médail F 2004 Mediterranean forest ecosystems, in *Encyclopedia of forest science*, eds. Burley J, Evans J, Youngquist JA (London), 1403–1414. [https://www.researchgate.net/profile/Thierry-Gauquelin/publication/311480684\\_The\\_Mediterranean\\_region\\_under\\_climate\\_change\\_a\\_scientific\\_update/links/584fe7d008aecb6bd8d1dee8/The-Mediterranean-region-under-climate-change-a-scientific-update.pdf](https://www.researchgate.net/profile/Thierry-Gauquelin/publication/311480684_The_Mediterranean_region_under_climate_change_a_scientific_update/links/584fe7d008aecb6bd8d1dee8/The-Mediterranean-region-under-climate-change-a-scientific-update.pdf)
- Fagerholm N, Torralba M, Burgess PJ, Plieninger T 2016 A systematic map of ecosystem services assessments around European agroforestry. *Ecol. Indic.* 62, 47–65. doi: [10.1016/J.ECOLIND.2015.11.016](https://doi.org/10.1016/J.ECOLIND.2015.11.016)
- Fagnano M, Fiorentino N 2018 The EcoRemed protocol for



- an integrated agronomic approach to characterization and remediation of contaminated soils. *Ital. J. Agron. Agron.* 13, 1–68. doi: [10.4081/ija.2018.1348](https://doi.org/10.4081/ija.2018.1348)
- Fahrig L 2003 Effects of Habitat Fragmentation on Biodiversity. *Annu. Rev. Ecol. Evol. Syst.* 34, 487–515. doi: [10.1146/annurev.ecolsys.34.011802.132419](https://doi.org/10.1146/annurev.ecolsys.34.011802.132419)
- Faluccci A, Maiorano L, Boitani L 2007 Changes in land-use/land-cover patterns in Italy and their implications for biodiversity conservation. *Landsc. Ecol.* 22, 617–631. doi: [10.1007/s10980-006-9056-4](https://doi.org/10.1007/s10980-006-9056-4)
- Falkenberg LJ, Bellerby RGJ, Connell SD, Fleming LE, Maycock B et al. 2020 Ocean Acidification and Human Health. *Int. J. Environ. Res. Public Health* 17, 4563. doi: [10.3390/ijerph17124563](https://doi.org/10.3390/ijerph17124563)
- Fang X, Hou X, Li X, Hou W, Nakaoka M et al. 2018 Ecological connectivity between land and sea: a review. *Ecol. Res.* 33, 51–61. doi: [10.1007/s11284-017-1549-x](https://doi.org/10.1007/s11284-017-1549-x)
- FAO 2016 Optimiser la production des biens et services par les écosystèmes boisés méditerranéens dans un contexte de changements globaux. Agents et causes de la déforestation et dégradation dans les sites pilotes du projet.
- FAO 2017 FAOSTAT.
- FAO, Plan Bleu 2013 State of Mediterranean Forests 2013. Rome, Italy. <http://www.fao.org/3/CA2081EN/ca2081en.PDF>
- FAO, Plan Bleu 2018 State of Mediterranean Forests 2018. Rome <http://www.fao.org/3/CA2081EN/ca2081en.PDF>
- Fatorić S, Chelleri L 2012 Vulnerability to the effects of climate change and adaptation: The case of the Spanish Ebro Delta. *Ocean Coast. Manag.* 60, 1–10. doi: [10.1016/j.ocecoaman.2011.12.015](https://doi.org/10.1016/j.ocecoaman.2011.12.015)
- Fauvel P 1937 Les fonds de pêche près d' Alexandrie. XI. Annélides Polychètes. Le Caire
- Feagin RA, Sherman DJ, Grant WE 2005 Coastal erosion, global sea-level rise, and the loss of sand dune plant habitats. *Front. Ecol. Environ.* 3, 359–364. doi: [10.1890/1540-9295\(2005\)003\[0359:CEGSRA\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2005)003[0359:CEGSRA]2.0.CO;2)
- Feio MJ, Aguiar FC, Almeida SFP, Ferreira J, Ferreira MT et al. 2014 Least Disturbed Condition for European Mediterranean rivers. *Sci. Total Environ.* 476–477, 745–756. doi: [10.1016/j.scitotenv.2013.05.056](https://doi.org/10.1016/j.scitotenv.2013.05.056)
- Feio MJ, Coimbra CN, Graça MAS, Nichols SJ, Norris RH 2010 The influence of extreme climatic events and human disturbance on macroinvertebrate community patterns of a Mediterranean stream over 15 y. *J. North Am. Benthol. Soc.* 29, 1397–1409. doi: [10.1899/09-158.1](https://doi.org/10.1899/09-158.1)
- Feki M, Ben Brahim M, Feki-Sahnoun W, Mahfoudi M, Sammari C et al. 2016 Seasonal and daily fluctuation of dinoflagellates during spring tide periods in Kerkennah islands (southern coast of Tunisia). [https://www.semanticscholar.org/paper/SEASon-AL-AnD-DAILY-FLUCTUATION-of-DINOFLAGELLATES-\[-Feki-Brahim/38c19b91b5fd6285d825de155d-98ab84dc9eec26](https://www.semanticscholar.org/paper/SEASon-AL-AnD-DAILY-FLUCTUATION-of-DINOFLAGELLATES-[-Feki-Brahim/38c19b91b5fd6285d825de155d-98ab84dc9eec26) [Accessed September 9, 2019]
- Fenu G, Cogoni D, Ulian T, Bacchetta G 2013 The impact of human trampling on a threatened coastal Mediterranean plant: The case of *Anchusa littorea* Moris (Boraginaceae). *Flora - Morphol. Distrib. Funct. Ecol. Plants* 208, 104–110. doi: [10.1016/j.flora.2013.02.003](https://doi.org/10.1016/j.flora.2013.02.003)
- Fernández C, Vega JA, Jiménez E, Vieira DCS, Merino A et al. 2012 Seeding and mulching + seeding effects on post-fire runoff, soil erosion and species diversity in Galicia (NW Spain). *L. Degrad. Dev.* 23, 150–156. doi: [10.1002/ldr.1064](https://doi.org/10.1002/ldr.1064)
- Fernández Calzado MR, Molero Mesa J, Merzouki A, Casares Porcel M 2012 Vascular plant diversity and climate change in the upper zone of Sierra Nevada, Spain. *Plant Biosyst.* 146, 1044–1053. doi: [10.1080/11263504.2012.710273](https://doi.org/10.1080/11263504.2012.710273)
- Fernández de Puelles ML, Molinero JC 2007 North Atlantic climate control on plankton variability in the Balearic Sea, western Mediterranean. *Geophys. Res. Lett.* 34. doi: [10.1029/2006GL028354](https://doi.org/10.1029/2006GL028354)
- Fernández R, Bertrand A, Reis R, Mourato MP, Martins LL et al. 2013 Growth and physiological responses to cadmium stress of two populations of *Dittrichia viscosa* (L.) Greuter. *J. Hazard. Mater.* 244–245, 555–562. doi: [10.1016/j.jhazmat.2012.10.044](https://doi.org/10.1016/j.jhazmat.2012.10.044)
- Ferreira JG, Andersen JH, Borja Á, Bricker SB, Camp J et al. 2011 Overview of eutrophication indicators to assess environmental status within the European Marine Strategy Framework Directive. *Estuar. Coast. Shelf Sci.* 93, 117–131. doi: [10.1016/J.ECSS.2011.03.014](https://doi.org/10.1016/J.ECSS.2011.03.014)
- Ferreira MT, Albuquerque A, Aguiar FC, Catarino LF 2001 Seasonal and yearly variations of macrophytes in a southern Iberian river. *Int. Vereinigung für Theor. und Angew. Limnol. Verhandlungen*, 27, 3833–3837. doi: [10.1080/03680770.1998.11901701](https://doi.org/10.1080/03680770.1998.11901701)
- Ferreira MT, Albuquerque A, Aguiar FC, Sidorkewicz N 2002 Assessing reference sites and ecological quality of river plant assemblages from an Iberian basin using a multivariate approach. *Fundam. Appl. Limnol.* 155, 121–145. doi: [10.1127/archiv-hydrobiol/155/2002/121](https://doi.org/10.1127/archiv-hydrobiol/155/2002/121)
- Ferreira T, Globevnik L, Schinegger R 2019 Water Stressors in Europe: New Threats in the Old World. *Mult. Stress. River Ecosyst.*, 139–155. doi: [10.1016/b978-0-12-811713-2.00008-x](https://doi.org/10.1016/b978-0-12-811713-2.00008-x)
- Filipe AF, Lawrence JE, Bonada N 2013 Vulnerability of stream biota to climate change in mediterranean climate regions: a synthesis of ecological responses and conservation challenges. *Hydrobiologia* 719, 331–351. doi: [10.1007/s10750-012-1244-4](https://doi.org/10.1007/s10750-012-1244-4)
- Fine M, Tsadok R, Meron D, Cohen S, Milazzo M 2017 Environmental sensitivity of *Neogoniolithon brassica-florida* associated with vermetid reefs in the Mediterranean Sea. *ICES J. Mar. Sci.* 74, 1074–1082. doi: [10.1093/icesjms/fsw167](https://doi.org/10.1093/icesjms/fsw167)
- Flecha S, Pérez FF, García-Lafuente J, Sammartino S, Ríos AF et al. 2015 Trends of pH decrease in the Mediterranean Sea through high frequency observational data: indication of ocean acidification in the basin. *Sci. Rep.* 5, 16770. doi: [10.1038/srep16770](https://doi.org/10.1038/srep16770)

- Forès E 1992 Nutrient loading and drainage channel response in a ricefield system. *Hydrobiologia* 230, 193–200. doi: [10.1007/BF00036565](https://doi.org/10.1007/BF00036565)
- Forey E, Chapelet B, Vitasse Y, Tilquin M, Touzard B et al. 2008 The relative importance of disturbance and environmental stress at local and regional scales in French coastal sand dunes. *J. Veg. Sci.* 19, 493–502. doi: [10.3170/2008-8-18392](https://doi.org/10.3170/2008-8-18392)
- Fortibuoni T, Giovanardi O, Pranovi F, Raicevich S, Solidoro C et al. 2017 Analysis of Long-Term Changes in a Mediterranean Marine Ecosystem Based on Fishery Landings. *Front. Mar. Sci.* 4. doi: [10.3389/fmars.2017.00033](https://doi.org/10.3389/fmars.2017.00033)
- Fossi MC, Panti C 2017 Sentinel Species of Marine Ecosystems. *Oxford Res. Encycl. Environ. Sci.* doi: [10.1093/acrefore/9780199389414.013.110](https://doi.org/10.1093/acrefore/9780199389414.013.110)
- Fossi MC, Panti C, Bainsi M, Lavers JL 2018 A review of plastic-associated pressures: cetaceans of the Mediterranean Sea and Eastern Australian Ssearwaters as case studies. *Front. Mar. Sci.* 5, 1–10. doi: [10.3389/fmars.2018.00173](https://doi.org/10.3389/fmars.2018.00173)
- Fossi MC, Panti C, Guerranti C, Coppola D, Giannetti M et al. 2012 Are baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale (*Balaenoptera physalus*). *Mar. Pollut. Bull.* 64, 2374–2379. doi: [10.1016/j.marpolbul.2012.08.013](https://doi.org/10.1016/j.marpolbul.2012.08.013)
- Fossi MC, Panti C, Marsili L, Maltese S, Spinsanti G et al. 2013 The Pelagos Sanctuary for Mediterranean marine mammals: Marine Protected Area (MPA) or marine polluted area? The case study of the striped dolphin (*Stenella coeruleoalba*). *Mar. Pollut. Bull.* 70, 64–72. doi: [10.1016/j.marpolbul.2013.02.013](https://doi.org/10.1016/j.marpolbul.2013.02.013)
- Foster N, Sciberras M, Jackson E, Ponge B, Toison V et al. 2014 Assessing the ecological coherence of the channel MPA network.
- Fouache E, Ghilardi M, Vouvalidis K, Syrides G, Styllas M et al. 2008 Contribution on the Holocene reconstruction of Thessaloniki coastal plain, Greece. *J. Coast. Res.* 24, 1161–1218. <https://www.jstor.org/stable/40065156> [Accessed December 21, 2018]
- Fouchy K, McClain ME, Conallin J, O'Brien G 2019 Multiple Stressors in African Freshwater Systems. *Mult. Stress. River Ecosyst.*, 179–191. doi: [10.1016/B978-0-12-811713-2.00010-8](https://doi.org/10.1016/B978-0-12-811713-2.00010-8)
- Foufopoulos J, Kilpatrick AM, Ives AR 2011 Climate change and elevated extinction rates of reptiles from Mediterranean islands. *Am. Nat.* 177, 119–129. doi: [10.1086/657624](https://doi.org/10.1086/657624)
- Fraga H, Malheiro AC, Moutinho-Pereira J, Santos JA 2012 An overview of climate change impacts on European viticulture. *Food Energy Secur.* 1, 94–110. doi: [10.1002/fes3.14](https://doi.org/10.1002/fes3.14)
- Fraixedas S, Galewski T, Ribeiro-Lopes S, Loh J, Blondel J et al. 2019 Estimating biodiversity changes in the Camargue wetlands: An expert knowledge approach. *PLoS One* 14, e0224235. doi: [10.1371/journal.pone.0224235](https://doi.org/10.1371/journal.pone.0224235)
- Franco A, Franzoi P, Malavasi S, Riccato F, Torricelli P et al. 2006 Use of shallow water habitats by fish assemblages in a Mediterranean coastal lagoon. *Estuar. Coast. Shelf Sci.* 66, 67–83. doi: [10.1016/j.ecss.2005.07.020](https://doi.org/10.1016/j.ecss.2005.07.020)
- Frantar P, Hrvatin M 2006 Pretočni režimi v Sloveniji med letoma 1971 in 2000. *Geogr. Vestn.* 77, 115–127.
- Frigola J, Moreno A, Cacho I, Canals M, Sierro FJ et al. 2008 Evidence of abrupt changes in Western Mediterranean Deep Water circulation during the last 50kyr: A high-resolution marine record from the Balearic Sea. *Quat. Int.* 181, 88–104. doi: [10.1016/j.quaint.2007.06.016](https://doi.org/10.1016/j.quaint.2007.06.016)
- Fuentes VL, Purcell JE, Condon RH, Lombard F, Lucas CH 2018 Jellyfish blooms: advances and challenges. *Mar. Ecol. Prog. Ser.* 591, 3–5. doi: [10.3354/meps12536](https://doi.org/10.3354/meps12536)
- Funes I, Aranda X, Biel C, Carbó J, Camps F et al. 2016 Future climate change impacts on apple flowering date in a Mediterranean subbasin. *Agric. Water Manag.* 164, 19–27. doi: [10.1016/J.AGWAT.2015.06.013](https://doi.org/10.1016/J.AGWAT.2015.06.013)
- Fyllas NM, Troumbis AY 2009 Simulating vegetation shifts in north-eastern Mediterranean mountain forests under climatic change scenarios. *Glob. Ecol. Biogeogr.* 18, 64–77. doi: [10.1111/j.1466-8238.2008.00419.x](https://doi.org/10.1111/j.1466-8238.2008.00419.x)
- Gabrié C, Lagabrielle E, Bissery C, Crochelet E, Meola B et al. 2012 The Status of Marine Protected Areas in the Mediterranean Sea.
- Gabrielides GP, Golik A, Loizides L, Marino MG, Bingel F et al. 1991 Man-made garbage pollution on the Mediterranean coastline. *Mar. Pollut. Bull.* 23, 437–441. doi: [10.1016/0025-326X\(91\)90713-3](https://doi.org/10.1016/0025-326X(91)90713-3)
- Gaget E, Galewski T, Jiguet F, Le Viol I 2018 Waterbird communities adjust to climate warming according to conservation policy and species protection status. *Biol. Conserv.* 227, 205–212. doi: [10.1016/j.biocon.2018.09.019](https://doi.org/10.1016/j.biocon.2018.09.019)
- Galgani F, Claro F, Depledge M, Fossi C 2014 Monitoring the impact of litter in large vertebrates in the Mediterranean Sea within the European Marine Strategy Framework Directive (MSFD): Constraints, specificities and recommendations. *Mar. Environ. Res.* 100, 3–9. doi: [10.1016/j.marenvres.2014.02.003](https://doi.org/10.1016/j.marenvres.2014.02.003)
- Galil BS 2000 A Sea Under Siege – Alien Species in the Mediterranean. *Biol. Invasions* 2, 177–186. doi: [10.1023/A:1010057010476](https://doi.org/10.1023/A:1010057010476)
- Galil BS, Marchini A, Occhipinti-Ambrogi A 2018 East is east and West is west? Management of marine bioinvasions in the Mediterranean Sea. *Estuar. Coast. Shelf Sci.* 201, 7–16. doi: [10.1016/j.ecss.2015.12.021](https://doi.org/10.1016/j.ecss.2015.12.021)
- Galili E, Zviely D, Weinstein-Evron M 2005 Holocene sea-level changes and landscape evolution on the northern Carmel coast (Israel). *Méditerranée. Rev. géographique des pays méditerranéens / J. Mediterr. Geogr.*, 79–86. doi: [10.4000/mediterranee.1912](https://doi.org/10.4000/mediterranee.1912)
- Galli G, Solidoro C, Lovato T 2017 Marine heat waves hazard 3D maps and the risk for low motility organisms in a warming Mediterranean Sea. *Front. Mar. Sci.* 4, 136. doi: [10.3389/fmars.2017.00136](https://doi.org/10.3389/fmars.2017.00136)

- Gallisai R, Peters F, Volpe G, Basart S, Baldasano JM 2014 Saharan Dust Deposition May Affect Phytoplankton Growth in the Mediterranean Sea at Ecological Time Scales. *PLoS One* 9, e110762. doi: [10.1371/journal.pone.0110762](https://doi.org/10.1371/journal.pone.0110762)
- Gambi C, Pusceddu A, Benedetti-Cecchi L, Danovaro R 2014 Species richness, species turnover and functional diversity in nematodes of the deep Mediterranean Sea: searching for drivers at different spatial scales. *Glob. Ecol. Biogeogr.* 23, 24–39. doi: [10.1111/geb.12094](https://doi.org/10.1111/geb.12094)
- Gamfeldt L, Snäll T, Bagchi R, Jonsson M, Gustafsson L et al. 2013 Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat. Commun.* 4, 1340. doi: [10.1038/ncomms2328](https://doi.org/10.1038/ncomms2328)
- Gao K, Gao G, Wang Y, Dupont S 2020 Impacts of ocean acidification under multiple stressors on typical organisms and ecological processes. *Mar. Life Sci. Technol.* 2, 279–291. doi: [10.1007/s42995-020-00048-w](https://doi.org/10.1007/s42995-020-00048-w)
- García-Comas C, Stemmann L, Ibañez F, Berline L, Maz-zocchi MG et al. 2011 Zooplankton long-term changes in the NW Mediterranean Sea: Decadal periodicity forced by winter hydrographic conditions related to large-scale atmospheric changes? *J. Mar. Syst.* 87, 216–226. doi: [10.1016/j.jmarsys.2011.04.003](https://doi.org/10.1016/j.jmarsys.2011.04.003)
- García-Gómez H, Garrido JL, Vivanco MG, Lassaletta L, Rábago I et al. 2014 Nitrogen deposition in Spain: Modeled patterns and threatened habitats within the Natura 2000 network. *Sci. Total Environ.* 485–486, 450–460. doi: [10.1016/j.scitotenv.2014.03.112](https://doi.org/10.1016/j.scitotenv.2014.03.112)
- García-Nieto AP, García-Llorente M, Iniesta-Arandia I, Martín-López B 2013 Mapping forest ecosystem services: From providing units to beneficiaries. *Ecosyst. Serv.* 4, 126–138. doi: [10.1016/j.ecoser.2013.03.003](https://doi.org/10.1016/j.ecoser.2013.03.003)
- García-Nieto AP, Geijzendorffer IR, Baró F, Roche PK, Bondeau A et al. 2018 Impacts of urbanization around Mediterranean cities: Changes in ecosystem service supply. *Ecol. Indic.* 91, 589–606. doi: [10.1016/j.ecolind.2018.03.082](https://doi.org/10.1016/j.ecolind.2018.03.082)
- García-Romero A, Muñoz J, Andrés N, Palacios D 2010 Relationship between climate change and vegetation distribution in the Mediterranean mountains: Manzanares Head valley, Sierra De Guadarrama (Central Spain). *Clim. Change* 100, 645–666. doi: [10.1007/s10584-009-9727-7](https://doi.org/10.1007/s10584-009-9727-7)
- García-Ruiz JM, Lana-Renault N 2011 Hydrological and erosive consequences of farmland abandonment in Europe, with special reference to the Mediterranean region - A review. *Agric. Ecosyst. Environ.* 140, 317–338. doi: [10.1016/j.agee.2011.01.003](https://doi.org/10.1016/j.agee.2011.01.003)
- García-Ruiz JM, López Moreno JI, Vicente-Serrano SM, Lasanta-Martínez T, Beguería S 2011 Mediterranean water resources in a global change scenario. *Earth-Science Rev.* 105, 121–139. doi: [10.1016/j.earscirev.2011.01.006](https://doi.org/10.1016/j.earscirev.2011.01.006)
- García C, Amengual A, Homar V, Zamora A 2017 Losing water in temporary streams on a Mediterranean island: Effects of climate and land-cover changes. *Glob. Planet. Change* 148, 139–152. doi: [10.1016/j.gloplacha.2016.11.010](https://doi.org/10.1016/j.gloplacha.2016.11.010)
- Garde RJ 2006 *River morphology*. , ed. New Age International.
- Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R et al. 2013 Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science (80-. )*. 339, 1608–1611. doi: [10.1126/science.1230200](https://doi.org/10.1126/science.1230200)
- Garofano-Gomez V, Martínez-Capel F, Peredo-Parada M, Olaya Marin EJ, Muñoz Mas R et al. 2011 Assessing hydromorphological and floristic patterns along a regulated Mediterranean river: The Serpis River (Spain). *Limnetica* 30, 307–328.
- Garófano Gómez V 2013 Riparian vegetation patterns according to hydrogeomorphological factors at different spatial and temporal scales in Mediterranean rivers. doi: [10.4995/thesis/10251/29395](https://doi.org/10.4995/thesis/10251/29395)
- Garrabou J, Ballesteros E, Zabala M 2002 Structure and dynamics of north-western Mediterranean rocky benthic communities along a depth gradient. *Estuar. Coast. Shelf Sci.* 55, 493–508. doi: [10.1006/ecss.2001.0920](https://doi.org/10.1006/ecss.2001.0920)
- Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P et al. 2009 Mass mortality in Northwestern Mediterranean rocky benthic communities: Effects of the 2003 heat wave. *Glob. Chang. Biol.* 15, 1090–1103. doi: [10.1111/j.1365-2486.2008.01823.x](https://doi.org/10.1111/j.1365-2486.2008.01823.x)
- Garrabou J, Gómez-Gras D, Ledoux J-B, Linares C, Bensoussan N et al. 2019 Collaborative Database to Track Mass Mortality Events in the Mediterranean Sea. *Front. Mar. Sci.* 6, 707. doi: [10.3389/fmars.2019.00707](https://doi.org/10.3389/fmars.2019.00707)
- Garrabou J, Perez T, Sartoretto S, Harmelin JG 2001 Mass mortality event in red coral *Corallium rubrum* populations in the Provence region (France, NW Mediterranean). *Mar. Ecol. Prog. Ser.* 217, 263–272. doi: [10.3354/meps217263](https://doi.org/10.3354/meps217263)
- Garrido P, Elbakidze M, Angelstam P, Plieninger T, Pulido F et al. 2017 Stakeholder perspectives of wood-pasture ecosystem services: A case study from Iberian dehesas. *Land use policy* 60, 324–333. doi: [10.1016/j.landusepol.2016.10.022](https://doi.org/10.1016/j.landusepol.2016.10.022)
- Gasith A, Resh VH 1999 Streams in Mediterranean Climate Regions: Abiotic Influences and Biotic Responses to Predictable Seasonal Events. *Annu. Rev. Ecol. Syst.* 30, 51–81. doi: [10.1146/annurev.ecolsys.30.1.51](https://doi.org/10.1146/annurev.ecolsys.30.1.51)
- Gatti G, Bianchi CN, Parravicini V, Rovere A, Peirano A et al. 2015 Ecological change, sliding baselines and the importance of historical data: lessons from combing observational and quantitative data on a temperate reef over 70 years. *PLoS One* 10, e0118581. doi: [10.1371/journal.pone.0118581](https://doi.org/10.1371/journal.pone.0118581)
- Gauquelin T, Bertaudiere V, Montes N, Badri W, Asmode JF 1999 Endangered stands of thuriferous juniper in the western Mediterranean basin: Ecological status, conservation and management. *Biodivers. Conserv.* 8, 1479–1498. doi: [10.1023/A:1008966808796](https://doi.org/10.1023/A:1008966808796)

- Gauquelin T, Michon G, Joffre R, Duponnois R, Génin D et al. 2018 Mediterranean forests, land use and climate change: a social-ecological perspective. *Reg. Environ. Chang.* 18, 623–636. doi: [10.1007/s10113-016-0994-3](https://doi.org/10.1007/s10113-016-0994-3)
- Gazeau F, Alliouane S, Bock C, Bramanti L, López Correa M et al. 2014 Impact of ocean acidification and warming on the Mediterranean mussel (*Mytilus galloprovincialis*). *Front. Mar. Sci.* 1, 62. doi: [10.3389/fmars.2014.00062](https://doi.org/10.3389/fmars.2014.00062)
- Gazol A, Camarero JJ, Vicente-Serrano SM, Sánchez-Salguero R, Gutiérrez E et al. 2018 Forest resilience to drought varies across biomes. *Glob. Chang. Biol.* 24, 2143–2158. doi: [10.1111/gcb.14082](https://doi.org/10.1111/gcb.14082)
- Geijzendorffer IR, Beltrame C, Chazee L, Gaget E, Galewski T et al. 2019a A More Effective Ramsar Convention for the Conservation of Mediterranean Wetlands. *Front. Ecol. Evol.* 7. doi: [10.3389/fevo.2019.00021](https://doi.org/10.3389/fevo.2019.00021)
- Geijzendorffer IR, Galewski T, Guelmami A, Perennou C, Popoff N et al. 2019b Mediterranean Wetlands: A Gradient from Natural Resilience to a Fragile Social-Ecosystem. *Atlas Ecosyst. Serv.*, 83–89. doi: [10.1007/978-3-319-96229-0\\_14](https://doi.org/10.1007/978-3-319-96229-0_14)
- Gendaszek AS, Magirl CS, Czuba CR 2012 Geomorphic response to flow regulation and channel and floodplain alteration in the gravel-bedded Cedar River, Washington, USA. *Geomorphology* 179, 258–268. doi: [10.1016/j.geomorph.2012.08.017](https://doi.org/10.1016/j.geomorph.2012.08.017)
- Genev M 2003 Patterns of runoff change in Bulgaria. *IAHS-AISH Publ.*, 79–85.
- Gennari G, Tamburini F, Ariztegui D, Hajdas I, Spezzaferri S 2009 Geochemical evidence for high-resolution variations during deposition of the Holocene S1 sapropel on the Cretan Ridge, Eastern Mediterranean. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 273, 239–248. doi: [10.1016/j.palaeo.2008.06.007](https://doi.org/10.1016/j.palaeo.2008.06.007)
- Gentilella T, Camarero JJ, Colangelo M, Nolè A, Ripullone F 2017 Drought-induced oak decline in the western mediterranean region: An overview on current evidences, mechanisms and management options to improve forest resilience. *IForest* 10, 796–806. doi: [10.3832/ifer2017-010](https://doi.org/10.3832/ifer2017-010)
- Gharib SM, El-Sherif ZM, Abdel-Halim AM, Radwan AA 2011 Phytoplankton and environmental variables as a water quality indicator for the beaches at Matrouh, south-eastern Mediterranean Sea, Egypt: an assessment. *Oceanologia* 53, 819–836. doi: [10.5697/oc.53-3.819](https://doi.org/10.5697/oc.53-3.819)
- Giakoumakis SG, Baloutsos G 1997 Investigation of trend in hydrological time series of the Evinos River basin. *Hydrol. Sci. J.* 42, 81–88. doi: [10.1080/02626669709492007](https://doi.org/10.1080/02626669709492007)
- Giakoumi S, Hermoso V, Carvalho SB, Markantonatou V, Dags M et al. 2019 Conserving European biodiversity across realms. *Conserv. Lett.* 12, e12586. doi: [10.1111/conl.12586](https://doi.org/10.1111/conl.12586)
- Giakoumi S, Sini M, Gerovasileiou V, Mazor T, Beher J et al. 2013 Ecoregion-based conservation planning in the Mediterranean: Dealing with large-scale heterogeneity. *PLoS One* 8, e76449. doi: [10.1371/journal.pone.0076449](https://doi.org/10.1371/journal.pone.0076449)
- Gili JM, Sardá R, Madurell T, Rossi S 2014 Zoobenthos, in *The Mediterranean Sea*, eds. Goffredo S, Dubinsky Z (Dordrecht, Netherlands: Springer). doi: [10.1007/978-94-007-6704-1\\_12](https://doi.org/10.1007/978-94-007-6704-1_12)
- Giuggiola A, Bugmann H, Zingg A, Dobbertin M, Rigling A 2013 Reduction of stand density increases drought resistance in xeric Scots pine forests. *For. Ecol. Manage.* 310, 827–835. doi: [10.1016/j.foreco.2013.09.030](https://doi.org/10.1016/j.foreco.2013.09.030)
- Glibert PM, Anderson DM, Gentien P, Granéli E, Sellner K 2005 The global, complex phenomena of Harmful Algal Blooms. *Oceanography* 18, 136–147. doi: [10.5670/oceanog.2005.49](https://doi.org/10.5670/oceanog.2005.49)
- Gober P 2010 Desert urbanization and the challenges of water sustainability. *Curr. Opin. Environ. Sustain.* 2, 144–150. doi: [10.1016/j.cosust.2010.06.006](https://doi.org/10.1016/j.cosust.2010.06.006)
- Goffart A, Collignon A, Lejeune P, Hecq J-H 2017 Thresholds of plankton community change in a Mediterranean coastal area : results from a long-term (1979–2014) time series. in *IMBIZO V : Marine biosphere research for a sustainable ocean : Linking ecosystems, future states and resource management* <https://orbi.uliege.be/handle/2268/219919> [Accessed September 9, 2019].
- Goffart A, Hecq J-H, Legendre L 2002 Changes in the development of the winter-spring phytoplankton bloom in the Bay of Calvi (NW Mediterranean) over the last two decades: a response to changing climate? *Mar. Ecol. Prog. Ser.* 236. doi: [10.3354/meps236045](https://doi.org/10.3354/meps236045)
- Gómez-Aparicio L, Zamora R, Castro J, Hódar JA 2008 Facilitation of tree saplings by nurse plants: Microhabitat amelioration or protection against herbivores? *J. Veg. Sci.* 19, 161–172. doi: [10.3170/2008-8-18347](https://doi.org/10.3170/2008-8-18347)
- Gómez-Gras D, Linares C, de Caralt S, Cebrian E, Frleta-Valiç M et al. 2019 Response diversity in Mediterranean coralligenous assemblages facing climate change: Insights from a multispecific thermotolerance experiment. *Ecol. Evol.* 9, 4168–4180. doi: [10.1002/ece3.5045](https://doi.org/10.1002/ece3.5045)
- González E, González-Sanchis M, Cabezas Á, Comín FA, Muller E 2010 Recent Changes in the Riparian Forest of a Large Regulated Mediterranean River: Implications for Management. *Environ. Manage.* 45, 669–681. doi: [10.1007/s00267-010-9441-2](https://doi.org/10.1007/s00267-010-9441-2)
- Gordo O, Sanz JJ 2010 Impact of climate change on plant phenology in Mediterranean ecosystems. *Glob. Chang. Biol.* 16, 1082–1106. doi: [10.1111/j.1365-2486.2009.02084.x](https://doi.org/10.1111/j.1365-2486.2009.02084.x)
- Gori A, Viladrich N, Gili J-M, Kotta M, Cucio C et al. 2012 Reproductive cycle and trophic ecology in deep versus shallow populations of the Mediterranean gorgonian *Eunicella singularis* (Cap de Creus, northwestern Mediterranean Sea). *Coral Reefs* 31, 823–837. doi: [10.1007/s00338-012-0904-1](https://doi.org/10.1007/s00338-012-0904-1)
- Gottfried M, Pauli H, Futschik A, Akhalkatsi M, Barančok

- P et al. 2012 Continent-wide response of mountain vegetation to climate change. *Nat. Clim. Chang.* 2, 111–115. doi: [10.1038/nclimate1329](https://doi.org/10.1038/nclimate1329)
- Goyet C, Hassoun AER, Gemayel E, Touratier F, Abboud-Abi Saab M et al. 2016 Thermodynamic forecasts of the Mediterranean Sea acidification. *Mediterr. Mar. Sci.* 17, 508–518. doi: [10.12681/mms.1487](https://doi.org/10.12681/mms.1487)
- Grantham TE, Figueroa R, Prat N 2013 Water management in mediterranean river basins: a comparison of management frameworks, physical impacts, and ecological responses. *Hydrobiologia* 719, 451–482. doi: [10.1007/s10750-012-1289-4](https://doi.org/10.1007/s10750-012-1289-4)
- Greenhill L, Kenter JO, Dannevig H 2020 Adaptation to climate change–related ocean acidification: An adaptive governance approach. *Ocean Coast. Manag.* 191, 105176. doi: [10.1016/j.ocecoaman.2020.105176](https://doi.org/10.1016/j.ocecoaman.2020.105176)
- Grelaud M, Marino G, Ziveri P, Rohling EJ 2012 Abrupt shoaling of the nutricline in response to massive freshwater flooding at the onset of the last interglacial sapropel event. *Paleoceanography* 27. doi: [10.1029/2012PA002288](https://doi.org/10.1029/2012PA002288)
- Griffin MT, Montz BE, S. Arrigo J 2013 Evaluating climate change induced water stress: A case study of the Lower Cape Fear basin, NC. *Appl. Geogr.* 40, 115–128. doi: [10.1016/j.apgeog.2013.02.009](https://doi.org/10.1016/j.apgeog.2013.02.009)
- Griffith AW, Gobler CJ 2020 Harmful algal blooms: A climate change co-stressor in marine and freshwater ecosystems. *Harmful Algae* 91, 101590. doi: [10.1016/j.hal.2019.03.008](https://doi.org/10.1016/j.hal.2019.03.008)
- Grizzetti B, Lique C, Pistocchi A, Vigiak O, Zulian G et al. 2019 Relationship between ecological condition and ecosystem services in European rivers, lakes and coastal waters. *Sci. Total Environ.* 671, 452–465. doi: [10.1016/j.scitotenv.2019.03.155](https://doi.org/10.1016/j.scitotenv.2019.03.155)
- Grouillet B, Fabre J, Ruelland D, Dezetter A 2015 Historical reconstruction and 2050 projections of water demand under anthropogenic and climate changes in two contrasted Mediterranean catchments. *J. Hydrol.* 522, 684–696. doi: [10.1016/j.jhydrol.2015.01.029](https://doi.org/10.1016/j.jhydrol.2015.01.029)
- Grove AT, Rackham O 2003 *The nature of Mediterranean Europe: an ecological history*. New Haven: Yale University Press.
- Gubbay S, Sanders N, Haynes T, Janssen JAM, Rodwell JR et al. 2016 *European Red list of habitats. Part 1. Marine habitats*. Luxembourg: Publications Office of the European Union.
- Gudka M, Davies J, Poulsen L, Schulte-Herbrüggen B, MacKinnon K et al. 2014 Conserving dryland biodiversity: a future vision of sustainable dryland development. *Biodiversity* 15, 143–147. doi: [10.1080/14888386.2014.930716](https://doi.org/10.1080/14888386.2014.930716)
- Guerra CA, Metzger MJ, Maes J, Pinto-Correia T 2016 Policy impacts on regulating ecosystem services: looking at the implications of 60 years of landscape change on soil erosion prevention in a Mediterranean silvo-pastoral system. *Landsc. Ecol.* 31, 271–290. doi: [10.1007/s10980-015-0241-1](https://doi.org/10.1007/s10980-015-0241-1)
- Guerrero E, Gili J-M, Grinyó J, Raya V, Sabatés A 2018 Long-term changes in the planktonic cnidarian community in a mesoscale area of the NW Mediterranean. *PLoS One* 13, e0196431. doi: [10.1371/journal.pone.0196431](https://doi.org/10.1371/journal.pone.0196431)
- Gugliemetti F, Cinquepalmi F, Astiaso Garcia D 2007 The use of environmental sensitivity indices (ESI) maps for the evaluation of oil spill risk in Mediterranean coastlines and coastal waters. *WIT Trans. Ecol. Environ.* 102, 8. doi: [10.2495/sdp070572](https://doi.org/10.2495/sdp070572)
- Guidetti P, Bianchi CN, La Mesa G, Modena M, Morri C et al. 2002 Abundance and size structure of *Thalassoma pavo* (Pisces: Labridae) in the western Mediterranean Sea: variability at different spatial scales. *J. Mar. Biol. Assoc. United Kingdom* 82, 495–500. doi: [10.1017/S0025315402005775](https://doi.org/10.1017/S0025315402005775)
- Guijarro M, Madrigal J, Hernando C, Sánchez de Ron D, Vázquez de la Cueva A 2017 Las repoblaciones y los incendios forestales, in *La Restauración Forestal de España: 75 Años de Una Ilusión*, eds. Permán García J, Iriarte Goñi I, Lario Leza FJ (Madrid, España: Ministerio de Agricultura y Pesca, Alimentación y Medio Ambiente), 343–371.
- Guiot J, Cramer W 2016 Climate change: The 2015 Paris Agreement thresholds and Mediterranean basin ecosystems. *Science (80-. )*. 354, 4528–4532. doi: [10.1126/science.aah5015](https://doi.org/10.1126/science.aah5015)
- Guiot J, Kaniewski D 2015 *The Mediterranean Basin and Southern Europe in a warmer world: What can we learn from the past?*. doi: [10.3389/feart.2015.00028](https://doi.org/10.3389/feart.2015.00028)
- Günlü A, Kadioğulları AI, Keleş S, Başkent EZ 2009 Spatio-temporal changes of landscape pattern in response to deforestation in Northeastern Turkey: A case study in Rize. *Environ. Monit. Assess.* 148, 127–137. doi: [10.1007/s10661-007-0144-y](https://doi.org/10.1007/s10661-007-0144-y)
- Guyot V, Castagnérol B, Vialatte A, Deconchat M, Jactel H 2016 Tree diversity reduces pest damage in mature forests across Europe. *Biol. Lett.* 12, 20151037. doi: [10.1098/rsbl.2015.1037](https://doi.org/10.1098/rsbl.2015.1037)
- Haddeland I, Heinke J, Biemans H, Eisner S, Flörke M et al. 2014 Global water resources affected by human interventions and climate change. *Proc. Natl. Acad. Sci. U. S. A.* 111, 3251–3256. doi: [10.1073/pnas.1222475110](https://doi.org/10.1073/pnas.1222475110)
- Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M et al. 2008 Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* 454, 96–99. doi: [10.1038/nature07051](https://doi.org/10.1038/nature07051)
- Hall DC, Hall J V, Murray SN 2002 Contingent Valuation of Marine Protected Areas: Southern California Rocky Intertidal Ecosystems. *Nat. Resour. Model.* 15, 335–368. doi: [10.1111/j.1939-7445.2002.tb00093.x](https://doi.org/10.1111/j.1939-7445.2002.tb00093.x)
- Hallegatte S, Billé R, Magnan AK, Gemenne F 2009 *La Méditerranée au futur : des impacts du changement climatique aux enjeux de l'adaptation*.
- Halpern BS, Ebert CM, Kappel C V., Madin EMP, Micheli F et al. 2009 Global priority areas for incorporating land-sea connections in marine conservation. *Con-*

- serv. Lett.* 2, 189–196.  
doi: [10.1111/j.1755-263x.2009.00060.x](https://doi.org/10.1111/j.1755-263x.2009.00060.x)
- Hamidov A, Helming K, Bellocchi G, Bojar W, Dalgaard T et al. 2018 Impacts of climate change adaptation options on soil functions: A review of European case-studies. *L. Degrad. Dev.* doi: [10.1002/ldr.3006](https://doi.org/10.1002/ldr.3006)
- Hannah DM, Demuth S, van Lanen HAJ, Looser U, Prudhomme C et al. 2011 Large-scale river flow archives: Importance, current status and future needs. *Hydrol. Process.* 25, 1191–1200. doi: [10.1002/hyp.7794](https://doi.org/10.1002/hyp.7794)
- Hannah L, Roehrdanz PR, Ikegami M, Shepard A V., Shaw MR et al. 2013 Climate change, wine, and conservation. *Proc. Natl. Acad. Sci. U. S. A.* 110, 6907–6912. doi: [10.1073/pnas.1210127110](https://doi.org/10.1073/pnas.1210127110)
- Hansen VD, Reiss KC 2015 Chapter 16 - Threats to Marsh Resources and Mitigation, in *Coastal and Marine Hazards, Risks, and Disasters*, eds. Shroder JF, Ellis JT, Sherman DJ (Boston: Elsevier), 467–494. <http://www.sciencedirect.com/science/article/pii/B9780123964830000169>
- Harland AD, Davies PS, Fixter LM 1992 Lipid content of some Caribbean corals in relation to depth and light. *Mar. Biol.* 113, 357–361. doi: [10.1007/bf00349159](https://doi.org/10.1007/bf00349159)
- Harley CDG, Randall Hughes A, Hultgren KM, Miner BG, Sorte CJB et al. 2006 The impacts of climate change in coastal marine systems. *Ecol. Lett.* 9, 228–241. doi: [10.1111/j.1461-0248.2005.00871.x](https://doi.org/10.1111/j.1461-0248.2005.00871.x)
- Harris SE 2012 Cyprus as a degraded landscape or resilient environment in the wake of colonial intrusion. *Proc. Natl. Acad. Sci. U. S. A.* 109, 3670–3675. doi: [10.1073/pnas.1114085109](https://doi.org/10.1073/pnas.1114085109)
- Harsch MA, Hulme PE, McGlone MS, Duncan RP 2009 Are treelines advancing? A global meta-analysis of treeline response to climate warming. *Ecol. Lett.* 12, 1040–1049. doi: [10.1111/j.1461-0248.2009.01355.x](https://doi.org/10.1111/j.1461-0248.2009.01355.x)
- Hassoun AER, Fakhri M, Raad N, Abboud-Abi Saab M, Gemayel E et al. 2019 The carbonate system of the Eastern-most Mediterranean Sea, Levantine Sub-basin: Variations and drivers. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 164, 54–73.
- Hassoun AER, Gemayel E, Krasakopoulou E, Goyet C, Abboud-Abi Saab M et al. 2015 Acidification of the Mediterranean Sea from anthropogenic carbon penetration. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 102, 1–15. doi: [10.1016/j.dsr.2015.04.005](https://doi.org/10.1016/j.dsr.2015.04.005)
- Hassoun AER, Ujević I, Mahfouz C, Fakhri M, Roje-Busatto R et al. 2021 Occurrence of domoic acid and cyclic imines in marine biota from Lebanon-Eastern Mediterranean Sea. *Sci. Total Environ.* 755, 142542. doi: [10.1016/j.scitotenv.2020.142542](https://doi.org/10.1016/j.scitotenv.2020.142542)
- Heinrich I, Touchan R, Dorado Liñán I, Vos H, Helle G 2013 Winter-to-spring temperature dynamics in Turkey derived from tree rings since AD 1125. *Clim. Dyn.* 41, 1685–1701. doi: [10.1007/s00382-013-1702-3](https://doi.org/10.1007/s00382-013-1702-3)
- Hejazi M, Edmonds J, Clarke L, Kyle P, Davies E et al. 2014 Long-term global water projections using six socioeconomic scenarios in an integrated assessment modeling framework. *Technol. Forecast. Soc. Change* 81, 205–226. doi: [10.1016/j.techfore.2013.05.006](https://doi.org/10.1016/j.techfore.2013.05.006)
- Helman D, Osem Y, Yakir D, Lensky IM 2017 Relationships between climate, topography, water use and productivity in two key Mediterranean forest types with different water-use strategies. *Agric. For. Meteorol.* 232, 319–330. doi: [10.1016/j.agrformet.2016.08.018](https://doi.org/10.1016/j.agrformet.2016.08.018)
- Hendriks IE, Olsen YS, Duarte CM 2017 Light availability and temperature, not increased CO<sub>2</sub>, will structure future meadows of *Posidonia oceanica*. *Aquat. Bot.* 139, 32–36. doi: [10.1016/j.aquabot.2017.02.004](https://doi.org/10.1016/j.aquabot.2017.02.004)
- Henle K, Dick D, Harpke A, Kühn I, Schweiger O et al. 2010 Climate change impacts on European amphibians and reptiles, in *Biodiversity and climate change: Reports and guidance developed under the Bern Convention - Volume I (Nature and Environment N°156)* (Strasbourg, France: Council of Europe Publishing), 225–305.
- Hennekam R, Jilbert T, Schnetger B, de Lange GJ 2014 Solar forcing of Nile discharge and sapropel S1 formation in the early to middle Holocene eastern Mediterranean. *Paleoceanography* 29, 343–356. doi: [10.1002/2013pa002553](https://doi.org/10.1002/2013pa002553)
- Henry BK, Eckard RJ, Beauchemin KA 2018 Review: Adaptation of ruminant livestock production systems to climate changes. *animal* 12, s445–s456. doi: [10.1017/S1751731118001301](https://doi.org/10.1017/S1751731118001301)
- Herbert ER, Boon PI, Burgin AJ, Neubauer SC, Franklin RB et al. 2015 A global perspective on wetland salinization: ecological consequences of a growing threat to freshwater wetlands. *Ecosphere* 6, art206. doi: [10.1890/ES14-00534.1](https://doi.org/10.1890/ES14-00534.1)
- Hernández-Morcillo M, Burgess PJ, Mirck J, Pantera A, Plieninger T 2018 Scanning agroforestry-based solutions for climate change mitigation and adaptation in Europe. *Environ. Sci. Policy* 80, 44–52. doi: [10.1016/J.ENVSCI.2017.11.013](https://doi.org/10.1016/J.ENVSCI.2017.11.013)
- Herrera AM, Dudley TL 2003 Reduction of riparian arthropod abundance and diversity as a consequence of giant reed (*Arundo donax*) invasion. *Biol. Invasions* 5, 167–177. doi: [10.1023/A:1026190115521](https://doi.org/10.1023/A:1026190115521)
- Herrmann M, Estournel C, Adloff F, Diaz F 2014 Impact of climate change on the northwestern Mediterranean Sea pelagic planktonic ecosystem and associated carbon cycle. *JGR Ocean.* 119, 5815–5836. doi: [10.1002/2014JC010016](https://doi.org/10.1002/2014JC010016)
- Heywood V 2017 The nature and composition of urban plant diversity in the Mediterranean. *Flora Mediterranea* 27, 195–220. doi: [10.7320/FlMedit27.195](https://doi.org/10.7320/FlMedit27.195)
- Hirich A, Choukr-Allah R, Jacobsen S-E 2014 Quinoa in Morocco - Effect of Sowing Dates on Development and Yield. *J. Agron. Crop Sci.* 200, 371–377. doi: [10.1111/jac.12071](https://doi.org/10.1111/jac.12071)
- Hočvar S, Malej A, Boldin B, Purcell JE 2018 Seasonal fluctuations in population dynamics of *Aurelia aurita* polyps in situ with a modelling perspective. *Mar. Ecol. Prog. Ser.* 591, 155–166. doi: [10.3354/meps12387](https://doi.org/10.3354/meps12387)
- Hódar JA, Zamora R 2004 Herbivory and climatic warm-

- ing: a Mediterranean outbreaking caterpillar attacks a relict, boreal pine species. *Biodivers. Conserv.* 13, 493–500. doi: [10.1023/b:bioc.0000009495.95589.a7](https://doi.org/10.1023/b:bioc.0000009495.95589.a7)
- Hooke JM 2006 Human impacts on fluvial systems in the Mediterranean region. *Geomorphology* 79, 311–335. doi: [10.1016/j.geomorph.2006.06.036](https://doi.org/10.1016/j.geomorph.2006.06.036)
- Howes EL, Joos F, Eakin CM, Gattuso J-P 2015 An updated synthesis of the observed and projected impacts of climate change on the chemical, physical and biological processes in the oceans. *Front. Mar. Sci.* 2, 36.
- Huang J, Yu H, Guan X, Wang G, Guo R 2016 Accelerated dryland expansion under climate change. *Nat. Clim. Chang.* 6, 166–171. doi: [10.1038/nclimate2837](https://doi.org/10.1038/nclimate2837)
- Hulme P 2004 Invasions, islands and impacts: A Mediterranean perspective, in *Island ecology*, eds. Fernandez Palacios JM, Morici C (Asociación Española de Ecología Terrestre, La Laguna), 337–361.
- Hulme PE 2017 Climate change and biological invasions: evidence, expectations, and response options. *Biol. Rev.* 92, 1297–1313. doi: [10.1111/brv.12282](https://doi.org/10.1111/brv.12282)
- Huntley B 2001 The nature of Mediterranean Europe: an ecological history edited by A. T. Grove and Oliver Rackham, Yale University Press, New Haven, 2001. No. of pages: 384. Price: £45. ISBN 0 300 08443 9. *Earth Surf. Process. Landforms* 26, 908–909. doi: [10.1002/esp.247](https://doi.org/10.1002/esp.247)
- Hurni H, Giger M, Liniger H, Mekdaschi Studer R, Messerli P et al. 2015 Soils, agriculture and food security: the interplay between ecosystem functioning and human well-being. *Curr. Opin. Environ. Sustain.* 15, 25–34. doi: [10.1016/J.COSUST.2015.07.009](https://doi.org/10.1016/J.COSUST.2015.07.009)
- Iakovoglou V, Zaimis GN, Gounaridis D 2013 Riparian areas in urban settings: Two case studies from Greece. *Int. J. Innov. Sustain. Dev.* 7, 271–288. doi: [10.1504/IJISD.2013.056944](https://doi.org/10.1504/IJISD.2013.056944)
- Iglesias A, Garrote L, Flores F, Moneo M 2007 Challenges to Manage the Risk of Water Scarcity and Climate Change in the Mediterranean. *Water Resour. Manag.* 21, 775–788. doi: [10.1007/s11269-006-9111-6](https://doi.org/10.1007/s11269-006-9111-6)
- Iglesias E, Báez K, Díaz-Ambrosio CH 2016 Assessing drought risk in Mediterranean Dehesa grazing lands. *Agric. Syst.* 149, 65–74. doi: [10.1016/J.AGSY.2016.07.017](https://doi.org/10.1016/J.AGSY.2016.07.017)
- Im U, Christensen JH, Geels C, Hansen KM, Brandt J et al. 2018 Influence of anthropogenic emissions and boundary conditions on multi-model simulations of major air pollutants over Europe and North America in the framework of AQMEII3. *Atmos. Chem. Phys.* 18, 8929–8952. doi: [10.5194/acp-18-8929-2018](https://doi.org/10.5194/acp-18-8929-2018)
- Incarbona A, Martrat B, Mortyn PG, Sprovieri M, Ziveri P et al. 2016 Mediterranean circulation perturbations over the last five centuries: Relevance to past Eastern Mediterranean Transient-type events. *Sci. Rep.* 6, 29623. doi: [10.1038/srep29623](https://doi.org/10.1038/srep29623)
- Incarbona A, Ziveri P, Sabatino N, Manta DS, Sprovieri M 2011 Conflicting coccolithophore and geochemical evidence for productivity levels in the Eastern Mediterranean sapropel S1. *Mar. Micropaleontol.* 81, 131–143. doi: [10.1016/j.marmicro.2011.09.003](https://doi.org/10.1016/j.marmicro.2011.09.003)
- Ingrosso G, Bensi M, Cardin V, Giani M 2017 Anthropogenic CO<sub>2</sub> in a dense water formation area of the Mediterranean Sea. *Deep. Res. Part I Oceanogr. Res. Pap.* 123, 118–128. doi: [10.1016/j.dsr.2017.04.004](https://doi.org/10.1016/j.dsr.2017.04.004)
- Ioannides D, Apostolopoulos Y, Sonmez S 2001 *Mediterranean islands and sustainable tourism development: practices management and policies*. London: Continuum Publishers.
- IPBES 2016 Summary for policymakers of the assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production.
- IPBES 2018 *The IPBES regional assessment report on biodiversity and ecosystem services for Africa.*, eds. Archer E, Dziba L, Mulongoy KJ, Maoela MA, Walters M Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services doi: [10.5281/zenodo.3236178](https://doi.org/10.5281/zenodo.3236178)
- IPBES 2019 *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.*, eds. Brondizio ES, Settele J, Díaz S, Ngo HT Bonn, Germany.
- IPCC 2014 *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change.*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press <http://www.citeulike.org/group/15400/article/13497155> [Accessed March 11, 2017]
- IPCC 2019 *Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems.*, eds. Shukla PR, Skea J, Buendia EC, Masson-Delmotte V, Pörtner H-O et al.
- Isnard E, Tournois J, McKenzie DJ, Ferraton F, Bodin N et al. 2015 Getting a good start in life? A comparative analysis of the quality of lagoons as juvenile habitats for the Gilthead Seabream *Sparus aurata* in the Gulf of Lions. *Estuaries and Coasts* 38, 1937–1950. doi: [10.1007/s12237-014-9939-6](https://doi.org/10.1007/s12237-014-9939-6)
- Ison RL 2010 *Systems practice: how to act in a climate change world*. Springer in association with the Open University.
- Ivajnsič D, Lipej L, Škornik I, Kaligarič M 2017 The sea level rise impact on four seashore breeding birds: the key study of Sečovlje Salina Nature Park. *Clim. Change* 140, 549–562. doi: [10.1007/s10584-016-1854-3](https://doi.org/10.1007/s10584-016-1854-3)
- Izquierdo-Muñoz A, Díaz-Valdés M, Ramos-Esplá AA 2009 Recent non-indigenous ascidians in the Mediterranean Sea. *Aquat. Invasions, Spec. issue "Proceedings 2nd Int. Invasive Sea Squirt Conf. (October 2-4, 2007, Prince Edward Island, Canada)* 4, 59–64. doi: [10.3391/ai](https://doi.org/10.3391/ai)

- Izrael Y 1991 Climate change impact studies: The IPCC Working Group II Report. in *Climate change: science, impacts and policy. Proceedings of the second world climate conference*, eds. Jager J, Ferguson HL (Cambridge University Press), 83–86.
- Jacobsen S-E 2014 New Climate-Proof Cropping Systems in Dry Areas of the Mediterranean Region. *J. Agron. Crop Sci.* 200, 399–401. doi: [10.1111/jac.12080](https://doi.org/10.1111/jac.12080)
- Jactel H, Bauhus J, Boberg J, Bonal D, Castagneyrol B et al. 2017 Tree Diversity Drives Forest Stand Resistance to Natural Disturbances. *Curr. For. Reports* 3, 223–243. doi: [10.1007/s40725-017-0064-1](https://doi.org/10.1007/s40725-017-0064-1)
- Jalali B, Sicre M-A, Bassetti M-A, Kallel N 2016 Holocene climate variability in the North-Western Mediterranean Sea (Gulf of Lions). *Clim. Past* 12, 91–101. doi: [10.5194/cp-12-91-2016](https://doi.org/10.5194/cp-12-91-2016)
- Jalali B, Sicre M-A, Kallel N, Azuara J, Combourieu-Nebout N et al. 2017 High-resolution Holocene climate and hydrological variability from two major Mediterranean deltas (Nile and Rhône). *The Holocene* 27, 1158–1168. doi: [10.1177/0959683616683258](https://doi.org/10.1177/0959683616683258)
- Janssen JAM, Rodwell JS, Criado MG, Arts GHP, Bijlsma RJ et al. 2016 *European Red List of Habitats. Part 2. Terrestrial and freshwater habitats*. Luxembourg: Publications Office of the European Union.
- Jemaa S, Bacha M, Khalaf G, Amara R 2015a Evidence for population complexity of the European anchovy (*Engraulis encrasicolus*) along its distributional range. *Fish. Res.* 168, 109–116. doi: [10.1016/j.fishres.2015.04.004](https://doi.org/10.1016/j.fishres.2015.04.004)
- Jemaa S, Bacha M, Khalaf G, Dessailly D, Rabhi K et al. 2015b What can otolith shape analysis tell us about population structure of the European sardine, *Sardina pilchardus*, from Atlantic and Mediterranean waters? *J. Sea Res.* 96, 11–17. doi: [10.1016/j.seares.2014.11.002](https://doi.org/10.1016/j.seares.2014.11.002)
- Jenhani ABR, Fathalli A, Naceur H Ben, Hayouni D, Aouani J et al. 2019 Screening for alien and harmful planktonic species in the Gulf of Gabes (Tunisia, South-eastern Mediterranean Sea). *Reg. Stud. Mar. Sci.* 27, 100526. doi: [10.1016/j.rsma.2019.100526](https://doi.org/10.1016/j.rsma.2019.100526)
- Jiménez-Espejo FJ, Martínez-Ruiz F, Sakamoto T, Iijima K, Gallego-Torres D et al. 2007 Paleoenvironmental changes in the western Mediterranean since the last glacial maximum: High resolution multiproxy record from the Algero–Balearic basin. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 246, 292–306. doi: [10.1016/j.palaeo.2006.10.005](https://doi.org/10.1016/j.palaeo.2006.10.005)
- Jiménez-Ruiz J, Santín-Montanyá MI 2016 An approach to the integrated management of exotic invasive weeds in riparian zones, in *Riparian Zones. Characteristics, Management Practices, and Ecological Impacts*, ed. Pokrovsky OS (Nova Science), 99–124.
- Johnston MW, Purkis SJ 2014 Are lionfish set for a Mediterranean invasion? Modelling explains why this is unlikely to occur. *Mar. Pollut. Bull.* 88, 138–147. doi: [10.1016/j.marpolbul.2014.09.013](https://doi.org/10.1016/j.marpolbul.2014.09.013)
- Jones CG, Lawton JH, Shachak M 1994 Organisms as Ecosystem Engineers. *Oikos* 69, 373. doi: [10.2307/3545850](https://doi.org/10.2307/3545850)
- Jongman RHG, Pungetti G 2004 *Ecological Networks and Greenways Concept, Design, Implementation*. Cambridge University Press. doi: [10.1017/CBO9780511606762](https://doi.org/10.1017/CBO9780511606762)
- Jorda-Capdevila D, Gampe D, Huber García V, Ludwig R, Sabater S et al. 2018 Impact and mitigation of global change on freshwater-related ecosystem services in Southern Europe. *Sci. Total Environ.* 651, 895–908. doi: [10.1016/j.scitotenv.2018.09.228](https://doi.org/10.1016/j.scitotenv.2018.09.228)
- Jordà G, Marbà N, Duarte CM 2012 Mediterranean seagrass vulnerable to regional climate warming. *Nat. Clim. Chang.* 2, 821–824. doi: [10.1038/nclimate1533](https://doi.org/10.1038/nclimate1533)
- Jorissen FJ, Nardelli MP, Almogi-Labin A, Barras C, Bergamin L et al. 2018 Developing Foram-AMBI for biomonitoring in the Mediterranean: Species assignments to ecological categories. *Mar. Micropaleontol.* 140, 33–45. <https://doi.org/10.1016/j.marmicro.2017.12.006>
- Jovan S, Riddell J, Padgett PE, Nash TH 2012 Eutrophic lichens respond to multiple forms of N: implications for critical levels and critical loads research. *Ecol. Appl.* 22, 1910–1922. doi: [10.1890/11-2075.1](https://doi.org/10.1890/11-2075.1)
- Jribi I, Bradai MN 2012 First record of the Lessepsian migrant species *Lagocephalus sceleratus* (Gmelin, 1789) (Actinopterygii: Tetraodontidae) in the Central Mediterranean. *Biol. Invasions Rec.* 1, 49–52. doi: [10.3391/bir.2012.1.1.11](https://doi.org/10.3391/bir.2012.1.1.11)
- Juárez-Escario A, Conesa JA, Solé-Senan XO 2017 Management as a driver of functional patterns and alien species prominence in weed communities of irrigated orchards in Mediterranean areas. *Agric. Ecosyst. Environ.* 249, 247–255. doi: [10.1016/j.agee.2017.07.042](https://doi.org/10.1016/j.agee.2017.07.042)
- Jucker T, Bouriaud O, Avacaritei D, Coomes DA 2014 Stabilizing effects of diversity on aboveground wood production in forest ecosystems: Linking patterns and processes. *Ecol. Lett.* 17, 1560–1569. doi: [10.1111/ele.12382](https://doi.org/10.1111/ele.12382)
- Jump AS, Hunt JM, Peñuelas J 2006 Rapid climate change-related growth decline at the southern range edge of *Fagus sylvatica*. *Glob. Chang. Biol.* 12, 2163–2174. doi: [10.1111/j.1365-2486.2006.01250.x](https://doi.org/10.1111/j.1365-2486.2006.01250.x)
- Jump AS, Mátyás C, Peñuelas J 2009 The altitude-for-latitude disparity in the range retractions of woody species. *Trends Ecol. Evol.* 24, 694–701. doi: [10.1016/j.tree.2009.06.007](https://doi.org/10.1016/j.tree.2009.06.007)
- Jump AS, Peñuelas J 2005 Running to stand still: Adaptation and the response of plants to rapid climate change. *Ecol. Lett.* 8, 1010–1020. doi: [10.1111/j.1461-0248.2005.00796.x](https://doi.org/10.1111/j.1461-0248.2005.00796.x)
- Kadioğullari AI, Başkent EZ 2008 Spatial and temporal dynamics of land use pattern in Eastern Turkey: a case study in Gümüşhane. *Environ. Monit. Assess.* 138, 289–303. doi: [10.1007/s10661-007-9798-8](https://doi.org/10.1007/s10661-007-9798-8)
- Kahiluoto H, Kaseva J, Balek J, Olesen JE, Ruiz-Ramos M et al. 2019 Decline in climate resilience of European wheat. *Proc. Natl. Acad. Sci. U. S. A.* 116, 123–128.



- doi: [10.1073/pnas.1804387115](https://doi.org/10.1073/pnas.1804387115)
- Kahya E, Kalayci S 2004 Trend analysis of streamflow in Turkey. *J. Hydrol.* 289, 128–144. doi: [10.1016/j.jhydrol.2003.11.006](https://doi.org/10.1016/j.jhydrol.2003.11.006)
- Kairis O, Karavitis C, Salvati L, Kounalaki A, Kosmas K 2015 Exploring the Impact of Overgrazing on Soil Erosion and Land Degradation in a Dry Mediterranean Agro-Forest Landscape (Crete, Greece). *Arid L. Res. Manag.* 29, 360–374. doi: [10.1080/15324982.2014.968691](https://doi.org/10.1080/15324982.2014.968691)
- Kallel N, Paterne M, Duplessy J-C, Vergnaud-Grazzini C, Pujol C et al. 1997 Enhanced rainfall in the Mediterranean region during the last sapropel event. *Oceanol. Acta* 20, 697–712. <http://archimer.ifremer.fr/doc/00093/20427/> [Accessed September 25, 2014]
- Kallimanis AS, Bergmeier E, Panitsa M, Georghiou K, Delipetrou P et al. 2010 Biogeographical determinants for total and endemic species richness in a continental archipelago. *Biodivers. Conserv.* 19, 1225–1235. doi: [10.1007/s10531-009-9748-6](https://doi.org/10.1007/s10531-009-9748-6)
- Kambouroglou V, Nicolaidou A 2006 A new alien species in Hellenic waters: *Pseudonereis anomala* (Polychaeta, Nereididae) invades harbors in the Eastern Mediterranean. *Aquat. Invasions* 1, 97–98. doi: [10.3391/ai.2006.1.2.8](https://doi.org/10.3391/ai.2006.1.2.8)
- Kaniewski D, Marriner N, Morhange C, Faivre S, Otto T et al. 2016 Solar pacing of storm surges, coastal flooding and agricultural losses in the Central Mediterranean. *Sci. Rep.* 6, 25197. doi: [10.1038/srep25197](https://doi.org/10.1038/srep25197)
- Kaniewski D, Paulissen E, de Laet V, Dossche K, Waelkens M 2007 A high-resolution Late Holocene landscape ecological history inferred from an intramontane basin in the Western Taurus Mountains, Turkey. *Quat. Sci. Rev.* 26, 2201–2218. doi: [10.1016/j.quascirev.2007.04.015](https://doi.org/10.1016/j.quascirev.2007.04.015)
- Kaniewski D, Van Campo E, Guiot J, Le Burel S, Otto T et al. 2013a Environmental roots of the late bronze age crisis. *PLoS One* 8, e71004. doi: [10.1371/journal.pone.0071004](https://doi.org/10.1371/journal.pone.0071004)
- Kaniewski D, Van Campo E, Morhange C, Guiot J, Zviely D et al. 2013b Early urban impact on Mediterranean coastal environments. *Sci. Rep.* 3, 3540. doi: [10.1038/srep03540](https://doi.org/10.1038/srep03540)
- Kaniewski D, Van Campo E, Morhange C, Guiot J, Zviely D et al. 2014 Vulnerability of mediterranean ecosystems to long-term changes along the coast of Israel. *PLoS One* 9. doi: [10.1371/journal.pone.0102090](https://doi.org/10.1371/journal.pone.0102090)
- Kapsenberg L, Cyronak T 2019 Ocean acidification refugia in variable environments. *Glob. Chang. Biol.* 25, 3201–3214. doi: [10.1111/gcb.14730](https://doi.org/10.1111/gcb.14730)
- Kara MH, Ben Lamine E, Francour P 2015 Range expansion of an invasive pufferfish, *Lagocephalus sceleratus* (Actinopterygii: Tetraodontiformes: Tetraodontidae), to the south-western Mediterranean. *Acta Ichthyol. Piscat.* 45, 103–108. doi: [10.3750/aip2015.45.1.13](https://doi.org/10.3750/aip2015.45.1.13)
- Karavani A, Boer MM, Baudena M, Colinas C, Díaz-Sierra R et al. 2018a Fire-induced deforestation in drought-prone Mediterranean forests: drivers and unknowns from leaves to communities. *Ecol. Monogr.* 88, 141–169. doi: [10.1002/ecm.1285](https://doi.org/10.1002/ecm.1285)
- Karavani A, de Cáceres M, Martínez de Aragón J, Bonet JA, de Miguel Magaña S 2018b Effect of climatic and soil moisture conditions on mushroom productivity and related ecosystem services in Mediterranean pine stands facing climate change. *Agric. For. Meteorol.* 248, 432–440. doi: [10.1016/j.agrformet.2017.10.024](https://doi.org/10.1016/j.agrformet.2017.10.024)
- Kark S, Levin N, Grantham HS, Possingham HP 2009 Between-country collaboration and consideration of costs increase conservation planning efficiency in the Mediterranean Basin. *Proc. Natl. Acad. Sci. U. S. A.* 106, 15368–15373. doi: [10.1073/pnas.0901001106](https://doi.org/10.1073/pnas.0901001106)
- Katona K, Coetsee C 2019 Impacts of Browsing and Grazing Ungulates on Faunal Biodiversity, in (Springer, Cham), 277–300. doi: [10.1007/978-3-030-25865-8\\_12](https://doi.org/10.1007/978-3-030-25865-8_12)
- Katsanevakis S, Coll M, Piroddi C, Steenbeek J, Ben Rais Lasram F et al. 2014a Invading the Mediterranean Sea: biodiversity patterns shaped by human activities. *Front. Mar. Sci.* 1. doi: [10.3389/fmars.2014.00032](https://doi.org/10.3389/fmars.2014.00032)
- Katsanevakis S, Wallentinus I, Zenetos A, Leppäkoski E, Çınar ME et al. 2014b Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Aquat. Invasions* 9, 391–423. doi: [10.3391/ai.2014.9.4.01](https://doi.org/10.3391/ai.2014.9.4.01)
- Katsanevakis S, Zenetos A, Belchior C, Cardoso AC 2013 Invading European Seas: Assessing pathways of introduction of marine aliens. *Ocean Coast. Manag.* 76, 64–74. doi: [10.1016/j.ocecoaman.2013.02.024](https://doi.org/10.1016/j.ocecoaman.2013.02.024)
- Katselidis KA, Schofield G, Stamou G, Dimopoulos P, Pantis JD 2014 Employing sea-level rise scenarios to strategically select sea turtle nesting habitat important for long-term management at a temperate breeding area. *J. Exp. Mar. Bio. Ecol.* 450, 47–54. doi: [10.1016/j.jembe.2013.10.017](https://doi.org/10.1016/j.jembe.2013.10.017)
- Kauserud H, Heegaard E, Büntgen U, Halvorsen R, Egli S et al. 2012 Warming-induced shift in European mushroom fruiting phenology. *Proc. Natl. Acad. Sci. U. S. A.* 109, 14488–14493. doi: [10.1073/pnas.1200789109](https://doi.org/10.1073/pnas.1200789109)
- Kawasaki T 1991 Effects of global change on marine ecosystem and fisheries, in *Climate changes science, impact and policy*, eds. Jager J, Fergusson HL (Cambridge, UK: Cambridge University Press), 291–299.
- Kazakis G, Ghosn D, Vogiatzakis IN, Papanastasis VP 2007 Vascular plant diversity and climate change in the alpine zone of the Lefka Ori, Crete. *Biodivers. Conserv.* 16, 1603–1615. doi: [10.1007/s10531-006-9021-1](https://doi.org/10.1007/s10531-006-9021-1)
- Keeley ATH, Ackerly DD, Cameron DR, Heller NE, Huber PR et al. 2018 New concepts, models, and assessments of climate-wise connectivity. *Environ. Res. Lett.* 13, 073002. doi: [10.1088/1748-9326/aacb85](https://doi.org/10.1088/1748-9326/aacb85)
- Keenan RJ, Reams GA, Achard F, de Freitas J V., Grainger A et al. 2015 Dynamics of global forest area: Results from the FAO Global Forest Resources Assessment 2015. *For. Ecol. Manage.* 352, 9–20. doi: [10.1016/j.foreco.2015.06.014](https://doi.org/10.1016/j.foreco.2015.06.014)

- Keenan TF, Maria Serra J, Lloret F, Ninyerola M, Sabate S 2011 Predicting the future of forests in the Mediterranean under climate change, with niche- and process-based models: CO<sub>2</sub> matters! *Glob. Chang. Biol.* 17, 565–579. doi: [10.1111/j.1365-2486.2010.02254.x](https://doi.org/10.1111/j.1365-2486.2010.02254.x)
- Kennedy CM, Lonsdorf E, Neel MC, Williams NM, Ricketts TH et al. 2013 A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecol. Lett.* 16, 584–599. doi: [10.1111/ele.12082](https://doi.org/10.1111/ele.12082)
- Khedhri I, Atoui A, Ibrahim M, Afli A, Aleya L 2016 Assessment of surface sediment dynamics and response of benthic macrofauna assemblages in Boughrara Lagoon (SW Mediterranean Sea). *Ecol. Indic.* 70, 77–88. doi: [10.1016/j.ecolind.2016.06.011](https://doi.org/10.1016/j.ecolind.2016.06.011)
- Kiernan JD, Moyle PB, Crain PK 2012 Restoring native fish assemblages to a regulated California stream using the natural flow regime concept. *Ecol. Appl.* 22, 1472–1482. doi: [10.1890/11-0480.1](https://doi.org/10.1890/11-0480.1)
- Kingsford MJ, Becken S, Bordehore C, Fuentes VL, Pitt KA et al. 2018 Empowering stakeholders to manage stinging jellyfish: A perspective. *Coast. Manag.* 46, 1–18. doi: [10.1080/08920753.2018.1405326](https://doi.org/10.1080/08920753.2018.1405326)
- Kingsford RT, Thomas RF 2004 Destruction of Wetlands and Waterbird Populations by Dams and Irrigation on the Murrumbidgee River in Arid Australia. *Environ. Manage.* 34, 383–396. doi: [10.1007/s00267-004-0250-3](https://doi.org/10.1007/s00267-004-0250-3)
- Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA et al. 2006 Importance of pollinators in changing landscapes for world crops. *Proc. R. Soc. B Biol. Sci.* 274, 303–313. doi: [10.1098/rspb.2006.3721](https://doi.org/10.1098/rspb.2006.3721)
- Klein T, Cahanovitch R, Sprintsin M, Herr N, Schiller G 2019 A nation-wide analysis of tree mortality under climate change: Forest loss and its causes in Israel 1948–2017. *For. Ecol. Manage.* 432, 840–849. doi: [10.1016/j.foreco.2018.10.020](https://doi.org/10.1016/j.foreco.2018.10.020)
- Kleitou DC, Hall-Spencer JM, Kleitou P 2016 A lionfish (*Pterois miles*) invasion has begun in the Mediterranean Sea. *Mar. Biodivers. Rec.* 9. doi: [10.1186/s41200-016-0065-y](https://doi.org/10.1186/s41200-016-0065-y)
- Klippel L, Krusic PJ, Brandes R, Hartl C, Belmecheri S et al. 2018 A 1286-year hydro-climate reconstruction for the Balkan Peninsula. *Boreas* 47, 1218–1229. doi: [10.1111/bor.12320](https://doi.org/10.1111/bor.12320)
- Kondolf GM 1997 PROFILE: Hungry Water: Effects of Dams and Gravel Mining on River Channels. *Environ. Manage.* 21, 533–551. doi: [10.1007/s002679900048](https://doi.org/10.1007/s002679900048)
- Konter O, Krusic PJ, Trouet V, Esper J 2017 Meet Adonis, Europe's oldest dendrochronologically dated tree. *Dendrochronologia* 42, 12. doi: [10.1016/j.dendro.2016.12.001](https://doi.org/10.1016/j.dendro.2016.12.001)
- Kontsiotis V, Zaimis GN, Tsiptsis S, Kiourtziadis P, Bakaloudis D 2019 Assessing the influence of riparian vegetation structure on bird communities in agricultural Mediterranean landscapes. *Agrofor. Syst.* 93, 675–687. doi: [10.1007/s10457-017-0162-x](https://doi.org/10.1007/s10457-017-0162-x)
- Koutroulis AG 2019 Dryland changes under different levels of global warming. *Sci. Total Environ.* 655, 482–511. doi: [10.1016/J.SCITOTENV.2018.11.215](https://doi.org/10.1016/J.SCITOTENV.2018.11.215)
- Koutroulis AG, Papadimitriou L V., Grillakis MG, Tsanis IK, Warren R et al. 2019 Global water availability under high-end climate change: A vulnerability based assessment. *Glob. Planet. Change* 175, 52–63. doi: [10.1016/J.GLOPLACHA.2019.01.013](https://doi.org/10.1016/J.GLOPLACHA.2019.01.013)
- Koutroulis AG, Tsanis IK, Daliakopoulos IN 2010 Seasonality of floods and their hydrometeorologic characteristics in the island of Crete. *J. Hydrol.* 394, 90–100. doi: [10.1016/J.JHYDROL.2010.04.025](https://doi.org/10.1016/J.JHYDROL.2010.04.025)
- Koutroulis AG, Tsanis IK, Daliakopoulos IN, Jacob D 2013 Impact of climate change on water resources status: A case study for Crete Island, Greece. *J. Hydrol.* 479, 146–158. doi: [10.1016/j.jhydrol.2012.11.055](https://doi.org/10.1016/j.jhydrol.2012.11.055)
- Kramer K, Leinonen I, Loustau D 2000 The importance of phenology for the evaluation of impact of climate change on growth of boreal, temperate and Mediterranean forests ecosystems: An overview. *Int. J. Biometeorol.* 44, 67–75. doi: [10.1007/s004840000066](https://doi.org/10.1007/s004840000066)
- Krause A, Pugh TAM, Bayer AD, Arneith A 2016 Impacts of land-use history on the recovery of ecosystems after agricultural abandonment. *Earth Syst. Dyn.* 7, 745–766. doi: [10.5194/esd-7-745-2016](https://doi.org/10.5194/esd-7-745-2016)
- Krchnak KM, Smith DM, Deutz A 2011 Putting Nature in the Nexus: Investing in Natural Infrastructure to Advance Water-Energy- Food Security. in *Bonn2011 Conference: The Water, Energy and Food Security Nexus – Solutions for the Green Economy Background Papers for the Stakeholder Engagement Process.*
- Kroeker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L et al. 2013 Impacts of ocean acidification on marine organisms: Quantifying sensitivities and interaction with warming. *Glob. Chang. Biol.* 19, 1884–1896. doi: [10.1111/gcb.12179](https://doi.org/10.1111/gcb.12179)
- Krupa S V. 2003 Effects of atmospheric ammonia (NH<sub>3</sub>) on terrestrial vegetation: A review. *Environ. Pollut.* 124, 179–221. doi: [10.1016/S0269-7491\(02\)00434-7](https://doi.org/10.1016/S0269-7491(02)00434-7)
- Kuhnt T, Schmiedl G, Ehrmann W, Hamann Y, Andersen N 2008 Stable isotopic composition of Holocene benthic foraminifers from the Eastern Mediterranean Sea: Past changes in productivity and deep water oxygenation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 268, 106–115. doi: [10.1016/j.palaeo.2008.07.010](https://doi.org/10.1016/j.palaeo.2008.07.010)
- Kummu M, Fader M, Gerten D, Guillaume JHA, Jalava M et al. 2017 Bringing it all together: linking measures to secure nations' food supply. *Curr. Opin. Environ. Sustain.* 29, 98–117. doi: [10.1016/j.cosust.2018.01.006](https://doi.org/10.1016/j.cosust.2018.01.006)
- Kushmaro A, Rosenberg E, Fine M, Haim YB, Loya Y 1998 Effect of temperature on bleaching of the coral *Oculina patagonica* by *Vibrio* AK-1. *Mar. Ecol. Prog. Ser.* 171, 131–137.
- Kuvan Y 2010 Mass tourism development and deforestation in Turkey. *Anatolia* 21, 155–168. doi: [10.1080/13032917.2010.9687096](https://doi.org/10.1080/13032917.2010.9687096)
- La Jeunesse I, Cirelli C, Aubin D, Larrue C, Sellami H et al. 2016 Is climate change a threat for water uses in the

- Mediterranean region? Results from a survey at local scale. *Sci. Total Environ.* 543, 981–996. doi: [10.1016/j.scitotenv.2015.04.062](https://doi.org/10.1016/j.scitotenv.2015.04.062)
- La Jeunesse I, Cirelli C, Sellami H, Aubin D, Deidda R et al. 2015 Is the governance of the Thau coastal lagoon ready to face climate change impacts? *Ocean Coast. Manag.* 118, 234–246. doi: [10.1016/j.ocecoaman.2015.05.014](https://doi.org/10.1016/j.ocecoaman.2015.05.014)
- Laanisto L, Niinemets Ü 2015 Polytolerance to abiotic stresses: How universal is the shade-drought tolerance trade-off in woody species? *Glob. Ecol. Biogeogr.* 24, 571–580. doi: [10.1111/geb.12288](https://doi.org/10.1111/geb.12288)
- Lacetera N 2019 Impact of climate change on animal health and welfare. *Anim. Front.* 9, 26–31. doi: [10.1093/af/vfy030](https://doi.org/10.1093/af/vfy030)
- Lacoue-Labarthe T, Nunes PALD, Ziveri P, Cinar M, Gazeau F et al. 2016 Impacts of ocean acidification in a warming Mediterranean Sea: An overview. *Reg. Stud. Mar. Sci.* 5, 1–11. doi: [10.1016/j.rsma.2015.12.005](https://doi.org/10.1016/j.rsma.2015.12.005)
- Laforteza R, Tanentzap AJ, Elia M, John R, Sanesi G et al. 2015 Prioritizing fuel management in urban interfaces threatened by wildfires. *Ecol. Indic.* 48, 342–347. doi: [10.1016/j.ecolind.2014.08.034](https://doi.org/10.1016/j.ecolind.2014.08.034)
- Lake PS 2003 Ecological effects of perturbation by drought in flowing waters. *Freshw. Biol.* 48, 1161–1172. doi: [10.1046/j.1365-2427.2003.01086.x](https://doi.org/10.1046/j.1365-2427.2003.01086.x)
- Lakkis S, Novel-Lakkis V 2007 Diversity and distribution of macrophytes along the coast of Lebanon (Levantine Basin, Eastern Mediterranean). *Rapp. la Comm. Int. pour l'Exploration Sci. la Mer Méditerranée* 38, 526.
- Lambeck K, Woodroffe CD, Antonioli F, Anzidei M, Gehrels WR et al. 2010 Paleoenvironmental Records, Geophysical Modeling, and Reconstruction of Sea-Level Trends and Variability on Centennial and Longer Timescales, in *Understanding Sea-Level Rise and Variability* (John Wiley & Sons, Ltd), 61–121. <https://onlinelibrary.wiley.com/doi/abs/10.1002/9781444323276.ch4>
- Langer G, Nehrke G, Probert I, Ly J, Ziveri P 2009 Strain-specific responses of *Emiliania huxleyi* to changing seawater carbonate chemistry. *Biogeosciences* 6, 2637–2646. doi: [10.5194/bg-6-2637-2009](https://doi.org/10.5194/bg-6-2637-2009)
- Langridge P 2019 Innovation in Breeding and Biotechnology, in *Agriculture and Food systems*, eds. Serraj R, Pingali P (Singapore: World Scientific), 245–284. doi: [10.1142/9789813278356\\_0008](https://doi.org/10.1142/9789813278356_0008)
- Laran S, Pettex E, Authier M, Blanck A, David L et al. 2017 Seasonal distribution and abundance of cetaceans within French waters- Part I: The North-Western Mediterranean, including the Pelagos sanctuary. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 141, 20–30. doi: [10.1016/j.dsr2.2016.12.011](https://doi.org/10.1016/j.dsr2.2016.12.011)
- Lasanta-Martínez T, Vicente-Serrano SM, Cuadrat-Prats JM 2005 Mountain Mediterranean landscape evolution caused by the abandonment of traditional primary activities: a study of the Spanish Central Pyrenees. *Appl. Geogr.* 25, 47–65. doi: [10.1016/j.apgeog.2004.11.001](https://doi.org/10.1016/j.apgeog.2004.11.001)
- Lasanta T, Arnáez J, Pascual N, Ruiz-Flaño P, Errea MP et al. 2017 Space-time process and drivers of land abandonment in Europe. *Catena* 149, 810–823. doi: [10.1016/j.catena.2016.02.024](https://doi.org/10.1016/j.catena.2016.02.024)
- Lasanta T, Nadal-Romero E, Arnáez J 2015 Managing abandoned farmland to control the impact of re-vegetation on the environment. The state of the art in Europe. *Environ. Sci. Policy* 52, 99–109. doi: [10.1016/J.ENVSCL.2015.05.012](https://doi.org/10.1016/J.ENVSCL.2015.05.012)
- Lasker HR, Coffroth MA 1999 Responses of clonal reef taxa to environmental change. *Am. Zool.* 39, 92–103. doi: [10.1093/icb/39.1.92](https://doi.org/10.1093/icb/39.1.92)
- Lastrucci L, Lazzaro L, Dell'Olmo L, Foggi B, Cianferoni F 2018 Impacts of *Myriophyllum aquaticum* invasion in a Mediterranean wetland on plant and macro-arthropod communities. *Plant Biosyst.* 152, 427–435. doi: [10.1080/11263504.2017.1303002](https://doi.org/10.1080/11263504.2017.1303002)
- Lavini A, Pulvento C, D'Andria R, Riccardi M, Choukr-Allah R et al. 2014 Quinoa's Potential in the Mediterranean Region. *J. Agron. Crop Sci.* 200, 344–360. doi: [10.1111/jac.12069](https://doi.org/10.1111/jac.12069)
- Lavorel S, Touzard B, Lebreton JD, Clément B 1998 Identifying functional groups for response to disturbance in an abandoned pasture. *Acta Oecologica* 19, 227–240. doi: [10.1016/S1146-609X\(98\)80027-1](https://doi.org/10.1016/S1146-609X(98)80027-1)
- Lawler JJ, Olden JD 2011 Reframing the debate over assisted colonization. *Front. Ecol. Environ.* 9, 569–574. doi: [10.1890/100106](https://doi.org/10.1890/100106)
- Lawrence JE, Lunde KB, Mazor RD, Bêche LA, McElravy EP et al. 2010 Long-term macroinvertebrate responses to climate change: Implications for biological assessment in mediterranean-climate streams. *J. North Am. Benthol. Soc.* 29, 1424–1440. doi: [10.1899/09-178.1](https://doi.org/10.1899/09-178.1)
- Lazar B, Gračan R 2011 Ingestion of marine debris by loggerhead sea turtles, *Caretta caretta*, in the Adriatic Sea. *Mar. Pollut. Bull.* 62, 43–47. doi: [10.1016/j.marpolbul.2010.09.013](https://doi.org/10.1016/j.marpolbul.2010.09.013)
- Lazzari P, Mattia G, Solidoro C, Salon S, Crise A et al. 2014 The impacts of climate change and environmental management policies on the trophic regimes in the Mediterranean Sea: Scenario analyses. *J. Mar. Syst.* 135, 137–149. doi: [10.1016/j.jmarsys.2013.06.005](https://doi.org/10.1016/j.jmarsys.2013.06.005)
- Le Houérou HN 1990 Agroforestry and sylvopastoralism to combat land degradation in the Mediterranean Basin: old approaches to new problems. *Agric. Ecosyst. Environ.* 33, 99–109. doi: [10.1016/0167-8809\(90\)90236-7](https://doi.org/10.1016/0167-8809(90)90236-7)
- Le Pape O, Modéran J, Beaunée G, Riera P, Nicolas D et al. 2013 Sources of organic matter for flatfish juveniles in coastal and estuarine nursery grounds: A meta-analysis for the common sole (*Solea solea*) in contrasted systems of Western Europe. *J. Sea Res.* 75, 85–95. doi: [10.1016/j.seares.2012.05.003](https://doi.org/10.1016/j.seares.2012.05.003)
- Lefèvre F, Fady B 2016 Introduction to Mediterranean Forest Systems: Mediterranean Basin, in *Insects and Diseases of Mediterranean Forest Systems.*, eds. Paine TD, Lieutier F (Springer International Publishing Switzerland), 7–28. doi: [10.1007/978-3-319-24744-1](https://doi.org/10.1007/978-3-319-24744-1)
- Legrand C, Casotti R 2009 Climate-induced changes and

- Harmful Algal Blooms in the Mediterranean: perspectives on future scenarios. in *CIESM Workshop Monographs n°40, Phytoplankton Response to Mediterranean Environmental Change*, 63–66.
- Lejeusne C, Chevaldonné P, Pergent-Martini C, Boudouresque C-F, Pérez T 2010 Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. *Trends Ecol. Evol.* 25, 250–260. doi: [10.1016/j.tree.2009.10.009](https://doi.org/10.1016/j.tree.2009.10.009)
- Lelieveld J, Hadjinicolaou P, Kostopoulou E, Chenoweth J, El Maayar M et al. 2012 Climate change and impacts in the Eastern Mediterranean and the Middle East. *Clim. Change* 114, 667–687. doi: [10.1007/s10584-012-0418-4](https://doi.org/10.1007/s10584-012-0418-4)
- Lelièvre F, Seddaiu G, Ledda L, Porqueddu C, Volaire F 2011 Water use efficiency and drought survival in Mediterranean perennial forage grasses. *F. Crop. Res.* 121, 333–342. doi: [10.1016/J.FCR.2010.12.023](https://doi.org/10.1016/J.FCR.2010.12.023)
- Lenoir J, Gégout JC, Marquet PA, de Ruffray P, Brisse H 2008 A significant upward shift in plant species optimum elevation during the 20th century. *Science (80-. )*. 320, 1768–71. doi: [10.1126/science.1156831](https://doi.org/10.1126/science.1156831)
- Levers C, Butsic V, Verburg PH, Müller D, Kuemmerle T 2016 Drivers of changes in agricultural intensity in Europe. *Land use policy* 58, 380–393. doi: [10.1016/j.landusepol.2016.08.013](https://doi.org/10.1016/j.landusepol.2016.08.013)
- Levinsky I, Skov F, Svenning JC, Rahbek C 2007 Potential impacts of climate change on the distributions and diversity patterns of European mammals. *Biodivers. Conserv.* 16, 3803–3816. doi: [10.1007/s10531-007-9181-7](https://doi.org/10.1007/s10531-007-9181-7)
- Leydet KP, Grupstra CGB, Coma R, Ribes M, Hellberg ME 2018 Host-targeted RAD-Seq reveals genetic changes in the coral *Oculina patagonica* associated with range expansion along the Spanish Mediterranean coast. *Mol. Ecol.* 27, 2529–2543. doi: [10.1111/mec.14702](https://doi.org/10.1111/mec.14702)
- Li W, Li X, Zhao Y, Zheng S, Bai Y et al. 2018 Ecosystem structure, functioning and stability under climate change and grazing in grasslands: current status and future prospects This review comes from a themed issue on System dynamics and sustainability. *Curr. Opin. Environ. Sustain.* 33, 124–135. doi: [10.1016/j.cosust.2018.05.008](https://doi.org/10.1016/j.cosust.2018.05.008)
- Licandro P, Conway DVP, Daly Yahia MN, Fernández de Puelles ML, Gasparini S et al. 2010 A blooming jellyfish in the northeast Atlantic and Mediterranean. *Biol. Lett.* 6, 688–691. doi: [10.1098/rsbl.2010.0150](https://doi.org/10.1098/rsbl.2010.0150)
- Liébault F, Piégay H 2001 Assessment of channel changes due to long-term bedload supply decrease, Roubion River, France. *Geomorphology* 36, 167–186. doi: [10.1016/S0169-555X\(00\)00044-1](https://doi.org/10.1016/S0169-555X(00)00044-1)
- Light T, Marchetti MP 2007 Distinguishing between invasions and habitat changes as drivers of diversity loss among California's freshwater fishes. *Conserv. Biol.* 21, 434–446. doi: [10.1111/j.1523-1739.2006.00643.x](https://doi.org/10.1111/j.1523-1739.2006.00643.x)
- Linares C, Coma R, Díaz D, Zabala M, Hereu B et al. 2005 Immediate and delayed effects of a mass mortality event on gorgonian population dynamics and benthic community structure in the NW Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 305, 127–137. doi: [10.3354/meps305127](https://doi.org/10.3354/meps305127)
- Linares C, Coma R, Zabala M 2008 Effects of a mass mortality event on gorgonian reproduction. *Coral Reefs* 27, 27–34. doi: [10.1007/s00338-007-0285-z](https://doi.org/10.1007/s00338-007-0285-z)
- Linares JC, Camarero JJ, Carreira JA 2010 Competition modulates the adaptation capacity of forests to climatic stress: insights from recent growth decline and death in relict stands of the Mediterranean fir *Abies pinsapo*. *J. Ecol.* 98, 592–603. doi: [10.1111/j.1365-2745.2010.01645.x](https://doi.org/10.1111/j.1365-2745.2010.01645.x)
- Linares JC, Delgado-Huertas A, Carreira JA 2011 Climatic trends and different drought adaptive capacity and vulnerability in a mixed *Abies pinsapo*-*Pinus halepensis* forest. *Clim. Change* 105, 67–90. doi: [10.1007/s10584-010-9878-6](https://doi.org/10.1007/s10584-010-9878-6)
- Lindner M, Calama R 2013 Climate change and the need for adaptation in Mediterranean forests., in *Forest management of Mediterranean forests under the new context of climate change. Building alternative for the coming future.*, ed. Lucas Borja M. (New York: Nova Science Publishers, Inc.), 13–28.
- Lindner M, Maroschek M, Netherer S, Kremer A, Barbati A et al. 2010 Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *For. Ecol. Manage.* 259, 698–709. doi: [10.1016/j.foreco.2009.09.023](https://doi.org/10.1016/j.foreco.2009.09.023)
- Lipkin Y 1975 *Halophila stipulacea*, a review of a successful immigration. *Aquat. Bot.* 1, 203–215. doi: [10.1016/0304-3770\(75\)90023-6](https://doi.org/10.1016/0304-3770(75)90023-6)
- Lloret F, de la Riva EG, Pérez-Ramos IM, Marañón T, Saura-Mas S et al. 2016 Climatic events inducing die-off in Mediterranean shrublands: are species' responses related to their functional traits? *Oecologia* 180, 961–973. doi: [10.1007/s00442-016-3550-4](https://doi.org/10.1007/s00442-016-3550-4)
- Lloret F, Martínez-Vilalta J, Serra-Díaz JM, Ninyerola M 2013 Relationship between projected changes in future climatic suitability and demographic and functional traits of forest tree species in Spain. *Clim. Change* 120, 449–462. doi: [10.1007/s10584-013-0820-6](https://doi.org/10.1007/s10584-013-0820-6)
- Lloret F, Siscart D, Dalmases C 2004 Canopy recovery after drought dieback in holm-oak Mediterranean forests of Catalonia (NE Spain). *Glob. Chang. Biol.* 10, 2092–2099. doi: [10.1111/j.1365-2486.2004.00870.x](https://doi.org/10.1111/j.1365-2486.2004.00870.x)
- Lloret J, Demestre M, Casadevall M, Muñoz M 2008 Towards new approaches to fisheries management in the Mediterranean Sea, in *Fisheries: Management, Economics and perspectives*, eds. McManys NF, Bellinghouse DS (Nova Science Publishers), 93–125.
- Lo Y-H, Blanco JA, González de Andrés E, Imbert JB, Castillo FJ 2019 CO<sub>2</sub> fertilization plays a minor role in long-term carbon accumulation patterns in temperate pine forests in the southwestern Pyrenees. *Ecol. Modell.* 407, 108737. doi: [10.1016/j.ecolmodel.2019.108737](https://doi.org/10.1016/j.ecolmodel.2019.108737)
- Lobera G, Batalla RJ, Vericat D, López-Tarazón JA, Tena A

- 2016 Sediment transport in two mediterranean regulated rivers. *Sci. Total Environ.* 540, 101–113. doi: [10.1016/j.scitotenv.2015.08.018](https://doi.org/10.1016/j.scitotenv.2015.08.018)
- Loepfe L, Martinez-Vilalta J, Piñol J 2012 Management alternatives to offset climate change effects on Mediterranean fire regimes in NE Spain. *Clim. Change* 115, 693–707. doi: [10.1007/s10584-012-0488-3](https://doi.org/10.1007/s10584-012-0488-3)
- Loizidou XI, Loizides MI, Orthodoxou DL 2018 Persistent marine litter: small plastics and cigarette butts remain on beaches after organized beach cleanups. *Environ. Monit. Assess.* 190, 1–10. doi: [10.1007/s10661-018-6798-9](https://doi.org/10.1007/s10661-018-6798-9)
- Longobardi L, Bavestrello G, Betti F, Cattaneo-Vietti R 2017 Long-term changes in a Ligurian infralittoral community (Mediterranean Sea): A warning signal? *Reg. Stud. Mar. Sci.* 14, 15–26. doi: [10.1016/j.rsma.2017.03.011](https://doi.org/10.1016/j.rsma.2017.03.011)
- López-Moreno JI 2005 Recent variations of snowpack depth in the central Spanish Pyrenees. *Arctic, Antarct. Alp. Res.* 37, 253–260. doi: [10.1657/1523-0430\(2005\)037\[0253:RVOSDI\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0253:RVOSDI]2.0.CO;2)
- López-Moreno JI, Beniston M, García-Ruiz JM 2008 Environmental change and water management in the Pyrenees: Facts and future perspectives for Mediterranean mountains. *Glob. Planet. Change* 61, 300–312. doi: [10.1016/j.gloplacha.2007.10.004](https://doi.org/10.1016/j.gloplacha.2007.10.004)
- López-Moreno JI, García-Ruiz JM 2004 Influence of snow accumulation and snowmelt on streamflow in the central Spanish Pyrenees. *Hydrol. Sci. J.* 49, 787–802. doi: [10.1623/hysj.49.5.787.55135](https://doi.org/10.1623/hysj.49.5.787.55135)
- López-Poma R, Orr BJ, Bautista S 2014 Successional stage after land abandonment modulates fire severity and post-fire recovery in a Mediterranean mountain landscape. *Int. J. Wildl. Fire* 23, 1005. doi: [10.1071/WF13150](https://doi.org/10.1071/WF13150)
- López-Tarazón JA, Batalla RJ, Vericat D, Balasch JC 2010 Rainfall, runoff and sediment transport relations in a mesoscale mountainous catchment: The River Isábena (Ebro basin). *Catena* 82, 23–34. doi: [10.1016/j.catena.2010.04.005](https://doi.org/10.1016/j.catena.2010.04.005)
- Lowder SK, Skoet J, Raney T 2016 The Number, Size, and Distribution of Farms, Smallholder Farms, and Family Farms Worldwide q. *World Dev.* 87, 16–29. doi: [10.1016/j.worlddev.2015.10.041](https://doi.org/10.1016/j.worlddev.2015.10.041)
- Luan Y, Cui X, Ferrat M 2013 Historical trends of food self-sufficiency in Africa. *Food Secur.* 5, 393–405. doi: [10.1007/s12571-013-0260-1](https://doi.org/10.1007/s12571-013-0260-1)
- Lucas CH, Gelicich S, Uye S-I 2014 Living with Jellyfish: Management and Adaptation Strategies, in *Jellyfish Blooms*, eds. Pitt K, Lucas C (Dordrecht, Netherlands: Springer), 129–150. doi: [10.1007/978-94-007-7015-7\\_6](https://doi.org/10.1007/978-94-007-7015-7_6)
- Luna GM, Bianchelli S, Decembrini F, de Domenico E, Danovaro R et al. 2012 The dark portion of the Mediterranean Sea is a bioreactor of organic matter cycling. *Global Biogeochem. Cycles* 26. doi: [10.1029/2011GB004168](https://doi.org/10.1029/2011GB004168)
- Luterbacher J, García-Herrera R, Akcer-On S, Allan R, Alvarez-Castro M-C et al. 2012 A review of 2000 years of paleoclimatic evidence in the Mediterranean, in *The climate of the Mediterranean region*, ed. Lionello P (Oxford: Elsevier), 87–185. <http://www.sciencedirect.com/science/article/pii/B9780124160422000021>
- Lytle DA, Poff NL 2004 Adaptation to natural flow regimes. *Trends Ecol. Evol.* 19, 94–100. doi: [10.1016/j.tree.2003.10.002](https://doi.org/10.1016/j.tree.2003.10.002)
- Maas GS, Macklin MG 2002 The impact of recent climate change on flooding and sediment supply within a Mediterranean mountain catchment, southwestern Crete, Greece. *Earth Surf. Process. Landforms* 27, 1087–1105. doi: [10.1002/esp.398](https://doi.org/10.1002/esp.398)
- Mabrouk L, Hamza A, Mahfoudhi M, Bradai MN 2012 Spatial and temporal variations of epiphytic *Ostreopsis siamensis* on *Posidonia oceanica* (L.) Delile leaves in Mahdia (Tunisia). *Cah. Biol. Mar.* 53, 419–427. <http://www.ramoge.org/fr/Ostreopsis/201-Spatial-and-temporal-variations-of-epiphytic-Ostreopsis-siamensis-on-Posidonia-oceanica-L-Delile-leaves-in-Mahdia-Tunisia.html> [Accessed September 10, 2019]
- Macías DM, García-Gorriç E, Stips A 2015 Productivity changes in the Mediterranean Sea for the twenty first century in response to changes in the regional atmospheric forcings. *Front. Mar. Sci.* 2. doi: [10.3389/fmars.2015.00079](https://doi.org/10.3389/fmars.2015.00079)
- Macías DM, García-Gorriç E, Piroddi C, Stips A 2014 Biogeochemical control of marine productivity in the Mediterranean Sea during the last 50 years. *Global Biogeochem. Cycles* 28, 897–907. doi: [10.1002/2014GB004846](https://doi.org/10.1002/2014GB004846)
- Mack L, Andersen HE, Beklioglu M, Bucak T, Couture R-M et al. 2019 The future depends on what we do today – Projecting Europe’s surface water quality into three different future scenarios. *Sci. Total Environ.* 668, 470–484. doi: [10.1016/j.scitotenv.2019.02.251](https://doi.org/10.1016/j.scitotenv.2019.02.251)
- Madlung A, Comai L 2004 The effect of stress on genome regulation and structure. *Ann. Bot.* 94, 481–495. doi: [10.1093/aob/mch172](https://doi.org/10.1093/aob/mch172)
- Maes J, Teller A, Erhard M, Grizzetti B, Barredo JI et al. 2018 Mapping and Assessment of Ecosystems and their Services An analytical framework for mapping and assessment of ecosystem condition in EU. Publications office of the European Union, Luxembourg. doi: [10.2779/055584](https://doi.org/10.2779/055584)
- Maestre FT, Callaway RM, Valladares F, Lortie CJ 2009 Refining the stress-gradient hypothesis for competition and facilitation in plant communities. *J. Ecol.* 97, 199–205. doi: [10.1111/j.1365-2745.2008.01476.x](https://doi.org/10.1111/j.1365-2745.2008.01476.x)
- Maestre FT, Cortina J 2003 Small-scale spatial variation in soil CO<sub>2</sub> efflux in a Mediterranean semiarid steppe. *Appl. Soil Ecol.* 23, 199–209. doi: [10.1016/s0929-1393\(03\)00050-7](https://doi.org/10.1016/s0929-1393(03)00050-7)
- Maestre FT, Cortina J 2005 Remnant shrubs in Mediterranean semi-arid steppes: effects of shrub size, abiotic factors and species identity on understorey richness and occurrence. *Acta Oecologica* 27, 161–169. doi: [10.1016/j.actao.2004.11.003](https://doi.org/10.1016/j.actao.2004.11.003)

- Magalhães MF, Beja P, Schlosser IJ, Collares-Pereira MJ 2007 Effects of multi-year droughts on fish assemblages of seasonally drying Mediterranean streams. *Freshw. Biol.* 52, 1494–1510. doi: [10.1111/j.1365-2427.2007.01781.x](https://doi.org/10.1111/j.1365-2427.2007.01781.x)
- Magdaleno F, Martinez R 2014 Evaluating the quality of riparian forest vegetation: the Riparian Forest Evaluation (RFV) index. *For. Syst.* 23, 259–272. doi: [10.5424/fs/2014232-04488](https://doi.org/10.5424/fs/2014232-04488)
- Maggi E, Bertocci I, Vaselli S, Benedetti-Cecchi L 2009 Effects of changes in number, identity and abundance of habitat-forming species on assemblages of rocky seashores. *Mar. Ecol. Prog. Ser.* 381, 39–49. doi: [10.3354/meps07949](https://doi.org/10.3354/meps07949)
- Maggio A, de Pascale S, Fagnano M, Barbieri G 2011 Saline agriculture in Mediterranean environments. *Ital. J. Agron.* 6, 7. doi: [10.4081/ija.2011.e7](https://doi.org/10.4081/ija.2011.e7)
- Magnan A, Garnaud B, Billé R, Gemenne F, Hallegatte S 2009 The future of the Mediterranean : From impacts of climate change to adaptation issues.
- Magris RA, Andreollo M, Pressey RL, Mouillot D, Dalongeville A et al. 2018 Biologically representative and well-connected marine reserves enhance biodiversity persistence in conservation planning. *Conserv. Lett.* 11, e12439. doi: [10.1111/conl.12439](https://doi.org/10.1111/conl.12439)
- Magris RA, Pressey RL, Weeks R, Ban NC 2014 Integrating connectivity and climate change into marine conservation planning. *Biol. Conserv.* 170, 207–221. doi: [10.1016/j.biocon.2013.12.032](https://doi.org/10.1016/j.biocon.2013.12.032)
- Mahmoud A 2017 Production of quinoa (*Chenopodium quinoa*) in the marginal environments of South Mediterranean Region: Nile Delta, Egypt. *Egypt. J. Soil Sci.* 0, 0–0. doi: [10.21608/ejss.2017.436.1062](https://doi.org/10.21608/ejss.2017.436.1062)
- Makowski D, Marajo-Petizon E, Durand JL, Ben-Ari T 2020 Quantitative synthesis of temperature, CO<sub>2</sub>, rainfall, and adaptation effects on global crop yields. *Eur. J. Agron.* 115, 126041. doi: [10.1016/j.eja.2020.126041](https://doi.org/10.1016/j.eja.2020.126041)
- Malavasi M, Carranza ML, Moravec D, Cutini M 2018 Reforestation dynamics after land abandonment: a trajectory analysis in Mediterranean mountain landscapes. *Reg. Environ. Chang.* 18, 2459–2469. doi: [10.1007/s10113-018-1368-9](https://doi.org/10.1007/s10113-018-1368-9)
- Malcolm JR, Liu C, Neilson RP, Hansen L, Hannah L 2006 Global warming and extinctions of endemic species from biodiversity hotspots. *Conserv. Biol.* 20, 538–548. doi: [10.1111/j.1523-1739.2006.00364.x](https://doi.org/10.1111/j.1523-1739.2006.00364.x)
- Malek Ž, Verburg PH 2017 Mediterranean land systems: Representing diversity and intensity of complex land systems in a dynamic region. *Landsc. Urban Plan.* 165, 102–116. doi: [10.1016/j.landurbplan.2017.05.012](https://doi.org/10.1016/j.landurbplan.2017.05.012)
- Malek Ž, Verburg PH 2018 Adaptation of land management in the Mediterranean under scenarios of irrigation water use and availability. *Mitig. Adapt. Strateg. Glob. Chang.* 23, 821–837. doi: [10.1007/s11027-017-9761-0](https://doi.org/10.1007/s11027-017-9761-0)
- Malek Ž, Verburg PH, Geijzendorffer IR, Bondeau A, Cramer W 2018 Global change effects on land management in the Mediterranean region. *Glob. Environ. Chang.* 50, 238–254. doi: [10.1016/j.gloenvcha.2018.04.007](https://doi.org/10.1016/j.gloenvcha.2018.04.007)
- Mallo M, Ziveri P, Mortyn PG, Schiebel R, Grelaud M 2017 Low planktic foraminiferal diversity and abundance observed in a spring 2013 west–east Mediterranean Sea plankton tow transect. *Biogeosciences* 14, 2245–2266. doi: [10.5194/bg-14-2245-2017](https://doi.org/10.5194/bg-14-2245-2017)
- Mallo M, Ziveri P, Reyes-García V, Rossi S 2019 Historical record of *Corallium rubrum* and its changing carbon sequestration capacity: A meta-analysis from the North Western Mediterranean. *PLoS One* 14, e0223802. doi: [10.1371/journal.pone.0223802](https://doi.org/10.1371/journal.pone.0223802)
- Manea E, Bianchelli S, Fanelli E, Danovaro R, Gissi E 2020 Towards an Ecosystem-Based Marine Spatial Planning in the deep Mediterranean Sea. *Sci. Total Environ.* 715, 136884. doi: [10.1016/j.scitotenv.2020.136884](https://doi.org/10.1016/j.scitotenv.2020.136884)
- Mangialajo L, Bertolotto R, Cattaneo-Vietti R, Chiantore M, Grillo C et al. 2008a The toxic benthic dinoflagellate *Ostreopsis ovata*: Quantification of proliferation along the coastline of Genoa, Italy. *Mar. Pollut. Bull.* 56, 1209–1214. doi: [10.1016/j.marpolbul.2008.02.028](https://doi.org/10.1016/j.marpolbul.2008.02.028)
- Mangialajo L, Chiantore M, Cattaneo-Vietti R 2008b Loss of furoid algae along a gradient of urbanisation, and structure of benthic assemblages. *Mar. Ecol. Prog. Ser.* 358, 63–74. doi: [10.3354/meps07400](https://doi.org/10.3354/meps07400)
- Mannino AM, Balistreri P, Deidun A 2017 The Marine Biodiversity of the Mediterranean Sea in a Changing Climate: The Impact of Biological Invasions, in *Mediterranean Identities - Environment, Society, Culture* doi: [10.5772/intechopen.69214](https://doi.org/10.5772/intechopen.69214)
- Mannino MA, Talamo S, Tagliacozzo A, Fiore I, Nehlich O et al. 2015 Climate-driven environmental changes around 8,200 years ago favoured increases in cetacean strandings and Mediterranean hunter-gatherers exploited them. *Sci. Rep.* 5, 16288. doi: [10.1038/srep16288](https://doi.org/10.1038/srep16288)
- Mansourian S, Rossi M, Vallauri D 2013 Ancient Forests in the Northern Mediterranean: Neglected High Conservation Value Areas. Marseille, France doi: [10.13140/2.1.5170.4640](https://doi.org/10.13140/2.1.5170.4640)
- Marba N, Duarte CM 2010 Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality. *Glob. Chang. Biol.* 16, 2366–2375.
- Marbà N, Jordà G, Agustí S, Girard C, Duarte CM 2015 Footprints of climate change on Mediterranean Sea biota. *Front. Mar. Sci.* 2. doi: [10.3389/fmars.2015.00056](https://doi.org/10.3389/fmars.2015.00056)
- Marcos M, Tsimplis MN 2008 Comparison of results of AOGCMs in the Mediterranean Sea during the 21<sup>st</sup> century. *JGR Ocean.* 113. doi: [10.1029/2008JC004820](https://doi.org/10.1029/2008JC004820)
- Marić D, Kraus R, Godrijan J, Supić N, Djakovac T et al. 2012 Phytoplankton response to climatic and anthropogenic influences in the north-eastern Adriatic during the last four decades. *Estuar. Coast. Shelf Sci.* 115, 98–112. doi: [10.1016/j.ecss.2012.02.003](https://doi.org/10.1016/j.ecss.2012.02.003)
- Marino G, Ziveri P 2013 Palaeo-carbonate chemistry of the Mediterranean Sea. in *40th CIESM Congress* (Marseille, France).
- Martín-Alcón S, Ameztegui A, Coll L 2017 Diversificación

- o naturalización de las repoblaciones forestales, in *La Restauración Forestal de España: 75 Años de Una Ilusión*, eds. Permán García J, Iriarte Goñi I, Lario Leza FJ (Madrid, España: Ministerio de Agricultura y Pesca, Alimentación y Medio Ambiente), 401–411.
- Martín-Alcón S, Coll L 2016 Unraveling the relative importance of factors driving post-fire regeneration trajectories in non-serotinous *Pinus nigra* forests. *For. Ecol. Manage.* 361, 13–22. doi: [10.1016/j.foreco.2015.11.006](https://doi.org/10.1016/j.foreco.2015.11.006)
- Martín-Alcón S, Coll L, Ameztegui A 2016 Diversifying sub-Mediterranean pinewoods with oak species in a context of assisted migration: Responses to local climate and light environment. *Appl. Veg. Sci.* 19, 254–266. doi: [10.1111/avsc.12216](https://doi.org/10.1111/avsc.12216)
- Martín-Benito D, del Río M, Heinrich I, Helle G, Cañellas I 2010 Response of climate-growth relationships and water use efficiency to thinning in a *Pinus nigra* afforestation. *For. Ecol. Manage.* 259, 967–975. doi: [10.1016/j.foreco.2009.12.001](https://doi.org/10.1016/j.foreco.2009.12.001)
- Martín-Ortega P, Picard N, García-Montero LG, Río S, Penas Á et al. 2018 Importance of Mediterranean forests, in *State of Mediterranean Forests*, eds. Bourlion N, Garavaglia V, Picard N (Marseille, France: FAO & Plan Bleu), 31–50.
- Martin R, Müller B, Linstädter A, Frank K 2014 How much climate change can pastoral livelihoods tolerate? Modelling rangeland use and evaluating risk. *Glob. Environ. Chang.* 24, 183–192. doi: [10.1016/j.gloenvcha.2013.09.009](https://doi.org/10.1016/j.gloenvcha.2013.09.009)
- Martin S, Gattuso J-P 2009 Response of Mediterranean coralline algae to ocean acidification and elevated temperature. *Glob. Chang. Biol.* 15, 2089–2100. doi: [10.1111/j.1365-2486.2009.01874.x](https://doi.org/10.1111/j.1365-2486.2009.01874.x)
- Martínez-Fernández J, Esteve MA, Carreño MF, Palazón JA 2009 Dynamics of land use change in the Mediterranean : Implications for sustainability, land use planning and nature conservation, in *Land Use Policy* (Nova Science Publishers), 101–145.
- Martínez-Gómez C, Fernández B, Robinson CD, Campillo JA, León VM et al. 2017 Assessing environmental quality status by integrating chemical and biological effect data: The Cartagena coastal zone as a case. *Mar. Environ. Res.* 124, 106–117. doi: [10.1016/j.marenvres.2016.04.008](https://doi.org/10.1016/j.marenvres.2016.04.008)
- Martínez-Valderrama J, Ibáñez J, del Barrio G, Alcalá FJ, Sanjuán ME et al. 2018 Doomed to collapse: Why Algerian steppe rangelands are overgrazed and some lessons to help land-use transitions. *Sci. Total Environ.* 613–614, 1489–1497. doi: [10.1016/j.scitotenv.2017.07.058](https://doi.org/10.1016/j.scitotenv.2017.07.058)
- Martínez-Vilalta J, Lloret F 2016 Drought-induced vegetation shifts in terrestrial ecosystems: The key role of regeneration dynamics. *Glob. Planet. Change* 144, 94–108. doi: [10.1016/j.gloplacha.2016.07.009](https://doi.org/10.1016/j.gloplacha.2016.07.009)
- Marty J-C, Chiavérini J 2010 Hydrological changes in the Ligurian Sea (NW Mediterranean, DYFAMED site) during 1995–2007 and biogeochemical consequences. *Biogeosciences* 7, 2117–2128. doi: [10.5194/bg-7-2117-2010](https://doi.org/10.5194/bg-7-2117-2010)
- Marty J-C, Chiavérini J, Pizay M-D, Avril B 2002 Seasonal and interannual dynamics of nutrients and phytoplankton pigments in the western Mediterranean Sea at the DYFAMED time-series station (1991–1999). *Deep Sea Res. Part II Top. Stud. Oceanogr.* 49, 1965–1985. doi: [10.1016/S0967-0645\(02\)00022-X](https://doi.org/10.1016/S0967-0645(02)00022-X)
- Masiero M, Calama R, Lindner M, Pettenella D 2013 Forests in the Mediterranean region, in *Forest management of Mediterranean forests under the new context of climate change. Building alternative for the coming future.*, ed. Lucas-Borja ME (New York: Nova Science Publishers, Inc.), 13–30.
- Mateo G, Benito M, Laguna E 2013 *Flora Valentina. Flora Vasculare de la Comunitat Valenciana. Angiospermae (III). Berberidaceae-Compositae*. Valencia: Fundación de la Comunidad Valenciana para el Medio Ambiente.
- Matesanz S, Valladares F 2014 Ecological and evolutionary responses of Mediterranean plants to global change. *Environ. Exp. Bot.* 103, 53–67. doi: [10.1016/j.envexpbot.2013.09.004](https://doi.org/10.1016/j.envexpbot.2013.09.004)
- Mauri L, Sallustio L, Tarolli P 2019 The geomorphologic forcing of wild boars. *Earth Surf. Process. Landforms* 44, 2085–2094. doi: [10.1002/esp.4623](https://doi.org/10.1002/esp.4623)
- Mazurek H 2018 Les modèles de ville durable en question, in *L'urbanisation du monde*, eds. Dorier E, Lecoquierre M (La Documentation Française). <https://hal.archives-ouvertes.fr/hal-02547104> [Accessed August 14, 2020]
- Mazzoleni S, di Pasquale G, Mulligan M, di Martino P, Rego F 2004 *Recent dynamics of the Mediterranean vegetation and landscape*. John Wiley & Sons, Ltd. doi: [10.1002/0470093714](https://doi.org/10.1002/0470093714)
- McFadden L, Spencer T, Nicholls RJ 2007 Broad-scale modelling of coastal wetlands: what is required? *Hydrobiologia* 577, 5–15. doi: [10.1007/s10750-006-0413-8](https://doi.org/10.1007/s10750-006-0413-8)
- McNeill JR 1992 *The Mountains of the Mediterranean World*. Cambridge University Press. doi: [10.1017/cbo9780511529023](https://doi.org/10.1017/cbo9780511529023)
- MEA 2005 *Ecosystems and human well-being: synthesis*. Island Press, Washington, DC. <https://www.cifor.org/library/1888/>
- MED-EUWI 2007 *Mediterranean Groundwater Report - Technical report on groundwater management in the Mediterranean and the Water Framework Directive*.
- Médail F 2013 The unique nature of Mediterranean island floras and the future of plant conservation. in *Proceedings of the 2nd Botanical Conference in Menorca "Islands and plants: preservation and understanding of flora on Mediterranean islands,"* 325–350.
- Médail F 2017 The specific vulnerability of plant biodiversity and vegetation on Mediterranean islands in the face of global change. *Reg. Environ. Chang.*, 1–16. doi: [10.1007/s10113-017-1123-7](https://doi.org/10.1007/s10113-017-1123-7)
- Médail F, Diadema K 2009 Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *J.*

- Biogeogr.* 36, 1333–1345.  
doi: [10.1111/j.1365-2699.2008.02051.x](https://doi.org/10.1111/j.1365-2699.2008.02051.x)
- Médail F, Quézel P 1997 Hotspots analysis for conservation of plant biodiversity in the Mediterranean Basin. *Ann. Missouri Bot. Gard.* 84, 112–127. doi: [10.2307/2399957](https://doi.org/10.2307/2399957)
- Médail F, Quézel P 1999 Biodiversity Hotspots in the Mediterranean Basin: Setting Global Conservation Priorities. *Conserv. Biol.* 13, 1510–1513.  
doi: [10.1046/j.1523-1739.1999.98467.x](https://doi.org/10.1046/j.1523-1739.1999.98467.x)
- Mediterranean Wetlands Observatory 2012 Biodiversity–Status and trends of species in Mediterranean wetlands. Thematic collection, 1.
- Mediterranean Wetlands Observatory 2018 *Mediterranean Wetland Outlook 2: Solutions for sustainable Mediterranean Wetlands.*, eds. Geijzenborffer IR, Chazée L, Gaget E, Galewski T, Guelmami A et al. Arles, France <https://tourduvalat.org/en/actions/les-zones-humides-mediterraneennes-enjeux-et-perspectives-2-solutions-pour-des-zones-humides-mediterraneennes-durables/>
- MEDSEA 2015 Mediterranean Sea Acidification in a changing climate. Final report.  
<http://www.medsea-project.eu>
- Meier KJS, Beaufort L, Heussner S, Ziveri P 2014 The role of ocean acidification in *Emiliania huxleyi* coccolith thinning in the Mediterranean Sea. *Biogeosciences Discuss.* 11, 2857–2869. doi: [10.5194/bg-11-2857-2014](https://doi.org/10.5194/bg-11-2857-2014)
- Melki T, Kallel N, Jorissen FJ, Guichard F, Dennielou B et al. 2009 Abrupt climate change, sea surface salinity and paleoproductivity in the western Mediterranean Sea (Gulf of Lion) during the last 28 kyr. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 279, 96–113.  
doi: [10.1016/j.palaeo.2009.05.005](https://doi.org/10.1016/j.palaeo.2009.05.005)
- Mena C, Reglero P, Hidalgo M, Sintés E, Santiago R et al. 2019 Phytoplankton community structure is driven by stratification in the oligotrophic Mediterranean Sea. *Front. Microbiol.* 10, 1698.  
doi: [10.3389/fmicb.2019.01698](https://doi.org/10.3389/fmicb.2019.01698)
- Menció A, Mas-Pla J 2010 Influence of groundwater exploitation on the ecological status of streams in a Mediterranean system (Selva Basin, NE Spain). *Ecol. Indic.* 10, 915–926. doi: [10.1016/j.ecolind.2010.02.001](https://doi.org/10.1016/j.ecolind.2010.02.001)
- Mercader M, Rider M, Cheminée A, Pastor J, Zawadzki A et al. 2018 Spatial distribution of juvenile fish along an artificialized seascape, insights from common coastal species in the Northwestern Mediterranean Sea. *Mar. Environ. Res.* 137, 60–72.  
doi: [10.1016/j.marenvres.2018.02.030](https://doi.org/10.1016/j.marenvres.2018.02.030)
- Merheb M, Moussa R, Abdallah C, Colin F, Perrin C et al. 2016 Hydrological response characteristics of Mediterranean catchments at different time scales: a meta-analysis. *Hydrol. Sci. J.* 61, 2520–2539.  
doi: [10.1080/02626667.2016.1140174](https://doi.org/10.1080/02626667.2016.1140174)
- Merlivat L, Boutin J, Antoine D, Beaumont L, Golbol M et al. 2018 Increase of dissolved inorganic carbon and decrease in pH in near-surface waters in the Mediterranean Sea during the past two decades. *Biogeosciences* 15, 5653–5662.  
doi: [10.5194/bg-15-5653-2018](https://doi.org/10.5194/bg-15-5653-2018)
- Merlo M, Croitoru L 2005 *Valuing Mediterranean forest: towards total economic value.* Cabi Publishing
- Meyer J, Riebesell U 2015 Reviews and Syntheses: Responses of coccolithophores to ocean acidification: a meta-analysis. *Biogeosciences* 12, 1671–1682.  
doi: [10.5194/bg-12-1671-2015](https://doi.org/10.5194/bg-12-1671-2015)
- Middleton N 2018 Rangeland management and climate hazards in drylands: dust storms, desertification and the overgrazing debate. *Nat. Hazards* 92, 57–70.  
doi: [10.1007/s11069-016-2592-6](https://doi.org/10.1007/s11069-016-2592-6)
- Middleton N, Thomas D 1997 *World atlas of desertification.* Routledge; 2nd Edition
- Midolo G, Alkemade R, Schipper AM, Benítez-López A, Perring MP et al. 2019 Impacts of nitrogen addition on plant species richness and abundance: A global meta-analysis. *Glob. Ecol. Biogeogr.* 28, 398–413.  
doi: [10.1111/geb.12856](https://doi.org/10.1111/geb.12856)
- Migliavacca M, Dosio A, Camia A, Hobourg R, Houston-Durrant T et al. 2013 Modeling biomass burning and related carbon emissions during the 21<sup>st</sup> century in Europe. *JGR Biogeosciences* 118, 1732–1747.  
doi: [10.1002/2013jg002444](https://doi.org/10.1002/2013jg002444)
- Milano M, Ruelland D, Fernandez S, Dezetter A, Fabre J et al. 2013 Current state of Mediterranean water resources and future trends under climatic and anthropogenic changes. *Hydrol. Sci. J.* 58, 498–518.  
doi: [10.1080/02626667.2013.774458](https://doi.org/10.1080/02626667.2013.774458)
- Milazzo A, Giles LC, Zhang Y, Koehler AP, Hiller JE et al. 2016 The effect of temperature on different *Salmonella* serotypes during warm seasons in a Mediterranean climate city, Adelaide, Australia. *Epidemiol. Infect.* 144, 1231–1240. doi: [10.1017/S0950268815002587](https://doi.org/10.1017/S0950268815002587)
- Milazzo M, Rodolfo-Metalpa R, Chan VBS, Fine M, Alessi C et al. 2014 Ocean acidification impairs vermetid reef recruitment. *Sci. Rep.* 4, 4189. doi: [10.1038/srep04189](https://doi.org/10.1038/srep04189)
- Milisenda G, Rossi S, Vizzini S, Fuentes VL, Purcell JE et al. 2018 Seasonal variability of diet and trophic level of the gelatinous predator *Pelagia noctiluca* (Scyphozoa). *Sci. Rep.* 8. doi: [10.1038/s41598-018-30474-x](https://doi.org/10.1038/s41598-018-30474-x)
- Miller TE, Gornish ES, Buckley HL 2009 Climate and coastal dune vegetation: disturbance, recovery, and succession. *Plant Ecol.* 206, 97.  
doi: [10.1007/s11258-009-9626-z](https://doi.org/10.1007/s11258-009-9626-z)
- Milly PCD, Dunne KA, Vecchia A V. 2005 Global pattern of trends in streamflow and water availability in a changing climate. *Nature* 438, 347–50.  
doi: [10.1038/nature04312](https://doi.org/10.1038/nature04312)
- Mina M, Bugmann H, Cordonnier T, Irauschek F, Klopčič M et al. 2017 Future ecosystem services from European mountain forests under climate change. *J. Appl. Ecol.* 54, 389–401. doi: [10.1111/1365-2664.12772](https://doi.org/10.1111/1365-2664.12772)
- Mittermeier RA, Gil PR, Hoffman M, Pilgrim J, Brooks T et al. 2005 *Hotspots Revisited. Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions.* Chicago: University of Chicago Press for Conservation Interna-



- tional. doi: [10.1017/s0376892901270088](https://doi.org/10.1017/s0376892901270088)
- Mojtahid M, Manceau R, Schiebel R, Hennekam R, de Lange GJ 2015 Thirteen thousand years of southeastern Mediterranean climate variability inferred from an integrative planktic foraminiferal-based approach. *Paleoceanography* 30, 402–422. doi: [10.1002/2014pa002705](https://doi.org/10.1002/2014pa002705)
- Molina AC, Montefalcone M, Vassallo P, Morri C, Bianchi CN et al. 2016 Combining literature review, acoustic mapping and in situ observations: an overview of coralligenous assemblages in Liguria (NW Mediterranean Sea). *Sci. Mar.* 80, 7–16. doi: [10.3989/scimar.04235.23A](https://doi.org/10.3989/scimar.04235.23A)
- Molina JA, Casermeiro MA, Moreno PNS 2003 Vegetation composition and soil salinity in a Spanish Mediterranean coastal ecosystem. *Phytocoenologia* 33, 475–494. doi: [10.1127/0340-269x/2003/0033-0475](https://doi.org/10.1127/0340-269x/2003/0033-0475)
- Molinero JC, Ibanez F, Nival P, Buecher E, Souissi S 2005 North Atlantic climate and northwestern Mediterranean plankton variability. *Limnol. Oceanogr.* 50, 1213–1220. doi: [10.4319/lo.2005.50.4.1213](https://doi.org/10.4319/lo.2005.50.4.1213)
- Molinero JC, Ibañez F, Souissi S, Buecher E, Dallot S et al. 2008 Climate control on the long-term anomalous changes of zooplankton communities in the Northwestern Mediterranean. *Glob. Chang. Biol.* 14, 11–26. doi: [10.1111/j.1365-2486.2007.01469.x](https://doi.org/10.1111/j.1365-2486.2007.01469.x)
- Monioudi IN, Velegarakis AF, Chatzipavlis AE, Rigos A, Karambas T et al. 2017 Assessment of island beach erosion due to sea level rise: the case of the Aegean archipelago (Eastern Mediterranean). *Nat. Hazards Earth Syst. Sci.* 17, 449–466. doi: [10.5194/nhess-17-449-2017](https://doi.org/10.5194/nhess-17-449-2017)
- Montefalcone M, Morri C, Parravicini V, Bianchi CN 2015 A tale of two invaders: divergent spreading kinetics of the alien green algae *Caulerpa taxifolia* and *Caulerpa cylindracea*. *Biol. Invasions* 17, 2717–2728. doi: [10.1007/s10530-015-0908-1](https://doi.org/10.1007/s10530-015-0908-1)
- Montero-Serra I, Linares C, García M, Pancaldi F, Frleta-Valiç M et al. 2015 Harvesting Effects, Recovery Mechanisms, and Management Strategies for a Long-Lived and Structural Precious Coral. *PLoS One* 10, e0117250. doi: [10.1371/journal.pone.0117250](https://doi.org/10.1371/journal.pone.0117250)
- Monti M, Minocci M, Beran A, Iveša L 2007 First record of *Ostreopsis* cfr. *ovata* on macroalgae in the Northern Adriatic Sea. *Mar. Pollut. Bull.* 54, 598–601. doi: [10.1016/j.marpolbul.2007.01.013](https://doi.org/10.1016/j.marpolbul.2007.01.013)
- Morán-Ordóñez A, Roces-Díaz J V., Otsu K, Ameztegui A, Coll L et al. 2019 The use of scenarios and models to evaluate the future of nature values and ecosystem services in Mediterranean forests. *Reg. Environ. Chang.* 19, 415–428. doi: [10.1007/s10113-018-1408-5](https://doi.org/10.1007/s10113-018-1408-5)
- Morán-Tejeda E, Ceballos-Barbancho A, Llorente-Pinto JM 2010 Hydrological response of Mediterranean headwaters to climate oscillations and land-cover changes: The mountains of Duero River basin (Central Spain). *Glob. Planet. Change* 72, 39–49. doi: [10.1016/j.gloplacha.2010.03.003](https://doi.org/10.1016/j.gloplacha.2010.03.003)
- Moreira F, Arianoutsou M, Corona P, de Las Heras J 2011 *Post-Fire Management and Restoration of Southern European Forests*. Springer Netherlands doi: [10.1007/978-94-007-2208-8](https://doi.org/10.1007/978-94-007-2208-8)
- Moreno G, Aviron S, Berg S, Crous-Duran J, Franca A et al. 2018 Agroforestry systems of high nature and cultural value in Europe: provision of commercial goods and other ecosystem services. *Agrofor. Syst.* 92, 877–891. doi: [10.1007/s10457-017-0126-1](https://doi.org/10.1007/s10457-017-0126-1)
- Moriondo M, Bindi M, Kundzewicz ZW, Szwed M, Chorynski A et al. 2010 Impact and adaptation opportunities for European agriculture in response to climatic change and variability. *Mitig. Adapt. Strateg. Glob. Chang.* 15, 657–679. doi: [10.1007/s11027-010-9219-0](https://doi.org/10.1007/s11027-010-9219-0)
- Moriondo M, Good P, Durao R, Bindi M, Giannakopoulos C et al. 2006 Potential impact of climate change on fire risk in the Mediterranean area. *Clim. Res.* 31, 85–95. doi: [10.3354/cr031085](https://doi.org/10.3354/cr031085)
- Moritz MA, Batllori E, Bradstock RA, Gill AM, Handmer J et al. 2014 Learning to coexist with wildfire. *Nature* 515, 58–66. doi: [10.1038/nature13946](https://doi.org/10.1038/nature13946)
- Morritt D 1988 Osmoregulation in littoral and terrestrial talitroidean amphipods (Crustacea) from Britain. *J. Exp. Mar. Bio. Ecol.* 123, 77–94. doi: [10.1016/0022-0981\(88\)90110-4](https://doi.org/10.1016/0022-0981(88)90110-4)
- Mosquera-Losada MR, Santiago-Freijanes JJ, Pisanelli A, Rois-Díaz M, Smith J et al. 2018 Agroforestry in the European common agricultural policy. *Agrofor. Syst.* 92, 1117–1127. doi: [10.1007/s10457-018-0251-5](https://doi.org/10.1007/s10457-018-0251-5)
- Moullec F, Barrier N, Drira S, Guilhaumon F, Marsaleix P et al. 2019 An End-to-End Model Reveals Losers and Winners in a Warming Mediterranean Sea. *Front. Mar. Sci.* 6. doi: [10.3389/fmars.2019.00345](https://doi.org/10.3389/fmars.2019.00345)
- Moullec F, Benedetti F, Saraux C, VAN BEVEREN E, SHIN Y-J 2016. Climate change induces bottom-up changes in the food webs of the Mediterranean Sea. In “The Mediterranean Region under Climate Change” IRD-Marseille, 2016. ISBN : 978, in
- Moustadraf J, Razack M, Sinan M 2008 Evaluation of the impacts of climate changes on the coastal Chaouia aquifer, Morocco, using numerical modeling. *Hydrogeol. J.* 16, 1411–1426. doi: [10.1007/s10040-008-0311-4](https://doi.org/10.1007/s10040-008-0311-4)
- Movilla J, Calvo EM, Pelejero C, Coma R, Serrano E et al. 2012 Calcification reduction and recovery in native and non-native Mediterranean corals in response to ocean acidification. *J. Exp. Mar. Bio. Ecol.* 438, 144–153. doi: [10.1016/j.jembe.2012.09.014](https://doi.org/10.1016/j.jembe.2012.09.014)
- Movilla J, Orejas C, Calvo EM, Gori A, López-Sanz À et al. 2014 Differential response of two Mediterranean cold-water coral species to ocean acidification. *Coral Reefs* 33, 675–686. doi: [10.1007/s00338-014-1159-9](https://doi.org/10.1007/s00338-014-1159-9)
- Moy AD, Howard WR, Bray SG, Trull TW 2009 Reduced calcification in modern Southern Ocean planktonic foraminifera. *Nat. Geosci.* 2, 276–280. doi: [10.1038/ngeo460](https://doi.org/10.1038/ngeo460)
- Moyle PB 1995 Conservation of native freshwater fishes in

- the Mediterranean-type climate of California, USA: A review. *Biol. Conserv.* 72, 271–279. doi: [10.1016/0006-3207\(94\)00089-9](https://doi.org/10.1016/0006-3207(94)00089-9)
- Mualla W 2018 Water Demand Management Is a Must in MENA Countries...But Is It Enough? *J. Geol. Resour. Eng.* 6, 59–64. doi: [10.17265/2328-2193/2018.02.002](https://doi.org/10.17265/2328-2193/2018.02.002)
- Mullan M, Kingsmill N, Agrawala S, Kramer AM 2015 *National Adaptation Planning: Lessons from OECD Countries*. ed. W. Leal Filho Berlin-Heidelberg: Springer-Verlag. doi: [10.1007/978-3-642-38670-1\\_38](https://doi.org/10.1007/978-3-642-38670-1_38)
- Mullin M, Smith MD, McNamara DE 2019 Paying to save the beach: effects of local finance decisions on coastal management. *Clim. Change* 152, 275–289. doi: [10.1007/s10584-018-2191-5](https://doi.org/10.1007/s10584-018-2191-5)
- Murciego AM, Sánchez AG, González MAR, Gil EP, Gordillo CT et al. 2007 Antimony distribution and mobility in topsoils and plants (*Cytisus striatus*, *Cistus ladanifer* and *Dittrichia viscosa*) from polluted Sb-mining areas in Extremadura (Spain). *Environ. Pollut.* 145, 15–21. doi: [10.1016/j.envpol.2006.04.004](https://doi.org/10.1016/j.envpol.2006.04.004)
- Murfin J, Spiegel M 2020 Is the risk of sea level rise capitalized in residential real estate? *Rev. Financ. Stud.* 33, 1217–1255. doi: [10.1093/rfs/hhz134](https://doi.org/10.1093/rfs/hhz134)
- Murray PJ, Cook R, Currie AF, Dawson LA, Gange AC et al. 2006 Interactions between fertilizer addition, plants and the soil environment: Implications for soil faunal structure and diversity. *Appl. Soil Ecol.* 33, 199–207. doi: [10.1016/j.apsoil.2005.11.004](https://doi.org/10.1016/j.apsoil.2005.11.004)
- Murray SN, Denis TG, Kido JS, Smith JR 1999 Human visitation and the frequency and potential effects of collecting on rocky intertidal populations in Southern California marine reserves. *Reports Calif. Coop. Ocean. Fish. Investig.* 40, 101–106. <https://pdfs.semanticscholar.org/2858/caa50f2eea96ba0ff-705f7ad978e31529332.pdf>
- Mutke S, Gordo J, Gil L 2005 Variability of Mediterranean Stone pine cone production: Yield loss as response to climate change. *Agric. For. Meteorol.* 132, 263–272. doi: [10.1016/j.agrformet.2005.08.002](https://doi.org/10.1016/j.agrformet.2005.08.002)
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J 2000 Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858.
- Myers PG, Haines K, Rohling EJ 1998 Modeling the paleocirculation of the Mediterranean: The Last Glacial Maximum and the Holocene with emphasis on the formation of sapropelS1. *Paleoceanography* 13, 586–606. doi: [10.1029/98pa02736](https://doi.org/10.1029/98pa02736)
- Nadal-Sala D, Hartig F, Gracia CA, Sabaté S 2019 Global warming likely to enhance black locust (*Robinia pseudoacacia* L.) growth in a Mediterranean riparian forest. *For. Ecol. Manage.* 449, 117448. doi: [10.1016/j.foreco.2019.117448](https://doi.org/10.1016/j.foreco.2019.117448)
- Nader MR, Indary S, Boustany L 2012 The puffer fish *Lagocephalus scleratus* (Gmelin, 1789) in the Eastern Mediterranean, in *EastMed Technical Documents (FAO)*, 34.
- Nagelkerken I, Russell BD, Gillanders BM, Connell SD 2016 Ocean acidification alters fish populations indirectly through habitat modification. *Nat. Clim. Chang.* 6, 89–93. doi: [10.1038/nclimate2757](https://doi.org/10.1038/nclimate2757)
- Naiman RJ, Latterell JJ, Pettit NE, Olden JD 2008 Flow variability and the biophysical vitality of river systems. *Comptes Rendus Geosci.* 340, 629–643. doi: [10.1016/j.crte.2008.01.002](https://doi.org/10.1016/j.crte.2008.01.002)
- Nainggolan D, de Vente J, Boix-Fayos C, Termansen M, Hubacek K et al. 2012 Afforestation, agricultural abandonment and intensification: Competing trajectories in semi-arid Mediterranean agro-ecosystems. *Agric. Ecosyst. Environ.* 159, 90–104. doi: [10.1016/J.AGEE.2012.06.023](https://doi.org/10.1016/J.AGEE.2012.06.023)
- Nastasi A 2010 Algal and Jellyfish Blooms in the Mediterranean and Black Sea: a brief review. in *GFCM Workshop on Algal and Jellyfish Blooms in the Mediterranean and Black Sea (FAO/GFCM-6th/8th October 2010)* (Istanbul, Turkey).
- Navarro-Ortega A, Acuña V, Bellin A, Burek P, Cassiani G et al. 2015 Managing the effects of multiple stressors on aquatic ecosystems under water scarcity. The GLOBAQUA project. *Sci. Total Environ.* 503–504, 3–9. doi: [10.1016/j.scitotenv.2014.06.081](https://doi.org/10.1016/j.scitotenv.2014.06.081)
- Newton A, Brito AC, Icely JD, Derolez V, Clara I et al. 2018 Assessing, quantifying and valuing the ecosystem services of coastal lagoons. *J. Nat. Conserv.* 44, 50–65. doi: [10.1016/j.jnc.2018.02.009](https://doi.org/10.1016/j.jnc.2018.02.009)
- Ngatia L, Grace III JM, Moriasi D, Taylor R 2019 Nitrogen and phosphorus eutrophication in marine ecosystems, in *Monitoring of Marine Pollution*. doi: [10.5772/intechopen.81869](https://doi.org/10.5772/intechopen.81869)
- Nguyen TPL, Mula L, Cortignani R, Seddaiu G, Dono G et al. 2016 Perceptions of present and future climate change impacts on water availability for agricultural systems in the western mediterranean region. *Water* 8. doi: [10.3390/w8110523](https://doi.org/10.3390/w8110523)
- Nicholls RJ, Hoozemans FMJ 1996 The Mediterranean: Vulnerability to coastal implications of climate change. *Ocean Coast. Manag.* 31, 105–132. doi: [10.1016/S0964-5691\(96\)00037-3](https://doi.org/10.1016/S0964-5691(96)00037-3)
- Nieto-Romero M, Oteros-Rozas E, González J a., Martín-López B 2014 Exploring the knowledge landscape of ecosystem services assessments in Mediterranean agroecosystems: Insights for future research. *Environ. Sci. Policy* 37, 121–133. doi: [10.1016/j.envsci.2013.09.003](https://doi.org/10.1016/j.envsci.2013.09.003)
- Nieto A, Roberts SPM, Kemp J, Rasmont P, Kuhlmann M et al. 2015 *European Red List of Bees*. doi: [10.2779/77003](https://doi.org/10.2779/77003)
- Nilsson C, Berggren K 2000 Alterations of Riparian Ecosystems Caused by River Regulation. *Bioscience* 50, 783. doi: [10.1641/0006-3568\(2000\)050\[0783:aorecb\]2.0.co;2](https://doi.org/10.1641/0006-3568(2000)050[0783:aorecb]2.0.co;2)
- Nilsson C, Reidy CA, Dynesius M, Revenga C 2005 Fragmentation and Flow Regulation of the World's Large River Systems. *Science (80-. )*. 308, 405–408. doi: [10.1126/science.1107887](https://doi.org/10.1126/science.1107887)
- Nilsson GE, Dixon DL, Domenici P, McCormick MI, Sørensen C et al. 2012 Near-future carbon dioxide

- levels alter fish behaviour by interfering with neurotransmitter function. *Nat. Clim. Chang.* 2, 201–204.
- Ninčević Gladan Ž, Matic F, Arapov J, Skejić S, Bužančić M et al. 2020 The relationship between toxic phytoplankton species occurrence and environmental and meteorological factors along the Eastern Adriatic coast. *Harmful Algae* 92, 101745. doi: [10.1016/j.hal.2020.101745](https://doi.org/10.1016/j.hal.2020.101745)
- Nizamuddin M 1991 *The green marine algae of Libya*. Bern: Elga Publishers.
- Noce S, Collalti A, Valentini R, Santini M 2016 Hot spot maps of forest presence in the Mediterranean basin. *iForest - Biogeosciences For.* 9, 766–774. doi: [10.3832/ifer1802-009](https://doi.org/10.3832/ifer1802-009)
- Nocentini S, Coll L 2013 Mediterranean forests: human use and complex adaptive systems, in *Managing Forests as Complex Adaptive Systems. Building resilience to the challenge of Global Change*, eds. Messier C, Puettmann K, Coates K (New York, NY: Routledge).
- Nogués-Bravo D, Araújo MB, Lasanta T, Moreno JIL 2008 Climate change in Mediterranean mountains during the 21<sup>st</sup> century. *Ambio* 37, 280–285. doi: [10.1579/0044-7447\(2008\)37\[280:ccimmd\]2.0.co;2](https://doi.org/10.1579/0044-7447(2008)37[280:ccimmd]2.0.co;2)
- Notarbartolo di Sciarra G 2014 Sperm whales, *Physeter macrocephalus*, in the Mediterranean Sea: a summary of status, threats, and conservation recommendations. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 24, 4–10. doi: [10.1002/aqc.2409](https://doi.org/10.1002/aqc.2409)
- Noto AE, Shurin JB 2017 Early stages of sea-level rise lead to decreased salt marsh plant diversity through stronger competition in Mediterranean-climate marshes. *PLoS One* 12, e0169056. doi: [10.1371/journal.pone.0169056](https://doi.org/10.1371/journal.pone.0169056)
- Noulas C, Tziouvalakas M, Vlachostergios D, Baxevanos D, Karyotis T et al. 2017 Adaptation, Agronomic Potential, and Current Perspectives of Quinoa Under Mediterranean Conditions: Case Studies from the Lowlands of Central Greece. *Commun. Soil Sci. Plant Anal.* 48, 2612–2629. doi: [10.1080/00103624.2017.1416129](https://doi.org/10.1080/00103624.2017.1416129)
- Noy-Meir I 1973 Desert Ecosystems: Environment and Producers. *Annu. Rev. Ecol. Syst.* 4, 25–51. doi: [10.1146/annurev.es.04.110173.000325](https://doi.org/10.1146/annurev.es.04.110173.000325)
- Nunes A, Köbel M, Pinho P, Matos P, Costantini EAC et al. 2019 Local topographic and edaphic factors largely predict shrub encroachment in Mediterranean drylands. *Sci. Total Environ.* 657, 310–318. doi: [10.1016/j.scitotenv.2018.11.475](https://doi.org/10.1016/j.scitotenv.2018.11.475)
- Ochoa-Hueso R, Bell MD, Manrique E 2014 Impacts of increased nitrogen deposition and altered precipitation regimes on soil fertility and functioning in semiarid Mediterranean shrublands. *J. Arid Environ.* 104, 106–115. doi: [10.1016/j.jaridenv.2014.01.020](https://doi.org/10.1016/j.jaridenv.2014.01.020)
- Ochoa-Hueso R, Munzi S, Alonso R, Arróniz-Crespo M, Avila A et al. 2017 Ecological impacts of atmospheric pollution and interactions with climate change in terrestrial ecosystems of the Mediterranean Basin: Current research and future directions. Elsevier <https://www.sciencedirect.com/science/article/pii/S0269749116320760?via%3Dihub> [Accessed March 30, 2019].
- Oliveira MA, Tomlinson SJ, Carnell EJ, Dore AJ, Serrano HC et al. 2020 Nitrogen and sulfur deposition over a region in SW Europe based on a regional atmospheric chemical transport model. *Atmos. Environ.* 223, 117290. doi: [10.1016/j.atmosenv.2020.117290](https://doi.org/10.1016/j.atmosenv.2020.117290)
- Oliveira V, Lauw A, Pereira H 2016 Sensitivity of cork growth to drought events: insights from a 24-year chronology. *Clim. Change* 137, 261–274. doi: [10.1007/s10584-016-1680-7](https://doi.org/10.1007/s10584-016-1680-7)
- Olsson L, Barbosa H, Bhadwal S, Cowie AL, Delusca K et al. 2019 Land Degradation, in *Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems*, eds. Shukla PR, Skea J, Buendia EC, Masson-Delmotte V, Pörtner H-O et al. (In press).
- Omezzine F, Rinez A, Ladhari A, Farooq M, Haouala R 2011 Allelopathic potential of *Inula viscosa* against crops and weeds. *Int. J. Agric. Biol.* 13, 841–849. <http://www.fsublishers.org> [Accessed August 12, 2020].
- Ondiviela B, Losada IJ, Lara JL, Maza M, Galván C et al. 2014 The role of seagrasses in coastal protection in a changing climate. *Coast. Eng.* 87, 158–168. doi: [10.1016/j.coastaleng.2013.11.005](https://doi.org/10.1016/j.coastaleng.2013.11.005)
- Ontoria Y, Gonzalez-Guedes E, Sanmartí N, Bernardeau-Esteller J, Ruiz JM et al. 2019 Interactive effects of global warming and eutrophication on a fast-growing Mediterranean seagrass. *Mar. Environ. Res.* 145, 27–38. doi: [10.1016/j.marenvres.2019.02.002](https://doi.org/10.1016/j.marenvres.2019.02.002)
- Ordoñez A, Wangpraseurt D, Lyndby NH, Kühl M, Díaz-Pulido G 2019 Elevated CO<sub>2</sub> leads to enhanced photosynthesis but decreased growth in early life stages of reef building coralline algae. *Front. Mar. Sci.* 5, 495. doi: [10.3389/fmars.2018.00495](https://doi.org/10.3389/fmars.2018.00495)
- Ortega-Reig M, Sanchis-Ibor C, Palau-Salvador G, García-Mollá M, Avellà-Reus L 2017 Institutional and management implications of drip irrigation introduction in collective irrigation systems in Spain. *Agric. Water Manag.* 187, 164–172. doi: [10.1016/J.AGWAT.2017.03.009](https://doi.org/10.1016/J.AGWAT.2017.03.009)
- Ortega M, Velasco J, Millán A, Guerrero C 2004 An ecological integrity index for littoral wetlands in agricultural catchments of semiarid mediterranean regions. *Environ. Manage.* 33, 412–430. doi: [10.1007/s00267-003-3059-6](https://doi.org/10.1007/s00267-003-3059-6)
- OSPAR Commission 2007 Background document to support the assessment of whether the OSPAR Network of Marine Protected Areas is ecologically coherent. *Biodivers. Ser.* <http://www.ospar.org/documents?v=7077>
- Otero I, Marull J, Tello E, Diana GL, Pons M et al. 2015 Land abandonment, landscape, and biodiversity: Questioning the restorative character of the forest

- transition in the Mediterranean. *Ecol. Soc.* 20. doi: [10.5751/ES-07378-200207](https://doi.org/10.5751/ES-07378-200207)
- Ouba A, Abboud-Abi Saab M, Stemmann L 2016 Temporal Variability of Zooplankton (2000–2013) in the Levantine Sea: Significant Changes Associated to the 2005–2010 EMT-like Event? *PLoS One* 11, e158484. doi: [10.1371/journal.pone.0158484](https://doi.org/10.1371/journal.pone.0158484)
- Oviedo A, Ziveri P, Álvarez M, Tanhua T 2015 Is coccolithophore distribution in the Mediterranean Sea related to seawater carbonate chemistry? *Ocean Sci.* 11, 13–32. doi: [10.5194/os-11-13-2015](https://doi.org/10.5194/os-11-13-2015)
- Owojori OJ, Reinecke AJ, Rozanov AB 2008 Effects of salinity on partitioning, uptake and toxicity of zinc in the earthworm *Eisenia fetida*. *Soil Biol. Biochem.* 40, 2385–2393. doi: [10.1016/j.soilbio.2008.05.019](https://doi.org/10.1016/j.soilbio.2008.05.019)
- Owojori OJ, Waszak K, Roembke J 2014 Avoidance and reproduction tests with the predatory mite *Hypoaspis aculeifer*: effects of different chemical substances. *Environ. Toxicol. Chem.* 33, 230–237. doi: [10.1002/etc.2421](https://doi.org/10.1002/etc.2421)
- Ozment S, DiFrancesco K, Gartner T 2015 The role of natural infrastructure in the water, energy and food nexus. Gland, Switzerland. doi: [10.2305/IUCN.CH.2015.NEX.4.en](https://doi.org/10.2305/IUCN.CH.2015.NEX.4.en)
- Paine TD, Lieutier F 2016 *Insects and diseases of Mediterranean forest systems*. Springer International Publishing Switzerland [https://www.researchgate.net/profile/Edmundo\\_Sousa/publication/309052281\\_The\\_Pine\\_Wood\\_Nematode\\_and\\_Its\\_Local\\_Vectors\\_in\\_the\\_Mediterranean\\_Basin/links/5a6724830f7e9b76ea8d6812/The-Pine-Wood-Nematode-and-Its-Local-Vectors-in-the-Mediterranean-Basin.pdf](https://www.researchgate.net/profile/Edmundo_Sousa/publication/309052281_The_Pine_Wood_Nematode_and_Its_Local_Vectors_in_the_Mediterranean_Basin/links/5a6724830f7e9b76ea8d6812/The-Pine-Wood-Nematode-and-Its-Local-Vectors-in-the-Mediterranean-Basin.pdf)
- Pairaud IL, Bensoussan N, Garreau P, Faure V, Garrabou J 2014 Impacts of climate change on coastal benthic ecosystems: assessing the current risk of mortality outbreaks associated with thermal stress in NW Mediterranean coastal areas. *Ocean Dyn.* 64, 103–115. doi: [10.1007/s10236-013-0661-x](https://doi.org/10.1007/s10236-013-0661-x)
- Palahi M, Mavsar R, Gracia C, Birot Y 2008 Mediterranean forests under focus. *Int. For. Rev.* 10, 676–688. doi: [10.1505/ifer.10.4.676](https://doi.org/10.1505/ifer.10.4.676)
- Palau-Salvador G, de Luis A, Pérez JJ, Sanchis-Ibor C 2019 Greening the post crisis. Collectivity in private and public community gardens in València (Spain). *Cities* 92, 292–302. doi: [10.1016/j.cities.2019.04.005](https://doi.org/10.1016/j.cities.2019.04.005)
- Palmiéri J, Orr JC, Dutay J-C, Béranger K, Schneider A et al. 2015 Simulated anthropogenic CO<sub>2</sub> storage and acidification of the Mediterranean Sea. *Biogeosciences* 12, 781–802. doi: [10.5194/bg-12-781-2015](https://doi.org/10.5194/bg-12-781-2015)
- Palombo C, Chirici G, Marchetti M, Tognetti R 2013 Is land abandonment affecting forest dynamics at high elevation in Mediterranean mountains more than climate change? *Plant Biosyst.* 147, 1–11. doi: [10.1080/11263504.2013.772081](https://doi.org/10.1080/11263504.2013.772081)
- Panayotov M, Bebi P, Trouet Y, Yurukov S 2010 Climate signal in tree-ring chronologies of *Pinus peuce* and *Pinus heldreichii* from the Pirin Mountains in Bulgaria. *Trees* 24, 479–490. doi: [10.1007/s00468-010-0416-y](https://doi.org/10.1007/s00468-010-0416-y)
- Panigada S, Pesante G, Zanardelli M, Capoulade F, Gannier A et al. 2006 Mediterranean fin whales at risk from fatal ship strikes. *Mar. Pollut. Bull.* 52, 1287–1298. doi: [10.1016/j.marpolbul.2006.03.014](https://doi.org/10.1016/j.marpolbul.2006.03.014)
- Papadopoulos AG 2015 The Impact of the CAP on Agriculture and Rural Areas of EU Member States. *Agrar. South J. Polit. Econ.* 4, 22–53. doi: [10.1177/2277976015574054](https://doi.org/10.1177/2277976015574054)
- Papanastasis VP 2012 Land use changes, in *Mediterranean mountain environments*, ed. Vogiatzakis IN (Wiley–Blackwell), 159–184.
- Papastergiadou ES, Retalis A, Kalliris P, Georgiadis T 2007 Land use changes and associated environmental impacts on the Mediterranean shallow Lake Stymfalia, Greece. in *Hydrobiologia* (Springer), 361–372. doi: [10.1007/s10750-007-0606-9](https://doi.org/10.1007/s10750-007-0606-9)
- Parihar P, Singh S, Singh R, Singh VP, Prasad SM 2015 Effect of salinity stress on plants and its tolerance strategies: a review. *Environ. Sci. Pollut. Res.* 22, 4056–4075. doi: [10.1007/s11356-014-3739-1](https://doi.org/10.1007/s11356-014-3739-1)
- Parravicini V, Mangialajo L, Mousseau L, Peirano A, Morri C et al. 2015 Climate change and warm-water species at the north-western boundary of the Mediterranean Sea. *Mar. Ecol.* 36, 897–909. doi: [10.1111/maec.12277](https://doi.org/10.1111/maec.12277)
- Pascual M, Rossetto M, Ojea E, Milchakova N, Giakoumi S et al. 2016 Socioeconomic impacts of marine protected areas in the Mediterranean and Black Seas. *Ocean Coast. Manag.* 133, 1–10. doi: [10.1016/j.ocecoaman.2016.09.001](https://doi.org/10.1016/j.ocecoaman.2016.09.001)
- Patton M 1996 *Islands in time: island sociogeography and Mediterranean prehistory*. London: Routledge
- Pauli H, Gottfried M, Dullinger S, Abdaladze O, Akhalkatsi M et al. 2012 Recent Plant Diversity Changes on Europe's Mountain Summits. *Science* (80-. ). 336, 353–355. <http://science.sciencemag.org/content/336/6079/353.abstract>
- Pausas JG, Llovet J, Rodrigo A, Vallejo R 2009 Are wildfires a disaster in the Mediterranean basin? - A review. *Int. J. Wildl. Fire* 17, 713–723. doi: [10.1071/wf07151](https://doi.org/10.1071/wf07151)
- Pautasso M, Döring TF, Garbelotto M, Pellis L, Jeger MJ 2012 Impacts of climate change on plant diseases—opinions and trends. *Eur. J. Plant Pathol.* 133, 295–313. doi: [10.1007/s10658-012-9936-1](https://doi.org/10.1007/s10658-012-9936-1)
- Pavlopoulos K, Karkanas P, Triantaphyllou M, Karymbalis E, Tsourou T et al. 2006 Paleoenvironmental Evolution of the Coastal Plain of Marathon, Greece, during the Late Holocene: Depositional Environment, Climate, and Sea Level Changes. *J. Coast. Res.*, 424–438. doi: [10.2112/03-0145.1](https://doi.org/10.2112/03-0145.1)
- Pemán J, Serrada R 2017 El Plan general de Repoblación de España de 1939, in *La Restauración Forestal de España: 75 Años de Una Ilusión*, eds. Permán García J, Iriarte Goñi I, Lario Leza FJ (Madrid, España: Ministerio de Agricultura y Pesca, Alimentación y Medio Ambiente), 119–134.

- Pennino MG, Arcangeli A, Prado Fonseca V, Campana I, Pierce GJ et al. 2017 A spatially explicit risk assessment approach: Cetaceans and marine traffic in the Pelagos Sanctuary (Mediterranean Sea). *PLoS One* 12, e0179686. doi: [10.1371/journal.pone.0179686](https://doi.org/10.1371/journal.pone.0179686)
- Pennino MG, Pérez Roda MA, Pierce GJ, Rotta A 2016 Effects of vessel traffic on relative abundance and behaviour of cetaceans: the case of the bottlenose dolphins in the Archipelago de La Maddalena, north-western Mediterranean sea. *Hydrobiologia* 776, 237–248. doi: [10.1007/s10750-016-2756-0](https://doi.org/10.1007/s10750-016-2756-0)
- Peñuelas J, Boada M 2003 A global change-induced biome shift in the Montseny mountains (NE Spain). *Glob. Chang. Biol.* 9, 131–140. doi: [10.1046/j.1365-2486.2003.00566.x](https://doi.org/10.1046/j.1365-2486.2003.00566.x)
- Peñuelas J, Canadell JG, Ogaya R 2011 Increased water-use efficiency during the 20<sup>th</sup> century did not translate into enhanced tree growth. *Glob. Ecol. Biogeogr.* 20, 597–608. doi: [10.1111/j.1466-8238.2010.00608.x](https://doi.org/10.1111/j.1466-8238.2010.00608.x)
- Peñuelas J, Filella I, Comas PE 2002 Changed plant and animal life cycles from 1952 to 2000 in the Mediterranean region. *Glob. Chang. Biol.* 8, 531–544. doi: [10.1046/j.1365-2486.2002.00489.x](https://doi.org/10.1046/j.1365-2486.2002.00489.x)
- Peñuelas J, Filella I, Llorens L, Ogaya R, Lloret F et al. 2004 Complex spatiotemporal phenological shifts as a response to rainfall changes. *New Phytol.* 161, 837–846. doi: [10.1111/j.1469-8137.2003.01003.x](https://doi.org/10.1111/j.1469-8137.2003.01003.x)
- Peñuelas J, Ogaya R, Boada M, Jump AS 2007 Migration, invasion and decline: changes in recruitment and forest structure in a warming-linked shift of European beech forest in Catalonia (NE Spain). *Ecography (Cop.)* 30, 829–837. doi: [10.1111/j.2007.0906-7590.05247.x](https://doi.org/10.1111/j.2007.0906-7590.05247.x)
- Peñuelas J, Sardans J, Filella I, Estiarte M, Llusà J et al. 2017 Impacts of global change on Mediterranean forests and their services. *Forests* 8, 463. doi: [10.3390/f8120463](https://doi.org/10.3390/f8120463)
- Peñuelas J, Sardans J, Filella I, Estiarte M, Llusà J et al. 2018 Assessment of the impacts of climate change on Mediterranean terrestrial ecosystems based on data from field experiments and long-term monitored field gradients in Catalonia. *Environ. Exp. Bot.* 152, 49–59. doi: [10.1016/j.envexpbot.2017.05.012](https://doi.org/10.1016/j.envexpbot.2017.05.012)
- Pereira SC, Lopes I, Abrantes I, Sousa JP, Chelinho S 2019 Salinization effects on coastal ecosystems: a terrestrial model ecosystem approach. *Philos. Trans. R. Soc. B Biol. Sci.* 374, 20180251. doi: [10.1098/rstb.2018.0251](https://doi.org/10.1098/rstb.2018.0251)
- Pérez-Ruzafa A, Marcos C, Pérez-Ruzafa IM 2011 Mediterranean coastal lagoons in an ecosystem and aquatic resources management context. *Phys. Chem. Earth, Parts A/B/C* 36, 160–166. doi: [10.1016/j.pce.2010.04.013](https://doi.org/10.1016/j.pce.2010.04.013)
- Perez T, Garrabou J, Sartoretto S, Harmelin JG, Francour P et al. 2000 Massive mortality of marine invertebrates: an unprecedented event in northwestern Mediterranean. *C. R. Acad. Sci. III.* 323, 853–865. doi: [10.1016/S0764-4469\(00\)01237-3](https://doi.org/10.1016/S0764-4469(00)01237-3)
- Petanidou T, Kallimanis AS, Tzanopoulos J, Sgardelis SP, Pantis JD 2008 Long-term observation of a pollination network: fluctuation in species and interactions, relative invariance of network structure and implications for estimates of specialization. *Ecol. Lett.* 11, 564–575. doi: [10.1111/j.1461-0248.2008.01170.x](https://doi.org/10.1111/j.1461-0248.2008.01170.x)
- Pétillon J, Ysnel F, Canard A, Lefeuvre JC 2005 Impact of an invasive plant (*Elymus athericus*) on the conservation value of tidal salt marshes in western France and implications for management: Responses of spider populations. *Biol. Conserv.* 126, 103–117. doi: [10.1016/j.biocon.2005.05.003](https://doi.org/10.1016/j.biocon.2005.05.003)
- Phalan B, Onial M, Balmford A, Green RE 2011 Reconciling Food Production and Biodiversity Conservation: Land Sharing and Land Sparing Compared. *Science (80-. )* 333, 1289–1291. doi: [10.1126/science.1208742](https://doi.org/10.1126/science.1208742)
- Philippart CJM, Anadón R, Danovaro R, Dippner JW, Drinkwater KF et al. 2011 Impacts of climate change on European marine ecosystems: Observations, expectations and indicators. *J. Exp. Mar. Bio. Ecol.* 400, 52–69. doi: [10.1016/j.jembe.2011.02.023](https://doi.org/10.1016/j.jembe.2011.02.023)
- Piazzi L, Balata D, Cecchi E, Cinelli F, Sartoni G 2009 Species composition and patterns of diversity of macroalgal coralligenous assemblages in the north-western Mediterranean Sea. *J. Nat. Hist.* 44, 1–22. doi: [10.1080/00222930903377547](https://doi.org/10.1080/00222930903377547)
- Piazzi L, Gennaro P, Balata D 2012 Threats to macroalgal coralligenous assemblages in the Mediterranean Sea. *Mar. Pollut. Bull.* 64, 2623–2629. doi: [10.1016/j.marpolbul.2012.07.027](https://doi.org/10.1016/j.marpolbul.2012.07.027)
- Pinho P, Augusto S, Martins-Loução MA, Pereira MJ, Soares A et al. 2008 Causes of change in nitrophytic and oligotrophic lichen species in a Mediterranean climate: Impact of land cover and atmospheric pollutants. *Environ. Pollut.* 154, 380–389. doi: [10.1016/j.envpol.2007.11.028](https://doi.org/10.1016/j.envpol.2007.11.028)
- Pinho P, Bergamini A, Carvalho P, Branquinho C, Stofer S et al. 2012a Lichen functional groups as ecological indicators of the effects of land-use in Mediterranean ecosystems. *Ecol. Indic.* 15, 36–42. doi: [10.1016/j.ecolind.2011.09.022](https://doi.org/10.1016/j.ecolind.2011.09.022)
- Pinho P, Branquinho C, Cruz C, Tang YS, Dias T et al. 2009 Assessment of critical levels of atmospheric ammonia for lichen diversity in cork-oak woodland, Portugal, in *Atmospheric Ammonia: Detecting Emission Changes and Environmental Impacts* (Springer Netherlands), 109–119. doi: [10.1007/978-1-4020-9121-6\\_10](https://doi.org/10.1007/978-1-4020-9121-6_10)
- Pinho P, Dias T, Cordovil CM d. S, Dragosits U, Dise NB et al. 2018 Mapping Portuguese Natura 2000 sites in risk of biodiversity change caused by atmospheric nitrogen pollution. *PLoS One* 13, e0198955. doi: [10.1371/journal.pone.0198955](https://doi.org/10.1371/journal.pone.0198955)
- Pinho P, Dias T, Cruz C, Sim Tang Y, Sutton MA et al. 2011 Using lichen functional diversity to assess the effects of atmospheric ammonia in Mediterranean woodlands. *J. Appl. Ecol.* 48, 1107–1116. doi: [10.1111/j.1365-2664.2011.02033.x](https://doi.org/10.1111/j.1365-2664.2011.02033.x)

- Pinho P, Theobald MR, Dias T, Tang YS, Cruz C et al. 2012b Critical loads of nitrogen deposition and critical levels of atmospheric ammonia for semi-natural Mediterranean evergreen woodlands. *Biogeosciences* 9, 1205–1215. doi: [10.5194/bg-9-1205-2012](https://doi.org/10.5194/bg-9-1205-2012)
- Pittalis D, Carletti A, Ghiglieri G, Celico F 2016 The influence of hydrogeological properties, seawater intrusion and refreshing on the quality of groundwater used for irrigation in an agricultural coastal plain in North Sardinia, Italy. *Environ. Earth Sci.* 75, 963. doi: [10.1007/s12665-016-5770-7](https://doi.org/10.1007/s12665-016-5770-7)
- Piva A, Asioli A, Trincardi F, Schneider RR, Vigliotti L 2008 Late-Holocene climate variability in the Adriatic Sea (Central Mediterranean). *The Holocene* 18, 153–167. doi: [10.1177/0959683607085606](https://doi.org/10.1177/0959683607085606)
- Pivotto ID, Nerini D, Masmoudi M, Kara H, Chaoui L et al. 2015 Highly contrasted responses of Mediterranean octocorals to climate change along a depth gradient. *R. Soc. Open Sci.* 2, 140493–140493. doi: [10.1098/rsos.140493](https://doi.org/10.1098/rsos.140493)
- Plieninger T, Gaertner M, Hui C, Huntsinger L 2013 Does land abandonment decrease species richness and abundance of plants and animals in Mediterranean pastures, arable lands and permanent croplands? *Environ. Evid.* 2, 3. doi: [10.1186/2047-2382-2-3](https://doi.org/10.1186/2047-2382-2-3)
- Poff NL, Allan JD, Bain MB, Karr JR, Prestegard KL et al. 1997 The natural flow regime. *Bioscience* 47, 769–784. doi: [10.2307/1313099](https://doi.org/10.2307/1313099)
- Poher Y, Ponel P, Médail F, Tachikawa K, Guiter F 2018 Response of two Mediterranean coastal wetlands (Corsica, France) to Holocene relative sea-level rise and land-use changes revealed by fossil Coleoptera records. in *IPA-IAL 2018 – Joint Meeting – Unravelling the Past and Future of Lakes* <https://hal-amu.archives-ouvertes.fr/hal-01910973/document>
- Poláková J, Tucker G, Hart K, Dwyer J, Rayment M 2011 Addressing biodiversity and habitat preservation through Measures applied under the Common Agricultural Policy. London.
- Poloczanska ES, Brown CJ, Sydeman WJ, Kiessling W, Schoeman DS et al. 2013 Global imprint of climate change on marine life. *Nat. Clim. Chang.* 3, 919–925. doi: [10.1038/nclimate1958](https://doi.org/10.1038/nclimate1958)
- Ponti L, Gutierrez AP, Ruti PM, Dell’aquila A 2014a Fine-scale ecological and economic assessment of climate change on olive in the Mediterranean Basin reveals winners and losers. *Proc. Natl. Acad. Sci. U. S. A.* 111, 5598–5603. doi: [10.1073/pnas.1314437111](https://doi.org/10.1073/pnas.1314437111)
- Ponti M, Grech D, Mori M, Perlini RA, Ventra V et al. 2016 The role of gorgonians on the diversity of vagile benthic fauna in Mediterranean rocky habitats. *Mar. Biol.* 163, 1–14. doi: [10.1007/s00227-016-2897-8](https://doi.org/10.1007/s00227-016-2897-8)
- Ponti M, Perlini RA, Ventra V, Grech D, Abbiati M et al. 2014b Ecological shifts in Mediterranean coralligenous assemblages related to gorgonian forest loss. *PLoS One* 9, e102782. doi: [10.1371/journal.pone.0102782](https://doi.org/10.1371/journal.pone.0102782)
- Ponti M, Turicchia E, Ferro F, Cerrano C, Abbiati M 2018 The understory of gorgonian forests in mesophotic temperate reefs. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 28, 1153–1166. doi: [10.1002/aqc.2928](https://doi.org/10.1002/aqc.2928)
- Popa I, Kern Z 2009 Long-term summer temperature reconstruction inferred from tree-ring records from the Eastern Carpathians. *Clim. Dyn.* 32, 1107–1117. doi: [10.1007/s00382-008-0439-x](https://doi.org/10.1007/s00382-008-0439-x)
- Por F 2009 Tethys returns to the Mediterranean: Success and limits of tropical re-colonization. *BioRisk* 3, 5–19. doi: [10.3897/biorisk.3.30](https://doi.org/10.3897/biorisk.3.30)
- Porqueddu C, Ates S, Louhaichi M, Kyriazopoulos AP, Moreno G et al. 2016 Grasslands in “Old World” and “New World” Mediterranean-climate zones: Past trends, current status and future research priorities. *Grass Forage Sci.* 71, 1–35. doi: [10.1111/gfs.12212](https://doi.org/10.1111/gfs.12212)
- Porte C, Escartín E, García de la Parra LM, Biosca X, Albaigés J 2002 Assessment of coastal pollution by combined determination of chemical and biochemical markers in *Mullus barbatus*. *Mar. Ecol. Prog. Ser.* 235, 205–216. doi: [10.3354/meps235205](https://doi.org/10.3354/meps235205)
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O et al. 2010 Global pollinator declines: Trends, impacts and drivers. *Trends Ecol. Evol.* 25, 345–353. doi: [10.1016/j.tree.2010.01.007](https://doi.org/10.1016/j.tree.2010.01.007)
- Potts SG, Vulliamy B, Dafni A, Ne’eman G, O’Toole C et al. 2003 Response of plant-pollinator communities to fire: Changes in diversity, abundance and floral reward structure. *Oikos* 101, 103–112. doi: [10.1034/j.1600-0706.2003.12186.x](https://doi.org/10.1034/j.1600-0706.2003.12186.x)
- Powell EJ, Tyrrell MC, Milliken A, Tirpak JM, Staudinger MD 2019 A review of coastal management approaches to support the integration of ecological and human community planning for climate change. *J. Coast. Conserv.* 23, 1–18. doi: [10.1007/s11852-018-0632-y](https://doi.org/10.1007/s11852-018-0632-y)
- Powley HR, Krom MD, Van Cappellen P 2016 Circulation and oxygen cycling in the Mediterranean Sea: Sensitivity to future climate change. *JGR Ocean.* 121, 8230–8247. doi: [10.1002/2016JC012224](https://doi.org/10.1002/2016JC012224)
- Prada F, Caroselli E, Mengoli S, Brizi L, Fantazzini P et al. 2017 Ocean warming and acidification synergistically increase coral mortality. *Sci. Rep.* 7, 40842. doi: [10.1038/srep40842](https://doi.org/10.1038/srep40842)
- Prat N, Ibáñez C 1995 Effects of water transfers projected in the Spanish National Hydrological Plan on the ecology of the lower River Ebro (N.E. Spain) and its delta. *Water Sci. Technol.* 31, 79–86. doi: [10.1016/0273-1223\(95\)00359-U](https://doi.org/10.1016/0273-1223(95)00359-U)
- Prenda J, Clavero M, Blanco-Garrido F, Menor A, Hermoso V 2006 Threats to the conservation of biotic integrity in Iberian fluvial ecosystems. *Limnetica* 25, 377–388.
- Prestele R, Alexander P, Rounsevell MDA, Arneth A, Calvin K V. et al. 2016 Hotspots of uncertainty in land-use and land-cover change projections: a global-scale model comparison. *Glob. Chang. Biol.* 22, 3967–3983. doi: [10.1111/gcb.13337](https://doi.org/10.1111/gcb.13337)
- Pretzsch H, Schütze G, Uhl E 2013 Resistance of European

- tree species to drought stress in mixed versus pure forests: Evidence of stress release by inter-specific facilitation. *Plant Biol.* 15, 483–495. doi: [10.1111/j.1438-8677.2012.00670.x](https://doi.org/10.1111/j.1438-8677.2012.00670.x)
- Prieto L, Macías DM, Peliz A, Ruiz J 2015 Portuguese Man-of-War (*Physalia physalis*) in the Mediterranean: A permanent invasion or a casual appearance? *Sci. Rep.* 5. doi: [10.1038/srep11545](https://doi.org/10.1038/srep11545)
- Príncipe A, Matos P, Sarris D, Gaiola G, do Rosário L et al. 2019 In Mediterranean drylands microclimate affects more tree seedlings than adult trees. *Ecol. Indic.* 106, 105476. doi: [10.1016/j.ecolind.2019.105476](https://doi.org/10.1016/j.ecolind.2019.105476)
- Prisco I, Carboni M, Acosta ATR 2013 The fate of threatened coastal dune habitats in Italy under climate change scenarios. *PLoS One* 8, e68850. doi: [10.1371/journal.pone.0068850](https://doi.org/10.1371/journal.pone.0068850)
- Prisco I, Carboni M, Jucker T, Acosta ATR 2016 Temporal changes in the vegetation of Italian coastal dunes: identifying winners and losers through the lens of functional traits. *J. Appl. Ecol.* 53, 1533–1542. doi: [10.1111/1365-2664.12684](https://doi.org/10.1111/1365-2664.12684)
- Prosperi P, Allen T, Padilla M, Peri I, Cogill B 2014 Sustainability and Food & Nutrition Security: A Vulnerability Assessment Framework for the Mediterranean Region. *SAGE Open* 4, 2158244014539169. doi: [10.1177/2158244014539169](https://doi.org/10.1177/2158244014539169)
- Provansal M, Dufour S, Sabatier F, Anthony EJ, Raccasi G et al. 2014 The geomorphic evolution and sediment balance of the lower Rhône River (southern France) over the last 130 years: Hydropower dams versus other control factors. *Geomorphology* 219, 27–41. doi: [10.1016/j.geomorph.2014.04.033](https://doi.org/10.1016/j.geomorph.2014.04.033)
- Psarra S, Tselepides A, Ignatiades L 2000 Primary productivity in the oligotrophic Cretan Sea (NE Mediterranean): seasonal and interannual variability. *Prog. Oceanogr.* 46, 187–204. doi: [10.1016/S0079-6611\(00\)00018-5](https://doi.org/10.1016/S0079-6611(00)00018-5)
- Psarra S, Zohary T, Krom MD, Mantoura RFC, Polychronaki T et al. 2005 Phytoplankton response to a Lagrangian phosphate addition in the Levantine Sea (Eastern Mediterranean). *Deep Sea Res. Part II Top. Stud. Oceanogr.* 52, 2944–2960. doi: [10.1016/j.dsr2.2005.08.015](https://doi.org/10.1016/j.dsr2.2005.08.015)
- Puettmann KJ, Coates KD, Messier C 2008 *A Critique of Silviculture - Managing for Complexity*. Washington, USA: Island Press, Washington, DC.
- Puigdefábregas J, Mendizabal T 1998 Perspectives on desertification: western Mediterranean. *J. Arid Environ.* 39, 209–224. doi: [10.1006/jare.1998.0401](https://doi.org/10.1006/jare.1998.0401)
- Pulina A, Lai R, Salis L, Seddaiu G, Roggero PP et al. 2017 Modelling pasture production and soil temperature, water and carbon fluxes in Mediterranean grassland systems with the Pasture Simulation model. *Grass Forage Sci.*, 1–12. doi: [10.1111/gfs.12310](https://doi.org/10.1111/gfs.12310)
- Qadir M, Sharma BR, Bruggeman A, Choukr-Allah R, Karajeh F 2007 Non-conventional water resources and opportunities for water augmentation to achieve food security in water scarce countries. *Agric. Water Manag.* 87, 2–22. doi: [10.1016/j.agwat.2006.03.018](https://doi.org/10.1016/j.agwat.2006.03.018)
- Quézel P, Médail F 2003 *Ecology and biogeography of Mediterranean Basin forests*. Elsevier M. Paris
- Quintana-Seguí P, Martin E, Sánchez E, Zribi M, Vennetier M et al. 2016 Drought: Observed trends, future projections, in *The Mediterranean Region under Climate Change. A Scientific Update*, eds. Thiébaud S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 123–132.
- Raji O, Dezileau L, Von Grafenstein U, Niazi S, Snoussi M et al. 2015 Extreme sea events during the last millennium in the northeast of Morocco. *Nat. Hazards Earth Syst. Sci.* 15, 203–211. doi: <https://doi.org/10.5194/nhess-15-203-2015>
- Ramieri E, Bocci M, Markovic M 2019 Linking integrated coastal zone management to maritime spatial planning: The mediterranean experience, in *Maritime Spatial Planning: past, present, future* (Springer International Publishing), 271–294. doi: [10.1007/978-3-319-98696-8\\_12](https://doi.org/10.1007/978-3-319-98696-8_12)
- Ramírez-Romero E, Molinero JC, Sommer U, Salhi N, Kéfi - Daly Yahia O et al. 2020 Phytoplankton size changes and diversity loss in the southwestern Mediterranean Sea in relation to long-term hydrographic variability. *Estuar. Coast. Shelf Sci.* 235, 106574. doi: [10.1016/j.ecss.2019.106574](https://doi.org/10.1016/j.ecss.2019.106574)
- Ramírez F, Rodríguez C, Seoane J, Figuerola J, Bustamante J 2018 How will climate change affect endangered Mediterranean waterbirds? *PLoS One* 13, e0192702. doi: [10.1371/journal.pone.0192702](https://doi.org/10.1371/journal.pone.0192702)
- Ramsar Convention on Wetlands 2018 Global Wetland Outlook: State of the World's Wetlands and their Services to People. Gland, Switzerland.
- Rana G, Ferrara RM 2019 Air cooling by tree transpiration: A case study of *Olea europaea*, *Citrus sinensis* and *Pinus pinea* in Mediterranean town. *Urban Clim.* 29, 100507. doi: [10.1016/j.uclim.2019.100507](https://doi.org/10.1016/j.uclim.2019.100507)
- Range P, Chícharo MA, Ben-Hamadou R, Piló D, Fernandez-Reiriz MJ et al. 2014 Impacts of CO<sub>2</sub>-induced seawater acidification on coastal Mediterranean bivalves and interactions with other climatic stressors. *Reg. Environ. Chang.* 14, 19–30. doi: [10.1007/s10113-013-0478-7](https://doi.org/10.1007/s10113-013-0478-7)
- Rault PAK, Koundouri P, Akinsete E, Ludwig R, Huber-Garcia V et al. 2019 Down scaling of climate change scenarios to river basin level: A transdisciplinary methodology applied to Evrotas river basin, Greece. *Sci. Total Environ.* 660, 1623–1632. doi: [10.1016/j.scitotenv.2018.12.369](https://doi.org/10.1016/j.scitotenv.2018.12.369)
- Raviv O, Zemah Shamir S, Izhaki I, Sagie H, Negev M et al. 2020 The socioeconomic value of multiple ecosystem types at a biosphere reserve as a baseline for one holistic conservation plan. *Ecosyst. Serv.* 41, 101043. doi: [10.1016/j.ecoser.2019.101043](https://doi.org/10.1016/j.ecoser.2019.101043)
- Regato P, Salman R 2008 *Mediterranean Mountains in a Changing World: Guidelines for developing action plans*. Malaga, Spain: IUCN Centre for Mediterranean

- Cooperation <https://www.researchgate.net/publication/228610010>
- Regos A, Aquilué N, López I, Codina M, Retana J et al. 2016 Synergies Between Forest Biomass Extraction for Bioenergy and Fire Suppression in Mediterranean Ecosystems: Insights from a Storyline-and-Simulation Approach. *Ecosystems* 19, 786–802. doi: [10.1007/s10021-016-9968-z](https://doi.org/10.1007/s10021-016-9968-z)
- Rejdali M 2004 Forest cover changes in the Maghreb countries with special reference to Morocco, in *Recent Dynamics of the Mediterranean Vegetation and Landscape*, eds. Mazzoleni S, Pasquale G Di, Mulligan M, Di Martino P, Rego F (John Wiley & Sons, Ltd), 21–31. doi: [10.1002/0470093714.ch3](https://doi.org/10.1002/0470093714.ch3)
- Rennenberg H, Loreto F, Polle A, Brilli F, Fares S et al. 2006 Physiological responses of forest trees to heat and drought. *Plant Biol.* 8, 556–571. doi: [10.1055/s-2006-924084](https://doi.org/10.1055/s-2006-924084)
- Rescia AJ, Willaarts BA, Schmitz MF, Aguilera PA 2010 Changes in land uses and management in two Nature Reserves in Spain: Evaluating the social-ecological resilience of cultural landscapes. *Landsc. Urban Plan.* 98, 26–35. doi: [10.1016/j.landurbplan.2010.07.007](https://doi.org/10.1016/j.landurbplan.2010.07.007)
- Resh VH, Beche LA, Lawrence JE, Mazor RD, McElravy EP et al. 2013 Long-term population and community patterns of benthic macroinvertebrates and fishes in Northern California Mediterranean-climate streams. *Hydrobiologia* 719, 93–118. doi: [10.1007/s10750-012-1373-9](https://doi.org/10.1007/s10750-012-1373-9)
- Reynard E, Bonriposi M, Graefe O, Homewood C, Huss M et al. 2014 Interdisciplinary assessment of complex regional water systems and their future evolution: how socioeconomic drivers can matter more than climate. *Wiley Interdiscip. Rev. Water* 1, 413–426. doi: [10.1002/wat2.1032](https://doi.org/10.1002/wat2.1032)
- Reynolds JF, Smith DMS, Lambin EF, Turner BL, Mortimore M et al. 2007 Global Desertification: Building a Science for Dryland Development. *Science (80-. )*. 316, 847–851. doi: [10.1126/science.1131634](https://doi.org/10.1126/science.1131634)
- Ribera d'Alcalà M, Conversano F, Corato F, Licandro P, Mangoni O et al. 2004 Seasonal patterns in plankton communities in a pluriannual time series at a coastal Mediterranean site (Gulf of Naples): an attempt to discern recurrences and trends. *Sci. Mar.* 68, 65–83. doi: [10.3989/scimar.2004.68s165](https://doi.org/10.3989/scimar.2004.68s165)
- Richon C, Dutay J-C, Bopp L, Le Vu B, Orr JC et al. 2019 Biogeochemical response of the Mediterranean Sea to the transient SRES-A2 climate change scenario. *Biogeosciences* 16, 135–165. doi: [10.5194/bg-16-135-2019](https://doi.org/10.5194/bg-16-135-2019)
- Richon C, Dutay J-C, Dulac F, Wang R, Balkanski Y 2018a Modeling the biogeochemical impact of atmospheric phosphate deposition from desert dust and combustion sources to the Mediterranean Sea. *Biogeosciences* 15, 2499–2524. doi: [10.5194/bg-15-2499-2018](https://doi.org/10.5194/bg-15-2499-2018)
- Richon C, Dutay J-C, Dulac F, Wang R, Balkanski Y et al. 2018b Modeling the impacts of atmospheric deposition of nitrogen and desert dust-derived phosphorus on nutrients and biological budgets of the Mediterranean Sea. *Prog. Oceanogr.* 163, 21–39. doi: [10.1016/j.pocean.2017.04.009](https://doi.org/10.1016/j.pocean.2017.04.009)
- Ricketts TH, Dinerstein E, Boucher T, Brooks TM, Butchart SHM et al. 2005 Pinpointing and preventing imminent extinctions. *Proc. Natl. Acad. Sci. U. S. A.* 102, 18497–18501. doi: [10.1073/pnas.0509060102](https://doi.org/10.1073/pnas.0509060102)
- Rilov G 2016 Multi-species collapses at the warm edge of a warming sea. *Sci. Rep.* 6, 36897. doi: [10.1038/srep36897](https://doi.org/10.1038/srep36897)
- Rivas BL, Koleva-Lizama I 2005 Influence of climate variability on water resources in the Bulgarian South Black Sea basin. *IAHS-AISH Publ.* 296, 81–88.
- Rivetti I, Frascchetti S, Lionello P, Zambianchi E, Boero F 2014 Global Warming and Mass Mortalities of Benthic Invertebrates in the Mediterranean Sea. *PLoS One* 9, e115655. doi: [10.1371/journal.pone.0115655](https://doi.org/10.1371/journal.pone.0115655)
- Rixen M, Beckers J-M, Levitus S, Antonov J, Boyer T et al. 2005 The Western Mediterranean Deep Water: A proxy for climate change. *Geophys. Res. Lett.* 32, n/a-n/a. doi: [10.1029/2005GL022702](https://doi.org/10.1029/2005GL022702)
- Rizzi J, Gallina V, Torresan S, Critto A, Gana S et al. 2016 Regional Risk Assessment addressing the impacts of climate change in the coastal area of the Gulf of Gabes (Tunisia). *Sustain. Sci.* 11, 455–476. doi: [10.1007/s11625-015-0344-2](https://doi.org/10.1007/s11625-015-0344-2)
- Rizzo L, Pusceddu A, Stabili L, Alifano P, Frascchetti S 2017 Potential effects of an invasive seaweed (*Caulerpa cylindracea*, Sonder) on sedimentary organic matter and microbial metabolic activities. *Sci. Rep.* 7. doi: [10.1038/s41598-017-12556-4](https://doi.org/10.1038/s41598-017-12556-4)
- Roberts C, Hawkins JP, Fletcher J, Hands S, Raab K et al. 2010 Guidance on the size and spacing of marine protected areas in England. Natural England Commissioned Report NECR037.
- Robertson M, Murray-Prior R 2016 Five reasons why it is difficult to talk to Australian farmers about the impacts of, and their adaptation to, climate change. *Reg. Environ. Chang.* 16, 189–198. doi: [10.1007/s10113-014-0743-4](https://doi.org/10.1007/s10113-014-0743-4)
- Robinson C, Tocker K, Ward J 2002 The fauna of dynamic riverine landscapes. *Freshw. Biol.* 47, 661–677. doi: [10.1046/j.1365-2427.2002.00921.x](https://doi.org/10.1046/j.1365-2427.2002.00921.x)
- Robinson RA, Learmonth JA, Hutson AM, Macleod CD, Sparks TH et al. 2005 Climate Change and Migratory Species. Thetford, Norfolk.
- Roca E, Gamboa G, Tàbara JD 2008 Assessing the multidimensionality of coastal erosion risks: public participation and multicriteria analysis in a Mediterranean coastal system. *Risk Anal.* 28, 399–412. doi: [10.1111/j.1539-6924.2008.01026.x](https://doi.org/10.1111/j.1539-6924.2008.01026.x)
- Röder A, Udelhoven T, Hill J, del Barrio G, Tsiourlis G 2008 Trend analysis of Landsat-TM and -ETM+ imagery to monitor grazing impact in a rangeland ecosystem in Northern Greece. *Remote Sens. Environ.* 112, 2863–2875. doi: [10.1016/j.rse.2008.01.018](https://doi.org/10.1016/j.rse.2008.01.018)
- Rodolfo-Metalpa R, Hoogenboom MO, Rottier C, Ra-



- mos-Esplá A, Baker AC et al. 2014 Thermally tolerant corals have limited capacity to acclimatize to future warming. *Glob. Chang. Biol.* 20, 3036–3049. doi: [10.1111/gcb.12571](https://doi.org/10.1111/gcb.12571)
- Rodolfo-Metalpa R, Houlbrèque F, Tambutté É, Boisson F, Baggini C et al. 2011 Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nat. Clim. Chang.* 1, 308–312. doi: [10.1038/nclimate1200](https://doi.org/10.1038/nclimate1200)
- Rodolfo-Metalpa R, Lombardi C, Cocito S, Hall-Spencer JM, Gambi MC 2010 Effects of ocean acidification and high temperatures on the bryozoan *Myriapora truncata* at natural CO<sub>2</sub> vents. *Mar. Ecol.* 31, 447–456. doi: [10.1111/j.1439-0485.2009.00354.x](https://doi.org/10.1111/j.1439-0485.2009.00354.x)
- Rodrigues LC, Van den Bergh JCJM, Massa F, Theodorou JA, Ziveri P et al. 2015 Sensitivity of Mediterranean bivalve mollusc aquaculture to climate change, ocean acidification, and other environmental pressures: findings from a producer survey. *J. Shellfish Res.* 34, 1161–1176. doi: [10.2983/035.034.0341](https://doi.org/10.2983/035.034.0341)
- Rodríguez-Díaz JA, Pérez-Urrestarazu L, Camacho-Poyato E, Montesinos P 2011 The paradox of irrigation scheme modernization: more efficient water use linked to higher energy demand. *Spanish J. Agric. Res.* 4, 1000–1008. doi: [10.5424/sjar/20110904-492-10](https://doi.org/10.5424/sjar/20110904-492-10)
- Rodríguez-Ortega T, Bernués A, Olaizola AM, Brown MT 2017 Does intensification result in higher efficiency and sustainability? An emergy analysis of Mediterranean sheep-crop farming systems. *J. Clean. Prod.* 144, 171–179. doi: [10.1016/J.JCLEPRO.2016.12.089](https://doi.org/10.1016/J.JCLEPRO.2016.12.089)
- Rodríguez-Ortega T, Olaizola AM, Bernués A 2018 A novel management-based system of payments for ecosystem services for targeted agri-environmental policy. *Ecosyst. Serv.* 34, 74–84. doi: [10.1016/J.ECOSER.2018.09.007](https://doi.org/10.1016/J.ECOSER.2018.09.007)
- Rodriguez-Ramirez N, Santonja M, Baldy V, Ballini C, Montès N 2017 Shrub species richness decreases negative impacts of drought in a Mediterranean ecosystem. *J. Veg. Sci.* doi: [10.1111/jvs.12558](https://doi.org/10.1111/jvs.12558)
- Rodríguez-Rodríguez D, Rees S, Mannaerts G, Sciberras M, Pirie C et al. 2015 Status of the marine protected area network across the English channel (La Manche): Cross-country similarities and differences in MPA designation, management and monitoring. *Mar. Policy* 51, 536–546. doi: [10.1016/j.marpol.2014.09.021](https://doi.org/10.1016/j.marpol.2014.09.021)
- Roether W, Manca BB, Klein B, Bregant D, Georgopoulos D et al. 1996 Recent Changes in Eastern Mediterranean Deep Waters. *Science (80-. )*. 271, 333–335. doi: [10.1126/science.271.5247.333](https://doi.org/10.1126/science.271.5247.333)
- Rohling EJ 1994 Review and new aspects concerning the formation of eastern Mediterranean sapropels. *Mar. Geol.* 122, 1–28. doi: [10.1016/0025-3227\(94\)90202-X](https://doi.org/10.1016/0025-3227(94)90202-X)
- Rojas-Downing MM, Nejadhashemi AP, Harrigan T, Woznicki SA 2017 Climate change and livestock: Impacts, adaptation, and mitigation. *Clim. Risk Manag.* 16, 145–163. doi: [10.1016/J.CRM.2017.02.001](https://doi.org/10.1016/J.CRM.2017.02.001)
- Rolo V, Moreno G 2019 Shrub encroachment and climate change increase the exposure to drought of Mediterranean wood-pastures. 660, 550–558. <https://www.sciencedirect.com/science/article/pii/S004896971930035X> [Accessed May 26, 2019].
- Romanazzi A, Gentile F, Polemio M 2015 Modelling and management of a Mediterranean karstic coastal aquifer under the effects of seawater intrusion and climate change. *Environ. Earth Sci.* 74, 115–128. doi: [10.1007/s12665-015-4423-6](https://doi.org/10.1007/s12665-015-4423-6)
- Romano JC, Bensoussan N, Younes WA, Arlhac D 2000 Thermal anomaly in waters of the Gulf of Marseilles during the summer of 1999. A partial explanation of the mortality of certain fixed invertebrates? *Comptes Rendus l'Académie des Sci. - Ser. III - Sci. la Vie* 323, 415–427. doi: [10.1016/S0764-4469\(00\)00141-4](https://doi.org/10.1016/S0764-4469(00)00141-4)
- Roques A, Rousselet J, Avci M, Avtzis DN, Basso A et al. 2015 Climate warming and past and present distribution of the processionary moths (*Thaumetopoea* spp.) in Europe, Asia Minor and North Africa, in *Processionary Moths and Climate Change: An Update*, ed. Roques A (Dordrecht: Springer Netherlands), 81–161. doi: [10.1007/978-94-017-9340-7\\_3](https://doi.org/10.1007/978-94-017-9340-7_3)
- Ros JD, Romero J, Ballesteros E, Gili JM 1985 Diving in blue water. The benthos, in *Western Mediterranean*, ed. Margalef R (Oxford: Pergamon Press), 233–295.
- Rosa S, Pansera M, Granata A, Guglielmo L 2013 Interannual variability, growth, reproduction and feeding of *Pelagia noctiluca* (Cnidaria: Scyphozoa) in the Straits of Messina (Central Mediterranean Sea): Linkages with temperature and diet. *J. Mar. Syst.* 111–112, 97–107. doi: [10.1016/j.jmarsys.2012.10.001](https://doi.org/10.1016/j.jmarsys.2012.10.001)
- Rossetti I, Bagella S, Cappai C, Caria MC, Lai R et al. 2015 Isolated cork oak trees affect soil properties and biodiversity in a Mediterranean wooded grassland. *Agric. Ecosyst. Environ.* 202. doi: [10.1016/j.agee.2015.01.008](https://doi.org/10.1016/j.agee.2015.01.008)
- Rossi S, Bramanti L, Gori A, Orejas C 2017a An overview of the animal forests of the world, in *Marine animal forests: the ecology of benthic biodiversity hotspots*, eds. Rossi S, Bramanti L, Gori A, Orejas C, 1–28.
- Rossi S, Coppari M, Viladrich N 2017b Benthic-Pelagic Coupling: New Perspectives in the Animal Forests, in *Marine Animal Forests* (Springer International Publishing), 855–885. doi: [10.1007/978-3-319-21012-4\\_23](https://doi.org/10.1007/978-3-319-21012-4_23)
- Rossi S, Isla E, Bosch-Belmar M, Galli G, Gori A et al. 2019 Changes of energy fluxes in marine animal forests of the Anthropocene: factors shaping the future seascape. *ICES J. Mar. Sci.* 76, 2008–2019. doi: [10.1093/icesjms/fsz147](https://doi.org/10.1093/icesjms/fsz147)
- Rossi S, Tsounis G 2007 Temporal and spatial variation in protein, carbohydrate, and lipid levels in *Corallium rubrum* (Anthozoa, Octocorallia). *Mar. Biol.* 152, 429–439. doi: [10.1007/s00227-007-0702-4](https://doi.org/10.1007/s00227-007-0702-4)
- Rossi S, Tsounis G, Orejas C, Padrón T, Gili JM et al. 2008 Survey of deep-dwelling red coral (*Corallium rubrum*) populations at Cap de Creus (NW Mediterranean). *Mar. Biol.* 154, 533–545. doi: [10.1007/s00227-008-0947-6](https://doi.org/10.1007/s00227-008-0947-6)
- Rosignol-Strick M, Nesteroff W, Olive P, Vergnaud-Grazz-

- ini C 1982 After the deluge: Mediterranean stagnation and sapropel formation. *Nature* 295, 105–110. doi: [10.1038/295105a0](https://doi.org/10.1038/295105a0)
- Rouis-Zargouni I, Turon J-L, Londeix L, Essallami L, Kallel N et al. 2010 Environmental and climatic changes in the central Mediterranean Sea (Siculo-Tunisian Strait) during the last 30ka based on dinoflagellate cyst and planktonic foraminifera assemblages. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 285, 17–29. doi: [10.1016/j.palaeo.2009.10.015](https://doi.org/10.1016/j.palaeo.2009.10.015)
- Rouissi M, Muller SD, Ben Haj Jilani I, Ghrabi-Gammar Z, Paradis L et al. 2018 History and conservation of Tunisia's largest freshwater wetland: Garâa Sejenane. *Rev. Palaeobot. Palynol.* 257, 43–56. doi: [10.1016/j.revpalbo.2018.06.008](https://doi.org/10.1016/j.revpalbo.2018.06.008)
- Rubio-Portillo E, Gago JF, Martínez-García M, Vezzulli L, Rosselló-Móra R et al. 2018 *Vibrio* communities in scleractinian corals differ according to health status and geographic location in the Mediterranean Sea. *Syst. Appl. Microbiol.* 41, 131–138. doi: [10.1016/j.syapm.2017.11.007](https://doi.org/10.1016/j.syapm.2017.11.007)
- Ruiz-Benito P, Cuevas JA, Bravo R, García-del-Barrio JM, Zavala MA 2012 Land use change in a Mediterranean metropolitan region and its periphery: assessment of conservation policies through CORINE Land Cover data and Markov models. *For. Syst.* 19, 315. doi: [10.5424/fs/2010193-8604](https://doi.org/10.5424/fs/2010193-8604)
- Ruiz-Labourdette D, Nogués-Bravo D, Ollero HS, Schmitz MF, Pineda FD 2012 Forest composition in Mediterranean mountains is projected to shift along the entire elevational gradient under climate change. *J. Biogeogr.* 39, 162–176. doi: [10.1111/j.1365-2699.2011.02592.x](https://doi.org/10.1111/j.1365-2699.2011.02592.x)
- Ruiz-Peinado R, Bravo-Oviedo A, López-Senespleda E, Bravo F, del Río M 2017 Forest management and carbon sequestration in the Mediterranean region: A review. *For. Syst.* 26, 10. doi: [10.5424/fs/2017262-11205](https://doi.org/10.5424/fs/2017262-11205)
- Saadi S, Todorovic M, Tanasijevic L, Pereira LS, Pizzigalli C et al. 2015 Climate change and Mediterranean agriculture: Impacts on winter wheat and tomato crop evapotranspiration, irrigation requirements and yield. *Agric. Water Manag.* 147, 103–115. doi: [10.1016/j.agwat.2014.05.008](https://doi.org/10.1016/j.agwat.2014.05.008)
- Sabaté S, Gracia CA 2002 Likely effects of climate change on growth of *Quercus ilex*, *Pinus halepensis*, *Pinus pinaster*, *Pinus sylvestris* and *Fagus sylvatica* forests in the Mediterranean region. *For. Ecol. Manage.* 162, 23–37.
- Sabater F, Guasch H, Marti E, Armengol J, Sabater S 1992 The Ter, a Mediterranean river system in Spain. *Limnetica* 8, 141–149.
- Sabatés A, Martín P, Lloret J, Raya V 2006 Sea warming and fish distribution: The case of the small pelagic fish, *Sardinella aurita*, in the western Mediterranean. *Glob. Chang. Biol.* 12, 2209–2219. doi: [10.1111/j.1365-2486.2006.01246.x](https://doi.org/10.1111/j.1365-2486.2006.01246.x)
- Safriel U 2006 *Dryland development, desertification and security in the Mediterranean*. NATO Secur. , eds. Kepner WG, Rubio JL, Mouat DA, Pedrazzini F Springer doi: [10.1007/1-4020-3760-0](https://doi.org/10.1007/1-4020-3760-0)
- Sala E, Ballesteros E, Dendrinis P, Franco A Di, Ferretti F et al. 2012 The Structure of Mediterranean Rocky Reef Ecosystems across Environmental and Human Gradients, and Conservation Implications. *PLoS One* 7, e32742. doi: [10.1371/journal.pone.0032742](https://doi.org/10.1371/journal.pone.0032742)
- Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J et al. 2000 Global biodiversity scenarios for the year 2100. *Science (80-. )*. 287, 1770–1774. doi: [10.1126/science.287.5459.1770](https://doi.org/10.1126/science.287.5459.1770)
- Salameh E, Abdallat G, van der Valk M, Salameh E, Abdallat G et al. 2019 Planning Considerations of Managed Aquifer Recharge (MAR) Projects in Jordan. *Water* 11, 182. doi: [10.3390/w11020182](https://doi.org/10.3390/w11020182)
- Salvati L, Ferrara A, Tombolini I, Gemmiti R, Colantoni A et al. 2015 Desperately Seeking Sustainability: Urban Shrinkage, Land Consumption and Regional Planning in a Mediterranean Metropolitan Area. *Sustainability* 7, 11980–11997. doi: [10.3390/su70911980](https://doi.org/10.3390/su70911980)
- San-Miguel-Ayán J, Moreno JM, Camia A 2013 Analysis of large fires in European Mediterranean landscapes: Lessons learned and perspectives. *For. Ecol. Manage.* 294, 11–22. doi: [10.1016/j.foreco.2012.10.050](https://doi.org/10.1016/j.foreco.2012.10.050)
- Sánchez-Arcilla A, Jiménez JA, Valdemoro HI, Gracia V 2008 Implications of Climatic Change on Spanish Mediterranean Low-Lying Coasts: The Ebro Delta Case. *J. Coast. Res.* 2008, 306–316. <http://www.bioone.org/doi/abs/10.2112/07A-0005.1>
- Sánchez-Pinillos M, Ameztegui A, Kitzberger T, Coll L 2018 Relative size to resprouters determines post-fire recruitment of non-serotinous pines. *For. Ecol. Manage.* 429, 300–307. doi: [10.1016/j.foreco.2018.07.009](https://doi.org/10.1016/j.foreco.2018.07.009)
- Sánchez-Pinillos M, Coll L, de Cáceres M, Ameztegui A 2016 Assessing the persistence capacity of communities facing natural disturbances on the basis of species response traits. *Ecol. Indic.* 66, 76–85. doi: [10.1016/j.ecolind.2016.01.024](https://doi.org/10.1016/j.ecolind.2016.01.024)
- Sánchez-Salguero R, Camarero JJ, Dobbertin M, Fernández-Cancio Á, Vilà-Cabrera A et al. 2013 Contrasting vulnerability and resilience to drought-induced decline of densely planted vs. natural rear-edge *Pinus nigra* forests. *For. Ecol. Manage.* 310, 956–967. doi: [10.1016/j.foreco.2013.09.050](https://doi.org/10.1016/j.foreco.2013.09.050)
- Sanchez A, Abdul Malak D, Guelmami A, Perennou C 2015 Development of an indicator to monitor Mediterranean wetlands. *PLoS One* 10, e0122694. doi: [10.1371/journal.pone.0122694](https://doi.org/10.1371/journal.pone.0122694)
- Santinelli C, Hansell DA, Ribera d'Alcalà M 2013 Influence of stratification on marine dissolved organic carbon (DOC) dynamics: The Mediterranean Sea case. *Prog. Oceanogr.* 119, 68–77. doi: [10.1016/j.pocean.2013.06.001](https://doi.org/10.1016/j.pocean.2013.06.001)
- Santos A, P. Godinho D, Vizinho A, Alves F, Pinho P et al. 2018 Artificial lakes as a climate change adaptation strategy in drylands: evaluating the trade-off on non-target ecosystem services. *Mitig. Adapt. Strateg.*

- Glob. Chang.* 23, 887–906.  
doi: [10.1007/s11027-017-9764-x](https://doi.org/10.1007/s11027-017-9764-x)
- Santos MJ 2010 Encroachment of upland Mediterranean plant species in riparian ecosystems of southern Portugal. *Biodivers. Conserv.* 19, 2667–2684.  
doi: [10.1007/s10531-010-9866-1](https://doi.org/10.1007/s10531-010-9866-1)
- Sanz-Elorza M, Dana ED, González-Moreno A, Sobrino E 2003 Changes in the high-mountain vegetation of the central Iberian Peninsula as a probable sign of global warming. *Ann. Bot.* 92, 273–280.  
doi: [10.1093/aob/mcg130](https://doi.org/10.1093/aob/mcg130)
- Sapes G, Serra-Diaz JM, Lloret F 2017 Species climatic niche explains drought-induced die-off in a Mediterranean woody community. *Ecosphere* 8, 1–15.  
doi: [10.1002/ecs2.1833](https://doi.org/10.1002/ecs2.1833)
- Sarà G, de Pirro M 2011 Heart beat rate adaptations to varying salinity of two intertidal Mediterranean bivalves: The invasive *Brachidontes pharaonis* and the native *Mytilaster minimus*. *Ital. J. Zool.* 78, 193–197.  
doi: [10.1080/11250001003657360](https://doi.org/10.1080/11250001003657360)
- Sarà G, Kearney M, Helmuth B 2011 Combining heat-transfer and energy budget models to predict thermal stress in Mediterranean intertidal mussels. *Chem. Ecol.* 27, 135–145. doi: [10.1080/02757540.2011.552227](https://doi.org/10.1080/02757540.2011.552227)
- Sarà G, Romano C, Widdows J, Staff FJ 2008 Effect of salinity and temperature on feeding physiology and scope for growth of an invasive species (*Brachidontes pharaonis* - Mollusca: Bivalvia) within the Mediterranean sea. *J. Exp. Mar. Bio. Ecol.* 363, 130–136.  
doi: [10.1016/j.jembe.2008.06.030](https://doi.org/10.1016/j.jembe.2008.06.030)
- Sardà R, Pinedo S, Gremare A, Taboada S 2000 Changes in the dynamics of shallow sandy-bottom assemblages due to sand extraction in the Catalan Western Mediterranean Sea. *ICES J. Mar. Sci.* 57, 1446–1453.  
doi: [10.1006/jmsc.2000.0922](https://doi.org/10.1006/jmsc.2000.0922)
- Sarris D, Christodoulakis D, Körner C 2011 Impact of recent climatic change on growth of low elevation eastern Mediterranean forest trees. *Clim. Change* 106, 203–223. doi: [10.1007/s10584-010-9901-y](https://doi.org/10.1007/s10584-010-9901-y)
- Sartoretto S, Schohn T, Bianchi CN, Morri C, Garrabou J et al. 2017 An integrated method to evaluate and monitor the conservation state of coralligenous habitats: The INDEX-COR approach. *Mar. Pollut. Bull.* 120, 222–231.  
doi: [10.1016/j.marpolbul.2017.05.020](https://doi.org/10.1016/j.marpolbul.2017.05.020)
- Satta A, Puddu M, Venturini S, Giupponi C 2017 Assessment of coastal risks to climate change related impacts at the regional scale: The case of the Mediterranean region. *Int. J. Disaster Risk Reduct.* 24, 284–296.  
doi: [10.1016/j.ijdrr.2017.06.018](https://doi.org/10.1016/j.ijdrr.2017.06.018)
- Saunders DL, Meeuwig JJ, Vincent ACJ 2002 Freshwater Protected Areas: Strategies for Conservation. *Conserv. Biol.* 16, 30–41. doi: [10.1046/j.1523-1739.2002.99562.x](https://doi.org/10.1046/j.1523-1739.2002.99562.x)
- Saura S, Bertzy B, Bastin L, Battistella L, Mandrici A et al. 2018 Protected area connectivity: Shortfalls in global targets and country-level priorities. *Biol. Conserv.* 219, 53–67. doi: [10.1016/j.biocon.2017.12.020](https://doi.org/10.1016/j.biocon.2017.12.020)
- Scalco E, Brunet C, Marino F, Rossi R, Soprano V et al. 2012 Growth and toxicity responses of Mediterranean *Ostreopsis cf. ovata* to seasonal irradiance and temperature conditions. *Harmful Algae* 17, 25–34.  
doi: [10.1016/j.hal.2012.02.008](https://doi.org/10.1016/j.hal.2012.02.008)
- Scapini F, Innocenti Degli E, Defeo O 2019 Behavioral adaptations of sandy beach macrofauna in face of climate change impacts: A conceptual framework. *Estuar. Coast. Shelf Sci.* 225, 106236.  
doi: [10.1016/j.ecss.2019.05.018](https://doi.org/10.1016/j.ecss.2019.05.018)
- Schatz B 2017 The orchids species of Cavallo island (Lavezzi archipelago, Corsica): an amazing abundance of the protected species *Gennaria diphylla*. *Ecol. Mediterr.* 43, 159–170.
- Schembri PJ, Deidun A, Mallia A, Mercieca L 2005 Rocky shore biotic assemblages of the Maltese Islands (Central Mediterranean) : a conservation perspective.  
doi: [10.2112/03-0043R.1](https://doi.org/10.2112/03-0043R.1)
- Schewe J, Levermann A 2017 Non-linear intensification of Sahel rainfall as a possible dynamic response to future warming. *Earth Syst. Dyn.* 8, 495–505.  
doi: [10.5194/esd-8-495-2017](https://doi.org/10.5194/esd-8-495-2017)
- Schilling J, Freier KP, Hertig E, Scheffran J 2012 Climate change, vulnerability and adaptation in North Africa with focus on Morocco. *Agric. Ecosyst. Environ.* 156, 12–26. doi: [10.1016/J.AGEE.2012.04.021](https://doi.org/10.1016/J.AGEE.2012.04.021)
- Schilman B, Almogi-Labin A, Bar-Matthews M, Labeyrie L, Paterne M et al. 2001a Long- and short-term carbon fluctuations in the Eastern Mediterranean during the late Holocene. *Geology* 29, 1099–1102.
- Schilman B, Bar-Matthews M, Almogi-Labin A, Luz B 2001b Global climate instability reflected by Eastern Mediterranean marine records during the late Holocene. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 176, 157–176. doi: [10.1016/S0031-0182\(01\)00336-4](https://doi.org/10.1016/S0031-0182(01)00336-4)
- Schils R, Olesen JE, Kersebaum KC, Rijk B, Oberforster M et al. 2018 Cereal yield gaps across Europe. *Eur. J. Agron.* 101, 109–120. doi: [10.1016/j.eja.2018.09.003](https://doi.org/10.1016/j.eja.2018.09.003)
- Schmiedl G, Hemleben C, Keller J, Segl M 2010a Impact of climatic changes on the benthic foraminiferal fauna in the Ionian Sea during the last 330,000 years. *Paleoceanography* 13, 447–458. doi: [10.1029/98PA01864](https://doi.org/10.1029/98PA01864)
- Schmiedl G, Kuhnt T, Ehrmann W, Emeis K-C, Hamann Y et al. 2010b Climatic forcing of eastern Mediterranean deep-water formation and benthic ecosystems during the past 22 000 years. *Quat. Sci. Rev.* 29, 3006–3020.  
doi: [10.1016/j.quascirev.2010.07.002](https://doi.org/10.1016/j.quascirev.2010.07.002)
- Schroeder K, Chiggiato J, Bryden HL, Borghini M, Ben Ismail S 2016 Abrupt climate shift in the Western Mediterranean Sea. *Sci. Rep.* 6. doi: [10.1038/srep23009](https://doi.org/10.1038/srep23009)
- Schubert N, Brown D, Rossi S 2017 Symbiotic Versus Non-symbiotic Octocorals: Physiological and Ecological Implications. *Mar. Anim. For.*, 887–918.  
doi: [10.1007/978-3-319-21012-4\\_54](https://doi.org/10.1007/978-3-319-21012-4_54)
- Schwacke LH, Gulland FM, White S 2013 Sentinel species in oceans and human health, in *Environmental Toxicology. Selected Entries from the Encyclopedia of Sustain-*

- ability Science and Technology, ed. Laws EA (Springer New York), 503–528.  
doi: [10.1007/978-1-4614-5764-0\\_18](https://doi.org/10.1007/978-1-4614-5764-0_18)
- Schwaiger F, Poschenrieder W, Biber P, Pretzsch H 2019 Ecosystem service trade-offs for adaptive forest management. *Ecosyst. Serv.* 39, 100993.  
doi: [10.1016/j.ecoser.2019.100993](https://doi.org/10.1016/j.ecoser.2019.100993)
- Sciandrello S, Tomaselli V 2014 Coastal salt-marshes plant communities of the Salicornietea fruticosae class in Apulia (Italy). *Biologia (Bratisl.)*. 69, 53–69.  
doi: [10.2478/s11756-013-0283-2](https://doi.org/10.2478/s11756-013-0283-2)
- Sciberras M, Rodriguez-Rodriguez D, Ponge B, Jackson E 2013 Criteria for assessing ecological coherence of MPA networks: A review.
- Sedano Vera F, Marquina D, Espinosa Torre F 2014 Usefulness of meiofauna at high taxonomic levels as a tool to assess harbor quality status. Marina del Este Harbor (Granada, Spain) as a case study. *Rev. ciencias Mar. y costeras* 6, 103–113. doi: [10.15359/revmar.6.7](https://doi.org/10.15359/revmar.6.7)
- Seddaiu G, Bagella S, Pulina A, Cappai C, Salis L et al. 2018 Mediterranean cork oak wooded grasslands: synergies and trade-offs between plant diversity, pasture production and soil carbon. *Agrofor. Syst.*  
doi: [10.1007/s10457-018-0225-7](https://doi.org/10.1007/s10457-018-0225-7)
- Seddaiu G, Porcu G, Ledda L, Roggero PP, Agnelli A et al. 2013 Soil organic matter content and composition as influenced by soil management in a semi-arid Mediterranean agro-silvo-pastoral system. *Agric. Ecosyst. Environ.* 167, 1–11. doi: [10.1016/J.AGEE.2013.01.002](https://doi.org/10.1016/J.AGEE.2013.01.002)
- Sefelnasr A, Sherif M 2014 Impacts of Seawater Rise on Seawater Intrusion in the Nile Delta Aquifer, Egypt. *Groundwater* 52, 264–276. doi: [10.1111/gwat.12058](https://doi.org/10.1111/gwat.12058)
- Seidl R, Schelhaas M-J, Rammer W, Verkerk PJ 2014 Increasing forest disturbances in Europe and their impact on carbon storage. *Nat. Clim. Chang.* 4, 806–810.  
doi: [10.1038/nclimate2318](https://doi.org/10.1038/nclimate2318)
- Selina MS, Morozova T V, Vyshkvartsev DI, Orlova TY 2014 Seasonal dynamics and spatial distribution of epi-phytic dinoflagellates in Peter the Great Bay (Sea of Japan) with special emphasis on *Ostreopsis* species. *Harmful Algae* 32, 1–10. doi: [10.1016/j.hal.2013.11.005](https://doi.org/10.1016/j.hal.2013.11.005)
- Serpa D, Nunes JP, Santos J, Sampaio E, Jacinto R et al. 2015 Impacts of climate and land use changes on the hydrological and erosion processes of two contrasting Mediterranean catchments. *Sci. Total Environ.* 538.  
doi: [10.1016/j.scitotenv.2015.08.033](https://doi.org/10.1016/j.scitotenv.2015.08.033)
- Serraj R, Pingali P 2019 *Agriculture & Food Systems to 2050*. World Scientific doi: [10.1142/11212](https://doi.org/10.1142/11212)
- Serrano E, Coma R, Ribes M, Weitzmann B, García M et al. 2013 Rapid Northward Spread of a Zooxanthellate Coral Enhanced by Artificial Structures and Sea Warming in the Western Mediterranean. *PLoS One* 8, e52739. doi: [10.1371/journal.pone.0052739](https://doi.org/10.1371/journal.pone.0052739)
- Settele J, Hammen V, Hulme P, Karlson U, Klotz S et al. 2005 ALARM: assessing large scale environmental Risks for biodiversity with tested methods. *GAIA* 14, 69–72.
- Settele J, Kudrna O, Harpke A, Kühn I, van Swaay C et al. 2008 Climatic Risk Atlas of European Butterflies. *BioRisk* 1, 1–712. doi: [10.3897/biorisk.1](https://doi.org/10.3897/biorisk.1)
- Sghaier YR, Zakhama-Sraieb R, Benamer I, Charfi-Cheikhrouha F 2011 Occurrence of the seagrass *Halophila stipulacea* (Hydrocharitaceae) in the southern Mediterranean Sea. *Bot. Mar.* 54.  
doi: [10.1515/bot.2011.061](https://doi.org/10.1515/bot.2011.061)
- Shaban A 2009 Indicators and aspects of hydrological drought in Lebanon. *Water Resour. Manag.* 23, 1875–1891. doi: [10.1007/s11269-008-9358-1](https://doi.org/10.1007/s11269-008-9358-1)
- Shackelford GE, Kelsey R, Robertson RJ, Williams DR, Dicks L V. 2017 Sustainable Agriculture in California and Mediterranean Climates: Evidence for the effects of selected interventions. Cambridge, UK.  
[https://www.conservationevidence.com/data/index-?synopsis\\_id%5B%5D=22](https://www.conservationevidence.com/data/index-?synopsis_id%5B%5D=22)
- Shakesby RA 2011 Post-wildfire soil erosion in the Mediterranean: Review and future research directions. *Earth-Science Rev.* 105, 71–100.  
doi: [10.1016/j.earscirev.2011.01.001](https://doi.org/10.1016/j.earscirev.2011.01.001)
- Shaltout M, Tonbol K, Omstedt A 2015 Sea-level change and projected future flooding along the Egyptian Mediterranean coast. *Oceanologia* 57, 293–307.  
doi: [10.1016/j.oceano.2015.06.004](https://doi.org/10.1016/j.oceano.2015.06.004)
- Shanks AL, Grantham BA, Carr MH 2003 Propagule dispersal distance and the size and spacing of marine reserves. *Ecol. Appl.* 13, S159–S169.
- Sherif MM, Singh VP 1999 Effect of climate change on sea water intrusion in coastal aquifers. *Hydrol. Process.* 13, 1277–1287. doi: [10.1002/\(sici\)1099-1085\(19990615\)13:8<1277::aid-hyp765>3.3.co;2-n](https://doi.org/10.1002/(sici)1099-1085(19990615)13:8<1277::aid-hyp765>3.3.co;2-n)
- Sicre M-A, Jalali B, Martrat B, Schmidt S, Bassetti M-A et al. 2016 Sea surface temperature variability in the North Western Mediterranean Sea (Gulf of Lion) during the Common Era. *Earth Planet. Sci. Lett.* 456, 124–133. doi: [10.1016/j.epsl.2016.09.032](https://doi.org/10.1016/j.epsl.2016.09.032)
- Sifleet S, Pendleton L, Murray BC 2011 State of the Science on Coastal Blue Carbon A Summary for Policy Makers.
- Silanikove N, Koluman DN 2015 Impact of climate change on the dairy industry in temperate zones: Predications on the overall negative impact and on the positive role of dairy goats in adaptation to earth warming. *Small Rumin. Res.* 123, 27–34.  
doi: [10.1016/j.smallrumres.2014.11.005](https://doi.org/10.1016/j.smallrumres.2014.11.005)
- Šilc U, Dajić-Stevanović Z, Ibraliu A, Luković M, Stešević D 2016 Human impact on sandy beach vegetation along the southeastern Adriatic coast. *Biologia (Bratisl.)*. 71.  
doi: [10.1515/biolog-2016-0111](https://doi.org/10.1515/biolog-2016-0111)
- Silenzi S, Antonioli F, Chemello R 2004 A new marker for sea surface temperature trend during the last centuries in temperate areas: Vermetid reef. *Glob. Planet. Change* 40, 105–114.  
doi: [10.1016/S0921-8181\(03\)00101-2](https://doi.org/10.1016/S0921-8181(03)00101-2)
- Simas T, Nunes JP, Ferreira JG 2001 Effects of global climate change on coastal salt marshes. *Ecol. Modell.*

- 139, 1–15. doi: [10.1016/S0304-3800\(01\)00226-5](https://doi.org/10.1016/S0304-3800(01)00226-5)
- Simon A, Rinaldi M 2006 Disturbance, stream incision, and channel evolution: The roles of excess transport capacity and boundary materials in controlling channel response. *Geomorphology* 79, 361–383. doi: [10.1016/j.geomorph.2006.06.037](https://doi.org/10.1016/j.geomorph.2006.06.037)
- Simonini R, Ansaloni I, Bonvicini Pagliai AM, Cavallini F, Iotti M et al. 2005 The effects of sand extraction on the macrobenthos of a relict sands area (northern Adriatic Sea): results 12 months post-extraction. *Mar. Pollut. Bull.* 50, 768–777. doi: [10.1016/j.marpolbul.2005.02.009](https://doi.org/10.1016/j.marpolbul.2005.02.009)
- Sirami C, Nespoulous A, Cheylan JP, Marty P, Hvenegaard GT et al. 2010 Long-term anthropogenic and ecological dynamics of a Mediterranean landscape: Impacts on multiple taxa. *Landsc. Urban Plan.* 96, 214–223. doi: [10.1016/j.landurbplan.2010.03.007](https://doi.org/10.1016/j.landurbplan.2010.03.007)
- Sisma-Ventura G, Guzner B, Yam R, Fine M, Shemesh A 2009 The reef builder gastropod *Dendropoma petreum* - A proxy of short and long term climatic events in the Eastern Mediterranean. *Geochim. Cosmochim. Acta* 73, 4376–4383. doi: [10.1016/j.gca.2009.04.037](https://doi.org/10.1016/j.gca.2009.04.037)
- Sivapalan M, Takeuchi K, Franks SW, Gupta VK, Karambiri H et al. 2003 IAHS Decade on Predictions in Ungauged Basins (PUB), 2003–2012: Shaping an exciting future for the hydrological sciences. *Hydrol. Sci. J.* 48, 857–880. doi: [10.1623/hysj.48.6.857.51421](https://doi.org/10.1623/hysj.48.6.857.51421)
- Skliris N, Marsh R, Josey SA, Good SA, Liu C et al. 2014 Salinity changes in the World Ocean since 1950 in relation to changing surface freshwater fluxes. *Clim. Dyn.* 43, 709–736. doi: [10.1007/s00382-014-2131-7](https://doi.org/10.1007/s00382-014-2131-7)
- Smayda TJ 2006 Harmful Algal Bloom Communities in Scottish Coastal Waters: Relationships to Fish Farming and Regional. Comparisons – A Review. <http://www.scotland.gov.uk/resource/doc/92174/0022031.pdf>
- Snoussi M, Ouchani T, Niaz S 2008 Vulnerability assessment of the impact of sea-level rise and flooding on the Moroccan coast: The case of the Mediterranean eastern zone. *Estuar. Coast. Shelf Sci.* 77, 206–213. doi: [10.1016/j.ecss.2007.09.024](https://doi.org/10.1016/j.ecss.2007.09.024)
- Sohn JA, Gebhardt T, Ammer C, Bauhus J, Häberle KH et al. 2013 Mitigation of drought by thinning: Short-term and long-term effects on growth and physiological performance of Norway spruce (*Picea abies*). *For. Ecol. Manage.* 308, 188–197. doi: [10.1016/j.foreco.2013.07.048](https://doi.org/10.1016/j.foreco.2013.07.048)
- Sohn JA, Hartig F, Kohler M, Huss J, Bauhus J 2016a Heavy and frequent thinning promotes drought adaptation in *Pinus sylvestris* forests. *Ecol. Appl.* 26, 2190–2205. doi: [10.1002/eap.1373](https://doi.org/10.1002/eap.1373)
- Sohn JA, Saha S, Bauhus J 2016b Potential of forest thinning to mitigate drought stress: A meta-analysis. *For. Ecol. Manage.* 380, 261–273. doi: [10.1016/j.foreco.2016.07.046](https://doi.org/10.1016/j.foreco.2016.07.046)
- Sokos CK, Mamolos AP, Kalburtji KL, Birtsas PK 2013 Farming and wildlife in Mediterranean agroecosystems. *J. Nat. Conserv.* 21, 81–92. doi: [10.1016/j.jnc.2012.11.001](https://doi.org/10.1016/j.jnc.2012.11.001)
- Sowers J, Vengosh A, Weinthal E 2011 Climate change, water resources, and the politics of adaptation in the Middle East and North Africa. *Clim. Change* 104, 599–627. doi: [10.1007/s10584-010-9835-4](https://doi.org/10.1007/s10584-010-9835-4)
- Stagnari F, Maggio A, Galieni A, Pisante M 2017 Multiple benefits of legumes for agriculture sustainability: an overview. *Chem. Biol. Technol. Agric.* 4, 2. doi: [10.1186/s40538-016-0085-1](https://doi.org/10.1186/s40538-016-0085-1)
- Stanhill G, Kurtzman D, Rosa R 2015 Estimating desalination requirements in semi-arid climates: A Mediterranean case study. *Desalination* 355, 118–123. doi: [10.1016/J.DESAL.2014.10.035](https://doi.org/10.1016/J.DESAL.2014.10.035)
- Stanisci A, Pelino G, Blasi C 2005 Vascular plant diversity and climate change in the alpine belt of the central Apennines (Italy). *Biodivers. Conserv.* 14, 1301–1318. doi: [10.1007/s10531-004-9674-6](https://doi.org/10.1007/s10531-004-9674-6)
- Stefanaki A, Kantsa A, Tscheulin T, Charitonidou M, Petanidou T 2015 Lessons from Red Data Books: Plant Vulnerability Increases with Floral Complexity. *PLoS One* 10, e0138414. doi: [10.1371/journal.pone.0138414](https://doi.org/10.1371/journal.pone.0138414)
- Stellmes M, Röder A, Udelhoven T, Hill J 2013 Mapping syndromes of land change in Spain with remote sensing time series, demographic and climatic data. *Land use policy* 30, 685–702. doi: [10.1016/j.landusepol.2012.05.007](https://doi.org/10.1016/j.landusepol.2012.05.007)
- Stergiou KI, Somarakis S, Triantafyllou G, Tsiaras KP, Giannoulaki M et al. 2016 Trends in productivity and biomass yields in the Mediterranean Sea Large Marine Ecosystem during climate change. *Environ. Dev.* 17, 57–74. doi: [10.1016/j.envdev.2015.09.001](https://doi.org/10.1016/j.envdev.2015.09.001)
- Stevens CJ, Dupr C, Dorland E, Gaudnik C, Gowing DJG et al. 2011 The impact of nitrogen deposition on acid grasslands in the Atlantic region of Europe. *Environ. Pollut.* 159, 2243–2250. doi: [10.1016/j.envpol.2010.11.026](https://doi.org/10.1016/j.envpol.2010.11.026)
- Stoate C, Báldi A, Beja P, Boatman ND, Herzon I et al. 2009 Ecological impacts of early 21st century agricultural change in Europe – A review. *J. Environ. Manage.* 91, 22–46. doi: [10.1016/j.jenvman.2009.07.005](https://doi.org/10.1016/j.jenvman.2009.07.005)
- Stoate C, Boatman ND, Borralho RJ, Carvalho CR, Snoo GR d. et al. 2001 Ecological impacts of arable intensification in Europe. *J. Environ. Manage.* 63, 337–365. doi: [10.1006/jema.2001.0473](https://doi.org/10.1006/jema.2001.0473)
- Storelli MM, Marcotrigiano GO 2005 Bioindicator organisms: Heavy metal pollution evaluation in the Ionian Sea (Mediterranean Sea-Italy). *Environ. Monit. Assess.* 102, 159–166. doi: [10.1007/s10661-005-6018-2](https://doi.org/10.1007/s10661-005-6018-2)
- Strefitaris N, Zenetos A 2006 Alien Marine Species in the Mediterranean - the 100 'Worst Invasives' and their Impact. *Mediterr. Mar. Sci.* 7, 87–118. doi: [10.12681/mms.180](https://doi.org/10.12681/mms.180)
- Stromberg JC, Boudell JA, Hazelton AF 2008 Differences in seed mass between hydric and xeric plants influence seed bank dynamics in a dryland riparian ecosystem.

- Funct. Ecol.* 22, 205–212.  
doi: [10.1111/j.1365-2435.2007.01375.x](https://doi.org/10.1111/j.1365-2435.2007.01375.x)
- Suaria G, Avio CG, Mineo A, Lattin GL, Magaldi MG et al. 2016 The Mediterranean plastic soup: synthetic polymers in Mediterranean surface waters. *Sci. Rep.* 6, 37551. doi: [10.1038/srep37551](https://doi.org/10.1038/srep37551)
- Sugden HE, Underwood AJ, Hawkins SJ 2009 The aesthetic value of littoral hard substrata and consideration of ethical frameworks for their investigation and conservation, in *Marine Hard Bottom Communities*, ed. Wahl M (Springer-Verlag), 409–421.  
<https://eprints.soton.ac.uk/187905/>
- Sweetman AK, Thurber AR, Smith CR, Levin LA, Mora C et al. 2017 Major impacts of climate change on deep-sea benthic ecosystems. *Elem Sci Anth* 5, 4.  
doi: [10.1525/elementa.203](https://doi.org/10.1525/elementa.203)
- Tanasijevic L, Todorovic M, Pereira LS, Pizzigalli C, Lionello P 2014 Impacts of climate change on olive crop evapotranspiration and irrigation requirements in the Mediterranean region. *Agric. Water Manag.* 144, 54–68.  
doi: [10.1016/j.agwat.2014.05.019](https://doi.org/10.1016/j.agwat.2014.05.019)
- Tardy V, Spor A, Mathieu O, Lévêque J, Terrat S et al. 2015 Shifts in microbial diversity through land use intensity as drivers of carbon mineralization in soil. *Soil Biol. Biochem.* 90, 204–213.  
doi: [10.1016/j.soilbio.2015.08.010](https://doi.org/10.1016/j.soilbio.2015.08.010)
- Tarjuelo JM, Rodríguez-Díaz JA, Abadía R, Camacho E, Rocamora C et al. 2015 Efficient water and energy use in irrigation modernization: Lessons from Spanish case studies. *Agric. Water Manag.* 162, 67–77.  
doi: [10.1016/j.agwat.2015.08.009](https://doi.org/10.1016/j.agwat.2015.08.009)
- Tasser E, Leitinger G, Tappeiner U 2017 Climate change versus land-use change—What affects the mountain landscapes more? *Land use policy* 60, 60–72.  
doi: [10.1016/j.landusepol.2016.10.019](https://doi.org/10.1016/j.landusepol.2016.10.019)
- Taviani M, Angeletti L, Beuck L, Campiani E, Canese S et al. 2016 On and Off the Beaten Track: Megafaunal Sessile Life and Adriatic Cascading Processes. *Mar. Geol.* 375, 146–160. doi: [10.1016/j.margeo.2015.10.003](https://doi.org/10.1016/j.margeo.2015.10.003)
- Teixidó N, Casas E, Cebrián E, Linares C, Garrabou J 2013 Impacts on Coralligenous Outcrop Biodiversity of a Dramatic Coastal Storm. *PLoS One* 8, e53742.  
doi: [10.1371/journal.pone.0053742](https://doi.org/10.1371/journal.pone.0053742)
- Telesca L, Belluscio A, Criscoli A, Ardizzone G, Apostolaki ET et al. 2015 Seagrass meadows (*Posidonia oceanica*) distribution and trajectories of change. *Sci. Rep.* 5.  
doi: [10.1038/srep12505](https://doi.org/10.1038/srep12505)
- Ternon E, Guieu C, Ridame C, L'Helguen S, Catala P 2011 Longitudinal variability of the biogeochemical role of Mediterranean aerosols in the Mediterranean Sea. *Biogeosciences* 8, 1067–1080.  
doi: [10.5194/bg-8-1067-2011](https://doi.org/10.5194/bg-8-1067-2011)
- Terrado M, Acuña V, Ennaanay D, Tallis H, Sabater S 2014 Impact of climate extremes on hydrological ecosystem services in a heavily humanized Mediterranean basin. *Ecol. Indic.* 37, 199–209.  
doi: [10.1016/j.ecolind.2013.01.016](https://doi.org/10.1016/j.ecolind.2013.01.016)
- The MerMex Group, de Madron XD, Guieu C, Sempéré R, Conan P et al. 2011 Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean. *Prog. Oceanogr.* 91, 97–166.  
doi: [10.1016/j.pocean.2011.02.003](https://doi.org/10.1016/j.pocean.2011.02.003)
- Theocharis A, Klein B, Nittis K, Roether W 2002 Evolution and status of the Eastern Mediterranean Transient (1997–1999). *J. Mar. Syst.* 33–34, 91–116.  
doi: [10.1016/S0924-7963\(02\)00054-4](https://doi.org/10.1016/S0924-7963(02)00054-4)
- Thiébaud S, Moatti J-P, Annesi-Maesano I, Aumeerud-Thomas Y, Barouki R et al. 2016 *The Mediterranean Region under Climate Change: A scientific update*. Marseille, France: Institut de Recherche pour le Développement <http://www.editions.ird.fr/produit/433/9782709922197/The-Mediterranean-Region-under-Climate-Change>
- Thomas OP, Sarrazin V, Ivanisevic J, Amade P, Pérez T 2007 Sponge chemical defenses in stress conditions: The case study of the last disease outbreak observed in the NW Mediterranean. in *Proceedings of the 5th European Conference on Marine Natural Products; 16–21 September 2007* (Ischia, Italy), 61.
- Thompson JD 2005 *Plant evolution in the Mediterranean*. Oxford Uni. Oxford, UK.
- Thorndycraft VR, Benito G 2006 The Holocene fluvial chronology of Spain: evidence from a newly compiled radiocarbon database. *Quat. Sci. Rev.* 25, 223–234.  
doi: [10.1016/j.quascirev.2005.07.003](https://doi.org/10.1016/j.quascirev.2005.07.003)
- Thuiller W, Lavorel S, Araújo MB, Sykes MT, Prentice IC 2005 Climate change threats to plant diversity in Europe. *Proc. Natl. Acad. Sci. U. S. A.* 102, 8245–8250.  
doi: [10.1073/pnas.0409902102](https://doi.org/10.1073/pnas.0409902102)
- Tobarra MA, López LA, Cadarso MA, Gómez N, Cazarro I 2018 Is Seasonal Households' Consumption Good for the Nexus Carbon/Water Footprint? The Spanish Fruits and Vegetables Case. *Environ. Sci. Technol.* 52, 12066–12077. doi: [10.1021/acs.est.8b00221](https://doi.org/10.1021/acs.est.8b00221)
- Todd VLG, Todd IB, Gardiner JC, Morrin ECN, MacPherson NA et al. 2015 A review of impacts of marine dredging activities on marine mammals. *ICES J. Mar. Sci.* 72, 328–340. doi: [10.1093/icesjms/fsu187](https://doi.org/10.1093/icesjms/fsu187)
- Tomas J, Guitart R, Mateo R, Raga JA 2002 Marine debris ingestion in loggerhead sea turtles, *Caretta caretta*, from the Western Mediterranean. *Mar. Pollut. Bull.* 44, 211–216. doi: [10.1016/S0025-326X\(01\)00236-3](https://doi.org/10.1016/S0025-326X(01)00236-3)
- Torralba M, Fagerholm N, Burgess PJ, Moreno G, Plieninger T 2016 Do European agroforestry systems enhance biodiversity and ecosystem services? A meta-analysis. *Agric. Ecosyst. Environ.* 230, 150–161.  
doi: [10.1016/j.agee.2016.06.002](https://doi.org/10.1016/j.agee.2016.06.002)
- Toseland A, Daines SJ, Clark JR, Kirkham A, Strauss J et al. 2013 The impact of temperature on marine phytoplankton resource allocation and metabolism. *Nat. Clim. Chang.* 3, 979–984. doi: [10.1038/nclimate1989](https://doi.org/10.1038/nclimate1989)
- Totti C, Romagnoli T, Accoroni S, Coluccelli A, Pellegrini M et al. 2019 Phytoplankton communities in the north-western Adriatic Sea: Interdecadal variability over a

- 30-years period (1988–2016) and relationships with meteorological drivers. *J. Mar. Syst.* 193, 137–153. doi: [10.1016/j.jmarsys.2019.01.007](https://doi.org/10.1016/j.jmarsys.2019.01.007)
- Tovar-Sánchez A, Sánchez-Quiles D, Rodríguez-Romero A 2019 Massive coastal tourism influx to the Mediterranean Sea: The environmental risk of sunscreens. *Sci. Total Environ.* 656, 316–321. doi: [10.1016/j.scitotenv.2018.11.399](https://doi.org/10.1016/j.scitotenv.2018.11.399)
- Townsend DW, Cammen LM, Holligan PM, Campbell DE, Pettigrew NR 1994 Causes and consequences of variability in the timing of spring phytoplankton blooms. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 41, 747–765.
- Traveset A, Riera N 2005 Disruption of a Plant-Lizard Seed Dispersal System and Its Ecological Effects on a Threatened Endemic Plant in the Balearic Islands. *Conserv. Biol.* 19, 421–431. doi: [10.1111/j.1523-1739.2005.00019.x](https://doi.org/10.1111/j.1523-1739.2005.00019.x)
- Tsikliras AC 2008 Climate-related geographic shift and sudden population increase of a small pelagic fish (*Sardinella aurita*) in the eastern Mediterranean Sea. *Mar. Biol. Res.* 4, 477–481. doi: [10.1080/17451000802291292](https://doi.org/10.1080/17451000802291292)
- Tsounis G, Rossi S, Grigg R, Santangelo G, Bramanti L et al. 2010 The Exploitation and Conservation of Precious Corals. doi: [10.1201/EBK1439821169-c3](https://doi.org/10.1201/EBK1439821169-c3)
- Tunin-Ley A, Ibañez F, Labat J-P, Zingone A, Lemée R 2009 Phytoplankton biodiversity and NW Mediterranean Sea warming: Changes in the dinoflagellate genus *Ceratium* in the 20<sup>th</sup> century. *Mar. Ecol. Prog. Ser.* 375, 85–99. doi: [10.3354/meps07730](https://doi.org/10.3354/meps07730)
- Tunin-Ley A, Lemée R 2013 The genus *Neoceratium* (planktonic dinoflagellates) as a potential indicator of ocean warming. *Microorganisms* 1, 58–70. doi: [10.3390/microorganisms1010058](https://doi.org/10.3390/microorganisms1010058)
- Turco M, Jerez S, Doblas-Reyes FJ, AghaKouchak A, Llasat MC et al. 2018a Skilful forecasting of global fire activity using seasonal climate predictions. *Nat. Commun.* 9, 2718. doi: [10.1038/s41467-018-05250-0](https://doi.org/10.1038/s41467-018-05250-0)
- Turco M, Rosa-Cánovas JJ, Bedía J, Jerez S, Montávez JP et al. 2018b Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models. *Nat. Commun.* 9, 3821. doi: [10.1038/s41467-018-06358-z](https://doi.org/10.1038/s41467-018-06358-z)
- Turhan E, Zografos C, Kallis G 2015 Adaptation as biopolitics: Why state policies in Turkey do not reduce the vulnerability of seasonal agricultural workers to climate change. *Glob. Environ. Chang.* 31, 296–306. doi: [10.1016/j.gloenvcha.2015.02.003](https://doi.org/10.1016/j.gloenvcha.2015.02.003)
- Turki S, Dhib A, Fertouna-Bellakhal M, Frossard V, Balti N et al. 2014 Harmful algal blooms (HABs) associated with phycotoxins in shellfish: What can be learned from five years of monitoring in Bizerte Lagoon (Southern Mediterranean Sea)? *Ecol. Eng.* 67, 39–47. doi: [10.1016/j.ecoleng.2014.03.028](https://doi.org/10.1016/j.ecoleng.2014.03.028)
- Turki S, Harzallah A, Sammari C 2006 Occurrence of harmful dinoflagellates in two different Tunisian ecosystems: the lake of Bizerte and the gulf of Gabes. *Cah. Biol. Mar.* 47, 253–259.
- Tzanopoulos J, Vogiatzakis IN 2011 Processes and patterns of landscape change on a small Aegean island: The case of Sifnos, Greece. *Landsc. Urban Plan.* 99, 58–64. doi: [10.1016/j.landurbplan.2010.08.014](https://doi.org/10.1016/j.landurbplan.2010.08.014)
- Tzatzanis M, Wrbka T, Sauberer N 2003 Landscape and vegetation responses to human impact in sandy coasts of Western Crete, Greece. *J. Nat. Conserv.* 11, 187–195. doi: [10.1078/1617-1381-00047](https://doi.org/10.1078/1617-1381-00047)
- UN 2013 World Population Prospects: The 2012 Revision, Highlights and Advance Tables.
- Underwood EC, Viers JH, Klausmeyer KR, Cox RL, Shaw MR 2009 Threats and biodiversity in the Mediterranean biome. *Divers. Distrib.* 15, 188–197. doi: [10.1111/j.1472-4642.2008.00518.x](https://doi.org/10.1111/j.1472-4642.2008.00518.x)
- UNEP-MAP-RAC/SPA 2010 Impact of climate change on marine and coastal biodiversity in the Mediterranean Sea: Current state of knowledge. Tunis
- UNEP/MAP-RAC/SPA 2009 Sub-regional report on vulnerability and impacts of climate change on marine and coastal biological diversity in the Mediterranean Arab countries. <https://wedocs.unep.org/handle/20.500.11822/1264>
- UNEP/MAP/PAP 2008 Protocol on integrated coastal zone management in the Mediterranean. Split.
- UNEP/MAP/PAP 2015 Guidelines for Adapting to Climate Variability and Change along the Mediterranean Coast. Split.
- UNEP/MAP 2012 State of the Mediterranean Marine and Coastal Environment. UNEP/MAP – Barcelona Convention, Athens.
- UNEP/MAP 2016 Background document to the Regional Climate Change Adaptation Framework.
- Uys MC, O’Keeffe JH 1997 Simple words and fuzzy zones: early directions for temporary river research in South Africa. *Environ. Manage.* 21, 517–531. doi: [10.1007/s002679900047](https://doi.org/10.1007/s002679900047)
- Väänänen PJ, Osem Y, Cohen S, Grünzweig JM 2020 Differential drought resistance strategies of co-existing woodland species enduring the long rainless Eastern Mediterranean summer. *Tree Physiol.* 40, 305–320. doi: [10.1093/TREEPHYS/TPZ130](https://doi.org/10.1093/TREEPHYS/TPZ130)
- Vacchi M, Marriner N, Morhange C, Spada G, Fontana A et al. 2016 Multiproxy assessment of Holocene relative sea-level changes in the western Mediterranean: Sea-level variability and improvements in the definition of the isostatic signal. *Earth-Science Rev.* 155, 172–197. doi: [10.1016/j.earscirev.2016.02.002](https://doi.org/10.1016/j.earscirev.2016.02.002)
- Vadrucci M, Sabetta L, Fiocca A, Mazziotti C, Silvestri C et al. 2008 Statistical evaluation of differences in phytoplankton richness and abundance as constrained by environmental drivers in transitional waters of the Mediterranean basin. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 18, S88–S104. doi: [10.1002/aqc.951](https://doi.org/10.1002/aqc.951)
- Vafeidis AT, Neumann B, Zimmermann J, Nicholls RJ 2011 MR9: Analysis of land area and population in the low-elevation coastal zone (LECZ).

- <https://eprints.soton.ac.uk/207617/>
- Valbuena-Carabaña M, de Heredia UL, Fuentes-Utrilla P, González-Doncel I, Gil L 2010 Historical and recent changes in the Spanish forests: A socio-economic process. *Rev. Palaeobot. Palynol.* 162, 492–506. doi: [10.1016/j.revpalbo.2009.11.003](https://doi.org/10.1016/j.revpalbo.2009.11.003)
- Valladares F, Benavides R, Rabasa SG, Díaz M, Pausas JG et al. 2014a Global change and Mediterranean forests: current impacts and potential responses, in *Forests and Global Change*, eds. Coomes DA, Burslem DFRP, Simonson WD (Cambridge, UK: Cambridge University Press), 47–76.
- Valladares F, Matesanz S, Guilhaumon F, Araújo MB, Balaguer L et al. 2014b The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecol. Lett.* 17, 1351–1364. doi: [10.1111/ele.12348](https://doi.org/10.1111/ele.12348)
- Vallejo R, Alloza JA 1998 The restoration of burned lands: the case of eastern Spain, in *Large forest fires*, ed. Moreno JM (Leiden, The Netherlands: Backhuys Publishers), 91–108.
- Valls A, Coll M, Christensen V 2015 Keystone species: toward an operational concept for marine biodiversity conservation. *Ecol. Monogr.* 85, 29–47. doi: [10.1890/14-0306.1](https://doi.org/10.1890/14-0306.1)
- Van Auken OW 2000 Shrub Invasions of North American Semiarid Grasslands. *Annu. Rev. Ecol. Syst.* 31, 197–215. doi: [10.1146/annurev.ecolsys.31.1.197](https://doi.org/10.1146/annurev.ecolsys.31.1.197)
- Van den Broeck M, Waterkeyn A, Rhazi L, Grillas P, Brendonck L 2015 Assessing the ecological integrity of endorheic wetlands, with focus on Mediterranean temporary ponds. *Ecol. Indic.* 54, 1–11. doi: [10.1016/j.ecolind.2015.02.016](https://doi.org/10.1016/j.ecolind.2015.02.016)
- van Franeker JA, Blaize C, Danielsen J, Fairclough K, Gollan J et al. 2011 Monitoring plastic ingestion by the northern fulmar *Fulmarus glacialis* in the North Sea. *Environ. Pollut.* 159, 2609–2615. doi: [10.1016/j.envpol.2011.06.008](https://doi.org/10.1016/j.envpol.2011.06.008)
- Van Harten D 1987 Ostracodes and the early Holocene anoxic event in the Eastern Mediterranean — Evidence and implications. *Mar. Geol.* 75, 263–269. doi: [10.1016/0025-3227\(87\)90108-3](https://doi.org/10.1016/0025-3227(87)90108-3)
- van Leeuwen C, Darriet P 2016 The Impact of Climate Change on Viticulture and Wine Quality. *J. Wine Econ.* 11, 150–167. doi: [10.1017/jwe.2015.21](https://doi.org/10.1017/jwe.2015.21)
- Van Steeter MM, Pitlick J 1998 Geomorphology and endangered fish habitats of the upper Colorado River: 1. Historic changes in streamflow, sediment load, and channel morphology. *Water Resour. Res.* 34, 287–302. doi: [10.1029/97wr02766](https://doi.org/10.1029/97wr02766)
- Vaquer-Sunyer R, Duarte CM 2013 Experimental Evaluation of the Response of Coastal Mediterranean Planktonic and Benthic Metabolism to Warming. *Estuaries and Coasts* 36, 697–707. doi: [10.1007/s12237-013-9595-2](https://doi.org/10.1007/s12237-013-9595-2)
- Varela-Ortega C, Blanco-Gutiérrez I, Swartz CH, Downing TE 2011 Balancing groundwater conservation and rural livelihoods under water and climate uncertainties: An integrated hydro-economic modeling framework. *Glob. Environ. Chang.* 21, 604–619. doi: [10.1016/j.gloenvcha.2010.12.001](https://doi.org/10.1016/j.gloenvcha.2010.12.001)
- Varela MR, Patricio AR, Anderson K, Broderick AC, DeBell L et al. 2019 Assessing climate change associated sea-level rise impacts on sea turtle nesting beaches using drones, photogrammetry and a novel GPS system. *Glob. Chang. Biol.* 25, 753–762. doi: [10.1111/gcb.14526](https://doi.org/10.1111/gcb.14526)
- Vargas-Yáñez M, Zunino P, Benali A, Delpy M, Pastre F et al. 2010 How much is the western Mediterranean really warming and salting? *JGR Atmos.* 115, C04001. doi: [10.1029/2009jc005816](https://doi.org/10.1029/2009jc005816)
- Vázquez-Luis M, Álvarez E, Barrajón A, García-March JR, Grau A et al. 2017 S.O.S. *Pinna nobilis*: A mass mortality event in Western Mediterranean Sea. *Front. Mar. Sci.* 4. doi: [10.3389/fmars.2017.00220](https://doi.org/10.3389/fmars.2017.00220)
- Velasco Ayuso S, Giraldo Silva A, Nelson C, Barger NN, Garcia-Pichel F 2017 Microbial Nursery Production of High-Quality Biological Soil Crust Biomass for Restoration of Degraded Dryland Soils. *Appl. Environ. Microbiol.* 83, e02179-16. doi: [10.1128/AEM.02179-16](https://doi.org/10.1128/AEM.02179-16)
- Velegrakis AF, Trygonis V, Chatzipavlis AE, Karambas T, Vousdoukas MI et al. 2016 Shoreline variability of an urban beach fronted by a beachrock reef from video imagery. *Nat. Hazards* 83, 201–222. doi: [10.1007/s11069-016-2415-9](https://doi.org/10.1007/s11069-016-2415-9)
- Verdura J, Linares C, Ballesteros E, Coma R, Uriz MJ et al. 2019 Biodiversity loss in a Mediterranean ecosystem due to an extreme warming event unveils the role of an engineering gorgonian species. *Sci. Rep.* 9. doi: [10.1038/s41598-019-41929-0](https://doi.org/10.1038/s41598-019-41929-0)
- Vergés A, Steinberg PD, Hay ME, Poore AGB, Campbell AH et al. 2014 The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B Biol. Sci.* 281, 20140846. doi: [10.1098/rspb.2014.0846](https://doi.org/10.1098/rspb.2014.0846)
- Vericat D, Batalla RJ 2006 Sediment transport in a large impounded river: The lower Ebro, NE Iberian Peninsula. *Geomorphology* 79, 72–92. doi: [10.1016/j.geomorph.2005.09.017](https://doi.org/10.1016/j.geomorph.2005.09.017)
- Verlaque M, Afonso-Carrillo J, Candelaria Gil-Rodríguez M, Durand C, Boudouresque CF et al. 2004 Blitzkrieg in a marine invasion: *Caulerpa racemosa* var. *cylindracea* (bryopsidales, chlorophyta) reaches the Canary Islands (north-east Atlantic). *Biol. Invasions* 6, 269–281. doi: [10.1023/B:BINV.0000034589.18347.d3](https://doi.org/10.1023/B:BINV.0000034589.18347.d3)
- Verlaque M, Durand C, Huisman JM, Boudouresque C-F, Le Parco Y 2003 On the identity and origin of the Mediterranean invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta). *Eur. J. Phycol.* 38, 325–339. doi: [10.1080/09670260310001612592](https://doi.org/10.1080/09670260310001612592)
- Verlaque M, Ruitton S, Mineur F, Boudouresque CF 2015 *CIESM Atlas of exotic species in the Mediterranean. Vol. 4. Macrophytes.*, ed. Briand F Monaco: CIESM Publishers.



- Vermeulen SJ, Challinor AJ, Thornton PK, Campbell BM, Eriyagama N et al. 2013 Addressing uncertainty in adaptation planning for agriculture. *Proc. Natl. Acad. Sci. U. S. A.* 110, 8357–8362. doi: [10.1073/pnas.1219441110](https://doi.org/10.1073/pnas.1219441110)
- Vermeulen SJ, Dinesh D, Howden SM, Cramer L, Thornton PK 2018 Transformation in Practice: A Review of Empirical Cases of Transformational Adaptation in Agriculture Under Climate Change. *Front. Sustain. Food Syst.* 2, 65. doi: [10.3389/fsufs.2018.00065](https://doi.org/10.3389/fsufs.2018.00065)
- Versteegh GJM, de Leeuw JW, Taricco C, Romero A 2007 Temperature and productivity influences on  $U_{37}^{K'}$  and their possible relation to solar forcing of the Mediterranean winter. *Geochemistry, Geophys. Geosystems* 8. doi: [10.1029/2006GC001543](https://doi.org/10.1029/2006GC001543)
- Vesperinas ES, Moreno AG, Elorza MS, Sánchez ED, Mata DS et al. 2001 The expansion of thermophilic plants in the Iberian Peninsula as a sign of climatic change, in “Fingerprints” of Climate Change, eds. Walther GR, Burga CA, Edwards PJ (Boston, MA: Springer), 163–184. doi: [10.1007/978-1-4419-8692-4](https://doi.org/10.1007/978-1-4419-8692-4)
- Vezzulli L, Colwell RR, Pruzzo C 2013 Ocean Warming and Spread of Pathogenic Vibrios in the Aquatic Environment. *Microb. Ecol.* 65, 817–825. doi: [10.1007/s00248-012-0163-2](https://doi.org/10.1007/s00248-012-0163-2)
- Vicente-Serrano SM, Lasanta T, Romo A 2004 Analysis of Spatial and Temporal Evolution of Vegetation Cover in the Spanish Central Pyrenees: Role of Human Management. *Environ. Manage.* 34, 802–818. doi: [10.1007/s00267-003-0022-5](https://doi.org/10.1007/s00267-003-0022-5)
- Vicente C, Espada M, Vieira P, Mota M 2012 Pine Wilt Disease: A threat to European forestry. *Eur. J. Plant Pathol.* 133, 89–99. doi: [10.1007/s10658-011-9924-x](https://doi.org/10.1007/s10658-011-9924-x)
- Viedma O, Moreno JM, Rieiro I 2006 Interactions between land use/land cover change, forest fires and landscape structure in Sierra de Gredos (Central Spain). *Environ. Conserv.* 33, 212–222. doi: [10.1017/S0376892906003122](https://doi.org/10.1017/S0376892906003122)
- Viegas DX 1998 Forest fire propagation. *Philos. Trans. R. Soc. London. Ser. A Math. Phys. Eng. Sci.* 356, 2907–2928. doi: [10.1098/rsta.1998.0303](https://doi.org/10.1098/rsta.1998.0303)
- Vilà-Cabrera A, Coll L, Martínez-Vilalta J, Retana J 2018 Forest management for adaptation to climate change in the Mediterranean basin: A synthesis of evidence. *For. Ecol. Manage.* 407, 16–22. doi: [10.1016/j.foreco.2017.10.021](https://doi.org/10.1016/j.foreco.2017.10.021)
- Vila M, Abós-Herrándiz R, Isern-Fontanet J, Álvarez J, Berdalet E 2016 Establishing the link between *Ostreopsis* cf. *ovata* blooms and human health impacts using ecology and epidemiology. *Sci. Mar.* 80, 107–115. doi: [10.3989/scimar.04395.08A](https://doi.org/10.3989/scimar.04395.08A)
- Vila M, Garcés E, Masó M 2001 Potentially toxic epiphytic dinoflagellate assemblages on macroalgae in the NW Mediterranean. *Aquat. Microb. Ecol.* 26, 51–60. doi: [10.3354/ame026051](https://doi.org/10.3354/ame026051)
- Viladrich N, Bramanti L, Tsounis G, Chocarro B, Martínez-Quitana A et al. 2016 Variation in lipid and free fatty acid content during spawning in two temperate octocorals with different reproductive strategies: surface versus internal brooder. *Coral Reefs* 35, 1033–1045. doi: [10.1007/s00338-016-1440-1](https://doi.org/10.1007/s00338-016-1440-1)
- Viladrich N, Bramanti L, Tsounis G, Martínez-Quitana A, Ferrier-Pagès C et al. 2017 Variation of lipid and free fatty acid contents during larval release in two temperate octocorals according to their trophic strategy. *Mar. Ecol. Prog. Ser.* 573, 117–128. doi: [10.3354/meps12141](https://doi.org/10.3354/meps12141)
- Vimal R, Geniaux G, Pluvinet P, Napoléone C, Lepart J 2012 Detecting threatened biodiversity by urbanization at regional and local scales using an urban sprawl simulation approach: Application on the French Mediterranean region. *Landsc. Urban Plan.* 104, 343–355. doi: [10.1016/j.landurbplan.2011.11.003](https://doi.org/10.1016/j.landurbplan.2011.11.003)
- Visciano P, Schirone M, Berti M, Milandri A, Tofalo R et al. 2016 Marine Biotoxins: Occurrence, Toxicity, Regulatory Limits and Reference Methods. *Front. Microbiol.* 7. doi: [10.3389/fmicb.2016.01051](https://doi.org/10.3389/fmicb.2016.01051)
- Visconti P, Elias V, Sousa Pinto I, Fischer M, Ali-Zade V et al. 2018 Status, trends and future dynamics of biodiversity and ecosystems underpinning nature’s contributions to people., in *The IPBES regional assessment report on biodiversity and ecosystem services for Europe and Central Asia*, ed. Rounsevell, M., Fischer, M., Torre-Marín Rando, A., Mader A (Bonn, Germany), 187–382.
- Visser ME, Both C, Lambrechts MM 2004 Global Climate Change Leads to Mistimed Avian Reproduction. *Adv. Ecol. Res.* 35, 89–110. doi: [10.1016/S0065-2504\(04\)35005-1](https://doi.org/10.1016/S0065-2504(04)35005-1)
- Vlamiš A, Katikou P 2015 Human impact in Mediterranean coastal ecosystems and climate change: emerging toxins, in *Climate Change and Marine and Freshwater Toxins*, eds. Botana LM, Louzao C, Vilariño N (Walter de Gruyter GmbH), 237–269.
- Vogiatzakis IN 2012 *Mediterranean Mountain Environments*. John Wiley & Sons, Ltd doi: [10.1002/9781119941156](https://doi.org/10.1002/9781119941156)
- Vogiatzakis IN, Griffiths GH 2008 Island Biogeography and Landscape Ecology, in *Mediterranean island landscapes: natural and cultural approaches*, eds. Vogiatzakis IN, Pungetti G, Mannion A (Dordrecht, Netherlands: Springer), 61–81.
- Vogiatzakis IN, Mannion AM, Sarris D 2016 Mediterranean island biodiversity and climate change: the last 10,000 years and the future. *Biodivers. Conserv.* 25, 2597–2627. doi: [10.1007/s10531-016-1204-9](https://doi.org/10.1007/s10531-016-1204-9)
- Vogiatzakis IN, Pungetti G, Mannion A 2008 *Mediterranean island landscapes: natural and cultural approaches*. Dordrecht: Springer.
- Vogt-Schilb H, Pradel R, Geniez P, Hugot L, Delage A et al. 2016 Responses of orchids to habitat change in Corsica over 27 years. *Ann. Bot.* 118, 115–123. doi: [10.1093/aob/mcw070](https://doi.org/10.1093/aob/mcw070)
- Vohník M, Borovec O, Kolařík M 2016 Communities of cultivable root mycobionts of the seagrass *Posidonia oceanica* in the northwest Mediterranean Sea are

- dominated by a hitherto undescribed Pleosporalean dark septate endophyte. *Microb. Ecol.* 71, 442–451. doi: [10.1007/s00248-015-0640-5](https://doi.org/10.1007/s00248-015-0640-5)
- Volpe G, Nardelli BB, Cipollini P, Santoleri R, Robinson IS 2012 Seasonal to interannual phytoplankton response to physical processes in the Mediterranean Sea from satellite observations. *Remote Sens. Environ.* 117, 223–235.
- Volpi I, Bosco S, Nasso Di Nasso N, Triana F, Roncucci N et al. 2016 Nitrous oxide emissions from clover in the Mediterranean environment. *Ital. J. Agron.* 11, 133. doi: [10.4081/ija.2016.728](https://doi.org/10.4081/ija.2016.728)
- Vousdoukas MI, Voukouvalas E, Annunziato A, Giardino A, Feyen L 2016 Projections of extreme storm surge levels along Europe. *Clim. Dyn.* 47, 3171–3190. doi: [10.1007/s00382-016-3019-5](https://doi.org/10.1007/s00382-016-3019-5)
- Wacquant JP 1990 Biogeographical and physiological aspects of the invasion by *Dittrichia* (ex-*Inula*) *viscosa* W. Greuter, a ruderal species in the Mediterranean Basin, in *Biological Invasions in Europe and the Mediterranean Basin. Monographiae Biologicae, vol 65*, eds. di Castri F, Hansen AJ, Debussche M (Dordrecht, Netherlands: Springer), 353–364. doi: [10.1007/978-94-009-1876-4\\_21](https://doi.org/10.1007/978-94-009-1876-4_21)
- Wacquant JP, Bouab N 1983 Nutritional differentiation within the species *Dittrichia viscosa* W. Greuter, between a population from a calcareous habitat and another from an acidic habitat, in *Genetic Aspects of Plant Nutrition. Developments in Plant and Soil Sciences, vol. 8*, eds. Sarić MR, Loughman BC (Dordrecht, Netherlands: Springer). doi: [10.1007/978-94-009-6836-3\\_35](https://doi.org/10.1007/978-94-009-6836-3_35)
- Wacquant JP, Picard JB 1992 Nutritional differentiation among populations of the mediterranean shrub *Dittrichia viscosa* (Asteraceae) in siliceous and calcareous habitats. *Oecologia* 92, 14–22. doi: [10.1007/BF00317257](https://doi.org/10.1007/BF00317257)
- Waterkeyn A, Vanschoenwinkel B, Grillas P, Brendoncka L 2010 Effect of salinity on seasonal community patterns of Mediterranean temporary wetland crustaceans: A mesocosm study. *Limnol. Oceanogr.* 55, 1712–1722. doi: [10.4319/lo.2010.55.4.1712](https://doi.org/10.4319/lo.2010.55.4.1712)
- Watts K, Handley P 2010 Developing a functional connectivity indicator to detect change in fragmented landscapes. *Ecol. Indic.* 10, 552–557. doi: [10.1016/j.ecolind.2009.07.009](https://doi.org/10.1016/j.ecolind.2009.07.009)
- Whittaker R, Fernandez-Palacios J 2007 *Island biogeography*. 2nd edn. Oxford: Oxford University Press
- Wichern J, Wichern F, Joergensen RG 2006 Impact of salinity on soil microbial communities and the decomposition of maize in acidic soils. *Geoderma* 137, 100–108. doi: [10.1016/j.geoderma.2006.08.001](https://doi.org/10.1016/j.geoderma.2006.08.001)
- Williams MI, Dumroese RK 2013 Preparing for Climate Change: Forestry and Assisted Migration. *J. For.* 111, 287–297. doi: [10.5849/jof.13-016](https://doi.org/10.5849/jof.13-016)
- Winter TC 1999 Relation of streams, lakes, and wetlands to groundwater flow systems. *Hydrogeol. J.* 7, 28–45. doi: [10.1007/s100400050178](https://doi.org/10.1007/s100400050178)
- Wise RM, Fazey I, Stafford Smith M, Park SE, Eakin HC et al. 2014 Reconceptualising adaptation to climate change as part of pathways of change and response. *Glob. Environ. Chang.* 28, 325–336. doi: [10.1016/J.GLOENVCHA.2013.12.002](https://doi.org/10.1016/J.GLOENVCHA.2013.12.002)
- Woessner WW 2000 Stream and fluvial plain ground water interactions: rescaling hydrogeologic thought. *Groundwater* 38, 423–429. doi: [10.1111/j.1745-6584.2000.tb00228.x](https://doi.org/10.1111/j.1745-6584.2000.tb00228.x)
- Wong PP, Losada IJ, Gattuso J-P, Hinkel J, Khattabi A et al. 2014 Coastal Systems and Low-Lying Areas, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 361–409. [https://www.ipcc.ch/pdf/assessment-report/ar5/wg2/WGI-AR5-Chap5\\_FINAL.pdf](https://www.ipcc.ch/pdf/assessment-report/ar5/wg2/WGI-AR5-Chap5_FINAL.pdf) [Accessed March 11, 2017].
- Xanthopoulos G 2015 Forest Fires in Greece: Past, Present and Future, in *Wildland Fires: A Worldwide Reality*, eds. Bento-Gonçalves AJ, Vieira AAB (Nova Science Publishers), 141–151.
- Xoplaki E, Luterbacher J, Wagner S, Zorita E, Fleitmann D et al. 2018 Modelling Climate and Societal Resilience in the Eastern Mediterranean in the Last Millennium. *Hum. Ecol.* 46, 363–379. doi: [10.1007/s10745-018-9995-9](https://doi.org/10.1007/s10745-018-9995-9)
- Yasuhara M, Danovaro R 2016 Temperature impacts on deep-sea biodiversity. *Biol. Rev.* 91, 275–287. doi: [10.1111/brv.12169](https://doi.org/10.1111/brv.12169)
- Yıldız G 2018 Physiological Responses of the Mediterranean Subtidal Alga *Peyssonnelia squamaria* to Elevated CO<sub>2</sub>. *Ocean Sci. J.* 53, 691–698. doi: [10.1007/s12601-018-0044-9](https://doi.org/10.1007/s12601-018-0044-9)
- Yokeş MB, Andreou V, Bakiu R, Bonanomi S, Camps J et al. 2018 New Mediterranean Biodiversity Records (November 2018). *Mediterr. Mar. Sci.* 19, 673–689. doi: [10.12681/mms.19386](https://doi.org/10.12681/mms.19386)
- Zaimes GN, Emmanouloudis D 2012 Sustainable management of the freshwater resources of Greece. *J. Eng. Sci. Technol. Rev.* 5, 77–82.
- Zaimes GN, Gounaridis D, Symeonakis E 2019 Assessing the impact of dams on riparian and deltaic vegetation using remotely-sensed vegetation indices and Random Forests modelling. *Ecol. Indic.* 103, 630–641. doi: [10.1016/j.ecolind.2019.04.047](https://doi.org/10.1016/j.ecolind.2019.04.047)
- Zaimes GN, Gounaridis D, Fotakis D 2011 Assessing riparian land-uses/vegetation cover along the Nestos River in Greece. *Fresenius Environ. Bull.* 20, 3217–3225.
- Zaimes GN, Iakovoglou V, Emmanouloudis D, Gounaridis D 2010 Riparian areas of Greece: their definition and characteristics. *J. Eng. Sci. Technol. Rev.* 3, 176–183. doi: [10.25103/jestr.031.29](https://doi.org/10.25103/jestr.031.29)
- Zamir R, Alpert P, Rilov G 2018 Increase in Weather Patterns Generating Extreme Desiccation Events: Impli-

- cations for Mediterranean Rocky Shore Ecosystems. *Estuaries and Coasts* 41, 1868–1884. doi: [10.1007/s12237-018-0408-5](https://doi.org/10.1007/s12237-018-0408-5)
- Zeder MA 2008 Domestication and early agriculture in the Mediterranean Basin: origins, diffusion, and impact. *Proc. Natl. Acad. Sci. U. S. A.* 105, 11597–11604. doi: [10.1073/pnas.0801317105](https://doi.org/10.1073/pnas.0801317105)
- Zenetos A, Gofas S, Morri C, Rosso A, Violanti D et al. 2012 Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterr. Mar. Sci.* 13, 328–352. doi: [10.12681/mms.327](https://doi.org/10.12681/mms.327)
- Zeng Z, Peng L, Piao S 2018 Response of terrestrial evapotranspiration to Earth's greening. *Curr. Opin. Environ. Sustain.* 33, 9–25. doi: [10.1016/J.COSUST.2018.03.001](https://doi.org/10.1016/J.COSUST.2018.03.001)
- Zhu K, Woodall CW, Clark JS 2012 Failure to migrate: Lack of tree range expansion in response to climate change. *Glob. Chang. Biol.* 18, 1042–1052. doi: [10.1111/j.1365-2486.2011.02571.x](https://doi.org/10.1111/j.1365-2486.2011.02571.x)
- Ziveri P, Delpiazzi E, Bosello F, Eboli F, van den Bergh JCM 2017 Adaptation policies and strategies as a response to ocean acidification and warming in the Mediterranean Sea, in *Handbook on the Economics and Management of Sustainable Oceans*, eds. Nunes PALD, Svensson LE, Markandya A (Edward Elgar Publishing), 339–352.
- Ziveri P, Passaro M, Incarbona A, Milazzo M, Rodolfo-Metalpa R et al. 2014 Decline in coccolithophore diversity and impact on coccolith morphogenesis along a natural CO<sub>2</sub> gradient. *Biol. Bull.* 226, 282–290. <https://doi.org/10.1086/BBLv226n3p282>
- Zomeni M, Vogiatzakis IN 2014 Roads and Roadless areas in Cyprus: Implications for the natura 2000 network. *J. Landsc. Ecol.* 7, 75–90. doi: [10.2478/jlecol-2014-0010](https://doi.org/10.2478/jlecol-2014-0010)
- Zunino S, Canu DM, Bandelj V, Solidoro C 2017 Effects of ocean acidification on benthic organisms in the Mediterranean Sea under realistic climatic scenarios: A meta-analysis. *Reg. Stud. Mar. Sci.* 10, 86–96. doi: [10.1016/j.rsma.2016.12.011](https://doi.org/10.1016/j.rsma.2016.12.011)
- Zviely D, Kit E, Rosen B, Galili E, Klein M 2009 Shoreline migration and beach-nearshore sand balance over the last 200 years in Haifa Bay (SE Mediterranean). *Geo-Marine Lett.* 29, 93–110. doi: [10.1007/s00367-008-0126-2](https://doi.org/10.1007/s00367-008-0126-2)

## Information about authors

### Coordinating Lead Authors

Mario Balzan:

*Malta College of Arts, Science and Technology (MCAST), Paola, Malta*

Abed El Rahman Hassoun:

*National Council for Scientific Research (CNRS- L), National Center for Marine Sciences, Batroun, Lebanon*

### Lead Authors

Najet Aroua:

*Polytechnic School of Architecture and Urbanism, Algiers/ Laboratory of Architecture, Urbanism and Environmental Design, Department of Architecture, University of Biskra, Biskra, Algeria*

Virginie Baldy:

*Mediterranean Institute of Biodiversity and Ecology (IMBE, Aix-Marseille University, Avignon University, CNRS, IRD), Marseille, France*

Sana Ben Ismail:

*National Institute of Marine Sciences and Technologies (INSTM), Tunis, Tunisia*

Magda Bou Dagher:

*Faculty of Science, Saint-Joseph University, Beirut, Lebanon*

Cristina Branquinho:

*Centre for Ecology, Evolution and Environmental Changes, Faculty of Sciences, University of Lisbon, Lisbon, Portugal*

Jean-Claude Dutay:

*Laboratory of Climate and Environmental Sciences (LSCE), Pierre Simon Laplace Institute (IPSL), Gif-sur-Yvette, France*

Monia El Bour:

*National Institute of Marine Sciences and Technologies (INSTM), Carthage - Salammbô, Tunisia*

Frédéric Médail:

*Mediterranean Institute of Biodiversity and Ecology (IMBE, Aix-Marseille University, Avignon University, CNRS, IRD), Marseille, France*

Meryem Mojtahid:

*University of Angers, Angers, France*

Alejandra Morán Ordóñez:

*Forest Science and Technology Centre of Catalonia (CTFC), Solsona, Spain*

Pier Paolo Roggero:

*Department of Agricultural Sciences, Desertification Research Centre, University of Sassari, Sassari, Italy*

Sergio Rossi Heras:

*Department of Biological and Environmental Sciences (DiSTeBA), University of Salento, Salento, Italy*

Bertrand Schatz:

*Center of Functional and Evolutionary Ecology (CEFE), French National Centre for Scientific Research (CNRS), Montpellier, France*

Ioannis Vogiatzakis:

*Open University of Cyprus, Nicosia, Cyprus*

George N. Zaimis:

*UNESCO Chair Con-E-Ect, International Hellenic University, Drama, Greece*

Patrizia Ziveri:

*Universitat Autònoma de Barcelona (UAB), Barcelona, Spain*

### Contributing Authors

Marie Abboud-Abi Saab:

*Lebanese National Council for Scientific Research, Batroun, Lebanon*

Aitor Ameztegui:

*University of Lleida, Lleida, Spain*

Margaretha Breil:

*Euro-Mediterranean Centre on Climate Change (CMCC), Venice, Italy*

Thierry Gauquelin:

*Mediterranean Institute of Biodiversity and Ecology (IMBE, Aix-Marseille University, Avignon University, CNRS, IRD), Marseille, France*

Ilse Geijzendorffer:

*Louis Bolk Institute, Bunnik, the Netherlands*

Aristeidis Koutroulis:

*Technical University of Crete, Chania, Greece*

Juerg Luterbacher:

*World Meteorological Organization (WMO), Geneva, Switzerland*

Mohammad Merheb:

*Lebanese University, Tripoli, Lebanon*

César Terrer:

*Lawrence Livermore National Laboratory, California, United States of America*

Marco Turco:

*University of Murcia, Murcia, Spain*

Elena Xoplaki:

*Department of Geography & Center for international Development and Environmental Research, Justus Liebig University, Giessen, Germany*



# SOCIETY 1-DEVELOPMENT

**Coordinating Lead Authors:**

Maria dos Santos (Portugal), Stefano Moncada (Malta)


**Lead Authors:**

Antonietta Elia (Spain/Italy), Manolis Grillakis (Greece), Nathalie Hilmi (Monaco)

**Contributing Authors:**

Shekoofeh Farahmand (Iran), Walid Marrouch (Lebanon), Alain Safa (France), Brice Teisserenc (France)

*This chapter should be cited as: Dos Santos M, Moncada S, Elia A, Grillakis M, Hilmi N 2020 Development. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 469-492.*



## Table of contents

<b>5.1 Development</b> .....	<b>472</b>
Executive summary .....	<b>472</b>
<b>5.1.1 Past trends and current situation</b> .....	<b>472</b>
5.1.1.1 <i>Sustainable development</i> .....	<b>472</b>
5.1.1.2 <i>The institutional framework</i> .....	<b>473</b>
5.1.1.3 <i>The economic dimension of sustainable development</i> .....	<b>473</b>
<i>Economic development and employment</i> .....	<b>473</b>
<i>Tourism</i> .....	<b>474</b>
<i>Agriculture, fisheries and food security</i> .....	<b>474</b>
<i>Economic activities and their impact on inequality, gender and poverty</i> .....	<b>475</b>
5.1.1.4 <i>The social dimension of sustainable development</i> .....	<b>477</b>
<b>5.1.2 Economic vulnerabilities and risks</b> .....	<b>478</b>
<b>5.1.3 Adaptation</b> .....	<b>479</b>
5.1.3.1 <i>National legal and policy framework for adaptation</i> .....	<b>480</b>
5.1.3.2 <i>Economic and financial tools to promote environmental management and climate change adaptation in the Mediterranean region</i> .....	<b>481</b>
<b>Box 5.1.1 Development indicators and terms</b> .....	<b>484</b>
<i>Poverty</i> .....	<b>484</b>
<i>Inequality</i> .....	<b>484</b>
<b>Box 5.1.2 Gender-related development indicators and term</b> .....	<b>485</b>
<i>Gender</i> .....	<b>485</b>
<b>Box 5.1.3 Vulnerability</b> .....	<b>485</b>
<b>References</b> .....	<b>486</b>
<b>Information about authors</b> .....	<b>492</b>

## 5.1 Development

### Executive summary

Sustainable development seeks to address the needs of current and future generations with the objective of increasing well-being by balancing economic, social and environmental dimensions. Current unsustainable development patterns, such as poverty, increasing population pressure, agricultural intensification, land degradation, and air, land, rivers and ocean pollution, will be further exacerbated by climate change impacts.

Environmental and climate change impacts are likely to have an effect on all economic sectors in the Mediterranean Basin, increasing production challenges and costs, affecting low-income cohorts increasingly disproportionately, and generally delaying the achievement of the Sustainable Development Goals (SDGs).

Rising temperatures, with more intense and longer heatwaves, and decreasing rainfall patterns, can exert a further strain on important sectors, such as agriculture and tourism, which represent important cultural, economic and heritage assets for Mediterranean economies and societies.

The existence of poverty, inequalities and gender imbalances relate both directly and indirectly to the challenges faced by Mediterranean countries in achieving the SDGs, with current and predicted environmental and climate change impacts threatening the progress made to date in many areas of the Mediterranean Basin. To address these challenges, a new approach to development must be sought in order to eliminate poverty, sustain economic growth and ensure social protection, while safeguarding environmental standards and integrating mainstream climate change adaptation into policy making.

The growing challenges of environmental and climate change impacts on economies and societies require an enhanced institutional response, at a local, national and international level. Effective preventive and restorative policies, including the promotion of context-specific climate change mitigation and adaptation measures, and economic instruments encouraging behavioral changes, can ensure long-term sustainable development in the Mediterranean Basin.

### 5.1.1 Past trends and current situation

#### 5.1.1.1 Sustainable development

Sustainable development seeks to address the needs of current and future generations, making it possible to access and use natural resources in an equitable manner both now and in the years to come (Zidanšek 2007; Szopik-Depczyńska et al. 2017; Kilkis 2018). It sets the framework for securing viable and lasting development and decent livelihoods for all. It aims to promote a dynamic economy with high levels of employment and education, health protection, social and territorial cohesion and environmental protection in a peaceful and secure environment, while respecting cultural diversity (Bontoux and Bengtsson 2015, 2016). Successful societies are known to be built on economic, social and environmental sustainability (Dos Santos 2018; Dos Santos and Mota 2019), ensuring long-term sustainable development.

Traditionally, development has been associated exclusively with increases in income levels, and the conventional manner in which countries, including those located in the Mediterranean Basin, measure progress in development represented by Gross Domestic Product (GDP). While this measurement gives a good indication of the monetary value of the goods and services produced in a specific year, it fails to sufficiently capture other important dimensions of development, especially social and environmental ones. This is mostly due to the fact that GDP, among other things: i) does not capture inequalities in the distribution of income; ii) leaves out some activities (volunteer work) and does not record harmful activities (pollution, climate change); and iii) does not seem to improve well-being beyond certain levels of income (Briguglio 2019). The concept of development, and its measurement, vary. Among these, there is the Human Development Index (HDI), which captures three dimensions: income per capita, health and education (UNDP 2019). Another approach is the OECD better-life index which, along with traditional economic measurements includes other aspects such as quality of health and environmental services (OECD 2017). A further approach is that of measuring happiness. The United Nations Sustainable Development Solutions Network publishes the World Happiness Report annually (Helliwell et al. 2018). The report considers variables for measuring Gross National Happiness (GNH) which are: GDP per capita, social support, healthy life expectancy, freedom to



make life choices, generosity, and freedom from corruption. Its ultimate goal is happiness and to ensure a good quality life for the people through people-centric development initiatives. Each variable measured reveals a populated-weighted average score on a scale running from 0 to 10 that is tracked over time and compared against other countries. The ranking of Mediterranean countries for their GNH in 2018 varies significantly, from the highest for Israel (7.190), Malta (6.627) and France (6.488) to the lowest for Tunisia (4.592), Egypt (4.419) and Syria (3.466).

Using sustainability development indicators, it appears that for sustainable development, no sacrifices in happiness are required in the interest of future generations, as it is possible to design strategies that improve happiness and sustainability simultaneously (Zidanšek 2007).

The United Nations defines the sustainable development goals as the blueprint to achieve a better and more sustainable future for all. They address the global challenges we face, including those related to poverty, inequality, climate change, environmental degradation, peace and justice. The 17 Goals are all interconnected, and share the goal “to leave no one behind” (UN 2015).

All the European Union (EU) Member States (MS) in general, and all the countries around the Mediterranean Basin in particular, are facing increasing economic, social, environmental and institutional challenges. The way in which countries are responding to these challenges vary, also according to resource and governance levels. The EU has a considerable set of laws and regulations aimed at addressing environmental and climate change concerns, often making them a priority for all MS (Queralt et al. 2017). The heavily regulated environmental legislation within the EU has often been replicated in bilateral and regional agreements with countries in the Mediterranean Basin, in an attempt to commonly address transboundary concerns. Furthermore, the EU carbon emission reductions targets of 40% by 2030 and 80% by 2050, are increasingly producing spillover effects in other countries in the Mediterranean Basin, prompting the adoption of newer technology and the overall efficient use of energy and an improvement in carbon efficiency (Queralt et al. 2017).

### 5.1.1.2 The institutional framework

The growing challenges to harmonious development in the Mediterranean Basin have been duly

recognized by national and international bodies, prompting an institutional response that eventually led to the creation of the Mediterranean Commission for Sustainable Development (UNEP-MCSD) in 1996. The aim of the MCSD is to provide a bridge between the desire to pursue sustainable development and its effective implementation. It offers a framework in which to define a Mediterranean Strategy for Sustainable Development (MSSD). The MSSD has the objective of pursuing sustainable development goals so as to strengthen peace, stability, and prosperity. The strategy is structured around objectives and interlinked priority fields of action. Specific indicators are also identified to properly monitor and evaluate the strategy. The strategy is regularly reviewed by the parties, and renewed every five years.

### 5.1.1.3 The economic dimension of sustainable development

#### *Economic development and employment*

Two out of three people are already living in the urban areas of Mediterranean countries, which is higher than the global average. The United Nations Human Settlements Programme predicts that by around 2050, the urban population will grow to around 170 million in the countries on the northern shore (140 million in 2005) and to over 300 million to the south and east where the population was 151 million in 2005 (UNEP/MAP 2016).

Studies demonstrate several projected negative impacts of climate change on economic growth in Mediterranean countries. For instance, based on the severity of the Spanish drought of 1990, it is estimated that economic damages caused by droughts will exceed damages caused by earthquakes or floods (Handmer et al. 2012). Summer crops are particularly vulnerable (Giannakopoulos et al. 2009). As winters become milder while summers become warmer and longer, more cooling by air conditioning is needed in summer, that would increase the demand for electricity generation in most Mediterranean countries (*Section 3.3.3.6*) (Jacob et al. 2014; Kovats et al. 2014). Given that the existing infrastructure was implemented assuming a stable climate around the Mediterranean Basin (Scott et al. 2016a), climate change is making it a challenge for the economic infrastructure to adapt fast enough. This fact points out to the need for investments in adaptive infrastructure in the coming decades. Research and development might reduce the cost of adaptation (Arent et al. 2014).

The main economic sectors driving development in the Mediterranean coastal regions are resource-based activities (i.e., fisheries, aquaculture, forestry, agriculture, and primary industries), secondary industries (e.g., food processing, housing and construction) and services, especially tourism (UNEP/MAP-Plan Bleu 2009). Current and predicted environmental and climate change impacts are expected, especially in the absence of adaptation measures, to increase production costs and reduce productivity (Teotónio et al. 2020) in key sectors, exerting further pressure on economic development trajectories and employment levels in all countries of the Mediterranean Basin.

### **Tourism**

The Mediterranean has a rich history as well as exceptional natural and cultural landscapes. Over 360 million tourists travelled to the Mediterranean region in 2017 – more than double the number recorded in 1995 (Mediterranean Growth Initiative 2017). In the past 20 years the contribution of the tourism sector to GDP has steadily increased by 60% in Mediterranean countries. It should be mentioned that while most Mediterranean countries have experienced significant economic growth in the sector, the vast majority of economic growth corresponds to north-western Mediterranean countries, such as Spain, France and Italy. However, the countries where the tourism sector contributes the highest percentage to the national GDP are Malta, Montenegro, Greece and Morocco (UNEP/MAP 2016). France, Spain and Italy account for 17% of inbound tourism worldwide, corresponding to 234 million people for these three countries alone, as they remain the most attractive of the Mediterranean in terms of numbers of visitors (UNWTO 2019).

In the recent past, Mediterranean coastal regions have been characterized as ideal in terms of climate comfort for outdoor activities, especially during the June to August period (Amelung et al. 2007; Grillakis et al. 2016b). Changes in climate can impact tourism flows, directly by affecting the thermal comfort for outdoor recreational activities (Salata et al. 2018), or indirectly by affecting the natural resources of the destination, such as coastal erosion due to sea level rise (Jiménez et al. 2017), or the reduction of freshwater availability. Given that most Mediterranean tourism is based, and marketed, on the basis of the "sun, sea and sand" model (Koutra and Karyopouli 2013), the socio-economic consequences of a drop in tourist numbers due to climate change impacts can be severe. As sea-level rise leads to coastal retreat,

anticipation is needed in order to adapt to, and mitigate the economic impacts of sea-level rise on tourism and populations living in the affected areas (Enríquez et al. 2017).

Environmental degradation caused by climate change and human pressure can have serious impacts on Mediterranean tourism and, eventually decrease the economic benefits arising from this sector (Dogru et al. 2016). Some Mediterranean countries have started to adapt to the changes brought about by these negative impacts, by, for example, developing tourism strategies that try to attract visitors in the "shoulder months", and not focusing entirely on periods (i.e., Summer) where these impacts can affect demand (Niavis 2020). Overall, the vulnerability of tourism is higher in countries with lower adaptive capacity in terms of economic, social and political conditions (Dogru et al. 2016).

Tourism can also directly or indirectly be a driver of environmental and climate change impacts. This is the case, for example, when the lack of energy and water efficiency measures are put into place, exerting further pressure on local ecosystems (Drius et al. 2019) and increasing the cost of infrastructural maintenance. However, tourism could potentially play an important role in both creating jobs and fostering sustainable development in the Mediterranean, assuming that the right set of policies are adopted and well implemented for comprehensive achievement of the SDGs. According to the World Tourism Organization (UNWTO 2018), 64 countries submitted their Voluntary National Reviews (VNRs) in 2016 and 2017 on the SDGs. In these reports, tourism appears to be largely recognized as a high-impact sector with potential to advance all SDGs. 41 VNRs (64%) make one or more direct references to tourism. Tourism is most commonly mentioned in relation to SDG 8 (Decent work and economic growth), SDG 12 (Responsible consumption and production) and SDG 17 (Partnership for the goals) in VNRs on the SDGs among which Mediterranean Cyprus, France, Italy, Monaco, Slovenia, Montenegro and Egypt (UNWTO 2018).

### **Agriculture, fisheries and food security**

International political and economic organizations have become aware that high and volatile food prices and deregulated markets put food security at risk and seriously affect global economic, social and political stability (FAO / IFAD / WFP 2011). The financial crisis in 2007 in the USA affected Mediterranean countries in 2008 and has led to instability in agricultural markets and a rise in

the prices of these goods (Dos Santos 2018; Dos Santos and Mota 2019).

Climate change is expected to threaten food security (see *Section 3.2*), especially livestock production and fisheries. Livestock production is an important contributor to the economy. Countries with a higher risk of livestock production being impacted by climate change (e.g., increase in diseases and consequences of higher temperature on animal health) are those which have lower adaptive capacity (Godber and Wall 2016).

Fisheries play an important role in the economy of Mediterranean countries. Total fish landings account for more than €3 billion yearly in the Mediterranean Sea, and including all the ancillaries services, this industry can reach an estimated value of around €10 billion yearly (Sacchi 2011). These values are likely to have been underestimated as significant portion of Mediterranean fish catches are not sold through regulated market outlets (Piante and Ody 2015), not to mention the cultural and tourism value that such an industry adds to local economies.

### ***Economic activities and their impact on inequality, gender and poverty***

The presence of poverty and income inequality is interconnected with economic growth (Galor and Zeira 1993; Ncube et al. 2014; Bruckner and Lederman 2015). According to the expectations of the UN SDGs, poverty, in both developed and developing countries, should be abolished by 2030. A priority of governments in this direction is the implementation of policies that enhance economic growth. However, given the tendency of the economic growth process to disproportionately exclude the lower income cohorts from accessing the benefits of new wealth, more equitable approaches to such wealth and social protection systems should be implemented, or enhanced.

These tendencies are confirmed by the data from the UNDP HDI (2019) when adjusted for inequality (*Box 5.1.1; Table 5.1*), showing that when considering the distribution of income, the HDI ranking for some countries drops and for others improves, which is likely to be associated with policies that try to address the problem of inequality.

The World Bank (2018) has used poverty lines to determine the headcount ratio and poverty gap. Four bases have been considered for poverty lines in order to determine the headcount ratio and poverty gap: 1.90 USD income per day,

3.20 USD income per day, 5.50 USD income per day, in addition to national poverty lines. Since there are many missing poverty data for some national poverty lines, the data are interpreted and countries are compared using standard poverty lines. Firstly, considering 1.90 USD a day as the poverty line, on average about 0.6% of the Mediterranean people are poor. The highest percent of the poor are in Italy, while the highest poverty gap is in Syria. Regarding this poverty line, there are no poor people in countries like France, Malta, Montenegro, Slovenia, Cyprus, and Lebanon. By increasing the poverty line to 3.20 USD, the highest percentage of the poor is in Egypt (16.1%), Syria (15.3%), and Albania (7.7%). In contrast, Slovenia and Cyprus have no such poor people and the percentage of people under this poverty line is low in France and Malta (0.2%). The largest poverty gaps are seen in Syria, Italy, and Albania. When 5.50 USD is used as the poverty line, both poverty percentages and poverty gap increase considerably in the region. Accordingly, the highest percentage of the poor lives in Egypt, Albania, and Morocco. Figures show that more than 60% of Egyptians have income of less than 5.50 USD per day. Also, most poverty gaps belong to the same countries. However, the percentage of people under the poverty line is low in France, Malta, and Slovenia.

Gender inequality indicators for Mediterranean countries, listed in UNDP human development reports, show a varied and complex situation. In the sample countries of the Mediterranean (*Box 5.1.2; Table 5.2*), Slovenia has the highest Gender Development Index (GDI), at 1.003 in 2017, which ranks it 18th out of 164 international countries in the 2017 index. This means that men and women have relatively the same achievement in three basic dimensions of human development. Croatia, France, and Cyprus are in the next positions in the region and have ranked 31st, 39th and 45th in the world. The last rank in the region belongs to Syria (ranked 159th in the world). The GDI value for Syria demonstrates that there is inequality in human development in favor of men. In fact, the human development index for Syrian men is almost 21% higher than for Syrian women.

For the Gender Inequality Index (GII), the highest value in the region is Slovenia (0.054), ranking it 7th out of 160 countries in the 2017 index. After Slovenia, Spain and France have the least gender inequality in the region and are ranked 15th and 16th in the world. According to the indicator, Syria has again the lowest place in the region and shows the highest gender inequality.

Country	HDI		INEQUALITY			
	HDI Value <sup>a</sup> (2018)	HDI Rank <sup>a</sup> (2018)	IHDI Value <sup>a</sup> (2018)	20:20 Ratio <sup>b</sup> (2010-2017)	Palma Ratio <sup>b</sup> (2011-2017)	Gini Index <sup>b</sup> (2011-2017)
<b>SOUTHERN EUROPE</b>						
Albania	0.791	67	0.705	4.25	1.0	29
Bosnia and Herzegovina	0.769	75	0.656	5.43	1.3	33
Croatia	0.837	46	0.768	5.26	1.1	31.1
France	0.891	26	0.809	5.18	1.3	32.7
Gibraltar	-	-	-	-	-	-
Greece	0.872	32	0.766	7.09	1.5	36
Italy	0.883	29	0.776	7.00	1.4	35.4
Malta	0.885	28	0.815	4.48	1.1	29.4
Monaco	-	-	-	-	-	-
Montenegro	0.816	52	0.746	4.77	1.2	31.9
Portugal	0.850	40	0.742			
Slovenia	0.902	24	0.858	3.66	0.9	25.4
Spain	0.893	25	0.765	7.26	1.5	36.2
Turkey	0.806	59	0.675	8.47	2.1	41.9
<b>LEVANTINE REGION</b>						
Cyprus	0.873	31	0.788	5.33	1.4	34
Israel	0.906	22	0.809	8.50	2.0	38.9
Jordan	0.723	102	0.617			
Lebanon	0.730	93	..	5.06	1.2	31.8
Palestine	0.690	119	0.597	..	1.4	34.4
Syrian Arab Republic	0.549	154	..	..	..	..
<b>NORTHERN AFRICA</b>						
Algeria	0.759	82	0.604	3.96	1.0	27.6
Egypt	0.700	116	0.492	4.56	1.3	31.8
Libya	0.708	110	-	-	-	-
Mauritania	0.527	161	0.358			
Morocco	0.671	121	-	7.02	2.0	39.5
Tunisia	0.739	91	0.585	5.24	1.5	32.8

**Table 5.1 | Inequality indicators for Mediterranean countries**

<sup>a</sup> Source: UNDP 2019 - <sup>b</sup> Source: World Bank 2019

The existence of poverty, inequalities and gender imbalances relate both directly and indirectly to the achievement of sustainable development goals in Mediterranean countries. Significant theoretical and applied research has shown that the presence of these imbalances, both relative and absolute, are obstacles to the expansion of economic development, de facto blocking parts of society from potentially enjoying the benefits of higher standards of living (Sen 1999; Sachs

2005). Moreover, the traditional way of measuring economic progress by only taking GDP into account, does not capture the problem in the first place, and the extent to which these imbalances permeate societies. The absence of this specific indicator, especially when measuring economic progress, does not bring about reaction or prevention from policy systems, therefore, overlooking the problems related to these distortions of the market economy.

Country	GDI			GII		
	Value <sup>a</sup> (2017)	Rank in Med	Rank in the world	Value <sup>b</sup> (2017)	Rank in Med	Rank in the world
<b>SOUTHERN EUROPE</b>						
Albania	0.970	7	67	0.238	13	52
Bosnia and Herzegovina	0.924	13	117	0.166	10	37
Croatia	0.991	2	31	0.124	8	29
France	0.987	3	39	0.083	3	16
Gibraltar	-	-	-	-	-	-
Greece	0.964	9	80	0.120	7	26
Italy	0.967	8	73	0.087	5	18
Malta	0.960	10	83	0.216	12	45
Monaco	-	-	-	-	-	-
Montenegro	0.956	11	88	0.132	9	32
Slovenia	1.003	1	18	0.054	1	7
Spain	0.979	5	51	0.080	2	15
Turkey	0.922	14	118	0.317	15	69
<b>LEVANTINE REGION</b>						
Cyprus	0.984	4	45	0.085	4	17
Israel	0.975	6	62	0.098	6	21
Lebanon	0.889	16	129	0.0381	16	85
Palestine (Gaza Strip)	0.877	17	132	-	-	-
Syrian Arab Republic	0.788	21	159	0.547	20	136
<b>NORTHERN AFRICA</b>						
Algeria	0.861	19	142	0.442	17	100
Egypt	0.873	18	135	0.449	18	101
Libya	0.929	12	112	0.170	11	38
Mauritania						
Morocco	0.838	20	151	0.482	19	119
Tunisia	0.897	15	125	0.298	14	63

**Table 5.2 | Gender indicators for Mediterranean countries** (UNDP 2019).

<sup>a</sup> Source: UNDP 2019 - <sup>b</sup> Source: World Bank 2019

#### **5.1.1.4 The social dimension of sustainable development**

Education is a fundamental prerequisite for addressing all issues related to Sustainable Development (SD). It creates the necessary enabling environment to enhance skills as well as individual and collective social commitment for the desired transformations, by also allowing for the creation of more sustainable societies (Voegtlin and Scherer 2017). Education can also support the development of better strategies for mitigating and adapting to climate change, thus promoting sustainable development (Anderson 2012).

Education for sustainable development (ESD) is an approach to teaching and learning based on the ideals and principles that underlie sustainability and applicable to all types, levels and settings of education. As such, ESD promotes multi-stakeholder social learning, emphasizes the empowerment of communities and citizens, engages with key issues such as human rights, poverty reduction, sustainable livelihoods, environmental education and gender equality in an integral way and encourages changes in behavior that will create a more sustainable future (Voegtlin and Scherer 2017).

A further important factor in the social dimension of sustainable development is participation. More active participation of the community, especially children and youth, as agents of change, can increase public authorities' understanding of problems, and facilitate the implementation of solutions among communities (Anderson 2012).

The removal of social imbalances, such as gender gaps in education and in salaries, can increase productivity and facilitate economic growth. The main results highlight that the educational gender gap hinders economic growth and development as a whole (Tansel and Güngör 2016; Minasyan et al. 2019). The majority of the results confirm a positive effect of female education on economic growth and development (Forbes 2000; Tansel and Güngör 2016).

### 5.1.2 Economic vulnerabilities and risks

Expected extreme climate conditions and pollution can enhance economic vulnerabilities and risks in the Mediterranean Basin (*Chapter 2*). In recent decades, a growing number of publications have identified and assessed how natural hazards occurring in the Mediterranean Region interact with its society and economy. This evidence is being produced on a sectoral level, with assessments of biodiversity, agriculture and cultural heritage systems (Palatnik and Lourenço Dias Nunes 2015; Fatorić and Seekamp 2017), according to the type of hazard (Llasat et al. 2013; Iglesias and Garrote 2014; Oliveira et al. 2018), or with a specific geographical scope (Schilling et al. 2012; Radhouane 2013; Monioudi et al. 2017). An underlying common denominator in the available literature seems to point towards the Mediterranean region experiencing a higher intensity of, and associated risks related to specific natural hazards than other European regions.

This seems to be the case of disastrous flash-floods, which are much more recurrent in some areas of the Mediterranean Basin, when compared to the rest of Europe (*Section 3.1.3.3*) (Llasat et al. 2010). These trends are confirmed by research that looked at flood event mortality in the eastern Mediterranean, which, when accounting for "high" number of casualties is higher, and for "no-deaths" is lower, than central Europe (Doocy et al. 2013). In selected areas of the Mediterranean Basin, the economic sectors more prone to be directly affected by floods are agriculture, followed by commerce and artisan trades, tourism, and industry (Llasat et al. 2013).

Current observations show an increase in drought events, and reduced soil moisture and groundwater availability (*Sections 3.1.3.1 and 3.1.3.4*). These impacts interact negatively, especially with the agricultural sector in the Mediterranean, threatening food security in rainfed yields (Tigkas and Tsakiris 2015) (*Section 3.1.2.2*), economic performance in terms of reduced wheat exports (Dellai and McCarl 2010) and livestock production (Blauhut et al. 2015). Combined with increasing population in Mediterranean countries, these impacts could intensify the problem of food security.

Changes to precipitation patterns and increased temperature (*Section 2.2.5.3*), can also affect the quantity and quality of grazing areas, directly impacting farmers' income, with a higher negative impact in non-EU Mediterranean countries, especially at the small-scale level (Abdul Malak et al. 2017).

The increase in sea temperatures and ocean acidification will likely have a negative impact on the fishery industry (*Section 3.2.2.2*), with these phenomena already linked to mass mortality events in the Mediterranean (Coma et al. 2009), affecting aquaculture by reducing available space to operate businesses (Bird et al. 2016), and potentially increasing mortality rates of the species cultivated, especially due to the increase in heat waves in summer (Rodrigues et al. 2015).

The economic vulnerabilities associated with sea-level rise and coastal erosion have received considerable attention in the Mediterranean region, also given the economic implications of tourism, which for some economies represents more than 30% of the aggregate GDP (Koutroulis et al. 2018a). Summer tourism in Mediterranean countries, which is based on beach holidays, can be threatened by hotter and drier summers, in turn affecting the comfort levels of tourists (Koutroulis et al. 2018b). However, the degree to which Mediterranean countries' tourism sectors might be affected by climate change is often a function of income levels, with the highest levels of vulnerability coming from the lowest income and least resilient countries (Dogru et al. 2019). The effect of sea level rise, together with changes in storm features can seriously affect port operations, slowing down trade operations and productivity levels (Sánchez-Arcilla et al. 2016).

Climate change is expected to cast a shadow of uncertainty over tourism in the Mediterranean. Uncertainties in the assessment of tourism de-

mand under future emission and socio-economic trajectories are subject to factors that affect the visitors' sensitivity to thermal comfort (age, type of tourism, country of origin) (Dubois and Ceron 2006; Dubois et al. 2016). Additional uncertainties stem from the adaptation and mitigation response to climate change (Koutroulis et al. 2018a), as well as the lack of integrated assessment that considers cross sectoral interactions to climate change (Scott et al. 2016b). Mediterranean summertime thermal comfort is expected to generally negatively affect tourism flows in the core summer tourism months of June to August (Amelung et al. 2007; Grillakis et al. 2016a). Improvement in the climate resource is expected for the same regions in the spring and autumn season (Amengual et al. 2014; Grillakis et al. 2016b).

Limited research has quantified the effect of global warming on the net economic impacts on tourism in the Mediterranean. A tourism climate index has been correlated to the total overnight stays in European summer tourism, quantifying the effect of climate change on future overnights stays under 1.5°C and 2°C of global warming above preindustrial levels (Jacob et al. 2018). It was found that 1.5°C of global warming will have an impact on European Mediterranean summer tourism comfort in the July to August period. For the May to October period, marginal positive changes are projected over the majority of the European region, while for the June to August period, a negative effect over southern Spain and Cyprus and for most coastal regions of the Mediterranean is projected. These comfort changes may have a direct impact on the number of overnight stays, with Cyprus and Greece to face a potential decrease of 8% and 2%, respectively. In Spain and Italy, the decrease in comfort over the southern regions of the countries could be compensated by an increase in the north, with a possible northward shift of tourism activity. At 2°C of global warming, this pattern of change is expected to augment further. From an economic perspective, northern Mediterranean regions could exhibit climate induced tourism revenue decreases up to -0.45% of their GDP per year by 2100 (Barrios and Ibañez 2015). A regional temperature increase of 2.2°C in selected regions of Sardinia and Tunisia (Cap Bon) is expected to improve in the shoulder (spring - autumn) season, while increased heat stress may cause a decline in tourism demand in summer (Köberl et al. 2016). The annual net effect is expected to be marginally positive in terms of overnight stays, however the net profit might be less than the present due to the potential increase in the cost of water.

### 5.1.3 Adaptation

Climate change adaptation can be defined as the process of adjustment to actual or expected climate change and its effects (Smit and Wandel 2006). Climate change adaptation has been identified by the international community as an essential policy response, and its integration into development planning is a key measure for the effective achievement of sustainable development goals. However, the limited resource base of some countries tends to hinder climate change adaptation measures, which are also unlikely to occur automatically in response to observed or expected changes. This kind of response is highly dependent on the specific characteristics of a system, or community, affected by the impacts. The overall long-term adaptive capacity of a population is, in fact, shaped by existing developmental deficits, by exposure to sensitive risks, and by the strategies employed by individuals and communities to cope with these deficits and risks. Existing developmental deficits in some Mediterranean countries, such as endemic poverty, limited infrastructure and technology, ecosystem degradation, conflicts and poor health, among others, challenge the capacity to cope with emerging climate change, in turn affecting the way socio-economic adaptation strategies are applied.

Promoting climate change adaptation is believed to be a win-win strategy. The concept of adaptation has become increasingly associated with what can be considered good development. Such actions to improve climate change adaptation span over a spectrum of initiatives involving investments in, for example, human capital, such as increasing levels of education, skills, and the health status of poor households, or physical capital, such as climate proofing infrastructure projects and investing in self-sustaining renewable energy projects. A lack of human capital, or having poor infrastructure status, can also have a direct impact on labor productivity, and the related capacity to provide secure livelihoods, both in more industrialized Mediterranean countries by slowing down economic development, and in less developed Mediterranean countries by also increasing inequalities. Predicted climate change impacts will only exacerbate such circumstances, especially if nothing is done to account for these impacts (Mavromatidi et al. 2018).

Although applied research is increasing (Cramer et al. 2018), how specific regions or communities in Mediterranean countries may be affected by climate change is still partially unknown. A lack of data and research capacity remains a major

problem in some Mediterranean countries. Generally, the most impoverished and vulnerable communities are frequently neglected in impacts and adaptation research (Satterthwaite 2013; Moncada et al. 2018). Additionally, any available data is frequently interpolated over large spatial or temporal scales. While such data can provide some initial insight into potential risks, the data may not be meaningful at the scale at which people live. Failure to identify specific local impacts may lead to adaptation activities that are poorly targeted, inadequate, or even maladaptive (Albizua et al. 2019). Maladaptation refers to actions that might lead to increased risks of adverse climate-related outcomes, increased vulnerability to climate change, or diminished welfare, now or in the future (adapted from Agard et al. 2014).

The promotion of climate change adaptation can be achieved by either acting on those constraints that impede the attainment of human needs (Sen 1999), or by directly improving the management of specific climate stressors (Pelling 2011). In this regard, a study by McGray et al. (2007) screened more than 100 projects considered to be climate change adaptation initiatives that mostly occurred in low- and middle-income countries. They subsequently found that the beneficial outcomes of the projects had little difference to what can be considered as good development (Klein 2010). McGray et al. (2007) identify a continuum of actions that can be undertaken in order to address climate change impacts. These range from pure development actions, with usually no intentions to tackle climate change adaptation, to purposely designed adaptation efforts. When the first set of actions, aimed at addressing vulnerabilities, can have a positive effect on adaptation, this is known as no-regret, win-win options (Kelly and Adger 2000). On the other hand, the actions targeted to tackle specific climate change impacts might not have any effect on development, unless they are effective at tackling climate change adaptation. In between “lies a broad spectrum of activities with gradations of emphasis on vulnerability and impacts” (Bapna and McGray 2008).

An additional aspect in which climate change adaptation affects development relates to the financial implications for countries to invest in climate change adaptation. External financial and technical assistance is needed in order to cope with the impacts of climate change. Interestingly, the tendency of the majority of the international community has been to finance adaptation mostly through tackling climate change impacts, and largely ignoring the benefit

of addressing baseline vulnerabilities/deficits, and effectively enhancing long-term adaptive capacities. Increasing the adaptive capacity of a given system, thus raising the overall level of development, reduces the undesirable impacts of climate change, by allowing a system to better cope with changing conditions, risks or opportunities related to climate change (Smit and Wandel 2006). It has therefore been argued that supporting short-term adaptive capacity, especially in poor communities, is an urgent priority (McGray et al. 2007), as well as being increasingly recognized as an essential element of development (Ayers and Dodman 2010).

This is especially true for investments that address context-specific vulnerabilities, as highlighted in the previous section. Adaptation, therefore, must permeate all policy areas and not only the environment. Investment decisions that do not consider mitigation and adaptation can block regional and national development for many years. Thus, the success of adaptation strategies will also involve adapting actions to specific regional climatic conditions, in sectoral, political and socio-economic contexts by ensuring dialogue between stakeholders, through cooperative structures and knowledge transfer and monitoring progress to support regular reviews of policy objectives and the inclusion of new scientific information when it becomes available (di Gregorio et al. 2017).

Although the links between climate change impacts, climate action and sustainable development are broadly accepted, there has been limited structured investigation in terms of specific SDG Targets, synergies and trade-offs. The Intergovernmental Panel on Climate Change (IPCC) special report on Global Warming of 1.5°C features a chapter that investigates links between certain climate mitigation and adaptation actions and the 17 SDGs (Roy et al. 2018), but it does not assess the specific synergies and trade-offs between climate impacts, climate action and all 169 individual targets of the 2030 Agenda (di Gregorio et al. 2017).

### **5.1.3.1 National legal and policy framework for adaptation**

In accordance with the existing EU framework, EU member States adopted a series of legal measures and a strategic plan of actions to promote adaptation. These measures are related to multiple sectors potentially impacting the environment and aim to reduce and/or mitigate actions affecting climate change. Between them, Spain,



Portugal, and Italy, have adopted a few relevant measures. For example, Spain adopted the Royal Decree-Law n. 15/2018, aiming to accelerate the integration of renewable energies into the economy and to promote energy efficiency. Relevant measures adopted by Portugal include the Law-Decree n. 4/2018 introducing incentives for urban electric mobility (similarly to the Spanish Royal Decree of 16th June 2017, n. 617/2017), Decree-Law n. 64/2017 creating a legal framework for the implementation of biomass plants by municipalities, the Portuguese Strategic Framework for Climate Policy was adopted through the Resolution of the Council of Ministers n. 56/2015, including the National Program for Climate Change 2020/2030 (in line with the EU law and strategy) and the National Strategy for Adaptation to Climate Change. The Italian case is more complex, because of the legislative powers divided between the central State and Regions (Art. 117 of the Italian Constitution). Even if the Italian Climate Adaptation Strategy was adopted in 2015 by Decree of the Ministry of Environment, local governments are responsible for its implementation.

Outside the EU, Israel adopted its National Plan for Implementation of the Greenhouse Gas Emissions Reduction Targets and for Energy Efficiency on 10th April 2016, by Government Decision n. 1403/2016. In relation to water resources and the public hydraulic domain, Morocco adopted Law n. 36-15 in 2016, including a total of 163 articles and 12 chapters. Other countries, like Lebanon, adopted two separate national strategies: one related to energy efficiency (with a mid-term scope) and a second broad document related to renewable energy (both in 2016).

The variants of sustainable urban growth (smart cities, green cities, resilient cities, low carbon cities, sustainable cities) have brought renewed opportunities to create pathways for sustainable urban development (Rodriguez et al. 2018). However, the proliferation of all these different concepts, often meaning the same thing, has also created competing agendas and confusion for local decision makers, planners, stakeholders, and business communities. Therefore, progress can be made by focusing on key opportunities that create precedents for transformative and sustainable urban development.

According to the IPCC Fifth Assessment Report (AR5) the majority of cities' adaptation plans and strategies are based on the construction of defensive infrastructure. Although defensive infrastructure is a relevant element of Climate

Change Adaptation (CCA) pathways and building resilient cities, making it the center of adaptation plans limits opportunities (Mimura et al. 2014). According to Rodriguez et al. (2018), SDGs can help create recognition for the wider social, cultural, economic, political, institutional and normative elements of adaptation that can lead to the construction of multidimensional operational approaches on the ground.

### **5.1.3.2 Economic and financial tools to promote environmental management and climate change adaptation in the Mediterranean region**

The use of economic instruments to achieve environmental goals and natural resource management (including the management of water quantity-typically extraction charges or taxes), fisheries (taxes, fees and transferable quotas), forestry (charges and subsidies) and wetlands (financial assistance to owners) has increased significantly since the 1970s. The most common market tools in the Mediterranean European countries are charges/taxes (France, Greece, Italy, Spain and Turkey), tradeable permits (France), deposit-refund systems (Italy and Turkey), non-compliance fees (Greece and Turkey) and subsidies (France, Greece and Turkey) (Bartels et al. 2016; Carreño 2019).

Pollution and climate change control instruments can be classified into three categories: 1) institutional approaches to facilitate internalization of externalities; 2) command and control instruments; and 3) economic incentive (market-based) instruments (Table 5.3). Each specific category includes different approaches to achieve the goal of environmental management and climate change adaptation. The institutional approach, where institutions use pollution control instruments to prevent damages to third parties (externalities) or to charge the polluters for the damage that has been produced (internalization of externalities), comprises three specific approaches, as follows:

- Bargaining between generators and victims of pollution could reduce pollution below the critical threshold, but it requires some Institutional intervention, because bargaining often fails to alleviate the targeted pollution. In fact, it is difficult to identify all affected parties, to place importance on future generations with current generations, etc. (Perman et al. 2003).

INSTRUMENT	DESCRIPTION
<b>Institutional approaches to facilitate internalization of externalities</b>	
Facilitation of bargaining	Cost of, or impediments to bargaining are reduced
Specification of liability	Codification of liability for environmental damage
Development of social responsibility	Education and socialization programs promoting citizenship
<b>Command and control instruments</b>	
Input controls over quantity and/or mix of inputs	Requirements to use particular inputs, or prohibitions/restrictions on use of others
Technology controls	Requirements to use particular methods or standards
Output controls: Output quotas or prohibitions	Non-transferable ceilings on product outputs
Emissions licenses	Non-transferable ceilings on emission quantities
Location controls (zoning, planning controls, relocation)	Regulations relating to admissible location of activities
<b>Economic incentive (market-based) instruments</b>	
Emissions charges/taxes	Direct charges based on quantity and/or quality of a pollutant
User charges/fees/natural resource taxes	Payment for cost of collective services (charges), or for use of a natural resource (fees or resource taxes)
Product charges/taxes	Charges/taxes applied to polluting products
Emissions abatement and resource management subsidies	Financial payments designed to reduce damaging emissions or conserve scarce resources
Marketable emissions permits	Two systems: those based on emissions reduction credits (ERCs) or cap-and-trade
Deposit-refund systems	A fully or partially reimbursable payment incurred at purchase of a product
Non-compliance fees	Payments made by polluters or resource users for non-compliance, usually proportional to damage or to profit gains
Performance bonds	A deposit paid, repayable on achieving compliance
Liability payments	Payments as compensation for damage
Loans	Loans available to enterprises to implement pollution control projects
Subsidies	Subsidies paid by the government to firms or consumers for per unit reductions in pollution
Payment for ecosystem services	Payments for environmental services or benefits made by a beneficiary to the provider of the service
Clean development mechanism	Allows a country with an emission-reduction or emission-limitation commitment to implement an emission-reduction project in developing countries
Voluntary emission reduction	Actions that allow the polluter to take advantage of voluntary efforts to reduce greenhouse gas emissions by following certain regulations and standards

**Table 5.3 | Classification of finance tools to protect the environment and promote sustainable development**  
(Perman et al. 2003)

- The liability principle, which says that the polluter pays to prevent and remedy environmental damage, and includes the use of direct control tools over polluters, such as mandatory obligations or restrictions on the behavior of firms and individuals. It is the most dominant method for protecting the environment. It is related to property rights and is currently implemented in France (Boivin and Emorine 2019), Italy (Chilosi et al. 2019), Spain (Almenar et al. 2019), Turkey (Perman et al. 2003; Mavioglu et al. 2019), and Slovenia (Justice and Environment 2012).

- Development of social responsibility, which creates incentives for polluters to voluntarily change their behavior. In many - but not all - circumstances, economic incentive-based instruments are more cost-effective than command and control instruments (Perman et al. 2003). This approach includes raising public awareness (UNECE 2013 in Cyprus; UNEP 2015) and environmental education, which is an effective part of the European Union's environment policy (Stokes et al. 2001), and is also implemented in Slovenia (Kraus 1998), Algeria (Environmental Rights Database) and Tunisia (MESD 2018).

The command and control instruments have five key tools, as highlighted in *Table 5.3*. Regulations regarding direct control may apply to:

- outputs of emissions themselves and to the quantity of final production, e.g., fishing quotas in France, when consumption rates exceed 70%, and Spain (OECD 2003),
  - the production techniques used, such as regulating industrial emissions within European Union countries, including Spain, under the Industrial Emissions Directive (IED) and Plan AIRE which regulates small installations' emissions in Spain (UNEP 2015),
  - the level and/or mix of productive inputs, e.g., input laws in agriculture in Croatia (Grgi et al.); bans on the use of phthalates in toys in France, Greece, and Italy; over 60 PVC-free cities and restrictions on PVC-packing in Spain (Center for Health Environment and Justice (undated)); restrictions on the use of asbestos in 15 Mediterranean countries (Kazan-Allen 2019); restricted financial support by French banks for coal mining and coal powered generation projects (Littlecott 2015); plans to shut all of France's coal-fired power plants by 2021 (White 2018; Climate Transparency 2019), as well as reducing fossil fuel use by 30% by 2030 (Littlecott 2015); and the coal phase-out plan compatible with the limit of global warming below 1.5°C in France and Italy and completing full decarbonization by 2050 (Climate Transparency 2019),
  - emission licenses (e.g., in Turkey) (Mavioglu et al. 2019),
  - and even to the location of emission sources, e.g., natural regional parks (Salanié and Coisson 2016) and Zones de Conservation Halieutique (ZCH) in France (OECD 2018); acoustic zoning in Italy (Prašević et al. 2012); and environmental zoning in Bosnia and Herzegovina (Zahumenská et al. 2015).
- Economic incentive (market-based) instruments, include many tools, which interface with prices and markets, which can be summarized as follows:
- Emission charges/taxes such as carbon taxes (in France and Slovenia) (CTC (Carbon Tax Center) 2018), as well as SO<sub>2</sub> charges and noise pollution charges (Perman et al. 2003).
  - User charges/fees/natural resource taxes such as congestion pricing in France, which does not require a complex system of monitoring, enforcement and compliance, incentivize responsible use of resources and promotes investments (Milewska 2019).
  - Product charges/taxes such as costs for plastic bags in Italy (Perman et al. 2003), taxes on energy products in Italy, which earns more than 0.5% of its GDP from taxation of energy products (OECD 2013), taxes on lubricants in France and Italy, and taxes on cadmium batteries in Italy (Barde 1994).
  - Marketable emissions permits have been used in some Mediterranean countries like Italy (Recchini 2016). They are allocated by selling them (e.g., by auction) or by giving them away (Devlin and Grafton 1996). Permits are not financially sustainable instruments and had weak performance in the EU (Pettinger 2017).
  - Deposit-refund systems are recommended as components of an overall socially optimal set of policies. They can efficiently control pollution in almost the same way as Pigouvian tax (Walls 2011).
  - Non-compliance fees are implemented in Greece (Perman et al. 2003), Bosnia and Herzegovina (UNECE 2018), France (Bianco et al. 2015), and Slovenia (OECD 2012b).
  - Liability payments are used in different forms in countries like France, Italy (OECD 2013), Spain (Justice and Environment 2017), Bosnia and Herzegovina (UNECE 2018), Slovenia (OECD 2012b, 2012a) and Turkey (OECD 2019). Pursuing liability claims is very costly and the outcome is highly uncertain (Anderson 2002).
  - Green loans are used in France (Zakhartchouk 2019), Italy (Lewenhak 2012), Croatia (UNECE 2014), and Lebanon (SwitchMed 2017). Also, Slovenia has plans to reduce water pollutants through loans (GEF 2019).
  - Payments for ecosystem services (PES) schemes are implemented in the European Mediterranean countries. While PES are a rapidly proliferating mechanism for natural resource management, their use is sometimes based on an incomplete understanding of their social and economic impacts (DIE 2014). Early PES experiences reveal some positive equity impacts like improved tenure security, community empowerment, organizational and social capital development (Richards and Jenkins 2007). Clean development mechanisms (CDM), e.g., the Concentrated Solar Power plant project in Morocco (ADB and African Development Bank 2011); and CDM or

Joint Implementation projects in several countries including Croatia, France, Greece, Italy, Spain, and Turkey (UNFCCC 2012).

- Voluntary emission reduction is a tool employed by France, Italy, Spain, and Greece, and some for-prof-

it organizations in France, Italy, Spain, and Greece, which reported voluntary carbon offsets (Hamrick and Brotto 2017). Voluntary emission reductions of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O have also been reported in Croatia (Delija-Ružić 2017).

## BOX 5.1.1

### Development indicators and terms

#### Poverty

There is growing consensus among development experts that poverty is multidimensional. In this context, the Multidimensional Poverty Index (MPI) was developed in 2010 by the Oxford Poverty & Human Development Initiative (OPHI) and the United Nations Development Programme (UNDP). The Multidimensional Poverty Index (MPI) provides a sound gauge of poverty. The index measures poverty in three dimensions, i.e., education, health and standard of living. This index is computed for Less Developed Countries (LDCs) and developing countries (UNDP 2018e). According to the MPI, there were 66 million poor in Arab countries in 2018 or around 15% of the total Arab population. Intricately related to poverty is income inequality, which is measured by Inequality-adjusted Human Development Index (IHDI).

The poverty headcount ratio is an index which measures the percentage of poor people whose income is less than the absolute or relative poverty line (World Bank 2018). The poverty gap is a ratio showing the average shortfall of the total population from the poverty line. In other words, it reflects the intensity of poverty in a nation (World Bank 2019). The poverty line is the minimum level of income required to secure the basic necessities for survival.

#### Inequality

The loss to human development due to inequality over the past few years (2010 to 2017) is consistently more significant in southern Mediterranean countries than northern Mediterranean countries (UNDP 2018d). There are many indicators for measuring income inequalities. Considering a combination of indices would help to better understand the income distribution, because each index is not complete and each of them has strengths as well as weaknesses: 20:20 ratio, Palma ratio, the Gini index, Human Development Index (HDI) and Inequality-adjusted Human Development Index (IHDI).

An inequality measure is often a function that ascribes a value to a specific distribution of income in a way that allows direct and objective comparisons across different distributions. The "20:20 ratio" compares the ratio of the average income of the richest 20 percent of the population to the average income of the poorest 20 percent of the population. In UN reports, it is called "income quintile ratio" (UNDP 2019). The Palma ratio is defined as the ratio of the richest

10% of the population's share of gross national income divided by the poorest 40%'s share (Cobham and Sumner 2013). The Gini index measures the extent to which the distribution of income (or, in some cases, consumption expenditure) among individuals or households within an economy deviates from a perfectly equal distribution. The Gini index measures the area between the Lorenz curve, indicating the inequality in income spread, and the hypothetical line of absolute equality, expressed as a percentage of the maximum area under the line. A Gini index of zero represents perfect equality and 100 perfect inequality (OECD 2006; UNDP 2018d).

The Human Development Index (HDI) is a summary measure of average achievement in key dimensions of human development: a long and healthy life, being knowledgeable and have a decent standard of living. The HDI is the geometric mean of normalized indices for each of the three dimensions. The health dimension is assessed by life expectancy at birth, the education dimension is measured by mean of years of schooling for adults aged 25 years and more and expected years of schooling for children of school entering age. The standard of living dimension is measured by gross national income per capita. The HDI uses the logarithm of income, to reflect the diminishing importance of income with increasing Gross National Income (GNI). The scores for the three HDI dimension indices are then aggregated into a composite index using geometric mean (UNDP 2018c).

The Inequality-adjusted Human Development Index (IHDI) is a viable measure of inequality produced by the UNDP (2018c). The IHDI combines a country's average achievements in health, education and income with how those achievements are distributed among a country's population by "discounting" each dimension's average value according to its level of inequality. Thus, the IHDI is a distribution-sensitive average level of HD. Two countries with different distributions of achievements can have the same average HDI value.

Under perfect equality, the IHDI is equal to the HDI, but falls below the traditional Human Development Index (HDI) when inequality rises (UNDP 2018c). The difference between the IHDI and HDI is the human development cost of inequality, also termed – the loss to human development due to inequality. The IHDI allows a direct link to inequalities in dimensions, it can inform policy makers on how to reduce inequality, and leads to a better understanding of inequalities across populations and their contribution to the overall human development cost.

**BOX 5.1.2****Gender-related development indicators and term****Gender**

The Gender Development Index (GDI) measures gender gaps in human development achievements by accounting for disparities between women and men in three basic dimensions of human development—health, knowledge and living standards - using the same component indicators as in the HDI. The GDI is the ratio of the HDIs calculated separately for females and males using the same methodology as in the HDI. It is a direct measure of gender gap showing the female HDI as a percentage of the male HDI (UNDP 2018a).

The Gender Inequality Index (GII) measures the importance of gender in inequality. Gender inequality remains a major barrier to human development. Girls and women have made major strides since 1990, but they have not yet

gained gender equity. The disadvantages facing women and girls are a major source of inequality. All too often, women and girls are discriminated against in health, education, political representation or the labor market—with negative consequences for the development of their capabilities and their freedom of choice. The GII measures gender inequalities in three important aspects of human development: i) reproductive health, measured by maternal mortality ratio and adolescent birth rates; ii) empowerment, measured by proportion of parliamentary seats occupied by females and proportion of adult females and males aged 25 years and older with at least some secondary education; and iii) economic status, expressed as labor market participation and measured by the labor force participation rate of female and male populations aged 15 years and older. It measures the human development costs of gender inequality. Thus, the higher the GII value the more disparities between females and males and the more loss to human development (UNDP 2018b).

**BOX 5.1.3****Vulnerability**

The term vulnerability relates to the negative consequences of natural hazards, and is used in economics, hazard and disaster management in different ways (Karagiorgos et al. 2016). Economic vulnerability, at country level, may be defined as inherent proneness to exogenous shocks over which the country can exert little or no control (Briguglio et al. 2009). A widely used measurement

of economic vulnerability has been proposed by Briguglio (2010), through an index which attempts to quantify the factors that lead to exposure to economic shocks, which include, among other variables, proneness to disasters, or natural hazards. There are few comprehensive studies on natural hazards for the entire Mediterranean Region (Lionello 2012; Lionello et al. 2014), and the fragmentation of available data does not always allow comparative studies that can extend applications to the whole Basin (González Tánago et al. 2016).



## References

- Abdul Malak D, McGlade K, Pascual D, Pla E 2017 *Adapting to Climate Change. An Assessment of Vulnerability and Risks to Human Security in the Western Mediterranean Basin*. First. SpringerBriefs in Environmental Science. doi: [10.1007/978-3-319-51680-6\\_5](https://doi.org/10.1007/978-3-319-51680-6_5)
- ADB, African Development Bank 2011 Examples Of CDM Projects. [https://www.afdb.org/fileadmin/uploads/afdb/Documents/Generic-Documents/P2\\_Examples\\_of\\_CDM\\_projects\\_AfDB\\_Dba\\_210911.pdf](https://www.afdb.org/fileadmin/uploads/afdb/Documents/Generic-Documents/P2_Examples_of_CDM_projects_AfDB_Dba_210911.pdf) [Accessed February 1, 2020].
- Agard J, Schipper ELF, Birkmann J, Campos M, Dubeux C et al. 2014 Glossary, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 1757–1758.
- Albizua A, Corbera E, Pascual U 2019 Farmers' vulnerability to global change in Navarre, Spain: large-scale irrigation as maladaptation. *Reg. Environ. Chang.* 19, 1147–1158. doi: [10.1007/s10113-019-01462-2](https://doi.org/10.1007/s10113-019-01462-2)
- Almenar J, Alcaraz C, Canseco O 2019 Environmental law and practice in Spain: overview. Thomson Reuters Practical Law. [https://uk.practicallaw.thomsonreuters.com/0-521-6274?\\_lrTS=20200415181111583&transitionType=Default&contextData=%28sc.Default%29](https://uk.practicallaw.thomsonreuters.com/0-521-6274?_lrTS=20200415181111583&transitionType=Default&contextData=%28sc.Default%29)
- Amelung B, Nicholls S, Viner D 2007 Implications of Global Climate Change for Tourism Flows and Seasonality. *J. Travel Res.* 45, 285–296. doi: [10.1177/0047287506295937](https://doi.org/10.1177/0047287506295937)
- Amengual A, Homar V, Romero R, Ramis C, Alonso S 2014 Projections for the 21<sup>st</sup> century of the climate potential for beach-based tourism in the Mediterranean. *Int. J. Climatol.* 34, 3481–3498. doi: [10.1002/joc.3922](https://doi.org/10.1002/joc.3922)
- Anderson A 2012 Climate Change Education for Mitigation and Adaptation. *J. Educ. Sustain. Dev.* 6, 191–206. doi: [10.1177/0973408212475199](https://doi.org/10.1177/0973408212475199)
- Anderson RC 2002 Incentive-based policies for environmental management in developing countries.
- Arent DJ, Tol RS, Faust E, Hella JP, Kumar S et al. 2014 Key economic sectors and services, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 659–708.
- Ayers J, Dodman D 2010 Climate change adaptation and development I: the state of the debate. *Prog. Dev. Stud.* 10, 161–168. doi: [10.1177/146499340901000205](https://doi.org/10.1177/146499340901000205)
- Bapna M, McGray H 2008 Financing Adaptation: Opportunities for innovation and experimentation. Washington, DC.
- Barde JP 1994 Economic Instruments in Environmental Policy. Organisation for Economic Co-Operation and Development (OECD). doi: [10.1787/754416133402](https://doi.org/10.1787/754416133402)
- Barrios S, Ibañez JN 2015 Time is of the essence: adaptation of tourism demand to climate change in Europe. *Clim. Change* 132, 645–660. doi: [10.1007/s10584-015-1431-1](https://doi.org/10.1007/s10584-015-1431-1)
- Bartels W, Kurznack L, Briau L, Krimphoff J 2016 Mainstreaming the green bond market: Pathways towards common standards. The Netherlands <https://assets.kpmg/content/dam/kpmg/lu/pdf/mainstreaming-the-green-bond-market-kpmg-wwf-2016.pdf>.
- Bianco F, Lucifora A, Vagliasindi GM 2015 Fighting Environmental Crime in France: A Country Report. Catania. [https://efface.eu/sites/default/files/EFFACE\\_Fighting\\_Environmental\\_Crime\\_in\\_France.pdf](https://efface.eu/sites/default/files/EFFACE_Fighting_Environmental_Crime_in_France.pdf)
- Bird DN, Benabdallah S, Gouda N, Hummel F, Koeberl J et al. 2016 Modelling climate change impacts on and adaptation strategies for agriculture in Sardinia and Tunisia using AquaCrop and value-at-risk. *Sci. Total Environ.* 543, 1019–1027. doi: [10.1016/j.scitotenv.2015.07.035](https://doi.org/10.1016/j.scitotenv.2015.07.035)
- Blauhut V, Gudmundsson L, Stahl K 2015 Towards pan-European drought risk maps: quantifying the link between drought indices and reported drought impacts. *Environ. Res. Lett.* 10, 14008. doi: [10.1088/1748-9326/10/1/014008](https://doi.org/10.1088/1748-9326/10/1/014008)
- Boivin JP, Emorine A 2019 Environmental law and practice in France: overview. Thomson Reuters Practical Law. <https://uk.practicallaw.thomsonreuters.com/w-010-5542?transitionType=Default&contextData=%28sc.Default%29>
- Bontoux L, Bengtsson AD 2015 2035: Paths towards a Sustainable EU Economy Sustainable Transitions and the Potential of Eco-Innovation for Jobs and Economic Development in EU Eco-Industries 2035. doi: [10.2760/738600](https://doi.org/10.2760/738600)
- Bontoux L, Bengtsson AD 2016 Using Scenarios to Assess Policy Mixes for Resource Efficiency and Eco-Innovation in Different Fiscal Policy Frameworks. *Sustainability* 8, 309. doi: [10.3390/su8040309](https://doi.org/10.3390/su8040309)
- Briguglio LP 2010 Defining and assessing the risk of being harmed by climate change. *Int. J. Clim. Chang. Strateg. Manag.* 2, 23–34. doi: [10.1108/17568691011020238](https://doi.org/10.1108/17568691011020238)
- Briguglio LP, Cordina G, Farrugia N, Vella S 2009 Economic Vulnerability and Resilience: Concepts and Measurements. *Oxford Dev. Stud.* 37, 229–247. doi: [10.1080/13600810903089893](https://doi.org/10.1080/13600810903089893)
- Briguglio M 2019 Wellbeing: an economics perspective, in *Perspectives on wellbeing*, eds. Vella S, Falzon R, Azzopardi A (Brill | Sense), 145–157. doi: [10.1163/9789004394179\\_012](https://doi.org/10.1163/9789004394179_012)
- Bruckner M, Lederman D 2015 Effects of income inequality on economic growth.
- Carreño B 2019 Spain may issue green bonds to finance environmental plan. *Reuters, Environ.* <https://www.reuters.com/article/us-spain-economy/spain-may-is->

- [sue-green-bonds-to-finance-environmental-plan-minister-says-idUSKCN1R81XT](#)
- Center for Health Environment and Justice (undated) PVC Policies Across the World. [http://www.chej.org/pvfactsheets/PVC\\_Policies\\_Around\\_The\\_World.html](http://www.chej.org/pvfactsheets/PVC_Policies_Around_The_World.html) [Accessed May 30, 2020].
- Chilosi M, Martelli A, Miranti A 2019 Environmental law and practice in Italy: overview. *Thomson Reuters Pract. Law*. <https://uk.practicallaw.thomsonreuters.com/1-503-2608?transitionType=Default&context-Data={sc.Default}ac&firstPage=true&bhcp=1>
- Climate Transparency 2019 Managing the coal phase-out – a comparison of actions in G20 countries. Berlin, Germany.
- Cobham A, Sumner A 2013 Putting the Gini Back in the Bottle? “The Palma” as a Policy-Relevant Measure of Inequality. London: King’s College. doi: [10.2139/ssrn.2366974](https://doi.org/10.2139/ssrn.2366974)
- Coma R, Ribes M, Serrano E, Jiménez E, Salat J et al. 2009 Global warming-enhanced stratification and mass mortality events in the Mediterranean. *Proc. Natl. Acad. Sci. U. S. A.* 106, 6176–6181. doi: [10.1073/pnas.0805801106](https://doi.org/10.1073/pnas.0805801106)
- Cramer W, Guiot J, Fader M, Garrabou J, Gattuso J-P et al. 2018 Climate change and interconnected risks to sustainable development in the Mediterranean. *Nat. Clim. Chang.* 8, 972–980. doi: [10.1038/s41558-018-0299-2](https://doi.org/10.1038/s41558-018-0299-2)
- CTC (Carbon Tax Center) 2018 Where carbon is taxed? *Carbon Tax Cent.* <https://www.carbontax.org/where-carbon-is-taxed/> [Accessed February 2, 2020].
- Delija-Ružić W 2017 Report on Implementation of Policies and Measures that Reduce Greenhouse Gas Emissions by Sources or Enhance Removals by Sinks. Republic of Croatia.
- Dellai I, McCarl BA 2010 The economic impacts of drought on agriculture: The case of Turkey, in *Economics of Drought and Drought Preparedness in a Climate Change Context*, eds. López-Francos A, López-Francos A (Zaragoza), 169–174. <http://om.iamm.fr/om/pdf/a95/00801342.pdf> [Accessed April 7, 2019].
- Devlin RA, Grafton RQ 1996 Marketable emission permits: Efficiency, profitability and substitutability. *Can. J. Econ.* 29, S260–S264. doi: [10.2307/135997](https://doi.org/10.2307/135997)
- di Gregorio M, Nurrochmat DR, Paavola J, Sari IM, Fatorelli L et al. 2017 Climate policy integration in the land use sector: Mitigation, adaptation and sustainable development linkages. *Environ. Sci. Policy* 67, 35–43. doi: [10.1016/j.envsci.2016.11.004](https://doi.org/10.1016/j.envsci.2016.11.004)
- DIE 2014 Why Power Matters in Payments for Environmental Services (PES). [https://www.die-gdi.de/uploads/media/BP\\_9.2014.pdf](https://www.die-gdi.de/uploads/media/BP_9.2014.pdf)
- Dogru T, Bulut U, Sirakaya-Turk E 2016 Theory of Vulnerability and Remarkable Resilience of Tourism Demand to Climate Change: Evidence from the Mediterranean Basin. *Tour. Anal.* 21, 645–660. doi: [10.3727/108354216X14713487283246](https://doi.org/10.3727/108354216X14713487283246)
- Dogru T, Marchio EA, Bulut U, Suess C 2019 Climate change: Vulnerability and resilience of tourism and the entire economy. *Tour. Manag.* 72, 292–305. doi: [10.1016/J.TOURMAN.2018.12.010](https://doi.org/10.1016/J.TOURMAN.2018.12.010)
- Doocy S, Daniels A, Murray S, Kirsch TD 2013 The Human Impact of Floods: a Historical Review of Events 1980–2009 and Systematic Literature Review. *PLoS Curr.* 5. doi: [10.1371/CURRENTS.DIS.F4DEB457904936B07C-09DAA98EE8171A](https://doi.org/10.1371/CURRENTS.DIS.F4DEB457904936B07C-09DAA98EE8171A)
- Dos Santos MJPL 2018 Nowcasting and forecasting aquaponics by Google Trends in European countries. *Technol. Forecast. Soc. Change* 134, 178–185. doi: [10.1016/j.techfore.2018.06.002](https://doi.org/10.1016/j.techfore.2018.06.002)
- Dos Santos MJPL, Mota M 2019 Sustainable and Smart Cities: The Case Study of African Cities. in *The 4th International Conference on Organization and Management. The 6th Corporate Social Responsibility (CSR), Ethics, Governance, and Sustainability*. (Abu Dhabi, UAE, UAE on 12th and 13th of June 2019: College of Abu Dhabi University).
- Drius M, Bongiorno L, Depellegrin D, Menegon S, Pugnetti A et al. 2019 Tackling challenges for Mediterranean sustainable coastal tourism: An ecosystem service perspective. *Sci. Total Environ.* 652, 1302–1317. doi: [10.1016/j.scitotenv.2018.10.121](https://doi.org/10.1016/j.scitotenv.2018.10.121)
- Dubois G, Ceron J-P 2006 Tourism and Climate Change: Proposals for a Research Agenda. *J. Sustain. Tour.* 14, 399–415. doi: [10.2167/jost539.0](https://doi.org/10.2167/jost539.0)
- Dubois G, Ceron J-P, Gössling S, Hall CM 2016 Weather preferences of French tourists: lessons for climate change impact assessment. *Clim. Change* 136, 339–351. doi: [10.1007/s10584-016-1620-6](https://doi.org/10.1007/s10584-016-1620-6)
- Enríquez AR, Marcos M, Álvarez-Ellacuría A, Orfila A, Gomis D 2017 Changes in beach shoreline due to sea level rise and waves under climate change scenarios: application to the Balearic Islands (western Mediterranean). *Nat. Hazards Earth Syst. Sci.* 17, 1075–1089. doi: [10.5194/nhess-17-1075-2017](https://doi.org/10.5194/nhess-17-1075-2017)
- Environmental Rights Database Environmental Education Programme. <http://environmentalrightsdatabase.org/environmental-education-programme/>.
- FAO / IFAD / WFP 2011 The State of Food Insecurity in the World. Rome, Italy.
- Fatorić S, Seekamp E 2017 Are cultural heritage and resources threatened by climate change? A systematic literature review. *Clim. Change* 142, 227–254. doi: [10.1007/s10584-017-1929-9](https://doi.org/10.1007/s10584-017-1929-9)
- Forbes KJ 2000 A Reassessment of the Relationship Between Inequality and Growth. *Am. Econ. Rev.* 90, 869–887. doi: [10.1257/aer.90.4.869](https://doi.org/10.1257/aer.90.4.869)
- Galor O, Zeira J 1993 Income Distribution and Macroeconomics. *Rev. Econ. Stud.* 60, 35. doi: [10.2307/2297811](https://doi.org/10.2307/2297811)
- GEF 2019 EBRD/GEF Environmental Credit Facility (formerly entitled Slovenia: National Pollution Reduction Project). <https://www.thegef.org/project/ebdref-environmental-credit-facility-formerly-entitled-slovenia-national-pollution>

- Godber O, Wall R 2016 Mediterranean goat production systems: vulnerability to population growth and climate change. *Mediterr. J. Biosci.* 1, 160–168.
- González Tánago I, Urquijo J, Blauhut V, Villarroya F, de Stefano L 2016 Learning from experience: a systematic review of assessments of vulnerability to drought. *Nat. Hazards* 80, 951–973. doi: [10.1007/s11069-015-2006-1](https://doi.org/10.1007/s11069-015-2006-1)
- Grgi Z, Frani R, Kisi I Country report on the present environmental situation in agriculture. *REU Tech. Ser.* <http://www.fao.org/3/x3413e12.htm> [Accessed May 30, 2020].
- Grillakis MG, Koutroulis AG, Seiradakis KD, Tsanis IK 2016a Implications of 2 °C global warming in European summer tourism. *Clim. Serv.* 1, 30–38. doi: [10.1016/j.cliser.2016.01.002](https://doi.org/10.1016/j.cliser.2016.01.002)
- Grillakis MG, Koutroulis AG, Tsanis IK 2016b The 2 °C global warming effect on summer European tourism through different indices. *Int. J. Biometeorol.* 60, 1205–1215. doi: [10.1007/s00484-015-1115-6](https://doi.org/10.1007/s00484-015-1115-6)
- Hamrick K, Brotto L 2017 State of European markets 2017 voluntary carbon. [https://www.forest-trends.org/wp-content/uploads/2017/07/doc\\_5591.pdf](https://www.forest-trends.org/wp-content/uploads/2017/07/doc_5591.pdf) [Accessed November 1, 2019].
- Handmer J, Honda Y, Kundzewicz ZW, Arnell NW, Benito G et al. 2012 Changes in impacts of climate extremes: human systems and ecosystems, in *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation* A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC)., eds. Field CB, Barros V, Stocker TF, Qin D, Dokken DJ et al. (Cambridge, United Kingdom and New York, NY, USA), 231–290.
- Helliwell JF, Layard R, Sachs JD, Helliwell, JF, Layard, R Sachs J 2018 World Happiness report. Sustainable Development Solution Network. A global initiative for the United Nations. UN. doi: [10.18356/6aa6d28a-en](https://doi.org/10.18356/6aa6d28a-en)
- Iglesias A, Garrote L 2014 Drought in the Light of Climate Change in the Mediterranean Area, in *Hydrometeorological Hazards*, ed. Quevauviller P (Chichester, UK: John Wiley & Sons, Ltd), 203–225. doi: [10.1002/9781118629567.ch3c](https://doi.org/10.1002/9781118629567.ch3c)
- Jacob D, Kotova L, Teichmann C, Sobolowski SP, Vautard R et al. 2018 Climate Impacts in Europe Under +1.5°C Global Warming. *Earth's Futur.* 6, 264–285. doi: [10.1002/2017EF000710](https://doi.org/10.1002/2017EF000710)
- Jacob D, Petersen J, Eggert B, Alias A, Christensen OB et al. 2014 EURO-CORDEX: new high-resolution climate change projections for European impact research. *Reg. Environ. Chang.* 14, 563–578. doi: [10.1007/s10113-013-0499-2](https://doi.org/10.1007/s10113-013-0499-2)
- Jiménez JA, Valdemoro HI, Bosom E, Sánchez-Arcilla A, Nicholls RJ 2017 Impacts of sea-level rise-induced erosion on the Catalan coast. *Reg. Environ. Chang.* 17, 593–603. doi: [10.1007/s10113-016-1052-x](https://doi.org/10.1007/s10113-016-1052-x)
- Justice and Environment 2012 Slovenia. Environmental Liability: National toolkit on the practical application of the ELD and its national equivalents. 1–6. [http://www.justiceandenvironment.org/files/file/2012/2012 ELD report Slovenia.pdf](http://www.justiceandenvironment.org/files/file/2012/2012%20ELD%20report%20Slovenia.pdf)
- Justice and Environment 2017 Spain. Environmental Liability: National toolkit on the practical application of the ELD and its national equivalents. 1–6. [http://www.justiceandenvironment.org/fileadmin/user\\_upload/Publications/2017/ELD\\_toolkit\\_ES\\_EN.pdf](http://www.justiceandenvironment.org/fileadmin/user_upload/Publications/2017/ELD_toolkit_ES_EN.pdf)
- Karagiorgos K, Thaler T, Heiser M, Hübl J, Fuchs S 2016 Integrated flash flood vulnerability assessment: Insights from East Attica, Greece. *J. Hydrol.* 541A, 553–562. doi: [10.1016/j.jhydrol.2016.02.052](https://doi.org/10.1016/j.jhydrol.2016.02.052)
- Kazan-Allen L 2019 Chronology of asbestos bans and restrictions. *Int. Ban Asbestos Secr.* [http://www.ibasecretariat.org/chron\\_ban\\_list.php](http://www.ibasecretariat.org/chron_ban_list.php)
- Kelly PM, Adger WN 2000 Theory and practice in assessing vulnerability to climate change and facilitating adaptation. *Clim. Change* 47, 325–352. doi: [10.1023/a:1005627828199](https://doi.org/10.1023/a:1005627828199)
- Kilkiş Ş 2018 Benchmarking South East European Cities with the Sustainable Development of Energy, Water and Environment Systems Index. *J. Sustain. Dev. Energy, Water Environ. Syst.* 6, 162–209. doi: [10.13044/j.sdewes.d5.0179](https://doi.org/10.13044/j.sdewes.d5.0179)
- Klein RJT 2010 Mainstreaming Climate Adaptation into Development: A Policy Dilemma. *Clim. Gov. Dev. Berlin Work. Ser.*, 35–52. doi: [10.1596/9780821379943\\_ch01](https://doi.org/10.1596/9780821379943_ch01)
- Köberl J, Prettenhaler F, Bird DN 2016 Modelling climate change impacts on tourism demand: A comparative study from Sardinia (Italy) and Cap Bon (Tunisia). *Sci. Total Environ.* 543, 1039–1053. doi: [10.1016/j.scitotenv.2015.03.099](https://doi.org/10.1016/j.scitotenv.2015.03.099)
- Koutra C, Karyopoulou S 2013 Cyprus' image-a sun and sea destination-as a detrimental factor to seasonal fluctuations. Exploration into motivational factors for holidaying in Cyprus. *J. Travel Tour. Mark.* 30, 700–714. doi: [10.1080/10548408.2013.827548](https://doi.org/10.1080/10548408.2013.827548)
- Koutroulis AG, Grillakis MG, Tsanis IK, Jacob D 2018a Mapping the vulnerability of European summer tourism under 2 °C global warming. *Clim. Change* 151, 157–171. doi: [10.1007/s10584-018-2298-8](https://doi.org/10.1007/s10584-018-2298-8)
- Koutroulis AG, Papadimitriou L V., Grillakis MG, Tsanis IK, Wyser K et al. 2018b Simulating Hydrological Impacts under Climate Change: Implications from Methodological Differences of a Pan European Assessment. *Water* 10, 1331. doi: [10.3390/w10101331](https://doi.org/10.3390/w10101331)
- Kovats RS, Valentini R, Bouwer LM, Georgopoulou E, Jacob D et al. 2014 Europe, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 1267–1326. doi: [10.1017/cbo9781107415386.003](https://doi.org/10.1017/cbo9781107415386.003)
- Kraus H 1998 Environment Policy in Slovenia. [https://www.europarl.europa.eu/workingpapers/envi/pdf/brief6en\\_en.pdf](https://www.europarl.europa.eu/workingpapers/envi/pdf/brief6en_en.pdf)



- Lewenhak S 2012 The Role of the European Investment Bank (RLE Banking & Finance). doi: [10.4324/9780203108246](https://doi.org/10.4324/9780203108246)
- Lionello P 2012 *The Climate of the Mediterranean Region: From the Past to the Future*. Elsevier Science doi: [10.1016/C2011-0-06210-5](https://doi.org/10.1016/C2011-0-06210-5)
- Lionello P, Abrantes FG, Gačić M, Planton S, Trigo RM et al. 2014 The climate of the Mediterranean region: research progress and climate change impacts. *Reg. Environ. Chang.* 14, 1679–1684. doi: [10.1007/s10113-014-0666-0](https://doi.org/10.1007/s10113-014-0666-0)
- Littlecott C 2015 Snapshot of France Coal Phase Out Progress. <https://www.e3g.org/library/snapshot-of-france-coal-phase-out-progress>
- Llasat MC, Llasat-Botija M, Petrucci O, Pasqua AA, Rosselló J et al. 2013 Towards a database on societal impact of Mediterranean floods within the framework of the HYMEX project. *Nat. Hazards Earth Syst. Sci.* 13, 1337–1350. doi: [10.5194/nhess-13-1337-2013](https://doi.org/10.5194/nhess-13-1337-2013)
- Llasat MC, Llasat-Botija M, Prat MA, PorcúPorc F, Price C et al. 2010 High-impact floods and flash floods in Mediterranean countries: the FLASH preliminary database. [www.adv-geosci.net/23/47/2010/](http://www.adv-geosci.net/23/47/2010/) [Accessed April 2, 2019]
- Mavioglu OY, Varol P, Tamtürk D 2019 Environmental law and practice in Turkey: overview. *Thomson Reuters Pract. Law.* <https://uk.practicallaw.thomsonreuters.com/Document/I203079191cb611e38578f7ccc38dcbee/>
- Mavromatidi A, Briche E, Claeys C 2018 Mapping and analyzing socio-environmental vulnerability to coastal hazards induced by climate change: An application to coastal Mediterranean cities in France. *Cities* 72, 189–200. doi: [10.1016/J.CITIES.2017.08.007](https://doi.org/10.1016/J.CITIES.2017.08.007)
- McGray H, Hammil I. A, Bradley R, Schipper ELF, Parry J 2007 *Weathering the storm: options for framing adaptation and development*, ed. World Resources Institute Washington, DC.
- Mediterranean Growth Initiative 2017 Tourism in the Mediterranean. <https://www.mgi.online/content/2017/8/4/tourism-in-the-mediterranean>
- MESD 2018 Education for sustainable development in Tunisia. <http://www.environnement.gov.tn/index.php?id=101&L=1>
- Milewska M 2019 Resource Taxation as an Alternative Method for Diversifying the Revenue of Developing Countries. <https://www.progress.org/articles/resource-taxation-as-an-alternative-method-for-diversifying-the-revenue-of-developing-countries>
- Mimura N, Pulwarty RS, Minh Duc D, Elshinnawy I, Redsteer MH et al. 2014 Adaptation Planning and Implementation, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 869–898. doi: [10.1017/CBO9781107415379.020](https://doi.org/10.1017/CBO9781107415379.020)
- Minasyan A, Zenker J, Klases S, Vollmer S 2019 Educational gender gaps and economic growth: A systematic review and meta-regression analysis. *World Dev.* 122, 199–217. doi: [10.1016/j.worlddev.2019.05.006](https://doi.org/10.1016/j.worlddev.2019.05.006)
- Moncada S, Briguglio LP, Bambrick H, Kelman I 2018 Guest editorial for the special issue on 'Development and Climate Change in Small Island Developing States'. *Int. J. Clim. Chang. Strateg. Manag.* 10, 214–216. doi: [10.1108/ijccsm-03-2018-184](https://doi.org/10.1108/ijccsm-03-2018-184)
- Monioudi IN, Velegarakis AF, Chatzipavlis AE, Rigos A, Karambas T et al. 2017 Assessment of island beach erosion due to sea level rise: the case of the Aegean archipelago (Eastern Mediterranean). *Nat. Hazards Earth Syst. Sci.* 17, 449–466. doi: [10.5194/nhess-17-449-2017](https://doi.org/10.5194/nhess-17-449-2017)
- Ncube M, Anyanwu JC, Hausken K 2014 Inequality, Economic Growth and Poverty in the Middle East and North Africa (MENA). *African Dev. Rev.* 26, 435–453. doi: [10.1111/1467-8268.12103](https://doi.org/10.1111/1467-8268.12103)
- Niavis S 2020 Evaluating the spatiotemporal performance of tourist destinations: the case of Mediterranean coastal regions. *J. Sustain. Tour.* 28, 1310–1331. doi: [10.1080/09669582.2020.1736087](https://doi.org/10.1080/09669582.2020.1736087)
- OECD 2003 *Review of Fisheries in OECD countries: Policies and summary statistics*. OECD Publication Service.
- OECD 2006 Glossary of statistical terms. <https://stats.oecd.org/glossary/detail.asp?ID=4842>.
- OECD 2012a Liability for environmental damage in Eastern Europe, Caucasus and Central Asia (EECCA): Implementation of good international practices. <http://www.oecd.org/env/outreach/50244626.pdf>
- OECD 2012b OECD Environmental Performance Reviews: Slovenia 2012. *OECD Publ.*
- OECD 2013 OECD Environmental Performance Reviews: Italy 2012. *OECD Publ.*
- OECD 2017 Measuring wellbeing and progress wellbeing research. <http://www.oecd.org/statistics/measuring-well-being-and-progress.htm>
- OECD 2018 OECD Review for Fisheries 2017: General Survey of Fisheries Policies. [http://www.oecd.org/official-documents/publicdisplaydocumentpdf/?cote=TAD/FI\(2017\)14/FINAL&docLanguage=En](http://www.oecd.org/official-documents/publicdisplaydocumentpdf/?cote=TAD/FI(2017)14/FINAL&docLanguage=En)
- OECD 2019 OECD Environmental Performance Reviews: Turkey 2019. *OECD Publ.*
- Oliveira S, Félix F, Nunes A, Lourenço L, Laneve G et al. 2018 Mapping wildfire vulnerability in Mediterranean Europe. Testing a stepwise approach for operational purposes. *J. Environ. Manage.* 206, 158–169. doi: [10.1016/J.JENVMAN.2017.10.003](https://doi.org/10.1016/J.JENVMAN.2017.10.003)
- Palatnik RR, Lourenço Dias Nunes PA 2015 Economic valuation of climate change-induced biodiversity impacts on agriculture: results from a macro-economic application to the Mediterranean basin. *J. Environ. Econ. Policy* 4, 45–63. doi: [10.1080/21606544.2014.963165](https://doi.org/10.1080/21606544.2014.963165)
- Pelling M 2011 *Adaptation to climate change: from resilience to transformation*. First edn. Routledge, New York, USA.

- Perman R, Ma Y, McGilvray J, Common M 2003 Natural Resource and Environmental Economics. *Mar. Resour. Econ.* 7. doi: [10.1086/mre.7.4.42629040](https://doi.org/10.1086/mre.7.4.42629040)
- Pettinger T 2017 Pollution Permits. <https://www.economicshelp.org/micro-economic-essays/marketfailure/pollution-permits/>
- Piante C, Ody D 2015 Blue Growth in the Mediterranean Sea: The Challenge of Good Environmental Status. <http://www.developpement-durable.gouv.fr/>
- Praščević M, Cvetković D, Mihajlov D, Holoček N 2012 The acoustic zoning - a comparison of legislation and experiences in Italy and Serbia. in *23rd National Conference & 4th International Conference: Noise and Vibration, 17-19.10.2012* (University of Nis (Serbia), "Politehnica" University of Timisoara (Romania)). <http://www.mfkv.kg.ac.rs/urbanoise/media/1210>  
Nis Prascevic 21-28.pdf
- Queralt A, Llasat MDC, Serena JM, Pont I 2017 The 3rd Report on Climate Change in Catalonia: a joint initiative to bridge the gap between Science and Decision-Making. In EGU General Assembly. in *EGU General Assembly Conference Abstracts 19:14658*
- Radhouane L 2013 Climate change impacts on North African countries and on some Tunisian economic sectors. *J. Agric. Environ. Int. Dev.* 107, 101-113. doi: [10.12895/JAEID.20131.123](https://doi.org/10.12895/JAEID.20131.123)
- Recchini E 2016 Estimating emission permits in Italy. in *Joint OECD/UNECE Seminar on Implementation of SEEA, 3-4 October 2016* (Geneva, Switzerland). [https://uk.practicallaw.thomsonreuters.com/1-503-2608?transitionType=Default&contextData=\(sc.Default\)&firstPage=true&bhcp=1](https://uk.practicallaw.thomsonreuters.com/1-503-2608?transitionType=Default&contextData=(sc.Default)&firstPage=true&bhcp=1)
- Richards M, Jenkins M 2007 Potential and challenges of payments for ecosystem service from tropical forests.
- Rodrigues LC, Van den Bergh JCJM, Massa F, Theodorou JA, Ziveri P et al. 2015 Sensitivity of Mediterranean bivalve mollusc aquaculture to climate change, ocean acidification, and other environmental pressures: findings from a producer survey. *J. Shellfish Res.* 34, 1161-1176. doi: [10.2983/035.034.0341](https://doi.org/10.2983/035.034.0341)
- Rodriguez RS, Üрге-Vorsatz D, Barau AS 2018 Sustainable Development Goals and climate change adaptation in cities. *Nat. Clim. Chang.* 8, 181-183. doi: [10.1038/s41558-018-0098-9](https://doi.org/10.1038/s41558-018-0098-9)
- Roy J, Tschakert P, Waisman H, Abdul Halim S, Antwi-Agyei P et al. 2018 Sustainable development, poverty eradication and reducing inequalities, in *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, eds. Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), in press.
- Sacchi J 2011 Analysis of economic activities in the Mediterranean: Fishery and aquaculture sectors. Valbonne [https://planbleu.org/sites/default/files/upload/files/analyse\\_activites\\_econ\\_pecheEN.pdf](https://planbleu.org/sites/default/files/upload/files/analyse_activites_econ_pecheEN.pdf)
- Sachs J 2005 *The End of Poverty: How We Can Make It Happen in Our Lifetime*. Penguin Bo. Wiley. doi: [10.1111/j.1468-0351.2006.00262.x](https://doi.org/10.1111/j.1468-0351.2006.00262.x)
- Salanié J, Coisson T 2016 Environmental Zoning and Urban Development: Natural Regional Parks in France. doi: [10.1787/5jlsk97vpwtd-en](https://doi.org/10.1787/5jlsk97vpwtd-en)
- Salata F, Golasi I, Treiani N, Plos R, de Lieto Vollaro A 2018 On the outdoor thermal perception and comfort of a Mediterranean subject across other Koppen-Geiger's climate zones. *Environ. Res.* 167, 115-128. doi: [10.1016/j.envres.2018.07.011](https://doi.org/10.1016/j.envres.2018.07.011)
- Sánchez-Arcilla A, Sierra JP, Brown S, Casas-Prat M, Nicholls RJ et al. 2016 A review of potential physical impacts on harbours in the Mediterranean Sea under climate change. *Reg. Environ. Chang.* 16, 2471-2484. doi: [10.1007/s10113-016-0972-9](https://doi.org/10.1007/s10113-016-0972-9)
- Satterthwaite D 2013 The political underpinnings of cities' accumulated resilience to climate change. *Environ. Urban.* 25, 381-391. doi: [10.1177/0956247813500902](https://doi.org/10.1177/0956247813500902)
- Schilling J, Freier KP, Hertig E, Scheffran J 2012 Climate change, vulnerability and adaptation in North Africa with focus on Morocco. *Agric. Ecosyst. Environ.* 156, 12-26. doi: [10.1016/J.AGEE.2012.04.021](https://doi.org/10.1016/J.AGEE.2012.04.021)
- Scott D, Hall CM, Gössling S 2016a A review of the IPCC Fifth Assessment and implications for tourism sector climate resilience and decarbonization. *J. Sustain. Tour.* 24, 8-30. doi: [10.1080/09669582.2015.1062021](https://doi.org/10.1080/09669582.2015.1062021)
- Scott D, Ruty M, Amelung B, Tang M 2016b An Inter-Comparison of the Holiday Climate Index (HCI) and the Tourism Climate Index (TCI) in Europe. *Atmosphere (Basel)*. 7, 80. doi: [10.3390/atmos7060080](https://doi.org/10.3390/atmos7060080)
- Sen A 1999 *Development as freedom*. Oxford Uni. Oxford India paperbacks.
- Smit B, Wandel J 2006 Adaptation, adaptive capacity and vulnerability. *Glob. Environ. Chang.* 16, 282-292. doi: [10.1016/j.gloenvcha.2006.03.008](https://doi.org/10.1016/j.gloenvcha.2006.03.008)
- Stokes E, Edge A, West A 2001 Environmental education in the educational systems of the European Union. <http://ibdigital.uib.es/greenstone/collect/cd2/import/ue/ue0002.pdf>
- SwitchMed 2017 Lebanon: Loans available to enterprise to implement pollution control projects. <https://www.switchmed.eu/en/country-hubs/lebanon/actions/policy-maker?c=policy&q=lebanon>
- Szopik-Depczyńska K, Cheba K, Bąk I, Kiba-Janiak M, Saniuk S et al. 2017 The application of relative taxonomy to the study of disproportions in the area of sustainable development of the European Union. *Land use policy* 68, 481-491. doi: [10.1016/j.landusepol.2017.08.013](https://doi.org/10.1016/j.landusepol.2017.08.013)
- Tansel A, Güngör ND 2016 Gender Effects of Education on Economic Development in Turkey: The Role of Socio-demographics, Entrepreneurship and Public Policies. *Women, Work Wellf. Middle East North Africa*, 57-86. doi: [10.1142/9781783267347\\_0003](https://doi.org/10.1142/9781783267347_0003)
- Teotónio C, Rodríguez M, Roebeling P, Fortes P 2020 Water

- competition through the 'water-energy' nexus: Assessing the economic impacts of climate change in a Mediterranean context. *Energy Econ.* 85, 104539. doi: [10.1016/j.eneco.2019.104539](https://doi.org/10.1016/j.eneco.2019.104539)
- Tigkas D, Tsakiris G 2015 Early Estimation of Drought Impacts on Rainfed Wheat Yield in Mediterranean Climate. *Environ. Process.* 2, 97–114. doi: [10.1007/s40710-014-0052-4](https://doi.org/10.1007/s40710-014-0052-4)
- UN 2015 Transforming our world: the 2030 Agenda for Sustainable Development. New York. [https://www.un.org/ga/search/view\\_doc.asp?symbol=A/RES/70/1&Lang=E](https://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E)
- UNDP 2018a Gender Development Index (GDI). *United Nations Dev. Program.* <http://hdr.undp.org/en/content/gender-development-index-gdi>
- UNDP 2018b Gender Inequality Index (GII). *United Nations Dev. Program.* <http://hdr.undp.org/en/content/gender-inequality-index-gii>
- UNDP 2018c Human Development Index (HDI). *United Nations Dev. Program.* <http://hdr.undp.org/en/content/human-development-index-hdi>
- UNDP 2018d Inequality-adjusted Human Development Index (IHDI). *United Nations Dev. Program.* <http://hdr.undp.org/en/content/inequality-adjusted-human-development-index-ihdi>
- UNDP 2018e The 2018 Global Multidimensional Poverty Index (MPI). *United Nations Dev. Program.* <http://hdr.undp.org/en//2018-MPI>
- UNDP 2019 Human Development Report.
- UNECE 2013 Strategies and Policies for Air Pollution Abatement. 2010 review prepared under the Convention on Long-range Transboundary Air Pollution. New York and Geneva. [http://www.unece.org/fileadmin/DAM/env/documents/2013/air/FINAL\\_version\\_of\\_the\\_publication\\_as\\_of\\_19\\_Feb\\_2014.pdf](http://www.unece.org/fileadmin/DAM/env/documents/2013/air/FINAL_version_of_the_publication_as_of_19_Feb_2014.pdf)
- UNECE 2014 Environmental Performance Reviews: Croatia. New York and Geneva. [https://www.unece.org/fileadmin/DAM/env/epr/epr\\_studies/ECE\\_CEP\\_172\\_En.pdf](https://www.unece.org/fileadmin/DAM/env/epr/epr_studies/ECE_CEP_172_En.pdf)
- UNECE 2018 Environmental Performance Reviews: Bosnia and Herzegovina. New York and Geneva, Environmental Performance Reviews Series No. 48. [https://www.unece.org/fileadmin/DAM/env/epr/epr\\_studies/ECE\\_CEP.184.Eng.pdf](https://www.unece.org/fileadmin/DAM/env/epr/epr_studies/ECE_CEP.184.Eng.pdf)
- UNEP/MAP-Plan Bleu 2009 State of the Environment and Development in the Mediterranean. Athens, Greece.
- UNEP/MAP 2016 Mediterranean Strategy for Sustainable Development 2016-2025. Valbonne.
- UNEP 2015 Spain Air Quality Policies. <https://wedocs.unep.org/bitstream/handle/20.500.11822/17112/Spain.pdf?sequence=1&isAllowed=y>
- UNFCCC 2012 Benefits of the clean development mechanism. [https://unfccc.int/resource/docs/publications/abc\\_2012.pdf](https://unfccc.int/resource/docs/publications/abc_2012.pdf).
- UNWTO 2018 UNWTO Tourist Highlights 2017. *World Tour. Organ.* <https://www.e-unwto.org/doi/pdf/10.18111/9789284419876> [Accessed December 11, 2019].
- UNWTO 2018 Tourism and Sustainable Development Goals - Journey to 2030. <https://www.e-unwto.org/doi/pdf/10.18111/9789284419401>
- UNWTO 2019 International Tourism Highlights. <https://www.e-unwto.org/doi/pdf/10.18111/9789284421152>
- Voegtlin C, Scherer AG 2017 Responsible Innovation and the Innovation of Responsibility: Governing Sustainable Development in a Globalized World. *J. Bus. Ethics* 143, 227–243. doi: [10.1007/s10551-015-2769-z](https://doi.org/10.1007/s10551-015-2769-z)
- Walls M 2011 Deposit-Refund Systems in Practice and Theory. doi: [10.2139/ssrn.1980142](https://doi.org/10.2139/ssrn.1980142)
- White HB 2018 France to shut all coal-fired power stations by 2021, Macron declares. *Independent.* <https://www.independent.co.uk/news/world/europe/france-coal-power-station-emmanuel-macron-davos-shut-2021-a8176796.html>
- World Bank 2018 Poverty gap at \$1.90 a day (2011 PPP) (%). <https://data.worldbank.org/indicator/SI.POV.GAPS>
- World Bank 2019 World Development Indicators. <https://databank.worldbank.org/source/world-development-indicators>
- Zahumenská V, Lemeš S, Delalić M, Zatloukalová K, Sobotková J et al. 2015 Environmental Democracy in Bosnia and Herzegovina: Limping Along. Zenica, Prague [https://books.google.com.lb/books?id=RhacCwAAQBA-J&pg=PA80&lpg=PA80&dq=environmental+zoning+in+bosnia+and+herzegovina&source=bl&ots=D9RjoNpn0l&sig=ACfU3U0PTIC4LYBxjOrXMByp4Rz-l34aog&hl=en&sa=X&ved=2ahUKewjw7\\_hpdPkAhXU-BUIHakABXcQ6AEwCHoECAkQAQ#v=onepage&q=e](https://books.google.com.lb/books?id=RhacCwAAQBA-J&pg=PA80&lpg=PA80&dq=environmental+zoning+in+bosnia+and+herzegovina&source=bl&ots=D9RjoNpn0l&sig=ACfU3U0PTIC4LYBxjOrXMByp4Rz-l34aog&hl=en&sa=X&ved=2ahUKewjw7_hpdPkAhXU-BUIHakABXcQ6AEwCHoECAkQAQ#v=onepage&q=e)
- Zakhartchouk A, 2019 Scaling up green finance in France. [https://www.cape4financeministry.org/sites/cape/files/inline-files/Session4-Scalingup\\_Green\\_Finance\\_in\\_France.pdf](https://www.cape4financeministry.org/sites/cape/files/inline-files/Session4-Scalingup_Green_Finance_in_France.pdf)
- Zidanšek A, 2007 Sustainable development and happiness in nations. *Energy* 32, 891–897. doi: [10.1016/j.energy.2006.09.016](https://doi.org/10.1016/j.energy.2006.09.016)

## Information about authors

### Coordinating Lead Authors

Maria Dos Santos:

*Polytechnical Institute of Lisbon, University Institute of Lisbon (ISCTE-IUL), Lisbon, Portugal*

Stefano Moncada:

*University of Malta, Msida, Malta*

### Lead Authors

Antonietta Elia:

*University of Santiago de Compostela, Santiago de Compostela, Spain*

Manolis Grillakis:

*Institute of Mediterranean Studies (Institute of Technology & Research – Hellas), Rethymno, Crete, Greece*

Nathalie Hilmi:

*Monaco Scientific Centre (MSC), Monaco*

### Contributing Authors

Shekoofeh Farahmand:

*Department of Economics, University of Isfahan, Isfahan, Iran*

Walid Marrouch:

*Lebanese American University, Beirut, Lebanon*

Alain Safa:

*University of Nice Sophia-Antipolis, EDHEC Business School, Nice, France*

Brice Teisserenc:

*University of Toulon, La Garde, France*




# 5 SOCIETY 2-HEALTH

**Coordinating Lead Authors:**  
Shlomit Paz (Israel)

**Lead Authors:**  
Cristina Linares (Spain), Julio Díaz (Spain), Maya Negev (Israel), Gerardo Sánchez Martínez (Denmark)

**Contributing Authors:**  
Roberto Debono (Malta)

*This chapter should be cited as: Linares C, Paz S, Díaz J, Negev M, Sánchez Martínez G 2020 Health. In: Climate and Environmental Change in the Mediterranean Basin - Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 493-514.*



# Table of contents

<b>5.2 Health</b> .....	<b>496</b>
Executive summary.....	<b>496</b>
<b>5.2.1 Introduction</b> .....	<b>496</b>
5.2.1.1 <i>Effects of climate and environmental change on human health</i> .....	<b>496</b>
5.2.1.2 <i>Multi-factorial changes in health attributed to environmental change</i> .....	<b>496</b>
<b>5.2.2 Vulnerability and health risk – main causes</b> .....	<b>497</b>
5.2.2.1 <i>Aging</i> .....	<b>497</b>
5.2.2.2 <i>Gender</i> .....	<b>497</b>
5.2.2.3 <i>Geographic location</i> .....	<b>497</b>
5.2.2.4 <i>Socio-economic status</i> .....	<b>497</b>
5.2.2.5 <i>Level of acclimation</i> .....	<b>497</b>
5.2.2.6 <i>Occupational health</i> .....	<b>498</b>
5.2.2.7 <i>National public health infrastructures, warning systems</i> .....	<b>498</b>
5.2.2.8 <i>Influence of urban landscape</i> .....	<b>498</b>
<b>5.2.3 Health impacts: recent and current situation</b> .....	<b>498</b>
5.2.3.1 <i>Heat-related impacts</i> .....	<b>498</b>
5.2.3.2 <i>Cold-related impacts</i> .....	<b>499</b>
5.2.3.3 <i>Vector-borne diseases</i> .....	<b>500</b>
<i>West Nile virus</i> .....	<b>500</b>
<i>Chikungunya</i> .....	<b>501</b>
<i>Rift Valley fever</i> .....	<b>501</b>
<i>Leishmaniasis</i> .....	<b>501</b>
5.2.3.4 <i>Food- and water-borne diseases</i> .....	<b>501</b>
<b>5.2.4 Indirect impacts: recent and current situation</b> .....	<b>502</b>
5.2.4.1 <i>Air quality</i> .....	<b>502</b>
5.2.4.2 <i>Mineral dust and forest fires</i> .....	<b>502</b>
5.2.4.3 <i>Mental health</i> .....	<b>503</b>
5.2.4.4 <i>Migration</i> .....	<b>503</b>
<b>5.2.5 Projections for global warming of 1.5°C, 2°C and more</b> .....	<b>503</b>
5.2.5.1 <i>Vulnerabilities and risks under different warming scenarios</i> .....	<b>503</b>
5.2.5.2 <i>Heat-related impacts</i> .....	<b>503</b>
5.2.5.3 <i>Cold-related impacts</i> .....	<b>505</b>
5.2.5.4 <i>Vector- food- and water-borne diseases</i> .....	<b>505</b>
5.2.5.5 <i>Air quality</i> .....	<b>505</b>
<b>5.2.6 Resilience, preparedness and adaptation</b> .....	<b>506</b>
5.2.6.1 <i>Health preparedness and adaptation measures</i> .....	<b>506</b>
5.2.6.2 <i>Regional coordination and collaboration</i> .....	<b>506</b>
<b>References</b> .....	<b>507</b>
<b>Information about authors</b> .....	<b>514</b>

## 5.2 Health<sup>39</sup>

### Executive summary

Climate and environmental change cause a wide range of impacts on human health in Mediterranean countries. The vulnerability of people to the impacts of climate and environmental change is strongly influenced by population density, level of economic development, food availability, income level and distribution, local environmental conditions, pre-existing health status, and the quality and availability of public health care. Poorer countries, particularly in North Africa and the Levant, are at highest risk.

Heat waves have the potential to cause very high rates of premature deaths, especially in large cities and among older people. Heat-related morbidity and mortality have been reduced in the region over recent years thanks to Heat-Health Action Plans (HHAPs). Despite the rise in mean temperature, cold waves are not expected to disappear with increased future climate variability.

Recent climate and landscape changes in the Mediterranean Basin may create suitable environments for mosquitoes, ticks, and other climate-sensitive vectors, and may exacerbate vector-borne diseases.

Every year, around one million fatalities are attributed to outdoor and indoor air pollution in the European and Eastern Mediterranean regions. There are synergistic effects between ozone levels, particulate matter concentrations and climate variables, especially during heat wave days, with high temporal and spatial variability. An increase in mortality of 1.66-2.1% is observed for each 1°C temperature increase.

Climate change and extreme events have a negative impact on mental health for people who experience loss of homes, destruction of settlements and damage to community infrastructure.

Future changes in the vulnerability of the Mediterranean Basin to vector-borne diseases transmission vary geographically, modifying significantly the extent and transmission patterns of these diseases. For example, by 2050, West Nile virus high-risk areas are expected to expand further, and transmission seasons will extend significantly.

It is important for prevention plans to be implemented. Most adaptation measures offer “win-win solutions” from a public health perspective, including the reduction of air pollution or providing shade. Additionally, Mediterranean countries have the potential to enhance cross-border collaboration for adaptation to many health risks.

### 5.2.1 Introduction

#### 5.2.1.1 Effects of climate and environmental change on human health

Climate change is a complex phenomenon that threatens all aspects of human society, including increasing risks to human life and health (WHO 2018). Most climate-related health impacts are mediated by complex ecological, environmental and social processes, while impacts vary in magnitude, scale and timing as a function of local environmental conditions and the vulnerability of the human population (Shuman 2010; Smith et al. 2014; Crowley 2016). Climate change impacts human health directly, through exposure to extreme heat and cold events, droughts or storms, or indirectly by changes in air quality, water availability, food availability and quality, and other stressors. The main health effects are related to extreme weather events (including floods and extreme temperatures), changes in the distribution of climate-sensitive diseases (such as vector-, water- and food-borne diseases), and changes in environmental and social conditions (EU Climate Policy).

#### 5.2.1.2 Multi-factorial changes in health attributed to environmental change

The Mediterranean Basin has been undergoing a warming trend with longer and warmer summers, an increase in the frequency, duration and severity of heat waves, and a reduction in rainfall. With significant gradients in socio-economic levels among Mediterranean countries, particularly between the North and the South, together with population growth and migration (World Bank 2017) (Section 5.3.2.3), increased water demand (Section 3.1.2), decreased water availability (Section 3.1.1) and quality (Section 3.1.3.5) (Bucak et al. 2017), ecosys-

<sup>39</sup> Parts of this chapter have been published by Linares et al. (2020).



tem degradation (Section 4.3) [e.g., Coll et al. 2010] and increased risk of forest fires (Section 4.3.2.1) [e.g., Turco et al. 2014], the vulnerability of the Mediterranean population to human health risks is increasing significantly.

## 5.2.2 Vulnerability and health risk – main causes

Population vulnerability to the impacts of environmental and climate change is strongly influenced by population density, level of economic development, food availability, income level and distribution, local environmental conditions, pre-existing health status, and the quality and availability of public health care (Woodward et al. 2000). Although socio-economic and demographic factors may vary geographically, there are some commonalities across populations in terms of risk factors (UNEP 2018). Characteristics that differentiate populations with particular health risks from environmental change include age, gender, geographic location, socio-economic status, acclimation, occupation, health infrastructure and the (often urban) housing situation (Smith et al. 2014).

### 5.2.2.1 Aging

Older populations are at particular risk of adverse climate change impacts due to decreased mobility and changes in physiology, as well as limited access to resources. These conditions may limit adaptive capacity among older and more vulnerable people (EASAC 2019). More specifically, with heat-related impacts, such as heat waves, elderly population groups are at particular risk due to dysfunctional thermoregulatory mechanisms, chronic dehydration and medications. People with pre-existing medical conditions, especially cardiovascular or pulmonary illnesses (Mayrhuber et al. 2018) and those with chronic diseases like diabetes are more vulnerable (Yardley et al. 2013), as are those who are obese and have cognitive impairment (Bouchama et al. 2007; Linares et al. 2016).

### 5.2.2.2 Gender

In addition to differences of a collective nature (such as body size, physical condition and state of acclimatization to heat), there are social factors such as differences in social isolation, that tend to be greater among men than women, and may prove a risk factor e.g., during heat waves (Canoui-Poitrine et al. 2006). There are factors of a physiological nature, such as women's tendency to sweat less than men (Gagnon et al. 2013), a natural thermoregulation mechanism that might explain the greater im-

pact of heat on women. Also single-parent women (De'Donato et al. 2018) are cited as more vulnerable. Moreover, for pregnant women and babies in gestation, extreme heat is a risk factor for adverse birth outcomes, such as low birth weight and premature birth (Arroyo et al. 2016).

### 5.2.2.3 Geographic location

Most studies show that there is important variability in the effects of climate change on morbidity and mortality related to geographic location and the sensitivity of populations to extreme values such as extreme heat or cold, urbanization level, and distance to health system infrastructures (Allen and Sheridan 2018). For example, rural populations will be at a high risk of vector-borne diseases related to climate warming.

### 5.2.2.4 Socio-economic status

Population vulnerability to high temperatures will be affected not only by climate change but also by socio-economic factors (Semenza et al. 2008). In socially disadvantaged groups, the effects are particularly pronounced among the poor, socially isolated, substance abusers and homeless (Nicolay et al. 2016). Migrants, refugees and internally displaced people may have pre-existing and post-displacement vulnerabilities such as malnutrition, untreated chronic medical conditions from limited access to health care, and lack of shelters that provide adequate protection, predisposing them to decompensation caused by heat or other extreme events (McMichael et al. 2012).

### 5.2.2.5 Level of acclimation

Climate change will affect an increasingly aging population, a larger percentage of whom have chronic diseases, and are therefore more susceptible to the effects of increasingly extreme temperatures (changes in the population susceptibility framework). Population effects are quantified through the increased number of people over 65, which is the target population for heat impacts, as well as those in energy poverty or living in older building structures. On the other hand, there are factors that should result in a decrease in the impact of heat in the future. These include, for example, the existence of an active adaptation process within the population (both autonomously by individuals and families, and by the authorities and institutions), due to multiple factors from the so called "culture of heat" (Bobb et al. 2014), to the implementation of prevention plans (Schifano et al. 2012), improvements in health services (van

Loenhout et al. 2016), and improvements in socio-economic circumstances and infrastructure of homes, as well as an increase in the number of air conditioning units (Díaz et al. 2018b), among others.

### 5.2.2.6 Occupational health

Extreme heat and cold waves have been linked to an increased risk of occupational injuries. Studies report significant losses in work capacity and productivity linked to climate warming. Several mechanisms are thought to be behind the link between ambient temperatures and risk of injury in the workplace (Martínez-Solanas et al. 2018). Exposure to high temperatures can lead to physiological and psychological changes associated with heat strain, which in turn can decrease workers' performance and lead to impaired concentration, increased distractibility and fatigue (Zander et al. 2015). Sectors with a high percentage of outdoor workers, such as agriculture and construction or police and security, have the highest risk of seeing heat stroke or even heat stress develop (Martínez-Solanas et al. 2018). Additionally, despite the rise in mean temperatures, cold waves are not going to disappear. Therefore, factors related to working in cold environments, such as thermal discomfort, hypothermia, or reduced mobility while wearing protective clothing are associated with decreased dexterity and performance among workers and can also trigger occupational injuries (Mäkinen et al. 2009).

### 5.2.2.7 National public health infrastructures, warning systems

Better surveillance and improved warning systems are needed for vulnerable population groups. Increased urbanization increases the level of population exposure and can put pressure on water management and energy infrastructure, social care and health systems, so as to make them inefficient or unable to adopt necessary measures and prevention plans (Environmental Audit Committee 2018). Prevention plans and early warning systems began in the European region after the heat wave of 2003 and their efficiency is under evaluation. In France, the implementation of the prevention plan and alert system after the heat wave of 2003 is considered to have contributed to a reduction of around 4,400 fatalities during the heat wave of 2006, especially benefitting people over 75 years of age (Fouillet et al. 2008). In recent years, Heat-Health Action Plans (HHAPs) led to a decline in mortality during heat waves (Martinez et al. 2019).

Currently, early warning systems for heat and cold waves are active in Mediterranean countries, such as the Egyptian Meteorological Authority (EMA).

### 5.2.2.8 Influence of urban landscape

Urban Heat Islands (UHI) are considered to be one of the greatest twentieth century problems facing humanity, and they are the result of urbanization and industrialization (Rizwan et al. 2008). Temperature differences between cities and rural areas due to the UHI effect can reach up to 10°C in large cities. The effect of heat in urban areas increases with population density, extensive economic activities and city expansion (Burkart et al. 2016; Milojevic et al. 2016). Factors that amplify the UHI effect include household characteristics such as the age of buildings, residence in the highest floor of a building, the presence of a bedroom immediately beneath the roof (due to the concentration of heat that accumulates during the day), and lack of good thermal isolation (Vandentorren et al. 2006; López-Bueno et al. 2019).

## 5.2.3 Health impacts: recent and current situation

### 5.2.3.1 Heat-related impacts

Heat waves have very high mortality rates in Europe causing tens of thousands of premature deaths, especially during the 2003 heat wave. Mediterranean cities like Athens, Barcelona and Rome have all experienced strong impacts of extremely high temperatures (De' Donato et al. 2015). Despite the aging European population and continuously increasing temperatures, Mediterranean cities (and also other areas with commonly high temperatures) have recently shown a reduction in heat-related morbidity and mortality (Díaz et al. 2018a). This reduction is attributable to several factors, such as the existence of Heat-Health Action Plans (HHAPs) and the implementation of prevention plans (Morabito et al. 2012), improvements in health services, infrastructure and housing, changes in patterns of susceptibility of the population and increased awareness of the effects of severe heat (Ragettli et al. 2017), which in turn may be a result of improved communication and media coverage (De' Donato et al. 2015).

An important decrease in heat-related mortality has been observed among children and elderly people, although at a lower rate for the elderly (Schifano et al. 2012; Díaz et al. 2015). The reduction in heat-related mortality does not show consistent differences by age group (De' Donato et al.

2015) or gender (Allen and Sheridan 2018) and is spatially variable (Toloo et al. 2013; Linares et al. 2015b). Plans and alert systems have helped raise awareness among the population about the risk, but they have probably not been enough to provoke changes in population behavior so as to lead people to take measures to protect themselves (Wolf et al. 2010). Health promotion and behavior studies suggest that people who are most likely to adopt these measures are also those who have a high risk perception, but the opposite is true for the most vulnerable groups.


The cost of investing in protection measures against heat is one of the barriers that prevent vulnerable populations from taking action. The negative environmental impact of air conditioning must also be considered (Chapter 3.3). Educational programs are required to inform the population, especially the most vulnerable and their caregivers, about the risks of exposure to high temperatures. In the European part of the Mediterranean Basin, the increase in temperatures will affect both warmer and temperate countries, and it is therefore important that prevention plans are implemented in those countries that currently have no prevention plans in place, and that they are improved where

they already exist. Ultimately, the effectiveness of prevention plans depends on the capacity (of the health sector, the local community, etc.) to adapt to the changes, the plan's ability to incorporate climate change into research frameworks on adaptation and implementation, and the knowledge generated in the field (Hess and Ebi 2016).

There is insufficient research on public awareness regarding climate change and health in the Mediterranean. People who perceive climate change as a risk to public health are more supportive of mitigation policies and show more willingness to take individual measures to mitigate climate change (Debono et al. 2012).

### 5.2.3.2 Cold-related impacts

Winter mortality is associated with low temperatures, extremely low temperatures or cold waves (The Eurowinter Group 1997). This phenomenon has attracted less attention than the analysis of heat waves, though its impact on mortality is higher and up to an order of magnitude greater than those related to heat (Vardoulakis et al. 2014). By the end of the 21st century, southern European regions are expected to experience a clear decline in cold-relat-



**Figure 5.1 | Distribution of the tiger mosquito, *Aedes albopictus*, in 2019 - a known vector of chikungunya and dengue viruses [from ECDC 2019].**

ed mortality opposite to the increase in heat-related mortality (Gasparrini et al. 2017). These conclusions are based on the dual assumption that there will be no population acclimatization processes to such extreme temperatures and no changes in mortality rate. Other studies indicate populations may adapt to heat (Oudin Åström et al. 2018). Despite the rise in mean temperatures (Section 2.2.4.2), cold waves are not expected to disappear. Added to the fact that the impact of cold-related mortality is greater than that of heat-related mortality (Carmona et al. 2016), from a public health standpoint it is essential that the climate impact on mortality risk is analyzed by considering the impacts of both heat and cold waves together.

### 5.2.3.3 Vector-borne diseases

One of the main impacts of environmental and climate changes on human health is the influence of warmer climates and changing rainfall patterns on vector-borne disease (VBD) transmission which, together with anthropogenic changes in landscapes may create hospitable environments for mosquitoes, ticks, and other climate-sensitive vectors (Crowley 2016) (Fig. 5.1).

Long-term anthropogenic climate change interacts with natural variability, influencing the transmission of VBDs from shorter (seasonal, annual) to longer (decadal) time scales, with variable effects and complex interactions at different times and locations (Campbell-Lendrum et al. 2015). These impacts are complex and may involve non-linear feedback inherent in the dynamics of many infections (Metcalf et al. 2017), including impacts of other environmental drivers such as biodiversity loss or changes in land use (Reisen et al. 2006; Marcantonio et al. 2015; Paz 2015) (Chapter 2). It is therefore expected that VBD outbreaks will be exacerbated in the region.

Most cities in the Mediterranean Basin are compact and densely populated. Air conditioning is used in regions with sufficient resources, but windows often remain open even during the hottest months. Many activities, particularly social gatherings, occur in outdoor locations such as shaded balconies, courtyards, and outdoor restaurants - all ideal places for contact with the vector. While warmer summers extend the potential season of the disease throughout the basin, poorer countries, particularly in North Africa and the Levant, are at highest risk (Negev et al. 2015).

Currently, the main vector-borne diseases in the Mediterranean basin, transmitted by insects and

potentially influenced by the changing climate are West Nile virus, Chikungunya and Leishmaniasis.

#### West Nile virus

West Nile virus (WNV) is a vector-borne pathogen of global importance since it is the most widely distributed virus of the encephalitic *Flavivirus* spp. Mosquito species from the genus *Culex* (family *Culicidae*) are the primary amplification vectors and also act as bridge vectors. The enzootic cycle is driven by continuous virus transmission to susceptible bird species through adult mosquito blood meal feeding, which results in virus amplification (Paz and Semenza 2013; Petersen et al. 2013).

The establishment of WNV in new regions is facilitated by warmer conditions. Ambient warming increases the growth rates of mosquito populations, decreases the interval between blood meals and shortens the incubation time in mosquitoes (Paz 2015; Moirano et al. 2018). Clear associations have been found between warm conditions and WNV outbreaks in Mediterranean countries (Paz et al. 2013; Tran et al. 2014; Marcantonio et al. 2015; Moirano et al. 2018).

Since the unprecedented WNV outbreak in 2010 in southern and eastern Europe, which was accelerated by extreme temperatures (Paz et al. 2013), outbreaks occur every summer (2011-2019) and there is evidence of ongoing transmission in Euro-Mediterranean countries (Semenza and Suk 2018). During the last decade, WNV outbreaks in humans erupted in many Mediterranean countries including France, Italy, Croatia, Slovenia, Greece, Turkey, Israel and the Mediterranean islands. During the 2018 transmission season, a higher number of cases was reported compared with previous years (ECDC 2018).

The impact of changes in rainfall patterns on disease incidence is influenced by precipitation levels (increased and extreme precipitation, e.g., Moirano et al. 2018, floods or droughts), depending on local conditions, the landscape (e.g. wetlands, Tran et al. 2014) and the differences in the ecology and sensitivity of mosquito species (Paz 2015). In the Mediterranean area, increased rainfall and humidity together with high temperatures probably favored the multiplication of *Culex* species, which led to numerous cases of WNV human infections in northern Greece in summer 2010 (Papa et al. 2010). Climate parameters were found as key predictors of WNV outbreaks including high precipitation in late winter/

early spring and summer drought (Marcantonio et al. 2015).

### **Chikungunya**

Chikungunya is a viral disease transmitted by *Aedes* mosquitoes to humans. The most common symptoms are fever and severe joint pain. In 2007, an outbreak of chikungunya virus infections took place for the first time in Italy, indicating the possibility of mosquito-borne outbreaks by *Aedes albopictus* in the Euro-Mediterranean area. In 2010 and 2014, autochthonous cases were reported in France. The risk of chikungunya spreading in the EU and the Mediterranean is high due to importation through infected travelers, presence of competent vectors in many countries (particularly around the Mediterranean coast) and population susceptibility (ECDC). In August-September 2017, local transmission of chikungunya was confirmed in southeastern France (WHO 2017a) and in Italy, six transmissions were reported in Rome and eight in the coastal area of Anzio in the Lazio Region (WHO 2017b).

### **Rift Valley fever**

Rift Valley fever is a mosquito-borne zoonotic climate-sensitive disease closely associated with high-rainfall conditions (e.g., after prolonged excessive rainfall in sub-Saharan Africa). However, large outbreaks have also occurred in the dry and low-rainfall climate of Egypt (Linthicum et al. 2016).

### **Leishmaniasis**

Leishmaniasis is a vector-borne disease with three main clinical forms: Visceral (Kala-azar), Cutaneous Leishmaniasis (CL), and Visceral Leishmaniasis (VL) caused by infection of *Leishmania* parasites and transmitted by the bite of infected females of *Phlebotomine* spp. sandflies. *Leishmania* genus includes about 20 species, widely distributed in more than 85 endemic countries, with 0.7-1.2 million new cases of CL every year, of which about a third occur in the Mediterranean region (Alvar et al. 2012). In the eastern Mediterranean Basin, two CL species, which manifest as skin sores, are common: *Leishmania major* and *Leishmania tropica*. *Leishmania tropica*, transmitted by the *Phlebotomus sergenti* sandfly, was first discovered in Israel in the early 1990s. Since the late 1990s, rapid unexpected outbreaks occurred in new urban and rural foci in Israel, Jordan and the Palestinian Authority (Al-Jawabreh et al. 2017; Waitz et al. 2018).

### **5.2.3.4 Food- and water-borne diseases**

Climate change increases the risks of food- and water-borne diseases (Ebi et al. 2018). For example, the survival and multiplication of salmonellosis in the environment and in food is influenced by high temperatures (Miraglia et al. 2009; Milazzo et al. 2016). It was shown for ten European countries that temperature influences infection transmission in about 35% of all cases of salmonellosis, while the greatest effect was apparent for temperature one week before the onset of illness (Kovats et al. 2004).

*Campylobacter* species have emerged as leading bacterial causes of gastroenteritis and food-borne infections in high-income countries (EFSA and ECDC 2015). The incidence of campylobacteriosis varies seasonally and geographically, and tends to be highest during the summer months (Bassal et al. 2016). While the temperature may directly affect the rate of replication of pathogens and their survival in the environment, increased ambient temperatures may increase bacterial contamination at various points along the food chain and also influence people's behavior which, in turn, may be translated into more risky patterns of food consumption (Lake et al. 2009). A recent retrospective study in Israel found that higher temperatures across seasons, prior to or around the time of food purchasing, played a role in human infection (Rosenberg et al. 2018).

Leptospirosis, caused by *Leptospira interrogans*, is a highly infectious emerging water-borne zoonosis of global significance. A study in Croatia showed strong influence of climate conditions on incidence of human leptospirosis at annual level. In the years 2010 and 2014 that were characterized as warm/extremely warm and wet/extremely wet, a significant temporal increase in incidence was observed. Increased risk for human infections is related to season, gender and age with peaks in incidence occurring cyclically and associated with extreme weather conditions. The influence of weather should not be considered without taking into account the wider impact of climate change on domestic, peridomestic and wild animals (Habus et al. 2017). In Israel, human leptospirosis is uncommon, but in summer 2018, a large outbreak of human leptospirosis was linked to contaminated water bodies in northern Israel after years of severe drought conditions which had resulted in particularly low water levels in the region (Dadon et al. 2018).

## 5.2.4 Indirect impacts: recent and current situation

### 5.2.4.1 Air quality

Climate and environmental change, anthropogenic activities, urbanization, industrialization, etc. affect air quality and impact human health, through several pathways, such as changes in atmospheric circulation, ventilation and dilution of air pollutants, removal processes, stratosphere–troposphere O<sub>3</sub> (ozone) exchange (e.g., Akritidis et al. 2016) and increase in the frequency of wildfires (Fiore et al. 2015) (see *Section 2.3.2*). Every year, around one million fatalities are attributed to outdoor and indoor air pollution in the European and eastern Mediterranean regions (WHO Regional Office for Europe and OECD 2015). In Lebanon, the prevalence of cardiovascular disease has been linked to exposure to pollution (Salameh et al. 2019). In Europe the cardiovascular diseases burden from ambient air pollution is substantially higher than previously assumed, though subject to some uncertainty (Lelieveld et al. 2019).

In Europe, 90% of citizens are exposed to levels of fine particulate matter (PM) that exceed World Health Organization (WHO) air quality guidelines. There are synergistic effects between ozone levels, PM concentrations and climate variables (Analitis et al. 2018), especially on heat wave days (Katsouyanni et al. 2009), together with a large variability on both temporal and geographical scales, likely connected to local climate characteristics, activity patterns and physical adaptation (de Sario et al. 2013). An increase in mortality of 1.66% was observed for each 1°C temperature increase on low ozone level days, and an increase of up to 2.1% on high ozone level days (Analitis et al. 2018). There is a positive relationship between cardiovascular mortality and the concentrations of nitrogen dioxide (NO<sub>2</sub>), the main precursor of tropospheric ozone (Nuvolone et al. 2013). Out of 524,000 pollution-related premature deaths, 432,000 are estimated to be attributable to PM<sub>2.5</sub>, 17,000 to O<sub>3</sub> and 75,000 to NO<sub>2</sub>. Reducing exposure to PM improves the life expectancy of Europeans by about 8 months (WHO Regional Office for Europe and OECD 2015). As the main emission source of these pollutants is vehicle traffic, largest impacts are concentrated in large cities due to microcirculation.

Atmospheric pollutants that are linked to climate change are considered major contributors to the large rise in the number of people affected during the allergy season. Airborne allergens chemical-

ly modified by the presence of NO<sub>2</sub> and O<sub>3</sub>, seem to increase their potency. Airborne allergies are thus becoming more common in combination with global climate change. Together with global warming, increased pollen production and earlier spring phenology, this leads to earlier, more frequent and more widespread pollen allergies (American Chemical Society 2015).

### 5.2.4.2 Mineral dust and forest fires

The main health impacts associated with PM occur in densely populated urban areas where the principal component is from anthropogenic emissions (Karanasiou et al. 2012). In southern European urban areas, these account for approx. 80% of PM and aerosol emissions, while the remaining 20% are of natural origin, mainly from advections of desert dust (Viana et al. 2014), sea spray (O'Dowd and de Leeuw 2007), volcanic emissions (von Glasow et al. 2009), and aerosols from wildfires, with Saharan dust intrusions and PM advection from wildfires being the primary sources.

Impacts of PM due to wildfires on human health are mainly respiratory problems (Mirabelli et al. 2009) or exacerbations of previous respiratory diseases (Martin et al. 2013), while exposure to forest fire smoke has also been linked to cardiac diseases (Weichenthal et al. 2017). Medium-size fires are found to increase daily mortality in Athens (Analitis et al. 2012), while the accumulative impact of PM<sub>10</sub> (PM with diameter less than 10 µm) during forest fires smoke advection exacerbates mortality for different age groups and causes (Faustini et al. 2015; Linares et al. 2015a).

Long-distance transport generates a change in the respective atmospheric concentration of the different sized particles and in the chemical composition of the particles present in the air (Pérez et al. 2012), while there is evidence that desert dust itself transports biological allergens or irritants (Garrison et al. 2006; Griffin 2007; Polymenakou et al. 2008). Non-biological compounds in dust may cause adverse health effects, or local conditions may modify the toxicological properties of the dust. The two circumstances of change, i.e., in PM concentration and chemical composition, are related to clearly differentiated morbidity-mortality patterns, which are observable on days with desert dust intrusions (Jiménez et al. 2010).

The human health effects of dust storms range from respiratory disorders (including asthma, tracheitis, pneumonia, allergic rhinitis and silicosis),

to cardiovascular disorders (including stroke), conjunctivitis, skin irritations, valley fever, diseases associated with toxic algal blooms, and mortality and injuries related to transport accidents (Goudie 2014). Spatial and temporal variability of the PM effects on human health due to Sahara dust intrusions in Euro-Mediterranean countries stems from co-existing PM concentrations due to traffic sources and their higher toxicity (Samoli et al. 2011; Stafoggia et al. 2016), the varying impacts on the different age groups (Zauli-Sajani et al. 2011), the specific causes of health impacts and PM groups (Neophytou et al. 2013).

### 5.2.4.3 Mental health

Climate change and extreme events have a negative impact on mental health in several ways. Floods, droughts and sea level rise have long-lasting impacts on societies who have experienced loss of homes, destruction of settlements and damage to community infrastructure. These impacts on mental health include anxiety, depression and post-traumatic stress disorders (Watts et al. 2015). However, there is insufficient research on the mental health impacts of climate change, both internationally (Watts et al. 2018) and also regionally in Mediterranean countries. The few studies that have been conducted in the Mediterranean found negative impacts of high temperature on mental health. For example, in Thessaloniki, Greece, high temperatures may be associated with increased male suicide rates. In the context of the recent economic crisis in Greece, a multilinear regression showed that high temperatures explain 51% of the variance in male suicides, while unemployment was insignificant (Fountoulakis et al. 2016). Another study in Northern Italy found a strong positive association between the number of daily emergency psychiatric visits and mean daily air temperature (Cervellin et al. 2014). More research is needed regarding the impact of different extreme climate events on diverse mental health outcomes.

There is some evidence that climate change may intensify violence in the Mediterranean, across all levels. Climate change-induced water shortage and food insecurity may intensify conflicts in the eastern Mediterranean (Brown and Crawford 2009), especially in counties that lack adaptive capacity (Feitelson and Tubi 2017). At the domestic level, a study in Madrid found an association between heat waves and increase in domestic violence, including an increase in police reports and helpline calls three days after the heat wave (Sanz-Barbero et al. 2018).

### 5.2.4.4 Migration

Displacement related to environmental change and climate disasters is not a new phenomenon in the Mediterranean (*Chapter 5.3*) (Charef and Dorai 2016). Migration potential has increased over recent decades due to the increasing impacts of climate change, frequent and more intense extreme events, especially in areas with high population density and areas at risk, with direct and indirect impacts on the well-being, livelihood and security of populations (*Chapter 5.3*). Displacement may lead to adverse health outcomes, especially for vulnerable population groups as well as those suffering from chronic diseases (Schütte et al. 2018). In lower-income countries hosting refugees in particular, it may undermine national health systems and diminish access to health care for domestic, as well as migrant, populations (Gostin and Roberts 2015). In Egypt, re-emerging diseases such as tuberculosis have been reported as an indirect effect of climate change linked to the crowdedness of slum areas due to irregular internal migration of farmers who have lost their land (Girardi et al. 2017).


## 5.2.5 Projections for global warming of 1.5°C, 2°C and more

### 5.2.5.1 Vulnerabilities and risks under different warming scenarios

The IPCC special report on 1.5°C global warming (IPCC 2018) shows that global warming of 2°C poses substantially greater risks to human health than 1.5°C, with actual risk levels varying regionally (Hoegh-Guldberg et al. 2018). The risks may be particularly high for heat-related morbidity and mortality, heat stress, ground-level O<sub>3</sub>, and malnutrition. For vector-borne diseases, the risks are more variable because warmer temperatures may result in some regions becoming too hot and/or too dry for a vector (Ebi et al. 2018).

### 5.2.5.2 Heat-related impacts

Many projections around heat-related mortality are made without considering the socio-economic conditions of the population. In order to show the contribution of changes in socio-economic and climate conditions to mortality due to heat in the European population, a study was developed that combined socio-economic scenarios with greenhouse gas emissions (RCP) (Mayrhuber et al. 2018). The percentage of the European population at risk of thermic stress is expected to increase constantly over coming years, and could increase from the



**Figure 5.2 | Attributable fraction of heat-related deaths during summer by country in European sub-region, A) RCP4.5 in 2050; B) RCP8.5 in 2050, C) RCP4.5 in 2085 and d) RCP8.5 in 2085 (from Kendrovski et al. 2017).**

current 0.4% to 20.3%, 32.6% or even 48.4% in 2050 depending on the scenario combination – unless substantial political changes occur rapidly and steadily shift the current socio-economic development pathway towards sustainability (Rohat et al. 2019). However, the impact of heat on mortality will be influenced more by socio-economic factors that enhance vulnerability than by exposure to high temperatures. Effects of heat-related mortality in Europe will vary considerably, with the Mediterranean region being the most affected (Mayrhuber et al. 2018).

Heat wave changes under the RCP4.5 and RCP8.5 scenarios will mainly affect the countries of Mediterranean Europe and eastern Europe (Kendrovski et al. 2017). More specifically, for the middle of the 21st century, 2035-2064, annual attributable mortality will increase by a factor 1.8 and 2.6 for RCP4.5 and RCP8.5, respectively, compared to the period 1971-2000. Heat wave attributable mortality at the end of the century will increase by around 3

and 7 times under RCP4.5 and RCP8.5 (Kendrovski et al. 2017). *Fig. 5.2* presents the mean warm season attributable fraction (the fraction of deaths attributable to mean apparent temperature above the threshold) based on the SMHI RCA4/HadGEM2 ES r1 (MOHC) climate model.

By the end of the 21st century (2090–2099), southern European countries are expected to experience a temperature increase of 1.9°C (1.3-2.2°C) and 4.5°C (3.0-5.1°C) respectively for RCP4.5 and RCP8.5, compared to the mean of 2010–2019 in a GCM-ensemble assuming no population changes (Gasparrini et al. 2017) (*Section 2.2*). The greatest changes in heat-related excess mortality are projected for southern Europe, with a 10.5% increase (IC95%: 5.6-17.3). For mortality related to heat waves, although the largest temperature increases are expected in southern Europe (for the period 2031–2080, changes in the 95th percentile: Italy 1.7°C and 3.2°C and Spain 1.4°C and 3.0°C, under RCP4.5 and RCP8.5, respectively),



the area will not experience the greatest excess mortality linked to heat waves, as a result of population adaptation and prevention plans (Guo et al. 2018).

### 5.2.5.3 Cold-related impacts

As global warming progresses, a decreasing trend in cold extremes is expected. However, highly variable future climates may retain cold wave hazards as a locally important threat. Cold-related mortality is also expected to increase with expected demographic changes in European cities (Smid et al. 2019), but the effect of cold waves on aging populations is not as pronounced as the heat, since its relationship with respiratory diseases affects younger age groups. At European level, cold-related mortality is projected to decrease by the 2080s as much as heat-related mortality is expected to increase for the same period (EEA 2017). Better social, economic and housing conditions in many European countries may encourage the estimated decreasing risk, despite the expected higher variability.

### 5.2.5.4 Vector- food- and water-borne diseases

The rise in temperature may lead to a geographic expansion of Euro-Mediterranean areas that are currently climatically suitable for WNV, and also to an extension of the transmission season, with the extent and pattern of changes varying depending on the location and degree of warming (Semenza et al. 2016). The transmission risk for WNV in Euro-Mediterranean areas varies spatially as well as temporally (Conte et al. 2015). Early summer months will provide suitable climatic conditions in Tunisia, Libya and Egypt, while in the European continent suitable conditions prevail only from July. From August to October, significantly increased transmission risk will characterize Italy, France, Spain, the Balkan countries, Morocco, northern Tunisia, and all along the Mediterranean coast of Africa and the Middle East. In November, with the exception of limited European coastal areas of the Mediterranean, the risk will be very low, while in North African and Middle Eastern coastal areas the same is valid from December. Projections for Europe indicate a continuous extension of regions with an increased risk of WNV infections, mainly on the fringes of the regions of transmission (Conte et al. 2015). Predictions for 2025 show an elevated risk in northeastern Greece, eastern Croatia and northwestern Turkey, while in 2050 high-risk areas expand further (Semenza et al. 2016; Semenza and Suk 2018).

Southernmost parts of Europe do not generally provide climatically suitable areas for Chikungunya transmission in the 21st century, except for restricted areas in France and North Italy at the end of the century (Fischer et al. 2013). A significant reduction of habitat suitability for *Aedes albopictus* is projected for the middle of the 21st century in southern Europe and the Mediterranean, related to significant increase of summer temperatures (Caminade et al. 2012; Proestos et al. 2015). Similar results are found for the end of the century (Tjaden et al. 2017) (Section 2.2.4.2).

Since populations are exposed to variability in weather patterns and increasingly warm temperatures, there is high confidence for increased risks of food- and water-borne diseases, such as diarrheal diseases and *Salmonella* spp. (Smith et al. 2014). With rising average temperatures and an increase in the frequency and length of heat waves, a rising number of cases of food-borne illness are expected in a business-as-usual scenario unless education, epidemiological surveillance and enforcement (related to food safety) are intensified. This will be compounded in the event of power outage due to peak energy demand (e.g., during heat waves) that may lead to malfunction of food preservation practices (refrigeration) (The Malta Resources Authority 2017).

### 5.2.5.5 Air quality

Climate change alters the dispersion of primary pollutants, particularly particulate matter, and intensifies the formation of secondary pollutants. According to global estimates, the number of days with ozone concentrations exceeding the thresholds for protection of human health are expected to increase. In polluted areas with high levels of nitrogen oxides (NO<sub>x</sub>), high surface temperature and humidity (Section 2.2.4.2) may generate an increase of surface O<sub>3</sub> concentrations, especially in southern Europe (Doherty et al. 2017). By the middle of the 21st century in southern Europe, climate change is expected to lead to an increased summer mean O<sub>3</sub> (0–3 ppb) and increased summer daily maximum O<sub>3</sub> (3–6 ppb) (Langner et al. 2012; Doherty et al. 2013, 2017; Colette et al. 2015). For RCP8.5, the Mediterranean Basin may experience an annual average difference in stratospheric origin ozone concentration at sea level of above 5 ppb by the end of the 21st century (Meul et al. 2018).

Regional projections indicate an increase of 10 to 14% in ozone-related morbidity and mortality from 2021 to 2050 in several countries, including France, Spain and Portugal. For 2050 a 8 to 11% increase in

non-accidental mortality is expected, and for 2080 a 15 to 16% increase, compared to the year 2000 (O<sub>3</sub> and PM<sub>2.5</sub> combined) in Europe (Orru et al. 2017). Changes in PMs under climate change still require further study, and important uncertainties remain with regard to the impact of temperature change on PM components, together with still uncertain precipitation patterns (Doherty et al. 2017). The relationships between climate change, air pollution and air pollution-related health impacts depend highly on the climate change scenario used, and on projections of future air pollution emissions, with relatively high uncertainty. Further studies focusing on effects on morbidity are needed (Orru et al. 2017).

## 5.2.6 Resilience, preparedness and adaptation

### 5.2.6.1 Health preparedness and adaptation measures

Health preparedness and adaptation to climate change includes adaptation of health systems to access morbidity due to extreme events, and adaptation of the built environment in order to reduce the burden of extreme climate. The quality of health systems and accessibility to healthcare is different across countries in the Mediterranean, largely along the North/South division. Health systems in the Mediterranean face climate change in the context of an increasingly elderly population, which is particularly vulnerable to heat waves, an increase in vector-borne diseases, and an increase in climate migration to Mediterranean countries. For health systems in the southern Mediterranean, another challenge is the declining resources for this sector (Sanderson et al. 2018). While investment in climate-related adaptation health systems appears to be cost-effective (Jeuken et al. 2016), health adaptation in the region is lacking, with only one out of 22 countries in the eastern Mediterranean having a Heat-Health Action Plan (UNEP 2018). Increasing preparedness of health systems in the Mediterranean Basin may be supported by the following measures:

- Implementation of heat early warning systems,
- Preparedness of emergency medicine professionals for treating morbidity related to extreme climate events, such as heat waves, cold spells and floods,
- Monitoring climate-related morbidity and mortality and designing interventions,
- Monitoring and surveillance of vector-borne diseases, including across borders with neighboring countries,

- Prevention of water-borne and food-borne diseases,
- Provision of access to healthcare, including mental health, to climate migrants,
- Training health professionals, including physicians, nurses and administrative staff regarding the health impacts of climate change, preparedness and adaptation in the health system,
- Increase public awareness of climate change-related health risks, and recommended prevention and mitigation of negative health outcomes, including behavior during heat waves, elimination of habitats for vectors, etc.,
- Expand urban green infrastructure including protecting inside areas and settlements.

### 5.2.6.2 Regional coordination and collaboration

The Mediterranean Basin, particularly its eastern and southern regions, is an area troubled by internal and cross-border conflicts, limited cross-border collaborations and limited links to the international frameworks for the whole Mediterranean. There are EU-funded regional frameworks such as Climate ADAPT Mediterranean area, that covers the southern part of Portugal, Mediterranean areas of Spain and France, almost all of Italy and the whole extension of Slovenia, Croatia, Greece, Malta and Cyprus, Albania, Bosnia-Herzegovina and Montenegro (<https://climate-adapt.eea.europa.eu/>). ClimaSouth covers Algeria, Egypt, Israel, Jordan, Lebanon, Libya, Morocco, Palestine and Tunisia (<http://www.climasouth.eu/en>). While both frameworks emphasize on increasing resilience, human health is not at the center of these programs. Other frameworks focus on infectious diseases, e.g., the Middle East Consortium on Infectious Disease Surveillance (MECIDS) that coordinates between Israel, Jordan and the Palestinian Authority (<http://www.mecidsnetwork.org/>). Climate change is expected to affect the Mediterranean, with heat and drought impacts on morbidity and migration, and vector-borne diseases spreading across borders. It is a challenge for countries that lack diplomatic relations to collaborate, but health agencies prove that it is possible (e.g., in the case of MECIDS), and regional collaboration at the Mediterranean level should be a priority for health agencies in the region.

## References

- Akritidis D, Pozzer A, Zanis P, Tyrllis E, Škerlak B et al. 2016 On the role of tropopause folds in summertime tropospheric ozone over the eastern Mediterranean and the Middle East. *Atmos. Chem. Phys.* 16, 14025–14039. doi: [10.5194/acp-16-14025-2016](https://doi.org/10.5194/acp-16-14025-2016)
- Al-Jawabreh A, Dumaidi K, Erekat S, Al-Jawabreh H, Nasereddin A et al. 2017 Molecular epidemiology of human cutaneous leishmaniasis in Jericho and its vicinity in Palestine from 1994 to 2015. *Infect. Genet. Evol.* 50, 95–101. doi: [10.1016/j.meegid.2016.06.007](https://doi.org/10.1016/j.meegid.2016.06.007)
- Allen MJ, Sheridan SC 2018 Mortality risks during extreme temperature events (ETEs) using a distributed lag non-linear model. *Int. J. Biometeorol.* 62, 57–67. doi: [10.1007/s00484-015-1117-4](https://doi.org/10.1007/s00484-015-1117-4)
- Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P et al. 2012 Leishmaniasis worldwide and global estimates of its incidence. *PLoS One* 7, e35671. doi: [10.1371/journal.pone.0035671](https://doi.org/10.1371/journal.pone.0035671)
- American Chemical Society 2015 Air pollutants could boost potency of common airborne allergens. <https://www.acs.org/content/acs/en/pressroom/newsreleases/2015/march/air-pollutants-could-boost-potency-of-common-airborne-allergens.html>
- Analitis A, De' Donato FK, Scortichini M, Lanki T, Basagaña X et al. 2018 Synergistic effects of ambient temperature and air pollution on health in Europe: results from the PHASE project. *Int. J. Environ. Res. Public Health* 15, 1856. doi: [10.3390/ijerph15091856](https://doi.org/10.3390/ijerph15091856)
- Analitis A, Georgiadis I, Katsouyanni K 2012 Forest fires are associated with elevated mortality in a dense urban setting. *Occup. Environ. Med.* 69, 158–162. doi: [10.1136/oem.2010.064238](https://doi.org/10.1136/oem.2010.064238)
- Arroyo V, Díaz J, Carmona R, Ortiz C, Linares C 2016 Impact of air pollution and temperature on adverse birth outcomes: Madrid, 2001–2009. *Environ. Pollut.* 218, 1154–1161. doi: [10.1016/j.envpol.2016.08.069](https://doi.org/10.1016/j.envpol.2016.08.069)
- Bassal R, Lerner L, Valinsky L, Agmon V, Peled N et al. 2016 Trends in the epidemiology of campylobacteriosis in Israel (1999–2012). *Foodborne Pathog. Dis.* 13, 448–455. doi: [10.1089/fpd.2015.2096](https://doi.org/10.1089/fpd.2015.2096)
- Bobb JF, Peng RD, Bell ML, Dominici F 2014 Heat-related mortality and adaptation to heat in the United States. *Environ. Health Perspect.* 122, 811–816. doi: [10.1289/ehp.1307392](https://doi.org/10.1289/ehp.1307392)
- Bouchama A, Dehbi M, Mohamed G, Matthies F, Shoukri M et al. 2007 Prognostic Factors in Heat Wave-Related Deaths: a meta-analysis. *Arch. Intern. Med.* 167, 2170–2176. doi: [10.1001/archinte.167.20.ira70009](https://doi.org/10.1001/archinte.167.20.ira70009)
- Brown O, Crawford A 2009 Rising Temperatures, Rising Tensions: Climate change and the risk of violent conflict in the Middle East. International Institute for Sustainable Development (IISD).
- Bucak T, Trolle D, Andersen HE, Thodsen H, Erdoğan Ş et al. 2017 Future water availability in the largest freshwater Mediterranean lake is at great risk as evidenced from simulations with the SWAT model. *Sci. Total Environ.* 581–582, 413–425. doi: [10.1016/J.SCITOTENV.2016.12.149](https://doi.org/10.1016/J.SCITOTENV.2016.12.149)
- Burkart K, Meier F, Schneider A, Breitner S, Canário P et al. 2016 Modification of Heat-Related Mortality in an Elderly Urban Population by Vegetation (Urban Green) and Proximity to Water (Urban Blue): Evidence from Lisbon, Portugal. *Environ. Health Perspect.* 124, 927–934. doi: [10.1289/ehp.1409529](https://doi.org/10.1289/ehp.1409529)
- Caminade C, Medlock JM, Ducheyne E, McIntyre KM, Leach S et al. 2012 Suitability of European climate for the Asian tiger mosquito *Aedes albopictus*: recent trends and future scenarios. *J. R. Soc. Interface* 9, 2708–2717. doi: [10.1098/rsif.2012.0138](https://doi.org/10.1098/rsif.2012.0138)
- Campbell-Lendrum D, Manga L, Bagayoko M, Sommerfeld J 2015 Climate change and vector-borne diseases: what are the implications for public health research and policy? *Philos. Trans. R. Soc. B Biol. Sci.* 370. doi: [10.1098/rstb.2013.0552](https://doi.org/10.1098/rstb.2013.0552)
- Canoui-Poitrine F, Cadot E, Spira A, Spira A 2006 Excess deaths during the August 2003 heat wave in Paris, France. *Rev. Epidemiol. Sante Publique* 54, 127–135. doi: [10.1016/S0398-7620\(06\)76706-2](https://doi.org/10.1016/S0398-7620(06)76706-2)
- Carmona R, Díaz J, Mirón IJ, Ortiz C, León I et al. 2016 Geographical variation in relative risks associated with cold waves in Spain: The need for a cold wave prevention plan. *Environ. Int.* 88, 103–111. doi: [10.1016/J.ENVINT.2015.12.027](https://doi.org/10.1016/J.ENVINT.2015.12.027)
- Cervellin G, Comelli I, Lippi G, Comelli D, Rastelli G et al. 2014 The number of emergency department visits for psychiatric emergencies is strongly associated with mean temperature and humidity variations. Results of a nine year survey. *Emerg. Care J.* 10. doi: [10.4081/ECJ.2014.2271](https://doi.org/10.4081/ECJ.2014.2271)
- Charef M, Dorai K 2016 Human migration and climate change in the Mediterranean region, in *The Mediterranean region under climate change. A scientific update*, eds. Thiébaud S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 439–444.
- Colette A, Andersson C, Baklanov A, Bessagnet B, Brandt J et al. 2015 Is the ozone climate penalty robust in Europe? *Environ. Res. Lett.* 10, 084015. doi: [10.1088/1748-9326/10/8/084015](https://doi.org/10.1088/1748-9326/10/8/084015)
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Ben Rais Lasram F et al. 2010 The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS One* 5, e11842. doi: [10.1371/journal.pone.0011842](https://doi.org/10.1371/journal.pone.0011842)
- Conte A, Candeloro L, Ippoliti C, Monaco F, de Massis F et al. 2015 Spatio-Temporal Identification of Areas Suitable for West Nile Disease in the Mediterranean Basin and Central Europe. *PLoS One* 10,

- e0146024. doi: [10.1371/journal.pone.0146024](https://doi.org/10.1371/journal.pone.0146024)
- Crowley RA 2016 Climate change and health: a position paper of the American College of Physicians. *Ann. Intern. Med.* 164, 608–610. doi: [10.7326/M15-2766](https://doi.org/10.7326/M15-2766)
- Dadon Y, Haas EJ, Kaliner E, Anis E, Singer SR et al. 2018 Outbreak of human leptospirosis linked to contaminated water bodies in Northern Israel, June to August 2018. *Euro Surveill.* 23, 1800486. doi: [10.2807/1560-7917.ES.2018.23.38.1800486](https://doi.org/10.2807/1560-7917.ES.2018.23.38.1800486)
- De' Donato FK, Leone M, Scortichini M, de Sario M, Katsouyanni K et al. 2015 Changes in the effect of heat on mortality in the last 20 years in nine European cities. Results from the PHASE project. *Int. J. Environ. Res. Public Health* 12, 15567–15583. doi: [10.3390/ijerph121215006](https://doi.org/10.3390/ijerph121215006)
- De' Donato FK, Scortichini M, de Sario M, de Martino A, Michelozzi P 2018 Temporal variation in the effect of heat and the role of the Italian heat prevention plan. *Public Health* 161, 154–162. doi: [10.1016/j.puhe.2018.03.030](https://doi.org/10.1016/j.puhe.2018.03.030)
- de Sario M, Katsouyanni K, Michelozzi P 2013 Climate change, extreme weather events, air pollution and respiratory health in Europe. *Eur. Respir. J.* 42, 826–843. doi: [10.1183/09031936.00074712](https://doi.org/10.1183/09031936.00074712)
- Debono R, Vincenti K, Calleja N 2012 Risk communication: climate change as a human-health threat, a survey of public perceptions in Malta. *Eur. J. Public Health* 22, 144–149. doi: [10.1093/eurpub/ckj181](https://doi.org/10.1093/eurpub/ckj181)
- Díaz J, Carmona R, Mirón IJ, Luna MY, Linares C 2018a Time trend in the impact of heat waves on daily mortality in Spain for a period of over thirty years (1983–2013). *Environ. Int.* 116, 10–17. doi: [10.1016/j.envint.2018.04.001](https://doi.org/10.1016/j.envint.2018.04.001)
- Díaz J, Carmona R, Mirón IJ, Ortiz C, Linares C 2015 Comparison of the effects of extreme temperatures on daily mortality in Madrid (Spain), by age group: The need for a cold wave prevention plan. *Environ. Res.* 143, 186–191. doi: [10.1016/j.envres.2015.10.018](https://doi.org/10.1016/j.envres.2015.10.018)
- Díaz J, López IA, Carmona R, Mirón IJ, Luna MYY et al. 2018b Short-term effect of heat waves on hospital admissions in Madrid: Analysis by gender and comparison with previous findings. *Environ. Pollut.* 243, 1648–1656. doi: [10.1016/j.envpol.2018.09.098](https://doi.org/10.1016/j.envpol.2018.09.098)
- Doherty RM, Heal MR, O'Connor FM 2017 Climate change impacts on human health over Europe through its effect on air quality. *Environ. Heal.* 16, 118. doi: [10.1186/s12940-017-0325-2](https://doi.org/10.1186/s12940-017-0325-2)
- Doherty RM, Wild O, Shindell DT, Zeng G, MacKenzie IA et al. 2013 Impacts of climate change on surface ozone and intercontinental ozone pollution: A multi-model study. *JGR Atmos.* 118, 3744–3763. doi: [10.1002/jgrd.50266](https://doi.org/10.1002/jgrd.50266)
- EASAC 2019 The imperative of climate action to protect human health in Europe.
- Ebi KL, Hasegawa T, Hayes K, Monaghan A, Paz S et al. 2018 Health risks of warming of 1.5 °C, 2 °C, and higher, above pre-industrial temperatures. *Environ. Res. Lett.* 13, 063007. doi: [10.1088/1748-9326/aac4bd](https://doi.org/10.1088/1748-9326/aac4bd)
- ECDC Factsheet about chikungunya. <https://ecdc.europa.eu/en/chikungunya/facts/factsheet> [Accessed January 31, 2020].
- ECDC 2018 Epidemiological update: West Nile virus transmission season in Europe, 2018. *Eur. Cent. Dis. Prev. Control.* <https://ecdc.europa.eu/en/news-events/epidemiological-update-west-nile-virus-transmission-season-europe-2018> [Accessed June 10, 2019].
- ECDC 2019 Mosquito maps. *Eur. Cent. Dis. Prev. Control.* <https://ecdc.europa.eu/en/disease-vectors/surveillance-and-disease-data/mosquito-maps>
- EEA 2017 Climate change, impacts and vulnerability in Europe 2016. An indicator-based report. doi: [10.2800/534806](https://doi.org/10.2800/534806)
- EFSA and ECDC 2015 The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2014. *EFSA J.* 13, 4329.
- Environmental Audit Committee 2018 Heatwaves: adapting to climate change. Ninth Report of Session 2017–19. <https://publications.parliament.uk/pa/cm201719/cmselect/cmenvaud/826/826.pdf>
- EU Climate Policy Adaptation policy in the EU – an overview. <https://climatepolicyinfohub.eu/adaptation-policy-eu---overview> [Accessed January 31, 2020].
- Faustini A, Alessandrini ER, Pey J, Perez N, Samoli E et al. 2015 Short-term effects of particulate matter on mortality during forest fires in Southern Europe: results of the MED-PARTICLES Project. *Occup. Environ. Med.* 72, 323–329. doi: [10.1136/oemed-2014-102459](https://doi.org/10.1136/oemed-2014-102459)
- Feitelson E, Tubi A 2017 A main driver or an intermediate variable? Climate change, water and security in the Middle East. *Glob. Environ. Chang.* 44, 39–48. doi: [10.1016/J.GLOENVCHA.2017.03.001](https://doi.org/10.1016/J.GLOENVCHA.2017.03.001)
- Fiore AM, Naik V, Leibensperger EM 2015 Air quality and climate connections. *J. Air Waste Manag. Assoc.* 65, 645–685. doi: [10.1080/10962247.2015.1040526](https://doi.org/10.1080/10962247.2015.1040526)
- Fischer D, Thomas SM, Suk JE, Sudre B, Hess A et al. 2013 Climate change effects on Chikungunya transmission in Europe: geospatial analysis of vector's climatic suitability and virus' temperature requirements. *Int. J. Health Geogr.* 12, 51. doi: [10.1186/1476-072X-12-51](https://doi.org/10.1186/1476-072X-12-51)
- Fouillet A, Rey G, Wagner V, Laaidi K, Empereur-Bissonnet P et al. 2008 Has the impact of heat waves on mortality changed in France since the European heat wave of summer 2003? A study of the 2006 heat wave. *Int. J. Epidemiol.* 37, 309–317. doi: [10.1093/ije/dym253](https://doi.org/10.1093/ije/dym253)
- Fountoulakis KN, Savopoulos C, Zannis P, Apostolopoulou M, Fountoukidis I et al. 2016 Climate change

- but not unemployment explains the changing suicidality in Thessaloniki Greece (2000-2012). *J. Affect. Disord.* 193, 331–338. doi: [10.1016/j.jad.2016.01.008](https://doi.org/10.1016/j.jad.2016.01.008)
- Gagnon D, Crandall CG, Kenny GP 2013 Sex differences in postsynaptic sweating and cutaneous vasodilation. *J. Appl. Physiol.* 114, 394–401. doi: [10.1152/jappphysiol.00877.2012](https://doi.org/10.1152/jappphysiol.00877.2012)
- Garrison VH, Foreman WT, Genualdi S, Griffin DW, Kellogg CA et al. 2006 Saharan dust - A carrier of persistent organic pollutants, metals and microbes to the Caribbean? *Rev. Biol. Trop. Int. J. Trop. Biol. Conserv.* 54, 9–21. doi: [10.15517/RBT.V54I3.26867](https://doi.org/10.15517/RBT.V54I3.26867)
- Gasparrini A, Guo Y, Sera F, Vicedo-Cabrera AM, Huber V et al. 2017 Projections of temperature-related excess mortality under climate change scenarios. *Lancet Planet. Heal.* 1, e360–e367. doi: [10.1016/S2542-5196\(17\)30156-0](https://doi.org/10.1016/S2542-5196(17)30156-0)
- Girardi E, Sañé Schepisi M, Goletti D, Bates M, Mwaba P et al. 2017 The global dynamics of diabetes and tuberculosis: the impact of migration and policy implications. *Int. J. Infect. Dis.* 56, 45–53. doi: [10.1016/j.ijid.2017.01.018](https://doi.org/10.1016/j.ijid.2017.01.018)
- Gostin LO, Roberts AE 2015 Forced migration the human face of a health crisis. *JAMA - J. Am. Med. Assoc.* 314, 2125–2126. doi: [10.1001/jama.2015.14906](https://doi.org/10.1001/jama.2015.14906)
- Goudie AS 2014 Desert dust and human health disorders. *Environ. Int.* 63, 101–113. doi: [10.1016/J.ENVINT.2013.10.011](https://doi.org/10.1016/J.ENVINT.2013.10.011)
- Griffin DW 2007 Atmospheric movement of microorganisms in clouds of desert dust and implications for human health. *Clin. Microbiol. Rev.* 20, 459–477. doi: [10.1128/CMR.00039-06](https://doi.org/10.1128/CMR.00039-06)
- Guo Y, Gasparrini A, Li S, Sera F, Vicedo-Cabrera AM et al. 2018 Quantifying excess deaths related to heatwaves under climate change scenarios: A multi-country time series modelling study. *PLoS Med.* 15, e1002629. doi: [10.1371/journal.pmed.1002629](https://doi.org/10.1371/journal.pmed.1002629)
- Habus J, Persic Z, Spicic S, Vince S, Stritof Z et al. 2017 New trends in human and animal leptospirosis in Croatia, 2009–2014. *Acta Trop.* 168, 1–8. doi: [10.1016/j.actatropica.2017.01.002](https://doi.org/10.1016/j.actatropica.2017.01.002)
- Hess JJ, Ebi KL 2016 Iterative management of heat early warning systems in a changing climate. *Ann. N. Y. Acad. Sci.* 1382, 21–30. doi: [10.1111/nyas.13258](https://doi.org/10.1111/nyas.13258)
- Hoegh-Guldberg O, Jacob D, Taylor M, Bindi M, Brown S et al. 2018 Impacts of 1.5°C of global warming on natural and human systems, in *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, eds. Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J et al. (Cambridge, United Kingdom and New York, NY, USA: In press), 175–311.
- IPCC 2018 *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, eds. Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J et al. In press.
- Jeuken A, Bouwer LM, Burzel A, Bosello F, Decian E et al. 2016 Bottom-up climate adaptation strategies towards a sustainable Europe. EU-wide economic evaluation of adaptation to climate change. <https://base-adaptation.eu/sites/default/files/D.6.3.pdf>
- Jiménez E, Linares C, Martínez D, Díaz J 2010 Role of Saharan dust in the relationship between particulate matter and short-term daily mortality among the elderly in Madrid (Spain). *Sci. Total Environ.* 408, 5729–5736. doi: [10.1016/j.scitotenv.2010.08.049](https://doi.org/10.1016/j.scitotenv.2010.08.049)
- Karanasiou A, Moreno N, Moreno T, Viana M-M, de Leeuw F et al. 2012 Health effects from Sahara dust episodes in Europe: Literature review and research gaps. *Environ. Int.* 47, 107–114. doi: [10.1016/J.ENVINT.2012.06.012](https://doi.org/10.1016/J.ENVINT.2012.06.012)
- Katsouyanni K, Samet JM, Anderson HR, Atkinson RW, Le Tertre A et al. 2009 Air pollution and health: a European and North American approach (APHE-NA). *Res. Rep. Health. Eff. Inst.*, 5–90. <http://www.ncbi.nlm.nih.gov/pubmed/20073322> [Accessed April 26, 2019].
- Kendrovski V, Baccini M, Martinez G, Wolf T, Paunovic E et al. 2017 Quantifying Projected Heat Mortality Impacts under 21<sup>st</sup>-Century Warming Conditions for Selected European Countries. *Int. J. Environ. Res. Public Health* 14, 729. doi: [10.3390/ijerph14070729](https://doi.org/10.3390/ijerph14070729)
- Kovats RS, Edwards SJ, Hajat S, Armstrong BG, Ebi KL et al. 2004 The effect of temperature on food poisoning: a time-series analysis of salmonellosis in ten European countries. *Epidemiol. Infect.* 132, 443–453. doi: [10.1017/s0950268804001992](https://doi.org/10.1017/s0950268804001992)
- Lake IR, Gillespie IA, Bentham G, Nichols GL, Lane C et al. 2009 A re-evaluation of the impact of temperature and climate change on foodborne illness. *Epidemiol. Infect.* 137, 1538–1547. doi: [10.1017/S0950268809002477](https://doi.org/10.1017/S0950268809002477)
- Langner J, Engardt M, Baklanov A, Christensen JH, Gauss M et al. 2012 A multi-model study of impacts of climate change on surface ozone in Europe. *Atmos. Chem. Phys.* 12, 10423–10440. doi: [10.5194/acp-12-10423-2012](https://doi.org/10.5194/acp-12-10423-2012)
- Lelieveld J, Klingmüller K, Pozzer A, Pöschl U, Fnais M et al. 2019 Cardiovascular disease burden from ambient air pollution in Europe reassessed using novel hazard ratio functions. *Eur. Heart J.* 40, 1590–1596. doi: [10.1093/eurheartj/ehz135](https://doi.org/10.1093/eurheartj/ehz135)
- Linares C, Carmona R, Tobías A, Mirón IJ, Díaz J 2015a Influence of advections of particulate matter from biomass combustion on specific-cause mortality in

- Madrid in the period 2004–2009. *Environ. Sci. Pollut. Res.* 22, 7012–7019.  
doi: [10.1007/s11356-014-3916-2](https://doi.org/10.1007/s11356-014-3916-2)
- Linares C, Díaz J, Negev M, Martínez GS, Debono R et al. 2020 Impacts of climate change on the public health of the Mediterranean Basin population - Current situation, projections, preparedness and adaptation. *Environ. Res.* 182, 109107.  
doi: [10.1016/j.envres.2019.109107](https://doi.org/10.1016/j.envres.2019.109107)
- Linares C, Martínez-Martin P, Rodríguez-Blázquez C, Forjaz MJ, Carmona R et al. 2016 Effect of heat waves on morbidity and mortality due to Parkinson's disease in Madrid: A time-series analysis. *Environ. Int.* 89–90, 1–6.  
doi: [10.1016/j.envint.2016.01.017](https://doi.org/10.1016/j.envint.2016.01.017)
- Linares C, Sánchez R, Mirón IJ, Díaz J 2015b Has there been a decrease in mortality due to heat waves in Spain? Findings from a multicity case study. *J. Integr. Environ. Sci.* 12, 153–163.  
doi: [10.1080/1943815X.2015.1062032](https://doi.org/10.1080/1943815X.2015.1062032)
- Linthicum KJ, Britch SC, Anyamba A 2016 Rift Valley Fever: An Emerging Mosquito-Borne Disease\*. *Annu. Rev. Entomol.* 61, 395–415.  
doi: [10.1146/annurev-ento-010715-023819](https://doi.org/10.1146/annurev-ento-010715-023819)
- López-Bueno JA, Díaz J, Linares C 2019 Differences in the impact of heat waves according to urban and peri-urban factors in Madrid. *Int. J. Biometeorol.* 63, 371–380. doi: [10.1007/s00484-019-01670-9](https://doi.org/10.1007/s00484-019-01670-9)
- Mäkinen TM, Juvonen R, Jokelainen J, Harju TH, Peitso A et al. 2009 Cold temperature and low humidity are associated with increased occurrence of respiratory tract infections. *Respir. Med.* 103, 456–62.  
doi: [10.1016/j.rmed.2008.09.011](https://doi.org/10.1016/j.rmed.2008.09.011)
- Marcantonio M, Rizzoli A, Metz M, Rosà R, Marini G et al. 2015 Identifying the Environmental Conditions Favouring West Nile Virus Outbreaks in Europe. *PLoS One* 10, e0121158.  
doi: [10.1371/journal.pone.0121158](https://doi.org/10.1371/journal.pone.0121158)
- Martin KL, Hanigan IC, Morgan GG, Henderson SB, Johnston FH 2013 Air pollution from bushfires and their association with hospital admissions in Sydney, Newcastle and Wollongong, Australia 1994–2007. *Aust. N. Z. J. Public Health* 37, 238–243.  
doi: [10.1111/1753-6405.12065](https://doi.org/10.1111/1753-6405.12065)
- Martínez-Solanas È, López-Ruiz M, Wellenius GA, Gasparrini A, Sunyer J et al. 2018 Evaluation of the impact of ambient temperatures on occupational Injuries in Spain. *Environ. Health Perspect.* 126, 067002. doi: [10.1289/EHP2590](https://doi.org/10.1289/EHP2590)
- Martínez GS, Linares C, Ayuso A, Kendrovski V, Boeckmann M et al. 2019 Heat-health action plans in Europe: Challenges ahead and how to tackle them. *Environ. Res.* 176.  
doi: [10.1016/j.envres.2019.108548](https://doi.org/10.1016/j.envres.2019.108548)
- Mayrhuber EA-S, Dücker ML, Wallner P, Arnberger A, Allex B et al. 2018 Vulnerability to heatwaves and implications for public health interventions – A scoping review. *Environ. Res.* 166, 42–54.  
doi: [10.1016/J.ENVRES.2018.05.021](https://doi.org/10.1016/J.ENVRES.2018.05.021)
- McMichael C, Barnett J, McMichael AJ 2012 An ill wind? Climate change, migration, and health. *Environ. Health Perspect.* 120, 646–654.  
doi: [10.1289/ehp.1104375](https://doi.org/10.1289/ehp.1104375)
- Metcalf CJE, Walter KS, Wesolowski A, Buckee CO, Shevliakova E et al. 2017 Identifying climate drivers of infectious disease dynamics: recent advances and challenges ahead. *Proc. R. Soc. B Biol. Sci.* 284, 20170901. doi: [10.1098/rspb.2017.0901](https://doi.org/10.1098/rspb.2017.0901)
- Meul S, Langematz U, Kröger P, Oberländer-Hayn S, Jöckel P 2018 Future changes in the stratosphere-to-troposphere ozone mass flux and the contribution from climate change and ozone recovery. *Atmos. Chem. Phys.* 18, 7721–7738. doi: [10.5194/acp-18-7721-2018](https://doi.org/10.5194/acp-18-7721-2018)
- Milazzo A, Giles LC, Zhang Y, Koehler AP, Hiller JE et al. 2016 The effect of temperature on different *Salmonella* serotypes during warm seasons in a Mediterranean climate city, Adelaide, Australia. *Epidemiol. Infect.* 144, 1231–1240.  
doi: [10.1017/S0950268815002587](https://doi.org/10.1017/S0950268815002587)
- Milojevic A, Armstrong BG, Gasparrini A, Bohnenstengel SI, Barratt B et al. 2016 Methods to estimate acclimatization to urban heat island effects on heat- and cold-related mortality. *Environ. Health Perspect.* 124, 1016–1022.  
doi: [10.1289/ehp.1510109](https://doi.org/10.1289/ehp.1510109)
- Mirabelli MC, Künzli N, Avol E, Gilliland FD, Gauderman WJ et al. 2009 Respiratory symptoms following wildfire smoke exposure. *Epidemiology* 20, 451–459. doi: [10.1097/EDE.0b013e31819d128d](https://doi.org/10.1097/EDE.0b013e31819d128d)
- Miraglia M, Marvin HJP, Kleter GA, Battilani P, Brera C et al. 2009 Climate change and food safety: An emerging issue with special focus on Europe. *Food Chem. Toxicol.* 47, 1009–1021.  
doi: [10.1016/j.fct.2009.02.005](https://doi.org/10.1016/j.fct.2009.02.005)
- Moirano G, Gasparrini A, Acquafredda F, Fratianni S, Merletti F et al. 2018 West Nile Virus infection in Northern Italy: Case-crossover study on the short-term effect of climatic parameters. *Environ. Res.* 167, 544–549. doi: [10.1016/j.envres.2018.08.016](https://doi.org/10.1016/j.envres.2018.08.016)
- Morabito M, Profili F, Crisci A, Francesconi P, Gensini GF et al. 2012 Heat-related mortality in the Florentine area (Italy) before and after the exceptional 2003 heat wave in Europe: an improved public health response? *Int. J. Biometeorol.* 56, 801–810.  
doi: [10.1007/s00484-011-0481-y](https://doi.org/10.1007/s00484-011-0481-y)
- Negev M, Paz S, Clermont A, Pri-Or NG, Shalom U et al. 2015 Impacts of climate change on vector borne diseases in the Mediterranean Basin — implications for preparedness and adaptation policy. *Int. J. Environ. Res. Public Health* 12, 6745–6770.  
doi: [10.3390/ijerph120606745](https://doi.org/10.3390/ijerph120606745)
- Neophytou AM, Yiallourous P, Coull BA, Kleanthous S, Pavlou P et al. 2013 Particulate matter concen-

- trations during desert dust outbreaks and daily mortality in Nicosia, Cyprus. *J. Expo. Sci. Environ. Epidemiol.* 23, 275–280. doi: [10.1038/jes.2013.10](https://doi.org/10.1038/jes.2013.10)
- Nicolay M, Brown LM, Johns R, Ialynytchev A 2016 A study of heat related illness preparedness in homeless veterans. *Int. J. Disaster Risk Reduct.* 18, 72–74. doi: [10.1016/j.ijdr.2016.05.009](https://doi.org/10.1016/j.ijdr.2016.05.009)
- Nuvolone D, Balzi D, Pepe P, Chini M, Scala D et al. 2013 Ozone short-term exposure and acute coronary events: a multicities study in Tuscany (Italy). *Environ. Res.* 126, 17–23. doi: [10.1016/j.envres.2013.08.002](https://doi.org/10.1016/j.envres.2013.08.002)
- O'Dowd CD, de Leeuw G 2007 Marine aerosol production: a review of the current knowledge. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 365, 1753–1774. doi: [10.1098/rsta.2007.2043](https://doi.org/10.1098/rsta.2007.2043)
- Orru H, Ebi KL, Forsberg B 2017 The interplay of climate change and air pollution on health. *Curr. Environ. Heal. Reports* 4, 504–513. doi: [10.1007/s40572-017-0168-6](https://doi.org/10.1007/s40572-017-0168-6)
- Oudin Åström D, Åström C, Forsberg B, Vicedo-Cabrera AM, Gasparrini A et al. 2018 Heat wave-related mortality in Sweden: A case-crossover study investigating effect modification by neighbourhood deprivation. *Scand. J. Public Health*, 140349481880161. doi: [10.1177/1403494818801615](https://doi.org/10.1177/1403494818801615)
- Papa A, Danis K, Baka A, Bakas A, Dougas G et al. 2010 Ongoing outbreak of West Nile virus infections in humans in Greece, July – August 2010. *Eurosurveillance* 15, 19644. doi: [10.2807/ese.15.34.19644-en](https://doi.org/10.2807/ese.15.34.19644-en)
- Paz S 2015 Climate change impacts on West Nile virus transmission in a global context. *Philos. Trans. R. Soc. B Biol. Sci.* 370, 20130561–20130561. doi: [10.1098/rstb.2013.0561](https://doi.org/10.1098/rstb.2013.0561)
- Paz S, Malkinson D, Green MS, Tsioni G, Papa A et al. 2013 Permissive Summer Temperatures of the 2010 European West Nile Fever Upsurge. *PLoS One* 8, e56398. doi: [10.1371/journal.pone.0056398](https://doi.org/10.1371/journal.pone.0056398)
- Paz S, Semenza JC 2013 Environmental drivers of West Nile fever epidemiology in Europe and Western Asia—a review. *Int. J. Environ. Res. Public Health* 10, 3543–3562. doi: [10.3390/ijerph10083543](https://doi.org/10.3390/ijerph10083543)
- Pérez L, Tobías A, Querol X, Pey J, Alastuey A et al. 2012 Saharan dust, particulate matter and cause-specific mortality: a case-crossover study in Barcelona (Spain). *Environ. Int.* 48, 150–155. doi: [10.1016/j.envint.2012.07.001](https://doi.org/10.1016/j.envint.2012.07.001)
- Petersen LR, Brault AC, Nasci RS 2013 West Nile virus: review of the literature. *JAMA* 310, 308–315. doi: [10.1001/jama.2013.8042](https://doi.org/10.1001/jama.2013.8042)
- Polymenakou PN, Mandalakis M, Stephanou EG, Tselepidis A 2008 Particle Size Distribution of Airborne Microorganisms and Pathogens during an Intense African Dust Event in the Eastern Mediterranean. *Environ. Health Perspect.* 116, 292–296. doi: [10.1289/ehp.10684](https://doi.org/10.1289/ehp.10684)
- Proestos Y, Christophides GK, Ergüler K, Tanarhte M, Waldock J et al. 2015 Present and future projections of habitat suitability of the Asian tiger mosquito, a vector of viral pathogens, from global climate simulation. *Philos. Trans. R. Soc. B Biol. Sci.* 370, 1–16. doi: [10.1098/rstb.2013.0554](https://doi.org/10.1098/rstb.2013.0554)
- Ragettli MS, Vicedo-Cabrera AM, Schindler C, Rösli M 2017 Exploring the association between heat and mortality in Switzerland between 1995 and 2013. *Environ. Res.* 158, 703–709. doi: [10.1016/j.envres.2017.07.021](https://doi.org/10.1016/j.envres.2017.07.021)
- Reisen WK, Fang Y, Martinez VM 2006 Effects of temperature on the transmission of west nile virus by *Culex tarsalis* (Diptera: Culicidae). *J. Med. Entomol.* 43, 309–317. doi: [10.1603/0022-2585\(2006\)043\[0309:EOTOTT\]2.0.CO;2](https://doi.org/10.1603/0022-2585(2006)043[0309:EOTOTT]2.0.CO;2)
- Rizwan AM, Dennis LYC, Liu C 2008 A review on the generation, determination and mitigation of Urban Heat Island. *J. Environ. Sci.* 20, 120–128. doi: [10.1016/S1001-0742\(08\)60019-4](https://doi.org/10.1016/S1001-0742(08)60019-4)
- Rohat G, Flacke J, Dosio A, Pedde S, Dao H et al. 2019 Influence of changes in socioeconomic and climatic conditions on future heat-related health challenges in Europe. *Glob. Planet. Change* 172, 45–59. doi: [10.1016/j.gloplacha.2018.09.013](https://doi.org/10.1016/j.gloplacha.2018.09.013)
- Rosenberg A, Weinberger M, Paz S, Valinsky L, Agmon V et al. 2018 Ambient temperature and age-related notified *Campylobacter* infection in Israel: A 12-year time series study. *Environ. Res.* 164, 539–545. doi: [10.1016/j.envres.2018.03.017](https://doi.org/10.1016/j.envres.2018.03.017)
- Salameh P, Zeidan RK, Hallit S, Farah R, Chahine M et al. 2019 Cardiovascular diseases and long-term self-reported exposure to pollution: results of a national epidemiological study in Lebanon. *J. Cardiopulm. Rehabil. Prev.* 39, 43–49. doi: [10.1097/HCR.0000000000000378](https://doi.org/10.1097/HCR.0000000000000378)
- Samoli E, Nastos PT, Paliatatos AG, Katsouyanni K, Priftis KN 2011 Acute effects of air pollution on pediatric asthma exacerbation: Evidence of association and effect modification. *Environ. Res.* 111, 418–424. doi: [10.1016/j.envres.2011.01.014](https://doi.org/10.1016/j.envres.2011.01.014)
- Sanderson H, Hildén M, Russel D, Penha-Lopes G, Capriolo A 2018 *Adapting to climate change in Europe: exploring sustainable pathways, from local measures to wider policies*. Elsevier
- Sanz-Barbero B, Linares C, Vives-Cases C, González JL, López-Ossorio JJ et al. 2018 Heat wave and the risk of intimate partner violence. *Sci. Total Environ.* 644, 413–419. doi: [10.1016/j.scitotenv.2018.06.368](https://doi.org/10.1016/j.scitotenv.2018.06.368)
- Schifano P, Leone M, de Sario M, De'Donato FK, Bargagli AM et al. 2012 Changes in the effects of heat on mortality among the elderly from 1998–2010: results from a multicenter time series study in Italy. *Environ. Heal.* 11, 58. doi: [10.1186/1476-069X-11-58](https://doi.org/10.1186/1476-069X-11-58)
- Schütte S, Gemenne F, Zaman M, Flahault A, Depoux A 2018 Connecting planetary health, climate change,

- and migration. *Lancet Planet. Heal.* 2, e58–e59. doi: [10.1016/S2542-5196\(18\)30004-4](https://doi.org/10.1016/S2542-5196(18)30004-4)
- Semenza JC, Suk JE 2018 Vector-borne diseases and climate change: a European perspective. *FEMS Microbiol. Lett.* 365. doi: [10.1093/femsle/fnx244](https://doi.org/10.1093/femsle/fnx244)
- Semenza JC, Tran A, Espinosa L, Sudre B, Domanovic D et al. 2016 Climate change projections of West Nile virus infections in Europe: implications for blood safety practices. *Environ. Heal.* 15, S28. doi: [10.1186/s12940-016-0105-4](https://doi.org/10.1186/s12940-016-0105-4)
- Semenza JC, Wilson DJ, Parra J, Bontempo BD, Hart M et al. 2008 Public perception and behavior change in relationship to hot weather and air pollution. *Environ. Res.* 107, 401–411. doi: [10.1016/j.envres.2008.03.005](https://doi.org/10.1016/j.envres.2008.03.005)
- Shuman EK 2010 Global climate change and infectious diseases. *N. Engl. J. Med.* 362, 1061–1063. doi: [10.1056/NEJMp0912931](https://doi.org/10.1056/NEJMp0912931)
- Smid M, Russo S, Costa AC, Granell C, Pebesma E 2019 Ranking European capitals by exposure to heat waves and cold waves. *Urban Clim.* 27, 388–402. doi: [10.1016/j.uclim.2018.12.010](https://doi.org/10.1016/j.uclim.2018.12.010)
- Smith KR, Woodward A, Campbell-Lendrum D, Chadee DD, Honda Y et al. 2014 Human Health: Impacts, Adaptation, and Co-benefits, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al., 709–754.
- Stafoggia M, Zauli-Sajani S, Pey J, Samoli E, Alessandrini ER et al. 2016 Desert dust outbreaks in Southern Europe: contribution to daily PM10 concentrations and short-term associations with mortality and hospital admissions. *Environ. Health Perspect.* 124, 413–419. doi: [10.1289/ehp.1409164](https://doi.org/10.1289/ehp.1409164)
- The Eurowinter Group 1997 Cold exposure and winter mortality from ischaemic heart disease, cerebrovascular disease, respiratory disease, and all causes in warm and cold regions of Europe. *Lancet (London, England)* 349, 1341–1346. <http://www.ncbi.nlm.nih.gov/pubmed/9149695> [Accessed April 26, 2019].
- The Malta Resources Authority 2017 Seventh national communication of Malta to the United Nations Framework Convention on Climate Change.
- Tjaden NB, Suk JE, Fischer D, Thomas SM, Beierkuhnlein C et al. 2017 Modelling the effects of global climate change on Chikungunya transmission in the 21 st century. *Sci. Rep.* 7. doi: [10.1038/s41598-017-03566-3](https://doi.org/10.1038/s41598-017-03566-3)
- Toloo G (Sam), FitzGerald G, Aitken P, Verrall K, Tong S 2013 Are heat warning systems effective? *Environ. Heal.* 12. doi: [10.1186/1476-069x-12-27](https://doi.org/10.1186/1476-069x-12-27)
- Tran A, Sudre B, Paz S, Rossi M, Desbrosse A et al. 2014 Environmental predictors of West Nile fever risk in Europe. *Int. J. Health Geogr.* 13, 26. doi: [10.1186/1476-072X-13-26](https://doi.org/10.1186/1476-072X-13-26)
- Turco M, Llasat MC, von Hardenberg J, Provenzale A 2014 Climate change impacts on wildfires in a Mediterranean environment. *Clim. Change* 125, 369–380. doi: [10.1007/s10584-014-1183-3](https://doi.org/10.1007/s10584-014-1183-3)
- UNEP 2018 The Adaptation Gap Report 2018.
- van Loenhout JAF, Rodriguez-Llanes JM, Guha-Sapir D 2016 Stakeholders' Perception on National Heatwave Plans and Their Local Implementation in Belgium and The Netherlands. *Int. J. Environ. Res. Public Health* 13. doi: [10.3390/ijerph13111120](https://doi.org/10.3390/ijerph13111120)
- Vandentorren S, Bretin P, Zeghnoun A, Mandereau-Bruno L, Croisier A et al. 2006 August 2003 heat wave in France: Risk factors for death of elderly people living at home. *Eur. J. Public Health* 16, 583–591. doi: [10.1093/eurpub/ckl063](https://doi.org/10.1093/eurpub/ckl063)
- Vardoulakis S, Dear K, Hajat S, Heaviside C, Eggen B et al. 2014 Comparative assessment of the effects of climate change on heat- and cold-related mortality in the United Kingdom and Australia. *Environ. Health Perspect.* 122, 1285–1292. doi: [10.1289/ehp.1307524](https://doi.org/10.1289/ehp.1307524)
- Viana M-M, Pey J, Querol X, Alastuey A, de Leeuw F et al. 2014 Natural sources of atmospheric aerosols influencing air quality across Europe. *Sci. Total Environ.* 472, 825–833. doi: [10.1016/j.scitotenv.2013.11.140](https://doi.org/10.1016/j.scitotenv.2013.11.140)
- von Glasow R, Bobrowski N, Kern C 2009 The effects of volcanic eruptions on atmospheric chemistry. *Chem. Geol.* 263, 131–142. doi: [10.1016/J.CHEMGEO.2008.08.020](https://doi.org/10.1016/J.CHEMGEO.2008.08.020)
- Waitz Y, Paz S, Meir D, Malkinson D 2018 Temperature effects on the activity of vectors for *Leishmania tropica* along rocky habitat gradients in the Eastern Mediterranean. *J. Vector Ecol.* 43, 205–214. doi: [10.1111/jvec.12304](https://doi.org/10.1111/jvec.12304)
- Watts N, Adger WN, Agnolucci P, Blackstock J, Byass P et al. 2015 Health and climate change: policy responses to protect public health. *Lancet* 386, 1861–1914. doi: [10.1016/S0140-6736\(15\)60854-6](https://doi.org/10.1016/S0140-6736(15)60854-6)
- Watts N, Amann M, Ayeb-Karlsson S, Belesova K, Bouley T et al. 2018 The Lancet Countdown on health and climate change: from 25 years of inaction to a global transformation for public health. *Lancet* 391, 581–630. doi: [10.1016/S0140-6736\(17\)32464-9](https://doi.org/10.1016/S0140-6736(17)32464-9)
- Weichenthal S, Kulka R, Lavigne E, van Rijswijk D, Brauer M et al. 2017 Biomass Burning as a Source of Ambient Fine Particulate Air Pollution and Acute Myocardial Infarction. *Epidemiology* 28, 329–337. doi: [10.1097/EDE.0000000000000636](https://doi.org/10.1097/EDE.0000000000000636)
- WHO 2017a Chikungunya – France. <https://www.who.int/csr/don/25-august-2017-chikungunya-france/en/>
- WHO 2017b Chikungunya – Italy. <https://www.who.int/csr/don/15-september-2017-chikungunya-italy/en/>



- WHO 2018 COP24 special report: health and climate change. World Health Organisation. Licence: CC BY-NC-SA 3.0 IGO. Geneva
- WHO Regional Office for Europe, OECD 2015 Economic cost of the health impact of air pollution in Europe: Clean air, health and wealth. Copenhagen, Denmark.
- Wolf J, Adger WN, Lorenzoni I 2010 Heat waves and cold spells: an analysis of policy response and perceptions of vulnerable populations in the UK. *Environ. Plan. A Econ. Sp.* 42, 2721–2734. doi: [10.1068/a42503](https://doi.org/10.1068/a42503)
- Woodward A, Hales S, Litidamu N, Phillips D, Martin J 2000 Protecting human health in a changing world: The role of social and economic development. *Bull. World Health Organ.* 78, 1148–1155. doi: [10.1590/S0042-96862000000900010](https://doi.org/10.1590/S0042-96862000000900010)
- World Bank 2017 Middle East & North Africa Data. <https://data.worldbank.org/>
- Yardley JE, Stapleton JM, Sigal RJ, Kenny GP 2013 Do heat events pose a greater health risk for individuals with type 2 diabetes? *Diabetes Technol. Ther.* 15, 520–529. doi: [10.1089/dia.2012.0324](https://doi.org/10.1089/dia.2012.0324)
- Zander KK, Botzen WJW, Oppermann E, Kjellstrom T, Garnett ST 2015 Heat stress causes substantial labour productivity loss in Australia. *Nat. Clim. Chang.* 5, 647. doi: [10.1038/nclimate2623](https://doi.org/10.1038/nclimate2623)
- Zauli-Sajani S, Miglio R, Bonasoni P, Cristofanelli P, Marinoni A et al. 2011 Saharan dust and daily mortality in Emilia-Romagna (Italy). *Occup. Environ. Med.* 68, 446–451. doi: [10.1136/oem.2010.058156](https://doi.org/10.1136/oem.2010.058156)

## Information about authors

### Coordinating Lead Authors

Shlomit Paz:

*University of Haifa, Haifa, Israel*

### Lead Authors

Cristina Linares:

*National School of Public Health, Carlos III Institute of Health, Madrid, Spain*

Julio Díaz:

*National School of Public Health, Carlos III Institute of Health, Madrid, Spain*

Maya Negev:

*School of Public Health, University of Haifa, Haifa, Israel*

Gerardo Sanchez-Martinez:

*United Nations Environment Programme-Technical University of Denmark (UNEP-DTU Partnership), Copenhagen, Denmark*

### Contributing Authors

Roberto Debono:

*Ministry for Health, WHO/UNECE National Focal Point, Malta*



# SOCIETY 3-HUMAN SECURITY


**Coordinating Lead Authors:**

**Vally (Vassiliki) Koubi** *(Greece/Switzerland)*

**Lead Authors:**

**Mohamed Behnassi** *(Morocco)*, **Antonietta Elia** *(Spain/Italy)*, **Manolis Grillakis** *(Greece)*, **Ethemcan Turhan** *(Turkey/The Netherlands)*

*Koubi V, Behnassi M, Elia A, Grillakis M, Turhan E 2020 Human security. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 515-538.*



## Table of contents

<b>5.3 Human security</b> .....	<b>518</b>
Executive summary.....	<b>518</b>
5.3.1 Concepts and evidence.....	<b>518</b>
5.3.1.1 <i>Definition and scope of human security in this report</i> .....	<b>518</b>
5.3.2 Past trends and current situation.....	<b>519</b>
5.3.2.1 <i>Livelihood, culture, human rights</i> .....	<b>519</b>
5.3.2.2 <i>Human rights</i> .....	<b>520</b>
5.3.2.3 <i>Environmental change and migration</i> .....	<b>522</b>
5.3.2.4 <i>Conflict and collapse of civilizations</i> .....	<b>523</b>
5.3.3 Projections, vulnerabilities and risks.....	<b>526</b>
5.3.3.1 <i>Livelihoods, culture, human rights</i> .....	<b>526</b>
5.3.3.2 <i>Conflict</i> .....	<b>527</b>
5.3.4 Adaptation.....	<b>528</b>
5.3.4.1 <i>Livelihoods, culture, human rights</i> .....	<b>528</b>
5.3.5 Knowledge gaps.....	<b>530</b>
<b>Box. 5.3.1 Climate change and the Syrian conflict</b> .....	<b>530</b>
<b>References</b> .....	<b>531</b>
<b>Information about authors</b> .....	<b>538</b>



## 5.3 Human security

### Executive summary

Climate change is a risk to human security (*medium confidence*). Climate change threatens human security by a) undermining livelihood, culture, and human rights (Sections 5.3.2.1; 5.3.2.2); b) increasing migration (Section 5.3.2.3); and c) indirectly influencing violent conflict (Section 5.3.2.4).

The impact of climate change on human security depends not only on climate events but also on country's contextual factors, such as geographical, social, cultural, economic, and political conditions, resulting in a substantial heterogeneous effect among Mediterranean countries (Section 5.3.1.1).

Climate and environmental changes interfere with the realization and enjoyment of fundamental, internationally recognized human rights such as life, self-determination, health, and education (Section 5.3.3.2).

Climate change and extreme events influence migration behavior and patterns (*medium confidence*). The majority of migration associated with climate change impacts is internal (mostly within southern and eastern countries of the Mediterranean Basin), but also international migration (primarily between the South and North). While fast-onset environmental and climatic events such as floods and storms lead to more sudden, involuntary, and short-term and short-distance movements, slow-onset events such as drought, desertification and salinity by allowing for adaptation, usually tend to result in immobility or in migration that is generally perceived as being voluntary and often predominantly economically motivated. Migrants themselves may be vulnerable to climate change impacts in destination areas, particularly in urban centers (Section 5.3.2.3).

Climate change could lead to conflict (*low confidence*). Climatic changes have likely played a role in the decline or collapse of ancient civilizations around the Mediterranean Basin. The evidence on the effect of climate change and variability on violent conflict is contested. Although there is little agreement about a direct relationship, there is evidence that climate change increases the risk of violent conflict indirectly through declining human well-being, especially in countries which are poor and are characterized by pre-existing tensions and conflict. Higher food prices caused

by climate changes have led to urban social unrest in MENA countries. The relationship between climate, migration, and conflict is highly complex as it depends on the social, political, cultural, and economic conditions of a specific country (Section 5.3.2.4).

Climate change poses a severe threat to many UNESCO World Heritage sites (*high confidence*). A large number of cultural sites are located in low-lying coastal areas of the Mediterranean, which are predicted to experience severe floods and coastal erosion in the future, indicating an urgent need for protective measures (Section 5.3.3.1).

Climate change is likely to act as a threat multiplier in the MENA region (*low confidence*). Establishing a strong link between future climate change and intrastate conflicts in MENA countries is challenging due to the lack of research for this region, which makes future research essential. Climate change is likely to affect inter-state relations concerning shared water resources, if existing water institutions are not able to accommodate change (Section 5.3.3.2).

Culture is a key factor for successful adaptation policies to environmental change in the highly diverse multicultural setting of the Mediterranean Basin. Climate adaptation policies have the potential to infringe on human rights in the Mediterranean region if they are disconnected from justice, equity, poverty alleviation, social inclusion, and redistribution. To successfully implement adaptation policies, many lower-income and climate-vulnerable Mediterranean countries need sufficient financial resources and science and technology transfers (Section 5.3.4.1).

### 5.3.1 Concepts and evidence

#### 5.3.1.1 Definition and scope of human security in this report

This chapter assesses what is known about the risks climate change poses to individuals and communities, including risks to livelihoods, culture, human rights, migration and political stability in the Mediterranean region. For this assessment, human security is considered a condition that exists when the vital core of human lives is protected, and where people have the freedom and capacity to live with dignity (IPCC 2014, AR5 Chapter 12:759). Human security encompasses univer-

sal (e.g., health and food), and culturally-specific (e.g., religion), material (e.g., clean water), and non-material (e.g., social recognition) elements necessary for survival, sustainable livelihoods, and dignity. Human rights, politically and legally legitimized by the international community, are a specific means of defining limits, benchmarks and social processes that provide human security.

Much research on human security focuses on various short-term threats to the vital core of people's lives, including economic crises, epidemics, extreme events, and violent conflict. There are also social and environmental threats that are more incremental in nature, for example slow economic development or land degradation. In addition, a part of existing research is biased because it disregards threats to human security, which are more pertinent to other social contexts, or tries to advance generalizable findings, which are context-specific (i.e., relevant research done either in northern or southern Mediterranean States).

This chapter specifically assesses research that investigates the ways in which climate change may exacerbate threats such as water security (*Chapter 3.1*), food security (*Chapter 3.2*), health (*Chapter 5.2*) and others. There are underlying processes that reduce the freedom and capacity of individuals and groups to adequately respond to these threats, including poor health, poverty, and restricted access to economic, social, political, and natural resources. The chapter also assesses research on the interaction between state security and human security that suggests that the increased human insecurity that arises from an inability to adapt to climate change may in turn create risks to national security through large-scale migration and an increased risk of violent conflicts. It also assesses countries' role in protecting human security in the presence of climate change.

Human security is an analytical lens that focuses attention on the ways in which cultural, demographic, economic, and political forces interact with climate change in ways that affect individuals and communities to different degrees. The focus is at the local level, but the analysis concerns drivers of change across multiple scales and sectors, including climate, culture, economic and political institutions, and population. Consequently, understanding the effects of climate change on human security requires evidence about social and environmental processes across multiple scales and sectors. This process-based evidence is collected through a wide array of methods (e.g., ethnographic techniques and large datasets) used in a

wide range of academic disciplines including environmental sciences, economics, political science, and law.

This chapter includes assessment of empirical studies from the social sciences using both qualitative and quantitative research designs. Most of these studies examine the interactions between environmental changes and social processes to explain social outcomes. While very few studies are explicitly about climate change and human security (since they mostly focus on climate variability and extreme weather events), all provide insights that could be used to make inferences about the effects of climate change on human security (Koubi 2019). Given the complexity of the processes that link environmental change to human security, uncertainties about the biophysical dimensions of change and the nature of the social science evidence thus far, high-confidence statements about the general effects of environmental change on all aspects of human security are not possible (Scheffran et al. 2012). Yet, there is strong evidence about some aspects of the links between environmental change and human security. While the impacts of environmental change on human security will be experienced the most in developing countries (including many MENA countries), human security is at risk for vulnerable populations throughout the Mediterranean region (IPCC 2014, 2018a).

### 5.3.2 Past trends and current situation

#### 5.3.2.1 Livelihood, culture, human rights

Although anthropogenic climate change may be new, significant local and regional variations in climate have occurred throughout history. Prehistoric modern humans had experienced repeated periods of abrupt and severe climate change, albeit at a slower pace than presently, which was often global in nature, and they responded and adapted to this change with varying degrees of success and a variety of different outcomes (Heyd and Brooks 2009). Many studies have shown that climate change threatens cultural dimensions of lives and livelihoods including the material and psychological aspects of culture, identity, community cohesion and sense of place (e.g., Hess et al. 2008; Brace and Geoghegan 2011; Adger et al. 2012). Since culture, differing widely between countries around the Mediterranean, is embedded in the dominant modes of production, consumption, lifestyles and social organization that give rise to emissions of greenhouse gases, the impacts of

these emissions are often given meaning through cultural interpretations of science and risk (Douglas and Wildavsky 1982; Shove 2003; Hulme 2008). Culture plays an important role in mediating human responses to environmental change and these responses depend heavily on the extent to which societies see themselves as separate from, or as part of the wider physical or "natural" environment (Heyd and Brooks 2009). The cultural dimensions of climate change risks and responses play a role in framing environmental change as a phenomenon of concern to society and they are not less important because they might inform adaptation planning (Adger et al. 2012). Countries require a diversity of adaptation measures very much depending on individual circumstances (UNFCCC 2007). In this perspective, Heyd and Brooks (2009) analyzed the influence of culture on conceptions of, and behavior towards natural systems and processes in a non-western context, and compare this example with the mainstream of western societies (this comparison may be relevant to the Mediterranean context, especially regarding its southern and northern shores). They conclude that culture may serve as a resource in two ways, in relation to the "management" of the non-human sphere and in relation to the development of governance processes, and that a deeper understanding of the cultural mediation of responses to environmental dynamism may be of significant value in the development of resilience to accelerating climate change.

For Richardson et al. (2009), no environmental policy will receive the support it needs, either formally in the political arena or at the pragmatic day-to-day level, unless cultures, values and world perspectives are considered from the outset – this also applies to the Mediterranean context. There are two reasons for this. First, individuals/societies do not perceive sophisticated science-based information and risk assessments in the same way, as those who produce them. Secondly, in order to be effective, policies need to consider the socio-culturally shaped setting that pre-dates the attempt to implement the policies. The following points underscore the significance of this main finding: information about climate change and local interpretations of risk assessments are culturally mediated through particular emotional ways of reasoning, typical meaning-making processes, specific conceptions of landscape and climate variability and change, and idiosyncratic notions of mitigation of risk. Local religious and spiritual beliefs, knowledge systems, understanding of nature-society relationships, and values and ethics influence how individuals and commu-

nities perceive and respond to climate change; and the implementation of adaptation strategies can raise issues that cut across power relations in existing situations of inequality, which may have unforeseen long-term effects for individuals and communities. Therefore, research on the role of culture, values, and worldviews in both the generation of and responses to climate change should become a top priority (Richardson et al. 2009).

Most regions in the Mediterranean Basin (and the rest of the world) have undergone environmental change since their first occupation by humans, and understanding how people who settled in these places chose to cope and/or adapt (or failed to adapt) may play a vital role in sustainable planning of our modern societies (Latorre et al. 2016). By affecting the availability of food and water to people, environmental and climate drivers are key factors for understanding this process. It is, however, an oversimplification to state that these drivers are directly responsible for livelihood and cultural changes. Although there are many examples in the literature of large civilizations that suffered some degree of collapse, to attribute these social changes solely on shifting climatic conditions disregards other complex processes associated with how human societies make decisions regarding resource constraints. Latorre et al. (2016) assert that the effects of climate change on past societies cannot be understated and there are often multiple feedbacks. Case by case comparative studies can bring out commonalities across livelihoods and cultures, varying geographies and global climate shifts.

### 5.3.2.2 Human rights

Climate change affects the rights of individuals and communities. The IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels (IPCC 2018a) recently assessed that climate change constitutes a massive threat to the enjoyment of economic, social and cultural rights. Following the release of this report, the Committee on Economic, Social and Cultural Rights adopted a statement on "Climate change and the International Covenant on Economic, Social and Cultural Rights" in which it welcomed the voluntary national commitments made under the climate change regime, and also emphasized that States have human rights obligations that should guide them in the design and implementation of measures to address climate change (CESCR 2018).

The current scientific and policy debate on links between environment and human rights increas-



CLIMATE CHANGE	IMPACT ON PEOPLE	RIGHTS IMPLICATED
<b>SEA LEVEL RISE</b> <ul style="list-style-type: none"> <li>• Flooding</li> <li>• Sea surges</li> <li>• Erosion</li> <li>• Salinization of land and water</li> </ul>	<ul style="list-style-type: none"> <li>• Loss of land</li> <li>• Drowning, injury</li> <li>• Lack of clean water, disease</li> <li>• Damage to coastal infrastructure, homes, and property</li> <li>• Loss of agricultural lands</li> <li>• Threat to tourism, beach loss</li> <li>• Quality of agricultural food products</li> <li>• Fishing and marine life that may affect the quality and quantities of sea foods</li> </ul>	<ul style="list-style-type: none"> <li>• Self-determination (ICCPR; ICESCR, 1)</li> <li>• Life (ICCPR, 6)</li> <li>• Health (ICESCR, 12)</li> <li>• Water (CEDAW, 14; ICRC, 24)</li> <li>• Means of subsistence (ICESCR, 1)</li> <li>• Adequate standard of living (ICESCR, 12)</li> <li>• Adequate housing (ICESCR, 12)</li> <li>• Culture (ICCPR, 27)</li> <li>• Property (UDHR, 17)</li> </ul>
<b>TEMPERATURE INCREASE</b> <ul style="list-style-type: none"> <li>• Change in the pattern of diseases</li> <li>• Coral bleaching</li> <li>• Impacts on fisheries</li> </ul>	<ul style="list-style-type: none"> <li>• Spread of disease</li> <li>• Changes in traditional fishing livelihoods and commercial fishing</li> <li>• Threat to tourism, lost coral and fish diversity</li> </ul>	<ul style="list-style-type: none"> <li>• Life (ICCPR, 6)</li> <li>• Health (ICESCR, 12)</li> <li>• Means of subsistence (ICESCR, 1)</li> <li>• Adequate standard of living (ICESCR, 12)</li> </ul>
<b>EXTREME WEATHER EVENTS</b> <ul style="list-style-type: none"> <li>• High intensity</li> <li>• Storms</li> <li>• Sea surges</li> </ul>	<ul style="list-style-type: none"> <li>• Displacement of populations</li> <li>• Contamination of water supply</li> <li>• Damage to infrastructure: delays in medical treatment, food crisis</li> <li>• Psychological distress</li> <li>• Increased transmission of disease</li> <li>• Damage to agricultural lands</li> <li>• Disruption in educational services</li> <li>• Damage to tourism sector</li> <li>• Massive property damage</li> </ul>	<ul style="list-style-type: none"> <li>• Life (ICCPR, 6)</li> <li>• Health (ICESCR, 12)</li> <li>• Water (CEDAW, 14; ICRC, 24)</li> <li>• Means of subsistence (ICESCR, 1)</li> <li>• Adequate standard of living (ICESCR, 12)</li> <li>• Adequate and secure housing (ICESCR, 12)</li> <li>• Education (ICESCR, 14)</li> <li>• Property (UDHR, 17)</li> </ul>
<b>PRECIPITATION CHANGES</b> <ul style="list-style-type: none"> <li>• Change in disease vectors</li> <li>• Erosion</li> <li>• Change in water safety</li> </ul>	<ul style="list-style-type: none"> <li>• Outbreak of disease</li> <li>• Depletion of agricultural soils</li> </ul>	<ul style="list-style-type: none"> <li>• Life (ICCPR, 6)</li> <li>• Health (ICESCR, 12)</li> <li>• Means of subsistence (ICESCR, 1)</li> </ul>

**Table 5.4 | Climate change impacts on selected human rights** (source: submission by the Maldives to the Office of the High Commissioner for Human Rights (OHCHR) in September 2008, as part of OHCHR's consultative study on the relationship between climate change and human rights, reproduced in CIEL (2013)).

ingly perceives climate change as a "risk multiplier" and a key cross-cutting issue. Climate and environmental change thereby are seen as one of the greatest challenges to the promotion of and effective implementation of fundamental rights since its human implications are already serious and alarming. Climate change is thus rapidly becoming an issue of justice and inequality for millions of people around the world and for future generations, which will suffer severe loss and damage as well. The international community's past failure – and insufficient current actions – to mitigate and adapt to climate change according to available scientific evidence further threaten human rights (CIEL and CARE 2015). More specifically, the rights of vulnerable peoples, who are already experiencing the adverse effects of

climate change, are more threatened than ever (CIEL 2013).

Recent studies and assessments have demonstrated that climate change is putting development and human security at risk (IPCC 2007, 2013, 2014) and the United Nations Human Rights Council has acknowledged in many statements (i.e., UNHRC 2008) that the environmental changes brought about by global warming can interfere with the realization and enjoyment of fundamental, internationally recognized human rights – including those protected by the International Covenants on Civil and Political Rights (ICCPR) and Economic, Social, and Cultural Rights (ICESCR).

Many Human Rights Judicial bodies, both at the universal and regional level, have underlined the link between environment and human rights effectiveness, and State-related obligations in different cases brought before them. For example, the European Court of Human Rights (the ECtHR) has developed its jurisprudence related to human rights and environment in recent years. The ECtHR emphasized on several occasions the impact of a “severe degradation of environment” on the breach of the right to life. Environmental degradation is thus increasingly recognized to threaten the human rights of individuals and minority groups. Therefore, states have the duty to respect, promote, protect, and fulfill human rights, and to secure a healthy environment by adopting positive actions to prevent environmental degradation, in order to guarantee a sustainable environment.

The “right to the highest attainable standard of health” is considered indispensable for the enjoyment of other human rights, and it is widely protected in international and regional instruments, and under national constitutions as well (Elia 2016). Climate change is a serious challenge that undermines and places the fundamental determinants of good health (such as clean air, fresh water, food security, and freedom from disease) at risk (health, *Chapter 5.2*). In the Mediterranean Basin, health impacts on populations vary significantly in terms of the risks individual countries are exposed to, available health services and socio-economic status. The most severe problems are likely to be faced by those countries that already have the biggest problems today – notably those in North Africa and the near East. Climate change could further increase the divide between northern and southern Mediterranean countries. Addressing environmental change is, therefore, perceived as a huge health opportunity that includes reducing the over 6.5 million annual deaths (WHO 2016) from air pollution (most victims are poor, elderly, women and children including rural households and people living in burgeoning low-income cities) (*Sections 2.3 and 5.2.4.1*).

Direct climate impacts, such as extreme weather events (e.g., floods) and rising sea levels, threaten millions of people in coastal and low-lying areas of the Mediterranean Basin and elsewhere. Sea-water intrusion contaminates groundwater in coastal communities, negatively affecting agricultural production and potable water availability. Ocean acidification and changes in weather patterns alter ecosystems and their capacity to provide services to human communities. Increasing weather extremes constrain food security and ac-

cess to nutritious forms of food while changing the prices of commodities in global markets, making food more expensive and harder to access for the poorest people (*Table 5.4*). The links between environmental change and human rights are being increasingly documented (*Table 5.4*), and therefore becoming less controversial and seem beyond dispute (Behnassi et al. 2019).

### 5.3.2.3 Environmental change and migration

Migration has been a defining feature of human populations in the Mediterranean for millennia, well before the emergence of nation states. Environmental changes, both in their rapid and slow-onset versions, have likely been influential to some extent in these migration flows. The question of a direct link between migration and environmental factors has therefore been debated intensely (de Haas 2011; Boas et al. 2019). Nevertheless, the empirical evidence from a large body of recent research, which employs macro-level and micro-level data as well as a diverse range of approaches, shows that, on the one hand, sudden-onset climatic events such as floods and storms lead to more sudden, involuntary, and short-term and short-distance movements (McLeman and Gemenne 2018). Empirical knowledge regarding the effects of slow-onset events such as droughts, on the other hand, remains varied (Adger et al. 2015; Hunter et al. 2015; IPCC 2018a). That is, there is no conclusive evidence on the direction and magnitude of the influence of gradual events on migration and their effects are multidimensional and heterogeneous. Still, it seems that slow-onset events usually tend to result either in migration that is generally perceived as voluntary and often predominantly economically-motivated or in immobility (Hunter et al. 2015; Koubi et al. 2016; Cattaneo et al. 2019).

On a century time-scale, the reduced dependence of Mediterranean people on subsistence agriculture during the 20th century has likely reduced possible direct causalities between environmental change and migration. This shift has been uneven both geographically along North-South and East-West axes of the Mediterranean and has been experienced at different times. Thus in a region whose migration patterns are marked with high uncertainty, environmental factors may “play a certain but rather indirect role” together with demographic, economic and political factors (de Haas 2011). While the effects of environmental and climate change are only “beginning to shape a new and more urgent need for the human se-

curity paradigm” (Behnassi and McGlade 2017), evidence about the relationship between climate change and migration in the Mediterranean is often disputed. The region is often referred to as being among the regions whose human security is most threatened by climate change (Brauch 2012).

The bulk of migration in Northern Africa (White 2011) and the eastern Mediterranean (Weinthal et al. 2015) happens locally and regionally. The effect of climatic factors on migration is, however, contentious. For instance, while some authors argue that the 2006 to 2010 drought in northeastern Syria led a large number of individuals to migrate (Kelley et al. 2015; Ash and Obradovich 2020), others take issue with both the number of migrants and the extent to which climate factors affected migration (Selby et al. 2017). Thus, “whether the drought was a relevant cause of rural livelihood loss and whether this livelihood loss facilitated a significant migration to urban areas remains contested, although the majority of studies support these claims” (Ide 2018a). Other studies on the case of Sahelian migration to Italy between 1995 and 2009, offer explanations for variance by climate factors and yields in migration data, where average annual temperatures are suggested to be the dominant factor in explaining migrations (Pasini and Amendola 2019). A recent global study suggests that climate change will not likely impact asylum seeking patterns everywhere, but probably only in countries undergoing political transformation when faced with immediate impacts of climate change due to inefficient policy response to the latter (Abel et al. 2019). Some authors also suggest that migration in the Mediterranean may be towards higher environmental risk rather than away from it due to pressing challenges of rapid urbanization, depopulation of rural areas, and the decline of traditional livelihoods (Geddes 2015). The challenge, therefore, is to better prepare and adapt infrastructure to sustain and protect migrants in their destinations.

A part of this framing can likely be credited to the work of NGOs and think tanks (Felli 2013). The eastern and southern shores of the Mediterranean Basin (MENA) in particular are frequently “cast as hotspots of mixing social, political and ecological problems, populated by racialized, helpless and passive victims, exposed to erratic and dangerous climate change” (Methmann and Rothe 2014). Rothe (2017) details how the Mediterranean came to be seen as the hotspot of climate migration with “interventions coming mainly from either international organizations or scientists and think tanks in northern industrialized countries.”

#### 5.3.2.4 Conflict and collapse of civilizations

There is a recent increase in research examining the effects of climate change and violent conflict, reflecting policy discourses that climate change impacts and resource scarcity could lead to conflict. While changes in climate could bring groups of people into conflict over scarce resources (Homer-Dixon 1999), there is also evidence that in specific circumstances resource scarcity may drive adaptation and innovation (Butzer 2012). While, on a decadal time-scale, violent conflict appears to have become less common and less intense after the end of the Cold War, since 2013, there have been upsurges in both the number of armed conflicts in the Mediterranean region, and also in the number of battle deaths - mainly as a result of the civil wars in Syria and Iraq (Dupuy et al. 2017). During the same period, there has been an increase in the occurrence and severity of drought events (*Chapter 2*), which by affecting agricultural production and crop yields (*Chapter 3*) (Durigon and de Jong van Lier 2013; Siebert et al. 2014; Schauburger et al. 2017), raise the question whether these factors could contribute to social unrest and conflict (discussed in *Box 5.3.1*).

A few studies explore the relationship between longer-term climate variations and the collapse of past civilizations and empires around the Mediterranean Basin using statistical analysis and data derived from archeological and other historical records (Holmgren et al. 2016; Kaniewski et al. 2018). Major changes in weather patterns, in particular drought conditions, have coincided with the collapse of several previously powerful civilizations in the Aegean and Eastern Mediterranean (Kaniewski et al. 2015), as well as of empires such as the Akkadian (Weiss et al. 1993; Carolin et al. 2019), and the Ottoman empires (Kaniewski et al. 2012). Regarding the Akkadian Empire, which ruled Mesopotamia from the headwaters of the Tigris-Euphrates Rivers to the Persian Gulf (all in what is now Iraq, Syria and parts of southern Turkey) during the late third millennium BC, archeological evidence has shown that this highly developed civilization collapsed abruptly around 4,200 years ago, possibly due to a shift to more arid conditions (Weiss et al. 1993). A recent study using speleothem geochemical records from northern Iran identified two major drought periods, which started 4,510 and 4,260 years ago, and lasted 110 and 290 years respectively. The latter event occurred precisely at the time of the Akkadian Empire’s collapse (Carolin et al. 2019). Similarly, a study coupling climate proxies with archaeo-

logical-historical data and a pollen-based record of agriculture shows that an abrupt shift to drier conditions at ca. AD 1,400 could have contributed to the Ottoman Empire's decline (Kaniewski et al. 2018).

Several studies also provide evidence for a climate-conflict relationship via reduced agricultural production across many centuries. The Little Ice Age appears to have been associated with more cases of political upheaval and warfare in the Ottoman (White 2011; Haldon et al. 2014) and Byzantine (Xoplaki et al. 2016) empires. In particular, the Gelali rebellions between the years 1,550 and 1,603 were reactions to social and economic crisis stemming in part from climatic hardship associated with the Little Ice Age. This evidence from historic antecedents, however, cannot be taken to mean that future changes in climate would lead to large-scale political collapse mainly due to globalization in the contemporary world (Butzer 2012).

Most of the recent research on the connections between climate change and conflict focuses on the effects of climate variability, mainly temperature, precipitation, and, to a lesser extent, natural disasters as proxies for the kinds of longer-term changes that might occur due to climate change on interstate conflict, e.g., civil conflict/war, communal violence as well as low intensity conflict such as demonstrations, protests, and riots. There is very limited research on the climate-conflict nexus in the Mediterranean region. Studies using global datasets have failed to uncover a robust direct relationship between climate variability or natural disasters and civil conflict. On the one hand, some studies report a positive effect of temperature on conflict onset or incidence at the global level (e.g., Landis 2014; Bollfrass and Shaver 2015). In particular, in countries with climates that have strong seasonality such as in most countries in the Mediterranean basin, civil conflict is more likely to occur when warm weather is prolonged (Landis 2014). Deviations in temperature, whether they are positive or negative, as well as changes in mean precipitation show no statistical association with the onset of civil war. On the other hand, others find that precipitation, in particular more rainfall, rather than temperature anomalies are associated with increases in organized political violence—especially in less developed countries (Salehyan and Hendrix 2014). Yet, other studies do not find any effect.

For the Mediterranean region, Böhmelt et al. (2014) report that climate variability measured as

a deviation of the current level of precipitation and temperature, respectively, from their past long-run levels, did not affect the onset of armed conflict during the 1997-2009 period. In addition, the authors show that population pressure, agricultural productivity, and economic development are likely to have a stronger impact on water conflict risk than climate variability (Bernauer et al. 2012; Theisen et al. 2013; Buhaug et al. 2014). They conclude that violent water conflicts are extremely rare, and that factors conducive to restraint, such as stable political conditions, may stimulate cooperation. Higher temperature, though, seems to increase the likelihood of low-level conflicts such as political instability in the form of irregular leader transitions (i.e., coups) (Dell et al. 2012) and incumbents' electoral losses potentially speeding democratic turnover (Obradovich 2017).

A growing body of research examines the connections between natural disasters, i.e., storms, floods or droughts and conflict. By damaging public and private infrastructure, destroying crops and killing livestock, these events can cause or worsen scarcity, subsequently leading to conflict. The direct association between natural disasters and armed conflict is contested (Nel and Righarts 2008; Bergholt and Lujala 2012). However, there is some evidence that natural disasters increase the outbreak of armed conflict in highly ethnically fractionalized countries (Schleussner et al. 2016), lengthen civil conflict and communal conflict (Eastin 2015; Detges 2016; Ghimire and Ferreira 2016) and increase state-sponsored repression (Wood and Wright 2016) and the likelihood of transnational terrorism (Paul and Bagchi 2018). Feitelson and Tubi (2017) examine the effect of the extreme 2007–2010 drought on armed conflict in the Euphrates and lower Jordan River basins and find that, with the exception of Syria, and the consequent spillover into Iraq, droughts do not constitute a main driver of armed conflict in the Middle East. They conclude that droughts may lead to conflict when more fundamental factors, especially adaptive capacity, have been compromised. There is also some evidence that widespread disasters generate solidarity and cooperation rather than conflict (Theisen 2012; Nardulli et al. 2015; Tubi and Feitelson 2016).

A limited body of research examines the connections between climate change and small-scale violent conflicts, i.e., non-state conflict or communal violence. There is some agreement that both decreased rainfall (resource scarcity) and increased rainfall (strategic advantage) in resource dependent societies increase the risk of localized

violence, particularly in pastoral societies in Africa (Fjelde and von Uexkull 2012; Theisen 2012; Detges 2014; Ember et al. 2014; Maystadt et al. 2015; Nordkvelle et al. 2017). Tubi and Feitelson (2016) examine the conflictive and cooperative interactions between Muslim Bedouin herders and Jewish agricultural settlements in Israel's semi-arid northern Negev region during the 1957–1963 drought. They find that while the interactions between these two groups ranged from violent clashes to extensive cooperation and assistance, conflict was limited and that the measures taken by state institutions to directly reduce frictions and to provide relief assistance were central to the overall limited level of conflict.

In response to the challenges of finding a direct association between climate changes and violent conflict, most recent research examines the effects of climate changes on conflict via their effects on well-known drivers of conflict, in particular economic conditions and migration. Starting with the economic conditions channel, research has not established a robust link between climate variables, economic growth, and conflict (Miguel et al. 2004; Ciccone 2011; Bergholt and Lujala 2012; Koubi et al. 2012; Hodler and Raschky 2014; van Weezel 2015). However, recent studies provide consistent evidence that climate changes, via their negative effect on agricultural production, livestock prices, and incomes, affect various types and dynamics of conflict (Bazzi and Blattman 2014; Maystadt and Ecker 2014; Fjelde 2015). Moreover, negative weather shocks occurring during the growing season of local crops increase the continuation and intensity rather than the outbreak of civil conflict (Jun 2017; Harari and Ferrara 2018), especially in regions with agriculturally dependent and politically excluded groups (von Uexkull 2014; Schleussner et al. 2016; von Uexkull et al. 2016). As an example of such shocks, dry spells during the wet season have been found to have serious economic consequences in rain-fed agriculture, which represents a significant fraction in many Mediterranean countries (>90% of cultivated regions in Algeria, Morocco and Tunisia, 57% in Turkey, 64% in Italy or 56% in Portugal (Tramblay and Hertig 2018). As an example, Morocco's 2007 drought reduced wheat production by 76% (Schilling et al. 2012), which has been shown to play a vital socio-economic role in Mediterranean countries (Páscoa et al. 2017).

Several recent studies identify a causal link between higher food prices caused by climate changes and urban social unrest in Africa (Berazneva and Lee 2013; Smith 2014; Bellemare 2015;

Raleigh et al. 2015), while Buhaug et al. (2015) disagree. Rising food prices are considered to have played a significant role in fomenting the Arab Spring unrest across North Africa and the Middle East in 2011 (Johnstone and Mazo 2011; Sternberg 2012; Newman 2020). However, Sneyd et al. (2013) show that such forms of violence are triggered by a complex set of political and economic factors rather than by higher food prices.

By reducing the supply of water in transboundary river basins, climate events can increase the likelihood of interstate conflict. Despite the menace of "water wars" prominently discussed since the early 1990's, shared water, although it does lead to tensions, threats, and even to some localized violence, it does not lead to war (Wolf 2007). Joint precipitation scarcity reduces the likelihood of interstate militarized disputes (Devlin and Hendrix 2014), and water scarcity enhances the incentives of riparian states to cooperate rather than to fight (Dinar et al. 2011). In addition, research shows that riparian states' liberal institutional and economic structures (Kalbhenn 2011; Feitelson and Tubi 2017), commercial treaties (Dinar et al. 2015), the behavior of upper riparian (Feitelson and Tubi 2017) and upstream-downstream relations (Munia et al. 2016), the existence of transboundary treaties (Tir and Stinnett 2012), the number of agreements between riparian states (Dinar et al. 2019), the specific design of international water agreements (Dinar et al. 2015) and institutional frameworks for flexible but specific water allocation mechanisms (Dinar et al. 2015; Oktav 2017) further mitigate the risk of conflict. While hydrological disputes in the Jordan, Nile, and Tigris-Euphrates river basins are mostly seen as a part of territorial security and development-oriented concerns rather than a genuine water issue on its own, climate change is likely to further affect inter-state relations and even threaten the stability of existing water institutions, if these institutions are not able to accommodate change (Dinar et al. 2015; de Stefano et al. 2017).

Regarding the channel of migration, the literature argues that environmental migration can cause conflict in the receiving areas by increasing competition for resources and jobs between immigrants and the native population, upsetting ethnic balance when immigrants are of a different ethnicity than the native population; and by exacerbating fault lines, such as between highly employed and unemployed segments of society, rural and urban areas, etc. (Goldstone 2002; Reuveny 2007). Recent research shows that although residents in receiving areas, in particular urban centers in developing countries, view climate

events (e.g., floods and storms) and conditions (e.g., drought and water/soil salinity) as legitimate reasons to migrate, yet environmental migrants are not seen as more deserving than economic migrants (Spilker et al. 2020).

However, despite the surge in the number of studies on the potential link between environmental change, migration, and conflict, this literature is still far from reaching a consensus on this relationship. For example, while some scholars provide evidence that mass population movements induced by climate shocks were an important factor leading to Syria's uprising and subsequent civil war (Kelley et al. 2015; Ash and Obradovich 2020), others conclude that the occurring drought had little if any impact on the outbreak of conflict (Fröhlich 2016; Selby et al. 2017) (see *Box 5.3.1*). Moreover, limited and often ambiguous empirical evidence based on large-N studies further adds to the difficulty to derive conclusive statements (e.g., Brzoska and Fröhlich 2016). Reuveny (2007), for instance, shows that migration caused by extreme events can induce more conflict in receiving communities, while Raleigh et al. (2008) do not find a significant effect. Bhavnani and Lacina (2015) find that greater rates of internal migration due to climate shocks are associated with a higher risk of riots in Indian states. Ghimire and Ferreira (2016), however, report that disaster-induced migration lengthens the duration of an existing civil conflict, but it does not affect the risk of new conflict outbreaks. Climate changes have also been shown to increase intercommunal violence by affecting pastoralists' copying strategies for access to water and foliage (Adano et al. 2012; Detges 2014). Finally, while long-term climate events, such as droughts, seem to enhance environmental migrants' conflict perceptions in their destination location (Koubi et al. 2018), they may not add to their willingness to support violence (Linke et al. 2018). Migrants who experienced short-term climate events, such as storms or heavy rains, on the other hand, are more likely to support violence if they were themselves victims of violence (Linke et al. 2018). Moreover, migrants to urban areas who had experienced both sudden and gradual climate events, e.g., a drought and a storm, at their location of origin are more likely to join and participate in social movements about migrant rights such as joining a migrant interest group and participating in protest rallies organized by the group even if these may cause violent clashes (Koubi et al. 2021).

Beyond economic and ethno-political conditions, as well as migration flows, the state of social and

political institutions is often found to shape the likelihood of collective violence. Various aspects of the strength and, more often, weaknesses of governments but also conflict-related institutions have been repeatedly demonstrated in recent research (Butler and Gates 2012; Koubi et al. 2012; Böhmelt et al. 2014; O'Loughlin et al. 2014; Linke et al. 2015, 2018; Tubi and Feitelson 2016; Wood and Wright 2016; Detges 2016; Fair et al. 2017; Jones et al. 2017; Ide 2018b).

In summary, there is some evidence that climate change may increase the risk of armed conflict in countries and/or regions, which are poor and highly dependent on agriculture, have few capabilities to cope with climate changes, and are characterized by pre-existing tensions and conflict (Ide et al. 2014; Koubi 2019). However, there is very limited knowledge regarding the mechanisms that connect climate change to conflict (Buhaug 2015; Koubi 2019). There is also very little research on the effects of climate changes on conflict in the Mediterranean region. Consequently, there is an urgent need for research and data collection that can be used to study the processes that could lead from changes in climate to conflict in the Mediterranean Basin.

### 5.3.3 Projections, vulnerabilities and risks

#### 5.3.3.1 Livelihoods, culture, human rights

Countries with high carbon emission potential directly and massively contribute to global warming, which is associated with substantial harms to poor and vulnerable populations, including indigenous people, through their multiple impacts and associated risks (i.e., increasing frequent extreme-weather events, spread of tropical diseases, desertification, rising sea levels, biodiversity loss, decrease of crop yield and food security, etc.) (Jodoin and Lofts 2013; IPCC 2018a). Poor people are much more vulnerable to these impacts and risks because they tend to live in the most exposed areas, typically cannot protect themselves, and lack the means to cope once a threat has materialized (Hallegatte et al. 2016; Hallegatte and Rozenberg 2017; IPCC 2018a; Wallemacq et al. 2018). Assuming that the global distribution of income and wealth will remain as uneven as today, present excess emissions are likely to cause vastly greater harms to poor and vulnerable populations in the future, than they are causing today (Burke et al. 2018; IPCC 2018a; Pretis et al. 2018; Tol 2018). In addition, climate change impacts threaten traditional knowledge about livelihoods in ways that

endanger culture and sense of place attachment (Adger et al. 2012; Tucker et al. 2015; Tschakert et al. 2019). Climate change could increase migration flows. While it is challenging to project the scale of future migration flows as complex interactions between economic, political, social, demographic, and environmental factors shape people's movements (Black et al. 2011), experts agree that millions of people, especially in Sub-Saharan Africa, South Asia, and Latin America, could be forced to move within their countries in the medium term due to climate changes (Rigaud et al. 2018). The scale and geographic scope of this type of population displacement could be one of the greatest human rights challenges of our time. Most countries do not have any governance framework to manage the internal movement of people living within their boundaries and no binding international human rights instrument exists to guide national governments to prepare and respond, creating an enormous protection gap for hundreds of millions of people (McAdam 2016).

Climate change also poses a severe threat to many cultural heritage sites (IUCN 2017). Mediterranean UNESCO World Heritage sites are highly likely to be impacted by climate change over the coming decades, notably from coastal hazards due to sea-level rise. A basin-wide study investigated risks for coastal World Heritage Sites from flooding and erosion under four sea level rise scenarios until 2100. Of 49 sites located in low-lying coastal areas of the Mediterranean, 37 are at risk from a 100-year flood and 42 from coastal erosion, already today (Reimann et al. 2018). The results indicate the urgent need to better protect those sites that are already at risk. Risks will exacerbate in the course of the twenty-first century and beyond (until 2100, flood risk may increase by 50% and erosion risk by 13% across the Mediterranean region), with their magnitude depending on global-scale mitigation efforts in the coming years. For these sites, adaptation can only be implemented to a limited degree because their specific cultural values could be compromised by adaptation measures. If no steps are taken, world heritage sites could lose their values during the coming centuries and will consequently be removed from the UNESCO World Heritage list.

### 5.3.3.2 Conflict

The Mediterranean Basin is characterized by projections of extreme heat, drought and aridity conditions under climate change (*Chapter 2*). The IPCC report on 1.5°C has stated that 'climate-related risks to livelihoods including food security,

water supply and economic growth are projected to increase with global warming of 1.5°C and increase further with 2°C (Summary to Policymakers: B5) (IPCC 2018b).

In this context, climate change is likely to act as a threat multiplier (CNA 2007) in the MENA region by placing additional pressure on already scarce resources and by reinforcing preexisting threats such as poverty, unemployment, and political instability as well as competition over shared water resources along river basins, ultimately leading to violence and conflict. However, establishing a link between future climate change and conflicts in MENA countries is challenging due to the lack of research and consequently scientific knowledge regarding the connections between climate change and conflict in this region. Nevertheless, considering that although the whole region is subjected to climate change but very few countries are experiencing (have experienced) conflict, a few of the main findings of the climate-conflict literature might as well apply here: climate change is very likely to lead to conflict in MENA countries that are poor and are characterized by pre-existing tensions and conflict (Ide et al. 2014; von Uexkull et al. 2016; Waha et al. 2017; Koubi 2019).

Civil conflict is "development in reverse" (Collier et al. 2003). Armed conflict kills people, destroys infrastructure that affects development outcomes, and hampers economic growth that lifts people from poverty, thus reducing the incentives to fight (Chassang 2009; Cederman et al. 2013). Gates et al. (2012) estimate the effect of conflict on income growth. According to their simulations, five consecutive conflict years with more than 1,000 battle deaths reduce per capita incomes by 20% relative to a no-conflict counterfactual, whereas eight additional years of further conflict widen the gap by 5-10%. When it comes to assessing the speed and extent of post-conflict recovery, the analyses in Gates et al. (2012) reveal that the negative impact of short wars is reversed after roughly five peaceful years, whereas long wars are associated with a permanent reduction in per capita incomes of about 10%. Similarly, Costalli et al. (2017) employ synthetic control modelling for 20 conflict countries. On average, years of ongoing conflict see 17.5% lower per capita incomes than carefully constructed counterfactual peace years. Their analysis also reveals massive heterogeneity across conflicts, with estimates ranging from a 1.8% average effect in Egypt (1993-1998) to a staggering 74% for the Liberian civil war (1989-1997 and 1999-2003). Their analysis also shows that the GDP losses due to conflict are especially high in

ethnically fractionalized countries and attribute this effect to disruptions to inter-ethnic cooperation and trade.

Conflict does not only undermine the capacity of governments and non-governmental actors to provide adequate protection from the impacts of climate change, but it is also a major driver of climate vulnerability via its negative effects on economic growth, education, food security and environmental destruction. Broader socio-economic development, expressed by higher growth in education and poverty alleviation, could help in offsetting most of the conflict risk in developing countries associated with reduced economic growth due to implementation of policies to curb GHG emissions (Hegre et al. 2016). Ending violent conflict may be one of the most efficient and cost-effective ways to improve social resilience to climate change.

Regarding future interstate conflicts in the MENA region's major river basins, i.e., Nile, Jordan, Tigris and Euphrates, the limited historic evidence for 'water wars' should not lead to complacency. These river basins will face a strong increase in demand for water due to demographic pressures, industrialization, and urbanization. Simultaneously, while in many cases supply will recede due not only to earlier mismanagement, but also to the impacts of climate change through, for instance, changes in precipitation, increased evaporation, shifts in the seasonality of rain, or droughts. These changes could have security implications at the international level. Consequently, it will take coordinated efforts by all, and especially the major riparian states to adapt to climate change in order to avoid increasing conflict in the future. To the extent that each country securitizes water –i.e., transforms water into a key aspect of its national security and development whose protection justifies the use of particular means such as dams, without taking into considerations the needs of its neighbors and especially downstream states, conflict could be more likely under future climate changes (Feitelson and Tubi 2017). International water agreements and more institutionalized forms of cooperation therefore need to be flexible and robust enough to cope with the emerging threats of climate change (Dinar et al. 2015; Link et al. 2016; de Stefano et al. 2017). There is very little research on the effects of future climate changes on conflict in the Mediterranean region. Consequently, there is an urgent need for research and data collection that can be used to forecast conflict in the face of climate change in the Mediterranean Basin.

### 5.3.4 Adaptation

#### 5.3.4.1 Livelihoods, culture, human rights

Climate change acts directly to change natural weather patterns, but the effects cascade quickly through many sectors with implications for livelihoods (i.e., loss of revenues and subsistence assets), culture (cultural ties to the land, biodiversity and food patterns), and human rights (i.e., rights to food, water, health, etc.). For this reason, governments have a large stake in making adaptation a national priority (Tolba and Saab 2009), and this applies to the Mediterranean as well given the severity of climate impacts in this region, which is increasingly considered as hot spot in many IPCC reports (IPCC 2007, 2014, 2018a). In addition to prevention and mitigation, adaptation is increasingly becoming the focus of formal and informal discussions on human responses to climate change both on the international level and in the Mediterranean.

The effectiveness of different ways to address adaptation to climate change is dependent on the underlying cultural fabric of the human groups involved for their successful implementation (Heyd and Brooks 2009). There are important cultural dimensions to how societies respond and adapt to climate-related risks since culture mediates changes in the environment and changes in societies. Culture is no less central to understanding and implementing adaptation; the identification of risks, decisions about responses, and means of implementation are all mediated by culture. Cultures are dynamic and reflexive and are therefore in turn shaped by the idea of climate change. Hence culture, and its analysis, is central to understanding the causes and meaning of, and human responses to climate change (Douglas and Wildavsky 1982; Shove 2003; Hess et al. 2008; Hulme 2008; Brace and Geoghegan 2011; Adger et al. 2012). This is highly relevant to the Mediterranean context, which is a rich multicultural region composed of countries in Europe, the Middle East, and North Africa, with different, sometimes heterogeneous, cultural representations (The Anna Lindh Report 2018).

Heyd and Brooks (2009) define culture as comprising the ways of living, which involve values, beliefs, practices and material artefacts that condition the production of tangible as well as intangible goods and services needed for the satisfaction of a human group's needs and wants. The culture of any group has to be conceived of as



dynamic, subject to constant transformation and in regular interaction with that of other groups, especially given the interrelationships of human populations in today's increasingly globalized context. Moreover, any set of values, beliefs or practices common to a human group is mediated by power relations, and is not simply the result of adaptation to objective conditions of the natural environment. Nevertheless, particular cultural patterns are among the factors that distinguish human groups, and may play a crucial role in the ability of these groups to cope with environmental/climate changes. In line with this perspective, Rull et al. (2016) tried to explain in a case study the concurrence of conspicuous climatic, ecological and cultural changes during the last millennia and how natural and anthropogenic drivers of change, as well as their potential synergies, might have been influential in determining some cultural shifts.

Climate adaptation policies have the potential to infringe on human rights in the Mediterranean region if their conception and implementation are not adequate or disconnected from some concerns such as justice, equity, poverty alleviation, social inclusion, and redistribution (Behnassi 2019). For instance, adaptation actions, such as relocation in response to sea-level rise or other environmental factors, may affect the right to culture, particularly for indigenous peoples, local communities and other vulnerable groups. Undeniably, relocation can have a particular impact on the right to culture of indigenous peoples whose cultural and spiritual practices are tied to the land, or for local communities who might lose access to significant sites such as ancestral burial grounds (Jodoin and Lofts 2013). Adaptation policies in the Mediterranean may have human rights implications, such as those pertaining to food, water, forest, and the availability of other resources to support the adaptation needs of vulnerable populations.

Governments' legal duty under universally recognized, international human rights instruments to protect people from harm, implies the mainstreaming of human rights into adaptation policy and governance (Behnassi 2019). States must therefore ensure that appropriate adaptation measures are taken to protect and fulfill the rights of all persons, particularly those most threatened by negative climate impacts (e.g., individuals and communities living in vulnerable areas of the Mediterranean such as small islands, riparian and low-lying coastal zones, and arid regions). In the context of adaptation, the protection of cultur-

al rights requires that states avoid or minimize policies that could affect these rights. To this end, the protection of cultural rights is increasingly considered as part of adaptation – and even mitigation – policies, and appropriate scoping and risk assessment activities should be undertaken during the process.

To implement adaptation policies, Mediterranean countries need sufficient resources. Lower-income and climate-vulnerable countries, especially on the south shore of the Mediterranean, which are characterized by limited adaptive capacity due to poverty and political instability (Price 2017), are not generally in a financial position to efficiently deal with climate change and fully protect their populations from its adverse impacts. Their limited public budgets are usually dedicated to cover other vital sectors, such as infrastructure, health, nutrition, and education. Any attempt to allocate available resources to fund adaptation policies may negatively affect these sectors. The development and application of financial safeguards can prevent social and environmental harm and maximize participation, transparency, accountability, equity, and rights protection.

Although the UNFCCC established the Green Climate Fund (GCF), from which many Mediterranean countries are already benefiting to support their adaptation policies (Patel et al. 2016), mechanisms to ensure social and environmental safeguards are yet to be applied to the fund. To do so, institutions involved in funding climate-related activities are required to provide transparent processes, maintain policies and procedures that respect internationally recognized rights, and allow meaningful opportunities for public participation.

Many lower-income and climate-vulnerable Mediterranean countries lack the scientific and technological capacities to deal appropriately and efficiently with environmental change. Thus, science and technology transfer is increasingly considered critical to supporting sustainability and avoiding the shifting of polluting industries from developed countries to the developing world (CIEL 2008), including from northern Mediterranean countries to the southern ones. Establishing an institutional mechanism for science and technology transfer could help to implement a future climate framework in the region. In terms of effective implementation of adaptation strategies, the Sustainable Development Goals (SDG) framework can help ensure that scientific inputs required by the most vulnerable peoples and communities are systematically considered a priority (Behnassi et al. 2019).

### 5.3.5 Knowledge gaps

The evidence reviewed in this chapter shows that climate change poses risks to various dimensions of human security in the Mediterranean region, which arise through diverse causal processes, and which will manifest at different scales. However, many knowledge gaps remain.

The effects of climate change on migration are contingent upon the vulnerability of individuals and societies to such events. In turn, vulnerability is based on physical risk, political, economic and social characteristics, and individual factors such as gender, age, education, wealth, and social capital. In addition, government strategies designed to address vulnerability and to increase resilience can provide the basis for successful adaptation to climate change. Consequently, more widespread and rigorous research is needed in the following areas:

- The conditioning effect of socio-economic and political factors at different levels. Local contextual factors might even be more important in conditioning migration as vulnerability differs significantly across climate-affected communities/areas.
- Micro-level analysis aiming at identifying the climate and non-climate-related determinants of individual migration decisions.
- The compound effects of both slow-onset climate change and sudden-onset extreme events on migration.
- Collecting better data on migration and natural disasters as well as using better modelling techniques to predict future migration flows.
- The implications of migration on vulnerability, especially in the case of rural-to-urban migration.

One of the most critical problems faced by Mediterranean populations is violent conflict, which often triggers migration and loss of life. Climate change is likely to act as a “threat multiplier” in the Mediterranean region for these processes. However, our knowledge is limited regarding how natural disasters interact with and/or are conditioned by socio-economic, political, and demographic factors that cause conflict. Future research should examine:

- When and where are climate conditions and other environmental challenges likely to lead to conflict or social unrest?
- How and why is climate change associated with conflict, and how are other environmental change drivers involved?

- How will conflict patterns evolve over time under different scenarios of future anthropogenic climate and environmental change?
- Which and how could mitigation and adaptation policies amplify the likelihood of conflict?

### BOX 5.3.1

#### Climate change and the Syrian conflict

Climate variability is popularly reported to be a significant cause of the Syrian conflict that began in 2011. Long-term drought and vulnerability of the population to drought led to large-scale internal migration that contributed to the 2011 popular unrest, which spiraled into Syria’s civil war (Gleick 2014; Kelley et al. 2015; Werrell et al. 2015).

Several studies, however, dispute these claims (de Châtel 2014; Kelley et al. 2015; Hendrix 2017; Selby et al. 2017), and argue that the conflict in Syria was not caused by drought but rather was the result of several factors, including water resource degradation, income inequality and increasing poverty, pre-existing socio-economic grievances, long-standing natural resource mismanagement by the government, and the collapse of Syria’s agrarian and rentier model of state-building and development. These studies identify government practices as being far more influential drivers than drought, especially since similar drought conditions did not stimulate conflict in neighboring countries (Hendrix 2017).

Authors of the original study on the Fertile Crescent rebut these claims and insist that overlap of acute vulnerability and the long-term drying trend was compounded with “population growth, poor agricultural policies, aggressive economic liberalization and the influx of Iraqi refugees” to create the suggested link between climate change, drought, displacement and unrest (Kelley et al. 2017). However, other critics suggest that the driving mechanisms between weather-related risks, the resource base and the subsequent risks through which climate-conflict-migration interaction occurs, as well as the evidence for exacerbation of displacement due to climate factors are rather poor (Chalilior et al. 2018). In his review of the available literature on the Syrian civil war, migration and climate change relations, Ide et al. (2018a) conclude that despite the multidecadal drying trend and an exceptional drought before the onset of the conflict “comprehensive evidence through attribution studies of a (probabilistic) link to climate change is still lacking”, and that there is still a lack of knowledge regarding whether the drought induced “migration, intensified existing grievances and facilitated the onset of protests and the subsequent civil war.”

## References

- Abel GJ, Brottrager M, Cuaresma JC, Muttarak R 2019 Climate, conflict and forced migration. *Glob. Environ. Chang.* 54, 239–249. doi: [10.1016/j.gloenvcha.2018.12.003](https://doi.org/10.1016/j.gloenvcha.2018.12.003)
- Adano WR, Dietz T, Witsenburg K, Zaal F 2012 Climate change, violent conflict and local institutions in Kenya's drylands. *J. Peace Res.* 49, 65–80. doi: [10.1177/0022343311427344](https://doi.org/10.1177/0022343311427344)
- Adger WN, Arnell NW, Black R, Dercon S, Geddes A et al. 2015 Focus on environmental risks and migration: Causes and consequences. *Environ. Res. Lett.* 10, 060201. doi: [10.1088/17489326/10/6/060201](https://doi.org/10.1088/17489326/10/6/060201)
- Adger WN, Barnett J, Brown K, Marshall N, O'Brien K 2012 Cultural dimensions of climate change impacts and adaptation. *Nat. Clim. Chang.* 3, 112–117. doi: [10.1038/nclimate1666](https://doi.org/10.1038/nclimate1666)
- Ash K, Obradovich N 2020 Climatic stress, internal migration, and Syrian civil war onset. *J. Conflict Resolut.* 64, 3–31. doi: [10.1177/0022002719864140](https://doi.org/10.1177/0022002719864140)
- Bazzi S, Blattman C 2014 Economic Shocks and Conflict: Evidence from Commodity Prices. *Am. Econ. J. Macroecon.* 6, 1–38. doi: [10.1257/mac.6.4.1](https://doi.org/10.1257/mac.6.4.1)
- Behnassi M 2019 *Mainstreaming a Rights-Based Approach in the Global Climate Regime*. Springer, Cham. doi: [10.1007/978-3-319-92828-9\\_1](https://doi.org/10.1007/978-3-319-92828-9_1)
- Behnassi M, Gupta H, Pollmann O 2019 *Human and Environmental Security in the Era of Global Risks*. DE: Spring. Berlin-Heidelberg: Springer International Publishing. doi: [10.1007/978-3-319-92828-9](https://doi.org/10.1007/978-3-319-92828-9)
- Behnassi M, McGlade K 2017 *Environmental Change and Human Security in Africa and the Middle East*. Berlin, Heidelberg: Springer.
- Bellemare MF 2015 Rising Food Prices, Food Price Volatility, and Social Unrest. *Am. J. Agric. Econ.* 97, 1–21. doi: [10.1093/ajae/aau038](https://doi.org/10.1093/ajae/aau038)
- Berazneva J, Lee DR 2013 Explaining the African food riots of 2007–2008: An empirical analysis. *Food Policy* 39, 28–39. doi: [10.1016/j.foodpol.2012.12.007](https://doi.org/10.1016/j.foodpol.2012.12.007)
- Bergholt D, Lujala P 2012 Climate-related natural disasters, economic growth, and armed civil conflict. *J. Peace Res.* 49, 147–162. doi: [10.1177/0022343311426167](https://doi.org/10.1177/0022343311426167)
- Bernauer T, Böhmelt T, Koubi V 2012 Environmental changes and violent conflict. *Environ. Res. Lett.* 7, 15601. doi: [10.1088/1748-9326/7/1/015601](https://doi.org/10.1088/1748-9326/7/1/015601)
- Bhavnani RR, Lacina B 2015 The effects of weather-induced migration on sons of the soil violence in India. *World Polit.* 67, 760–794. doi: [10.1017/S0043887115000222](https://doi.org/10.1017/S0043887115000222)
- Black R, Adger WN, Arnell NW, Dercon S, Geddes A et al. 2011 The effect of environmental change on human migration. *Glob. Environ. Chang.* 21, S3–S11. doi: [10.1016/j.gloenvcha.2011.10.001](https://doi.org/10.1016/j.gloenvcha.2011.10.001)
- Boas I, Farbotko C, Adams H, Sterly H, Bush S et al. 2019 Climate migration myths. *Nat. Clim. Chang.* 9, 901–903. doi: [10.1038/s41558-019-0633-3](https://doi.org/10.1038/s41558-019-0633-3)
- Böhmelt T, Bernauer T, Buhaug H, Gleditsch NP, Tribaldos T et al. 2014 Demand, supply, and restraint: Determinants of domestic water conflict and cooperation. *Glob. Environ. Chang.* 29, 337–348. doi: [10.1016/j.gloenvcha.2013.11.018](https://doi.org/10.1016/j.gloenvcha.2013.11.018)
- Bollfrass A, Shaver A 2015 The effects of temperature on political violence: global evidence at the sub-national level. *PLoS One* 10, e0123505. doi: [10.1371/journal.pone.0123505](https://doi.org/10.1371/journal.pone.0123505)
- Brace C, Geoghegan H 2011 Human geographies of climate change: Landscape, temporality, and lay knowledges. *Prog. Hum. Geogr.* 35, 284–302. doi: [10.1177/0309132510376259](https://doi.org/10.1177/0309132510376259)
- Brauch HG 2012 Policy Responses to Climate Change in the Mediterranean and MENA Region during the Anthropocene. *Hexag. Ser. Hum. Environ. Secur. Peace*, 719–794. doi: [10.1007/978-3-642-28626-1\\_37](https://doi.org/10.1007/978-3-642-28626-1_37)
- Brzoska M, Fröhlich C 2016 Climate change, migration and violent conflict: vulnerabilities, pathways and adaptation strategies. *Migr. Dev.* 5, 190–210. doi: [10.1080/21632324.2015.1022973](https://doi.org/10.1080/21632324.2015.1022973)
- Buhaug H 2015 Climate-conflict research: some reflections on the way forward. *Wiley Interdiscip. Rev. Clim. Chang.* 6, 269–275. doi: [10.1002/wcc.336](https://doi.org/10.1002/wcc.336)
- Buhaug H, Benjaminsen TA, Sjaastad E, Magnus Theisen O 2015 Climate variability, food production shocks, and violent conflict in Sub-Saharan Africa. *Environ. Res. Lett.* 10, 125015. doi: [10.1088/1748-9326/10/12/125015](https://doi.org/10.1088/1748-9326/10/12/125015)
- Buhaug H, Nordkvelle J, Bernauer T, Böhmelt T, Brzoska M et al. 2014 One effect to rule them all? A comment on climate and conflict. *Clim. Change* 127, 391–397. doi: [10.1007/s10584-014-1266-1](https://doi.org/10.1007/s10584-014-1266-1)
- Burke M, Davis WM, Diffenbaugh NS 2018 Large potential reduction in economic damages under UN mitigation targets. *Nature* 557, 549–553. doi: [10.1038/s41586-018-0071-9](https://doi.org/10.1038/s41586-018-0071-9)
- Butler CK, Gates S 2012 African range wars: Climate, conflict, and property rights. *J. Peace Res.* 49, 23–34. doi: [10.1177/0022343311426166](https://doi.org/10.1177/0022343311426166)
- Butzer KW 2012 Collapse, environment, and society. *Proc. Natl. Acad. Sci. U. S. A.* 109, 3632–3639. doi: [10.1073/pnas.1114845109](https://doi.org/10.1073/pnas.1114845109)
- Carolin SA, Walker RT, Day CC, Ersek V, Sloan RA et al. 2019 Precise timing of abrupt increase in dust activity in the Middle East coincident with 4.2 ka social change. *Proc. Natl. Acad. Sci. U. S. A.* 116, 67–72. doi: [10.1073/pnas.1808103115](https://doi.org/10.1073/pnas.1808103115)
- Cattaneo C, Beine M, Fröhlich CJ, Kniveton D, Martinez-Zarzoso I et al. 2019 Human migration in the era of climate change. *Rev. Environ. Econ. Policy* 13, 189–206. doi: [10.1093/reep/rez008](https://doi.org/10.1093/reep/rez008)
- Cederman L-E, Gleditsch KS, Buhaug H 2013 *Inequality, Grievances, and Civil War*. Cambridge University

- Press. doi: [10.1017/cbo9781139084161](https://doi.org/10.1017/cbo9781139084161)
- CESCR 2018 Climate change and the International Covenant on Economic, Social and Cultural Rights. Statement of the Committee on Economic, Social and Cultural Rights.
- Challinor AJ, Adger WN, Benton TG, Conway D, Joshi M et al. 2018 Transmission of climate risks across sectors and borders. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376, 20170301. doi: [10.1098/rsta.2017.0301](https://doi.org/10.1098/rsta.2017.0301)
- Chassang S 2009 Economic Shocks and Civil War. *Quart. J. Polit. Sci.* 4, 211–228. doi: [10.1561/100.00008072](https://doi.org/10.1561/100.00008072)
- Cicchone A 2011 Economic Shocks and Civil Conflict: A Comment. *Am. Econ. J. Appl. Econ.* 3, 215–227. doi: [10.1257/app.3.4.215](https://doi.org/10.1257/app.3.4.215)
- CIEL 2008 A Rights-Based Approach to Climate Change Mitigation: The Clean Development Mechanism of the Kyoto Protocol. [prepared for the International Union for the Conservation of Nature].
- CIEL 2013 Climate Change and Human Rights: A Primer. Brill. doi: [10.1163/9789004322714\\_cclc\\_2015-0142-004](https://doi.org/10.1163/9789004322714_cclc_2015-0142-004)
- CIEL and CARE 2015 Climate change: Tackling the greatest human rights challenge of our time—Recommendations for effective action on climate change and human rights. [http://www.carefrance.org/ressources/themas/1/4566.CARE\\_and\\_CIEL\\_-\\_Climate\\_Change\\_and\\_.pdf](http://www.carefrance.org/ressources/themas/1/4566.CARE_and_CIEL_-_Climate_Change_and_.pdf)
- CNA 2007 National security and the threat of climate change.
- Collier P, Elliot VL, Hegre H, Hoeffler A, Reynal-Querol M et al. 2003 Breaking the Conflict Trap: Civil War and Development Policy. Washington, DC: World Bank Publications. doi: [10.1037/e504012013-001](https://doi.org/10.1037/e504012013-001)
- Costalli S, Moretti L, Pischedda C 2017 The economic costs of civil war. Synthetic counterfactual evidence and the effects of ethnic fractionalization. *J. Peace Res.* 54, 80–98. doi: [10.1177/0022343316675200](https://doi.org/10.1177/0022343316675200)
- de Châtel F 2014 The Role of Drought and Climate Change in the Syrian Uprising: Untangling the Triggers of the Revolution. *Middle East. Stud.* 50, 521–535. doi: [10.1080/00263206.2013.850076](https://doi.org/10.1080/00263206.2013.850076)
- de Haas H 2011 Mediterranean migration futures: Patterns, drivers and scenarios. *Glob. Environ. Chang.* 21, S59–S69. doi: [10.1016/j.gloenvcha.2011.09.003](https://doi.org/10.1016/j.gloenvcha.2011.09.003)
- de Stefano L, Petersen-Perlman JD, Sproles EA, Eyraud J, Wolf AT 2017 Assessment of transboundary river basins for potential hydro-political tensions. *Glob. Environ. Chang.* 45, 35–46. doi: [10.1016/j.gloenvcha.2017.04.008](https://doi.org/10.1016/j.gloenvcha.2017.04.008)
- Dell M, Jones BF, Olken BA 2012 Temperature shocks and economic growth: Evidence from the last half century. *Am. Econ. J. Macroecon.* 4, 66–95. doi: [10.1257/mac.4.3.66](https://doi.org/10.1257/mac.4.3.66)
- Detges A 2014 Close-up on renewable resources and armed conflict. *Polit. Geogr.* 42, 57–65. doi: [10.1016/j.polgeo.2014.06.003](https://doi.org/10.1016/j.polgeo.2014.06.003)
- Detges A 2016 Local conditions of drought-related violence in sub-Saharan Africa. *J. Peace Res.* 53, 696–710. doi: [10.1177/0022343316651922](https://doi.org/10.1177/0022343316651922)
- Devlin C, Hendrix CS 2014 Trends and triggers redux: Climate change, rainfall, and interstate conflict. *Polit. Geogr.* 43, 27–39. doi: [10.1016/j.polgeo.2014.07.001](https://doi.org/10.1016/j.polgeo.2014.07.001)
- Dinar S, Dinar A, Kurukulasuriya P 2011 Scarcity and cooperation along international rivers: an empirical assessment of bilateral treaties. *Int. Stud. Q.* 55, 809–833. doi: [10.1111/j.1468-2478.2011.00671.x](https://doi.org/10.1111/j.1468-2478.2011.00671.x)
- Dinar S, Katz D, de Stefano L, Blankespoor B 2015 Climate change, conflict, and cooperation: Global analysis of the effectiveness of international river treaties in addressing water variability. *Polit. Geogr.* 45, 55–66. doi: [10.1016/j.polgeo.2014.08.003](https://doi.org/10.1016/j.polgeo.2014.08.003)
- Dinar S, Katz D, de Stefano L, Blankespoor B 2019 Do treaties matter? Climate change, water variability, and cooperation along transboundary river basins. *Polit. Geogr.* 69, 162–172. doi: [10.1016/j.polgeo.2018.08.007](https://doi.org/10.1016/j.polgeo.2018.08.007)
- Douglas M, Wildavsky A 1982 Risk and culture: an essay on the selection of technological and environmental dangers. Oxford University Press (OUP). doi: [10.2307/3984511](https://doi.org/10.2307/3984511)
- Dupuy K, Gates S, Nygård HM, Rudolfson I, Rustad SA et al. 2017 Trends in Armed Conflict, 1946–2016. Oslo: PRIO. <https://www.prio.org/utility/DownloadFile.ashx?id=1373&type=publicationfile>
- Durigon A, de Jong van Lier Q 2013 Canopy temperature versus soil water pressure head for the prediction of crop water stress. *Agric. Water Manag.* 127, 1–6. doi: [10.1016/j.agwat.2013.05.014](https://doi.org/10.1016/j.agwat.2013.05.014)
- Eastin J 2015 Fuel to the Fire: Natural Disasters and the Duration of Civil Conflict. *Int. Interact.* 42, 322–349. doi: [10.1080/03050629.2016.1115402](https://doi.org/10.1080/03050629.2016.1115402)
- Elia A 2016 Guaranteeing Access to Palliative Care between National Law and Emerging International Legal Framework: An Overview of the Italian and Spanish Experiences. *Asia Pacific J. Heal. Law Ethics* 10, 71.
- Ember CR, Skoggard I, Adem TA, Faas AJ 2014 Rain and raids revisited: disaggregating ethnic group livestock raiding in the Ethiopian-Kenyan border region. *Civ. Wars* 16, 300–327. doi: [10.1080/13698249.2014.966430](https://doi.org/10.1080/13698249.2014.966430)
- Fair CC, Kuhn P, Malhotra NA, Shapiro J 2017 Natural disasters and political engagement: evidence from the 2010–11 Pakistani floods. *Stanford Univ. Grad. Sch. Bus. Res. Pap.* 17–42. doi: [10.2139/ssrn.2978047](https://doi.org/10.2139/ssrn.2978047)

- Feitelson E, Tubi A 2017 A main driver or an intermediate variable? Climate change, water and security in the Middle East. *Glob. Environ. Chang.* 44, 39–48. doi: [10.1016/J.GLOENVCHA.2017.03.001](https://doi.org/10.1016/J.GLOENVCHA.2017.03.001)
- Felli R 2013 Managing Climate Insecurity by Ensuring Continuous Capital Accumulation: "Climate Refugees" and "Climate Migrants." *New Polit. Econ.* 18, 337–363. doi: [10.1080/13563467.2012.687716](https://doi.org/10.1080/13563467.2012.687716)
- Fjelde H 2015 Farming or Ffighting? Agricultural price shocks and civil war in Africa. *World Dev.* 67, 525–534. doi: [10.1016/j.worlddev.2014.10.032](https://doi.org/10.1016/j.worlddev.2014.10.032)
- Fjelde H, von Uexkull N 2012 Climate triggers: Rain-fall anomalies, vulnerability and communal conflict in Sub-Saharan Africa. *Polit. Geogr.* 31, 444–453. doi: [10.1016/j.polgeo.2012.08.004](https://doi.org/10.1016/j.polgeo.2012.08.004)
- Fröhlich CJ 2016 Climate migrants as protestors? Dispelling misconceptions about global environmental change in pre-revolutionary Syria. *Contemp. Levant* 1, 38–50. doi: [10.1080/20581831.2016.1149355](https://doi.org/10.1080/20581831.2016.1149355)
- Gates S, Hegre H, Nygård HM, Strand H 2012 Development Consequences of Armed Conflict. *World Dev.* 40, 1713–1722. doi: [10.1016/j.worlddev.2012.04.031](https://doi.org/10.1016/j.worlddev.2012.04.031)
- Geddes A 2015 Governing migration from a distance: interactions between climate, migration, and security in the South Mediterranean. *Eur. Secur.* 24, 473–490. doi: [10.1080/09662839.2015.1028191](https://doi.org/10.1080/09662839.2015.1028191)
- Ghimire R, Ferreira S 2016 Floods and armed conflict. *Environ. Dev. Econ.* 21, 23–52. doi: [10.1017/s1355770x15000157](https://doi.org/10.1017/s1355770x15000157)
- Gleick PH 2014 Water, drought, climate change, and conflict in Syria. *Weather. Clim. Soc.* 6, 331–340. doi: [10.1175/wcas-d-13-00059.1](https://doi.org/10.1175/wcas-d-13-00059.1)
- Goldstone JA 2002 Population and security. How demographic change can lead to violent conflict. *J. Int. Aff.* 56, 3–22.
- Haldon J, Roberts N, Izdebski A, Fleitmann D, McCormick M et al. 2014 The climate and environment of Byzantine Anatolia: Integrating science, history, and archaeology. *J. Interdiscip. Hist.* 45, 113–161. doi: [10.1162/JINH{ }\\_a{ }00682](https://doi.org/10.1162/JINH{ }_a{ }00682)
- Hallegatte S, Bangalore M, Bonzanigo L, Fay M, Kane T et al. 2016 *Shock waves: managing the impacts of climate change on poverty*. The World. Washington, DC. doi: [10.1596/978-1-4648-0673-5](https://doi.org/10.1596/978-1-4648-0673-5)
- Hallegatte S, Rozenberg J 2017 Climate change through a poverty lens. *Nat. Clim. Chang.* 7, 250–256. doi: [10.1038/nclimate3253](https://doi.org/10.1038/nclimate3253)
- Harari M, Ferrara E La 2018 Conflict, climate, and cells: A disaggregated analysis. *Rev. Econ. Stat.* 100, 594–608. doi: [10.1162/rest\\_a\\_00730](https://doi.org/10.1162/rest_a_00730)
- Hegre H, Buhaug H, Calvin K V., Nordkvelle J, Waldhoff ST et al. 2016 Forecasting civil conflict along the shared socioeconomic pathways. *Environ. Res. Lett.* 11, 54002. doi: [10.1088/1748-9326/11/5/054002](https://doi.org/10.1088/1748-9326/11/5/054002)
- Hendrix CS 2017 A comment on "climate change and the Syrian civil war revisited." *Polit. Geogr.* 60, 251–252. doi: [10.1016/j.polgeo.2017.06.010](https://doi.org/10.1016/j.polgeo.2017.06.010)
- Hess JJ, Malilay JN, Parkinson AJ 2008 Climate Change: The importance of place. *Am. J. Prev. Med.* 35, 468–478. doi: [10.1016/j.amepre.2008.08.024](https://doi.org/10.1016/j.amepre.2008.08.024)
- Heyd T, Brooks N 2009 Exploring cultural dimensions of adaptation to climate change: Thresholds, values, governance, in *Adapting to Climate Change*, eds. Neil Adger W, Lorenzoni I, O'Brien K (Cambridge University Press), 269–282. doi: [10.1017/cbo9780511596667.018](https://doi.org/10.1017/cbo9780511596667.018)
- Hodler R, Raschky PA 2014 Economic shocks and civil conflict at the regional level. *Econ. Lett.* 124, 530–533. doi: [10.1016/j.econlet.2014.07.027](https://doi.org/10.1016/j.econlet.2014.07.027)
- Holmgren K, Gogou A, Izdebski A, Luterbacher J, Sicre M-A et al. 2016 Mediterranean Holocene climate, environment and human societies. *Quat. Sci. Rev.* 136, 1–4. doi: [10.1016/j.quascirev.2015.12.014](https://doi.org/10.1016/j.quascirev.2015.12.014)
- Homer-Dixon TF 1999 *Environment, Scarcity, and Violence*. Princeton: Princeton University Press.
- Hulme M 2008 The conquering of climate: discourses of fear and their dissolution. *Geogr. J.* 174, 5–16. doi: [10.1111/j.1475-4959.2008.00266.x](https://doi.org/10.1111/j.1475-4959.2008.00266.x)
- Hunter LM, Luna JK, Norton RM 2015 Environmental Dimensions of Migration. *Annu. Rev. Sociol.* 41, 377–397. doi: [10.1146/annurev-soc-073014-112223](https://doi.org/10.1146/annurev-soc-073014-112223)
- Ide T 2018a Climate War in the Middle East? Drought, the Syrian Civil War and the State of Climate-Conflict Research. *Curr. Clim. Chang. Reports* 4, 347–354. doi: [10.1007/s40641-018-0115-0](https://doi.org/10.1007/s40641-018-0115-0)
- Ide T 2018b Does environmental peacemaking between states work? Insights on cooperative environmental agreements and reconciliation in international rivalries. *J. Peace Res.* 55, 351–365. doi: [10.1177/0022343317750216](https://doi.org/10.1177/0022343317750216)
- Ide T, Schilling J, Link JSA, Scheffran J, Ngaruiya G et al. 2014 On exposure, vulnerability and violence: Spatial distribution of risk factors for climate change and violent conflict across Kenya and Uganda. *Polit. Geogr.* 43, 68–81. doi: [10.1016/j.polgeo.2014.10.007](https://doi.org/10.1016/j.polgeo.2014.10.007)
- IPCC 2007 *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* , eds. Parry ML, Canziani OF, Palutikof JP, Van der Linden PJ, Hanson CE Cambridge, United Kingdom and New York, NY, USA.
- IPCC 2013 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* , eds. Stocker TF, Qin D, Plat-

- tner G-K, Tignor M, Allen SK et al. Cambridge, United Kingdom and New York, NY, USA. IPCC 2014 *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change.*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press <http://www.citeulike.org/group/15400/article/13497155> [Accessed March 11, 2017].
- IPCC 2018a *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change.*, eds. Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J et al. In press.
- IPCC 2018b Summary for Policymakers, in *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, eds. Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J et al. (In press).
- IUCN 2017 IUCN World Heritage Outlook 2017. <https://worldheritageoutlook.iucn.org/results>
- Jodoin S, Lofts K 2013 Economic, social, and cultural rights and climate change: a legal reference guide. *New Haven, Ct. CISDL, GEM ASAP.*
- Johnstone S, Mazo J 2011 Global warming and the Arab Spring. *Survival (Lond)*. 53, 11–17. doi: [10.1080/00396338.2011.571006](https://doi.org/10.1080/00396338.2011.571006)
- Jones BT, Mattiacci E, Braumoeller BF 2017 Food scarcity and state vulnerability: Unpacking the link between climate variability and violent unrest. *J. Peace Res.* 54, 335–350. doi: [10.1177/0022343316684662](https://doi.org/10.1177/0022343316684662)
- Jun T 2017 Temperature, maize yield, and civil conflicts in sub-Saharan Africa. *Clim. Change* 142, 183–197. doi: [10.1007/s10584-017-1941-0](https://doi.org/10.1007/s10584-017-1941-0)
- Kalbhenn A 2011 Liberal peace and shared resources – A fair-weather phenomenon? *J. Peace Res.* 48, 715–735. doi: [10.1177/0022343311420459](https://doi.org/10.1177/0022343311420459)
- Kaniewski D, Guiot J, Van Campo E 2015 Drought and societal collapse 3,200 years ago in the Eastern Mediterranean: a review. *Wiley Interdiscip. Rev. Clim. Chang.* 6, 369–382. doi: [10.1002/wcc.345](https://doi.org/10.1002/wcc.345)
- Kaniewski D, Marriner N, Cheddadi R, Guiot J, Van Campo E 2018 The 4.2 ka BP event in the Levant. *Clim. Past* 14, 1529–1542. doi: [10.5194/cp-14-1529-2018](https://doi.org/10.5194/cp-14-1529-2018)
- Kaniewski D, Van Campo E, Weiss H 2012 Drought is a recurring challenge in the Middle East. *Proc. Natl. Acad. Sci. U. S. A.* 109, 3862–3867. doi: [10.1073/pnas.1116304109](https://doi.org/10.1073/pnas.1116304109)
- Kelley C, Mohtadi S, Cane MA, Seager R, Kushnir Y 2017 Commentary on the Syria case: Climate as a contributing factor. *Polit. Geogr.* 60, 245–247. doi: [10.1016/j.polgeo.2017.06.013](https://doi.org/10.1016/j.polgeo.2017.06.013)
- Kelley CP, Mohtadi S, Cane MA, Seager R, Kushnir Y 2015 Climate change in the Fertile Crescent and implications of the recent Syrian drought. *Proc. Natl. Acad. Sci. U. S. A.* 112, 3241–3246. doi: [10.1073/pnas.1421533112](https://doi.org/10.1073/pnas.1421533112)
- Koubi V 2019 Climate Change and Conflict. *Annu. Rev. Polit. Sci.* 22, 343–360. doi: [10.1146/annurev-polisci-050317-070830](https://doi.org/10.1146/annurev-polisci-050317-070830)
- Koubi V, Bernauer T, Kalbhenn A, Spilker G 2012 Climate variability, economic growth, and civil conflict. *J. Peace Res.* 49, 113–127. doi: [10.1177/0022343311427173](https://doi.org/10.1177/0022343311427173)
- Koubi V, Böhmelt T, Spilker G, Schaffer L 2018 The determinants of environmental migrants' conflict perception. *Int. Organ.* 72, 905–936. doi: [10.1017/S0020818318000231](https://doi.org/10.1017/S0020818318000231)
- Koubi V, Nguyen Q, Spilker G, Böhmelt T 2021 Environmental migrants and social movement participation. *J. Peace Res.* 58, xx–xx.
- Koubi V, Spilker G, Schaffer L, Böhmelt T 2016 The role of environmental perceptions in migration decision-making: evidence from both migrants and non-migrants in five developing countries. *Popul. Environ.* 38, 134–163. doi: [10.1007/s11111-016-0258-7](https://doi.org/10.1007/s11111-016-0258-7)
- Landis ST 2014 Temperature seasonality and violent conflict. *J. Peace Res.* 51, 603–618. doi: [10.1177/0022343314538275](https://doi.org/10.1177/0022343314538275)
- Latorre C, Wilmshurst J, von Gunten L 2016 Climate change and cultural evolution across the world. *Past Glob. Chang. Mag.* 24, 55. doi: [10.22498/pages.24.2.55](https://doi.org/10.22498/pages.24.2.55)
- Link PM, Scheffran J, Ide T 2016 Conflict and cooperation in the water-security nexus: a global comparative analysis of river basins under climate change. *Wiley Interdiscip. Rev. Water* 3, 495–515. doi: [10.1002/wat2.1151](https://doi.org/10.1002/wat2.1151)
- Linke AM, O'Loughlin J, McCabe JT, Tir J, Witmer FDW 2015 Rainfall variability and violence in rural Kenya: Investigating the effects of drought and the role of local institutions with survey data. *Glob. Environ. Chang.* 34, 35–47. doi: [10.1016/j.gloenvcha.2015.04.007](https://doi.org/10.1016/j.gloenvcha.2015.04.007)
- Linke AM, Witmer FDW, O'Loughlin J, McCabe JT, Tir J 2018 The consequences of relocating in response to drought: human mobility and conflict in contemporary Kenya. *Environ. Res. Lett.* 13, 094014. doi: [10.1088/1748-9326/aad8cc](https://doi.org/10.1088/1748-9326/aad8cc)
- Maystadt J-F, Calderone M, You L 2015 Local warming and violent conflict in North and South Sudan. *J. Econ. Geogr.* 15, 649–671. doi: [10.1093/jeg/lbu033](https://doi.org/10.1093/jeg/lbu033)
- Maystadt J-F, Ecker O 2014 Extreme weather and civil war: does drought fuel conflict in Somalia through

- livestock price shocks? *Am. J. Agric. Econ.* 96, 1157–1182. doi: [10.1093/ajae/aa010](https://doi.org/10.1093/ajae/aa010)
- McAdam J 2016 From the Nansen Initiative to the platform on disaster displacement: Shaping international approaches to climate change, disasters and displacement. *Univ. N. S. W. Law J.* 39, UNSW Law Research Paper No. 17-4.
- McLeman R, Gemenne F 2018 Environmental migration research: Evolution and current state of the science, in *Handbook of Environmental Displacement and Migration*, eds. McLeman R, Gemenne F (London: Routledge), 3–16.
- Methmann C, Rothe D 2014 Tracing the spectre that haunts Europe: the visual construction of climate-induced migration in the MENA region. *Crit. Stud. Secur.* 2, 162–179. doi: [10.1080/21624887.2014.909226](https://doi.org/10.1080/21624887.2014.909226)
- Miguel E, Satyanath S, Sergenti E 2004 Economic shocks and civil conflict: an instrumental variables approach. *J. Polit. Econ.* 112, 725–753. doi: [10.1086/421174](https://doi.org/10.1086/421174)
- Munia H, Guillaume JHA, Mirumachi N, Porkka M, Wada Y et al. 2016 Water stress in global trans-boundary river basins: significance of upstream water use on downstream stress. *Environ. Res. Lett.* 11, 14002. doi: [10.1088/1748-9326/11/1/014002](https://doi.org/10.1088/1748-9326/11/1/014002)
- Nardulli PF, Peyton B, Bajjalieh J 2015 Climate change and civil unrest: the impact of rapid-onset disasters. *J. Conflict Resolut.* 59, 310–335. doi: [10.1177/0022002713503809](https://doi.org/10.1177/0022002713503809)
- Nel P, Righarts M 2008 Natural Disasters and the Risk of Violent Civil Conflict. *Int. Stud. Q.* 52, 159–185. doi: [10.1111/j.1468-2478.2007.00495.x](https://doi.org/10.1111/j.1468-2478.2007.00495.x)
- Newman E 2020 Hungry, or hungry for change? Food riots and political conflict, 2005–2015. *Stud. Confl. Terror.* 43, 300–324. doi: [10.1080/1057610x.2018.1454042](https://doi.org/10.1080/1057610x.2018.1454042)
- Nordkvelle J, Rustad SA, Salmivalli M 2017 Identifying the effect of climate variability on communal conflict through randomization. *Clim. Change* 141, 627–639. doi: [10.1007/s10584-018-2303-2](https://doi.org/10.1007/s10584-018-2303-2)
- O'Loughlin J, Linke AM, Witmer FDW 2014 Effects of temperature and precipitation variability on the risk of violence in sub-Saharan Africa, 1980–2012. *Proc. Natl. Acad. Sci. U. S. A.* 111, 16712–16717. doi: [10.1073/pnas.1411899111](https://doi.org/10.1073/pnas.1411899111)
- Obradovich N 2017 Climate change may speed democratic turnover. *Clim. Change* 140, 135–147. doi: [10.1007/s10584-016-1833-8](https://doi.org/10.1007/s10584-016-1833-8)
- Oktav ÖZ 2017 Turkey's Water Policy in the Euphrates-Tigris Basin. *Environ. Chang. Hum. Secur. Africa Middle East*, 239–255. doi: [10.1007/978-3-319-45648-5\\_13](https://doi.org/10.1007/978-3-319-45648-5_13)
- Páscoa P, Gouveia CM, Russo A, Trigo RM 2017 The role of drought on wheat yield interannual variability in the Iberian Peninsula from 1929 to 2012. *Int. J. Biometeorol.* 61, 439–451. doi: [10.1007/s00484-016-1224-x](https://doi.org/10.1007/s00484-016-1224-x)
- Pasini A, Amendola S 2019 Linear and nonlinear influences of climatic changes on migration flows: a case study for the 'Mediterranean bridge.' *Environ. Res. Commun.* 1, 11005. doi: [10.1088/2515-7620/ab0464](https://doi.org/10.1088/2515-7620/ab0464)
- Patel S, Watson S, Schalatek L 2016 Climate Finance Regional Briefing: Middle East and North Africa, Climate Funds Update. *Clim. Financ. Fundam.* 9, 4. <https://www.odi.org/sites/odi.org.uk/files/resource-documents/11042.pdf>
- Paul JA, Bagchi A 2018 Does terrorism increase after a natural disaster? An analysis based upon property damage. *Def. Peace Econ.* 29, 407–439. doi: [10.1080/10242694.2016.1204169](https://doi.org/10.1080/10242694.2016.1204169)
- Pretis F, Schwarz M, Tang K, Haustein K, Allen MR 2018 Uncertain impacts on economic growth when stabilizing global temperatures at 1.5°C or 2°C warming. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376, 20160460. doi: [10.1098/rsta.2016.0460](https://doi.org/10.1098/rsta.2016.0460)
- Price RA 2017 Climate change and stability in North Africa. 18. [https://assets.publishing.service.gov.uk/media/5a7052bde915d266017b8aa/242\\_Climate\\_change\\_and\\_stability\\_in\\_Northern\\_Africa.pdf](https://assets.publishing.service.gov.uk/media/5a7052bde915d266017b8aa/242_Climate_change_and_stability_in_Northern_Africa.pdf)
- Raleigh C, Choi HJ, Kniveton D 2015 The devil is in the details: An investigation of the relationships between conflict, food price and climate across Africa. *Glob. Environ. Chang.* 32, 187–199. doi: [10.1016/j.gloenvcha.2015.03.005](https://doi.org/10.1016/j.gloenvcha.2015.03.005)
- Raleigh C, Jordan L, Salehyan I 2008 Assessing the Impact of Climate Change on Migration and Conflict. Washington, DC.
- Reimann L, Vafeidis AT, Brown S, Hinkel J, Tol RSJ 2018 Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-level rise. *Nat. Commun.* 9. doi: [10.1038/s41467-018-06645-9](https://doi.org/10.1038/s41467-018-06645-9)
- Reuveny R 2007 Climate change-induced migration and violent conflict. *Polit. Geogr.* 26, 656–673. doi: [10.1016/j.polgeo.2007.05.001](https://doi.org/10.1016/j.polgeo.2007.05.001)
- Richardson K, Steffen W, Schellnhuber H-J, Alcamo J, Barker T et al. 2009 *Synthesis Report. Climate change: global risks, challenges and decisions, Copenhagen, Denmark, 10-12 March, 2009*. Copenhagen: University of Copenhagen.
- Rigaud KK, de Sherbinin A, Jones B, Bergmann J, Clement V et al. 2018 Groundswell: Preparing for Internal Climate Migration. Washington, DC <https://openknowledge.worldbank.org/handle/10986/29461>
- Rothe D 2017 *Securitizing global warming: a climate of complexity*. Routledge. doi: [10.4324/9781315677514](https://doi.org/10.4324/9781315677514)
- Rull V, Cañellas-Boltà N, Margalef O, Pla-Rabes S,

- Sáez A et al. 2016 Climate changes and cultural shifts on Easter Island during the last three millennia. *Past Glob. Chang. Mag.* 24, 70–71. doi: [10.22498/pages.24.2.70](https://doi.org/10.22498/pages.24.2.70)
- Salehyan I, Hendrix CS 2014 Climate shocks and political violence. *Glob. Environ. Chang.* 28, 239–250. doi: [10.1016/j.gloenvcha.2014.07.007](https://doi.org/10.1016/j.gloenvcha.2014.07.007)
- Schauberger B, Archontoulis S, Arneith A, Balkovič J, Ciais P et al. 2017 Consistent negative response of US crops to high temperatures in observations and crop models. *Nat. Commun.* 8, 13931. doi: [10.1038/ncomms13931](https://doi.org/10.1038/ncomms13931)
- Scheffran J, Brzoska M, Brauch HG, Link PM, Schilling J 2012 *Climate Change, Human Security and Violent Conflict: Challenges for Societal Stability*. Springer. Berlin. <https://www.springer.com/gp/book/9783642286254>
- Schilling J, Freier KP, Hertig E, Scheffran J 2012 Climate change, vulnerability and adaptation in North Africa with focus on Morocco. *Agric. Ecosyst. Environ.* 156, 12–26. doi: [10.1016/J.AGEE.2012.04.021](https://doi.org/10.1016/J.AGEE.2012.04.021)
- Schleussner C-F, Donges JF, Donner R V., Schellnhuber H-J 2016 Armed-conflict risks enhanced by climate-related disasters in ethnically fractionalized countries. *Proc. Natl. Acad. Sci. U. S. A.* 113, 9216–9221. doi: [10.1073/pnas.1601611113](https://doi.org/10.1073/pnas.1601611113)
- Selby J, Dahi OS, Fröhlich C, Hulme M 2017 Climate change and the Syrian civil war revisited. *Polit. Geogr.* 60, 232–244. doi: [10.1016/j.polgeo.2017.05.007](https://doi.org/10.1016/j.polgeo.2017.05.007)
- Shove S 2003 *Comfort, Cleanliness and Convenience: The Social Organization of Normality*. Bloomsbury Academic. doi: [10.5040/9781474214605](https://doi.org/10.5040/9781474214605)
- Siebert S, Ewert F, Eyshi Rezaei E, Kage H, Graß R 2014 Impact of heat stress on crop yield—on the importance of considering canopy temperature. *Environ. Res. Lett.* 9, 044012. doi: [10.1088/1748-9326/9/4/044012](https://doi.org/10.1088/1748-9326/9/4/044012)
- Smith TG 2014 Feeding unrest: Disentangling the causal relationship between food price shocks and sociopolitical conflict in urban Africa. *J. Peace Res.* 51, 679–695. doi: [10.1177/0022343314543722](https://doi.org/10.1177/0022343314543722)
- Sneyd LQ, Legwegoh A, Fraser EDG 2013 Food riots: Media perspectives on the causes of food protest in Africa. *Food Secur.* 5, 485–497. doi: [10.1007/s12571-013-0272-x](https://doi.org/10.1007/s12571-013-0272-x)
- Spilker G, Nguyen Q, Koubi V, Böhmelt T 2020 Attitudes of urban residents towards environmental migration in Kenya and Vietnam. *Nat. Clim. Chang.* 10, 622–627. doi: [10.1038/s41558-020-0805-1](https://doi.org/10.1038/s41558-020-0805-1)
- Sternberg T 2012 Chinese drought, bread and the Arab Spring. *Appl. Geogr.* 34, 519–524. doi: [10.1016/j.apgeog.2012.02.004](https://doi.org/10.1016/j.apgeog.2012.02.004)
- The Anna Lindh Report 2018 Intercultural Trends and Social Change in the Euro-Mediterranean region. <https://www.interculturaltrendsreport.com/wp-content/uploads/2018/11/Anna-Lindh-Report-on-Intercultural-Trends.pdf>
- Theisen OM 2012 Climate clashes? Weather variability, land pressure, and organized violence in Kenya, 1989–2004. *J. Peace Res.* 49, 81–96. doi: [10.1177/0022343311425842](https://doi.org/10.1177/0022343311425842)
- Theisen OM, Gleditsch NP, Buhaug H 2013 Is climate change a driver of armed conflict? *Clim. Change* 117, 613–625. doi: [10.1007/s10584-012-0649-4](https://doi.org/10.1007/s10584-012-0649-4)
- Tir J, Stinnett DM 2012 Weathering climate change: Can institutions mitigate international water conflict? *J. Peace Res.* 49, 211–225. doi: [10.1177/0022343311427066](https://doi.org/10.1177/0022343311427066)
- Tol RSJ 2018 The Economic Impacts of Climate Change. *Rev. Environ. Econ. Policy* 12, 4–25. doi: [10.1093/reep/rex027](https://doi.org/10.1093/reep/rex027)
- Tolba MK, Saab N 2009 Impact of Climate Change on Arab Countries. [http://www.afedonline.org/afed-report09/Full English Report.pdf](http://www.afedonline.org/afed-report09/Full%20English%20Report.pdf)
- Tramblay Y, Hertig E 2018 Modelling extreme dry spells in the Mediterranean region in connection with atmospheric circulation. *Atmos. Res.* 202, 40–48. doi: [10.1016/j.atmosres.2017.11.015](https://doi.org/10.1016/j.atmosres.2017.11.015)
- Tschakert P, Ellis NR, Anderson C, Kelly A, Obeng J 2019 One thousand ways to experience loss: A systematic analysis of climate-related intangible harm from around the world. *Glob. Environ. Chang.* 55, 58–72. doi: [10.1016/j.gloenvcha.2018.11.006](https://doi.org/10.1016/j.gloenvcha.2018.11.006)
- Tubi A, Feitelson E 2016 Drought and cooperation in a conflict prone area: Bedouin herders and Jewish farmers in Israel's northern Negev, 1957–1963. *Polit. Geogr.* 51, 30–42. doi: [10.1016/j.polgeo.2015.11.009](https://doi.org/10.1016/j.polgeo.2015.11.009)
- Tucker J, Daoud M, Oates N, Few R, Conway D et al. 2015 Social vulnerability in three high-poverty climate change hot spots: What does the climate change literature tell us? *Reg. Environ. Chang.* 15, 783–800. doi: [10.1007/s10113-014-0741-6](https://doi.org/10.1007/s10113-014-0741-6)
- UNFCCC 2007 Climate Change: Impacts, Vulnerabilities and Adaptation in Developing Countries, Information Services of the UNFCCC secretariat. <https://unfccc.int/resource/docs/convkp/conveng.pdf>
- UNHRC 2008 Report of the Human Rights Council on its 7th session (A/HRC/7/78). <https://documents-dds-ny.un.org/doc/UNDOC/GEN/G08/146/63/PDF/G08146663.pdf?OpenElement>
- van Weezel S 2015 Economic shocks & civil conflict onset in Sub-Saharan Africa, 1981–2010. *Def. Peace Econ.* 26, 153–177. doi: [10.1080/10242694.2014.887489](https://doi.org/10.1080/10242694.2014.887489)
- von Uexkull N 2014 Sustained drought, vulnerability and civil conflict in Sub-Saharan Africa. *Polit. Geogr.* 43, 16–26. doi: [10.1016/j.polgeo.2014.10.003](https://doi.org/10.1016/j.polgeo.2014.10.003)
- von Uexkull N, Croicu M, Fjelde H, Buhaug H 2016 Civil conflict sensitivity to growing-season drought.



- Proc. Natl. Acad. Sci. U. S. A.* 113, 12391–12396.  
doi: [10.1073/pnas.1607542113](https://doi.org/10.1073/pnas.1607542113)
- Waha K, Krumpalauer L, Adams S, Aich V, Baarsch F et al. 2017 Climate change impacts in the Middle East and Northern Africa (MENA) region and their implications for vulnerable population groups. *Reg. Environ. Chang.* 17, 1623–1638.  
doi: [10.1007/s10113-017-1144-2](https://doi.org/10.1007/s10113-017-1144-2)
- Wallemacq P, Below R, McLean D 2018 Economic Losses, Poverty and Disasters 1998–2017.
- Weinthal E, Zawahri N, Sowers J 2015 Securitizing water, climate, and migration in Israel, Jordan, and Syria. *Int. Environ. Agreements Polit. Law Econ.* 15, 293–307. doi: [10.1007/s10784-015-9279-4](https://doi.org/10.1007/s10784-015-9279-4)
- Weiss H, Courty M-A, Wetterstrom W, Guichard F, Senior L et al. 1993 The Genesis and Collapse of Third Millennium North Mesopotamian Civilization. *Science (80-. )*. 261, 995–1004.  
doi: [10.1126/science.261.5124.995](https://doi.org/10.1126/science.261.5124.995)
- Werrell CE, Femia F, Sternberg T 2015 “Did We See It Coming?: State Fragility, Climate Vulnerability, and the Uprisings in Syria and Egypt.” *SAIS Rev. Int. Aff.* 35 (1), 29–46. doi: [10.1353/sais.2015.0002](https://doi.org/10.1353/sais.2015.0002)
- White S 2011 *The Climate of Rebellion in the Early Modern Ottoman Empire*. Cambridge University Press  
doi: [10.1017/cbo9780511844058](https://doi.org/10.1017/cbo9780511844058)
- WHO 2016 World Health Statistics 2016: Monitoring Health for the SDGs, sustainable development goals. [https://www.who.int/gho/publications/world\\_health\\_statistics/2016/en/](https://www.who.int/gho/publications/world_health_statistics/2016/en/)
- Wolf AT 2007 Shared Waters: Conflict and Cooperation. *Annu. Rev. Environ. Resour.* 32, 241–269.  
doi: [10.1146/annurev.energy.32.041006.101434](https://doi.org/10.1146/annurev.energy.32.041006.101434)
- Wood RM, Wright TM 2016 Responding to catastrophe: Repression dynamics following rapid-onset natural disasters. *J. Conflict Resolut.* 60, 1446–1472.  
doi: [10.1177/0022002715596366](https://doi.org/10.1177/0022002715596366)
- Xoplaki E, Fleitmann D, Luterbacher J, Wagner S, Hal-don JF et al. 2016 The Medieval Climate Anomaly and Byzantium: A review of the evidence on climatic fluctuations, economic performance and societal change. *Quat. Sci. Rev.* 136, 229–252.  
doi: [10.1016/j.quascirev.2015.10.004](https://doi.org/10.1016/j.quascirev.2015.10.004)



## Information about authors

### Coordinating Lead Authors

Vally (Vassiliki) Koubi:

*Center for Comparative and International Studies  
(CIS), ETH Zürich, Zürich, Switzerland*

### Lead Authors

Mohamed Behnassi:

*Center for Environment, Human Security and Governance,  
Ibn Zohr University of Agadir, Agadir, Morocco*

Antonietta Elia:

*University of Santiago de Compostela, Santiago de  
Compostela, Spain*

Manolis Grillakis:

*Institute of Mediterranean Studies (Institute of  
Technology & Research – Hellas), Rethymno, Crete,  
Greece*

Ethemcan Turhan:

*University of Groningen, Groningen, the Netherlands*



# MANAGING FUTURE RISKS AND BUILDING SOCIO-ECOLOGICAL RESILIENCE

**Coordinating Lead Authors:**

Athanasios T. Vafeidis (Germany/Greece)


**Lead Authors:**

Ameer A. Abdulla (Spain/Egypt), Alberte Bondeau (France), Lluís Brotons (Spain), Ralf Ludwig (Germany), Michelle Portman (Israel), Lena Reimann (Germany), Michalis Vousdoukas (Italy/Greece), Elena Xoplaki (Germany/Greece)

**Contributing Authors:**

Najet Aroua (Algeria), Lorine Behr (Germany), Francesco Dottori (Italy), Joaquim Garrabou (Spain), Christos Giannakopoulos (Greece), Guillaume Rohat (Switzerland/France), Elias Symeonakis (UK/Greece)

*This chapter should be cited as: Vafeidis AT, Abdulla AA, Bondeau A, Brotons L, Ludwig R, Portman M, Reimann L, Vousdoukas M, Xoplaki E 2020 Managing future risks and building socio-ecological resilience in the Mediterranean. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 539-588.*



# Table of contents

<b>6</b>	<b>Managing future risks and building socio-ecological resilience</b> .....	<b>543</b>
	Executive summary.....	543
<b>6.1</b>	<b>Introduction</b> .....	<b>543</b>
<b>6.2</b>	<b>Human health impacts</b> .....	<b>544</b>
6.2.1	Future health risks.....	544
6.2.2	Management approaches, governance, and adaptation for health risks.....	545
6.2.3	Case studies.....	546
6.2.4	Innovation.....	546
<b>6.3</b>	<b>Water security</b> .....	<b>547</b>
6.3.1	Future risks for water security.....	547
6.3.2	Management approaches, governance, and adaptation for water security.....	547
6.3.3	Case studies.....	548
6.3.4	Innovation.....	549
<b>6.4</b>	<b>Agricultural drought</b> .....	<b>550</b>
6.4.1	Future drought risks in agriculture.....	550
6.4.2	Management approaches, governance, and adaptation for agricultural drought.....	550
	<i>Adjusting irrigation water supply to satisfy water requirements</i> .....	550
	<i>Reducing water stress</i> .....	551
6.4.3	Case studies.....	552
6.4.4	Innovation.....	552
	<i>Mycorrhizal symbiosis</i> .....	552
	<i>Composting</i> .....	553
<b>6.5</b>	<b>Wildfires</b> .....	<b>553</b>
6.5.1	Future wildfire risks.....	553
6.5.2	Management approaches, governance, and adaptation for wildfires.....	554
6.5.3	Case studies.....	555
6.5.4	Innovation.....	555
<b>6.6</b>	<b>Soil erosion, degradation, and desertification</b> .....	<b>555</b>
6.6.1	Future risks for soils.....	555
6.6.2	Management approaches, governance, and adaptation for soil protection.....	556
6.6.3	Case studies.....	557
6.6.4	Innovation.....	557
<b>6.7</b>	<b>Heat waves</b> .....	<b>558</b>
6.7.1	Future heat wave risks.....	558
6.7.2	Management approaches, governance, and adaptation for heat wave risks.....	558
<b>6.8</b>	<b>River and pluvial flooding</b> .....	<b>559</b>
6.8.1	Future flood risk.....	559
6.8.2	Management approaches, governance, and adaptation for flood protection.....	559
6.8.3	Case studies and innovation.....	560

<b>6.9</b>	<b>Sea-level rise: coastal erosion and flooding, saltwater intrusion</b> .....	<b>560</b>
6.9.1	Future risk associated with sea-level rise.....	560
6.9.2	Management approaches, governance, and adaptation for coastal protection.....	561
6.9.3	Case studies.....	562
6.9.4	Innovation.....	562
<b>6.10</b>	<b>Seawater temperature anomalies and extremes</b> .....	<b>563</b>
6.10.1	Future risk of marine heat waves.....	563
6.10.2	Management approaches, governance, and adaptation for ocean warming.....	564
<b>6.11</b>	<b>Ocean acidification</b> .....	<b>564</b>
6.11.1	Future risk of ocean acidification.....	565
6.11.2	Management approaches, governance, and adaptation for ocean acidification.....	565
<b>6.12</b>	<b>Non-indigenous species: marine, freshwater, and terrestrial</b> .....	<b>566</b>
6.12.1	Future risks associated with non-indigenous species.....	566
6.12.2	Management approaches, governance, and adaptation for non-indigenous species.....	567
6.12.3	Innovation.....	568
<b>6.13</b>	<b>Interactions of hazards, synergies and trade-offs between adaptation strategies and mitigation</b> .....	<b>568</b>
	<b>References</b> .....	<b>572</b>
	<b>Information about authors</b> .....	<b>588</b>

## 6 Managing future risks and building socio-ecological resilience

### Executive summary

The Mediterranean Basin is experiencing major changes in environmental conditions, which can introduce new challenges to the resilience of its natural and human systems. This situation is combined with rapid and spatially diverse socio-economic development in the region, mainly in terms of demographic trends and settlement patterns, thus leading to higher exposure to environmental hazards. Furthermore, new risks are expected to emerge from interactions between drivers and impacts across sectors, thus increasing the vulnerability of natural systems and human populations.

Future risks in the Mediterranean region will be determined by hazard characteristics (intensity and frequency) and by developments in socio-economic conditions that determine a society's adaptive capacity to cope with these hazards. Current risks to human population, economies and ecosystems will increase as a result of changes in the patterns of droughts, wildfires, soil degradation, desertification, sea level rise, heat waves and river flooding, and other pressures, potentially leading to greater impacts. These impacts can be further exacerbated by the occurrence of compound and cumulative events, which can seriously challenge the adaptive capacity and resilience of biophysical and human subsystems. Coping with these risks, adapting to change and increasing the resilience of Mediterranean systems will be essential for ensuring sustainable development in the region.

Successful practices and initiatives for risk reduction and management, such as water-sensitive urban design, implementation of nature-based solutions, operational flood forecasting systems, or collaborations within cities' networks, are already being implemented across the region. However, these efforts are often slow in catching on or fail to consider the rising pressures in the light of

changing environmental conditions and developmental demands. Understanding these changes and demands is essential for managing future risks. In this context, Mediterranean-wide initiatives such as establishing long-term monitoring schemes to obtain data missing in many parts of the basin; accounting for differences in monitoring and reporting between northern (EU), eastern, and southern countries of the region; advancing (climate) modeling techniques for the short-term prediction of extreme events (e.g., heat, flooding), and improvements in seasonal forecasts are essential for supporting future management and adaptation strategies and for enhancing resilience. Furthermore, public participation in the development and implementation of these strategies is necessary in order to increase local relevance and acceptance of proposed strategies, and is particularly important for building a resilient society.

The level of future risk in the Mediterranean Basin will largely depend on the timing of adaptation and on how soon and how effectively sustainable development is pursued. In particular, addressing more pressing natural and socio-economic challenges in several countries in the Middle East and North Africa is essential for avoiding the widening of the development gap between northern, southern, and eastern countries of the region. Therefore, developing joint, region-wide, and integrated management and adaptation approaches that treat multiple hazards in a holistic manner is of utmost importance for sustainable development in the entire region. Nonetheless, no one-size-fits-all strategy exists, but each measure needs to be tailored to the respective local conditions. Regional co-operations, e.g. in the form of active participation in regional-to-global initiatives and networks concerned with building socio-ecological resilience, will be an important step forward in transferring knowledge on successful practices and innovation across the Mediterranean nations.

### 6.1 Introduction

Scenarios of environmental and socio-economic change for this century suggest severe challenges to the resilience of natural and human systems worldwide. For climate, such challenges will be particularly posed by extreme events, such as increased temperature anomalies (*Section 2.2.4*)

and potential changes in storm intensity (*Section 2.2.3*), as well as by slow onset events such as sea level rise (*Section 2.2.8*). From a management and policy perspective, this means that these changes increase the vulnerability of certain groups that are natural resource-dependent and increase the

need to enhance the resilience of ecosystems and human systems. Finally, they will also increase the need for efforts to reduce local stressors and identify adaptation options.

The Mediterranean Basin is experiencing major changes in environmental conditions, combined with rapid and spatially diverse socio-economic development. These factors combined are exerting increased pressure on natural systems and human societies in the region. At the same time, new risks may emerge from interactions between drivers (Section 2.6) and from the interactions of impacts across sectors (Cramer et al. 2018), which may result in greater impacts, while increasing the vulnerability of less resilient natural systems and populations. These risks can affect the provision of services from natural systems and lead to severe disruptions in social systems.

Chapter 6 deals with managing future risks, identifying adaptation options and building capacity for resilience to climate and environmental changes. Addressing this aim, the chapter discusses three key components of emerging policy needs in the basin. The first component is the current understanding on the trajectory, intensity and spatial extent of future risk for the principal hazards, and

associated policy considerations of the region. Secondly, the chapter outlines the current management and adaptation approaches and prevalent governance frameworks for coping with these risks. The third component critically reviews a range of examples of adaptation and mitigation for sectoral approaches, while considering case studies from Mediterranean-type environments.

Chapter 6 identifies a number of innovative and successful practices for achieving sound and sustainable development in countries of the Mediterranean Basin. Successful adaptation case studies involve stakeholder participation, structural political and economic change, gender considerations and weather-indexed insurance schemes. Successful mitigation involves options with clear societal benefits, such as energy cooperatives, energy efficiency, or regional cooperation. The final part of this section discusses potential interactions between hazards and sectors, which may lead to increased impacts. It further includes proposals to improve synergies between adaptation and mitigation practices and suggestions to promote Mediterranean cooperation and networking for building resilience, while also focusing on education and capacity-building.

## 6.2 Human health impacts

### 6.2.1 Future health risks

Environmental change can lead to a wide range of impacts on human health in Mediterranean countries (Sections 5.2.3 and 5.2.4). The most well-known impacts are direct impacts, e.g., extreme temperatures, cold and heat waves leading to cardiovascular and respiratory diseases and death (Gasparrini et al. 2017), wildfires leading to lethal injuries and respiratory diseases (Reid et al. 2016), and direct physical injuries and deaths resulting from extreme weather events, such as intense rainfall, river flooding, and storms (Forzieri et al. 2017). Impacts on human health can also be indirect, e.g., climate-related changes in food availability and quality that threaten food security (Deryng et al. 2016), increased variability of rainfall patterns that jeopardizes the availability and quality of freshwater (Koutroulis et al. 2016; Flörke et al. 2018), worsened air quality causing respiratory illnesses (de Sario et al. 2013; Doherty et al. 2017), and climate-driven increase in vector-borne diseases (Negev et al. 2015). The extent to which environmental change

affects human health largely depends on the vulnerability of the exposed populations, that is, their ability to face and cope with climate-related hazards (IPCC 2012). Just as, for example, climate change is altering the climate system, socio-economic development and demographic growth are shaping the future vulnerability of populations in the Mediterranean Basin, with contrasting trends depending on the type of socio-economic trajectory (O'Neill et al. 2017; Reimann et al. 2018a; Kok et al. 2019). Urban areas along the Mediterranean coast are especially affected by climate change impacts on health because these areas concentrate people and assets (Watts et al. 2015). Urban areas often intensify climate-related hazards, e.g., hotter temperatures during heat waves due to the urban heat island effect (Papalexiou et al. 2018) and increase in run-off and flooding during extreme precipitation events due to soil sealing (Romero Diaz et al. 2017).

Heat-related morbidity and mortality are projected to increase substantially throughout the Mediterranean countries, under all climate scenarios (Sec-



tion 5.2.5.2). Impacts are expected to be the greatest in urban areas where people are concentrated and where the urban heat islands lead to higher inner-city temperatures (Yang et al. 2016). Future heat-related health risks are well-documented, with a large number of case studies spread across the Mediterranean Basin. Examples include (but are not limited to) a ~3- to 9-fold increase in the heat-related mortality rate in Cyprus (Heaviside et al. 2016); a 50-fold increase in mortality (compared to the current situation) on average across southern Europe by the end of the century (Forzieri et al. 2017) and a substantial difference in the increase of mortality between RCP2.6/RCP4.5 and RCP8.5 scenarios (Gasparrini et al. 2017; Kendrovski et al. 2017) (Section 5.2.5.2). In contrast, cold-related mortality is projected to decrease under all scenarios (Forzieri et al. 2017) (Section 5.2.5.3). It is also worth noting that changes in socio-economic and demographic conditions such as urbanization, demographic growth, and aging are also expected to further increase the burden of heat-related morbidity and mortality in Mediterranean countries (Rohat et al. 2019b, 2019a).

In contrast to other parts of the world, climate change is expected to lead to an overall increase in ground-level ozone- and fine particulate matter-related mortality in Mediterranean countries (Silva et al. 2016), with the exception of high-end climate change scenario RCP8.5, which leads to an increase in the health burden of air pollution in most Mediterranean countries (Silva et al. 2017). However, the significant uncertainties that exist in the trend directions and in risk estimates, that are primarily linked to the uncertainty in future types of emissions must be noted (Doherty et al. 2017).

Temperature rise will expand the habitat suitability for vectors, such as mosquitoes and sandflies (Negev et al. 2015; Semenza and Suk 2018; Hertig 2019) to most of the Mediterranean Basin by the end of the century and increase the transmission risk of the diseases they can carry, such as dengue, West Nile Fever and leishmaniasis (Bouzig et al. 2014; Semenza et al. 2016; Liu-Helmersson et al. 2019) (Section 5.2.5.4). One exception is the reduction of habitat suitability for *Aedes albopictus* in the southernmost parts of Europe (Caminade et al. 2012; Proestos et al. 2015), leading to a reduction of climatically suitable areas for the transmission of Chikungunya (Fischer et al. 2013). Changes to the hydrological cycle caused by climate changes are expected to further amplify such health issues and lead to increased fatalities. Erratic precipitation and extreme events and floods could support the flourishing of bacteria, parasites and algal blooms,

including the protozoan parasites *Cryptosporidium*, hepatitis A viruses, *Escherichia coli* bacteria, and more than 100 other pathogens. The increase in human mobility also plays a crucial role in spreading vector-borne diseases throughout the Mediterranean Basin in newly suitable habitats (Thomas et al. 2014; Roche et al. 2015; Hertig 2019; Kraemer et al. 2019).

The combination of longer fire seasons and more frequent, large, and severe fires – triggered by increased droughts and land-use change (Turco et al. 2014; Knorr et al. 2016) – is projected to lead to greater fire risk and casualties, particularly in sub-urban areas (Forzieri et al. 2017) (Section 2.6.3.3). Similarly, more intense and frequent extreme precipitation events are expected to trigger a strong increase in flash flood-related injuries and mortalities throughout Mediterranean countries (Gaume et al. 2016) (Section 3.1.4.1). Floods can further damage water infrastructure, contaminate freshwater supplies, heighten the risk of water-borne diseases, and create breeding grounds for disease-carrying insects, especially threatening those with already limited access to water and sanitation (WHO 2017). The combination of demographic growth and changing diets is expected to lead to higher food demand across the region (Paciello 2015), while changes in extreme events such as droughts, heat waves, and extreme precipitation are projected to decrease crop and livestock yields substantially (Bernués et al. 2011; Deryng et al. 2016) (Section 3.2.2.1). This is particularly the case in the southern part of the Mediterranean Basin.

## 6.2.2 Management approaches, governance, and adaptation for health risks

National adaptation policies have been adopted in a large number of Mediterranean countries, often covering and acting on large-scale health topics such as extreme heat, air pollution, and vector-borne diseases (Negev et al. 2015). Although national governments have an important role to play in reducing the burden of climate change on human health, it is at the local scale that most actions and measures are taken (Paz et al. 2016). In fact, cities and municipalities in the Mediterranean Basin are at the forefront of climate change adaptation, particularly with regard to climate change impacts on human health, and often drive the regional effort to better anticipate and prepare for the adverse effects of climate change on human health and well-being (Reckien et al. 2018).

City-level adaptation is, more often than not, preferred to national-scale adaptation to decrease the

vulnerability of the local population. This is accomplished through measures that include (but are not limited to) the improvement of housing and infrastructure, the education and awareness-raising of the most vulnerable communities, the implementation of early warning systems, the strengthening of local emergency and healthcare services, and the general strengthening of the community's and local institutions' adaptive capacity (Larsen 2015; Liotta et al. 2018). City-level adaptation can also directly target the reduction of climate-related hazards, such as building multi-usage buffer zones to reduce flood risk and to decrease the urban heat island (Yang et al. 2016).

Interestingly, management approaches sometimes attempt to develop adaptation measures that also affect climate change mitigation and/or that trigger health benefits, such as using green roofs to retrofit existing buildings (Gagliano et al. 2016) and transforming the transportation systems to mitigate emissions and reduce air pollution (WHO Europe 2017).

### 6.2.3 Case studies

It is important to note that most adaptation actions are impact-, context- and place-specific and there is no one-size-fits-all adaptation measure to reduce climate impacts on human health. Adaptation measures can take a wide range of forms, be triggered by different events, operate on different spatial and temporal scales, and be associated with different implementation constraints (Fernandez Milan and Creutzig 2015; Holman et al. 2018).

A number of Mediterranean cities have developed adaptation plans that specifically target the reduction of human health impacts. A significant part of the actions depicted in such climate adaptation plans are broad and unspecific (Reckien et al. 2018), which can constitute a bad practice and often do not mention potential negative effects, such as the increase in inner-city temperature and air pollution due to the systematic installation of air conditioning (Salamanca et al. 2014). Certain adaptation plans depict context-specific and quantified actions, such as in the city of Barcelona, which for instance, plans to increase its urban green areas by 1 m<sup>2</sup> per city resident by 2030 in order to decrease the urban heat island in case of extreme heat and increase water infiltration in the event of flash flooding (Barcelona Sostenible 2015).

<sup>40</sup> <https://www.medilabsecure.com/home.html>

<sup>41</sup> <https://www.rockefellerfoundation.org/100-resilient-cities/>

<sup>42</sup> <https://www.c40.org/>

For regional climate-related hazards, such as vector-borne diseases, a multi-country and trans-boundary approach to adaptation is crucial (Negev et al. 2015) and has been implemented over the past decades. In its current form, the MediLabSecure project<sup>40</sup> covers all Mediterranean countries and aims at preventing vector-borne diseases in these countries through scientific research and concrete actions.

### 6.2.4 Innovation

Climate change vulnerability assessments with a strong focus on human health have been undertaken over the past few years for cities without dedicated adaptation plans, including case studies for Cairo (Katzan and Owsianowski 2017), Nicosia (Kaimaki et al. 2014), and Antalya (Antalya Metropolitan Municipality 2018). These studies provide a strong scientific basis for the design of context-specific adaptation plans in the years to come. Collaboration within networks of cities with the goal to act on climate change (including adaptation to human health impacts) is promising in terms of identifying and sharing knowledge on best practices and concrete actions (Román and Midttun 2010; Rosenzweig et al. 2018). For example, cities such as Tel Aviv, Rome, Thessaloniki, Ramallah and Byblos are members of the "100 Resilient Cities"<sup>41</sup> network, Venice is a member of the "C40 Cities"<sup>42</sup> network and its program for connecting delta cities, and numerous cities are members of the Global Covenant of Mayors for Climate and Energy.

The integration of climate adaptation and mitigation plans within a unique climate plan is rarely achieved, but appears to be an efficient way to design measures that benefit both adaptation and mitigation (Reckien et al. 2018). The city of Athens has recently entered the circle of cities to have done so, with results on the reduction of human health impacts expected to come in the next few years (City of Athens 2017).

## 6.3 Water security

The Sustainable Water Partnership (SWP)<sup>43</sup> defines water security as “the adaptive capacity to safeguard the sustainable availability of, access to, and safe use of an adequate, reliable and resilient quantity and quality of water for health, livelihoods, ecosystems and productive economies”. This embracing definition reveals the pivotal role water security plays on all levels for reaching the ambitious goals laid out by the UN’s Sustainable Development Goals (UN 2015; Bhaduri et al. 2016). Diametric to water security is water scarcity, a state reached when water demand can no longer be satisfied due to a lack of freshwater resources (Srinivasan et al. 2012). Physical water scarcity results in the depletion of water resources for both humans and natural systems and causes important transitions in the exploitation of different water compartments, e.g., from surface to groundwater sources, or even water transfers between basins. When excessive human consumption of water resources occurs under these circumstances, it may cause significant pressures on aquifers and surface waters, producing adverse effects on water quantity (over-exploitation) and quality (nutrient excess, pollution and lower biodiversity), which is detrimental to economic development and even compromises human health.

### 6.3.1 Future risks for water security

The Mediterranean Basin is particularly prone to limited water security due to its semi-arid to arid climates, especially as most important economies, such as tourism development (Section 3.1.2.3) and intensive agriculture (Section 3.1.2.2) are heavily water dependent and critically vulnerable to water scarcity and stress (Barceló and Sabater 2010). Thus, water security is at severe risk in the Mediterranean Basin. This susceptibility to scarcity is caused by strong human pressures, under the form of overexploitation, for agricultural, urban and industrial water uses, together with reduced availability of water due to climate change. Many Mediterranean water bodies, aquifer systems in particular (Section 3.1.3.4), show over-exploitation associated with high seasonal water demand, and suffer from salinization, particularly in coastal areas and regions of intense irrigation and soil degradation. High human water demands in the region concentrate when water availability is at the lowest and exhausted aquifers co-occur with transformation of watercourses from permanent

into intermittent. An increasingly common scenario in river basins includes headwaters becoming intermittent or even ephemeral, while lowlands bear aquifers that are depleted or contaminated by either salt or pollutants (Choukr-Allah et al. 2017). Growing human demands for water are leading to rapid increases in the frequency and severity of water scarcity, where there is insufficient water to simultaneously support both human and ecosystem water needs (Bond et al. 2019). With climate change and increasing demand for food and commercial services due to a growing population with higher demands, such patterns will very likely increase dramatically (Iglesias et al. 2012).

### 6.3.2 Management approaches, governance, and adaptation for water security

Observed trends and projections for the future indicate a strong susceptibility to changes in hydrological regimes, an increasing general shortage of water resources and consequent threats to water availability and management (Section 3.1.1.1). However, it must be clearly stated that current uncertainties in climate projections and subsequent impact models, a yet incomplete understanding of the impact of a climate change signal on economic mechanisms or the lack of an elaborate and integrated human security conceptual framework, are imposing strong limitations on water-related decision-making under climate change conditions (Section 3.1.5).

Climate, demographic and economic changes are expected to have strong impacts on the management of water resources, as well as on key strategic sectors of regional economies and their macroeconomic implications (Section 3.1.5). Such developments bare the capacity to exacerbate tensions, and even intra- and inter-state conflicts among social, political, ecological and economic actors. Meanwhile, it is widely agreed that effective adaptation and prevention measures need multi-disciplinary preparation, analysis, action and promotion of collaborative strategies.

The complexity of the water cycle contrasts strongly with the low data availability, which (a) limits the number of analysis techniques and methods available to researchers, (b) limits the accuracy of models and predictions, and (c) consequently

<sup>43</sup> <https://www.swpwater.org/>

challenges the capabilities to develop appropriate management measures to mitigate or adapt the environment to scarcity and drought conditions. The current potential to develop appropriate regional adaptation measures to climate change impacts suffers heavily from large uncertainties. These spread along a long chain of components, starting from the definition of emission scenarios to global and regional climate modeling to impact models and a subsequent variety of management options. Furthermore, the lack of awareness or understanding of the complex climate-resource-society dynamics often leads to inappropriate measures or no measures at all. Integrated water resources management is a holistic approach that focuses on both environmental as well as on socio-economic factors influencing water availability and supply, and seeks an efficient blend of all available conventional and unconventional water resources to meet the demands of the full range of water users, especially in agriculture, industry and tourism. However, the management approaches and solutions adopted, e.g., in form of decision support for specific water resources systems, are often highly specific for individual case studies (Section 3.1.5).

An inventory of international, national and regional policies dealing with responses to climate change, water resources management, responses to hazards and disasters, and security in the region, is essential for proposing a suitable policy framework to integrate security, climate change adaptation and water management issues and specific recommendations for policy streamlining at the UN, EU, national and regional levels.

Political, economic and social factors seem to be more important drivers of water-related conflicts than climate-related variables (Section 5.3.3.2). States and state-led adaptation play a prominent role in affecting human security: states can greatly facilitate adaptation, but policies are also prone to adverse effects. Adaptation can both increase and diminish water security for certain groups, although this depends to a great extent on factors such as power relations, marginalization and governance. There are also varying capacities of states to implement effective adaptation policies. Analyzing the political economy in an area or country helps to understand state-led adaptation. Impacts on key strategic sectors typically consider agriculture and tourism. These sectors show specific dependence on water security, which is of quintessential importance in the Mediterranean economy, with relatively high adaptation potential to strategic policies.

Most Mediterranean countries will likely face water shortages (Section 3.1.1.1). This can have significant implications in terms of agricultural productivity, income and welfare. However, the water gap in the Mediterranean area will be affected by different external drivers. In northern Mediterranean countries, this will be due to increased temperature and decreased precipitation. Middle East and North Africa economies will likely find it difficult to put aside precious water resources for the purpose of environmental preservation. In southern Mediterranean countries, the growing non-agricultural water needs (induced by strong economic and demographic development) will be an additional challenge to water security, demanding management improvements in water efficiency. Innovations include highly successful efforts to increase water use efficiency. Smart metering, for example, is being deployed to improve accuracy in billing, evaluate consumption and increase users' awareness of their own consumption (Revolve Water 2017).

### 6.3.3 Case studies

Due to the already high and expected increasing pressure on water resources in the Mediterranean, the efforts to counteract water scarcity and establish water security are manifold in scope, action and scale. As the challenges can be very site-specific, and triggered by both natural and anthropogenic drivers in various constellations, significant uncertainties remain in identifying suitable programs of measures, which would be generally applicable for being independent of region and scale. Thus, related activities can be embedded anywhere from pan-continental to national levels, but often basin-scale and even highly localized programs and case studies are implemented. The range of measures (Section 3.1.5) includes water-saving technologies, such as new equipment in irrigation agriculture and households, often complemented by improved water efficiency (e.g., by means of adapted water management procedures), as well as direct measures to increase water availability through additional multi-scale storage solutions (ranging from cisterns to large reservoirs) or through the use of unconventional water sources stemming from recharging wastewater or seawater desalination. The latter may however, cause environmental concerns due to soil contamination or energy consumption (IWA 2012).

All these aspects can be useful components of an integrated water resources management approach (Choukr-Allah et al. 2012) (Section 3.1.5.1). To date, there are several highly successful examples of

such an approach, but negative case studies also exist. This highlights the prevailing need for further research and transdisciplinary collaboration to reach and maintain water security in the Mediterranean (Ludwig et al. 2011; Ludwig and Roson 2016; Saladini et al. 2018).

Several success stories of water security measures related to wastewater re-use experiments on local scale applications exist. The case of Oueljet El Khoder, Tunisia, is an exemplary effort, which succeeded in establishing a sound system for water re-use to provide reliable water resources for irrigation and ensuring sustainable conditions for the underlying aquifer. In this case, the collaborative project SWIM Sustain Water MED<sup>44</sup> has introduced a tertiary treatment unit including a slow sand filter alongside a monitoring and early warning system for monitoring the quality of the treated wastewater. The installations resulted in an increased rate of re-use of reclaimed water and an extension of the agricultural irrigation area.

A main challenge, however, is the fact that despite the evidence of water scarcity being felt by stakeholders and end-users, the role of climate change and the related future exacerbation of water stress is often ignored and not perceived as a key issue for water uses and water security (La Jeunesse et al. 2015). In the course of the CLIMB project<sup>45</sup>, several circum-Mediterranean case studies (e.g., France, Italy, Turkey, Egypt and Tunisia) showed that the main response to increasing water demand in the Mediterranean region is a progressive externalization of water resources, with limits imposed by national borders and technological possibilities. This thinking, which does not consider climate change as a driving force, is not sustainable and prone to rising water conflicts.

### 6.3.4 Innovation

In recent years, great energy and investment has been placed in the modernization of installations and development of (sometimes integrated) water resources management (Section 3.1.5) (Cameira and Santos Pereira 2019). However, in many cases, these efforts seek to adapt to current state challenges and fail to consider the rising pressures in the light of climate change and growing domestic and industrial water demand. One of the expected consequences of climate change alone is a reduction in annual precipitation, paired with a very likely increase in rainfall variability and extremes

(Section 2.2.5). All of these factors contribute to increasing vulnerability and risk in potentially affected regions and can consequently jeopardize water security. Innovation is needed to reach beyond the current limitations of water resources management by introducing flexible mechanisms that not only include novel water (saving) technologies, but also build on targeted water system analysis and research (Section 3.1.5.2). Important elements of these types of systems start with the (re-)establishment of environmental monitoring networks, composed of a dense in-situ observational network paired with operational remote sensing applications (e.g., for spatial drought monitoring, including vegetation status, soil moisture, water levels). Based on such regular time series of data products, spatially-explicit and process-based models can be built with sufficient predictive power to support long-term planning and decision-making to adapt to the impacts of a gradual climate and global change.

Great innovation potential lies in the development of regionally specified and flexible response schemes to water scarcity that reach beyond the state-of-art and provide integrated solutions for increased water efficiency by combining improved water-saving technologies (Wang and Polcher 2019) with the provision of unconventional water resources (e.g., by managed aquifer recharge or saline water for irrigation (Reca et al. 2018; Tzoraki et al. 2018)), to avoid water stress (Section 3.1.5.2). It is further necessary to establish systems for short-term predictions of extreme events and seasonal forecasts that allow for extended reaction time of first responders (Haro-Monteagudo et al. 2017; Corral et al. 2019) and affected industries, such as agriculture (Martínez-Fernández et al. 2013; Kourgialas et al. 2019) or tourism (Hadjikakou et al. 2013; Toth et al. 2018). Water-sensitive urban design (WSUD) is approach to management that is starting to take hold in cities, although slow to catch on in the Mediterranean (Goulden et al. 2018). This paradigm is fueled by the interest in sustainable urban development and it aims to integrate best water management practices (many related to stormwater runoff), with mechanisms of urban planning. WSUD, developed in Australia, connects urban planning with stormwater management mainly for protecting groundwater in aquifers. In the United States, planners employ a similar approach, called low-impact development (LID), which focuses on maintaining a steady hydrological response (i.e., stormwater runoff vol-

<sup>44</sup> <https://www.swim-h2020.eu/>

<sup>45</sup> <https://cordis.europa.eu/project/id/244151>

ume and discharge rate leaving the spatial unit before and after development), but also seeks to view stormwater as a benefit to the environment, rather than only as a disturbance (Carlson et al. 2015). While both LID and WSUD aim to minimize the hydrological effects of urban development on the surrounding environment, WSUD puts more emphasis on maintaining a water balance that considers waterway erosion along with the management of groundwater, stream flows, and flood damage. In Israel, this approach has been considered and implemented, but mostly on a local, site-specific basis, through such practices as retention pools, but it has been less successfully

implemented to curb such problems as increased coastal erosion (Portman 2018).

In order to have practical impact, a crucial element in this endeavor is to fully take into consideration the political and institutional dimensions of dynamically changing priorities in water governance. This can be supported by novel ways of public participation and knowledge sharing between institutions and researchers (Bielsa and Cazcarro 2014), which in combination could and should lead to the development of smart water grids and efficient water licensing and metering.

## 6.4 Agricultural drought

### 6.4.1 Future drought risks in agriculture

Agricultural drought occurs when soil moisture availability to plants has dropped to such a level that it adversely affects crop and pasture growth (Mannochi et al. 2004). The Mediterranean region stands out as one meteorological drought hotspot where drought severity has increased in recent decades (Spinoni et al. 2019) (Section 2.2.5). Regarding agriculture, climate warming exacerbates the impact of meteorological droughts through the increasing evaporative demand (Quintana-Seguí et al. 2016). The analysis of climate model ensembles in the Mediterranean consistently projects future meteorological droughts that translate into stronger soil moisture anomalies (Planton et al. 2012; Orłowsky and Seneviratne 2013; Dubrovský et al. 2014; Ruosteenoja et al. 2018) (Section 3.1.4.1). More recently, Rojas et al. (2019) showed that climate models project negative precipitation trends outside the natural variability in the Mediterranean region in the mid-century, in all RCPs. A 10 to 30% decrease in precipitation is expected as early as 2040, in particular causing drier winters in northern Africa, and summer drying in southern Europe.

Already under “low” global warming levels of 1.5°C and 2°C, the exacerbation of drought conditions in the Mediterranean will be unprecedented since the last millennium (Guiot and Cramer 2016; Samaniego et al. 2018) (Section 3.1.4.1). Furthermore, as Mediterranean drought events also imply hot summers (Zampieri et al. 2009; Hirschi et al. 2011; Russo et al. 2018), they drive a positive feedback that again enhances the frequency and the severity of agriculture droughts, directly challenging both

crop and pasture management (e.g., Saadi et al. 2015; Scocco et al. 2016). Both rain-fed agriculture and irrigated agriculture are vulnerable to drought (García-Garizábal et al. 2014), because the availability of irrigation water may become limited by several factors, including depletion of overexploited groundwater (Famiglietti 2014), competition for water due to the expansion of irrigated agriculture (Khadra and Sagardoy 2019) or conflict with other water usages (e.g., Gössling et al. 2012) (Section 3.2.2.1).

### 6.4.2 Management approaches, governance, and adaptation for agricultural drought

Farmers, who have been historically exposed to variable climate conditions, such as in the Mediterranean region, tend to be more prepared to cope with climate change (Reidsma et al. 2009). When it comes to droughts, several options are considered for avoiding water-stress in crops/pastures. Two main strategies can be identified: either ensuring that the water requirements are fulfilled (e.g., Fader et al. 2016), or requiring less water by modifying the agricultural system and its management so that the crops/pastures can better endure drought (Section 3.2.3.1).

#### *Adjusting irrigation water supply to satisfy water requirements*

Rapid solutions for satisfying increasing water requirements, such as expanding irrigated areas or increasing groundwater and/or reservoir pumping, only have short-term effects and are often not sustainable when they lead to decreased ground-

water levels, as reported from many regions (see Richey 2014 for the groundwater depletion of the major aquifers of the MENA region) or when only limited surface reservoirs are available (*Section 3.1.2.2*). This affects all competitive water users, including the environment, e.g., the wetlands in the Upper Guadiana Basin in Spain (Carmona et al. 2011). Other solutions include the deployment of improved irrigation and conveyance systems, which have a large water-saving potential (e.g., sprinkler or drip). In recent years, governments of a few Mediterranean countries have subsidized pressurized irrigation systems (Daccache et al. 2014). According to the study by Fader et al. (2016), the Mediterranean region could save up to 35% of water by implementing such irrigation techniques (*Section 3.1.5.2*). Nevertheless, these techniques alone are insufficient to face the increasing water demand resulting from climate change, demography, and socio-economic change (Malek and Verburg 2018).

Increasing attention is being given to wastewater reclamation and re-use, with important projects developed in countries all over the Mediterranean Basin since the end of the 20th century (Angelakis et al. 1999). From different experiments, it appears that treated wastewater re-use in integrated water resources management systems may provide significant benefits for irrigated agriculture and could be implemented in most water-scarce regions (Kalavrouziotis et al. 2015) (*Section 3.1.5.2*). Even the use of poorly controlled treated wastewater does not damage the agronomic quality of soils. It even increases the soil organic matter (Cherfouh et al. 2018). Consequently, it is possible to expect both potential agronomical benefits and improved water supply from wastewater management.

Desalination of seawater for irrigation has high costs and many negative environmental impacts (Sadhvani et al. 2005). Furthermore high-level desalination removes ions that are essential for plant growth (Yermiyahu et al. 2007). The above studies concluded that desalination facilities for irrigation need revised treatment standards. An alternative strategy looks at crop performance under deficit irrigation. Promising results indicate an enhancement of water productivity, leading to proportionally lower yield reduction than water deficit. Furthermore, in the case of tomatoes, fruit quality is improved (Patanè et al. 2011).

### **Reducing water stress**

The development of intensive agriculture since the second half of the 20th century has changed soil

properties in several ways, including change of structure, decrease in soil organic matter, and decrease in biological activity (e.g., García-Orenes et al. 2012; Aguilera et al. 2013; Morugán-Coronado et al. 2015). In addition, many arable soils with cereals are left bare for extended periods (Kosmas et al. 1997), and bare soil beneath the rows is also a frequent feature of industrial perennial crops (Gómez et al. 2011). Both aspects impact the soil hydrological cycle, i.e., the water resources for crops/pastures. First, besides increasing erosion (which has reached dramatic levels in some Mediterranean cropping areas), bare soils favor evaporation and intercept precipitation water less well than vegetated or mulched soil (Monteiro and Lopes 2007). Second, low organic matter content, tillage practices, and the decline of biological activities all imply soil structure changes such as porosity (Pagliai et al. 2004) (*Section 3.2.3.2*). In particular, the micropore to macropore ratio is modified: the proportion of micropores, which are considered the most important both in soil-water-plant relationships and in maintaining a good soil structure (Pagliai et al. 1983), is decreased by tillage, with a significantly reduced capacity to store water (Lampurlanés et al. 2016). Since the end of the 20th century, these phenomena have been studied in the Mediterranean Basin, using experiments on the effects of conservation agriculture (no tillage / reduced tillage, cover crops / mulching) on the soil-water dynamics for most of the Mediterranean cropping systems. These strategies are particularly promising in dry areas, and their average effect on Mediterranean agro-ecosystems, including yields, is beneficial, especially during water-stressed periods, despite the existence of contradictory results that may occur for many reasons (Mrabet 2002; Álvaro-Fuentes et al. 2007, 2008; Mrabet et al. 2012; Tomaz et al. 2017) (*Section 3.2.3*).

These adaptation strategies also have benefits for climate mitigation, since conservation agriculture emits less greenhouse gases and generally leads to soil carbon sequestration (Kassam et al. 2012; Aguilera et al. 2013; García-Tejero et al. 2020) (*Sections 3.2.3.2 and 3.2.3.3*). The net effect of this is still debated and clearly depends on other factors as well (Govaerts et al. 2009). There is also a finite time horizon as the agro-ecosystem soil carbon maximum capacity is often reached after a 20-50-year period (Lal and Bruce 1999). In any case, incentives from different institutions now exist in several countries in order to promote agricultural management strategies that rely on key principles of conservation agriculture (Calatrava et al. 2011). Water stress can also be reduced if several crops (or crops and flower/grass strips) are grown in

combination on the same land, which may result in a deeper penetration of the roots into the soil. Such plasticity has been observed in vineyards with cover cropping, where the compensatory growth of the grapevine root system allowed the resources (e.g., water) of deeper soil layers to be exploited (Celette et al. 2008). In an agroforestry system mixing walnut trees and winter crops, the competition with the winter crops induces deeper rooting of the walnut trees (Cardinael et al. 2015). Besides revealing the adaptive capacity of plants, these agroforestry practices provide welcomed shade in summer in the Mediterranean Basin, which is beneficial for both crops and livestock (Sá-Sousa 2014). Agroforestry systems are multifunctional, currently re-discovered in temperate areas – only for the montado-dehesa system of the Iberian Peninsula, a savanna-like rangeland dominated by scattered Mediterranean evergreen oak trees, the positive role of the trees on the water balance has been shown since the 20th century (Joffre and Rambal 1993, 2006).

### 6.4.3 Case studies

Many studies on no tillage in the Mediterranean show that this practice has positive effects on the soil for keeping more water, therefore enhancing


yields, especially in water-stressed years. A few studies on the similar positive effects of agroforestry are shown in Fig. 6.1.

### 6.4.4 Innovation

#### *Mycorrhizal symbiosis*

In the Mediterranean Basin, the alleviation of drought stress by mycorrhizal symbiosis has been studied for more than 25 years (Sánchez-Díaz and Honrubia 1994). Using soils of different Mediterranean locations, controlled experiments regularly report the beneficial effect of arbuscular mycorrhizal symbiosis for crops in drought conditions: Meddich et al. (2000) for clover, Ruiz-Lozano et al. (2001) for soybean, Marulanda et al. (2007) for lavender, Navarro García et al. (2011) for cane-apple bush, Armada et al. (2015) for maize (using also drought-adapted autochthonous microorganisms), and Calvo-Polanco et al. (2016) for olive among others. Field experiments confirm that mycorrhizal inoculation alleviates water deficit impact (e.g., in Hungary, Bakr et al. 2018), but we are not aware of such field studies in the Mediterranean.

Considerable progress has been made in understanding the role of arbuscular mycorrhizal sym-



**Figure 6.1 | Mediterranean sites where the impacts of innovative agricultural practices have been surveyed.**

These practices include “conservation agriculture”, i.e., no or reduced tillage, organic amendments, cover crop, and two agroforestry sites. Dark green dots: sites where the impact of these practices on soil carbon have been measured, blue dots: sites where impacts on soil hydrology have been measured, orange circles: sites where the impacts on yields have been measured or surveyed.



biosis in reducing drought effects (Rapparini and Peñuelas 2014), but more studies are needed to elucidate the relevant mechanisms. Experiments are being carried out worldwide with different types of plants (Tyagi et al. 2017; Pavithra and Yapa 2018; Zhang et al. 2019), with particular focus on the efficient cooperation between nodulation and arbuscular mycorrhizal fungi (AMF) for legumes (Foyer et al. 2018). Understanding the AMF-mediated mechanisms that are important for regulating the establishment of mycorrhizal association and plant protective responses to unfavorable conditions will open up to new approaches to exploit AMF as a bioprotective tool against drought (Bahadur et al. 2019). Antagonistic interactions between barley and AMF have been observed under drought conditions, particularly at high AMF richness (Sendek et al. 2019) and suggest that unexpected alterations to plant-soil biotic interactions could occur under climate change.

Despite the benefits of AMF inoculation to crop production under water deficit, outcomes and challenges of AMF application for practical use in crop production may vary, e.g., in the event of possible colonization competition between the native populations of AMF in soils and the introduced symbionts (Posta and Duc 2019). Recent research has shown that compatible combination of AMF with other beneficial microbes such as plant growth-promoting bacteria offering syner-

gistic effects on plant tolerance to stressful environments including drought stress is a promising perspective (Rahimzadeh and Pirzad 2017). Studies on quantitative trait loci involved in mycorrhizal plant responses to drought stress are needed for breeding programs to create new cultivars with a combination of drought-tolerant traits and AMF benefits. Although biotechnology practices have already made the production of efficient arbuscular mycorrhizal fungal inoculants possible for the past 15 years (e.g., Barea et al. 2005), the farmers' awareness and acceptance of (relatively expensive) mycorrhizal inoculation remain low (Posta and Duc 2019). To conclude, while AMF inoculation in crop productions under water deficit seems promising, it has not yet proven its ability to be usable and successful for Mediterranean farming systems.

### Composting

Composting technology is a modern technology that can produce a stable humus complex, used as high quality compost, providing plants with all required nutrients and micro-elements. Producers claim that the structure of this humus may increase the water holding capacity of soils by up to 70%, and have established composting facilities with organic farms in the Egyptian desert (Bandel 2009). However, results regarding water holding capacity development and enhanced resistance to drought are currently limited.

## 6.5 Wildfires

Mediterranean-type ecosystems are characterized by hot and dry summers and strong seasonality (Olson et al. 2001). Cool wet winters promote biomass growth and extended summer drought favors the regular occurrence of wildfires (Batllori et al. 2013). Historically, fires started by lightning during wet or dry storms, which can be very common in many Mediterranean-type ecosystems (Pineda and Rigo 2017). The geographic location of Mediterranean regions also benefits the frequency of strong wind events that further exacerbate fire activity. These ecosystems are dominated by fire-adapted vegetation resulting from a long evolutionary association with fire (Pausas and Keeley 2009), where usually crown and high-intensity fires largely prevail (Keeley et al. 2012a; *Section 4.3.3.1*).

Ever since prehistoric times, natural fire regimes have been altered by human activity in a multitude of ways, by modifying fuel structure, igniting

new fires and extinguishing wildfires (Bowman et al. 2011; Keeley et al. 2012b). In highly populated areas, such as the Mediterranean Basin, it makes little sense to refer to a "natural" fire regime because the footprint of human dynamics has interacted with natural factors to mold fire regimes in time and space, and makes the characterization of a 'baseline' fire regime nearly impossible (Lloret and Vilà 2003). The alteration of ecosystems at unprecedented rates may lead to unidentified changes, making natural systems unable to persist within their natural variability regimes (Vitousek et al. 1997), potentially reaching no-return ecological states during this century (FAO 2013; Batllori et al. 2017).

### 6.5.1 Future wildfire risks

The present escalation of environmental changes is modifying fire regimes and producing new

challenges for conservation management. In Mediterranean-type ecosystems of the European countries, afforestation linked to rural abandonment has occurred in recent decades (*Section 4.3.1.2*) and has shifted the systems to weather-limited fire regimes (Moreira et al. 2001; Pausas and Fernández-Muñoz 2012), in which the occurrence of fire-weather conditions drives fire activity (Pausas 2004), increasing the uncontrollability of fire events. The increase of adverse weather events associated with warming climate has stimulated an unsustainable fire regime perceived as a threat by society. Urbanization of rural areas during the second half of the 20th century has further modified fire dynamics, aggravating fire hazards due to the increase in ignition sources in these areas and an increased exposure of human activities to fire effects (Lampin-Maillet et al. 2011).

Direct human fire actions have also altered fire regimes (Bowman et al. 2011; Loepfe et al. 2011; Oliveira et al. 2012; Brotons et al. 2013; Chergui et al. 2017; Costafreda-Aumedes et al. 2017). Besides altering the spatial distribution of fuel, humans have also directly affected fire regimes by boosting anthropic ignitions and by suppressing fires with investments in huge fire-fighting structures (*Section 4.3.3.1*). In European Mediterranean countries, fire management policies basically rely on the fire suppression principle, and the increasing effort made in this direction has strongly modified fire regimes (Brotons et al. 2013; Turco et al. 2013; Moreno et al. 2014; Otero and Nielsen 2017).

Climate change in the Mediterranean Basin is projected to increase summer heat wave events, extend fire seasons, increase yearly average temperatures and increase precipitation irregularities (*Section 2.2.5*) (Field et al. 2014). How these changes will impact wildfires is still being studied (Westerling et al. 2011; Batllori et al. 2013). While a warmer climate will upsurge fire activity by increasing water demand and decreasing fuel moisture, this increase in temperatures may also lead to a decline in ecosystem productivity and thus to an overall reduction of fuel biomass (Flannigan et al. 2009; Batllori et al. 2013), which can potentially counteract warming effects on fire activity. Climate change may also promote the occurrence of other disturbances (forest outbreaks, windstorms, non-indigenous etc.) that can result in new drivers of fire regime change (*Section 2.4.1.1*). There is still a significant gap in the understanding and projection of future climate shifts and its impacts on ecosystems (Schoennagel et al. 2017; *Section 4.3.2.1*).

### 6.5.2 Management approaches, governance, and adaptation for wildfires

Changing fire regimes are now one of the most significant risks to natural systems and societies in places such as the Mediterranean Basin (Pausas et al. 2009). A deeper understanding of fire dynamics is therefore needed to enhance possibilities of successful biodiversity conservation strategies at the ecosystem level. In addition, a comprehensive understanding of fire regime patterns and processes will help to transform our societies within the resilience paradigm (Tedim et al. 2016). In recent decades, a rise in urbanization at the wildland-urban interfaces has led to an increasing number of fatalities (Moritz et al. 2014). The political response has been directed towards trying to eliminate fire from the system, with very limited success anywhere in the world (San-Miguel-Ayanz et al. 2013; Moritz et al. 2014; Archibald 2016; Tedim et al. 2016). There is an ongoing effort to promote development under which people are less vulnerable and more resilient to fire impacts (*Section 5.1.3*).

The understanding on how the different drivers of change can further impact fire regimes is still limited (Flannigan et al. 2009; Westerling et al. 2011; Regos et al. 2014). However, there is no clear consensus on future land-cover change directions because they rely more on local economic drivers with high uncertainty in their long-term predictions (Rounsevell et al. 2006). In addition, the complex interactions of drivers, the cascading effects of sequential disturbances (Batllori et al. 2017), and the uncertainty of future conditions (Thompson and Calkin 2011) make the projection of future changes a major challenge. Fire research requires further tools and approaches that help to understand ongoing changes and provide solutions to help to make effective decisions.

Available evidence from recent decades show a steady increase in wildfire events leading to extreme wildfire events escaping from fire-fighting efforts, reaching acute fire intensities and often burning very large areas (San-Miguel-Ayanz et al. 2013) (*Section 2.6.3.3*). Extreme wildfires have more significant consequences for societies and ecosystem properties than small fires (Adams 2013; San-Miguel-Ayanz et al. 2013; Tedim et al. 2013), and their occurrence is based on outstanding environmental conditions (San-Miguel-Ayanz et al. 2013). In European countries from the Mediterranean Basin, the appearance of these wildfires has been related to an expansion of forests interacting with increasingly hotter and drier weather conditions (Tedim et al. 2013). The high fuel loads accumu-

lated in forests have resulted in intense fire behaviors (high flames, fire spotting capacity) that make them very difficult for fire-fighting brigades to control. Moreover, suppression systems often collapse when protecting dispersed human assets, diminishing direct fire suppression effectiveness. Under a climate change context, these extreme wildfires are projected to increase (Amatulli et al. 2013).

### 6.5.3 Case studies

Fire suppression strategies based on proactive opportunity search and advanced fire behavior (Castellnou et al. 2019) have been successful in some regions. However, increasing fuel loads and greater climate vulnerability make fire-fighting strategies prone to collapse in the event of extremely large or intense large fires, which has already happened in countries such as Greece and Portugal in recent years. Proactive systems may open the way for local stakeholders to participate in fire-fighting decisions (Otero et al. 2018). However, the key tractable factor behind potential reduction in future aggressive fire behavior is fuel availability. On these lines, different regions deploy prescribed fire techniques to decrease fuel loads in particular areas. However, contrary to other places with Mediterranean-type climate (such as Australia and California), deployment of prescribed fire over large tracts of land raises public concerns and is

difficult to implement in Mediterranean countries, particularly in areas with a high percentage of private property (Fernandes 2018). In these cases, a combination of prescribed fire with other forest management techniques (such as using fuel for energy biomass) may be used (Regos et al. 2016). On the other hand, large tracts of conifer and eucalyptus plantations may increase the overall fire risk at the landscape scale, especially in comparison with mature native forests or more open farmland-dominated landscapes (Bowman et al. 2019).

### 6.5.4 Innovation

The key to sustainable, fire resilient landscapes is the development of sustainable socio-economic activities that allow local communities to thrive while ensuring low overall landscape risk and ensuring the persistence of other natural values (Smith et al. 2016). Such nature-based solutions to fire risk management (Duane et al. 2019) arise as an area where innovation, especially social innovation, is expected to develop in the coming years (Chergui et al. 2017). Technological innovation is also rapidly being introduced into strategic and operative fire-fighting, especially in relation to the use of remote sensors for data acquisition and remote control to predict extreme weather events leading to high-risk conditions conducive to intense fires (Peterson et al. 2017).

## 6.6 Soil erosion, degradation and desertification

### 6.6.1 Future risks for soils

Soil erosion, by water or wind, is the most widespread form of soil degradation worldwide (Panagos et al. 2017b). It is widespread in the Mediterranean region and includes sheet wash, rill and gully erosion, shallow landsliding, and the development of large and active badlands in both sub-humid and semi-arid areas (García-Ruiz et al. 2013). Soil erosion significantly alters the composition of soils, has a direct impact on the biogeochemical cycles that are responsible for supporting life on Earth and significantly reduces the ecosystem services and the economic systems that rely on them (Cherlet et al. 2018). The susceptibility of Mediterranean soils to erosion, degradation and desertification under changing conditions is exacerbated by a number of factors, such as deforestation, frequent forest fires, the cultivation of steep slopes and overgrazing (García-Ruiz et al. 2013) (Section 2.4.1.2). According to the Unit-

ed Nations Convention to Combat Desertification (UNCCD, 2004), Portugal, Spain, Italy, Greece, Turkey and Morocco have a significant problem with desertification because of the occurrence of particular conditions over large areas. International and interdisciplinary research initiatives have come to support this statement and have provided ample documentation that large areas of the European Mediterranean region are being increasingly affected by desertification, e.g., the EU MEDALUS, DISMED, MEDACTION, LEDDRA projects (Kosmas et al. 1999; Drake and Vafeidis 2004; Kepner et al. 2006; Sommer et al. 2011) (Section 2.4.1.1).

The assessment of future degradation and desertification risk and whether it can be reversed with land conservation and management practices, is affected by our ability to accurately set a baseline (Behnke and Mortimore 2016) or even decide on what constitutes an alarming rate. With the very slow rate of soil formation, any soil loss of more

than  $1 \text{ t ha}^{-1} \text{ yr}^{-1}$  can be considered as irreversible within a time span of 50-100 years. However, the concept of variable tolerable rates of erosion should be noted and requires further definition (di Stefano and Ferro 2016).

Numerous efforts to estimate current erosion rates have been reported in the literature, most commonly using empirical models (e.g., RUSLE) or physical process-based models (e.g., PESERA). A recent attempt to quantify soil erosion by water over the European region using the RUSLE model has reported very high soil loss rates for European Mediterranean countries of commonly over  $50 \text{ t ha}^{-1} \text{ yr}^{-1}$ , mainly in southern Spain and Italy and to a lesser extent in Greece, Cyprus and France (Corsica) (Panagos et al. 2015). A more recent global modelling effort based on RUSLE by Borrelli et al. (2017) assessed the impact of land use change on soil erosion between 2001 and 2012. With regard to the Mediterranean Basin, it corroborated the previous findings and identified Morocco, northern Algeria, western Syria, Albania, Serbia, Montenegro and Bosnia Herzegovina as hotspots where erosion rates were predicted to increase according to a baseline scenario. Syria, Serbia, Croatia, Montenegro and Morocco were also projected to have increased soil erosion rates even with a conservation scenario. It is worth noting that soil erosion risk models contain erosivity and erodibility factors that reflect average-year rainfall. Therefore the currently available values for these factors may inadequately represent the more frequent and intense storms projected under most climate change scenarios (Jones et al. 2012). Moreover, Eekhout and de Vente (2019) have shown that applying different bias correction methods to contrasting Mediterranean conditions can lead to disparate soil erosion projections of either a future decrease or increase.

Other efforts have aimed at assessing the sensitivity of an area to degradation and desertification processes, using a system of indicators developed during the MEDALUS EU project (Kosmas et al. 1999), including soil erosivity, vegetation cover, climatic parameters (such as aridity), land use and land management. These studies have been applied in study sites throughout the Mediterranean, and have often identified hotspots of critical sensitivity to degradation and desertification (e.g., Lavado Contador et al. (2009) in Spain; Salvati and Bajocco (2011) in Italy; Symeonakis et al. (2014) in Greece; Kamel et al. (2015) in Lebanon; Boudjemline and Semar (2018) in Algeria, and Ait Lamqadem et al. (2018) in Morocco). Právělie et al. (2017) also applied this approach to the entire European Mediterranean for the years 2008 and 2017 and

found widespread increases in sensitivity to desertification: the amount of territory with a high or very high sensitivity to desertification had increased, in less than a decade, by 177,000  $\text{km}^2$ .

Adding to the complexity of assessing the future risks related to soil erosion and the reversibility of related degradation and desertification, climate change is expected to alter erosion rates in a complex, non-linear way. Rainfall changes (in either the intensity only or in the amounts as well), along with expected changes in temperature, solar radiation, and atmospheric  $\text{CO}_2$  concentrations, will have significant impacts on soil erosion rates (Nunes et al. 2013; Li and Fang 2016; Zare et al. 2016; Zhou et al. 2016; Guo et al. 2019). Kirkby et al. (2004) describe a non-linear spatial and temporal response to climate change, with relatively large increases in erosion during wet years compared to dry years, and sporadic increases locally. However, the processes involved in the impact of climate change on soil erosion by water are complex, involving the abovementioned changes in rainfall amounts and intensities, the number of days of precipitation, plant biomass production and residue decomposition rates, soil microbial activity, evapotranspiration rates, and shifts in land use necessary to accommodate the new climatic regime (Nearing et al. 2004). Projections of changes in factors related to desertification indicate significant exacerbation of desertification risk in southern Europe and particularly in Spain, southern Italy, and Greece (Panagos et al. 2017a; Samaniego et al. 2018).

### 6.6.2 Management approaches, governance, and adaptation for soil protection

Soil erosion is greatly affected by human-environment interactions, most notably land use and land use changes. However, overly simplistic cause and effect approaches to what leads to degradation and desertification have now been abandoned (Cherlet et al. 2018) as the complex nature of non-equilibrium systems has been identified and acknowledged (Reynolds et al. 2007; Behnke and Mortimore 2016). A more integrated land management approach is currently driving policy-making, including the development and implementation of adaptive practices of sustainable land management. The World Overview of Conservation Approaches and Technologies (WOCAT) is a network that develops, archives, shares and disseminates sustainable land management knowledge to improve human livelihoods and the environment (Lingiger et al. 2007), gaining broad appreciation from all involved stakeholders.

Sustainable land management approaches are continuously adapted in response to changing environmental conditions and human needs. From a list of hundreds of archived case studies of sustainable land management in the Mediterranean region, five types of measures are identified that can be taken to address land degradation (Sections 3.2.3.2 and 4.3.3.3): (i) agronomic measures: measures that improve soil cover (e.g., green cover), measures that enhance organic matter (e.g., manuring), soil-surface treatment (e.g., conservation tillage), and sub-surface treatment (e.g., ripping); (ii) vegetative measures: plantation of trees and shrubs (e.g., live fences), grasses and herbaceous plants (e.g., grass strips); (iii) structural measures: terraces, bunds, banks, dams, pans, ditches, walls, barriers, and palisades; (iv) management measures: change of land use type, change in management/intensity level, change in timing of activities, and control/change of species composition, and (v) combinations of the other four types (Liniger et al. 2007; Cherlet et al. 2018).

With regard to policy, at the moment, only a few EU Member States have specific legislation on soil protection. Soil is not subject to a comprehensive and coherent set of rules in the European Union. Existing EU policies in areas such as agriculture, water, waste, chemicals, and prevention of industrial pollution indirectly contribute to the protection of soils. However, as these policies have other aims and scopes of action, they are not sufficient to ensure an adequate level of protection for all soils in Europe<sup>46</sup>. A limited number of countries or Autonomous Regions have Soil Protection Plans (e.g., the Basque Autonomous Country (Landeta 1995), Italy (Law 97 of 1994)) while a much larger number have ratified the UNCCD and have prepared a National Programme to Combat Drought and Desertification or National Action Plan, namely, Algeria, Egypt, Greece, Italy, Lebanon, Morocco, Portugal, Spain, Tunisia and Turkey.

### 6.6.3 Case studies

Based on the WOCAT classification of measures that address soil erosion and land degradation, the following is a successful example of a structure measure from Spain. Rodrigo-Comino et al. (2017) assessed agri-spillways as a soil erosion protection measure in Mediterranean sloping vineyards in southern Spain. Their results showed a great capacity by rills to canalize large amounts of water

and sediments, and higher water flow speeds and sediment concentration rates than typically found in other Mediterranean areas and land uses (such as badlands, rangelands or extensive crops of olives and almonds). They concluded that agri-spillways can be a potential solution as an inexpensive method to protect the soil in sloping Mediterranean vineyards.

Another example for sustainable land management comes from Italy, a case of a vegetative measure. Bagagiolo et al. (2018) studied the effect of controlled grass cover on water and soil losses in different rain-fed sloping fields in northwestern Italy. Rainfall, runoff and erosion variables were monitored in hydraulically bounded vineyard plots, where the inter-rows were managed with tillage and grass cover. The grass cover proved to be effective in decreasing runoff and soil losses during most of the events, reducing soil losses especially when intense events occurred (i.e., during summer). Their results also showed the fundamental role of contour-slope row orientation in reducing runoff and soil losses, irrespective of the adopted inter-row soil management approach.

### 6.6.4 Innovation

Land Degradation Neutrality (LDN) is a new conceptual framework, introduced by the UNCCD to halt the loss of land due to unsustainable management and land use changes (Cowie et al. 2018). Its purpose is to maintain the land resource base so that it can continue to supply ecosystem services while enhancing the resilience of the communities that depend on the land (Metternicht et al. 2019). The LDN framework is designed to apply to all land uses and all types of land degradation. To achieve LDN, countries will need to assess the effect of land use decisions and undertake measures to restore degraded land so as to compensate anticipated losses (Cowie et al. 2018). The UNCCD suggests that countries should consider the social, economic and environmental outcomes of alternative land degradation and desertification mitigation options when planning LDN measures and should strive to engage relevant stakeholders. Some applications of the LDN framework have only just begun to materialize (e.g., in south-east Australia, Cowie et al. 2019), but none have yet been applied in Mediterranean countries or climates.

<sup>46</sup> [https://ec.europa.eu/environment/soil/index\\_en.htm](https://ec.europa.eu/environment/soil/index_en.htm)

## 6.7 Heat waves

Temperature extremes occur on different time scales and need temporally high-resolution data to accurately assess possible changes (IPCC 2012). Temperature is associated with different types of extremes. It is of importance to distinguish between maximum, minimum and daily mean, as well as between cold and warm extremes, as they have different impacts on human health (Sections 5.2.3 and 6.2), the physical environment (Section 2.3), ecosystems (Section 4.3), and energy consumption (Section 3.3). Increases in the intensity, number, and length of heat waves have been reported for Mediterranean summers since the 1960s (Kuglitsch et al. 2010; Efthymiadis et al. 2011; Lelieveld et al. 2012) (Section 2.2.4.1).

### 6.7.1 Future heat wave risks

Future projections for the Euro-Mediterranean area have shown spatial heterogeneity in increases in the intensity, frequency and duration of heat waves (Section 2.2.4.2). Major increases in warm temperature extremes are expected across the Mediterranean region (Jacob et al. 2014; Russo et al. 2015; Zittis et al. 2016; Pereira et al. 2017) including hot days ( $T_{\max} > 30^{\circ}\text{C}$ ) and tropical nights ( $T_{\min} > 20^{\circ}\text{C}$ ) (Giannakopoulos et al. 2009; Tolika et al. 2012). Larger increases in intensity and duration are projected for southern Europe where heat wave days are projected to increase 20-fold by 2100 (Fischer and Schär 2010). Other projections over the Mediterranean include dramatic increases in the frequency of hot temperature extremes and heat stress by the end of the 21st century (Section 2.2.4.2). Cities in southern Europe are expected to face longer heat waves (Guerreiro et al. 2018), thus increasing their vulnerability to climate impacts and the need for costly adaptation measures.

Projected changes in the characteristics of future heat waves are related to increasing risks in several sectors. Intense and long heat waves are related to increased morbidity and mortality in Mediterranean countries, especially in cities where the built environment amplifies the exposure to heat (Sections 5.2.2.8 and 6.2). Increasing temperatures affect overall energy demand for cooling, while heat waves may also affect peak demand that is mainly provided by electricity (EEA 2019a). The largest absolute increases in electricity peak demand are projected for Italy, Spain and France (Damm et al. 2017). The tourism sector plays an important role for the economic well-being and livelihoods of Mediterranean countries

(Section 5.1.1.3). Frequent heat waves may reduce tourist flows by the mid-21st century due to exceeded comfort levels (Hein et al. 2009) and could shift tourist demand outside the peak summer time (Perry 2003; Esteban-Talaya et al. 2005; Ciscar et al. 2009). Future increased extreme temperatures will increase the impact on transport infrastructure in the Mediterranean and will lead to damage to roads, rail, airports, and ports (Nemry and Demirel 2012; UNCTAD 2017; Vogel et al. 2017) with significant increases in adaptation costs (Nemry and Demirel 2012). High temperatures and drought will increase forest fire risk, which might lead to drastic damages in Mediterranean forests (Trigo et al. 2013; Gudmundsson et al. 2014; Turco et al. 2018) (Section 4.3.2.1). High future temperatures and heat waves have a direct impact on crop growth conditions, crop productivity and crop distribution, agricultural pests and diseases, and the conditions for livestock production in the Mediterranean (Section 3.2.1.4). These impacts will generate changing land-use patterns and will trigger economy-wide effects (Skuras and Psaltopoulos 2012). In southern Europe, yields for all the dominant (non-tropical) crops decreased by 5-60% because of climate change, depending on the country, the crop and the scenario (Section 3.2.2.1). The combined effect of extreme heat events and shorter growing seasons will result in a loss of land suitable for agriculture (Fraga et al. 2016; Resco et al. 2016; EEA 2019b) in southern Europe (Section 3.2.2.1). Furthermore, the Mediterranean agro-climate zone is expected to experience pronounced increases in the areas affected by mild to strong heat stress, which will occur earlier and will impact winter wheat (Ceglar et al. 2019).

### 6.7.2 Management approaches, governance, and adaptation for heat wave risks

Reducing the direct impacts of extreme temperatures requires focus on information and preparedness associated with early warning (PPRD East 2013). The need for the implementation of early warning systems has risen since the 2003 summer heat wave (García-Herrera et al. 2010). Adaptation for heat waves in cities is a major challenge in design and costs estimation (Guerreiro et al. 2018). Prevention in the long term must further ensure that the vulnerability of the population and relevant infrastructure are reduced by improving urban planning and architecture (e.g., increasing the canopy cover in urban areas, cooling open

public areas, adjustments in energy generation and transmission infrastructure), as well as through energy and transport policies (PPRD East 2013). Strategies are needed to reduce heat exposure of individuals and communities (especially vulnerable populations), to plan health and social services and infrastructure, and to provide timely information to the population (Future Earth 2019). Some of the adaptation measures for the project-

ed changes entail fundamental, and expensive re-engineering of each city or water resource system. In the Mediterranean, significant adaptation measures to climate extremes, primarily in the form of structural protection measures, have already been implemented in the framework of the adaptation plans at the city, regional and national levels across the Mediterranean Basin.

## 6.8 River and pluvial flooding

### 6.8.1. Future flood risk

The Mediterranean region is characterized by numerous water courses with small and steep river catchments (Tarolli et al. 2012; Trambly et al. 2019), although with notable exceptions, such as the Nile, Rhone, Ebro and Po rivers (*Section 3.1.1.1*). The steep orography surrounding the Mediterranean Sea favors the occurrence of intense precipitation events triggered by spatially confined convective processes (Amponsah et al. 2018), especially in autumn (Gaume et al. 2016) (*Section 2.2.5*). The resulting runoff can produce devastating flash floods in small river basins, i.e., less than 2,000–3,000 km<sup>2</sup> in size (Amponsah et al. 2018), especially where urbanized areas are located downstream of these small basins (Llasat et al. 2010; Gaume et al. 2016) (*Section 3.1.3.3*).

The magnitude and impact of floods vary significantly over the Mediterranean region, with more frequent and severe events in the western part (Llasat et al. 2010; Gaume et al. 2016). Some sub-regions in southwestern Europe, including Liguria and Piedmont in Italy, Cévennes-Vivarais-Roussillon in France, and Catalonia and the province of Valencia in Spain are particularly prone to extremely severe events, due to geographic and climatological conditions (Gaume et al. 2016). Floods in Morocco, Algeria and Tunisia are less frequent but they are often associated with high mortality, while European countries suffer the highest economic damages (Llasat et al. 2010).

Trends in annual maximum peak flow in European Mediterranean countries have been decreasing in the past decades (Blöschl et al. 2019). However, no significant trend in the frequency and magnitude of extreme floods has been found for the Mediterranean as a whole (Gaume et al. 2016), or for large regions such as Catalonia and southern France (Llasat et al. 2005, 2014; Trambly et al.

2019), even though local increasing trends have been observed (e.g., Genoa urban area, Faccini et al. 2018; *Section 3.1.3.3*). Future trends in flood patterns still appear unclear, with different studies reporting contrasting results (Kundzewicz et al. 2017), partly because of the limitations of regional- and global-scale models in representing small catchments (Trambly et al. 2019) (*Sections 3.1.4.1 and 3.1.4.2: Floods*).

### 6.8.2 Management approaches, governance, and adaptation for flood protection

Flash flood risk management presents several challenges with respect to other types of flooding processes. The triggering meteorological and hydrological processes are difficult to monitor with traditional hydro-meteorological networks, given the small spatial and temporal scales involved (Amponsah et al. 2018). Moreover, flash flood risk can be associated with other hazards, particularly in mountain settings (e.g., landslides and debris flows). This complicates the implementation of forecasting and early warning systems as well as the design of physical flood defense infrastructure (Borga et al. 2011). Preparedness strategies need to be structured in accordance with these and other characteristics, such as short to negligible warning lead times, immediate threat to life and properties requiring quick response times, as well as the need for refuges and safe places (Borga et al. 2011). This requires effective coordination of response management by authorities and public awareness.

Good practices in flash flood risk management reported in the literature and applied in several case studies include: post-event surveys to collect information on flood-generating processes and impacts (Kreibich et al. 2017; Amponsah et al. 2018), development of dedicated early warning systems

(EWS) based on gauge and radar networks, numerical weather and hydrological predictions [Corral et al. 2019], construction of check dams and reforestation in upstream areas [Kourgialas and Karatzas 2017], floodplain restoration and bank erosion protection [Kourgialas and Karatzas 2017; Cortès et al. 2018], suitable agricultural practices to retain water and reduce flood damage to crops [Kourgialas and Karatzas 2017], improvement of drainage systems in urbanized areas [Cortès et al. 2018], increased citizen awareness [Borga et al. 2011; Cortès et al. 2018; Faccini et al. 2018], emergency management plans [Kreibich et al. 2017], and viable insurance schemes for damage compensation [Faccini et al. 2018].

### 6.8.3 Case studies and innovation

At the European level, the European Directive on Floods (Directive 2007/60/CE, European Parliament 2007) regulates flood risk management plans, focusing on prevention, protection and preparation. The implementation of the Floods Directive has driven notable improvements, also in flash flood risk management. According to Kreibich et al. [2017], vulnerability to flash floods was greatly reduced in recent events in Italy and Spain as compared to similar events that occurred several decades ago, due to improved awareness, preparedness and emergency management.

Cortès et al. [2018] report that in the Metropolitan Area of Barcelona, the implementation of prevention measures such as constructing rainwater tanks, or the establishment of warning systems,

decreased the impacts of flood events between 1981 and 2015. Nowadays, different flash-flood forecasting systems are present in Catalonia (Spain), Liguria (Italy) and Southern France [Corral et al. 2019]. Notably, the European Flood Awareness System (EFAS)<sup>47</sup> provides different flash flood indicators [Raynaud et al. 2015; Corral et al. 2019], and has recently been extended to the entire Mediterranean Basin, therefore offering the first pan-Mediterranean forecasting system for river and flash floods. Finally, in Spain and France, dedicated national insurance schemes against natural disasters exist, which cover losses through economic compensation.

However, not all Mediterranean areas benefit from recent advances. Information on flood hazard and risk is missing or scarce in some southern and eastern Mediterranean countries (Llasat et al. 2010), as well as in small and ungauged catchments in Europe [Kourgialas and Karatzas 2017]. Adaptation plans in southern Europe suffer from a lack of funding in rural and low-populated areas [Aguar et al. 2018]. Challenges are still present even in large cities. For example, the city of Genoa, Italy, is particularly exposed to flash floods due to its geographical location, meteorological conditions and dense urbanization with inadequate planning (e.g., reduced or culverted river network in the river valleys) [Faccini et al. 2018]. While progress has been made in increasing citizen awareness and improving early warning systems, structural solutions (e.g., diversion channels, relocation of the most exposed properties) appear unfeasible due to the large areas involved.

## 6.9 Sea-level rise: coastal erosion and flooding, saltwater intrusion

### 6.9.1 Future risk associated with sea-level rise

Mediterranean mean sea levels are projected to rise by 21 to 27 cm by 2050, under RCP4.5 and RCP8.5 scenarios, respectively [Jackson and Jevrejeva 2016; Jevrejeva et al. 2016]. By the end of the century, the mean sea level would range between 20 cm and 110 cm above the present level (1980-1999), depending on the greenhouse gas emission scenario and the modeling system (Section 2.2.8.2). Such sea-level rise, combined with variations in extreme weather and thus waves and storm surges, will substantially increase the frequency of extreme events as

the present day event of the century is expected to occur every 10 years by 2050 and at least yearly by the end of the century [Vousdoukas et al. 2018b]. All the above changes are projected to expose Mediterranean societies to unprecedented levels of coastal flooding and losses. Without considering socio-economic development, a 6 to 8-fold increase in annual damage is expected by 2050 and at least 25 times more annual damage is expected by the end of the century if no further investments in coastal protection are undertaken [Vousdoukas et al. 2018a]. When climate change projections are combined with socio-economic scenarios, expected annual damage is projected to rise by 90 to 900 times, depending on

<sup>47</sup> <https://www.efas.eu/>



the scenario. Adaptation in the form of dykes can cut damage costs in half, with countries such as France, Spain, Greece, and Italy having the highest damage costs in absolute terms (Hinkel et al. 2010), and Egypt and Tunisia facing the highest damages relative to their annual Gross Domestic Product (GDP) (Hinkel et al. 2012). Accordingly, Italy and France have the largest length of coasts where protection would be economically beneficial (Vousdoukas et al. 2020).

Most coastal regions globally are exposed on a daily basis to tidal water level variations of more than 50 cm, and ocean waves, which require wider active beach zones to act as a buffer against the ocean's forces. This is not the case in the Mediterranean, which is a micro-tidal area where a significant part of the coastline is not exposed to harsh marine storms (Section 2.2.8). The above-mentioned characteristic makes the Mediterranean more susceptible to coastal hazards in view of climate change compared to other parts of the world. It is important to highlight that for many Mediterranean locations; the projected sea-level rise is of similar magnitude to the increase in sea levels during extreme events. At the same time, communities have developed lifestyles adapted to non-dynamic water levels, as several activities take place and infrastructure is located in close proximity to the sea (within few meters in many cases). This is also because apart from local-scale erosion, the coastline has been relatively stable for global standards with the exception of some cases of stronger shoreline retreat trends, observed in the Nile delta, Tunisia, Venice, and Albania (Luijendijk et al. 2018; Mentaschi et al. 2018).

Finally, interconnected hazards may exacerbate issues related to sea-level rise. For example, while the coastal environment encompasses particular characteristics distinct from general issues of water (such as shortages and drought) and precipitation (or lack thereof), there are numerous interconnections between water runoff, drainage and watershed management that are linked to hazards related to sea-level rise (O'Connor et al. 2009; Lichter and Felsenstein 2012; Portman 2018). Such hazards may result in compound effects that can lead to non-linear increases in the magnitude of individual hazards.

### 6.9.2 Management approaches, governance, and adaptation for coastal protection

It is important to highlight that coastal erosion in Mediterranean countries has been primarily driven by human interference with natural processes

(Section 4.2.1). For example, inadequate coastal management practices and, most importantly, unregulated construction have been reported in several regions (ERML 2012; de Leo et al. 2017; UNDP 2017). A recurring problem is the reduction or depletion of terrestrial sediment supply, that would naturally feed sandy beaches, resulting from the construction of upstream dams (Poulos and Collins 2002) (Section 4.2.1.2). Such examples include the Beni Khair and Dar Chaabane coasts and the Oued El Kebir river (Imen and Souissi 2018), Lesvos Island (Velegrakis et al. 2008), and Rhodope, Greece (Xeidakis et al. 2006).

Coastal adaptation practices can be classified into the following broad categories: protect, accommodate, advance, and retreat. Under protection practices, societies tend to "hold the line" by installing coastal protection elements. Traditionally these were mainly "hard structures" such as breakwaters and seawalls (Lamberti and Zanuttigh 2005). Dykes are another potential flood prevention solution, but they are very rare in the Mediterranean, as they are more common in meso-/macro-tidal environments. The same applies for surge barriers, with the only example being the MOSE system in Venice (CVN 2019). Submerged breakwaters reduce wave energy and mitigate erosion and have also become common practice along the Mediterranean coastline (Tomasicchio 1996; Sancho-García et al. 2013; Bouvier et al. 2017).

"Soft-protection", in the form of beach and shore nourishment as well as dune or wetland restoration, has become a more common alternative to hard structures in recent decades, with many examples, especially in France, Spain and Italy (Hamm et al. 1998; Hanson et al. 2002). Lately there is a tendency towards Ecosystem-based Adaptation (EbA) (Section 4.2.3.5), also referred to as "soft protection", using ecological features such as reefs and/or coastal vegetation as coastal protection elements. Among the few examples of EbA is the coastal protection service provided by the Étang de Vic coastal lagoon in France (Conservatoire du littoral)<sup>48</sup> and the coastal dune reconstruction at the natural protection area of the Bevano river mouth in Emilia Romagna (Italy) (Giambastiani et al. 2016).

Until recently, advances through land reclamation has been more related to the need for more space to accommodate human activities (Mentaschi et al. 2018), but is also being increasingly considered

<sup>48</sup> [http://www.conservatoire-du-littoral.fr/siteLittoral/106/28-etang-de-vic-34\\_herault.htm](http://www.conservatoire-du-littoral.fr/siteLittoral/106/28-etang-de-vic-34_herault.htm)

in the context of adaptation to sea-level rise. However, this practice is practically non-existent in the Mediterranean Sea. The same can be argued for accommodation i.e., increasing the resilience of infrastructure by making it less vulnerable to flooding. Recent studies have shown that flood fatalities have been reduced as societies are learning to live with flood hazards [Bouwer and Jonkman 2018], while there have been efforts to develop and implement Early Warning Systems for disaster risk reduction [Ciavola et al. 2011; Harley et al. 2011; Fernández-Montblanc et al. 2019]. However, there are very few, if any, examples of large-scale efforts to develop flood-resilient buildings around the Mediterranean coastline. The same applies to the retreat option in which exposure to coastal hazard is reduced by removing assets and people from potentially vulnerable areas.

### 6.9.3 Case studies

Hard protection structures can be found all along the Mediterranean coastline and in most cases they contribute to sustaining a safe and functional coastal zone [Iskander et al. 2007; Becchi et al. 2014]. However, as it has been already pointed out in other parts of the world [Cooper and Pilkey 2012], this comes at a price. Hard protection can alter nearshore sediment transport patterns and result in beach erosion. Such side effects have been observed in Greece and Cyprus [Tsoukala et al. 2015], Tunisia [Saiidi et al. 2012], and Egypt [Masria et al. 2015]. In addition, hard structures can affect the nearshore ecology, as they can act as habitats for species which normally thrive in rocky shores [Munari et al. 2011]. However such effects have been shown to depend on local conditions and not to be overwhelming [Colosio et al. 2007; Becchi et al. 2014].

There have been several beach nourishment projects along the Mediterranean coastline, some of which have been reported in the scientific literature [Hamm et al. 1998; Hanson et al. 2002; Masria et al. 2015] (*Section 4.2.1.1*). These initiatives are ecologically milder but can still come with negative impacts [Colosio et al. 2007]. For example, nourishment at Poniente Beach (Benidorm, Spain) has been shown to have caused the disappearance of the *Posidonia oceanica* meadows, which resulted in a strong beach erosion process [Aragonés et al. 2015]. However, there are several studies which report that small-scale beach nourishments appear to be an eco-sustainable

approach to combat coastal erosion [Borg et al. 2006; Danovaro et al. 2018]. Geotextiles have been installed in several locations as a soft protection practice, but information on their performance is limited in the scientific literature with a few exceptions, such as the positive outcome in Lido de Sete, in France [Balouin et al. 2015]. It is important to highlight that most of the literature shows that no universal solution exists and that robust planning and implementation is a prerequisite for any successful intervention.

### 6.9.4 Innovation

Risk and climate change adaptation efforts are inextricably linked. Having acknowledged risks, some countries have developed either “resilience toolkits” (e.g., US) or “adaptation toolkits” (e.g., Ireland) that address how civil society must prepare for hazards, with emphasis on coastal areas [Paterson et al. 2017; McDermott and Surminski 2018; Gardiner et al. 2019]. Most Mediterranean countries are lagging behind in this respect. Recently there has been significant work on at least assessment of future risks pertaining to air, water, and sea [Navarra and Tubiana 2013]. However, little has been done on the aspects of extreme hazards and the effects of climate change on society, which could encourage more resources (both human and financial) being dedicated to adaptation planning. Nevertheless, some examples of such actions exist. Countries such as Italy, France and Spain have established national and subnational initiatives on coastal adaptation and management<sup>49</sup> [Losada et al. 2019] while multi-national initiatives such as the Bologna Charter<sup>50</sup> have introduced action plans for the protection and sustainable development of coastal areas in the region through e.g., the establishment of a network of coastal observatories.

At the same time, interconnections between different types of hazards need to be addressed in research, planning and management for adaptation. To some extent, such interconnections are recognized and have led to initiatives. One example is the DANUBIUS-RI [Bradley et al. 2018], which is a platform designed to support interdisciplinary research on rivers and seas by facilitating biogeochemical monitoring while also spanning various aspects of environmental, social and economic sciences. These types of initiatives will no doubt support projects and future risk assessments related to climate change.

<sup>49</sup> [www.erosionecostiera.isprambiente.it](http://www.erosionecostiera.isprambiente.it)

<sup>50</sup> [www.bolognacharter.eu](http://www.bolognacharter.eu)

There is still a lack of information on the risks associated with the economic, livelihood and cultural consequences of coastal change (Reimann et al. 2018b) at the regional scale that would encourage progress towards the international and transboundary cooperation needed to address these challenges among Mediterranean countries. Transboundary cooperation is particularly difficult in the deep-sea areas, far from national jurisdiction. In these areas, cooperation is voluntary, often temporary and malleable at best, and non-existent at worst, even though it is compul-

sory for EU Member States based on Directive 2014/89/EU. Beyond the EU Mediterranean space, cooperation is voluntary. Much more oversight, accountability and especially monitoring is needed internationally (Neumann and Unger 2019), particularly in the Mediterranean. With regard to climate change, the “Our Ocean” Conference series, which has a strong topical relationship with SDG 14, has adopted climate change as one of its six areas of action (others are: marine protected areas, sustainable fisheries, marine pollution, sustainable blue economy, and maritime security).

## 6.10 Seawater temperature anomalies and extremes

### 6.10.1 Future risk of marine heat waves

Marine heat waves are periods of extremely warm sea surface temperature that persist from days to months and can extend up to thousands of kilometers (*Section 2.2.7.1*). Recently observed marine heat waves demonstrated the strong influence of extreme climate events on marine organisms, including mass mortalities and shifts in species ranges (Rosenzweig et al. 2008), but also economic impacts on fisheries and aquaculture (*Section 4.2.1.1*). In coastal areas at regional scales, little is known about the propagation at depth of a warming signal detected in sea temperature surface conditions. This is due to the scarcity of continuous observational data sets over the long-term (>10 years) from surface down through the water column (+40 m depth). Analysis of in situ temperature data available from different coastal sites confirmed warming trends in deeper layers consistent with those reported for surface waters (Bensoussan et al. 2019a). Thus, the warming is not limited to the surface, but propagates into deep coastal water layers (up to 80 m depth). Importantly, this warming displays significant variability along the depth gradient depending on local thermal regimes and seasonal stratification dynamics (Garrabou et al. 2019a). Likewise, marine heat waves have been recorded along depths with different intensity and duration depending on the years and concerned areas (Bensoussan et al. 2019b). Sustained observation in pilot sites will provide important information to validate models and track subsurface warming trends.

Like their atmospheric counterpart, Mediterranean marine heat waves are expected to increase in intensity, frequency and duration under anthropogenic climate change (*Section 2.2.7.2*)

(Coumou and Rahmstorf 2012; Oliver et al. 2018). Darmaraki et al. (2019) used ensemble set of fully coupled Regional Climate Models (RCMs) from the Med-CORDEX initiative and a multi-scenario approach of different representative concentration pathways (RCPs), where marine heat waves become stronger and more intense under RCP4.5 and RCP8.5 than RCP2.6 by the year 2100. Under RCP8.5, a long-lasting Mediterranean marine heat wave appears at least once every year. Therefore, future marine heat waves appear up to three months longer, about four times more intense and 42 times more severe than at present (*Section 2.2.7.2*) and will affect the entire basin, predominantly in the warm and dry season from June to October. The main trigger can be attributed to the increase in the mean sea surface temperature (SST) and the daily SST variability. However, there is a lack of information on future trajectories of temperature conditions in coastal waters (from surface to 50 m depth and beyond) mainly due to the lack of customized modelling for these hydrodynamically complex areas. The results that are available point to an unambiguous increase in mean temperatures and frequency of extreme events, consistent with results obtained at the regional level (Garrabou et al. 2019b).

Current and future climate change trajectories are considered one of the major concerns for the conservation of marine biodiversity (Hughes et al. 2017; Cramer et al. 2018). In the Mediterranean, observed warming is already significantly affecting marine ecosystems (*Sections 4.1.1 and 4.2.1*), resulting in two main impacts: i) the shift in species distribution (indigenous and non-indigenous) and ii) the occurrence of unprecedented mass mortality events (MMEs). Besides these major impacts, other effects associated with warming are

being reported as well, such as species proliferation and changes in species reproduction timing and migration patterns (Otero et al. 2013). Overall climate change is already dramatically affecting the abundance and distribution of species as well as the functioning of ecosystems (Sala et al. 2011; Givan et al. 2017; Cramer et al. 2018). It is difficult to foresee with precision to what extent the current climate trends will affect marine ecosystems and key species in the Mediterranean Sea in the coming decades. However, recent studies indicate that an increased extinction risk for endemic fauna, loss of habitat complexity and changes in ecosystem configurations is occurring (Ben Rais Lasram and Mouillot 2009; Ben Rais Lasram et al. 2010; Sala et al. 2011; Azzurro et al. 2019; Monteiro-Serra et al. 2019).

Three main patterns in species distribution associated with warming are being observed: i) northward expansion are extremely clear for warm-affinity native species such as the bluefish, *Pomatomus saltarix* (Dulčić et al. 2005; Sabaté et al. 2012), whose Mediterranean distribution was historically restricted to the southern and eastern sectors of the basin (Whitehead et al. 1986); ii) distribution contraction of cold-water affinity species in the northern areas such as the sprat *Sprattus sprattus* (Margonski et al. 2010), whose populations have drastically declined since the 1990s in the northern Adriatic and the Gulf of Lion (Lloret et al. 2001; Grbec et al. 2002; Hidalgo et al.

2020), and finally iii) west-eastward expansion of non-indigenous warm-adapted species of tropical origin, which are expanding their presence in the Mediterranean (Raitsos et al. 2010; Azzurro and Bariche 2017; Azzurro et al. 2019), for instance the case of the rabbitfish *Siganus luridus* and *S. rivulatus*, which are rapidly expanding their distribution and increase in abundance at the expense of their native counterpart *Sarpa salpa* (Marras et al. 2015) (Sections 2.5.1 and 4.1.1).

### 6.10.2 Management approaches, governance, and adaptation for ocean warming

Monitoring marine heat waves leads to a better understanding of their development, drivers and characteristics. Monitoring of near-time sea-surface temperature based on satellite data is possible, while the use of oceanographic arrays could provide information about heat penetration in deeper ocean layers. In the Mediterranean Sea, "T-MEDNet" was created in 2010 to develop an observation network on climate change effects and to spread standard monitoring protocols on seawater temperature and biological indicators. To date, continuous, quality checked temperature series are available at >70 sites and different ocean depths (5 to 40 m; T-MedNet 2019). They also evaluate satellite-derived sea-surface temperatures to track Mediterranean marine heat waves in near real-time.

## 6.11 Ocean acidification

Ocean acidification acts together with other global changes (e.g., warming, seawater expansion) and with local changes (e.g., pollution, eutrophication) (Section 2.2.9). These simultaneous pressures and stresses lead to interactive, complex and amplified impacts for species and ecosystems (Section 4.1.1.1). Globally, a pH change of -0.08 has occurred, on average, in the acidity of the oceans since the industrial age began (Section 2.2.9.1), i.e. a 30% increase in acidity. If we continue on our present course, this will lead to a -0.46 increase by the end of the century (Section 2.2.9.2), representing a 5-fold increase in acidity (Kolbert 2014). The term "ocean(s)" here is inclusive, encompassing marine and brackish water systems, from the open ocean to coastal waters, with the latter reflecting the immediate interface of land activities affecting the ocean, which has numerous implications for both eutrophication and acidification.

One of the issues generally underlined regarding research, and management to some extent, is the problem of ocean acidification being overshadowed by other more immediate, tangible and high-profile issues affecting the marine environment, such as marine litter (Tiller et al. 2019) (Section 2.3.2.3). This is also true in the Mediterranean region where the marine plastic and marine litter issue is quite acute and where there are tangible and significant effects on economic well-being (i.e., tourism), health and well-being (Portman and Brennan 2017; Portman et al. 2019).

It is difficult to carry out long-term realistic manipulations of CO<sub>2</sub> levels, and therefore scientists have used areas with naturally occurring high CO<sub>2</sub> levels to forecast the effects of ocean acidification. In an elaborate census offshore of Naples, Italy, divers collected data around deep-sea volcanic vents

to find out which species, habitats and processes are resilient to and/or adversely affected by ocean acidification. At several hundred meters from the vents, scientists observed seaweeds of different types, sea cucumbers and urchins (by counting both sedentary flora and fauna and observing the movements of creatures). Closer to the vents, they observed that the number of species dropped. As pH levels dropped in proximity to the vents (indicating higher acidity), macroalgal habitats were found to be significantly altered. Also, mollusks or limpets, which came close to the vents, exhibited dissolved shells (e.g., with holes in them) (Porzio et al. 2011). Similar work has also been carried out more recently at Mediterranean sea vents by Vizzini et al. (2019).

With regard to close-to-shore systems, there are high levels of uncertainty about how coastal ecosystems will be affected by rapid ocean acidification caused by anthropogenic CO<sub>2</sub>, due to a lack of data. However, further study is needed to investigate whether the observed response of macroalgal communities can be replicated in different seasons and from a range of geographical regions for incorporation into global modelling studies to predict the effects of CO<sub>2</sub> emissions on the Earth's ecosystems (Porzio et al. 2011).

### 6.11.1 Future risk of ocean acidification

On a global level, not specific to the Mediterranean, some effects of CO<sub>2</sub> absorption can be explored by researching conditions with lower pH (representing greater acidity) in waters near hydrothermal vents (Portman 2016). Hall-Spencer et al. (2008) found that typical rocky shore communities with abundant calcareous organisms shifted to communities lacking scleractinian corals with significant reductions in sea urchin and coralline algal abundance. To our knowledge, this is the first ecosystem-scale validation of predictions that these important groups of organisms are susceptible to elevated amounts of pCO<sub>2</sub>. Seagrass production was highest in an area at mean pH 7.6 (1,827  $\mu\text{atm pCO}_2$ ) where coralline algal biomass was significantly reduced and gastropod shells were dissolving due to periods of carbonate sub-saturation.

Some work in the Mediterranean region has translated expected changes in ocean chemistry into impacts, first on marine and coastal ecosystems and then, through effects on services provided by these ecosystems to humans, into socio-economic costs using economic market and non-market valuation techniques (Rodrigues et al. 2013; Peled

et al. 2018). Initial evaluations suggest that the important sectors affected are tourism and recreation, red coral extraction, and fisheries (both capture and aquaculture production) (Rodrigues et al. 2013) (*Section 4.1.2.1*).

One way to assess the future impacts of ocean acidification, especially socio-economic impacts, is through the assessment of ecosystem services. A number of general studies have looked at the effects of climate change including acidification. This includes studies by Canu et al. (2015) for the general Mediterranean and by Peled et al. (2018) for the eastern Mediterranean in particular. The advantage to such approaches is that they estimate the monetary value of maintaining elements of the environment that have the potential to reduce acidification. The problem is incorporating these approaches into policy so that there is practical application (Portman, 2013).

One of the most harmful effects of acidification will be on fisheries, which are increasingly important and threatened in the Mediterranean Sea. Lacoue-Labarthe et al. (2016) contend that ocean acidification should therefore be factored into fisheries and aquaculture management plans (*Section 4.1.3.4*). Recruitment and seed production present possible bottlenecks for shellfish aquaculture in the future since early life stages are vulnerable to acidification and warming. Although adult finfish seem able to withstand the projected increases in seawater CO<sub>2</sub>, degradation of seabed habitats and increases in harmful blooms of algae and jellyfish might adversely affect fish stocks (Lacoue-Labarthe et al. 2016).

### 6.11.2 Management approaches, governance, and adaptation for ocean acidification

One approach that has been applied to encourage actions that will counter acidification is that of ecosystem services assessment. This approach aims to encourage action by evaluating the costs of inaction. Peled et al. (2018) did such an evaluation for the Israeli Exclusive Economic Zone. One advantage to their approach is that they account for permanent and temporary carbon sequestration and the use of Social Cost of Carbon (SCC) values. Overall, they find that within the context of ecosystem services, the biological component within the oceanic carbon cycle acts as a sink, which in its hypothetical absence would cause higher levels of CO<sub>2</sub> outgassing back to the atmosphere, potentially leading to greater acidification once gases are reabsorbed (Peled et al. 2018) (*Section 4.2.2.2*).

Kelly et al. (2011) posit that ocean acidification can be curbed by focusing more attention on local and regional actions within terrestrial watersheds. Ramajo et al. (2019) and others have suggested that seagrasses may provide “refugia” from ocean

acidification for associated calcifying organisms, as their photosynthetic activity may raise pH above the thresholds for impacts on calcification and/or limit the time spent below some critical pH threshold.

## 6.12 Non-indigenous species: marine, freshwater, and terrestrial

### 6.12.1 Future risks associated with non-indigenous species

Non-indigenous species may be a significant threat to biodiversity, economies and human health globally (Early et al. 2016; Tobin 2018) (*Section 2.5*). Climate change and projected climate-driven biome and thermal niche shifts, along with increases in trade and mobility, are the main drivers of non-indigenous species expansion globally (Early et al. 2016) and in the Mediterranean.

Today, the highest numbers of non-indigenous species have been recorded in high Human Development Index (HDI) and economically developed countries, which are also able to collect the most information and mobilize the best efforts to manage them (Early et al. 2016). Studies show that countries which are the biggest agricultural producers (such as China and the United States) could be the main potential sources of non-indigenous species and experience the largest negative impacts from future non-indigenous species introductions (Paini et al. 2016).

Future trends in geographical distributions of non-indigenous species intrusions are likely to differ considerably from current patterns (*Section 2.5.1.3*) (Early et al. 2016). Although the level of non-indigenous species will remain high in developed countries in the coming decades, they will increase substantially in developing countries where biodiversity may be high but capacity to manage non-indigenous species is low. Developing countries, especially Sub-Saharan African countries, could be the most vulnerable to non-indigenous species expansion (Paini et al. 2016). In such places, non-indigenous species will increasingly threaten human livelihoods.

Water-borne infectious diseases are strongly associated with freshwater non-indigenous species that are linked to changes in environmental conditions produced by climate change (*Sections 5.2.3.3 and 5.2.3.4*). Some pathogens including West Nile Virus, dengue, yellow fever virus, chikungunya

fever virus, malaria sporozoan protists, filariasis and dirofilariasis nematodes, require aquatic arthropod vectors that are extending their range due to climate changes, at least on the northern rim of the Mediterranean (*Section 5.2.5.4*).

The number of non-indigenous plants (Doblas-Miranda et al. 2017) in the Mediterranean Basin seems to be lower than in other European regions (Vilà et al. 2007; Gassó et al. 2012), probably due to environmental constraints, the long history of anthropogenic disturbances and the lower economic development of the region until recently (Castrì et al. 1990; Vilà and Pujadas 2001). With regard to non-indigenous, the first vertebrates established in the Mediterranean Basin date back from the Neolithic period, although there has been an extraordinary increase in the rate of introduction of non-indigenous species since 1850 and especially in recent decades (Genovesi et al. 2009). Establishment success seems to be higher than in other Mediterranean-type climate regions of the world, at least for birds (Kark and Sol 2005). However, information related to non-native terrestrial invertebrates is largely unknown (Roques et al. 2009).

Introduction patterns of non-indigenous species differ considerably amongst groups, although they tend to mostly occupy anthropogenically modified habitats (*Section 2.5.2.1*), while contrary to other regions of the world, natural and semi-natural woody habitats are relatively resistant to non-indigenous species (Vilà et al. 2007; Kark et al. 2009; Roques et al. 2009; Arianoutsou et al. 2010). As in other regions of the world, the increase in the establishment of non-indigenous species in the Mediterranean Basin will continue due to the increasing rate of transport of goods and people. Delays in the management response therefore suggest that non-indigenous species will become of even greater concern in the future. Currently, the information available on non-indigenous species in the Basin is not complete and the number of non-indigenous species across taxonomic groups is underestimated (DAISIE 2009). Detailed information on their distribution and ecological

impacts is necessary to accurately determine the current status of non-indigenous species in the Mediterranean region.

The ecological and economic consequences of non-indigenous species introductions in terrestrial ecosystems of the Mediterranean Basin are beginning to emerge. Non-indigenous plants compete with indigenous species, decreasing local diversity and changing community composition (Vilà et al. 2006). Changes in ecosystem functioning have been less explored but include alterations in decomposition rates (Castro-Díez et al. 2009) and changes in soil carbon and nitrogen pools (Vilà et al. 2006). Even though the number of successful non-indigenous species seems to be higher in plants, the impacts of non-indigenous animals are not of lower magnitude. The presence of non-indigenous vertebrates poses severe threats to native biodiversity through competition for resources, predation and hybridization with native species, and economic impacts mainly through crop damage (Genovesi et al. 2009). Besides the lack of knowledge on the number of non-indigenous terrestrial invertebrates present in the Mediterranean Basin, most species established in Europe are known to be potential pests for agriculture and forestry products, while around 7% affect human and animal health (Roques et al. 2009). Their ecological consequences have received minor attention, although certain non-indigenous insect predators, such as *Linepithema humile* or *Harmonia axyridis*, are known to have a dramatic effect on native invertebrate communities (Angulo et al. 2011; Roy et al. 2011a, 2011b).

The Mediterranean Sea has a long history of anthropogenic activity and introduction of non-indigenous species and currently has a large number of them (Section 2.5.1). In recent years, the expansion of non-indigenous thermophilic species (that originally began started to enter the Mediterranean from the Indo-Pacific region during the 20th century) has been linked to climate-driven hydrographic changes. In the Mediterranean, non-indigenous thermophilic biota used to be restricted to the Levantine Basin, but are now found in the central and western basins (Occhipinti-Ambrogi and Galil 2010). The speed at which non-indigenous species are spreading in the Mediterranean Sea due to climate change is much faster than the actual increase in temperature, which is a great threat jeopardizing the future of biodiversity in the Mediterranean Sea (Raitsos et al. 2010).

Biodiversity hotspots are highly vulnerable to non-indigenous species given that many of the

nations that harbor them have low management capacity (Early et al. 2016). This is likely to be the case in eastern Mediterranean countries that have experienced a 150% increase in the mean annual rate of species introductions since 1924. Studies of long-term data since 1924 of 149 warm-water non-indigenous species in the Mediterranean Sea show that the Lessepsian introductions has been amplified by the warming of the eastern Mediterranean Sea (Raitsos et al. 2010).

The freshwater ecosystems of the Mediterranean Basin are considered a biodiversity hotspot with a high level of endemism and small natural ranges of native fish vulnerable to extinction (Ribeiro and Leunda 2012). Aquatic non-indigenous species have the potential to cause cascading disruption in entire food webs, cause biodiversity loss and do economic harm (Thomaz et al. 2014). The spreading of non-indigenous species in Iberian Peninsula freshwater rivers is a potent threat to native freshwater populations. Studies in the southwestern Iberian Peninsula freshwater rivers show that the quantities of non-indigenous species were the best forecaster of the decline of native fish species (Hermoso et al. 2011). In addition, the risk of exotic pathogens is threatening European Mediterranean countries through a continued introduction of non-indigenous disease vectors and changing climate and environments (Medlock et al. 2012) (Section 2.5.2.3).

STAGE OF INTRODUCTION	STRATEGY
ARRIVAL	<ul style="list-style-type: none"> <li>• Risk Analysis</li> <li>• International Standards</li> <li>• Inspection</li> </ul>
ESTABLISHMENT	<ul style="list-style-type: none"> <li>• Detection</li> <li>• Eradication</li> </ul>
SPREAD	<ul style="list-style-type: none"> <li>• Quarantine</li> <li>• Barrier Zone</li> </ul>
IMPACT	<ul style="list-style-type: none"> <li>• Suppression</li> <li>• Adaptation</li> </ul>

**Table 6.1 | Overview of stages of non-indigenous species introduction and potential management strategies** (based on Lockwood et al. 2007; Tobin 2018).

### 6.12.2 Management approaches, governance, and adaptation for non-indigenous species

Patterns of introduction, magnitude and expansion of non-indigenous species are currently at the most rapid rate of change ever recorded in

human history (Early et al. 2016). Only a minority of non-indigenous species succeed in establishing in their new locations and become a threat but those that do can result in billions of dollars in costs (Tobin 2018). As a result, management strategies continue to be an important element in global discussions on non-indigenous species. Central to best practice efforts in developing and implementing management frameworks is assessing the introduction stage of the species being addressed to identify the appropriate strategy (Table 6.1).

### 6.12.3 Innovation

Effective management strategies often involve preventing the arrival of non-indigenous species from the onset. Advances in risk analysis have led

to refined estimates of likely introduction pathways and the time at which the pathway is most likely to result in successful establishment (Gray 2016). This has led to more optimized allocation of limited inspection resources. Other advances in risk analysis include use of new technologies for detection and surveillance of non-indigenous species such as eDNA (Valentin et al. 2018) and utilizing bioeconomic models to formally consider ecological and economic links and dynamics that allow us to assess the costs of different management strategies (Lodge et al. 2016; Epanchin-Niell 2017). Finally, models of non-indigenous species distribution developed on their biological characteristics and climate suitability can potentially be used to predict susceptible areas (Mainali et al. 2015; Barbet-Massin et al. 2018).

## 6.13 Interactions of hazards, synergies and trade-offs between adaptation strategies and mitigation

The previous sections present the risks of the main hazards in the Mediterranean region, which are expected to increase in the future due to changes in environmental and societal conditions. Each section analyzes these hazards in isolation, without considering potential interactions. However, when two or more hazards occur at the same time, for example heavy precipitation coinciding with storm surge flooding, potential impacts increase due to compounding effects (Zscheischler et al. 2018), even in cases when none of the individual events is extreme. Also, cascading effects of hazards occurring in succession and overlapping temporally or spatially (de Ruiter et al. 2020), such as heavy precipitation triggering landslides, can lead to increased impacts (Gallina et al. 2016; Terzi et al. 2019). To cope with the impacts of compound and consecutive events, a holistic approach to future risk is needed that considers the interaction between hazards and identifies management and adaptation practices that can be successful in coping with a wide range of hazards. Such approaches build socio-ecological resilience, preparing society for future environmental change in a sustainable manner.

A large number of the management and adaptation measures discussed for a single hazard or sector present synergies with other hazards or sectors. For instance, the implementation of green roofs against heat stress (Section 6.2.2) additionally increases infiltration during flood events. Similarly,

managing agricultural drought by using agroforestry systems (Section 6.4.2) increases shade thus decreasing heat stress, decreases soil erosion due to a deeper penetration of roots, and has a positive effect on the water balance, which can counteract water scarcity. However, some strategies can lead to trade-offs with other hazards or sectors. While Ecosystem-based Adaptation (EbA) can be a successful strategy against sea-level rise-related hazards (Section 6.9.2) and can, at the same time, provide health benefits to the population, EbA measures have high space needs and are therefore only applicable to a limited degree in urban locations (Temmerman et al. 2013). Another example is the use of desalination plants for managing water scarcity (Section 6.4.2), which can lead to severe soil contamination. Examples of potential synergies and tradeoffs between adaptation measures are presented in Table 6.2.

The majority of strategies discussed above have positive effects on mitigation. Water-sensitive urban design (WSUD), sustainable land management and EbA, and other strategies have the potential to enhance CO<sub>2</sub> sequestration due to an increase in biomass. Such primarily nature-based strategies manage and protect ecosystems and their functions. Nature-based solutions can increase socio-ecological resilience in a wide range of contexts as these strategies, along with the concept of ecosystem services, further help to raise awareness regarding the importance of ecosys-



ADAPTATION STRATEGY	HAZARDS: SYNERGIES (+) & TRADE-OFFS (-)	SYNERGIES (+) & TRADE-OFFS (-) WITH MITIGATION
<b>URBAN PLANNING</b>		
Green roofs	+ Reduces heat stress + Increases infiltration during floods + Health benefits	+ Increases CO <sub>2</sub> sequestration in biomass
Increase in canopy cover in cities	+ Reduces heat stress + Increases infiltration during floods	+ Increases CO <sub>2</sub> sequestration in biomass
Water-sensitive urban design (WSUD), e.g., retention pools	+ Counteracts water scarcity + Counteracts salt water intrusion + Counteracts soil erosion + Increases infiltration during floods	+ Increases CO <sub>2</sub> sequestration due to more open/green space
Hard protection, e.g., sea walls	+ Protects from sea-level rise impacts - Potential increase in river/pluvial flood risk due to damming effects	- Energy intensive production
<b>NATURE-BASED SOLUTIONS</b>		
Conservation agriculture	+ Counteracts agricultural drought + Reduces soil erosion	+ Increases CO <sub>2</sub> sequestration in soils
Agroforestry systems	+ Counteracts agricultural drought + Shade reduces heat stress + Deeper penetration of roots counteracts soil erosion + Positive effect on water balance	+ Increases CO <sub>2</sub> sequestration in biomass
Sustainable land management, e.g., green cover	+ Counteracts soil erosion and desertification + Increases infiltration during floods + Increases water storage capacity	+ Increases CO <sub>2</sub> sequestration in biomass
Prescribed fire techniques	+ Reduce wildfire risk - Difficult to implement due to high amount of private property	+ Avoid large wildfires and so increase CO <sub>2</sub> sequestration potential in biomass
Reforestation in upstream areas	+ Reduces river flooding + Reduces soil erosion - Increases fuel biomass for wildfires	+ Increases CO <sub>2</sub> sequestration in biomass
Ecosystem-based Adaptation (EbA)	+ Protects from sea-level rise impacts + Health benefits - High space needs: applicable in selected locations only	+ Increases CO <sub>2</sub> sequestration in biomass
<b>ENGINEERED SOLUTIONS</b>		
Desalination of sea water	+ Counteracts water scarcity - Soil contamination	- Energy intensive process
<b>PUBLIC OUTREACH</b>		
Early warning systems (EWS), e.g., the European Flood Awareness System EFAS	+ Warn against multiple hazards, especially extremes, e.g., wildfires, coastal and river flooding, heat stress	
Awareness raising through ecosystem service assessment	+ Potential to reduce ocean acidification + Increases EbA via ecosystem conservation	- Increases CO <sub>2</sub> sequestration in biomass (if ecosystems conserved)

**Table 6.2 | Selected adaptation strategies** discussed in this chapter grouped by type of strategy, along with synergies and/or trade-offs with other hazards/sectors, and climate mitigation.

tems as an adaptation strategy, with positive effects on human well-being (Keesstra et al. 2018; Seddon et al. 2019). On the other hand, a number of adaptation strategies are energy-intensive and their implementation may lead to an increase in greenhouse gas emissions. Examples are the use of desalination plants or the construction of hard protection measures against sea-level rise.

The potential synergies and trade-offs of adaptation strategies with mitigation illustrate the importance of developing integrated policies for responding to future risks that incorporate adaptation and mitigation strategies (Section 5.1.3.1). This would allow synergies to be harnessed more strategically while, at the same time, avoiding potential trade-offs between mitigation and adaptation practices. In a study assessing adaptation and mitigation plans in European cities, Reckien et al. (2018) found that only a few Mediterranean cities have local climate plans that consider both mitigation and adaptation in a joint manner. These cities were primarily located in France, with a small number of cities in Spain. In most other European Mediterranean countries, the majority of cities have climate plans for mitigation only, very few for adaptation only, and some do not have any climate plans at all. Assuming that this finding can be transferred to southern and eastern Mediterranean countries, there is an urgent need for such local climate plans. Cities, in particular, need to become more resilient to environmental change as impacts will be disproportionately high in these locations due to the concentration of population and assets in combination with hazard-amplifying conditions (e.g., increased run-off through soil sealing, urban heat island effect (Rosenzweig et al. 2010).

A number of region-wide concerns and needs are raised across the chapter that, if addressed, can promote socio-ecological resilience and sustainable development in the entire region. Long-term monitoring data are missing in many parts of the basin, with particularly large differences in monitoring and reporting schemes between northern (EU), eastern, and southern countries of the region. There is also a need for advancing (climate) modeling techniques such as the representation of small river catchments, the short-term prediction of extreme events (e.g., heat, flooding), and improvements in seasonal forecasts. Furthermore, public participation in the development and implementation of management and adaptation strategies is important for their success. Stakeholders need to be involved in this process right from the start to increase local relevance and acceptance of the proposed strategies, thus facilitating imple-

mentation. Sharing and including local knowledge in the process is of prime importance in building a resilient society in a sustainable manner (Oppenheimer et al. 2019). Low-effort and low-cost strategies, e.g., promoting household-level adaptation, can play an important role in increasing resilience and coping with risk in the near future (Koerth et al. 2013b, 2013a).

Although national and local strategies are essential and successful in coping with risk and in increasing resilience, integrated management and adaptation approaches that treat multiple hazards in a holistic manner are required to address the above-stated concerns. Such approaches can be initiated in a top-down manner through region-wide policies such as the Barcelona Convention. The Barcelona Convention for the Protection of the Marine Environment and the Coastal Region of the Mediterranean, established in 1976, provides a suitable basis for devising Mediterranean-wide policies. It was updated in 1995 and sets the basis for the Mediterranean Action Plan that is part of the UNEP Regional Seas Programme. One of its goals is to promote integrated management of the Mediterranean coastal zone. Considerable efforts have been undertaken in recent years with the aim to facilitate basin-wide planning and management such as the Protocol on Integrated Coastal Zone Management in the Mediterranean (UNEP/MAP/PAP 2008), the Mediterranean Strategy for Sustainable Development 2016-2025 (UNEP/MAP 2016) and the Regional Climate Change Adaptation Framework for the Mediterranean Marine and Coastal Areas (UNEP/MAP 2017) (Section 5.1.1.2). These policy documents explicitly state the need for developing climate-resilient cities, acknowledging the importance of ecosystems for climate adaptation and mitigation, and enhancing regional and cross-border cooperation to promote sustainable development in the region (Benoit and Comeau 2005; UNEP/MAP 2012, 2016, 2017).

Active participation in regional-to-global initiatives and networks concerned with building socio-ecological resilience can be an additional important step forward. The "C40 Cities" network is concerned with achieving the goals of the Paris Agreement and currently has six members from the Mediterranean region (Barcelona, Rome, Venice, Athens, Istanbul, Tel Aviv). The "100 Resilient Cities" network aims to increase cities' resilience to a wide range of hazards, including drought, extreme heat, sea-level rise, but also other societal challenges such as corruption, demographic change, and poverty. Currently, ten Mediterranean cities are part of the network, including six from the

northern Mediterranean and four from the South and East. Such initiatives can foster knowledge exchange, provide funding for specific projects, and promote ambitious action against climate and environmental change.

Lastly, the transfer of scientific knowledge to policy-making needs to be facilitated, for instance with the help of policy briefs and so-called "resilience toolkits" (such as RISC-KIT for coastal resilience<sup>51</sup>) in order to support well-informed decisions. Similarly, knowledge transfer concerning environmental issues and sustainable development needs to be an integral part of the curriculum in primary and secondary education, therefore increasing awareness and establishing sustainable lifestyles as a social norm (Otto et al. 2020).

This chapter illustrates that future risks in the Mediterranean region will be determined by hazard characteristics (intensity and frequency) and by developments in socio-economic conditions that determine a society's adaptive capacity to cope with those hazards. The level of risk will largely depend on how soon and how effectively sustainable development is pursued. With the tourism sector being a large source of revenue in most parts of the region, transforming this sector will be particularly challenging. War and social unrest pose an additional, currently more pressing challenge in several countries in the Middle East and North Africa. These current developments may lead to a widening of the development gap between northern, southern, and eastern countries of the region. Therefore, developing joint, region-wide, and integrated management and adaptation approaches that treat multiple hazards in a holistic manner is of utmost importance for sustainable development in the entire region. Nonetheless, no one-size-fits-all strategy exists, and each measure needs to be tailored to the respective local conditions.

---

<sup>51</sup> [www.risckit.eu](http://www.risckit.eu)

## References

- Adams MA 2013 Mega-fires, tipping points and ecosystem services: Managing forests and woodlands in an uncertain future. *For. Ecol. Manage.* 294, 250–261. doi: [10.1016/j.foreco.2012.11.039](https://doi.org/10.1016/j.foreco.2012.11.039)
- Aguiar FC, Bentz J, Silva JMN, Fonseca AL, Swart R et al. 2018 Adaptation to climate change at local level in Europe: An overview. *Environ. Sci. Policy* 86, 38–63. doi: [10.1016/j.envsci.2018.04.010](https://doi.org/10.1016/j.envsci.2018.04.010)
- Aguilera E, Lassaletta L, Gattinger A, Gimeno BS 2013 Managing soil carbon for climate change mitigation and adaptation in Mediterranean cropping systems: A meta-analysis. *Agric. Ecosyst. Environ.* 168, 25–36. doi: [10.1016/j.agee.2013.02.003](https://doi.org/10.1016/j.agee.2013.02.003)
- Álvaro-Fuentes J, Arrúe JL, Gracia R, López MV 2007 Soil management effects on aggregate dynamics in semiarid Aragon (NE Spain). *Sci. Total Environ.* 378, 179–182. doi: [10.1016/j.scitotenv.2007.01.046](https://doi.org/10.1016/j.scitotenv.2007.01.046)
- Álvaro-Fuentes J, Arrúe JL, Gracia R, López M V. 2008 Tillage and cropping intensification effects on soil aggregation: Temporal dynamics and controlling factors under semiarid conditions. *Geoderma* 145, 390–396. doi: [10.1016/j.geoderma.2008.04.005](https://doi.org/10.1016/j.geoderma.2008.04.005)
- Amatulli G, Camia A, San-Miguel-Ayanz J 2013 Estimating future burned areas under changing climate in the EU-Mediterranean countries. *Sci. Total Environ.* 450–451, 209–222. doi: [10.1016/j.scitotenv.2013.02.014](https://doi.org/10.1016/j.scitotenv.2013.02.014)
- Amponsah W, Ayrat P-A, Boudevillain B, Bouvier C, Braud I et al. 2018 Integrated high-resolution dataset of high-intensity European and Mediterranean flash floods. *Earth Syst. Sci. Data* 10, 1783–1794. doi: [10.5194/essd-10-1783-2018](https://doi.org/10.5194/essd-10-1783-2018)
- Angelakis AN, Marecos Do Monte MHF, Bontoux L, Asano T 1999 The status of wastewater reuse practice in the Mediterranean basin: Need for guidelines. *Water Res.* 33, 2201–2217. doi: [10.1016/s0043-1354\(98\)00465-5](https://doi.org/10.1016/s0043-1354(98)00465-5)
- Angulo E, Caut S, Cerdá X 2011 Scavenging in Mediterranean ecosystems: Effect of the invasive Argentine ant. *Biol. Invasions* 13, 1183–1194. doi: [10.1007/s10530-011-9953-6](https://doi.org/10.1007/s10530-011-9953-6)
- Antalya Metropolitan Municipality 2018 Climate change risk analysis in Antalya: Project website. <http://antalyadeniziklim.org/en/project-aim/>
- Aragonés L, García-Barba J, García-Bleda E, López I, Serra JC 2015 Beach nourishment impact on *Posidonia oceanica*: Case study of Poniente Beach (Benidorm, Spain). *Ocean Eng.* 107, 1–12. doi: [10.1016/j.oceaneng.2015.07.005](https://doi.org/10.1016/j.oceaneng.2015.07.005)
- Archibald S 2016 Managing the human component of fire regimes: lessons from Africa. *Philos. Trans. R. Soc. B Biol. Sci.* 371. doi: [10.1098/rstb.2015.0346](https://doi.org/10.1098/rstb.2015.0346)
- Arianoutsou M, Delipetrou P, Celesti-Grapow L, Basnou C, Bazos I et al. 2010 Comparing naturalized alien plants and recipient habitats across an east-west gradient in the Mediterranean Basin. *J. Biogeogr.* 37, 1811–1823. doi: [10.1111/j.1365-2699.2010.02324.x](https://doi.org/10.1111/j.1365-2699.2010.02324.x)
- Armada E, Azcón R, López-Castillo OM, Calvo-Polanco M, Ruiz-Lozano JM 2015 Autochthonous arbuscular mycorrhizal fungi and *Bacillus thuringiensis* from a degraded Mediterranean area can be used to improve physiological traits and performance of a plant of agronomic interest under drought conditions. *Plant Physiol. Biochem.* 90, 64–74. doi: [10.1016/j.plaphy.2015.03.004](https://doi.org/10.1016/j.plaphy.2015.03.004)
- Azzurro E, Bariche M 2017 Local knowledge and awareness on the incipient lionfish invasion in the eastern Mediterranean Sea. *Mar. Freshw. Res.* 68, 1950. doi: [10.1071/MF16358](https://doi.org/10.1071/MF16358)
- Azzurro E, Sbragaglia V, Cerri J, Bariche M, Bolognini L et al. 2019 Climate change, biological invasions, and the shifting distribution of Mediterranean fishes: A large-scale survey based on local ecological knowledge. *Glob. Chang. Biol.* 25, 2779–2792. doi: [10.1111/gcb.14670](https://doi.org/10.1111/gcb.14670)
- Bagagiolo G, Biddoccu M, Rabino D, Cavallo E 2018 Effects of rows arrangement, soil management, and rainfall characteristics on water and soil losses in Italian sloping vineyards. *Environ. Res.* 166, 690–704. doi: [10.1016/j.envres.2018.06.048](https://doi.org/10.1016/j.envres.2018.06.048)
- Bahadur A, Batool A, Nasir F, Jiang S, Mingsen Q et al. 2019 Mechanistic Insights into Arbuscular Mycorrhizal Fungi-Mediated Drought Stress Tolerance in Plants. *Int. J. Mol. Sci.* 20, 4199. doi: [10.3390/ijms20174199](https://doi.org/10.3390/ijms20174199)
- Bakr J, Pék Z, Helyes L, Posta K 2018 Mycorrhizal Inoculation Alleviates Water Deficit Impact on Field-Grown Processing Tomato. *Polish J. Environ. Stud.* 27, 1949–1958. doi: [10.15244/pjoes/78624](https://doi.org/10.15244/pjoes/78624)
- Balouin Y, Longueville F, Colombet Y 2015 Video monitoring of soft coastal defenses at the Lido of Sète, France. *Ed. 3, Ferrare, Ital.* doi: [10.5150/cmcm.2015.038](https://doi.org/10.5150/cmcm.2015.038)
- Bandel T 2009 Capitalizing on the Competitive Advantage of Sustainable Agriculture in Egypt. Sekem and Soil & More – a partnership for sustainable development, in *Proceedings of the International Conference on Organic Agriculture and Climate Change* (Sofia, Bulgaria. 28–29 September 2009: Darko Znaor (ed)), 82–87.
- Barbet-Massin M, Rome Q, Villemant C, Courchamp F 2018 Can species distribution models really predict the expansion of invasive species? *PLoS One* 13. doi: [10.1371/journal.pone.0193085](https://doi.org/10.1371/journal.pone.0193085)
- Barceló D, Sabater S 2010 Water quality and assessment under scarcity: Prospects and challenges in Mediterranean watersheds. *J. Hydrol.* 383, 1–4. doi: [10.1016/j.jhydrol.2010.01.010](https://doi.org/10.1016/j.jhydrol.2010.01.010)
- Barcelona Sostenible 2015 Barcelona's commitment to the climate. Barcelona, Spain: Ajuntament de Barcelona
- Barea JM, Werner D, Azcón-Guilera C, Azcón R 2005 Interactions of Arbuscular Mycorrhiza and Nitrogen-Fixing Symbiosis in Sustainable Agriculture, in *Nitrogen Fixation in Agriculture, Forestry, Ecology, and the Environment*, eds. Werner D, Newton WE (Dordrecht: Springer Netherlands), 199–222. doi: [10.1007/1-4020-3544-6\\_10](https://doi.org/10.1007/1-4020-3544-6_10)
- Battlori E, de Cáceres M, Brotons L, Ackerly DD, Moritz MA et al. 2017 Cumulative effects of fire and drought in Mediterranean ecosystems. *Ecosphere* 8, e01906. doi: [10.1002/ecs2.1906](https://doi.org/10.1002/ecs2.1906)

- Batllori E, Parisien M-A, Krawchuk MA, Moritz MA 2013 Climate change-induced shifts in fire for Mediterranean ecosystems. *Glob. Ecol. Biogeogr.* 22, 1118–1129. doi: [10.1111/geb.12065](https://doi.org/10.1111/geb.12065)
- Becchi C, Ortolani I, Muir A, Cannicci S 2014 The effect of breakwaters on the structure of marine soft-bottom assemblages: A case study from a North-Western Mediterranean basin. *Mar. Pollut. Bull.* 87, 131–139. doi: [10.1016/j.marpolbul.2014.08.002](https://doi.org/10.1016/j.marpolbul.2014.08.002)
- Behnke R, Mortimore M 2016 *The End of Desertification ?*. Springer Berlin Heidelberg. doi: [10.1007/978-3-642-16014-1](https://doi.org/10.1007/978-3-642-16014-1)
- Ben Rais Lasram F, Guilhaumon F, Albouy C, Somot S, Thuiller W et al. 2010 The Mediterranean Sea as a “cul-de-sac” for endemic fishes facing climate change. *Glob. Chang. Biol.* 16, 3223–3245. <http://www.documentation.ird.fr/hor/fdi:010054015> [Accessed March 25, 2019].
- Ben Rais Lasram F, Mouillot D 2009 Increasing southern invasion enhances congruence between endemic and exotic Mediterranean fish fauna. *Biol. Invasions* 11, 697. doi: [10.1007/s10530-008-9284-4](https://doi.org/10.1007/s10530-008-9284-4)
- Benoit G, Comeau A 2005 *A Sustainable Future for the Mediterranean*. London: Routledge doi: [10.4324/9781849770323](https://doi.org/10.4324/9781849770323)
- Bensoussan N, Cebrian E, Dominici JM, Kersting D, Kipson S et al. 2019a Using CMEMS and the Mediterranean Marine Protected Areas sentinel network to track ocean warming effects in coastal areas. *J. Oper. Oceanogr.* 12, S65–S73.
- Bensoussan N, Chiggiato J, Buongiorno Nardelli B, Pisano A, Garrabou J 2019b Insights on 2017 Marine Heat Waves in the Mediterranean Sea. In: Copernicus Marine Service Ocean State Report, Issue 3. *J. Oper. Oceanogr.* 12, S26–S30. doi: [10.1080/1755876X.2019.1633075](https://doi.org/10.1080/1755876X.2019.1633075)
- Bernués A, Ruiz R, Olaizola A, Villalba D, Casasús I 2011 Sustainability of pasture-based livestock farming systems in the European Mediterranean context: Synergies and trade-offs. *Livest. Sci.* 139, 44–57. doi: [10.1016/j.livsci.2011.03.018](https://doi.org/10.1016/j.livsci.2011.03.018)
- Bhaduri A, Bogardi J, Siddiqi A, Voigt H, Vörösmarty C et al. 2016 Achieving Sustainable Development Goals from a Water Perspective. *Front. Environ. Sci.* 4, 211. doi: [10.3389/fenvs.2016.00064](https://doi.org/10.3389/fenvs.2016.00064)
- Bielsa J, Cazcarro I 2014 Implementing Integrated Water Resources Management in the Ebro River Basin: From Theory to Facts. *Sustainability* 7, 441–464. doi: [10.3390/su7010441](https://doi.org/10.3390/su7010441)
- Blöschl G, Hall J, Viglione A, Perdigão RAP, Parajka J et al. 2019 Changing climate both increases and decreases European river floods. *Nature* 573, 108–111. doi: [10.1038/s41586-019-1495-6](https://doi.org/10.1038/s41586-019-1495-6)
- Bond NR, Burrows RM, Kennard MJ, Bunn SE 2019 Water Scarcity as a Driver of Multiple Stressor Effects, in *Multiple Stressors in River Ecosystems* [Elsevier], 111–129. doi: [10.1016/b978-0-12-811713-2.00006-6](https://doi.org/10.1016/b978-0-12-811713-2.00006-6)
- Borg JA, Gauci MJ, Magro M, Micallef MA 2006 Environmental monitoring at St. George’s Bay (Malta) in connection with beach replenishment works. Gozo, Malta.
- Borga M, Anagnostou EN, Blöschl G, Creutin J-D 2011 Flash flood forecasting, warning and risk management: the HYDRATE project. *Environ. Sci. Policy* 14, 834–844. doi: [10.1016/j.envsci.2011.05.017](https://doi.org/10.1016/j.envsci.2011.05.017)
- Borrelli P, Robinson DA, Fleischer LR, Lugato E, Ballabio C et al. 2017 An assessment of the global impact of 21<sup>st</sup> century land use change on soil erosion. *Nat. Commun.* 8, 2013. doi: [10.1038/s41467-017-02142-7](https://doi.org/10.1038/s41467-017-02142-7)
- Boudjemline F, Semar A 2018 Assessment and mapping of desertification sensitivity with MEDALUS model and GIS – Case study: basin of Hodna, Algeria. *J. Water L. Dev.* 36, 17–26. doi: [10.2478/jwld-2018-0002](https://doi.org/10.2478/jwld-2018-0002)
- Bouvier C, Balouin Y, Castelle B 2017 Video monitoring of sandbar-shoreline response to an offshore submerged structure at a microtidal beach. *Geomorphology* 295, 297–305. doi: [10.1016/j.geomorph.2017.07.017](https://doi.org/10.1016/j.geomorph.2017.07.017)
- Bouwer LM, Jonkman SN 2018 Global mortality from storm surges is decreasing. *Environ. Res. Lett.* 13, 14008. doi: [10.1088/1748-9326/aa98a3](https://doi.org/10.1088/1748-9326/aa98a3)
- Bouzig M, Colon-Gonzalez FJ, Lung T, Lake IR, Hunter PR 2014 Climate change and the emergence of vector-borne diseases in Europe: case study of dengue fever. *BMC Public Health* 14, 781.
- Bowman DMJS, Balch JK, Artaxo P, Bond WJ, Cochrane MA et al. 2011 The human dimension of fire regimes on Earth. *J. Biogeogr.* 38, 2223–2236. doi: [10.1111/j.1365-2699.2011.02595.x](https://doi.org/10.1111/j.1365-2699.2011.02595.x)
- Bowman DMJS, Moreira-Munoz A, Kolden CA, Chavez RO, Munoz AA et al. 2019 Human-environmental drivers and impacts of the globally extreme 2017 Chilean fires. *Ambio* 48, 350–362.
- Bradley C, Bowes MJ, Brils J, Friedrich J, Gault J et al. 2018 Advancing integrated research on European river-sea systems: The DANUBIUS-RI project. *Int. J. Water Resour. Dev.* 34, 888–899. doi: [10.1080/07900627.2017.1399107](https://doi.org/10.1080/07900627.2017.1399107)
- Brotos L, Aquilué N, de Cáceres M, Fortin MJ, Fall A 2013 How Fire History, Fire Suppression Practices and Climate Change Affect Wildfire Regimes in Mediterranean Landscapes. *PLoS One* 8, e62392. doi: [10.1371/journal.pone.0062392](https://doi.org/10.1371/journal.pone.0062392)
- Calatrava J, Barberá GG, Castillo VM 2011 Farming practices and policy measures for agricultural soil conservation in semi-arid Mediterranean areas: The case of the Guadalentín basin in southeast Spain. *L. Degrad. Dev.* 22, 58–69. doi: [10.1002/ldr.1013](https://doi.org/10.1002/ldr.1013)
- Calvo-Polanco M, Sánchez-Castro I, Cantos M, García JL, Azcón R et al. 2016 Effects of different arbuscular mycorrhizal fungal backgrounds and soils on olive plants growth and water relation properties under well-watered and drought conditions. *Plant. Cell Environ.* 39, 2498–2514. doi: [10.1111/pce.12807](https://doi.org/10.1111/pce.12807)
- Cameira M do R, Santos Pereira L 2019 Innovation Issues in Water, Agriculture and Food. *Water* 11, 1230. doi: [10.3390/w11061230](https://doi.org/10.3390/w11061230)
- Caminade C, Medlock JM, Ducheyne E, McIntyre KM, Leach

- S et al. 2012 Suitability of European climate for the Asian tiger mosquito *Aedes albopictus*: recent trends and future scenarios. *J. R. Soc. Interface* 9, 2708–2717. doi: [10.1098/rsif.2012.0138](https://doi.org/10.1098/rsif.2012.0138)
- Canu DM, Ghermandi A, Nunes PALD, Lazzari P, Cossarini G et al. 2015 Estimating the value of carbon sequestration ecosystem services in the Mediterranean Sea: An ecological economics approach. *Glob. Environ. Chang.* 32, 87–95. doi: [10.1016/j.gloenvcha.2015.02.008](https://doi.org/10.1016/j.gloenvcha.2015.02.008)
- Cardinael R, Mao Z, Prieto I, Stokes A, Dupraz C et al. 2015 Competition with winter crops induces deeper rooting of walnut trees in a Mediterranean alley cropping agroforestry system. *Plant Soil* 391, 219–235. doi: [10.1007/s11104-015-2422-8](https://doi.org/10.1007/s11104-015-2422-8)
- Carlson C, Barreteau O, Kirshen P, Foltz K 2015 Storm Water Management as a Public Good Provision Problem: Survey to Understand Perspectives of Low-Impact Development for Urban Storm Water Management Practices under Climate Change. *J. Water Resour. Plan. Manag.* 141, 4014080. doi: [10.1061/\(asce\)wr.1943-5452.0000476](https://doi.org/10.1061/(asce)wr.1943-5452.0000476)
- Carmona G, Varela-Ortega C, Bromley J 2011 The Use of Participatory Object-Oriented Bayesian Networks and Agro-Economic Models for Groundwater Management in Spain. *Water Resour. Manag.* 25, 1509–1524. doi: [10.1007/s11269-010-9757-y](https://doi.org/10.1007/s11269-010-9757-y)
- Castellnou M, Prat-Guitart N, Arilla E, Larranaga A, Nebot E et al. 2019 Empowering strategic decision-making for wildfire management: avoiding the fear trap and creating a resilient landscape. *Fire Ecol.* 15.
- Castri F, Hansen AJ, Debussche M 1990 *Biological Invasions in Europe and the Mediterranean Basin*. Dordrecht: Springer Netherlands.
- Castro-Díez P, González-Muñoz N, Alonso AM, Gallardo A, Poorter L 2009 Effects of exotic invasive trees on nitrogen cycling: A case study in Central Spain. *Biol. Invasions* 11, 1973–1986. doi: [10.1007/s10530-008-9374-3](https://doi.org/10.1007/s10530-008-9374-3)
- Ceglar A, Zampieri M, Toreti A, Dentener FJ 2019 Observed Northward Migration of Agro-Climatic Zones in Europe Will Further Accelerate Under Climate Change. *Earth's Futur.* 7, 1088–1101. doi: [10.1029/2019ef001178](https://doi.org/10.1029/2019ef001178)
- Celette F, Gaudin R, Gary C 2008 Spatial and temporal changes to the water regime of a Mediterranean vineyard due to the adoption of cover cropping. *Eur. J. Agron.* 29, 153–162. doi: [10.1016/j.eja.2008.04.007](https://doi.org/10.1016/j.eja.2008.04.007)
- Cherfouh R, Lucas Y, Derridj A, Merdy P 2018 Long-term, low technicality sewage sludge amendment and irrigation with treated wastewater under Mediterranean climate: Impact on agronomical soil quality. *Environ. Sci. Pollut. Res.* 25, 35571–35581. doi: [10.1007/s11356-018-3463-3](https://doi.org/10.1007/s11356-018-3463-3)
- Chergui B, Fahd S, Santos X, Pausas JG 2017 Socioeconomic Factors Drive Fire-Regime Variability in the Mediterranean Basin. *Ecosystems* 21, 619–628. doi: [10.1007/s10021-017-0172-6](https://doi.org/10.1007/s10021-017-0172-6)
- Cherlet M, Hutchinson C, Reynolds J, Hill J, Sommer S et al. 2018 *World atlas of desertification: Rethinking land degradation and sustainable land management*. Publications Office of the European Union. doi: [10.2788/21872](https://doi.org/10.2788/21872)
- Choukr-Allah R, Ragab R, Bouchaou L, Barceló D 2017 The Souss-Massa River Basin, Morocco. *Handb. Environ. Chem.*, 355. doi: [10.1007/978-3-319-51131-3](https://doi.org/10.1007/978-3-319-51131-3)
- Choukr-Allah R, Ragab R, Rodriguez-Clemente R 2012 *Integrated Water Resources Management in the Mediterranean Region*. Dordrecht: Springer Netherlands
- Ciavola P, Ferreira O, Haerens P, Van Koningsveld M, Armario C 2011 Storm impacts along European coastlines. Part 2: lessons learned from the MICORE project. *Environ. Sci. Policy* 14, 924–933. doi: [10.1016/j.envsci.2011.05.009](https://doi.org/10.1016/j.envsci.2011.05.009)
- Ciscar J-C, Soria A, Goodess C, Christensen O, Iglesias A et al. 2009 Climate change impacts in Europe. Final report of the PESETA research project. doi: [10.2791/32500](https://doi.org/10.2791/32500)
- City of Athens G 2017 Athens becomes the first city in Greece with an integrated climate action plan. [https://www.c40.org/blog\\_posts/athens-becomes-the-first-city-in-greece-with-an-integrated-climate-change-action-plan](https://www.c40.org/blog_posts/athens-becomes-the-first-city-in-greece-with-an-integrated-climate-change-action-plan)
- Colosio F, Abbiati M, Airoldi L 2007 Effects of beach nourishment on sediments and benthic assemblages. *Mar. Pollut. Bull.* 54, 1197–1206. doi: [10.1016/j.marpolbul.2007.04.007](https://doi.org/10.1016/j.marpolbul.2007.04.007)
- Cooper JAG, Pilkey OH 2012 *Pitfalls of Shoreline Stabilization*. Springer Netherlands. doi: [10.1007/978-94-007-4123-2](https://doi.org/10.1007/978-94-007-4123-2)
- Corral C, Berenguer M, Sempere-Torres D, Poletti L, Silvestro F et al. 2019 Comparison of two early warning systems for regional flash flood hazard forecasting. *J. Hydrol.* 572, 603–619. doi: [10.1016/j.jhydrol.2019.03.026](https://doi.org/10.1016/j.jhydrol.2019.03.026)
- Cortès M, Llasat MC, Gilabert J, Llasat-Botija M, Turco M et al. 2018 Towards a better understanding of the evolution of the flood risk in Mediterranean urban areas: the case of Barcelona. *Nat. Hazards* 93, 39–60. doi: [10.1007/s11069-017-3014-0](https://doi.org/10.1007/s11069-017-3014-0)
- Costafreda-Aumedes S, Comas C, Vega-Garcia C 2017 Human-caused fire occurrence modelling in perspective: A review. *Int. J. Wildl. Fire* 26, 983. doi: [10.1071/wf17026](https://doi.org/10.1071/wf17026)
- Coumou D, Rahmstorf S 2012 A decade of weather extremes. *Nat. Clim. Chang.* 2, 491–496. doi: [10.1038/nclimate1452](https://doi.org/10.1038/nclimate1452)
- Cowie AL, Orr BJ, Castillo Sanchez VM, Chasek P, Crossman ND et al. 2018 Land in balance: The scientific conceptual framework for Land Degradation Neutrality. *Environ. Sci. Policy* 79, 25–35. doi: [10.1016/j.envsci.2017.10.011](https://doi.org/10.1016/j.envsci.2017.10.011)
- Cowie AL, Waters CM, Garland F, Orgill SE, Baumber A et al. 2019 Assessing resilience to underpin implementation of Land Degradation Neutrality: A case study in the rangelands of western New South Wales, Australia. *Environ. Sci. Policy* 100, 37–46. doi: [10.1016/j.envsci.2019.06.002](https://doi.org/10.1016/j.envsci.2019.06.002)
- Cramer W, Guiot J, Fader M, Garrabou J, Gattuso J-P et al. 2018 Climate change and interconnected risks to sustainable development in the Mediterranean. *Nat. Clim. Chang.* 8, 972–980. doi: [10.1038/s41558-018-0299-2](https://doi.org/10.1038/s41558-018-0299-2)
- CVN 2019 MO.S.E. <https://www.mosevenezia.eu/>
- Daccache A, Ciurana JS, Rodríguez-Díaz JA, Knox JW 2014 Water and energy footprint of irrigated agriculture in the Mediterranean region. *Environ. Res. Lett.* 9, 124014. doi: [10.1088/1748-9326/9/12/124014](https://doi.org/10.1088/1748-9326/9/12/124014)
- DAISIE 2009 *Handbook of Alien Species in Europe*. Springer Netherlands. doi: [10.1007/978-1-4020-8280-1](https://doi.org/10.1007/978-1-4020-8280-1)

- Damm A, Köberl J, Pretenthaler F, Rogler N, Töglhofer C 2017 Impacts of +2 °C global warming on electricity demand in Europe. *Clim. Serv.* 7, 12–30. doi: [10.1016/j.cliser.2016.07.001](https://doi.org/10.1016/j.cliser.2016.07.001)
- Danovaro R, Nepote E, Martire M Lo, Ciotti C, de Grandis G et al. 2018 Limited impact of beach nourishment on macrofaunal recruitment/settlement in a site of community interest in coastal area of the Adriatic Sea (Mediterranean Sea). *Mar. Pollut. Bull.* 128, 259–266. doi: [10.1016/j.marpolbul.2018.01.033](https://doi.org/10.1016/j.marpolbul.2018.01.033)
- Darmaraki S, Somot S, Sevault F, Nabat P, Cabos Narvaez WD et al. 2019 Future evolution of Marine Heatwaves in the Mediterranean Sea. *Clim. Dyn.* 53, 1371–1392. doi: [10.1007/s00382-019-04661-z](https://doi.org/10.1007/s00382-019-04661-z)
- de Leo F, Besio G, Zolezzi G, Bezzi M, Floqi T et al. 2017 Coastal erosion triggered by political and socio-economical abrupt changes: The case of Lalzit Bay, Albania. *Coast. Eng. Proc.* 1, 13. doi: [10.9753/icce.v35.management.13](https://doi.org/10.9753/icce.v35.management.13)
- de Ruiter MC, Couasnon A, van den Homberg MJC, Daniell JE, Gill JC et al. 2020 Why We Can No Longer Ignore Consecutive Disasters. *Earth's Futur.* 8, e2019EF001425. doi: [10.1029/2019EF001425](https://doi.org/10.1029/2019EF001425)
- de Sario M, Katsouyanni K, Michelozzi P 2013 Climate change, extreme weather events, air pollution and respiratory health in Europe. *Eur. Respir. J.* 42, 826–843. doi: [10.1183/09031936.00074712](https://doi.org/10.1183/09031936.00074712)
- Deryng D, Elliott J, Folberth C, Müller C, Pugh TAM et al. 2016 Regional disparities in the beneficial effects of rising CO<sub>2</sub> concentrations on crop water productivity. *Nat. Clim. Chang.* 6, 786–790. doi: [10.1038/nclimate2995](https://doi.org/10.1038/nclimate2995)
- di Stefano C, Ferro V 2016 Establishing soil loss tolerance: an overview. *J. Agric. Eng.* 47, 127–133. doi: [10.4081/jae.2016.560](https://doi.org/10.4081/jae.2016.560)
- Doblas-Miranda E, Alonso R, Arnán X, Bermejo V, Brotons L et al. 2017 A review of the combination among global change factors in forests, shrublands and pastures of the Mediterranean Region: Beyond drought effects. *Glob. Planet. Change* 148, 42–54. doi: [10.1016/j.gloplacha.2016.11.012](https://doi.org/10.1016/j.gloplacha.2016.11.012)
- Doherty RM, Heal MR, O'Connor FM 2017 Climate change impacts on human health over Europe through its effect on air quality. *Environ. Heal.* 16, 118. doi: [10.1186/s12940-017-0325-2](https://doi.org/10.1186/s12940-017-0325-2)
- Drake N, Vafeidis AT 2004 Review of spatial and temporal methods for assessing land degradation in the Mediterranean. *Adv. Environ. Monit. Model.* 1, 1–52.
- Duane A, Aquilué N, Canelles Q, Morán-Ordóñez A, de Cáceres M et al. 2019 Adapting prescribed burns to future climate change in Mediterranean landscapes. *Sci. Total Environ.* 677, 68–83. doi: [10.1016/j.scitotenv.2019.04.348](https://doi.org/10.1016/j.scitotenv.2019.04.348)
- Dubrovský M, Hayes M, Duce P, Trnka M, Svoboda M et al. 2014 Multi-GCM projections of future drought and climate variability indicators for the Mediterranean region. *Reg. Environ. Chang.* 14, 1907–1919. doi: [10.1007/s10113-013-0562-z](https://doi.org/10.1007/s10113-013-0562-z)
- Dulčić J, Kraljevic M, Pallaoro A, Glamuzina B 2005 Unusual catch of bluefish *Pomatomus saltatrix* (Pomatomidae) in Tarska cove (northern Adriatic). *Cybium* 29, 207–208.
- Early R, Bradley BA, Duker JS, Lawler JJ, Olden JD et al. 2016 Global threats from invasive alien species in the twenty-first century and national response capacities. *Nat. Commun.* 7, 12485. doi: [10.1038/ncomms12485](https://doi.org/10.1038/ncomms12485)
- EEA 2019a Adaptation challenges and opportunities for the European energy system. Building a climate-resilient low-carbon energy system.
- EEA 2019b Climate change adaptation in the agriculture sector in Europe.
- Eekhout JPC, Vente J 2019 The implications of bias correction methods and climate model ensembles on soil erosion projections under climate change. *Earth Surf. Process. Landforms* 44, 1137–1147. doi: [10.1002/esp.4563](https://doi.org/10.1002/esp.4563)
- Efthymiadis D, Goodess CM, Jones PD 2011 Trends in Mediterranean gridded temperature extremes and large-scale circulation influences. *Nat. Hazards Earth Syst. Sci.* 11, 2199–2214. doi: [10.5194/nhess-11-2199-2011](https://doi.org/10.5194/nhess-11-2199-2011)
- Epanchin-Niell RS 2017 Economics of invasive species policy and management. *Biol. Invasions* 19, 3333–3354. doi: [10.1007/s10530-017-1406-4](https://doi.org/10.1007/s10530-017-1406-4)
- ERML 2012 Improved Understanding, Management and Monitoring in the Coastal Zone. Marine Resources and Coastal Zone Management Program Institute of the Environment – University of Balamand.
- Esteban-Talaya A, López Palomeque F, Pérez EA 2005 Impacts in the touristic sector, in *A Preliminary Assessment of the Impacts in Spain due to the Effect of Climate Change*, ed. Moreno JM (Madrid, Spain: Ministry of Environment), 653–690.
- European Parliament 2007 Directive 2007/60/EC of the European Parliament and of the Council of 23 October 2007 on the assessment and management of flood risks.
- Faccini F, Luino F, Paliaga G, Sacchini A, Turconi L et al. 2018 Role of rainfall intensity and urban sprawl in the 2014 flash flood in Genoa City, Bisagno catchment (Liguria, Italy). *Appl. Geogr.* 98, 224–241. doi: [10.1016/j.apgeog.2018.07.022](https://doi.org/10.1016/j.apgeog.2018.07.022)
- Fader M, Shi S, Von Bloh W, Bondeau A, Cramer W 2016 Mediterranean irrigation under climate change: More efficient irrigation needed to compensate for increases in irrigation water requirements. *Hydrol. Earth Syst. Sci.* 20, 953–973. doi: [10.5194/hess-20-953-2016](https://doi.org/10.5194/hess-20-953-2016)
- Famiglietti JS 2014 The global groundwater crisis. *Nat. Clim. Chang.* 4, 945. doi: [10.1038/nclimate2425](https://doi.org/10.1038/nclimate2425)
- FAO 2013 *Climate-smart agriculture sourcebook*. Rome: Food and Agriculture Organization of the United Nations.
- Fernandes PM 2018 Scientific support to prescribed underburning in southern Europe: What do we know? *Sci. Total Environ.* 630, 340–348.
- Fernández-Montblanc T, Vousdoukas MI, Ciavola P, Voukouvalas E, Mentaschi L et al. 2019 Towards robust pan-European storm surge forecasting. *Ocean Model.* 133, 129–144. doi: [10.1016/j.ocemod.2018.12.001](https://doi.org/10.1016/j.ocemod.2018.12.001)

- Fernandez Milan B, Creutzig F 2015 Reducing urban heat wave risk in the 21<sup>st</sup> century. *Curr. Opin. Environ. Sustain.* 14, 221–231. doi: [10.1016/j.cosust.2015.08.002](https://doi.org/10.1016/j.cosust.2015.08.002)
- Field CB, Barros VR, Mach KJ, Mastrandrea MD, van Aalst MK et al. 2014 Technical Summary, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 35–94. [https://www.ipcc.ch/pdf/assessment-report/ar5/wg2/WGIIAR5-TS\\_FINAL.pdf](https://www.ipcc.ch/pdf/assessment-report/ar5/wg2/WGIIAR5-TS_FINAL.pdf) [Accessed March 11, 2017].
- Fischer D, Thomas SM, Suk JE, Sudre B, Hess A et al. 2013 Climate change effects on Chikungunya transmission in Europe: geospatial analysis of vector's climatic suitability and virus' temperature requirements. *Int. J. Health Geogr.* 12, 51. doi: [10.1186/1476-072X-12-51](https://doi.org/10.1186/1476-072X-12-51)
- Fischer EM, Schär C 2010 Consistent geographical patterns of changes in high-impact European heatwaves. *Nat. Geosci.* 3, 398–403. doi: [10.1038/ngeo866](https://doi.org/10.1038/ngeo866)
- Flannigan MD, Krawchuk MA, Groot WJ de, Wotton BM, Gowman LM 2009 Implications of changing climate for global wildland fire. *Int. J. Wildl. Fire* 18, 483. doi: [10.1071/wf08187](https://doi.org/10.1071/wf08187)
- Flörke M, Schneider C, McDonald RI 2018 Water competition between cities and agriculture driven by climate change and urban growth. *Nat. Sustain.* 1, 51–58. doi: [10.1038/s41893-017-0006-8](https://doi.org/10.1038/s41893-017-0006-8)
- Forzieri G, Cescatti A, Batista e Silva F, Feyen L 2017 Increasing risk over time of weather-related hazards to the European population: a data-driven prognostic study. *Lancet Planet. Heal.* 1, e200–e208. doi: [10.1016/s2542-5196\(17\)30082-7](https://doi.org/10.1016/s2542-5196(17)30082-7)
- Foyer CH, Nguyen HT, Lam H-M 2018 A seed change in our understanding of legume biology from genomics to the efficient cooperation between nodulation and arbuscular mycorrhizal fungi. *Plant. Cell Environ.* 41, 1949–1954. doi: [10.1111/pce.13419](https://doi.org/10.1111/pce.13419)
- Fraga H, García de Cortázar Aatauri I, Malheiro AC, Santos JA 2016 Modelling climate change impacts on viticultural yield, phenology and stress conditions in Europe. *Glob. Chang. Biol.* 22, 3774–3788. doi: [10.1111/gcb.13382](https://doi.org/10.1111/gcb.13382)
- Future Earth 2019 Heatwave Issue Brief. <https://futureearth.org/publications/issue-briefs-2/heatwaves/>
- Gagliano A, Detommaso M, Nocera F, Berardi U 2016 The adoption of green roofs for the retrofitting of existing buildings in the Mediterranean climate. *Int. J. Sustain. Build. Technol. Urban Dev.* 7, 116–129. doi: [10.1080/2093761X.2016.1172279](https://doi.org/10.1080/2093761X.2016.1172279)
- Gallina V, Torresan S, Critto A, Sperotto A, Glade T et al. 2016 A review of multi-risk methodologies for natural hazards: Consequences and challenges for a climate change impact assessment. *J. Environ. Manage.* 168, 123–132. doi: [10.1016/j.jenvman.2015.11.011](https://doi.org/10.1016/j.jenvman.2015.11.011)
- García-Garizábal I, Causapé J, Abrahao R, Merchan D 2014 Impact of Climate Change on Mediterranean Irrigation Demand: Historical Dynamics of Climate and Future Projections. *Water Resour. Manag.* 28, 1449–1462. doi: [10.1007/s11269-014-0565-7](https://doi.org/10.1007/s11269-014-0565-7)
- García-Herrera R, Díaz J, Trigo RM, Luterbacher J, Fischer EM 2010 A Review of the European Summer Heat Wave of 2003. *Crit. Rev. Environ. Sci. Technol.* 40, 267–306. doi: [10.1080/10643380802238137](https://doi.org/10.1080/10643380802238137)
- García-Orenes F, Roldán A, Mataix-Solera J, Cerdà A, Campoy M et al. 2012 Soil structural stability and erosion rates influenced by agricultural management practices in a semi-arid Mediterranean agro-ecosystem. *Soil Use Manag.* 28, 571–579. doi: [10.1111/j.1475-2743.2012.00451.x](https://doi.org/10.1111/j.1475-2743.2012.00451.x)
- García-Ruiz JM, Nadal-Romero E, Lana-Renault N, Beguería S 2013 Erosion in Mediterranean landscapes: Changes and future challenges. *Geomorphology* 198, 20–36. doi: [10.1016/j.geomorph.2013.05.023](https://doi.org/10.1016/j.geomorph.2013.05.023)
- García-Tejero IF, Carbonell R, Ordoñez R, Torres FP, Durán Zuazo VH 2020 Conservation agriculture practices to improve the soil water management and soil carbon storage in Mediterranean rainfed agro-ecosystems, in *Soil Health Restoration and Management*, ed. Meena RS (Singapore: Springer Singapore), 203–230. doi: [10.1007/978-981-13-8570-4\\_6](https://doi.org/10.1007/978-981-13-8570-4_6)
- Gardiner EP, Herring DD, Fox JF 2019 The U.S. Climate Resilience Toolkit: Evidence of progress. *Clim. Change* 153, 477–490. doi: [10.1007/s10584-018-2216-0](https://doi.org/10.1007/s10584-018-2216-0)
- Garrabou J, Bensoussan N, Azzurro E 2019a Monitoring Climate-related responses in Mediterranean Marine Protected Areas and beyond: Five Standard Protocols. doi: [10.20350/digitalCSIC/8612](https://doi.org/10.20350/digitalCSIC/8612)
- Garrabou J, Gómez-Gras D, Ledoux J-B, Linares C, Bensoussan N et al. 2019b Collaborative Database to Track Mass Mortality Events in the Mediterranean Sea. *Front. Mar. Sci.* 6, 707. doi: [10.3389/fmars.2019.00707](https://doi.org/10.3389/fmars.2019.00707)
- Gasparrini A, Guo Y, Sera F, Vicedo-Cabrera AM, Huber V et al. 2017 Projections of temperature-related excess mortality under climate change scenarios. *Lancet Planet. Heal.* 1, e360–e367. doi: [10.1016/S2542-5196\(17\)30156-0](https://doi.org/10.1016/S2542-5196(17)30156-0)
- Gassó N, Thuiller W, Pino J, Vilà M 2012 Potential distribution range of invasive plant species in Spain. *Neobiota* 12, 25–40. doi: [10.3897/neobiota.12.2341](https://doi.org/10.3897/neobiota.12.2341)
- Gaume E, Borga M, Llasat MC, Maouche S, Lang M et al. 2016 Mediterranean extreme floods and flash floods, in *The Mediterranean Region under Climate Change. A Scientific Update Coll. Synthèses.*, eds. Thiébaud S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 133–144. <https://hal.archives-ouvertes.fr/hal-01465740>
- Genovesi P, Bacher S, Kobelt M, Pascal M, Scalera R 2009 Alien mammals of Europe, in *Handbook of alien species in Europe*, ed. DAISIE (Springer Netherlands), 119–128. doi: [10.1007/978-1-4020-8280-1\\_9](https://doi.org/10.1007/978-1-4020-8280-1_9)
- Giambastiani BMS, Greggio N, Sistilli F, Fabbri S, Scarelli F et al. 2016 RIGED-RA project-Restoration and manage-



- ment of Coastal Dunes in the Northern Adriatic Coast, Ravenna Area-Italy. *IOP Conf. Ser. Earth Environ. Sci.* 44, 1–7. doi: [10.1088/1755-1315/44/5/052038](https://doi.org/10.1088/1755-1315/44/5/052038)
- Giannakopoulos C, Le Sager P, Bindi M, Moriondo M, Kostopoulou E et al. 2009 Climatic changes and associated impacts in the Mediterranean resulting from a 2°C global warming. *Glob. Planet. Change* 68, 209–224. doi: [10.1016/j.gloplacha.2009.06.001](https://doi.org/10.1016/j.gloplacha.2009.06.001)
- Givan O, Edelist D, Sonin O, Belmaker J 2017 Thermal affinity as the dominant factor changing Mediterranean fish abundances. *Glob. Chang. Biol.* 24, e80–e89. doi: [10.1111/gcb.13835](https://doi.org/10.1111/gcb.13835)
- Gómez JA, Llewellyn C, Basch G, Sutton PB, Dyson JS et al. 2011 The effects of cover crops and conventional tillage on soil and runoff loss in vineyards and olive groves in several Mediterranean countries. *Soil Use Manag.* 27, 502–514. doi: [10.1111/j.1475-2743.2011.00367.x](https://doi.org/10.1111/j.1475-2743.2011.00367.x)
- Gössling S, Peeters P, Hall CM, Ceron J-P, Dubois G et al. 2012 Tourism and water use: Supply, demand, and security. An international review. *Tour. Manag.* 33, 1–15. doi: [10.1016/j.tourman.2011.03.015](https://doi.org/10.1016/j.tourman.2011.03.015)
- Goulden S, Portman ME, Carmon N, Alon-Mozes T 2018 From conventional drainage to sustainable stormwater management: Beyond the technical challenges. *J. Environ. Manage.* 219, 37–45. doi: [10.1016/j.jenvman.2018.04.066](https://doi.org/10.1016/j.jenvman.2018.04.066)
- Govaerts B, Verhulst N, Castellanos-Navarrete A, Sayre KD, Dixon J et al. 2009 Conservation Agriculture and Soil Carbon Sequestration: Between Myth and Farmer Reality. *CRC. Crit. Rev. Plant Sci.* 28, 97–122. doi: [10.1080/07352680902776358](https://doi.org/10.1080/07352680902776358)
- Gray DR 2016 Risk Reduction of an Invasive Insect by Targeting Surveillance Efforts with the Assistance of a Phenology Model and International Maritime Shipping Routes and Schedules. *Risk Anal.* 36, 914–925. doi: [10.1111/risa.12474](https://doi.org/10.1111/risa.12474)
- Grbec B, Dulcic J, Morovic M 2002 Long-term changes in landings of small pelagic fish in the eastern Adriatic—possible influence of climate oscillations over the Northern Hemisphere. *Clim. Res.* 20, 241–252. doi: [10.3354/cr020241](https://doi.org/10.3354/cr020241)
- Gudmundsson L, Rego FC, Rocha M, Seneviratne SI 2014 Predicting above normal wildfire activity in southern Europe as a function of meteorological drought. *Environ. Res. Lett.* 9, 84008. doi: [10.1088/1748-9326/9/8/084008](https://doi.org/10.1088/1748-9326/9/8/084008)
- Guerreiro SB, Dawson RJ, Kilsby C, Lewis E, Ford A 2018 Future heat-waves, droughts and floods in 571 European cities. *Environ. Res. Lett.* 13, 34009. doi: [10.1088/1748-9326/aaaad3](https://doi.org/10.1088/1748-9326/aaaad3)
- Guiot J, Cramer W 2016 Climate change: The 2015 Paris Agreement thresholds and Mediterranean basin ecosystems. *Science (80-. ).* 354, 4528–4532. doi: [10.1126/science.aah5015](https://doi.org/10.1126/science.aah5015)
- Guo Y, Peng C, Zhu Q, Wang M, Wang H et al. 2019 Modelling the impacts of climate and land use changes on soil water erosion: Model applications, limitations and future challenges. *J. Environ. Manage.* 250, 109403. doi: [10.1016/j.jenvman.2019.109403](https://doi.org/10.1016/j.jenvman.2019.109403)
- Hadjikakou M, Chenoweth J, Miller G 2013 Estimating the direct and indirect water use of tourism in the eastern Mediterranean. *J. Environ. Manage.* 114, 548–556. doi: [10.1016/j.jenvman.2012.11.002](https://doi.org/10.1016/j.jenvman.2012.11.002)
- Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M et al. 2008 Seawater carbonate chemistry in Ischia, Italy, 2008. doi: [10.1594/pangaea.819633](https://doi.org/10.1594/pangaea.819633)
- Hamm L, Hanson H, Capobianco M, Dette HH, Lechuga A et al. 1998 Beach Fills in Europe – Projects, Practices, and Objectives, in *26th International Conference on Coastal Engineering*, ed. Edge BL, 3060–3073. doi: [10.1061/9780784404119.232](https://doi.org/10.1061/9780784404119.232)
- Hanson H, Brampton A, Capobianco M, Dette HH, Hamm L et al. 2002 Beach nourishment projects, practices, and objectives—a European overview. *Coast. Eng.* 47, 81–111. doi: [10.1016/s0378-3839\(02\)00122-9](https://doi.org/10.1016/s0378-3839(02)00122-9)
- Harley MD, Armaroli C, Ciavola P 2011 Evaluation of XBeach predictions for a real-time warning system in Emilia-Romagna, Northern Italy. *J. Coast. Res.*, 1861–1865.
- Haro-Monteagudo D, Daccache A, Knox JW 2017 Exploring the utility of drought indicators to assess climate risks to agricultural productivity in a humid climate. *Hydrol. Res.* 49, 539–551. doi: [10.2166/nh.2017.010](https://doi.org/10.2166/nh.2017.010)
- Heaviside C, Tsangari H, Paschalidou A, Vardoulakis S, Kasomenos P et al. 2016 Heat-related mortality in Cyprus for current and future climate scenarios. *Sci. Total Environ.* 569–570, 627–633. doi: [10.1016/j.scitotenv.2016.06.138](https://doi.org/10.1016/j.scitotenv.2016.06.138)
- Hein L, Metzger MJ, Moreno A 2009 Potential impacts of climate change on tourism; a case study for Spain. *Curr. Opin. Environ. Sustain.* 1, 170–178. doi: [10.1016/j.cosust.2009.10.011](https://doi.org/10.1016/j.cosust.2009.10.011)
- Hermoso V, Clavero M, Blanco-Garrido F, Prenda J 2011 Invasive species and habitat degradation in Iberian streams: an analysis of their role in freshwater fish diversity loss. *Ecol. Appl.* 21, 175–188. doi: [10.1890/09-2011.1](https://doi.org/10.1890/09-2011.1)
- Hertig E 2019 Distribution of Anopheles vectors and potential malaria transmission stability in Europe and the Mediterranean area under future climate change. *Parasit. Vectors* 12, 18. doi: [10.1186/s13071-018-3278-6](https://doi.org/10.1186/s13071-018-3278-6)
- Hidalgo M, Ligas A, Bellido JM, Bitetto I, Carbonara P et al. 2020 Size-dependent survival of European hake juveniles in the Mediterranean Sea. *Sci. Mar.* 83, 207. doi: [10.3989/scimar.04857.16a](https://doi.org/10.3989/scimar.04857.16a)
- Hinkel J, Brown S, Exner L, Nicholls RJ, Vafeidis AT et al. 2012 Sea-level rise impacts on Africa and the effects of mitigation and adaptation: An application of DIVA. *Reg. Environ. Chang.* 12, 207–224. doi: [10.1007/s10113-011-0249-2](https://doi.org/10.1007/s10113-011-0249-2)
- Hinkel J, Nicholls RJ, Vafeidis AT, Tol RSJ, Avagianou T 2010 Assessing risk of and adaptation to sea-level rise in the European Union: An application of DIVA. *Mitig. Adapt. Strateg. Glob. Chang.* 15, 703–719. doi: [10.1007/s11027-010-9237-y](https://doi.org/10.1007/s11027-010-9237-y)
- Hirschi M, Seneviratne SI, Alexandrov V, Boberg F, Boroneant C et al. 2011 Observational evidence for soil-moisture impact on hot extremes in southeastern Europe. *Nat.*

- Geosci.* 4, 17–21. doi: [10.1038/ngeo1032](https://doi.org/10.1038/ngeo1032)
- Holman IP, Brown C, Carter TR, Harrison PA, Rounsevell M 2018 Improving the representation of adaptation in climate change impact models. *Reg. Environ. Chang.* doi: [10.1007/s10113-018-1328-4](https://doi.org/10.1007/s10113-018-1328-4)
- Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD et al. 2017 Global warming and recurrent mass bleaching of corals. *Nature* 543, 373–377. doi: [10.1038/nature21707](https://doi.org/10.1038/nature21707)
- Iglesias A, Garrote L, Quiroga S, Moneo M 2012 A regional comparison of the effects of climate change on agricultural crops in Europe. *Clim. Change* 112, 29–46. doi: [10.1007/s10584-011-0338-8](https://doi.org/10.1007/s10584-011-0338-8)
- Imen T, Souissi R 2018 Coastal erosion in the south-eastern Mediterranean: case of beaches in North Tunisia. *Arab. J. Geosci.* 11. doi: [10.1007/s12517-018-3716-y](https://doi.org/10.1007/s12517-018-3716-y)
- IPCC 2012 *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change.*, eds. Field CB, Barros V, Stocker TF, Qin D, Dokken DJ et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- Iskander MM, Frihy OE, El Ansary AE, Abd El Mooty MM, Nagy HM 2007 Beach impacts of shore-parallel breakwaters backing offshore submerged ridges, Western Mediterranean Coast of Egypt. *J. Environ. Manage.* 85, 1109–1119. doi: [10.1016/j.jenvman.2006.11.018](https://doi.org/10.1016/j.jenvman.2006.11.018)
- IWA 2012 Long-term master plan for the national water sector: Part A. Israel Water Authority.
- Jackson LP, Jevrejeva S 2016 A probabilistic approach to 21<sup>st</sup> century regional sea-level projections using RCP and High-end scenarios. *Glob. Planet. Change* 146, 179–189. doi: [10.1016/j.gloplacha.2016.10.006](https://doi.org/10.1016/j.gloplacha.2016.10.006)
- Jacob D, Petersen J, Eggert B, Alias A, Christensen OB et al. 2014 EURO-CORDEX: new high-resolution climate change projections for European impact research. *Reg. Environ. Chang.* 14, 563–578. doi: [10.1007/s10113-013-0499-2](https://doi.org/10.1007/s10113-013-0499-2)
- Jevrejeva S, Jackson LP, Riva REM, Grinsted A, Moore JC 2016 Coastal sea level rise with warming above 2 °C. *Proc. Natl. Acad. Sci. U. S. A.* 113, 13342–13347. doi: [10.1073/pnas.1605312113](https://doi.org/10.1073/pnas.1605312113)
- Joffre RR, Rambal S 1993 How Tree Cover Influences the Water Balance of Mediterranean Rangelands. *Ecology* 74, 570–582. doi: [10.2307/1939317](https://doi.org/10.2307/1939317)
- Joffre RR, Rambal S 2006 Tree-grass interactions in the south-western Iberian Peninsula dehesas and montados. *Sécheresse* 17, 340–342.
- Jones PD, Lister DH, Osborn TJ, Harpham C, Salmon M et al. 2012 Hemispheric and large-scale land surface air temperature variations: an extensive review and an update to 2010. *JGR Atmos.* 117, D05127. doi: [10.1029/2011JD017139](https://doi.org/10.1029/2011JD017139)
- Kaimaki PS, Gkouvatso E, Fyllas N, Chistopoulou A, Papanousi F 2014 The Cyprus climate change risk assessment: Evidence report. Nicosia, Cyprus: Ministry of Agriculture, Rural Development and Environment, Department of Environment, Cyprus Government.
- Kalavrouziotis IK, Koukoulakis PH, Drakatos PA 2015 Water and wastewater management in antiquity in the context of an ethically oriented environmental protection. *Int. J. Glob. Environ. Issues* 14, 226. doi: [10.1504/ijgenvi.2015.071847](https://doi.org/10.1504/ijgenvi.2015.071847)
- Kamel A, Ali H, Ghaleb F, Mario M, Tony G 2015 GIS-based mapping of areas sensitive to desertification in a semi-arid region in Lebanon. *South-Eastern Eur. J. Earth Obs. Geomatics* 4, 91–103.
- Kark S, Sol D 2005 Establishment Success across Convergent Mediterranean Ecosystems: An Analysis of Bird Introductions. *Conserv. Biol.* 19, 1519–1527. doi: [10.1111/j.1523-1739.2005.004365.x](https://doi.org/10.1111/j.1523-1739.2005.004365.x)
- Kark S, Solarz W, Chiron F, Clergeau P, Shirley S 2009 Alien Birds, Amphibians and Reptiles of Europe, in *Handbook of Alien Species in Europe*, ed. DAISIE (Springer Netherlands), 105–118. doi: [10.1007/978-1-4020-8280-1\\_8](https://doi.org/10.1007/978-1-4020-8280-1_8)
- Kassam A, Friedrich T, Derpsch R, Lahmar R, Mrabet R et al. 2012 Conservation agriculture in the dry Mediterranean climate. *F. Crop. Res.* 132, 7–17. doi: [10.1016/j.fcr.2012.02.023](https://doi.org/10.1016/j.fcr.2012.02.023)
- Katzan J, Owsianowski S 2017 Protecting health from heat stress in informal settlements of the Greater Cairo Region. Bonn, Germany: GIZ, Federal Ministry for Economic Cooperation and Development.
- Keeley JE, Bond WJ, Bradstock RA, Pausas JG, Rundel PW 2012a *Fire in Mediterranean ecosystems: Ecology, evolution and management.* Cambridge, UK: Cambridge University Press. doi: [10.1017/cbo9781139033091](https://doi.org/10.1017/cbo9781139033091)
- Keeley JE, Fotheringham C, Rundel P 2012b Postfire Chaparral Regeneration Under Mediterranean and Non-Mediterranean Climates. *Madroño*, 109–127.
- Keesstra SD, Nunes JP, Novara A, Finger D, Avelar D et al. 2018 The superior effect of nature based solutions in land management for enhancing ecosystem services. *Sci. Total Environ.* 610–611, 997–1009. doi: [10.1016/j.scitotenv.2017.08.077](https://doi.org/10.1016/j.scitotenv.2017.08.077)
- Kelly RP, Foley MM, Fisher WS, Feely RA, Halpern BS et al. 2011 Oceans. Mitigating local causes of ocean acidification with existing laws. *Science (80-. )*. 332, 1036–1037. doi: [10.1126/science.1203815](https://doi.org/10.1126/science.1203815)
- Kendrovski V, Baccini M, Martinez G, Wolf T, Paunovic E et al. 2017 Quantifying Projected Heat Mortality Impacts under 21<sup>st</sup>-Century Warming Conditions for Selected European Countries. *Int. J. Environ. Res. Public Health* 14, 729. doi: [10.3390/ijerph14070729](https://doi.org/10.3390/ijerph14070729)
- Kepner WG, Rubio JL, Mouat DA, Pedrazzini F 2006 Desertification in the Mediterranean Region. A Security Issue. in *Proceedings of the NATO Mediterranean Dialogue Workshop held in Valencia, Spain, 2-5 December 2003* (Springer Science and Business Media).
- Khadra R, Sagardoy JA 2019 *Irrigation Governance Challenges in the Mediterranean Region: Learning from Experiences and Promoting Sustainable Performance.* Cham: Springer International Publishing.
- Kirkby MJ, Jones RJA, Irvine B, Gobin A, Govers G et al. 2004

- Pan-European soil erosion risk assessment: The PE-SERA Map. [https://publications.jrc.ec.europa.eu/repository/bitstream/JRC27736/EUR\\_21176\\_EN.pdf](https://publications.jrc.ec.europa.eu/repository/bitstream/JRC27736/EUR_21176_EN.pdf)
- Knorr W, Arneith A, Jiang L 2016 Demographic controls of future global fire risk. *Nat. Clim. Chang.* doi: [10.1038/nclimate2999](https://doi.org/10.1038/nclimate2999)
- Koerth J, Jones N, Vafeidis AT, Dimitrakopoulos PG, Melliou A et al. 2013a Household adaptation and intention to adapt to coastal flooding in the Axios – Loudias – Aliakmonas National Park, Greece. *Ocean Coast. Manag.* 82, 43–50. doi: [10.1016/j.ocecoaman.2013.05.008](https://doi.org/10.1016/j.ocecoaman.2013.05.008)
- Koerth J, Vafeidis AT, Hinkel J, Sterr H 2013b What motivates coastal households to adapt pro-actively to sea-level rise and increasing flood risk? *Reg. Environ. Chang.* 13, 897–909. doi: [10.1007/s10113-012-0399-x](https://doi.org/10.1007/s10113-012-0399-x)
- Kok K, Pedde S, Gramberger M, Harrison PA, Holman IP 2019 New European socio-economic scenarios for climate change research: Operationalising concepts to extend the shared socio-economic pathways. *Reg. Environ. Chang.* 19, 643–654. doi: [10.1007/s10113-018-1400-0](https://doi.org/10.1007/s10113-018-1400-0)
- Kolbert E 2014 *The sixth extinction: An unnatural history*. New York: Henry Holt and Company.
- Kosmas C, Danalatos N, Cammeraat LH, Chabart M, Diamantopoulos J et al. 1997 The effect of land use on runoff and soil erosion rates under Mediterranean conditions. *Catena* 29, 45–59. doi: [10.1016/s0341-8162\(96\)00062-8](https://doi.org/10.1016/s0341-8162(96)00062-8)
- Kosmas C, Kirkby MJ, Geeson N 1999 *The Medalus project. Mediterranean desertification and land use : manual on key indicators of desertification and mapping environmentally sensitive areas to desertification*. Brussels: Directorate-General Science, European Commission.
- Kourgialas NN, Karatzas GP 2017 A national scale flood hazard mapping methodology: The case of Greece – Protection and adaptation policy approaches. *Sci. Total Environ.* 601–602, 441–452. doi: [10.1016/j.scitotenv.2017.05.197](https://doi.org/10.1016/j.scitotenv.2017.05.197)
- Kourgialas NN, Koubouris GC, Dokou Z 2019 Optimal irrigation planning for addressing current or future water scarcity in Mediterranean tree crops. *Sci. Total Environ.* 654, 616–632. doi: [10.1016/j.scitotenv.2018.11.118](https://doi.org/10.1016/j.scitotenv.2018.11.118)
- Koutroulis AG, Grillakis MG, Daliakopoulos IN, Tsanis IK, Jacob D 2016 Cross sectoral impacts on water availability at +2 degrees C and +3 degrees C for east Mediterranean island states: The case of Crete. *J. Hydrol.* 532, 16–28. doi: [10.1016/j.jhydrol.2015.11.015](https://doi.org/10.1016/j.jhydrol.2015.11.015)
- Kraemer MUG, Reiner Jr. RC, Brady OJ, Messina JP, Gilbert M et al. 2019 Past and future spread of the arbovirus vectors *Aedes aegypti* and *Aedes albopictus*. *Nat. Microbiol.* 4, 854–863. doi: [10.1038/s41564-019-0376-y](https://doi.org/10.1038/s41564-019-0376-y)
- Kreibich H, di Baldassarre G, Vorogushyn S, Aerts JCJH, Apel H et al. 2017 Adaptation to flood risk: Results of international paired flood event studies. *Earth's Futur.* 5, 953–965. doi: [10.1002/2017ef000606](https://doi.org/10.1002/2017ef000606)
- Kuglitsch FG, Toreti A, Xoplaki E, Della-Marta PM, Zerefos CS et al. 2010 Heat wave changes in the eastern Mediterranean since 1960. *Geophys. Res. Lett.* 37, 1–5. doi: [10.1029/2009GL041841](https://doi.org/10.1029/2009GL041841)
- Kundzewicz ZW, Krysanova V, Dankers R, Hirabayashi Y, Kanae S et al. 2017 Differences in flood hazard projections in Europe – their causes and consequences for decision making. *Hydrol. Sci. J.* 62, 1–14. doi: [10.1080/02626667.2016.1241398](https://doi.org/10.1080/02626667.2016.1241398)
- La Jeunesse I, Cirelli C, Sellami H, Aubin D, Deidda R et al. 2015 Is the governance of the Thau coastal lagoon ready to face climate change impacts? *Ocean Coast. Manag.* 118, 234–246. doi: [10.1016/j.ocecoaman.2015.05.014](https://doi.org/10.1016/j.ocecoaman.2015.05.014)
- Lacoue-Labarthe T, Nunes PALD, Ziveri P, Cinar M, Gazeau F et al. 2016 Impacts of ocean acidification in a warming Mediterranean Sea: An overview. *Reg. Stud. Mar. Sci.* 5, 1–11. doi: [10.1016/j.rsma.2015.12.005](https://doi.org/10.1016/j.rsma.2015.12.005)
- Lal R, Bruce JP 1999 The potential of world cropland soils to sequester C and mitigate the greenhouse effect. *Environ. Sci. Policy* 2, 177–185. doi: [10.1016/s1462-9011\(99\)00012-x](https://doi.org/10.1016/s1462-9011(99)00012-x)
- Lamberti A, Zanuttigh B 2005 An integrated approach to beach management in Lido di Dante, Italy. *Estuar. Coast. Shelf Sci.* 62, 441–451. doi: [10.1016/j.ecss.2004.09.022](https://doi.org/10.1016/j.ecss.2004.09.022)
- Lampin-Maillet C, Long-Fournel M, Ganteaume A, Jappiot M, Ferrier JP 2011 Land cover analysis in wildland–urban interfaces according to wildfire risk: A case study in the South of France. *For. Ecol. Manag.* 261, 2200–2213. doi: [10.1016/j.foreco.2010.11.022](https://doi.org/10.1016/j.foreco.2010.11.022)
- Lampurlanés J, Plaza-Bonilla D, Álvaro-Fuentes J, Cantero-Martínez C 2016 Long-term analysis of soil water conservation and crop yield under different tillage systems in Mediterranean rainfed conditions. *F. Crop. Res.* 189, 59–67. doi: [10.1016/j.fcr.2016.02.010](https://doi.org/10.1016/j.fcr.2016.02.010)
- Lamqadem AA, Pradhan B, Saber H, Rahimi A 2018 Desertification Sensitivity Analysis Using MEDALUS Model and GIS: A Case Study of the Oases of Middle Draa Valley, Morocco. *Sensors (Basel)*. 18, 2230. doi: [10.3390/s18072230](https://doi.org/10.3390/s18072230)
- Landeta AA 1995 The Master Plan for Soil Protection of the Basque Autonomous Community, in *Contaminated Soil '95. Soil & Environment, vol 5.*, eds. Van Den Brink WJ, Bosman R, Arendt F (Dordrecht: Springer).
- Larsen L 2015 Urban climate and adaptation strategies. *Front. Ecol. Environ.* 13, 486–492. doi: [10.1890/150103](https://doi.org/10.1890/150103)
- Lavado Contador JF, Schnabel S, Gómez Gutiérrez A, Pulido Fernández M 2009 Mapping sensitivity to land degradation in Extremadura, SW Spain. *L. Degrad. Dev.* 20, 129–144. doi: [10.1002/ldr.884](https://doi.org/10.1002/ldr.884)
- Lelieveld J, Hadjinicolaou P, Kostopoulou E, Chenoweth J, El Maayar M et al. 2012 Climate change and impacts in the Eastern Mediterranean and the Middle East. *Clim. Change* 114, 667–687. doi: [10.1007/s10584-012-0418-4](https://doi.org/10.1007/s10584-012-0418-4)
- Li Z, Fang H 2016 Impacts of climate change on water erosion: A review. *Earth-Science Rev.* 163, 94–117. doi: [10.1016/j.earscirev.2016.10.004](https://doi.org/10.1016/j.earscirev.2016.10.004)
- Lichter M, Felsenstein D 2012 Assessing the costs of sea-level rise and extreme flooding at the local level: A GIS-based approach. *Ocean Coast. Manag.* 59, 47–62. doi: [10.1016/j.ocecoaman.2011.12.020](https://doi.org/10.1016/j.ocecoaman.2011.12.020)
- Liniger H, Critchley W, WOCAT 2007 *WOCAT 2007 : where the*

- land is greener : case studies and analysis of soil and water conservation initiatives worldwide. CTA, UNEP, FAO and CDE.
- Liotta G, Inzerilli MC, Palombi L, Madaro O, Orlando S et al. 2018 Social Interventions to Prevent Heat-Related Mortality in the Older Adult in Rome, Italy: A Quasi-Experimental Study. *Int. J. Environ. Res. Public Health* 15. doi: [10.3390/ijerph15040715](https://doi.org/10.3390/ijerph15040715)
- Liu-Helmersson J, Rocklöv J, Sewe M, Brännström Å 2019 Climate change may enable *Aedes aegypti* infestation in major European cities by 2100. *Env. Res* 172, 693–699. doi: [10.1016/j.envres.2019.02.026](https://doi.org/10.1016/j.envres.2019.02.026)
- Llasat MC, Barriendos M, Barrera A, Rigo T 2005 Floods in Catalonia (NE Spain) since the 14th century. Climatological and meteorological aspects from historical documentary sources and old instrumental records. in *Journal of Hydrology* (Elsevier), 32–47. doi: [10.1016/j.jhydrol.2005.02.004](https://doi.org/10.1016/j.jhydrol.2005.02.004)
- Llasat MC, Llasat-Botija M, Prat MA, PorcúPorc F, Price C et al. 2010 High-impact floods and flash floods in Mediterranean countries: the FLASH preliminary database. [www.adv-geosci.net/23/47/2010/](http://www.adv-geosci.net/23/47/2010/) [Accessed April 2, 2019].
- Llasat MC, Marcos R, Llasat-Botija M, Gilabert J, Turco M et al. 2014 Flash flood evolution in North-Western Mediterranean. *Atmos. Res.* 149, 230–243. doi: [10.1016/j.atmosres.2014.05.024](https://doi.org/10.1016/j.atmosres.2014.05.024)
- Lloret F, Vilà M 2003 Diversity patterns of plant functional types in relation to fire regime and previous land use in Mediterranean woodlands. *J. Veg. Sci.* 14, 387–398. doi: [10.1111/j.1654-1103.2003.tb02164.x](https://doi.org/10.1111/j.1654-1103.2003.tb02164.x)
- Lloret J, Lleonart J, Sole I, Fromentin J-M 2001 Fluctuations of landings and environmental conditions in the north-western Mediterranean Sea. *Fish. Oceanogr.* 10, 33–50. doi: [10.1046/j.1365-2419.2001.00151.x](https://doi.org/10.1046/j.1365-2419.2001.00151.x)
- Lockwood J, Hoopes M, Marchetti M 2007 *Invasion Ecology*. Blackwell Publishing.
- Lodge DM, Simonin PW, Burgiel SW, Keller RP, Bossenbroek JM et al. 2016 Risk Analysis and Bioeconomics of Invasive Species to Inform Policy and Management. *Annu. Rev. Environ. Resour.* 41, 453–488. doi: [10.1146/annurev-environ-110615-085532](https://doi.org/10.1146/annurev-environ-110615-085532)
- Loepfe L, Martinez-Vilalta J, Piñol J 2011 An integrative model of human-influenced fire regimes and landscape dynamics. *Environ. Model. Softw.* 26, 1028–1040. doi: [10.1016/j.envsoft.2011.02.015](https://doi.org/10.1016/j.envsoft.2011.02.015)
- Losada IJ, Toimil A, Muñoz A, Garcia-Fletcher AP, Diaz-Simal P 2019 A planning strategy for the adaptation of coastal areas to climate change: The Spanish case. *Ocean Coast. Manag.* 182, 104983. doi: [10.1016/j.ocecoaman.2019.104983](https://doi.org/10.1016/j.ocecoaman.2019.104983)
- Ludwig R, Roson R 2016 Climate change, water and security in the Mediterranean: Introduction to the special issue. *Sci. Total Environ.* 543, 847–850. doi: [10.1016/j.scitotenv.2015.10.142](https://doi.org/10.1016/j.scitotenv.2015.10.142)
- Ludwig R, Roson R, Zografos C, Kallis G 2011 Towards an inter-disciplinary research agenda on climate change, water and security in Southern Europe and neighboring countries. *Environ. Sci. Policy* 14, 794–803. doi: [10.1016/j.envsci.2011.04.003](https://doi.org/10.1016/j.envsci.2011.04.003)
- Luijendijk A, Hagenaars G, Ranasinghe R, Baart F, Donchyts G et al. 2018 The State of the World's Beaches. *Sci. Rep.* 8. doi: [10.1038/s41598-018-24630-6](https://doi.org/10.1038/s41598-018-24630-6)
- Mainali KP, Warren DL, Dhileepan K, McConnachie A, Strathie L et al. 2015 Projecting future expansion of invasive species: comparing and improving methodologies for species distribution modeling. *Glob. Chang. Biol.* 21, 4464–4480. doi: [10.1111/gcb.13038](https://doi.org/10.1111/gcb.13038)
- Malek Ž, Verburg PH 2018 Adaptation of land management in the Mediterranean under scenarios of irrigation water use and availability. *Mitig. Adapt. Strateg. Glob. Chang.* 23, 821–837. doi: [10.1007/s11027-017-9761-0](https://doi.org/10.1007/s11027-017-9761-0)
- Mannochi F, Todisco F, Vergni L 2004 Agricultural drought: Indices, definition and analysis, in *The Basis of Civilization - Water Science ?* (Rome, Italy, December 2003: Rodda JC, Ubertini L (eds)), 246–254. <https://iahs.info/Publications-News.do>
- Margonski P, Hansson S, Tomczak MT, Grzebielec R 2010 Climate influence on Baltic cod, sprat, and herring stock-recruitment relationships. *Prog. Oceanogr.* 87, 277–288. doi: [10.1016/j.pocean.2010.08.003](https://doi.org/10.1016/j.pocean.2010.08.003)
- Marras S, Cucco A, Antognarelli F, Azzurro E, Milazzo M et al. 2015 Predicting future thermal habitat suitability of competing native and invasive fish species: from metabolic scope to oceanographic modelling. *Conserv. Physiol.* 3, cou059–cou059. doi: [10.1093/conphys/cou059](https://doi.org/10.1093/conphys/cou059)
- Martínez-Fernández J, Esteve-Selma MA, Baños-González I, Carreño F, Moreno A 2013 Sustainability of Mediterranean irrigated agro-landscapes. *Ecol. Modell.* 248, 11–19. doi: [10.1016/j.ecolmodel.2012.09.018](https://doi.org/10.1016/j.ecolmodel.2012.09.018)
- Marulanda A, Porcel R, Barea JM, Azcón R 2007 Drought tolerance and antioxidant activities in lavender plants colonized by native drought-tolerant or drought-sensitive *Glomus* species. *Microb. Ecol.* 54, 543. doi: [10.1007/s00248-007-9237-y](https://doi.org/10.1007/s00248-007-9237-y)
- Masria A, Iskander M, Negm A 2015 Coastal protection measures, case study (Mediterranean zone, Egypt). *J. Coast. Conserv.* 19, 281–294. doi: [10.1007/s11852-015-0389-5](https://doi.org/10.1007/s11852-015-0389-5)
- McDermott TKJ, Surminski S 2018 How normative interpretations of climate risk assessment affect local decision-making: an exploratory study at the city scale in Cork, Ireland. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376. doi: [10.1098/rsta.2017.0300](https://doi.org/10.1098/rsta.2017.0300)
- Meddich A, Oihabi A, Abbas Y, Bizid E 2000 Rôle des champignons mycorhiziens à arbuscules de zones arides dans la résistance du trèfle (*Trifolium alexandrinum* L.) au déficit hydrique. *Agronomie* 20, 283–295. doi: [10.1051/agro:2000127](https://doi.org/10.1051/agro:2000127)
- Medlock JM, Hansford KM, Schaffner F, Versteirt V, Hendrickx G et al. 2012 A review of the invasive mosquitoes in Europe: Ecology, public health risks, and control options. *Vector-Borne Zoonotic Dis.* 12, 435–447. doi: [10.1089/vbz.2011.0814](https://doi.org/10.1089/vbz.2011.0814)
- Mentaschi L, Vousdoukas MI, Pekel J-F, Voukouvalas E, Feyen L 2018 Global long-term observations of coastal erosion

- and accretion. *Sci. Rep.* 8. doi: [10.1038/s41598-018-30904-w](https://doi.org/10.1038/s41598-018-30904-w)
- Metternicht G, Akhtar-Schuster M, Castillo V 2019 Implementing land degradation neutrality: From policy challenges to policy opportunities for national sustainable development. *Environ. Sci. Policy* 100, 189–191. doi: [10.1016/j.envsci.2019.07.010](https://doi.org/10.1016/j.envsci.2019.07.010)
- Monteiro A, Lopes CM 2007 Influence of cover crop on water use and performance of vineyard in Mediterranean Portugal. *Agric. Ecosyst. Environ.* 121, 336–342. doi: [10.1016/j.agee.2006.11.016](https://doi.org/10.1016/j.agee.2006.11.016)
- Montero-Serra I, Garrabou J, Doak DF, Ledoux J, Linares C 2019 Marine protected areas enhance structural complexity but do not buffer the consequences of ocean warming for an overexploited precious coral. *J. Appl. Ecol.* 56, 1063–1074. doi: [10.1111/1365-2664.13321](https://doi.org/10.1111/1365-2664.13321)
- Moreira F, Ferreira PG, Rego FC, Bunting S 2001 Landscape changes and breeding bird assemblages in northwestern Portugal: the role of fire. *Landsc. Ecol.* 16, 175–187. doi: [10.1023/a:1011169614489](https://doi.org/10.1023/a:1011169614489)
- Moreno J, Arianoutsou M, González-Cabán A, Mouillot F, Oechel W et al. 2014 Forest fires under climate, social and economic changes in Europe, the Mediterranean and other fire-affected areas of the world.
- Moritz MA, Batllori E, Bradstock RA, Gill AM, Handmer J et al. 2014 Learning to coexist with wildfire. *Nature* 515, 58–66. doi: [10.1038/nature13946](https://doi.org/10.1038/nature13946)
- Morugán-Coronado A, García-Orenes F, Cerdà A 2015 Changes in soil microbial activity and physicochemical properties in agricultural soils in Eastern Spain. *Spanish J. Soil Sci.* 5. doi: [10.3232/sjss.2015.V5.N3.02](https://doi.org/10.3232/sjss.2015.V5.N3.02)
- Mrabet R 2002 Wheat Yield And Water use efficiency Under Contrasting Residue and tillage Management Systems In A Semiarid Area of Morocco. *Exp. Agric.* 38, 237–248. doi: [10.1017/s0014479702000285](https://doi.org/10.1017/s0014479702000285)
- Mrabet R, Moussadek R, Fadlaoui A, van Ranst E 2012 Conservation agriculture in dry areas of Morocco. *F. Crop. Res.* 132, 84–94. doi: [10.1016/j.fcr.2011.11.017](https://doi.org/10.1016/j.fcr.2011.11.017)
- Munari C, Corbau C, Simeoni U, Mistri M 2011 Coastal defence through low crested breakwater structures: Jumping out of the frying pan into the fire? *Mar. Pollut. Bull.* 62, 1641–1651. doi: [10.1016/j.marpolbul.2011.06.012](https://doi.org/10.1016/j.marpolbul.2011.06.012)
- Navarra A, Tubiana L 2013 *Regional Assessment of Climate Change in the Mediterranean*. Berlin, Heidelberg, New York: Springer
- Navarro García A, del Pilar Bañón Árias S, Morte A, Sánchez-Blanco MJ 2011 Effects of nursery preconditioning through mycorrhizal inoculation and drought in *Arbutus unedo* L. plants. *Mycorrhiza* 21, 53–64. doi: [10.1007/s00572-010-0310-x](https://doi.org/10.1007/s00572-010-0310-x)
- Nearing M, Pruski FF, O'Neal MR 2004 Expected climate change impacts on soil erosion rates: A review. *J. Soil Water Conserv.* 59, 43–50.
- Negev M, Paz S, Clermont A, Pri-Or NG, Shalom U et al. 2015 Impacts of climate change on vector borne diseases in the Mediterranean Basin - implications for preparedness and adaptation policy. *Int. J. Environ. Res. Public Health* 12, 6745–6770. doi: [10.3390/ijerph120606745](https://doi.org/10.3390/ijerph120606745)
- Nemry F, Demirel H 2012 Impacts of Climate Change on Transport: A focus on road and rail transport infrastructures. doi: [10.2791/15504](https://doi.org/10.2791/15504)
- Neumann B, Unger S 2019 From voluntary commitments to ocean sustainability. *Science (80-. )*. 363, 35–36. doi: [10.1126/science.aav5727](https://doi.org/10.1126/science.aav5727)
- Nunes JP, Seixas J, Keizer JJ 2013 Modeling the response of within-storm runoff and erosion dynamics to climate change in two Mediterranean watersheds: A multi-model, multi-scale approach to scenario design and analysis. *CATENA* 102, 27–39. doi: [10.1016/j.catena.2011.04.001](https://doi.org/10.1016/j.catena.2011.04.001)
- O'Connor MC, Lymbery G, Cooper JAG, Gault J, McKenna J 2009 Practice versus policy-led coastal defence management. *Mar. Policy* 33, 923–929. doi: [10.1016/j.marpol.2009.03.007](https://doi.org/10.1016/j.marpol.2009.03.007)
- O'Neill BC, Kriegler E, Ebi KL, Kemp-Benedict E, Riahi K et al. 2017 The roads ahead: Narratives for shared socio-economic pathways describing world futures in the 21<sup>st</sup> century. *Glob. Environ. Chang.* 42, 169–180. doi: [10.1016/j.gloenvcha.2015.01.004](https://doi.org/10.1016/j.gloenvcha.2015.01.004)
- Occhipinti-Ambrogi A, Galil B 2010 Marine alien species as an aspect of global change. *Adv. Oceanogr. Limnol.* 1, 199–218. doi: [10.1080/19475721003743876](https://doi.org/10.1080/19475721003743876)
- Oliveira SLJ, Pereira JMC, Carreiras JMB 2012 Fire frequency analysis in Portugal (1975 - 2005), using Landsat-based burnt area maps. *Int. J. Wildl. Fire* 21, 48. doi: [10.1071/wf10131](https://doi.org/10.1071/wf10131)
- Oliver ECJ, Donat MG, Burrows MT, Moore PJ, Smale DA et al. 2018 Longer and more frequent marine heatwaves over the past century. *Nat. Commun.* 9, 1324. doi: [10.1038/s41467-018-03732-9](https://doi.org/10.1038/s41467-018-03732-9)
- Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN et al. 2001 Terrestrial Ecoregions of the World: A New Map of Life on Earth. *Bioscience* 51, 933. doi: [10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2)
- Oppenheimer M, Glavovic BC, Hinkel J, van de Wal R, Magann AK et al. 2019 Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities, in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, eds. Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M et al. (in press).
- Orlowsky B, Seneviratne SI 2013 Elusive drought: uncertainty in observed trends and short- and long-term CMIP5 projections. *Hydrol. Earth Syst. Sci.* 17, 1765–1781. doi: [10.5194/hess-17-1765-2013](https://doi.org/10.5194/hess-17-1765-2013)
- Otero I, Castellnou M, González I, Arilla E, Castell L et al. 2018 Democratizing wildfire strategies. Do you realize what it means? Insights from a participatory process in the Montseny region (Catalonia, Spain). *PLoS One* 13. doi: [ARTN e0204806 10.1371/journal.pone.0204806](https://doi.org/10.1371/journal.pone.0204806)
- Otero I, Nielsen JØ 2017 Coexisting with wildfire?: Achievements and challenges for a radical social-ecological transformation in Catalonia (Spain). *Geoforum* 85, 234–246. doi: [10.1016/j.geoforum.2017.07.020](https://doi.org/10.1016/j.geoforum.2017.07.020)
- Otero J, L'Abée-Lund JH, Castro-Santos T, Leonardsson K, Storvik GO et al. 2013 Basin-scale phenology and effects

- of climate variability on global timing of initial seaward migration of Atlantic salmon (*Salmo salar*). *Glob. Chang. Biol.* 20, 61–75. doi: [10.1111/gcb.12363](https://doi.org/10.1111/gcb.12363)
- Otto IM, Donges JF, Cremades R, Bhowmik A, Hewitt RJ et al. 2020 Social tipping dynamics for stabilizing Earth's climate by 2050. *Proc. Natl. Acad. Sci. U. S. A.* 117, 2354–2365. doi: [10.1073/pnas.1900577117](https://doi.org/10.1073/pnas.1900577117)
- Paciello MC 2015 *Building sustainable agriculture for food security in the Euro-Mediterranean area: Challenges and policy options*. Rome, Italy: Nuova Cultura.
- Pagliai M, Lamarca M, Lucamante G 1983 Micromorphometric and micromorphological investigations of a clay loam soil in viticulture under zero and conventional tillage. *J. Soil Sci.* 34, 391–403. doi: [10.1111/j.1365-2389.1983.tb01044.x](https://doi.org/10.1111/j.1365-2389.1983.tb01044.x)
- Pagliai M, Vignozzi N, Pellegrini S 2004 Soil structure and the effect of management practices. *Soil Tillage Res.* 79, 131–143. doi: [10.1016/j.still.2004.07.002](https://doi.org/10.1016/j.still.2004.07.002)
- Paini DR, Sheppard AW, Cook DC, de Barro PJ, Worner SP et al. 2016 Global threat to agriculture from invasive species. *Proc. Natl. Acad. Sci. U. S. A.* 113, 7575–7579. doi: [10.1073/pnas.1602205113](https://doi.org/10.1073/pnas.1602205113)
- Panagos P, Ballabio C, Meusburger K, Spinoni J, Alewell C et al. 2017a Towards estimates of future rainfall erosivity in Europe based on REDES and WorldClim datasets. *J. Hydrol.* 548, 251–262. doi: [10.1016/j.jhydrol.2017.03.006](https://doi.org/10.1016/j.jhydrol.2017.03.006)
- Panagos P, Borrelli P, Meusburger K, Yu B, Klik A et al. 2017b Global rainfall erosivity assessment based on high-temporal resolution rainfall records. *Sci. Rep.* 7, 4175. doi: [10.1038/s41598-017-04282-8](https://doi.org/10.1038/s41598-017-04282-8)
- Panagos P, Borrelli P, Poesen J, Ballabio C, Lugato E et al. 2015 The new assessment of soil loss by water erosion in Europe. *Environ. Sci. Policy* 54, 438–447. doi: [10.1016/j.envsci.2015.08.012](https://doi.org/10.1016/j.envsci.2015.08.012)
- Papalexioiu SM, AghaKouchak A, Trenberth KE, Fofoula-Georgiou E 2018 Global, Regional, and Megacity Trends in the Highest Temperature of the Year: Diagnostics and Evidence for Accelerating Trends. *Earth's Futur.* doi: [10.1002/2017ef000709](https://doi.org/10.1002/2017ef000709)
- Patanè C, Tringali S, Sortino O 2011 Effects of deficit irrigation on biomass, yield, water productivity and fruit quality of processing tomato under semi-arid Mediterranean climate conditions. *Sci. Hortic. (Amsterdam)*. 129, 590–596. doi: [10.1016/j.scienta.2011.04.030](https://doi.org/10.1016/j.scienta.2011.04.030)
- Paterson SK, Pelling M, Nunes LH, Araújo Moreira F de, Guida K et al. 2017 Size does matter: City scale and the asymmetries of climate change adaptation in three coastal towns. *Geoforum* 81, 109–119. doi: [10.1016/j.geoforum.2017.02.014](https://doi.org/10.1016/j.geoforum.2017.02.014)
- Pausas JG 2004 Changes in Fire and Climate in the Eastern Iberian Peninsula (Mediterranean Basin). *Clim. Change* 63, 337–350. doi: [10.1023/B:CLIM.0000018508.94901.9c](https://doi.org/10.1023/B:CLIM.0000018508.94901.9c)
- Pausas JG, Fernández-Muñoz S 2012 Fire regime changes in the Western Mediterranean Basin: from fuel-limited to drought-driven fire regime. *Clim. Change* 110, 215–226. doi: [10.1007/s10584-011-0060-6](https://doi.org/10.1007/s10584-011-0060-6)
- Pausas JG, Keeley JE 2009 A Burning Story: The Role of Fire in the History of Life. *Bioscience* 59, 593–601. doi: [10.1525/bio.2009.59.7.10](https://doi.org/10.1525/bio.2009.59.7.10)
- Pausas JG, Llovet J, Rodrigo A, Vallejo R 2009 Are wildfires a disaster in the Mediterranean basin? - A review. *Int. J. Wildl. Fire* 17, 713–723. doi: [10.1071/wf07151](https://doi.org/10.1071/wf07151)
- Pavithra D, Yapa N 2018 Arbuscular mycorrhizal fungi inoculation enhances drought stress tolerance of plants. *Groundw. Sustain. Dev.* 7, 490–494. doi: [10.1016/j.gsd.2018.03.005](https://doi.org/10.1016/j.gsd.2018.03.005)
- Paz S, Negev M, Clermont A, Green MS 2016 Health aspects of climate change in cities with Mediterranean climate, and local adaptation plans. *Int. J. Environ. Res. Public Health* 13. doi: [10.3390/ijerph13040438](https://doi.org/10.3390/ijerph13040438)
- Peled Y, Zemah Shamir S, Shechter M, Rahav E, Israel A 2018 A new perspective on valuating marine climate regulation: The Israeli Mediterranean as a case study. *Ecosyst. Serv.* 29, 83–90. doi: [10.1016/j.ecoser.2017.12.001](https://doi.org/10.1016/j.ecoser.2017.12.001)
- Pereira SC, Marta-Almeida M, Carvalho AC, Rocha A 2017 Heat wave and cold spell changes in Iberia for a future climate scenario. *Int. J. Climatol.* 37, 5192–5205. doi: [10.1002/joc.5158](https://doi.org/10.1002/joc.5158)
- Perry A 2003 Impacts of climate change on tourism in the Mediterranean, in *Climate Change in the Mediterranean: socio-economic perspectives of impacts, vulnerability and adaptation*, eds. Giupponi C, Shechter M (Edward Elgar Pub).
- Peterson DA, Fromm MD, Solbrig JE, Hyer EJ, Surratt ML et al. 2017 Detection and Inventory of Intense Pyroconvection in Western North America using GOES-15 Daytime Infrared Data. *J. Appl. Meteorol. Climatol.* 56, 471–493.
- Pineda N, Rigo T 2017 The rainfall factor in lightning-ignited wildfires in Catalonia. *Agric. For. Meteorol.* 239, 249–263. doi: [10.1016/j.agrformet.2017.03.016](https://doi.org/10.1016/j.agrformet.2017.03.016)
- Planton S, Lionello P, Artale V, Aznar R, Carrillo A et al. 2012 The climate of the Mediterranean region in future climate projections, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Oxford: Elsevier), 449–502. doi: [10.1016/B978-0-12-416042-2.00008-2](https://doi.org/10.1016/B978-0-12-416042-2.00008-2)
- Portman ME 2013 Ecosystem services in practice: Challenges to real world implementation of ecosystem services across multiple landscapes - A critical review. *Appl. Geogr.* 45, 185–192. doi: [10.1016/j.apgeog.2013.09.011](https://doi.org/10.1016/j.apgeog.2013.09.011)
- Portman ME 2016 *Environmental Planning for Oceans and Coasts: Methods, Tools, and Technologies*. Switzerland: Springer International Publishing. doi: [10.1007/978-3-319-26971-9](https://doi.org/10.1007/978-3-319-26971-9)
- Portman ME 2018 Policy Options for Coastal Protection: Integrating Inland Water Management with Coastal Management for Greater Community Resilience. *J. Water Resour. Plan. Manag.* 144, 5018005. doi: [10.1061/\(asce\)wr.1943-5452.0000913](https://doi.org/10.1061/(asce)wr.1943-5452.0000913)
- Portman ME, Brennan RE 2017 Marine litter from beach-based sources: Case study of an Eastern Mediterranean coastal town. *Waste Manag.* 69, 535–544. doi: [10.1016/j.wasman.2017.07.040](https://doi.org/10.1016/j.wasman.2017.07.040)
- Portman ME, Pasternak G, Yotam Y, Nusbaum R, Behar D

- 2019 Beachgoer participation in prevention of marine litter: Using design for behavior change. *Mar. Pollut. Bull.* 144, 1–10. doi: [10.1016/j.marpolbul.2019.04.071](https://doi.org/10.1016/j.marpolbul.2019.04.071)
- Porzio L, Buia M-C, Hall-Spencer JM 2011 Effects of ocean acidification on macroalgal communities. *J. Exp. Mar. Bio. Ecol.* 400, 278–287. doi: [10.1016/j.jembe.2011.02.011](https://doi.org/10.1016/j.jembe.2011.02.011)
- Posta K, Duc NH 2019 Benefits of Arbuscular Mycorrhizal Fungi Application to Crop Production under Water Scarcity, in *Drought (Aridity)* (IntechOpen). doi: [10.5772/intechopen.86595](https://doi.org/10.5772/intechopen.86595)
- Poulos SE, Collins MB 2002 Fluvial sediment fluxes to the Mediterranean Sea: a quantitative approach and the influence of dams. *Geol. Soc. London, Spec. Publ.* 191, 227–245. doi: [10.1144/gsl.sp.2002.191.01.16](https://doi.org/10.1144/gsl.sp.2002.191.01.16)
- PPRD East 2013 Good Practices in Disaster Prevention. Final Report. <https://climate-adapt.eea.europa.eu/metadata/publications/good-practices-in-disaster-prevention-1>
- Prävälje R, Patriche C, Bandoc G 2017 Quantification of land degradation sensitivity areas in Southern and Central Southeastern Europe. New results based on improving DISMED methodology with new climate data. *CATENA* 158, 309–320. doi: [10.1016/j.catena.2017.07.006](https://doi.org/10.1016/j.catena.2017.07.006)
- Proestos Y, Christophides GK, Ergüler K, Tanarhte M, Wal-dock J et al. 2015 Present and future projections of habitat suitability of the Asian tiger mosquito, a vector of viral pathogens, from global climate simulation. *Philos. Trans. R. Soc. B Biol. Sci.* 370, 1–16. doi: [10.1098/rstb.2013.0554](https://doi.org/10.1098/rstb.2013.0554)
- Quintana-Seguí P, Martin E, Sánchez E, Zribi M, Vennetier M et al. 2016 Drought: Observed trends, future projections, in *The Mediterranean Region under Climate Change. A Scientific Update*, eds. Thiébault S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 123–132.
- Rahimzadeh S, Pirzad AR 2017 Microorganisms (AMF and PSB) interaction on linseed productivity under water-deficit condition. *Int. J. Plant Prod.* 11, 259–274. [http://ijpp.gau.ac.ir/issue\\_488\\_501\\_Volume+11%2C+Issue+2%2C+Spring+2017%2C+Page+210-347.html](http://ijpp.gau.ac.ir/issue_488_501_Volume+11%2C+Issue+2%2C+Spring+2017%2C+Page+210-347.html)
- Raitsos DE, Beaugrand G, Georgopoulos D, Zenetos A, Pancucci-Papadopoulou AM et al. 2010 Global climate change amplifies the entry of tropical species into the eastern Mediterranean Sea. *Limnol. Oceanogr.* 55, 1478–1484. doi: [10.4319/lo.2010.55.4.1478](https://doi.org/10.4319/lo.2010.55.4.1478)
- Ramajo L, Lagos NA, Duarte CM 2019 Seagrass *Posidonia oceanica* diel pH fluctuations reduce the mortality of epiphytic forams under experimental ocean acidification. *Mar. Pollut. Bull.* 146, 247–254. doi: [10.1016/j.marpolbul.2019.06.011](https://doi.org/10.1016/j.marpolbul.2019.06.011)
- Rapparini F, Peñuelas J 2014 Mycorrhizal Fungi to Alleviate Drought Stress on Plant Growth, in *Use of Microbes for the Alleviation of Soil Stresses, Volume 1*, ed. Miransari M (New York, NY: Springer New York), 21–42. doi: [10.1007/978-1-4614-9466-9\\_2](https://doi.org/10.1007/978-1-4614-9466-9_2)
- Raynaud D, Thielen J, Salamon P, Burek P, Anquetin S et al. 2015 A dynamic runoff co-efficient to improve flash flood early warning in Europe: evaluation on the 2013 central European floods in Germany. *Meteorol. Appl.* 22, 410–418. doi: [10.1002/met.1469](https://doi.org/10.1002/met.1469)
- Reca J, Trillo C, Sánchez JA, Martínez J, Valera D 2018 Optimization model for on-farm irrigation management of Mediterranean greenhouse crops using desalinated and saline water from different sources. *Agric. Syst.* 166, 173–183. doi: [10.1016/j.agsy.2018.02.004](https://doi.org/10.1016/j.agsy.2018.02.004)
- Reckien D, Salvia M, Heidrich O, Church JM, Pietrapertosa F et al. 2018 How are cities planning to respond to climate change? Assessment of local climate plans from 885 cities in the EU-28. *J. Clean. Prod.* 191, 207–219. doi: [10.1016/j.jclepro.2018.03.220](https://doi.org/10.1016/j.jclepro.2018.03.220)
- Regos A, Aquilué N, López I, Codina M, Retana J et al. 2016 Synergies Between Forest Biomass Extraction for Bioenergy and Fire Suppression in Mediterranean Ecosystems: Insights from a Storyline-and-Simulation Approach. *Ecosystems* 19, 786–802. doi: [10.1007/s10021-016-9968-z](https://doi.org/10.1007/s10021-016-9968-z)
- Regos A, Aquilué N, Retana J, de Cáceres M, Brotons L 2014 Using unplanned fires to help suppressing future large fires in Mediterranean forests. *PLoS One* 9, e94906. doi: [10.1371/journal.pone.0094906](https://doi.org/10.1371/journal.pone.0094906)
- Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR et al. 2016 Critical Review of Health Impacts of Wildfire Smoke Exposure. *Env. Heal. Perspect* 124, 1334–1343. doi: [10.1289/ehp.1409277](https://doi.org/10.1289/ehp.1409277)
- Reidsma P, Ewert F, Oude Lansink A, Leemans R 2009 Vulnerability and adaptation of European farmers: A multi-level analysis of yield and income responses to climate variability. *Reg. Environ. Chang.* 9, 25–40. doi: [10.1007/s10113-008-0059-3](https://doi.org/10.1007/s10113-008-0059-3)
- Reimann L, Merckens J-L, Vafeidis AT 2018a Regionalized Shared Socioeconomic Pathways: Narratives and spatial population projections for the Mediterranean coastal zone. *Reg. Environ. Chang.* 18, 235–245. doi: [10.1007/s10113-017-1189-2](https://doi.org/10.1007/s10113-017-1189-2)
- Reimann L, Vafeidis AT, Brown S, Hinkel J, Tol RSJ 2018b Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-level rise. *Nat. Commun.* 9. doi: [10.1038/s41467-018-06645-9](https://doi.org/10.1038/s41467-018-06645-9)
- Resco P, Iglesias A, Bardají I, Sotés V 2016 Exploring adaptation choices for grapevine regions in Spain. *Reg. Environ. Chang.* 16, 979–993. doi: [10.1007/s10113-015-0811-4](https://doi.org/10.1007/s10113-015-0811-4)
- Revolve Water 2017 Water around the Mediterranean. <https://ufmsecretariat.org/wp-content/uploads/2018/01/water-report-2017.pdf>
- Reynolds JF, Smith DMS, Lambin EF, Turner BL, Mortimore M et al. 2007 Global Desertification: Building a Science for Dryland Development. *Science (80-. )*. 316, 847–851. doi: [10.1126/science.1131634](https://doi.org/10.1126/science.1131634)
- Ribeiro F, Leunda PM 2012 Non-native fish impacts on Mediterranean freshwater ecosystems: current knowledge and research needs. *Fish. Manag. Ecol.* 19, 142–156. doi: [10.1111/j.1365-2400.2011.00842.x](https://doi.org/10.1111/j.1365-2400.2011.00842.x)
- Richey AS 2014 Stress and Resilience in the World's Largest Aquifer Systems. <https://escholarship.org/uc/item/4701986p>

- Roche B, Leger L, L'Ambert G, Lacour G, Foussadier R et al. 2015 The spread of *Aedes albopictus* in metropolitan France: Contribution of environmental drivers and human activities and predictions for a near future. *PLoS One* 10, e0125600. doi: [10.1371/journal.pone.0125600](https://doi.org/10.1371/journal.pone.0125600)
- Rodrigo-Comino J, Wirtz S, Brevik EC, Ruiz-Sinoga JD, Ries JB 2017 Assessment of agri-spillways as a soil erosion protection measure in Mediterranean sloping vineyards. *J. Mt. Sci.* 14, 1009–1022. doi: [10.1007/s11629-016-4269-8](https://doi.org/10.1007/s11629-016-4269-8)
- Rodrigues LC, Van den Bergh JCJM, Ghermandi A 2013 Socio-economic impacts of ocean acidification in the Mediterranean Sea. *Mar. Policy* 38, 447–456. doi: [10.1016/j.marpol.2012.07.005](https://doi.org/10.1016/j.marpol.2012.07.005)
- Rohat G, Flacke J, Dosio A, Dao H, Van Maarseveen M 2019a Projections of human exposure to dangerous heat in African cities under multiple socioeconomic and climate scenarios. *Earth's Futur.*
- Rohat G, Flacke J, Dosio A, Pedde S, Dao H et al. 2019b Influence of changes in socioeconomic and climatic conditions on future heat-related health challenges in Europe. *Glob. Planet. Change* 172, 45–59. doi: [10.1016/j.gloplacha.2018.09.013](https://doi.org/10.1016/j.gloplacha.2018.09.013)
- Rojas M, Lambert F, Ramirez-Villegas J, Challinor AJ 2019 Emergence of robust precipitation changes across crop production areas in the 21<sup>st</sup> century. *Proc. Natl. Acad. Sci. U. S. A.*, 201811463. doi: [10.1073/pnas.1811463116](https://doi.org/10.1073/pnas.1811463116)
- Román M, Midttun A 2010 Governing from the middle: the C40 Cities Leadership Group. *Corp. Gov. Int. J. Bus. Soc.* 10, 73–84. doi: [10.1108/14720701011021120](https://doi.org/10.1108/14720701011021120)
- Romero Diaz A, Caballero Pedraza A, Pérez Morales A 2017 Urbanisation and tourism in the Campo Carategna-Mar Menor area (Murcia, Spain): Impact on soil sealing. *Cuad. Tur.* 39, 521–546.
- Roques A, Rabitsch W, Rasplus J-Y, Vaamonde C, Nentwig W et al. 2009 Alien Terrestrial Invertebrates of Europe, in *Handbook of Alien Species* (Dordrecht: Springer Netherlands), 63–79. doi: [10.1007/978-1-4020-8280-1\\_5](https://doi.org/10.1007/978-1-4020-8280-1_5)
- Rosenzweig C, Karoly D, Vicarelli M, Neofotis P, Wu Q et al. 2008 Attributing physical and biological impacts to anthropogenic climate change. *Nature* 453, 353–357. doi: [10.1038/nature06937](https://doi.org/10.1038/nature06937)
- Rosenzweig C, Solecki W, Hammer SA, Mehrotra S 2010 Cities lead the way in climate-change action. *Nature* 467, 909–911. doi: [10.1038/467909a](https://doi.org/10.1038/467909a)
- Rosenzweig C, Solecki W, Romero-Lankao P, Mehrotra S, Dhakal S et al. 2018 *Second Assessment Report on Climate Change and Cities of the Urban Climate Change Research Network*. New York, USA: Cambridge University Press.
- Rounsevell MDA, Reginster I, Araújo MB, Carter TR, Denoncker N et al. 2006 A coherent set of future land use change scenarios for Europe. *Agric. Ecosyst. Environ.* 114, 57–68. doi: [10.1016/j.agee.2005.11.027](https://doi.org/10.1016/j.agee.2005.11.027)
- Roy HE, de Clercq P, Lawson Handley L-J, Poland RL, Sloggett JJ et al. 2011a Alien arthropod predators and parasitoids: An ecological approach. *BioControl* 56, 375–382. doi: [10.1007/s10526-011-9388-0](https://doi.org/10.1007/s10526-011-9388-0)
- Roy HE, Rhule E, Harding S, Lawson Handley L-J, Poland RL et al. 2011b Living with the enemy: Parasites and pathogens of the ladybird *Harmonia axyridis*. *BioControl* 56, 663–679. doi: [10.1007/s10526-011-9387-1](https://doi.org/10.1007/s10526-011-9387-1)
- Ruiz-Lozano JM, Collados C, Barea JM, Azcón R 2001 Arbuscular mycorrhizal symbiosis can alleviate drought-induced nodule senescence in soybean plants. *New Phytol.* 151, 493–502. doi: [10.1046/j.0028-646x.2001.00196.x](https://doi.org/10.1046/j.0028-646x.2001.00196.x)
- Ruosteenoja K, Markkanen T, Venäläinen A, Räisänen P, Peltola H 2018 Seasonal soil moisture and drought occurrence in Europe in CMIP5 projections for the 21<sup>st</sup> century. *Clim. Dyn.* 50, 1177–1192. doi: [10.1007/s00382-017-3671-4](https://doi.org/10.1007/s00382-017-3671-4)
- Russo A, Gouveia C, Dutra E, Soares PMM, Trigo RM 2018 The synergy between drought and extremely hot summers in the Mediterranean. *Environ. Res. Lett.* 14. doi: [10.1088/1748-9326/aaf09e](https://doi.org/10.1088/1748-9326/aaf09e)
- Russo S, Sillmann J, Fischer EM 2015 Top ten European heatwaves since 1950 and their occurrence in the coming decades. *Environ. Res. Lett.* 10, 124003. doi: [10.1088/1748-9326/10/12/124003](https://doi.org/10.1088/1748-9326/10/12/124003)
- Sá-Sousa P 2014 The Portuguese montado: Conciliating ecological values with human demands within a dynamic agroforestry system. *Ann. For. Sci.* 71, 1–3. doi: [10.1007/s13595-013-0338-0](https://doi.org/10.1007/s13595-013-0338-0)
- Saadi S, Todorovic M, Tanasijevic L, Pereira LS, Pizzigalli C et al. 2015 Climate change and Mediterranean agriculture: Impacts on winter wheat and tomato crop evapotranspiration, irrigation requirements and yield. *Agric. Water Manag.* 147, 103–115. doi: [10.1016/J.AGWAT.2014.05.008](https://doi.org/10.1016/J.AGWAT.2014.05.008)
- Sabatés A, Martin P, Raya V, Sabatés A, Martín P et al. 2012 Changes in life-history traits in relation to climate change: bluefish (*Pomatomus saltatrix*) in the northwestern Mediterranean. *ICES J. Mar. Sci.* 69, 1000–1009. doi: [10.1093/icesjms/fss053](https://doi.org/10.1093/icesjms/fss053)
- Sadhvani JJ, Veza JM, Santana C 2005 Case studies on environmental impact of seawater desalination. *Desalination* 185, 1–8. doi: [10.1016/j.desal.2005.02.072](https://doi.org/10.1016/j.desal.2005.02.072)
- Saidi H, Souissi R, Zargouni F 2012 Environmental impact of detached breakwaters on the Mediterranean coastline of Soliman (North-East of Tunisia). *Rend. Lincei* 23, 339–347. doi: [10.1007/s12210-012-0191-3](https://doi.org/10.1007/s12210-012-0191-3)
- Sala E, Kizilkaya Z, Yildirim D, Ballesteros E 2011 Alien Marine Fishes Deplete Algal Biomass in the Eastern Mediterranean. *PLoS One* 6, e17356. doi: [10.1371/journal.pone.0017356](https://doi.org/10.1371/journal.pone.0017356)
- Saladini F, Betti G, Ferragina E, Bouraoui F, Cupertino S et al. 2018 Linking the water-energy-food nexus and sustainable development indicators for the Mediterranean region. *Ecol. Indic.* 91, 689–697. doi: [10.1016/j.ecolind.2018.04.035](https://doi.org/10.1016/j.ecolind.2018.04.035)
- Salamanca F, Georgescu M, Mahalov A, Moustaoui M, Wang M 2014 Anthropogenic heating of the urban environment due to air conditioning. *JGR Atmos.* 119, 5949–5965. doi: [10.1002/2013JD021225](https://doi.org/10.1002/2013JD021225)
- Salvati L, Bajocco S 2011 Land sensitivity to desertification across Italy: Past, present, and future. *Appl. Geogr.* 31,



- 223–231. doi: [10.1016/j.apgeog.2010.04.006](https://doi.org/10.1016/j.apgeog.2010.04.006)
- Samaniego L, Thober S, Kumar R, Wanders N, Rakovec O et al. 2018 Anthropogenic warming exacerbates European soil moisture droughts. *Nat. Clim. Chang.* 8, 421–426. doi: [10.1038/s41558-018-0138-5](https://doi.org/10.1038/s41558-018-0138-5)
- San-Miguel-Ayanz J, Moreno JM, Camia A 2013 Analysis of large fires in European Mediterranean landscapes: Lessons learned and perspectives. *For. Ecol. Manage.* 294, 11–22. doi: [10.1016/j.foreco.2012.10.050](https://doi.org/10.1016/j.foreco.2012.10.050)
- Sánchez-Díaz M, Honrubia M 1994 Water relations and alleviation of drought stress in mycorrhizal plants, in *Impact of Arbuscular Mycorrhizas on Sustainable Agriculture and Natural Ecosystems*, eds. Gianinazzi S, Schüepp H (Basel: Birkhäuser Basel), 167–178. doi: [10.1007/978-3-0348-8504-1\\_13](https://doi.org/10.1007/978-3-0348-8504-1_13)
- Sancho-García A, Guillén J, Ojeda E 2013 Storm-induced readjustment of an embayed beach after modification by protection works. *Geo-Marine Lett.* 33, 159–172. doi: [10.1007/s00367-012-0319-6](https://doi.org/10.1007/s00367-012-0319-6)
- Schoennagel T, Balch JK, Brenkert-Smith H, Dennis on PE, Harvey BJ et al. 2017 Adapt to more wildfire in western North American forests as climate changes. *Proc. Natl. Acad. Sci. U. S. A.* 114, 4582–4590. doi: [10.1073/pnas.1617464114](https://doi.org/10.1073/pnas.1617464114)
- Scocco P, Piermarteri K, Malfatti A, Tardella FM, Catorci A 2016 Increase of drought stress negatively affects the sustainability of extensive sheep farming in sub-Mediterranean climate. *J. Arid Environ.* 128, 50–58. doi: [10.1016/j.jaridenv.2016.01.006](https://doi.org/10.1016/j.jaridenv.2016.01.006)
- Seddon N, Turner B, Berry PM, Chausson A, Girardin CAJ 2019 Grounding nature-based climate solutions in sound biodiversity science. *Nat. Clim. Chang.* 9, 84–87. doi: [10.1038/s41558-019-0405-0](https://doi.org/10.1038/s41558-019-0405-0)
- Semenza JC, Suk JE 2018 Vector-borne diseases and climate change: a European perspective. *FEMS Microbiol. Lett.* 365. doi: [10.1093/femsle/fnx244](https://doi.org/10.1093/femsle/fnx244)
- Semenza JC, Tran A, Espinosa L, Sudre B, Domanovic D et al. 2016 Climate change projections of West Nile virus infections in Europe: implications for blood safety practices. *Environ. Heal.* 15, S28. doi: [10.1186/s12940-016-0105-4](https://doi.org/10.1186/s12940-016-0105-4)
- Sendek A, Karakoç C, Wagg C, Domínguez-Begines J, do Couto GM et al. 2019 Drought modulates interactions between arbuscular mycorrhizal fungal diversity and barley genotype diversity. *Sci. Rep.* 9, 9650. doi: [10.1038/s41598-019-45702-1](https://doi.org/10.1038/s41598-019-45702-1)
- Silva RA, West JJ, Lamarque JF, Shindell DT, Collins WJ et al. 2016 The effect of future ambient air pollution on human premature mortality to 2100 using output from the ACCMIP model ensemble. *Atmos. Chem. Phys.* 16, 9847–9862. doi: [10.5194/acp-16-9847-2016](https://doi.org/10.5194/acp-16-9847-2016)
- Silva RA, West JJ, Lamarque JF, Shindell DT, Collins WJ et al. 2017 Future Global Mortality from Changes in Air Pollution Attributable to Climate Change. *Nat. Clim. Chang.* 7, 647–651. doi: [10.1038/nclimate3354](https://doi.org/10.1038/nclimate3354)
- Skuras D, Psaltopoulos D 2012 A broad overview of the main problems derived from climate change that will affect agricultural production in the Mediterranean area, in *Building resilience for adaptation to climate change in the agriculture sector: Proceedings of a Joint FAO/OECD Workshop*, 217–260. <http://www.fao.org/docrep/017/i3084e/i3084e.pdf>
- Smith AMS, Kolden CA, Paveglio TB, Cochrane MA, Bowman DMJS et al. 2016 The Science of Firescapes: Achieving Fire-Resilient Communities. *Bioscience* 66, 130–146.
- Sommer S, Zucca C, Grainger A, Cherlet M, Zougmore R et al. 2011 Application of indicator systems for monitoring and assessment of desertification from national to global scales. *L. Degrad. Dev.* 22, 184–197. doi: [10.1002/ldr.1084](https://doi.org/10.1002/ldr.1084)
- Spinoni J, Barbosa P, Jager A de, McCormick N, Naumann G et al. 2019 A new global database of meteorological drought events from 1951 to 2016. *J. Hydrol. Reg. Stud.* 22, 100593. doi: [10.1016/j.ejrh.2019.100593](https://doi.org/10.1016/j.ejrh.2019.100593)
- Srinivasan V, Lambin EF, Gorelick SM, Thompson BH, Rozelle S 2012 The nature and causes of the global water crisis: Syndromes from a meta-analysis of coupled human-water studies. *Water Resour. Res.* 48, 1. doi: [10.1029/2011wr011087](https://doi.org/10.1029/2011wr011087)
- Symeonakis E, Karathanasis N, Koukoulas S, Panagopoulos G 2014 Monitoring Sensitivity to Land Degradation and Desertification with the Environmentally Sensitive Area Index: The Case of Lesvos Island. *L. Degrad. Dev.* 27, 1562–1573. doi: [10.1002/ldr.2285](https://doi.org/10.1002/ldr.2285)
- T-MedNet 2019 Marine heatwaves. <http://www.t-mednet.org/t-resources/marine-heatwaves> [Accessed April 26, 2020].
- Tarolli P, Borga M, Morin E, Delrieu G 2012 Analysis of flash flood regimes in the North-Western and South-Eastern Mediterranean regions. *Nat. Hazards Earth Syst. Sci.* 12, 1255–1265. doi: [10.5194/nhess-12-1255-2012](https://doi.org/10.5194/nhess-12-1255-2012)
- Tedim F, Leone V, Xanthopoulos G 2016 A wildfire risk management concept based on a social-ecological approach in the European Union: Fire Smart Territory. *Int. J. Disaster Risk Reduct.* 18, 138–153. doi: [10.1016/j.ijdrr.2016.06.005](https://doi.org/10.1016/j.ijdrr.2016.06.005)
- Tedim F, Remelgado R, Borges C, Carvalho S, Martins J 2013 Exploring the occurrence of mega-fires in Portugal. *For. Ecol. Manage.* 294, 86–96. doi: [10.1016/j.foreco.2012.07.031](https://doi.org/10.1016/j.foreco.2012.07.031)
- Temmerman S, Meire P, Bouma TJ, Herman PMJ, Ysebaert T et al. 2013 Ecosystem-based coastal defence in the face of global change. *Nature* 504, 79–83. doi: [10.1038/nature12859](https://doi.org/10.1038/nature12859)
- Terzi S, Torresan S, Schneiderbauer S, Critto A, Zebisch M et al. 2019 Multi-risk assessment in mountain regions: A review of modelling approaches for climate change adaptation. *J. Environ. Manage.* 232, 759–771. doi: [10.1016/j.jenvman.2018.11.100](https://doi.org/10.1016/j.jenvman.2018.11.100)
- Thomas SM, Tjaden NB, van den Bos S, Beierkuhnlein C 2014 Implementing cargo movement into climate based risk assessment of vector-borne diseases. *Int. J. Environ. Res. Public Health* 11, 3360–3374. doi: [10.3390/ijerph110303360](https://doi.org/10.3390/ijerph110303360)

- Thomaz SM, Kovalenko KE, Havel JE, Kats LB 2014 Aquatic invasive species: general trends in the literature and introduction to the special issue. *Hydrobiologia* 746, 1–12. doi: [10.1007/s10750-014-2150-8](https://doi.org/10.1007/s10750-014-2150-8)
- Thompson MP, Calkin DE 2011 Uncertainty and risk in wild-land fire management: a review. *J. Environ. Manage.* 92, 1895–1909. doi: [10.1016/j.jenvman.2011.03.015](https://doi.org/10.1016/j.jenvman.2011.03.015)
- Tiller R, Arenas F, Galdies C, Leitão F, Malej A et al. 2019 Who cares about ocean acidification in the Plasticene? *Ocean Coast. Manag.* 174, 170–180. doi: [10.1016/j.ocecoaman.2019.03.020](https://doi.org/10.1016/j.ocecoaman.2019.03.020)
- Tobin PC 2018 Managing invasive species. *F1000Research* 7. doi: [10.12688/F1000RESEARCH.15414.1](https://doi.org/10.12688/F1000RESEARCH.15414.1)
- Tolika CK, Zanis P, Anagnostopoulou C 2012 Regional climate change scenarios for Greece: Future temperature and precipitation projections from ensembles of RCMs. *Glob. Nest J.* 14, 407–421. doi: [10.30955/gnj.000776](https://doi.org/10.30955/gnj.000776)
- Tomasicchio U 1996 Submerged Breakwaters for the Defence of the Shoreline at Ostia Field Experiences, Comparison, in *25th International Conference on Coastal Engineering*, ed. Edge BL, 2404–2417. doi: [10.1061/9780784402429.186](https://doi.org/10.1061/9780784402429.186)
- Tomaz A, Pacheco CA, Coletto Martinez JM 2017 Influence of cover cropping on water uptake dynamics in an irrigated Mediterranean vineyard. *Irrig. Drain.* 66, 387–395. doi: [10.1002/ird.2115](https://doi.org/10.1002/ird.2115)
- Toth E, Bragalli C, Neri M 2018 Assessing the significance of tourism and climate on residential water demand: Panel-data analysis and non-linear modelling of monthly water consumptions. *Environ. Model. Softw.* 103, 52–61. doi: [10.1016/j.envsoft.2018.01.011](https://doi.org/10.1016/j.envsoft.2018.01.011)
- Tramblay Y, Mimeau L, Neppel L, Vinet F, Sauquet E 2019 Detection and attribution of flood trends in Mediterranean basins. *Hydrol. Earth Syst. Sci.* 23, 4419–4431. doi: [10.5194/hess-23-4419-2019](https://doi.org/10.5194/hess-23-4419-2019)
- Trigo RM, Sousa PM, Pereira MG, Rasilla D, Gouveia CM 2013 Modelling wildfire activity in Iberia with different atmospheric circulation weather types. *Int. J. Climatol.* 36, 2761–2778. doi: [10.1002/joc.3749](https://doi.org/10.1002/joc.3749)
- Tsoukala VK, Katsardi V, Hadjibiros K, Moutzouris CI 2015 Beach Erosion and Consequential Impacts Due to the Presence of Harbours in Sandy Beaches in Greece and Cyprus. *Environ. Process.* 2, 55–71. doi: [10.1007/s40710-015-0096-0](https://doi.org/10.1007/s40710-015-0096-0)
- Turco M, Llasat MC, Tudela A, Castro X, Provenzale A 2013 Decreasing fires in a Mediterranean region (1970–2010, NE Spain). *Nat. Hazards Earth Syst. Sci.* 13, 649–652. doi: [10.5194/nhess-13-649-2013](https://doi.org/10.5194/nhess-13-649-2013)
- Turco M, Llasat MC, von Hardenberg J, Provenzale A 2014 Climate change impacts on wildfires in a Mediterranean environment. *Clim. Change* 125, 369–380. doi: [10.1007/s10584-014-1183-3](https://doi.org/10.1007/s10584-014-1183-3)
- Turco M, Rosa-Cánovas JJ, Bedía J, Jerez S, Montávez JP et al. 2018 Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models. *Nat. Commun.* 9, 3821. doi: [10.1038/s41467-018-06358-z](https://doi.org/10.1038/s41467-018-06358-z)
- Tyagi J, Varma A, Pudake RN 2017 Evaluation of comparative effects of arbuscular mycorrhiza (*Rhizophagus intraradices*) and endophyte (*Piriformospora indica*) association with finger millet (*Eleusine coracana*) under drought stress. *Eur. J. Soil Biol.* 81, 1–10. doi: [10.1016/j.ejsobi.2017.05.007](https://doi.org/10.1016/j.ejsobi.2017.05.007)
- Tzoraki O, Dokou Z, Christodoulou G, Gaganis P, Karatzas G 2018 Assessing the efficiency of a coastal Managed Aquifer Recharge (MAR) system in Cyprus. *Sci. Total Environ.* 626, 875–886. doi: [10.1016/j.scitotenv.2018.01.160](https://doi.org/10.1016/j.scitotenv.2018.01.160)
- UN 2015 Transforming our world: the 2030 Agenda for Sustainable Development. New York. [https://www.un.org/ga/search/view\\_doc.asp?symbol=A/RES/70/1&Lang=E](https://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E)
- UNCTAD 2017 Port Industry Survey on Climate Change Impacts and Adaptation.
- UNDP 2017 Addressing climate change vulnerabilities and risks in vulnerable coastal areas of Tunisia. <http://adaptation-undp.org/projects/sccf-tunisia>
- UNEP/MAP/PAP 2008 Protocol on integrated coastal zone management in the Mediterranean. Split
- UNEP/MAP 2012 State of the Mediterranean Marine and Coastal Environment. UNEP/MAP – Barcelona Convention, Athens.
- UNEP/MAP 2016 Mediterranean Strategy for Sustainable Development 2016–2025. Valbonne.
- UNEP/MAP 2017 Regional Climate Change Adaptation Framework for the Mediterranean Marine and Coastal Areas. Athens, Greece.
- Valentin RE, Fonseca DM, Nielsen AL, Leskey TC, Lockwood JL 2018 Early detection of invasive exotic insect infestations using eDNA from crop surfaces. *Front. Ecol. Environ.* 16, 265–270. doi: [10.1002/fee.1811](https://doi.org/10.1002/fee.1811)
- Velegrakis AF, Voudoukas MI, Andreadis OP, Adamakis G, Pasakalidou E et al. 2008 Influence of Dams on Downstream Beaches: Eressos, Lesbos, Eastern Mediterranean. *Mar. Georesources Geotechnol.* 26, 350–371. doi: [10.1080/10641190802425598](https://doi.org/10.1080/10641190802425598)
- Vilà M, Pino J, Font X 2007 Regional assessment of plant invasions across different habitat types. *J. Veg. Sci.* 18, 35–42. doi: [10.1111/j.1654-1103.2007.tb02513.x](https://doi.org/10.1111/j.1654-1103.2007.tb02513.x)
- Vilà M, Pujadas J 2001 Land-use and socio-economic correlates of plant invasions in European and North African countries. *Biol. Conserv.* 100, 397–401. doi: [10.1016/s0006-3207\(01\)00047-7](https://doi.org/10.1016/s0006-3207(01)00047-7)
- Vilà M, Tessier M, Suehs CM, Brundu G, Carta L et al. 2006 Local and regional assessments of the impacts of plant invaders on vegetation structure and soil properties of Mediterranean islands. *J. Biogeogr.* 33, 853–861. doi: [10.1111/j.1365-2699.2005.01430.x](https://doi.org/10.1111/j.1365-2699.2005.01430.x)
- Vitousek PM, Mooney HA, Lubchenco J, Melillo J 1997 Human Domination of Earth's Ecosystems. *Science (80-. ).* 277, 193–207.
- Vizzini S, Apostolaki ET, Ricevuto E, Polymenakou P, Mazzola A 2019 Plant and sediment properties in seagrass meadows from two Mediterranean CO<sub>2</sub> vents: Implica-

- tions for carbon storage capacity of acidified oceans. *Mar. Environ. Res.* 146, 101–108. doi: [10.1016/j.marenvres.2019.03.001](https://doi.org/10.1016/j.marenvres.2019.03.001)
- Vogel MM, Orth R, Cheruy F, Hagemann S, Lorenz R et al. 2017 Regional amplification of projected changes in extreme temperatures strongly controlled by soil moisture-temperature feedbacks. *Geophys. Res. Lett.* 44, 1511–1519. doi: [10.1002/2016gl071235](https://doi.org/10.1002/2016gl071235)
- Vousdoukas MI, Mentaschi L, Hinkel J, Ward PJ, Mongelli I et al. 2020 Economic motivation for raising coastal flood defenses in Europe. *Nat. Commun.* 11, 1–11. doi: [10.1038/s41467-020-15665-3](https://doi.org/10.1038/s41467-020-15665-3)
- Vousdoukas MI, Mentaschi L, Voukouvalas E, Bianchi A, Dottori F et al. 2018a Climatic and socioeconomic controls of future coastal flood risk in Europe. *Nat. Clim. Chang.* 8, 776–780. doi: [10.1038/s41558-018-0260-4](https://doi.org/10.1038/s41558-018-0260-4)
- Vousdoukas MI, Mentaschi L, Voukouvalas E, Verlaan M, Jevrejeva S et al. 2018b Global probabilistic projections of extreme sea levels show intensification of coastal flood hazard. *Nat. Commun.* 9. doi: [10.1038/s41467-018-04692-w](https://doi.org/10.1038/s41467-018-04692-w)
- Wang F, Polcher J 2019 Assessing the freshwater flux from the continents to the Mediterranean Sea. *Sci. Rep.* 9, 8024. doi: [10.1038/s41598-019-44293-1](https://doi.org/10.1038/s41598-019-44293-1)
- Watts N, Adger WN, Agnolucci P, Blackstock J, Byass P et al. 2015 Health and climate change: policy responses to protect public health. *Lancet* 386, 1861–1914. doi: [10.1016/S0140-6736\(15\)60854-6](https://doi.org/10.1016/S0140-6736(15)60854-6)
- Westerling AL, Turner MG, Smithwick EAH, Romme WH, Ryan MG 2011 Continued warming could transform Greater Yellowstone fire regimes by mid-21<sup>st</sup> century. *Proc. Natl. Acad. Sci. U. S. A.* 108, 13165–13170. doi: [10.1073/pnas.1110199108](https://doi.org/10.1073/pnas.1110199108)
- Whitehead PJP, Bauchot ML, Hureau JC, Nielsen J, Tortonese E 1986 *Fishes of the North-eastern Atlantic and the Mediterranean*. UNESCO.
- WHO 2017 Flooding: Managing Health Risks in the WHO European Region. Copenhagen, Denmark.
- WHO Europe 2017 Protecting health in Europe from climate change: 2017 update. Copenhagen, Denmark: WHO Regional Office for Europe.
- Xeidakis GS, Delimani PK, Skias SG 2006 Sea cliff erosion in the Eastern part of the North Aegean coastline, Northern Greece. *J. Environ. Sci. Heal. Part A* 41, 1989–2011. doi: [10.1080/10934520600780610](https://doi.org/10.1080/10934520600780610)
- Yang L, Qian F, Song D-X, Zheng K-J 2016 Research on Urban Heat-Island Effect. *Procedia Eng.* 169, 11–18. doi: [10.1016/j.proeng.2016.10.002](https://doi.org/10.1016/j.proeng.2016.10.002)
- Yermiyahu U, Tal A, Ben-Gal A, Bar-Tal A, Tarchitzky J et al. 2007 Rethinking Desalinated Water Quality and Agriculture. *Science (80-. ).* 318, 920. doi: [10.1126/science.1146339](https://doi.org/10.1126/science.1146339)
- Zampieri M, D'Andrea F, Vautard R, Ciais P, de Noblet-Ducoudré N et al. 2009 Hot European summers and the role of soil moisture in the propagation of Mediterranean drought. *J. Clim.* 22, 4747–4758. doi: [10.1175/2009JCLI2568.1](https://doi.org/10.1175/2009JCLI2568.1)
- Zare M, Nazari Samani AA, Mohammady M, Teimurian T, Bazrafshan J 2016 Simulation of soil erosion under the influence of climate change scenarios. *Environ. Earth Sci.* 75. doi: [10.1007/s12665-016-6180-6](https://doi.org/10.1007/s12665-016-6180-6)
- Zhang Z, Zhang J, Xu G, Zhou L, Li Y 2019 Arbuscular mycorrhizal fungi improve the growth and drought tolerance of *Zenia insignis* seedlings under drought stress. *New For.* 50, 593–604. doi: [10.1007/s11056-018-9681-1](https://doi.org/10.1007/s11056-018-9681-1)
- Zhou J, Fu B, Gao G, Lü Y, Liu Y et al. 2016 Effects of precipitation and restoration vegetation on soil erosion in a semi-arid environment in the Loess Plateau, China. *CATENA* 137, 1–11. doi: [10.1016/j.catena.2015.08.015](https://doi.org/10.1016/j.catena.2015.08.015)
- Zittis G, Hadjinicolaou P, Fnais M, Lelieveld J 2016 Projected changes in heat wave characteristics in the eastern Mediterranean and the Middle East. *Reg. Environ. Chang.* 16, 1863–1876. doi: [10.1007/s10113-014-0753-2](https://doi.org/10.1007/s10113-014-0753-2)
- Zscheischler J, Westra S, Van Den Hurk BJJM, Seneviratne SI, Ward PJ et al. 2018 Future climate risk from compound events. *Nat. Clim. Chang.* 8, 469–477. doi: [10.1038/s41558-018-0156-3](https://doi.org/10.1038/s41558-018-0156-3)

## Information about authors

### Coordinating Lead Authors

Athanasios Vafeidis:

*Christian-Albrechts University, Kiel, Germany*

### Lead Authors

Ameer Abdulla:

*Global Change Institute, Centre for Biodiversity and Conservation Science, University of Queensland, Australia*

Alberte Bondeau:

*Mediterranean Institute of Biodiversity and Ecology (IMBE, Aix-Marseille University, Avignon University, CNRS, IRD), Aix-en-Provence, France*

Lluís Brotons:

*Forest Science and Technology Centre of Catalonia (CTFC), Solsona, Spain*

Ralf Ludwig:

*Department of Geography, Ludwig Maximilian University of Munich, Munich Germany*

Michelle Portman:

*MarCoast Ecosystem Integration Lab, Technion - Israel Institute of Technology, Haifa, Israel*

Lena Reimann:

*Christian-Albrechts University, Kiel, Germany*

Michalis Voutsoukas:

*European Commission, Joint Research Centre, Ispra, Italy*

Elena Xoplaki:

*Department of Geography & Center for international Development and Environmental Research, Justus Liebig University, Giessen, Germany*

### Contributing Authors

Najet Aroua:

*Polytechnic School of Architecture and Urbanism, Algiers/ Laboratory of Architecture, Urbanism and Environmental Design, Department of Architecture, University of Biskra, Biskra, Algeria*

Lorine Behr:

*Center for International Development and Environmental Research, Justus-Liebig-University in Giessen, Germany*

Francesco Dottori:

*European Commission, Joint Research Centre, Ispra, Italy*

Joaquim Garrabou:

*Institute of Marine Sciences-CSIC, Barcelona, Spain*

Christos Giannakopoulos:

*National Observatory of Athens (NOA), Athens, Greece*

Guillaume Rohat:

*University of Geneva, Switzerland & University of Twente, the Netherlands*

Elias Symeonakis:

*Manchester Metropolitan University, United Kingdom*



# APPENDIX

## Appendix to Chapter 1 Introduction

### Coordinating Lead Authors:

Manfred A. Lange (Cyprus), Maria Carmen Llasat (Spain), Maria Snoussi (Morocco)

### Lead Authors:

Arnault Graves (Spain/France), Julien Le Tellier (Greece/France), Arnau Queralt (Spain), Grazia Maria Vagliasindi (Italy)

### Contributing Authors:

Elen Lemaitre-Curri (France), Piero Lionello (Italy), Katarzyna Marini (France), Cyril Moulin (France)

*This chapter should be cited as: Lange MA, Llasat MC, Snoussi M, Graves A, Le Tellier J, Queralt A, Vagliasindi GM 2020 Introduction. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 589-598.*

## A.1 MedECC Partners

In December 2017 the Secretariat of the Union for the Mediterranean (UfM) signed an agreement with the Plan Bleu Regional Activity Centre (UNEP/MAP) to jointly support MedECC. The MedECC Secretariat is supported and funded by UfM, through a grant provided by the Swedish International Development Cooperation Agency (SIDA), hosted by Plan Bleu in Marseille, France.

MedECC is also supported by: the French Agency for Ecological Transition (ADEME), Mediterranean Integrated Studies at Regional And Local Scales (MISTRALS), the Principality of Monaco, the Advisory Council for the Sustainable Development of Catalonia of the Government of Catalonia (CADS, Spain), Métropole Aix-Marseille Provence (France), Laboratory of Excellence OT-Med, Aix-Marseille University (France), the French National Research Institute for Sustainable Development (IRD), Association for Innovation and Research in Climate (AIR Climat, France) et ACTERRA Consulting (France).

**The Union for the Mediterranean (UfM)** is an intergovernmental institution created in 2008 which brings together all 28 countries of the European Union and 15 countries of the southern and eastern Mediterranean. UfM's mission is to enhance regional cooperation, dialogue and the implementation of projects and initiatives with tangible impact on citizens, addressing three strategic objectives: stability, human development and integration. The UfM Climate Change Expert Group (UfM CCEG) was established at the first UfM Ministerial Meeting on Environment and Climate Change in May 2014 in Athens, Greece. The UfM Ministerial Declaration of Athens expressed the need for a regional vulnerability assessment regarding climate change impacts in the Mediterranean. The role of the UfM CCEG is to advance discussions on climate change priority actions and accelerate the identification and development of concrete projects and initiatives. UfM CCEG has approved a Work Program (2017-2022) in which a specific activity "To promote a regional science-based consensus on climate impacts in the UfM region especially on the Mediterranean Basin" was included. The work undertaken by MedECC responds to this activity and is therefore supported by UfM under its Ministerial mandate.

Administered by the **United Nations Environment Programme (UNEP)** for more than four decades as the first Convention of its Regional Seas Programme and as a regional Multilateral Environ-

mental Agreement, the **Mediterranean Action Plan (MAP)** was approved in 1975 by the Mediterranean States and the European Union (EU) and was amended in 1995 after the Rio Summit to better reflect the integrated approach and indivisible nature of sustainable development. The Convention for the Protection of the Marine Environment and the Coastal Region of the Mediterranean (Barcelona Convention) was adopted in its original form in 1976 and amended in 1995 as a response to the MAP's revision. Today the UNEP/MAP – Barcelona Convention system represents the unique legally binding regime dedicated to the protection, depollution and preservation of the Mediterranean Sea and coast, and to sustainable development of its coastal area. The Barcelona Convention's 22 Contracting Parties adopted the Mediterranean Strategy for Sustainable Development (MSSD) in February 2016 at their 19th Meeting held in Greece, Athens, as a strategic guiding document for all stakeholders and partners to translate the 2030 Agenda for Sustainable Development (2030 Agenda) at regional, sub-regional and national levels. The MSSD provides an integrative policy framework for securing a sustainable future for the Mediterranean region consistent with the Sustainable Development Goals (SDGs). The creation of MedECC responds to needs and intentions expressed by the MSSD, in particular under the Flagship Initiative of its Objective 3 "Addressing climate change as a priority issue for the Mediterranean", which reads as follows: "Establish a regional science-policy interface mechanism, including the social and behavioral sciences, endorsed by all the Contracting Parties to the Barcelona Convention, with a view to preparing consolidated regional scientific assessments and guidance on climate change trends, impacts and adaptation and mitigation options".

**Plan Bleu** is one the UNEP/MAP Regional Activity Centers, located in Sophia-Antipolis and Marseille (France). Its program of work is approved every two years by the Contracting Parties to the Barcelona Convention. The main objective of Plan Bleu is to raise awareness of Mediterranean stakeholders and decision makers regarding environment and sustainable development issues in the region, through the following activities: a) Observing interactions between environment and development to support decision makers; b) Shaping possible futures for sustainable development (prospective studies, scenario building); c) Monitoring the implementation of the Mediterranean Strategy for

Sustainable Development; d) Integrating climate change as a priority; e) Supporting the transition towards a green and blue economy; f) Offering a socio-economic perspective for the appropriate management of Mediterranean resources.

MedECC has taken an active role as co-lead of the Climate Change chapter in the recent State of the Environment and Development Report (SoED 2019) coordinated by Plan Bleu.

The French **Research Institute for Sustainable Development** (IRD – *Institut de Recherche pour le Développement*) is a key French player on the international development agenda working primarily in partnership with Mediterranean and inter-tropical countries. It is based on the principle that scientific progress is necessary to further sustainable and human development, and uses an original model: equitable scientific partnership with developing countries to co-design solutions adapted to the challenges faced by humans and the planet, including pandemics, climate change, humanitarian and political crises, etc. It has been able to make an important contribution to MedECC thanks to its network and presence in fifty or so countries encouraging science and innovation as key drivers in their development.

The **Advisory Council for Sustainable Development of Catalonia** of the Government of Catalonia (CADS) is the advisory body on sustainability of the Catalan Government, whose main aim is to be an effective and successful interface between scientists, policymakers and stakeholders. The council is an active member of the European Environment and Sustainable Development Advisory Councils (EEAC), a network of advisory bodies established by national or regional governments or parliaments. Since 2005, CADS has overseen the elaboration of the periodic Report on Climate Change in Catalonia. The 3rd edition was published in September 2016 and involved 150 experts and more than 40 reviewers. A 4th edition is under preparation.

The **Ministry of Foreign Affairs and Cooperation of the Principality of Monaco** manages public policy in the following areas: immunity, diplomatic channels and consular affairs, European affairs, international and multilateral affairs and international environment.

The **French Agency for Ecological Transition (ADEME)** is active in the implementation of public policy in the areas of the environment, energy and sustainable development. ADEME provides

expertise and advisory services to businesses, local authorities and communities, government bodies and the public at large, to enable them to establish and consolidate their environmental action. As part of this work the agency helps finance projects, from research to implementation, in its areas of action.

The **Association for Innovation and Research in Climate (AIR Climat)** aims to contribute to awareness-raising on climate change issues and to help, through research and innovation, to implement new solutions in the field of carbon economy, lifestyles and their evolution. AIR Climat brings together climate change scientists and technicians. The association leads the Regional Group of Experts on Climate in the “South Region” of France (Région Sud – Provence-Alpes-Cote d’Azur), GREC-SUD, which aims to centralize, transcribe and share scientific knowledge on climate and climate change in the region. The priority objective of the group is to inform decision makers (elected representatives, local authorities) of the territory, so that scientific results are considered in public policies. Eight thematic booklets on climate change in the region have been published so far.

**ACTERRA** is a consultancy firm dedicated to environmental policy and climate change. ACTERRA has expertise in designing and implementing adaptation measures at different scales: strategic approaches and institutions at international and national level, regional and interregional initiatives, research and consulting for local authorities, operational actions at local level, etc., in particular in southern Mediterranean countries.

**Labex OT-Med** (*Objectif Terre – Bassin Méditerranéen*) is a “Laboratoire d’excellence” (LABEX) selected by the French “Investissements d’Avenir” program that brings together 10 research laboratories and 1 research federation specialized in different fields: a) environmental sciences, law, economy and social sciences; b) global change and natural hazards in the Mediterranean Basin.

MedECC has also obtained support from the **MISTRALS** program (Mediterranean Integrated STudies at Regional And Local Scales). The main objective of this program is to bring together French researchers and to reinforce international collaboration between Mediterranean countries. MISTRALS is led by CNRS, with strong partnerships with several other French research institutions. The MedECC network of experts allows the dissemination of scientific results to stakeholders, policymakers and managers to help them address

societal, environmental and economic challenges for the sustainable development of Mediterranean countries. MISTRALS addresses the following scientific questions, all of them included in the scientific objectives of MedECC: a) Links between past climate variability and evolution of Mediterranean civilizations and societies; b) Hydrological cycle in the Mediterranean and extreme rainfall events; c) Evolution of marine biogeochemistry under climate change and anthropogenic pressure; impacts on marine ecosystems; d) Change in atmospheric composition and air quality under climate change and anthropogenic pressure and impacts on health; e) Monitoring continental and marine biodiversity and their sensitivity to climate change and anthropogenic pressure; f) Changes in continental surfaces, from urban to countryside regions under climate change and anthropogenic

pressure. Action b, in particular, is the objective of the research program HYMEX (Hydrological and Mediterranean Experiment) that has a strong relationship with MedECC, with some members of both corresponding steering committees and authors of this MAR1 report in common.

A similar relationship is maintained with **MEDCLIVAR** (Mediterranean Climate Variability and Predictability), which is a scientific network endorsed by the international CLIVAR Office, to promote better communication among different scientific disciplines and develop a multidisciplinary vision of the evolution of the Mediterranean climate through studies that integrate atmospheric, marine, and terrestrial climate components at time scales ranging from paleo-reconstructions to future climate scenarios.

## A.2 Research activities in the Mediterranean region

A number of past and ongoing research programs and projects have objectives relating to the MedECC assessment process. The following compilation lists some of them, without attempting to be complete.

- The Mediterranean Experiment (MEDEX) is focused on cyclones in the Mediterranean, their impact and forecasting (Jansa et al. 2014).
- The Hydrological Mediterranean Experiment (HyMEX) aims to elucidate the hydrological cycle in the Mediterranean, with emphasis on extreme weather events, inter-annual to decadal variability of the coupled Mediterranean system, and associated trends in the context of global change (Drobinski et al. 2014).
- The Med-CORDEX initiative attempts to down-scale global climate scenarios in the Mediterranean (Ruti et al. 2016).

Other large collaborative scientific projects and networks do exist and are actively engaged in enhancing the necessary scientific knowledge base, often through a multidisciplinary and integrated approach (i.e. CIRCE-Climate Change and Impact Research: the Mediterranean Environment). The MedCLIVAR network is a case in point. Results of the investigations have been published in three books (Bolle 2003; Lionello et al. 2006; Lionello 2012) and numerous scientific articles. While often conceived as scientific platforms for better

communication and cooperation among scientists, these projects, networks and initiatives and their results are usually not easily accessible to decision- and policymakers.

The recent assessment reports of the Intergovernmental Panel on Climate Change (IPCC) and the Intergovernmental Science Policy Platform on Biodiversity and Ecosystem Services (IPBES) provide useful knowledge but they have not provided an integrated view on the Mediterranean Basin. These assessments cover only parts of the region in disconnected chapters or only some of the relevant topics (e.g. climate variability). In close cooperation with MedECC, the forthcoming Sixth Assessment Report of the IPCC will contain, for the first time, a “Cross-Chapter Paper” dedicated to the Mediterranean Basin.



### A.3 Institutional context of MedECC

The origin of the MedECC initiative can be traced back to a meeting of the MedCOP 21 in Marseille, France, on 4-5 June 2015. In the context of the “Agenda of Positive Solutions”, the idea of a network of Mediterranean Experts on Climate and environmental Change (MedECC) was first put forward and very positively received during this meeting. MedECC was envisioned as a Mediterranean derivative of both IPCC and IPBES. Just one month later, on July 9, 2015, MedECC was more officially established in the context of a side event at the international scientific conference “Our Common Future under Climate Change” (CFCC) in Paris under the auspices of the International Council of Scientific Unions (ICSU), Future Earth, the United Nations Educational Scientific and Cultural Organization (UNESCO) and a number of French research institutions. CFCC was the key scientific event to prepare for the UNFCCC COP-21 held in Paris on November 30 – December 12, 2015. The Mediterranean side event at CFCC brought together about 40 scientists and representatives of the initial supporting institutions, including Plan Bleu (United Nations Environment Programme/Mediterranean Action Plan Regional Activity Centre), Labex OT-Med, MISTRALS, UfM, the World Bank Center for Mediterranean Integration (CMI), Regional Group of Experts on Climate in the South Provence-Alpes-Cote d’Azur (GREC-SUD), French National Research Institute for Sustainable Development (IRD) and Advisory Council for the Sustainable Development of Catalonia of the Government of Catalonia (CADS) as well as decision- and policymakers.

Since 2015, MedECC has developed its network of voluntary contributions, aiming to contribute to a science-policy interface for Mediterranean sustainable development. Major steps in this context have included:

- Regular contributions to the meetings of the Union for the Mediterranean Climate Change Expert Group (UfM CCEG) since its meeting in Barcelona, Spain, on October 1 and 2 2015.
- Side events organized by MedECC during the 21st and 22nd Sessions of the Conference of the Parties to the United Nations Framework Convention on Climate Change (COP 21 and COP 22 – UNFCCC) respectively in Paris, France, in December 2015, and in Marrakesh, Morocco, in November 2016.
- A substantive contribution to the implementation of the Mediterranean Strategy for Sustainable Development (MSSD) 2016-2025, approved during the 19th Meeting of the Contracting Parties to the Convention for the Protection of the Marine Environment and the Coastal Region of the Mediterranean (Barcelona Convention) in Athens, Greece, in February 2016.
- Following the adoption of the 2017-2022 work program of the UfM CCEG, it was decided to rely primarily on MedECC to provide an assessment of the impacts of climate and environmental change in the Mediterranean Basin. Given the cross-sectorial scope of MedECC and its emphasis on environmental issues, it was decided that any possible UfM contribution to MedECC activities will be inclusive and involve Environment and Climate Change representatives of the UfM Member States, and will be pursued in coordination with all relevant UfM structures.
- The signature of an agreement in December 2017 between the Secretariat of the UfM and Plan Bleu Regional Activity Centre (UNEP/MAP) to jointly support MedECC, with the establishment of a MedECC Secretariat – financed in 2018-2020 by UfM through a grant from the Swedish International Development Cooperation Agency (SIDA) and hosted by Plan Bleu in Marseille, France, since May 2018.

### A.4 The path towards the first assessment (MAR1)

#### A.4.1 Main steps in report preparation

The Secretariat of the UfM hosted the first meeting of the MedECC ad hoc SC on April 7 and 8 2016 in Barcelona, Spain. The purpose of this meeting, to which a few key scientists and representatives of the main user institutions were invited, was to debate the group’s governance system and the

foundations of the First Mediterranean Assessment Report (MAR1). This meeting was followed by a workshop in Aix-en-Provence, France, in October 2016, where participants from 16 countries, 50 scientists and 18 representatives of end-user organizations, met to develop the general structure and outline of MAR1. From May 2017 to March 2018, thematic workshops were organized in order

to discuss the topics to be included in MAR1. More specifically the thematic workshops comprised: (i) a workshop on “Water, food, energy” (May 2017, Rabat, Morocco), (ii) a workshop on “Development, health and human security” (June 2017, Palermo, Italy), (iii) a workshop on “Ecosystems and ecosystem services” (July 2017, Marseille, France), (iv) a workshop on human impacts on Mediterranean marine ecosystems and the economy (October 2017, Monaco) and (v) a workshop on drivers of climate and environmental change (March 2018, Aix-en-Provence, France).

A call for self-nominations of Coordinating Lead Authors (CLAs) and Lead Authors (LAs) for MAR1 was widely distributed on May 31, 2018, with a deadline extended to June 22, 2018. This call was circulated widely in various scientific networks. MedECC invited self-nominations from experts from the full range of scientific, technical and socio-economic views and backgrounds linked with the Mediterranean. MedECC received 161 self-nominations from 24 countries (39% female and 61% male).

MedECC CLAs were appointed by the MedECC SC, giving priority to recognized scientific capacity and ensuring, to the highest degree possible, diversity in terms of scientific discipline, country of origin and gender. During a second phase LAs were appointed by CLAs in collaboration with the MedECC SC, using the same criteria. Contributing Authors (CAs) were selected by the CLAs, with the approval of the MedECC Coordinators. The drafting of MedECC MAR1 can thus be considered a highly participatory process.

All appointed authors are widely recognized experts who represent a broad range of subject areas and opinions in order to achieve scientific excellence. CLAs and LAs interact frequently with MedECC Assessment Coordinators, the MedECC Secretariat and the MedECC SC. In 2020, the final editing of the report and the Summary for Policymakers was undertaken by an ad hoc Editorial Committee consisting of MedECC coordinators and several CLAs and/or SC members.

In October 2018, a few scientists representing the MedECC network published the first synthesis of multiple changes in the environment that impact the livelihoods of people in the entire Mediterranean Basin (Cramer et al. 2018).

In December 2018, scientists representing the MedECC network, in collaboration with decision makers, published a preliminary assessment of risks associated with climate and environmental chang-

es in the Mediterranean region (MedECC booklet) (MedECC 2018).

An event organized on 10 October 2019 by the UfM Secretariat as part of the IV UfM Regional Forum was a good occasion for presenting this MedECC booklet with updated information and draft key messages. The event was a very good opportunity for bringing the report to the attention of the representatives of major stakeholders, all sectors of society, and especially policymakers from across the region, ahead of its finalization and communication to the ministers in charge of environment and climate change at the upcoming UfM Ministerial meeting dedicated to these issues. It had a huge impact in the media across the region, and helped raising awareness.

#### A.4.2 Tasks and responsibilities

Role attribution in report drafting and definitions are generally based on IPCC procedures, with modifications due to more limited resources than IPCC.

##### *Assessment Coordinators*

The role of coordinating the assessment is shared between two scientists. An assessment coordinator’s role is to assume responsibility for overseeing the preparation of an assessment report, as well as its Summary for Policymakers (SPM) and generally ensuring that the report is completed to a high standard, that the chapters feed into each other and that their messages are not contradicting.

The coordinators are both senior experts in their field and have experience in coordinating the work of experts. Besides overseeing the development of the assessment, the coordinators also contributed text to chapters.

The coordinators of MAR1 were Joël Guiot (CEREGE, CNRS, France) and Wolfgang Cramer (IMBE, CNRS, France).

##### *Coordinating Lead Authors (CLAs)*

CLAs take overall responsibility for coordinating major sections of an assessment report. CLAs have similar roles as LAs with the added responsibility of ensuring that major sections of the report are completed to a high standard, collated and delivered to the MedECC Secretariat in a timely manner. CLAs play a leading role in ensuring that any cross-cutting scientific or technical issues

which may involve several sections of a report, are addressed in a complete and coherent manner and reflect the latest information available.

### **Lead Authors (LAs)**

LAs are responsible for the production of designated sections on the basis of the best scientific, technical and socio-economic information available. LAs typically work in small groups, which have responsibility for ensuring that the various components of their sections are brought together in time, are of uniformly high quality and conform to any overall standards of style set for the document as a whole. During the final stages of the report preparation, when the workload may be particularly heavy, LAs are dependent upon each other to read and edit material, and to promptly agree on any changes deemed necessary.

The essence of the LAs' task is the synthesis of material drawn from all available literature. LAs are also required to take account of expert and government review comments when revising text. LAs must have the ability to develop text that is scientifically, technically and socio-economically sound and that faithfully represents, as much as possible, contributions by a wide variety of experts. LAs are required to record in the report views which cannot be reconciled with a consensus view, but which are nonetheless scientifically or technically valid.

### **Contributing Authors (CAs)**

CAs are asked to prepare technical information in the form of text, graphs or data for integration by the LAs into the draft section. Input from a wide range of contributors is a key element in the success of the MedECC assessment report. Contributed material may be edited, merged and if necessary, amended, in the course of developing the overall draft text.

### **Expert Reviewers**

Expert Reviewers provide comments on the accuracy and completeness of the scientific, technical and socio-economic content and the overall balance of the drafts. Expert reviewers comment on the text according to their own knowledge and experience.

### **MedECC Secretariat**

The MedECC Secretariat assists with all matters related to the preparation of the MedECC MAR1,

including communication, exchange of scientific information, management of documents and drafts, the review process and other matters. The MedECC Secretariat is also responsible for collaborating and coordinating with the CLAs to ensure that the chapters are delivered in a timely manner and to a high standard. The Secretariat collaborates closely with MedECC Coordinators and has been funded by UfM thanks to SIDA and based in Plan Bleu premises in Marseille, France, since May 2018.

The MedECC Secretariat is currently composed of the MedECC Science Officer, Katarzyna Marini. MedECC will seek to expand this support structure.

### **MedECC Steering Committee (SC)**

The MedECC SC decides on the functioning of MedECC and focuses on strengthening the science-policy dialogue, as well as the visibility and credibility of MedECC. The SC participated in the development and validation of the structure and outline of MAR1, reviewed the CLA and LA nominations and validated the final list of CLAs and LAs.

At the time of MAR1 the SC included:

- **Magda Bou Dagher Kharrat** (*Saint Joseph University, Beirut, Lebanon*)
- **Ghani Chehbouni** (*IRD, Rabat, Morocco*)
- **Wolfgang Cramer** (*CNRS, IMBE, Aix-en-Provence, France*)
- **Marianela Fader** (*International Centre for Water Resources and Global Change (UNESCO), Federal Institute of Hydrology, Koblenz, Germany*)
- **Carlo Giupponi** (*Ca' Foscari University and Venice International University, Italy*)
- **Arnault Graves** (*Union for the Mediterranean, Barcelona, Spain*)
- **Samir Grimes** (*National High School of Marine Sciences and Coastal Management, Algiers, Algeria*)
- **Joël Guiot** (*CNRS, CEREGE, Aix-en-Provence, France*)
- **Manfred A. Lange** (*The Cyprus Institute, Nicosia, Cyprus*)
- **Elen Lemaitre-Curri** (*Plan Bleu, Marseille, France*), until July 2020 ; **François Guerquin** (*Plan Bleu, Marseille, France*), after July 2020
- **Julien Le Tellier** (*UNEP/MAP – Barcelona Convention Secretariat, Athens, Greece*)
- **Piero Lionello** (*University of Salento, Lecce, Italy*)
- **Maria Carmen Llasat** (*University of Barcelona, Spain*)

- **Cyril Moulin** (*National Institute for Earth Sciences and Astronomy (INSU), CNRS, MISTRALS, Paris, France*)
- **Shlomit Paz** (*University of Haifa, Israel*)
- **Arnau Queralt Bassa** (*Advisory Council for the Sustainable Development of Catalonia (CADS), Barcelona, Spain*)
- **Maria Snoussi** (*Mohammed V University, Rabat, Morocco*)
- **Andrea Toreti** (*European Commission – Joint Research Centre (JRC), Ispra, Italy*)
- **Ethemcan Turhan** (*Environmental Humanities Lab, Kungliga Tekniska Högskolan (KTH), Stockholm, Sweden*)
- **Elena Xoplaki** (*Justus-Liebig Universität Gießen, Germany*)

#### **A.4.2 Key stages of MedECC report production**

##### ***Development of the detailed outline of the report – March 2018***

The overall structure of the report was discussed and validated during the scoping workshop, which took place in Aix-en-Provence, France, in October 2016. Detailed outlines of chapters were developed during thematic workshops: (i) on “Water, food, energy” (May 2017, Rabat, Morocco), (ii) on “Development, health and human security” (June 2017, Palermo, Italy), (iii) on “Ecosystems and ecosystem services” (July 2017, Marseille, France) and (iv) on drivers of climate and environmental change (March 2018, Aix-en-Provence, France).

##### ***Development of the First Order Draft (FOD) – April 2019***

Numerous on-line and on-site meetings between the MedECC SC members were held, starting with the Barcelona meeting in 2017. The MedECC Secretariat was established in May 2018. The first LA on-line meeting (on-line) was held in October 2018 to discuss the report production procedure and timeline. The 1st physical Mediterranean Assessment Report (MAR1) LA meeting was held in Milan (Italy), on March 4-7 2019. The meeting gathered 58 participants from 16 countries: MAR1 CLAs and LAs, MedECC SC Members and MedECC Partners. The main objective of this meeting was to work together on the 1st Mediterranean Assessment Report (MAR1). Authors discussed and further developed the contents of each chapter. The discussions in cross-chapter groups made it possible to verify the consistency of information provided across the whole report and identify overlaps.

The First Order Draft (FOD) of all chapters was finalized in April 2019. The FOD had at least 70% completed text for all major sections. It underwent a review internal to the assessment (Co-coordinators, CLAs, LAs, SC) in May 2019, which provided an opportunity to understand where the overlaps are between chapters, and gaps in text and expertise.

##### ***Development of the Second Order Draft (SOD) – September 2019***

Between June and September 2019 LAs held numerous on-line meetings. The SOD was the first complete draft of the technically and scientifically balanced assessment. Each chapter was required to include an Executive Summary. At this stage, authors had thought about graphics and had either identified existing graphics for inclusion or identified where graphics will be developed. Authors were also asked to be mindful of the language used in the preparation of the SOD and to present the range of scientific, technical and socio-economic evidence clearly and concisely.

In preparing the SOD and at subsequent stages of revision after review, CLAs were requested to clearly identify disparate views for which there is significant scientific or technical support, together with the relevant arguments.

The Executive Summary located at the beginning of each chapter of the report outlines the key findings arising from the assessment process. The summaries are crucial in how the outcomes of MedECC assessment are communicated to its primary audience. They are not abstracts, but a synthesis, analysis and collective expert judgment of the chapter findings. A key statement in the SPM should be readily traceable back to an Executive Summary statement(s) which in turn must be readily traceable back to a section(s) of the chapter text, which in turn should be traceable where appropriate to the primary literature through references.

##### ***Peer-review of SOD by scientific experts – October-November 2019***

An open call for independent reviewers was launched. The MedECC Coordinators and the Secretariat selected the reviewers. The reviewers come from a variety of institutions but are required to have a scientific background in the field of the assessment. The role of the reviewers was to comment on the accuracy and completeness of the scientific, technical or socio-economic contents

and the overall scientific, technical or socio-economic balance of the draft report. Expert reviewers provided the comments to the CLAs through the MedECC Secretariat in an agreed format. The MedECC Secretariat received 113 reviews.

Comments were collated and sent to authors of the assessment. Upon request, the Secretariat made available any material that is referenced in the document being reviewed that is not available in the international published literature. Authors needed to have this material available in case a request is made.

### ***MedECC Steering Committee and Coordinating Lead Authors meeting – December 2019***

The objective of this meeting was to discuss the results of the external peer review by scientific experts, develop the Summary for Policymakers and to plan next steps for producing the Final Draft (FD). The SPM (of about 20 pages) was produced by a team consisting of Coordinators, CLAs and selected LAs. The SPM is primarily based on the chapters' Executive Summaries. It contains the main policy-relevant, but policy-neutral findings of the assessment in synthesized and less technical language, generally in the form of top key messages and presented without reference to the main chapters. These messages represent the highest level of synthesis of the assessment and may be structured differently from the set of main findings in the SPM.

### ***Development of the Final Draft (FD) – February-April 2020***

The FD incorporated further development of the assessment by chapter teams as well as the results of the peer-review of SOD by scientific experts. All review comments require a written response – the resulting tables will be made publicly available upon request. Where authors reject a comment, they will provide written justification for doing so.

### ***Review of the draft SPM by stakeholders – May-June 2020***

The draft of the SPM was reviewed by governments and other relevant stakeholders through an open and transparent process. Drafts of report chapters were also provided as a source of the findings provided in the SPM.

The SPM approval procedure is organized with the UNEP/MAP – Barcelona Convention Secretariat and its Plan Bleu Regional Activity Centre,

through their Focal Points and/or the Members of the Mediterranean Commission on Sustainable Development (MCSD), as well as with UfM Member State representatives within the regional Climate Change Expert Group (CCEG). The exact procedure has been established in close cooperation with UNEP/MAP and UfM.

During review and approval by policymakers, the scientific content of the report and the clarity of its presentation were discussed.

### ***Finalization of draft Assessment Report and SPM for Plenary discussion – July-August 2020***

The final draft will take into consideration all comments from the review by stakeholders and policymakers. Where authors reject a comment, they will provide written justification for doing so. A final draft of the SPM including key messages and graphics was also developed during this period. The preparation of the final version of the report, considering all stakeholder and expert comments, was undertaken by the Coordinators, CLAs and LAs in consultation with the MedECC Editorial Committee. This stage is critical for the coordination of key findings and policy relevant messages in the SPM, for developing graphics, for quality assurance of chapters and ensuring consistency and traceability of confidence statements between the SPM and the chapters.

The final draft should reflect comments made by policymakers, stakeholders and scientific experts. If necessary, authors, together with review editors and reviewers can try to resolve areas of major differences of opinion. Reports should describe different, possibly controversial, scientific, technical and socio-economic views on a given subject, particularly if they are relevant to the policy debate. The final report will credit all Coordinators, CLAs, LAs, CAs, reviewers and MedECC Editorial Committee and other contributors, as appropriate, by name and affiliation.

### ***Plenary consultation on the SPM – 22 September 2020***

The revised Summary for Policymakers (SPM) of the First Mediterranean Assessment Report (MAR1) was the subject of the plenary consultation with policymakers, governments, decision makers and stakeholders, which took place on 22 September 2020 (in Marseille, France and on-line). Due to the health crisis, the plenary consultation was predominantly held virtually. The particular aim of the plenary consultation was to ascertain that

MAR1 findings, as presented in the SPM, are fully comprehensible and unambiguous and that the remarks from the on-line consultation were well integrated. This meeting gathered more than 100 participants. It was attended by representatives from 15 countries. The agreed changes were implemented in the SPM, which should no longer be changed (except for editorial and technical modifications). At the end of the meeting the conclusions and recommendations were adopted. Participants in the plenary session were: Focal Points of the

Union for the Mediterranean Climate Change Expert Group (UfM CCEG) and the UfM Environment Task Force, Focal Points of Plan Bleu/Regional Activity Centre (United Nations Environment Programme / Mediterranean Action Plan (UNEP/MAP)), Members of the Steering Committee of the Mediterranean Commission on Sustainable Development (MCSD), MedECC Coordinators, Secretariat, Steering Committee members and MAR1 Coordinating Lead Authors, Plan Bleu, UNEP/MAP and UfM Secretariat representatives.




# B APPENDIX

## Maps of seasonal temperature and precipitation changes for the Mediterranean Basin

**Authors:**

George Zittis (Cyprus), Samuel Somot (France) and Filippo Giorgi (Italy)

*This chapter should be cited as: Zittis G, Somot S, Giorgi F 2020 Appendix B – Maps of seasonal temperature and precipitation changes for the Mediterranean Basin. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guot J, Marini K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 599-612.*





## Table of contents

<b>Introduction and scope</b> .....	<b>602</b>
<b>Data and methods</b> .....	<b>602</b>
<b>Projected temperatures</b> .....	<b>604</b>
Winter temperature .....	604
Spring temperature .....	605
Summer temperature .....	606
Autumn temperature .....	607
<b>Precipitation</b> .....	<b>608</b>
Winter precipitation .....	608
Spring precipitation .....	609
Summer precipitation .....	610
Autumn precipitation .....	611
<b>References</b> .....	<b>612</b>



## Introduction and scope

Projected changes of key meteorological variables (for example, precipitation and temperature) are not, in terms of magnitude, expected to be consistent throughout the Mediterranean region. Similarly, according to global and regional model projections and due to several climatic feedbacks, these changes will not be uniformly distributed throughout the year. While changes in seasonal temperature and precipitation were also discussed in *Chapter 2*

of the First Mediterranean Assessment Report (MAR1), here we present a complementary but more informative assessment that could provide useful information for impact and adaptation studies and also motivate mitigation actions. This is presented in the form of an atlas that includes all the Mediterranean countries, a range of future periods and the two greenhouse gas emission pathways (RCP2.6 and RCP8.5) that were discussed primarily in MAR1.

## Data and methods

At the time of writing, the most comprehensive and populated ensemble of regional climate projections, that adequately includes most of the Mediterranean region, is the EURO-CORDEX (Jacob et al. 2020), which is the European initiative of the Coordinated Regional Downscaling Experiment (Giorgi and Gutowski 2015). This set of state-of-the-art regional projections, available in a horizontal resolution of about 12 km is the basis of the present assessment. The full list of experiments taken into account as well as the availability per variable and scenario are presented in *Table 1*.

When assessing future climate change, it is important to specify the reference period to which climate projections are compared, along with future "time slices" of particular interest. In MAR1 and the present atlas, we use 20-year periods. This length is sufficient to smooth part of the high-frequency natural climate variability that may otherwise mask the forced trend, but it is short enough to assume that climate does not change much during the 20 years covered. For the reference period, we chose the last decades of the 20th century (1980-1999). For the future, we kept 20-year time slices in order to sample the same level of internal variability as in the reference period. We divided the 21st century into 20-year time slices with a near-future period (2020-2039), a mid-term period centred in 2050 (2040-2059) and a far-future period close to the end of the 21st century (2080-2099). The mid-21st century period is arguably of particular interest for many stakeholders,

especially for mid-term adaptation. The end of the 21st century period is also of interest for stakeholders working on mitigation targets and involved in very long-term planning (e.g., for the design and planning of dams, forests or cities).

In terms of seasons, we use the boreal hemisphere definition for winter (December-February), spring (March-May), summer (June-August) and autumn (September-November), for analysis of both temperature and precipitation. Maps of future projections on an annual basis are discussed in the main text of MAR1 and are therefore not presented here.

For future climate, an important part of the uncertainty is related to the evolution of socio-economic development. To be able to propose future climate projections according to various possible socio-economic and climate policy trajectories, we follow the Representative Concentration Pathways or RCPs, defined in *Box 2.1* of MAR1 (Meinshausen et al. 2011). Here, we focus mostly on two of such pathways which encompass the range of IPCC-AR5, CMIP5 and CORDEX simulations: the 'business as usual' scenario of high emissions (RCP8.5) and a more optimistic pathway closest to meeting the UN-FCCC Paris Agreement main targets (RCP2.6). These scenarios have been chosen also due to model projection availability constraints at the regional scale.

GLOBAL MODEL	REGIONAL MODEL	TEMPERATURE AT 2M		PRECIPITATION	
		RCP2.6	RCP8.5	RCP2.6	RCP8.5
CNRM-CERFACS-CNRM-CM5 (r1i1p1)	CLMcom-CCLM4-8-17_v1		•		•
CNRM-CERFACS-CNRM-CM5 (r1i1p1)	CNRM-ALADIN53_v1	•	•		
CNRM-CERFACS-CNRM-CM5 (r1i1p1)	CNRM-ALADIN63_v2	•	•	•	•
CNRM-CERFACS-CNRM-CM5 (r1i1p1)	DMI-HIRHAM5_v2		•		•
CNRM-CERFACS-CNRM-CM5 (r1i1p1)	KNMI-RACMO22E_v2	•	•	•	•
CNRM-CERFACS-CNRM-CM5 (r1i1p1)	RMIB-UGent-ALARO-0_v1	•	•	•	•
CNRM-CERFACS-CNRM-CM5 (r1i1p1)	SMHI-RCA4_v1		•		•
ICHEC-EC-EARTH (r12i1p1)	CLMcom-CCLM4-8-17_v1	•	•	•	•
ICHEC-EC-EARTH (r12i1p1)	DMI-HIRHAM5_v1		•		•
ICHEC-EC-EARTH (r12i1p1)	KNMI-RACMO22E_v1	•	•	•	•
ICHEC-EC-EARTH (r12i1p1)	SMHI-RCA4_v1	•	•	•	•
ICHEC-EC-EARTH (r3i1p1)	KNMI-RACMO22E_v1		•		•
ICHEC-EC-EARTH (r3i1p1)	SMHI-RCA4_v1		•		•
IPSL-IPSL-CM5A-LR (r1i1p1)	GERICS-REM02015_v1	•		•	
IPSL-IPSL-CM5A-MR (r1i1p1)	SMHI-RCA4_v1		•		•
MOHC-HadGEM2-ES (r1i1p1)	CLMcom-CCLM4-8-17_v1		•		•
MOHC-HadGEM2-ES (r1i1p1)	DMI-HIRHAM5_v1		•		•
MOHC-HadGEM2-ES (r1i1p1)	KNMI-RACMO22E_v2	•	•	•	•
MOHC-HadGEM2-ES (r1i1p1)	SMHI-RCA4_v1	•	•	•	•
MPI-M-MPI-ESM-LR (r1i1p1)	CLMcom-CCLM4-8-17_v1		•		•
MPI-M-MPI-ESM-LR (r1i1p1)	MPI-CSC-REM02009_v1	•	•	•	•
MPI-M-MPI-ESM-LR (r1i1p1)	SMHI-RCA4_v1	•	•	•	•
NCC-NorESM1-M (r1i1p1)	DMI-HIRHAM5_v2		•		•
NCC-NorESM1-M (r1i1p1)	GERICS-REM02015_v1		•		•
NCC-NorESM1-M (r1i1p1)	KNMI-RACMO22E_v1		•		
NCC-NorESM1-M (r1i1p1)	SMHI-RCA4_v1	•	•	•	•
NOAA-GFDL-GFDL-ESM2G (r1i1p1)	GERICS-REM02015_v1	•		•	
	TOTAL	14	25	13	23


**Table B.1 | List of EURO-CORDEX experiments taken into account in the presented assessment and availability of variables.**

## Projected temperatures

### Winter temperature

Projected winter temperature changes for the Mediterranean are presented in *Figure B.1*. For pathway RCP2.6, the EURO-CORDEX multi-model ensemble suggests a relatively mild increase that is not expected to exceed 1°C-2°C for all sub-periods (*Fig. B.1 – left panels*). Particularly for the middle and late-21st century, the projected winter warming is slightly higher over the eastern part of the Mediterranean. Nevertheless, the regional differences are not so evident. For the business-as-usual RCP8.5, the near-future winter temperature projections are of the same magnitude as the worst-case

ones for RCP2.6 (*Fig. B.1 – top right*). Already by mid-century, winter warming is expected to reach 3°C, with respect to the historical reference, in many parts of the region. This is the case mainly in high-elevation regions, such as the Atlas Mountains, the Alps, Anatolia and parts of the Balkan Peninsula, highlighting that winter warming could be enhanced by positive snow-albedo feedbacks. For the end of the current century (*Fig. B.1 – bottom right*), warming is projected to intensify and exceed 4°C in most of the region. In the hotspot mountainous areas, this warming is projected to reach 6°C with respect to the reference period.




**Figure B.1** | Projected changes in winter (December, January, February) temperatures between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (left panels) and RCP8.5 (right panels).

**Spring temperature**

The projected changes during boreal spring are presented in Fig. B.2. Under pathway RCP2.6 and for near-future and mid-century, the projected spring warming is somehow higher than during the winter season. For southern Mediterranean regions, such as the Maghreb, this warming will reach 2.5°C-3°C. By the end of the century,

the warming is not projected to exceed 2°C in most of the Mediterranean, with the exception of Anatolia. As expected, under the business-as-usual pathway (Fig. B.2 - right panels), the spring warming is projected to follow the same spatial patterns. The EURO-CORDEX ensemble suggests warming between 4°C and 5°C, with higher values in North Africa and the mountainous regions of the Mediterranean.




**Figure B.2 | Projected changes in spring (March, April, May) precipitation between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (left panels) and RCP8.5 (right panels).**

**Summer temperature**

As discussed in MAR1, the projected summer temperature increase in the Mediterranean region, mainly in the South, is particularly high. This is the case mostly for RCP8.5 (Fig. B.3 – right panels). By mid-century, summer warming will likely exceed 3°C in many parts of the region. The late-21st century RCP8.5 projections suggest that this warming will further intensify and

locally exceed levels of 6°C-6.5°C. This is expected mainly for southern latitudes and regions such as the Maghreb, the Iberian Peninsula and Anatolia, as well as the Alps. Soil-atmosphere interactions have been found to play a role in this summer warming amplification (e.g., Zittis et al. 2014). In contrast, under the more moderate RCP2.6 pathway, summer temperature changes will likely be less than 2°C throughout the Mediterranean (Fig. B.3 – left panels).




**Figure B.3 | Projected changes in summer (June, July, August) precipitation between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (left panels) and RCP8.5 (right panels).**

### Autumn temperature

Maps of projected changes for the transitional season of autumn are presented in Fig. B.4. The spatial patterns are very similar to those for the summer season, however, the magnitude of warming is lower. For RCP2.6, future changes range between 1°C and 1.5°C for all future periods (Fig. B.4 – left panels). The only exception is the western part of the Mediterranean and middle century projections that are expected

to be somehow higher (up to 2°C). Under the high-emission pathway (Fig. B.4 – right panels), the near future changes are comparable to those for RCP2.6, while the middle century projections indicate that the autumn warming, with respect to the reference period, is not expected to exceed 3°C. The end-of-century projections under RCP8.5 indicate further warming of up to 5°C–6°C, expected mainly for the southern Mediterranean (for example, the Maghreb, the Iberian Peninsula and Anatolia), as well as the Alps.




**Figure B.4** | Projected changes in autumn (September, October, November) precipitation between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (left panels) and RCP8.5 (right panels).

## Precipitation

### Winter precipitation

Projected changes for boreal winter (December, January and February) precipitation are presented in Fig. B.5 as percentage differences from the historical reference period. Under both emission pathways, and all future periods a North-South gradient of the climate change signal is evident. For southern Europe and the northern Mediterranean territories, winter precipitation is expected to change slightly or in-

crease up to 10%-30%. In contrast, for the drier southern parts of the region, winter precipitation, which is more critical for replenishing water resources, is projected to decrease between 20% and 50%. For the Maghreb region, which is a hotspot of drying, the projected winter precipitation decrease could even exceed 60%. The projected changes are higher for the end of the current century and this is likely the case for both pathways under investigation.




**Figure B.5 | Projected changes in winter (December, January, February) precipitation between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (Left panels) and RCP8.5 (right panels).**



### Spring precipitation

For spring (March to May) precipitation changes, the North-South gradient is still evident, however, this pattern is less pronounced (*Fig. B.6*). Noteworthy, for the next two decades (i.e., the near future sub-period), the projected changes are higher for RCP2.6, indicating that even under low-emission pathways, global warming could introduce changes with high impact at

regional scales. This is the case mainly for the Maghreb region (*Fig. B.6 – top panels*). For the mid-21st century, the EURO-CORDEX ensemble suggests similar changes for both pathways. Small changes are expected for southern Europe, while for North Africa, a decrease of 10%-30% in spring precipitation is expected. Under RCP8.5, seasonal drying is expected to intensify towards the end of the century.




**Figure B.6 | Projected changes in spring (March, April, May) precipitation between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (left panels) and RCP8.5 (right panels).**

**Summer precipitation**

The projected precipitation changes of the summer season are presented in Fig. B.7. For the northern Mediterranean and southern Europe, the projected changes are relatively mild and were found to range between  $\pm 10\%$ . For parts of the southern Mediterranean, a precipitation increase is projected, however, this is not always significant in actual precipitation amounts since

summer precipitation is limited (not shown). This increase, that varies in magnitude between the different pathways and time periods, is likely related to a northward expansion of the inter-tropical convergence zone (Evans 2010). A strong summer precipitation decline is evident only for the end of the current century and the RCP8.5 pathway (Fig. B.7 – bottom right). This is apparent for most of the Mediterranean and southern Europe.




**Figure B.7 | Projected changes in summer (June, July, August) precipitation between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (left panels) and RCP8.5 (right panels).**

### Autumn precipitation

For the northern Mediterranean, the projected change in autumn precipitation is limited ( $\pm 10\%$ ). For other regions (e.g., Southwest Mediterranean), the future change signal is not consistent between the two pathways. For the Maghreb,

precipitation increases [10%-30%] have been projected under RCP2.6 (Fig. B.8 – left panels). On the contrary under RCP8.5, this region is projected to be subject to strong drying trends (up to 40%-50%), particularly towards the end of the current century (Fig. B.8 – right panels).



**Figure B.8** | Projected changes in autumn (September, October, November) precipitation between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean results of the EURO-CORDEX high-resolution simulations for pathways RCP2.6 (Left panels) and RCP8.5 (right panels).

## References

- Evans JP (2010) Global warming impact on the dominant precipitation processes in the Middle East. *Theor Appl Climatol* 99:389–402. doi: [10.1007/s00704-009-0151-8](https://doi.org/10.1007/s00704-009-0151-8)
- Giorgi F, Gutowski WJ (2015) Regional Dynamical Downscaling and the CORDEX Initiative. *Annu Rev Environ Resour* 40:467–490. doi: [10.1146/annurev-environ-102014-021217](https://doi.org/10.1146/annurev-environ-102014-021217)
- Jacob D, Teichmann C, Sobolowski S, et al (2020) Regional climate downscaling over Europe: perspectives from the EURO-CORDEX community. *Reg Environ Chang* 20:. doi: [10.1007/s10113-020-01606-9](https://doi.org/10.1007/s10113-020-01606-9)
- Meinshausen M, Smith SJ, Calvin K, et al (2011) The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Clim Change* 109:213–241. doi: [10.1007/s10584-011-0156-z](https://doi.org/10.1007/s10584-011-0156-z)
- Zittis G, Hadjinicolaou P, Lelieveld J (2014) Role of soil moisture in the amplification of climate warming in the eastern Mediterranean and the Middle East. *Clim Res* 59:27–37. doi: [10.3354/cr01205](https://doi.org/10.3354/cr01205)



# APPENDIX

**List of acronyms,  
chemical symbols  
and scientific units**

%	Percent, part per hundred
‰	Per-mil, part per thousand
°C	Degree Celsius
\$	American dollar
€	Euro
<b>A1</b>	IPCC SRES storyline and scenario family: a future world of very rapid economic growth, global population that peaks in mid-century and declines thereafter, and rapid introduction of new and more efficient technologies
<b>A2</b>	IPCC SRES storyline and scenario family: a very heterogeneous world with continuously increasing global population and regionally oriented economic growth that is more fragmented and slower than in other storylines
<b>ACP</b>	Asbestos Cement Pipeline
<b>AD</b>	Anno Domini
<b>ADEME</b>	French Agency for Ecological Transition
<b>AGCM</b>	Atmospheric General Circulation Model
<b>AI</b>	Aridity Index
<b>AIR Climat</b>	Association for Innovation and Research in Climate
<b>Al</b>	Aluminum
<b>AMF</b>	Arbuscular mycorrhiza fungi
<b>AMO</b>	Atlantic Multidecadal Oscillation
<b>Approx.</b>	Approximately
<b>AR3</b>	IPCC Third Assessment Report
<b>AR4</b>	IPCC Fourth Assessment Report
<b>AR5</b>	IPCC Fifth Assessment Report
<b>AR6</b>	IPCC Sixth Assessment Report
<b>Art.</b>	Article
<b>As</b>	Arsenic
<b>a.s.l.</b>	above sea level
<b>ASP</b>	Amnesic Shellfish Poisoning
<b>atm</b>	standard atmosphere (unit of pressure)
<b>B</b>	Boron
<b>B1</b>	IPCC SRES storyline and scenario family: a convergent world with the same global population as in the A1 storyline but with rapid changes in economic structures toward a service and information economy, with reductions in material intensity, and the introduction of clean and resource-efficient technologies
<b>B2</b>	IPCC SRES storyline and scenario family: a world in which the emphasis is on local solutions to economic, social, and environmental sustainability, with continuously increasing population (lower than A2) and intermediate economic development.
<b>BC</b>	Before Christ
<b>BiOS</b>	Bimodal Oscillating System
<b>BP</b>	Before Present
<b>C</b>	Carbon
<b>CA</b>	Contributing Author
<b>ca.</b>	circa, meaning approximately
<b>CAAGR</b>	Compounded Average Annual Growth Rate
<b>CaCO<sub>3</sub></b>	Calcium carbonate
<b>CADS</b>	Advisory Council for the Sustainable Development of the Government of Catalonia
<b>cal</b>	calibrated years

<b>cap</b>	capita
<b>CAP</b>	Common Agricultural Policy
<b>CBD</b>	Convention on Biological Diversity
<b>CCA</b>	Climate Change Adaptation
<b>CCEG</b>	Climate Change Expert Group
<b>Cd</b>	Cadmium
<b>CDM</b>	Clean Development Mechanisms
<b>CE</b>	Common Era or Current Era
<b>CEDAW</b>	Convention on the Elimination of All Forms of Discrimination against Women
<b>CFCC</b>	Our Common Future under Climate Change
<b>CH<sub>4</sub></b>	Methane
<b>CIMPAL</b>	Cumulative IMPacts of invasive ALien species
<b>CIRCE</b>	Climate change and impact research: the Mediterranean environment
<b>CL</b>	Cutaneous Leishmaniasis
<b>CLA</b>	Coordinating Lead Author
<b>cm</b>	centimeter
<b>CMEMS</b>	Copernicus Marine Environment Monitoring Service
<b>CMIP3</b>	Couples Model Intercomparison Project Phase 3
<b>CMIP5</b>	Couples Model Intercomparison Project Phase 5
<b>CNR-ISMAR</b>	Italian National Research Council, Institute for Marine Science
<b>CNRM</b>	French National Centre for Meteorological Research
<b>CNRS</b>	French National Centre of Scientific Research
<b>Co</b>	Cobalt
<b>CO<sub>2</sub></b>	Carbon dioxide
<b>CO<sub>2e</sub></b>	Carbon dioxide equivalent
<b>CO<sub>3</sub><sup>2-</sup></b>	Carbonate ion
<b>COP</b>	Conference of the Parties
<b>CORDEX</b>	Coordinated Regional Climate Downscaling Experiment
<b>CP</b>	Contracting Party
<b>CPRCM</b>	Convection-Permitting Regional Climate Models
<b>Cr</b>	Chromium
<b>CRI-MED</b>	Coastal Risk Index
<b>CRU</b>	Climatic Research Unit
<b>CRU TS</b>	Climatic Research Unit Timeseries
<b>CS</b>	“Conservative” Scenario
<b>CSO</b>	Civil Society Organisation
<b>CSP</b>	Concentrated Solar Power
<b>CT</b>	Conventional Tillage
<b>CTD</b>	conductivity-temperature-depth
<b>Cu</b>	Copper
<b>DAISIE</b>	Delivering Alien Invasive Species Inventories for Europe
<b>DI</b>	Development Index
<b>Dii</b>	Desert industrial energy initiative
<b>DJF</b>	December-January-February
<b>DSP</b>	Diarrheic Shelfish Poisoning
<b>e.g.</b>	for example
<b>EC</b>	Emerging Contaminants
<b>EC</b>	European Commission
<b>ECtHR</b>	European Court of Human Rights
<b>ED</b>	Endocrine Disruptors
<b>eDNA</b>	environmental deoxyribonucleic acid
<b>EEA</b>	European Environment Agency

<b>EEAC</b>	European Environment and Sustainable Development Advisory Councils
<b>EF</b>	Emission Factor
<b>E</b>	Exa ( $10^{18}$ )
<b>ELA</b>	Equilibrium Line Altitude
<b>EMT</b>	Eastern Mediterranean Transient
<b>ENSO</b>	El Niño Southern Oscillation
<b>EPPO</b>	European and Mediterranean Plant Protection Organization
<b>ESD</b>	education for sustainable development
<b>etc.</b>	et cetera meaning “and other similar things”
<b>EU</b>	European Union
<b>EWG</b>	Expert Working Groups
<b>EWS</b>	Early Warning Systems
<b>F</b>	Fishing mortality rate i.e. the catch relative to the size of the stock (the proportion of fish caught and removed by fishing).
<b>FAO</b>	Food and Agriculture Organisation
<b>FD</b>	Final Draft
<b>Fe</b>	Iron
<b>Fig.</b>	Figure
<b>FMSY</b>	The maximum rate of fishing mortality (the proportion of a fish stock caught and removed by fishing)
<b>FOD</b>	First Order Draft
<b>FPS</b>	Flagship Pilot Study
<b>FRA</b>	Fisheries Restricted Area
<b>G</b>	Giga ( $10^9$ )
<b>g</b>	gram
<b>GCM</b>	Global Climate Model
<b>GDI</b>	Gender Development Index
<b>GDP</b>	Gross Domestic Product
<b>GFCM</b>	General Fisheries Commission for the Mediterranean
<b>GHG</b>	Greenhouse Gases
<b>GII</b>	Gender Inequality Index
<b>GIS</b>	Geographic Information System
<b>GNH</b>	Gross National Happiness
<b>GNI</b>	Gross National Income
<b>GOA-ON</b>	Global Ocean Acidification-Observing Network
<b>GOLT</b>	Gill-Oxygen Limitation Theory
<b>GRACE</b>	NASA's Gravity Recovery and Climate Experiment
<b>GREC-SUD</b>	Regional Group of Experts on Climate in the “South Region” of France (Provence-Alpes-Cote d'Azur)
<b>GWL</b>	Global Warming Level
<b>h</b>	hour
<b>H<sup>+</sup></b>	Hydrogen ion
<b>ha</b>	hectare
<b>HAB</b>	Harmful Algal Bloom
<b>HCO<sub>3</sub><sup>-</sup></b>	Bicarbonate ion
<b>HD</b>	Human Development
<b>HDPE</b>	High-Density Polyethylene
<b>Hg</b>	Mercury
<b>i.e.</b>	that is
<b>IBT</b>	Inter-Basin Transfer
<b>ICCPR</b>	International Covenant on Civil and Political Rights



<b>ICESCR</b>	International Covenant on Economic, Social and Cultural Rights
<b>ICRC</b>	Convention on the Rights of the Child
<b>IGO</b>	Intergovernmental Organization
<b>IHDI</b>	Inequality-adjusted Human Development Index
<b>IIASA</b>	International Institute for Applied Systems Analysis
<b>IPBES</b>	Intergovernmental Science Policy Platform on Biodiversity and Ecosystem Services
<b>IPCC</b>	Intergovernmental Panel on Climate Change
<b>IRD</b>	French National Research Institute for Sustainable Development
<b>ISBA</b>	Interactions between Soil, Biosphere, and Atmosphere
<b>IUCN</b>	International Union for Conservation of Nature
<b>IWRM</b>	Integrated Water Resources Management
<b>J</b>	Joule
<b>JJA</b>	June-July-August
<b>K</b>	Potassium
<b>ka</b>	kilo annum, thousand calendar years ago
<b>k</b>	kilo (10 <sup>3</sup> )
<b>l</b>	liter
<b>LA</b>	Lead Author
<b>LabEx OT- Med</b>	Laboratory of Excellence "Objectif Terre - Bassin Méditerranéen"
<b>LCOE</b>	levelized Cost Of Energy
<b>LDC</b>	Less Developed Country
<b>LDN</b>	Land Degradation Neutrality
<b>LID</b>	low-impact development
<b>LIW</b>	Levantine Intermediate Water
<b>LMPA</b>	Large Marine Protected Area
<b>LPI</b>	Living Planet Index
<b>M</b>	Mega (10 <sup>6</sup> )
<b>m</b>	meter
<b>m</b>	milli (10 <sup>-3</sup> )
<b>MAES</b>	Mesophotic Assemblages Ecological Status
<b>MAP</b>	Mediterranean Action Plan
<b>MAR</b>	Managed Aquifer Recharge
<b>MAR1</b>	1st Mediterranean Assessment Report
<b>max</b>	maximum
<b>MCSd</b>	Mediterranean Commission on Sustainable Development
<b>MECIDS</b>	Middle East Consortium on Infectious Disease Surveillance
<b>Med</b>	Mediterranean
<b>MedCLIVAR</b>	Mediterranean CLImate VARIability and Predictability
<b>MedCOF</b>	Mediterranean Climate Outlook Forum
<b>MedECC</b>	Mediterranean Experts on Climate and environmental Change
<b>MEDENER</b>	Mediterranean Association of National Agencies for Energy Management
<b>MEDREG</b>	Association of Mediterranean Energy Regulators
<b>MENA</b>	Middle East and North Africa
<b>MEP</b>	Mediterranean Energy Perspectives
<b>min</b>	minimum
<b>MISTRALS</b>	Mediterranean Integrated STudies at Regional And Local Scales
<b>MMEs</b>	Mass Mortalities Events
<b>Mn</b>	Manganese
<b>MP</b>	Microplastics (plastic particles with a longest dimension <5 mm)
<b>MPA</b>	Marine Protected Area



<b>MPI</b>	Multidimensional Poverty Index
<b>MS</b>	Mediterranean Sea
<b>MS</b>	Member States
<b>MSFD</b>	Marine Strategy Framework Directive
<b>MSSD</b>	Mediterranean Strategy for Sustainable Development
<b>Mt</b>	Mountain
<b>MTE</b>	Metal Trace Elements
<b>n</b>	nano ( $10^{-9}$ )
<b>N</b>	Nitrogen
<b>N<sub>2</sub></b>	molecular nitrogen
<b>N<sub>2</sub>O</b>	Nitrous oxide
<b>NAO</b>	North Atlantic Oscillation
<b>NDC</b>	Nationally Determined Contributions
<b>NGO</b>	Non-Governmental Organization
<b>NH<sub>3</sub></b>	Ammonia
<b>NH<sub>4</sub><sup>+</sup></b>	Ammonium ion
<b>Ni</b>	Nickel
<b>NI</b>	Nitrification and urease inhibitors
<b>NIS</b>	Non-Indigenous Species
<b>no</b>	number
<b>NO</b>	Nitric oxide
<b>NO<sub>2</sub></b>	Nitrogen dioxide
<b>NO<sub>3</sub></b>	Nitrate
<b>NO<sub>3</sub><sup>-</sup></b>	Nitrate ion
<b>NO<sub>x</sub></b>	Nitrogen oxide
<b>Nr</b>	Reactive nitrogen
<b>NRW</b>	Non-Revenue Water
<b>NT</b>	No Tillage
<b>NW</b>	NorthWestern
<b>NWFP</b>	Non-Wood Forest Product
<b>O<sub>3</sub></b>	Ozone
<b>OAR</b>	Ocean Acidification Refugia
<b>OCP</b>	Organochlorinated phenyl
<b>OECD</b>	Organisation for Economic Co-operation and Development
<b>OHCHR</b>	Office of the High Commissioner for Human Rights
<b>OME</b>	Mediterranean Energy Observatory
<b>OPHI</b>	Oxford Poverty & Human Development Initiative
<b>Osm</b>	Osmole
<b>P</b>	Penta ( $10^{15}$ )
<b>P</b>	Phosphorus
<b>p</b>	probability value (significance)
<b>PAE</b>	Phthalic Acid Ester
<b>PAH</b>	Polycyclic Aromatic Hydrocarbon
<b>PAI</b>	Pesticide Active Ingredients
<b>Pb</b>	Lead
<b>PCB</b>	Polychlorinated biphenyl
<b>pCO<sub>2</sub></b>	Partial pressure of carbon dioxide
<b>PES</b>	Payments for Ecosystem Services
<b>PET</b>	Potential Evapotranspiration
<b>pH</b>	Figure expressing the acidity or alkalinity of a solution on a logarithmic scale on which 7 is neutral, lower values are more acid and higher values more alkaline

<b>PI</b>	Precipitation Index
<b>PM</b>	Particulate Matter
<b>PM10</b>	PM with diameter below 10 µm
<b>PM2.5</b>	PM with diameter below 2.5 µm
<b>POP</b>	Persistent Organic Pollutant
<b>ppb</b>	parts per billion
<b>PPP</b>	Public-Private Partnership
<b>PRA</b>	Pest Risk Analysis
<b>PS</b>	Proactive Scenario
<b>PSP</b>	Paralytic Shellfish Poisoning
<b>Psu</b>	Practical salinity unit
<b>PV</b>	Photovoltaic
<b>PVC</b>	Polyvinyl chloride
<b>R&amp;D</b>	Research and Development
<b>RCM</b>	Regional Climate Model
<b>RCP</b>	Representative Concentration Pathway
<b>RCSM</b>	Regional Climate System Model
<b>RES</b>	Renewable Energy Sources
<b>resp.</b>	respectively
<b>RS</b>	Reference Scenario
<b>s</b>	second
<b>SC</b>	Steering Committee
<b>SD</b>	Sustainable Development
<b>SDG</b>	Sustainable Development Goal
<b>SDM</b>	Species Distribution Models
<b>Se</b>	Selenium
<b>SE</b>	SouthEastern
<b>SEMC</b>	Southern and Eastern Mediterranean country
<b>SIDA</b>	Swedish International Development Cooperation Agency
<b>SLM</b>	Sustainable Land Management
<b>SLP</b>	Sea-Level Pressure
<b>SO<sub>2</sub></b>	Sulfur dioxide
<b>SOC</b>	Soil Organic Carbon
<b>SOD</b>	Second Order Draft
<b>SoED 2019</b>	State of the Environment and Development Report coordinated by Plan Bleu
<b>SOM</b>	Soil Organic Matter
<b>SPI</b>	Standardized Precipitation Index
<b>SPM</b>	Summary for Policymakers
<b>SR1.5</b>	IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways
<b>SRES</b>	IPCC Special Report on Emissions Scenarios
<b>SREX</b>	IPCC Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation
<b>SROCC</b>	IPCC Special Report on the Ocean and Cryosphere in a Changing Climate
<b>SSP</b>	Shared Socio-Economic Pathway
<b>SST</b>	Sea Surface Temperature
<b>T</b>	Tera (10 <sup>12</sup> )
<b>t</b>	tonne
<b>Ti</b>	Titanium
<b>toe</b>	tonne of oil equivalent

<b>TS</b>	Transition Scenario
<b>TWW</b>	Treated Waste Water
<b>UDHR</b>	Universal Declaration of Human Rights
<b>UfM</b>	Union for the Mediterranean
<b>UfM CCEG</b>	Union for the Mediterranean Climate Change Expert Group
<b>UHI</b>	Urban Heat Islands
<b>UN</b>	United Nations
<b>UNCCD</b>	United Nations Convention to Combat Desertification
<b>UNDP</b>	United Nations Development Programme
<b>UNEP</b>	United Nations Environment Programme
<b>UNESCO</b>	United Nations Educational, Scientific and Cultural Organization
<b>UNFCCC</b>	United Nations Framework Convention on Climate Change
<b>US</b>	United States
<b>USA</b>	United States of America
<b>USD</b>	American dollar
<b>VBD</b>	Vector-Borne Disease
<b>VL</b>	Visceral Leishmaniasis
<b>VNR</b>	Voluntary National Review
<b>vs.</b>	versus
<b>VW</b>	Virtual Water
<b>W</b>	Watt
<b>WCRP</b>	World Climate Research Programme
<b>WDM</b>	Water Demand Management
<b>WeMO</b>	Western Mediterranean Oscillation
<b>WF</b>	Water Footprint
<b>WFD</b>	Water Framework Directive
<b>WFP</b>	Wood Forest Products
<b>WFPS</b>	Water Filled Pore Space
<b>WHO</b>	World Health Organization
<b>WMT</b>	Western Mediterranean Transition
<b>WNV</b>	West Nile virus
<b>WOCAT</b>	World Overview of Conservation approaches and Technologies
<b>WSUD</b>	Water-Sensitive Urban Design
<b>WUE</b>	Water-Use Efficiency
<b>WWTP</b>	Wastewater Treatment Plant
<b>yr</b>	year
<b>Zn</b>	Zinc
<b><math>\delta^{11}\text{B}</math></b>	Boron isotope
<b><math>\delta^{13}\text{C}</math></b>	Carbon isotope
<b><math>\mu</math></b>	micro
<b><math>\mu\text{M}</math></b>	micromolar, micromole per liter



# D APPENDIX

**ISO2  
country codes**

Iso Code	Country	Iso Code	Country
AL	Albania	LB	Lebanon
AT	Austria	LT	Lithuania
BA	Bosnia and Herzegovina	LU	Luxembourg
BE	Belgium	LV	Latvia
BG	Bulgaria	LY	Libya
CH	Switzerland	MA	Morocco
CY	Cyprus	MC	Monaco
CZ	Czech Republic	ME	Montenegro
DE	Germany	MK	The Republic of North Macedonia
DK	Denmark	MR	Mauritania
DZ	Algeria	MT	Malta
EE	Estonia	NL	The Netherlands
EG	Egypt	NO	Norway
ES	Spain	PL	Poland
FI	Finland	PS	State of Palestine
FR	France	PT	Portugal
GB	The United Kingdom of Great Britain and Northern Ireland	RO	Romania
GR	Greece	RS	Serbia
HR	Croatia	SE	Sweden
HU	Hungary	SI	Slovenia
IE	Ireland	SK	The Slovak Republic
IL	Israel	SY	Syrian Arab Republic
IT	Italy	TN	Tunisia
JO	Jordan	TR	Turkey

Table D.1 | ISO country codes



# APPENDIX

**Lists of Figures,  
Tables and Boxes**

## FIGURES

Figure 1.1	Structure and functioning of MedECC.
Figure 1.2	Distribution of MAR1 Coordinating Lead Authors and Lead Authors by country.
Figure 1.3	Geography, physiography and landscapes of the Mediterranean Basin.
Figure 1.4	The basis for the confidence level is given as a combination of evidence (limited, medium, robust) and agreement (low, medium and high).
Figure 2.1	Mediterranean coastline, topography over land and bathymetry over the sea plus the box definition.
Figure 2.2	Trends in sea level pressure (SLP).
Figure 2.3	Projected changes (%) in surface wind speed.
Figure 2.4	Past and future evolution of Surface Downwelling Shortwave Radiation (RSDS in W/m <sup>2</sup> ), Total Cloud cover (CLT in %) and Aerosol Optical Depth (AOD).
Figure 2.5	Observed temperature trends and time-series of temperature over land for the Mediterranean based on the Climatic Research Unit (CRU) gridded observations.
Figure 2.6	Projected changes in annual temperature between the recent past reference period three future sub-periods.
Figure 2.7	Time-series of simulated mean annual, summer and winter temperature averaged over the Mediterranean based on EURO-CORDEX 0.11° simulations.
Figure 2.8	Observed annual, DJF, JJA precipitation trends from the CRU dataset.
Figure 2.9	Time series of annual, DJF, and JJA precipitation over land from the CRU dataset.
Figure 2.10	Maps showing EURO-CORDEX-based change in annual, DJF and JJA precipitation change.
Figure 2.11	Time-series of simulated mean annual, wet and dry season over the Mediterranean land areas based on EURO-CORDEX 0.11° simulations.
Figure 2.12	Local amplification factor of the Mediterranean SST warming using 1980-1999 as reference period. Fraction of the Mediterranean Sea surface (in %) experiencing a given SST change value (in °C).
Figure 2.13	Warming stripes in the Intermediate Water (from east to west) and the Deep Water (in the Gulf of Lion).
Figure 2.14	Mediterranean sea level rise averaged in (2080-2099) with respect to present climate (1980-1999) under scenario RCP8.5.
Figure 2.15	PM10 concentration above the annual limit value of 40 µg m <sup>-3</sup> .
Figure 2.16	Averaged 2013–2017 map of plastic debris concentration (g km <sup>-2</sup> ) at the sea surface.
Figure 2.17	Fertilizer use and nitrogen release in Mediterranean Sea.
Figure 2.18	Modelled nitrogen deposition for the Mediterranean region.
Figure 2.19	Nitrogen dioxide (NO <sub>2</sub> ) concentrations above the annual limit value of 40 µg m <sup>-3</sup> .
Figure 2.20	Number of days (more than 25) above ozone (O <sub>3</sub> ) limit value of 120 µg m <sup>-3</sup> .
Figure 2.21	Mean concentrations of principles trace metals in coastal sediments of the Mediterranean Basin.
Figure 2.22	Potential sources and pathways for grounds and surface water pollution.
Figure 2.23	Common sources of endocrine disruptors in the environment.
Figure 2.24	Harmful algal blooms.
Figure 2.25	Number of new non-indigenous per 6 years in the Mediterranean since 1988.
Figure 2.26	Change in the number of new non-indigenous species per 6 years in the Mediterranean MSFD regions.



Figure 2.27	Ecosystem shift from algal forests to barrens due to the overgrazing activity of invasive herbivore rabbitfish.
Figure 2.28	Number and proportion of terrestrial non-indigenous plant, invertebrate and vertebrate species per country.
Figure 2.29	Number of non-indigenous woody plant pathogens per country.
Figure 2.30	Mediterranean population projections under the different Shared Socioeconomic Pathways (SSPs).
Figure 2.31	Mediterranean Gross Domestic Product (GDP) projections under the different Shared Socioeconomic Pathways (SSP).
Figure 2.32	Spatially explicit population projections produced for the Mediterranean Coastal SSPs.
Figure 2.33	Time series of the annual average temperature of the Mediterranean Basin.
Figure 3.1	Major river basins draining into the Mediterranean.
Figure 3.2	Water demand per sectoral use as percentage of total water demand.
Figure 3.3	Sources of water supply as percentage of total water supply.
Figure 3.4	Trends in municipal water withdrawal.
Figure 3.5	Potentially renewable water resources.
Figure 3.6	Regional patterns of changes in multi-model mean simulated annual runoff.
Figure 3.7	Same as Fig. 3.6 for the relative changes in multi-model mean simulated annual runoff at 4°C above pre-industrial.
Figure 3.8	Potential wastewater reuse per sector.
Figure 3.9	Example for a large-scale IBT project is the Southern Conveyor Project on the island of Cyprus.
Figure 3.10	Number of large dams in Mediterranean countries.
Figure 3.11	Net virtual water imports of countries.
Figure 3.12	Water-use efficiency per country.
Figure 3.13	Total agricultural land in the Mediterranean countries.
Figure 3.14	Total irrigated land in the Mediterranean countries.
Figure 3.15	Meat and fish consumption (kg capita <sup>-1</sup> yr <sup>-1</sup> ) in Mediterranean countries.
Figure 3.16	Cereal trade patterns.
Figure 3.17	Trade patterns in fodder and feeding products.
Figure 3.18	Net protein fluxes.
Figure 3.19	Total fish landings (tonnes) in the Mediterranean Sea.
Figure 3.20	Total landings (t).
Figure 3.21	Crop yield changes in the MENA region.
Figure 3.22	Evolution of Primary Energy Consumption.
Figure 3.23	Evolution of per capita Energy Intensity Index.
Figure 3.24	Evolution of primary energy production.
Figure 3.25	Evolution of primary energy consumption.
Figure 3.26	Energy weight in trade balance.
Figure 3.27	Energy consumption.
Figure 3.28	Impacts of streamflow drought and high water temperature on utilisation rates of hydropower.
Figure 3.29	Distribution of wave energy flux.
Figure 3.30	Potential locations for offshore wind farms.
Figure 3.31	Firewood production.
Figure 3.32	Forest area and production forests available for industrial use including biomass for energy purposes.
Figure 3.33	Electricity production from biogas and solid biomass.
Figure 3.34	Projected constraints to the establishment of bioenergy plantations by 2050.
Figure 3.35	Mediterranean patterns with number of days in the given year with streamflow drought, high water temperature, and that both events coincide.

Figure 3.36	Voluntary commitments of SEMC.
Figure 3.37	Primary energy demand by region.
Figure 3.38	Primary energy resources in the Mediterranean energy mix.
Figure 3.39	Electricity generation mix by fuel type.
Figure 3.40	Demographic and energy projections in the Mediterranean in 2040.
Figure 3.41	Benefits of the implementation of the Proactive Scenario (PS) compared to the Reference Scenario (RS) at the horizon 2040.
Figure 3.42	Future changes in national wind power, solar PV power, hydropower and thermoelectric power production under +1.5°C global warming, 2°C and 3°C.
Figure 3.43	Impacts of climate change on annual mean streamflow and water temperature.
Figure 3.44	Impacts of climate and water resources change on annual mean usable capacity of current hydropower and thermoelectric power plants.
Figure 3.45	Regional share of biomass in the primary energy use in the IPCC 1.5°C scenarios.
Figure 3.46	Global fossil fuels, biomass, and agricultural products financial flows in 2010 and in 2100.
Figure 3.47	Percentage change in average daily peak electric load from 2006–2012 to 2080–2099 for projected daily maximum temperatures under RCP4.5 and RCP8.5 climate change scenarios.
Figure 3.48	Impacts of adaptation options on power-generation vulnerability to water constraints under climate change.
Figure 3.49	Economic and financial contexts around the Mediterranean.
Figure 3.50	Change in infrastructure spending for a 2°C scenario in the Mediterranean region, percentage change in expenditure over 2015–2030 compared to Business-as-usual.
Figure 3.51	International public climate finance by sector in SEMCs.
Figure 3.52	International public climate financial instruments by SEMCs recipient countries.
Figure 3.53	Energy subsidies in SEMCs.
Figure 3.54	Cross-border physical electricity flows.
Figure 3.55	Mediterranean interconnection projects.
Figure 3.56	Strategy of Mediterranean regulators for regional integration.
Figure 3.57	Illustration of the optimal, or Pareto, frontiers for two objectives – maximizing the mean and minimizing the standard deviation of the total wind-PV penetration. PV and wind optimal capacity distribution.
Figure 4.1	Summary of interactions between large marine vertebrates and marine litter.
Figure 4.2	The trajectory and spatial pattern of chondrichthyan (cartilaginous fishes that include sharks, skates, rays and chimaeras) fisheries catch landings and fin exports.
Figure 4.3	Temperature trends across the Mediterranean Basin.
Figure 4.4	Different drivers potentially affecting marine pelagos and benthos in the Mediterranean Sea.
Figure 4.5	Coastal Risk Index (CRI-MED) map of the Mediterranean.
Figure 4.6	Cross-section of a sandy dune system.
Figure 4.7	Typical salt marsh zonation.
Figure 4.8	Expected geographical species shifts for the 288 coastal Mediterranean fish species.
Figure 4.9	Nile Delta, Egypt.
Figure 4.10	Mediterranean land systems.
Figure 4.11	Map of Global Dryland Assessment (GDA) plots.
Figure 4.12	Change in the percentage of land-use change from 1961 to present in relation to desertification and land degradation.

<b>Figure 4.13</b> .....	Reduction in freshwater discharge flows into the Mediterranean for all rivers.
<b>Figure 4.14</b> .....	Tree- and ecosystem-level responses in Mediterranean forests.
<b>Figure 4.15</b> .....	Distribution of drylands and their subtypes based on observations. Distribution of projected drylands transitions according to RCP8.5 for three Global Warming Levels.
<b>Figure 5.1</b> .....	Distribution of the tiger mosquito, <i>Aedes albopictus</i> .
<b>Figure 5.2</b> .....	Attributable fraction of heat-related deaths during summer by country in European sub-region.
<b>Figure 6.1</b> .....	Mediterranean sites where the impacts of innovative agricultural practices have been surveyed.
<b>Figure B1</b> .....	Projected changes in winter temperatures between the recent past reference period and three future sub-periods.
<b>Figure B2</b> .....	Projected changes in spring precipitation between the recent past reference period and three future sub-periods.
<b>Figure B3</b> .....	Projected changes in summer precipitation between the recent past reference period and three future sub-periods.
<b>Figure B4</b> .....	Projected changes in autumn precipitation between the recent past reference period and three future sub-periods.
<b>Figure B5</b> .....	Projected changes in winter precipitation between the recent past reference period and three future sub-periods.
<b>Figure B6</b> .....	Projected changes in spring precipitation between the recent past reference period and three future sub-periods.
<b>Figure B7</b> .....	Projected changes in summer precipitation between the recent past reference period and three future sub-periods.
<b>Figure B8</b> .....	Projected changes in autumn precipitation between the recent past reference period and three future sub-periods.

## TABLES

<b>Table 1.1</b> .....	Likelihood terms associated with outcomes used in MAR1.
<b>Table 2.1</b> .....	Climate change as a function of time period and Representative Concentration Pathway for the land sub-regions in Fig. 2.1 and the whole Mediterranean Sea area.
<b>Table 2.2</b> .....	Metal concentrations ( $\mu\text{g g}^{-1}$ dry weight) in marine sediment of Taranto Gulf (Ionian Sea, southern Italy).
<b>Table 2.3</b> .....	Main interactions among drivers.
<b>Table 2.4</b> .....	Overview of selected socioeconomic scenarios that cover Mediterranean countries.
<b>Table 3.1</b> .....	Available and exploitable water resources in the Mediterranean region per country.
<b>Table 3.2</b> .....	Percentage of electricity generated from hydropower in selected Mediterranean countries.
<b>Table 3.3</b> .....	Municipal water withdrawal in absolute values, in percentage of total withdrawal and per capita.
<b>Table 3.4</b> .....	Major environmental problems for water quality along the coastal zone of Mediterranean countries.
<b>Table 3.5</b> .....	Projected changes (%) of soil moisture for the different Mediterranean regions.
<b>Table 3.6</b> .....	Simulated changes in spatially averaged multi-model mean annual runoff.
<b>Table 3.7</b> .....	Characteristics and relative changes in runoff and discharge under high-end climate change.

<b>Table 3.8</b> .....	Water loss definitions and classifications.
<b>Table S3.1</b> .....	Sources of water supply.
<b>Table 3.9</b> .....	Production of cereals, fruit, vegetables, meat and milk in the Mediterranean countries.
<b>Table 3.10</b> .....	Reported installed wind and solar (photovoltaic, PV) power in Mediterranean countries in 2019.
<b>Table 3.11</b> .....	Levels of primary solid biofuels.
<b>Table 5.1</b> .....	Inequality indicators for Mediterranean countries.
<b>Table 5.2</b> .....	Gender indicators for Mediterranean countries.
<b>Table 5.3</b> .....	Classification of finance tools to protect the environment and promote sustainable development.
<b>Table 5.4</b> .....	Climate change impacts on selected human rights.
<b>Table 6.1</b> .....	Overview of stages of non-indigenous species introduction and potential management strategies.
<b>Table 6.2</b> .....	Selected adaptation strategies.
<b>Table B1</b> .....	List of EURO-CORDEX experiments taken into account in the presented assessment and availability of variables.
<b>Table D1</b> .....	ISO2 country codes.

## BOXES

<b>Box 2.1</b> .....	How much has the Mediterranean Basin warmed since the pre-industrial period ?
<b>Box 2.2</b> .....	Representative Concentration Pathways (RCPs).
<b>Box 3.1.1</b> .....	Impacts of structural aging and climate change on water infrastructure.
<b>Box 3.1.2</b> .....	Water use and the specific Mediterranean diet.
<b>Box 3.3.1</b> .....	Climate variability and energy planning.
<b>Box 3.3.2</b> .....	Energy issues for Mediterranean islands.
<b>Box 4.1</b> .....	Bio-indicators for the assessment of changes in Mediterranean marine ecosystems.
<b>Box 4.2</b> .....	Urban biodiversity in the Mediterranean Region.
<b>Box 4.3</b> .....	Nitrogen deposition and ecosystems.
<b>Box 4.4</b> .....	Mediterranean islands.
<b>Box 5.1.1</b> .....	Development indicators and terms.
<b>Box 5.1.2</b> .....	Gender-related development indicators and term.
<b>Box 5.1.3</b> .....	Vulnerability.
<b>Box 5.3.1</b> .....	Climate change and the Syrian conflict.









[www.medecc.org/first-mediterranean-assessment-report-mar1/](http://www.medecc.org/first-mediterranean-assessment-report-mar1/)  
[enquiries: marini@medecc.org](mailto:enquiries:marini@medecc.org)

**MedECC**  
Mediterranean Experts on Climate  
and environmental Change



Union for the Mediterranean  
Union pour la Méditerranée  
الاتحاد من أجل المتوسط

**UN**  
environment  
programme



Mediterranean  
Action Plan  
Barcelona  
Convention



ISBN: 978-2-9577416-0-1