

Table 1. Location, site abbreviation (Abbr), sample year, sample size (*n*), and life stage (A = adult, J = juvenile) of fish sampled from 32 putative coho salmon populations in seven regions of Alaska.

Region and population ^{a,b}	Abbr.	Year	<i>n</i>	Life Stage
<i>Northwest Alaska</i>				
Kisaralik R	NW1	1997	95	A
Nome R	NW2	1998	89	A
Clearwater R	NW3	1997	56	A
Nenana R	NW4	1997	56	A
<i>Southwest Alaska</i>				
Gertrude Ck	SW1	1997	96	A
Alagnak R	SW2	1997	89	A
Kulukak R	SW3	1997	91	A
<i>Aleutian Is./AK Pen.</i>				
Finger Ck	AP1	1997	180	J
NavFac Ck	AP2	1997	150	J
Joshua Green Ck	AP3	1995	50	A
<i>Kodiak Archipelago</i>				
Akalura R	KO1	1997	43	A
Ayakulik R	KO2	1997	82	A
Bear Ck	KO3	1997	96	A
Buskin R	KO4	1998	69	A
Midway R	KO5	1997	94	A
Pauls L	KO6	1997	50	A
Spiridon R	KO7	1998	73	A
Zachery R	KO8	1997	77	A
<i>Cook Inlet</i>				
Chuitna R	CI1	1992	52	A
Crescent R	CI2	1998	95	A
Killey R	CI3	2000	79	A
Kenai R KL	CI4	1999	75	A
Kenai R SL	CI5	1999	55	A
Quartz Ck	CI6	1998	69	A
Snow R	CI7	1998	64	A
Tern L	CI8	2000	81	A
<i>Prince William Sound</i>				
Gun Boat Ck	PW1	1997	62	J
Otter Ck	PW2	1997	64	J
Red Ck	PW3	1997	85	J
<i>Southeast Alaska</i>				
Alsek R	SE1	1995	49	A
Margaret Ck	SE2	1997	69	A
Pavlof R	SE3	1998	91	J

^a NW3 and NW4 are Yukon River tributaries. CI3-CI8 are Kenai River tributaries.

^b CI4 and CI5 spawn in late winter. All other populations spawn in late summer and fall.