




# New distribution record of *Hoya sipitangensis* Kloppenb. & Wiberg (Apocynaceae, Asclepiadoideae) from Palawan, Philippines

Marjorie D. delos Angeles<sup>1\*</sup>, Cristian C. Lucañas<sup>2</sup>, Annalee S. Hadsall<sup>3</sup>

<sup>1</sup> Plant Biology Division, Institute of Biological Sciences, College of Arts and Sciences, University of the Philippines Los Baños, Los Baños, Philippines • [mddelosangeles1@up.edu.ph](mailto:mddelosangeles1@up.edu.ph)  <https://orcid.org/0000-0003-3729-2230>

<sup>2</sup> Entomology Collections, Museum of Natural History, University of the Philippines Los Baños, Los Baños, Philippines • [cclucanas@up.edu.ph](mailto:cclucanas@up.edu.ph)  <https://orcid.org/0000-0002-2143-4057>

<sup>3</sup> Botanical Herbarium, Museum of Natural History, University of the Philippines Los Baños, Los Baños, Philippines • [ashadsall@up.edu.ph](mailto:ashadsall@up.edu.ph)  <https://orcid.org/0000-0002-8884-3904>

\* Corresponding author

## Abstract

We report *Hoya sipitangensis* Kloppenb. & Wiberg for the first time on Palawan Island, Philippines. This record extends its distribution northward from Borneo to the Philippines. Additional taxonomic information and photographs from field collections are provided.

## Keywords

Forests over limestone, Puerto Princesa Subterranean River National Park, range extension, wax plant

**Academic editor:** Nik Fadzly N. Rosely | Received 27 March 2021 | Accepted 6 June 2021 | Published 14 June 2021

**Citation:** delos Angeles MD, Lucañas CC, Hadsall AS (2021) New distribution record of *Hoya sipitangensis* Kloppenb. & Wiberg (Apocynaceae, Asclepiadoideae) from Palawan, Philippines. Check List 17 (3): 917–922. <https://doi.org/10.15560/17.3.917>

## Introduction


The genus *Hoya* R.Br. (Brown 1810: 459) contains 500 or more described species of mostly epiphytic flowering plants. Distinct characters of this genus include milky sap, umbelliform inflorescence, and a star-shaped corona (Kidyoo and Wai 2009). It is distributed throughout Southeast Asia, China, New Guinea, Australia, and some Pacific Islands (Hadsall et al. 2015), with the highest diversity in the Philippines, Borneo, and New Guinea (Pelser et al. 2011; Lobb and Rodda 2016; Juhonewe and Rodda 2017). In the Philippines, 202 species and 39 subspecies of *Hoya* have been reported, of which 197 are reported to be endemics (Pelser et al. 2011).

Palawan Island, Philippines, is a distinct biogeographic zone due to its historical connections with the

Greater Sunda Shelf (Van-Wright 1990). Santiago and Buot (2017) noted 17 species of *Hoya* from Palawan Island, four of which are island endemics and two are shared with the Sunda shelf. The low endemism record of Palawan compared to the overall high rate of Philippine *Hoya* endemism may be attributed to a dearth in data as a result of scarce botanical expeditions focusing on this genus on Palawan. in the province.

## Methods

During a recent expedition to Puerto Princesa, Palawan, a small *Hoya* population was observed hanging among branches of trees in a lowland forest. The specimens



**Figure 1.** *Hoya sipitangensis* Kloppenberg & Wiberg. **A.** Growth habit. **B.** Leaf (adaxial view). **C.** Leaf (abaxial view). **D.** Inflorescence. **E.** Fruit. (Photo credits: RAA Bustamante (A); CC Lucanas (B, C) MD delos Angeles (D, E).

were collected using the Gratuitous Permit (2021-02) issued by the Palawan Council for Sustainable Development (PCSD) and Protected Area Management Board (PAMB) of Puerto Princesa. Several plant samples were preserved in 95% ethyl alcohol, then dried and mounted

on herbarium sheets. Measurements were made using digital calipers, and ImageJ. Multiple images of dissected parts were photographed using a Nikon D3100 camera and were focus stacked using CombineZM. Distribution maps were generated using Natural Earth Data in QGIS

v. 2.18.0 (QGIS Development Team 2009).

The specimens were compared with original descriptions of *H. sipitangensis* Kloppenb. & Wiberg and *H. yapiana* Kloppenb from UC herbaria (the acronym follow Thiers 2021). The voucher specimen was deposited at the Botanical Herbarium (CAHUP), University of the Philippines Los Baños Museum of Natural History. The description below of this species is based on the collected Philippine materials.

## Results

### *Hoya sipitangensis* Kloppenb. & Wiberg

Fraterna 15(3): 4. 2002 (Kloppenb and Wiberg 2002). Sipitang, Sarawak, Borneo (neotype UC [UC1784973]).

**New record.** PHILIPPINES – Palawan Island • Puerto Princesa, Brgy. Cabayugan, foot of St. Paul’s; 10°09’ 21”N, 118°53’04”E; 10 m a.s.l.; 24 Feb. 2021; MDDelos Angeles & RABustamante leg.; Epiphytic flora found on a tree near an agricultural field; CAHUP 074067 (Figs. 1, 2).

**Additional specimens examined.** BRUNEI • originally from Belait, Seria, Tutong District and cultivated in Singapore; 21 Jan. 2008. (holotype SING 0124341; isotype BRUN n.v., K [K000898014]).

**Identification.** A small climbing epiphyte. Stem slender, up to 2.39 mm in diameter, yellow-green, glabrous. Leaves: petiole round and glabrous, up to 6.62 mm long; blade ovate to elliptic, 69.5 long, 31.08 mm wide and 0.85 mm thick, apex acute, base round to cuneate; midrib and venations obscure. Inflorescence: axillary, positively geotropic, 20–28-flowered, flowers with sweet scent; peduncle 29.77 mm long, glabrous, rachis 23 mm long. Pedicels vary in length, 7.16–11.02 mm long, glabrous. Corolla round, 3.56 mm in diameter, revolute, creamy white on the outer surface, pink towards the inner surface, minutely pubescent. Corona 1.6 mm in diameter; inner lobes pink to carmine; center yellow; outer lobes long and acuminate, translucent. Pollinarium: pollinia 0.25 mm long. Fruit: follicle 51.23 mm long, borne on a stalk 6.73 mm long.

## Discussion


On account of its revolute corolla lobes, boat-shaped corona scales with two tooth-like projections below, and the presence of an unsulcated pentamerous skirt (annulus) beneath the corona scales, *Hoya sipitangensis* is placed in the section *Otostemma* Blume (Kloppenb 1990). It can be differentiated from *H. obscura* Elmer ex C.M.Burton and *H. lacunosa* Blume on the basis of its larger leaves and flowers where the corolla is pubescent and not velvety-villous (Kloppenb and Wiberg 2002). The inflorescence of *H. sipitangensis* has a sweet scent reminiscent of *H. lacunosa*. According to Kloppenb and Siar (2010), *H. sipitangensis* is also similar

**Table 1.** Comparisons of *Hoya sipitangensis* Kloppenb. & Wiberg from three localities.


Characters (in mm)	<i>H. sipitangensis</i> (Puerto Princesa, Philippines)	<i>H. sipitangensis</i> (Sipitang, Borneo)	<i>H. sipitangensis</i> (= <i>H. yapianum</i> ; Brunei)
Pedicel			
Length	4.0	8.0	15
Diameter	1	0.9	10
Sepal			
Length	0.736	0.8	0.8
Diameter	0.708	2.6	0.8
Ovaries			
Length	0.7	—	—
Corolla			
Diameter flattened	7.065	12.4	5.1
Center to apex	3.462	6.2	—
Apex-sinus	1.735	4	—
Sinus-center	2.11	2.5	—
Sinus-sinus	2.485	3.3	—
Corona			
Apex to apex	1.783	2.3	—
Apex-center	1.12	2.6	—
Widest	0.7	1.1	0.7
Pollinia			
Length	0.289	0.36	0.25
Width	0.137	0.09	0.12
Retinaculum			
Length	0.073	0.06	0.06
Head	0.006	—	—
Shoulders	0.031	0.04	—
Waist	0.012	0.02	—
Hips	0.034	0.05	—
Extensions	0.036	0.04	0.04
Translator			
Length	0.145	0.13	0.10
Width	0.040	0.01	0.04
Caudical			
Length	0.045	0.04	—

in appearance to *H. yapianum* Kloppenb. & Siar, but the latter differs in the apex of its sepals which: “are obtuse and not acute; the pedicels are glabrous, not puberulent; the inner apex of the corona is acute, not spatulate; the foliage is larger and the pollinia are about 1/3 shorter” Kloppenb and Siar, (2010: 328). However, Rodda (2016) considered *H. yapianum* a synonym of *H. sipitangensis* on the basis that the differences are part of the natural variation of the latter. The dimension of individual plants of *H. sipitangensis* that we found on Palawan are notably smaller compared to the type description by Kloppenb and Wiberg (2002) (Table 1).

*Hoya sipitangensis*, and its synonym *H. yapianum*, have only been previously reported from their type localities, Sipitang, Sabah, Borneo and Seria, Belait, Brunei, respectively. It has been found on coastal to lowland forests at 300 m a.s.l. Here, we report it as a component of the epiphytic flora in a lowland forest in Puerto Princesa, Palawan (Fig. 3). Several clusters of *H. sipitangensis* were spotted hanging among branches of three Lythraceae tree species. The host tree species forms dense canopies creating shady and moist understories. This



**Figure 2.** *Hoya sipitangensis* Kloppenberg & Wiberg. **A, B.** Floret: **(A)** dorsal view; **(B)** ventral view. **C, D.** Pedicel and calyx: **(C)** lateral view ovary (with arrow), **(D)** ventral view. **E–H.** Corona: **(E)** dorsal; **(F)** lateral; **(G)** ventral; **(H)** dorsal. **I.** Pollinarium.



**Figure 3.** Distribution of *Hoya sipitangensis* Kloppenburg & Wiberg. Blue circle = known reports (data after Lobb and Rodda 2016); red circle = new record in the Philippines.

supports the previous observations on the influence of Bornean biogeography on Palawan Island (Heaney 1985; Wikramanayake et al. 2002; Atkins and Cronk 2001).

## Acknowledgements

We extend our sincere gratitude to the following: Park Superintendent Elizabeth A. Maclang for the support she has given us throughout the expedition; forest rangers: Augusto A. Asis, Jr., Ernesto B. Lapuz, Jr., Arnold Magallanes, and Francisco Dela Cruz for their unwavering help during our fieldwork; Dr. Aimee-Lynn B. Dupo for the use of laboratory space and equipment; and Dr. Michele Rodda for his valuable comments and insights. The field collection is made possible under GP 2021-02 granted by the Palawan Council for Sustainable Development (PCSD) with the Transport permit no: PPC-LTP-AO12-2021-316 and Certificate of Inspection no. 1614239583882.

## Author Contributions

Conceptualization: MDDA. Formal Analysis: MDDA and CCL. Project Administration: MDDA. Writing – original draft: MDDA. Data curation: MDDA and CCL. Methodology: MDDA. Software: MDDA and CCL. Validation: MDDA and CCL. Writing-review and editing: MDDA, CCL, and ASH.

## References

- Atkins H, Cronk QCB (2001) The genus *Cyrtandra* (Gesneriaceae) in Palawan, Philippines. *Edinburgh Journal of Botany* 58 (3): 443–458. <https://doi.org/10.1017/S0960428601000762>
- Aurigue FB (2013) Collection of Philippine Hoyas and their culture. DOST-PCAARRD, Laguna, Philippines, 195 pp.
- Brown, RB (1810) *Hoya*. In: *Prodromus Florae Novae Hollandiae et Insulae Van-Diemen: exhibens characteres plantarum quas annis 1802–1805 .... Vol. I.* Richardi Taylor et socii, London, 459–460. <https://doi.org/10.5962/bhl.title.3678>
- Co's digital flora of the Philippines (2011) <http://www.philippineplants.org>. Accessed on 2021-3-21.
- Hadsall AS, Cajano MAO, Lambio IAF (2015) A new distribution record of *Hoya espadoniana* Kloppenburg, Siar & Cajano (Section *Acanthostemma*: Apocynaceae). *Asia Life Sciences* 24 (2): 469–476.
- Heaney LR (1985) Zoogeographic evidence for middle and late Pleistocene land bridges to the Philippine Islands. *Modern Quaternary Research in Southeast Asia* 9: 127–144.
- Kidyoo M, Wai JS (2009) A new record of *Hoya* (Asclepiadaceae) from southern Thailand. *The Natural History Journal of Chulalongkorn University* 9 (2): 235–238.
- Kloppenburger D (1990) *Hoya* sections: a complete study with modifications and additions. Fresno, California, USA, 51 pp.
- Kloppenburger D, Siar SV (2010) New species of *Hoya* (Apocynaceae) from Brunei and the Philippines. *Gardens' Bulletin Singapore* 61 (2): 327–333.
- Kloppenburger D, Wiberg EK (2002) New *Hoya* species from Sipitang, Sarawak, Borneo. *Fraterna* 15 (3): 4.
- Lamb A, Rodda M, Gokusing L, Bosuang S, Rahayu S (2016) A guide to hoyas of Borneo. Natural History Publications, Kota Kinabalu, Malaysia, 204 pp.

- QGIS Development Team, (2009) QGIS Geographic Information System. Open Source Geospatial Foundation. <http://qgis.org>. Accessed on 2021-3-18.
- Rodda M (2017) Index of names and types of *Hoya* (Apocynaceae: Asclepiadoideae) of Borneo. *Garden's Bulletin Singapore* 69 (1): 33–65.
- Santiago JO, Buot IE Jr. (2017) Checklist of *Hoya* species on Palawan Island, Philippines. *Journal of Nature Studies* 16 (1): 34–44.
- Siar SV, Philippine Council for Agriculture, Forestry and Natural Resources Research and Development—Department of Science and Technology (2005) Philippine hoyas. Philippines Council for Agriculture and Aquatic Resources Research and Development, Laguna, Philippines, 23 pp.
- Thiers B (2021) Index herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium, New York, USA. <http://sweetgum.nybg.org/science/ih/>. Accessed on: 2021-03-21.
- Vane-Wright D (1990) The Philippines—key to the biogeography of Wallacea. In: Knight WJ, Holloway JD (Eds.) *Insects and the rain forests of South East Asia (Wallacea)*. Royal Entomological Society of London, London, UK, 19–34.
- Wikramanayake ED, Dinerstein E, Loucks CJ, Olson DM, Morrison J, Lamoreux J, McKnight M, Hedao P (2002) *Terrestrial ecoregions of the Indo-Pacific: a conservation assessment*. Island Press, Washington, DC, USA, 643 pp.