



Five new records of nematodes from East Antarctica

P. Bohra¹, A.K. Sanyal², A. Hussain³ & B. Mitra⁴

¹Zoological Survey of India, Desert Regional Centre, Jhalamand, Pali Road, Jodhpur, Rajasthan 342005, India.

^{2&4} Prani Vigyan Bhawan, Zoological Survey of India, M-Block, 535 New Alipore, Kolkata, West Bengal 700053, India.

³Zoological Survey of India, Northern Regional Centre, 218, Kaulagarh Road, Dehra Dun, Uttarakhand 248195, India.

Email: ¹bohrapadma@gmail.com

The terrestrial invertebrate fauna of Antarctica comprise of Protozoa, Rotifera, Tardigrada, Nematoda and Arthropoda. A handful of soil contains thousands of the microscopic nematode worms, many of them parasites of insects, plants or animals. Free-living species are abundant, including nematodes that feed on bacteria, fungi, and other nematodes.

The Antarctic region has been subdivided into three main ecological zones viz., the Continental Antarctic, the Maritime Antarctic and the Sub-Antarctic with distinctive climatic and biotic characteristics (Holdgate 1970). The free-living soil, freshwater and moss inhabiting nematodes of these regions have been studied by de Man (1904), Krijanova (1958), Timm (1971), Gray (1975), Loof (1975), Heyns (1993), Kito et al. (1996), and others. Andrassy (1998) has reported nematodes as the most abundant group in the Antarctic region. He listed 43 species comprising 29 species from the maritime region and 14 from Antarctica proper. Besides, he has also discussed survival strategies and geographic distribution

of nematodes. He has also provided a key to the species of Plectidae and Qudsianematidae. Bostrom (2005) added three species of nematodes from Maudland, East Antarctica and Ghosh et al. (2005) reported one new species *Antarctenchus motillius* from East Antarctica. Andrassy (2008) described six new species of the genus *Eudorylaimus* from continental Antarctica.

The moss and soil samples were collected from different sites at the periphery of Priyadarshini Lake (PDL) located at Schirmacher Oasis, East Antarctica. The nematodes were killed and fixed in hot 4% formalin and mounted in anhydrous glycerin. The specimens are deposited in the National Zoological Collection of Desert Regional Centre, Zoological Survey of India, Jodhpur.

Results

The analysis of samples yielded five species belonging to three genera and three families of the Orders Tylenchida, Dorylaimida and Araeolaimida. *Helicotylenchus dihystra*, *H. diagonicus*, *H. exallus*, *Eudorylaimus sabulophilus* and *Plectus telekii* are being reported for the first time from East Antarctica.

Systematic account

Order: Tylenchida Thorne, 1949

Superfamily: Hoplolaimoidea Filipjev, 1934 (Paramonov, 1967)

Family: Hoplolaimidae, Filipjev, 1934 (Wieser, 1953)

Genus: *Helicotylenchus* Steiner, 1945

***Helicotylenchus diagonicus* Perry in Perry, Darling & Thorne, 1959 (Fig. 1)**

Material examined: 3 females, 14.i.1999, coll. A.K. Sanyal, Reg.No. IV/1923.

Host: Moss at the bank of PDL.

Measurements: Female (3): L = 0.49-0.90 mm; a = 24-26; b = 5.5-6.0; c = 50-55; V = $10^{-11}56-62^{9-10}$

Description:

Female: Body spiral shaped. Lip region truncated with 3-4 annules. Stylet 20-22 μ m long, basal knobs flattened slightly indented anteriorly. Spermatheca offset without sperms. Phasmids at the level of anus. Tail more curved dorsally with slight ventral projection.

Abbreviations: L = body length (mm/ μ m); a = body length / maximum body width; b = body length / oesophageal length; C = body length / tail length; c' = tail length / body width at anus; V = distance from head end to vulva x 100 / body length

Date of publication (online): 26 June 2010
Date of publication (print): 26 June 2010
ISSN 0974-7907 (online) | 0974-7893 (print)

Editor: Anwar L. Bilgrami

Manuscript details:

Ms # o2342

Received 09 November 2009

Final received 11 May 2010

Finally accepted 15 May 2010

Citation: Bohra, P., A.K. Sanyal, A. Hussain & B. Mitra (2010). Five new records of nematodes from East Antarctica. *Journal of Threatened Taxa* 2(6): 974-977.

Copyright: © P. Bohra, A.K. Sanyal, A. Hussain & B. Mitra 2010. Creative Commons Attribution 3.0 Unported License. JoTT allows unrestricted use of this article in any medium for non-profit purposes, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Acknowledgement: The authors are highly thankful to Dr. Ramakrishna, Director, Zoological Survey of India, Kolkata for providing necessary facilities to carry out the research work.

OPEN ACCESS | FREE DOWNLOAD






Figure 1. *Helicotylenchus diagonicus*
A - Female Anterior region; B - Posterior region.

Male: Not found.

Distribution: Canada, Greece, India, Poland, USA.

Remarks: First report from Antarctica.

***Helicotylenchus exallus* Sher, 1966**
(Fig. 2)

Material examined: 5 females, 30.xii.1999, coll. A.K. Sanyal, Reg.No. IV/1924.

Host: Moss at the bank of PDL.

Measurements: Females (5): L = 0.61-0.65 mm; a = 26-28; b = 6.0-6.2; c = 40-45; c' = 0.7-1.2; V = ¹⁰⁻¹¹60-62⁸⁻¹⁰

Description

Female: Body spiral shaped. Lip region hemispherical marked by four annules. Stylet 24-26 µm long, basal knobs with flattened anterior surface. Spermatheca large, offset with sperms. Phasmid 2-3 annules anterior to anus. Tail more curved dorsally with slight ventral projection.

Male: Not found.




Figure 2. *Helicotylenchus exallus*
A - Female Anterior region; B - Posterior region.

Distribution: India, USA.

Remarks: First report from Antarctica.

***Helicotylenchus dihystra* (Cobb, 1893) Sher, 1961**
(Fig. 3)

Material examined: 5 females, 03.i.1999, coll. B. Mitra, Reg.No. IV/1921.

Host: Wet Moss along with soil at the bank of Priyadarshini Lake (PDL).

Measurements: Females (5): L = 0.54-0.83 mm; a = 20-32; b = 4.5-6.4; c = 44-46; c' = 1.0-1.3; V = ¹²⁻¹⁵60-66¹¹⁻¹³

Description

Female: Body spiral. Cuticle with distinct transverse striae. Lip region continuous with body, hemispherical bearing four annules. Cephalic framework conspicuous. Lateral fields marked by four incisures. Stylet 24-26 µm long, basal knobs concave. Oesophagus with gland overlapping ventrally. Spermatheca rounded offset, without sperms. Tail dorsally convex-conoid usually with slight ventral projection.

Male: Not found.

Distribution: This is a cosmopolitan and widely distributed species.




Figure 3. *Helicotylenchus dihystra*
A - Female Anterior region; B - Posterior region.




Figure 4. *Eudorylaimus sabulophilus*
A - Female Anterior region; B - Female Posterior region.

Remarks: First time reported from Antarctica.

Order: Dorylaimida Pearse, 1942
Superfamily: Dorylaimoidea De Man, 1876
Family: Qudsianematidae Jairajpuri, 1965

***Eudorylaimus sabulophilus* Tijepkema, Ferris & Ferris, 1971**
(Fig. 4)

Material examined: 3 females, 23.ii.1999, coll. A. Hussain, Reg.No. IV/2240

Host: Moss and algae at the bank of PDL.

Measurement: Females (3): L = 0.87-0.98mm; a = 17-22; b = 3.47-3.92; c = 17.3-21.7; V = 56-59.

Description:

Female: Body slightly curved ventrally upon fixation. Lips well separated and set off from body by constriction. Odontostyle 17-18 μm long; its aperture 30% of odontostyle length. Odontophore rod-like 22-24 μm long. Oesophageal expansion gradual. Basal expanded part of oesophagus occupies 45-47% of oesophageal length. Reproductive system amphidelphic. Tail dorsally-convex with subacute or narrow rounded tip. Tail tip slightly bent ventrally.

Male: Not found.

Distribution: USA.

Remarks: First time reported from Antarctica.




Figure 5. *Plectus telekii*
A - Female Anterior region; B - Female Posterior region.

Order: Araeolaimida De Coninck & Schuurmans Stekhoven, 1933
Superfamily: Plectoidea Örley, 1880
Family: Plectidae Örley, 1880

***Plectus telekii* Mulk & Coomans, 1978**
(Fig. 5)

Material examined: 6 females, 23.ii.1999, coll. A. Hussain, Reg.No. IV/1925-26.

Host: Moss at the bank of PDL.

Female (5): L = 669 μm -794 μm a = 18.0-19.0, b = 4.0-4.5, c = 8.0-8.8, c' = 3.7-5.0, V = 45.2-49.7, Pharynx = 165-186 μm , Anal body diameter = 18-22 μm , Tail length = 79-97 μm

Description:

Female: Body medium-sized, arcuate, open C shape upon fixation. Lip region continuous with body contour. Lips elevated 2.5-3.0 μm long. Lip region 7-11 μm wide. Cephalic sensilla setose, 2.0-2.5 μm long. Amphids located 11-14 μm from anterior end. Stoma 19-30 μm long, cheilostom cuticularised. Nerve ring encircles isthmus. Excretory pore inconspicuous. Intestine granular. Female reproductive system amphidelphic. Ovary reflexed. Vulva slit like. Uterus without egg. Vulva-anus distance three times tail length. Tail cylindrical, arcuate. Spinneret about 2 μm long.

Male: Not found.

Distribution: Rajasthan, India

Remarks: First report from Antarctica.

REFERENCES

- Andrassy, I. (1998).** Nematodes in Sixth continent. *Journal of Nematode Morphology and Systematics* 1: 107-186.
- Andrassy, I. (2008).** *Eudorylaimus* species (Nematoda: Dorylaimida) of continental Antarctica. *Journal of Nematode Morphology & Systematic* 11(1): 49-66.
- Bostrom, S. (2005).** Nematodes from Sirmacher Oasis, Dronning, Maud Land, East Antarctica. *Russian Journal of Nematology* 13: 43-54.
- Christy, J.R. & V.G. Perry (1951).** Removing nematodes from soil. *Proceedings Helminthological Society, Washington* 18: 106-108.
- Cobb, N.A. (1918).** Estimating the Nema Population of the Soil. Agricultural Technology Circular I. Bureau of Plant Industry, United States, Department of Agriculture, 48pp.
- De Man, J.G. (1904).** Nematodes libres. Result. Voyage S.Y. *Belegica, Zoologie* 55pp.
- Ghosh, S.C, A.C. Bulgarin & J. De (2005).** *Antarctenchus motilius* sp. n. (Nematoda: Tylenchida) from Schirmacher Oasis, East Antarctica. *Journal of Interacademia* 9(3): 367-371.
- Gray, N.F. (1975).** A preliminary survey of Antarctic nematophagous fungi with reference to temperature tolerance. BSc. Degree Project Report No. 745/1138. The Polytechnic of Central London, 92pp (unpublished).
- Heyns, J. (1993).** *Eudorylaimus nudicaudatus* sp. n. from Antarctica (Nematoda: Dorylaimoidea). *South African Journal of Antarctic Research* 23: 33-36.
- Holdgate, M.W. (ed.) (1970).** *Antarctic Ecology*. London and New York; Academic Press, 998pp (2 volumes). [Proceedings of 2nd SCAR Symposium on Antarctic Biology]
- Kito, K., Y. Shishida & Y. Ohyama (1996).** A new species of the genus *Eudorylaimus* Andrassy, 1959 (Nematoda: Qudisianematidae) from East Antarctica. *Polar Biology* 16: 163-169.
- Krijanova, E.S. (1958).** Antarkticheskie predsataviteli predstaviteli presnovodnykh nematodroela *Plectus* Bastian (Nematodes, Plectidae) (Antarctic specimens of freshwater nematodes of the genus *Plectus* Bastian (Nematoda, plectida). *Inf. Byull. Sov. Antarkt. Eksped.* No. 3 : 101-03 [English translation Vol. 1, 1964]
- Loof, P.A.A. (1975).** Dorylaimoidea from some subantarctic islands. *Nematologica* 21: 219-255.
- Seinhorst, J.W. (1959).** A rapid method for the transfer of nematodes from fixative to anhydrous glycerine. *Nematologica* 4: 67-69.
- Timm, R.W. (1971).** Antarctic soil and freshwater nematodes from the Mc Murdo Sound Region. *Proceedings of the Helminthological Society of Washington* 38: 42-52.

