

Linzer biol. Beitr.	52/2	1267-1306	Februar 2021
---------------------	------	-----------	--------------

Annotated supplements to catalogues of the family Scathophagidae (Diptera) in the world, with new taxonomic data, notes on some species and new list of species

Marco V. BERNASCONI & František ŠIFNER

A b s t r a c t : The last catalogues of the species of the family Scathophagidae for the individual zoogeographical regions were published in the following years: for the Afrotropical Region in 1976; for the Nearctic Region in 1965; for the Neotropical Region in 1984 and, partly, in 2010; for the Oriental Region in 1977 and for Palaearctic Region in 2008. The changes or additions published since the last catalogue form four basic groups: (I) new genera described (5 in Nearctic Region, 1 in Oriental Region, 8 in Palaearctic Region), (II) new species described (4 in Nearctic Region, 4 in Oriental Region, and 51 in Palaearctic Region), (III) new established synonyms (2 in Afrotropical Region, 1 in Nearctic Region, 4 in Neotropical Region, 4 in Oriental Region, 22 in Palaearctic Region), and (IV) new proposed combinations (1 in Afrotropical Region, 6 in Nearctic Region, 3 in Neotropical Region, 14 in Palaearctic Region).

K e y w o r d s : Diptera, Scathophagidae, taxonomy, zoogeographical regions, distribution, catalogue.

Introduction

The knowledge on the species of the family Scathophagidae is summarised in the catalogues dealing with the fauna of the individual zoogeographical regions. The following catalogues were published in the last fifty years: VOCKEROTH (1980) for the Afrotropical Region, VOCKEROTH (1965) for the Nearctic Region, ALBUQUERQUE (1984) and VOCKEROTH (2010) for the Neotropical Region, VOCKEROTH (1977) for the Oriental Region, and GORODKOV (1986) and ŠIFNER (2008) for the Palaearctic Region. The publication of Volume 4 of the new Manual of Afrotropical Diptera, containing also the Chapter dedicated to the Scathophagidae of this region should be available in 2022 (<https://www.nhm.ac.uk/our-science/our-work/biodiversity/manual-afrtropical-diptera.html>). The number of species recorded in the particular regions or subregions, including the individual countries, varies greatly and depends on the level of the performed faunistic and taxonomic research. The fauna of some regions or subregions are nearly to completely unknown (e. g. the Oriental Region and the Mediterranean Subregion of the Palaearctic Region).

This paper summarises the taxonomic changes made since the publication of the particular regional catalogues. Among the most important changes we must stress the changes based on revision of the oldest literature and verified by examination of the type specimens from Meigen's collection in Musée National d'Histoire Naturelle, Paris, and in Becker's collection in Museum für Naturkunde, Berlin and in Zoological Institute, St. Petersburg

(see OZEROV, 2014a,b and OZEROV & KRIVOSHEINA 2011c). Fundamental knowledge on the phylogenetic position of the family Scathophagidae among Muscoidea, on the phylogenetic relationships within Scathophagidae, and within *Scathophaga* in particular, as well as on the taxonomic status of some species, have been investigated by using various genetic markers (BERNASCONI et al. 1999, BERNASCONI 2000, BERNASCONI et al. 2000a,b, 2001, 2010; KUTTY et al. 2007). Therefore, the relationships in this family are now generally well understood (BERNASCONI et al. 2010).

Material and methods

In this paper, two subfamilies are recognised: the Delininae and the Scathophaginae (cf. COLLIN 1958). For the Palaearctic Region, we use the system of eight tribes proposed by ŠIFNER (2003, 2008). The text is arranged according to zoogeographical regions, and within each region the tribes, genera and species are arranged alphabetically.

All localities from the Czech Republic are accompanied with grid mapping codes according to PRUNER & MÍKA (1992). For the species of the Palaearctic Region, only the system proposed by Šifner (2003, 2008) is applied.

The following abbreviations of collections are used in the text:

- CNC.....Canadian National Collection, Ottawa, Canada;
- CAS.....Californian Academy of Sciences, San Francisco, USA;
- FSCP.....František Šifner privat collection, Praha (Chyňava), Czech Republic;
- ISEA.....Institute of Systematics and Ecology of Animals, Novosibirsk, Russia;
- MHNG.....Musée d'Histoire Naturelle, Genève, Switzerland;
- MNHN.....Musée National d'Histoire Naturelle, Paris, France;
- MZLU.....Museum and Department of Zoology, Lund University, Lund, Sweden;
- NHMW.....Naturhistorisches Museum, Wien, Austria;
- NMRB.....National Museum, Rio de Janeiro, Brasil;
- OU.....Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Hokkaido, Japan;
- SDEI.....Senckenberg Deutschen Entomologisches Institut, Münchenberg, Germany;
- SZMN.....Siberian Zoological Museum, Novosibirsk, Russia;
- TAUI.....Tel Aviv University, Tel Aviv, Israel;
- ZISP.....Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia;
- ZMUM.....Zoological Institute Moscow State University, Moscow, Russia;
- ZMHU.....Museum für Naturkunde der Humboldt University, Berlin, Germany.

Taxonomy

Afrotropical Region

The Afrotropical fauna of the family Scathophagidae is very limited and it is unambiguously influenced by the level of faunistic research. Only five species are known from this region (VOCKEROTH 1980), including *Scathophaga soror* (WIEDEMAN, 1800). The latter species was commonly but erroneously considered as a synonym of *Scathophaga stercoraria* (LINNAEUS, 1758). The details on *Scathophaga soror* are mentioned by OZEROV (2009a, 2017a), BERNASCONI et al. 2010, and by VOCKEROTH (1976, 1980).

***Scathophaga alata* (BECKER, 1894)**

Scopeuma alata (BECKER, 1894): 127

L e c t o t y p e : male, Mt. Kenya vers't Ouest, 2800-3200 m, Janv. 1912 (MNHN), designated by OZEROV (2009a), for detail OZEROV (2009a: 300-301 and 2017a: 187-190).

D i s t r i b u t i o n : Kenya (BECKER 1914, OZEROV 2009a).

Scathophaga foetulecta* (SÉGUY, 1966), as *Scatophaga

L e c t o t y p e : male, Cameroun, Mt. Babouto (MNHN), designated by OZEROV (2009a); for detail see OZEROV (2009a: 304 and 2017a: 193).

D i s t r i b u t i o n : Cameroon (SÉGUY 1966).

***Scathophaga longiaculeata* OZEROV, 2009a**

H o l o t y p e : male, Ethiopia, Shawa, Menageska forest, 3050 m a.s.l., 9°02'N 38°35'E, 19.IV.2005, A. Friedburg leg. (TAU).

D i s t r i b u t i o n : Ethiopia, Uganda (OZEROV 2009a).

***Scathophaga socia* (BECKER, 1894)**

Scopeuma socia (BECKER, 1894: 127)

L e c t o t y p e : male, Tanzania, Bismarck-Hügel, 2700-2800 m, IV.1912 (MNHN), designated by OZEROV (2009a); for detail OZEROV (2009a: 303 and 2017a: 191).

D i s t r i b u t i o n : Kenya, Tanzania (OZEROV 2009a).

Scathophaga soror* (WIEDEMAN, 1818), as *Scatophaga

This species was restituted as valid species by WERNER et al. 2006: 147.

Scatophaga capensis ROBINEAU-DESOVIDY, 1830: 625; OZEROV (2017: 192)

Scatophaga hottentota MACQUART, 1843: 342; OZEROV (2017: 192)

L e c t o t y p e : male, in Macquart collection under No. 1000 and locality "Cap" (MNHN), designated by OZEROV (2017a); for detail OZEROV (2017a: 192).

= *Scatophaga heleneae* (THOMSON, 1869); OZEROV (2012: 14-16).

Nearctic Region

The last catalogue of the Nearctic Region was compiled by VOCKEROTH (1965); the author listed 145 species and one subspecies. VOCKEROOTH (1987) described six monotypic genera which were not published with the term "gen.nov." or "nov.sp." as required Article 13 of the International Code of Zoological Nomenclature. These names, *Brokssiella*, *Dromogaster*, *Huckettia*, *Neorthocheta*, *Peratomyia* and *Synchysa*, were validated by VOCKEROOTH (1995). ŠIFNER (1977) described a new species, *Leptopa hendeli*, from the territory of United States; this species and the new genus record for the Nearctic Region were omitted by VOCKEROOTH (1987).

***Brockssiella* VOCKEROTH, 1995**

Type species: *Microprosopa varicornis* CURRAN, 1927: 257
Brockssiella VOCKEROOTH, 1987, nomen nudum

***Brockssiella varicornis* (CURRAN, 1927)**

new comb. by VOCKEROOTH (1995: 235)

Distribution: Canada (VOCKEROTH 1955).

***Coniosternum lapponicum* RINGDAHL, 1920**

Scathophaga minuta (MALLOCH, 1935); ŠIFNER (2019: 41)

***Coniosternum masneri* ŠIFNER, 2017**

Holotype: male, Canada, Nunavut, Chesterfield, N.W.T., 28.VII.1950, J. G. Chillcot leg.
(FSPC). Distribution: Canada (Šifner: 2017).

***Coniosternum molle* (BECKER, 1894)**

Comment: The transfer of *Scathophaga mollis* to the *Coniosternum* is based on the shape of the male terminalia which are typical for the genus *Coniosternum* (ŠIFNER 2019: 42).

***Coniosternum nigripalpis* (BECKER, 1907)**

Comment: The transfer of *Scathophaga nigripalpis* to the *Coniosternum* is based on the shape of the male terminalia which are typical for the genus *Coniosternum* (ŠIFNER 2019: 43).

***Dromogaster* VOCKEROTH, 1995**

Type species: *Dromogaster incompta* VOCKEROTH, 1995: 732;
Dromogaster VOCKEROOTH, 1987: 1094, nomen nudum

***Dromogaster incompta* VOCKEROTH, 1995**

Holotype: male, Canada: Billy, Alta. [= Alberta], 1.VI.1924, D. Bryant leg. (CAS).

Distribution: Canada: Alberta, Saskatschevan (VOCKEROTH 1995).

***Huckettia* VOCKEROOTH, 1995**

Huckettia VOCKEROOTH, 1987: 1096, nomen nudum

***Huckettia nearctica* VOCKEROOTH, 1995**

Type species: *Huckettia nearctica* VOCKEROOTH, 1995: 733

Holotype: male, Canada: Beaver Hill Lake, N.W.T., 66°44N 104°20W, 29.VI.1966, G. E. Shewell leg. (CNC)

Distribution: USA - California, Los Angeles (VOCKEROTH 1995)

***Leptopa hendeli* ŠIFNER, 1977**

Holotype: male, USA: Colorado, without data, with Hendel's handwritten label "*Leptopa* sp.n.", det. Hendel (NHMW).

Distribution: USA: Colorado (ŠIFNER 1977).

***Neorthochaeta* VOCKEROOTH, 1995**

Type species: *Ortachaeta dissimilis* MALLOCH, 1924: 174

Neorthochaeta VOCKEROOTH, 1987: 1096, nomen nudum

Neorthochaeta dissimilis (MALLOCH, 1924): 194, VOCKEROOTH (1995: 733).

Distribution: Canada, USA (VOCKEROTH 1995).

***Peratomyia* VOCKEROOTH, 1995**

Type species: *Hexamitocera vittata* COQUILLET, 1898: 165

Peratomyia VOCKEROOTH, 1987: 1096, nomen nudum

Peratomyia vittata (COQUILLET, 1898): 165, VOCKEROOTH (1995: 733).

Distribution: Canada, USA (VOCKEROTH 1995).

***Scathophaga islandica* (BECKER, 1894)**

status restituted by ŠIFNER (2018: 36)

Comment: The validity of this species was doubted by COLLIN (1931) and VOCKEROOTH (1965). JAMES (1950) first synonymized this species with *Scathophaga litorea*. Restored as valid species by ŠIFNER (2018: 36).

***Synchysa* VOCKEROOTH, 1995**

Type species: *Coenosia tricincta* LOEW, 1869: 177

Synchysa VOCKEROOTH, 1987: 1096, nomen nudum

***Synchysa tricincta* (LOEW, 1869)**

new comb. by VOCKEROOTH (1995: 734)

Distribution: Canada (VOCKEROTH 1995).

Neotropical Region

The first catalogue was compiled by VAN DER WULP (1897), the last catalogues by

ALBUQUERQUE (1984) and VOCKEROOTH (2010), with only few notes published subsequently. Only three genera and eight species are known from this region: *Scatogera* ALBUQUERQUE, 1984 (1 species), *Scatomyza* FALLÉN 1810 (1 species) and *Scathophaga* MEIGEN, 1803 (6 species).

***Scatomyza* FALLÉN, 1810**

C o m m e n t : *Scatomyza* was restored as valid genus by OZEROV & KRIVOSHEINA (2011a) based on the presence of a postmetacoxal bridge. This character was first introduced by VOCKEROOTH (2010) but without any taxonomic consequences.

***Scatomyza reses* (GIGLIO-TOS, 1893)**

Scatophaga reses (GIGLIO-TOS, 1893): 3

D i s t r i b u t i o n : Belize (former British Honduras) (GIGLIO-TOS 1893), Guatemala, Mexico (VOCKEROOTH 1965), Bolivia (ALBUQUERQUE 1984).

***Scatogera primogenita* ALBUQUERQUE, 1984**

D i s t r i b u t i o n : Ecuador (ALBUQUERQUE 1984) and Colombia (WOLFF et al. 2016).

***Scathophaga furcata* (SAY, 1823)**

Pyropa furcata (SAY, 1823): 15

D i s t r i b u t i o n : Mexico (ALBUQUERQUE 1984, VOCKEROOTH 1965), Guatemala, Costa Rica, Peru (VOCKEROOTH 2010).

Scathophaga chilensis* (ALBUQUERQUE, 1953), as *Scathophaga

D i s t r i b u t i o n : Chile (ALBUQUERQUE 1953).

***Scathophaga peruvensis* (LOPEZ & COURI, 1986), new combination**

Scopeuma peruvensis (LOPEZ & COURI, 1986): 314, new comb. by ŠIFNER in this paper.

H o l o t y p e : male, Peru, Mina Manuclita, Boja, without further data (NMRB).

C o m m e n t : This species was omitted in previous catalogues (ALBUQUERQUE 1984, VOCKEROOTH 2010). The transfer in the genus is realised by Šifner in this paper.

***Scathophaga stercoraria* (LINNAEUS, 1758)**

Musca stercoraria (LINNAEUS, 1758): 591

D i s t r i b u t i o n : Mexico, Haiti (ALBUQUERQUE 1984), Guatemala, Costa Rica (VOCKEROOTH 2010). Portorico, Uruguay (CUNY 1983).

Scathophaga tropicalis* (MALLOCH, 1931), as *Scatophaga

D i s t r i b u t i o n : Chile, Peru (MALLOCH 1931, CUNY 1983), Argentina, Bolivia (SZILADY 1926).

C o m m e n t : *Scathophaga tropicalis* (MALLOCH, 1931) is *nomen novum* for

Scatophaga lanata SZILADY, 1926: 595. This species name has been previously used in the genus by LUNDBECK for *Scatophaga lanata* LUNDBECK, 1901: 294 (type locality: Greenland), it was synonymized by VOCKEROTH (1965: 834) (see also ŠIFNER 2008a: 160) with *Scathophaga apicalis* (CURTIS, 1833). The species *Scatophaga lanata* sensu SZILADY (1926) was originally described from Bolivia: "Typen aus Cororica und "Cillumbicara... ein Männchen aus Rimconada, in 4.000 m Höhe"; 7-10 mm. Type material was deposited in Naturhistorisches Museum in Budapest (Hungary) and was destroyed in 1956. MALLOCH (1931: 443) noted: "...I have seen it from Peru in the collection of the United States National Musem. MALLOCH (1935) noted: "*Scatophaga tropicalis* is very similar to *stercoraria*. The only South America species of this genus."

***Scatomyza reses* (GIGLIO-TOS, 1893)**

Cordylura brevifrons VAN DER WULP, 1897; CUNY (1983: 12)
Cordylura vulpina VAN DER WULP, 1897; CUNY (1983: 12)
Scatophaga suilla var. *americana* MALLOCH, 1935; CUNY (1983: 13)
Scathophaga furcata (SAY, 1823)
Scatophaga vicina VAN DER WULP, 1897; CUNY (1983: 13)

Oriental Region

The Oriental Region reaches northward to 23° to 25° N with a transition area in the southwestern lowland of China, southward to Malaysia and eastward approximately to Wallace line. The first data related to the family Scathophagidae were published in the WALKER's (1860) catalogue and subsequently by VAN DER WULP (1896); both the authors cited only one species, *Cordylura bisignata* WALKER, 1860, from Sulawesi. VOCKEROTH (1977) mentioned four described and seven unnamed species in six genera.

***Cordilura bisignata* (WALKER, 1860)**

Cordylura bisgnata (WALKER, 1860): 142

D i s t r i b u t i o n : Sulawesi (in original text Macassar) (WALKER 1860).

N o t e : The validity of this species is unclair and the revision of the taxon is necessary; OOSTERBROEK (1998) wrote it "...does not belong to the Scathophagidae".

***Cordilura pulchra* OZEROV & KRIVOSHEINA, 2012**

H o l o t y p e : male, Vietnam, Lai Chau Province, Hoang Lien National Park, 22.347948°N 103.769714°E, 1900 m [a.s.l.], 19.iv.2012, A. L. Ozerov leg. (ZMUM).

D i s t r i b u t i o n : Vietnam (OZEROV & KRIVOSHEINA 2012).

Lubomyia Šifner, 2010

T y p e s p e c i e s : *Lubomyia orientalis* ŠIFNER, 2010

***Lubomyia orientalis* ŠIFNER, 2010**

H o l o t y p e : male, N.E. Burma, Kombaiti, 7000n. [?], 9.iv.1934, R. Malaise [leg?] (CNC).

C o m m e n t : This specis was mentioned by VOCKEROTH (1977: 436) as an unnamed species of the genus *Acerocnema*.

***Parallelolomma merzi* OZEROV, 2011a**

Distribution: Thailand, Vietnam (OZEROV & KRIVOSHEINA 2011a).

Comment: MERZ (2006: 224) cited one male of *Parallelolomma* sp. from MHNG as "...first genus of this family recorded from Thailand". This specimen was later described by OZEROV & KRIVOSHEINA (2011a) as a new species.

***Scatomyza curtipilata* (FENG, 2002)**

Scatomyza fascipaniscola OZEROV & KRIVOSHEINA, 2011a: 5; OZEROV & KRIVOSHEINA (2018: 2)

Heteroterra fajingensis WEI, 2006: 531, OZEROV & KRIVOSHEINA (2018: 2)

Scatomyza mellipes* (COQUILLETT, 1899), as *Scatophaga

Scatophaga chinensis MALLOCH, 1935: 260, OZEROV & KRIVOSHEINA (2018: 7)

Scathophaga eoa OZEROV, 2007: 2, OZEROV & KRIVOSHEINA (2018: 7)

Distribution: China, India (VOCKEROTH, 1977), Vietnam, Japan, Nepal, Russia, Korea (OZEROV & KRIVOSHEINA 2011a).

***Scatomyza scybalaria* (LINNAEUS, 1758)**

Musca scybalaria (LINNAEUS, 1758): 599

Distribution: Vietnam (OZEROV & KRIVOSHEINA 2011a: 8).

***Scathophaga stercoraria* (LINNAEUS, 1758)**

Musca stercoraria (LINNAEUS, 1758): 599.

Distribution: China, India (VOCKEROTH 1977, ŠIFNER 2010)

Palaearctic Region

The last catalogue of Palaearctic Scathophagidae was compiled by ŠIFNER (2008a), with some additions and corrections supplemented by ŠIFNER (2009); it contains 38 genera, 221 valid species, 198 synonyms and 12 doubtful species.

Subfamily Delininae ROBINEAU-DESOVIDY, 1830**Tribe Delinini ŠIFNER, 2003*****Americina caucasica* (OZEROV, 2009b)**

Parallelolomma caucasica (OZEROV, 2009b): 314

Holotype: male, Russia, Adygeya, N Lagonaki Mt., 44:093° 40.019°E, 1725 a.s.l., *Picea* forest, 28.VI.2009, coll. K. Tomkovish (ZMUM).

***Americina hostae* (HERING, 1955)**

Chylizosoma hostae (HERING, 1955): 6

Comment: This species was synonymized by ŠIFNER (1978: 290) with *Americina*

vittata (MEIGEN, 1826) (cf. ŠIFNER 2008a); revalidated by OZEROV (2009d: 312 in Russian, 317 in English) as a valid species.

***Micropselapha basovi* OZEROV, 2010b**

Micropsalapha bohemica BARTÁK & KUBÍK 2012: 101, OZEROV (2014a: 320), (cf. ŠIFNER 2018: 1641, Note 1).

H o l o t y p e : male, Russia, Tatarstan, Kamskiy Reserve, Lake Raifa, 55.897357°N 48.733022°E, 13.v.2005, V. Basov leg. (ZMUM).

***Mirekiana* ŠIFNER, 2012**

Mirekiana anthrax (SCHINER, 1864): 16, ŠIFNER (2012: 17).

= *Cleigastra carbonaria* POKORNY, 1887: 441, STROBL (1898: 182).

T y p e s p e c i e s : *Cleigastra anthrax* SCHINER, 1864

H o l o t y p u s : male, Czech Rep., Krkonoše Mts., (National Park), Úpské rašeliniště (peat bog) 21.-22.vii.1966, Šifner leg.

***Neochirosia nigriceps* (BECKER, 1894)**

C o m m e n t : The transfer *Delina nigriceps* (BECKER, 1894) in the genus *Neochirosia* MALLOCH, 1919 was made by OZEROV (2009c: 238).

***Neochirosia nikita* OZEROV & KRIVOSHEINA, 2015**

H o l o t y p e : male, Serbia, Stara Planina, 43.37°N 22.60°E, 1.-8. V. 2015, N. Vikhrev (ZMUM).

***Neochirosia pechorica* OZEROV 2009c**

H o l o t y p e : male, Russia, Archangelskaya oblast [= region], Nametskiy Reserve, 68:175957°N 53.645394°E, 19. viii. 2008, A. L. Ozerov leg. (ZMUM).

***Neochirosia veratri* (HENDEL, 1925)**

C o m m e n t s : *Clidogastra veratri* was synonymized by ŠIFNER (2003: 63, 2008a: 175) with *Delina nigriceps* (BECKER, 1894); it was revalidated and transferred in the genus *Neochirosia* by OZEROV (2009c: 239). ŠIFNER (2018: 1637) mentioned *Neochirosia cepelaki* as valid in the genus *Neochirosia* by mistake.

S u b f a m i l y Scathophaginae ROBINEAU-DESVOIDY, 1830

T r i b e Amaurosomini ŠIFNER, 2003

C o m m e n t s : The use of the name *Amaurosoma* BECKER, 1894 or *Nanna* STROBL, 1894 was interpreted correctly by PÜCHEL (1999): the name *Nanna* STROBL, 1894 is a synonym of *Amaurosoma* BECKER, 1894. This problem was resolved in detail by ŠIFNER (2003).

Amaurosoma amurensis* (OZEROV, 2010b), as *Nanna

H o l o t y p e : male, Russia, Amurskaya oblast, town Zeya, 53.7485°N 127.2614°E, 14.vi.1978, A. Shatalkin leg. (ZMUM).

***Amaurosoma articulatum* BECKER, 1894**

L e c t o t y p e : male, Number 29936 in Meigen's collection, 6.v. (MNHN), designated by OZEROV (2014b: 35) (see also ŠIFNER 2008a: 151).

***Amaurosoma cryophila* (OZEROV & KRIVOSHEINA, 2015), new combination**

Nanna cryophila (OZEROV & KRIVOSHEINA, 2015): 228. The transfer in the genus *Amaurosoma* was made by ŠIFNER in this paper.

H o l o t y p e : male, Russia, Khabarovsk Region, Khicka R., 690 m, 49.05°N 137.43°E, 10.vi.2014, N. Vikhrev leg. (ZMUM).

***Amaurosoma fasciatum* (MEIGEN, 1826)**

Cordylura fasciatum (MEIGEN, 1826): 238

H o l o t y p e : male, Number 2153 in Meigen's collection (MNHN), labelled by MEIGEN as *fasciata* (MNHN), for detail see OZEROV (2017a: 190).

***Amaurosoma longicorne* (VON ROSER, 1840)**

Cordylura longicorne (VON ROSER, 1840)

H o l o t y p e (?): Germany, environs of Württemberg (cf. ŠIFNER 2008a: 154).

C o m m e n t : The redescription of this species based on one male and two females from Karelia by OZEROV (2010a: 3-5). The redescription of this species by ŠIFNER (2008b: 104) was a misidentification; the figures 1 to 3 (p. 104) belong to *Amaurosoma obenbergeri* ŠIFNER (2016).

***Amaurosoma obenbergeri* ŠIFNER, 2016**

Amaurosoma longicorne (VON ROSER, 1840) (misidentification): ŠIFNER (2016: 114).

H o l o t y p e : male, Czech Republic, Bohemia mer., Malšice (6653), 4.x.1976, Jan Máca leg. (FSPC).

***Amaurosoma obscuripes* (BECKER, 1915)**

Nanna indotatum ENGELMARK, 1999: 163, OZEROV & KRIVOSHEINA (2015: 144).

C o m m e n t : The species *Acerocnema obscuripes* BECKER, 1915 was considered by ŠIFNER (2008a: 198) as a junior synonym of *Acerocnema macrocera* (MEIGEN, 1826). However, OZEROV & KRIVOSHEINA (2015) confirmed it as valid species and included this species in the genus *Nanna* STROBL, 1894.

***Amaurosoma rossolimoae* (OZEROV, 2010b), new combination**

Nanna rossolimoae (OZEROV, 2010b): 161; The transfer in the genus *Amaurosoma* was made by ŠIFNER in this paper.

H o l o t y p e : male, Russia, Amurskaya oblast, town Zeya, 53.7485°N 127.2614°E, A. Ozerov leg. (ZMUM).

***Gabreta* ŠIFNER, 2015**

T y p e s p e c i e s : *Gabreta macai* ŠIFNER, 2015: 75

***Gabreta macai* ŠIFNER, 2015**

H o l o t y p e : male, Czech Republic, Bohemia mer., Modrava (6946), Modravská Medvědí Mt., 1200 m, beertrap, 1.vi.2014, Jan Máca leg. (FSPC).

***Julienomyia* ŠIFNER, 2015**

T y p e s p e c i e s : *Julienomyia miroslavi* ŠIFNER, 2015: 75

***Julienomyia miroslavi* ŠIFNER, 2015**

H o l o t y p e : male, Czech Republic, Bohemia mer., Sezimovo Ústí (6654), Luna Natur Monument, 400 m, 1.vi.2001, Jan Máca leg. (FSPC).

***Miroslava knajfli* ŠIFNER, 2010**

H o l o t y p e : male, Nepal, 27°57'N 84°59'E, Malaise trap No. 5, Canadian Nepal Expedition leg., as *Megaphthalmoides* sp. [det.?] (CNS).

***Gonatherus planiceps* (FALLÉN, 1826)**

Gonatherus fumipennis HENDEL, 1930: 7, OZEROV (1930: 6)

Tribe Cleigastrini ŠIFNER, 2003

OZEROV & KRIVOSHEINA (2015) based on the variability of chaetotaxy and the shape of some sternites proposed the following nomenclatorial acts: The genera *Gonarcticus* BECKER, 1894, *Gonatherus* RONDANI 1856, *Orthacheta* BECKER, 1894, *Spathephilus* BECKER, 1894 and *Amaurosoma* BECKER, 1894 (= *Nanna* STROBL, 1894 sensu OZEROV & KRIVOSHEINA 2010a) are proposed as synonyms of *Cleigastra* MACQUART, 1835. These authors recommend three subgenera: *Cleigastra* MACQUART s. str., *Gonatherus* RONDANI and *Orthacheta* BECKER. We do not follow this opinion in this paper.

***Acerocnema arctica* OZEROV, 2013**

H o l o t y p e : male, Russia, Krasnoyarskiy Kray, Lake Taymyr, Baykuranera Bay, 74.117°N 102.95°E, 30. vii.-7. viii.1976, A. Ranitsin leg. (ZMUM).

***Acerocnema breviseta* (ZETTERSTEDT, 1846)**

Cordylura breviseta ZETTERSTEDT, 1846: 2022

Acerocnema tiefi BECKER, 1894: 155, ŠIFNER (2003: 47).

Acerocnema pokornyi BECKER, 1894: 157, ŠIFNER 2003: 47.

H o l o t y p e : female, Denmark: "circa Hafniam" [Copenhagen env.], labelled: "[18]62, as "breviseta Staeger 41", as holotype designated by J. R. Vockeroth in 1954 (MZLU).

C o m m e n t : Confirmed as valid species by OZEROV (2014a: 317).

***Acerocnema flavifrons* IWASA, 2013**

H o l o t y p e : male, Japan, Nakasatsunai, Tokachi, Hokkaido, 26.v.2013, M. Iwasa leg. (OU).

***Acerocnema kishimotoae* IWASA, 2013**

H o l o t y p e : male, Japan, Toyokoro, Tokachi, Hokkaido, 4.v.2013, M. Iwasa leg. (OU).

***Acerocnema sinuata* IWASA, 2013**

H o l o t y p e : male, Japan, Nakasatsunai, Tokachi, Hokkaido, 26.v.2013, M. Iwasa leg. (OU).

***Acerocnema richterae* OZEROV, 2015**

H o l o t y p e : male, Russia, Buryatia, Chamney, 50.402°N 103.8681°E, bottom land of Shibertuy River, 29.iv.971, V. Richter leg. (ZISP).

***Cleigastra intermedia* OZEROV & KRIVOSHEINA, 2016**

H o l o t y p e : male, Russia, Novosibirsk Region, Solair Range, 54.55°N 84.86°E, 19.v.2015, O. Kosterin leg. (ZMUM).

***Cleigastra maritima* OZEROV & KRIVOSHEINA, 2016**

H o l o t y p e : male, Russia, Promorskiy Kray, Gribnoe, 44.2574°N 132.6817°E, 10 km SE of Chernigorka, 10.v.1979, A Zinoviev leg. (ZISP).

***Cleigastra sundukovi* OZEROV, 2013**

H o l o t y p e : male, Russia, Kuril Islands, Shikotan I., Traskovnaya Bay, 43.75°N 146.70°E, 10.-14.vi. 2002, Yu. Sundukov leg. (ZMUM).

Hajekiana ŠIFNER, 2016

T y p e s p e c i e s : *Hajekiana orlicensis* ŠIFNER, 2016.

***Hajekiana orlicensis* ŠIFNER, 2016a**

H o l o t y p e : male, Czech Republic, Orlické hory Mts., National reserve Bukačka (5664), 18.vi.-5.viii.1994, yellow pan trap., Hájek leg. (FSPC).

***Spathophilus nigriventris* (LOEW, 1864)**

Cordylura nigriventris (LOEW, 1864): 19

L e c t o t y p e : female, labelled "23/5 43, 11392", coll. Loew, Type *C. nigriventris* m., type locality "Posen" [Poland, Poznań] (ZMHB).

= *Cordylura breviventris* LOEW, 1873: 250, OZEROV (2010a: 5, new synonymy).

C o m m e n t : This species was cited by ŠIFNER (1968: 167, 2008a: 150-151) as *Spathophilus breviventris* (LOEW, 1873); the species *Cordylura nigriventris* was considered erroneously as a synonym of *Amaurosoma flavipes* (FALLÉN, 1819) (cf. ŠIFNER 2008a: 152-153).

Tribe Cordilurini ŠIFNER, 2003***Cordilura aberrans* (BECKER, 1894)**

Cordylura rubifromata BECKER, 1894: 91, OZEROV (2014b: 41).

***Cordilura albilabris* (FABRICIUS, 1805)**

Ocyptera albilabris (FABRICIUS, 1805): 315

C o m m e n t : This species was formerly commonly placed in the genus *Phrosia* ROBINEAU-DESOUDY, 1830. Nowadays, BERNASCONI & al. (2000) and KUTTY & al (2007), based on DNA sequences, proposed it placed among of *Cordilura* species.

***Cordilura bezzi* (SACK, 1937)**

Cordylura bezzi (SACK, 1937): 18

L e c t o t y p e : male, Livrio, without data, designated by VOCKEROOTH 1954 (SDEI).

C o m m e n t : This species was placed by ŠIFNER (2008) in the genus *Bucephalina* MALLOCH, 1909.

***Cordilura ciliata* (MEIGEN, 1826)**

Cordylura ciliata (MEIGEN, 1826): 231

L e c t o t y p e : male, Number 2137 in Meigen's collection (MNHN), designated by Ozerov; for detail see OZEROV (2017a: 190).

***Cordilura fulvifrons* OZEROV, 1997**

C o m m e n t : OZEROV & KRIVOSHEINA (2014a) completed the original data on the type locality: "This species was described from Boitsova 46.9823°N 134.3246°E. ŠIFNER (2008a) included this species in the genus *Scoliaphleps* BECKER, 1894; by OZEROV (1997) it is placed in the genus *Cordilura*.

***Cordilura fuscipes* (ZETTERSTEDT, 1838)**

Cordylura apicata HENDEL, 1830: 4, OZEROV (2007: 6).

Cordilura grunini OZEROV & KRIVOSHEINA, 2017a

H o l o t y p e : male, Russia, Primorskiy kray, valley of the river Serebryanka, 45.04878°N 136.63558°E, 17.iv.1937, Grunin leg. (ZISP).

***Cordilura kosterini* OZEROV & KRIVOSHEINA, 2014a**

H o l o t y p e : male, Russia, Primorskiy Kray, Lotus Lake, 42:46°N 130.64°E, 1.-3.viii.2014, O. Kosterin leg. (ZMUM).

***Cordilura monochroma* OZEROV & KRIVOSHEINA, 2014a**

H o l o t y p e : male, Russia, Sachalin oblast, Kuril Islands, Kunashir I., 43.026°N 145.862°E, 6.vii.2013, Yu. Sundukov leg. (ZMUM).

***Cordilura negrobovi* OZEROV & KRIVOSHEINA, 2017a**

H o l o t y p e : male, Russia, Republic of Buryatia, East Sayan, Arschan, Tagarlay, 51.8786°N 192.3927°E, 3.vii.1965, Negrobov leg. (ZISP).

***Cordilura picticornis* (LOEW, 1864)**

Codylura picticornis (LOEW, 1864): 22

Cordylura latigenis HENDEL, 1930: 3, OZEROV (1007: 7).

L e c t o t y p e : female, Number 11369, designated by Vockeroth 19[54] (ZMHB), for detail see OZEROV (2014b: 40).

Cordilura rufipes* (MEIGEN, 1826)Codylura rufipes* (MEIGEN, 1826): 242

H o l o t y p e : Number 2144, in Meigen's collection (MNHN); for detail see OZEROV (2017a: 193).

***Cordilura sidorenkovi* OZEROV & KRIVOSHEINA, 2012a**

H o l o t y p e : male, Russia, Primorskiy Kray, Larorskij reserve, 43.0052°N 134.1238°E (ZMUM).

Milania ŠIFNER, 2010T y p e s p e c i e s : *Norellisoma agrion* SÉGUY, 1948: 169.***Milania agrion* (SÉGUY, 1948)**

new comb. by ŠIFNER (2010: 610)

Norellisoma agrion* SÉGUY, 1948**Milania longiabdomina* (SUN, 1992)***Norellia longiabdomina* (SUN, 1992), ŠIFNER (2010: 612).***Mixocordylura parva* OZEROV, in OZEROV & KRIVOSHEINA, 2011b**

H o l o t y p e : male, Russia, Altay, Kurayski range, 2500 m a.s.l., 50.33°N 87.75°E, mountain tundra, 3.viii.2008, A. Barkalov leg. (SZMN).

Norellisoma armipes* (MEIGEN, 1826)Cordylura armipes* (MEIGEN, 1826): 234

L e c t o t y p e : male, number 2142 in Meigen's collection (MNHN); for detail see OZEROV (2017a: 190).

C o m m e n t : The male genitalia and abdominal sternites 5 and 4 were dissected and illustrated by OZEROV (2009b). This species was erroneously synonymized as *Norellisoma spinimanum* by ŠIFNER (2008a: 130).***Norellisoma flavicorne* (MEIGEN, 1826)***Cordylura flavicorne* (MEIGEN, 1826): 239

L e c t o t y p e : male, Number 2157 in Meigen's collection (MNHN), originally labelled by Meigen (Dr. L. Matile, pers. comm. to F. ŠIFNER in 1970); for detail see OZEROV (2017a: 191).

C o m m e n t : The validity of this species is under debate (cf. OZEROV 2017a: 191). However, the examined specimens from the Czech Republic including their variability confirm unambiguously *N. flavicorne* as a valid species (ŠIFNER, unpubl.)***Norellisoma flavostriatum* OZEROV 2008a**

H o l o t y p e : male, Russia, Sochi, environs Estosadok, Mt. Psechatsa, subalp. (43°41'28"N 40°22"E), 14.-18.vi.2008, K. Tomkovich leg. (ZMUM).

Norellisoma lituratum* (MEIGEN, 1826)Cordylura lituratum* (MEIGEN, 1826): 238

L e c t o t y p e : male, Number 2141 im Meigen's collection (MNHN); for detail OZEROV (2017a: 192).

= *Norellisoma altaicum* (OZEROV, 2008): 2 as *Norellia*, OZEROV & KRIVOSHEINA (2020b: 110).***Norellisoma nervosum* (MEIGEN, 1826)***Cordylura nervosum* (MEIGEN, 1826: 234)

L e c t o t y p e : male, Number 2141 in Meigen's collection (MNHN); for detail see OZEROV (2017a: 192).

***Norellisoma nigrovenosum* OZEROV, 2008a**

H o l o t y p e : male, Azerbaijan, Lenkoran Distr., Burzaly, 38°65'55"N 48°77'9"E, 27.x.2008, D. Daryushin leg. (ZMUM).

Norellisoma oreinum* Ozerov, 2009b**H o l o t y p e : male, Russia, Adygea, N Lagonaki Mt., Arich cave env, 44°09'3"N 40°01'9"E, 1725 m a.s.l., *Picea* forest, 26.-28.vi.2009, coll. K. Tomkovich (ZMUM).Norellisoma spinimanum* (FALLÉN, 1819)***Norellisoma septentrionale* HENDEL, 1930: 2, OZEROV (2007: 7).***Norellisoma striolatum* (MEIGEN, 1826)***Cordylura striolatum* (MEIGEN, 1826): 235

L e c t o t y p e : male, Number 2146 in Meigen's collection (MNHN); for detail see OZEROV (2017a: 194).

***Norellisoma tomkovichi* OZEROV, 2010b**

H o l o t y p e : male, Russia, Adygea, N Lagonaki Mt., 1830 m, 44°05'0"N 40°01'8"E, meadow, 20.-23.8.2009, coll. K. Tomkovich (ZMUM).

Norellisoma vockerothi* OZEROV, 2013**H o l o t y p e : male, Switzerland, St. Moritz, 3/7 1057 D, 46°58'N 9°83"E, det J. R. Vockeroth 1954 as *Norellisoma* n.sp. (ZMUM).Parallelomma albipes* (FALLÉN, 1819)***Cordylura unicolor* LOEW, 1864: 16, OZEROV (2014b: 41)L e c t o t y p e : *C. unicolor*, female, Number 11373, designated by Vockeroth (SDEI).C o m m e n t : *Cordylura unicolor* was erroneously synonymized by ŠIFNER (2008a: 114) with *Americina vittata* (MEIGEN, 1826).***Parallelomma fuscipes* (ZETTERSTEDT, 1838)***Cordylura apicata* HENDEL, 1930: 4, OZEROV (2007: 6)*Cordylura nigrithorax* HENDEL, 1930: 5, OZEROV (2007: 7)

Cordylura ochracea HENDEL, 1930: 6, OZEROV (2007: 7)

***Suwaia* ŠIFNER, 2009**

Type species: *Parallelomma longicornis* HENDEL, 1913: 77.

***Suwaia longicornis* (HENDEL, 1913)**

new comb. by ŠIFNER (2009: 290)

***Parallelomma longicornis* HENDEL, 1913**

Comment: This species was placed by VOCKEROTH (1977: 437) in the genus *Megaphthalma* BECKER, 1894; for details see OZEROV (2014b: 38).

Tribus Gimnomerini ŠIFNER, 2003

***Gimnomera* RONDANI 1866**

Langechristia OZEROV, 1999: 510, OZEROV (2019a: 197).

***Gimnomera amica* (OZEROV, 1999)**

comb.nov., des. by OZEROV (2014: 34)

Comment: The holotype of this species is a female; ŠIFNER (2008a: 135) erroneously cited it as a male.

***Gimnomera kirgizica* OZEROV, 2008a**

Holotype: male, Kyrg. [yzstan], Issykkul area, Ala Bush pass, 2365 m a.s.l., 13.vii.[19]94, Milko leg. (MHNG).

***Gimnomera montana* OZEROV & KIROVHEINA, 2013**

Holotype: male, Russia (North Ossetia-Allama, range Bakhty Laparyrag, (42.9383°N 44.2872°E), 1720m. (ZISP)

***Gimnomera nigricorpus* OZEROV, 2019b**

Holotype: male, Russia, Krasnoyarski Krai, Taimyr, Agapa (ca. 71.412°N 89.2689°E), the River Pyasia, 14.vii. 167, Gorodkov (ZISP).

***Gimnomera novgorodae* (OZEROV, 2006)**

Cochliarum novgorodae (OZEROV, 2006): 334

Cochliarium sorokiniae OZEROV, 2006: 335, OZEROV (2019b: 207)

***Gimnomera nudipedis* OZEROV, 2019b**

Holotype: male, Russia, Krasnoyarski Krai, Taimyr, Agaps (ca: 71.412°N 89.2689°E), the River Pyasia, 14.vii.1967, Gorodkov (ZISP).

***Gimnomera palliseta* OZEROV, 2019b**

H o l o t y p e : Male, RUSSIA, Magadan Oblast, Sokol env. (59°92'N 150°71'E), 11-19. vii. 2014, N. Vikhrev (ZMUM).

***Gimnomera speciosa* (OZEROV, 1999)**

comb. nov., des. by OZEROV (2014: 41)

***Gimnomera tibialis* (MALLOCH, 1919),**

Dasypleuron tibialis (MALLOCH, 1919)

Cochliarium sibiricum ENGELMARK, 1999: 164, OZEROV (2019b: 212).

***Gimnomera ziegleri* (OZEROV, 1999)**

comb.nov., des by Ozerov (2014: 42)

***Norellia altaica* OZEROV 2008b**

H o l o t y p e : male, Russia, Altai Kray, 45 km E Ust-Ulagan, the upper reaches of the Old River, 50.5°N 88.57°E, 2020 m, 17.vii.2005, coll. V. Sorokina (ZMUM). Female unknown.

***Norellia spinipes* (MEIGEN, 1826)**

Corylura spinipes (MEIGEN, 1826): 237

L e c t o t y p e : male, with disc. no."2349/40" in Meigen's collection (MNHN); designated by OZEROV (2017a: 193) as lectotype.

***Norellia tipularia* (FABRICIUS, 1794)**

Musca tipularia (FABRICIUS, 1794): 337

C o m m e n t : *Musca tipularia* was considered nomen dubium (ŠIFNER 2003: 32, 2008a: 179); DE JONG (1985: 21) cited *Norellia* (= *Musca*) *tipularia* as a valid species; *Norellia tipularia* was redescribed by OZEROV & FREIDBERG (2010). The type specimen of *Musca tipularia* is not present in the collection of FABRICIUS (ZIEMSEN 1964) (cf. ŠIFNER 2003a: 32).

***Norellisoma yolduense* (OZEROV, 2008b)**

H o l o t y p e : male, Russia, Altai Kray, 45 km E Ust-Ulagan, the upper reaches of the Old River, 50°N 88.57°E, 2020 m, 19.vii.2005 (SZMN).

T r i b e *Hydromyzini* ŠIFNER, 2003

The genus *Okeniella* HENDEL, 1907 was synonymized with the genus *Pogonota* ZETTERSTEDT, 1860 by OZEROV (2016). The basic differences between species of both genera are based on the shape of cerci: in species of *Okeniella* those are separated and fused only basally, in species of *Pogonota* those are fused almost at the whole length; there are differences in the morphology of fore legs and the shape of male abdominal sternites 4 to 5. KUTTY & al. (2007), based on DNA sequences, suggested the both genera as synonyms.

***Bostichopyga crassipes* (ZETTERSTEDT, 1838)**

Bostichopyga borealis HENDEL, 1930: 385, OZEROV (2014b: 87).

H o l o t y p e of *B. borealis*: male, Norway, Austad, 58.090°N 7.0442°E, Norway (Salt) Strand, red label *Bostichopyga* sp.n., Type. *Bostichopyga borealis*, Mus. Caes. Vidobob. (NHMW). For details see OZEROV & KRIVOSHEINA (2014b: 87).

C o m m e n t : In the original HENDEL's (1930) paper the data on the type of *B. borealis* were only fragmentary (cf. ŠIFNER 2008a: 141).

***Chaetosa punctipes* (MEIGEN, 1826)**

L e c t o t y p e : male, with a disc. no."2352/40" in Meigen's collection (MNHN), designated by OZEROV (2017a: 193) as lectotype.

***Pleurochaetella barkalovi* OZEROV & KRIVOSHEINA, 2012b**

H o l o t y p e : male, Russia, Krasnoyarsk Kray, NW Taimyr pen., 12.5 km S. Oreon. settl., 73°24'N 80°39'E, bank of Lemberova River, 6.-26.vii.2012, coll. A. Barkalov (ISEA).

***Spaziphora tomkovichi* OZEROV, 2011**

H o l o t y p e : male, Russia, Chukotka, Amadyr. Distr., Meynipylgino, 62.576°N 177.023°E, lake morene, 10.-13.vii.2011, P. S. Tomovich leg. (ZMUM).

T r i b e *Microprosopini* ŠIFNER, 2003

OZEROV (2017a) included the genus *Allomyella* MALLOCH, 1923 as a subgenus in the genus *Microprosopa* BECKER, 1894 based on the presence or absence of the hairs on anepimeron and the shape of male abdominal sternite 5.

ŠIFNER (2013a) reclassified the genus *Microprosopa* and the subgenus *Paramicroprosopa* sensu RINGDAHL, 1936 which he raised to the generic rank, providing key to genera of the *Microprosopini*. In the genus *Paramicroprosopa* he included the following species: *P. hoherlandti* (ŠIFNER, 1981), *P. bartaki* (ŠIFNER, 1999), *P. frontata* (ZETTERSTEDT, 1838), *P. ozerovi* (ŠIFNER, 2008b), *P. pokornyi* (ŠIFNER, 2011).

***Acanthocnema sternalis* SUWA, 1986**

Ancanthocnema vikhrevi OZEROV & KRIVOSHEINA, 2014b: 204 as *Acanthocnema*. OZEROV (2019a: 330).

***Allomyella unguiculata* (MALOCH, 1919)**

Allomyia unguiculata (MALOCH, 1919):80

Allomyella borealis CURRAN, 1927: 260, OZEROV & BARKALOV (2014: 469), by author's first discovery in the Palaearctic Region.

***Megaphthalmoides japonicus* OZEROV, 2008a**

H o l o t y p e : male, Japan, Osaka Prefecture, 800 m, Izumi Katwiagi, 24.ix.1993, A. Friedberg & F. Kaplan leg. (TAUI).

***Microprosopa gruberi* ŠIFNER, 2013a**

H o l o t y p e : male, Slovakia, Malá Fatra Mts., Štefanová - Stohové pol'any (= mountain meadows), 17.v.1975, Šifner leg. (FSPC).

***Microprosopa paveli* OZEROV & KRIVOSHEINA, 2013b**

H o l o t y p e : male, Russia, Chukotka, Meynipygino, 62.5385°N 177.0519°E (ZMUM).

***Microprosopa preisleri* ŠIFNER, 2013a**

H o l o t y p e : male, Czech Republic, Krkonoše Mts. (5660), Modrý důl (= valley), 10.vii.1963, Šifner leg. (FSPC).

***Microprosopa taymirica* OZEROV, 2017b**

H o l o t y p e : male, Russia, Krasnodarskiy Kray, Lake Engel'gardt, 75.101°N 100.2349°E, 2.viii.1967, Gorodkov leg. (ZISP).

***Microprosopa zemani* ŠIFNER, 2013a**

H o l o t y p e : female, Czech Republic, Krkonoše Mts. (5659), Sedmidolí, Erlenbach bouda (= chalet), 8.-18.vi.1968, Doskočil leg. (FSPC).

***Microprosopa zlobini* OZEROV, 2008a**

H o l o t y p e : male, Russia, North Ossetia, environs of Buron, Tsey gorge, 42.793055°N 43.92216°E, 1764 m (ZMUM).

Paramicroprosopa bartaki (ŠIFNER, 1999), as *Microprosopa*

Paramicroprosopa frontata (ZETTERSTEDT, 1838), as *Scatomyza*

Paramicroprosopa hoberlandti (ŠIFNER, 1981), as *Microprosopa*

Paramicroprosopa ozerovi (ŠIFNER, 2008), as *Microprosopa*

Paramicroprosopa pokornyi (ŠIFNER, 2011), as *Microprosopa*

H o l o t y p e : male, Czech Republic, Chyňava (5950), PLA Křivoklátsko, 50.040°N, 14.037°E, 11.-14.vi.2010, Šifner leg. (FSPC).

***Trichopalpus fraternus* (MEIGEN, 1826)**

Cordylura fraternus (MEIGEN, 1826): 243

L e c t o t y p e : male, Number 2162 in Meigen's collection ((MNHN); designated by OZEROV (2017a: 191) (see also ŠIFNER 2008a: 140).

T r i b e Scathophagini ŠIFNER, 2003***Coniosternum alashanicum* (OZEROV & KRIVOSHEINA, 2019c), new combination**

as *Scathophaga*, comb.nov. by ŠIFNER, in this paper.

H o l o t y p e : male, China, [Alashan or Alxa, 38.8442°N 105.7070°E], 30.v.1908, Kozlov (ZISP).

***Coniosternum decipiens* (HALIDAY, 1832)**

Coniosternum fluviale (RONDANI, 1867): 113, OZEROV & KRIVOSHEINA (2020a: 227)
 Comment: This nomenclatorial act is unclear; it is not figured the pregonite.

***Coniosternum dvoraki* ŠIFNER, 2016**

Holo type: male, Czech Republic, Slavkovský les (= forest), Lazy, Lesný (5941), smrčina (= spruce forest), pivní past (= beer trap), 12.-30.vi.2004, Dvořák leg. (FSPC).

***Coniosternum kovari* ŠIFNER, 2013b**

Holo type: male, Tajikistan, Hissar Mts., Vaahab distr., Magov, 38°40N 69°06E, 18.-21.ix.1989, I. Kovář leg. (NMPC).

***Coniosternum lapponicum* RINGDAHL, 1920**

Scatophaga minuta MALLOCH, 1935: 255, ŠIFNER (2018b: 41).

Comment: The species *S. minuta* was erroneously synonymized with *Coniosternum obscurum* (FALLÉN, 1819) by VOCKEROTH (1965: 839).

***Scatomyza* FALLÉN, 1810**

Type species: *Musca scybalaria* LINNAEUS, 1758: 591

Comment: The genus *Scatomyza* FALLÉN, 1810 was revalidated by OZEROV & KRIVOSHEINA (2011a: 3). It differ from *Scathophaga* MEIGEN, 1803 by the presence of a metacoxal bridge (cf. VOCKEROTH 2010: 1268, Fig. 7).

***Scatomyza amplipennis* (PORTSCHINSKIY, 1887)**

Scatophaga amplipennis (PORTSCHINSKIY, 1887): 199, OZEROV & KRIVOSHEINA (2011a: 4).

***Scatomyza magnipennis* (PORTSCHINSKIY, 1887)**

Scatophaga magnipennis (PORTSCHINSKIY, 1887): 198, OZEROV & KRIVOSHEINA (2011a: 4).

Lectotype: male, Kyrgyzstan, loc. Sankul (= Sankul Lake), designated by OZEROV & KRIVOSHEINA (2013: 4)

***Scatomyza medvedevi* OZEROV & KRIVOSHEINA, 2018a**

Holo type: male, Nepal, Rasuwa Disom, Buddha Mandir env., 28.08.79°N 85.39.00°E, 4215 m, 13.-14.iii.2017, coll. A. Medvedev (ZMUM).

***Scatomyza mellipes* (COQUILLETT, 1899)**

Scatophaga mellipes (COQUILLETT, 1899): 335, OZEROV & KRIVOSHEINA (2011a: 8).

Scatophaga chinensis MALLOCH, 1935: 260, OZEROV & KRIVOSHEINA (2011a: 8).

Scathophaga eoa OZEROV, 2007: 2, OZEROV & KRIVOSHEINA (2011a: 8).

***Scatomyza nigrolineata* OZEROV & KRIVOSHEINA, 2018a**

Holo type: male, Nepal, Rasuwa disom, Buddha Mandir env., 28.08.79°N 85.39.00°E, 4215 m, 13.-14.iii.2017, coll. A. Medvedev (ZMUM).

***Scatomyza scybalaria* (LINNAEUS, 1758): 599**

Musca scybalaria (LINNAEUS, 1758): 599 OZEROV & KRIVOSHEINA (2011a: 8).

***Scatomyza sinensis* (SUN, 1996)**

Scathophaga sinensis (SUN, 1996): 630, OZEROV & KRIVOSHEINA (2018a: 11).

Scathophaga calceata OZEROV, 2009: 425, OZEROV & KRIVOSHEINA (2018a: 11).

***Scathophaga analis* (MEIGEN, 1826)**

Cordylura analis (MEIGEN, 1826): 251

Scathophaga bohemiae ŠIFNER, 2000: 193, ŠIFNER (2018: 1644).

***Scathophaga annae* ŠIFNER, 2019**

H o l o t y p e : male, Norway, Svalbard, Spitzbergen Island, locality Pyramiden, 78°39'22"N 16°19'30"E, 15.vii.2015, Anna Mácová leg. (FSPC).

***Scathophaga incompleta* OZEROV & KRIVOSHEINA, 2020a**

H o l o t y p e : male, Russia, Kalmykia, Priutnoe env. (46.1°N 43.5°E), 2.-3.v. 2013, N. Vikhrev (ZMUM).

***Scathophaga intermedia* (WALKER, 1849)**

Scathophaga hadleyi OZEROV 2013: 88, OZEROV (2019a: 330).

Scathophaga buryatica OZEROV & KRIISHEINA, 2019c

H o l o t y p e : male, Russia, East Sayan, Samarta (52:0944°N, 101:1372°E), 40km NO of Mondy, 1.viii.1965, Gorodkov (ZMUM).

***Scathophaga karellica* OZEROV, 2013**

H o l o t y p e : male, Russia, Karelia, Poyakonda, 66.589220°N 32.828671°E, 30.vi.2010, A. L. Ozerov leg. (ZMUM).

***Scathophaga karmazini* ŠIFNER, 2020**

H o l o t y p e : male, Austria, Hohe Tauern, Stubnerkogel Mt., 2240 m a.s.l., 27.vii.2001, Šifner leg. (FSPC).

***Scathophaga merdaria* (FABRICIUS, 1794)**

Musca merdaria (FABRICIUS, 1794): 344

C o m m e n t : This species was confirmed as valid species and distinguished from *Scathophaga stercoraria* (LINNAEUS, 1758) based on the differences of males and females terminalia by ŠIFNER (2020: 736, Figs 37-40).

***Scathophaga moraviensis* ŠIFNER, 2011**

H o l o t y p e : male, Czech Republic, Moravia, Bílé Karpaty Mts., Malá Vrbka (7170), Čertoryje, 48.859°N 17.431°E, 360 m, Malaise trap, 25. v.-15.vi.2910, Macek & Chvojka leg. (FSPC).

***Scathophaga simaceki* ŠIFNER, 2018**

H o l o t y p e : male, Norway, Svalbard, Spitzbergen Island, locality Pyramiden, 78°39'22"N 16°19'30"E, 15.vii.2015, Anna Mácová leg. (FSPC).

***Scathophaga stercoraria* (LINNAEUS, 1758)**

Musca stercoraria (LINNAEUS, 1758)

Chione ichneumonea ROBINEAU-DESOVIDY, 1830, OZEROV (2017a: 192).

C o m m e n t : *Chione ichneumonea* was mentioned by ŠIFNER (2008a: 177) as *nomen dubium*.

***Scathophaga yakutica* OZEROV, 2017c**

H o l o t y p e : male, Russia, Yakutia, Aldan Plateau, 60 km SE of Tommot, right bank of Russkaya River, ca. 58.56°N 126.56°E, 15.vii.2007, A. A. Popov (ZMUM).

Table 1. A list of the world fauna of species of the family Scathophgidae to 2020.

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
D e l i n i n a e ROBINEAU - DESVOIDY, 1830					
<i>Americina</i> MALLOCH, 1923					
<i>caucasica</i> (OZEROV, 2009)					X
<i>hostae</i> (HERING 1955),					X
<i>media</i> (MEIGEN, 1826)					X
<i>sellata</i> (HACKMAN, 1956)					X
<i>vittata</i> (MEIGEN, 1826)					X
<i>Delina</i> ROBINEAU-DESOVIDY, 1830					
<i>nigrita</i> (FALLÉN, 1819)		X			X
<i>Leptopa</i> ZETTERSTEDT, 1838					
<i>filiformis</i> ZETTERSTEST, 1838					X
<i>hendeli</i> ŠIFNER, 1977		X			
<i>Micripselapha</i> BECKER, 1894					
<i>basovi</i> OZEROV, 2010					X
<i>filiformis</i> (ZETTERSTEDT, 1846)					X
<i>Mirekiana</i> ŠIFNER, 2012					
<i>anthrax</i> ŠIFNER, 2012					X
<i>Neochirosia</i> MALLOCH, 1917					
<i>atrifrons</i> (COQUILLETT, 1910)	X				
<i>nigriceps</i> (BECKER, 1894)					X

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>nikita</i> OZEROV & KRIVOSHEINA, 2015					X
<i>nuda</i> MALLOCH 1922	X				
<i>pechorica</i> OZEROV, 2009					X
<i>setiger</i> MALLOCH, 1917	X				
<i>veratri</i> (HENDEL, 1925)					X
S c a t h o p h a g i n a e					
ROBINEAU-DESOVIDY, 1830					
<i>Acanthocnema</i> BECKER, 1894					
<i>albibarba</i> (LOEW, 1869)		X			
<i>capillata</i> (LOEW, 1872)		X			
<i>glaucescens</i> (LOEW, 1864)		X			X
<i>himalaica</i> SUWA, 1986					X
<i>longispina</i> SUWA, 1986					X
<i>nigrimana</i> (ZETTERSTEDT, 1846)					X
<i>ruficauda</i> CURRAN, 1929		X			
<i>sternalis</i> SUWA, 1986					X
Acerocnema BECKER, 1894					
<i>barkalovii</i> OZEROV, 2006					X
<i>lobanovi</i> OZEROV, 2006					X
<i>macrocera</i> (MEIGEN, 1826)					X
<i>paradoxopyga</i> STACKELBERG, 1952					X
Allomyella MALLOCH, 1923					
<i>albibennis</i> (ZETTERSTEDT, 1838)		X			X
<i>borealis</i> CURRAN, 1927		X			
<i>brevipennis</i> MALLOCH, 1923		X			
<i>crinipes</i> (RINGDAGHL, 1928)		X			X
<i>frigida</i> (HOLMGREN, 1883)		X			X
<i>portenkoi</i> (STACKELBERG, 1952)		X			X
<i>unguiculata</i> (MALLOCH, 1919)		X			
Amaurosoma BECKER, 1894					
<i>amurensis</i> OZEROV, 2010)					X
<i>armillatum</i> (ZETTERSTEDT, 1846)		X			X
<i>articulatum</i> BECKER, 1894					X
<i>atripes</i> MALLOCH, 1931		X			
<i>bernasconi</i> ŠIFNER, 2008					X
<i>bispinosa</i> MALLOCH, 1920		X			
<i>brevifrons</i> (ZETTERSTEDT, 1838)					X

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>bruneicosta</i> JOHNSON, 1927		X			X
<i>cryophila</i> (OZEROV & KRIVOSHEINA, 2015)					X
<i>fasciatum</i> (MEIGEN, 1826)					X
<i>flavipes</i> (FALLÉN, 1819)					X
<i>inerme</i> BECKER, 1894					X
<i>katmaiensis</i> MALLOCH, 1920		X			
<i>leucostoma</i> (ZETTERSTEDT, 1846)					X
<i>loewi</i> (BECKER, 1894)		X			X
<i>longicorne</i> (VON ROSER, 1840)					X
<i>nigrifrontatum</i> BECKER, 1894					
<i>obenbergeri</i> ŠIFNER, 2016					X
<i>pallidipes</i> MALLOCH, 1922		X			
<i>puberulum</i> BECKER, 1894					X
<i>rossolimoae</i> (OZEROV, 2010)					X
<i>similis</i> (COQUILLET, 1902)		X			
<i>tibiellum</i> (ZETTERSTEDT, 1838)					X
<i>truncatum</i> (FAN, 1976)					X
<i>unispinosa</i> MALLOCH, 1920		X			
<i>Bostrichopyga</i> BECKER, 1894					
<i>crassipes</i> (ZETTERSTEDT, 1838)					X
<i>Brockssiella</i> VOCKEROTH, 1955					
<i>varicornis</i> (CURRAN, 1927)		X			
<i>Ceratinostoma</i> MEADE, 1885					
<i>ostiorum</i> (CURTIS, 1832)		X			X
<i>Chaetosa</i> COQUILLET, 1898					
<i>churchilli</i> MALLOCH, 1931		X			
<i>punctipes</i> (MEIGEN, 1826)		X			X
<i>Cleigastra</i> MACQUART, 1835					
<i>apicalis</i> (MEIGEN, 1826)					X
<i>intermedia</i> OZEROV & KRIVOSHEINA, 2016					X
<i>maritima</i> OZEROV & KRIVOSHEINA, 2016					X
<i>sundukovi</i> OZEROV, 2013					X
<i>Coniosternum</i> BECKER, 1894					

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>alashanicum</i> (OZEROV & KRIVOSHEINA, 2019)					X
<i>decipiens</i> (HALIDAY, 1832)					X
<i>dvoraki</i> ŠIFNER, 2016					X
<i>fluviale</i> RONDANI, 1867					X
<i>kovari</i> ŠIFNER, 2013					X
<i>infumatum</i> BECKER, 1907					X
<i>jezeki</i> ŠIFNER, 1981					X
<i>kaszabi</i> ŠIFNER, 1975					X
<i>lapponicum</i> RINGDAHL, 1920	X				X
<i>masneri</i> ŠIFNER, 2017	X				
<i>mihalyi</i> ŠIFNER, 1975					X
<i>milani</i> ŠIFNER, 1981					X
<i>moceki</i> ŠIFNER, 2004					X
<i>molle</i> (BECKER, 1894)					X
<i>nelsoni</i> ŠIFNER, 2003					X
<i>nigripalpe</i> (BECKER, 1907)					X
<i>nigrohirtum</i> CZERNY, 1909					X
<i>obscurum</i> (FALLÉN, 1819)	X				X
<i>tinctinerve</i> BECKER, 1894					X
Cordilura FALLÉN, 1810					
<i>aberrans</i> BECKER, 1894					X
<i>adrogans</i> CRESSON, 1918	X				
<i>aemula</i> COLLIN, 1958					X
<i>albertha</i> (CURRAN, 1927)	X				
<i>albicoxa</i> JAMES, 1955	X				
<i>albilabris</i> (FABRICIUS, 1805)					X
<i>angustifrons</i> LOEW, 1863	X				
<i>amurensis</i> OZEROV, 2007					X
<i>atrata</i> ZETTERSTEDT, 1846	X				X
<i>atripennis</i> JAMES, 1955	X				
<i>banksi</i> JAMES, 1955	X				
<i>bezzi</i> (SACK, 1937)					X
<i>bicoloripes</i> OZEROV, 1997					X
<i>ciliata</i> (MEIGEN, 1826)					X
<i>ciliatipes</i> JAMES, 1955	X				
<i>confusa</i> LOEW, 1863	X				
<i>criddlei</i> CURRAN, 1929	X				
<i>cuspidata</i> SASAKAWA, 1986					X
<i>dimidiata</i> (CRESSON, 1918)	X				
<i>deceptiva</i> MALLOCH, 1923	X				

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>emarginata</i> (MALLOCH, 1923)		X			
<i>fasciventris</i> CURRAN, 1927					X
<i>flavovenosa</i> BECKER, 1894		X			X
<i>femoralis</i> SUN, 1993					X
<i>fulvifrons</i> OZEROV, 1997					X
<i>gagatina</i> LOEW, 1969		X			
<i>glabra</i> LOEW, 1869		X			
<i>grunini</i> OZEROV & KRIVOSHEINA, 2014					X
<i>gracilipes</i> LOEW, 1869		X			
<i>impudica</i> RONDANI, 1866					X
<i>intermedia</i> (CURRAN, 1927)		X			
<i>kakaberans</i> OZEROV, 1997					X
<i>kosterini</i> OZEROV & KRIVOSHEINA, 2014					X
<i>krocha</i> OZEROV, 2007					X
<i>latifrons</i> LOEW, 1869		X			
<i>loewi</i> JAMES, 1955		X			
<i>luteola</i> MALLOCH, 1924		X			
<i>marginata</i> (MALLOCH, 1931)		X			
<i>mundula</i> LOEW, 1869		X			
<i>monochroma</i> OZEROV & KRIVOSHEINA, 2014					X
<i>negrobovi</i> OZEROV & KRIVOSHEINA, 2017					X
<i>nigrifrons</i> SUN, 1993					X
<i>nigriseta</i> RONDANI, 1867					X
<i>nubecula</i> SASAKAWA, 1986					X
<i>ontario</i> CURRAN, 1929		X			
<i>passiva</i> CURRAN, 1929		X			
<i>picipes</i> MEIGEN, 1826					X
<i>picticornis</i> Loew, 1864		X			X
<i>pilosella</i> (COQUILLET, 1898)		X			
<i>pleuritica</i> LOEW, 1863		X			
<i>praeusta</i> (LOEW, 1864)		X			
<i>proboscoidea</i> ZETTERSTEDT, 1838		X			X
<i>pudica</i> MEIGEN, 1826		X			X
<i>pulchra</i> OZEROV & KRIVOSHEINA, 2012				X	
<i>remota</i> OZEROV, 1997					X
<i>rufimana</i> MEIGEN, 1826		X			X
<i>rufipes</i> MEIGEN, 1826		X			X
<i>sagittifera</i> GORODKOV, 1974					X

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>scapularis</i> LOEW, 1869		X			
<i>setosa</i> LOEW, 1860		X			
<i>shataikini</i> OZEROV, 1997					X
<i>sidorenkovi</i> OZEROV & KRIVOSHEINA, 2012					X
<i>sibirica</i> GORODKOV, 1974					X
<i>sifneri</i> OZEROV, 2007					X
<i>similis</i> SIEBKЕ, 1873					X
<i>socialis</i> BECKER, 1894					X
<i>tartariana</i> OZEROV, 2007					X
<i>tarsalis</i> (MALLOCH, 1923)		X			
<i>umbrosa</i> LOEW, 1873					X
<i>varitibia</i> LOEW, 1876		X			
<i>varicornis</i> CURRAN, 1929		X			
<i>zaitzevi</i> GORODKOV, 1974					X
<i>Cordylurella</i> MALLOCH, 1919					
<i>nana</i> (LOEW, 1864)		X			
<i>nebulosa</i> (COQUILLETT, 1898)		X			
<i>rufula</i> CURRAN, 1927		X			
<i>Cosmetopus</i> BECKER, 1894					
<i>dentimanus</i> (ZETTERSTEDT, 1838)					X
<i>longus</i> (WALKER, 1849)		X			
<i>ringdahli</i> ANDERSON, 1974					X
<i>Dromogaster</i> VOCKEROTH, 1995					
<i>incompta</i> VOCKEROTH, 1995		X			
<i>Ernoneura</i> BECKER, 1894					
<i>argus</i> (ZETTERSTEDT, (1818))		X			X
<i>Gabreta</i> ŠIFNER, 2015					
<i>macai</i> ŠIFNER, 2015					X
<i>Gimnomera</i> RONDANI, 1867					
<i>alanica</i> OZEROV, 1999					X
<i>albipila</i> (ZETTERSTEDT, 1846)					X
<i>alpina</i> ŠIFNER, 2003					X
<i>amica</i> (OZEROV, 1999)					X
<i>castanipes</i> (BECKER, 1894)		X			X
<i>cerea</i> (COQUILLETT, 1908)		X			

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>cuneiventris</i> (ZETTERSTEDT, 1846)		X			
<i>dorsata</i> (ZETTERSTEDT, 1838)					X
<i>fasciventris</i> MALLOCH, 1920		X			
<i>hirta</i> HENDEL, 1930					X
<i>incisurata</i> MALLOCH, 1920		X			
<i>kirkizica</i> OZEROV, 2008					X
<i>lasiostoma</i> (BECKER, 1894)					X
<i>novgorodovae</i> (OZEROV, 2006)					X
<i>slovaca</i> ŠIFNER, 2003					X
<i>speciosa</i> (OZEROV, 1999)					X
<i>subvittata</i> (MALLOCH, 1919)		X			X
<i>tarsea</i> (FALLÉN, 1819)					X
<i>tatrica</i> ŠIFNER, 2003					X
<i>tukuringa</i> (OZEROV, 1999)					X
<i>ziegleri</i> (OZEROV, 1999)					X
<i>Hajekiana</i> ŠIFNER, 2015					
<i>orlicensis</i> ŠIFNER, 2015					X
<i>Hexmitocera</i> BECKER 1894					
<i>loxocerata</i> (FALLÉN, 1826)					X
<i>martineki</i> ŠIFNER, 2003					X
<i>vockerothi</i> ŠIFNER, 2004					X
<i>Huckettia</i> VOCKEROTH, 1995					
<i>nearctica</i> VOCKEROTH, 1995		X			
<i>Hydromyza</i> FALLÉN, 1823					
<i>confluens</i> LOEW, 1863		X			
<i>livens</i> (FABRICIUS, 1794)					X
<i>Jezekia</i> ŠIFNER, 2009					
<i>kmenti</i> ŠIFNER, 2009					X
<i>Julienomyia</i> ŠIFNER, 2015					
<i>miroslavi</i> ŠIFNER, 2015					X
<i>Lasioscelus</i> BECKER, 1894					
<i>immundus</i> (ZETTERSTEDT, 1838)		X			X
<i>nigricans</i> (LOEW, 1873)					X
<i>sahlbergi</i> BECKER, 1900		X			X

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>Lubomiyia</i> ŠIFNER, 2010					
<i>orientalis</i> ŠIFNER, 2010				X	
<i>Megaphthalma</i> BECKER, 1894					
<i>palida palida</i> (FALLÉN, 1819)					X
<i>palida americana</i> MALLOCH, 1924		X			
<i>Megaphthalmoides</i> RINGDAHL, 1936					
<i>unilineatus</i> (ZETTERSTEDT, 1838)		X			X
<i>japonicus</i> OZEROV, 2008					X
<i>Microprosopa</i> BECKER, 1894					
<i>diversipes</i> CURRAN, 1927		X			
<i>frontata</i> (ZETTERSTEDT, 1838)					X
<i>gruberi</i> ŠIFNER, 2013					X
<i>haemorrhoidalis</i> (MEIGEN, 1826)					X
<i>heteromyzina</i> (ZETTERSTEDT, 1938)					X
<i>lacteipennis</i> RINGDAHL, 1920					X
<i>lineata</i> (ZETTERSTEDT, 1938)		X			X
<i>pallidicauda</i> (ZETTERSTEDT, 1838)		X			X
<i>paveli</i> OZEROV & KRIVOSHEINA, 2013)					X
<i>preisleri</i> ŠIFNER, 2013					X
<i>taymirica</i> OZEROV, 2017					X
<i>varicornis</i> CURRAN, 1927		X			X
<i>zemani</i> ŠIFNER, 2013					X
<i>zlobini</i> OZEROV, 2008					X
<i>Milania</i> ŠIFNER, 2010					
<i>agrion</i> (SÉGUY, 1948)					X
<i>longiabdomina</i> (SUN, 1992)					X
<i>Miroslava</i> ŠIFNER, 2016					
<i>knaifli</i> ŠIFNER, 2016					X
<i>jitkae</i> ŠIFNER, 1999					X
<i>montana</i> ŠIFNER, 1999					X
<i>Neorthochaeta</i> VOCKEROTH, 1995					
<i>dissimilis</i> (MALLOCH, 1924)		X			
<i>Norellia</i> ROBINEAU - DESVOIDY, 1830					

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>spinipes</i> (MEIGEN, 1826)					X
<i>tipularia</i> (FABRICIUS, 1794)					X
<i>Norellisoma</i> WAHLGREN, 1917					
<i>alpestre</i> (SCHINER, 1864)					X
<i>armipes</i> (MEIGEN, 1826)					X
<i>caucasicum</i> (OZEROV, 1993)					X
<i>femorale</i> (LOEW, 1864)					X
<i>flavicornе</i> (MEIGEN, 1826)					X
<i>flavostriatum</i> OZEROV, 2008					X
<i>insulare</i> (OZEROV, 1993)					X
<i>ivanae</i> ŠIFNER, 2003					X
<i>japonicum</i> HIRONAGA & SUWA 2005					X
<i>jelineki</i> ŠIFNER, 2006					X
<i>lesgiae</i> (BECKER, 1894)					X
<i>lituratum</i> (MEIGEN, 1826)					X
<i>mireki</i> ŠIFNER, 1977					X
<i>mirusae</i> ŠIFNER, 1974					X
<i>montanopratense</i> (OZEROV, 1993)					X
<i>nervosum</i> (MEIGEN, 1826)					X
<i>orientale</i> (OZEROV, 1993)					X
<i>seguyi</i> ŠIFNER, 1973					X
<i>spinimanum</i> (FALLÉN, 1819)	X				X
<i>striolatum</i> (MEIGEN, 1826)					X
<i>sylviae</i> ŠIFNER, 1999					X
<i>triangulum</i> (SUN, 1992)					X
<i>vockerothi</i> OZEROV, 2013					X
<i>vonickai</i> ŠIFNER, 2008					X
<i>yolduense</i> (OZEROV, 2008)					X
<i>Okeniella</i> HENDEL, 1907					
<i>caudata</i> (ZETTERSTEDT, 1838)					X
<i>dasyprocta</i> (LOEW, 1864)	X				
<i>gorodkovi</i> OZEROV, 2006					X
<i>stackelbergi</i> GORODKOV, 1967					X
<i>Orthacheta</i> BECKER, 1894					
<i>cornuta</i> (LOEW, 1863)	X				
<i>pilosa</i> ZETTERSTEDT, 1838)					X
<i>Paracosmetopus</i> HACKMAN, 1956					
<i>helleni</i> HACKMAN, 1956					X

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>Paramicroprosopa</i> RINGDAHL, 1936					
<i>bartaki</i> ŠIFNER, 1999					X
<i>frontata</i> (ZETTERSTEDT, 1838)					X
<i>hoberlandti</i> (ŠIFNER, 1981)					X
<i>ozerovi</i> (ŠIFNER, 2008)					X
<i>pokornyi</i> (ŠIFNER, 2011)					X
<i>Parallelomma</i> BECKER,					
<i>albipes</i> (FALLÉN, 1819)					X
<i>fuscipes</i> (ZETTERSTEDT, 1846)		X			X
<i>lauterereri</i> ŠIFNER, 2002					X
<i>merzi</i> OZEROV, 2009				X	
<i>Phrosia</i> ROBINEAU - DESVOIDY, 1830					
<i>remmi</i> (ELBERG, 1972)					X
<i>Peratomyia</i> VOCKEROTH, 1995					
<i>vittata</i> (COQUILLET, 1898)		X			
<i>Plethochaeta</i> COQUILLET, 1901					
<i>testacea</i> (MALLOCH, 1931)		X			
<i>varicolor</i> COQUILLET, 1901		X			
<i>Pleurochaetella</i> VOCKEROTH, 1965					
<i>barkalovi</i> OZEROV & KRIVOSHEINA, 2012					X
<i>simplicipes</i> (BECKER, 1900)		X			X
<i>Pogonota</i> ZETTERSTEDT, 1860					
<i>barbata</i> , ZETTERSTEDT, 1838		X			X
<i>gilvipes</i> (LOEW, 1863)		X			
<i>pallida</i> MALLOCH, 1931		X			
<i>Scatomyza</i> FALLÉN, 1810					
<i>amplipennis</i> PORTSCHINSKIY, 1887					X
<i>curtipilata</i> (FENG, 2002)				X	
<i>magnipennis</i> PORTSCHINSKIY, 1887					X
<i>medvedevi</i> OZEROV & KRIVOSHEINA, 2018					X
<i>mellipes</i> (COQUILLET, 1899)				X	X
<i>reses</i> (GIGLIO-TOS, 1893)			X		
<i>scybalaria</i> (LINNAEUS, 1758)					X

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>nigrolineata</i> OZEROV & KRIVOSHEINA, 2018					X
<i>sinensis</i> (SUN, 1996)					X
<i>Scatogera</i> ALBUQUERQUE, 1984					
<i>primogenita</i> ALBUQUERQUE, 1984			X		
<i>Scathophaga</i> MEIGEN, 1803					
<i>alata</i> (Becker, 1894)	X				
<i>albidohirta</i> (BECKER, 1907)					X
<i>aldrichi</i> (MALLOCH, 1920)		X			
<i>analis</i> (MEIGEN, 1826)					X
<i>annae</i> ŠIFNER, 2018					X
<i>apicalis</i> CURTIS, 1935		X			X
<i>calida</i> (HALIDAY, 1832)					X
<i>chilensis</i> (ALBUQUERQUE, 1953)			X		
<i>cineraria</i> (MEIGEN, 1826)					X
<i>cordylurina</i> (HOLMGREN, 1883)					X
<i>crinita</i> (COQUILLET, 1901)		X			X
<i>dasythrix</i> (BECKER, 1894)					X
<i>exalata</i> OZEROV, 1996					X
<i>foetulecta</i> (SÉGUY, 1966)	X				
<i>frigida</i> COQUILLET, 1900		X			
<i>furcata</i> (SAY, 1823)		X	X		X
<i>gigantea</i> (ALDRICH, 1923)					
<i>grisea</i> MALLOCH, 1920		X			
<i>hiemalis</i> (JAMES, 1950)		X			
<i>incola</i> BECKER, 1900		X			X
<i>inquinata</i> (MEIGEN, 1826)					X
<i>intermedia</i> WALKER, 1849		X			
<i>islandica</i> BECKER, 1894		X			X
<i>jizerensis</i> ŠIFNER, 2004					X
<i>karellica</i> OZEROV, 2013					X
<i>karmazini</i> ŠIFNER, 2020					X
<i>litorea</i> (FALLÉN, 1819)		X			X
<i>longiaculeata</i> OzErov, 2009	X				
<i>lutaria</i> (FABRICIUS, 1794)					X
<i>merdaria</i> (FABRICUS, 1794)					X
<i>monticola</i> MALLOCH, 1924		X			
<i>moraviensis</i> ŠIFNER, 2011					X
<i>multisetosa</i> (HOLMGREN, 1883)					X
<i>nigrolimbata</i> CRESSON, 1918		X			

	Afrotropical Region	Nearctic Region	Neotropical Region	Oriental Region	Palaearctic Region
<i>obscurinervis</i> (BECKER, 1900)		X			X
<i>odontosternita</i> FENG, 1999					X
<i>orcasae</i> , MALLOCH, 1935		X			
<i>peruensis</i> (LOPEZ & COURI, 1986)			X		
<i>pictipennis</i> (OLDENBERG, 1923)		X			X
<i>simaceki</i> ŠIFNER, 2018					X
<i>socia</i> BECKER, 1894	X				
<i>soror</i> WIEDEMAN, 1818	X		X		X
<i>staryi</i> ŠIFNER, 2000					X
<i>stercaria</i> (LINNAEUS, 1758)		X	X	X	X
<i>suilla</i> (FABRICIUS, 1794)		X			X
<i>taeniopa</i> (RONDANI, 1867)					X
<i>tropicalis</i> (MALLOCH, 1931)			X		
<i>varipes</i> (HOLMGREN, 1883)					X
<i>villosiventris</i> (RINGDAHL, 1937)					X
<i>vlastae</i> ŠIFNER, 2000					X
 <i>Scoliaphleps</i> BECKER, 1894					
<i>fulvifrons</i> (OZEROV, 1997)					X
<i>ustulata</i> (ZETTERSTEDT, 1838)		X			X
 <i>Spathophilus</i> BECKER, 1894					
<i>nigriventris</i> LOEW, 1864)					X
 <i>Spaziphora</i> RONDANI, 1856					
<i>cincta</i> (LOEW, 1863)		X			
<i>hydromyzina</i> (FALLÉN, 1819)					X
 <i>Staegeria</i> RONDANI, 1856					
<i>kunzei</i> (ZETTERSTEDT, 1821)					X
<i>unicornuta</i> DZIEDZICKI, 1887					X
 <i>Synchysa</i> VOCKEROTH, 1995					
<i>tricincta</i> (LOEW, 1869)		X			
 <i>Trichopalpus</i> RONDANI, 1856					
<i>fraternus</i> (MEIGEN, 1826)					X
<i>nigribasis</i> CURRAN, 1927		X			
<i>obscrellus</i> (ZETTERSTEDT, 1846)		X			X

Acknowledgement

We would like to thanks to Petr Kment (National Museum Praha, Czech Republic) and Thibault Lachat (Bern University of Applied Sciences, Zollikofen, Switzerland) for critical comments on the manuscript and Mirek ŠIFNER (Praha, Czech Republic) for his continuing large help.

Summary

In this paper, we summarise nomenclatorial changes in the family Scathophagidae during the last 30-40 years. Moreover, the list of species of the family Scathophagidae summarise the number of valid species in the word fauna by the end 2020.

(I) The world fauna of this family contains 362 valid species in 65 genera; the number of genera or subgenera depends on the criterias applied by the single authors. (II) The Afrotropical Region contains 5 species in a single genus. (III) The Nearctic Region contains 84 species in 35 genera. (IV) The Neotropical Region contains 9 species in two genera. (V) The Oriental Region contains 8 described species in 5 genera. (VI) The Palaearctic Region contains 256 species in 44 genera.

In the arctic to temperate zone of the Palaearctic and Nearctic regions, 39 species were found, which is only 10.2 % of all species occurring the Holarctic Region. The differences in the number of species among the individual regions depend on the level of faunistic and taxonomic research.

Two species, *Scatomyza janmayeni* Séguy, 1938 from Palaearctic Region and *Cordilura bissignata* (Walker, 1860) from Oriental Region, should be considered as doubtful species.

Zusammenfassung

Diese Arbeit beinhaltet grundlegende neue Erkenntnisse über die Forschung der Familie Scathophagidae in den letzten 30-40 Jahren und das Verzeichnis der Arten gibt die Übersicht der anerkannten Arten bis ins Jahr 2020. Über derzeitige taxonomische Kriterien und das Wissen über die zoogeografische Verbreitung der Arten dieser Familie kann man folgende Angaben geben:

(I) In der Weltfauna der Familie Scathophagidae wurden insgesamt 362 gültige Arten in 65 Gattungen festgestellt; die Anzahl der Gattungen oder Untergattungen sind von den angewendeten Kriterien der einzelnen Autoren abhängig. (II) Afrotropische Region: 5 Arten in einer Gattung. (III) Nearktische Region: 84 Arten in 35 Gattungen. (IV) Neotropische Region: 9 Arten in drei Gattungen. (V) Orientalische Region: 8 Arten in 5 Gattungen. (VI) Paläarktische Region: 256 Arten in 44 Gattungen.

In den arktischen und gemäßigten Zonen der Paläarktischen und Nearktischen Regionen wurden nur 39 Arten (10,2%) gemeinsam für beide Regionen festgestellt. Unterschiede in Zahl der Arten zwischen den Regionen sind von der faunistischen und taxonomischen Forschung abhängig.

Die zwei Arten, *Scatomyza janmayeni* SÉGUY, 1938 aus die Paläarktische Region und *Cordilura bissignata* (WALKER, 1860) aus der Orientalische Region sind als fraglich zu betrachten.

References

- ALBUQUERQUE D. & O. ALBUQUERQUE (1953): Contribuição ao conhecimento dos Scathophagidae Neotropicais, com descrição de uma espécie nova (Diptera: Scathophagidae). — Bolletino Museu Nacional, Rio de Janeiro, Zoologia **121**: 1-6.
- ALBUQUERQUE D. & O. ALBUQUERQUE (1984): 96 B. Family Scathophagidae. Pp. 96B1-96B4. — In: PAPAVERO N. (ed.): A catalogue of the Diptera of the Americas south of the United States. Departamento de Zoologia, Secretaria da Agricultura, São Paulo.

- BARTÁK M. & Š. KUBÍK (2012): A redefinition of *Micropselapha* Becker, 1894 (Diptera, Scathophagidae) with description of a new species from Central Europe. — Entomological News **123**: 101-109.
- BECKER T. (1914): Diptères nouveaux récoltés par Ch. Alluaud e R. Jeannel en Afrique oriental. — Annales de la Société Entomologique de France **83**: 120-130.
- BERNASCONI M.V., VALSANGIACOMO C., PIFFARETTI J.-C. & P.I. WARD (1999): The use of mitochondrial DNA sequences in insect taxonomy: examples from the Scathophagidae (Diptera, Calypratae). Bollettino della Società Ticinese di Scienze naturali **87**: 45-48.
- BERNASCONI M.V. (2000): Mitochondrial DNA sequences in insect phylogeny and taxonomy: Applications to the systematics of the Scathophagidae (Diptera, Calypratae), Ph.D. thesis, pp. 1-57. — Zoologisches Museum der Universität Zürich, Switzerland.
- BERNASCONI M.V., PAWLowski J., VALSANGIACOMO C., PIFFARETTI J.-C. & P.I. WARD (2000a): Phylogeny of the Scathophagidae (Diptera, Calypratae) based on Mitochondrial DNA sequences. — Molecular Phylogenetics and Evolution **16**: 308-315.
- BERNASCONI M.V., VALSANGIACOMO C., PIFFARETTI J.-C. & P.I. WARD 2000b: Phylogenetic relationships among Muscoidea (Diptera: Calypratae) base on mitochondrial DNA sequences. — Insect Molecular Biology **9**(1): 67-74.
- BERNASCONI M.V., PAWLISKI J., VALSANGIACOMO C., PIFFARETTI, J.-C. & P.I. WARD (2001): Phylogeny of the genus *Scathophaga* (Diptera: Scathophagidae) inferred from mitochondrial DNA sequences. — Canadian Journal of Zoology **79**: 517-524.
- BERNASCONI M.V., BERGER D. & W.U. BLANCKENHORN (2010): Systematic ambiguity in the well-established model system insect *Scathophaga stercoraria* (Diptera: Scathophagidae): sister species *S. soror* revealed by molecular evidence. — Zootaxa **2441**: 27-40.
- COLLIN J.E. (1931): The Oxford University Expedition to Greenland, 1923. Diptera Orthorrhapha Brachycera and Cyclorrhapha) from Greenland. — Annals and Magazine of Natural History, Series 10, **7**: 67-91.
- COLLIN J.E. (1958): A short synopsis of the British Scatophagidae (Diptera). — Transactions of the Society for British Entomology **13**: 37-56.
- CUNY R. (1983): Las Moscas Escatófagos (Scathophagidae, Diptera) de Latinoamérica. — Revista Peruana de Entomología **26**: 9-19.
- GIGLIO-TOS E. (1993): Dagnosi di nuovi generi e di nuove specie di Ditteri. IX. — Bollettino dei Musei di Biologia e Anatomia Comparata della Reale Università di Torino **8**: 1-14.
- GORODKOV K.B. (1986): Family Scathophagidae. Pp. 11-41. — In: SOÓS A. & PAPP L. (eds): Catalogue of Palaearctic Diptera. Vol. 7. Akadémiai Kiadó, Budapest, 346 pp.
- IWASA M. (2013): Three new species of the genus *Acerocnema* Becker (Diptera: Scathophagidae) from Japan, with a key to Palaearctic species. — Studia Dipterologica **20**: 175-183.
- JAMES M.T. (1950): The genus *Scopeuma* in western United States and southwestern Canada. — Annals of the Entomological Society of America **43**: 243-253.
- KUTTY S.N., BERNASCONI M.V., ŠIFNER F. & R. MEIER (2007): Sensitivity analysis, molecular systematic and natural history evolution of Scathophagidae (Diptera: Cyclorrhapha Calypratae). — Cladistics **23**: 64-83.
- LOPEZ S.M. & M. SOUTO CURI (1986): Description of new species of *Scopeuma* Meigen, 1800 (Diptera: Scatophagidae) from Peru. — Revista Brasiliera de Biología **46**: 313-316.
- LUNDBECK W. (1901): Diptera groenlandica. — Videnskabelige Meddelelser fra den Naturhistorisk Forening København, Serie 6, **2**: 281-316.
- MALLOCH J.R. (1931): Exotic Muscaridae (Diptera). — Annals and Magazin of Natural History, Series 10, **8**: 425-446.
- MALLOCH J.R. (1935): Exotic Muscaridae (Diptera). — Annals and Magazin of Natural History, Series 10, **15**: 242-266.

- OSTERBROEK P. (1998): The families of Diptera of the Maly Archipelage. — Fauna Malesiana Handbook. Vol. 1. Brill, Leiden, I-XII + 1-287 pp.
- OZEROV A.L. (2007): On Scathophagidae species (Insecta: Diptera), described by F. Hendel from the materials of the Swedish Kamchatka Expedition 1920-1922. — Far Eastern Entomologist **174**: 5-8.
- OZEROV A.L. (2008a): New species of Scathophagidae (Diptera). — Russian Entomological Journal **17**: 419-427.
- OZEROV A.L. (2008b): Two new species of the genus *Norellia* ROBINEAU-DESOVIDY, 1830 (Diptera, Scathophagidae). — Far Eastern Entomologist **184**: 1-4.
- OZEROV A.L. (2009a): Review of Afrotropical Scathophagidae (Diptera). — Russian Entomological Journal **18**: 299-308.
- OZEROV A.L. (2009b): On the Palaearctic fauna of *Norellisoma* WAHLGREN, 1917 (Diptera: Scathophagidae). — Russian Entomological Journal **18**: 229-234.
- OZEROV A.L. (2009c): A review of Palaearctic species of the genera *Delina* Robineau-Desvoidy, 1830 and *Neochirosia* Malloch, 1917 (Diptera, Scathophagidae). — Russian Entomological Journal **18**: 235-241.
- OZEROV A.L. (2009d): Flies of the genus *Paraleolmma* BECKER in STROBL, 1894 (Diptera, Scathophagidae) of Russia. — Russian Entomological Journal **18**: 309-317.
- OZEROV A.L. (2010a): A review of Palaearctic species of the genus *Nanna* STROBL, 1894 (Diptera, Scathophagidae) with shining stripes or spots on scutum. — Far Eastern Entomologist **214**: 1-8.
- OZEROV A.L. (2010b): Five new species of Scathophagidae (Diptera) from Russia. — Russian Entomological Journal **19**: 157-166.
- OZEROV A.L. (2011): A review of the World species of the genus *Spaziphora* RONDANI, 1856 (Diptera: Scathophagidae). — Russian Entomological Journal **20**: 423-430.
- OZEROV A.L. (2012): "Scathophaga helenae" (THOMSON, 1869), a New Synonym of *Scathophaga soror* (WIEDEMANN, 1818) (Diptera: Scathophagidae). — Far Eastern Entomologist **175**: 14-16.
- OZEROV A.L. (2013): Description of five new species and notes on taxonomy of Scathophagidae (Diptera). — Russian Entomological Journal **22**: 81-89.
- OZEROV A.L. (2014a): Notes on types of three species of Scathophagidae (Diptera) described by J. W. Zetterstedt. — Russian Entomological Journal **23**: 317-320.
- OZEROV A.L. (2014b): The primary types of Scathophagidae (Diptera) in the Museum für Naturkunde Berlin, and Senckenberg Deutschen Entomologisches Institut, Germany. — Zoosystematics and Evolution **90**: 33-43.
- OZEROV A.L. (2015): A new species of the genus *Acerocnema* BECKER, 1894 (Diptera: Scathophagidae) from Buryatia, Russia. — Caucasian Entomological Bulletin **11**: 211-213.
- OZEROV A.L. (2016): A review of the genus *Pogonota* ZETTERSTEDT, 1860 (Diptera: Scathophagidae) in Russia. — Russian Entomological Journal **25**: 185-207.
- OZEROV A.L. (2017a): The types of Scathophagidae (Diptera) in the Muséum national d'Histoire naturelle, Paris, France. — Russian Entomological Journal **26**: 189-194.
- OZEROV A.L. (2017b): A review of the genus *Microprosopa* BECKER, 1894 (Diptera: Scathophagidae) of Russia. — Russian Entomological Journal **25**: 71-99.
- OZEROV A.L. (2017c): A new species of the genus *Scathophaga* MEIGEN, 1803 (Diptera, Scathophagidae) from Yakutia (Russia). — Entomological Review **97** (1): 132-135.
- OZEROV A.L. (2019a): Morphological and nomenclatural notes on some Scathophagidae (Diptera). Russian Entomological Journal **28**(3): 328-330.
- OZEROV A.L. (2019b): A review of the genus *Gimnomera* RONDANI, 1866 (Diptera: Scathophagidae) of Russia. Russian Entomological Journal **28**(2): 195-218.

- OZEROV A.L. & A.V. BARKALOV (2014): On the fauna of Scathophagidae (Diptera) of Taimyr Peninsula (Russia: Krasnoyarskiy Kray). — Euroasian Entomological Journal **3**: 560-565.
- OZEROV, A. L. & FREIDBERG A. (2010): The Scathophagidae (Diptera) of Israel. — Israel Journal of Entomology **40**: 169-185.
- OZEROV A.L. & M.G. KRIVOSHEINA (2011a): To the knowledge of dung flies (Diptera, Scathophagidae) of Vietnam, with the description of a new species. — Far Eastern Entomologist **228**: 1-10.
- OZEROV A.L. & M.G. KRIVOSHEINA (2011b): A revision of dung flies of the genus *Mixocordylura* Hendel (Diptera: Scathophagidae), with the description of new species From Altai. — Russian Entomological Journal **20**: 431-435.
- OZEROV A.L. & M.G. KRIVOSHEINA (2011c): A revision of the types of two Scathophagidae (Diptera), described by Becker and deposited in the Zoological Institute, St.-Petersburg, Russia. — Russian Entomological Journal **20**: 105-108.
- OZEROV A.L. & M.G. KRIVOSHEINA (2012a): To the knowledge of the genus *Cordilura* FALLÉN (Diptera, Scathophagidae), with description of a new species from the Russian Far East. — Far East Entomologist **241**: 1-4.
- OZEROV A.L. & M.G. KRIVOSHEINA (2012b): A review of World species of the genus *Pleurochaetella* Vockeroth, 1965 (Diptera: Scathophagidae). — Russian Entomological Journal **21**: 435-440.
- OZEROV A.L. & M.G. KRIVOSHEINA (2013a): Types of Scathophagidae (Diptera) at the Zoological Institute of Russian Academy of Sciences, St. Petersburg. — Zoosystematica Rossica **22**: 141-150.
- OZEROV A.L. & M.G. KRIVOSHEINA (2013b): New and little known Palaearctic species of Scathophagidae (Diptera). — Far Eastern Entomologist **270**: 1-6.
- OZEROV A.L. & M.G. KRIVOSHEINA (2014a): The fauna of dung flies (Diptera: Scathophagidae) of Russian Far East. — Russian Entomological Journal **23**: 203-222.
- OZEROV A.L. & M.G. KRIVOSHEINA (2014b): A review of species of the genera *Bostrichopyga* BECKER, 1894 and *Paracosmetopus* HACKMAN, 1956 (Diptera: Scathophagidae). — Russian Entomological Journal **23**: 85-90.
- OZEROV A.L. & M.G. KRIVOSHEINA (2015a): A new species of the genus *Neochirosia* MALLOCH, 1917 (Diptera, Scathophagidae) from Serbia. — Russian Entomological Journal **24**: 145-147.
- OZEROV A.L. & M.G. KRIVOSHEINA (2015b): A review of the genera *Cleigastra* Macquart, *Gonarcticus* Becker, *Gonatherus* Rondani, *Hexamitocera* BECKER, *Nanna* STROBL, *Othacheta* BECKER and *Spathephilus* BECKER (Diptera, Scathophagidae) of Russia. — Zootaxa **4012** (2): 201-258.
- OZEROV A.L. & M.G. KRIVOSHEINA (2016): To taxonomy of the genus *Cleigastra* MACQUART, 1835 (Diptera: Scathophagidae) with description of two new species. — Russian Entomological Journal **25**: 97-102.
- OZEROV A.L. & M.G. KRIVOSHEINA (2017): Two new species of the genus *Cordilura* FALLÉN, 1810 (Diptera: Scathophagidae) from Buryatia and Primorskiy Krai, Russia. — Far Eastern Entomologist **335**: 1-6.
- OZEROV A.L. & M.G. KRIVOSHEINA (2018): To the knowledge of the genus *Scatomyza* FALLÉN, 1810 (Diptera: Scathophagidae) of Nepal. — Far Eastern Entomologist **353**: 1-16.
- OZEROV A.I. & M.G. KRIVOSHEINA (2019): Two new species of the genus *Scathophaga* MEIGEN, 1803 (Diptera: Scathophagidae) from Asia. — Russian Entomological Journal **28** (1): 102-106.
- OZEROV A.I. & M.G. KRIVOSHEINA (2020a): The taxonomy of the genus *Scathophaga* MEIGEN 1803 (Diptera: Scathophagidae) with description of a new species. — Russian Entomol. J. **29** (2): 227-231.

- OZEROV A.I. & M.G. KRIVOSHEINA (2020b): A review of the genus *Norellisoma* WAHLGREN, 1917 (Diptera: Scathophagidae) of Russia. Russian Entomol. J. **29**(1): 103-120.
- PAPP L., MERZ B. & FÖLDVÁRI M. (2006): Diptera of Thailand. — Acta Zoologica Academiae Scientiarum Hungaricae **52**: 97-269.
- PRUNER L. & P. MÍKA (1996): Seznam obcí a jejich částí v České republice s čísly mapových polí pro síťové mapování fauny. (List of settlements in the Czech Republic with associated map field codes for faunistic mapping system). — Klapalekiana **32** (Supplement): 1-175 (in Czech, English summary).
- PÜCHEL F. (1999): Scathophagidae. Pp. 186-187. — In: STARK A. & F. MENZEL (eds): Dipterologische Monographien. Studia Dipterologica, Supplement **2**: 1-354.
- SACK P. (1937): 62a. Cordyluridae. — In: LINDNER E. (ed.): Die Fliegen der Palaearktischen Region. Vol. 7. E. Schweizerbart'sche Verlagsbuchhandlung (Erwin Nägele), Stuttgart, 130 pp + 6 pls.
- STROBL G. (1898): Die Dipteren von Steiermark (IV. Theil). — Mitteilungen des Naturwissenschaftlichen Vereins für Steiermark **34**: 192-298.
- SZILADY Z. (1926): Dipterenstudien II. Bemerkungen über einige Scatophagiden des Ungarischen National Museum. — Annales Historico-Naturales Musei Nationalis Hungarici **24**: 593-597.
- ŠIFNER F. (1968): Description de la femelle de *Spathephilus breviventris* (LOEW) de la Tchécoslovaquie (Diptera, Scathophagidae). — Acta Entomologica Bohemoslovaca **65**: 467-469.
- ŠIFNER F. (1977): Les nouvelles indications taxonomique de la famille Scathophagidae (Diptera). — In: Diptera Bohemoslovaca II. Acta Universitatis Carolinae, Biologica **1977**: 397-402.
- ŠIFNER F. (2003): The family Scathophagidae (Diptera) of the Czech and Slovak Republics with notes on selected Palaearctic taxa. — Acta Musei Nationalis Pragae, Series B, Historia Naturalis **59**: 1-90.
- ŠIFNER F. (2008a): A catalogue of the Scathophagidae (Diptera), of the Palaearctic region, with notes on their taxonomy and faunistics. — Acta Entomologica Musei Nationalis Pragae **48**: 111-196.
- ŠIFNER F. (2008b): Three new species of the family Scathophagidae (Diptera) from the Palaearctic region with a redescription of the male of *Amaurosoma longicorne*. — Acta Entomologica Musei Nationalis Pragae **48**: 103-109.
- ŠIFNER F. (2009b): Addition and corrections to the catalogue of the Scathophagidae (Diptera) of the Palaearctic region. — Acta Entomologica Musei Nationalis Pragae **49**: 293-295.
- ŠIFNER F. (2009c): Two new genera of the family Scathophagidae (Diptera). — Acta Entomologica Musei Nationalis Pragae **49**: 287-292.
- ŠIFNER F. (2010): Two new genera and species of the family Scathophagidae (Diptera) from Palaearctic and Oriental region, with additional faunistic records. — Acta Entomologica Musei Nationalis Pragae **50**: 609-618.
- ŠIFNER F. (2011): Two new species of Scathophagidae (Insecta: Diptera) from the Czech Republic, with description of the males of *Scathophaga bohemiae* and *Norellisoma vonickai*. — Journal of the National Museum (Prague), Natural History Series **180**: 31-38.
- ŠIFNER F. (2012): *Mirekiana* gen.nov., a new genus of dung flies (Diptera: Scathophagidae) from the Czech Republic. — Journal of the National Museum (Prague), Natural History Series **181**: 15-19.
- ŠIFNER F. (2013a): Three new species of the family Scathophagidae (Diptera) from the Czech and Slovakia. — Journal of the National Museum (Prague), Natural History Series **182**: 35-50.
- ŠIFNER F. (2013b): *Coniosternum kovari* nov.sp. from Tajikistan, with the key to Palaearctic species of *Coniosternum* (Diptera: Scathophagidae). — Klapalekiana **49**: 205-218.

- ŠIFNER F. (2015): Two new genera of the family Scathophagidae (Diptera) from the Czech Republic. — Journal of the National Museum (Prague), Natural History Series **184**: 75-80.
- ŠIFNER F. (2016a): One new genus and two new species of the family Scathophagidae (Diptera) from the Czech Republic. — Journal of the National Museum (Prague), Natural History Series **185**: 1-6.
- ŠIFNER F. (2016b): *Amaurosoma obenbegeri* nov.sp., a new species of the family Scathophagidae (Insecta: Diptera) from the Czech Republic. — Journal of the National Museum (Prague), Natural History Series **185**: 21-24.
- ŠIFNER F. (2017): *Coniosternum masneri* sp. n., a new species of the family Scathophagidae from Canada. — Journal of the National Museum (Prague), Natural History Series **136**: 113-116.
- ŠIFNER F. (2018a): Annotated checklist of the family Scathophagidae (Diptera) in Central Europe, with new faunistic data on some species. — Linzer biol. Beitr. **50** (2): 1635-1655.
- ŠIFNER F. (2018b): Two new species of the genus *Scathophaga* (Diptera: Scathophagidae) from Svalbard, with a key to selected northern species and comments on this taxonomy. — Journal of the National Museum (Prague), Natural History Series **187** [2018]: 27-46.
- ŠIFNER F. (2020): *Scathophaga karmazini* nov.sp., a new species from Austria, with a key of species of the genus *Scathophaga* MEIGEN, 1803 from Central Europe (Diptera, Scathophagidae). — Linzer biol. Beitr. **52** (1): 731-757.
- VOCKEROTH J.R. (1965): Subfamily Scatophaginae. Pp. 826-842. — In: STONE A., SABROWSKY C.W., WIRTH W.W., FOOTE R.H. & J.R. COULSON (eds): A catalog of the Diptera of America North of Mexico. — Agricultural Handbook No. **267**. Agricultural Research Service, United States Department of Agriculture, Washington D.C., 1696 pp.
- VOCKEROTH J.R. (1976): La faune terrestre de l'Île de Saint-Hélène. Troisième partie. Musée Royal de l'Afrique Centrale, Tervuren, Belgique. — Annales de Sciences Zoologique **15**: 120-121.
- VOCKEROTH J.R. (1977): Superfamily Muscoidea. Family Scathophagidae. Pp. 436-438. — In: DELFINADO M.D. & HARDY D.E. (eds): A catalog of the Diptera of the Oriental Region. **Vol. 3**. The University Press of Hawaii, Honolulu.
- VOCKEROTH J.R. (1980): 82. Family Scathophagidae. Pp. 714. — In: CROSSKEY R.W. (ed.): Catalogue of the Diptera of the Afrotropical Region. British Museum, Natural History, London.
- VOCKEROTH J.R. (1987): Scathophagidae. Vol. 2. Pp. 1085-1097. — In: MCALPIN J.R. & al. (eds): Manual of Nearctic Diptera. Research Branch Agriculture Canada, Ottawa. Agricultural Canadian Monograph No. **28**, vi + 675 + 332 pp.
- VOCKEROTH J.R. (1995): Validation of nomina nuda of Nearctic Tethinidae, Scathophagidae, and Muscidae proposed in Manual of Nearctic Diptera. — Proceedings of the Entomological Society of Washington **97**: 732-734.
- VOCKEROTH J.R. (2010): Scathophagidae (Dung flies). Pp. 1267-1269. — In: BROWN R.V. & al. (eds): Manual of Central Americana Diptera Vol. 2. NAC Research Press, Ottawa, 728 pp.
- WALKER F. (1860): Catalogue of Dipterous Insects collected at Makessar in Celebes by A. R. Wallace with description of a new species. — Journal of the Proceedings of Linnean Society of London, Zoology **4**: 99-114.
- WERNER D., MANN D.J. & A.C. PONT (2006): Notes on predation by Scathophagidae (Diptera) on Simuliidae (Diptera). The Entomologist Monthly Magazine, **142**:143-150.
- WOLFF M., GRISALES D. & S.S. NIHEI (2016): Family Scathophagidae. — Zootaxa **4122** (1): 854-855.
- WULP F.M. VAN DER (1896): Catalogue of the described Diptera from South Asia. — Dutch Entomological Society, Hague, 211 pp.

WULP F.M. VAN DER (1897): Insecta. Diptera (part). Pp. 345-30. — In: GODMAN F.D. & O. SALVIN (eds): Biologia Centrali Americana or contributions to the knowledge of the fauna and flora of Mexico and Central Americana. Zoologia. Class Insecta. Order Diptera. Volumen 2, London, 489 pp.

Authors' addresses: Marco V. BERNASCONI
Natur-Museum Luzern
Kasernenplatz 6
CH-6063 Luzern, Switzerland

František ŠIFNER
Všíhláč 1311
CZ-14200 Praha 4, Czech Republic
E-mail: miroslav.sifner@email.cz