

Aphyllorchis yachangensis (Orchidaceae), a new holomycotrophic orchid from China

Ying Qin¹, Hailing Chen¹, Zhenhai Deng², Yan Liu¹

1 Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin, Guangxi, 541006, China **2** Management Center of Yachang Orchid National Nature Reserve, Baise, Guangxi, 533200, China

Corresponding author: Yan Liu (gxibly@163.com)

Academic editor: M. Simo-Droissart | Received 5 February 2021 | Accepted 4 June 2021 | Published 5 July 2021

Citation: Qin Y, Chen HL, Deng ZH, Liu Y (2021) *Aphyllorchis yachangensis* (Orchidaceae), a new holomycotrophic orchid from China. *PhytoKeys* 179: 91–97. <https://doi.org/10.3897/phytokeys.179.63994>

Abstract

Aphyllorchis yachangensis, a new holomycotrophic orchid from Guangxi, southern China is described and illustrated here. This new species is similar to *A. caudata* but differs from the latter mainly by the sepals acute at the apex, the hypochile with 2 smaller and semicircular wings, the epichile adaxially smooth, acute, the lateral lobes triangular-ovate and the column clavate.

Keywords

Aphyllorchis, Guangxi Province, new taxa, saprophytic Orchidaceae, taxonomy

Introduction

Aphyllorchis Blume (1825) includes about 30 species, and is mainly distributed in tropical Asia and the Himalayas, extending to the north of Japan and to the south of Australia (Chen and Stephan 2009; Tian et al. 2013). The species of *Aphyllorchis* are holomycotrophic herbs, with short rhizomes; fleshy roots; erect, unbranched stems, with sheaths; racemose inflorescences; petals similar to sepals, free; lip 2-partite, hypochile smaller than epichile, sometimes without hypochile; staminodes 2; pollinia 2, granular-farinaceous, caudicles absent (Chen and Stephan 2009). In China, six species and one variety are reported from the South, Southwest and Taiwan Island, including *Aphyllorchis alpina* King & Pantling, *A. caudata* Rolfe ex Downie, *A. gollanii* Duthie, *A. montana* H. G. Reichenbach var. *montana*, *A. montana* H. G. Reichenbach var.

membranacea T. C. Hsu, *A. pallida* Blume and *A. simplex* Tang & F. T. Wang (Chen and Stephan 2009; Fan et al. 2011; Hsieh et al. 2013; Huang et al. 2014; Hsu and Chung 2017; Lin 2017).

In late July 2018, during a botanic expedition in the Yachang Orchid National Nature Reserve, Guangxi Province in China, the first author collected a specimen (*Ying Qin* et al. YC3639) belonging to *Aphyllorchis*, with residual flowers. The narrow hypochile with 2 unobvious wings at base and purple epichile of the species drew the author's attention. However, due to the incomplete structure of sepals, petals and column, it was not accurately identified at that point. Fortunately, after two further botanic expeditions, one in July 2019 and the other in July 2020, we successfully collected two specimens (*Ying Qin* et al. QY20190719001 and *Ying Qin* et al. QYYC20200703009) belonging to the same taxon, but on these occasions with complete flowers. After dissecting and examining those flowers, and consulting the relevant literature (Reichenbach 1876; Downie 1925; Fukuyama 1934; Pearce and Cribb 1999; Roy et al. 2009; Aravindhan et al. 2013; Merckx et al. 2013; Tian et al. 2013; Campbell 2014; Sun et al. 2017; Suetsugu et al. 2018), we finally confirmed that it is a new species, which is described here.

Material and methods

From July 2018 to July 2020, we examined four specimens of *Aphyllorchis* in IBK on the field, and also examined forty specimens of *Aphyllorchis* in PE, HITBC, KUN, AU, IMDY, SZG, etc. through CVH (<https://www.cvh.ac.cn/index.php>). Except for *Ying Qin* et al. YC3639, *Ying Qin* et al. QY20190719001 and *Ying Qin* et al. QYYC20200703009 collected by the first author and kept in IBK, none of the specimens belong to *Aphyllorchis yachangensis*. Photographs of plants and flowers were taken using a Canon PowerShot G16. Morphological characters of the new species were measured with a ruler on living plants in the wild. The terminologies used to describe parts of the new species, such as rhizome, ovary, sepals, petals, lip, hypochile, epichile, column, etc. come from Flora of China (Chen and Stephan 2009).

Taxonomic Treatment

Aphyllorchis yachangensis Ying Qin & Yan Liu, sp. nov.

urn:lsid:ipni.org:names:77218063-1

Figures 1, 2

Diagnosis. *Aphyllorchis yachangensis* is similar to *A. caudata* but differs from the latter mainly by its hypochile with two wings of 1–1.4 × 0.4–0.6 mm (vs. 2–3 × ca. 4 mm), epichile adaxially smooth (vs. densely papillose), lateral lobes triangular-ovate (vs. semicircular), sepals acute at apex (vs. long cuspidate) and column clavate (vs.




Figure 1. *Aphyllorchis yachangensis* **A** flowering habit **B** inflorescence **C** bract **D** flower in side view **E** flower in front view **F** column in top view **G** column in ventral view **H** lip, dorsal sepal, lateral sepals, petals, column and ovary **I** column in side view **J** lip in side view **K** lip in top view **L** pollinaria **M** anther cap in top view **N** rhizome and roots. Photographed by Ying Qin from *Ying Qin et al. QY20190719001* (holotype: IBK!).

approximately cylindrical). Detailed morphological comparisons between *A. yachangensis* and *A. caudata* are provided in Table 1.

Type. CHINA. Guangxi, Baise city, Leye county, Yachang Orchid National Nature Reserve, 1862 m, 19 July 2019, *Ying Qin* et al. QY20190719001 (**holotype:** IBK! isotype: IBK!).

Description. Herb holomycotrophic, leafless, 107–113 cm tall. Rhizome ca. 10.8 cm long, 4.9–6.2 mm in diameter, densely noded; internodes 2.4–12 mm long. Roots spreading, stout, 3.4–6.4 mm in diameter. Stem 61–71 cm tall, 4.5–7.3 mm in diameter, yellow-green, usually with many dark purple stripes and spots, with 1.4–4 cm long sheaths; the uppermost sheath lanceolate; other sheaths tubular. Inflorescence terminal, racemose, 42–47 cm long, with 19–23 well-spaced flowers; rachis sparsely glandular puberulent, yellow-green, usually with many dark purple stripes and spots. Bracts lanceolate, shorter than pedicel and ovary, 1.2–3 cm long, 2.8–6.8 mm wide, yellow-green, usually with many dark purple stripes and spots on abaxial surface, sparsely glandular puberulent. Ovary cylindrical, yellow-green, usually with many dark purple stripes and spots, sparsely glandular puberulent, 2.2–2.8 mm in diameter, including pedicel to 2.3–3.5 cm long. Sepals yellow-green, usually with many dark purple stripes and spots on abaxial surface, abaxially sparsely glandular puberulent; dorsal sepal lanceolate, slightly concave, 2.5–4 cm long, 7.9–10.7 mm wide, apex acute, reflexed; lateral sepals lanceolate, slightly oblique at base, 2.5–4 cm long, 6.1–8.2 mm wide, apex acute, reflexed. Petals lanceolate, reflexed, 2.2–2.7 cm long, 5.6–7.2 mm wide, yellow-green, pale purple to dark purple at base, apex acute. Lip 2-partite; hypochile narrow, concave, 6.6–7.1 mm long, 3.6–4.5 mm wide, yellow or reddish brown, with 2 unobvious wings at base; wings semicircular, 1–1.4 mm wide at base, 0.4–0.6 mm tall; epichile ca. 1.9–2.2 cm long, 1.3–1.5 cm wide when flattened, 3-lobed, yellow with pale purple, or purple to dark purple, adaxially smooth, margin slightly erose; lateral lobes triangular-ovate, 8.2–8.9 mm long, 3.6–4.3 mm wide; mid-lobe narrowly triangular, 1.1–1.3 cm long, 5.6–5.9 mm wide, acute at apex, margin slightly involute. Column clavate, arcuate, 1.4–1.6 cm long, 1.8–3.3 mm in diameter in lower part, 4.3–6.1 mm in diameter in upper part, yellow-green, usually purple from middle to base; stigma ovate, slightly concave; staminodes 2, on both sides of apex, yellow-white to pale yellow-green; pollinia 2, ovate, granular-farinaceous, 4 mm long, yellow; anther cap ovate, slightly laterally compressed, 4 mm long, pale yellow.

Distribution, habitat and ecology. *Aphyllorchis yachangensis* was discovered in the Yachang Orchid National Nature Reserve, Leye county, Baise city, Guangxi Province, China. The holotype subpopulation is distributed in a subtropical evergreen and deciduous broad-leaved mixed forest, and is found growing with *Chimonobambusa quadrangularis* (Franceschi) Makino (Gramineae), *Lithocarpus glaber* (Thunberg) Nakai (Fagaceae), *Manglietia fordiana* Oliver (Magnoliaceae), *Juglans mandshurica* Maximowicz (Juglandaceae), *Dysosma majorensis* (Gagnep.) Ying (Berberidaceae), *Ophiopogon bockianus* Diels (Liliaceae), *Disporopsis fuscipicta* Hance (Liliaceae), *Calanthe brevicornu* Lindley (Orchidaceae), *Goodyera biflora* (Lindley) J. D. Hooker (Orchidaceae), *G. velutina* Maximowicz ex Regel (Orchidaceae), etc.




Figure 2. *Aphyllorchis yachangensis* (an individual with dark colors in the same environment with *Ying Qin* et al. QY20190719001) **A** flowering habit **B** bud **C** flower in back view **D** flower in front view. Photographed by Ying Qin from *Ying Qin* et al. QYYC20200703009 (paratype: IBK!).

Table I. Morphological comparison of *Aphyllorchis yachangensis* and *A. caudata*.

Characters	<i>A. yachangensis</i>	<i>A. caudata</i>
Plant size	107–113 cm	100 cm
Inflorescence	42–47 cm	rachis to 50 cm
Bracts	lanceolate, 12–30 × 2.8–6.8 mm	narrowly lanceolate, 40–48 × 5–6 mm
Dorsal sepal	lanceolate, 25–40 × 7.9–10.7 mm, apex acute	linear-lanceolate or lanceolate, 30–35 × ca. 8.0 mm, apex long cuspidate
Lateral sepals	lanceolate, 6.1–8.2 mm wide, apex acute	linear-lanceolate or lanceolate, 6–7 mm wide, apex long cuspidate
Petals	lanceolate, 22–27 × 5.6–7.2 mm	lanceolate, ca. 20 × 6–7 mm
Hypochile	6.6–7.1 mm long, with 2 wings; wings semicircular, 1–1.4 × 0.4–0.6 mm	2–3 mm long, with 2 wings; wings ligulate, 2–3 × ca. 4 mm
Epichile	1.9–2.2 cm long, adaxially smooth; lateral lobes triangular-ovate	ca. 1.2 cm long, adaxially densely papillose; lateral lobes semicircular
Column	1.4–1.6 cm long, clavate (apex obviously inflated)	1.1–1.4 cm long, approximately cylindrical
Flowering period	late June and July	July and August

Phenology. Flowering in late June and July, capsules not seen.

Etymology. The specific epithet is derived from the type locality, Yachang Orchid National Nature Reserve.

Chinese name. 雅长无叶兰 (Ya Chang Wu Ye Lan)

Additional specimens examined (paratypes). CHINA: Guangxi: Baise city, Leye county, Yachang Orchid National Nature Reserve, 1862 m, 28 July 2018, *Ying Qin* et al. YC3639 (IBK!); Guangxi: Baise city, Leye county, Yachang Orchid National Nature Reserve, 1859 m, 03 July 2020, *Ying Qin* et al. QYYC20200703009 (IBK!).

Key to *Aphyllorchis* taxa in China

- 1 Lip similar to lateral petals *A. simplex*
- Lip distinctly different from petals 2
- 2 Bracts longer than pedicel and ovary 3
- Bracts shorter than pedicel and ovary 4
- 3 Flowers yellowish green; bracts linear to linear-lanceolate, 3–4 mm wide; lip contracted at middle into hypochile and epichile *A. alpina*
- Flowers pale purplish brown; bracts ovate to elliptic-lanceolate, 6–8 mm wide; lip contracted near base into hypochile and epichile *A. gollanii*
- 4 Sepals equal to or longer than 25 mm 5
- Sepals equal to or shorter than 11 mm 6
- 5 Sepals apex long cuspidate; epichile adaxially densely papillose *A. caudata*
- Sepals apex acute; epichile adaxially smooth *A. yachangensis*
- 6 Sepals 4–5 mm long *A. pallida*
- Sepals 9–11 mm long 7
- 7 Lip fleshy *A. montana* var. *montana*
- Lip membranous *A. montana* var. *membranacea*

Acknowledgements

We thank Yu-Liang Jiang and Jin-Quan Huang from IBK for their valuable help with the botanic expeditions. This study was supported by Orchidaceae Resources Survey Project in Guangxi (2019073012, 2019073013) and The National Natural Science Foundation of China (41661012).

References

- Aravindhan V, Mahendran G, Rajendran A, Narmatha BN (2013) *Aphyllorchis montana* Rchb. f. (Orchidaceae), a new distributional record for Peninsular India. *Annals of Plant Sciences* 2: 153–155.

- Blume CL (1825) *Aphyllorchis*. Tabellen en platen voor de Javaansche orchideën, Ter Lands, Batavia, t. 16.
- Campbell F (2014) A summary of Holomycotrophic orchids. The MIOS Journal 15(4): 6–17.
- Chen SC, Stephan WG (2009) *Aphyllorchis*. In: Wu ZY, Raven PH, Hong DY (Eds) Flora of China, Vol. 25. Science Press, Beijing, 177–179.
- Downie DG (1925) Contributions to the flora of Siam. Additamentum XVII. Bulletin of Miscellaneous Information 1925: 404–423. <https://doi.org/10.2307/4115102>
- Fan J, Jin XH, Xiang XG (2011) *Aphyllorchis pallida*, a new record of orchidaceae from China. Zhiwu Kexue Xuebao 29: 647–648.
- Fukuyama N (1934) Studia orchidacearum Japonicarum III. Botanical Magazine Tokyo 48(571): 429–442. <https://doi.org/10.15281/jplantres1887.48.429>
- Hsieh S, Leou C, Yu S, Lee C, Yeh C (2013) *Aphyllorchis rotundatipetala* (Orchidaceae), a new species from Taiwan. Annales Botanici Fennici 50(3): 179–182. <https://doi.org/10.5735/085.050.0309>
- Hsu TC, Chung SW (2017) Illustrated Flora of Taiwan, Vol. 1. Owl Publishing House, Taipei, 295–296.
- Huang MZ, Liu ZL, Wang QL, Yang GS (2014) Two genus and eight species of orchidaceae, newly recorded in hainan. Redai Zuowu Xuebao 35(1): 138–141.
- Lin TP (2017) Newly discovered native orchids of Taiwan (IX). Taiwania 62(2): 205–210.
- Merckx VSFT, Freudenstein JV, Kissling J, Christenhusz MJM, Stotler RE, Crandall-Stotler B, Wickett N, Rudall PJ, Maas-van de Kamer H, Maas PJM (2013) Taxonomy and classification. In: Merckx VSFT (Ed.) Mycoheterotrophy: the biology of plants living on fungi. Springer, New York, 19–102. https://doi.org/10.1007/978-1-4614-5209-6_2
- Pearce N, Cribb P (1999) Notes relating to the flora of Bhutan: XXXVII. New species and records of Orchidaceae from Bhutan and India (Sikkim). Edinburgh Journal of Botany 56(2): 273–284. <https://doi.org/10.1017/S096042860000113X>
- Reichenbach HG (1876) Orchideae Roezlianae novae seu criticae. Linnaea 41: 1–98.
- Roy M, Watthana S, Stier A, Richard F, Vessabutr S, Selosse M (2009) Two mycoheterotrophic orchids from Thailand tropical dipterocarpacean forests associate with a broad diversity of ectomycorrhizal fungi. BMC Biology 7(1): e51. <https://doi.org/10.1186/1741-7007-7-51>
- Suetsugu K, Suleiman M, Anthony F, Tsukaya H (2018) *Aphyllorchis maliauensis* (Orchidaceae), a new species from the Maliau Basin, Sabah, Borneo. Phytotaxa 367(1): 85–90. <https://doi.org/10.11646/phytotaxa.367.1.10>
- Sun Y, Li B, Guo SX (2017) Research progress of saprophytic orchids. Guihaia 37(2): 191–203.
- Tian HZ, Li XL, Hu C (2013) A new synonym and lectotypification of *Aphyllorchis caudata* (Orchidaceae). Kew Bulletin 68(2): 341–344. <https://doi.org/10.1007/s12225-013-9451-3>