

Article

Checklist of oribatid mites (Acari: Oribatida) of the Central Black Sea basin of Turkey with new records for the country

Maka Murvanidze^{1*}, Cihan Cilbircioğlu², Esengül Özdemir³ and Emre Inak³

1. Institute of Entomology, Agricultural University of Georgia. 240, D. Aghmashenebeli Alley, 0131 Tbilisi, Georgia; E-mail: m.murvanidze@agruri.edu.ge

2. Department of Organic Agriculture, Taşköprü Vocational School, Kastamonu University, Kastamonu, Turkey; E-mail: cihancilbirci@hotmail.com

3. Department of Plant Protection, Faculty of Agriculture, Ankara University, Ankara, Turkey; E-mail: esengul.ozdemir@ankara.edu.tr

* Corresponding author

ABSTRACT

Fifty two species/subspecies of oribatid mites are identified from forest areas in the central Black Sea basin and Central Anatolia. Seventeen species among them are new records for the country. For each species GPS coordinates, habitat, regional and global distribution data are provided and for new records additional remarks with taxonomic, ecological and/or regional peculiarities are given.

KEY WORDS: Biodiversity; forest; new records; Turkish mite fauna; zoogeography.

PAPER INFO.: Received: 21 May 2020, Accepted: 20 June 2020, Published: 15 July 2020

INTRODUCTION

Turkish Black Sea basin is stretched from west to east along the southern part of the Black Sea. Mountainous landscape from the east and the Black Sea from the north, create a transitional climatic profile with precipitation between 500 and 1000 mm; rich flora with intact forests are characterized by dominance of woody species such as *Fagus orientalis* Lipsky, *Quercus* sp., *Carpinus betulus* L., *Pinus nigra* Arn. (Zeydanli 2020). Starting from north eastern Black Sea coast (which is a part of tertiary Colchis refugia) towards the east, climate gets drier supporting a wealth of habitat varieties and vegetation types (Williams *et al.* 2006). This landscape and habitat diversity is also reflected in soil animal communities which are also very high for any well studied animal taxa (Schütt 2005; Murvanidze and Mumladze 2016; Murvanidze *et al.* 2016). While oribatid fauna of the rest of Turkey is relatively poorly studied, from the Black Sea area more than 120 species are known (Ayyildiz 1988; Ayyildiz and Luxton 1989a, b; Dik *et al.* 1995, 1999; Baran and Ayyildiz 2000; Ayyildiz *et al.* 2011a; Toluk and Ayyildiz 2011; Yalçın *et al.* 2013; Akman *et al.* 2018; Baran and Merve 2019); however, given the diversity of landscapes and habitats, many more species are expected to be found in this region. Here we report the species recorded in the central

How to cite: Murvanidze, M., Cilbircioğlu, C., Özdemir, E. & Inak, E. (2020) Checklist of oribatid mites (Acari: Oribatida) of the Central Black Sea basin of Turkey with new records for the country. *Persian Journal of Acarology*, 9(3): 255–277.

and eastern Black Sea areas of Turkey collected in soil and arboricolous habitats of pure pine stands and mixed deciduous forests covering three national parks close to the capital city Ankara and intact forest areas of Kastamonu region.

MATERIAL AND METHODS

Study sites

Study area including Aluçdağı, Tekkedağı and Şahinler national parks is located in transition zone of Central Anatolia and the Black Sea region with climate characteristic for the region (see above). Aluçdağı National Park was created in 2005 and covers a 90 ha area. Woody plants are represented by *Pinus nigra* Arn., *P. brutia* Ten., *Quercus* sp., *Juniperus* sp. Tekkedağı National Park was established in 2011 with a 100 ha area in total. Woody plants are represented mainly by *Pinus nigra*, *P. sylvestris* L., *Quercus* sp., *Juniperus* sp. Şahinler National Park was established in 2011 with a 40 ha area in total. Woody plants are represented mainly by *Pinus nigra*, *P. sylvestris*, *Quercus* sp., *Juniperus* sp., *Fagus* sp.

In Kastamonu region sampling was conducted in Taşköprü, Küre and Daday forests between September 2017 and December 2018. Forests are spread along transition zone of the Black sea and Küre mountain range and therefore share climate characteristics of both regions. Taşköprü forest covers 113.500 ha area, Daday forest occupies 63867 ha and Küre forest 73.693 ha area. Küre forests belong to Küre National Park, which was created in 7 July 2000 and is known as one of the diversity hotspots of Turkey because of its 930 plant, 129 bird, 48 mammal, 8 reptile and 9 amphibian registered species (Anonymous 2014).

Sampling procedures and laboratory treatment

Forty-five soil samples were taken in Aluçdağı National Park (Fig. 1), 50 samples were collected in Tekke dağı and 20 samples were collected in Şahinler by the Turkish co-authors. Each soil sample was collected randomly using a hand trowel at a depth of 0–30 cm; moss cover was taken together with soil where available. Each sample consisted of approximately 1.5 kg of soil collected by combining 3–4 sub-samples randomly taken mostly around trees. In total 115 soil samples were collected. For each site, GPS coordinates and sampling habitat description are provided.

In forests of Kastamonu region dead wood and parts of tree bark of *P. nigra* and *P. sylvestris* together with lichens (where available) were collected on 1.5 m height from the forest floor. In addition, several *P. abies* (L.) H. Karst. trunks were sampled from forest of Trabzon which is located in the Eastern Black Sea Region (Fig. 1).

All samples were placed in plastic bags, labeled and transported to the laboratory in refrigerators. Mites were extracted using Berlese-Tullgren funnels for one week and individuals were stored in 80% ethanol. For species identification temporary slides were prepared using lactic acid.

Species identification and characteristics

Only adult mites were identified to species level using mainly keys of Ghilarov and Krivolutsky (1975) and Weigmann (2006). Ecology for each species is provided after Weigmann (2006) and Murvanidze and Mumladze (2016), while global distribution data - after Subías (2004, 2020 update). Specific synonyms of the species are listed only as far as cited in Turkish literature. For each species, locations in Turkey are given based on existing literature. We could not refer to the early checklists of Turkish oribatid mites (Özkan *et al.* 1994; Erman *et al.* 2007), since these papers only indicate presence of listed species in Turkey, without any information about their location. Remarks referring to ecological, taxonomic and/or regional peculiarities are provided for each new record. For new records we also provide measurements of body length and width, rostral

(*ro*), lamellar (*le*), interlamellar (*in*), notogastral (*ng*) setae and sensilli (*ss*) in case these measures contradicted with available descriptions. Measurements were taken dorsally under Accu-Scope 3000-Led Series microscope.




Figure 1. Map of study area with the points of data collection.

We followed systematic statement of Schatz *et al.* (2011). Species names are mostly provided after Weigmann (2006), for all ptyctimous mites catalogue of Niedbala and Liu (2018) was used. The new records for Turkey are marked with an asterisk (*)

RESULTS

Superfamily Epilohmannioidea Oudemans, 1923 Family Epilohmanniidae Oudemans, 1923

Epilohmannia (Epilohmannia) cylindrica (Berlese, 1904)

Material – One specimen in Saraycık forest processing area, Taşköprü forest, N 41° 20' 46.25", E 34° 33' 15.72", alt. = 1223 m, 20 June 2018, dead *P. nigra*.

Previous records in Turkey – Erzurum, Konya (Dik *et al.* 1999), Kayseri (Toluk and Ayyildiz 2008a), Kocaeli (Yaça and Baran 2019), Sakarya (Baran *et al.* 2015).

Global distribution – Cosmopolitan.

Ecology – Dry soils.

Superfamily Euphthiracaroidea Jacot, 1930 Family Euphthiracaridae Jacot, 1930

Acrotritia ardua (C.L. Koch, 1841)

Syn.: *Rhysotritia ardua* (C.L. Koch, 1841) *sensu* Baran and Ayyildiz (2000)

Material – One specimen in Taşköprü forest, N $41^{\circ} 20' 09.64''$, E $34^{\circ} 25' 58.98''$, alt. = 1327 m, 23 June 2018; one specimen in Taşköprü forest, N $41^{\circ} 22' 29.96''$, E $34^{\circ} 25' 57.29''$, alt. = 1200 m, 17 July 2018, tree bark and debris of *P. nigra*.

Previous records in Turkey – Artvin Province (Ayyildiz *et al.* 2011a), Erzincan, Ankara (Baran and Ayyildiz 2000).

Global distribution – Cosmopolitan.

Ecology – All types of habitats.

Superfamily Phthiracaroidea Perty, 1841

Family Phthiracaridae Perty, 1841

Phthiracarus (Phthiracarus) ferrugineus* (C.L. Koch, 1841)

Material – New record for Turkey. One specimen in Trabzon, N $40^{\circ} 58' 24.75''$, E $39^{\circ} 30' 25.55''$, alt. = 348 m, Yeşiltepe Forest Area, Akçabat. 06 June 2018, trunk of *P. abies*.

Global distribution – Holarctic.

Ecology – Forest soils.

Note – The species is known to be generally associated with forest soils (Weigmann 2006), however, is recorded for arboricolar habitats as well. It was found abundantly on twigs and leaves of *Rhododendron ponticum* L. (Murvanidze and Arabuli 2015).

Steganacarus (Tropacarus) carinatus carinatus* (C.L. Koch, 1841)

Material – New record for Turkey. Two specimens in Aluçdağı National Park, N $40^{\circ} 49' 75.37''$, E $32^{\circ} 58' 26.46''$, alt. = 1402 m, 17 July 2018, soil.

Global distribution – Palaearctic.

Ecology – All types of habitats.

Superfamily Crotonioidea Thorell, 1876

Family Crotoniidae Thorell, 1876

***Camisia (Camisia) horrida* (Hermann, 1804)**

Material – One specimen in Aluçdağı National Park, N $40^{\circ} 49' 7537''$, E $32^{\circ} 58' 2646''$, alt. = 1402 m, 17 July 2018, soil.

Previous record in Turkey – Sivas (Toluk and Akın 2017).

Global distribution – Holarctic.

Ecology – Soil, moss, frequent on tree trunks and canopy.

Family Nothridae Berlese, 1896

***Nothrus borussicus* Sellnick, 1928**

Material – One specimen in Aluçdağı National Park, N $40^{\circ} 49' 75.37''$, E $32^{\circ} 58' 26.46''$, alt. = 1402 m, 17 July 2018, soil.

Previous record in Turkey – Konya (Zeytun *et al.* 2017).

Global distribution – Palaearctic.

Ecology – Alpine meadows and mountain forests.

Superfamily Neoliodoidea Sellnick, 1928
Family Neolididae Sellnick 1928

Poroliodes farinosus (C.L. Koch, 1839)

Material – One specimen in Taşköprü forest, N 41° 15' 36.46'', E 34° 13' 11.96'', alt. = 1236 m, 15 May 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Uzunulok forest in Erzurum (Yalçın *et al.* 2013).

Global distribution – Palaearctic.

Ecology – Arboricolous, rare in forest and meadow soils.

Superfamily Plateremaeoidea Trägårdh, 1931
Family Aleurodamaeidae Paschoal & Johnston, 1984

Aleurodamaeus setosus (Berlese, 1883)

Material – One specimen in Çiftlik Village Tekçam Area, Taşköprü forest, N 41° 15' 19.75'', E 34° 14' 06.85'', alt. = 1329 m, 15 May 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Cat forest, Sivas Province (Toluk and Akin 2017).

Global distribution – Palaearctic.

Ecology – Forest soils.

Superfamily Cepheoidea Berlese, 1896
Family Cepheidae Berlese, 1896

Eupterotegaeus ornatissimus (Berlese, 1908)

Material – One specimen in Aluçdağı National Park, N 40° 49' 75.37'', E 32° 58' 26.46'', alt. = 1402 m, 15 July 2018, soil.

Previous record in Turkey – Sakarya (Susyal *et al.* 2018).

Global distribution – Holarctic.

Ecology – Forest soils.

Superfamily Ameroidea Bulanova-Zachvatkina, 1957
Family Eremulidae Grandjean, 1965

Eremulus flagellifer Berlese, 1908

Material – One specimen in Daday forest, N 41° 23' 55.18'', E 33° 13' 09.44'', alt. = 1136 m, 19 October 2018, tree bark and debris from *P. nigra*.

Previous record in Turkey – Yozgat Province (Per *et al.* 2018).

Global distribution – Cosmopolitan.

Ecology – Forest soils and meadows.

Superfamily Zetorchostoidea Michael, 1898
Family Eremaeidae Oudemans, 1900

Eueremaeus oblongus quadrilamellatus (Hammer, 1952)*

Material – New record for Turkey. One specimen in Aluç Dağı National Park, Ankara, N 40° 49' 75.37'', E 32° 58' 26.46'', alt = 1402 m, 15 July 2018, soil.

Global distribution – Holarctic.

Ecology – Forest soils and litter. It was described from thick litter made of leaves and spruce needles (Hammer 1952).

Superfamily Gustavioidea Oudemans, 1900
Family Astegistidae Balogh, 1961

***Cultroribula bicultrata* (Berlese, 1905)**

Material – One specimen in Taşköprü forest, Karadedeoğlu Village, N 41° 23' 08.05'', E 34° 25' 07.52'', alt. = 1126 m, 24 July 2018, tree bark and debris of *P. nigra*; one specimen in Çamkonak Forest Processing Area, Daday forest. N 41° 23' 55.18'', E 33° 13' 09.44'', alt. = 1136 m, 30 May 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Kocaeli (Baran and Merve 2019; Yaça and Baran 2019).

Global distribution – Holarctic.

Ecology – Forest soils.

Family Liacaridae Sellnick, 1928

***Adoristes* (*Adoristes*) *ovatus* (C.L. Koch, 1839)**

Material – One specimen in Çiftlik Village Tekçam Area, Taşköprü forest, N 41° 14' 39.56'', E 34° 12' 19.57'', alt. = 1422 m, 15 May 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Erzurum (Akman et al. 2018).

Global distribution – Holarctic.

Ecology – All types of soils with preference of forest habitats.

***Liacarus* (*Liacarus*) *brevilamellatus* Mihelčič, 1955**

Material – One specimen in Çamkonak Forest Processing Area, Daday forest, N 41° 23' 25.96'', E 33° 13' 58.20'', alt. = 1271 m, 19 October 2018, dead *P. nigra*; one specimen in Saraycık Forest Processing Area, Taşköprü Forest, N 41° 22' 51.20'', E 34° 32' 18.21'', alt. = 1128 m, 15 July 2018, tree bark and debris of *P. nigra*.

Previous records in Turkey – Erzurum (Ocak et al. 2007), Kocaeli (Yaça and Baran 2019).

Global distribution – Palaearctic.

Ecology – Forest soils, moss, litter

***Liacarus* (*Liacarus*) *coracinus* (C.L. Koch, 1841)**

Material – One specimen in Aluç Dağı (mountain) Natural Park, Ankara, N 40° 49' 75.37'', E 32° 58' 26.46'', alt. = 1402 m, 15 July 2018, soil.

Previous records in Turkey – Giresun, Mersin, Ordu and Samsun, Trabzon (Grobler et al. 2003), Kocaeli (Yaça and Baran 2019).

Global distribution – Palaearctic.

Ecology – Forest soils, moss, litter.

Family Peloppiidae Balogh, 1943

Ceratoppia quadridentata (Haller, 1882)

Material – One specimen in Saraycık Forest Processing Area, Taşköprü forest, N 41° 22' 51.20'', E 34° 32' 18.21'', alt. = 1128 m, 15 July 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Erzurum (Akman *et al.* 2018).

Global distribution – Holarctic.

Ecology – Forest soils.

Superfamily Carabodoidea C.L. Koch, 1837
Family Carabodidae C.L. Koch, 1837

Carabodes (Carabodes) labyrinthicus (Michael, 1879)

Material – One specimen in Çiftlik Village Tekçam Area, Taşköprü forest, N 41° 14' 39.56'', E 34° 12' 19.57'', alt. = 1422 m, 15 May 2018; one individual in Alasökü Village, Taşköprü forest, N 41° 20' 09.64'', E 34° 25' 58.98'', alt. = 1327 m, 10 June 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Bolu (Toluk and Ayyıldız 2016).

Global distribution – Holarctic.

Ecology – Forest soils, moss, litter, tree bark and canopy.

Superfamily Oppioidea Grandjean, 1951
Family Epimerellidae Ayyıldız & Luxton, 1989

Epimerella smirnovi longisetosa Kulijev, 1967

Material – One specimen in Tekke Dağı Natural Park, Ankara, N 40° 16' 96.96'', E 31° 91' 72.68'', alt. = 1424 m, 25 July 2018, soil.

Previous record in Turkey – Amanos mountains (Ay and Ayyıldız 2019).

Global distribution – Mediterranean.

Ecology – Dry meadows and urban soils.

Remarks – Kulijev (1962) reported a longer size for sensilli for *E. smirnovi smirnovi* Kulijev, 1962 (94 µm), while in paper published in 1967, the size of sensilli is indicated as 54 µm, but no size of sensilli is indicated for *E. smirnovi longisetosa*. We measured sensilli for Turkish individual which equals 80 µm. For *Epimerella smirnovi smirnovi* found in Georgia, size of sensilli is between 75–80 µm and matches the size of Turkish specimen.

Family Oppiidae Grandjean, 1954

Graptoppia (Graptoppia) paraanalisis Subias and Rodriguez, 1985*

Material – New record for Turkey. One specimen in Çiftlik Village, Tekçam Area, Taşköprü forest, Kastamonu, N 41° 27' 39.53'', E 34° 19' 42.99'', alt. = 898 m, 5 June 2018; one specimen in Yeşiltepe Area, Akçaabat, Trabzon, 27 May 2018, N 40° 58' 24.75'', E 39° 30' 25.55'', alt. = 348 m; seven specimens in Alasökü Village, Taşköprü forest, Kastamonu, 16 June 2018, N 41° 20' 09.64'', E 34° 25' 58.98'', alt. = 1327 m; three specimens in Çamkonak Forests, Daday, Kastamonu, 09 October 2018, N 41° 23' 54.24'', E 33° 13' 13.73'', alt. = 1147 m. Tree bark and debris of *P. abies* and *P. nigra*.

Global distribution – Palaearctic.

Ecology – Litter and soil.

Remarks – Turkish individuals are larger ($285 \times 155 \mu\text{m}$) than reported by Subías and Rodriguez (1985) for Spanish and Italian specimens ($232\text{--}245 \mu\text{m}$). All other characters match the original description (Subías and Rodriguez 1985).

***Lauroppia fallax* (Paoli, 1908)**

Material – One specimen in Aluç Dağı Natural Park, Ankara, N $40^\circ 49' 75.37''$, E $32^\circ 58' 26.46''$, alt. = 1402 m, 15 July 2018, soil.

Previous records in Turkey – Locality unknown (Toluk and Ayyıldız 2008).

Global distribution – Semicosmopolitan.

Ecology – All types of habitats.

***Ramusella (Insculptoppia) insculpta* (Paoli, 1908)**

Material – One specimen in Saraycık Forest Processing Area, Taşköprü, N $41^\circ 18' 10.00''$, E $34^\circ 25' 48.64''$, alt. = 1277 m, 10 July 2018, *P. sylvestris*, dead tree; one specimen in Karadedeoğlu Village Tekkeyaylası Area, Taşköprü forest, N $41^\circ 22' 55.26''$, E $34^\circ 24' 19.06''$, alt. = 1177 m, 2 August 2018, *P. nigra*, dead tree; three specimens in Yeşiltepe Area, Akçaabat, Trabzon, N $41^\circ 20' 09.64''$, E $34^\circ 25' 58.98''$, alt. = 1327 m, 2 June 2018, tree trunk of *P. abies*.

Previous record in Turkey – Erzurum (Baran and Ayyıldız 2004), Trabzon, Samsun (Ayyıldız 1989), Ankara (Çobanoğlu and Bayram 1998).

Global distribution – Palaearctic.

Ecology – All types of habitats.

Superfamily Tectocepheoidea Grandjean, 1954
Family Tectocepheidae Grandjean, 1954

Tectocepheus punctulatus* Djaparidze, 1985

Material – New record for Turkey. Five specimens in Aluç Dağı Natural Park, Ankara. N $40^\circ 49' 75.37''$, E $32^\circ 58' 26.46''$, alt. = 1402 m. 15 July 2018, Soil.

Global distribution – Palaearctic.

Ecology – All types of habitats.

***Tectocepheus velatus sarekensis* Trägårdh, 1910**

Syn.: *T. sarekensis* sensu Dik et al. (1999)

Material – One specimen in Alasökü Village, Taşköprü forest, N $41^\circ 20' 09.64''$, E $34^\circ 25' 58.98''$, alt. = 1327 m, 2 July 2018; one specimen in Çamkonak Forest Processing Area, Daday, N $41^\circ 23' 54.24''$, E $33^\circ 13' 13.73''$, alt. = 1147 m, 19 October 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Konya (Dik et al. 1999).

Global distribution – Cosmopolitan.

Ecology – All types of habitats.

Superfamily Cymbaeremaeoidea Sellnick, 1928
Family Cymbaeremaeidae Sellnick, 1928

***Cymbaeremaeus cymba* (Nicolet, 1855)**

Material – One specimen in Karadedeoğlu Village, Taşköprü forest, N $41^\circ 23' 08.05''$, E $34^\circ 25'$

07.52'', alt. = 1126 m, 17 July 2018, tree bark and debris of *P. nigra*.

Previous records in Turkey – Ankara (Grobler *et al.* 2004), Izmir (Çobanoğlu 2009).

Global distribution – Palaearctic.

Ecology – All types of habitats. Frequent on tree barks and canopy.

Superfamily Licneremaoidea Grandjean, 1931
Family Licneremaeidae Grandjean, 1931

***Licneremaeus licnophorus* (Michael, 1882)**

Material – One specimen in Karadedeoğlu Village Tekkeyaylası Area, Taşköprü forest, N 41° 22' 24.45'', E 34° 23' 25.15'', alt. = 1224 m, 2 August 2018, tree trunk of *P. nigra*; one specimen in Karadedeoğlu Village, Taşköprü forest, N 41° 22' 55.11'', E 34° 24' 03.56'', alt. = 1201 m, 17 July 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Bolu (Toluk and Ayyıldız 2011).

Global distribution – Holarctic.

Ecology – Moss, tree trunks.

Family Passalozetidae Grandjean, 1954

***Passalozetes africanus* Grandjean, 1932**

Material – One specimen in Kalecik, N 40° 06' 44.50'', E 33° 25' 43.30'', alt. = 732 m, 20 July 2018, soil with litter in vineyard.

Previous record in Turkey – Erzurum (Ayyıldız 1988).

Global distribution – Palaearctic.

Ecology – Dry meadow soil.

Superfamily Phenopelopoidea Petrunkevitch, 1955
Family Phenopelopidae Petrunkevitch, 1955

***Eupelops acromios* (Hermann, 1804)**

Material – Two specimens in Çiftlik Village, Tekçam Area, Taşköprü forest, N 41° 14' 39.56'', E 34° 12' 19.57'', alt. = 1422 m, 15 May 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Konya (Dik *et al.* 1999).

Global distribution – Semicosmopolitan.

Ecology – All types of habitats.

Eupelops plicatus* (C.L. Koch, 1835)

Material – New record for Turkey. One specimen in Ankara university campus, N 39° 96' 35.51'', E 32° 86' 11.78'', alt. = 858 m, 25 July 2018, mushrooms.

Global distribution – Holarctic.

Ecology – All types of habitats.

Remarks – Measurements: The size of Turkish specimen (500 × 345 µm) is at the lower range provided in the literature (Weigmann 2006). All characters match previous descriptions (Ghilarov and Krivolutsky 1975; Beck and Woas 1991; Weigmann 2006).

***Peloptulus (Peloptulus) phaeonotus* (C.L. Koch, 1844)**

Material – One specimen in Köçekli Village, Taşköprü forest, N $41^{\circ} 24' 05.84''$, E $34^{\circ} 22' 30.42''$, alt. = 1193 m, 10 August 2018, tree bark and debris of *P. sylvestris*.

Previous record in Turkey – Kayseri (Seniczak et al. 2014)

Global distribution – Palaearctic.

Ecology – All types of habitats.

Superfamily Achipteroidea Thor, 1929
Family Achipteriidae Thor, 1929

***Parachipteria fanzagoi* (Jacot, 1929)**

Syn.: *Parachipteria willmanni* van der Hammen, 1952 *sensu* Per and Ercan (2018)

Material – Two specimens in Aluç Dağı Natural Park, Ankara, N $40^{\circ} 49' 75.37''$, E $32^{\circ} 58' 26.46''$, alt. = 1402 m, 15 July 2018, soil.

Previous record in Turkey – Kayseri (Per and Ercan 2015)

Global distribution – Holarctic.

Ecology – Forest moss, litter and soils.

Remarks – In the world catalogue of Subías (2020 update) this species is listed as *Campachipteria (Triachipteria) fanzagoi* (Jacot, 1929). In the checklist of Georgian oribatid mites (Murvanidze and Mumladze 2016), we explained the reasons of disagreements with this placement with the main reason of *Campachipteria* species having monodactylous legs (Aoki 1995) and *P. fanzagoi* being the tridactylous species. The name of the species is provided after Weigmann (2006).

Family Tegoribatidae Grandjean, 1954

***Lepidozetes singularis* Berlese, 1910**

Material – Two specimens in Çiftlik Village Tekçam Area, Taşköprü forest, N $41^{\circ} 14' 39.56''$, E $34^{\circ} 12' 19.57''$, alt. = 1422 m, 15 May 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Erzurum (Ayyıldız et al. 2011b).

Global distribution – Holarctic.

Ecology – forest soils, moss, frequent on trees.

Superfamily Oripodoidea Jacot, 1925
Family Haplozetidae Grandjean, 1936

***Protoribates (Protoribates) capucinus* (Berlese, 1908)**

Material: Two specimens in Saraycık Forest Processing Area, Taşköprü forest, N $41^{\circ} 22' 51.20''$, E $34^{\circ} 32' 18.21''$, alt. = 1128 m, 3 July 2018; one specimen in Çamkonak Forest Processing Area, Daday, N $41^{\circ} 23' 54.24''$, E $33^{\circ} 13' 13.73''$, alt. = 1147 m, 30 May 2018, tree trunk of *P. nigra*.

Previous records in Turkey – Erzurum (Ayyıldız 1988), Kocaeli (Yaça and Baran 2019), Konya (Dik et al. 1999).

Global distribution – Cosmopolitan.

Ecology – All types of habitats with preference of humid forest soils.

Family Oribatulidae Thor, 1929

***Oribatula (Zygoribatula) exilis exilis* (Nicolet, 1855)**

Material – Two specimens in Karadedeoğlu Village, Tekkeyaylası Area, Taşköprü forest, N 41° 22' 24.45", E 34° 23' 25.15", alt. = 1224 m, 2 August 2018, tree trunk of *P.nigra*; 8 specimens in Çiftlik Village, Tekçam Area, Taşköprü forest, 15 May 2018, N 41° 15' 19.75", E34° 14' 06.85", alt. = 1329 m, tree bark and debris of *P. nigra*.

Previous record in Turkey – Ankara (Grobler *et al.* 2005).

Global distribution – Holarctic.

Ecology – All types of habitats.

***Oribatula (Zygoribatula) frisiae* (Oudemans, 1900)**

Material – Twenty specimens in Güdül, N 40° 2' 03.34", E 32° 15' 23.28", alt. = 729 m, 20 July 2018, soil with litter in vineyard.

Previous records in Turkey – Ankara (Kumral and Cobanoğlu 2016), Kocaeli (Yaça and Baran 2019).

Global distribution – Holarctic.

Ecology – Forest and urban soils.

***Oribatula (Zygoribatula) nicora* (Djaparidze, 1985)**

Material – Twelve specimens in Karadedeoğlu Village, Taşköprü forest, N 41° 22' 55.11", E 34° 24' 03.56", alt. = 1204 m, 24 July 2018, tree bark and debris of *P. nigra*; 21 specimens in Yeşiltepe Area, Akçaabat, Trabzon, N 40° 58' 24.75", E 39° 30' 25.55", alt. = 348 m, 22 May 2018, tree bark and debris of *P. abies*; one specimen in Alasökü Village, Taşköprü, N 41° 19' 56.66", E 34° 24' 24.72", alt. = 1468 m, 16 June 2018, tree bark and debris of *P. sylvestris*; three specimens in Çamkonak Forest Processing Area, Daday forest, N 41° 23' 3.74", E 33° 14' 02.03", alt. = 1382 m, 19 October 2018, tree bark and debris of *P. nigra*.

Previous records in Turkey – Halaçlı, Taşköprü Kabalar, Merkez Uzunkavak, Taşköprü Kızılcaören, Taşköprü Ağcıkışi, Merkez Çavundur, Hanönü Kornapa, Taşköprü Ağcıkışi villages in Kastamonu region (Murvanidze *et al.* 2018).

Global distribution – Caucasus.

Ecology – All types of habitats.

Oribatula (Zygoribatula) propinqua* (Oudemans, 1900)

Material – New record for Turkey. One specimen in Şahinler Natural Park, Ankara. Woody plants are represented by *Pinus nigra*, *P. sylvestris* L., *Quercus* sp., *Fagus* sp., N 40° 61' 74.65", E 32° 45' 55.46", alt. = 1660 m, 16 June 2018, soil with moss.

Global distribution – Palaearctic.

Ecology – On the trees, rare in dry soils.

Phauloppia rauschenensis* (Sellnick, 1908)

Material – New record for Turkey. Three specimens in Karadedeoğlu Village, Tekkeyaylası Forest Area, Taşköprü-Kastamonu, N 41° 22' 23.26", E 34° 24' 22.11", alt. = 1187 m, 17 July 2018, tree bark from *P. nigra* and *P. sylvestris*.

Global distribution – Palaearctic.

Ecology – Forest soils, frequent in canopy.

Remarks – Measurements: Body size: 400–430 × 195–215 µm. *ro* 45 µm, *in* 40 µm, *le* 45 µm, *ss* 25 µm, all *ng* setae except *dp* and *lp* 30–35 µm, *dp* and *lp* 40 µm.

Body size of Turkish individuals is at the upper range of the reported size, i.e. 310–390 µm (Weigmann 2006) or larger. Notogastral sculpture (foveolae) is evident at the sides of the notogaster. *Areae porosae* are round and small. Sensilli with short stalk and rounded head. Notogastral setae about 30–40 µm in length versus reported 25 µm (Weigmann 2006).

Species prefers canopy habitats (Weigmann 2006). This habitat preference is proved by Turkish findings on the bark of the pine trees. *Phauloppia rauschenensis* was found abundantly on branches of *Picea orientalis* and in lower numbers on *Fagus orientalis* in mixed forests of Borjom-Kharagauli National Park in Georgia (Murvanidze and Mumladze 2014). Presence of this species on beech bark was also reported by Wunderle *et al.* (1990). This shows that *P. rauschenensis* prefers canopy habitats in general, with no difference between coniferous or broad-leaved tree species.

Siculobata (Paraleius) leontonycha (Berlese, 1910)*

Material – New record for Turkey. Four specimens in Karadedeoğlu Village, Tekkeyaylaı Forest Area, Taşköprü-Kastamonu, N 41° 22' 23.26", E 34° 24' 22.11", alt. = 1187 m, 23 August 2018; two specimens in Alasökü Village Forests, Taşköprü-Kastamonu N 41° 20' 09.64", E 34° 25' 58.98", alt. = 1327 m, 10 June 2018, tree bark and debris of *P. nigra* and *P. sylvestris*.

Global distribution – Holarctic.

Ecology – Bark debris around the trunks, in pathways of woodborer beetles.

Remarks – Measurements: Weigmann (2006) reports size of *S. leontonycha* as 435–500 µm. Travé (1960), Wunderle *et al.* (1990) and Perez-Iñigo (1993) also indicate larger individuals (440–500 µm, 435–480 µm and 440–500 µm respectively), while Ghilarov and Krivolutsky (1975) and Mahunka (1996) do not indicate body size at all. All Turkish individuals are smaller (360–365 × 215–220 µm) than reported. The rest of the characters match with provided descriptions.

The species is known to be arboricolous (Weigmann 2006). Turkish records prove this habitat preference by finding all individuals on bark of pine tree species. Checklist of oribatid mites of Georgia (Murvanidze and Mumladze 2016) report presence of this species in soils of alpine meadow after the catalogue of Caucasian oribatids of Shtanchaeva and Subias (2010), however, this report is a mistake, because the catalogue does not list *S. leontonycha* at all and Georgian record can be discarded.

Family Scheloribatidae Grandjean, 1933

Liebstadia (Liebstadia) similis (Michael, 1888)

Material – One specimen in Yeşiltepe Area, Akçaabat, Trabzon, N 40° 58' 24.75", E 39° 30' 25.55", alt. = 348 m, 27 May 2018, tree trunk of *P. abies*; five specimens in Saraycık Forest Processing Area, Taşköprü, 15 July 2018, tree bark and debris of *P. nigra*.

Previous records in Turkey – Erzurum (Ayyıldız 1988), Ankara (Grobler *et al.* 2004).

Global distribution – Holarctic.

Ecology – All types of habitats.

Scheloribates (Scheloribates) barbatulus Mihelčić, 1956*

Material – New record for Turkey. Two specimens in Çamkonak Forest Processing Area, Daday forest, N 41° 23' 54.24", E 33° 13' 13.73", alt. = 1147 m, 30 May 2018, tree bark of *P. nigra*.

Global distribution – Palaearctic.

Ecology – Forest soils.

Remarks – Although the species is known not to be adapted to arboricolar habitats (Perez-Iñigo 1993) we found them on bark of the pine tree. Other representative of genus *Scheloribates* [*S. latipes* (C.L. Koch, 1841)] has been reported in minor numbers on twigs of spruce and beech trees (Murvanidze and Mumladze 2014).

***Scheloribates (Scheloribates) laevigatus* (C.L. Koch, 1835)**

Material – Sixteen specimens in Aluç Dağı Natural Park, Ankara, 19 July 2018, N 40° 49' 75.37'', E 32° 58' 26.46'', alt. = 1402 m, soil.

Previous records in Turkey – Erzincan (Dik *et al.* 1995), Konya (Dik *et al.* 1999), Kocaeli (Yaça and Baran 2019).

Global distribution – Semicosmopolitan.

Ecology – All types of habitats.

***Scheloribates (Scheloribates) latipes* (C.L. Koch, 1844)**

Material – One specimen in Çiftlik Village, Tekçam Area, Taşköprü forest, N 41° 22' 14.15'', E 34° 22' 36.05'', alt. = 1272 m, 5 September 2018, tree trunk of *P. nigra*; three specimens in Karadedeoğlu Village, Taşköprü forest, N 41° 22' 55.11'', E 34° 24' 03.56'', alt. = 1204 m, 24 July 2018, tree trunk of *P. nigra*; one specimen in Çiftlik Village, Tekçam Area, Taşköprü forest, N 41° 15' 19.75'', E 34° 14' 06.85'', alt. = 1329 m, 15 May 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Konya (Dik *et al.* 1999).

Global distribution – Semicosmopolitan.

Ecology – All types of habitats.

Superfamily Ceratozetoidea Jacot, 1925
Family Ceratozetidae Jacot, 1925

Ceratozetes (Ceratozetes) colchicus* Murvanidze & Weigmann, 2003

Material – New record for Turkey. One specimen in Alasökü Village, Taşköprü forest, N 41° 19' 55.68'', E 34° 25' 16.18'', alt. = 1369 m, 16 June 2018, tree trunk of *P. sylvestris*; one specimen in Şahinler Natural park, Ankara. N 40° 61' 74.65'', E 32° 45' 55.46'', alt. = 1670 m, 16 June 2018, soil.

Global distribution – Caucasus and Turkey.

Ecology – Humid forest soils.

Remarks – Measurements: Body size: 500 × 320 µm. *ro* 65 µm, *in* 130 µm, *le* 75 µm, *ss* 95 µm, *ng* vestigial.

Turkish specimens show smaller size compared to the original description (Murvanidze and Weigmann 2003): 1) body length in Turkish specimens is about 500–510 µm, while in description 540–580 µm is indicated; 2) lamellar setae 75 µm vs reported 85 µm; 3) *ss* 95 µm vs 150 µm; 4) length of lamellar cuspides 50 µm vs 80 µm. The rest of the characters match the provided description and paratypes available in the collection of the Georgian Agricultural University.

Ceratozetes (Ceratozetes) conjunctus* Mihelčić, 1956

Material – New record for Turkey. One specimen in Alasökü Village, Taşköprü forest, N 41° 19' 55.68'', E 34° 25' 16.18'', alt. = 1369 m, 16 June 2018; one specimen in Alasökü Village, Taşköprü forest, N 41° 19' 55.16'', E 34° 25' 08.12'', alt. = 1408 m, 2 July 2018; one specimen in Çamkonak

forests, Daday, Kastamonu, N $41^{\circ} 23' 43.63''$, E $33^{\circ} 13' 38.96''$, alt. = 1215 m, 19 October 2018, tree trunk of *P. sylvestris*.

Global distribution – Palaearctic.

Ecology – Alpine meadow and forest soils.

Ceratozetes (Ceratozetes) minutissimus* Willmann, 1951

Material – New record for Turkey. One specimen in Yeşiltepe forest area, Akçabat, Trabzon, N $40^{\circ} 58' 24.75''$, E $39^{\circ} 30' 25.55''$, alt. = 348 m, 22 May 2018, soil under *P. sylvestris*.

Global distribution – Palaearctic.

Ecology – Forest and meadow soils.

Jugatala angulata* (C.L. Koch, 1839)

Material – New record for Turkey. One specimen in Karadedeoğlu Village, Tekkeyaylasi forest area, Taşköprü-Kastamonu, N $41^{\circ} 22' 23.26''$, E $34^{\circ} 24' 22.11''$, alt. = 1187 m, 17 July 2018, tree bark of *P. nigra*.

Global distribution – Europe.

Ecology – Arboricolous, mostly on coniferous trees (spruce and pine).

Remarks – Measurements: Body size: $600 \times 370 \mu\text{m}$.

All characters match the description provided by Weigmann (2006), however, Bayartogtokh and Schatz (2008) report a smaller size ($486 \mu\text{m}$) for individuals found in Austria. Up to now *J. angulata* was known from Europe only (Weigmann 2006; Bayartogtokh and Schatz 2008). Our record broadens the area of occurrence for this species. We confirm habitat preference of *J. angulata* by finding it on the bark of the pine tree as originally described, however, Sobek *et al.* (2008) indicate presence of the species on broadleaved trees (beech and oak) as well.

Trichoribates (Trichoribates) myricus* Gjelstrup and Solhøy, 1994

Material – New record for Turkey. Two specimens in Aluç Dağı Natural Park, Ankara, N $40^{\circ} 49' 75.37''$, E $32^{\circ} 58' 26.46''$, alt. = 1402 m, 15 June 2018, soil.

Global distribution – Euroatlantic.

Ecology – Humid meadows, forest soils.

Remarks – Measurements: Body size of Turkish individuals ($570 \times 345 \mu\text{m}$) is smaller than provided in original description ($650 \times 420 \mu\text{m}$) (Gjelstrup and Solhøy 1994), however Weigmann (2006) reports smaller size ($555\text{--}670 \mu\text{m}$), so that our findings match within the given range. Weigmann (2006) also indicates size of nogastral setae up to $60 \mu\text{m}$, while for Turkish individuals these setae are even shorter, $45 \mu\text{m}$.

This is the first record of the species out of Europe. Previous findings were known from Iceland (Gjelstrup and Solhøy 1994) and Germany (Weigmann 2006). The species was described from bogs and swamps in Island and was additionally found in Schleswig-Holstein and Odertal in Germany (Weigmann 2006). Finding site in forested area of Turkey adds more information about habitat preferences for this species.

Family Mycobatidae Grandjean, 1954

***Puncitoribates (Puncitoribates) punctum* (C.L. Koch, 1839)**

Material – Nine specimens in Saraycık Forest Processing Area, Taşköprü, N $41^{\circ} 19' 15.36''$, E $34^{\circ} 26' 57.46''$, alt. = 1122 m, 27 June 2018; one individual in Karadedeoğlu Village, Taşköprü, N 41°

22' 23.26'', E 34° 24' 22.11'', alt. = 1187 m, 2 August 2018; one specimen in Saraycık Forest Processing Area, Taşköprü, N 41° 19' 15.36'', E 34° 26' 57.46'', alt. = 1122 m, 15 July 2018, tree trunk of *P. sylvestris*.

Previous records in Turkey – Demirli, Çarsamba (Bayartogtokh *et al.* 2002).

Global distribution – Cosmopolitan.

Ecology – All types of habitats.

Superfamily Galumnoidea Jacot, 1925
Family Galumnidae Jacot, 1925

Galumna (Galumna) alata* (Hermann, 1804)

Material – New record for Turkey. One specimen in Tekke Dağı (mountain) Natural Park, Ankara, N 40° 16' 96.96'', E 31° 91' 72.68'', alt. = 1485 m, 25 July 2018.

Global distribution – Semicosmopolitan.

Ecology – All types of habitats.

Remarks – Measurements: Body size of Turkish individual is a little smaller (450 × 345 µm) than reported by Perez-Iñigo (1993) and Weigmann (2006), 500–630 µm. All other characters match available descriptions. The species is easy to identify by shape of sensilli, rounded areae porosae which are almost similar in size and group of irregular spots on posterior part of notogaster.

Galumna (Galumna) lanceata* (Oudemans, 1900)

Material – New record for Turkey. One specimen in Saraycık forest area, Taşköprü forests, Kastamonu, N 41° 18' 10.00'', E 34° 25' 48.64'', alt. = 1277 m, 10 July 2018, tree bark and debris of *P. nigra*.

Global distribution – Palaearctic.

Ecology – Humid forest soils.

***Pergalumna (Pergalumna) nervosa* (Berlese, 1914)**

Material – One specimen in Aluç dağı natural park, Ankara, N 40° 49' 75.37'', E 32° 58' 26.46'', alt. = 1402 m, 17 July 2018, soil.

Previous record in Turkey – Konya (Dik *et al.* 1999).

Global distribution – Holarctic.

Ecology – Forest and meadow soils. Frequent on dumps and reclaimed sites.

***Pilogalumna crassiclava* (Berlese, 1914)**

Material – One specimen in Saraycık forest processing area, Taşköprü, N 41° 22' 51.20'', E 34° 32' 18.21'', alt. = 1128 m, 15 July 2018, tree bark and debris of *P. nigra*.

Previous record in Turkey – Ankara (Grobler *et al.* 2004).

Global distribution – Palaearctic.

Ecology – All types of habitats.

DISCUSSION

Investigations dealing with the oribatid diversity of Turkish fauna are generally concentrated around the northern part of the country adjacent to the Black Sea region (Ayyıldız 1988; Ayyıldız and

Luxton 1989a, b; Dik *et al.* 1995, 1999; Baran and Merve 1999; Baran and Ayyildiz 2000; Ayyildiz *et al.* 2011a; Toluk and Ayyildiz 2011; Yalçın *et al.* 2013; Akman *et al.* 2018); while the main part of the country still remains uninvestigated. Most samplings include soil, moss and litter habitats. Specific habitats like arboricolar ones are severely understudied. There is only one paper dealing with the records of oribatid mites inhabiting tree (*P. nigra*) trunks (Toluk and Akin 2017), whereas arboricolar habitats include twigs, leaves, suspended soil, moss, lichens, etc. (André 1985; Proctor *et al.* 2002; Lindo and Winchester 2006) and generally harbor high oribatid diversity (Behan-Pelletier and Winchester 1998; Winchester *et al.* 1999, 2008; Murvanidze and Arabuli 2015).

Here, 52 species of oribatid mites were identified while 17 species of them are new for the country. New records comprise 33% of total diversity and three of them (*Ph. rauschenensis*, *S. (Paraleius) leontonycha*, *J. angulata*) are specific to arboricolar habitats (Weigmann 2006). Canopy specific oribatids (10 species) make 19% of investigated fauna, while most of the identified species (19) are known as ecological ubiquists, found in all types of habitats, 16 species are forest specific and five are mostly found in meadows.

Generally, twigs and trunk of coniferous trees are known to be favorable for oribatid diversity due to cracks, grooves and furrows on them offering more food and shelter for mites (Prinzing 1997; Sobek *et al.* 2008). *Poroliodes farinosus*, *C. labyrinthicus* and *P. rauschensis* were found in abundance on twigs of spruce trees [*Picea orientalis* (Lipsky)] in Borjomi-Kharagauli National Park in Georgia (Murvanidze and Mumladze 2014), while *C. cymba*, *C. labyrinthicus*, *P. rauschensis* and *J. angulata* were abundant on larch (*Larix decidua* Mill.) and spruce (*Picea abies* L.) branches in Central European mixed forests in Switzerland (Sobek *et al.* 2008).

Among the new records there are some widespread and common species such as *P. ferrugineus*, *S. carinatus carinatus*, *E. flagellifer*, *E. plicatus*, *G. alata*, but also some quite rare species like *E. oblongus quadrilamellatus*, *J. angulosa* and *T. myrica* which were only known from Europe up to now. All these indicate the necessity and importance of continued sampling around the country including the wide range of possible habitats.

ACKNOWLEDGEMENTS

The first author would like to express cordial gratitude to Prof. Levan Mumladze for help in creating map of sampling locations and for proofreading early version of the manuscript. We also thank the editor, Prof. Alireza Saboori, Dr. Sergey Ermilov and two other anonymous reviewers for their valuable comments. This work was supported by Kastamonu University (The Coordinator of Scientific Research Projects) in Turkey (Grant No: KÜ-BAP01/2018-26).

REFERENCES

- Akman, N., Aydin, D. & Ayyildiz, N. (2018) Gustavioid mites of the province of Erzurum (Acari, Oribatida, Gustavioidea). *Plant Protection Bulletin*, 58(2): 55–62 (In Turkish).
DOI: 10.16955/bitkorb.349179.
- André, H.M. (1985) Associations between corticolous microarthropod communities and epiphytic cover on bark. *Ecography*, 8(2): 113–119.
DOI: 10.1111/j.1600-0587.1985.tb01161.x.
- Anonymous (2014) Kure Mountains National Park. Available from: <http://nationalparksofturkey.com/kure-mountains-national-park/> (Accessed on 25 December 2019).
- Aoki, J. (1995) Oribatid mites of high altitude forests of Taiwan. *Special Bulletin of Japanese Society of Coleopterologists*, 4: 123–130.
- Ay, Y. & Ayyildiz, N. (2019) Investigations on oppioid oribatid mites of Amanos Mountains (Turkey). *Plant Protection Bulletin*, 59(3): 15–23 (In Turkish).

- Ayyildiz, N. (1988) Systematic investigations on the oribatid mites (Acari: Oribatida) of the Erzurum Plain. II. Higher oribatids. *Turkish Journal of Zoology*, 12(2): 131–144 (In Turkish).
- Ayyildiz, N. (1989) Mites of the family Oppiidae (Acari: Oribatida) from Turkey. *Journal of Natural History*, 23(6): 1373–1379.
DOI: 10.1080/00222938900770741.
- Ayyildiz, N. & Luxton, M. (1989a) New and unrecorded mites (Acari) from Turkey. *Zoologischer Anzeiger*, 222(5/6): 294–300.
- Ayyildiz, N. & Luxton, M. (1989b) Epimerellidae (Acari, Oribatida), a new mite family. *Journal of Natural History*, 23(6): 1381–1386.
- Ayyildiz, N., Aydin Yüksel, H. & Toluk, A. (2011a) The euphthiracarid mites detected in Artvin Province (Acari: Oribatida). *Plant Protection Bulletin*, 51(1): 1–15.
- Ayyildiz, N., Per, S. & Taşdemir, A. (2011b) A new record for the oribatid mite fauna of Turkey: *Lepidozetes singularis* Berlese, 1910 (Acari, Oribatida, Tegoribatidae). *Çankaya University Journal of Science and Engineering*, 8(2): 183–187.
- Balogh, J. (1943) Sistematische studien über Sieberbürgische Moosmilben. *Annales Musei historico-naturalis Hungari*, 36: 32–42.
- Balogh, J. (1961) Identification keys of world oribatid (Acari) families and genera. *Acta zoologica academiae scientiarum Hungaricae*, 7(3–4): 243–344.
- Baran, Ş. & Ayyildiz, N. (2000) Systematic studies on *Rhysotritia ardua* (C.L. Koch) (Acari, Oribatida) in Erzincan and Erzurum plains. *Turkish Journal of Zoology*, 24: 231–236.
- Baran, Ş. & Ayyildiz, N. (2004) First records of *Ramusella* Hammer, 1962 (Acari: Oribatida: Oppiidae) species in Turkey. *Turkish Journal of Entomology*, 28(1): 39–44.
- Baran, Ş., Ayyildiz, N. & Türer, G. (2015) Second record of *Epilohmannia imreorum* (Acari: Oribatida). *Journal of Acarological Society of Japan*, 24(2): 63–69.
DOI: 10.2300/acari.24.63.
- Baran, Ş. & Merve, Y. (1999) Oribatid mite fauna of Kocaeli City Forest (Kocaeli, Turkey). *Turkish Journal of Entomology*, 43(1): 41–56.
DOI: 10.16970/entoted.508099.
- Bayartogtokh, B., Çobanoğlu, S. & Ozman, S.K. (2002) Oribatid mites of the superfamily Ceratozetoidea (Acari: Oribatida) from Turkey. *Acarina*, 10(1): 3–23.
- Bayartogtokh, B. & Schatz, H. (2008) *Trichoribates* and *Jugatala* (Acari: Oribatida: Ceratozetidae) from the Central and Southern Alps, with notes on their distribution. *Zootaxa*, 1948: 1–35.
DOI: 10.11646/zootaxa.1948.1.1.
- Beck, L. & Woas, S. (1991) Die Oribatiden-Arten (Acari) eines Südwestdeutschen Buchenwaldes I. *Carolinea*, 49: 37–82.
- Behan-Pelletier, V. & Winchester, N. (1998) Arboreal oribatid mite diversity: Colonizing the canopy. *Applied Soil Ecology*, 9(1–3): 45–51.
DOI: 10.1016/S0929-1393(98)00052-3.
- Berlese, A. (1883) Sopra due generi di Acari Italiani. Lettura fatta alla R. Accademia di Padova, 33: 45–52.
- Berlese, A. (1896) *Acari, Myriapoda et Scorpiones hucusque in Italia Reperta. Ordo Cryptostigmata II (Oribatidae)*. Portici, Padova, 98 pp.
- Berlese, A. (1904) Spicilegia Zoologica. Acari Nuovi. *Redia*, 2: 10–32.
- Berlese, A. (1905) Acari nuovi. Materiali del manipulus V. *Redia*, 2: 231–261.
- Berlese, A. (1908) Elenco di generi e specie nuove di Acari. *Redia*, 5(1): 1–16.
- Berlese, A. (1910) Brevi diagnosi di generi e specie nuovi di Acari. *Redia*, 6(2): 346–388.
- Berlese, A. (1914) Acari Nuovi. *Redia*, 10(1/2): 113–150.
- Bulanova-Zachvatkina, E.M. (1957) Mites of the family Damaeidae Berl. (Acariformes, Oribatei) 1st information. *Zoologicheskii Zhurnal*, 36: 1167–1186 (In Russian).

- Çobanoğlu, S. (2009) Mite population density analyses of stored dried apricots in Turkey. *International Journal of Acarology*, 35(1): 67–75.
DOI: 10.1080/01647950902884645.
- Çobanoğlu, S. & Bayram, Ş. (1998) Mites (Acari) and flies (Insecta: Diptera) from natural edible mushrooms (Morchella: Ascomycetes) in Ankara, Turkey. *Bulletin et Annales de la Société Royale Belge d'Entomologie*, 134: 187–197.
- Djaparidze, N. (1985a) New findings of beetle mites (Acariformes, Oribatei) in the Georgian SSR. In: Cholokava, A. (Ed.), *Fauna i ekologija nekotorikh nasekomikh I kleshchei Gruzii*, Tbilisi, Metsniereba, pp. 58–71 (In Russian).
- Djaparidze, N. (1985b) A new species of the genus *Tectocepheus* Berlese, 1913 and the specific independence of *Tectocepheus velatus* var. *sarekensis* Trägardh, 1910. In: Cholokava, A. (Ed.), *Fauna i ekologija nekotorikh nasekomikh I kleshchei Gruzii*, Tbilisi, Metsniereba, pp. 72–81 (In Russian).
- Dik, B., Starý, J., Güçlü, F., Cantoray, R. & Gülbahçe, S. (1995) Oribatid mites (Acari: Oribatida) from Konya Province, Central Anatolia, Turkey. *Turkish Journal of Parasitology*, 19(4): 592–596.
- Dik, B., Güçlü, F., Cantoray, F., Cülbahçe, S. & Starý, J. (1999) Oribatid mites (Acari, Oribatida): Faunistic list, seasonal density and intermediate hosts of *Moniezia* sp. in province of Konya. *Turkish Journal of Veterinary and Animal Sciences*, 23(2): 385–391.
- Erman, O., Özkan, M., Ayyıldız, N. & Doğan, S. (2007) Checklist of the mites (Arachnida: Acari) of Turkey. Second supplement. *Zootaxa*, 1532: 1–21.
- Ghilarov, M.S. & Krivolutsky, D.A. (1975) *Identification keys of soil inhabiting mites, Sarcoptiformes*. Moscow, Nauka, 491 pp. (In Russian).
- Gjelstrup, P. & Solhøy, T. (1994) Oribatid mites. In: Jonasson, P.M. (Ed.), *The zoology of Iceland*, 3(57): 1–78.
- Gandjean, F. (1931) Le genre *Licneremaeus* Paoli (Acaiens). *Bulletin de la Société zoologique*, 56: 221–250.
- Gandjean, F. (1932) Observations sur les Oribates (3e serie). *Bulletin du museum national d'histoie naturelle, Paris*, 4(2): 292–306.
- Gandjean, F. (1936) Observations sur les Oribates (10e serie). *Bulletin du museum national d'histoie naturelle, Paris*, 2(8): 246–253.
- Grandjean, F. (1954) Observations sur les Oribates (28e serie). *Bulletin du museum national d'histoie naturelle, Paris*, 26(2): 204–211.
- Grandjean, F. (1965) Complement a mon travail de 1953 sur la classification des Oribatides. *Acarologia*, 7(4): 713–734.
- Grobler, L., Ozman, S.K. & Çobanoğlu S. (2003) The genera *Liacarus*, *Stenoxenillus* and *Xenillus* (Oribatida: Gustavioidea) from Turkey. *Acarologia*, 43(1): 131–147.
- Grobler, L., Bayram, Ş. & Çobanoğlu, S. (2004) Two new species and new records of oribatid mites from Turkey. *International Journal of Acarology*, 30(4): 351–358.
DOI: 10.1080/01647950408684405.
- Grobler, L., Bayram, Ş. & Çobanoğlu, S. (2005) Two new records of *Oribatula* (*Zygoribatula*) species (Acari: Oribatida) from Turkey with redescriptions. *Zoological Science*, 22(12): 1347–1351.
DOI: 10.2108/zsj.22.1347.
- Hammenn, van der L. (1952) The Oribatei (Acari) of the Netherlands. *Zoologische Verhandelingen*, 17: 1–139.
- Hammer, M. (1952) *Investigations on the microfauna of northern Canada. Part I. Oribatida*. Copenhagen, Ejnar Munksgaard, 108 pp.
- Hermann, F.L. (1804) Memoire apterologique. *Strasbourg, de l'imprimerie de F.G. Levraut*, 12: 87–95.

- Jacot, A.P. (1925) Phylogeny in the Oribatoidea. *The American Naturalist*, 59(662): 272–279.
- Jacot, A.P. (1929) Genera of Pterogasterine Oribatidae (Acarina). *Transactions of the American Microscopical Society*, 48(4): 416–430.
- Jacot, A.P. (1930) Oribatid mites of the subfamily Phthiracarinae of the Noartheastern United States. *Proceedings of the Boston Society of Natural History*, 39(6): 209–261.
- Koch, C.L. (1835) Deutschlands Crustaceen, Myriapoden und Arachiden. *Regensburg*, 1: 1–57.
- Koch, C.L. (1837) Deutschlands Crustaceen, Myriapoden und Arachiden. *Regensburg*, 10: 1–24.
- Koch, C.L. (1839) Deutschlands Crustaceen, Myriapoden und Arachiden. *Regensburg*, 24: 1–24.
- Koch, C.L. (1841) Deutschlands Crustaceen, Myriapoden und Arachiden. *Regensburg*, 32: 1–24.
- Koch, C.L. (1844a) Deutschlands Crustaceen, Myriapoden und Arachiden. *Regensburg*, 38: 1–24.
- Koch, C.L. (1844b) Deutschlands Crustaceen, Myriapoden und Arachiden. *Regensburg*, 39: 1–23.
- Kulijev, K.A. (1962) Fifteen new species of oribatid mites (Acariformes, Oribatei) of genera *Oppia* and *Ceratozetes*. *Trudy Azerbajianskogo Nauchno Issledovatel'skogo Instituta*, 13: 250–268 (In Russian).
- Kulijev, K.A. (1967) Two new genera of oribatid mites from Azerbaijan. *Dokladi Akademii Nauk Azerbajianskoi SSR*, 23: 85–91 (In Russian).
- Kumral, N.A. & Çobanoğlu, S. (2016) The mite (Acari) biodiversity and population fluctuation of predominant species in Eggplant. *Journal of Agricultural Science*, 22(2): 261–274.
- Lindo, Z. & Winchester, N.N. (2006) A comparison of microarthropod assemblages with emphasis on oribatid mites in canopy suspended soils and forest floors associated with ancient western redcedar trees. *Pedobiologia*, 50(1): 31–41.
DOI: 10.1016/j.pedobi.2005.09.002
- Mahunka, S. (1996) Oribatids of the Bükk National Park (Acari: Oribatida). In: Mahunka, S. (Ed.), *The Fauna of the Bükk National Park II*. Budapest: Hungarian Natural History Museum, pp. 491–532.
- Michael, A.D. (1879) A contribution to the knowledge of British Oribatidae. *Journal of the Royal Microscopical Society*, 12: 224–251.
- Michael, A.D. (1882) Further notes on British Oribatidae. *Journal of the Royal Microscopical Society*, 2: 1–18.
- Michael, A.D. (1888) *British Oribatidae. Vol. II*. Ray Society, London, pp. 337–657.
- Michael, A.D. (1898) Oribatidae. In: Lohmann, H. (Ed.), *Das Tierreich. Eine Zusammenstellung und Kennzeichnung der rezenten Tierformen*. Berlin, Friedlander und Sohn, pp. 1–93.
- Mihelčíč, F. (1955) Oribatiden Südeuropas. *Zoologischer Anzeiger*, 155(9–10): 244–248.
- Mihelčíč, F. (1956a) Oribatiden Südeuropas IV. *Zoologischer Anzeiger*, 156(7/8): 205–226.
- Mihelčíč, F. (1956b) Oribatiden Südeuropas V. *Zoologischer Anzeiger*, 157(7/8): 154–174.
- Murvanidze, M. & Arabuli, T. (2015) Oribatid mite diversity in *Rhododendron ponticum* L. canopy along an altitudinal gradient in Mtirala National Park. *Acarologia*, 55(2): 219–230.
DOI: 10.1051/acarologia/20152162
- Murvanidze, M., Çobanoglu, S. & Cilbircioğlu, C. (2018) New record of *Oribatula* (*Zygoribatula*) *nicora* comb. nov. from Turkey with a redescription of species. *Turkish Journal of Zoology*, 42: 601–607.
DOI: 10.3906/zoo-1806-24.
- Murvanidze, M. & Mumladze, L. (2014) Oribatid mite (Acari: Oribatida) diversity in different forest stands of Borjomi-Kharagauli National Park (Georgia). *Persian Journal of Acarology*, 3(4): 257–276.
DOI: 10.22073/pja.v3i4.10169
- Murvanidze, M. & Mumladze, L. (2016) Annotated checklist of Georgian oribatid mites. *Zootaxa*, 4089(1): 1–81.
DOI: 10.11646/zootaxa.4089.1.1.

- Murvanidze, M., Mumladze, L., Arabuli, T., Barjadze, Sh. & Salakaia, M. (2016) Oribatida diversity in different microhabitats of Mtirala National Park. *Journal of Acarological Society of Japan*, 25(S1): 35–49.
- Murvanidze M. & Weigmann G. (2003) Contribution to the oribatid mite fauna of Georgia. 1. New species of Poronota (Acari, Oribatida). *Spixiana*, 26(2): 165–170.
- Niedbala, W. & Liu, D. (2018) Catalogue of ptyctimous mites (Acari, Oribatida) of the world. *Zootaxa*, 4393(1): 1–238.
DOI: 10.11646/zootaxa.4393.1.1
- Nicolet, M.H. (1855) Histoire naturelles des Acariens qui se trouvent aux environs de Paris. *Archives du museum d'histoire naturelle*, 7: 381–482 (In French).
- Ocak, I., Doğan, S., Ayyıldız, N. & Hasenkoglu, I. (2007) The external mycoflora of the oribatid mites (Acari) in Turkey, with three new mite records. *Archives des Sciences*, 61(1): 1–6.
- Oudemans, A.C. (1900) Further notes on Acari. *Tijdschrift voor Entomologie*, 43: 109–128.
- Oudemans, A.C. (1923) Studie over de sedert 1877 ontworpen systemen de Acari; niuwe classificatie; phylogenetische beschouwingen. *Tijdschrift voor Entomologie*, 66: 4–33 (In Dutch).
- Özkan, M., Ayyıldız, N. & Erman, O. (1994) Checklist of the Acari of Turkey. First supplement. *EURAAC Newsletter*, 6: 4–12.
- Paschoal, A.D. & Jonston, D.E. (1985) Aleurodamaeidae (Acari: Oribatei), a new family of oribatid, with a description of *Aleurodamaeus hungaricus*, sp. n. *Revista Brasileira de Entomologia*, 29(1): 21–26.
- Paoli, G. (1908) Monographia del genre *Damaeosoma* Berl. A generi affini. *Redia*, 5: 1–90.
- Per, S. & Ercan, F.S. (2015) Comparison of three methods of DNA extraction from *Parachipteria willmanni* (Acari: Oribatida) collected in Turkey. *Journal of Biotech Research*, 1(4): 16–20.
- Per, S., Toluk, A. & Ayyıldız, N. (2018) A new record for the Turkish oribatid mite (Acari) fauna from Yozgat Province: *Eremulus flagellifer* Berlese, 1908. *International Ecology Symposium*, 19–23 June 2018, Kastamonu, Turkey, p. 701.
- Perez-Iñigo, C. (1993) Acari: Oribatei: Poronota. In: Ramos, M.A. (Ed.), *Fauna Iberica. Vol. 3. Museo Nacional de Ciencias naturales*, Madrid, pp. 1–320.
- Perty, M. (1841) Allgemeine Naturgeschichte, als philosophische und Humanitätswissenschaft für Naturforscher, Philosophen und das höher gebildete Publikum, 3: 873–875.
- Petrunkewitch, A. (1955) Arthropoda 2. Chelicerata with sections on Pycnogonida and Palaeoisopus. In: Moore, R.C. (Ed.), *Treatise on invertebrate Palaeontology*. Geological Society of America and University of Kansas Press, pp. 97–99.
- Prinzing, A. (1997) Spatial and temporal use of microhabitats as a key strategy for the colonization of tree bark by *Entomobrya anivalis* L. (Collembola: Entomobryidae). In: Stork, N.E., Adis, J. & Didham, R.K. (Eds.), *Canopy arthropods*. Chapman & Hall, pp. 453–476.
- Proctor, H.C., Montgomery, K.M., Rosen, K.E. & Kitching, R.L. (2002) Are tree trunks habitats or highways? A comparison of oribatid mite assemblages from hooppine bark and litter. *Australian Journal of Entomology*, 41: 294–299.
DOI: 10.1046/j.1440-6055.2002.00309.x.
- Schatz, H., Behan-Pelletier, V.M., OConnor, B.M. & Norton, R.A. (2011) Suborder Oribatida van der Hammen, 1968. In: Zhang, Z.-Q. (Ed.), *Animal biodiversity: An outline of higher level classification and survey of taxonomic richness*. Zootaxa, 3148: 141–148.
- Schütt, H. (2005) *The Turkish land snails, 1758–2000: Vollständig Revidierte und Erweiterte Auflage*. Nature und Wissenschaft, 559 pp.
- Sellnick, M. (1908) Die Tardigraden und Oribatiden der ostpreussischen Moosrasen. *Schriften der Physikalisch-Ökonomischen Gesellschaft*, 49: 317–350.
- Sellnick, M. (1928) Formenkreis: Hornmilben, Oribatei. In: Brohmer, P., Ehrman, P. & Ulmer, G. (Eds.), *Die Tierwelt Mitteleuropas*. Leipzig, Quelle und Meyer, Vol. 3, Part 9, pp. 1–42.

- Seniczak, A., Seniczak, S., Kaczmarek, S. & Kowalski, J. (2014) Ontogeny of morphological traits in Phenopelopidae (Acari: Oribatida). *International Journal of Acarology*, 40(8): 611–637.
DOI: 10.1080/01647954.2014.975744.
- Shtanchaeva, U.Ya. & Subias, L.S. (2010) *The catalogue of Caucasian oribatid mites*. Nauka, Makhachkala, 276 pp. (In Russian).
- Sobek, S., Kampichler, Ch. & Weigmann, G. (2008) Oribatid mites (Acari: Oribatida) in the canopy of a Central European mixed forest: Species richness and species similarity between tree species and habitat types. In: Floren, A. & Schmidl, J. (Eds.), *Canopy arthropod research in Europe*. Nuremberg, Bioform Entomology, pp. 339–354.
- Subías, L.S. (2004 updated in 2020) Listado sistemático, sinonímico y biogeográfico de los oribatidas (Acariformes, Oribatida) del mundo (1758–2002). 2004. *Graelsia*, 60 (numero extraordinario) Available from: <http://www.ucm.es/info/zoo/Artropodos/Catalog.pdf>. (Access-ed on 21 February 2020).
- Subías, L.S. & Rodriguez, P. (1985) Oppiidae (Acari, Oribatida) de los sabinares (*Juniperus thurifera*) de España III. *Graptoppia* a. str. Balogh. *Cuadernos de Investigación Biológica, Bilbao*, 8: 69–76.
- Susyal, B., Ayyildiz, N. & Baran, Ş. (2018) A new species of *Eupterotegaeus* (Acari, Oribatida, Compactozetidae) from Turkey. *Entomological News*, 128(1): 65–71.
DOI: 10.3157/021.128.0109.
- Thor, S. (1929) Über die phylogenie und systematik der acarina. Mit meiträgen zur ersten entwicklunsgeschichte einzelner gruppen. *Nyt Magazine for Naturvidenskaberne*, 67: 145–210.
- Thorell, T. (1876) Sopra alcuni Opilioni (Phalangidea) d'Europa e dell' Asia occidentale, con un quadro dei generi europei di quest' Ordine. *Annali del museo civico di storia naturale di Genova*, 8: 1–155.
- Toluk, A. & Ayyıldız, N. (2008) The systematic studies on oppiid oribatid mites of Yozgat pine grove national park (Acari: Oribatida). *Erciyes Üniversitesi Fen Bilimleri Enstitüsü Fen Bilimleri Dergisi*, 24(1–2): 52–81.
- Toluk, A., Ayyıldız, N. (2008a) Ali Dağı' ndan (Kayseri) kaydedilen iki ilkel Oribatid akar: *Sphaerochthonius splendidus* (Berlese, 1904) ve *Epilohmannia cylindrica*. *Erciyes Üniversitesi Fen Bilimleri Enstitüsü Fen Bilimleri Dergisi*, 24(1–2): 101–111 (In Turkish).
- Toluk, A. & Ayyıldız, N. (2011) Contributions to the Turkish oribatid fauna (Acari: Oribatida) from Bolu Province. *Turkish Journal of Zoology*, 35(1): 63–70.
DOI: 10.3906/zoo-0905-1.
- Toluk, A. & Ayyıldız, N. (2016) New records of the genus *Carabodes* (Acari, Oribatida, Carabodidae) from Turkey. *8th Symposium of the European Association of Acarologists (EURAAC 2016)*, 11–15 July, Valencia, p. 78.
- Toluk, A. & Akin, A.T. (2017) Oribatid mite fauna (Acari) of Çat Forest, Sivas Province, Turkey. *Turkish Journal of Entomology*, 41: 293–307.
- Trägårdh, I. (1910) Acariden aus dem Sarekgebirge. In: Hamberg, A. (Ed.), *Naturwissenschaftliche Untersuchungen Sarekgebirges in Schwedisch-Lappland*. Berlin, Fiedländer & Sohn, pp. 375–586 (In German).
- Trägårdh, I. (1931) Acarina from the Juan Fernan-dez Islands. In: Skottsberg, C. (Ed.), The natural history of Juan Fernandez and Easter Island. *Uppsala*, 3(4): 553–628.
- Travé, J. (1960) Contribution à l'étude de la faune de la Massane. (3^e note) Oribates (Acarines), 2^e Partie I. *Vie et Milieu*, 11(2): 209–233 (In French).
- Weigmann, G. (2006) Hornmilben (Oribatida). *Die Tierwelt Deutschlands. 76 Teil*. Goecke and Evers, Keltern, 520 pp.
- Williams, L., Zazanashvili, N., Sanadiradze, G. & Kandaurov, A. (2006) *An ecoregional conservation plan for Caucasus*. WWF, Brussels, Belgium, 220 pp.

- Willmann, C. (1951) Untersuchungen über die terrestrische Milbenfauna im pannonicischen Klimagebiet Österreichs. *Sitzungsberichte Österreichischen Akademie der Wissenschaften fuer Mathematik und Naturwissenschaften*, 160(1/2): 91–176.
- Winchester, N.N., Behan-Pelletier, V.M. & Ring, R.A. (1999) Arboreal specificity, diversity and abundance of canopy-dwelling oribatid mites (Acari: Oribatida). *Pedobiologia*, 43: 391–400.
- Winchester, N.N., Lindo, Z. & Behan-Pelletier, V.M. (2008) Oribatid mite communities in the canopy of montane *Abies amabilis* and *Tsuga heterophylla* trees on Vancouver Island, British Columbia. *Environmental Entomology*, 37(2): 464–471.
DOI: 10.1093/ee/37.2.464.
- Wunderle, I., Beck, L. & Woas, S. (1990) Ein Beitrag zur Taxonomie und Okologie der Oribatulidae und Scheloribatidae (Acari, Oribatei) in Südwestdeutschland. *Andrias*, 7: 15–60.
- Yaça, M. & Baran, Ş. (2019) Oribatid mite fauna of Kocaeli City Forest (Kocaeli, Turkey). *Turkish Journal of Entomology*, 43(1): 41–56.
DOI: 10.16970/entoted.508099.
- Yalçın, S., Doğan, S. & Ayyıldız, N. (2013) Some oribatid mites living in Uzunoluk forest (Erzurum) and microfungi isolated from them. *Turkish Journal of Entomology*, 37(1): 117–131.
- Zeytun, E., Doğan, S. & Ayyıldız, N. (2017) Two newly recorded species of oribatid mites (Acari, Oribatida) for the Turkish fauna, from the house dust. *The 3rd International Symposium on EuroAsian Biodiversity, 05–08 July 2017*, Minsk, Belarus, p. 314.
- Zeydanli, U. (2020) Western Asia: Northern Turkey extending along the southern Black Sea region. Available from: <https://www.worldwildlife.org/ecoregions/pa0515> (Accessed on 5 February 2020).

COPYRIGHT

 Murvanidze et al. Persian Journal of Acarology is under a free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

چکلیست کنه‌های اریباتید (Acari: Oribatida) حوزه مرکزی دریای سیاه ترکیه همراه با گزارش‌های جدید برای آن کشور

مکه مروانیدزه^{۱*}، سیهان سیلیبر سیو غلو^۲، اسنگول او زدمیز^۳ و امره اینک^۳

- ۱- موسسه حشره‌شناسی، دانشگاه کشاورزی گرجستان، ۰۱۳۱، ۰۲۴۰، کوچه دی آقمانشبلی، تفلیس، گرجستان؛ رایانامه: m.murvanidze@agruni.edu.ge
- ۲- گروه کشاورزی ارگانیک، دانشکده وکیشنال تاشکورپو، دانشگاه کاستامونو، کاستامونو، ترکیه؛ رایانامه: cihancilbirci@hotmail.com
- ۳- گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه آنکارا، آنکارا، ترکیه؛ رایانامه: esengul.ozdemir@ankara.edu.tr

* نویسنده مسئول

چکیده

پنجاه و دو گونه/زیرگونه از کنه‌های اریباتید از مناطق جنگلی حوزه مرکزی دریای سیاه و آناتولیای مرکزی شناسایی شدند. هفده گونه از بین آنها برای ترکیه گزارش جدید محسوب می‌شوند. برای هر گونه، اطلاعات جغرافیایی ثبت شده با جی‌پی‌اس، زیستگاه، پراکندگی منطقه‌ای و جهان تهیه و برای گزارش‌های جدید بحث‌های آرایه‌شناختی، اکولوژیک و/یا ویژگی‌های منطقه‌ای ارایه شده است.

واژگان کلیدی: تنوع زیستی؛ جنگل؛ گزارش‌های جدید؛ فون کنه‌های ترکیه؛ جغرافیای جانوری.

اطلاعات مقاله: تاریخ دریافت: ۱۳۹۹/۲/۱، تاریخ پذیرش: ۱۳۹۹/۲/۳۱، تاریخ چاپ: ۱۳۹۹/۴/۲۵