

# *Pseudostellaria wuyishanensis*, a new species of Caryophyllaceae from Fujian, China

Xiao Luo<sup>1,2\*</sup>, Qi-Yi Yang<sup>2\*</sup>, Zhe Zhang<sup>3</sup>, Pan Zhu<sup>4</sup>, Liang Ma<sup>5</sup>,  
Xin-Yan Chen<sup>6</sup>, Shu-Yi Lin<sup>7</sup>, Shi-Pin Chen<sup>2</sup>

**1** Fujian Provincial Forestry Survey and Design Institute, Fuzhou 350002, China **2** College of Forestry, Fujian Agriculture and Forestry University, Fuzhou 350002, China **3** College of Forestry, Southwest Forestry University, Kunming 650224, China **4** College of Horticulture and Landscape Architecture, Southwest University, Chongqing 400700, China **5** Key Laboratory of National Forestry and Grassland Administration for Orchid Conservation and Utilization at College of Landscape Architecture, Fujian Agriculture and Forestry University, Fuzhou 350002, China **6** Sanming City Garden Center, Sanmin 365000, China **7** Middlebury College, Vermont 05753, USA

Corresponding author: Shi-Pin Chen ([fjcsp@126.com](mailto:fjcsp@126.com))

Academic editor: G. P. Giusso del Galdo | Received 16 April 2021 | Accepted 16 July 2021 | Published 25 August 2021

**Citation:** Luo X, Yang Q-Y, Zhang Z, Zhu P, Ma L, Chen X-Y, Lin S-Y, Chen S-P (2021) *Pseudostellaria wuyishanensis*, a new species of Caryophyllaceae from Fujian, China. *PhytoKeys* 181: 21–28. <https://doi.org/10.3897/phytokeys.181.67436>

## Abstract

*Pseudostellaria wuyishanensis*, a new species from the Wuyishan National Park, Fujian, China, is described and illustrated. Morphologically, *Pseudostellaria wuyishanensis* resembles *P. heterantha*. However, the new species can be distinguished by presence of stolons, 1 line of hairs on the stem, smaller leaf blades, shorter pedicels, and ovary with 2 styles.

## Keywords

Caryophyllaceae, Fujian, *Pseudostellaria*, Wuyishan National Park

## Introduction

*Pseudostellaria* Pax is a small genus that belongs to the tribe Alsineae in Caryophyllaceae (Bittrich 1993; Tang et al. 1996). This genus can be easily distinguished from other genera in Caryophyllaceae from the presence of the flesh root tuber. In addition, the vast majority

\* These authors contributed equally to this work

of species in the genus have cleistogamous flower and chasmogamous flowers that have petals with two sections (Zeng et al. 2016). Some recent molecular studies show that this group is non-monophyletic which includes a new described genus *Hartmaniella* and 2 species *Stellaria americana* (Porter & B.L.Rob.) Standl. and *Arenaria przewalskii* Maxim. nested within *Pseudostellaria* (Greenberg and Donoghue 2011; Zhang et al. 2017). Russian botanist Turczaninow (1842) first used *Krascheninikovia* Turcz. ex Fenzl for this genus, but this name was in fact a previous synonym for *Eurotia* Adans and did not comply with the international nomenclature regulations. *Pseudostellaria* was established by Pax in 1934 as the new name, which has been used until now (Schischkin and Komarov 1936; Ohwi 1937; Mizushima 1965). Currently, the genus is represented by ca.22 accepted species that are widely distributed all over the world, with 20 species in eastern and northern Asia, 1 species in Europe, and 1 species in North America. (Zeng et al. 2016; Zhang et al. 2017).

Since the turn of the 21<sup>st</sup> century, 3 new species of *Pseudostellaria* have been established in China. Jin and Ding (2003) described *P. zhejiangensis* X.F Jin & B.Y Ding from the Zhejiang province based on its decumbent creeping stems, obtuse petals, and compressed seeds with a narrow wing. Lian (2009) described *P. polymorpha* W. Z. Di & Y. Ren based on the regular variation in its floral morphology from stem apex to base. Xia et al. (2011) described *P. tianmushansis* Xia et al. based on its several tubers in a row, obovate with a bi-lobed apex petal and tubercles awned seeds. 12 species of *Pseudostellaria* have been recorded in China out of which 5 species are endemic.

During an investigation of wild plants in Fujian Province, southeastern China, that took place in May 2019 and October 2020, an unknown species of *Pseudostellaria* was collected from the deciduous broad-leaved forest in Wuyishan National Park. We found that it resembles *P. heterantha* Pax but has stolons, 1 line of hairs in the stem, smaller leaves, and shorter pedicels. Therefore, we established it as a new species.

## Material and methods

All general morphological data were obtained by observation of specimens during field-works and AU, FJIDC, IBSC, KUN, LE herbaria. Terminologies used in the present study follows the *Flora of China* (Lu and Rabeler 2001) and additional consultation of online databases, including Chinese Field Herbarium and Plant Photo Bank of China.

## Taxonomy

***Pseudostellaria wuyishanensis* X. Luo & Q.Y. Yang, sp. nov.**


urn:lsid:ipni.org:names:77219367-1

Figs 1, 2

**Type.** China. Fujian: Wuyishan National Park, on rocks along a stream, ca.1700 m a.s.l, 1 May 2019, Xiao Luo et Qiyi Yang20190501 (**holotype:** FAFU!; **isotype:** FAFU!)


**Table 1.** Morphological comparison of *Pseudostellaria wuyishanensis*, *P. heterantha*.

Characters	<i>P. wuyishanensis</i>	<i>P. heterantha</i>
Stem	has stolons, branched at apex, 6–7 cm tall, with 1 line of hairs	no stolon, branched at base, 8–15 cm tall, with 2 lines of hairs
Leaf blade	1–1.6 × 0.5–0.7 cm	2–2.5 × 0.8–1.2 cm
Pedicel	ca. 2 cm long	3–3.5 cm long
Sepal	abaxially pilose, margin glabrous	abaxially pilose, margin ciliate
Ovary	2 styles	2 or 3 styles
Distribution	Fujian (Southeastern China)	Northern and Southwestern China

**Figure 1.** *Pseudostellaria wuyishanensis* sp. nov. **A** plant **B** stem with one line of hair **C** flower **D** flower in side view **E** calyx **F** petal **G** tuber **H** gynoecium of chasmogamous flower.

**Diagnosis.** *Pseudostellaria wuyishanensis* can be distinguished from *P. heterantha* by several morphological features and distribution (Table 1). *P. wuyishanensis* has stolons (vs. no stolon in *P. heterantha*), 1–1.6 × 0.5–0.7 cm (vs. 2–2.5 × 0.8–1.2 cm in *P. heterantha*) leaf blade, ca. 2 cm long (vs. 3–3.5 cm long in *P. heterantha*) pedicel and is 6–7 cm tall with 1 line of hairs (vs. 8–15 cm tall with 2 lines of hairs in *P. heterantha*). *P. wuyishanensis* only distribute in Wuyishan National Park, Southeastern China (vs. Northern and Southwestern China in *P. heterantha*)

**Description.** Plants perennial. Root tubers green, fusiform, 0.4–0.6 × 0.2–0.3 cm. Stem erect, 6–7 cm tall, slender, unbranched at base, apex false dichotomous branched, stoloniferous, with 1 line of hairs. Leaves opposite, entire, 1–1.6 × 0.5–0.7 cm; proximal middle leaves oblanceolate, base attenuate into a petiole, apex acute; distal leaves ovate, shortly petiolate, membranous, both surfaces glabrous, the adaxial green, the



**Figure 2.** *Pseudostellaria wuyishanensis* X. Luo, Q.Y. Yang **A** habitat **B** habit **C** tuber **D** plant **E** flowering plant **F** flower **G** all parts of flower.

abaxial viridescent, apex acute, usually with mucro ca.0.5 mm, sparsely ciliate at base, pinnately veined, lateral veins 3–4 pairs, inconspicuous. Chasmogamic flowers terminal or axillary, solitary; pedicel erect, ca. 2 cm long, pilose; sepals 5, green, lanceolate, ca. 3 mm, abaxially slightly pilose, margin membranous, glabrous; petal 5, oblong, slightly longer than sepals, ca. 4 mm, apically emarginate, base with a short claw;

stamens 10, shorter than petals, ca. 4 mm; filament glabrous; anthers purple-red, reniform; ovary coniform, ca.  $2 \times 0.9$  mm, with 2 thin styles to 3 mm, revolute, longer than the ovary, ovules numerous. Cleistogamous flowers and fruits not seen.

**Distribution and habitat.** The new species is endemic to the Wuyishan National Park, Fujian Province. The plant grows in the deciduous broad-leaved forest at 2000 m in elevation. The dominant species of the community include *Pinus taiwanensis* Hayata (Pinaceae), *Lithocarpus harlandii* (Hance) Rehder (Fagaceae), *Buxus sinica* var. *parvifolia* M. Cheng (Buxaceae), *Veratrum schindleri* Loes (Melanthiaceae), and *Dichocarpum franchetii* (Finet&Gagnepain) W.T. Wang & Hsiao (Ranunculaceae).

**Phenology.** Flowers were observed in June.

**Conservation status.** There is only one known location and fewer than 50 individuals of *P. wuyishanensis* found during our fieldworks in the Wuyishan National Park in both 2019 and 2020. But the investigation has not been through enough to fully understand the species natural distribution. According to IUCN Red List criteria (2012), this new species should be assessed as Data Deficient (DD; criteria B1ab(i-v) + 2ab(i-v)).

**Etymology.** The specific epithet ‘*wuyishanensis*’ refers to Wuyishan National Park, the locality of the type collection.

## Discussion

The new species morphologically resembles *P. heterantha* in the leaf shape, terminal chasmogamous flowers with pilose pedicel, and white emarginate petal. The two taxa differ in that the stem of *P. wuyishanensis* is shorter, conspicuously stoloniferous, apex false dichotomous branched, and only has 1 line of hairs, while that of *P. heterantha* is longer, solitary, branched at base, and has 2 line of hairs; the leaf blades of *P. wuyishanensis* is smaller and the pedicel is shorter.(Table 1).

Ohwi (1937) regarded *P. maximowicziana* (Franch. & Sav.) Pax and *P. himalaica* (Franchet) Pax as the synonym of *P. heterantha*. The view was also approved by Mizushima (1965) and Lu (1998). However, some research results published in recent years do not support such a view (Chen et al. 2014; Zeng et al. 2016; Zhang et al. 2017). Zeng et al. (2016) suggested taking *P. maximowicziana*, *P. himalaica*, and *P. heterantha* as independent species respectively. None of the 3 species were collected in Fujian province or the surrounding area.

We only found two *Pseudostellaria* sp. specimens, IBSC 0149273 and IBSC 0149274 (Fig. 3) were collected in the Wuyishan National Park. Former researchers have identified them as *P. rupestris* (Turczaninow) Pax or *P. heterophylla* (Zeng et al. 2016). Morphologically, we found that the arrangement, shape and hairs of the leaves of these specimens were completely different from those two species, and the morphology of each part was consistent with *P. wuyishanensis*. In addition, the distribution location of *P. rupestris* was far away from the collection site. Considering all the factors, we believe that these specimens are in fact *P. wuyishanensis*.




Figure 3. Specimens of *Pseudostellaria wuyishanensis* collected in 1980s.

Key to the Chinese species of *Pseudostellaria*

- 1 Seeds with persistent anchor-shaped barb ..... *Pseudostellaria rupestris*
- Seeds with awned tubercles ..... 2
- 2 Stems with apical 2 pairs of leaves larger, approximate, decussate ..... *P. heterophylla*
- Stems not as above ..... 3
- 3 Chasmogamic flowers with petals apex 2-lobed ..... 4
- Chasmogamic flowers with petals apex entire, sometimes emarginate ..... 7
- 4 Root tubers several in a row ..... 5
- Root tubers solitary ..... 6
- 5 All leaves linear or lanceolate-linear, sessile ..... *P. sylvatica*
- All leaves narrow elliptic-lanceolate, with short petiole .... *P. tianmushanensis*
- 6 Chasmogamic flowers with sepals 4, petals 4, stamens 8 .... *P. helanshanensis*
- Chasmogamic flowers with sepals 5, petals 5, stamens 10 ..... *P. japonica*
- 7 Chasmogamic flowers with sepals glabrous ..... *P. tibetica*
- Chasmogamic flowers with sepals abaxially pubescent ..... 8
- 8 Stem repent ..... 9
- Stem erect ..... 10

- 9 Leaves pubescent in both side; seeds flat, with narrow wings.....*P. zhejiangensis*  
 .....*P. zhejiangensis*
- Leaves ciliate; seeds reniform or subglobose ..... *P. davidii*
- 10 Stem pubescent; Leaves both surfaces pubescent ..... *P. himalaica*
- Stems with 1 or 2 line of hairs; base of the leaves sparsely ciliate ..... 11
- 11 Stem has stolons, with 1 line of hairs; pedicel short, ca. 2 cm.....  
 ..... *P. wuyishanensis* (sp. nov.)
- Stem has 2 line of hairs; pedicel longer than 3 cm..... 12
- 12 Petals spatulate or obovate ..... *P. maximowicziana*
- Petals oblong-oblancoate ..... *P. heterantha*

## Acknowledgements

We thank the reviewers for their extensively helpful comments. This study was supported by the National Special Fund for Chinese medicine resources Research in the Public Interest of China (No. 2019-39) and Wuyishan forest germplasm resources investigation[[350782]FJXW[GK]2018004].

## References

- Bittrich V (1993) Caryophyllaceae. In: Kubitzki K, Rohwer JG, Bittrich V (Eds) The families and genera of vascular plants. Springer, Berlin, 206–236. [https://doi.org/10.1007/978-3-662-02899-5\\_21](https://doi.org/10.1007/978-3-662-02899-5_21)
- Chen XB, Meng SY, Zhang XF, Han YB, Liu QR (2014) Numerical Taxonomic Analysis of *Stellaria* and *Pseudostellaria* (Caryophyllaceae). *Zhiwu Xuebao* 49(4): 432–439. <https://doi.org/10.3724/SP.J.1259.2014.00432>
- Greenberg AK, Donoghue MJ (2011) Molecular systematics and character evolution in Caryophyllaceae. *Taxon* 60(6): 1637–1652. <https://doi.org/10.1002/tax.606009>
- IUCN (2012) IUCN Red List Categories and Criteria, Version 3.1. 2<sup>nd</sup> edn., IUCN, Gland.
- Jin XF, Ding BY (2003) A new species of *Pseudostellaria* (Caryophyllaceae) from Zhejiang. *Yunnan Zhi Wu Yan Jiu* 25(6): 639–640.
- Lian YS (2009) A new species of *Pseudostellaria* (Caryophyllaceae) from Gansu, China. *Novon* 19(2): 191–193. <https://doi.org/10.3417/2004104>
- Lu DQ (1998) Some revisions and supplements on Caryophyllaceae from China. *Bulletin of Botanical Research* (1): 1–3.
- Lu DQ, Rabeler RK (2001) *Pseudostellaria*. In: Wu ZY, PH Raven, Hong DY (Eds) *Flora of China*. St. Louis: Science Press, Missouri Botanical Garden Press, Beijing, 7–10.
- Mizushima M (1965) Critical studies on Japanese plants, II: The genus *Pseudostellaria* Pax in Japan. *Bulletin of the Botanical Survey of India* 7: 62–72.
- Ohwi J (1937) The Revision of the genus *Pseudostellaria*. *Shokubutsu Kenkyu Zasshi* 9: 95–105.

- Schischkin BK, Komarov VL (1936) Flora of the USSR, vol. 6. Akademiya Nauk SSSR, Moskva-Leningrad, 326–330.
- Tang CL, Ke P, Lu DQ, Zhou LH, Wu ZY (1996) Caryophyllaceae. In: Tang CL (Ed.) Flora of China. St. Louis: Science Press, Missouri Botanical Garden Press, Beijing, 47–448.
- Turczaninow NS (1842) Krascheninnikowia. Bulletin de la Société Impériale des Naturalistes de Moscou 15: 609.
- Xia GH, Liu CH, Xie WY, Li GY (2011) *Pseudostellaria tianmushanensis* sp. nov. (Caryophyllaceae) from Zhejiang, China. Nordic Journal of Botany 29(2): 204–207. <https://doi.org/10.1111/j.1756-1051.2010.00933.x>
- Zeng XQ, Zhang ML, Lei Y (2016) Classification outline and geographical distribution of *Pseudostellaria* Pax in the world. Journal of Plant Resources and Environment 25: 92–99.
- Zhang ML, Zeng XQ, Li C, Sanderson SC, Byalt VV, Lei Y (2017) Molecular phylogenetic analysis and character evolution in *Pseudostellaria* (Caryophyllaceae) and description of a new genus, *Hartmaniella*, in North America. Botanical Journal of the Linnean Society 184(4): 444–456. <https://doi.org/10.1093/botlinnean/box036>