


Acarologia



A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr



**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2021 (Volume 61): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2020): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)



Acarologia is under **free license** and distributed under the terms of the Creative Commons-BY.

Mites on Annonaceae species in northeast Brazil and in the state of Pará

Josilene Maria de SOUSA^{1*}, Manoel G.C. GONDIM JR.¹, Antônio Carlos LOFEGO²
and Gilberto José de MORAES³

(Received 08 August 2014; accepted 07 November 2014; published online 30 March 2015)

¹ Departamento de Agronomia, Área Fitossanidade, Universidade Federal Rural de Pernambuco. Av. Dom Manoel de Medeiros s/n, Dois Irmãos, 52171-900 Recife, Pernambuco, Brasil. podos@ig.com.br, mguedes@depa.ufrpe.br (* Corresponding author)

² UNESP – Universidade Estadual Paulista, Departamento de Zootecnia e Botânica, Rua Cristóvão Colombo, 2265, 15054-000, São José do Rio Preto, SP, Brasil. aclofego@ig.com.br

³ Departamento de Entomologia, Fitopatologia e Zoologia Agrícola, Escola Superior de Agricultura Luiz de Queiroz, 13418-900 Piracicaba, São Paulo, Brasil. gjmoraes@carpa.ciagri.usp.br

ABSTRACT — The family Annonaceae Juss. comprises over 120 genera and 2000 species with origins in the American, Asian and African tropical regions. Many of these species, such as those of the genera *Annona*, *Rollinia*, *Duguetia*, *Uvaria* and *Asimira*, are edible. Some pests have been reported in soursop, sugar apple and atemoya, the main annonas grown in Brazil. However, few mites are mentioned as pests of this plant family. This study aimed to identify mite species associated with the Annonaceae in northeastern Brazil and in the state of Pará. A survey was conducted, collecting samples of *Annona muricata* L. (soursop), *Annona squamosa* L. (sugar apples), *Annona cherimola* × *A. squamosa* (atemoya) and *Annona coriacea* Mart. (araticum) leaves. *Annona coriacea* is a wild Annonaceae widely distributed in northeastern Brazil. For each sample, a total of 100 leaves were collected from five individuals of the same species at each site. The samples were taken to the laboratory, where they were processed and the mites were mounted and identified. The mite species found belonged to the families Ascidae, Bdellidae, Phytoseiidae, Stigmeidae, Tarsonemidae, Tetranychidae, Tenuipalpidae and Tydeidae. Phytoseiidae, Tetranychidae and Tydeidae had the highest diversity. Five of the reported species were found on all four of the studied cultures: *Amblyseius aerialis* (Muma), *Tetranychus mexicanus* (McGregor), *Parapronematus acaciae* Baker, *Pronematus ubiquitus* (McGregor) and *Agistemus floridanus* González.

KEYWORDS — Acari; diversity; mites; *Annona muricata*; *Annona squamosa*

INTRODUCTION

The family Annonaceae Juss. comprises over 120 genera and 2000 species with origins in the American, Asian and African tropical regions. Many of these species are of interest for their edible fruits and are distributed among the genera *Annona*, *Rollinia*, *Duguetia*, *Uvaria* and *Asimira*. However, only the first two, especially the *Annona*, are of economic importance (Mabberley 1997; Nogueira

et al. 2005). The genus *Annona* has approximately 90 species, and its main representatives are *Annona reticulata* L. (wild sweetsop), *Annona muricata* L. (soursop), *Annona squamosa* L. (sugar apples), *Annona cherimola* Mill. (cherimoya) and the hybrid *A. cherimola* × *A. squamosa* (atemoya). Commercial Annonaceae cultivation occurs regionally according to the climatic requirements of each species. The Northeast of Brazil is the main producing region of Annonaceae, especially soursop and sugar apple

(Leon 1987; Sobrinho *et al.* 1998; Cardoso and Sousa 2000; Nogueira *et al.* 2005).

Several pests, especially insects, are reported in annonas grown in Brazil. However, few pest mites, such as *Oligonychus anoneae* Paschoal (Tetranychidae) and *Aculops flechtmanni* Keifer (Eriophyidae), have been found to date; both of these species were described in Southeast Brazil (Flechtmann 1985). This study aimed to describe the existing

mite fauna in the main cultivated and native Annonaceae species in Northeast Brazil and the state of Pará.

MATERIALS AND METHODS

The study was conducted at the laboratory of Agricultural Acarology in the Agronomy Department of the Federal Rural University of Pernambuco (Uni-

TABLE 1: State, county, coordinates and host plants.

| State | County | Coordinates | Host plants |
|---------------------|----------------------|---------------------------|--|
| Alagoas | Miai de Baixo | 36°13'59.7"W 10°13'59.7"S | <i>A. muricata</i> |
| Bahia | Juazeiro | 40°24'43.6"W 09°23'19.8"S | <i>A. squamosa; A. muricata</i> <i>A. cherimola x A. squamosa</i> |
| | Una | 39°03'51.2"W 15°17'13.8"S | <i>A. muricata</i> |
| Ceará | Acaraú | 40°07'18.5"W 02°52'19.6"S | <i>A. squamosa</i> |
| | Barroquinha | 41°08'22.3"W 03°01'14.2"S | <i>A. muricata</i> |
| Maranhão | Paraipaba | 39°13'49.2"W 03°27'49.4"S | <i>A. muricata</i> |
| | Junco do Maranhão | 46°03'01.7"W 01°44'52.9"S | <i>A. muricata</i> |
| Pará | Viana | 44°59'36.1"W 03°12'11.8"S | <i>A. squamosa</i> |
| | Tutóia | 42°22'43.0"W 02°47'52.7"S | <i>A. squamosa</i> |
| Pernambuco | Rosário | 44°13'09.5"W 02°57'09.6"S | <i>A. squamosa</i> |
| | Capanema | 47°12'33.6"W 01°00'23.0"S | <i>A. squamosa</i> |
| Paraíba | Castanhal | 47°56'57.1"W 01°18'03.2"S | <i>A. muricata</i> |
| | Água Preta | 35°31'51.0"W 08°42'25"S | <i>A. squamosa ; A. muricata</i> |
| Rio Grande do Norte | Bonito | 35°35'00.0"W 08°40'00"S | <i>A. squamosa; A. muricata</i> <i>A. cherimola x A. squamosa</i> |
| | Chã Grande | 35°30'59.0"W 08°09'23.0"S | <i>A. muricata</i> |
| Sergipe | Goiâna | 34°57'07.6"W 07°38'19.8"S | <i>A. coriacea ; A. muricata</i> |
| | Recife | 34°56'41.4"W 08°01'07.0"S | <i>A. coriacea ; A. squamosa</i> <i>A. muricata</i> |
| Rio Grande do Norte | Moreno | 35°05'32.0"W 08°07'07.0"S | <i>A. muricata ;</i> <i>A. cherimola x A. squamosa</i> |
| | Pitimbú | 34°48'27.3"W 07°28'15.7"S | <i>A. muricata</i> |
| Sergipe | João Câmara | 35°50'24.1"W 05°31'01"S | <i>A. squamosa</i> |
| | Muriú | 35°15'13.5"W 05°35'12.9"S | <i>A. muricata</i> |
| Sergipe | Touros | 35°33'00.6"W 05°14'56.7"S | <i>A. squamosa</i> |
| | Santa Cruz dos Abais | 37°17'15.1"W 11°19'03.0"S | <i>A. muricata</i> |
| | Neópolis | 36°38'28.1"W 10°19'03.0"S | <i>A. muricata</i> |

versidade Federal Rural de Pernambuco – UFRPE). The sampling sites are listed in Table 1.

At each sampling site, five plants were chosen. Twenty leaves were sampled from each plant. The samples were taken to the laboratory in plastic bags for further processing. Mites were removed using a stereoscope and a brush, placed in plastic containers containing 70% alcohol and labeled. Subsequently, the mites were mounted in Hoyer's medium and identified by optical microscopy to at least the genus level.

RESULTS

The adult mites identified in this survey belong to eight families, twenty-five genera and thirty-eight species. The families Ascidae, Bdellidae, Phytoseiidae and Stigmeidae were composed mainly of predator and a few generalist species; the families Tarsonemidae, Tenuipalpidae and Tetranychidae were comprised of phytophagous and one mycophagous species (*T. confusus*), and Tydeidae included species of polyphagous habits. Among all of the species reported, four (*A. aerialis*, *T. mexicanus*, *P. acaciae* and *P. ubiquitus*) were found on all the sampled cultures.

A total of 1011 mite specimens were sampled and identified, of which 500 belonged to the Tydeidae family, 162 to Tenuipalpidae, 138 to Tetranychidae, 136 to Phytoseiidae, 47 to Stigmeidae, 25 to Tarsonemidae, 2 to Bdellidae and 1 to the Ascidae family.

FAMILY PHYTOSEIIDAE BERLESE SUBFAMILY AMBLYSEINAE MUMA

Genus *Amblyseius* Berlese

Amblydromalus manihoti (Moraes, 1994)

Amblyseius manihoti Moraes et al. 1994: 211.
Typhlodromalus manihoti Gondim Jr. and Moraes 2001: 82; Zacarias and Moraes 2001: 582; Moraes et al. 2004: 200.
Amblydromalus manihoti Chant and McMurtry 2005: 207.

Specimens examined — Pernambuco, Recife, 16 September 2005 (2 ♀; 2 ♂), 16 December 2005 (1 ♀) and 18 January 2006 (1 ♀) on *Annona squamosa*.

Amblyseius aerialis (Muma, 1955)

Amblyseiopsis aerialis Muma 1955: 264; Garman 1958: 75.

Typhlodromus (Amblyseius) aerialis Chant 1959: 88; Muma 1961: 287.

Amblyseius aerialis De Leon 1966: 91; Moraes and Mesa 1988: 71; Denmark and Muma 1989: 15; Kreiter and Moraes 1997: 377; Feres and Moraes 1998: 126.

Specimens examined — Pernambuco, Recife, 8 August 2005 (3 ♀) and 16 September 2005 (1 ♀); Rio Grande do Norte, João Câmara, 6 September 2005 (1 ♀) on *Annona squamosa*. Pernambuco, Chã Grande, 8 July 2005 (1 ♀), Água Preta, 4 September 2005 (1 ♀) on *Annona muricata*. Pernambuco, Moreno, 5 September 2005 (1 ♀) and 23 November 2005 (1 ♀; 2 ♂) on *Annona cherimolia* × *Annona squamosa*. Pernambuco, Recife, 12 December 2005 (1 ♀) on *Annona coriacea*.

Amblyseius tamatavensis Blommers, 1974

Amblyseius tamatavensis Blommers 1974: 144; Denmark and Muma 1989: 13; Moraes et al. 1991: 119.

Amblyseius maai Tseng 1976: 123 (synonymy according to Denmark and Muma, 1989).

Amblyseius aegyptiacus Matthysse and Denmark 1981: 343.

Specimens examined — Pernambuco, Bonito, 14 August 2005 (1 ♀), 4 September 2005 (1 ♀; 2 ♂) and 23 October 2005 (6 ♀; 4 ♂); Água Preta, 10 September 2005 (1 ♀); Ceará, Paraipaba, 5 September 2005 (1 ♀) on *Annona muricata*. Pernambuco, Bonito, 14 August 2005 (13 ♀; 9 ♂) on *Annona squamosa*.

Genus *Euseius* Wainstein

Euseius alatus De Leon, 1966

Euseius alatus De Leon 1966: 87; Denmark and Muma 1973: 262; Moraes and McMurtry 1983: 137; Feres and Moraes 1998: 127.

Euseius paraguayensis Denmark and Muma 1970:

224 (synonymy according to Moraes and McMurtry 1983: 137)

Specimens examined — Pernambuco, Recife, 4 July 2005 (1 ♀), 12 December 2005 (1 ♀; 1 ♂) and 18 January 2005 (2 ♀; 1 ♂); Goiana, 8 July 2005 (1 ♂) on *Annona coriacea*. Pernambuco, Recife, 8 August 2005 (1 ♀; 1 ♂), 4 October 2005 (2 ♀) and 10 November 2005 (2 ♂) on *Annona squamosa*. Pernambuco, Goiana, 17 August 2005 (1 ♂) on *Annona muricata*.

Euseius citrifolius Denmark and Muma, 1970

Euseius citrifolius Denmark and Muma 1970: 222; Moraes and McMurtry 1983: 138; Moraes et al. 1991: 131; Feres and Moraes 1998: 127; Lofego et al. 2004: 4; Lofego et al. 2009: 43.

Specimens examined — Bahia, Juazeiro, 18 July 2005 (2 ♀) on *Annona muricata*. Bahia, Juazeiro, 18 July 2005 (2 ♀) on *Annona cherimolia × Annona squamosa*.

Euseius concordis (Chant, 1959)

Typhlodromus (Amblyseius) concordis Chant 1959: 69. *Amblyseius (Iphiseius) concordis* Muma 1961: 288. *Amblyseius concordis* Chant and Baker 1965: 22; Moraes and McMurtry 1983: 138. *Euseius flechtmanni* Denmark and Muma 1970: 223; 1973: 261 (synonymy according to Moraes et al. 1982: 18). *Euseius concordis* Denmark and Muma 1973: 264; Moraes and Oliveira 1982: 317; Moraes and McMurtry 1983: 138; Feres and Moraes 1998: 127; Ferla and Moraes 2002a: 870; Lofego et al. 2004: 5; 2009: 44.

Specimens examined — Pernambuco, Chã Grande, 8 July 2005 (2 ♀); Bahia, Juazeiro, 18 July 2005 (1 ♀) on *Annona muricata*. Bahia, Juazeiro, 18 July 2005 (1 ♂); Pernambuco, Recife, 16 December 2005 (1 ♀); Maranhão, Tutóia, 3 September 2005 (1 ♀) on *Annona squamosa*.

Euseius ho (De Leon, 1965)

Amblyseius (Euseius) ho De Leon 1965a: 125. *Euseius ho* Denmark and Muma 1973: 262; Moraes and McMurtry 1983: 139; Moraes et al. 1991: 132.

Specimens examined — Pernambuco, Recife, 8 September 2005 (1 ♂), 16 September 2005 (1 ♂) and

4 September 2005 (1 ♂) on *Annona squamosa*. Pernambuco, Moreno, 5 September 2005 (1 ♂) and 23 November 2005 (1 ♀) on *Annona cherimolia × Annona squamosa*. Pernambuco, Recife, 5 October 2005 (1 ♂), 12 December 2005 (1 ♀), 18 January 2006 (1 ♀) on *Annona coriacea*.

Euseius sibelius (De Leon, 1962)

Amblyseius (Typhlodromalus) sibelius De Leon 1962: 21.

Euseius sibelius Muma et al. 1970: 98; Moraes and McMurtry 1983: 140; Moraes et al. 1986: 54; 2004: 83; Moraes and Mesa 1988: 81; Feres and Moraes 1998: 128; Chant and McMurtry 2005: 216; 2007: 123; Lofego et al. 2004: 6; 2009: 45; Feres et al. 2009: 467; Demite et al. 2011: 42.

Euseius subalatus De Leon 1965a: 127 (synonymy according to Muma et al. 1970).

Specimens examined — Bahia, Juazeiro, 18 July 2005 (13 ♀) on *Annona cherimolia × Annona squamosa*.

Genus *Iphiseiodes* De Leon

Iphiseiodes zuluagai Denmark and Muma, 1972

Iphiseiodes zuluagai Denmark and Muma 1972: 23; Aponte and McMurtry 1995: 165; Kreiter and Moraes 1997: 377; Feres and Moraes 1998: 127; Lofego et al. 2004: 7; 2009: 45.

Amblyseius zuluagai Moraes and Mesa 1988: 79; Moraes et al. 1991: 125.

Specimens examined — Pernambuco, Bonito, 14 August 2005 (1 ♂), Água Preta, 14 September 2005 (1 ♂) and 4 September 2005 (1 ♂) on *Annona muricata*. Pernambuco, Recife, 4 July 2005 (1 ♀), 5 October 2005 (1 ♂) and 8 August 2005 (1 ♂) on *Annona coriacea*.

Genus *Proprioseiopsis* Muma

Proprioseiopsis cannaensis (Muma, 1962)

Amblyseiulus cannaensis Muma 1962: 4.

Amblyseius cannaensis Moraes and McMurtry 1983: 132; Moraes and Mesa 1988: 77; Moraes et al. 1991: 126; Lofego et al. 2009: 53.

Proprioseiopsis cannaensis Muma et al. 1970: 38; Kreiter and Moraes 1997: 379.

Specimens examined — Maranhão, Junco do Maranhão, 12 September 2005 (1 ♀) on *Annona muricata*.

SUBFAMILY PHYTOSEIINAE BERLESE

Genus *Phytoseius* Ribaga

Phytoseius guianensis De Leon 1965

Phytoseius guianensis De Leon 1965b: 18; Denmark 1966: 23; Denmark and Muma 1973: 269; Moraes and McMurtry 1983: 144; Lofego *et al.* 2004: 11.

Specimens examined — Bahia, Juazeiro, 18 July 2005 (1 ♀) on *Annona cherimolia* × *Annona squamosa*.

SUBFAMILY TYPHLODROMINAE SCHEUTEN

Genus *Leonseius* Chant and McMutry

Leonseius regularis (De Leon, 1965)

Typhloseiopsis regularis De Leon 1965a: 122; Moraes *et al.* 1991: 136.

Diadromus regularis De Leon 1966: 100.

Chanteius regularis De Leon 1967: 16.

Typhlodromus regularis Chant and Yoshida-Shaul 1983: 1034; El-Benawy 1984: 139.

Specimens examined — Pernambuco, Goiana, 8 July 2005 (4 ♀), 17 August 2005 (4 ♀) and 17 September (3 ♀; 1 ♂); Recife, 8 July 2005 (6 ♀; 1 ♂) on *Annona coriacea*. Pernambuco, Goiana, 5 October 2005 (2 ♀; 1 ♂) on *Annona muricata*.

Typhlodromus (Anthoseius) transvaalensis (Nesbitt, 1951)

Kampimodromus transvaalensis Nesbitt 1951: 55.

Neoseiulus transvalensis Muma 1961: 295.

Typhlodromus transvaalensis Chant 1955: 498; Ferla and Moraes 2002b: 1020.

Specimens examined — Bahia, Juazeiro, 18 July 2005 (1 ♀) on *Annona cherimolia* × *Annona squamosa*.

Genus *Typhlodromus* Scheuten

Typhlodromus (Anthoseius) ornatus (Denmark and Muma, 1973)

Amblydromella ornata Denmark and Muma 1973: 270.

Typhlodromus (Anthoseius) ornatus Gondim Jr. and Moraes 2001: 89; Moraes *et al.* 2004: 341; Lofego *et al.* 2004: 13.

Specimens examined — Bahia, Juazeiro, 18 July 2005 (1 ♀) on *Annona cherimolia* × *Annona squamosa*.

Genus *Typhlodromina* Muma

Typhlodromina subtropica Muma and Denmark, 1969

Typhlodromina subtropica Muma and Denmark 1969: 412; Moraes and McMurtry 1983: 142.

Specimens examined — Pernambuco, Recife, 4 July 2005 (1 ♀) on *Annona coriacea*.

FAMILY ASCIDAE VOIGTS AND OUDEMANS

Genus *Proctolaelaps* Berlese

Proctolaelaps bickleyi (Bram, 1956)

Garmania bickleyi Bram 1956: 292.

Proctolaelaps striata Westerboer 1963: 356. Synonymy according to Lindquist and Hunter (1965).

Garmania domestica Nesbitt 1951: 44.

Proctolaelaps bickleyi Chant 1963: 269; Lindquist and Hunter 1965: 16; McGraw and Farrier 1969: 72; Karg 1985: 192; 1988: 448; Halliday *et al.* 1998: 31.

Specimens examined — Pernambuco, Chã Grande, 8 July 2005 (1 ♀) on *Annona muricata*.

FAMILY TETRANYCHIDAE DONNADIEU SUBFAMILY TETRANYCHINAE BERLESE

Genus *Atrichoprocus* Flechtmann

Atrichoprocus uncinatus Flechtmann, 1967

Atrichoprocus uncinatus Flechtmann 1967: 323; Flechtmann and Baker 1970: 157; Flechtmann and Baker 1975: 116; Feres 2000: 166.

Specimens examined — Pernambuco, Água Preta, 23 February 2005 (4 ♀) on *Annona squamosa*.

Genus *Oligonychus* Berlese

Oligonychus punicae (Hirst, 1926)

Paratetranychus punicae Hirst 1926: 830.

Oligonychus punicae Pritchard and Baker 1955: 335.

Specimens examined — Pernambuco, Recife, 16 September 2005 (7 ♀; 1 ♂) on *Annona squamosa*.

Genus *Tetranychus* Dufour

Tetranychus desertorum Banks, 1900

Tetranychus desertorum Banks 1900: 76; Pritchard and Baker 1955: 403; Tuttle and Baker 1968: 4; Paschoal 1970a: 45.

Specimens examined — Pernambuco, Goiana, 8 July 2005 (16 ♀; 4 ♂) on *Annona coriacea*. Bahia, Juazeiro, 18 July 2005 (1 ♂) on *Annona muricata*.

Tetranychus mexicanus (McGregor, 1950)

Septanychus mexicanus McGregor 1950: 323.

Tetranychus mexicanus Pritchard and Baker 1955: 411.

Specimens examined — Pernambuco, Goiana, 19 September 2005 (1 ♂); Água Preta, 24 October 2005 (1 ♀), 23 November 2005 (3 ♀; 4 ♂) and 5 December 2005 (4 ♀; 2 ♂) on *Annona muricata*. Pernambuco, Moreno, 23 November 2005 (9 ♀; 5 ♂) on *Annona cherimolia* × *Annona squamosa*. Pernambuco, Água Preta, 24 October 2005 (8 ♀; 2 ♂) and 23 November 2005 (17 ♀; 13 ♂) on *Annona squamosa*. Pernambuco, Recife 18 January 2006 (4 ♀; 2 ♂) on *Annona coriacea*.

Tetranychus neocaledonicus André, 1933

Tetranychus neocaledonicus André 1933: 302; Pritchard and Baker 1955: 430; Flechtmann and Baker 1970: 162; Flechtmann and Bastos 1972: 86.

Specimens examined — Pernambuco, Bonito, 4 September 2005 (19 ♀; 6 ♂) on *Annona cherimolia* × *Annona squamosa*.

***Tetranychus urticae* Koch, 1836**

Tetranychus urticae Koch 1836: 10; Boudreux and Dosse 1963: 363; Tuttle and Baker 1968: 129; Flechtmann and Baker 1970: 161; Paschoal 1970a: 49; Flechtmann and Bastos 1972: 86; Flechtmann and Baker 1975: 120.

Specimens examined — Pernambuco, Goiana, 8 July 2005 (4 ♀; 1 ♂) on *Annona coriacea*.

FAMILY TARSONEMIDAE CANESTRINI

AND FANZAGO

SUBFAMILY TARSONEMINAE KRAMER

Genus *Daidalotarsonemus* De Leon

Daidalotarsonemus folisetae Lofego et al. 2005

Daidalotarsonemus folisetae Lofego et al. 2005: 2.

Specimens examined — Sergipe, Neópolis, 21 July 2005 (17 ♀) on *Annona muricata*.

Daidalotarsonemus fossae De Leon, 1956

Daidalotarsonemus fossae De Leon 1956: 165; Suski 1968: 777; Smiley 1972: 91.

Specimens examined — Pernambuco, Goiana, 8 July 2005 (1 ♀), 15 July 2005 (1 ♀) on *Annona coriacea*.

Daidalotarsonemus tessellatus De Leon, 1956

Daidalotarsonemus tessellatus De Leon 1956: 163; Smiley 1972: 91; Lofego et al. 2005: 6.

Specimens examined — Pernambuco, Moreno, 5 September 2005 (1 ♀); Sergipe, Neópolis, 21, July 2005 (2 ♀) on *Annona cherimolia* × *Annona squamosa*. Sergipe, Neópolis, 21 July 2005 (2 ♀) on *Annona muricata*.

Genus *Tarsonemus* Canestrini and Fanzago

Tarsonemus confusus Ewing, 1939

Tarsonemus confusus Ewing 1939: 26; Beer 1954: 1173; Smiley 1969: 221; Kaliszewski 1993: 40; Feres et al. 2005: 6; Lofego et al. 2005: 24.

Specimens examined — Pernambuco, Água Preta, 14 August 2005 (1 ♀) on *Annona muricata*.

FAMILY TENUIPALPIDAE BERLESE
SUBFAMILY BREVIPALPINAE RECK

Genus *Brevipalpus* Donnadiet

Brevipalpus phoenicis (Geijskes, 1939)

Tenuipalpus phoenicis Geijskes 1939: 23.

Brevipalpus phoenicis Sayed 1946: 99; Pritchard and Baker 1958: 233; De Leon 1961: 48; Baker et al. 1975: 18; Gonzalez 1975: 82; Meyer 1979: 87; Baker and Tuttle 1987: 98.

Brevipalpus yothersi Baker 1949: 374.

Brevipalpus mcbridei Baker 1949: 374.

Brevipalpus papayensis Baker 1949: 379.

Specimens examined — Pernambuco, Água Preta, 4 September 2005 (8 ♀), 23 December 2005 (1 ♀); Bonito, 14 August 2005 (6 ♀); Ceará, Acaraú, 4 September 2005 (1 ♀); Maranhão, Tutóia, 3 September 2005 (1 ♀); Pará, Capanema, 31 July 2005 (1 ♀) on *Annona squamosa*. Pernambuco, Chã Grande, 8 July 2005 (4 ♀); Bonito, 14 August 2005 (9 ♀), 4 September 2005 (2 ♀), 23 October 2005 (2 ♀); Água Preta, 4 September 2005 (1 ♀), 10 September 2005 (1 ♀), 5 December 2005 (1 ♀); Goiana, 8 December 2005 (1 ♀); Sergipe, Neópolis, 21 July 2005 (2 ♀); Alagoas, Miao de Baixo, 22 July 2005 (2 ♀); Ceará, Barroquinha, 4 September 2005 (2 ♀); Rio Grande do Norte, Muriú, 7 September 2005 (1 ♀) on *Annona muricata*. Bahia, Juazeiro, 18 July 2005 (12 ♀); Pernambuco, Bonito, 4 September 2005 (5 ♀), 23 October 2005 (1 ♀) on *Annona cherimolia* × *Annona squamosa*.

SUBFAMILY TENUIPALPINAE
MITROFANOV

Genus *Tenuipalpus* Donnadiet

Tenuipalpus annonae De Leon, 1957

Tenuipalpus annonae De Leon 1957: 91.

Specimens examined — Bahia, Juazeiro, 18 July 2005 (85 ♀); Pernambuco, Moreno, 23 November 2005 (6 ♀) on *Annona cherimolia* × *Annona squamosa*. Maranhão, Rosário, 3 September 2005 (5 ♀); Rio Grande do Norte, João Câmara, 5 September 2005 (2 ♀) on *Annona squamosa*.

FAMILY TYDEIDAE KRAMER
SUBFAMILY PRONEMATINAE ANDRÉ

Genus *Homeopronematus* André

Homeopronematus anconai (Baker, 1943)

Pronematus anconai Baker 1944a: 188.

Specimens examined — Pernambuco, Bonito, 14 August 2005 (4 ♀), 17 August 2005 (1 ♀), 19 September 2005 (1 ♀), 5 October 2005 (1 ♀), 12 December 2005 (2 ♀; 1 ♂); Água Preta, 23 November 2005 (1 ♀); Maranhão, Junco do Maranhão, 12 September 2005 (4 ♀) on *Annona muricata*. Pernambuco, Bonito, 14 August 2005 (3 ♀; 1 ♂); Água Preta, 4 September 2005 (1 ♀), 24 October 2005 (11 ♀; 2 ♂) on *Annona squamosa*. Pernambuco, Recife, 8 August 2005 (23 ♀; 12 ♂), 5 October 2005 (10 ♀; 3 ♂), 12 December 2005 (15 ♀; 2 ♂), 18 January 2006 (15 ♀; 3 ♂) on *Annona coriacea*.

Genus *Parapronematus* Baker

Parapronematus acaciae Baker, 1965

Parapronematus acaciae Baker 1965: 116; Zacarias 2001: 74.

Specimens examined — Pernambuco, Goiana, 5 October 2005 (2 ♀); Água Preta, 24 October 2005 (1 ♂) on *Annona muricata*. Pernambuco, Recife, 8 August 2005 (4 ♀; 2 ♂), 16 September 2005 (1 ♀), 4 October 2005 (1 ♀), 10 November 2005 (2 ♀; 4 ♂), 16 December 2005 (1 ♀); Bonito, 14 August 2005 (14 ♀; 3 ♂) on *Annona squamosa*. Pernambuco, Recife, 8 August 2005 (4 ♀), 5 October 2005 (10 ♀; 5 ♂), 12 December 2005 (5 ♀; 2 ♂) on *Annona coriacea*. Pernambuco, Bonito, 23 October 2005 (1 ♂); Moreno, 23 November 2005 (9 ♀; 7 ♂) on *Annona cherimolia* × *Annona squamosa*.

Genus *Pronematus* Canestrini

Pronematus ubiquitus (McGregor, 1932)

Tydeus ubiquitus McGregor 1932: 62.

Pronematus ubiquitus Thor 1933: 46; Baker 1939: 273; Baker 1946: 255; Meyer and Rodriguez 1965: 19; Baker 1968b: 1093.

Specimens examined — Pernambuco, Goiana, 19 September 2005 (2 ♀; 1 ♂), 5 October 2005 (1 ♀); Sergipe, Neópolis, 21 July 2005 (1 ♀); Bahia,

Juazeiro, 18 July 2005 (9 ♀; 1 ♂) on *Annona muricata*. Pernambuco, Recife, 8 August 2005 (7 ♀), 16 September 2005 (2 ♀), 10 November 2005 (28 ♀; 20 ♂), 16 December 2005 (21 ♀; 6 ♂), 18 January 2006 (34 ♀; 2 ♂); Bonito, 14 August 2005 (2 ♀; 4 ♂); Água Preta, 23 November 2005 (1 ♀); Rio Grande do Norte, Touros, 7 September 2005 (1 ♂); Bahia, Juazeiro, 18 July 2005 (3 ♀) on *Annona squamosa*. Pernambuco, Recife, 8 August 2005 (1 ♀; 1 ♂), 5 October 2005 (2 ♀; 1 ♂), 12 December 2005 (1 ♀) on *Annona coriacea*. Pernambuco, Bonito, 23 October 2005 (10 ♀; 8 ♂) on *Annona cherimolia* × *Annona squamosa*.

SUBFAMILY TYDEINAE ANDRÉ

Genus *Lorryia* Oudemans

Lorryia formosa Cooreman, 1958

Lorryia formosa Cooreman 1958: 6; Baker 1968a: 995; Feres et al. 2005: 5

Specimens examined — Bahia, Una, 19 July 2005 (1 ♀); Sergipe, Neópolis, 21 July 2005 (6 ♀); Ceará, Paraipaba, 5 September 2005 (4 ♀) on *Annona muricata*. Pernambuco, Recife, 8 August 2005 (12 ♀; 9 ♂), 16 September 2005 (3 ♀; 3 ♂), 4 October 2005 (3 ♂), 10 November 2005 (2 ♂), 16 December 2005 (1 ♀); Bonito, 14 August 2005 (5 ♀) on *Annona squamosa*. Pernambuco, Recife, 4 July 2005 (13 ♀), 8 August 2005 (4 ♀; 5 ♂), 5 October 2005 (1 ♂), 12 December 2005 (2 ♀; 13 ♂), 18 January 2006 (11 ♀; 11 ♂) on *Annona coriacea*.

Lorryia turrialbensis Baker, 1968

Lorryia turrialbensis Baker 1968a: 999; Salviejo 1969: 266; Kazmierski 1998: 298

Specimens examined — Pernambuco, Goiana, 12 December 2005 (1 ♀); Alagoas, Santa Cruz dos Abais, 21 July 2005 (2 ♀) on *Annona muricata*. Pernambuco, Goiana, 17 August 2005 (10 ♀), 17 September 2005 (2 ♀; 1 ♂) on *Annona coriacea*. Pernambuco, Moreno, 23 November 2005 (7 ♂) on *Annona cherimolia* × *Annona squamosa*.

Genus *Neolorryia* André

Neolorryia boycei (Baker, 1944)

Retetydeus boycei Baker 1944b: 78.

Lorryia boycei Baker 1968a: 1004; Feres 2000: 163.

Neolorryia boycei André 1980: 127; Kazmierski 1998: 350.

Specimens examined — Bahia, Juazeiro, 18 July 2005 (5 ♀) on *Annona cherimolia* × *Annona squamosa*.

FAMILY BDELLIDAE DUGES SUBFAMILY BDELLINAE GRANDJEAN

Genus *Bdella* Latreille

Bdella aff. captiosa Atyeo, 1963

Bdella aff. captiosa Atyeo 1963: 170.

Specimens examined — Alagoas, Miao de Baixo, 22 July 2005 (1 ♀) on *Annona muricata*. Pernambuco, Recife, 16 December 2005 (1 ♀) on *Annona squamosa*.

FAMILY STIGMAEIDAE OUDEMANS

Genus *Agistemus* Summers

Agistemus floridanus González, 1965

Agistemus floridanus Gonzalez 1965: 38; Matioli et al. 2002: 103; Arruda Filho and Moraes 2003: 52.

Specimens examined — Pernambuco, Água Preta, 14 August 2005 (1 ♀), 23 November 2005 (1 ♀); Goiana, 5 October 2005 (3 ♀; 3 ♂), 12 December 2005 (4 ♀; 1 ♂); Bonito, 23 October 2005 (6 ♀; 6 ♂) on *Annona muricata*. Pernambuco, Bonito, 14 August 2005 (11 ♀; 5 ♂) on *Annona squamosa*. Bahia, Juazeiro, 18 July 2005 (5 ♀) on *Annona cherimolia* × *Annona squamosa*.

Genus *Zetzellia* Oudemans

Zetzellia languida González, 1965

Zetzellia languida Gonzalez 1965: 21; Arruda Filho and Moraes 2003: 56

Specimens examined — Pernambuco, Goiana, 5 October 2005 (1 ♂) on *Annona muricata*.

DISCUSSION

The phytoseiid mites had the highest species diversity. Their presence in these microsystems likely indicates environmental balance because most mites are predators that feed on phytophagous mites and insect eggs, while others feed on pollen and are generalists. Most of the mites in this family are studied for use in biological control. Some Phytoseiidae species are widely used in Europe and North America, and new species are still being researched (Moraes 2002). Ten phytoseiid mite species were sampled on *A. muricata*, six on *A. squamosa*, seven on *A. cherimola* × *A. squamosa* and six on *A. coriacea*.

In Brazil, many phytoseiid species were recorded on Annonaceae such as *A. aerialis*, *Amblyseius acalyphus* Denmark and Muma, *Amblyseius chiapensis* De Leon, *Amblyseius largoensis* (Muma), *Amblyseius neochiapensis* Lofego, Moraes and McMurtry, *E. alatus*, *Euseius brazilli* (EL-Benawy), *E. citrifolius*, *E. concordis*, *E. sibelius*, *Galendromus (Galendromus) annectens* De Leon, *I. zuluagai*, *Neoseiulus tunus* (De Leon), *P. guianensis*, *Phytoseius kaapre* Demite, Lofego and Feres, *Phytoseius nahuatlensis* De Leon, *Proprioseiopsis dominigos* (El-Banawy), *Ricoseius loxocheles* (De Leon) and *Typhlodromalus peregrinus* (Muma) (Denmark and Muma 1973; Moraes and McMurtry 1983; Ferla and Moraes 2002b; Vasconcelos *et al.* 2006; Demite *et al.* 2009; Demite *et al.* 2011; Rezende & Lofego 2011; Nuvoloni *et al.* 2011). Among these species, *A. aerialis*, *E. alatus*, *E. citrifolius*, *E. concordis*, *E. sibelius*, *I. zuluagai* and *P. guianensis* were also recorded in the present study, in addition to another eight species that were also recorded for the first time on annonas.

The tetranychids *A. uncinatus* and *T. mexicanus* and the tenuipalpid *B. phoenicis* were already reported on annonas species in Brazil (Paschoal 1970b; Flechtmann and Baker 1975; Tuttle *et al.* 1977; Moraes and Flechtmann 1981). The species *B. phoenicis* was the most commonly found mite in the sampling sites (12). Among the studied species, this species was not recorded only in *A. coriacea*. Some Tenuipalpidae mites had already been reported on annonas. Among these mites, the species *B. phoenicis* was found in Angola (Africa) on *A. cherimola*, *Annona* sp. and *Rollinia* sp. (Meyer 1979). In

Brazil, Flechtmann (1976) observed this species on the same host, and Nuvoloni *et al.* (2011) found this mite on *Xilopia aromatic* (Lam.). The species *T. annonae* was described in Mexico on *Annona* sp. (De Leon, 1957). In this study, *T. annonae* was found on *A. squamosa* and *A. squamosa* × *A. cherimolia*. In Panamá, Ochoa *et al.* (1994) observed the species *Brevipalpus pseudostriatus* Ochoa and Salas on *A. cherimola*.

A total of six Tydeidae species were reported, five of which were never recorded on annonas in Brazil. Only the species *L. formosa* was observed on *X. aromatic* by Nuvoloni *et al.* (2011). Two of the three species of the Pronematinae (= Iolinidae) subfamily were found on the four studied hosts, while *H. anconai* was detected on all of the studied plants except atemoya. Among the species belonging to the subfamily Tydeinae, *L. formosa* occurred on sugar apples, soursop and *A. coriacea*. *Lorryia turrialbensis* occurred on all the hosts except for sugar apples, and *N. boycei* was detected only on atemoya. These mites are not considered plant pests. Some of these mites are mycophagous and may reduce the development of fungi populations, some appear to be alternative prey to Phytoseiidae predator mites, and others are predators. Certain mite species are known for their importance in ecological chains in agroecosystems. Groups of generalist species, such as species of the Tydeidae family, are important in maintaining populations of efficient predator mites and are also alternative food when the preferred prey level is low (Moraes and Flechtmann 2008).

Nuvoloni *et al.* (2011) found *Daidalotarsonemus* sp. and *Tarsonemus* sp. on *X. aromatic*. In the present study, *D. folissetae* and *T. confusus* were recorded on *A. muricata*, *D. fossae* was found on *A. coriacea*, and *D. tesselatus* was found on atemoya. In the Tarsonemidae family, the genus *Tarsonemus* is composed of micophagous species, and the genus *Daidalotarsonemus* is composed by species with food habit not well defined. Lindquist (1986) considered that *Daidalotarsonenus* are probably algophagous or fungivorous, rather than phytophagous, however, Lofego *et al.* (2005) observed *D. tesselatus* feed on leaves of *Psidium guajava* L. In addition to the mentioned species, three new tarsonemid species of

were also found on the sampled Annonaceae. Two of these species belong to the genus *Daidalotarsonomus* and one to *Fungitarsonemus*, and both of these groups will soon be described.

The family Tetranychidae is composed of strictly phytophagous organisms. Among the reported species, *T. mexicanus* stood out for occurring on all four of the studied hosts. The species *T. neocaledonicus* was reported on *A. muricata* in Australia by Gutierrez and Schicha (1983). Additionally, *T. mexicanus* and *B. phoenicis* were found on *A. muricata* (i.e., soursop) in Água Preta County. These mites are found on leaves in small populations and without causing damage. However, high populations are found on fruits, causing intense darkening, size reduction and induration of healthy fruits. Moraes and Flechtmann (2008) reported these same species in association with damage to the soursop fruit.

The great mite diversity and apparent species equilibrium in Annonaceae detected in this study may be due to the usage of little or no acaricides in these culture systems and to the essentially cultural management of annonas. There are few studies reporting mite species and the damage caused by them on Annonaceae in Brazil. Usually, these mites are identified through surveys of areas involving various plant species of several families. This study demonstrated the great diversity of mites on the plant species belonging to the family Annonaceae in Northeast Brazil and the state of Pará. However, this fauna seems to be in balance, and high populations were not observed, except for *T. mexicanus* and *B. phoenicis* on soursop.

ACKNOWLEDGEMENTS

We thank the Brazilian Federal Agency for the Support and Evaluation of Graduate Education (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES) and the Pernambuco Science and Technology Support Foundation (Fundação de Amparo a Ciência e Tecnologia do estado de Pernambuco – FACEPE) for granting the first author's scholarship and research funding, and we thank the National Council for Scientific and Technological Development (Conselho Nacional de Desen-

volvimento Científico e Tecnológico – CNPq) for granting the research productivity scholarship to the other authors.

REFERENCES

- André H.M. 1980 — A generic revision of the family Tydeidae (Acaria: Actinedida). IV. Generic descriptions, keys and conclusions — Bull. Ann. Soc. R. Entomol. Belg., 116: 103-167.
- André M. 1933 — Note sur un Tetranique nuisible au Cotonier en Nouvelle-Caledonie — Bull. Mus. Nat. Hist. Natur. Paris (Ser.2), 5: 302-308.
- Aponte O., McMurtry J.A. 1995 — Revision of the genus *Iphiseiodes* De Leon (Acaria: Phytoseiidae) — Int. J. Acarol., 21: 165-183.
[doi:10.1080/01647959508684056](https://doi.org/10.1080/01647959508684056)
- Arruda F.G.P., Moraes G.J. de 2003 — Stigmeidae mites (Acaria: Raphgnathoidea) from Arecaceae of the Atlantic Forest in São Paulo State, Brazil — Neotrop. Entomol., 32: 49-57.
[doi:10.1590/S1519-566X2003000100007](https://doi.org/10.1590/S1519-566X2003000100007)
- Atyeo Warren T. 1963 — The Bdellidae of the Australian Realm. Part. I. New Zealand, Campbell Island, and the Auckland Islands. Part. II. Australia and Tasmania — Bull. Univ. Nebraska St. Mus., 4: 113-210.
- Baker E.W. 1939 — The fig mite, *Eriophyes ficus* Cotte, and other mites of the fig tree, *Ficus carica* Linn. — Calif. Dep. Agr., 28: 266-275.
- Baker E.W. 1944a — Nuevos tydeidos mexicanos (Acarina) — Rev. Soc. Mex. de Hist. Natur., 4: 181-189.
- Baker E.W. 1944b — Tideideos mexicanos (Acari, Tydeidae) — Rev. Soc. Mex. Hist. Natur., 5: 73-81.
- Baker E.W. 1946 — Some Tydeidae from the fig tree (*Ficus carica* L.) — Anales Escuela Nac. Cienc. Biol. Mex., 4: 255-261.
- Baker E.W. 1949 — The genus *Brevipalpus* (Acarina:Pseudoleptidae) — Am. Midl. Nat., 42: 350-402.
[doi:10.2307/2422013](https://doi.org/10.2307/2422013)
- Baker E.W. 1965 — A review of the genera of the family Tydeidae (Acarina) — In: Advances of Acarology. Cornell Univ. Press, Ithaca. 2: pp. 95-120.
- Baker E.W. 1968a — The genus *Lorryia* — Ann. Entomol. Soc. Amer., 61: 986-1008.
[doi:10.1093/aesa/61.4.986](https://doi.org/10.1093/aesa/61.4.986)
- Baker E.W. 1968b — The genus *Pronematus* — Ann. Entomol. Soc. Am., 61: 1091-1097.
[doi:10.1093/aesa/61.5.1091](https://doi.org/10.1093/aesa/61.5.1091)
- Baker E.W., Tuttle D.M. 1987 — The false spider mites of Mexico (Tenuipalpidae: Acari) — US. Agric. Res. Serv. Tech. Bul., 1706: 1-237.

- Baker E.W., Tuttle D.M., Abbatiello M.J. 1975 — The false spider mites of northwestern and north central Mexico (Acarina:Tenuipalpidae) — Smith. Contrib. Zool., 194: 1-23.
[doi:10.5479/si.00810282.194](https://doi.org/10.5479/si.00810282.194)
- Banks N. 1900 — The red spiders of the United States (*Tetranychus* and *Stigmaeus*) — U.S., Dept. Div. Agr. Div. Ent. Tech. Ser., 8: 65-77.
- Beer R.E. 1954. A revision of the Tarsonemidae of the Western Hemisphere (Ordem Acarina). — Kansas Univ. Sci. Bull. 36, Pt. 2: 1091-1387.
- Blommers L. 1974 — Species of the genus *Amblyseius* Berlese, 1914, from Tamatave, east Madagascar (Acarina: Phytoseiidae) — Bull. Zool. Mus. Univ. Amst., 3: 143-155.
- Boudreux H.B., Dosse G. 1963 — Concerning the names of some common spider mites — Adv. Acarol., 1: 350-364.
- Bram R.A. 1956 — A new predatory mite from insect culture — Proc. Ent. Soc. Wash., 58: 292-294.
- Cardoso C.E.L., Sousa J.S. 2000 — Fruticultura tropical: Perspectivas e tendências. — Rev. Econ. Nordeste., 31: 84-95.
- Chant D.A. 1955 — Notes on mites of the genus *Typhlodromus* Scheuten, 1857 (Acarina: Laelaptidae), with descriptions of the males of some species and the female of a new species — Can. Entomol., 87: 496-503.
[doi:10.4039/Ent87496-11](https://doi.org/10.4039/Ent87496-11)
- Chant D.A. 1959 — Phytoseiid mites (Acarina: Phytoseiidae). Part I. Bionomics of seven species in southeastern England. Part II. A taxonomic review of family Phytoseiidae, with descriptions of 38 new species — Can. Entomol. Suppl., 91: 1-166.
- Chant D.A. 1963 — The subfamily Blattisocinae Garman (= Aceosejinae Evans) (Acarina: Blattisocidae Garman) (=Aceosejidae Baker and Warton) in North America, with descriptions of new species — Can. J. Zool., 41: 243-305
[doi:10.1139/z63-025](https://doi.org/10.1139/z63-025)
- Chant D.A., Baker E.W. 1965 — The Phytoseiidae (Acarina) of Central America — Mem. Entomol. Soc. Can., 41: 1-56.
[doi:10.4039/entm9741fv](https://doi.org/10.4039/entm9741fv)
- Chant D.A., McMurtry J.A. 2005 — A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part VI. The tribe Euseini n. tribe, subtribes Typhlodromalina n. subtribe, Euseiina n. subtribe, and Ricoseiina n. subtribe. — Int. J. Acarol., 31: 187-224.
[doi:10.1080/01647950508684424](https://doi.org/10.1080/01647950508684424)
- Chant D.A., McMurtry J.A. 2007 — Illustrated Keys and Diagnoses for the Genera and Subgenera of the Phytoseiidae of the World (Acari: Mesostigmata). — Indra Publishing House, 220 pp.
- Chant D.A., Yoshida-Shaul E. 1983 — A world review of five similar species groups in the genus *Typhlodromus* Scheuten: Part I. The *regularis* and *theodoliticus* groups (Acarina: Phytoseiidae) — Can. J. Zool., 61: 1032-1040.
[doi:10.1139/z83-137](https://doi.org/10.1139/z83-137)
- Cooreman J. 1958 — Notes et observations sur les Acairiens. VII — *Photia graeca* n. sp. (Acaridia, Canestriniidae) et *Lorryia formosa* n. sp. (Stomatostigmata, Tydeyidae) — Bull. Inst. Roy. Sci. Nat., 34: 1-10.
- De Leon D. 1956 — Some mites from lychee. Description of two genera and five new species of Tarsonemidae — Fla. Entomol., 39: 163-174.
[doi:10.2307/3492593](https://doi.org/10.2307/3492593)
- De Leon D. 1957 — The genus *Tenuipalpus* in Mexico (Acarina: Tenuipalpidae) — Fla. Entomol., 40: 81-93.
[doi:10.2307/3492507](https://doi.org/10.2307/3492507)
- De Leon D. 1961 — The genus *Brevipalpus* in Mexico. Part II (Acarina: Tenuipalpidae) — Fla. Entomol., 44: 41-52.
[doi:10.2307/3492547](https://doi.org/10.2307/3492547)
- De Leon D. 1962 — Twenty-three new phytoseiids, mostly from southeastern United States (Acarina: Phytoseiidae) — Fla. Entomol., 45: 11-27
[doi:10.2307/3492899](https://doi.org/10.2307/3492899)
- De Leon D. 1965a — Phytoseiid mites from Puerto Rico with descriptions of new species (Acarina: Mesostigmata) — Fla. Entomol., 48: 121-131.
[doi:10.2307/3493102](https://doi.org/10.2307/3493102)
- De Leon D. 1965b — Ten new species of *Phytoseius (Pennaseius)* from Mexico, Trinidad, and British Guiana with a key to species (Acarina: Phytoseiidae) — Entomol. News, 76: 11-21.
- De Leon D. 1966 — Phytoseiidae of British Guyana with keys to species (Acarina: Mesostigmata) — In: Studies on the fauna of Suriname and other Guyanas. 8: 81-102.
- De Leon D. 1967 — Some mites of the Caribbean Area. Part I. Acarina on plants in Trinidad, West Indies — Lawrence: Allen Press Inc. 66 pp.
- Demite P.R., Feres R.J.F., Lofego A.C., Oliveira A.R. 2009 — Plant inhabiting mites (Acari) from Cerrado biome of Mato Grosso State, Brazil — Zootaxa, 2061: 45-60.
- Demite P.R., Lofego A.C., Feres R.J.F. 2011 — Phytoseiidae (Acari) in forest fragments in the State of São Paulo, Brazil — Zootaxa, 3086: 31-56.
- Denmark H.A. 1966 — Revision of the genus *Phytoseius* Ribaga, 1904 (Acarina: Phytoseiidae) — Fla. Dep. Agric. Bull., 6: 1-105.
- Denmark H.A., Muma M.H. 1970 — Some phytoseiid mites of Paraguay (Phytoseiidae: Acarina) — Fla. Entomol., 53: 219-227.
[doi:10.2307/3493192](https://doi.org/10.2307/3493192)
- Denmark H.A., Muma M.H. 1972 — Some Phytoseiidae of Colombia (Acarina: Phytoseiidae) — Fla. Entomol.,

- 55: 19-29.
[doi:10.2307/3493637](https://doi.org/10.2307/3493637)
- Denmark H.A., Muma M.H. 1973 — Phytoseiidae mites of Brazil (Acarina, Phytoseiidae) — Rev. Bras. Biol., 33: 235-276.
- Denmark H.A., Muma M.H. 1989 — A revision of the genus *Amblyseius* Berlese, 1914 (Acari: Phytoseiidae) — Occas. Pap. Fla. State Collect. Arthropods, 4: 1-149.
- El-Benawy E.M. 1984 — Description of some phytoseiid mites from Brazil (Acarina: Phytoseiidae) — Acarologia., 25: 125-144.
- Ewing H.E. 1939 — A revision of the mites of the subfamily Tarsoneminae of North America, the West Indies and Hawaiian Islands — U. S. Dep. Agric., Tech. Bull., 653: 1-63.
- Feres R.J.F. 2000 — Levantamento e observações naturalísticas da acarofauna (Acari, Arachnida) de seringueiras cultivadas (*Hevea* spp., Euphorbiaceae) no Brasil — Rev. Bras. Zool., 17: 157-173.
- Feres R.J.F., Lofego A.C., Oliveira A.R. 2005 — Ácaros plantícolas (Acari) da "Estação Ecológica do Noroeste Paulista", estado de São Paulo — Biota Neotrop., 5: 1-14.
- Feres R.J.F., Moraes G.J. 1998 — Phytoseiid mites (Acari: Phytoseiidae) from woody areas in the State of São Paulo, Brazil — Syst. Appl. Acarol., 3: 125-132.
- Feres R.J.F., Vieira M.R., Daud R.D., Pereira Jr. E.G., Oliveira G.F., Dourado C.L. 2009 — Ácaros (Arachnida, Acari) de plantas ornamentais na região noroeste do estado de São Paulo, Brasil: inventário e descrição dos sintomas causados pelos fitófagos — Rev. Bras. Entomol., 53: 466-474.
[doi:10.1590/S0085-56262009000300024](https://doi.org/10.1590/S0085-56262009000300024)
- Ferla N.J., Moraes G.J. 2002a — Ácaros (Arachnida, Acari) da seringueira (*Hevea brasiliensis* Muell. Arg.) no Estado do Mato Grosso, Brasil — Rev. Bras. Zool., 19: 867-888.
[doi:10.1590/S0101-81752002000300025](https://doi.org/10.1590/S0101-81752002000300025)
- Ferla N.J., Moraes G.J. 2002b — Ácaros predadores (Acari) em plantas nativas e cultivadas do Estado do Rio Grande do Sul — Rev. Bras. Zool., 19: 1011-1031.
[doi:10.1590/S0101-81752002000400006](https://doi.org/10.1590/S0101-81752002000400006)
- Flechtmann C.H.W. 1967 — New Tetranychidae (Acarina) from Brazil — An. Esc. Super Agric. "Luiz de Queiroz", 24: 323-331.
[doi:10.1590/S0071-12761967000100030](https://doi.org/10.1590/S0071-12761967000100030)
- Flechtmann C.H.W. 1976 — Preliminary report on the false spider mites (Acari: Tenuipalpidae) from Brazil and Paraguay — Proc. Entomol. Soc. Wash., 78: 58-64.
- Flechtmann C.H.W. 1985 — Ácaros de Importância Agrícola — São Paulo, Nobel, 189 pp.
- Flechtmann C.H.W., Baker E.W. 1970 — A preliminary report on the Tetranychidae (Acarina) of Brazil — Ann. Entomol. Soc. Am., 63: 156-163.
[doi:10.1093/aesa/63.1.156](https://doi.org/10.1093/aesa/63.1.156)
- Flechtmann C.H.W., Baker E.W. 1975 — A report on the Tetranychidae (Acari) of Brazil — Rev. Bras. Entomol., 19: 111-121.
- Flechtmann C.H.W., Bastos J.A.M. 1972 — Ácaros Tetranychoidea do estado do Ceará, Brasil — Ciênc. Agron., 2: 83-90.
- Garman P. 1958 — New species belonging to the genera *Amblyseius* and *Amblyseiopsis* with keys to *Amblyseius*, *Amblyseiopsis* and *Phytoseiulus* — Ann. Entomol. Soc. Am., 51: 69-79.
[doi:10.1093/aesa/51.1.69](https://doi.org/10.1093/aesa/51.1.69)
- Geijkse D.C. 1939 — Beitraege zur Kenntnis der europäischen Spinnmilben (Acari, Tetranychidae), mit besonderer Berücksichtigung der niederländischen Arten. Meded. Landbouwhoogesch — Wageningen., 42: 1-68.
- Gondim Jr. M.G.C., Moraes G.J. 2001 — Phytoseiid mites (Acari: Phytoseiidae) associated with palm trees (Arecaceae) in Brazil — Syst. Appl. Acarol., 6: 65-94.
- Gonzalez R.H. 1965 — A taxonomic study of the genera *Mediolata*, *Zetzellia* and *Agistemus* (Acarina: Stigmaeidae) — Univ. Calif. Public. Entomol., 41: 1-64.
- Gonzalez R.H. 1975 — Revision of the *Brevipalpus phoenicus* "complex", with descriptions of new species from Chile and Thailand (Acarina, Tenuipalpidae) — Acarologia., 17: 82-91.
- Gutierrez J., Schicha E. 1983 — The spider mite family Tetranychidae (Acari) in New South Wales — Int. J. Acarol., 9: 99-1983.
- Halliday R.B., Walter D.E. and Lindquist E.E. 1998 — Revision of the Australian Ascidae (Acarina: Mesostigmata) — Invert. Taxon., 12: 1-54.
- Hirst S. 1926 — Descriptions of new mites including four new species of "red spider" — Proc. Zool. Soc., 96: 825-841.
[doi:10.1111/j.1469-7998.1926.tb07130.x](https://doi.org/10.1111/j.1469-7998.1926.tb07130.x)
- Kaliszewski M. 1993 — Key to Palearctic females of the genus *Tarsonemus*. Wydawnictwo Naukowe Uniwersytetu Im. Adama Mickiewicza w Poznaniu, Poznań — Seria Zoologia, 14: pp. 204.
- Karg W. 1985 — Die Raubmilbgattung *Proctolaelaps* Berlese, 1923 — Zool. Jahr. Abt. Syst. Ökol., 112: 185-206.
- Karg W. 1988 — Zur Kenntnis der Gattung *Proctolaelaps* Berlese, 1923 (Acarina, Parasitiformes) — Zool. Jahr. Abt. Syst. Ökol., 115: 441-454.
- Kazmierski A. 1998 — Tydeidae of the world: generic relationships, new and redescribed taxa and keys to all species. A revision of the subfamilies Pretydeinae and

- Tydeinae (Acari: Actinedida: Tydeidae) — Acta Zool. Cracov., 41: 283-455.
- Koch C.L. 1836 — Deutschen Crustacea, Miriapoda, Arachnida — Fasc., 1: 10.
- Kreiter S., Moraes G.J. 1997 — Phytoseiid mites (Acari: Phytoseiidae) from Guadeloupe and Martinique — Fla. Entomol., 80: 376-382.
[doi:10.2307/3495770](https://doi.org/10.2307/3495770)
- Leon J. 1987 — Botânica de los cultivos tropicales — San José, Ilca, 444 pp.
- Lindquist E.E., Hunter P.E. 1965 — Some mites of the genus *Proctolaelaps* Berlese (Acarina: Blattisocidae) associated with forest insect pests — Can. Entomol., 97: 16-32.
[doi:10.4039/Ent9715-1](https://doi.org/10.4039/Ent9715-1)
- Lindquist E.E. 1986 — The world genera of Tarsonemidae (Acari: Heterostigmata): A morphological, phylogenetic, and systematic revision, with reclassification of family — group taxa in the Heterostigmata — Mem. Entomol. Soc. Can., 136: 1-517.
[doi:10.4039/entm118136fv](https://doi.org/10.4039/entm118136fv)
- Lofego A.C., Demite P.R., Kishimoto R.G., Moraes G.J. 2009 — Phytoseiid mites on grasses in Brazil (Acari: Phytoseiidae) — Zootaxa, 2240: 41-59.
- Lofego A.C., Moraes G.J. de, Castro L.A.S. 2004 — Phytoseiid mites (Acari: Phytoseiidae) on Myrtaceae in the State of São Paulo, Brazil — Zootaxa., 516: 1-18.
- Lofego A.C., Ochoa R. and Moraes G.J. 2005 — Some tarsonemid mites (Acari: Tarsonemidae) from the Brazilian "Cerrado" vegetation, with descriptions of three new species — Zootaxa., 823: 1-27.
- Mabberley D.J. 1997 — The plant-Book — New York, University Press, 858 pp.
- Mathysse J.G., Denmark H.A. 1981 — Some Phytoseiids of Nigeria (Acarina: Phytoseiidae) — Fla. Entomol., 64: 340-357.
[doi:10.2307/3494585](https://doi.org/10.2307/3494585)
- Matioli A.L., Ueckermann E.A. and Oliveira C.A.L. de 2002 — Some Stigmaeid and Eupalopsellid mites from citrus orchards in Brazil (Acari: Stigmaeidae and Eupalopsellidae) — Int. J. Acarol., 28: 99-120.
[doi:10.1080/01647950208684287](https://doi.org/10.1080/01647950208684287)
- McGraw J.R., Farrier M.H. 1969 — Mites of the superfamily Parasitoidea (Acarina: Mesostigmata) associated with *Dendroctonus* and *Ips* (Coleoptera: Scolytidae) — Agric. Exper. Stat. Tec., North Carolina. Bulletin, 192 pp.
- McGregor E.A. 1932 — The ubiquitous mite, a new species on citrus — Proc. Entomol. Soc. Wash., 34: 60-63.
- McGregor E.A. 1950 — Mites of the family Tetranychidae — Am. Midl. Nat., 44: 257-420.
[doi:10.2307/2421963](https://doi.org/10.2307/2421963)
- Meyer M.K.P.S. 1979 — The Tenuipalpidae (Acari) of Africawith keys to the world fauna — Entomology Mem. Dep.agric. tech. Serv. Repub. S. Afr. n° 50, 135 pp.
- Meyer M.K.P.S., Rodriguez M.C. 1965 — Acari associated with cotton in Southern Africa (with reference to other plants) — Garcia Orta. Ser. Zool., 13: 195-226.
- Moraes G.J. 2002 — Controle Biológico de ácaros fitófagos com ácaros predadores In Parra, J.R.P., Botelho, P.S.M., Ferreira, B.S.C., Bento, J.M.S. Controle Biológico no Brasil: Parasitóides e Predadores — Manole, pp. 225-237.
- Moraes G.J., Denmark H.A., Guerrero J.M. 1982 — Phytoseiid mites of Colombia (Acarina: Phytoseiidae) — Int. J. Acarol., 8: 15-22.
[doi:10.1080/01647958208683273](https://doi.org/10.1080/01647958208683273)
- Moraes G.J., Flechtmann C.H.W. 1981 — Ácaros fitófagos do Nordeste do Brasil — Pesqui. Agropecu. Bras., 6: 177-186.
- Moraes G.J., Flechtmann C.H.W. 2008 — Manual de Acarologia: Acarologia básica e ácaros de plantas cultivadas no Brasil — Holos editora, 288 pp.
- Moraes G.J., McMurtry J.A. 1983 — Phytoseiid mites (Acarina) of northeastern Brazil with descriptions of four new species — Int. J. Acarol., 9: 131-148.
[doi:10.1080/01647958308683326](https://doi.org/10.1080/01647958308683326)
- Moraes G.J., McMurtry J.A., Denmark H.A. 1986 — A catalog of the mite family Phytoseiidae: references to taxonomy, synonymy, distribution and habitat — Brasilia. Embrapa-DDT., 353 pp.
- Moraes G.J., McMurtry J.A., Denmark H.A., Campos C.B. 2004 — A revised catalog of the mite family Phytoseiidae — Zootaxa., 434: 1-494.
- Moraes G.J., Mesa N.C. 1988 — Mites of the family Phytoseiidae (Acari) in Colômbia, with descriptions of three new species — Int. J. Acarol., 14: 71-88.
[doi:10.1080/01647958808683790](https://doi.org/10.1080/01647958808683790)
- Moraes G.J., Mesa N.C., Braun A. 1991 — Some Phytoseiid mites of Latina America (Acari: Phytoseiidae) — Int. J. Acarol., 17: 117-139.
[doi:10.1080/01647959108683892](https://doi.org/10.1080/01647959108683892)
- Moraes G.J., Mesa N.C., Braun A., Melo E.L. 1994 — Definition of the *Amblyseius limonicus* species group (Acari: Phytoseiidae), with descriptions of two new species and new records — Int. J. Acarol., 20: 209-217.
[doi:10.1080/01647959408684019](https://doi.org/10.1080/01647959408684019)
- Moraes G.J., Oliveira J.V. 1982 — Phytoseiid mites of coastal Pernambuco in northeastern Brazil — Acarologia., 23: 315-318.
- Muma M.H. 1955 — Phytoseiidae (Acarina) associated with citrus in Florida — Ann. Entomol. Soc. Am.,

- 48: 262-272.
[doi:10.1093/aesa/48.4.262](https://doi.org/10.1093/aesa/48.4.262)
- Muma M.H. 1961 — Subfamilies, genera and species of Phytoseiidae (Acarina: Mesostigmata) — Bull. Fla. Sta. Mus. Biol. Sci., 5: 267-302.
- Muma M.H. 1962 — New Phytoseiidae (Acarina: Mesostigmata) from Florida — Fla. Entomol., 45: 1-10.
[doi:10.2307/3492897](https://doi.org/10.2307/3492897)
- Muma M.H., Denmark H.A. 1969 — The *conspicua* species-group of *Typhlodromina* Muma, 1961 — Ann. Entomol. Soc. Am., 62: 406-413.
[doi:10.1093/aesa/62.2.406](https://doi.org/10.1093/aesa/62.2.406)
- Muma M.H., Denmark H.A., De Leon D. 1970 — Phytoseiidae of the Florida. Arthropods of Florida and neighboring land areas — USA: Florida Department of Agriculture and Consumer Services, Division of Plant Industry., 150 pp.
- Nesbitt H.H.J. 1951 — A taxonomic study of the Phytoseiinae (Family Laelaptidae) predaceous upon Tetranychidae of economic importance — Zool. Verh., 12: 1-64.
- Nogueira E.A., Mello N.T.C., Maia M.L. 2005 — Produção e comercialização de anonáceas em São Paulo e Brasil — Informações Econômicas., 35: 51-54.
- Nuvoloni F.M., Feres R.J.F., Demite P.R. 2011 — Mites associated to *Xylopia aromatico* (Lam.) Mart. (Annonaceae) in urban and rural fragments of semideciduous forest — Rev. Bras. Entomol., 55: 571-577.
[doi:10.1590/S0085-56262011000400013](https://doi.org/10.1590/S0085-56262011000400013)
- Ochoa R., Aguilar H., Vargas C. 1994 — Phytophagous mites of Central America: An Illustrated Guide — CATIE, Turrialba, Costa Rica., 234 pp.
- Paschoal A.D. 1970a — Contribuição ao conhecimento da família Tetranychidae no Brasil (Arachnida: Acarina). (Tese de doutorado) — In: Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo, 116 pp.
- Paschoal A.D. 1970b — Revisão da família Tetranychidae no Brasil (Arachnida: Acarina) — An. Esc. Super. Agric. "Luiz de Queiroz", 27: 457-483.
- Pritchard A.E., Baker E.W. 1955 — A revision of the spider mite family Tetranychidae — Pac. Coast Entomol. Soc. Mem. ser. 2, 472 pp.
- Pritchard A.E., Baker E.W. 1958 — The false spider mites (Acarina: Tenuipalpidae) — Univ. Calif. Publ. Entomol., 14: 175-274.
- Rezende J.M., Lofego A.C. 2011 — Phytoseiidae (Acari: Mesostigmata) on plants of the central region of the Brazilian Cerrado — Acarologia, 51: 449-463.
- Salviejo P.B. 1969 — Some Philippine Tydeid Mites (Tydeidae: Acarina) — Philipp. Entomol., 1: 261-277.
- Sayed M.T. 1946 — Description of *Tenuipalpus granati* nov. spec. and *Brevipalpus pyri* nov. spec. — Bull. Soc. Fouad 1er Entomologie., 30: 99-104.
- Smiley R.L. 1969 — Further studies on the Tarsonemidae, II (Acarina) — Proc. Entomol. Soc. Wash., 71: 218-229.
- Smiley R.L. 1972 — A review of the genus *Daidalotarsonemus* De Leon (Acarina: Tarsonemidae) — Proc. Entomol. Soc. Wash., 74: 89-94.
- Sobrinho R.B., Cardoso J.E., Freire F.C.O. 1998 — Pragas de fruteiras tropicais de importância agroindustrial — EMBRAPA, Brasília, 209 pp.
- Suski Z.W. 1968 — Setation of legs I, II and III in some species of the mite family Tarsonemidae (Acarina: Heterostigmata) — Roczn. Nauk Roln., 93: 757-787.
- Thor S. 1933 — Acarina Tydeidae, Ereynetidae — Tierreich., 60: 1-82.
- Tseng Y.H. 1976 — Systematics of the mite family Phytoseiidae from Taiwan, with a revised key to genera of the world (II) — J. Agr. Assoc. China, 94: 85-128.
- Tuttle D.M., Baker E.W. 1968 — Spider mites of the Southwestern United States and a revision of the family Tetranychidae — Univ. Arizona Press, Tucson, 143 pp.
- Tuttle D.M., Baker E.W., Sales F.M. 1977 — Spider mites (Tetranychidae: Acarina) of the state of Ceará, Brazil — Int. J. Acarol., 3: 1-8.
- Vasconcelos G.J.N., Silva F.R., Barbosa D.G.F., Gondim Jr. M.G.C., Moraes G.J. 2006 — Diversidade de fitoseídeos (Acari: Phytoseiidae) em fruteira tropicais no estado de Pernambuco, Brasil — Magistra, 18: 90-101.
- Westerboer I. 1963 — Die Familie Podocinidae Berlese, 1916. In: Beiträge zur Systematik und Ökologie Mitteleuropäischer Acarina. Mesostigmata 1. (H.J. Stammer, ed.) — Akademische Verlagsgesellschaft Geest and Portig K.-G. Leipzig., pp. 179-450.
- Zacarias M.S. 2001 — Diversidade de ácaros (Arthropoda: Acari) em euforbiáceas (Euphorbiaceae) de três localidades do Estado de São Paulo. (Tese) — Escola Superior de Agricultura, Universidade de São Paulo, 154 pp.
- Zacarias M.S., Moraes G.J. 2001 — Phytoseiid Mites (Acari) associated with rubber trees and other euphorbiaceous plants in southeastern Brazil — Neotrop. Entomol., 30: 579-58
[doi:10.1590/S1519-566X2001000400011](https://doi.org/10.1590/S1519-566X2001000400011)

COPYRIGHT

 Sousa J.M. de et al. Acarologia is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.