

|                     |      |         |           |
|---------------------|------|---------|-----------|
| Linzer biol. Beitr. | 47/1 | 665-666 | 31.7.2015 |
|---------------------|------|---------|-----------|

## Notes on Palaearctic Leucospidae (Hymenoptera, Chalcidoidea), especially from Libya, Egypt, Iran and Pakistan

Michael MADL

This paper is an addition and correction to MADL & SCHWARZ (2014). The note of GIORDANI SOIKA (1934) is dealing with the red colour forms of the species mentioned below. Libya is the correct spelling of the country and not Lybia, which is the scientific genus name of the boxer crabs (Decapoda: Xanthidae).

### ***Leucospis brevicauda* FABRICIUS, 1804**

*Leucospis brevicauda* FABRICIUS, 1804: GIORDANI SOIKA 1934: 67 (footnote 1: tax., Egypt without locality, Libya without locality).

### ***Leucospis dorsigera* FABRICIUS, 1775**

*Leucospis dorsigera* FABRICIUS, 1775: GIORDANI SOIKA 1934: 67 (footnote 1: tax., Egypt without locality, Libya without locality).

*Leucospis dorsigera* FABRICIUS, 1775: HESAMI, AKRAMI & BAUR 2005: 66 (biol., Iran), 67 (figs 1, 2).

*Leucospis dorsigera* FABRICIUS, 1775: BAUR 2005: 34 (biol., Iran), 43 (host-parasite cat.), 70 (cat. biol.).

The page numbers of HESAMI, AKRAMI & BAUR (2005) cited in MADL & SCHWARZ (2014: 1572) are those of BAUR (2005). The latter paper is not listed in MADL & SCHWARZ (2014).

### ***Leucospis gigas* FABRICIUS, 1793**

*Leucospis gigas* FABRICIUS, 1793: GIORDANI SOIKA 1934: 67 (footnote 1: tax., Egypt without locality, Libya without locality).

\**Leucospis* sp.: NURSE 1903: 360 (Pakistan).

NURSE (1903) mentions only the genus *Leucospis* in his survey of the fauna of Quetta. Some years later two new species have been described by CAMERON (1906). As both species are considered as synonyms of *Leucospis gigas*, the note of NURSE is cited *Leucospis* sp.

### ***Leucospis intermedia* ILLIGER, 1807**

*Leucospis intermedia* ILLIGER, 1807: GIORDANI SOIKA 1934: 67 (footnote 1: tax., Egypt without locality, Libya without locality).

## Zusammenfassung

Berichtigungen und Ergänzungen zur Arbeit von MADL & SCHWARZ (2014) werden mitgeteilt. GIORDANI SOIKA (1934) weist auf rötliche Formen bei *Leucospis brevicauda* FABRICIUS, 1804, *Leucospis dorsigera* FABRICIUS, 1775 und *Leucospis gigas* FABRICIUS, 1793, die in Ägypten und Libyen gesammelt worden sind, hin.

## References

- BAUR H. (ed) (2005): Determination lists of entomophagous insects 14. — IOBC/WPRS Bulletin **28** (11): III, 79 pp.
- GIORDANI SOIKA A. (1934): Monografia degli *Odynerus* etiopici. Parte prima. — Annali del Museo Civico di Storia Naturale di Genova **57**: 23-83.
- HESAMI S.A., AKRAMI M.A. & H. BAUR (2005): *Leucospis dorsigera* (Hymenoptera, Leucospidae) as a hyperparasitoid of Cerambycidae (Coleoptera) through Xoridinae (Hymenoptera: Ichneumonidae). — Journal of Hymenoptera Research **14** (1): 66-67.
- MADL M. & M. SCHWARZ (2014): Notes on Palaearctic species of the family Leucospidae (Hymenoptera, Chalcidoidea), with new records from North Africa and Middle East. — Linzer biologische Beiträge **46** (2): 1569-1580.
- NURSE C.G. (1903): Notes on the Insects of Quetta. — Journal of the Bombay Natural History Society **15** (2): 359-362.

Author's address:

Michael MADL  
2. Zoologische Abteilung  
Naturhistorisches Museum  
Burgring 7  
A-1010 Wien, Austria  
E-mail: [michael.madl@nhm-wien.ac.at](mailto:michael.madl@nhm-wien.ac.at)