



CODEN [USA]: IAJ PBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Research Article

ASSESSMENT OF KNOWLEDGE OF FAMILY PLANNING AMONG NURSES AT SERVICES HOSPITAL LAHORE, PAKISTAN

¹Shaheen Kausar, ²Shabnam Kousar, ³Tahira Shaheen

¹Head Nurse, Services Hospital Lahore, Email ID: shaheenkousar1233@gmail.com

²Head Nurse, Services Hospital Lahore, Email ID: tweetyalone001@gmail.com

³Nursing Instructor

Post Graduate College of Nursing Punjab, Lahore, Email ID: ushnaumna@gmail.com

Article Received: August 2021

Accepted: August 2021

Published: September 2021

Abstract:

Introduction: In spite of the recent advances in telecommunication and globalization, the international community has failed to bridle the giant of population explosion. Newly married couples fear and are hesitant to adopt family planning approaches. That is the reason they not only feel shy to buy but are also ignorant of their practical importance. Most of the couples are surrounded by myths and lack of evidence regarding contraception. Since the healthcare

professionals especially the nurses are the frontline companions at any health provision facility therefore his study was designed to evaluate the knowledge about the family planning among nurses at Services Hospital Lahore.

Method: A self-structured questionnaire containing the sociodemographic data and the content of the study was handed over to the 100 nurses of

Services Hospital Lahore. **Result:** All the participants had heard about different available family planning methods. The key sources of the knowledge were their instructors (67%) and social media (16%), self study (13%) and miscellaneous (4%) Fig.1 . About 86% of the study

participants marked correct options regarding various types of contraceptives. **Conclusion:** Our study revealed that the level of knowledge was satisfactory regarding family planning among

the nurses working in Services Hospital Lahore. In order to imbibe positive attitude and enlighten the general public regarding the utilization of the birth control measures, the nurses working in different capacities across the country need to be trained and taught so that they can trickle down the benefits of reproductive health to their clients and decipher the outlook of the country.

Keywords: Family planning, population, contraceptives

Corresponding author:

Shaheen Kausar,

Head Nurse,

Services Hospital Lahore,

Email ID: shaheenkousar1233@gmail.com

QR code



Please cite this article in press Shaheen Kausar et al, Assessment Of Knowledge Of Family Planning Among Nurses At Services Hospital Lahore, Pakistan., Indo Am. J. P. Sci, 2021; 08(9).

INTRODUCTION:

WHO has defined family planning as a multi-dimensional process involving shared decision making among the couples to choose willfully the resources favourable to them according to their needs. [1] It involves practical approaches by the couples to brace the reproductive cycle and birth spacing by modifying the physiochemical anatomy of their reproductive systems.[2] Therefore an Ideal Family planning method should deal with both the general and reproductive health of the mother and the fetus, maintain the desired gap between two offsprings, check unwanted pregnancies and forced abortions, prevent sexually transmitted diseases, and enhance the quality of life of the mother, fetus, and family as a whole.[3]

Fertilization is the fusion of the male sperm and female egg during sexual intercourse inside the female genital tract. Contraception deals with either by intercepting with the ovulation or by keeping the egg and the sperm at distance so that they do not fuse and get implanted onto the uterine wall thus avoiding unwanted and unplanned pregnancies.[4] According to the latest survey the world population today has reached above 7.7 billion mark and is predicted to hit 9 billion in the next 20 years.[5] Rapidly growing population is a matter of great concern for the both the developing and developed world Pakistan is among the top 10 densely populated countries of the world with an estimated population above 22 crore and a growth rate of about 10 million a year.[4] Despite several awareness campaigns and extensive population welfare and reproductive health initiatives we are facing the music to uplift the socioeconomic indicators for improving the determinants of human development at the national level. [6]

Therefore it has become essential that the national resources should only be reserved for the planned and desired pregnancies who can develop into healthy and responsible citizens play their role in nation building. The contraceptives not only reduce risk of maternal mortality by 25% associated with unsafe abortions, unplanned and unbooked pregnancies but also protect the individuals from the sexually transmitted diseases such as AIDS(HIV), Chlamydia,


Syphilis, Gonorrhoea Genital Herpes and Hepatitis B etc.,[7] Developing countries mostly encounter the issues associated with the product information ,supply, cost and the usage of the contraceptives[8] Currently all public and private hospitals are offering both over the counter, short term and invasive permanent methods for birth control e.g vasectomy and tubal ligation.[9] According to study of Jimma, Ethiopia, envisioned that a sound knowledge is not enough to warrant to adopt the contraceptive practices.[10] WHO has issued guidelines regarding the delivery and administration of contraceptive methods for the health professionals according to their domain and expertise.[11] Pills and barrier method are usually offered by the Lady Health Visitors and trained nurses and the vasectomies and bilateral tubal ligations are performed by gynaecologists or trained physicians [12] Therefore it is of sublime importance that all healthcare workers especially nurses possess sufficient knowledge attitude and skills so that they can clear the doubts and motivate their clients for opting suitable contraception methods; [13][14] [15]. The myths in the minds of the young couples can only be cleared by delivering evidence-based information at their threshold [16,17] This study is aimed to evaluate the knowledge regarding family planning among nurses at Services Hospital Lahore.

MATERIALS AND METHODS:

After seeking formal approval from the ethical review committee of Services Institute of Medical Sciences (SIMS), Lahore. Purposive sampling was done for the sample size. A self-structured questionnaire containing the sociodemographic data and the content of the study was handed over to the 200 nurses working in different departments. This questionnaire contained the sociodemographic data and the variables to examine knowledge regarding the family planning control. All the participant included were married at the time of conducting this study. The data was analyzed by SPSS version 21.

Table.1 Sociodemographic data of the participants

Age groups	Frequency(n=200)	Percentage (%)
≤25	48	24
26-29	72	36
>30	80	40
Total	200	100
Income (Rupees)	Frequency(n=200)	Percentage (%)
<50,000	64	32
50,000-75,000	108	54
>75,000	28	14
Age at the time of marriage	Frequency(n=200)	Percentage (%)
≤20	68	34
20-25	72	36
>25	60	30
Length of marriage (years)	Frequency(n=200)	Percentage(%)
<5	96	48
5-10	64	32
>10	40	20
Number of children	Frequency(n=200)	Percentage(%)
0	52	26
1	46	23
2	56	28
3	26	13
>3	20	10
Number of trainings	Frequency(n=200)	Percentage(%)
≤5	144	72
>5	56	28
Years Of Experience	Frequency(n=200)	Percentage(%)
≤5	76	38
>5	124	62
Residence	Frequency(n=200)	Percentage(%)
Rural	72	36
Urban	128	64
Qualification	Frequency(n=200)	Percentage(%)
Diploma	124	62
Bachelors	48	24
Post Graduation	32	16

Figure.1 Source of knowledge for family planning**RESULTS:**

According to the sociodemographic details, out of 200 respondents 64% belonged to urban parts of the Lahore district while 36 were from rural parts of the province. About 62% of the participants were Diploma holders. The monthly family income of the majority (54.0%) of the participants had their salary between 50,000 and 75,000 PKR. Regarding the family size of the participants, majority (28.0%) of them had two children. The majority (36%) belonged to the age group 26-30 year. The mean age of participants was 29.2 ± 5.4 years [Table 1].

Knowledge status of participants

Almost all the respondents had heard about the various available methods of family planning. The key sources of knowledge were their instructors (67.0%) and social media (16.0%), self-study (13%) and miscellaneous (4.0%) Fig.1 About 86% of the study participants marked correct options regarding various types of contraceptives. About 87.0% marked correct response regarding over-the-counter short term contraceptive methods, 78.0% gave correct answers regarding long-term hormonal birth control methods, 94.0% answered correctly regarding

permanent method of contraception e.g vasectomy and tubal ligation and 84% gave correct response regarding natural/traditional methods used for family planning. About 76% could correctly name the different types of oral contraceptive pills. Only 64% can recall the adverse effects of OCPs.

DISCUSSION:

Despite several awareness campaigns and extensive population welfare and reproductive health initiatives we are facing the music to uplift the socioeconomic indicators for improving the determinants of human development at the national level. [6] Therefore it has become essential that the national resources should only be reserved for the planned and desired pregnancies who can develop into healthy and responsible citizens play their role in nation building. The contraceptives not only reduce risk of maternal mortality by 25% associated with unsafe abortions, unplanned and unbooked pregnancies but also protect the individuals from the sexually transmitted diseases such as AIDS(HIV), Chlamydia, Syphilis, Gonorrhea Genital Herpes and Hepatitis B etc.,[7] Developing countries mostly encounter the issues associated with the product information ,supply, cost and the usage of

the contraceptives[8] Currently all public and private hospitals are offering both over the counter, short term and invasive permanent methods for birth control e.g vasectomy and tubal ligation.[9] According to study of Jimma, Ethiopia, envisioned that a sound knowledge is not enough to warrant to adopt the contraceptive practices.[10] WHO has issued guidelines regarding the delivery and administration of contraceptive methods for the health professionals according to their domain and expertise. The results of this study showed that all of respondents had at least come across the term of family planning and their major source of information were their instructors (67%) and social media (16%), self study (13%) and miscellaneous (4%). Fig.1 About 86% of the study participants marked correct options regarding various types of contraceptives. These results are higher than a study conducted in Jimma zone, Southwest Ethiopia,[10] and another study done in Rohtak district, India.[20] The difference may be attributed to the sample population comprising only married/committed and professional women working in a tertiary care setup. However, the statistics were better than a study conducted in Northwest Ethiopia in 2018.[14] Previous studies encompassed gross differences between the reproductive health care offered by the community and professional organizations [21,22). Both these setups can perform their best only when they are provided with the technical support, funding and skill development for the efficient evidence based error free expertise.[23-29]. About 76% could correctly name the different types of oral contraceptive tablets. Only 64% can recall the adverse effects of OCPs.

Therefore, to implement the national family and reproductive health policy under the initiative of the federal and provincial government. It is extremely essential that the recommended guidelines of WHO and UNO should be endorsed in full swing for the provision of hassle-free doorstep reproductive and family health coverage for all sectors of the community. For this purpose the nurses ought to revise the fundamentals of family planning regularly so that they can act as facilitators in the community.

CONCLUSION AND RECOMMENDATION:

Our study revealed that although the extent of knowledge regarding the family planning approaches was satisfactory but only 64% could recall the potential side effects of the oral contraceptive pills. Our study revealed that the level of knowledge was satisfactory regarding family planning among the nurses working in Services Hospital Lahore. In order to imbibe positive attitude and enlighten the general

public regarding the utilization of the birth control measures, the nurses working in different capacities across the country need to be trained and taught so that they can trickle down the benefits of reproductive health to their clients and decipher the outlook of the country by imparting them user and health friendly advices.

REFERENCES:

1. World Health Organization. Standards for Maternal and Neonatal Care. Geneva: World Health Organization; 2007.
2. Survey H. Central Statistical Agency. Ethiopian Demographic and Health Survey 2016 Key Indicators Report. Addis Ababa and Maryland, Ethiopia, 2016.
3. World Health Organization. World Health Organization. Fact Sheets on Family Planning, World Health Organization.
4. Park K. Park's Textbook of Preventive and Social Medicine. 17th ed. Jabalpur: Banarasidas Bhanot; 2002. p. 325-58.
5. Van Bavel J. The world population explosion: Causes, backgrounds and -projections for the future. Facts Views Vis Obgyn 2013; 5:281-91.
6. Renjhen P, Kumar A, Pattanshetty S, Sagir A, Samarasinghe CM. A study on knowledge, attitude and practice of contraception among college students in Sikkim, India. J Turkish Ger Gynecol Assoc 2010; 11:78-81.
7. Weldegerima B, Deneke A. Women's knowledge, preferences, and practices of modern contraceptive methods in Woreta, Ethiopia. Res Social Adm Pharm 2008; 4:302-7.
8. Jahan U, Verma K, Gupta S, Gupta R, Mahour S, Kirti N, *et al*. Awareness, attitude and practice of family planning methods in a tertiary care hospital, Uttar Pradesh, India. Int J Reprod Contraception Obstet Gynecol 2017; 6:500-6.
9. Republic FD. National guideline for family planning Federal Democratic Republic of Ethiopia. 2011; 1-69.
10. Tilahun T, Coene G, Luchters S, Kassahun W, Leye E, Temmerman M, *et al*. Family Planning Knowledge, Attitude and Practice among Married Couples in Jimma Zone, Ethiopia. PLoS One 2013; 8:e61335.
11. Family Planning Global Handbook WHO 2018 [Internet]. [Cited 2019 Feb 04]. Available from: <https://www.fphandbook.org/sites/default/files/global-handbook-2018-full-web.pdf>.
12. Welfare UF, Fertility T, Goals D. Family Planning. 2002.
13. Mahadeen AI, Khalil AO, Hamdan-Mansour AM, Sato T, Imoto A. Knowledge, attitudes and

- practices towards family planning among women in the rural southern region of Jordan. *East Mediterr Health J* 2012; 18:567-72.
14. Semachew Kasa A, Tarekegn M, Embiale N. Knowledge, attitude and practice towards family planning among reproductive age women in a resource limited settings of Northwest Ethiopia. *BMC Res Notes* 2018; 11:7-12.
 15. White K, Hopkins K, Grossman D, Potter JE. Providing family planning services at primary care organizations after the exclusion of Planned Parenthood from publicly funded programs in Texas: Early qualitative evidence. *Health Serv Res* 2018; 53:2770-86.
 16. Ai O, Ol A, Aa A. Contraceptive practices among women in rural communities in South-Western Nigeria. *Global Journal of Medical Research* 2011; 11.
 17. Oyedokun AO. Determinants of contraceptive Usage: Lessons from women in Osun State, Nigeria. *J Humanit Soc Sci* 2007; 1:1-14.
 18. Santos JRA. Cronbach's alpha: A tool for assessing the reliability of scales. *J Extension* 1999; 37:1-5.
 19. Republic FD. National guideline for family planning Federal Democratic Republic of Ethiopia. 2011.
 20. Gupta V, Mohapatra D, Kumar V. Family planning knowledge, attitude, and practices among the currently married women (aged 15–45 years) in an urban area of Rohtak district, Haryana. 2016; 5:627-32.
 21. Frost JJ, Frohwirth LF, Blades N, Zolna MR, Douglas-Hall A, Bearak J. Guttmacher Institute 1 Publicly Funded Contraceptive Services at U.S [Internet]. *Clinics*. 2015 [cited 2019 Feb 04]. Available from: https://www.guttmacher.org/sites/default/files/report/publicly_funded_contraceptive_services_2015_3.pdf.
 22. Carter MW, Gavin L, Zapata LB, Bornstein M, Mautone-Smith N, Moskosky SB. Four aspects of the scope and quality of family planning services in US publicly funded health centers: Results from a survey of health center administrators. *Contraception* 2016; 94:340-7.
 23. Lohr PA, Schwarz EB, Gladstein JE, Nelson AL. Provision of contraceptive counseling by internal medicine residents. *J Womens Health (Larchmt)* 2009; 18:127-31.
 24. Akers AY, Gold MA, Borrero S, Santucci A, Schwarz EB. Providers' perspectives on challenges to contraceptive counseling in primary care settings. *J Women's Heal* 2010; 19:1163-70.
 25. Chuang CH, Martenis ME, Parisi SM, Delano RE, Sobota M, Nothnagle M, *et al*. Contraception and abortion coverage: What do primary care physicians think? *Contraception* 2012; 86:153-6.
 26. Biggs M, Rocca C, Brindis C, Hirsch H, Grossman D. UCSF UC San Francisco previously published works title did increase use of highly effective contraception contribute to declining abortions in Iowa? *Contraception* 2015; 91:167-73.
 27. Luchowski AT, Anderson BL, Power ML, Raglan GB, Espey E, Schulkin J. Obstetrician-gynecologists and contraception: Practice and opinions about the use of IUDs in nulliparous women, adolescents and other patient populations. *Contraception* 2014; 89:572-7.
 28. Biggs MA, Harper CC, Brindis CD. California family planning health care providers' challenges to same-day long-acting reversible contraception provision. *Obstet Gynecol* 2015; 126:338-45.
 29. Wood S, Beeson T, Bruen B, Goldberg DG, Mead H, Shin P, *et al*. Scope of family planning services available in federally qualified health centers. *Contraception* 2014; 89:85-90.
 30. Sreytouch V. Knowledge, Attitude and Practice (KAP) of Family Planning among Married Women in Banteay Meanchey, Cambodia.
 31. Quereishi MJ, Mathew AK, Sinha A. Knowledge, attitude and practice of family planning methods among the rural females of Bagbahara block Mahasamund district in Chhattishgarh. *Glob J Med Public Health* 2017; 6:1-7.
 32. Moizuddin MK, Shahabuddin S, Ajit GS. Study of Knowledge and Practice of Contraception in Urban Slum Community Mumbai. *International Journal of Current Medical and Applied Sciences* 2014; 3:35-41.
 33. Renjhen P, Gupta S, Barua A, Shipra J, Binita K. A study of knowledge, attitude and practice of family planning among the women of reproductive age group in Sikkim. *J Obstet Gynecol India* 2007; 63-7.
 34. Ashwini Nayak U, Ramakrishnan KG, Venkateswar KN, Vijayshree M. Assessing the knowledge, attitude and practice of contraception in rural India: A necessary step in achieving population control 2017; 6:3328-31.
 35. Gambhir P, Khaira R, Singh A, Raj H. A cross-sectional observational pilot study regarding status of contraceptive prevalence rate in family planning programme in rural practice area of government medical college Patiala. *Journal of Dental and Medical Sciences* 2018; 17:35-40.