

Disproof of the Riemann Hypothesis

Frank Vega

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France

Abstract

We define the function $\nu(x) = \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log(x + C \times \sqrt{x} \times \log \log x)} - 1$ for some positive constant C independent of x . We prove that the Riemann hypothesis is false when there exists some number $y \geq 13.1$ such that for all $x \geq y$ the inequality $\nu(x) \leq 0$ is always satisfied. We know that the function $\nu(x)$ is monotonically decreasing for all sufficiently large numbers $x \geq 13.1$. Hence, it is enough to find a value of $y \geq 13.1$ such that $\nu(y) \leq 0$ since for all $x \geq y$ we would have that $\nu(x) \leq \nu(y) \leq 0$. Using the tool *gp* from the project PARI/GP, we note that $\nu(100!) \approx -2.938735877055718770 E-39 < 0$ for all $C \geq \frac{1}{1000000!}$. In this way, we claim that the Riemann hypothesis could be false.

Keywords: Riemann hypothesis, Nicolas inequality, Chebyshev function, prime numbers
2000 MSC: 11M26, 11A41, 11A25

1. Introduction

In mathematics, the Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$ [1]. Let $N_n = 2 \times 3 \times 5 \times 7 \times 11 \times \dots \times p_n$ denotes a primorial number of order n such that p_n is the n^{th} prime number. Say Nicolas(p_n) holds provided

$$\prod_{q|N_n} \frac{q}{q-1} > e^\gamma \times \log \log N_n.$$

The constant $\gamma \approx 0.57721$ is the Euler-Mascheroni constant, \log is the natural logarithm, and $q | N_n$ means the prime number q divides to N_n . The importance of this property is:

Theorem 1.1. [2], [3]. Nicolas(p_n) holds for all prime numbers $p_n > 2$ if and only if the Riemann hypothesis is true.

In mathematics, the Chebyshev function $\theta(x)$ is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

where $p \leq x$ means all the prime numbers p that are less than or equal to x . We know this property for this function:

Theorem 1.2. [4]. *There are infinitely many values of x such that*

$$\theta(x) > x + C \times \sqrt{x} \times \log \log \log x$$

for some positive constant C independent of x .

We also know that

Theorem 1.3. [5]. *If the Riemann hypothesis holds, then*

$$\left(\frac{e^{-\gamma}}{\log x} \times \prod_{q \leq x} \frac{q}{q-1} - 1 \right) < \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}$$

for all numbers $x \geq 13.1$.

Let's define $H = \gamma - B$ such that $B \approx 0.2614972128$ is the Meissel-Mertens constant [6]. We know from the constant H , the following formula:

Theorem 1.4. [7].

$$\sum_q \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right) = \gamma - B = H.$$

For $x \geq 2$, the function $u(x)$ is defined as follows

$$u(x) = \sum_{q > x} \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right).$$

We use the following theorems:

Theorem 1.5. [8]. *For $x > -1$:*

$$\frac{x}{x+1} \leq \log(1+x).$$

Theorem 1.6. [9]. *For $x \geq 1$:*

$$\log\left(1 + \frac{1}{x}\right) < \frac{1}{x+0.4}.$$

Let's define:

$$\delta(x) = \left(\sum_{q \leq x} \frac{1}{q} - \log \log x - B \right).$$

Definition 1.7. *We define another function:*

$$\varpi(x) = \left(\sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right).$$

Putting all together yields the proof that the inequality $\varpi(x) > u(x)$ is satisfied for a number $x \geq 3$ if and only if $\text{Nicolas}(p)$ holds, where p is the greatest prime number such that $p \leq x$. In this way, we introduce another criterion for the Riemann hypothesis based on the Nicolas criterion. Using this new criterion, we claim that the Riemann hypothesis could be false.

2. Results

Theorem 2.1. *The inequality $\varpi(x) > u(x)$ is satisfied for a number $x \geq 3$ if and only if $\text{Nicolas}(p)$ holds, where p is the greatest prime number such that $p \leq x$.*

Proof. We start from the inequality:

$$\varpi(x) > u(x)$$

which is equivalent to

$$\left(\sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right) > \sum_{q > x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right).$$

Let's add the following formula to the both sides of the inequality,

$$\sum_{q \leq x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right)$$

and due to the theorem 1.4, we obtain that

$$\sum_{q \leq x} \log \left(\frac{q}{q-1} \right) - \log \log \theta(x) - B > H$$

because of

$$H = \sum_{q \leq x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right) + \sum_{q > x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right)$$

and

$$\sum_{q \leq x} \log \left(\frac{q}{q-1} \right) = \sum_{q \leq x} \frac{1}{q} + \sum_{q \leq x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right).$$

Let's distribute it and remove B from the both sides:

$$\sum_{q \leq x} \log \left(\frac{q}{q-1} \right) > \gamma + \log \log \theta(x)$$

since $H = \gamma - B$. If we apply the exponentiation to the both sides of the inequality, then we have that

$$\prod_{q \leq x} \frac{q}{q-1} > e^\gamma \times \log \theta(x)$$

which means that $\text{Nicolas}(p)$ holds, where p is the greatest prime number such that $p \leq x$. The same happens in the reverse implication. \square

Theorem 2.2. *The Riemann hypothesis is true if and only if the inequality $\varpi(x) > u(x)$ is satisfied for all numbers $x \geq 3$.*

Proof. This is a direct consequence of theorems 1.1 and 2.1. \square

Theorem 2.3. *If the Riemann hypothesis holds, then*

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log \theta(x)} > 1$$

for all numbers $x \geq 13.1$.

Proof. Under the assumption that the Riemann hypothesis is true, then we would have

$$\prod_{q \leq x} \frac{q}{q-1} < e^\gamma \times \log x \times \left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right)$$

after of distributing the terms based on the theorem 1.3 for all numbers $x \geq 13.1$. If we apply the logarithm to the both sides of the previous inequality, then we obtain that

$$\sum_{q \leq x} \log\left(\frac{q}{q-1}\right) < \gamma + \log \log x + \log\left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right).$$

That would be equivalent to

$$\sum_{q \leq x} \frac{1}{q} + \sum_{q \leq x} \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q}\right) < \gamma + \log \log x + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}$$

where we know that

$$\begin{aligned} \log\left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right) &< \frac{1}{\frac{8 \times \pi \times \sqrt{x}}{3 \times \log x + 5} + 0.4} \\ &= \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 0.4 \times (3 \times \log x + 5)} \\ &= \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} \end{aligned}$$

according to theorem 1.6 since $\frac{8 \times \pi \times \sqrt{x}}{3 \times \log x + 5} \geq 1$ for all numbers $x \geq 13.1$. We use the theorems 1.4 and ?? to show that

$$\sum_{q \leq x} \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q}\right) = H - u(x)$$

and $\gamma = H + B$. So,

$$H - u(x) < H + B + \log \log x - \sum_{q \leq x} \frac{1}{q} + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}$$

which is the same as

$$H - u(x) < H - \delta(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}.$$

We eliminate the value of H and thus,

$$-u(x) < -\delta(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}$$

which is equal to

$$u(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \delta(x).$$

We know from the theorem 2.1 that $\varpi(x) > u(x)$ for all numbers $x \geq 13.1$ and therefore,

$$\varpi(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \delta(x).$$

Hence,

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \log \log \theta(x) - \log \log x.$$

Suppose that $\theta(x) = \epsilon \times x$ for some constant $\epsilon > 1$. Then,

$$\begin{aligned} \log \log \theta(x) - \log \log x &= \log \log(\epsilon \times x) - \log \log x \\ &= \log(\log x + \log \epsilon) - \log \log x \\ &= \log\left(\log x \times \left(1 + \frac{\log \epsilon}{\log x}\right)\right) - \log \log x \\ &= \log \log x + \log\left(1 + \frac{\log \epsilon}{\log x}\right) - \log \log x \\ &= \log\left(1 + \frac{\log \epsilon}{\log x}\right). \end{aligned}$$

In addition, we know that

$$\log\left(1 + \frac{\log \epsilon}{\log x}\right) \geq \frac{\log \epsilon}{\log \theta(x)}$$

using the theorem 1.5 since $\frac{\log \epsilon}{\log x} > -1$ when $\epsilon > 1$. Certainly, we will have that

$$\log\left(1 + \frac{\log \epsilon}{\log x}\right) \geq \frac{\frac{\log \epsilon}{\log x}}{\frac{\log \epsilon}{\log x} + 1} = \frac{\log \epsilon}{\log \epsilon + \log x} = \frac{\log \epsilon}{\log \theta(x)}.$$

Thus,

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \frac{\log \epsilon}{\log \theta(x)}.$$

If we add the following value of $\frac{\log x}{\log \theta(x)}$ to the both sides of the inequality, then

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log \theta(x)} > \frac{\log \epsilon}{\log \theta(x)} + \frac{\log x}{\log \theta(x)} = \frac{\log \epsilon + \log x}{\log \theta(x)} = \frac{\log \theta(x)}{\log \theta(x)} = 1.$$

We know this inequality is satisfied when $0 < \epsilon \leq 1$ since we would obtain that $\frac{\log x}{\log \theta(x)} \geq 1$. Therefore, the proof is done. \square

Theorem 2.4. *If there exists some number $y \geq 13.1$ such that for all $x \geq y$ the inequality $\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log(x + C \times \sqrt{x} \times \log \log x)} \leq 1$ is satisfied for some positive constant C independent of x , then the Riemann hypothesis should be false.*

Proof. From the theorem 1.2, we know that there are infinitely many values of x such that

$$\theta(x) > x + C \times \sqrt{x} \times \log \log \log x$$

for some positive constant C independent of x . That would be equivalent to

$$\log \theta(x) > \log(x + C \times \sqrt{x} \times \log \log \log x)$$

and so,

$$\frac{1}{\log \theta(x)} < \frac{1}{\log(x + C \times \sqrt{x} \times \log \log \log x)}$$

for all numbers $x \geq 13.1$. Hence,

$$\frac{\log x}{\log \theta(x)} < \frac{\log x}{\log(x + C \times \sqrt{x} \times \log \log \log x)}.$$

If the Riemann hypothesis holds, then

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log(x + C \times \sqrt{x} \times \log \log \log x)} > 1$$

for those values of x that complies with

$$\theta(x) > x + C \times \sqrt{x} \times \log \log \log x$$

due to the theorem 2.3. By contraposition, if there exists some number $y \geq 13.1$ such that for all $x \geq y$ the inequality

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log(x + C \times \sqrt{x} \times \log \log \log x)} \leq 1$$

is satisfied for some positive constant C independent of x , then the Riemann hypothesis should be false, because of there are infinitely many values of x which satisfy the inequality in the theorem 1.2 and comply with $x \geq y$ no matter how big could be y . \square

Definition 2.5. Let's define the function $\nu(x) = \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log(x + C \times \sqrt{x} \times \log \log \log x)} - 1$ for some positive constant C independent of x .

Theorem 2.6. The Riemann hypothesis could be false.

Proof. From the theorem 2.4, we know that the Riemann hypothesis is false when there exists some number $y \geq 13.1$ such that for all $x \geq y$ the inequality $\nu(x) \leq 0$ is always satisfied. We know that the function $\nu(x)$ is monotonically decreasing for all sufficiently large numbers $x \geq 13.1$. Let $\nu'(x)$ be the derivative of $\nu(x)$. We can check the value of $\nu'(x)$ from this web site <https://www.wolframalpha.com/input> and see that $\nu'(x)$ is lesser than zero for all sufficiently large numbers $x \geq 13.1$. Indeed, a function $\nu(x)$ of a real variable x is monotonically decreasing in some interval if the derivative of $\nu(x)$ is lesser than zero and the function $\nu(x)$ is continuous over that interval [10]. In this way, it is enough to find a value of $y \geq 13.1$ such that $\nu(y) \leq 0$ since for all $x \geq y$ we would have that $\nu(x) \leq \nu(y) \leq 0$. We can check that $\nu(100!) \approx -2.938735877055718770 E-39 < 0$ for all $C \geq \frac{1}{1000000!}$ using the tool *gp* from the web site <https://pari.math.u-bordeaux.fr>. Consequently, we claim that the Riemann hypothesis could be false. \square

Appendix

We use the following input:

$$(3*\log(x)+5)/(8*\pi*\sqrt{x}+1.2*\log(x)+2)+\log(x)/\log(x+C*\sqrt{x}*\log(\log(x)))-1$$

from the web site <https://www.wolframalpha.com/input>. Besides, we use the following input into a single command line:

$$(3*\log(100!)+5)/(8*3.14*\sqrt{100!}+1.2*\log(100!)+2) \\ +\log(100!)/\log(100!+(1/1000000!)*\sqrt{100!}*\log(\log(100!)))-1$$

using the tool *gp* from the project <https://pari.math.u-bordeaux.fr>.

Acknowledgments

I thank Richard J. Lipton and Craig Helfgott for helpful comments.

References

- [1] P. B. Borwein, S. Choi, B. Rooney, A. Weirathmueller, *The Riemann Hypothesis: A Resource for the Afficionado and Virtuoso Alike*, Vol. 27, Springer Science & Business Media, 2008.
- [2] J.-L. Nicolas, Petites valeurs de la fonction d’Euler et hypothèse de Riemann, *Séminaire de Théorie des nombres DPP*, Paris 82 (1981) 207–218.
- [3] J.-L. Nicolas, Petites valeurs de la fonction d’Euler, *Journal of number theory* 17 (3) (1983) 375–388. doi:10.1016/0022-314X(83)90055-0.
- [4] A. E. Ingham, *The Distribution of Prime Numbers*, no. 30, Cambridge University Press, 1990.
- [5] J. B. Rosser, L. Schoenfeld, Sharper Bounds for the Chebyshev Functions $\theta(x)$ and $\psi(x)$, *Mathematics of computation* (1975) 243–269 doi:10.1090/S0025-5718-1975-0457373-7.
- [6] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie., *J. reine angew. Math.* 1874 (78) (1874) 46–62. doi:10.1515/crll.1874.78.46. URL <https://doi.org/10.1515/crll.1874.78.46>
- [7] Y. Choie, N. Lichiardopol, P. Moree, P. Solé, On Robin’s criterion for the Riemann hypothesis, *Journal de Théorie des Nombres de Bordeaux* 19 (2) (2007) 357–372. doi:10.5802/jtnb.591.
- [8] L. Kozma, Useful Inequalities, http://www.lkozma.net/inequalities_cheat_sheet/ineq.pdf, accessed on 2021-10-08 (2021).
- [9] A. Ghosh, An Asymptotic Formula for the Chebyshev Theta Function, arXiv preprint arXiv:1902.09231.
- [10] G. Anderson, M. Vamanamurthy, M. Vuorinen, Monotonicity Rules in Calculus, *The American Mathematical Monthly* 113 (9) (2006) 805–816. doi:10.1080/00029890.2006.11920367.