

A faunistic study on Scarabaeidae (Coleoptera: Scarabaeoidea) of Iran

Studium faunistyczne Scarabaeidae Iranu (Coleoptera: Scarabaeoidea)

Marek BUNALSKI¹, Najmeh SAMIN², Majid NAVAEIAN³, Javad AZADIYAN⁴,
Naser Kamali MASHHOOD⁵, Hamid SAKENIN⁶

¹ Department of Entomology and Environmental Protection, University of Life Sciences, Dąbrowskiego 159, 60-594 Poznań, Poland; email: marek.bunalski@up.poznan.pl; ORCID: 0000-0001-6969-7625

² Young Researchers and Elites Club, Science and Research Branch, Islamic Azad University, Tehran, Iran

³ Department of Biology, Yadegar-e Imam Khomeini (RAH) Shahre Rey Branch, Islamic Azad University, Tehran, Iran

⁴ Pars Gostaran Zharf Tourism Company, Tehran Jurassic Park, Tehran, Iran

⁵ Darabad Natural History Museum, First Municipal District, Tehran, Iran

⁶ Department of Plant Protection, Qaemshahr Branch, Islamic Azad University, Mazandaran, Iran

ABSTRACT: This paper deals with a faunistic survey on Scarabaeidae (Coleoptera) of Iran. In total, 16 species in 10 genera and five subfamilies Aphodiinae, Cetoniinae, Melolonthinae, Rutelinae and Scarabaeinae were identified. Four species are new records for the fauna of Iran: *Bodilopsis sordida sordida* (FABRICIUS, 1775) and *Phalacrothous biguttatus* (GERMAR, 1824) (Aphodiinae), *Onthophagus (Palaeonthophagus) basipustulatus* HEYDEN, 1889 and *Onthophagus (Palaeonthophagus) coenobita* (HERBST, 1783) (Scarabaeinae).

KEY WORDS: Aphodiinae, Cetoniinae, Melolonthinae, Rutelinae, Scarabaeinae, distribution, new records, Iran.

Introduction

The superfamily Scarabaeoidea is a very large taxon of beetles of which are known roughly 1,600 genera and 35,000 species (RATCLIFFE & PAULSEN 2008). The Scarabaeoidea first appear in the fossil record in the Middle Jurassic (KRELL 2000, 2007). The family Scarabaeidae is the largest family in this superfamily with over 30,000 species worldwide (GUNTER & al. 2016).

The fauna of Iranian Scarabaeoidea has been studied rather well (PETROVITZ 1954, 1958, 1980; BARAUD 1968; BORTESI & ZUNINO 1974; ZAIRI 1976; KABAKOV 1982; NIKODYM & KRÁL 1998; BARARI & al. 1998a, 1998b; BARARI 2001; KEITH & AHRENS 2002; MODARRES AWAL 2006; ZIANI & GUDENZI 2000, 2006, 2007, 2009; ZIANI 2006, 2011; MONTREULL & SERRI 2007; KEITH & ULIANA 2008; MOWLAVI & al. 2008, MORADI GHARAKHLOO & ZIANI 2009, 2010; ZIANI & MORADI GHARAKHLOO 2010; GHAHARI & al. 2011; YARMAND & al. 2012; SAKENIN & SAMIN 2015; BUNALSKI & GHAHARI 2015; BUNALSKI & al. 2016; DAVOODI & al. 2017).

The first checklist of Iranian Scarabaeidae was prepared by MODARRES AWAL (1997) with 148 species, and then MODARRES AWAL (2012) listed 178 species. The first checklist of Iranian Scarabaeoidea with 509 species was prepared by BUNALSKI & al. (2014a).

The purpose of this paper is to record the species of Scarabaeidae from some regions of Iran as part of ongoing faunistic studies of Scarabaeoidea in Iran.

Material and methods

The materials of this research were collected by sweeping net, light traps and knock down during occasional field surveys from some regions of Iran. The collected specimens were killed by ethyl acetate within killing bottle and then pinned, or were put in alcohol (ethanol 75%) and determined by the authors. Information concerning date of collection, locality, and number of specimens are given. Some of the specimens are preserved in Bugsland of Jurassic Park (Tehran, Iran) and private collection of the second author. The classification, nomenclature and distribution of scarab beetles suggested by LÖBL

& SMETANA (2006) and LÖBL & LÖBL (2016) have been followed.

Results

In total, 16 species in 10 genera of Scarabaeidae have been studied in this paper which four species, *Bodilopsis sordida sordida* (FABRICIUS, 1775), *Phalacronothus biguttatus* (GERMAR, 1824), *Onthophagus (Palaeonthophagus) basipustulatus* HEYDEN, 1889 and *Onthophagus (Palaeonthophagus) coenobita* (HERBST, 1783) are represent as new country records. The list of species is given below alphabetically.

Subfamily Aphodiinae LEACH, 1815

Tribe Aphodiini LEACH, 1815

Subtribe Aphodiina LEACH, 1815

Genus *Agrilinus* MULSANT & REY, 1870

1. *Agrilinus convexus* (ERICHSON, 1848)

Distribution: AFRICA: Algeria, Morocco, Tunisia, ASIA: Iran, Mongolia, Russia, Turkey, EUROPE: Armenia, Austria, Azerbaijan, Bosnia-Herzegovina, Bulgaria, Croatia, Czech Republic, France, Germany, Greece, Hungary, Italy, Netherlands, Poland, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Switzerland, Ukraine.

Material examined: Gilan prov., Talesh, V 2011, 2 exx.; East Azarbayjan prov., Absh-Ahmad, VII 2016, 3 exx.

Genus *Amidorus* MULSANT & REY, 1870

2. *Amidorus cribrarius* (BRULLÉ, 1832)

Distribution: ASIA: Cyprus, Israel, Iran (NADERIAN & al. 2021), Lebanon, Syria, Turkey, EUROPE: Albania, Azerbaijan, Bosnia-Herzegovina, Bulgaria, Croatia, Georgia, Greece, Macedonia, Russia, Serbia.

Material examined: East Azarbayjan prov., Arasbaran forest, IX 2012, 2 exx.

Genus *Bodilopsis* ÁDÁM, 1994

3. *Bodilopsis sordida sordida* (FABRICIUS, 1775)

Distribution: ASIA: China, Japan, Kazakhstan, Kyrgyzstan, North Korea, Russia, South Korea, Turkey; EUROPE: Albania, Armenia, Austria, Belgium, Bulgaria, Byelorussia, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Georgia, Great Britain, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Macedonia, Netherlands, Norway, Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Ukraine.

Material examined: Isfahan prov., Kashan, 3 exx., VI 2012; Golestan prov., Golestan National Park, 4 exx., IV 2013. These are the first records for Iran.

Genus *Otophorus* MULSANT, 1842

4. *Otophorus haemorrhoidalis* (LINNAEUS, 1758)

Distribution: AFRICA: Algeria, Morocco; ASIA: Bhutan, China, Cyprus, India, Iran, Japan, Kazakhstan, Kyrgyzstan, Nepal, North Korea, South Korea, Mongolia, Russia, Syria, Tadzhikistan, Turkey, Turkmenistan, Uzbekistan; EUROPE: Albania, Armenia, Austria, Azerbaijan, Belgium, Bosnia-Herzegovina, Bulgaria, Byelorussia, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Georgia, Great Britain, Greece, Hungary, Italy, Kosovo, Latvia, Lithuania, Luxembourg, Malta, Moldavia, Montenegro, Netherlands, Norway, Poland, Portugal, Romania, Russia, Serbia, Slovenia, Spain, Sweden, Switzerland, Ukraine.

Material examined: Semnan prov., Shahrud Jangale-Abr, VI 2011, 5 exx.; Mazandaran prov., Behshahr, VII 2017, 3 exx.

Genus *Phalacronothus* MOTSCHULSKY, 1860

5. *Phalacronothus biguttatus* (GERMAR, 1824)

Distribution: AFRICA: Algeria, Morocco, Tunisia; ASIA: Kazakhstan, Russia, Turkey; EUROPE: Albania, Armenia, Austria, Belgium, Bosnia-Herzegovina, Bulgaria, Croatia, Czech Republic, Germany, Georgia, Great Britain, Hungary, Italy, Latvia, Luxembourg, Macedonia, Moldavia, Netherlands, Poland, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Switzerland, Ukraine.

Material examined: Zanzjan prov., Abhar, V 2016, 3 exx. This is the first record for Iran.

Subfamily Cetoniinae LEACH, 1815

Tribe Cetoniini LEACH, 1815

Subtribe Cetoniina LEACH, 1815

Genus *Cetonia* FABRICIUS, 1775

6. *Cetonia (Cetonia) viridescens* REITTER, 1891

Distribution: ASIA: Iran, Syria.

Material examined: West Azarbayjan prov., Oshnavieh, VIII 2014, 1 ex.

Genus *Protaetia* BURMEISTER, 1842

7. *Protaetia (Philhelena) ignisternum ignisternum* (REITTER, 1891)

Distribution: ASIA: Iran, Iraq, Turkey, Turkmenistan; EUROPE: Azerbaijan.

Material examined: West Azarbayjan prov., Salmas, IX 2009, 2 exx.

Genus *Tropinota* MULSANT, 1842

8. *Tropinota (Epicometis) senicula*
(MÉNÉTRIÉS, 1832)

Distribution: ASIA: Iran, Turkey, EUROPE: Armenia, Azerbaijan, Georgia.

Material examined: East Azarbayjan prov., Arasbaran forest, IX 2012, 3 exx.

Tribe Valgini MULSANT, 1842

Genus *Valgus* SCRIBA, 1790

9. *Valgus hemipterus hemipterus*
(LINNAEUS, 1758)

Distribution: AFRICA: Algeria, Morocco, Tunisia; ASIA: Iran, Kazakhstan, Kyrgyzstan, Russia, Tadjikistan, Turkey, Uzbekistan; EUROPE: Albania, Armenia, Austria, Azerbaijan, Belgium, Bosnia-Herzegovina, Bulgaria, Byelorussia, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Georgia, Great Britain, Greece, Hungary, Italy, Kosovo, Latvia, Lithuania, Luxembourg, Macedonia, Moldavia, Montenegro, Netherlands, Norway, Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey, Ukraine.

Material examined: Isfahan prov., Najaf-Abad, V 2010, 3 exx.; Fars prov., Abadeh, VI 2017, 2 exx.

Subfamily Melolonthinae LEACH, 1819

Tribe Melolonthini LEACH, 1819

Genus *Polyphylla* HARRIS, 1841

10. *Polyphylla (Polyphylla) olivieri*
(LAPORTE, 1840)

Distribution: ASIA: Iran, Israel, Syria, Turkey, Turkmenistan; EUROPE: Armenia, Azerbaijan, Georgia, Greece, Russia.

Material examined: Ardabil prov., Aslandooz, VI 2012, 4 exx.; Kurdistan prov., Bijar, VIII 2015, 2 exx.; Mazandaran prov., Qaemshahr, V 2017, 3 exx.

Tribe Rhizotrogini BURMEISTER, 1855

Genus *Amphimallon* LATREILLE, 1825

11. *Amphimallon leuthneri* REITER, 1902

Distribution: ASIA: Iran, Syria, Turkey.

Material examined: Kurdistan prov., Baneh, IV 2013, 2 exx.

Subfamily Rutelinae W.S. MACLEAY, 1819

Tribe Adoretini BURMEISTER, 1844

Subtribe Adoretina BURMEISTER, 1844

Genus *Adoretus* DEJEAN, 1833

12. *Adoretus (Adoretus) nigrifrons*
(STEVEN, 1809)

Distribution: ASIA: Afghanistan, China, Iran, Kazakhstan, Kyrgyzstan, Russia, Tadjikistan, Turkmenistan, Uzbekistan; EUROPE: Azerbaijan, Georgia, Russia.

Material examined: East Azarbayjan prov., Maragheh, VII 2016, 2 exx.

Subfamily Scarabaeinae LATREILLE, 1802

Tribe Onitini LAPORTE, 1840

Genus *Onitis* FABRICIUS, 1798

13. *Onitis humerosus* (PALLAS, 1771)

Distribution: ASIA: Afghanistan, China, Iran, Israel, Jordan, Kazakhstan, Kyrgyzstan, Lebanon, Pakistan, Russia, Syria, Tadjikistan, Turkey, Turkmenistan, Uzbekistan; EUROPE: Armenia, Azerbaijan, Cyprus, Georgia, Russia.

Material examined: Semnan prov., Damghan, VI 2011, 1 ex.; Razavi Khorasan prov., Neyshaboor, VI 2012, 2 exx.

Tribe Onthophagini BURMEISTER, 1846

Genus *Onthophagus* LATREILLE, 1802

14. *Onthophagus (Palaeonthophagus) coenobita*
(HERBST, 1783)

Distribution: ASIA: Israel, Turkey, Turkmenistan; EUROPE: Albania, Armenia, Austria, Belgium, Bosnia-Herzegovina, Bulgaria, Byelorussia, Croatia, Czech Republic, Denmark, France, Germany, Georgia, Great Britain, Greece, Hungary, Italy, Kosovo, Latvia, Lithuania, Luxembourg, Macedonia, Moldavia, Montenegro, Netherlands, Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey, Ukraine.

Material examined: Gilan prov., Talesh, IX 2013, 2 exx. This is the first record for Iran.

15. *Onthophagus (Palaeonthophagus) basipustulatus* HEYDEN, 1889

Distribution: ASIA: Afghanistan, Kyrgyzstan, Syria, Tadjikistan, Turkey, Uzbekistan, EUROPE: Armenia, Azerbaijan.

Material examined: Northern Khorasan prov., Farooj, V 2011, 3 exx. This is the first record for Iran.

16. *Onthophagus (Furconthophagus) furcatus*
(FABRICIUS, 1781)

Distribution: ASIA: Iran, Iraq, Israel, Albania, Kazakhstan, Russia, Saudi Arabia, Syria, Turkey, Turkmenistan; EUROPE: Armenia, Austria, Azerbaijan, Belgium, Bosnia-Herzegovina, Bulgaria, Byelorussia, Croatia, Czech Republic, France,

Germany, Georgia, Great Britain, Greece, Hungary, Italy, Kosovo, Macedonia, Moldavia, Montenegro, Morocco, Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Switzerland, Turkey, Ukraine.

Material examined: Ardabil prov., Germe, VI 2012, 4 exx.; Hamadan prov., Razan, VI 2015, 2 exx.



Fig. Map of Iran with provincial boundaries, and distribution of Scarabaeidae of this research.
The numerals refer to species number in the paper.

Ryc. Mapa Iranu z podziałem na prowincje i zaznaczonymi stanowiskami badawczymi.
Numery odpowiadają numeracji gatunków w pracy

Discussion

The result of the present research indicates that the fauna of Iranian Scarabaeidae is rich and on the other hand unknown. In this faunistic investigation, 16 species distributed in 13 provinces (Fig.) were identified. With attention to four new records and the new contribution (SAKENIN & al. 2018; SAMIN & al. 2018; NADERIAN & al. 2021), the total number of Iranian Scarabaeidae species reached to 522. Since Iran is a large country (1,623,779 km²) with various geographical regions and climates, so we expect much more new data in different regions after faunistic surveys systematically.

Acknowledgements

The authors are grateful to Dr. B.C. RATCLIFFE (University of Nebraska, USA) and Dr. W.B. JĘDRYCKOWSKI (University of Ecology and Management, Warszawa, Poland) for cooperation in this research. This project was supported by Islamic Azad University and Pars Gostaran Zharf Tourism Company (Tehran, Iran).

STRESZCZENIE

Na podstawie analizy materiałów zgromadzonych w latach 2010-2016 w różnych prowincjach Iranu wykazano 16 gatunków Scarabaeidae należących do 10 rodzajów z 5 podrodziny: Aphodiinae, Cetoniinae, Melolonthinae, Rutelinae i Scarabaeinae. Cztery gatunki nie były do tej pory podawane z Iranu. Są to: *Bodilopsis sordida sordida* (FABRICIUS, 1775) i *Phalacrothous biguttatus* (GERMAR, 1824) z podrodziny Aphodiinae oraz *Onthophagus (Palaeonthophagus) basipustulatus* HEYDEN, 1889 i *Onthophagus (Palaeonthophagus) coenobita* (HERBST, 1783) z podrodziny Scarabaeinae. Zamieszczono mapkę przedstawiającą lokalizację stanowisk badawczych dla poszczególnych gatunków.

REFERENCES

- BARARI H. 2001. Insects of Iran. The list of Coleoptera in the insect collection of Plant Pests and Diseases Research Institute. Coleoptera (XXIV): Scarabaeoidea. group: Laparosticti. Fam.: 39, 40, 44, 46 (Geotrupidae, Hybosoridae, Orphnidae, Scarabaeidae and Aphodiidae). Plant Pests & Diseases Research Institute Tehran, 7: 1-64.
- BARARI H., KHARRAZI PAKDEL A., BOROUMAND H. 1998a. A faunistic survey of Laparosticti group (Col.: Scarabaeidae) in Karaj area. Proceedings of the 13th Iranian Plant Protection Congress: 202.
- BARARI H., KHARRAZI PAKDEL A., BOROUMAND H. 1998b. A faunistic survey of Pleurosticti group (Col.: Scarabaeidae) in Karaj area. Proceedings of the 13th Iranian Plant Protection Congress: 203.
- BARAUD J. 1968. Contribution à la faune de l'Iran. 10. Coleoptera: Scarabaeoidea. Annales de la Société Entomologique de France (n.s.), 4: 915-925.
- BORTESI O., ZUNINO M. 1974. Les résultats de l'expédition entomologique tchécoslovaque-iranienne l'Iran en 1970. No. 10: Les Onthophagus du sous-genre *Euonthophagus* Blath. (Coleoptera, Scarabaeoidea). Acta Entomologica Musei Nationalis Pragae, 6: 105-107.
- BUNALSKI M., GHAHARI H. 2015. Contribution to the knowledge Scarabaeoidea (Coleoptera) of Guilan province, Northern Iran. Wiadomości Entomologiczne, 34 (1) 19-28.
- BUNALSKI M., GHAHARI H. 2017. Contribution to the study of darkling beetles (Coleoptera: Tenebrionidae) from Golestan province, Northern Iran. Wiadomości Entomologiczne, 36 (1): 25-31.
- BUNALSKI M., SAMIN N., GHAHARI H., HAWKESWOOD T.J. 2014a. Contributions to the knowledge the scarab beetles of Golestan province, Northern Iran with checklist of Iranian Scarabaeoidea (Coleoptera). Polish Journal of Entomology, 83: 141-170.
- BUNALSKI M., SAMIN N., GHAHARI H. 2014b. Contribution to the knowledge of darkling beetles (Coleoptera: Tenebrionidae) from the cotton fields of Iran. Wiadomości Entomologiczne, 33 (2) 117-125.
- BUNALSKI M., SAMIN N., GHAHARI H. 2014c. Contributions to the knowledge on the darkling beetles (Coleoptera: Tenebrionidae) of Khorasan and Semnan Province, Iran. Wiadomości Entomologiczne, 33 (3): 188-193.
- BUNALSKI M., SAMIN N., GHAHARI H. 2016. A contribution to the faunistic study of Scarabaeoidea (Coleoptera) from Mazandaran province, northern Iran. Wiadomości Entomologiczne, 35 (1) 31-40.
- DAVOODI S., HOSSEINI R., HAJIZADEH J., BARIMANI H., MONTREUIL O. 2017. Report of eight species of dung beetles (Col.: Scarabaeidae) from Guilan province. Plant Pest Research, 7 (2): 83-86.
- GHAHARI H., BUNALSKI M. 2011. A study on darkling beetles (Coleoptera: Tenebrionidae) from south and southeastern Iran. Linzer Biologischer Beiträge, 43 (2): 1285-1290.
- GHAHARI H., BUNALSKI M., TABARI M., OSTOVAN H. 2010a. Contribution to the knowledge of darkling beetles (Coleoptera: Tenebrionidae) from Iranian rice fields and surrounding grasslands. Polish Journal of Entomology, 79 (1): 81-90.
- GHAHARI H., BUNALSKI M., HAVASKARY M., OSTOVAN H., SAMIN N. 2010b. Contributions to the knowledge the darkling beetles (Coleoptera: Tenebrionidae) of Arasbaran, Northwestern Iran. Polish Journal of Entomology, 79 (4): 463-468.
- GHAHARI H., BUNALSKI M., TABARI M., OSTOVAN H. 2011. Scarab beetles (Coleoptera: Scarabaeidae) collected in Iranian rice fields and surrounding grasslands. Polish Journal of Entomology, 80: 465-474.
- GHAHARI H., MAKHAN D., HAWKESWOOD T.J. 2012. A contribution to the knowledge of Carabidae, Staphylinidae and Tenebrionidae (Coleoptera) from the desert regions of Semnan province, Iran. Calodema, 207: 1-4.
- GUNTER N.L., WEIR T.A., ŚLIPIŃSKI A., BOCAK L., CAMERON S.L. 2016. If dung beetles (Scarabaeidae: Scarabaeinae) arose in association with dinosaurs, did they also suffer a mass co-extinction at the K-Pg boundary? PLOS ONE DOI:10.1371/journal.pone.0153570
- KABAKOV O.N. 1982. K faune plastinchatousykh zhukov roda *Onthophagus* Latr. (Coleoptera, Scarabaeidae) Sredney Azii, Afganistana i Irana. Trudy Zoologicheskogo Instituta Akademii Nauk SSSR, 110: 8-19.

- KEITH D., AHRENS D. 2002. Zur Kenntnis der Arten der Gattung *Maladera* Mulsant et Rey, 1871 aus dem Iran (Coleoptera: Scarabaeidae, Sericinae). *Entomologische Zeitschrift*, **112** (7): 194-197.
- KEITH D., ULIANA M. 2008. Contribution to the knowledge of Iranian *Pygopleurus* MOTSCHULSKY, with description of *P. rapuzzii* n. sp. (Coleoptera: Scarabaeoidea: Glaphyridae). *Zootaxa*, **1825**: 59-64.
- KRELL F.T. 2000. The fossil record of Mesozoic and Tertiary Scarabaeoidea (Coleoptera: Polyphaga). *Invertebrate Taxonomy*, **14**: 871-905.
- KRELL F.T. 2007. Catalogue of fossil Scarabaeoidea (Coleoptera: Polyphaga) of the Mesozoic and Tertiary – Version 2007. Denver Museum of Nature & Science Technical Report 2007, **8**: 1-79.
- LÖB, I., LÖBL D. 2016. Catalogue of Palaearctic Coleoptera. Volume 3, Scarabaeoidea, Scirtoidea, Dascilloidea, Buprestoidea, Byrrhoidea. Revised and updated edition. Leiden – Boston, Brill. 961 pp.
- LÖBL I., SMETANA A. 2006. Catalogue of Palaearctic Coleoptera. Volume 3. Scarabaeoidea – Scirtoidea – Dascilloidea – Buprestoidea – Byrrhoidea. Apollo Books, Stenstrup. 690 pp.
- MATTHEWS E.G., LAWRENCE J.F., BOUCHARD P., STEINER W.E., ŚLIPIŃSKI S.A. 2010. Tenebrionidae LATREILLE, 1802. (pp. 574-659). [In:] LESCHEN R.A.B, BEUTEL R.G., LAWRENCE J.F. (eds.): Handbook of zoology. Arthropoda: Insecta. Part 38. Coleoptera, beetles. Vol. 2. Morphology and systematics (Elateroidea, Bostrichiformia, Cucujiformia partim). Walter de Gruyter, Berlin.
- MODARRES AWAL M. 1997. Family Scarabaeidae. (pp. 186-192) [In:] MODARRES AWAL M. (ed.): List of agricultural pests and their natural enemies in Iran. Second edition. Ferdowsi University of Mashhad Press. 429 pp.
- MODARRES AWAL M. 2006. Preliminary studies on Scarabaeoidea (Coleoptera) fauna of Razavi Khorasan province of Iran. *Türkiye Entomoloji Dergisi*, **30** (3): 163-172.
- MODARRES AWAL M. 2012. Family Scarabaeidae. (pp. 357-366) [In:] MODARRES AWAL M. (ed.): List of agricultural pests and their natural enemies in Iran. Third edition. Ferdowsi University of Mashhad Press. 759 pp.
- MONTREULL O., SERRI S. 2007. Description of a new *Pygopleurus* species (Col.: Glaphyridae), with a list of recorded species of the genera *Pygopleurus* and *Eulasia* from Iran. *Journal of Entomological Society of Iran*, **26** (2): 1-6.
- MORADI GHARAKHLOO M., ZIANI S. 2009. Occurrence of scarab beetles inside rodent burrows in some parts of Iran (Coleoptera Scarabaeidae). *Zoology in the Middle East*, **46**: 95-98.
- MORADI GHARAKHLOO M., ZIANI 2010. Iranian dung beetles (Coleoptera, Scarabaeidae) associated with rodent burrows: list of collected species and some additional comments. *Boletín de la Sociedad Entomológica Aragonesa*, **47**: 165-170.
- MOWLAVI G., MIKAEILI E., MOBEDI I., KIA E.B., MASOOMI L., VATANDOOST H. 2008. Scarab beetle (Coleoptera: Scarabaeidae) fauna in Ardabil province, north west Iran. *Journal of Arthropod-Borne Diseases*, **2** (2): 35-41.
- NADERIAN H., SAKENIN H., HAWKESWOOD T.J., SAMIN N., JĘDRYCKOWSKI W.B. 2021. A faunistic study on some families of Coleoptera (Insecta) from Iran. *Calodema*, **871**: 1-2.
- NIKODYM M., KRÁL D. 1998. Results of the Czech Biological Expedition to Iran. Part 3. Coleoptera: Glaphyridae and Scarabaeidae: Sericinae. *Acta Societatis Zoologicae Bohemoslovenicae*, **62**: 343-351.
- PETROVITZ R. 1954. Ergebnisse der Österreichischen Iran-Expedition 1949/50. Coleoptera, IV. Teil. Scarabaeidae. Sitzungsberichte der Österreichischen Akademie der Wissenschaften, **163**: 261-275.
- PETROVITZ R. 1958. Beitrag zur Kenntnis der Scarabaeiden-Fauna des Iran. *Stuttgarter Beiträge zur Naturkunde*, **8**: 1-12.
- PETROVITZ R. 1980. Österreichische entomologische expeditionen nach Persien und Afghanistan. Beiträge zur Coleopteren-fauna. Teil XII. Weiteres über Lamellicornia aus Iran. *Annalen des Naturhistorischen Museum in Wien*, **83**: 597-638.
- RATCLIFFE B.C., PAULSEN M.J. 2008. The Scarabaeoid beetles of Nebraska. *Bulletin of the University of Nebraska State Museum*, **22**: 1-569.
- SAKENIN H., SAMIN N. 2015. A study on the fauna of Scarabaeoidea (Coleoptera) from Golestan province, northern Iran. *Entomofauna*, **36**: 513-520.
- SAKENIN H., SAMIN N., HÁVA J., BUNALSKI M., NADERIAN H., KUBISZ D. 2018. A contribution to the study of Iranian Coleoptera with some new records. *Wiadomości Entomologiczne*, **37** (2): 96-109.
- SAMIN N., BUNALSKI M., KUBISZ D., HÁVA J., NAVAEIAN M., OTERO J.C., HAWKESWOOD T.J., ZHOU H., SAKENIN H., JĘDRYCKOWSKI W.B. 2018. Contributions to the knowledge of the distribution of select Coleoptera families from Iran. *Wiadomości Entomologiczne*, **37** (4): 197-209.
- YARMAND H., MEHRABI A., MONTREULL O. 2012. Faunistic survey of the some scarab beetles in Ghazvin province. *Proceedings of the 20th Iranian Plant Protection Congress*: 129.
- ZAIRI M. 1976. List of Scarabaeidae from province of Fars. Part 4. *Journal of Entomological Society of Iran*, **3** (1-2): 125-127.
- ZIANI S. 2006. Remarks on some near eastern *Euonthophagus* species with the description of two new species from Iran (Insecta Coleoptera Scarabaeidae: Onthophagini). *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, **23**: 95-130.
- ZIANI S. 2011. A new Iranian *Onthophagus* species belonging to the *O. semicornis* group (Coleoptera: Scarabaeidae: Onthophagini). *Zootaxa*, **2907**: 47-50.
- ZIANI S., GUDENZI I. 2000. About some *Onthophagus* Latreille, 1802 (sensu lato) collected during a trip to Iran (Insecta, Coleoptera, Scarabaeidae). *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, **13** (Suppl.): 17-26.
- ZIANI S., GUDENZI I. 2006. Studies on Palearctic *Onthophagus* associated with burrows of small mammals. I. *O. furciceps*, *O. kindermanni*, *O. vitulus* and closely related species (Coleoptera Scarabaeidae). *Bollettino della Società Entomologica Italiana*, **138** (3): 207-248.
- ZIANI S., GUDENZI I. 2007. Studies on Palearctic *Onthophagus* associated with burrows of small mammals. II. *O. semicornis* and closely related species (Coleoptera: Scarabaeidae: Onthophaginae). *Folia Heyrovskyana, Ser. A*, **15** (2): 89-114.
- ZIANI S., GUDENZI I. 2009. Studies on Palearctic *Onthophagus* associated with burrows of small mammals. III. *O. aerarius* and closely related species (Coleoptera Scarabaeidae). *Bollettino della Società Entomologica Italiana*, **141** (1): 29-44.
- ZIANI S., MORADI GHARAKHLOO M. 2010. Studies on Palearctic *Onthophagus* associated with burrows of small mammals. IV. A new Iranian species belonging to the furciceps group (Coleoptera, Scarabaeidae, Onthophagini). *ZooKeys*, **34**: 33-40.