

Superficially described and ignored for 92 years, rediscovered and emended: *Apodera angatakere* (Amoebozoa: Arcellinida: Hyalospheniformes) is a new flagship testate amoeba taxon from Aotearoa (New Zealand)

Clément Duckert¹  | Quentin Blandenier¹  | Michelle McKeown^{2,3} | Holden Hohaia⁴ | Stefan Luketa^{1,5}  | Janet Wilmshurst^{2,6}  | Enrique Lara⁷  | Edward A. D. Mitchell^{1,8} 

¹Laboratory of Soil Biodiversity, University of Neuchâtel, Neuchâtel, Switzerland

²Manaaki Whenua - Landcare Research, Lincoln, New Zealand

³Wallbridge Gilbert Aztec, Christchurch, New Zealand

⁴Manaaki Whenua/Landcare Research, Wellington, New Zealand

⁵Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Novi Sad, Serbia

⁶School of Environment, The University of Auckland, Auckland, New Zealand

⁷Real Jardín Botánico, CSIC, Madrid, Spain

⁸Jardin Botanique de Neuchâtel, Neuchâtel, Switzerland

Correspondence

Clément Duckert, Laboratory of Soil Biodiversity, University of Neuchâtel, Neuchâtel, Switzerland.
Email: clement.duckert@unine.ch

Funding information

Comunidad de Madrid, Grant/Award Number: 2017-T1/AMB-5210; Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung, Grant/Award Number: 31003A_163254; MYXOTROPIC VI, Grant/Award Number: PGC2018-094660-B-I00

Abstract

Eukaryotic microbial diversity is known to be extensive but remains largely undescribed and uncharted. While much of this unknown diversity is composed of inconspicuous flagellates and parasites, larger and morphologically distinct protists are regularly discovered, most notably from poorly studied regions. Here we report a new flagship species of hyalospheniid (Amoebozoa; Arcellinida; Hyalospheniformes) testate amoeba from New Zealand and an unusual story of overlooked description under a preoccupied name and subsequent oversight for nearly one century. Through a process involving The Māori Language Commission, we named the species *Apodera angatakere*, meaning “a shell with a keel.” This species resembles *Apodera vas* but differs by the presence of a distinctive hollow keel. Cytochrome Oxidase Subunit 1 (COI) sequence data show that this species forms a distinct clade nested within genus *Apodera*. This conspicuous species is so far known only from New Zealand and is restricted to peatlands. It is one of the few examples of endemic microorganisms from this biodiversity hotspot and biogeographer's paradise. As over 90% of New Zealand's peatlands have been lost since European colonization and much of the remaining surfaces are threatened, *Apodera angatakere* could be a flagship species not only for microbial biogeography but also for island biodiversity conservation.

KEYWORDS

Apodera vas, biodiversity conservation, biogeography, *Gibbocarina*, Gondwana, Hyalospheniidae, Māori language and culture, microbial diversity, peatlands, *Sphagnum*, taxonomy

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Journal of Eukaryotic Microbiology* published by Wiley Periodicals LLC on behalf of International Society of Protistologists.

INTRODUCTION

Anyone with a sense for adventure may dream about the age of discovery (15th–17th century period), during which time the New World was (re)discovered and many new inventions made that changed our perception of the world. Traveling back in time to experience such thrill first hand is, unfortunately, not possible. However, when we consider the lack of knowledge of the microbial world, every sample can be an unexplored island for the curious protistologist with an inclination for natural history.

The invention of the microscope in the late 16th century paved a new era of observation and discovery by making the microbial world accessible to scientific researchers. With subsequent improvements in optics, along with the exploration of various habitats and regions of the World a broad range of microbes was gradually described, gaining momentum in the 19th and early 20th century by pioneers such as Christian Gottfried Ehrenberg, Ernst Haeckel, Josef Leidy and Eugène Penard (Ehrenberg, 1838; Haeckel, 1899–1904; Leidy, 1879; Penard, 1902). More recent developments such as high throughput sequencing of environmental DNA (eDNA) are allowing large amounts of genetic data to be generated revealing a huge, mostly unknown, diversity of microorganisms including bacteria (Delgado-Baquerizo et al., 2018), fungi (Nilsson et al., 2019), protists (Mahé et al., 2017) and micro-Metazoa (van den Hoogen et al., 2019). The magnitude of protist diversity is now believed to exceed that of bacteria (Delgado-Baquerizo et al., 2018) and fungi (Tedersoo et al., 2014), at least in the ocean (de Vargas et al., 2015) but recent work also suggests that this may be the case for terrestrial soil (Geisen et al., 2018). This diversity is confirmed by numerous examples of cryptic diversity reported in classical DNA barcoding studies (Fontaneto et al., 2008; Kosakyan et al., 2012; Kumar & Foissner, 2016; Mann, 2010; Pawlowski et al., 2008), as was first documented for insects (Hebert et al., 2004).

These newly available molecular identification tools have stimulated another golden age of protist discovery, comparable to that triggered by the invention of the microscope (Geisen et al., 2015; Heger et al., 2014; Mitchell, 2015). Novel protist diversity can now be found virtually anywhere. DNA barcoding studies are regularly revealing the existence of numerous species within known groups, questioning long-held ideas about their geographical distribution or ecology that were formerly based on only morphological similarities (Singer et al., 2015; de Vargas et al., 1999). Taxa known from ancient morphological descriptions are now being redescribed and placed in the tree of life using the full range of molecular and detailed microscopy techniques (Burki et al., 2019; Dumack et al., 2016, 2018; Lax et al., 2018).

Finding novel protist diversity can be expected when exploring under-studied regions (Yubuki et al., 2009) or poorly studied groups (Dumack et al., 2018; Siemensma et al., 2020). Although there is a strong sampling bias in favor of Europe and North America where most protistologists work, detailed studies of other regions frequently report many new species, as illustrated in South America, Australia, Africa, and Antarctica (Fernández et al., 2015; Foissner, 1997). Twenty-four years ago, Foissner estimated that at least 70% of the world soil ciliates remained to be described (Foissner, 1997), which is still an underestimation of their true diversity as revealed by eDNA sequencing (Boscaro et al., 2018).

New Zealand is a biogeographer's paradise but has barely been explored for soil protists and especially testate amoebae. Eugène Penard reported 45 testate amoeba species, including three new taxa, *Nebela certesi*, *Hyalosphenia cockayni*, and *Nebela griseola*, from samples collected during the 1907–09 British Antarctic Expedition (Penard, 1911). All three species have since been transferred to new genera, *Certesella certesi*, *Alocodera cockayni*, and *Physochila griseola*. Since then, to our knowledge 19 studies were published on the diversity and ecology of testate amoebae of New Zealand, reviewed in McKeown et al. (2021). A limited number of studies described new species for science, new records for New Zealand or, more recently, focused primarily on the community ecology of testate amoebae and their application in paleoecology (Bamforth, 2015; Charman, 1997; Deevey, 1955; Hoogenraad & de Groot, 1948; McGlone & Wilmshurst, 1999; McKeown et al., 2019; van Oye, 1956; Stout, 1984; Wilmshurst et al., 2002, 2003; Winterbourn & Brown, 1967). However, there has been no modern study on testate amoeba taxonomy that incorporates molecular, morphometrical and ecological data, which are required to accurately describe new species (Lara et al., 2020). As a consequence, it is most likely that many New Zealand species remain overlooked or undescribed.

Here we report *Nebela penardi* from *Sphagnum* peatlands in New Zealand for the first time since its description (Brehm, 1928), a highly conspicuous species that had been superficially described and completely overlooked since then. Based on Cytochrome Oxidase Subunit 1 (COI) data we determine its phylogenetic position within genus *Apodera* and discuss its potential as flagship species for both biogeographical and ecological surveys. Furthermore, as *Nebela penardi* is an invalid name and there is no known available synonym we formally rename it *Apodera angatakere* in collaboration with Te Taura Whiri I te Reo Māori (The Māori Language Commission).

TABLE 1 Samples collected in New Zealand containing *Apodera angatakere*

Sample codes	Coord decimal ° Lat (dec °)	Coord decimal ° Long (dec °)	Elevation (m)	Sampling location	Habitat	Sample type
EM-2540	-39.2550°	174.0431°	921	Egmont National Park, Mt. Taranaki, Pouakai Circuit	Large peatland in saddle between Henry Peak and Taranaki; hollow	<i>Sphagnum</i>
EM-2616	-41.6986°	172.1594°	1185	Old Ghost Road trail, shore of "Ghost Lake," km 30, Westland	<i>Sphagnum</i> peatland on lake shore	<i>Sphagnum</i>
EM-2638	-50.8126°	166.0653°	85	Auckland Island	Low <i>Metrosideros</i> forest - moss tuft	Mosses
EM-2661	-45.3892°	167.5763°	1214	Kepler Track, clockwise, day 2	Alpine terrace with stream, ponds, fens and <i>Sphagnum</i> patches on side (slightly elevated). Collected from lower, more minerotrophic micro-site	Brown mosses
EM-2670	-44.7271°	168.1728°	1266	Routeburn Great Walk trail—pass above Lake Harris, outlet of upper of the two "larger" lakes	<i>Sphagnum</i> patch on side of lake outlet	<i>Sphagnum</i>
EM-2699; X18/109/ SST1	-43.2173°	170.2843°	120	Okarito Lagoon, Westland	Ombrotrophic bog in dense forest	<i>Sphagnum</i>
EM-2705; X18/10/ SST1	-43.5989°	169.6154°	5	Bruce Bay, South Westland	<i>Sphagnum</i> lawn	<i>Sphagnum</i>
EM-2765	-45.7958°	170.4829°	710	Swampy Summit, NW of Dunedin	Tarn edge: <i>Sphagnum</i> growing in water	<i>Sphagnum</i>
EM-2785	-44.8618°	167.8245°	288	Milford Track—day 2—Hidden lake, Fiordland	Wetland near lake with bushes, <i>Sphagnum</i> and other mosses	<i>Sphagnum</i> and other mosses
EM-2693	-42.9071°	171.5587°	914	Arthur's Pass, South Island	Wetland area with ponds, fens and patches of <i>Sphagnum</i>	<i>Sphagnum</i>
X20/8	-42.9156°	171.5524°	920	Margaret's Tarn, off the Mt Bealey Track between Arthur's Pass and Mt. Rolleston, South Island	Tarn (small closed-basin lake)	<i>Sphagnum cristatum</i> & <i>S. falcatulum</i>

MATERIAL AND METHODS

Sampling, single cell isolation, and morphometry

Samples of *Sphagnum*, other mosses (unidentified brown mosses from wetlands) and forest litter were collected from numerous forests, peatlands, and fens across New Zealand's North and South Islands and Campbell and Auckland Islands as well as Macquarie Island as part of a wider project on the diversity of testate amoebae in New Zealand and peri-Antarctic islands. Testate amoebae that could be assigned to *Apodera angatakere* were found in 11 samples (Table 1). Testate amoebae were isolated for further analyses from four samples (Figs S1–S8):

1. EM 2540: *Sphagnum* mosses from a hollow in Ahukawakawa Swamp, along the Pouakai crossing trail, on the saddle between Taranaki Maunga and Pouakai Hut, New Zealand's North Island. Coord: Lat -39.255058° , Long 174.043106° , Elevation: 921 m a.s.l.
2. EM 2543: Forest litter and humus from the Northern foothills of Taranaki Maunga, New Zealand's North Island. Coord. Lat -39.22505° , Long 174.122842° , Elevation: 478 m a.s.l.
3. EM 2935: *Sphagnum cristatum* and *S. falcatulum* from Margaret's Tarn, off the Mt Bealey Track between Arthur's Pass and Mt. Rolleston on the South Island. Coordinates: Lat -42.915655° , Long 171.552487° , Elevation: 920 m a.s.l.
4. EM 2764: *Sphagnum falcatulum* from the north shore of Macquarie Island, collected along Featherbed Track, between Handspike Corner and West Beach. Coordinates: Lat -54.49928200° , Long 158.90512300° , Elevation ~ 5 m a.s.l.

The samples were placed in closed plastic sampling bags. Once in the laboratory the samples were sub-sampled and material was placed in a container and shaken in tap water for 1–2 min and then filtered over a 200- μ m mesh filter. The filtrate was observed under an inverted microscope and testate amoeba specimens that could be assigned to genus *Apodera* were isolated with a narrow diameter pipette and transferred to a new slide with distilled water to remove any unwanted material. Pictures were taken for each individual specimen used for DNA barcoding, which was then transferred to a PCR tube with 50 μ l of guanidinium thiocyanate solution (Chomczynski & Sacchi, 1987). Additional living or dead individuals were used for scanning electron microscopy (SEM) imaging. These specimens were further cleaned in ethanol and transferred to a SEM stub on which a carbonate filter had been fixed. The samples were then sputter coated with gold. Images were taken with a Hitachi TM 3030 plus scanning electron microscope operating at 5–15 kV. SEM images were taken at the electron microscopy facility of Manaaki Whenua/Landcare Research, Lincoln, New Zealand.

Measurements were taken on light microscopy and SEM images for each observed specimen: test length, test width, aperture long axis, length of the neck, width at the base of the neck, maximum width of the neck, and length of the main (aboral) part of the test (Figure 1). Depth could only be measured accurately on four individuals resting on their side, but it could be estimated based on the height of the focus plane for individuals resting flat.

DNA extraction, PCR amplification, and DNA sequencing

DNA extraction was performed with a guanidinium thiocyanate solution following the protocol of Duckert et al. (2018) adapted from Chomczynski and Sacchi (1987). PCR of partial mitochondrial cytochrome oxidase I (COI) gene were processed using a nested PCR protocol. The first PCR was conducted with the eukaryote-general primers LCO1490 (GGT CAA CAA ATC ATA AAG ATA TTG G) and HCO2198 (TAA ACT TCA GGG TGA CCA AAA AAT CA) (Folmer et al., 1994) under the following conditions: Denaturation step at 95°C for 5 min, then 45 cycles with a denaturation step at 95°C for 15 s, an hybridization step at 43°C for 15 s, an elongation step at 72°C for 1 min, and a final elongation at 72°C for 10 min. *Apodera vas* specimens EM2543-15, EM2543-22 and *A. angatakere* EM2540-14, EM2540-19 were then amplified with LCO1490 and ApocoxR (Kosakyan et al., 2012) under the following conditions: Denaturation step at 95°C for 5 min, then 45 cycles with a denaturation step at 95°C for 30 s, an hybridization step at 50°C for 30 s, an elongation step at 72°C for 1 min, and a final elongation at 72°C for 10 min. On the other hand, *A. angatakere* EM2540-11 and *A. vas* EM2764-2 were amplified using Arcelcox1F (Kosakyan et al., 2012) and Apo2F (a newly designed forward primer: TGG AAT TAG CAT ATC MGG AAT T) respectively, with HCO2198 under the same condition as the first PCR, with the only exception of an hybridization step of 45°C for the former taxa. Positive PCR products were directly purified using Milipore kit and sent for sequencing with an ABI3730XL DNA sequencer (Applied Biosystems) at Macrogen, Amsterdam NL. Sequences




FIGURE 1 Schematic drawing of *Apodera angatakere* and codes of the measurements taken. (A) Length of the test; (B) width of test; (C) width of the pseudostome; (D) length of the neck; (E) maximal width of the neck; (F) width of the test at the constriction; (G) length of the test without the neck

were deposited in GenBank with the accession numbers MZ615186–MZ615191. Light microscopy pictures of the corresponding cells are shown in Figure 2.

Phylogenetic analysis

All sequences obtained were aligned manually using BioEdit (Hall, 1999) together with an exhaustive reference database composed of three previously published sequences of genus *Alocodera* and three of genus *Padaungiella* (Kosakyan et al., 2012) retrieved from the GenBank database (Table S2). The tree was rooted with 10 Hyalospheniformes sequences, corresponding to eight different genera, published in four previous studies (i.e., Duckert et al., 2018; Kosakyan et al., 2012, 2013; Singer et al., 2015). Phylogenetic reconstruction was conducted on the complete LCO-HCO fragment (i.e., 655 nucleotides) using the CIPRES Science Gateway V. 3.3 (Miller et al., 2010). A maximum likelihood phylogenetic tree was built using the RAxML v.8.2.10 algorithm (Stamatakis, 2014) with the GTR + GAMMA model and automatic bootstrapping halt. We further evaluated node robustness using a Bayesian approach with MrBayes (Ronquist & Huelsenbeck, 2003) using the GTR + GAMMA model with default settings on two independent runs sampled every 100 generations. The analysis was automatically stopped when convergence was reached after 260, 100 generations resulting in 5202 trees of which 25% were discarded as the burn-in.

Collaborating with Māori in the naming process

Naming species that are discovered in regions where indigenous people live should respect the cultural principles. According to the Waitangi tribunal Wai 262 report (Waitangi Tribunal Report Wai 11, 1986), Māori retain “kaitiakitanga” (guardianship) over “taonga” (treasured) species and the cultural relationship between kaitiaki (guardian) and taonga (treasured) species is entitled to reasonable protection. According to Veale et al. (2019), this implies that there is an imperative to conduct processes such as formally naming and describing species of cultural relevance to Māori in partnership with Māori. In agreement with this principle, as a new species name needed to be found, this was done in partnership with Te Taura Whiri I te Reo Māori (The Māori Language Commission). Our aim was to find a name that would be appropriate for the organism and ideally reflect the specific morphology: a flattened gourd with a constriction at the base of the neck and a hollow “keel.” The choice of a name was an iterative process involving Ngahiwi Apanui from Te Taura Whiri I te Reo Māori and all authors.




FIGURE 2 Top half: *Apodera angatakere* n. gen. n. sp. (A–C, E and F), five specimens from Ahukawakawa swamp, Taranaki Maunga, New Zealand's North Island (sample EM-2540): (A–C) three barcoded individuals, (D) Brehm's original drawing of *Apodera angatakere* (described as *Nebela penardi*) from Margaret's Tarn, Arthur's Pass, New Zealand's South Island, (E and F), two individuals from sample EM-2540 (LM and SEM, respectively). E is the holotype. Note the presence of a ca. 10 μm wide keel. All specimens illustrated here as well as in Figures S2–S8 were used for morphometrical analyses (Figure 1). Scale bars (20, 50, or 100 μm) are shown for all specimen but were not provided in the original description. Bottom half: *Apodera* *vas*. (G) barcoded specimen from Macquarie Island (sample EM-2764), (H–J) three specimens from forest litter collected on the lower slopes of Taranaki Maunga, New Zealand's North Island (sample EM-2543). (H and I) Two barcoded specimen, (J) SEM of a third individual; note the absence of a keel. The codes of the barcoded specimens are the same as in the phylogenetic tree (Figure 3)

TABLE 2 Summary morphometric statistics of *Apodera angatakere*. A: Length of the test; B: Width of test; C: Width of the pseudostome; D: Length of the neck; E: Maximal width of the neck; F: Width of the test at the constriction; G: Length of the test without the neck. Full details are given in Table S1

	a	b	c	d	e	f	g	a/b ratio	g/b ratio
<i>n</i>	62	60	59	62	61	61	63	60	60
Average	208.2	147.0	44.0	70.5	75.6	71.0	137.7	1.42	0.94
Median	209.0	148.0	44.0	70.7	75.9	72.0	137.7	1.42	0.94
Min	186.0	120.0	39.1	61.0	60.4	58.4	125.0	1.19	0.77
Max	225.9	167.0	50.2	79.3	87.6	79.1	148.9	1.67	1.10
SD	8.59	8.55	2.04	4.40	4.29	4.49	5.87	0.07	0.06
CV	4.12	5.82	4.64	6.23	5.68	6.32	4.26	5.19	5.95




FIGURE 3 Maximum likelihood phylogenetic tree of the Hyalospheniformes with a focus on *Apodera*, *Alocodera*, and *Padaungiella* based on COI gene sequences. Bootstrap values (bs) and Bayesian posterior probabilities (p.p.) are indicated respectively between branches. COI sequences from genera other than *Apodera* were retrieved from GenBank

RESULTS AND DISCUSSION

Morphological characterization of the tests

We found two distinct morphotypes that could be assigned to genus *Apodera* based on their lageniform shell composed of recycled xenosomes, with a clear constriction that separates the neck from the body. These two morphotypes could easily be distinguished based on their size and the presence of a hollow keel on the larger morphotype (Figure 2). The smaller morphotype (mean length: 147 μ m, data not shown) corresponds well to the description of *Apodera vas* (Certes) and our measurements fall in the middle of the range recorded for this taxon (Zapata & Fernandez, 2008). It must be noted that the high degree of variability in dimensions and shape reported for *Apodera vas* suggests the existence of a complex of distinct species (Penard, 1911; Zapata & Fernandez, 2008). Although, the size of the larger morphotype (mean length: 208 μ m) (Table 2) falls within the range recorded for *Apodera vas*, it can be easily differentiated

by the presence of a distinct hollow keel surrounding the main body, as in genus *Gibbocarina*. The margins of the neck range from almost straight to slightly concave with a bulge at the base. The margins of the neck are compressed, except in the basal bulge region, as an extension of the hollow keel. This is easily visible on SEM images but is less obvious, and possibly overlooked in light microscopy. This morphology matches well with the line drawing of the original description of *Nebela penardi* Brehm, 1928 (hereafter *Apodera angatakere*), which unfortunately lacked a scale or indication of size. The tests of *Apodera vas* were found in forest litter samples, brown mosses, and *Sphagnum* whereas tests of *Apodera angatakere* were almost exclusively found in *Sphagnum* samples collected in peatlands but also brown mosses in an alpine fen and mosses in low *Metrosideros* forest on Auckland Island.

Phylogenetic placement within Hyalospheniformes and parphyly of *Apodera vas*

We obtained partial COI sequences of 655 nucleotides from three specimens of *Apodera vas* and three of *Apodera angatakere*. Sequences of genus *Apodera* form a moderately supported clade (bs: 66; pp: 58; Figure 3) and together with the genera *Alocodera* and *Padaungiella* they form a highly supported clade that is sister to the clade formed by all the other genera of Hyalospheniformes used to build the tree.

While sequences of *Apodera angatakere* form a highly supported monophyletic clade, *A. vas* appears to be paraphyletic with one sequence branching at the base of *A. angatakere* and two other sequences forming a distinct clade within *Apodera*. Interestingly the two specimens of *A. vas* branching together were found in the same sample from Taranaki Maunga, New Zealand's North Island, whereas the one that branches at the base of *A. angatakere* was found on Macquarie Island. This strongly supports the hypothesis that *Apodera vas* is currently a complex of distinct species, possibly with distinct distribution patterns.

Taxonomic status of *Apodera angatakere* (*Nebela penardi*)

Apodera angatakere was first briefly reported as *Nebela penardi* in a publication focusing mostly on copepods (Brehm, 1928). Only two drawings were shown (Figure 2D) and no indication of size or scale was provided. Brehm had sent a moss sample containing testate amoebae to Eugène Penard who described the finding as follows: “It is similar to *Nebela vas* Certes (now *Apodera vas*), but differs therefrom in that the shell possesses a broad, hollow, and characteristic keel, which is always present and might well be regarded as a sufficient character to distinguish a new species. I have not been able to identify it with any known species.” (Brehm, 1928). As a result, Brehm proposed to name the species *Nebela penardi*.

To our knowledge there is no further mention of this species in the scientific literature and the species has been overlooked since then. Notably, Deflandre did not mention it in his 1936 monograph of the genus *Nebela* (Deflandre, 1936). Moreover, at the time of its description, the name *Nebela penardi* Brehm, 1928 was preoccupied by *Nebela penardi* Heinis, 1914. It is therefore a junior homonym of this species, which implies that *Nebela penardi* Brehm was unavailable and thus invalid.

As there is no known synonym for this taxon, we therefore rename it in agreement with ICZN article 60.3, and as we transfer it to a new genus both a new generic name and a new species epithet are required. Furthermore, as a side note, Heinis (Heinis, 1914) used the name *Nebela penardi* to erroneously rename a taxon that, according to Deflandre (1936), corresponds to *Nebela martiali* (now *Certesella martiali*). *Nebela penardi* Heinis is therefore a junior synonym of *Certesella martiali*.

A very similar species was described under the name *Nebela kenyana* by Didier Chardez from mountain lakes in Kenya at over 4000 m a.s.l. (Chardez, 1982). This species has a similar size and overall morphology including a hollow keel surrounding the fundus of the test and a distinct neck with swollen sides. However, it lacks the characteristic constriction at the base of the neck which defines *Apodera* and, in that, it is more similar to genus *Padaungiella*.

Choice of a species name—an iterative process involving the Māori Language Commission

The choice of a name was an iterative process involving all authors in collaboration with Te Taura Whiri I te Reo Māori. We aimed to find a species name that would be appropriate for the organism and ideally reflect the specific morphology which could be described in general words as a flattened gourd with a constriction at the base of the neck and a hollow “keel.” We also aimed to represent the Mauri (life force) of this unique species that was named by Māori for them as kaitiaki (guardian) of Aotearoa (New Zealand).

The options considered were:

1. An amoeba with a hard shell. According to the Reed Dictionary of Modern Māori, the word “amoeba” is translated as “pūora hurikē” and a “hard shell” is translated as “anga.” Combining the two with the word “whai” meaning “in possession of” leads to the name “Pūora hurikē whaianga.”
2. Considering that a shelled amoeba is analogous to a miniature snail, the second option was “A snail with a keel-like shell.” Snail and keel being respectively translated as “ngata” and “takere,” this would then form “Ngata anga whaitakere.”
3. An amoeba with a keel-like hull: Pūora anga whaitakere.
4. An amoeba with a keel: Pūora whaitakere.
5. A shell with a keel: Anga whaitakere.

The fifth option was chosen both for the way it sounded and the fact that the name would describe well the morphology of the amoeba. However, in a final discussion, Mr. Ngahiwi Apanui from the Māori Language Commission suggested that the “whai” could be dropped to shorten the name. The final chosen name was then «Angatakere», to be pronounced æŋætakerɛ (AN-GAH-TAH-KEH-REH).

Implications for biodiversity estimates and conservation

It is estimated that 10% of the New Zealand mainland was covered by wetlands before human arrival around 780 cal. Yr BP (Ausseil et al., 2011; McGlone, 2009), and while testate amoebae were too small to be recognized by Māori until the introduction of the microscope, the wetland habitats that supported these organisms were highly valued by Māori for centuries, as mahinga kai/hauanga kai (food gathering areas), rongoā (gathering plants for medicinal use) and for material resources. Wetland extent was minimally affected by early Māori settlers, although extensive deforestation of dryland forest by burning transformed the landscape (Argiriadis et al., 2018; McWethy et al., 2010; Perry et al., 2014). Wetland loss rapidly accelerated after European arrival in the 1800s, largely for agricultural development, and is now only 10% of the original extent (Ausseil et al., 2011). It is estimated that the 250,000 ha of wetlands that remain are under increasing pressure from drainage, area loss, fragmentation, grazing, fire, pollution, and climate change (Meyer et al., 2013; Robertson et al., 2019). The fact that we can still find novel diversity or diversity that has been very poorly described in such habitats makes them even more precious and worthier of conservation.

Describing the still mostly unknown diversity of protists requires a major effort in basic taxonomy (Heger et al., 2014). The magnitude of land-use changes and natural habitat destruction occurring throughout the world and the now well-established existence of restricted geographical distribution patterns in free-living protists (Foissner, 2008) implies that a large proportion of protist diversity will likely disappear before it can be described, and thus the conservation of protists should indeed be a priority (Cotterill et al., 2008; Qin et al., 2016). *Apodera angatakere* is a highly conspicuous genus of testate amoeba and has to date only been found in New Zealand. Large species are more likely to have restricted geographical distribution as shown empirically for terrestrial and subaquatic testate amoebae in the southern temperate and Antarctic zones (Wilkinson, 1994), at the global scale (Wilkinson, 2001; Yang et al., 2010) and confirmed by an atmospheric circulation modeling study (Wilkinson et al., 2012). Given its large size it is likely an endemic taxon and as such could represent the first documented microbial species for which New Zealand has a conservation responsibility at the global scale. *Apodera angatakere* could therefore be considered as a flagship species for microbial biogeography and conservation.

To discover, or not to discover: that is the question

Columbus thought he had discovered a new world hitherto unknown to Europeans. But it was later established that the Vikings had already made this discovery before him. And of course, the sheer notion of this major discovery disregarded the fact that native populations had already colonized the Americas millennia before Europeans were even able to conceive of the idea of sailing across the ocean.

This history is mirrored in the story of *Apodera angatakere*: Just like Columbus one of us (EM) thought he had discovered a new species only to be brought to the attention by another one of us (SL) that it had been previously described in a publication on microcrustaceans which had escaped the attention of previous researchers studying testate amoebae in New Zealand. But just as the Vikings did not establish a permanent settlement in America, Brehm's discovery was lost to science, or almost so.

The microbial world remains unknown for most of the people but charismatic groups such as testate amoebae are useful as messengers of the invisible dimension of nature's wonders and our impact on the biosphere. The fact

that *A. angatakere* is known only from New Zealand and is restricted to ecosystems that have been almost entirely destroyed since European colonization is a perfect illustration of the fact that many species are being lost before we even have a chance to describe them.

Taxonomic actions

Taxonomic summary:

Amorphea Adl et al. 2021

Amoebozoa Lhe 1913, sensu Cavalier-Smith 1998

Tubulinea Smirnov et al. 2005

Elardia Kang et al. 2017

Arcellinida Kent 1880

Diffflugina Meisterfeld 2002, sensu Kosakyan et al., 2016

Hyalospheniformes Lahr et al., 2019

Hyalospheniidae (Schultze 1877) Kosakyan et Lara 2012

Apodera angatakere (Brehm, 1928) Mitchell, Blandenier & Duckert 2021

1928 *Nebela penardi* Brehm

Icon: Brehm, 1928 Fig. 52

Description: Test composed of two clearly distinct parts, a subcircular, oval, or ellipsoidal, compressed posterior part (body) and a neck. The two parts separated by a deep constriction around the entire base of the neck. Sides of the neck straight to slightly concave with a bulge at the base in broad view. The margins of the neck sometimes compressed. Body almost circular. Dimensions based on 63 specimens: 53 individuals from Taranaki Maunga, North Island, four from the Old Ghost Road, South Island, and six from the Keppler Track, South Island: Length (min.–average–max.): 186–208–226 μm , width: 120–148–167 μm , pseudostome: 39–44–50 μm . Circa 80 μm in breadth.

Etymology: In the Mori language “angatakere” can be translated to “a keeled shell,” referring to the conspicuous keel present on the outline of the test.

Neotype: Brehm did not preserve any specimen, and the original type material is only represented by two simple drawings without indication of size. Because small variations in the morphology of the test can be used to distinguish closely related species, only high magnification microphotographs can be used to accurately represent a species and reliably distinguish it from taxa yet to be described (Duckert et al., 2020; Kosakyan et al., 2016). For this reason, we designate the specimen in pictures Figure 2E as the neotype. As we were unable to find unfractured tests at the previous type locality (Margaret's tarn, Mt. Rolleston near Arthur's pass, New Zealand South Island) the neotype has been designated among a population from Taranaki Maunga, New Zealand's North Island. However, tests from the previous type locality and the ones from Taranaki Maunga were similar in all points. A permanent slide has been deposited at the Natural History Museum of Neuchtel (Switzerland) with the ID 95-1.

New type locality: Ahukawakawa swamp, along the Pouakai crossing trail, on the saddle between Taranaki Maunga and Pouakai Hut, New Zealand's North Island. Coord: -39.255058° , 174.043106° , Elevation: 921 m a.s.l.

Geographical distribution: Known from New Zealand North and South Islands and Auckland Island. Likely also Campbell and Chatham Islands. We did not find it in Macquarie Island where all *Apodera* specimens lacked the characteristic hollow keel.

Habitat: *Sphagnum* and brown mosses in peatlands and alpine wetlands in New Zealand's North and South Islands, mosses in low *Metrosideros* forest (Auckland Island).

Remarks: We did not succeed in obtaining DNA sequences from material collected from the previous type locality (Margaret's tarn, Mt. Rolleston near Arthur's pass, New Zealand South Island) and found only fractured test unfit to be designated as the neotype, we thus chose a specimen from Taranaki Maunga, New Zealand's North Island as the neotype. Given that 13 distinct molecular clades were recorded within *Hyalosphenia papilio*, a common but smaller species commonly found in Holarctic *Sphagnum* peatlands (Heger et al., 2013; Singer et al., 2019), it is possible that several cryptic or pseudo-cryptic species exist within *A. angatakere* and that specimens from Margaret's tarn constitute a distinct species. If this were the case a new species would need to be described from Margaret's tarn, New Zealand South Island with its own type locality. The Auckland Island record is based only on microscopic observation (not illustrated) and is considered valid given the characteristic morphology. Nevertheless, it would need to be further confirmed by molecular data.

Three COI gene sequences of *Apodera angatakere* and *Apodera vas* (352–655 bp) were deposited in GenBank under the number MZ615186–MZ615188 and MZ615189–MZ615191, respectively.

ZooBank registration number: urn:lsid:zoobank.org:act:921D8CE1-EF0F-4839-B264-97ED438B5694.

ACKNOWLEDGMENTS

This work was initiated during the scientific leave of EM in New Zealand. EM thanks Manaaki Whenua—Landcare Research for support during this stay and the many colleagues, especially of the Long-Term Ecology group but also many others for highly stimulating and friendly discussions. He also thanks Phil Novis for allowing him to use the SEM facility, Rowan Buxton and Chris Morse for collecting samples from Margaret’s Tarn, Alex Fergus for collecting the sample from Macquarie Island, Linnéa Mitchell for assistance in collecting samples from Mt. Taranaki, and Carina Davis for performing some of the molecular work. We thank the Tasmanian Department of Primary Industries, Parks, Water and Environment, for granting the sampling permit for Macquarie (authority no. FL 17338 to EM) and the New Zealand Ministry for Primary Industries—Manatū Ahu Matua for delivering a Biosecurity clearance certificate for importing samples from Australia (BACC No.: B2018/110263). The samples from New Zealand were collected on public conservation estate under Manaaki Whenua—Landcare Research’s global concession (permit Number CA-31615-OTH). This work was also supported by the Swiss National Science Foundation 31003A_163254 to EL and QB. EL was also supported by an “Atracción de talento investigador” grant from the Community of Madrid, project 2017-T1/AMB-5210 and by the project MYXOTROPIC VI (PGC2018-094660-B-I00). Last but certainly not least, we would like to express our warmest thanks to Ngahiwi Apanui from The Māori Language Commission for fruitful discussion leading to the choice of the name “angatakere.”

ORCID

Clément Duckert  <https://orcid.org/0000-0001-9386-2950>
 Quentin Blandenier  <https://orcid.org/0000-0002-4297-0262>
 Janet Wilmshurst  <https://orcid.org/0000-0002-4474-8569>
 Enrique Lara  <https://orcid.org/0000-0001-8500-522X>
 Edward A. D. Mitchell  <https://orcid.org/0000-0003-0358-506X>

REFERENCES

- Argiriadis, E., Battistel, D., McWethy, D.B., Vecchiato, M., Kirchgeorg, T., Kehrwald, N.M. et al. (2018) Lake sediment fecal and biomass burning biomarkers provide direct evidence for prehistoric human-lit fires in New Zealand. *Scientific Reports*, 8, 12113.
- Ausseil, A.-G.-E., Dymond, J.R. & Weeks, E.S. (2011) Provision of natural habitat for biodiversity: quantifying recent trends in New Zealand. In: Grillo, O. & Venora, G. (Eds.) *Biodiversity loss in a changing planet*. London, UK: IntechOpen, pp. 201–220.
- Bamforth, S.S. (2015) Composition of soil testate amoebae communities: their structure and modifications in the temperate rain forests of New Zealand and Tasmania. *Journal of Eukaryotic Microbiology*, 62, 217–226.
- Boscaro, V., Santoferrara, L.F., Zhang, Q., Gentekaki, E., Syberg-Olsen, M.J., del Campo, J. et al. (2018) EukRef-Ciliophora: a manually curated, phylogeny-based database of small subunit rRNA gene sequences of ciliates. *Environmental Microbiology*, 20, 2218–2230.
- Brehm, V. (1928) Fresh-water fauna of New Zealand, contribution to a knowledge of. Part 1. On the microfauna of a pond in a Sphagnum bog on Mount Rolleston. *Transactions and proceedings of the New Zealand Institute*, 59, 779–812.
- Burki, F., Roger, A.J., Brown, M.W. & Simpson, A.G.B. (2019) The new tree of eukaryotes. *Trends in Ecology & Evolution*, 35, 43–55.
- Chardez, D. (1982) Thécamoébiens récoltés dans quelques lacs du Kenya. *Revue Vervétoise d’Histoire Naturelle*, 7, 57–60.
- Charman, D.J. (1997) Modelling hydrological relationships of testate amoebae (Protozoa : Rhizopoda) on New Zealand peatlands. *Journal of the Royal Society of New Zealand*, 27, 465–483.
- Chomczynski, P. & Sacchi, N. (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Annales of Biochemistry*, 162, 156–159.
- Cotterill, F.P.D., Al-Rasheid, K.A.S. & Foissner, W. (2008) Conservation of protists: is it needed at all? *Biodiversity and Conservation*, 17, 427–443.
- de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahe, F., Logares, R. et al. (2015) Eukaryotic plankton diversity in the sunlit ocean. *Science*, 348, 1261605.
- Deevey, E.S. (1955) Paleolimnology of the upper swamp deposit. *Pyramid Valley. Rec. Canterbury Mus*, 6, 291–344.
- Deflandre, G. (1936) Etude monographique sur le genre *Nebela* Leidy. *Annales de Protistologie*, 5, 201–286.
- Delgado-Baquerizo, M., Oliverio, A.M., Brewer, T.E., Benavent-González, A., Eldridge, D.J., Bardgett, R.D. et al. (2018) A global atlas of the dominant bacteria found in soil. *Science*, 359, 320–325.
- Duckert, C., Blandenier, Q., Kupferschmid, F.A.L., Kosakyan, A., Mitchell, E.A.D., Lara, E. et al. (2018) En garde! Redefinition of *Nebela militaris* (Arcellinida, Hyalospheniidae) and erection of *Alabasta* gen. nov. *European Journal of Protistology*, 66, 156–165.
- Duckert, C., Blandenier, Q., Kupferschmid, F.A.L., Kosakyan, A., Mitchell, E.A.D., Lara, E. et al. (2020) Case 3782 – *Nebela militaris* Penard, 1890 (Arcellinida, HYALOSPHEINIIDAE): proposed conservation of the specific name by giving it precedence over *Nebela bursella* Taranek, 1881. *Bulletin of Zoological Nomenclature*, 77, 22–28.
- Dumack, K., Schuster, J., Bass, D. & Bonkowski, M. (2016) A Novel Lineage of ‘Naked Filose Amoebae’: *Kraken carinae* gen. nov sp nov (Cercozoa) with a remarkable locomotion by disassembly of its cell body. *Protist*, 167, 268–278.
- Dumack, K., Siemensma, F. & Bonkowski, M. (2018) Rediscovery of the testate amoeba genus *penardeugenia* (Thaumatomonadida, Imbricatea). *Protist*, 169, 29–42.
- Ehrenberg, G.C. (1838) *Die Infusionsthierchen als vollkommene Organismen. Ein Blick in das tiefere organischen Leben der Natur*. L. Voss, Leipzig.
- Fernández, L.D., Lara, E. & Mitchell, E.A.D. (2015) Checklist, diversity and distribution of testate amoebae in Chile. *European Journal of Protistology*, 51, 409–424.
- Foissner, W. (1997) Global soil ciliate (Protozoa, Ciliophora) diversity: a probability-based approach using large sample collections from Africa, Australia and Antarctica. *Biodiversity and Conservation*, 6, 1627–1638.
- Foissner, W. (2008) Protist diversity and distribution: some basic considerations. *Biodiversity and Conservation*, 17, 235–242.

- Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294–299.
- Fontaneto, D., Boschetti, C. & Ricci, C. (2008) Cryptic diversification in ancient asexuals: evidence from the bdelloid rotifer *Philodina flaviceps*. *Journal of Evolutionary Biology*, 21, 580–587.
- Geisen, S., Mitchell, E.A.D., Adl, S., Bonkowski, M., Dunthorn, M., Ekelund, F. et al. (2018) Soil protists: a fertile frontier in soil biology research. *FEMS Microbiology Reviews*, 42, 293–323.
- Geisen, S., Rosengarten, J., Koller, R., Mulder, C., Urich, T. & Bonkowski, M. (2015) Pack hunting by a common soil amoeba on nematodes. *Environmental Microbiology*, 17, 4538–4546.
- Haeckel, E. (1899–1904) *Kunstformen der Natur*. Leipzig & Wien: Bibliographisches Institut.
- Hall, T.A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41, 95–98.
- Hebert, P.D.N., Penton, E.H., Burns, J.M., Janzen, D.H. & Hallwachs, W. (2004) Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 14812–14817.
- Heger, T.J., Edgcomb, V.P., Kim, E., Lukeš, J., Leander, B.S. & Yubuki, N. (2014) A resurgence in field research is essential to better understand the diversity, ecology, and evolution of microbial eukaryotes. *Journal of Eukaryotic Microbiology*, 61, 214–223.
- Heger, T.J., Mitchell, E.A. & Leander, B.S. (2013) Holarctic phylogeography of the testate amoeba *Hyalosphenia papilio* (Amoebozoa: Arcellinida) reveals extensive genetic diversity explained more by environment than dispersal limitation. *Molecular ecology*, 22, 5172–5184.
- Heinis, F. (1914) Die Moosfauna Columbiens: in: Voyage d'exploration scientifique en Colombie. *Mémoires de la Société Neuchateloise des Sciences Naturelles*, 5, 675–730.
- van den Hoogen, J., Geisen, S., Routh, D., Ferris, H., Traunspurger, W., Wardle, D.A. et al. (2019) Soil nematode abundance and functional group composition at a global scale. *Nature*, 572, 194–198.
- Hoogenraad, H.R. & de Groot, A.A. (1948) Thecamoebous moss-rhizopods from New Zealand. *Hydrobiologia*, 1, 28–43.
- Kosakyan, A., Gomaa, F., Mitchell, E.A.D., Heger, T.J. & Lara, E. (2013) Using DNA-barcoding for sorting out protist species complexes: a case study of the *Nebela tinctoria-collaris-bohemica* group (Amoebozoa; Arcellinida, Hyalospheniidae). *European Journal of Protistology*, 49, 222–237.
- Kosakyan, A., Heger, T.J., Leander, B.S., Todorov, M., Mitchell, E.A.D. & Lara, E. (2012) COI Barcoding of Nebelid Testate Amoebae (Amoebozoa: Arcellinida): extensive cryptic diversity and redefinition of the hyalospheniidae schultze. *Protist*, 163, 415–434.
- Kosakyan, A., Lahr, D.J.G., Mulot, M., Meisterfeld, R., Mitchell, E.A.D. & Lara, E. (2016) Phylogenetic reconstruction based on COI reshuffles the taxonomy of hyalosphenid shelled (testate) amoebae and reveals the convoluted evolution of shell plate shapes. *Cladistics*, 32, 606–623.
- Kumar, S. & Foissner, W. (2016) High cryptic soil ciliate (Ciliophora, Hypotrichida) diversity in Australia. *European Journal of Protistology*, 53, 61–95.
- Lara, E., Dumack, K., García-Martín, J.M., Kudryavtsev, A. & Kosakyan, A. (2020) Amoeboid protist systematics: a report on the “Systematics of amoeboid protists” symposium at the VIIIth ECOP/ISOP meeting in Rome, 2019. *European Journal of Protistology*, 76, 125727.
- Lax, G., Eglit, Y., Eme, L., Bertrand, E.M., Roger, A.J. & Simpson, A.G.B. (2018) Hemimastigophora is a novel supra-kingdom-level lineage of eukaryotes. *Nature*, 564, 410–414.
- Leidy, J. (1879) *Freshwater Rhizopods of North America*. Washington: United States Geological Survey of the Territories.
- Mahé, F., de Vargas, C., Bass, D., Czech, L., Stamatakis, A., Lara, E. et al. (2017) Parasites dominate hyperdiverse soil protist communities in Neotropical rainforests. *Nature Ecology & Evolution*, 1, 1–8.
- Mann, D.G. (2010) Discovering diatom species: is a long history of disagreements about species-level taxonomy now at an end? *Plant Ecology and Evolution*, 143, 251–264.
- McGlone, M.S. (2009) Postglacial history of New Zealand wetlands and implications for their conservation. *New Zealand Journal of Ecology*, 33, 1–23.
- McGlone, M.S. & Wilmshurst, J.M. (1999) A Holocene record of climate, vegetation change and peat bog development, east Otago, South Island, New Zealand. *Journal of Quaternary Science*, 14, 239–254.
- McKeown, M.M., Mitchell, E.A.D., Amesbury, M.J., Blandenier, Q., Charman, D., Duckert, C. et al. (2021) The testate amoebae of New Zealand: a checklist, identification key and assessment of biogeographic patterns. *European Journal of Protistology*, 125789. <https://doi.org/10.1016/j.ejop.2021.125789>
- McKeown, M.M., Wilmshurst, J.M., Duckert, C., Wood, J.R. & Mitchell, E.A.D. (2019) Assessing the ecological value of small testate amoebae (<45 µm) in New Zealand peatlands. *European Journal of Protistology*, 68, 1–16.
- McWethy, D.B., Whitlock, C., Wilmshurst, J.M., McGlone, M.S., Fromont, M., Li, X. et al. (2010) Rapid landscape transformation in South Island, New Zealand, following initial Polynesian settlement. *Proceedings of the National Academy of Sciences*, 107, 21343–21348.
- Meyer, M.D., Davis, C.A. & Bidwell, J.R. (2013) Assessment of two methods for sampling invertebrates in shallow vegetated wetlands. *Wetlands*, 33, 1063–1073.
- Miller, M.A., Pfeiffer, W. & Schwartz, T. (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In *Proceedings of the Gateway Computing Environments Workshop (GCE)*, 14 Nov. 2010, pp. 1–8. New Orleans, LA.
- Mitchell, E.A.D. (2015) Pack hunting by minute soil testate amoebae: nematode hell is a naturalist's paradise. *Environmental Microbiology*, 17, 4145–4147.
- Nilsson, R.H., Anslan, S., Bahram, M., Wurzbacher, C., Baldrian, P. & Tedersoo, L. (2019) Mycobiome diversity: high-throughput sequencing and identification of fungi. *Nature Reviews Microbiology*, 17, 95–109.
- van Oye, P. (1956) On the thecamoeban fauna of New Zealand with a description of four new species and biogeographical discussion. *Hydrobiologia*, 8, 16–37.
- Pawlowski, J., Majewski, W., Longet, D., Guiard, J., Cedhagen, T., Gooday, A.J. et al. (2008) Genetic differentiation between Arctic and Antarctic monothalamous foraminiferans. *Polar Biology*, 31, 1205–1216.
- Penard, E. (1902) *Faune rhizopodique du bassin du Léman*. Genève: Kündig.
- Penard, E. (1911) Part VI - Sarcodina - Rhizopodes d'eau douce. In: J. Murray (Ed.) *British Antarctic Expedition 1907-09 under the command of Sir. E. H. Shackleton, C.V.O. Reports of the Scientific Investigations. Vol. I - biology*, pp. 203–263. London: William Heinemann.

- Perry, G.L., Wilmshurst, J.M. & McGlone, M.S. (2014) Ecology and long-term history of fire in New Zealand. *New Zealand Journal of Ecology*, 38, 157–176.
- Qin, Y.M., Man, B.Y., Kosakyan, A., Lara, E., Gu, Y.S., Wang, H.M. et al. (2016) *Nebela jiuhuensis* nov sp (Amoebozoa; Arcellinida; Hyalospheniidae): A new member of the *Nebela saccifera* - *equicalceus* - *ansata* Group Described from Sphagnum Peatlands in South-Central China. *Journal of Eukaryotic Microbiology*, 63, 558–566.
- Robertson, H.A., Ausseil, A.-G., Rance, B.D., Betts, H. & Pomeroy, E. (2019) Loss of wetlands since 1990 in Southland, New Zealand. *New Zealand Journal of Ecology*, 43, 3355.
- Ronquist, F. & Huelsenbeck, J.P. (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19, 1572–1574.
- Siemensma, F., Holzmann, M., Apothéoz-Perret-Gentil, L., Clauß, S., Voelcker, E., Bettighofer, W. et al. (2020) Broad sampling of monothalamids (Rhizaria, Foraminifera) gives further insight into diversity of non-marine Foraminifera. *European Journal of Protistology*, 77, 125744.
- Singer, D., Kosakyan, A., Pillonel, A., Mitchell, E.A.D. & Lara, E. (2015) Eight species in the *Nebela collaris* complex: *Nebela gimlii* (Arcellinida, Hyalospheniidae), a new species described from a Swiss raised bog. *European Journal of Protistology*, 51, 79–85.
- Singer, D., Mitchell, E.A.D., Payne, R.J., Blandenier, Q., Duckert, C., Fernández, L.D. et al. (2019) Dispersal limitations and historical factors determine the biogeography of specialized terrestrial protists. *Molecular Ecology*, 28, 3089–3100.
- Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30, 1312–1313.
- Stout, J.D. (1984) The protozoan fauna of a seasonally inundated soil under grassland. *Soil Biology and Biochemistry*, 16, 121–125.
- Tedersoo, L., Bahram, M., Pölme, S., Kõljalg, U., Yorou, N.S., Wijesundera, R. et al. (2014) Global diversity and geography of soil fungi. *Science*, 346, 1256688.
- de Vargas, C., Norris, R., Zaninetti, L., Gibb, S.W. & Pawlowski, J. (1999) Molecular evidence of cryptic speciation in planktonic foraminifers and their relation to oceanic provinces. *Proceedings of the National Academy of Sciences of the United States of America*, 96, 2864–2868.
- Veale, A.J., de Lange, P., Buckley, T.R., Cracknell, M., Hohaia, H., Parry, K. et al. (2019) Using te reo Māori and ta re Mori in taxonomy. *New Zealand Journal of Ecology*, 43, 3388.
- Waitangi Tribunal Report Wai 11 (1986) Report of the Waitangi Tribunal on the Te Reo Maori claim. pp. 56.
- Wilkinson, D.M. (1994) A review of the biogeography of the protozoan genus *Nebela* in the southern temperate and Antarctic zones. *Area*, 26, 150–157.
- Wilkinson, D.M. (2001) What is the upper size limit for cosmopolitan distribution in free-living microorganisms? *Journal of Biogeography*, 28, 285–291.
- Wilkinson, D.M., Koumoutsaris, S., Mitchell, E.A.D. & Bey, I. (2012) Modelling the effect of size on the aerial dispersal of microorganisms. *Journal of Biogeography*, 39, 89–97.
- Wilmshurst, J.M., McGlone, M.S. & Charman, D.J. (2002) Holocene vegetation and climate change in southern New Zealand: linkages between forest composition and quantitative surface moisture reconstructions from an ombrogenous bog. *Journal of Quaternary Science*, 17, 653–666.
- Wilmshurst, J.M., Wiser, S.K. & Charman, D.J. (2003) Reconstructing Holocene water tables in New Zealand using testate amoebae: differential preservation of tests and implications for the use of transfer functions. *The Holocene*, 13, 61–72.
- Winterbourn, M. & Brown, T. (1967) Observations on the faunas of two warm streams in the Taupo thermal region. *New Zealand Journal of Marine and Freshwater Research*, 1, 38–50.
- Yang, J., Smith, H.G., Sherratt, T.N. & Wilkinson, D.M. (2010) Is there a size limit for cosmopolitan distribution in free-living microorganisms? A biogeographical analysis of testate amoebae from polar areas. *Microbial Ecology*, 59, 635–645.
- Yubuki, N., Edgcomb, V.P., Bernhard, J.M. & Leander, B.S. (2009) Ultrastructure and molecular phylogeny of *Calkinsia aureus*: cellular identity of a novel clade of deep-sea euglenozoans with epibiotic bacteria. *BMC Microbiology*, 9, 22.
- Zapata, J. & Fernandez, L. (2008) Morphology and morphometry of *Apodera vas* (Certes, 1889) (Protozoa: Testacea) from two peatlands in Southern Chile. *Acta Protozoologica*, 47, 389–395.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Duckert, C., Blandenier, Q., McKeown, M., Hohaia, H., Luketa, S., Wilmshurst, J., et al (2021) Superficially described and ignored for 92 years, rediscovered and emended: *Apodera angatakere* (Amoebozoa: Arcellinida: Hyalospheniiformes) is a new flagship testate amoeba taxon from Aotearoa (New Zealand). *J. Eukaryot. Microbiol.* 00, e12867. <https://doi.org/10.1111/jeu.12867>