



RESEARCH ARTICLE

CRISPR-Cas12a Genome Editing at the Whole-Plant Level Using Two Compatible RNA Virus Vectors

Mireia Uranga, Marta Vazquez-Vilar, Diego Orzáez, and José-Antonio Daròs*

Abstract

The use of viral vectors that can replicate and move systemically through the host plant to deliver bacterial CRISPR components enables genome editing at the whole-plant level and avoids the requirement for labor-intensive stable transformation. However, this approach usually relies on previously transformed plants that stably express a CRISPR-Cas nuclease. Here, we describe successful DNA-free genome editing of *Nicotiana benthamiana* using two compatible RNA virus vectors derived from tobacco etch virus (TEV; genus *Potyvirus*) and potato virus X (PVX; genus *Potexvirus*), which replicate in the same cells. The TEV and PVX vectors respectively express a Cas12a nuclease and the corresponding guide RNA. This novel two-virus vector system improves the toolbox for transformation-free virus-induced genome editing in plants and will advance efforts to breed more nutritious, resistant, and productive crops.

Introduction

Systems derived from bacterial CRISPR and CRISPR-associated (Cas) proteins¹ have revolutionized biotechnology. In plants, CRISPR-Cas holds great promise for unprecedented genome engineering of both model species and crops.^{2–6} Most common CRISPR-Cas arrangements include a Cas endonuclease, such as *Streptococcus pyogenes* SpCas9, and a single-guide RNA (sgRNA), which specifically directs the nuclease to a sequence of interest in the genome. As in other taxonomic groups,^{7–10} virus-derived vectors have been reported as a powerful alternative to express the CRISPR-Cas components at the whole-plant level, avoiding the labor-intensive and time-consuming tissue culture approaches required for stable transformation. These strategies are commonly termed “virus-induced genome editing” (VIGE) and have focused on the delivery of one or more sgRNAs using RNA or DNA virus vectors in transgenic plants that stably express the Cas nuclease.^{11–19}

Expression of a Cas nuclease using a plant virus-derived vector able to move systemically through a plant was long considered unachievable due to cargo constraints. However, the innovative work by Ma *et al.*²⁰ demonstrated efficient genome editing by delivering both the sgRNA and

SpCas9 at the whole-plant level using a vector derived from sonchus yellow net virus (SYNV; family *Rhabdoviridae*). Despite this unprecedented achievement, additional virus-based systems for the co-expression of Cas nucleases and sgRNAs at the whole-plant level are still required to improve the current toolbox for crop engineering. Notably, each viral vector has its own unique properties, particularly a specific host range.

To expand the virus-based tools for transformation-free genome editing in plants, we co-expressed the Cas nuclease and the guide RNA using two compatible viral vectors that replicate in the same cells and coordinately move systemically through the whole plant. We chose a potyvirus vector to express the Cas nuclease. Potyviruses (genus *Potyvirus*) are the largest group of plus-strand RNA viruses, with more than 200 currently known species that collectively can infect a large range of host plants.²¹ We also focused on a Cas12a (formerly Cpf1) nuclease, which is a component of a class 2 type V CRISPR system, isolated from *Lachnospiraceae bacterium* ND2006 (LbCas12a).²² Genome editing using LbCas12a has been demonstrated in plants,^{23–26} and the coding sequence corresponding to this nuclease is smaller than that of SpCas9.

Instituto de Biología Molecular y Celular de Plantas (Consejo Superior de Investigaciones Científicas-Universitat Politècnica de València), Valencia, Spain.

*Address correspondence to: José-Antonio Daròs, PhD, IBMCP (CSIC-Universitat Politècnica de València), Avenida de los Naranjos s/n, 46022 Valencia, Spain, Email: jadaros@ibmcp.upv.es

© Mireia Uranga, et al. 2021; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons Attribution Noncommercial License [CC-BY-NC] (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are cited.

We also co-expressed the guide RNA using a recently described potato virus X (PVX, genus *Potexvirus*, family *Virgaviridae*) vector, which efficiently induces heritable gene editing in *Nicotiana benthamiana* plants that stably express SpCas9.²⁷ Our results demonstrated efficient DNA-free *N. benthamiana* genome editing using the two compatible RNA virus vectors.

Methods

Viral vectors

Guide RNAs to target *N. benthamiana* Flowering locus *T* (*NbFT*; SolGenomics Niben101Scf01519g10008.1) and *Xylosyl transferase 1* (*NbXT1*; Niben101Scf04205g03008.1) were selected using the CRISPR-P online tool as described by Bernabé-Orts *et al.*²⁸ (Supplementary Table S1). Nucleotide (nt) sequences of recombinant viral clones are shown in Supplementary Figures S1 and S2. These clones were built using the primers shown in Supplementary Tables S2 and S3.

Plant inoculation

N. benthamiana wild type and transformed plants expressing tobacco etch virus (TEV, genus *Potyvirus*) nuclear inclusion *b* (NIB) protein²⁹ were grown at 25°C under a 12 h/12 h day/night photoperiod. Plants that were 4–6 weeks old were agro-inoculated, as previously described.^{27,30} Tissue samples (approximately 100 mg) from the first symptomatic upper non-inoculated leaf were collected at different days post inoculation (dpi), as indicated, for virus progeny and plant genome-editing analyses.

Analysis of Viral Progeny

RNA was purified from leaf samples using silica gel columns.²⁷ cDNA was synthesized using RevertAid reverse transcriptase (Thermo Fisher Scientific) and primer D179 (Supplementary Table S4). Polymerase chain reaction (PCR) with *Thermus thermophilus* DNA polymerase (Biotools) was used to amplify the TEV coat protein (CP) cistron (primers D178 and D211; Supplementary Table S4) or a fragment of the LbCas12a open reading frame (ORF; primers D3604 and D3605; Supplementary Table S4). The presence of full-length LbCas12a coding sequence in the viral progeny was assayed by reverse transcription (RT)-PCR using primers D2567 (RT) and D570-D571 (PCR; Supplementary Table S4). PCR products were separated by electrophoresis in 1% agarose gel followed by staining with ethidium bromide.

Analysis of *N. benthamiana* genome editing

DNA from leaf samples was purified using silica gel columns.²⁷ *N. benthamiana* genome fragments were amplified by PCR using high-fidelity Phusion DNA

polymerase (Thermo Fisher Scientific; Supplementary Table S5). PCR products were separated by agarose gel electrophoresis, purified from the gel, and subjected to Sanger sequencing (Supplementary Table S5). The presence of sequence modifications was analyzed using the inference of CRISPR edits (ICE) software.

Results

A dual virus-based vector system to co-express Cas nucleases and guide RNAs in plants

Initially, we built a TEV recombinant clone in which the cDNA of a human codon-optimized LbCas12a replaced that of the viral NIB protein (TEVΔNIB::LbCas12a). Our previous work showed that this vector could express large exogenous sequences, such as a whole bacterial metabolic pathway to biosynthesize lycopene or the saffron carotenoid cleavage dioxygenase.^{29,31} However, since the virus lacks the viral RNA-dependent RNA polymerase NIB, it replicates only in plants that express this protein.³¹ In our recombinant clone, the sequence coding for LbCas12a replaced most of the NIB cistron and was flanked by the native nuclear inclusion *a* protease (NIAPro) cleavage sites that mediate the release of the nuclease from the viral polyprotein (Fig. 1A and Supplementary Fig. S1).

Next, we built a recombinant version of PVX in which the LbCas12a CRISPR RNA (crRNA) was expressed under the control of the viral CP promoter, and the 29 initial codons of PVX CP were deleted to improve the stability of the recombinant clone.³² The viral CP was expressed from a heterologous promoter derived from that of the CP of bamboo mosaic virus (BaMV; genus *Potexvirus*). Based on previous work by Bernabé-Orts *et al.*²⁸ assessing the efficiency of the Cas12a-mediated gene editing of several *N. benthamiana* loci, *NbFT* was selected as the target gene (PVX::crFT; Fig. 1B and Supplementary Fig. S2). The 65 nt crRNA was cloned downstream of the PVX CP promoter and consisted of a 23 nt protospacer sequence specific to the target gene flanked on both 5' and 3' ends by a conserved 21 nt scaffold, also known as a direct repeat.


N. benthamiana plants constitutively expressing TEV NIB under the control of cauliflower mosaic virus (CaMV) 35S promoter and terminator²⁹ were co-inoculated with a 1:1 mix of two cultures of *Agrobacterium tumefaciens* C58C1 transformed with plasmids harboring TEVΔNIB::LbCas12a and PVX::crFT constructs. Controls included inoculation of TEVΔNIB::LbCas12a alone and inoculation of TEVΔNIB::crtB, which allows visual tracking of virus systemic movement due to the yellow pigmentation of infected tissue induced by *Pantoea ananatis* phytoene synthase (crtB).³¹ At 7 dpi, typical symptoms

of TEV infection emerged in the upper non-inoculated leaves of all plants. Notably, at 14 dpi, irregular chlorotic spots appeared in the leaves of plants co-inoculated with TEVΔNlb::LbCas12a and PVX::crFT but not in those inoculated with TEVΔNlb::LbCas12a alone (Fig. 2). PVX infection is characterized by the appearance of vein banding, ring spots, and leaf atrophy.³³ Therefore, the observed phenotypic alteration could have been due to LbCas12a-mediated editing of *NbFT* or an effect of the co-infection. Samples from the first systemically infected upper leaf were collected at 7 and 14 dpi, and TEVΔNlb progeny was studied by RT-PCR analysis (Fig. 1C). An 804 bp specific region of the TEV genome corresponding to the CP cistron was amplified in all virus-inoculated plants, confirming the presence of the virus (Fig. 1C, top panel). However, for viral vectors carrying big cargos, genomes larger than wild type are likely to recombine to smaller sizes, thus triggering the loss of heterologous genes.³⁴ An additional RT-PCR analysis was performed, and a 600 bp cDNA corresponding to a fragment of the LbCas12a coding sequence was exclusively amplified from plants inoculated with TEVΔNlb::LbCas12a alone or co-inoculated with TEVΔNlb::LbCas12a and PVX::crFT, regardless of sampling time (Fig. 1C, bottom

panel). A complementary RT-PCR analysis using primers hybridizing on the flanking borders of the viral vector confirmed the presence of full-length LbCas12a coding sequence in the viral progeny at 14 dpi, although it also indicated that viral progeny partially lost the cargo at this time point (Supplementary Fig. S3). Taken together, these results suggest that the expression of LbCas12a nuclease expands from the onset of viral infection until an undetermined time point in which the viral vector completely loses the heterologous gene.

Next, DNA was purified from leaf samples collected at 7 and 14 dpi, and a 550 bp fragment of the *NbFT* gene covering the LbCas12a target site was amplified by PCR. Sanger sequencing of the PCR products and ICE analysis revealed robust gene editing at 14 dpi in plants co-inoculated with TEVΔNlb::LbCas12a and PVX::crFT, reaching an insertion and deletion (indel) percentage of up to 75% (Fig. 1D and Supplementary Fig. S4). No significant differences were observed at 21 and 28 dpi. The indel distribution was consistent with the deletion-enriched mutagenesis profile characteristic of Cas12a activity, being mainly 5–10 bp deletions.²⁸ These results indicate that the simultaneous delivery of CRISPR-Cas12a components

FIG. 1. DNA-free gene editing in *Nicotiana benthamiana* based on virally delivered LbCas12a and crRNA. **(A)** Schematic representation of recombinant virus TEVΔNlb::LbCas12a. Cas12a ORF from *Lachnospiraceae bacterium* ND2006 (LbCas12a), containing a carboxy-terminal nucleoplasmic nuclear localization signal (NLS; gray box), replaces the TEV Nlb cistron (light blue box) and is flanked by proteolytic processing sites of viral NlaPro. TEV cistrons P1, HC-Pro, P3, P3N-PIPO, 6K1, CI, 6K2, VPg, NlaPro, Nlb, and CP are represented by boxes. 5' and 3' untranslated regions (UTRs) are represented by lines. **(B)** Schematic representation of recombinant clone PVX::crFT. RNA-dependent RNA polymerase (RdRp), triple gene block (TGB), and coat protein (CP) are represented by open boxes. Heterologous *Bamboo mosaic virus* CP promoter (CP_{BaMV}) is represented by a black arrow. 5' and 3' UTRs are represented by black lines. crRNA consists of a gene-specific 23 nt protospacer (pink diamond) and conserved 21 nt direct repeats (gray boxes). CP_{BaMV}, protospacer and scaffold are not shown at scale. **(C)** Reverse transcription (RT) polymerase chain reaction (PCR) analysis of TEVΔNlb progeny at 7 and 14 days post inoculation (dpi) in 35S::Nlb *N. benthamiana* plants inoculated with TEVΔNlb::LbCas12a alone or co-inoculated with TEVΔNlb::LbCas12a and PVX::crFT. Amplification products were separated by electrophoresis in an agarose gel stained with ethidium bromide. Lane 1, 1 kb ladder DNA marker with the length of some components (in bp) indicated on the left; lane 2, non-inoculated plant; lanes 3–7, plants inoculated with TEVΔNlb::crtB (lane 3) or TEVΔNlb::LbCas12a (lanes 4 and 6), and co-inoculated with TEVΔNlb::LbCas12a and PVX::crFT (lanes 5 and 7). The amplification products corresponding to cDNA regions of TEV CP (804 bp; top) or LbCas12a (600 bp; bottom) are indicated by arrows. **(D)** Inference of CRISPR edits (ICE) analysis of the first systemically infected upper leaf of *N. benthamiana* plants co-inoculated with TEVΔNlb::LbCas12a and PVX::crFT at the indicated dpi (*n* = 6). Box plot represents indels (%) and standard deviation (*n* = 6). TEVΔNlb::crtB was used as a negative control. **(E)** Schematic representation of recombinant clone PVX::crXT1:crFT. Protospacers for *NbXT1* and *NbFT* are represented by orange and pink diamonds, respectively. Other details are indicated above. **(F)** ICE analysis of the first systemically infected upper leaf of *N. benthamiana* plants co-inoculated with TEVΔNlb::LbCas12a and PVX::crXT1:crFT at the indicated dpi. Box plot represents indels (%) and standard deviation (*n* = 6).



through two compatible viral vectors (i.e., TEVΔNIB and PVX) allows highly efficient targeted mutagenesis in *N. benthamiana*.

Multiplex genome editing using the dual virus-based vector system in plants

A key advantage of CRISPR-Cas genome editing is the capacity to target several loci at once by the simultaneous expression of several guide RNAs (i.e., multiplexing). Based on our observations in Cas9-expressing plants,²⁷

we wondered whether the PVX vector could allow the delivery of multiple, functional crRNAs for Cas12a-mediated editing. To investigate this, we selected *NbXT1* as the second target gene. As Cas12a can self-process crRNAs due to its RNase III activity, *NbXT1* and *NbFT* crRNAs were arranged in tandem under the control of the same CP promoter, thus creating the PVX::crXT1:crFT construct (Fig. 1E and Supplementary Fig. S2). *N. benthamiana* plants constitutively expressing TEV NIB were co-inoculated with a 1:1 mix of two A.




FIG. 2. 35S::NIB *N. benthamiana* plants and representative leaves from these plants inoculated with TEVΔNIB::LbCas12a (left) or co-inoculated with TEVΔNIB::LbCas12a and PVX::crFT (right) at 14 dpi. Chlorotic spots in leaves from co-inoculated plants are indicated with red arrows.

tumefaciens cultures carrying TEVΔNIB::LbCas12a and PVX::crXT1:crFT. *A. tumefaciens* transformed with TEVΔNIB::crtB was used as a control in this assay. The first systemically infected leaf was sampled at 14, 21, and 28 dpi, following extraction of genomic DNA and PCR amplification of the target sites. ICE analysis revealed efficient gene editing on both *NbXT1* and *NbFT*, with average indel percentages ranging from 76% to 88%, which was maintained regardless of the sampling time (Fig. 1F). In addition, the absence of statistically relevant differences in gene editing among *NbXT1* and *NbFT* suggested that LbCas12a can efficiently self-process tandemly arrayed crRNAs. Time-course comparison with the single crRNA construct, in the case of *NbFT*, also revealed that multiplexing does not affect the editing efficiency, since the mutation rates achieved with both strategies were similar (compare Fig. 1D and F).

Dual vector CRISPR-Cas genome editing in wild-type plants

The fact that TEVΔNIB infectivity depends on the supplementation of viral NIB from a transgene may be perceived as a limitation of this genome-editing system, as the approach is still bound to a previously transformed plant. An alternative strategy for supplying NIB activity consists of the co-inoculation of TEVΔNIB with a recombinant PVX expressing NIB.³⁰ We wondered whether a single PVX vector could: (1) provide NIB activity for the replication of TEVΔNIB, and (2) deliver the crRNA for LbCas12a-mediated gene editing. Thus, the coding sequence of the TEV NIB cistron plus an additional amino-terminal Met was inserted within the expression cassette in PVX. *NbFT*-specific crRNA was added downstream of NIB without any linker sequence, so that both NIB and crFT expression were under the control of the

PVX CP promoter (PVX::NIB:crFT; Fig. 3A and Supplementary Fig. S2). Wild-type *N. benthamiana* plants were co-inoculated with a 1:1 mix of two cultures of *A. tumefaciens* carrying TEVΔNIB::LbCas12a and PVX::NIB:crFT. *A. tumefaciens* transformed with TEV::crtB was this time used as a control. At 7 dpi, the apical leaves of the co-inoculated *N. benthamiana* plants became symptomatic (i.e., leaf curling). At 14 dpi, necrotic spotting and interveinal mottling were observed in systemic leaves, which became more noticeable over time (Fig. 4). Samples from the first systemically infected upper leaf were collected at 14 dpi, and the 550 bp fragment of the *NbFT* gene covering the LbCas12a target site was amplified by PCR (Fig. 3B). ICE analysis of the PCR products exhibited 20% indels in plants co-inoculated with TEVΔNIB::LbCas12a and PVX::NIB:crFT (Fig. 3C). These results indicate that a single PVX vector can supply the viral NIB activity that allows TEVΔNIB to replicate and systemically spread in wild-type *N. benthamiana*, as well as to perform crRNA delivery for LbCas12a-mediated genome editing.

Discussion

In this study, we describe the engineering of a dual RNA virus system for the delivery of CRISPR-Cas12a components for genome editing in plants. The system consists of two compatible plus-strand RNA viruses, specifically the potyvirus TEV and the potexvirus PVX, to express the Cas nuclease and the guide RNA, respectively. Notably, the TEV vector (TEVΔNIB) contains the deletion of the RNA-dependent RNA polymerase NIB in order to accommodate the large coding sequence of the Cas nuclease. Our previous efforts to express a functional Cas nuclease using a full-length potyvirus vector in plants had been unsuccessful. However, recent works have reported




FIG. 3. Engineering of a single PVX vector for complementation of defective TEVΔNib and expression of LbCas12a crRNA. **(A)** Schematic representation of recombinant virus PVX::Nib:crFT. TEV Nib cistron is represented by a *light blue box*. Other details are as described in the legend to Figure 1. **(B)** RT-PCR analysis of TEVΔNib progeny at 14 dpi in 35S::Nib *N. benthamiana* plants co-inoculated with TEVΔNib::LbCas12a and PVX::crFT or wild-type *N. benthamiana* plants co-inoculated with TEVΔNib::LbCas12a and PVX::Nib:crFT. Amplification products were separated by electrophoresis in an agarose gel that was stained with ethidium bromide. Lane 1, DNA marker ladder with the length of some components (in bp) indicated on the *left*; lane 2, non-inoculated plant; lanes 3–5, plants inoculated with TEVΔNib::crtB (lane 3) and co-inoculated with TEVΔNib::LbCas12a and PVX::crFT (lane 4) or with TEVΔNib::LbCas12a and PVX::Nib:crFT (lane 5). The amplification products corresponding to cDNA regions of TEV CP (804 bp; *top*) or LbCas12a (600 bp; *bottom*) are indicated by *arrows*. **(C)** ICE analysis of the first systemically infected upper leaf of *N. benthamiana* plants co-inoculated with TEVΔNib::LbCas12a and PVX::Nib:crFT at 14 dpi (n=6). TEVΔNib::crtB was used as a negative control. *Box plot* represents indels (%) and standard deviation (n=6).

successful Cas9 expression and DNA-free genome editing using single vectors in *N. benthamiana*.^{20,35–37} These contrasting results emphasize the diverse properties of vectors derived from viruses belonging to different genera and families, and the necessity for a large spectrum of molecular tools, operating in a wide range of host species, to tackle challenging VIGE goals in crop

plants. Notably, SYNIV is a minus-strand RNA virus, the inoculation of which entails some complexity,³⁸ and as with each plant virus, it exhibits a particular host range.³⁹

In this work, we aimed to express a Cas nuclease using a potyvirus vector for DNA-free plant genome editing. The genus *Potyvirus* is the largest among the plant



FIG. 4. Representative leaves from wild-type *N. benthamiana* plants inoculated with TEVΔNlb::LbCas12a alone (left), PVX::Nlb:crFT alone (middle), or co-inoculated with TEVΔNlb::LbCas12a and PVX::Nlb:crFT (right) at 14 dpi.

RNA viruses and comprises more than 200 species that infect a wide range of host plants from many different botanical families.²¹ Therefore, it offers a wealth of genetic resources for VIGE. Although we were unable to express the Cas nuclease successfully using a full-length potyvirus vector, we demonstrated that TEVΔNlb allows the transient expression of LbCas12a and that PVX can perform both single or multiple crRNA delivery as well as providing the Nlb activity. Our two-virus delivery system resulted in efficient targeted editing in both Nlb-expressing and wild-type *N. benthamiana* plants, reaching indel percentages of up to 80% and 20%, respectively (Figs. 1 and 3). This dual vector system not only incorporates the enormous genetic resources of potyviruses for VIGE, but also demonstrates that compatible RNA virus vectors can be used to deliver several CRISPR-Cas components simultaneously. This could be useful for more sophisticated arrangements in plant genome editing and gene expression regulation studies. However, caution must be taken when implementing these designs because positive-strand RNA viruses, such as potyviruses and potexviruses, are prone to homologous recombination.⁴⁰ Homology must be avoided in flanking sequences of transgenes and heterologous genes, such as Nlb in this work, to avoid undesired acquisition by viral vectors (Supplementary Fig. S5). What is also of importance is that our dual RNA virus system is based on two positive-strand viral species that belong to different families and are able to co-infect the same cells. When designing systems with multiple viral vectors, superinfection exclusion must be taken into consideration because phylogenetically related viruses are generally excluded to co-infect the same cells.⁴¹

Recently, we developed a PVX vector to express multiplex SpCas9 sgRNAs efficiently at the whole-plant level.²⁷ Current results with LbCas12a confirm that PVX can be easily engineered for the simultaneous delivery of guide RNAs regardless of the nature of the Cas nuclease, which highlights its usefulness in a variety of

multiplexing approaches. Efficient genome editing in our dual-virus system implies that LbCas12a is able to find and process the crRNA either in the whole PVX genome, in viral subgenomic transcripts or in both. Particularly in the first case, LbCas12a-mediated cleavage will produce replication-incompetent viruses. The observation that LbCas12a expression has no effect on PVX vector infectivity suggests that either the whole viral genome is not the preferred target or cleavage is not that quantitative to abolish infectivity. We obtained similar results when flanked Cas9 guide RNAs with cleavable tRNA sequences in the PVX vector.²⁷

In contrast to previous reports of DNA-free VIGE that used SpCas9,^{20,36} we selected Cas12a. This was due to the following unique features of the nuclease⁴²: (1) the cleavage of target DNA is directed by a single crRNA shorter than that of SpCas9 sgRNA; (2) the protospacer adjacent motif (PAM) is T-rich (5'-TTTN-3'); (3) DNA cleavage results in cohesive ends with 4 or 5 nt overhangs, which might facilitate homology-directed repair (HDR); (4) it exhibits RNase activity useful to facilitate multiplex gene editing; and (5) it is smaller than SpCas9 (3.8 vs. 4.2 kb), which is important in terms of viral delivery. Moreover, the unique characteristics of the CRISPR-Cas12a system, unlike that of Cas9, such as the recognition of a T-rich PAM and the induction of staggered ends that facilitate homologous recombination, limit the range of target sequences and thus reduce off-target activity.⁴² Cas12a orthologues from *Francisella novicida* U112 (FnCas12a), *Acidaminococcus* sp. BV3L6 (AsCas12a), and LbCas12a were first experimentally validated in mammalian cells.^{43–45} In plants, targeted mutagenesis was achieved in rice and tobacco using any of the orthologues.^{23–26} A DNA-free approach based on the delivery of AsCas12a or LbCas12a loaded with crRNA was also validated in wild tobacco and soybean protoplasts.⁴⁶ Here, we focused on LbCas12a, since previous works reported that this nuclease possesses higher efficiency than FnCas12a or AsCas12a.^{24,28}

We showed here that LbCas12a is also effective for plant genome editing when virally delivered.

In conclusion, our dual RNA virus-based system broadens the current toolbox for DNA-free VIGE and will contribute to applications in plant functional genomics and crop improvement.

Acknowledgment

The authors thank Verónica Aragonés for excellent technical assistance.

Author Disclosure Statement

No competing financial interests exist.

Funding Information

This research was supported by grants BIO2017-83184-R and PID2019-108203RB-I00 from the Ministerio de Ciencia e Innovación (Spain) through the Agencia Estatal de Investigación (co-financed by the European Regional Development Fund) and H2020-760331 Newcotiana from the European Commission. M.U. and M.V.-V. are the recipients of fellowships FPU17/05503 from the Ministerio de Ciencia e Innovación (Spain) and APOSTD/2020/096 from the Generalitat Valenciana (Spain), respectively.

Supplementary Material

Supplementary Figure S1
Supplementary Figure S2
Supplementary Figure S3
Supplementary Figure S4
Supplementary Figure S5
Supplementary Table S1
Supplementary Table S2
Supplementary Table S3
Supplementary Table S4
Supplementary Table S5

References

- Cong L, Ran FA, Cox D, et al. Multiplex genome engineering using CRISPR-Cas systems. *Science* 2013;339:819–823. DOI: 10.1126/science.1231143.
- Nekrasov V, Staskawicz B, Weigel D, et al. Targeted mutagenesis in the model plant *Nicotiana benthamiana* using Cas9 RNA-guided endonuclease. *Nat Biotechnol* 2013;31:691–693. DOI: 10.1038/nbt.2655.
- Zhang Y, Liang Z, Zong Y, et al. Efficient and transgene-free genome editing in wheat through transient expression of CRISPR/Cas9 DNA or RNA. *Nat Commun* 2016;7:12617. DOI: 10.1038/ncomms12617.
- Li J, Zhang H, Si X, et al. Generation of thermosensitive male-sterile maize by targeted knockout of the *ZmTMS5* gene. *J Genet Genomics* 2017;44:465–468. DOI: 10.1016/j.jgg.2017.02.002.
- Zhu H, Li C, Gao C. Applications of CRISPR-Cas in agriculture and plant biotechnology. *Nat Rev Mol Cell Biol* 2020;21:661–677. DOI: 10.1038/s41580-020-00288-9.
- Huang TK, Puchta H. Novel CRISPR/Cas applications in plants: from prime editing to chromosome engineering. *Transgenic Res* 2021 Mar 1 [Epub ahead of print]; DOI: 10.1007/s11248-021-00238-x.
- Platt RJ, Chen S, Zhou Y, et al. CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell* 2014;159:440–455. DOI: 10.1016/j.cell.2014.09.014.
- Senís E, Fatouros C, Große S, et al. CRISPR/Cas9-mediated genome engineering: an adeno-associated viral (AAV) vector toolbox. *Biotechnol J* 2014;9:1402–1412. DOI: 10.1002/biot.201400046.
- Lau CH, Suh Y. *In vivo* genome editing in animals using AAV-CRISPR system: applications to translational research of human disease. *F1000Research* 2017;6:2153. DOI: 10.12688/f1000research.11243.1.
- Xu CL, Ruan MZC, Mahajan VB, et al. Viral delivery systems for CRISPR. *Viruses* 2019;11:28. DOI: 10.3390/v11010028.
- Ali Z, Abul-faraj A, Li L, et al. Efficient virus-mediated genome editing in plants using the CRISPR/Cas9 system. *Mol Plant* 2015;8:1288–1291. DOI: 10.1016/j.molp.2015.02.011.
- Yin K, Han T, Liu G, et al. A geminivirus-based guide RNA delivery system for CRISPR/Cas9 mediated plant genome editing. *Sci Rep* 2015;5:14926. DOI: 10.1038/srep14926.
- Cody WB, Scholthof HB, Mirkov TE. Multiplexed gene editing and protein overexpression using a tobacco mosaic virus viral vector. *Plant Physiol* 2017;175:23–35. DOI: 10.1104/pp.17.00411.
- Ali Z, Eid A, Ali S, et al. Pea early-browning virus-mediated genome editing via the CRISPR/Cas9 system in *Nicotiana benthamiana* and *Arabidopsis*. *Virus Res* 2018;244:333–337. DOI: 10.1016/j.virusres.2017.10.009.
- Hu J, Li S, Li Z, et al. A barley stripe mosaic virus-based guide RNA delivery system for targeted mutagenesis in wheat and maize. *Mol Plant Pathol* 2019;20:1463–1474. DOI: 10.1111/mpp.12849.
- Jiang N, Zhang C, Liu JY, et al. Development of Beet necrotic yellow vein virus-based vectors for multiple-gene expression and guide RNA delivery in plant genome editing. *Plant Biotechnol J* 2019;17:1302–1315. DOI: 10.1111/pbi.13055.
- Ellison EE, Nagalakshmi U, Gamo ME, et al. Multiplexed heritable gene editing using RNA viruses and mobile single guide RNAs. *Nat Plants* 2020;6:620–624. DOI: 10.1038/s41477-020-0670-y.
- Lei J, Dai P, Li Y, et al. Heritable gene editing using FT mobile guide RNAs and DNA viruses. *Plant Methods* 2021;17:20. DOI: 10.1186/s13007-021-00719-4.
- Mei Y, Beernink BM, Ellison EE, et al. Protein expression and gene editing in monocots using foxtail mosaic virus vectors. *Plant Direct* 2019;3:e00181. DOI: 10.1002/pld3.181.
- Ma X, Zhang X, Liu H, et al. Highly efficient DNA-free plant genome editing using virally delivered CRISPR-Cas9. *Nat Plants* 2020;6:773–779. DOI: 10.1038/s41477-020-0704-5.
- Wylie SJ, Adams M, Chalam C, et al. ICTV virus taxonomy profile: Potyviridae. *J Gen Virol* 2017;98:352–354. DOI: 10.1099/jgv.0.000740.
- Zetsche B, Gootenberg JS, Abudayyeh OO, et al. Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell* 2015;163:759–771. DOI: 10.1016/j.cell.2015.09.038.
- Endo A, Masafumi M, Kaya H, et al. Efficient targeted mutagenesis of rice and tobacco genomes using Cpf1 from *Francisella novicida*. *Sci Rep* 2016;6:38169. DOI: 10.1038/srep38169.
- Tang X, Lowder LG, Zhang T, et al. A CRISPR-Cpf1 system for efficient genome editing and transcriptional repression in plants. *Nat Plants* 2017;3:17103. DOI: 10.1038/nplants.2017.103.
- Wang M, Mao Y, Lu Y, et al. Multiplex gene editing in rice using the CRISPR-Cpf1 system. *Mol Plant* 2017;10:1011–1013. DOI: 10.1016/j.molp.2017.03.001.
- Xu R, Qin R, Li H, et al. Generation of targeted mutant rice using a CRISPR-Cpf1 system. *Plant Biotechnol J* 2017;15:713–717. DOI: 10.1111/pbi.12669.
- Uranga M, Aragonés V, Selma S, et al. Efficient Cas9 multiplex editing using unspaced sgRNA arrays engineering in a potato virus X vector. *Plant J* 2021;106:555–565. DOI: 10.1111/tppj.15164.
- Bernabé-Orts JM, Casas-Rodrigo I, Minguet EG, et al. Assessment of Cas12a-mediated gene editing efficiency in plants. *Plant Biotechnol J* 2019;17:1971–1984. DOI: 10.1111/pbi.13113.
- Marti M, Diretto G, Aragonés V, et al. Efficient production of saffron crocins and picrocrocin in *Nicotiana benthamiana* using a virus-driven system. *Metab Eng* 2020;61:238–250. DOI: 10.1016/j.jymben.2020.06.009.
- Bedoya L, Martínez F, Rubio L, et al. Simultaneous equimolar expression of multiple proteins in plants from a disarmed potyvirus vector. *J Biotechnol* 2010;150:268–275. DOI: 10.1016/j.jbiotec.2010.08.006.
- Majer E, Llorente B, Rodríguez-Concepción M, et al. Rewiring carotenoid biosynthesis in plants using a viral vector. *Sci Rep* 2017;7:41645. DOI: 10.1038/srep41645.
- Dickmeis C, Fischer R, Commandeur U. Potato virus X-based expression vectors are stabilized for long-term production of proteins and larger inserts. *Biotechnol J* 2014;9:1369–1379. DOI: 10.1002/biot.201400347.
- Loebenstein G, Gaba V. Chapter 6—Viruses of potato. *Adv Virus Res* 2012;84:209–246. DOI: 10.1016/B978-0-12-394314-9.00006-3.

34. Gilbertson RL, Sudarshana M, Jiang H, et al. Limitations on geminivirus genome size imposed by plasmodesmata and virus-encoded movement protein: insights into DNA trafficking. *Plant Cell* 2003;15:2578–2591. DOI: 10.1105/tpc.015057.
35. Gao Q, Xu WY, Yan T, et al. Rescue of a plant cytorhabdovirus as versatile expression platforms for planthopper and cereal genomic studies. *New Phytol* 2019;223:2120–2133. DOI: 10.1111/nph.15889.
36. Ariga H, Toki S, Ishibashi K. Potato virus X vector-mediated DNA-free genome editing in plants. *Plant Cell Physiol* 2020;61:1946–1953. DOI: 10.1093/pcp/pcaa123.
37. Zhang X, Kang L, Zhang Q, et al. An RNAi suppressor activates in planta virus-mediated gene editing. *Funct Integr Genomics* 2020;20:471–477. DOI: 10.1007/s10142-019-00730-y.
38. Peng X, Ma X, Lu S, et al. A versatile plant rhabdovirus-based vector for gene silencing, miRNA expression and depletion, and antibody production. *Front Plant Sci* 2021;11:627880. DOI: 10.3389/fpls.2020.627880.
39. Jackson AO, Christie SR. Purification and some physicochemical properties of sonchus yellow net virus. *Virology* 1977;77:344–355. DOI: 10.1016/0042-6822(77)90431-7.
40. Lai MM. RNA recombination in animal and plant viruses. *Microbiol Rev* 1992;56:61–79. DOI: 10.1128/mr.56.1.61-79.1992.
41. Zhang XF, Zhang S, Guo Q, et al. A new mechanistic model for viral cross protection and superinfection exclusion. *Front Plant Sci* 2018;9:40. DOI: 10.3389/fpls.2018.00040.
42. Zaidi SS-E-A, Mahfouz MM, Mansoor S. CRISPR-Cpf1: a new tool for plant genome editing. *Trends Plant Sci* 2017;22:550–553. DOI: 10.1016/j.tplants.2017.05.001.
43. Kim D, Kim J, Hur JK, et al. Genome-wide analysis reveals specificities of Cpf1 endonucleases in human cells. *Nat Biotechnol* 2016;34:863–868. DOI: 10.1038/nbt.3609.
44. Kim H, Kim ST, Ryu J, et al. CRISPR/Cpf1-mediated DNA-free plant genome editing. *Nat Commun* 2017;8:14406. DOI: 10.1038/ncomms14406.
45. Zetsche B, Heidenreich M, Mohanraju P, et al. Multiplex gene editing by CRISPR-Cpf1 using a single crRNA array. *Nat Biotechnol* 2017;35:31–34. DOI: 10.1038/nbt.3737.
46. Kim HK, Song M, Lee J, et al. *In vivo* high-throughput profiling of CRISPR-Cpf1 activity. *Nat Methods* 2017;14:153–159. DOI: 10.1038/nmeth.4104.