

**A new species of the genus *Archaeoheilus*
Legalov, 2018 (Coleoptera: Curculionidae)
from the Paleocene of Menat (France)**

Andrei A. LEGALOV, Alexander G. KIREJTSHUK & André NEL



DIRECTEURS DE LA PUBLICATION / PUBLICATION DIRECTORS :
Bruno David, Président du Muséum national d'Histoire naturelle
Étienne Ghys, Secrétaire perpétuel de l'Académie des sciences

RÉDACTEURS EN CHEF / EDITORS-IN-CHIEF: Michel Laurin (CNRS), Philippe Taquet (Académie des sciences)

ASSISTANTE DE RÉDACTION / ASSISTANT EDITOR: Adenise Lopes (Académie des sciences; cr-palevol@academie-sciences.fr)

MISE EN PAGE / PAGE LAYOUT: Audrina Neveu (Muséum national d'Histoire naturelle; audrina.neveu@mnhn.fr)

RÉVISIONS LINGUISTIQUES DES TEXTES ANGLAIS / ENGLISH LANGUAGE REVISIONS: Kevin Padian (University of California at Berkeley)

RÉDACTEURS ASSOCIÉS / ASSOCIATE EDITORS (*, *took charge of the editorial process of the article/a pris en charge le suivi éditorial de l'article*):

Micropaléontologie/*Micropalaeontology*

Maria Rose Petrizzo (Università di Milano, Milano)

Paléobotanique/*Palaeobotany*

Cyrille Prestianni (Royal Belgian Institute of Natural Sciences, Brussels)

Métazoaires/*Metazoa*

Annalisa Ferretti* (Università di Modena e Reggio Emilia, Modena)

Paléochthyologie/*Palaeoichthyology*

Philippe Janvier (Muséum national d'Histoire naturelle, Académie des sciences, Paris)

Amniotes du Mésozoïque/*Mesozoic amniotes*

Hans-Dieter Sues (Smithsonian National Museum of Natural History, Washington)

Tortues/*Turtles*

Juliana Sterli (CONICET, Museo Paleontológico Egidio Feruglio, Trelew)

Lépidosauromorphes/*Lepidosauromorphs*

Hussam Zaher (Universidade de São Paulo)

Oiseaux/*Birds*

Eric Buffetaut (CNRS, École Normale Supérieure, Paris)

Paléomammalogie (mammifères de moyenne et grande taille)/*Palaeomammalogy (large and mid-sized mammals)*

Lorenzo Rook (Università degli Studi di Firenze, Firenze)

Paléomammalogie (petits mammifères sauf Euarchontoglires)/*Palaeomammalogy (small mammals except for Euarchontoglires)*

Robert Asher (Cambridge University, Cambridge)

Paléomammalogie (Euarchontoglires)/*Palaeomammalogy (Euarchontoglires)*

K. Christopher Beard (University of Kansas, Lawrence)

Paléoanthropologie/*Palaeoanthropology*

Roberto Macchiarelli (Université de Poitiers, Poitiers)

Archéologie préhistorique/*Prehistoric archaeology*

Marcel Otte (Université de Liège, Liège)

RÉFÉRÉS / REVIEWERS: <https://sciencepress.mnhn.fr/periodiques/comptes-rendus-palevol/referes-du-journal>

COUVERTURE / COVER:

Made from the Figures of the article.

Comptes Rendus Palevol est indexé dans / *Comptes Rendus Palevol is indexed by:*

- Cambridge Scientific Abstracts
- Current Contents® Physical
- Chemical, and Earth Sciences®
- ISI Alerting Services®
- Geoabstracts, Geobase, Georef, Inspec, Pascal
- Science Citation Index®, Science Citation Index Expanded®
- Scopus®.

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Comptes Rendus Palevol* sont référencés par / *Articles and nomenclatural novelties published in Comptes Rendus Palevol are registered on:*

- ZooBank® (<http://zoobank.org>)

Comptes Rendus Palevol est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris et l'Académie des sciences, Paris
Comptes Rendus Palevol is a fast track journal published by the Museum Science Press, Paris and the Académie des sciences, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish:*

Adansonia, Geodiversitas, Zoosystema, Anthropolozologica, European Journal of Taxonomy, Naturae, Cryptogamie sous-sections *Algologie, Bryologie, Mycologie.*

L'Académie des sciences publie aussi / *The Académie des sciences also publishes:*

Comptes Rendus Mathématique, Comptes Rendus Physique, Comptes Rendus Mécanique, Comptes Rendus Chimie, Comptes Rendus Géoscience, Comptes Rendus Biologies.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél. : 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <https://sciencepress.mnhn.fr>

Académie des sciences, Institut de France, 23 quai de Conti, 75006 Paris.

© This article is licensed under the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>)
ISSN (imprimé / print): 1631-0683/ ISSN (électronique / electronic): 1777-571X

A new species of the genus *Archaeoheilus* Legalov, 2018 (Coleoptera: Curculionidae) from the Paleocene of Menat (France)

Andrei A. LEGALOV

Institute of Systematics and Ecology of Animals SB RAS,
Frunze street, 11, 630091 Novosibirsk (Russia)
and Biological Institute, Tomsk State University, Lenina Prospekt 36, 634050 Tomsk (Russia)
and Institut de Systématique, Évolution, Biodiversité (ISYEB),
Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles
case postale 50, 57 rue Cuvier, F-75231 Paris cedex 05 (France)
fossilweevils@gmail.com (corresponding author)

Alexander G. KIREJTSHUK

Zoological Institute RAS, Universitetskaya nab., 1, 199034 St. Petersburg (Russia)
agk@zin.ru
kirejtshuk@gmail.com

André NEL

Institut de Systématique, Évolution, Biodiversité (ISYEB),
Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles
case postale 50, 57 rue Cuvier, F-75231 Paris cedex 05 (France)
anel@mnhn.fr

Submitted on 30 July 2020 | Accepted on 15 December 2020 | Published on 4 April 2022

urn:lsid:zoobank.org:pub:AA8B4F8F-A2B0-4D1D-8561-4982443352E3

Legalov A. A., Kirejtshuk A. G. & Nel A. 2022. — A new species of the genus *Archaeoheilus* Legalov, 2018 (Coleoptera: Curculionidae) from the Paleocene of Menat (France). *Comptes Rendus Palevol* 21 (12): 245-251. <https://doi.org/10.5852/cr-palevol2022v21a12>

ABSTRACT

A new molytine species, *Archaeoheilus gallicus* n. sp. from the Paleocene of Menat (France) is described and illustrated. The new species is similar to *A. provectus* (Scudder, 1876), *A. scudderi* Legalov, 2018, *A. ovalis* Legalov, 2018 from the early-middle Eocene Green River Formation, and *A. lacoei* (Scudder, 1893) from the late Eocene Florissant Formation. But it differs from them in the narrow elytral interstriae. The new species differs also from *A. scudderi* in the thicker rostrum, larger body size, and wider tarsomeres 2 and 3; and also from *A. provectus*, *A. ovalis*, and *A. lacoei* in the thinner rostrum; from *A. provectus* and *A. lacoei* in the smaller body size, and from *A. ovalis* in the larger body size and antennal scrobes reaching eyes. The key to species of the genus *Archaeoheilus* Legalov, 2018 is given. The links between the weevil faunas of America and Europe in the Paleogene are discussed.

KEY WORDS

Molytinae,
diagnostics,
key to species,
France,
new species.

RÉSUMÉ

Une nouvelle espèce du genre Archaeoheilus Legalov, 2018 (Coleoptera: Curculionidae) dans le Paléocène de Menat (France).

La nouvelle espèce de Molytinae *Archaeoheilus gallicus* n. sp. est décrite et illustrée du Paléocène de Menat (France). Elle est similaire à *A. provectus* (Scudder, 1876), *A. scudderi* Legalov, 2018, *A. ovalis*

MOTS CLÉS
Molytinae,
diagnostiques,
clef des espèces,
France,
nouvelle espèce.

Legalov, 2018 de la formation du Green River (Éocène inférieur à moyen), et *A. lacoei* (Scudder, 1893) de la formation de Florissant (Éocène terminal). Mais elle en diffère par les interstries élytrales étroites. La nouvelle espèce diffère également d'*A. scudderi* par le rostre plus épais, la taille du corps plus grande et les tarsomères 2 et 3 plus larges; et aussi de *A. provectus*, *A. ovalis* et *A. lacoei* par son rostre plus fin; de *A. provectus* et *A. lacoei* par son plus petit corps, de *A. ovalis* par sa plus grande taille et les scrobes antennaires atteignant les yeux. Une clef des espèces du genre *Archaeoheilus* Legalov, 2018 est proposée. Les liens entre les faunes de charançons d'Amérique et d'Europe au Paléogène sont discutés.

INTRODUCTION

The weevil fauna of the Paleocene is the least studied among those of the Paleogene. This is primarily because there are so few Paleocene localities with fossil remnants accessible for study. Up to now Paleocene weevils have been found in five localities (Heer 1870; Cockerell 1925, 1926, 1936; Piton 1940; Williams 1943; Legalov 2010; Legalov *et al.* 2017), in France (Menat), Starostin (Norway, Svalbard), Arkhara (south of the Russian Far East), and Hong Kong (Ping Chau Formation, dated from the Paleocene by Li & Lee [1997], and from the early Cretaceous by Wang *et al.* [2019]), and Sunchal (Argentina). In total, 27 weevil species have been described from the Paleocene, but among them 18 species known from Starostin, Sunchal, and Mirs isolated elytra (Heer 1870; Cockerell 1925, 1926, 1936; Cockerell *in* Williams 1943; Legalov 2020b). Only eight species represented by complete impressions have been described from Menat, Selandian-Thanetian of France (Piton 1940; Legalov *et al.* 2017), where representatives of the families Ithyceridae, Attelabidae, Brentidae, and Curculionidae are recorded.

The new species described here is the first representative of the Molytinae in the fauna of Menat. At the same time, it is also the first species of the genus *Archaeoheilus* Legalov, 2018 known from fossil faunas of the Eastern Hemisphere.

MATERIAL AND METHODS

The holotype of the new species is deposited in the Menat Town Museum, Village of Menat, Puy-de-Dôme (further MTM). The specimen was studied using a stereomicroscope Olympus SCX9 in MNHN.

SITE DESCRIPTION

The middle Paleocene Menat fossil site (Menat Basin, Puy-de-Dôme, France) is a volcanic maar containing a paleolake *c.* 1 km in diameter (46°06'N, 2°54'E), which contains sedimentary rocks (spongo-diatomites) with remains of diverse aquatic and terrestrial flora and fauna (Piton 1940; Vincent *et al.* 1977; Nel 1989, 2008; etc.). The composition of faunal and floral remains suggests that the lake was surrounded by a mainly deciduous forest and that the palaeoenvironment

was warm and humid (Wedmann *et al.* 2018). Following the pollen, mammalian stratigraphic and radiometric K/Ar analyses, the age of Menat outcrop was estimated as 59 Ma (Kedves & Russel 1982; Nel 2008). However, the new estimate based on macroflora postulated its age within 60–61 Ma (Wappler *et al.* 2009). Some preliminary results of studies on the beetle fauna from Menat outcrop were published by Nabozhenko & Kirejtshuk (2014, 2017), Kirejtshuk *et al.* (2016), Legalov *et al.* (2017, 2019a) and Kirejtshuk & Nel (2018).

SYSTEMATICS

Family CURCULIONIDAE Latreille, 1802
Subfamily MOLYTINAE Schoenherr, 1823
Tribe MOLYTINI Schoenherr, 1823
Subtribe HYLOBIINA W. Kirby, 1837
Genus *Archaeoheilus* Legalov, 2018

Archaeoheilus gallicus n. sp.
(Figs 1; 2)

[urn:lsid:zoobank.org:act:E1B359E5-AC9E-4FBD-8B05-1300604A7AB9](https://zoobank.org/act:E1B359E5-AC9E-4FBD-8B05-1300604A7AB9)

ETYMOLOGY. — The name derives from the Latin name of France (Gallia).

TYPE LOCALITY AND STRATUM. — Menat Basin, Puy-de-Dôme, France. Middle Paleocene (46°06'N, 2°54'E), collected in a new, small outcrop near the south-east of the village of Menat.

HOLOTYPE. — MNT-Nel 2002 (well preserved on the piece of rock; the specimen shows the main parts of body sclerites in dorso-lateral view but antennae are missing), stored in the collection of the town of Menat.

DIAGNOSIS. — Body glabrous; rostrum curved, longer than pronotum; eyes transverse oval; antennal scrobes lateral, directed to eye; antennae inserted subapically; elytra convex, with narrow interstriae; prosternum without rostral channel; pygidium covered by elytra; metacoxae transverse elongate; tibiae almost uncinatate.

DESCRIPTION OF HOLOTYPE

Body black and glabrous. Rostrum rather thick, slightly curved, slightly longer than pronotum, finely punctured, 4.2 times as long as wide at apex, 3.2 times as long as wide in middle and at base, 1.2 times as long as pronotum. Forehead

slightly narrower than rostrum base wide, densely punctured. Eyes large, transverse oval, 0.5 times as long as wide. Temples subequal to eye length. Antennal scrobes long, lateral, oblique, directed to basal third of eye, reaching one. Antennae inserted in apical part of rostrum. Pronotum about 0.8 times as long as wide at apex, about 0.6 times as long as wide in middle and at base. Pronotal disc flattened, densely punctured. Elytra weakly convex, 3.8 times as long as pronotum. Elytral base weakly concave. Humeral calli convex. Striae quite wide, with large oval points. Interstriae narrow, weakly convex, punctured, 1.2–1.7 times as long as wide of striae. Prosternum with postorbital lobes, without rostral channel. Precoxal part of prosternum long. Postcoxal part of prosternum narrow. Mesepimera medium-sized. Metaventrite 2.8 times as long as metacoxal length, flattened, densely punctured. Metepisternum about 3.5 times as long as wide in middle. Abdomen convex. Ventrite 1 1.6 times as long as metacoxa. Ventrite 2 as long as ventrite 1. Ventrite 3 0.5 times as long as ventrite 2. Ventrite 4 0.8 times as long as ventrite 3. Ventrite 5 2.7 times as long as ventrite 4. Pygidium not exposed. Legs long. Procoxae conical. Metacoxae transverse elongate. Femora thickened, toothless. Profemora about 3.7 times as long as maximal wide. Mesofemora about 3.0 times as long as maximum wide. Metafemora about 3.3 times as long as maximum wide. Tibiae almost straight, with large uncus at apex. Protibiae about 4.8 times as long as wide. Mesotibiae about 6.9 times as long as wide. Metatibiae about 6.3 times as long as wide, reaching ventrite 3. Tarsi long. Tarsomere 1 conical. Tarsomere 2 wide conical. Tarsomere 3 bilobed. Tarsomere 5 long. Metatarsomere 1 about 1.5 times as long as wide at apex. Metatarsomere 2 0.8 times as long as wide at apex, 0.7 times as long as and 1.3 times as wide as tarsomere 1. Metatarsomere 3 0.7 times as long as wide at apex, subequal in length with and 1.2 times as wide as metatarsomere 2. Metatarsomere 5 6.0 times as long as wide at apex, 2.3 times as long as and 0.3 times as wide as metatarsomere 3.

Measurements

Body length (without rostrum) 4.0 mm. Rostrum length 1.0 mm.

COMPARISON WITH OTHER *ARCHAEOHEILUS* SPECIES

Archaeoheilus gallicus n. sp. is similar to *A. provectus* (Scudder, 1876), *A. scudderi* Legalov, 2018, *A. ovalis* Legalov, 2018 from Green River, and *A. lacoei* (Scudder, 1893) from Florissant with transverse eyes, which width is about two times longer than length. However, it differs generally from all compared species in the narrow elytral interstriae. But the new species also differs from individual species in the following features: from *A. scudderi* – in the thicker rostrum, larger body sizes, and wider tarsomere 2 and 3 –; from *A. provectus*, *A. ovalis* and *A. lacoei* – in the thinner rostrum –; from *A. provectus* and *A. lacoei* – in the smaller body sizes –, from *A. ovalis* – in the larger body sizes and antennal scrobes reaching eyes. *Archaeoheilus gallicus* n. sp. is also distinguished from *A. packardii* (Scudder, 1893) from Green River in the transverse eyes, thinner rostrum, narrow

elytral interstriae and smaller body size and from *A. deleticius* (Scudder, 1893) from White River in the thinner rostrum, less transverse eyes, and narrow elytral interstriae.

ADDITIONAL NOTES

Archaeoheilus gallicus n. sp. belongs to the subfamily Molytinae, because its tibiae have a large uncus, its rostrum is rather thick, and also its antennae are inserted at the apical third of the rostrum. The prosternum without any rostral channel, the convex body, the conical tarsomere 2, the elytral base weakly concave, the mesepimera not enlarged, the pronotum less than half of elytral length, the pygidium not exposed, the antennal scrobes oblique, the metacoxae transverse elongate, and the large transverse eyes a placement in the tribe Molytini. Distinct humeri allow it to be placed in the subtribe Hylobiina. Antennae are missing in the type specimen and the structure of its club not observable, which is one of the characters for separating Plintini and Molytini, but the Plintini are characterized by the rounded or not large transverse eyes, which makes a position in this tribe unlikely. *Archaeoheilus gallicus* n. sp. should be put in the genus *Archaeoheilus* because of its toothless femora, quite short and thick rostrum, and small body size. The small body size is not characteristic of the genera of Molytini. The genus *Archaeoheilus* with a body length within 3.4 and 5.8 mm is the smallest in this tribe.

DISCUSSION AND CONCLUSIONS

Piton (1940) described six Curculionidae from Menat, viz. *Phytonomus punctatus* Piton, 1940, *Rhysosternum punctatolineatum* Piton, 1940, *Curculio elegans* Piton, 1940, *Centrinus longipes* Piton, 1940, *Hipporhinus ventricosus* Piton, 1940, and *Lixus ligniticus* Piton, 1940. The study of the holotypes of *Curculio elegans* and *Hipporhinus ventricosus* showed that they belong to the tribes Curculionini (Curculioninae) and Naupactini (Entiminae) respectively (Legalov *et al.* 2017). The types of the other species have not yet been found, but the evaluation of the original descriptions, drawings, and materials from Menat made it possible to conclude that “*Phytonomus punctatus*” belongs to the family Attelabidae, “*Rhysosternum punctatolineatum*” to the family Rhynchitidae, “*Centrinus longipes*” to the subfamily Curculioninae of the family Curculionidae, and that the systematic position of *Lixus ligniticus* was correctly identified in the original description. Thousands of additional specimens of Curculionoidea were recently found in the Menat outcrop, which wait to be studied.

The subfamily Molytinae is a diverse group of weevils (Lyal 2014; Legalov 2018c). The earliest records of the representatives of this subfamily are known from the Late Cretaceous (Maastrichtian). *Hylobiites cretaceus* Scudder, 1895 was found in Assiniboine, Canada and *Dorotheus guidensis* Kuschel, 1959 is documented in Magallanes, Chile (Scudder 1895; Kuschel 1959; Legalov 2015). Unfortunately, both of them are known only from isolated elytra and their systematic positions within the subfamily is poorly constrained. The Molytinae *Archaralites zherichini* Legalov, 2010, based on




Fig. 1.— *Archaeoheilus gallicus* n. sp. from Paleocene of Menat, holotype, body, lateral view. Scale bar: 1.0 mm.

an incomplete impression from the Priamurye, is the earliest Paleocene Molytinae (Legalov 2010). *Archaeoheilus gallicus* n. sp. is the second representative of Paleocene Molytinae. It is known from complete impression on which most of the diagnostic characters are visible. The Eocene Molytinae are more diverse (Legalov 2015). Representatives of the tribes Molytini (*Furhylobius* Legalov, 2015), Camptorhinini Lacordaire, 1865 (*Camptorrhinites* Britton, 1960, *Korystina* Britton, 1960), Cryptorhynchini Schoenherr, 1826 (*Oisecalles latosquamosus* Legalov, Kirejtshuk & Nel, 2019 and *Taylorius* Britton, 1960), and Sciabregmini Legalov, Kirejtshuk & Nel, 2019 (*Sciabregma* Scudder, 1893) were found in the early Eocene (Scudder 1893; Britton 1960; Legalov 2015; Legalov *et al.* 2019b). Two genera, *Lutago* Britton, 1960 and *Pissodites* Britton, 1960, of unknown tribal affinities, have been described from the early Eocene deposits of Great Britain (Britton 1960). The early-middle Eocene Molytinae from the Green River Formation are represented by species of the genera *Archaeoheilus*, *Rhysosternum* Scudder, 1893, *Sciabregma* Scudder, 1893, and also formally placed in the genera “*Magdalis*” and “*Cryptorhynchus*” (Scudder 1893; Cockerell 1921; Legalov 2015, 2018b). Thirteen species of Molytinae have been found in the middle and late Eocene deposits and seven species were recovered in Baltic amber (Scudder 1893; Wickham 1912; Hustache 1942; Haupt 1950; Zherikhin 1971; Legalov 2016, 2020a, 2020b; Bukejs *et al.* 2020).

The new species belongs to a genus previously known from the Green River Formation and the latest Eocene of Florissant (Scudder 1876, 1893; Legalov 2018b). It was not known from the Eocene deposits and amber of Europe. The genus *Archaeoheilus* is similar to the genus *Lutago*. The differences between them are mainly in the thin rostrum and small size (2.2–2.6 mm) of the latter genus.

The discovery of a new *Archaeoheilus* species allows us to discuss probable faunal links between Eastern and Western hemispheres. Such links are already known for many other insect orders, especially in the Odonata (see Garrouste & Nel 2019). Europe and America were joined in the Mesozoic (Boucot *et al.* 2013). However, the main divergences among the Curculionidae could have occurred only in the Late Cretaceous, since the first known representatives of this family are from the end of the early Cretaceous (Legalov 2014a). Only representatives of the most ‘primitive’ subfamily Eirrhiniinae were found in the Cenomanian and Turonian deposits (Legalov 2014b, 2018a; Legalov & Poinar 2015). Land bridges between North America through Greenland and Scandinavia, on the one hand, and the British Isles, on the other, existed in the Paleocene (Wen *et al.* 2016). The bridge through Greenland remained in the Eocene, but Scandinavia at that time was separated from Western Europe (Boucot *et al.* 2013; Wen *et al.* 2016). Menat and Green River were located the Boreo-tropical belt, but the




FIG. 2.— *Archaeoheilus gallicus* n. sp. from Paleocene of Menat, holotype, reconstruction drawing of body, counterpart, lateral view. Scale bar: 1.0 mm.

connection between America and Europe was by the warm temperature belt (Baltic amber was in this belt) in the Eocene (Boucot *et al.* 2013). Accordingly, Curculionidae could freely move between these continental plates. Species from four genera are shared by the past faunas of Europe and America: *Archaeoheilus* is found in the Paleocene of France (present study) and the Eocene of the United States (Legalov 2018b), *Sciabregma* in the early Eocene of France (Legalov *et al.* 2019b) and the early-middle Eocene of the United States

(Scudder 1893; Cockerell 1921), *Limalophus* in the middle Eocene amber of Europe (Legalov 2020a) and the early-late Eocene of the United States (Scudder 1893), *Toxorhynchus* in the middle Eocene of Europe (Bukejs & Legalov 2020) and the late Eocene of the United States (Scudder 1893; Legalov 2013), as well as in Miocene Dominican amber (Poinar & Legalov 2015) and the Recent fauna of the New World (Bukejs & Legalov 2020). The Neotropical subtribe Prionomerina and Oriental Zherichinixenina are found

KEY TO SPECIES OF THE GENUS *ARCHAEOHEILUS* LEGALOV, 2018

1. Eyes rounded. Body length 6.4 mm *A. packardii* (Scudder, 1893)
- Eyes transverse oval 2
2. Elytral interstriae narrow, 1.2-1.7 times as long as wide as striae. Body length 4.0 mm *A. gallicus* n. sp.
- Elytral interstriae wide, 3.0-5.8 times as long as wide as striae 3
3. Eye width/length ratio 1.7. Body length 3.8 mm *A. deleticius* (Scudder, 1893)
- Eye width/length ratio 1.9-2.1 4
4. Rostrum slender, 3.7 times as long as wide in the middle. Body length 5.8 mm *A. scudderi* Legalov, 2018
- Rostrum more robust, 2.4-2.9 times as long as wide in the middle 5
5. Smaller. Body length 3.4 mm *A. ovalis* Legalov, 2018
- Larger. Body length 5.0-7.3 mm 6
6. Smaller. Rostrum 2.9 times as long as wide in the middle. Body length 5.0 mm *A. provectus* (Scudder, 1876)
- Larger. Rostrum 2.4 times as long as wide in the middle. Body length 7.3 mm *A. lacoei* (Scudder, 1893)

in the Eocene of the United States (Scudder 1893) and Baltic amber (Legalov 2020a). The mostly American tribe Naupactini was found in the Paleocene of France (Legalov *et al.* 2017) and the middle Eocene amber (Yunakov & Kirejtshuk 2011; Bukejs & Legalov 2019). The presence of the Neotropical tribe Anypotactini can be noted in the middle Eocene of Europe (Yunakov & Kirejtshuk 2011; Legalov 2015; Legalov *et al.* 2019c). Its representatives have not yet been found either in the Eocene of the United States or in the Paleocene of Europe (Legalov 2015), but will probably be found in the future.

Acknowledgements

The authors had the pleasure to receive generous assistance from Clotilde Berger-Pompili and Mathilde Leygnac, directors of the EHPAD du Pays de Menat, for their generous help and authorisations to collect fossil insects in a small but rich outcrop near the village of Menat (Puy-de-Dôme, France). The authors thank also A. P. Rasnitsyn and A. G. Ponomarenko (Moscow) for providing comparative material from the collection of the Borissiak Paleontological Institute of the Russian Academy of Sciences. The study of the first author was partially supported by the Federal Fundamental Scientific Research Programme for 2021-2025 (grant number 0247-2021-0004), Muséum national d'Histoire naturelle (Paris), and the Russian Foundation for Basic Research (grant numbers 18-04-00243-a and 19-04-00465-a). The study of the second author was partly carried out under the auspices of the Russian state research project no. 1021051302540-6, programs of the Presidium of the Russian Academy of Sciences "Evolution of organic world and planet processes" and the Russian Foundation of Basic Research (grant numbers 18-04-00243-a and 19-04-00465-a). The authors thanks the reviewers and the associate editor, Annalisa Ferretti.

REFERENCES

BOUCOT A. J., CHEN X. & SCOTese C. R. 2013. — Phanerozoic paleoclimate: an atlas of lithologic indicators of climate. *SEPM Concepts in Sedimentology and Paleontology*, 478 p. <https://doi.org/10.2110/sepmcsp.11>

BRITTON E. B. 1960. — Beetles from the London clay (Eocene) of Bognor Regis, Sussex. *Bulletin of the British Museum (Natural History) Geology* 4 (2): 27-50. <https://doi.org/10.5962/p.313857>

BUKEJS A. & LEGALOV A. A. 2019. — First record of the tribe Naupactini (Coleoptera: Curculionidae) in Rovno amber. *Fossil Record* 22 (1): 25-30. <https://doi.org/10.5194/fr-22-25-2019>

BUKEJS A. & LEGALOV A. A. 2020. — The first record of Brentidae (Coleoptera) in Eocene Rovno amber with description of a new fossil species of *Toxorhynchus* Scudder, 1893. *Fossil Record* 23 (2): 169-177. <https://doi.org/10.5194/fr-23-169-2020>

BUKEJS A., ALEKSEEV V. I. & LEGALOV A. A. 2020. — A new Eocene genus of the subtribe Tylodina (Coleoptera: Curculionidae) and notes concerning local differences of Baltic amber in the Kaliningrad Region. *Fossil Record* 23 (1): 75-81. <https://doi.org/10.5194/fr-23-75-2020>

COCKERELL T. D. A. 1921. — Some Eocene insects from Colorado and Wyoming. *Proceedings of the United States National Museum* 49 (2358): 29-39. <https://doi.org/10.5479/si.00963801.59-2358.29>

COCKERELL T. D. A. 1925. — Tertiary insects from Argentina. *Proceedings of the United States National Museum* 68 (2602): 1-5. <https://doi.org/10.5479/si.00963801.68-2602.1>

COCKERELL T. D. A. 1926. — Some tertiary fossil insects. *Annals and Magazine of Natural History* 9 (18): 313-324. <https://doi.org/10.1080/00222932608633519>

COCKERELL T. D. A. 1936. — The fauna of the Sunchal (or Margas Verdes) Formation, Northern Argentina. *American Museum Novitates* 886: 1-9. <http://hdl.handle.net/2246/4185>

GARROUSTE R. & NEL A. 2019. — Alaskan Palaeogene insects: a challenge for a better knowledge of the Beringian 'route' (Odonata: Aeshnidae, Dysagrionidae). *Journal of Systematic Palaeontology* 17 (22): 1939-1946. <https://doi.org/10.1080/14772019.2019.1572235>

HAUPT H. 1950. — Die Käfer (Coleoptera) aus der Eozänen Braunkohle des Geiseltales. *Geologica* 6: 1-168.

HEER O. 1870. — Die miocene Flora and Fauna Spitsbergens. *Kungliga Svenska vetenskapsakademiens handlingar* 8: 1-98, I-XVI.

HUSTACHE A. 1942. — Un curculionide de l'ambre de la Baltique. *Bulletin mensuel de la Société linnéenne de Lyon* 11: 108-109.

KEDVES M. & RUSSEL D. E. 1982. — Palynology of the Thanetian layers of Menat. The Geology of the Menat basin, France. *Palaeontographica B* 182: 87-150.

KIREJTSHUK A. G. & NEL A. 2018. — Nitidulidae (Coleoptera) from the Paleocene of Menat (France). *Zootaxa* 4402 (1): 1-41. <https://doi.org/10.11646/zootaxa.4402.1.1>

KIREJTSHUK A. G., NEL A. & KIREJTSHUK P. A. 2016. — Taxonomy of the reticulate beetles of the subfamily Cupedinae (Coleoptera, Archostemata), with a review of the historical development. *Invertebrate Zoology* 13 (2): 61-190. <https://doi.org/10.15298/invertzool.13.2.01>

KUSCHEL G. 1959. — Un Curculiónido del Cretácao Superior, primer insecto fósil de Chile. *Invest. Zool. Chilenas* 5: 49-54.

LEGALOV A. A. 2010. — Checklist of Mesozoic Curculionoidea (Coleoptera) with description of new taxa. *Baltic Journal of Coleopterology* 10 (1): 71-101.

LEGALOV A. A. 2013. — New and little known weevils (Coleoptera: Curculionoidea) from the Paleogene and Neogene. *Historical Biology* 25 (1): 59-80. <https://doi.org/10.1080/08912963.2012.692681>

LEGALOV A. A. 2014a. — The oldest Brentidae and Curculionidae (Coleoptera: Curculionoidea) from the Aptian of Bon-Tsagaan. *Historical Biology* 26 (1): 6-15. <https://doi.org/10.1080/08912963.2012.751103>

LEGALOV A. A. 2014b. — New Nemonychidae, Brentidae and Curculionidae (Coleoptera: Curculionoidea) from the Turonian of Kzyl-Dzhar (Kazakhstan). *Historical Biology* 26 (5): 675-689. <https://doi.org/10.1080/08912963.2013.829826>

LEGALOV A. A. 2015. — Fossil Mesozoic and Cenozoic weevils (Coleoptera, Obrienioidea, Curculionoidea). *Paleontological Journal* 49 (13): 1442-1513. <https://doi.org/10.1134/S0031030115130067>

LEGALOV A. A. 2016. — New weevils (Curculionidae) in Baltic amber. *Paleontological Journal* 50 (9): 970-985. <https://doi.org/10.1134/S0031030116090057>

LEGALOV A. A. 2018a. — A new weevil, *Burmorhinus georgei* gen. et sp. nov. (Coleoptera; Curculionidae) from the Cretaceous Burmese amber. *Cretaceous Research* 84: 13-17. <https://doi.org/10.1016/j.cretres.2017.11.002>

LEGALOV A. A. 2018b. — New weevils (Coleoptera, Curculionoidea) from the Eocene of the Green River, United States: Part 1. *Paleontological Journal* 52 (3): 294-302. <https://doi.org/10.1134/S0031030118030061>

LEGALOV A. A. 2018c. — Annotated key to weevils of the world. Part 2. Subfamily Molytinae (Coleoptera, Curculionidae). *Ukrainian Journal of Ecology* 8 (4): 340-350.

LEGALOV A. A. 2020a. — A review of the Curculionoidea (Coleoptera) from European Eocene ambers. *Geosciences* 10 (1): 1-74. <https://doi.org/10.3390/geosciences10010016>

- LEGALOV A. A. 2020b. — Fossil history of Curculionoidea (Coleoptera) from the Paleogene. *Geosciences* 10 (9): 1-50. <https://doi.org/10.3390/geosciences10090358>
- LEGALOV A. A. & POINAR JR. G. 2015. — New tribes of the superfamily Curculionoidea (Coleoptera) in Burmese amber. *Historical Biology* 27 (5): 558-564. <https://doi.org/10.1080/08912963.2014.896908>
- LEGALOV A. A., NEL A. & KIREJTSHUK A. G. 2017. — New and little known weevils (Coleoptera: Curculionoidea) from the Paleocene of Menat (France). *Comptes Rendus Palevol* 16 (3): 248-256. <https://doi.org/10.1016/j.crpv.2016.10.007>
- LEGALOV A. A., KIREJTSHUK A. G. & NEL A. 2019a. — The oldest genus of the subfamily Sagrinae (Coleoptera: Chrysomelidae) from the Paleocene of Menat (France). *Comptes Rendus Palevol* 18 (2): 178-185. <https://doi.org/10.1016/j.crpv.2018.10.003>
- LEGALOV A. A., KIREJTSHUK A. G. & NEL A. 2019b. — New weevils (Coleoptera, Curculionoidea) from the earliest Eocene Oise amber. *Paleontological Journal* 53 (7): 729-751. <https://doi.org/10.1134/S0031030119070049>
- LEGALOV A. A., NAZARENKO V. YU. & PERKOVSKY E. E. 2019c. — New weevils (Coleoptera: Curculionidae) from Rovno amber. *Paleontological Journal* 53 (10): 1045-1059. <https://doi.org/10.1134/S0031030119100101>
- LIN Q.-B. & LEE C. M. 1997. — Early Tertiary insects from Peng Chau, Hong Kong. *Palaeontology and Stratigraphy of Hong Kong* 1: 185-199.
- LYAL C. H. C. 2014. — 3.7.7. Molytinae Schoenherr, 1823, in LESCHEN R. A. B. & BEUTEL R. G. (eds), *Handbook of Zoology. Arthropoda: Insecta. Tb. 40: Coleoptera (Beetles). Vol. 3: Morphology and Systematics (Phytophaga)*. De Gruyter, Berlin and Boston: 529-569.
- NABOZHENKO M. N. & KIREJTSHUK A. G. 2014. — *Cryptohelops menaticus* – a new genus and species of the tribe Helopini (Coleoptera: Tenebrionidae). *Comptes Rendus Palevol* 13 (2): 65-71. <https://doi.org/10.1016/j.crpv.2013.09.005>
- NABOZHENKO M. V. & KIREJTSHUK A. G. 2017. — The oldest opatrine terrestrial darkling beetle (Coleoptera: Tenebrionidae: Tenebrioninae) from the Paleocene of Menat (France). *Paläontologische Zeitschrift* 91: 307-313. <https://doi.org/10.1007/s12542-017-0368-2>
- NEL A. 1989. — Les Gyrinidae fossiles de France (Coleoptera). *Annales de la Société Entomologique de France (Nouvelle série)* 25 (3): 321-330.
- NEL A. 2008. — The oldest bee fly in the French Paleocene (Diptera: Bombyliidae). *Comptes Rendus Palevol* 7 (7): 401-405. <https://doi.org/10.1016/j.crpv.2008.08.001>
- PITON L. E. 1940. — *Paléontologie du gisement éocène de Menat (Puy-de-Dôme), flore et faune*. Mémoires de la Société d'Histoire Naturelle d'Auvergne, Clermont-Ferrand 1: 1-303.
- POINAR G. JR. & LEGALOV A. A. 2015. — New Apioninae (Coleoptera: Brentidae) in Dominican amber. *Historical Biology* 27 (2): 134-157. <https://doi.org/10.1080/08912963.2013.871010>
- SCUDDER S. H. 1876. — Fossil Coleoptera from the Rocky mountain tertiaries. *Bulletin of the United States Geological and Geographical Survey of the Territories* 2: 77-87.
- SCUDDER S. H. 1893. — Rhynchophorus coleoptera of the United States. *Monographs of the United States Geological Survey* 21: 1-206.
- SCUDDER S. H. 1895. — Canadian fossil insects. Vol. 2. The Coleoptera hitherto found fossil in Canada. *Contributions to Canadian Palaeontology by Geological Survey of Canada* 2 (1): 1-57.
- VINCENT P. M., AUBERT M., BOIVIN P., CANTAGREL J. M. & LENAT J. F. 1977. — Découverte d'un volcanisme paléocène en Auvergne: les maars de Menat et leurs annexes; étude géologique et géophysique. *Bulletin de la Société géologique de France* S7-19 (5): 1057-1070. <https://doi.org/10.2113/gssgfbull.S7-XIX.5.1057>
- WANG L., ZHAO L., TIAN M., WU F. & ZHANG X. 2019. — Late Jurassic to Early Cretaceous volcanism of Hong Kong: Insights from the Ping Chau Formation. *Geoscience Frontiers* 10 (2): 553-568. <https://doi.org/10.1016/j.gsf.2018.03.012>
- WAPPLER T., CURRANO E. D., WILF P., RUST J. & LABANDEIRA C. C. 2009. — No post-Cretaceous ecosystem depression in European forests? Rich insect feeding damage on diverse Middle Palaeocene plants, Menat, France. *Proceedings of the Royal Society B* 276 (1677): 4271-4277. <https://doi.org/10.1098/rspb.2009.1255>
- WEDMANN S., UHL D., LEHMANN T., GARROUSTE R., NEL A., GOMEZ B., SMITH K. & SCHAAL S. F. K. 2018. — The Konservat-Lagerstätte Menat (Paleocene; France) – an overview and new insights. *Geologica Acta* 16 (2): 189-213. <https://doi.org/10.1344/GeologicaActa2018.16.2.5>
- WEN J., NIE Z.-L. & ICKERT-BOND S. M. 2016. — Intercontinental disjunctions between eastern Asia and western North America in vascular plants highlight the biogeographic importance of the Bering land bridge from late Cretaceous to Neogene. *Journal of Systematics and Evolution* 54 (5): 469-490. <https://doi.org/10.1111/jse.12222>
- WICKHAM H. F. 1912. — On some fossil Rhynchophorus Coleoptera from Florissant Colorado. *Bulletin of the American Museum of Natural History* 31 (4): 41-55, I-IV. <http://hdl.handle.net/2246/1716>
- WILLIAMS M. Y. 1943. — The stratigraphy and palaeontology of Hong Kong and the new territories. *Transactions of the Royal Society of Canada* 3: 93-116.
- YUNAKOV N. N. & KIREJTSHUK A. G. 2011. — New genus and species of broad-nosed weevils from Baltic amber and notes on fossils of the subfamily Entiminae (Coleoptera, Curculionidae). *ZooKeys* 160: 73-96. <https://doi.org/10.3897/zookeys.160.2108>
- ZHERIKHIN V. V. 1971. — On weevils (Insecta, Coleoptera) from the Baltic amber. *Trudy Paleontologicheskogo Instituta Akademii Nauk SSSR* 130: 197-209.

Submitted on 30 July 2020;
accepted on 15 December 2020;
published on 4 April 2022.