

Two new species and records of *Neoperla* (Plecoptera, Perlidae) from Yunnan, China

Yingying Wang¹, Wenliang Li¹, Weihai Li²

1 College of Horticulture and Plant Protection, Henan University of Science and Technology, Luoyang, Henan 471023, China **2** Department of Plant Protection, Henan Institute of Science and Technology, Xinxiang, Henan 453003, China

Corresponding authors: Wenliang Li (wenliangli@haust.edu.cn), Weihai Li (lwh7969@163.com)

Academic editor: Sven Bradler | Received 16 November 2021 | Accepted 11 March 2022 | Published 4 April 2022

<http://zoobank.org/C5EABBE8-DEBF-4643-8321-C4334AE18987>

Citation: Wang Y, Li W, Li W (2022) Two new species and records of *Neoperla* (Plecoptera, Perlidae) from Yunnan, China. ZooKeys 1092: 1–18. <https://doi.org/10.3897/zookeys.1092.78069>

Abstract

Two new species of the stonefly genus *Neoperla*, *N. gaoligongshana* **sp. nov.** and *N. hajeki* **sp. nov.** are described from Yunnan Province of southwestern China based on the morphological and distributional data, and the new species are compared with two congeners, *N. wuzhishana* Chen & Du, 2016 and *N. orissa* Stark & Sivec, 2015. *Neoperla hubleyi* Stark & Sivec, 2008 is recorded from Yunnan Province for the first time.

Keywords

New record, new species, Perlinae, stoneflies, Yunnan

Introduction

Neoperla Needham, 1905 is the largest genus in Perlinae, with more than 270 species worldwide. In China, there are at least 110 valid species described by Chen and Du (2015, 2016a, 2016b, 2016c, 2017, 2018), Chu (1929), Claassen (1940), Dewalt et al. (2021), Du (1998, 1999, 2000a, 2000b), Du et al. (1999, 2001), Du and Sivec (2004, 2005), Du and Wang (2005, 2007), Huo et al. (2021), Kong et al. (2014), Kong and Li (2016), Li et al. (2011a, 2011b, 2012a, 2012b, 2013a, 2013b, 2014a, 2014b, 2017, 2019, 2020), Li et al. (2021), Li and Li (2013a, 2013b), Li and Liu (2019), Li and Wang (2011), Li and Zhang (2014), Mo et al. (2019, 2020a, 2020b, 2021a, 2021b), Murányi et al. (2015), Needham (1905), Qin et al. (2013), Sivec et

al. (1988), Sivec and Zwick (1987), Stark and Sivec (2008, 2015), Wang et al. (2013a, 2013b, 2014), Wu (1935, 1938, 1948, 1962, 1973), Wu and Claassen (1934), Yang et al. (2017), and Yang and Yang (1990, 1991, 1992, 1993, 1995a, 1995b, 1996, 1998), Yang and Li (2018), and Zwick (1973, 1977, 1983, 1986). Eight species of *Neoperla* are known from Yunnan Province: *N. cavaleriei* Navás, 1922, *N. diehli* Sivec, 1985, *N. libuae* Li & Murányi, 2014, *N. limbatella* Navás, 1933, *N. lui* Du, 2004, *N. yanlii* Li & Wang, 2014, *N. yunnana* Li & Wang, 2014, and *N. obscurolulva* Wu, 1962. Yunnan Province is located in southwestern China. The region is adjacent to Guizhou and Guangxi in the east, Tibet to the northwest, Myanmar to the west, Sichuan to the north, and Laos and Vietnam to the south. The total area of Yunnan Province is 3941 million square kilometers.

We recently examined a collection of stonefly specimens received from the National Museum Prague of the Czech Republic, and some of the results of our investigations into this material have been already published (Mo et al. 2019, 2020b, 2021a, 2021b; Li and Kong 2020; Li et al. 2019, 2020). Herein, we report our results on the *Neoperla* from Yunnan Province in this collection, including two new species and one new record from the province.

Materials and methods

Types are kept in the Insect Collection of Henan Institute of Science and Technology (HIST), Xinxiang of China and the National Museum Prague (NMP) of the Czech Republic, and the Collection of Smaller Insect Orders, Department of Zoology, Hungarian Natural History Museum (HNHM), Budapest, as indicated in the text. Specimens used in this study were collected using a light trap and stored in 75% ethanol. Specimens were examined with the aid of a Leica M420 dissecting microscope and the color photographs were taken with a Leica S8APO stereo microscope. Aedeagi were everted using the cold-maceration technique of Zwick (1983). The morphological terminology follows that of Stark and Sivec (2008). The map (Fig. 7) was prepared using a base map of Yunnan Province downloaded from DataV.GeoAtlas (Alibaba, China) and ACME Mapper 2.2 (<http://datav.aliyun.com/tools/atlas/index.html>; <http://mapper.acme.com/>).

Results and discussion

Neoperla gaoligongshana sp. nov.

<http://zoobank.org/69D93D93-861F-4F64-98BC-0F01F862F255>

Figs 1, 2

Material examined. Holotype: male (NMP), China: Yunnan Province, Baoshan City, Gaoligongshan National Nature Reserve, Baihualing Village, 6–8.VII.2016, 25°17.7'N, 98°48.1'E, 1535 m, light trap, leg. J. Hájek & J. Růžička.

Diagnosis. Males of this species are characterized by having the tergum 7 with a triangular process with sparse spines apically. The aedeagal tube is straight, and aedeagal sac is armed with spines.

Description. Adult habitus (Fig. 1). Body color brown. Head mostly yellowish brown, with a black marking covering ocellar triangle, the marking extended forward to pale M-line and getting brown, and a narrow triangular marking occurring forward of M-line. Head approximately as wide as the pronotum; compound eyes black; antenna and palpi yellow. Distance between ocelli slightly wider than the diameter of an ocellus. Pronotum disc brown, midline darker, margins pale (Fig. 1B, C). Wings subhyaline, veins brown; legs pale to yellowish brown, femorotibial joint dark brown. Abdomen brownish, cerci brownish (Fig. 1B, C).

Male (Figs 1, 2). Forewing length ca 10.9 mm, hindwing length ca 9.7 mm. Tergum 6 unmodified. Process of tergum 7 triangular, apex rounded and margined by sparse spines (Fig. 1B, D). Tergum 8 with a median sensilla basiconica patch on a trapezoidal sclerite. Tergum 9 with two lateral patches of several sensilla basiconica and long hairs. Hemitergal processes slender, straight in lateral aspect, apex obtuse (Fig. 1B, D).

Aedeagus (Fig. 2). Aedeagal tube weakly sclerotized ventrally and dorsally sclerotized strongly, apex with two ventral spinous lobes (Fig. 2A, B). Sac nearly straight, ca 2× as long as the tube. Spinose apex of sac slender, slightly ventrally curved, with an apical dorsolateral patch of black spines and a subapical ventral patch of spines (Fig. 2A, C); two wide rows of numerous smaller spines covering most of the dorsal surface of the sac, basal half of spinous rows interrupted medially (Fig. 2A, C).

Female. Unknown.

Etymology. The specific name refers to the Gaoligongshan National Nature Reserve in Yunnan Province, where the type locality is situated.

Distribution. China (Yunnan).

Ecology. Gaoligongshan National Nature Reserve is located in northwestern Yunnan Province and is the largest nature reserve of the province. At the same locality, accompanying stoneflies were *Neoperla hajeki* sp. nov. and *Tyloperla illiesi* Stark & Sivec, 2005.

Remarks. The new species is a member of the *N. montivaga* group. The aedeagal tube and terga 7–10 of the new species are similar to those of *Neoperla wuzhishana* Chen & Du, 2016, but *N. wuzhishana* can be distinguished from the new species primarily by the shape of the aedeagal sac and aedeagal armatures (Fig. 8C). In *N. wuzhishana*, the aedeagal sac is distinctly curved and expanded apically (Fig. 8C, present study) (which is obscure in the original drawing because of an apical damage in the type) and has at least four large dorsal spines subapically (see figs 7, 8 in Chen, Du 2016b); however, the aedeagal sac of *N. gaoligongshana* is nearly straight and the apex is constricted and has only small spines subapically on its dorsal side. In addition, the aedeagal sac of *N. wuzhishana* bears a dozen moderately long spines at mid-length which are absent in *N. gaoligongshana*. Besides, the color pattern including that of the head and legs of *N. wuzhishana* and *N. gaoligongshana* is different: *N. wuzhishana* has brown legs and the head is pale with a small dark spot between the posterior ocelli, while in *N. gaoligongshana*, the legs are pale to yellowish brown, the femorotibial joint is dark brown, and the head has a large, black marking covering the ocellar triangle, with this marking extending anterolaterally under the pale M-line (Fig. 1B).




Figure 1. *Neoperla gaoligongshana* sp. nov. (male) **A** head and pronotum, dorsal view **B** adult habitus, dorsal view **C** terminalia, dorsal view **D** terminalia, lateral view. Scale bars: 1 mm.



Figure 2. *Neoperla gaoligongshana* sp. nov. (male) **A** aedeagus with everted sac, dorsal view **B** aedeagus with everted sac, ventral view **C** aedeagus with everted sac, dorsolateral view. Scale bars: 1 mm.

We examined specimens of *N. wuzhishana* from Yinggeling, Hainan Province, and found slight intraspecific variations of head pattern and armatures of the aedeagal sac: the dark area between ocelli in males was slightly smaller than in females, which in both sexes are quite small (Fig. 8A, B); four large spines in types (both in the original

illustrations and descriptions) (see figs 7, 8 in Chen, Du 2016b), which can number seven to nine in our specimens (Fig. 8C, D). Therefore, the absence of large aedeagal spines in *N. gaoligongshana* is regarded as a distinguishing character, separating it from *N. wuzhishana*.

***Neoperla hajeki* sp. nov.**

<http://zoobank.org/2E0FD484-BE6F-4350-9C62-30AB5B651779>

Figs 3–5

Type material. *Holotype*: male (NMP), China: Yunnan Province, Baoshan City, Zizhi Village, 29.VI–2.VII.2016, 25°43.7'N, 98°34.1'E, 1995 m, light trap, leg. J. Hájek and J. Růžička. *Paratypes*: 9 females (NMP), 1 male and 3 females (HIST), 1 male and 2 females (HNHM), same date as holotype.

Diagnosis. This species is characterized by a small dark marking over the ocellar area and a dark brown stigma before the M-line. The male of new species is characterized by an S-shaped aedeagal tube and by a sac bearing a subapical triangular patch of spinules in dorsal aspect.

Description. Adult habitus (Figs 3, 5). General body color brown. Head general pale brown, a small dark marking covers ocellar area, with a dark brown stigma before M-line. Head slightly wider than pronotum; compound eyes blackish, antenna dark brown except the basal segment yellowish (Figs 3A, 5A), palpi pale brown. Distance between ocelli narrower than the diameter of ocellus and a small marking between ocelli dark brown. Pronotum yellow, with rugosities and a strip-like midline (Figs 3A, 5A). Wings hyaline, veins brown; legs brown, basal part of femur dark brown, with wider yellow bands in mid- and hind legs (Figs 3B, 5B). Cerci yellowish (Figs 3D, 5B).

Male (Figs 3, 4). Forewing length 9.1–9.3 mm, hind wing length 7.9–8.1 mm. Process of tergum 7 sclerotized and triangular, with a nipple-like apex and covered with small sensilla basiconica (Fig. 3B, D). Tergum 8 with a median plump mound and a few sensilla basiconica, mostly covered by the process of tergum 7. Tergum 9 with two separated mesal projections covered by sensilla basiconica and hairs. Hemitergal processes of tergum 10 finger-like in shape, slightly curved medially in dorsal aspect, apex acute (Fig. 3B, D).

Aedeagus. Aedeagal tube sclerotized, slightly S-shaped in lateral aspect. Sac short, ca 1/3 as long as the tube (Fig. 4). Aedeagal sac bearing a subapical triangular patch of spinules in dorsal aspect (Fig. 4A), remainder bald except a few ventral spinules (Fig. 4B, C).

Female (Fig. 5). Forewing length 9.3–9.6 mm, hind wing length 8.2–8.4 mm. General color pattern is similar to males. The subgenital plate of sternum 8 is not produced posteriorly. Vagina large and apically slender, spiral and incurved, apical round, full of scaly spots. Spermatheca small, its origin in the terminal vagina.

Egg (Fig. 6). Chorion length 346–348 μm , width 173–177 μm . Micropyles 3 with rims, placed ca 1/3 length near operculum, each located between striae (Fig. 6A, F). Collar not distinctly stalked, but slightly constricted at base; width ca 79.5 μm at collar




Figure 3. *Neoperla hajeki* sp. nov. (male) **A** head and pronotum, dorsal view **B** adult habitus, dorsal view **C** terminalia, dorsal view **D** terminalia, lateral view. Scale bars: 1 mm.



Figure 4. *Neoperla hajeki* sp. nov. (male) **A** aedeagus with everted sac, dorsal view **B** aedeagus with everted sac, ventral view **C** aedeagus with everted sac, lateral view. Scale bars: 1 mm.




Figure 5. *Neoperla bajeki* sp. nov. (male) **A** head and pronotum, dorsal view **B** terminalia, dorsal view **C** vagina, dorsal view **D** vagina, lateral view. Scale bars: 1 mm (**A–B**); 0.2 mm (**C–D**).

(Fig. 6D). Collar short, irregular in shape (Fig. 6D). Rim slightly flanged, margin irregularly scalloped. Chorion with striate on lid and collar. FCIs on lid distinct; cells with thin, smooth walls and floors punctuated (Fig. 6B, E).

Etymology. The species is named after Dr Jiri Hájek for collecting the specimens.

Distribution. China (Yunnan).

Ecology. See ecology of *Neoperla gaoligongshana* sp. nov.

Remarks. The new species is a member of the *N. clymene* group. Color pattern, pronotum, and male genital features are similar to *Neoperla orissa* Stark & Sivec, 2015 from India. The new species can be easily separated from *N. orissa* by the projection of tergum 7 and detail of the aedeagal sac armature. In *N. hajeki*, the projection of tergum 7 is pointed in dorsal aspect, and the dorsal spines of the aedeagal sac are triangularly arranged. In *N. orissa*, the slender and median projection of tergum 7 appears truncate in dorsal aspect, and the spines of the sac are arranged in several close-set rows.

Neoperla libuae Li & Murányi, 2014

Neoperla libuae Li and Murányi 2014: 2 (original description).

Material examined. 1 male and 3 females (NMP), 1 male and 1 female (HIST), China: Yunnan Province, Yingjiang County, Dehong Dai and Jingpo Autonomous Prefecture, Tongbiguan Town, at light in village near river, 24–26.VI.2016, 24°36.7'N, 97°39.4'E, 1340 m, leg. J. Hájek & J. Růžička.

Distribution. China: Yunnan.

Ecology. Tongbiguan Town is located in southwestern Yingjiang County, and it also belongs to the Gaoligongshan region. At the same locality, accompanying stoneflies were *Neoperla hubleyi* Stark & Sivec, 2008 and one unidentified female *Neoperla* sp.

Remarks. *Neoperla libuae* Li & Murányi, 2014 was originally described by Li et al. (2014a) from Xishuangbanna in Yunnan Province. Our specimens from Yingjiang County agree well with original description of the head pattern and terminalia, the aedeagal tube and sac.

Neoperla hubleyi Stark & Sivec, 2008

Neoperla hubleyi Stark and Sivec 2008: 30 (original description); Mo et al. 2020: 521 (new record for Guangxi Zhuang Autonomous Region).

Material examined. 1 male (HIST), China: Yunnan Province, Dehong Dai and Jingpo Autonomous Prefecture, Yingjiang County, Tongbiguan Town, at light in village near river, 24–26.VI.2016, 24°36.7'N, 97°39.4'E, 1340 m, leg. J. Hájek & J. Růžička.

Distribution. China: Yunnan and Guangxi Zhuang Autonomous Region; Vietnam.




Figure 6. *Neoperla hajeki* sp. nov. (egg) **A** egg **B** lid **C** collar **D** collar and lid **E** FCIs and lid **F** micropyles.




Figure 7. Distribution of *Neoperla* Needham, 1905 in cities of Yunnan. Yellow dots indicate location of the two new species, green dot indicates two new records.

Ecology. See ecology of *Neoperla lihuuae* Li & Murányi, 2014.

Remarks. Stark and Sivec (2008) originally described this species from Vietnam, and Mo et al. (2020a) described a new record in Guangxi Zhuang Autonomous Region. The terminalia and aedeagus of our specimen fit the original description of Stark and Sivec (2008).

Concluding remarks

Neoperla was divided into the *Neoperla clymene* group and the *Neoperla montivaga* group by Zwick (1983). The *N. clymene* species group currently includes more than 150 species, and the *N. montivaga* species group includes over 123 species worldwide (Huo et al. 2021; Mo et al. 2021a).

So far 11 species of *Neoperla* have been recorded from Yunnan Province, including the two new species and one species newly recorded in this paper. Among the six




Figure 8. *Neoperla wuzhishana* Chen & Du, 2016 (**A, C–E** male **B** female), all from Yinggeling, Hainan Province **A** head and pronotum, dorsal view **B** head and pronotum, dorsal view **C** Aedeagus, lateral view **D** Large spines of aedeagal sac, ventral view **E** legs, dorsal view. Scale bars: 1 mm (**A–C, E**); 0.1 mm (**D**).

endemic species of *Neoperla*, including the two new species listed below—*N. libuae*, *N. yanlii*, *N. obscurolulva*, *N. yunnana*, *N. hájeki*, and *N. gaoligongshana*—three of them are distributed in Baoshan City. Five species are widely distributed: *N. cavaleriei*, *N. diehli*, *N. hubleyi*, *N. limbatella*, and *N. lui*. *Neoperla binodosa* Wu, 1973 was transferred to the genus *Phanoperla* and is placed as a synonym of *P. pallipennis* Banks, 1938 by Mo et al. (2021c). Most species are distributed in western and southeastern Yunnan Province and central and northern Yunnan still needs to be surveyed (Fig. 7).

Acknowledgements

We are indebted to Dr J. Hájek and Dr J. Růžicka for collecting specimens and Ms Ruijun Liu for her assistance in SEM studies. We acknowledge the support of the Graduate Education Innovation Training Base Project of Henan Province in 2021 (no. 107020221005), the National Natural Science Foundation of China (no. 31970402), and the Program for Science & Technology Innovation Talents in Universities of Henan Province (no. 21HASTIT042).

References

- Chen ZT, Du YZ (2015) The first record of a species of Perlidae from Jiangsu Province, China: a new species of *Neoperla* (Plecoptera: Perlidae). *Zootaxa* 3974(3): 424–430. <https://doi.org/10.11646/zootaxa.3974.3.9>
- Chen ZT, Du YZ (2016a) A new species of *Neoperla* (Plecoptera: Perlidae) from Zhejiang Province of China. *Zootaxa* 4093(4): 589–594. <https://doi.org/10.11646/zootaxa.4093.4.12>
- Chen ZT, Du YZ (2016b) Contribution to the knowledge of *Neoperla* (Plecoptera: Perlidae) from Hainan Province of China. *Zootaxa* 4144(2): 243–252. <https://doi.org/10.11646/zootaxa.4144.2.6>
- Chen ZT, Du YZ (2016c) A new species of *Neoperla* from China, with a redescription of the female of *N. mnong* Stark, 1987 (Plecoptera, Perlidae). *ZooKeys* 616: 103–113. <https://doi.org/10.3897/zookeys.629.10167>
- Chen ZT, Du YZ (2017) Description of *Neoperla yingshana* sp. nov. (Plecoptera: Perlidae) from Hubei Province of China. *Zootaxa* 4362(2): 287–293. <https://doi.org/10.11646/zootaxa.4362.2.8>
- Chen ZT, Du YZ (2018) A checklist and adult key to the Chinese stonefly (Plecoptera) genera. *Zootaxa* 4375(1): 059–074. <https://doi.org/10.11646/zootaxa.4375.1.2>
- Chu YT (1929) Descriptions of four new species and one new genus of stoneflies in the family Perlidae from Hangchow. *China Journal* (Canberra, A.C.T.) 10: 88–92.
- Claassen PW (1940) A catalogue of the Plecoptera of the world. *Memoirs of the Cornell University Agricultural Experiment Station* 232: 1–235.
- DeWalt RE, Maehr MD, Hopkins H, Neu-Becker U, Stueber G (2021) Plecoptera Species File Online. Version 5.0/5.0. <http://Plecoptera.SpeciesFile.org> [accessed 18 August 2021]
- Du YZ (1998) Two new record species of genus *Neoperla* Needham (Plecoptera: Perlidae: Perlinae) from China. *Journal of Zhejiang Agricultural University* 24: 392–394.

- Du YZ (1999) Plecoptera. In: Huang BK (Ed.) Fauna of Insects in Fujian Province of China. Vol. 3. Fujian Science and Technology Publishing House, Fujian, 301–335.
- Du YZ (2000a) *Neoperla magisterchoui*, a new species of the genus *Neoperla* Needham (Plecoptera: Perlidae) from China. In: Chang YL (Ed.) Systematic and Faunistic Research on Chinese Insects. Proceedings of the 5th National Congress of Insect Taxonomy. China Agriculture Press, Beijing, 1–3.
- Du YZ (2000b) Two new species of the genus *Neoperla* Needham (Plecoptera: Perlidae: Perlinae) from Guizhou, China. *Entomotaxonomia* 22(1): 1–5.
- Du YZ, Sivec I (2004) Plecoptera: Perlidae, Nemouridae, Leuctridae. In: Yang XK (Ed.) Insects from Mt. Shiwandashan Area of Guangxi. China Forestry Publishing House, Beijing, 39–45.
- Du YZ, Sivec I (2005) Plecoptera. In: Yang XK (Ed.) Insect Fauna of Middle-West Qinling Range and South Mountains of Gansu Province. Science Press, Beijing, China, 38–54.
- Du YZ, Wang ZJ (2005) Plecoptera: Leuctridae, Nemouridae, Perlidae and Peltoperlidae. In: Yang MF, Jin DC (Eds) Insects from Dashahe Nature Reserve of Guizhou. Guizhou People Press, Guizhou, 51–57.
- Du YZ, Wang ZJ (2007) Nemouridae and Perlidae. In: Li ZZ, Yang MF, Jin DC (Eds) Insects from Mountain Leigongshan Landscape of Guizhou. Guizhou Science and Technology Publishing House, Guizhou, 84–90.
- Du YZ, Sivec I, He JH (1999) A checklist of the Chinese species of the family Perlidae (Plecoptera: Perloidea). *Acta Entomologica Slovenica* (Ljubljana) 7: 59–67.
- Du YZ, Sivec I, Zhao MS (2001) Plecoptera. In: Wu H, Pan CW (Eds) Insects of Tianmushan National Nature Reserve. Science Press, Beijing, 69–80.
- Huo QB, Du YZ, Zhu BQ, Yu L (2021) Notes on *Neoperla sinensis* Chu, 1928 and *Neoperla anjiensis* Yang Yang, 1998, with descriptions of new species of *Neoperla* from China (Plecoptera: Perlidae). *Zootaxa* 5004(2): 288–310. <https://doi.org/10.11646/zootaxa.5004.2.3>
- Kong FB, Li WH (2016) Type designation for *Neoperla furcomaculata* Kong, Li, 2014 (Plecoptera: Perlidae). *Illiesia* 12(10): e59.
- Kong FB, Lv JF, Li WH (2014) Three new species of *Neoperla* in the *montivaga* group (Plecoptera: Perlidae) from China. *Zootaxa* 3841(3): 429–438. <https://doi.org/10.11646/zootaxa.3841.3.7>
- Li WH, Li XP (2013a) A new species of the genus *Neoperla* (Plecoptera: Perlidae) from China. *Dong Wu Fen Lei Xue Bao* 38: 75–77.
- Li WH, Li XP (2013b) A replacement name for *Neoperla siveci* Li and Li (Plecoptera: Perlidae). *Illiesia* 9(9): e109.
- Li WH, Liu RJ (2019) A new species of *Neoperla* (Plecoptera: Perlidae) from southeastern Tibet of China with additions to the knowledge of *N. perspicillata* Zwick, 1980. *Zootaxa* 4686(2): 271–281. <https://doi.org/10.11646/zootaxa.4686.2.7>
- Li WH, Kong FB (2020) A new species and new record of *Tyloperla* (Plecoptera: Perlidae) from China. *Zootaxa* 4728 (2): 289–294. <https://doi.org/10.11646/zootaxa.4728.2.10>
- Li WH, Wang RF (2011) A new species of *Neoperla* (Plecoptera: Perlidae) from China. *Entomological News* 122(3): 261–264. <https://doi.org/10.3157/021.122.0308>
- Li WH, Zhang SQ (2014) Two new species of *Neoperla* (Plecoptera: Perlidae) from Dabie Mountains of China. *ZooKeys* 438: 45–55. <https://doi.org/10.3897/zookeys.438.8230>

- Li WH, Wang HL, Lu WY (2011a) Species of the genus *Neoperla* (Plecoptera: Perlidae) from Henan, China. *Zootaxa* 2735(1): 57–63. <https://doi.org/10.11646/zootaxa.2735.1.6>
- Li WH, Wu LM, Zhang HR (2011b) A new species of the genus *Neoperla* (Plecoptera: Perlidae) from Henan, China. *Dong Wu Fen Lei Xue Bao* 36: 33–35.
- Li WH, Wang GQ, Lu WY (2012a) Species of *Neoperla* (Plecoptera: Perlidae) from Hubei, China. *Zootaxa* 3478(1): 32–37. <https://doi.org/10.11646/zootaxa.3478.1.5>
- Li WH, Wang GQ, Li WL, Murányi D (2012b) Review of *Neoperla* (Plecoptera: Perlidae) from Guangdong Province of China. *Zootaxa* 3597(1): 15–24. <https://doi.org/10.11646/zootaxa.3597.1.2>
- Li WH, Liang HY, Li WL (2013a) Review of *Neoperla* (Plecoptera: Perlidae) from Zhejiang Province, China. *Zootaxa* 3652(3): 353–369. <https://doi.org/10.11646/zootaxa.3652.3.4>
- Li WH, Wang GQ, Qin XF (2013b) Two new species of *Neoperla* (Plecoptera: Perlidae) from China. *ZooKeys* 290: 21–30. <https://doi.org/10.3897/zookeys.290.4568>
- Li WH, Murányi D, Wang RF (2014a) Species of *Neoperla* (Plecoptera: Perlidae) from Yunnan Province, China. *Zootaxa* 3753(1): 1–9. <https://doi.org/10.11646/zootaxa.3753.1.1>
- Li WH, Li S, Feng GW, Wang YB (2014b) Species of *Neoperla* (Plecoptera: Perlidae) from Henan Province, China. *Zootaxa* 3838(2): 174–182. <https://doi.org/10.11646/zootaxa.3838.2.2>
- Li WH, Wang Y, Wang RF (2017) Taxonomic notes on *Neoperla* (Plecoptera: Perlidae) from Sichuan Province of China with the description of two new species. *Zootaxa* 4221(2): 191–204. <https://doi.org/10.11646/zootaxa.4221.2.3>
- Li WH, Mo RR, Wang GQ (2019) Additions to the genus *Hemacroneuria* Enderlein (Plecoptera: Perlidae) from China. *Zootaxa* 4652(2): 349–358. <https://doi.org/10.11646/zootaxa.4652.2.7>
- Li WH, Mo RR, Wang GQ (2020) Two new species of *Neoperla montivaga* group (Plecoptera: Perlidae) from Guangxi Zhuang Autonomous Region of southern China. *Zootaxa* 4718(4): 573–583. <https://doi.org/10.11646/zootaxa.4718.4.10>
- Li WL, Wang YY, Li WH (2021) A new species of *Neoperla* (Plecoptera: Perlidae) from Motuo County of Tibet, China and redescription of *Neoperla perspicillata* Zwick, 1980. *Zootaxa* 4964(1): 169–178. <https://doi.org/10.11646/zootaxa.4964.1.9>
- Mo RR, Yao G, Wang GQ, Li WH (2019) One new species and one new Chinese record of *Neoperla* (Plecoptera: Perlidae) from Mount Shiwandashan of the Guangxi Zhuang Autonomous Region of southern China. *Zootaxa* 4652(3): 497–506. <https://doi.org/10.11646/zootaxa.4652.3.6>
- Mo RR, Wang GQ, Li WH (2020a) A new species and two new records of the *Neoperla diehli* subgroup of the *N. montivaga* group (Plecoptera: Perlidae) from China. *Zootaxa* 4790(3): 515–524. <https://doi.org/10.11646/zootaxa.4790.3.6>
- Mo RR, Liu RJ, Li WH, Shi MW (2020b) Rediscovery of two species of *Neoperla* (Plecoptera: Perlidae) from China. *Zootaxa* 4758(2): 369–377. <https://doi.org/10.11646/zootaxa.4758.2.10>
- Mo RR, Wang GQ, Li WH, Murányi D (2021a) Two new species and one new record of *Neoperla* (Plecoptera, Perlidae) from Guangxi Zhuang Autonomous Region, China. *ZooKeys* 1056: 191–206. <https://doi.org/10.3897/zookeys.1056.61565>
- Mo RR, Wang GQ, Li WH (2021b) One new species and one poorly known species of *Neoperla* in the *montivaga* group (Plecoptera: Perlidae) from Hainan Province, China. *Zootaxa* 5004(2): 385–394. <https://doi.org/10.11646/zootaxa.5004.2.7>

- Mo RR, Wang Y, Wang GQ, Li WH (2021c) Review of *Phanoperla* (Plecoptera, Perlidae) from China. *ZooKeys* 1052: 71–81. <https://doi.org/10.3897/zookeys.1052.64060>
- Murányi D, Li WH, Jeon MJ, Hwang JM, Seo HY (2015) Korean species of the genus *Neoperla* Needham, 1905 (Plecoptera: Perlidae). *Zootaxa* 3918(1): 113–127. <https://doi.org/10.11646/zootaxa.3918.1.5>
- Needham JG (1905) New genera and species of Perlidae. *Proceedings of the Biological Society of Washington* 17: 107–110.
- Qin XF, Murányi D, Wang GQ, Li WH (2013) Stoneflies of the genus *Neoperla* (Plecoptera, Perlidae) from Wuyi Mountain National Nature Reserve, Fujian of China. *ZooKeys* 326: 1–16. <https://doi.org/10.3897/zookeys.326.5911>
- Sivec I, Zwick P (1987) Some *Neoperla* (Plecoptera) from Taiwan. *Beiträge zur Entomologie* 37: 391–405.
- Sivec I, Stark BP, Uchida S (1988) Synopsis of the world genera of Perlinae (Plecoptera: Perlidae). *Scopolia* 16: 1–66.
- Stark BP, Sivec I (2008) New species and records of *Neoperla* (Plecoptera: Perlidae) from Vietnam. *Illiesia* 4(3): 19–54.
- Stark BP, Sivec I (2015) New species and records of *Neoperla* (Plecoptera: Perlidae) from India. *Illiesia* 11(7): 75–91. <http://illiesia.speciesfile.org/papers/Illiesia11-07.pdf>
- Wang GQ, Li WH, Yang J (2013a) New species and records of the stonefly genus *Neoperla* (Plecoptera, Perlidae) from Jinhuacha Nature Reserve, Guangxi of China. *ZooKeys* 351: 37–48. <https://doi.org/10.3897/zookeys.351.6259>
- Wang HL, Wang GQ, Li WH (2013b) Two new species in the subfamily Perlinae (Plecoptera, Perlidae) from China. *ZooKeys* 313: 81–90. <https://doi.org/10.3897/zookeys.313.5460>
- Wang GQ, Murányi D, Li WH (2014) Two new species of dark *Neoperla* (Plecoptera, Perlidae) from Oriental Realm of China. *Zootaxa* 3872(1): 22–30. <https://doi.org/10.11646/zootaxa.3872.1.2>
- Wu CF (1935) Aquatic insects of China. Article XXI. New species of stoneflies from East and South China. (Order Plecoptera). *Peking Natural History Bulletin* 9: 227–243.
- Wu CF (1938) *Plecopterorum sinensium: a Monograph of Stoneflies of China* (Order Plecoptera). Yenching University, Beijing, 225 pp.
- Wu CF (1948) Fourth supplement to the stoneflies of China (order Plecoptera). *Peking Natural History Bulletin* 17: 75–82.
- Wu CF (1962) Results of the Zoologico-Botanical expedition to southwest China, 1955–1957 (Plecoptera). *Acta Entomologica Sinica* 11(Supplement): 139–153.
- Wu CF (1973) New species of Chinese stoneflies (order Plecoptera). *Acta Entomologica Sinica* 16: 97–118.
- Wu CF, Claassen PW (1934) Aquatic insects of China. Article XVIII. New species of Chinese stoneflies. (Order Plecoptera). *Peking Natural History Bulletin* 9: 111–129.
- Yang D, Li WH (2018) *Species Catalogue of China. Volume 2, Animals, Insecta (III), Plecoptera*. Science Press, Beijing, 71 pp.
- Yang CK, Yang D (1990) New and little-known species of Plecoptera from Guizhou Province (I). *Guizhou Science* 8(4): 1–4.
- Yang CK, Yang D (1991) New and little-known species of Plecoptera from Guizhou Province (II). *Guizhou Science* 9: 48–50.

- Yang D, Yang CK (1992) Plecoptera: Perlidae. In: Huang FS (Ed.) Insects of Wuling Mountains Area, Southwestern China. Science Press, Beijing, 62–64.
- Yang D, Yang CK (1993) New and little-known species of Plecoptera from Guizhou Province (III). *Entomotaxonomia* 15: 235–238.
- Yang D, Yang CK (1995a) Three new species of Plecoptera from Hainan Province. *Acta Agriculture Universitatis Pekinensis* 21: 223–225.
- Yang D, Yang CK (1995b) Plecoptera: Perlidae. In: Wu H (Ed.) Insects of Baishanzu Mountain, Eastern China. China Forestry Publishing House, Beijing, 59–60.
- Yang D, Yang CK (1996) Four new species of Plecoptera from Nei Mongol. *Zhongguo Nongye Daxue Xuebao* 1: 115–118.
- Yang D, Yang CK (1998) In: Wu H (Ed.) Insects of Longwangshan. China Forestry Publishing House, Beijing, 40–46.
- Yang J, Zhang SJ, Li WH (2017) Description of two new species of the *Neoperla montivaga* group (Plecoptera: Perlidae) from Guangxi Zhuang Autonomous Region, China. *Zootaxa* 4238(3): 385–394. <https://doi.org/10.11646/zootaxa.4238.3.5>
- Zwick P (1973) Insecta: Plecoptera phylogenetisches System und Katalog. *Das Tierreich* 94: 1–465.
- Zwick P (1977) Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel. *Entomologica Basiliensia* 2: 121–123.
- Zwick P (1983) The *Neoperla* of Sumatra and Java (Indonesia) (Plecoptera: Perlidae). *Spixiana* 6: 167–204.
- Zwick P (1986) The Bornean species of the stonefly genus *Neoperla* (Plecoptera: Perlidae). *Aquatic Insects* 8(1): 1–53. <https://doi.org/10.1080/01650428609361227>